

- (46) FROST, H. Investigations into the pathogenesis of arteriosclerosis: drug prophylaxis. In: Shimamoto, T., Numano, F., Addison, G.M., (Editors). *Atherogenesis, Volume II. Proceedings of the Second International Symposium on Atherogenesis, Thrombogenesis and Pyridinolcarbamate Treatment*, Tokyo, May 18-20, 1972. Amsterdam, Excerpta Medica, 1973, pp. 32-50.
- (47) GARRISON, R.J., KANNEL, W.B., FEINLEIB, M., CASTELLI, W.P., MCNAMARA, P.M., PADGETT, S.J. Cigarette smoking and HDL cholesterol. The Framingham Offspring Study. *Atherosclerosis* 30: 17-25, 1978.
- (48) GOLDBOURT, U., MEDALIE, J.H. Characteristics of smokers, nonsmokers and ex-smokers among 10,000 adult males in Israel. II. Physiologic, biochemical and genetic characteristics. *American Journal of Epidemiology* 105(1): 75-86, 1977.
- (49) GORDON, T., CASTELLI, W.P., HJORTLAND, M.C., KANNEL, W.B., DAWBER, T.R. High density lipoprotein as a protective factor against coronary heart disease. The Framingham study. *American Journal of Medicine* 62: 707-714, May 1977.
- (50) GORDON, T., KANNEL, W.B., MCGEE, D., DAWBER, T.R. Death and coronary attacks in men after giving up cigarette smoking. A report from the Framingham study. *Lancet* 2: 1345-1348, December 7, 1974.
- (51) GRAHAM, I., MULCAHY, R., HICKEY, N., SYNNOTT, M. Mode of death related to smoking in patients with coronary heart disease. *Journal of the Irish Medical Association* 70(7): 234-235, May 14, 1977.
- (52) GREENSPAN, K., EDMANDS, R.E., KNOEBEL, S.B., FISCH, C. Some effects of nicotine on cardiac automaticity, conduction, and inotrophy. *Archives of Internal Medicine* 123(6): 707-712, June 1969.
- (53) HAMMOND, E.C., GARFINKEL, L., SEIDMAN, H., LEW, E.A. "Tar" and nicotine content of cigarette smoke in relation to death rates. *Environmental Research* 12(3): 263-274, December 1976.
- (54) HARKAVY, J. Tobacco allergy in cardiovascular disease: A review. *Annals of Allergy* 26(8): 447-459, August 1968.
- (55) HAUST, M.D., MORE, R.H. Spontaneous lesions of the aorta in the rabbit. In Roberts, J.C., Jr., Straus, R. (Editors). *Comparative Atherosclerosis. The Morphology of Spontaneous and Induced Atherosclerotic Lesions in Animals and Its Relation to Human Disease*. New York, Harper and Row, 1965, pp. 255-275.
- (56) HAWKINS, L.H. Blood carbon monoxide levels as a function of daily cigarette consumption and physical activity. *British Journal of Industrial Medicine* 33(2): 123-125, May 1976.
- (57) HAYES, M.J., MORRIS, G.K., HAMPTON, J.R. Lack of effect of bed rest and cigarette smoking on development of deep venous thrombosis after myocardial infarction. *British Heart Journal* 38(9): 981-983, September 1976.
- (58) HOLLANDER, W., PRUSTY, S., KIRKPATRICK, B., PADDOCK, J. NAGRAJ S. Role of hypertension in ischemic heart disease and cerebral vascular disease in the cynomolgus monkey with coarctation of the aorta. *Circulation Research* 40(5, Supplement 1): I-70-I-83, May 1977.
- (59) HULLEY, S.B., COHEN, R., WIDDOWSON, G. Plasma high-density lipoprotein cholesterol level. Influence of risk factor intervention. *Journal of the American Medical Association* 238(21): 2269-2271, November 21, 1977.
- (60) JAIN, A.C., BOWYER, A.F., MARSHALL, R.J., ASATO, H. Left ventricular function after cigarette smoking by chronic smokers: Comparison of normal subjects and patients with coronary artery disease. *American Journal of Cardiology* 39(1): 27-31, January 1977.
- (61) JAIN, A.K. Cigarette smoking, use of oral contraceptives, and myocardial infarction. *American Journal of Obstetrics and Gynecology* 126(3): 301-307, October 1, 1976.

- (62) JANZON, L. Smoking cessation and peripheral circulation. A population study in 59-year-old men with plethysmography and segmental measurements of systolic blood pressure. *VASA* 4(3): 282-287, 1975.
- (63) JENKINS, C.D., ROSENMAN, R.H., ZYZANSKI, S.J. Cigarette smoking. Its relationship to coronary heart disease and related risk factors in the Western Collaborative Group Study. *Circulation* 38(6): 1140-1155, December 1968.
- (64) JENKINS, C.D., ZYZANSKI, S.J., ROSENMAN, R.H. Risk of new myocardial infarction in middle-aged men with manifest coronary heart disease. *Circulation* 53(2): 342-347, February 1976.
- (65) JENNINGS, R.B. Relationship of acute ischemia to functional defects and irreversibility. *Circulation* 53(3, Supplement 1): I-26—I-29, March 1976.
- (66) KAGAN, A.R., STERNBY, N.H., UEMURA, K., VANECEK, R., VIHERT, A.M., LIFSIC, A.M., MATOVA, E.E., ZAHOR, Z., ZDANOV, V.S. Atherosclerosis of the aorta and coronary arteries in five towns. *Bulletin of the World Health Organization* 53(5-6): 485-645, 1976.
- (67) KAHN, H.A., MEDALIE, J.H., NEUFELD, H.N., RISS, E., GOLDBOURT, U. The incidence of hypertension and associated factors: The Israel ischemic heart disease study. *American Heart Journal* 84(2): 171-182, August 1972.
- (68) KANNEL, W.B. Epidemiologic studies on smoking in cerebral and peripheral vascular disease. In: Wynder, E.L., Hoffmann, D., Gori, G.B., (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume I. *Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No. (NIH) 76-1221, 1976, pp. 257-274.
- (69) KANNEL, W.B. Some lessons in cardiovascular epidemiology from Framingham. *American Journal of Cardiology* 37: 269-282, February 1976.
- (70) KANNEL, W.B., CASTELLI, W.P. Significance of nicotine, carbon monoxide and other smoke components in the development of cardiovascular disease. U.S. Public Health Service, DHEW Publication No. (NIH) 76-1221, 1976, pp. 369-381.
- (71) KANNEL, W.B., DOYLE, J.T., MCNAMARA, P.M., QUICKENTON, P., GORDON, T. Precursors of sudden coronary death. Factors related to the incidence of sudden death. *Circulation* 51: 606-613, April 1975.
- (72) KJELDSEN, K., ASTRUP, P., WANSTRUP, J. Reversal of rabbit atheromatosis by hyperoxia. *Journal of Atherosclerosis Research* 10: 173-178, 1969.
- (73) KJELDSEN, K., THOMSEN, H.K. The effect of hypoxia on the fine structure of the aortic intima in rabbits. *Laboratory Investigation* 33(5): 533-543, 1975.
- (74) KJELDSEN, K., WANSTRUP, J., ASTRUP, P. Enhancing influence of arterial hypoxia on the development of atheromatosis in cholesterol-fed rabbits. *Journal of Atherosclerosis Research* 8: 835-845, 1968.
- (75) KOCH, A. Smoking and peripheral arterial disease. In: Wynder, E.L. Hoffmann, D., Gori, G.B. (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume I. *Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No. (NIH) 76-1221, 1976, pp. 281-283.
- (76) LAWSON, D.H., DAVIDSON, J.F., JICK, H. Oral contraceptive use and venous thromboembolism: Absence of an effect of smoking. *British Medical Journal* 2: 729-730, September 17, 1977.
- (77) LAWTON, G. Cigarette consumption and atherosclerosis. Their relationship in the aortic and iliac and femoral arteries. *British Journal of Surgery* 60(11): 873-876, November 1973.

- (78) LEVINE, P.H. An acute effect of cigarette smoking on platelet function. A possible link between smoking and arterial thrombosis. *Circulation* 48: 619-623, September 1973.
- (79) LIFSIC, A.M. Atherosclerosis in smokers. *Bulletin of the World Health Organization* 53(5-6): 631-638, 1976.
- (80) MALINOW, M.R., MCLAUGHLIN, P., DHINDSA, D.S., METCALFE, J., OCHSNER, A.J., III, HILL, J., MCNULTY, W.P. Failure of carbon monoxide to induce myocardial infarction in cholesterol-fed cynomolgus monkeys (*Macaca fascicularis*). *Cardiovascular Research* 10: 101-108, 1976.
- (81) MANN, J.I., INMAN, W.H.W., THOROGOOD, M. Oral contraceptive use in older women and fatal myocardial infarction. *British Medical Journal* 2: 445-447, August 21, 1976.
- (82) MANN, J.I., VESSEY, M.P., THOROGOOD, M., DOLL, R. Myocardial infarction in young women with special reference to oral contraceptive practice. *British Medical Journal* 2: 241-245, May 3, 1975.
- (83) MARKS, P., EMERSON, P.A. Increased incidence of deep vein thrombosis after myocardial infarction in nonsmokers. *British Medical Journal* 3(5925): 232-234, July 27, 1974.
- (84) MCGILL, H.C., JR. Atherosclerosis: Problems in pathogenesis. In: Paoletti, R., Gotto, A.M., Jr. (Editors). *Atherosclerosis Reviews*, Volume 2. New York, Raven Press, 1977, pp. 27-65.
- (85) MCGILL, H.C., JR., (Editor). General findings of the International Atherosclerosis Project. *Laboratory Investigation* 18(5): 498-502, May 1968.
- (86) MCGILL, H.C., JR., ROGERS, W.R., WILBUR, R.L., JOHNSON, D.E. Cigarette smoking baboon model: Demonstration of feasibility (40119). *Proceedings of the Society for Experimental Biology and Medicine* 157: 672-676, 1978.
- (87) MCKUSICK, J.A., HARRIS, W.S., OTTESEN, O.E., GOODMAN, R.M., SHELLEY, W.M., BLOODWELL, R.D. Buerger's disease: A distinct clinical and pathologic entity. *Journal of the American Medical Association* 181(1): 5-12, July 7, 1962.
- (88) MCMAHAN, C.A., RICHARDS, M.L., STRONG, J.P. Individual cigarette usage: Self-reported data as a function of respondent-reported data. *Atherosclerosis* 23(3): 477-488, May/June 1976.
- (89) MCMILLAN, G.C. Development of Arteriosclerosis. *The American Journal of Cardiology* 31: 542-546, May 1973.
- (90) MCMILLAN, G.C. Evidence for components other than carbon monoxide and nicotine as etiological factors in cardiovascular disease. In: Wynder, E.L., Hoffmann, D., Gori, G.B. (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume I. *Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No. (NIH) 76-1221, 1976, pp. 363-367.
- (91) MEDALIE, J.H., SNYDER, M., GROEN, J.J., NEUFELD, H.N., GOLDBOURT, U., RISS, E. Angina pectoris among 10,000 men. 5 year incidence and univariate analysis. *American Journal of Medicine* 55: 583-594, November 1973.
- (92) MILLER, N.E., FORDE, O.H., THELLE, D.S., MJOS, O.D. The Tromso Heart Study. High-density lipoprotein and coronary heart disease: A prospective case-control study. *Lancet* 1 (8019): 965-968, May 7, 1977.
- (93) MINICK, C.R., MURPHY, G.E. Experimental induction of atheroarteriosclerosis by the synergy of allergic injury to arteries and lipid-rich diet. II. Effect of a repeatedly injected foreign protein in rabbits fed a lipid-rich, cholesterol-poor diet. *American Journal of Pathology* 73(2): 265-300, November 1973.

- (94) MJOS, O.D., THELLE, D.S., FORDE, O.H., VIK-MO, H. Family study of high density lipoprotein cholesterol and the relation to age and sex. *Acta Medica Scandinavica* 201(4): 323-329, 1977.
- (95) MORRIS, J.N., CHAVE, S.P.W., ADAM, C., SIREY, C., EPSTEIN, L., SHEEHAN, D.J. Vigorous exercise in leisure-time and the incidence of coronary heart-disease. *Lancet* 1: 333-339, February 17, 1973.
- (96) MOSS, A.J., DECAMILLA, J., DAVIS, H. Cardiac Death in the first 6 months after myocardial infarction: Potential for mortality reduction in the early posthospital period. *American Journal of Cardiology* 39(6): 816-820, May 26, 1977.
- (97) MULCAHY, R., HICKEY, N., GRAHAM, I.M., MACAIRT, J. Factors affecting the 5-year survival rate of men following acute coronary heart disease. *American Heart Journal* 93(5): 556-559, May 1977.
- (98) NAEYE, R.L., TRUONG, L.D. Effects of cigarette smoking on intramyocardial arteries and arterioles in man. *American Journal of Clinical Pathology* 68(4): 493-498, October 1977.
- (99) NAKAYAMA, Y. Epidemiological research in Japan on smoking and cardiovascular diseases. In: Schettler, G., Goto, Y., Hata, Y., Klose, G. (Editors). *Atherosclerosis IV. Proceedings of the Fourth International Symposium, Tokyo, 1976*. Berlin, Springer-Verlag, 1977, pp. 149-153.
- (100) OBERMAN, A., RAY, M., TURNER, M.E., BARNES, G., GROOMS, C. Sudden death in patients evaluated for ischemic heart disease. *Circulation* 51/52 (Supplement III): 170-172, December 1975.
- (101) OMAE, T., TAKESHITA, M., HIROTA, Y. The Hisayama study and joint study on cerebrovascular diseases in Japan. In: Scheinberg, P. (Editor). *Cerebrovascular Diseases. Proceedings of the Tenth Princeton Conference, New Jersey, 1976*. New York, Raven Press, 1976, pp. 255-265.
- (102) ORY, H.W. Association between oral contraceptives and myocardial infarction. A review. *Journal of the American Medical Association* 237(24): 2619-2622, June 13, 1977.
- (103) PAUL, O. Discussion on Dr. Dawber's Paper (The interrelationship of tobacco smoke components to hyperlipidemia and other risk factors). In: Wynder, E.L., Hoffmann, D., Gori, G.B., (Editors). *Proceedings of the Third World Conference on Smoking and Health, New York, June 2-5, 1975. Volume I. Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No. (NIH) 76-1221, 1976, pp. 293-295.
- (104) PEARSON, T.A., DILLMAN, J.M., SOLEZ, K., HEPTINSTALL, R.H. Clonal markers in the study of the origin and growth of human atherosclerotic lesions. *Circulation Research* 43(1): 10-18, July 1978.
- (105) PEARSON, T.A., WANG, A., SOLEZ, K., HEPTINSTALL, R.H. Clonal characteristics of fibrous plaques and fatty streaks from human aortas. *American Journal of Pathology* 81(2): 379-387, November 1975.
- (106) PETITTI, D.B., WINGERD, J. Use of oral contraceptives, cigarette smoking, and risk of subarachnoid haemorrhage. *Lancet* 2: 234-236, July 29, 1978.
- (107) THE POOLING PROJECT RESEARCH GROUP. Relationship of blood pressure, serum cholesterol, smoking habit, relative weight and ECG abnormalities to incidence of major coronary events: Final report of the Pooling Project. *Journal of Chronic Diseases* 31(4): 201-306, April 1978.
- (108) REEVES, T.J., OBERMAN, A., JONES, W.B., SHEFFIELD, L.T. Natural history of angina pectoris. *American Journal of Cardiology* 33: 423-430, March 1974.
- (109) REICHENBACH, D.D., MOSS, N.S., MEYER, E. Pathology of the heart in sudden cardiac death. *The American Journal of Cardiology* 39(6): 865-872, May 26, 1977.

- (110) REID, D.D., HAMILTON, P.J.S., MCCARTNEY, P., ROSE, G., JARRETT, R.J., KEEN, H. Smoking and other risk factors for coronary heart-disease in British civil servants. *Lancet* 2(7993): 979-984, November 6, 1976.
- (111) RESEARCH COMMITTEE OF THE SCOTTISH SOCIETY OF PHYSICIANS. Ischaemic heart disease. A secondary prevention trial using clofibrate. Report by a Research Committee of the Scottish Society of Physicians. *British Medical Journal* 4: 775-784, December 25, 1971.
- (112) RHOADS, G.G., BLACKWELDER, W.C., STEMMERMAN, G.N., HAYASHI, T., KAGAN, A. Coronary risk factors and autopsy findings in Japanese-American men. *Laboratory Investigation* 39(3): 304-311, 1978.
- (113) ROBICSEK, F., DAUGHERTY, H.K., MULLEN, D.C. MASTERS, T.N., NARBAY, D., SANGER, P.W., COOK, J.W. The effect of continued cigarette smoking on the patency of synthetic vascular grafts in Leriche syndrome. *Collected Works on Cardio-Pulmonary Disease* 20: 62-70, December 1975.
- (114) ROCK, W., OALMANN, M., STRONG, J. Community pathology of myocardial lesions in men 25 to 44 years of age. *Laboratory Investigation* 32(3): 483, March 1975. (Abstract)
- (115) ROSE, G., REID, D.D., HAMILTON, P.J.S., MCCARTNEY, P., KEEN, H., JARRETT, R.J. Myocardial ischaemia, risk factors and death from coronary heart-disease. *Lancet* 1(8003): 105-109, January 15, 1977.
- (116) ROSENMAN, R.H., BRAND, R.J., SHOLTZ, R.I., FRIEDMAN, M. Multivariate prediction of coronary heart disease during 8.5 year follow-up in the Western Collaborative Group Study. *American Journal of Cardiology* 37(5): 903-910, May 1976.
- (117) ROSS, R., GLOMSET, J.A. The pathogenesis of atherosclerosis (Parts One and Two). *New England Journal of Medicine* 295(7): 369-377, August 12, 1976 and 295(8): 420-425, August 19, 1976.
- (118) RUBERMAN, W., WEINBLATT, E., GOLDBERG, J.D., FRANK, C.W., SHAPIRO, S. Ventricular premature beats and mortality after myocardial infarction. *The New England Journal of Medicine* 297(14): 750-757, October 6, 1977.
- (119) ST. CLAIR, R.W. Metabolism of the arterial wall and atherosclerosis. In: Paoletti, R., Gotto, A.M. Jr. (Editors). *Atherosclerosis Reviews*, Volume 1. New York, Raven Press, 1976, pp. 61-117.
- (120) SCHIEVELBEIN, H. The evidence for nicotine as an etiological factor in cardiovascular disease. In: Wynder, E.L., Hoffmann, D., Gori, G.B. (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume I. *Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No.(NIH) 76-1221, 1976, pp. 297-307.
- (121) SELTZER, C.C. Effect of smoking on blood pressure. *American Heart Journal* 87(5): 558-564, May 1974.
- (122) SHAPIRO, S., WEINBLATT, E., FRANK, C.W., SAGER, R.V. Incidence of coronary heart disease in a population insured for medical care (HIP). Myocardial infarction, angina pectoris, and possible myocardial infarction. *American Journal of Public Health* 59 (Supplement 6): 1-101, June 1969.
- (123) SKINHOJ, E., OLESEN, J., PAULSON, O.B. Influence of smoking and nicotine on cerebral blood flow and metabolic rate of oxygen in man. *Journal of Applied Physiology* 35(6): 820-822, 1973.
- (124) SLONE, D., SHAPIRO, S., ROSENBERG, L., KAUFMAN, D.W., HARTZ, S.C., ROSSI, A.C., STOLLEY, P.D., MIETTINEN, O.S. Relation of cigarette smoking to myocardial infarction in young women. *New England Journal of Medicine* 298(23): 1273-1276, June 8, 1978.

- (125) SMALL, D.M. Cellular mechanisms for lipid deposition in atherosclerosis (Parts One and Two). *The New England Journal of Medicine* 297(16): 873-877, October 20, 1977 and 297(17): 924-929, October 27, 1977.
- (126) SMITH, E.B., SMITH, R.H. Early changes in aortic intima. In: Paoletti, R., Gotto, A.M., Jr. (Editors). *Atherosclerosis Reviews*, Volume 1. New York, Raven Press, 1976, pp. 119-136.
- (127) SPAIN, D.M., BRADESS, V.A. Sudden death from coronary heart disease: Survival time, frequency of thrombi, and cigarette smoking. *Chest* 58(2): 107-110, August 1970.
- (128) SPAIN, D.M., SIEGEL, H., BRADESS, V.A. Women smokers and sudden death. The relationship of cigarette smoking to coronary disease. *Journal of the American Medical Association* 224(7): 1005-1007, May 14, 1973.
- (129) STAMLER, J., RHOMBERG, P., SCHOENBERGER, J.A., SHEKELLE, R.B., DYER, A., SHEKELLE, S., STAMLER, R., WANNAMAKER, J. Multivariate analysis of the relationship of seven variables to blood pressure: Findings of the Chicago Heart Association Detection Project in Industry, 1967-1972. *Journal of Chronic Diseases* 28(10): 527-548, November 1975.
- (130) STENDER, S., ASTRUP, P., KJELDSEN, K. The effect of carbon monoxide on cholesterol in the aortic wall of rabbits. *Atherosclerosis* 28(4): 357-367, December 1977.
- (131) STRONG, J.P., OMAE, P. (CHAIRMEN). Workshop 3. Epidemiology of atherosclerosis and geographic differences in risk factors. In: Schettler, G., Goto, Y., Hata, Y., Klose, G. (Editors). *Atherosclerosis IV. Proceedings of the Fourth International Symposium, Tokyo, 1976*. Berlin, Springer-Verlag, 1977, pp. 92-120.
- (132) STRONG, J.P., RICHARDS, M.L. Cigarette smoking and atherosclerosis in autopsied men. *Atherosclerosis* 23(3): 451-476, May/June 1976.
- (133) STRONG, J.P., SOLBERG, L.A., RESTREPO, C. Atherosclerosis in persons with coronary heart disease. *Laboratory Investigation* 18(5): 527-537, May 1968.
- (134) TALBOTT, E., KULLER, L.H., DETRE, K., PERPER, J. Biologic and psychosocial risk factors of sudden death from coronary disease in white women. *The American Journal of Cardiology* 39(6): 858-864, May 26, 1977.
- (135) THOMAS, W.A., FLORENTIN, R.A., REINER, J.M., LEE, W.M., LEE, K.T. Alterations in population dynamics of arterial smooth muscle cells during atherogenesis. IV. Evidence for a polyclonal origin of hypercholesterolemic diet-induced atherosclerotic lesions in young swine. *Experimental and Molecular Pathology* 24: 244-260, 1976.
- (136) TOPPING, D.L. Metabolic effects of carbon monoxide in relation to atherogenesis. *Atherosclerosis* 26(2): 129-137, February 1977.
- (137) TRUMP, B.F., MERGNER, W.J., KAHNG, M.W., SALADINO, A.J. Studies on the subcellular pathophysiology of ischemia. *Circulation* 53(3) Supplement 1: I-17-I-26, March 1976.
- (138) U.S. PUBLIC HEALTH SERVICE. *The Health Consequences of Smoking. A Reference Edition: 1976*. U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control, HEW Publication No. (CDC) 78-8357, 1976, 657 pp.
- (139) VESSELINOVITCH, D., WISSLER, R.W., FISHER-DZOGA, K., HUGHES, R., DUBIEN, L. Regression of atherosclerosis in rabbits. Part 1. Treatment with low-fat diet, hyperoxia, and hypolipidemic agents. *Atherosclerosis* 19: 259-275, 1974.
- (140) VESSEY, M.P., DOLL, R. Investigation of relation between use of oral contraceptives and thromboembolic disease. A further report. *British Medical Journal* 2(5658): 651-657, June 14, 1969.

- (141) VETERANS' ADMINISTRATION COOPERATIVE UROLOGICAL RESEARCH GROUP. Treatment and survival of patients with cancer of the prostate. *Surgery, Gynecology and Obstetrics* 124(5): 1011-1017, May 1967.
- (142) VISMARA, L.A., VERA, Z., FOERSTER, J.M., AMSTERDAM, E.A., MASON, D.T. Identification of sudden death risk factors in acute and chronic coronary artery disease. *American Journal of Cardiology* 39(6): 821-828, May 26, 1977.
- (143) VON AHN, B. Tobacco smoking, the electrocardiogram, and angina pectoris. *Annals of the New York Academy of Sciences* 90(1): 190-198, September, 1960.
- (144) WAGNER, W.D., ST. CLAIR, R.W., CLARKSON, T.B. Angiochemical and tissue cholesterol changes of *Macaca fascicularis* fed an atherogenic diet for 3 years. *Experimental and Molecular Pathology* 28(2): 140-153, April 1978.
- (145) WALD, N.J. Carbon monoxide as an aetiological agent in arterial disease—Some human evidence. In: Wynder, E.L., Hoffmann, D., Gori, G.B. (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume I. *Modifying the Risk for the Smoker*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No.(NIH) 76-1221, 1976, pp. 349-361.
- (146) WALD, N., HOWARD, S., SMITH, P.G., KJELDSEN, K. Association between atherosclerotic diseases and carboxyhaemoglobin levels in tobacco smokers. *British Medical Journal* 1: 761-765, March 31, 1973.
- (147) WANSTRUP, J., KJELDSEN, K., ASTRUP, P. Acceleration of spontaneous intimal-subintimal changes in rabbit aorta by a prolonged, moderate carbon monoxide exposure. *Acta Pathologica et Microbiologica Scandinavica* 75(3): 353-362, 1969.
- (148) WEBSTER, W.S., CLARKSON, T.B., LOFLAND, H.B. Carbon monoxide-aggravated atherosclerosis in the squirrel monkey. *Experimental and Molecular Pathology* 13: 36-50, 1970.
- (149) WEINBLATT, E., FRANK, C.W., SHAPIRO, S., SAGER, R.V. Prognostic factors in angina pectoris—A prospective study. *Journal of Chronic Diseases* 21: 231-245, July 1968.
- (150) WEINBLATT, E., SHAPIRO, S., FRANK, C.W., SAGER, R.V. Prognosis of men after first myocardial infarction: Mortality and first recurrence in relation to selected parameters. *American Journal of Public Health* 58(8): 1329-1347, August 1968.
- (151) WEINROTH, L.A., HERZSTEIN, J. Relation of tobacco smoking to arteriosclerosis obliterans in diabetes mellitus. *Journal of the American Medical Association* 131(3): 205-209, May 1946.
- (152) WEISS, N.S. Cigarette smoking and arteriosclerosis obliterans: An epidemiologic approach. *American Journal of Epidemiology* 95(1): 17-25, 1972.
- (153) WESSLER, S., MING, S.-C., GUREWICH, V., FREIMAN, D.G. A critical evaluation of thromboangiitis obliterans. The case against Buerger's disease. *New England Journal of Medicine* 262(23): 1150-1160, June 9, 1960.
- (154) WILHELMSSEN, L. Recent studies on smoking and CVD epidemiology: Scandinavia and some other Western European countries. In: Steinfeld, J., Griffiths, W., Ball, K., Taylor, R.M. (Editors). *Proceedings of the Third World Conference on Smoking and Health*, New York, June 2-5, 1975. Volume II. *Health Consequences, Education, Cessation Activities and Social Action*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Cancer Institute, DHEW Publication No. (NIH) 77-1413, 1977, pp. 171-177.
- (155) WISSLER, R.W., VESSELINOVITCH, D., GETZ, G.S. Abnormalities of the arterial wall and its metabolism in atherogenesis. *Progress in Cardiovascular Diseases* 18(5): 341-369, March/April 1976.

- (156) WOLINSKY, H. A new look at atherosclerosis. Cardiovascular Medicine: 41-54, September 1976.
- (157) WRAY, R., DEPALMA, R.G., HUBAY, C.H. Late occlusion of aortofemoral bypass grafts: Influence of cigarette smoking. Surgery 70(6): 969-973, December 1971.



## **5. CANCER.**

## CONTENTS

---

Introduction .....	9
<hr/>	
Lung Cancer .....	9
Trends in Lung Cancer Mortality.....	10
Epidemiological Studies .....	11
Dose-Response Relationships .....	12
Number of Cigarettes Smoked Per Day .....	12
Age at Which Smoking Began.....	13
Inhalation of Cigarette Smoke.....	14
Tar and Nicotine Content of Cigarettes .....	15
Lung Cancer in Women .....	16
Trends in Cigarette Consumption	
among Females .....	16
Epidemiological Studies.....	20
Dose-Response Relationships .....	21
Patterns of Cigarette Use.....	21
Twins.....	23
Lung Cancer and the Use of Other Forms	
of Tobacco .....	23
Histology of Lung Cancer.....	23
Cessation of Smoking.....	24
Lung Cancer and Air Pollution.....	25
Lung Cancer and Occupational Factors .....	27
Asbestos.....	28
Uranium Mining.....	28
Nickel.....	28
Chloromethyl Ethers .....	29
Animal Studies .....	29
Skin Painting and Subcutaneous Injections.....	29
Tracheobronchial Implantation and Instillation..	29
Inhalation Carcinogenesis .....	30
Nitrosamines.....	30
Phagocytosis .....	31
Conclusions.....	31
<hr/>	
Cancer of the Larynx.....	32
Epidemiological Studies .....	33
Asbestos .....	34
Animal Studies .....	34

Conclusions.....	36
<hr/>	
Oral Cancer .....	39
Epidemiological Studies .....	39
Other Forms of Tobacco.....	40
Other Risk Factors.....	40
Leukoplakia.....	41
Animal Studies .....	41
Conclusions.....	42
<hr/>	
Cancer of the Esophagus.....	42
Epidemiological Studies .....	42
Other Forms of Tobacco Use .....	43
Other Risk Factors.....	43
Autopsy Studies.....	44
Animal Studies .....	44
Conclusions.....	44
<hr/>	
Cancer of the Urinary Bladder and Kidney.....	45
Bladder Cancer .....	45
Epidemiological Studies.....	45
Other Risk Factors .....	47
Animal Studies.....	47
Kidney Cancer .....	47
Epidemiological Data.....	47
Conclusions.....	49
<hr/>	
Cancer of the Pancreas.....	50
Epidemiological Studies .....	50
Other Risk Factors .....	51
Animal Studies .....	51
Conclusions.....	53
<hr/>	
Mechanisms of Carcinogenesis .....	53
Smoke Composition.....	53
Experimental Models.....	53
Concepts of Carcinogenesis.....	54
Aryl Hydrocarbon Hydroxylase.....	57
Multi-Stage Model of Carcinogenesis.....	58
<hr/>	
References .....	59

## LIST OF FIGURES

---

Figure 1.—Relative risk of lung cancer for males, by number of cigarettes smoked per day and long-term use of filter (F) and nonfilter (NF) cigarettes..... 18

---

Figure 2.—Relative risk of lung cancer for females, by number of cigarettes smoked per day and long-term use of filter (F) and nonfilter (NF) cigarettes..... 19

---

Figure 3.—Lung cancer mortality in continuing cigarette smokers and nonsmokers as a percentage of the rate among ex-cigarette smokers at the time they stopped smoking..... 26

---

Figure 4.—Relative risk of developing larynx cancer for males, by number of cigarettes smoked per day and use of filter and nonfilter cigarettes..... 35

---

Figure 5.—Relative risk of developing larynx cancer for females, by number of cigarettes smoked per day and use of filter and nonfilter cigarettes..... 36

---

Figure 6.—Relative risk of developing larynx cancer for male ex-smokers, by years of smoking cessation..... 37

---

Figure 7.—Relative risk of developing larynx cancer for female ex-smokers, by years of smoking cessation..... 38

---

Figure 8.—Relative risk of pancreatic cancer in males, by number of cigarettes smoked ..... 52

## LIST OF TABLES

---

Table 1.—Lung cancer mortality ratios—prospective studies..... 12

Table 2.—Lung cancer mortality ratios for males, by current number of cigarettes smoked per day, from selected prospective studies .....	13
Table 3.—Lung cancer mortality ratios for males, by age began smoking, from selected prospective studies .....	14
Table 4.—Lung cancer mortality ratios for males, by degree of inhalation, from selected prospective studies.....	15
Table 5.—Age-adjusted lung cancer mortality ratios for males and females, by tar and nicotine in cigarettes smoked .....	16
Table 6.—Age-adjusted lung cancer mortality ratios for males and females, comparing those who smoked a few high T/N cigarettes with those who smoked many low T/N cigarettes.....	17
Table 7.—Mortality rates for lung cancer and cancer of the respiratory tract for white females in the United States per 100,000 population for selected years: 1940 to 1976.....	20
Table 8.—Percent of adult population who were current cigarette smokers in selected years in the United States .....	20
Table 9.—Percent of teenagers who were current cigarette smokers in selected years in the United States .....	21
Table 10.—Lung cancer mortality ratios for women—prospective studies.....	21
Table 11.—Lung cancer mortality ratios for females, by number of cigarettes smoked per day: A.C.S. 25-State Study .....	22
Table 12.—Lung cancer mortality ratios for females, by number of cigarettes smoked per day: Haenszel and Taeuber.....	22
Table 13.—Lung cancer mortality ratios for females, by duration of smoking: Swedish Study .....	22

Table 14.—Lung cancer mortality ratios for females, by degree of inhalation: A.C.S. 25-State Study.....	22
Table 15.—Lung cancer mortality ratios in ex-cigarette smokers, by number of years stopped smoking.....	25
Table 16.—Mortality ratios for cancer of the larynx—prospective studies.....	33
Table 17.—Mortality ratios for cancer of the oral cavity—prospective studies.....	40
Table 18.—Mortality ratios for cancer of the esophagus—prospective studies.....	43
Table 19.—Bladder cancer mortality ratios—prospective studies.....	46
Table 20.—Kidney cancer mortality, ratios and relative risks: selected prospective and retrospective studies .....	48
Table 21.—Kidney cancer mortality ratios, by amount smoked: U. S. Veterans Study.....	49
Table 22.—Pancreatic cancer mortality ratios—prospective studies.....	51
Table 23.—Mortality ratios for cancer of the pancreas among Swedish subjects, aged 18–69, by sex and amount smoked.....	52
Table 24.—Carcinogenic, promoting, and ciliotoxic agents in the gas phase of tobacco smoke.....	55
Table 25.—Carcinogenic agents in the particulate phase of tobacco smoke .....	56
Table 26.—Tumor promoters and co-carcinogens in the particulate phase of tobacco smoke .....	57

## **Introduction**

Cancer has been the second leading cause of death in the United States since 1937. There were an estimated 390,000 deaths from cancer in 1978 (4). The association between tobacco smoking and the development of lung cancer was first suggested in the 1920's and early 1930's (159, 206). In the early 1950's, more than a dozen retrospective studies were published which first generally alerted the medical and scientific community to the health hazards associated with cigarette smoking. The public was informed of the results of these studies, and as a consequence there was a significant, but brief, dip in the per capita consumption of cigarettes. The next decade brought an intensive worldwide investigation into the various diseases associated with cigarette smoking. The first official statement on smoking and health by the U.S. Government was contained in the Report of the Advisory Committee to the Surgeon General of the U.S. Public Health Service, which was released 15 years ago. The evidence available at that time warranted the conclusion that "Cigarette smoking is causally related to lung cancer in men; the magnitude of the effect of cigarette smoking far outweighs all other factors. The data for women, though less extensive, point in the same direction. The risk of developing lung cancer increases with the duration of smoking and the number of cigarettes smoked per day, and is diminished by discontinuing smoking" (217). In the 15 years since the 1964 Surgeon General's Report was published, these conclusions have been confirmed by numerous investigations in many countries. Cigarette smoking has also been implicated as a significant cause of cancer of the larynx, oral cavity, esophagus, urinary bladder, kidney, and pancreas. As data concerning the relationship of smoking to the development of cancer at various sites became available, they were summarized and published in the annual issues of the Health Consequences of Smoking (209, 210, 211, 212, 212a, 213, 214, 215, 216).

This chapter reviews the epidemiological and experimental data for each of the cancer sites associated with cigarette smoking. Discussions of the specific cancers are presented sequentially, based on the strength of the association with cigarette smoking: cancer of the lung, larynx, oral cavity, esophagus, urinary bladder, kidney, and pancreas.

## **Lung Cancer**

This year more people in the United States will die from lung cancer than from any other malignant disease. In 1950, when the nation first became generally aware that there was an association between smoking and lung cancer, there were 18,313 lung cancer deaths. In 1964, there were 45,838 deaths from lung cancer. The National Center for Health Statistics reported that in 1976 there were 86,267 deaths from lung cancer in the United States (150). It is estimated that there

were 92,400 deaths from lung cancer in 1978 (4). For every preventable death from highway accidents, there were approximately two deaths from lung cancer which could have been prevented if the individual had not smoked cigarettes. There are about 280 deaths from lung cancer each day in the United States.

This epidemic increase in lung cancer is reflected in rapidly changing mortality rates in both men and women. The mortality rate for men in 1950 was 19.9/100,000/year. This rose to 41.4 in 1964, and to 63.0 in 1976. The comparable figures for white females were 4.7 in 1950 and 8.0 in 1965, and climbing rapidly to 19.5 in 1976 (Table 7).

According to results from the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) Program, the mortality rates for black males and females are higher than for whites. In 1976, the lung cancer mortality rate for black males was 93.0, for black females it was 17.4 (154). Due to recent increases in death rates among females, the ratio of male to female mortality for lung cancer has dropped from 7:1 to less than 4:1.

While recent years have seen dramatic increases in relative survival rates for acute leukemias in children, Hodgkin's disease, multiple myeloma, and certain other malignancies, there has been little increase in survival rates for lung cancer. The 5-year survival rate for lung cancer in all states is 8 percent for males and 12 percent for females (151). The difference in survival rates between males and females can be explained by sex-specific differences in histology or stage of the disease.

#### **Trends in Lung Cancer Mortality**

In the United States there has been in the past few years a significant reduction in the percent of males and females who smoke cigarettes. As yet, there has not been a decline in the age-adjusted *total* mortality rates for lung cancer. When the lung cancer mortality rates by age are examined from 1950 through 1975, there is a continuing increase in older age groups for both males and females. This is probably due to the elevated risk experienced by older persons who use nonfiltered, high tar and nicotine cigarettes and who have done so for the majority of their lives. However, for female cohorts born in 1950-54 and male cohorts born in 1935-39 and 1940-44, the *age-specific* lung cancer mortality rates are below those of previous cohorts. This probably results from the reductions in cigarette consumption which have occurred in these groups.

There has been a change in the epidemic of lung cancer in England and Wales, as summarized by the International Union Against Cancer (UICC) workshop on the biology of cancer (243):

In England and Wales, lung cancer mortality stopped increasing in men under the age of 50 years during the 1950's and more recently has fallen in men under the age of 60 years. The death rate from



lung cancer in women ages 40 years and over has continued to rise, but has leveled, or fallen in younger women since the 1960's...The fall in lung cancer mortality among men under the age of 60 years is likely to be due to their reduced consumption since the end of the Second World War, and to the reduction in the tar yield of cigarettes since 1955; particularly with the change to filter cigarettes.

Although lung cancer mortality in women over 40 years has continued to increase along with their cigarette consumption, it is unlikely that the incidence of lung cancer will ever reach the high levels recorded in men, because the increasing cigarette consumption by women has been, and is continuing to be compensated for by a decrease in tar yield.

### **Epidemiological Studies**

The first comprehensive reviews of the effects of smoking on lung cancer were published in 1962 and 1964 by the Royal College of Surgeons of London and the Surgeon General of the United States, respectively (171, 217). They included data from studies on epidemiology, profiles of the consumption of tobacco, the composition and carcinogenicity of components of tobacco smoke, the effects of smoke on experimental animals, and the pathological changes observed in humans and animals. The conclusions reached in these assessments and by all of the periodic reviews that have followed at regular intervals (209, 210, 211, 212, 212a, 213, 214, 215, 216) are impressively uniform and consistent. So much so that it has been observed that the results of any one of the major studies might be taken to represent all of them.

There have been at least nine major prospective epidemiological studies which have examined the relationship between cigarette smoking and mortality from various causes. The results of eight of these studies are related to cigarette smoking and lung cancer and are presented in Table 1. The lowest mortality ratios are experienced by female smokers. The mortality ratios for male cigarette smokers are as low as 3.85 for Japanese males and as high as 14.0 for British doctors and Canadian veterans. Combining the data from the largest studies allows the conclusion that cigarette smokers on the average are 10 times as likely to develop lung cancer as nonsmokers. The mortality ratios are much higher for heavy cigarette smokers. This will be detailed in the section on dose-response relationships.

In the past 30 years, more than 50 retrospective studies on the relationship between cigarette smoking and lung cancer have been published. These data are too extensive for convenient summarization; they have been reviewed in recent issues of the Health Consequences of Smoking (212, 212a, 213, 214, 215).

**TABLE 1.—Lung cancer mortality ratios—prospective studies**

Population	Size	Number of deaths	Nonsmokers	Cigarette smokers
British doctors(47a)	34,000 males	441	1.00	14.0
Swedish study(32)	27,000 males 28,000 females	55 8	1.00 1.00	8.2 4.5
Japanese study(77a,78)	122,000 males 143,000 females	590 148	1.00 1.00	3.76 2.08
A.C.S. 25-State Study(65)	440,000 males 562,000 females	1,159 183	1.00 1.00	9.20 2.20
U.S. veterans(90)	239,000 males	1,256	1.00	12.14
Canadian veterans(20)	78,000 males	331	1.00	14.2
A.C.S. 9-State Study(68)	188,000 males	448	1.00	10.73
California males in 9 occupations(228)	68,000 males	368	1.00	7.61

### **Dose-Response Relationships**

An important factor in the causal relationship between smoking and lung cancer is the demonstration of dose-response relationships. In most epidemiological studies, dosage has been measured by the number of cigarettes smoked per day at the time of entry into the study. Other dose variables which have been examined include the maximum number of cigarettes smoked per day, the age an individual began smoking, the degree of inhalation of tobacco smoke, the total number of years an individual has smoked, the total lifetime number of cigarettes smoked, tar and nicotine levels of the brand of cigarettes used, the number of puffs per cigarette, the length of the unburned portion of the cigarette, and combinations of these variables into "dosage" scores. All of these variables have been shown in one study or another to contribute to the risk of developing lung cancer. Only a few representative samples of dosage variables as related to lung cancer mortality are examined in this section.

#### *Number of Cigarettes Smoked Per Day*

The risk of developing lung cancer increases with the number of cigarettes smoked per day. In the U.S. and British populations, the risk of developing lung cancer for individuals smoking more than two packs

**TABLE 2.—Lung cancer mortality ratios for males, by current number of cigarettes smoked per day, from selected prospective studies**

	Cigarettes smoked per day	Mortality ratio
A.C.S. 25- state study(65)	Nonsmoker	1.00
	1-9	4.62
	10-19	8.62
	20-39	14.69
	40+	18.77
British doctors(47a)	Nonsmoker	1.00
	1-14	7.80
	15-24	12.70
	25+	25.10
Swedish males(32)	Nonsmoker	1.00
	1-7	2.30
	8-15	8.80
	16+	13.90
Japanese males(78)	Nonsmoker	1.00
	1-9	1.90
	10-14	3.52
	15-24	4.11
	25-49	4.57
	50+	5.78

a day is approximately 20 times that of nonsmokers (47a, 65, 68, 80, 228). Data for Swedish males are of the same magnitude (32). Japanese males who smoke 50 or more cigarettes a day experience a risk which is 5.8 times greater than for nonsmokers. Hirayama noted that the slope of the dose-response curve for lung cancer was less in Japan than in the United States and that this was probably due to the lower percentage of regular deep inhalers, a lower level of environmental promoting conditions, and also a higher percentage of adenocarcinoma in Japan than in the United States (78). Table 2 presents lung cancer mortality ratios from selected prospective studies for males by the current number of cigarettes smoked per day.

#### *Age at which Smoking Began*

Lung cancer mortality ratios exhibit an inverse relationship with the age of initiation of the smoking habit. Lung cancer mortality ratios for males by age at which they began smoking are presented in Table 3. Most cigarette smokers began the habit while in high school and are at the greatest risk of developing lung cancer. Those who began smoking

**TABLE 3.—Lung cancer mortality ratios for males, by age began smoking, from selected prospective studies**

	Age began smoking in years	Mortality ratio
A.C.S. 25-State Study(65)	Nonsmoker	1.00
	25+	4.08
	20-24	10.08
	15-19	19.69
	under 15	16.77
Japanese study(78)	Nonsmoker	1.00
	25+	2.87
	20-24	3.85
	under 20	4.44
U.S. veterans(90)	Nonsmoker	1.00
	25+	5.20
	20-24	9.50
	15-19	14.40
	under 15	18.70

after the age of 25 have mortality ratios which are only 4 to 5 times greater than those of nonsmokers.

*Inhalation of Cigarette Smoke*

Inhalation of tobacco smoke is an important dosage variable. Inhalation of smoke well into the lungs is the major mechanism whereby lung tissue is exposed to the carcinogens which ultimately produce lung cancer. Techniques for quantitating the degree of tobacco smoke inhalation have been developed using carboxyhemoglobin levels or end expiratory carbon monoxide levels as an index of smoke inhalation. These objective methods of measuring inhalation have not been applied to studies of lung cancer mortality. In most investigations, the smoker was asked to report subjectively on his own inhalation practices. This is subject to considerable variation but is not as inaccurate as might be presumed. Available data show a strong dose-response relationship between self-reported inhalation of cigarette smoke and lung cancer mortality. Representative figures from selected prospective studies are presented in Table 4. These data suggest that cigarette smokers may underestimate the degree to which they inhale cigarette smoke. Those who report that they do not inhale cigarette smoke experience lung cancer mortality ratios which are 4 to 8 times greater than for nonsmokers. Deep inhalation results in mortality ratios which are as high as 17 times greater than for nonsmokers.

**TABLE 4.—Lung cancer mortality ratios for males, by degree of inhalation, from selected prospective studies**

	Degree of inhalation	Mortality ratio
A.C.S. 25-State Study(65)	Nonsmoker	1.00
	None	8.00
	Slight	8.92
	Moderate	13.08
	Deep	17.00
Swedish males(32)	Nonsmoker	1.00
	None	3.70
	Light inhalation	7.80
	Deep inhalation	9.20

*Tar and Nicotine Content of Cigarettes*

The major constituents of cigarette smoke that cause lung cancer are among the more than 2,000 different compounds found in cigarette smoke. Cigarette filters, first introduced during the mid-1950's, have the effect of trapping tar. Data presented by Maxwell (136) show that, in 1976, more than 600 billion cigarettes were smoked and that 88.4 percent of these were filtered. It has been known that the risk of developing lung cancer increased with the tar and nicotine content of cigarettes. Until recently, however, there has not been a great deal of evidence that individuals who switch to lower tar and nicotine cigarettes experience less lung cancer mortality (27). It has been argued that, if the tar and nicotine content of tobacco were reduced, individuals might increase the number of cigarettes smoked per day and thereby abolish any benefit that might be gained. Alternatively, those who switch to low tar and nicotine cigarettes might inhale the smoke more deeply than smokers of high tar and nicotine cigarettes, and thereby exposure to tar and nicotine might not be reduced. In a large prospective study by Hammond, et al. (67), these tar and nicotine relationships were examined with respect to lung cancer. The 897,825 men and women in 23 States were divided into 3 tar and nicotine categories. The high tar and nicotine (T/N) category was defined as 2.0 to 2.7 mg of nicotine and 25.8 to 35.7 mg of tar. The medium T/N category was defined as 1.2 to 1.9 mg of nicotine and 17.6 to 25.7 mg of tar. The low T/N category included cigarettes containing less than 1.2 mg of nicotine and less than 17.6 mg of tar. A matched-group analysis, similar to age standardization, was utilized. Individuals in each group were alike with respect to age, race, number of cigarettes smoked per day, age when they began to smoke cigarettes, place of residence,

**TABLE 5.—Age-adjusted lung cancer mortality ratios\* for males and females, by tar and nicotine in cigarettes smoked**

	Males	Females
High T/N	1.00	1.00
Medium T/N	0.95	0.79
Low T/N	0.81	0.60

\*The mortality ratio for the category with highest risk was made 1.00 so that the relative reductions in risk with the use of lower T/N cigarettes could be visualized.

SOURCE: Hammond, E.C. (67)

occupational exposure to dust fumes, chemicals, etc., education, prior history of lung cancer, and prior history of heart disease. Results of this analysis are presented in Table 5. The mortality ratio for the category with the highest risk was made 1.0 so that the relative reduction in risk with the use of lower T/N cigarettes could be visualized. For males smoking the same number of cigarettes per day, there appears to be a 20 percent reduction in risk of developing lung cancer with the use of low T/N cigarettes. For females, there was a 40 percent reduction in the risk of developing lung cancer with the use of low T/N cigarettes, keeping the number of cigarettes smoked per day constant. The amount of tar and nicotine taken into the body per day depends on the number of cigarettes smoked, as well as on the tar and nicotine content of each cigarette. Hammond conducted a second matched-group analysis comparing subjects who smoked 1 to 19 high T/N cigarettes per day and those who smoked 20 to 39 low T/N cigarettes per day. These results are presented in Table 6. The number of cigarettes smoked per day was a relatively more important variable than the tar and nicotine content of cigarettes. The mortality ratio was 1.6 for males and 2.1 for females who smoked 20 to 39 low T/N cigarettes a day, compared to individuals who smoked only 1 to 19 high T/N cigarettes per day.

Wynder and Stellman (253) conducted a large retrospective study of 1,034 white males and females with histologically proved cancer of the lung and larynx. Relative risks were consistently lower among long-term smokers of filter cigarettes, compared to smokers of nonfilter cigarettes. These groups were standardized for number of cigarettes smoked, duration of smoking, inhalation, and cigarette butt length. These dose-response relationships are presented in Figures 1 and 2.

### **Lung Cancer in Women**

#### *Trends in Cigarette Consumption Among Females*

In 1964, the Advisory Committee to the Surgeon General concluded that cigarette smoking was causally related to cancer in men, and that

**TABLE 6.—Age-adjusted lung cancer mortality ratios\* for males and females, comparing those who smoked a few high T/N cigarettes with those who smoked many low T/N cigarettes**

	1-19 high T/N cigarettes/day	20-39 low T/N cigarettes/day
Males	1.00	1.6
Females	1.00	2.1

\*The mortality ratio for the category with lowest risk was made 1.00 so the increase in risk with smoking more cigarettes/day could be illustrated.

SOURCE: Hammond, E. C. (67)


“the data for women though less extensive, point in the same direction” (217). Today, 15 years later, the lung cancer epidemic among women is well established. Several investigators had predicted sharp increases in lung cancer mortality among women. In 1966, Linden (118) examined lung cancer mortality in California women and predicted: “One can expect to see further increase in the number of lung cancer deaths and the death rates as the increasing proportions of women who smoke cigarettes reach the age when lung cancer is most likely to occur.”

In 1964, lung cancer was the fifth leading cause of death from cancer in women. It became the fourth leading cause in 1967 and moved to the third leading cause of death from cancer in 1969, passing cancer of the uterus. Projections for 1979 indicate that lung cancer is approaching cancer of the colon and rectum as the second leading cause of death from cancer in women. If present trends are not reversed, during the next decade lung cancer will become the leading cause of death from cancer in women, exceeding deaths from cancer of the breast.

In 1955, there were only 4,100 deaths from lung cancer in women. In 1976, the National Center for Health Statistics reported there were 20,455 deaths from lung cancer among females in the United States (150); the American Cancer Society estimated that in 1978 this increased to 21,900 deaths (4).

These increases are not due to increases in the population. Death rates for lung cancer have been steadily rising in women, especially in the past decade. The lung cancer mortality rate for white females in 1950 was 4.7 per 100,000; by 1976 this had risen to 19.5 per 100,000. This is more than a fourfold increase (Table 7).

The Surveillance, Epidemiology and End Results (SEER) Program of the National Cancer Institute recently reported that the lung cancer death rate for black females exceeded that of white females (16.8 blacks, 15.0 whites)(154). Data from this survey are collected from 10 geographic areas in the United States and therefore do not represent




**FIGURE 1.—Relative risk of lung cancer for males, by number of cigarettes smoked per day and long-term use of filter (F) or nonfilter (NF) cigarettes**  
SOURCE: Wynder, E.L. (259)

national trends per se. The lung cancer mortality rate (15.0 per 100,000) among black females in the general U.S. population is equal to that of whites.

Increases in lung cancer mortality among females cannot be explained by exposure to occupational carcinogens. Increases in cigarette consumption are responsible for these trends.





**FIGURE 2.—Relative risk of lung cancer for females, by number of cigarettes smoked per day and long-term use of filter (F) and nonfilter (NF) cigarettes**

SOURCE: Wynder, E.L. (253)

The epidemic of lung cancer in women has lagged behind that in men, primarily because of differences in patterns of cigarette smoking. There are fewer women smoking than men, but the gap is narrowing. Among teenagers in several age categories, girls are smoking more than boys (155). Table 8 shows the percentage of the U.S. adult population who are currently smoking cigarettes for selected years. In 1975, approximately 29 percent of adult females were smoking, whereas 39 percent of adult males were smoking (155). It should also be noted that, over the past decade, there has been a 2.6 percent