

# 1. Eastern Bering Sea Walleye Pollock

James N. Ianelli, Steve Barbeaux, Taina Honkalehto,  
Stan Kotwicki, Kerim Aydin and Neal Williamson

Alaska Fisheries Science Center  
National Marine Fisheries Service

## Executive Summary

The focus of this chapter is on the Eastern Bering Sea (EBS) region. The Aleutian Islands region (Chapter 1A) and the Bogoslof Island area (Chapter 1B) are presented as separate sections.

### Changes in the input data

The 2007 NMFS summer bottom-trawl survey (BTS) abundance at age estimates were computed and included for this assessment. The NOAA ship R/V Oscar Dyson was diverted from planned survey research in the Gulf of Alaska this summer to add a mid-water echo-integration trawl (EIT) survey for the EBS. This was the first complete survey conducted by this vessel in this region. Age composition estimates were derived from the population-at-length estimates using the 2007 BTS age-length key. The 2006 age composition estimates were updated using EIT age data (last year the BTS age-length key was used). The EIT survey also extended into the Russian zone and covered part of the Navarin Basin.

The BTS biomass estimate was 4.3 million t, up by 42% from the 2006 estimate of 3.0 million t but still only 87% of the long-term mean of the bottom-trawl survey since 1982. The 2007 EIT survey biomass estimate was 1.88 million t, up slightly from last year's survey but only 55% of the long-term mean for this survey (since 1979). Both surveys indicate that the 2006 year class is strong and that the 2005 year-class is now apparently below average.

Observer data for age and size composition and average weight-at-age were evaluated for the 2006 fishery and were included in the analyses. The mean weights-at-age assumed in last year's assessment (for ABC and the 2006 fishery) were significantly higher than the new data indicate. The catch-at-age data were reanalyzed for uncertainty using a bootstrap two-stage sampling approach for 1991-2006. This includes an evaluation in fishery mean body weight-at-age. Results from these analyses are presented in Attachment A1. The total pollock catch for 2006 was estimated from the NMFS Alaska Region data and the value for 2007 catch was assumed to be 1,340,000 t.

### Changes in the assessment model

The assessment model was modified to evaluate a broader range of options. The more significant of these changes are as follows: 1) an age-length transition matrix was estimated so that the current-year fishery length frequency data can be used; 2) the ability to split the age-1 values from the rest of the age compositions for the EIT survey was added so that the interaction between age-1 variability and other age groups was reduced; 3) total age 2 and older numbers-at-age from the surveys were used to tune the model (previously 1-year olds were included); and 4) The Tier 1 ABC estimation method now uses "fishable" biomass defined as the biomass of pollock available to the fishery as modified by the selectivity-at-age estimates; previously age 3+ biomass had been used.

### Changes in the assessment results

As was projected in last year's assessment, the stock is estimated to be below the  $B_{msy}$  level beginning in 2007 and into 2008. Under FMP amendment 56, this invokes a proportional adjustment to the harvest rate. Since the available data indicate the spawning biomass for 2008 is projected to be lower than expected in last year's assessment, the adjustment to the harvest rate drops from about 84% to about 72%. Combined with other factors (e.g., a lower than expected mean weight-at-age observed in the 2006 fishery, a poorer showing of the 2005 year class in this year's surveys) this results in an ABC that is about 13% lower than projected from last year's assessment for 2008. The maximum permissible ABC based

on the Tier 1b harmonic mean  $F_{msy}$  is 1,170 thousand t for 2008. The corresponding overfishing level (OFL) is estimated 1,443 thousand t. The 2009 projection indicates that although the stock appears to show positive signs of recruitment, the impact of 4-5 successively below-average year-classes in the spawning component of the stock results in a 2009 Tier 1b ABC value to 976 thousand t. To return spawning biomass exploitation rates to pre 2006 levels and to reduce the projected variability, an ABC of 1.0 million t is recommended for 2008. Alternatively, a value of 1,020 thousand t is projected to be equal to the 1990 exploitation rate (the highest rate observed since 1978).

#### Response to SSC and Plan Team comments

The following SSC comments were provided in its December 2006 minutes along with responses relevant to this assessment.

*From Juvenile weight-at-age: The SSC appreciates the effort to examine juvenile weight-at-age in the BASIS surveys and looks forward to seeing the results. In addition, the SSC would like to know if weight of age 1 fish in previous surveys could yield information of value to address ecosystem concerns.*

The BASIS program continues to evaluate biological patterns in fish relative to environmental conditions. To date, there are limited data from this survey available. However, patterns in mean weights at age for juveniles in fishery and survey data are presented and discussed below. There appears to be some pattern in growth rates by years, and for survey data, part of the trend could be explained by the average date of survey operation.

*Non-pollock backscatter: The SSC found this information intriguing and would like further evaluation of the composition of the backscatter. Could it be age – 0 pollock to any degree? Is there the possibility of using higher-frequency acoustics to measure zooplankton biomass?*

Age 0 pollock probably form part of the 38-kHz non-pollock backscatter although their contribution relative to that of other scatterers (abiotic and biotic) is unknown. (see Fig. 1.46 and text below). The primary trawl sampling gear used to identify echosign during the EIT surveys is assumed to provide representative samples of larger fishes (e.g., age 1+-sized pollock). Without the use of other specialized sampling gear (and additional vessel time) the information needed to identify components of this non-pollock backscatter is unavailable. However, the species composition of this backscatter may be diverse and likely varies in time and space. Additionally, estimates of target strength for the species that comprise this backscatter may also be unknown. This lack of information means extra caution is needed to interpret the significance of the raw non-pollock unscaled acoustic backscatter data presented in Figure 1.46 relative to zooplankton biomass. MACE Program scientists are currently evaluating the use of multi-frequency acoustics and have had some success identifying euphausiid aggregations using the 38- and 120-kHz backscatter data.

*Russian catches: The SSC appreciates the efforts to formally include Russian catches in the pollock model and to contact Russian scientists for information. The SSC encourages ongoing efforts to examine age composition, recognizing that standardization of ageing is still an issue.*

The ability to explicitly evaluate Russian fishery impacts by modifying how survey catchabilities are treated was added this year. While this fails to address the issue of Russian fishery age compositions and removals directly, coupled with the EIT surveys that extended into the Russian zone this year, helps to more explicitly account for components of EBS pollock that move into the Russian zone.

*Weight-length relationships: The SSC encourages the authors to see if weight-length relationships change from year to year in parallel with plankton abundance or other components of the ecosystem.*

An evaluation of within-year changes in mean weights given length is presented, along with detailed patterns of fishery stratum-specific mean weights-at-age. Results suggest that the 2006 value of mean weight at age is substantially lower than the mean, consistent with other observations that the productivity of the environment was lower.

*Arrowtooth flounder predation: The SSC encourages the authors to explore the sensitivity of the stock assessment model to this predation. This could be done in a variety of ways: (1) using time-varying  $M$  or (2) making  $M$  a function of arrowtooth flounder abundance or consumption of pollock by examining Jurado-Molina's multi-species VPA model.*

Advances on developing the statistical multi-species model continue. In the past year a technical workshop was held where methods for parameterizations and treatment of stomach content data were presented and a number of improvements suggested. Additionally, a University of Washington post-doc was hired to assist in exploring patterns of EBS arrowtooth flounder bio-energetics and foraging patterns.

## Introduction

Walleye pollock (*Theragra chalcogramma*) are broadly distributed throughout the North Pacific with the largest concentrations found in the Eastern Bering Sea. Also marketed under the name Alaska pollock, this species continues to represent over 40% of the global whitefish production with the market disposition split fairly evenly between fillets, whole (head and gutted), and surimi. An important component of the commercial production is the sale of roe from pre-spawning pollock. Pollock are considered a relatively fast growing and short-lived species and currently represents a major biological component of the Bering Sea ecosystem.

In the U.S. portion of the Bering Sea three stocks of pollock are identified for management purposes. These are: Eastern Bering Sea which consists of pollock occurring on the Eastern Bering Sea shelf from Unimak Pass to the U.S.-Russia Convention line; the Aleutian Islands Region encompassing the Aleutian Islands shelf region from 170°W to the U.S.-Russia Convention line; and the Central Bering Sea—Bogoslof Island pollock. These three management stocks undoubtedly have some degree of exchange. The Bogoslof stock forms a distinct spawning aggregation that has some connection with the deep water region of the Aleutian Basin. In the Russian EEZ, pollock are considered to form two stocks, a western Bering Sea stock centered in the Gulf of Olyutorski, and a northern stock located along the Navarin shelf from 171°E to the U.S.-Russia Convention line. There is some indication (based on contiguous surveys) that the fishery in the northern region may be a mixture of Eastern and western Bering Sea pollock with the former predominant. Bailey et al. (1999) present a thorough review of population structure of pollock throughout the north Pacific region. Genetic differentiation using microsatellite methods suggest that populations from across the North Pacific Ocean and Bering Sea were similar. However, weak differences were significant on large geographical scales and conform to an isolation-by-distance pattern (O'Reilly and Canino, 2004; Canino et al. 2005).

## Fishery

From 1954 to 1963, pollock were harvested at low levels in the Eastern Bering Sea and directed foreign fisheries began in 1964. Catches increased rapidly during the late 1960s and reached a peak in 1970-75 when they ranged from 1.3 to 1.9 million t annually (Fig. 1.1). Following a peak catch of 1.9 million t in 1972, catches were reduced through bilateral agreements with Japan and the USSR.

Since the advent of the U.S. EEZ in 1977 the annual average Eastern Bering Sea pollock catch has been 1.2 million t and has ranged from 0.9 million t in 1987 to nearly 1.5 million t in recent years (Fig. 1.1). Stock biomass has apparently ranged from a low of 4-5 million t to highs of 10-12 million t. United States vessels began fishing for pollock in 1980 and by 1987 they were able to take 99% of the quota. Since 1988, only U.S. vessels have been operating in this fishery. By 1991, the current NMFS observer program for north Pacific groundfish-fisheries was in place.

Foreign vessels began fishing in the mid-1980s in the international zone of the Bering Sea (commonly referred to as the “Donut Hole”). The Donut Hole is entirely contained in the deep water of the Aleutian Basin and is distinct from the customary areas of pollock fisheries, namely the continental shelves and slopes. Japanese scientists began reporting the presence of large quantities of pollock in the Aleutian Basin in the mid-to-late 1970's, but large scale fisheries did not occur until the mid-1980s. In 1984, the Donut Hole catch was only 181 thousand t (Fig. 1.1, Table 1.1). The catch grew rapidly and by 1987 the high seas catch exceeded the pollock catch within the U.S. Bering Sea EEZ. The extra-EEZ catch peaked in 1989 at 1.45 million t and has declined sharply since then. By 1991 the Donut Hole catch was 80% less than the peak catch, and data for 1992 and 1993 indicate very low catches (Table 1.1). A fishing moratorium was enacted in 1993 and only trace amounts of pollock have been harvested from the Aleutian Basin by resource assessment fisheries. During 2002-2005 the EBS region pollock catch has averaged 1.463 million tons while for the period 1982-2000, the average was 1.15 million tons. The

effect of this level of fishing continues to be closely monitored by resource assessment surveys and an extensive fishery observer program.

### **Fishery characteristics**

Pre-spawning aggregations of pollock are the focus of the so-called “A-season” which opens on January 20<sup>th</sup> and extends into early-mid April. This fishery produces highly valued roe which can comprise over 4% of the catch in weight. The second season presently opens on June 1<sup>st</sup> and extends through late October. Since the closure of the Bogoslof management district (INPFC area 518) to directed pollock fishing in 1992, the “A-season” (January – March) pollock fishery on the Eastern Bering Sea (EBS) shelf has been concentrated primarily north and west of Unimak Island (Ianelli *et al.* 1998). Depending on ice conditions and fish distribution, there has also been effort along the 100 m contour (and deeper) between Unimak Island and the Pribilof Islands. This pattern has been fairly similar during the period 2006 - 2007 (Fig. 1.2), especially compared to the 2005 winter fishery which was distributed farther north. The catch estimates by sex for the A-season compared to estimates for the entire season indicate that over time, the number of males and females has been fairly equal (Fig. 1.3). The length frequency information from the fishery shows that the size of pollock caught are generally larger than 40 cm with some smaller fish caught during years when a strong year class appeared (Fig. 1.4).

In 2006 and 2007 the summer fishing has concentrated more in the NW region (Fig. 1.5). Coupled with higher fuel prices, this was a concern for shore-based vessels that had much longer distances to travel to the prime fishing grounds. While both these years exhibited colder-than-usual bottom temperatures (see discussion of bottom trawl survey results below), it is unclear that these conditions are the major cause of this apparent shift in fish distribution. For contrast, historical foreign-reported data were recovered and evaluated to see if the shift in distribution was a unique event. In fact, prior to 1986, in about half of the years the majority of the summer fish were taken to the west of 170°W (the NW zone of the EBS). Since 1991, the summer pollock catches have been much more concentrated in the SE (east of 170°W) zone (Fig. 1.6).

The length frequency information from the fishery reveals a marked progression of the large 1989 year class growing over time and the appearance of the 1992 year class in 1996-97, the 1996 year class in 1998-2001, and subsequently the 2000 year class (Fig. 1.7). The 2003 fishery data show an unusually high mode of fish at around 40cm that advanced to 45 cm by 2004 and reached about 48 cm in 2005.

Barbeaux *et al.* (2005b) presented some results on the development of small-scale spatial patterns of pollock aggregations. This involved a subset of some 32,000 km (~17,300 nm) of tracked acoustic backscatter collected opportunistically aboard commercial vessels. They found that during the daytime pollock tend to form patchy, dense aggregations while at night they disperse to a few uniform low-density aggregations. Changes in trawl tow duration and search patterns coincide with these changes in pollock distributions. Qualitative results suggest that rapid changes in distributions and local densities of Alaska pollock aggregations occur in areas of high fishing pressure. Analyses of this type will continue to improve understanding on the dynamics of the pollock fishery and biological responses.

### **Fisheries Management**

Due to concerns over possible impacts groundfish fisheries may have on rebuilding populations of Steller sea lions, NMFS and the NPFMC have changed management of Atka mackerel (mackerel) and pollock fisheries in the Bering Sea/Aleutian Islands (BSAI) and Gulf of Alaska (GOA). These changes were designed to reduce the possibility of competitive interactions with Steller sea lions. For the pollock fisheries, comparisons of seasonal fishery catch and pollock biomass distributions (from surveys) by area in the Eastern Bering Sea led to the conclusion that the pollock fishery may have had disproportionately high seasonal harvest rates within critical habitat that *could* lead to reduced sea lion prey densities. Consequently, management measures redistributed the fishery both temporally and spatially according to pollock biomass distributions. The idea was that seasonal and spatially explicit exploitation rates should

be consistent with area-wide and annual exploitation rates for pollock. Three types of measures were implemented in the pollock fisheries: 1) pollock fishery exclusion zones around sea lion rookery or haulout sites; 2) phased-in reductions in the seasonal proportions of TAC that can be taken from critical habitat; and 3) additional seasonal TAC releases to disperse the fishery in time.

Prior to the management measures, the pollock fishery occurred in each of the three major fishery management regions of the north Pacific ocean managed by the NPFMC: the Aleutian Islands (1,001,780 km<sup>2</sup> inside the EEZ), the Eastern Bering Sea (968,600 km<sup>2</sup>), and the Gulf of Alaska (1,156,100 km<sup>2</sup>). The marine portion of Steller sea lion critical habitat in Alaska west of 150°W encompasses 386,770 km<sup>2</sup> of ocean surface, or 12% of the fishery management regions.

Prior to 1999, a total of 84,100 km<sup>2</sup>, or 22% of critical habitat, was closed to the pollock fishery. Most of this closure consisted of the 10 and 20 nm radius all-trawl fishery exclusion zones around sea lion rookeries (48,920 km<sup>2</sup> or 13% of critical habitat). The remainder was largely management area 518 (35,180 km<sup>2</sup>, or 9% of critical habitat) which was closed pursuant to an international agreement to protect spawning stocks of central Bering Sea pollock.

In 1999, an additional 83,080 km<sup>2</sup> (21%) of critical habitat in the Aleutian Islands was closed to pollock fishing along with 43,170 km<sup>2</sup> (11%) around sea lion haulouts in the GOA and Eastern Bering Sea. In 1998, over 22,000 t of pollock were caught in the Aleutian Island regions, with over 17,000 t caught in Aleutian Islands critical habitat region. Between 1998 and 2004 a directed fishery for pollock was prohibited. Consequently, a total of 210,350 km<sup>2</sup> (54%) of critical habitat was closed to the pollock fishery. The portion of critical habitat that remained open to the pollock fishery consisted primarily of the area between 10 and 20 nm from rookeries and haulouts in the GOA and parts of the Eastern Bering Sea foraging area. In 2000, phased-in reductions in the proportions of seasonal TAC that could be caught within the BSAI Steller sea lion Conservation Area (SCA) were implemented. Since 2005, a limited pollock fishery has been prosecuted in the Aleutian Islands but with less than 2,000 t of annual catch.

The Bering Sea/Aleutian Islands pollock fishery was also subject to changes in total catch and catch distribution. Disentangling the specific changes in the temporal and spatial dispersion of the EBS pollock fishery resulting from the sea lion management measures from those resulting from implementation of the American Fisheries Act (AFA) is difficult. The AFA reduced the capacity of the catcher/processor fleet and permitted the formation of cooperatives in each industry sector by the year 2000. Both of these changes would be expected to reduce the rate at which the catcher/processor sector (allocated 36% of the EBS pollock TAC) caught pollock beginning in 1999, and the fleet as a whole in 2000 when a large component of the onshore fleet also joined cooperatives. Because of some of its provisions, the AFA gave the industry the ability to respond efficiently to changes mandated for sea lion conservation that otherwise could have been more disruptive to the industry.

On the Eastern Bering Sea shelf, an estimate (based on observer at-sea data) of the proportion of pollock caught in the SCA has averaged about 38% annually. During the "A-season," the average is about 49% (since pollock are more concentrated in this area during this period). The proportion of pollock caught within the SCA varies considerably, presumably due to temperature regimes and population age structure. Since 2005 the proportion of catch within the SCA has dropped considerably with about 30% of the catch taken in this area. However, the proportion taken in the A-season reached 57% in 2007, the highest level since 1999:

| <b>Year</b> | <b>Months</b> | <b>Catch<br/>outside SCA</b> | <b>Catch<br/>Total</b> | <b>Percent catch<br/>inside SCA</b> |
|-------------|---------------|------------------------------|------------------------|-------------------------------------|
| 1998        | Jan-Jun       | 71                           | 385                    | 82%                                 |
|             | Jul-Dec       | 248                          | 403                    | 38%                                 |
|             | Jan-Dec       | 318                          | 788                    | 60%                                 |
| 1999        | Jan-Jun       | 155                          | 339                    | 54%                                 |
|             | Jul-Dec       | 360                          | 468                    | 23%                                 |
|             | Jan-Dec       | 515                          | 807                    | 36%                                 |
| 2000        | Jan-Jun       | 241                          | 375                    | 36%                                 |
|             | Jul-Dec       | 550                          | 572                    | 4%                                  |
|             | Jan-Dec       | 791                          | 947                    | 16%                                 |
| 2001        | Jan-Jun       | 357                          | 490                    | 27%                                 |
|             | Jul-Dec       | 367                          | 674                    | 46%                                 |
|             | Jan-Dec       | 724                          | 1,164                  | 38%                                 |
| 2002        | Jan-Jun       | 263                          | 566                    | 54%                                 |
|             | Jul-Dec       | 350                          | 690                    | 49%                                 |
|             | Jan-Dec       | 613                          | 1,256                  | 51%                                 |
| 2003        | Jan-Jun       | 336                          | 616                    | 45%                                 |
|             | Jul-Dec       | 397                          | 680                    | 42%                                 |
|             | Jan-Dec       | 733                          | 1,296                  | 43%                                 |
| 2004        | Jan-Jun       | 293                          | 531                    | 45%                                 |
|             | Jul-Dec       | 472                          | 711                    | 34%                                 |
|             | Jan-Dec       | 765                          | 1,242                  | 38%                                 |
| 2005        | Jan-Jun       | 293                          | 529                    | 45%                                 |
|             | Jul-Dec       | 558                          | 673                    | 17%                                 |
|             | Jan-Dec       | 851                          | 1,203                  | 29%                                 |
| 2006        | Jan-Jun       | 262                          | 533                    | 51%                                 |
|             | Jul-Dec       | 656                          | 764                    | 14%                                 |
|             | Jan-Dec       | 917                          | 1,298                  | 29%                                 |
| 2007        | Jan-Jun       | 209                          | 481                    | 57%                                 |
|             | Jul-Dec       | 540                          | 600                    | 10%                                 |
|             | Jan-Dec       | 749                          | 1,081                  | 31%                                 |

*Note: Pollock catches (thousands of tons) are as reported by at-sea observers only, 2007 data are preliminary.*

An additional goal to minimize potential adverse effects on sea lion populations is to disperse the fishery throughout more of the pollock range on the Eastern Bering Sea shelf. While the distribution of fishing during the A season is limited due to ice and weather conditions, there appears to be some dispersion to the northwest area (Fig. 1.2).

The fishery in recent years has undertaken measures to reduce bycatch of salmon. Recent bycatch levels for Chinook and chum salmon have been very high due in part to large runs of salmon and in part to restrictions on areas where pollock fishing may occur. Bycatch levels for chum salmon in 2005 were the highest on record but declined substantially in 2006 and remain low in 2007 to date. Bycatch for Chinook salmon, however, remains at high levels with bycatch to date in 2007 the highest on record. Given information indicating that large scale regulatory closures were potentially exacerbating the bycatch of these species, the Council acted and developed an extensive analysis leading to amendment 84 of the FMP to a regulatory exemption for vessels participating in a voluntary rolling hot spot (VRHS) closure system. This system is thought to be more responsive and dynamic to changing conditions in the fishery compared to static area closures. Additional salmon bycatch management measures including new regulatory closures and caps on the pollock fishery are currently under consideration by the Council.

## Catch data

Since 2001, the total allowable catch (TAC) for EBS pollock has been at record levels over 1.39 million t. This is roughly 22% above the average levels of catch from 1977-2004 (1.15 million t; Table 1.2).

Significant quantities of pollock are discarded and must be taken into account in estimation of population size and forecasts of yield. Observer length frequency observations indicated that discards include both large and small pollock. Since observers usually sample the catch prior to discarding, the size distribution of pollock sampled closely reflects that of the actual *total* catch. Discard data as compiled by the NMFS Alaska Regional Office have been included in estimates of total catch since 1990.

Pollock catch in the Eastern Bering Sea and Aleutian Islands by area from observer estimates of retained and discarded catch for 1991-2007 are shown in Table 1.3. Since 1991, estimates of discarded pollock have ranged from a high of 9.1% of total pollock catch in 1992 to recent lows of around 1%. These low values reflect the implementation of the Council's Improved Utilization and Improved Retention program. Discard levels are likely affected by the age-structure and relative abundance of the available population. For example, if the most abundant year class in the population is below marketable size, these smaller fish may be caught incidentally. With the implementation of the AFA in 1999, the fleets have more time to pursue the sizes of fish they desire since they are guaranteed a fraction of the quota. In addition, several vessels have made gear modifications to avoid retention of smaller pollock. In all cases, the magnitude of discards is accounted for within the population assessment and for management (to ensure the TAC is not exceeded). Presentation of bycatch of other non-target, target and prohibited species is presented in the section titled "Ecosystem Considerations" below.

The catch-at-age composition was estimated using the methods described by Kimura (1989) and modified by Dorn (1992). Briefly, length-stratified age data are used to construct age-length keys for each stratum and sex. These keys are then applied to randomly sampled catch length frequency data. The stratum-specific age composition estimates are then weighted by the catch within each stratum to arrive at an overall age composition for each year. Data were collected through shore-side sampling and at-sea observers. The three strata for the EBS were: *i*) January–June (all areas, but mainly east of 170°W); *ii*) INPFC area 51 (east of 170°W) from July–December; and *iii*) INPFC area 52 (west of 170°W) from July–December. This method was used to derive the age compositions from 1991-2004 (the period for which all the necessary information is readily available). Prior to 1991, we used the same catch - age composition estimates as presented in Weststad *et al.* (1996).

Recently a comprehensive development of statistical estimators for catch (including length- and age-specific quantities for groundfish fisheries was completed (Miller 2005). Estimators presented in this study hold promise for use in stock assessment purposes since rigorously developed variance estimates are currently unavailable. For the analyses completed on EBS pollock, the estimated variance of the total catch is consistent with the assumption used in the assessment model. Also, the values on total pollock catch estimated by NMFS staff at the Regional Office appear to be unbiased (based on limited comparisons). The coefficient of variation of total catch is specified to be 3% for this stock assessment. This value is a bit higher than the ~1% CVs estimated by Miller (2005) for pollock in the EBS.

In 2006, all age-composition estimates were re-evaluated. This year, the catch-age estimation method was further modified to allow bootstrap re-sampling of the data. This involved a two stage process whereby observed tows were first selected with replacement, followed by re-sampling actual lengths and age-data specimens. The results and application of this approach is presented in attachment A1. Most significantly, this provides a method to more objectively specify the effective sample size for fitting fishery age composition data within the assessment model. An added benefit of this approach was to carefully evaluate general patterns in growth and growth variability. As part of this analysis, the seasonal aspect of pollock condition factor was evaluated from the fishery data from 1991-2006. This involved simply standardizing monthly mean weights given length. Results show that within a year, the condition factor for pollock varies by more than 15% with the "fattest" pollock caught late in the year, from

October-December (although most fishing occurs during other times of the year) and the thinnest fish at length tend to occur in late winter (Fig. 1.8). Relative to recent information, the mean body weight (mass, kg) from the 2006 fishery is considerably lower than the long term mean and the 3-year average (2003-2005) used in last year's assessment (Fig. 1.9 lower panel). A similar decline has been seen in the mean length at age observed in the fishery during 2004-2006 (Fig. 1.9, upper panel). These patterns are partially due to the fact that the fishery shifted more towards the northwest where pollock tend to be smaller at age. Interestingly, the bottom trawl survey mean weight-at-age data are relatively stable; this lends some support to the hypothesis that the pattern seen in the fishery data is a function of geographic distribution.

The recent fishery age ranges appear to be focused primarily on pollock age 4-7 with the 2000 year class making up the majority of the catch until 2006 where the relative fraction of this year class drops considerably (Fig. 1.10). The 2006 fishery data show higher levels (proportionately) of the 2001 year class than in previous years. The corresponding values of catch-at-age used in the model are presented in Table 1.4.

Since 1999 the observer program adopted a new sampling strategy for lengths and age-determination studies (Barbeaux et al. 2005a). Under this scheme, more observers collect otoliths from a greater number of hauls (but far fewer specimens per haul). This has improved the geographic coverage but lowered the total number of otoliths collected. Previously, large numbers were collected but most were not aged. The sampling effort for lengths has decreased since 1999 but the number of otoliths processed for age-determinations increased (Tables 1.5 and 1.6). The sampling effort for pollock catch, length, and age samples by area has been shown to be relatively proportional (e.g., Fig. 1.8 in Ianelli et al. 2004).

## Resource surveys

Scientific research catches are reported to fulfill requirements of the Magnuson-Stevens Fisheries Conservation and Management Act. The annual research catches (1963 - 2007) from NMFS surveys in the Bering Sea and Aleutian Islands Region is given in Table 1.7. Since these values represent extremely small fractions of the total removals (~0.02%), they are not explicitly added to the total removals by the fishery.

### Bottom trawl surveys

Trawl surveys have been conducted annually by the AFSC to assess the abundance of crab and groundfish in the Eastern Bering Sea. Bottom trawl surveys are considered to assess pollock from the bottom to 3 m off bottom. Until 1975 the survey only covered a small portion of the pollock range. In 1975 and since 1979, the survey was expanded to encompass more of the EBS shelf occupied by pollock. The level of sampling for lengths and ages in the BTS is shown in Table 1.8.

Between 1983 and 1990 the BTS biomass estimates were relatively high and showed a slightly increasing trend (Table 1.9; Fig. 1.11). Between 1991 and 2005 the BTS biomass estimates ranged from 2.21 to 8.14 million t. The 2007 estimate is 4.33 million t, up from the 2.85 from 2006 survey but below the 2005 estimate of 5.13 million. This survey estimate is about 87% of the average of all BTS estimates since 1982 and reflects the apparent general decline in the stock since 2003. In 2007 the distribution of pollock from the BTS continued to show higher concentrations in the northwestern area of the surveys and closer to the shelf break compared to the 2005 pattern but also had some higher CPUE stations near Unimak Island in the south-east area of the survey (Fig. 1.12).

In general, the interannual variability of survey estimates is due to the effect of year class variability. Survey abundance-at-age estimates reflect the impact of this variability (Fig. 1.13). The survey operations generally catch pollock above 40 cm in length, and in some years lots of 1-year olds (with modal lengths around 10-15 cm; Fig. 1.14). Other sources of variability may be due to unaccounted-for variability in natural mortality and migration. For example, some strong year classes appear in the



surveys over several ages (e.g., the 1989 year class) while others appear at older ages (e.g., the 1992 year class). More recently, the estimate of the strength of the 1996 year class has waned compared to previous assessments. In 2003 the point estimate for this year class was 43 billion one-year olds whereas for the past two assessments, the estimate from the model has been around 32 billion. This could be due in part to emigration of this year-class outside of our main fishery and survey zones. Alternatively, this may reflect the effect of variable natural mortality rates. In the retrospective analyses presented in a last year's report (Ianelli et al. 2006), the characteristics of how strong year-class estimates change over time were illustrated.

Beginning in 1987 NMFS expanded the standard survey area farther to the northwest. For consistency, these extra strata (8 and 9) had traditionally been excluded for consideration within the model. The pollock biomass levels found in these non-standard regions were highly variable, ranging from 1% to 22% of the total biomass, and averaging about 6% (Table 1.10). Closer examination of the years where significant concentrations of pollock were found (1997 and 1998) revealed some stations with high catches of pollock. The variance estimates for these northwest strata were quite high in those years (CVs of 95% and 65% for 1997 and 1998 respectively). Nonetheless, since this region is contiguous with the Russian border, including these strata was considered important and better covered the range of the exploited stock of pollock. The use of the additional strata was evaluated in 2006 and accepted as appropriate by the Council's SSC. The estimated numbers-at-age from the BTS for the standard strata (1-6) and for the northern strata included are presented in Table 1.11.

In response to the SSC's request, an evaluation of body mass changes for juvenile pollock shows that there is a large degree of variability over time, particularly for 3-year old pollock (Fig. 1.15, top panel). As observed in the fishery, the mean weight is low in 2006 which may reflect feeding conditions. However, since the timing of surveys has varied (Ianelli et al. 2006), consideration on the mean date (allowing for growth) and temperature may be important (Fig. 1.15, bottom panel). Regarding conditions in 2007, it appears that the mean body weight at age is closer to or above recent averages and is unlike the 2006 observations (Fig. 1.16). This provides some justification to use a mean value for the 2007 fishery as opposed to using the 2006 estimates.

As in the past few assessments, an analysis using survey data alone was conducted to evaluate mortality patterns. Cotter et al. (2004) promote this type of analysis as having a simple and intuitive appeal which is independent of population scale. In this approach, log-abundance of age 6 and older pollock is regressed against age by cohort. The negative values estimated for the slope are estimates of total annual mortality. Age-6 was selected because younger pollock are still recruiting to the bottom trawl survey gear. A key assumption of this analysis is that all ages are equally available to the gear. Total mortality by cohort seems to be variable (unlike the example in Cotter et al., 2004) with lower mortality overall for cohorts during the 1990s followed by recent increases (Fig. 1.17). Total mortality estimates by cohort represent lifetime averages since harvest rates (and actual natural mortality) vary from year to year. The low values estimated from some year classes (e.g., the 1990-1992 cohorts) could be because these age groups have only recently become available to the survey (i.e., that the availability/selectivity to the survey gear has changed for these cohorts). Alternatively, it may suggest some net immigration into the survey area or a period of lower natural mortality. In general, these values are consistent with the types of values obtained from within the assessment models for total mortality (although the model values tend to be somewhat higher, averaging about 0.5 for recent cohorts).

For the past several years the effect of bottom temperature on pollock habitat relative to the standard survey area has been evaluated. Modeling survey availability as a function of temperature helps account for the observation that environmental conditions affect the distribution of pollock. Previously, temperature was shown to affect the proportion of the stock that lies within or outside of the standard survey area (or extended area, as the case may be). This summer, the bottom trawl survey data revealed a cold pool similar to that observed last year and in 1999 (Fig. 1.18). As noted last year, the timing of the survey can affect the estimate of "cold" years since surveys that occur earlier in the summer (such as in

1999) tend to exhibit colder temperatures on average. Last year a refinement to using mean temperature relative to survey catchability was attempted. This involved computing the surface area between two isotherms within the shelf region thought to cover the desired habitat and using that measure to modify catchability/availability of the survey index. This measure, as well as others relating mean temperatures to survey catchability have been poorly correlated and were omitted in this assessment. Spatially, a relationship between temperature and pollock density appears to exist. However, it could be that the variability (e.g., patchiness) of the pollock increases during colder conditions which may mask relationships between temperature effects on the pollock habitat relative to the fixed survey area. Planned work on this is to use station-by-station data within the model so that individual survey trawl operations can be modeled within the stock assessment. This feature has been implemented in the assessment model but at the time of writing, requires more testing.

### **Echo-integration trawl (EIT) surveys**

The EIT surveys are conducted biennially and are designed to estimate the off-bottom component of the pollock stock (compared to the BTS which are conducted annually and provide an abundance index of the near-bottom pollock).

In summer 2004 NMFS conducted an EIT survey that extended into the Russian zone (Honkalehto et al. 2005). The biomass estimate from this survey was 3.31 million t (U.S. zone only), down from 3.62 million t estimated in 2002 but close to the average estimated by this survey since 1982 (3.36 million t; Table 1.9).

The 2007 echo-integration trawl (EIT) survey also extended into the Russian zone. The U.S. EEZ biomass estimate was low (1.88 million t) and represents only 55% of the long-term mean value from this survey (since 1979; Table 1.9). However, this is a modest improvement over the 2006 estimate of 1.56 million t. The geographic concentration and extent of pollock from the 2007 survey shows highest concentrations in the northwestern region (Fig. 1.19) with 5% of the total survey-wide biomass occurring in the Russian EEZ. Size composition in the Russian zone was similar to that observed in the U.S. EEZ west of 170° W. Also, these figures show that the EIT survey appears to detect pollock concentrations in areas where they were less abundant in the bottom-trawl survey.

The number of trawl hauls and sampling quantities for lengths and ages from the EIT survey are presented in Table 1.12. Last year, to estimate the 2006 EIT survey population numbers at age estimates geographically split age-length keys (E and W of 170°W) were used. These keys were developed from the 2006 BTS study since there was insufficient time to conduct the age-determinations from samples collected on this research cruise. These data have been revised using age data from EIT samples that have been processed in the last year. Comparing results using one or two age-length keys suggests that one age length key from the bottom-trawl survey more closely approximated the age composition from the EIT survey (Fig. 1.20). Hence, for the 2007 EIT survey, the age composition was generated using a single age-length key from the bottom-trawl survey (Table 1.13 and also Fig. 1.21).

Proportions of pollock biomass estimated east vs. west of 170° W, and inside vs. outside the SCA, are about the same for summer EIT surveys conducted from 1994 to 2007 (Table 1.14). Compared to 2004, the relative abundance of pollock in 2006 and 2007 was much lower overall with the biggest difference being the relative lack of pollock in the region east of 170°W.

## **Analytic approach**

### **Model structure**

A statistical age-structured assessment model conceptually outlined in Fournier and Archibald (1982) and similar to Methot's (1990) extensions was applied over the period 1964-2007. A technical description is presented in the "Model Details" section. The analysis was first introduced in the 1996 SAFE report

(Ianelli 1996) and compared to the cohort analyses that had been used previously. The current model also was documented in the Academy of Sciences National Research Council (Ianelli and Fournier 1998). The model was implemented using automatic differentiation software developed as a set of libraries under the C++ language (AD Model Builder).

The main changes from last year's analyses include:

*New data:*

- The 2007 EBS bottom trawl survey estimate of population numbers-at-age was added.
- The 2007 EBS EIT survey estimate of population numbers-at-age were included using age-length keys from the 2007 BTS survey data.
- The 2006 EBS EIT survey estimate of population numbers-at-age were updated from last year's values by using age-length keys from the 2006 EIT survey data.
- The 2006 fishery age composition data were added.
- Length frequency data from the 2007 fishery was incorporated (and growth estimates to use in tuning the model). This is the first time length frequency data have been applied directly within the EBS pollock model.

*Changes to the model:*

- Multi-year forecast projections to facilitate Tier 1 OFL and ABC determinations. For the purpose of these projections, catch in 2008 (for deriving the 2009 ABC and OFL) was assumed to be equal to the maximum permissible Tier 1b level (1,170 t).
- The growth transition matrix (based on 2006 fishery data) was used to estimate the dispersion of pollock lengths given age (Fig. 1.22)
- Age 1 data from the EIT survey were treated as a separate recruitment index. Previously, these were treated as part of the multinomial age-sampling process.
- The frequency of fishery selectivity change was modified to occur every two years instead of in 3-year blocks.
- Flexibility was added to allow more extensive evaluation of survey catchabilities (past models used surveys exclusively as relative indices).
- The facility to use alternative mean-weights-at-age for spawning biomass calculations were added (in the past, fishery values were used).
- The ability to apply uncertainty in future mean weights-at-age was developed.
- The Tier 1 ABC estimation method now uses "fishable" biomass defined as the biomass of pollock available to the fishery as modified by the selectivity-at-age estimates. Previously age 3+ biomass had been used. This was changed because based on evaluation, the knife-edge aspect of age 3+ biomass unnecessarily increased the variability of yield at  $F_{msyr}$ .

Also, alternative output values for diagnostic purposes added last year include a "replay" of the estimated time series of spawning biomass and age 3+ biomass given recruitments as estimated and omitting the fishing mortality component and the projection aspect of the model was modified to more easily accommodate Tier 1, 2-year and beyond forecasts for ABC and OFL levels.

## **Parameters estimated independently**

*Natural mortality and maturity at age*

For the reference model fixed natural mortality-at-age were assumed ( $M=0.9$ ,  $0.45$ , and  $0.3$  for ages 1, 2, and 3+ respectively; Weststad and Terry 1984). These values have been applied to catch-age models and forecasts since 1982 and appear reasonable for pollock. Estimates of natural mortality are higher when predation (e.g., when consumption by Steller sea lions and Pacific cod) are explicitly considered

(Livingston and Methot 1998; Hollowed et al. 2000). The reference model values were selected because Clark (1999) found that specifying a conservative (lower) natural mortality rate is typically more precautionary when natural mortality rates are uncertain.

Pollock maturity-at-age (Smith 1981) values (tabulated with reference model values for natural mortality-at-age) are:

| Age          | 1     | 2     | 3     | 4     | 5     | 6     | 7     |
|--------------|-------|-------|-------|-------|-------|-------|-------|
| M            | 0.900 | 0.450 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 |
| Prop. Mature | 0.000 | 0.008 | 0.290 | 0.642 | 0.842 | 0.902 | 0.948 |

| Age          | 8     | 9     | 10    | 11    | 12    | 13    | 14    | 15    |
|--------------|-------|-------|-------|-------|-------|-------|-------|-------|
| M            | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 |
| Prop. Mature | 0.964 | 0.970 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |

These maturity-at-age values were reevaluated based on the studies of Stahl (2004). A total of 10,197 samples of maturity stage and gonad weight were collected during late winter and early spring of 2002 and 2003 from 16 different vessels. In addition, 173 samples were collected for histological determination of maturity state (Stahl 2004). In her study, maturity-at-length converted to maturity-at-age via a fishery-derived age-length key from the same seasons and areas suggest similar results to the maturity-at-age schedule used in this assessment but with some inter-annual variability.

Ianelli et al. (2005) investigated the inter-annual variability found by Stahl (2004). This involved using the fixed maturity-at-age levels presented above (for the reference model) to get estimates of total mature and immature numbers at age and then converting those to values at length using female mean-lengths at age (with an assumed natural variability about these means). Expected proportion mature-at-length for 2002 matched Stahl's data whereas for 2003, the model's expected values for maturity-at-length were shifted towards larger pollock. This result suggests that younger-than-currently-assumed pollock may contribute to the spawning stock.

#### *Length and Weight at Age*

Extensive length, weight, and age data have been collected and show that growth may differ by sex, area, and year class. Pollock in the northwest area typically are smaller at age than pollock in the southeast area. The differences in average weight-at-age are taken into account by stratifying estimates of catch-at-age by year, area, season and weighting estimates proportional to catch (Table 1.15).

#### **Parameters estimated conditionally**

A total of 544 parameters were estimated conditioned on data and model assumptions. Initial age composition, subsequent recruitment values and stock-recruitment parameters account for 67 parameters. This includes vectors describing mean recruitment and variability for the first year (as ages 2-15 in 1964, projected forward from 1949) and the recruitment mean and deviations (at age 1) from 1964-2007 and projected recruitment variability (using the variance of past recruitments) for five years (2008-2012). The two-parameter stock-recruitment curve is included in addition to a term that allows the average recruitment before 1964 (that comprises the initial age composition in that year) to have a mean value different from subsequent years.

Fishing mortality is parameterized to be semi-separable with year and age (selectivity) components. The age component is allowed to vary over time; changes are allowed every two years (previously this was done at three-year intervals). The two most recent years (2006-2007) forming the last "group" of estimates. The mean value of the age component is constrained to equal one and the last 4 age groups (ages 12-15) are specified to be equal. The annual components of fishing mortality result in 45 parameters and the age-time forms an 11x22 matrix of 242 parameters bringing the total fishing mortality parameters to 287.

Selectivity-at-age estimates for the bottom trawl survey are specified with age and year specific deviations in the average availability-at-age and a catchability coefficient totaling 81 parameters. For the EIT survey, which began in 1979, 105 parameters are used to specify age-time specific availability. Unlike in previous versions of this model, the estimates in EIT selectivity occur only in years when the survey was conducted. Time-varying survey selectivity is estimated to account for the changes in availability of pollock to the survey gear. Four catchability coefficients were estimated; one each for the early CPUE data, the early bottom trawl survey data (where only 6 strata were surveyed), the main bottom trawl survey data, and the EIT survey data.

For all models presented this year, catchability coefficients for are estimated as independent indices of abundance. In the past, these values have been reported but without discussion on the implications. This year, based on the work of Von Szalay et al. (2007) we introduce prior distributions on the sum of the EIT and BTS catchability coefficients. If the BTS survey covers the bottom-dwelling pollock (up to ~3 m above the bottom) and the EIT survey covers the remainder of the water column, then the catchabilities from both surveys should sum to unity. Values of this sum that are less than one imply that there are spatial aspects of the pollock stock that are missed whereas values greater than one imply that there are pollock on the shelf during the summer that could be considered as “visitors” perhaps originating (and returning to) other areas such as the Russian zone.

Additional fishing mortality rates used for recommending harvest levels are estimated conditionally on other outputs from the model. For example, the values corresponding to the  $F_{40\%}$ ,  $F_{35\%}$  and  $F_{msy}$  harvest rates are found by satisfying the constraint that given age specific population parameters (e.g., selectivity, maturity, mortality, weight-at-age), unique values exist that correspond to these fishing mortality rates.

The likelihood components can thus be partitioned into the following groups:

- Total catch biomass (Log normal,  $\sigma=0.05$ )
- Log-normal indices of abundance (bottom trawl surveys assume annual estimates sampling error, as represented in Fig. 1.11; for the EIT and CPUE indices values of  $\sigma=0.2$  were assumed)
- Fishery and survey proportions-at-age estimates (robust quasi-multinomial with effective sample sizes presented in Table 1.16).
- Age 1 index from the EIT survey
- Selectivity constraints: penalties/priors on age-age variability, time changes, and decreasing (with age) patterns
- Stock-recruitment: penalties/priors involved with fitting a stochastic stock-recruitment relationship within the integrated model.

## Model evaluation

### Last year’s assessment with data and assumption revisions

The developments in data revisions and model configurations introduced from last year's model were evaluated in a stepwise fashion. This was done to reveal how revised historical data impact subsequent analysis. The revisions to the 2006 assessment and data were as follows: 1) revised EIT age data; 2) revised mean weights used in the 2006 fishery (and subsequently used for ABC projections; 3) fishery age-composition sample size revisions based on bootstrap analysis in presented in attachment 1A; 4) accounting for weight-at-age variability as an explicit part of the uncertainty for  $F_{msy}$  calculations (see section on Model Details for an explanation); and 5) a model run with all revisions indicated in 1) - 4). Note that 4) is equivalent to using the long-term mean weight-at-age (as opposed to the 3-year average).

Results from this exercise showed that the largest impact (using exactly the same model code from last year's assessment) was from the revised mean weights-at-age (Fig. 1.23). This is because these values were much lower than the mean value from 2003-2005 used for 2006 in last year's assessment.

To gain some appreciation for the impact of mean weight-at-age differences, equilibrium stock conditions using mean age-specific survival from 1997-2006 were calculated to obtain age 3+ biomass values using last year's mean weight-at-age estimates and then again with the revised estimates of mean weights-at-age observed in 2006. This resulted in a 12% reduction in age 3+ biomass due to changes in mean body weight alone. However, for last year's 2006 estimated numbers at age the reduction is only 7% due to the age distribution of the population.

This impacts the assessment in two important ways. First, the biomass levels will be lower for Tier 1 calculations. Secondly, the mean weights at age used for the quota are based on the most recent information (which the SSC has accepted as the most recent three-years of data).

### Evaluation of the 2007 assessment model

Going forward with the addition of 2007 data, all historical data revisions from the previous section were included with one important exception: the assumed 2007 mean weights-at-age were set to the average observed during 2004-2006. This is significantly lower than what was assumed in last year's model for 2006, but isn't as low as the presently observed 2006 fishery mean weights at age. The survey data and evaluation of existing length-weight data for 2007 indicate that growth patterns and fish size have improved over the 2006 condition.

Following a similar approach to evaluating the influence of added data, models were constructed with combinations of including or excluding new data as follows:

|         | Shorthand   | Description                                       |
|---------|-------------|---|
| Model_1 | <b>C</b>    | 2007 total catch only included                    |
| Model_2 | <b>CA</b>   | Catch and 2006 fishery age data included          |
| Model_3 | <b>CB</b>   | Catch, and 2007 bottom-trawl survey data included |
| Model_4 | <b>CE</b>   | Catch and 2007 EIT survey included                |
| Model_5 | <b>CAB</b>  | Catch, age, and bottom-trawl survey               |
| Model_6 | <b>CAE</b>  | Catch, age, and EIT survey                        |
| Model_7 | <b>CBE</b>  | Catch, bottom-trawl survey, and EIT survey        |
| Model_8 | <b>CABE</b> | Catch, age, bottom-trawl survey, and EIT survey   |

In last year's assessment an extensive retrospective analysis was conducted. This work concluded that in general, there was a tendency to under-estimate terminal year biomass levels even though the estimated uncertainty is quite high. Also, it appears that some strong year-class estimates evolve quite differently as additional data are added in each year suggesting unaccounted for process errors.

## Results

Evaluating the influence of new data as they are introduced can reveal where consistencies with past predictions occur and where things may diverge. Adding the fishery catch-age data and other data affects the point estimates of the age 3+ biomass but reduces the uncertainty on the 2005 year class (Fig. 1.24). Closer examination of the age data that impact these results show how model CA (only new 2007 data include fishery catch and age compositions) results in particularly poor fits to the observed 2007 survey age compositions (Fig. 1.25). Similarly, if the 2006 fishery age composition data is omitted, (model C) the fit to the 2006 fishery age composition is also poor due to the lack of many age 6 pollock in the 2006 fishery (Fig. 1.26). This indicates that last year's model anticipated that the 2006 fishery would still have a strong showing of the 2000 year class. The new data contradict this prediction.

A variety of alternative model configurations have been explored in the past, and continued again this year. Alternative non-informative prior distributions on the aggregate “catchability” of the EIT and bottom trawl survey were examined. In particular, Von Szalay et al.’s (2007) results suggest that vertical herding of pollock (i.e., fish diving toward bottom and becoming vulnerable) was limited. This indicates that a rationale for having the combined catchability be closer to unity than the current estimated value of 1.54. However, alternatives lower than this number degraded the fit to the data substantially and represents a major departure from past assessments. Highlighting this fact does provide some added level of precaution since imposing an informative prior on the combined survey catchabilities to lower values would scale population to higher levels. A profile over this quantity indicates that the value is quite well determined and unlikely (given data and model structure assumptions) to be below a value of 1.3—which would increase the current stock-size estimated by 30% (Fig. 1.27). Thus for clarity and consistency, the model presented below represents that accepted in previous years with the modifications mentioned above. This model was used with all new data included (CABE) as a reasonable representation of stock status and associated uncertainty.

Comparing the current and projected age structure for this model relative to last year’s assessment indicates that the abundance estimate of 2008 numbers at age is substantially lower, particularly for the 2005 and 2004 year classes (3 and 4 year olds; Fig. 1.28, top panel). When summed by age for biomass, the difference relative to last year’s assessment is more dramatic indicating that the nominal age 3+ begin-year biomass is about 65% of last year’s estimate for 2008 (Fig. 1.28; bottom panel). Part of this difference is due to the difference in mean weights-at-age used this year.

The estimated selectivity pattern changes over time and reflects to some degree the extent that the fishery is focused on particularly prominent year-classes (Fig. 1.29). As noted above, the 2005 fishery apparently caught a proportionately large amount of 5 year olds but these failed to show in substantial numbers in the 2006 fishery. Part of this may be due to the increased level of fishing in the northwest region during the summer/fall season. The model fits the fishery age-composition data quite well under this form of selectivity (Fig. 1.30). The length frequency data for 2007 also fit reasonably well. The model fit to the early Japanese fishery CPUE data (Low and Ikeda, 1980) is consistent with the population trends for this period and is essentially unchanged since introduced to the assessment several years ago.

Bottom-trawl survey selectivity and fits to the numbers of age 2 and older pollock are shown in Fig. 1.31. The bottom trawl survey age composition data continue to indicate that the 2006 year class is large but that the 2005 and 2000 year-classes are less abundant than in recent years. (Fig. 1.32).

The EIT survey selectivity shows some inter-annual variability but has generally stabilized since the early 1990s as the echo-sounder and trawl methods became more standardized (Fig. 1.33; top panel). Of course this could also be due in part to changes in age-specific pollock distributions over time. The fit to the numbers of age 2 and older pollock in the EIT survey generally falls within the confidence bounds of the survey sampling distributions (here assumed to have a CV of 20%) with a fairly reasonable pattern of residuals (Fig. 1.33; bottom panel). As with the fishery and bottom trawl survey age composition data, the EIT age compositions consistently track large year classes through the population and the model fits these patterns reasonably well (Fig. 1.34). The EIT age-1 index component which was split from the age composition data this year, demonstrates the difficulty in having a highly precise pre-recruit index (Fig. 1.34; bottom panel).

The estimate of 2008 spawning stock size and corresponding estimates of  $B_{msy}$  have coefficients of variation that exceed 20% (Table 1.17). For 2008, the Tier 1 levels of yield are 1,170 thousand t from a fishable biomass estimated at around 4.77 million t (Table 1.18). Estimated numbers-at-age are presented in Table 1.19 and estimated catch-at-age presented in Table 1.20. Estimated summary biomass (age 3+), female spawning biomass, and age-1 recruitment is given in Table 1.21.

In addition to extensive model-specification evaluations, the selected model was used to examine the multidimensional parameter uncertainty by integrating approximations to the posterior distribution using

Monte-Carlo Markov Chain methods. This involved generating several million simulations drawn from the posterior distribution with the individual draws “weighted” by their density. This chain was then “thinned” to reduce potential serial correlation to 8,000 parameter vectors. Selected model parameters were then plotted pair-wise to provide some indication of the shape of the posterior distribution. In general, the model given the available data appears to be quite well behaved (no curved or skewed tear-drop shapes over critical parameters; Fig. 1.35). For a closer evaluation of how parameter uncertainty manifests in the posterior distribution, a subset of stock-recruitment “curves” from the posterior shows a broad range of values and shapes (Fig. 1.36).

The assessment results indicate that the spawning biomass will be well below  $B_{40\%}$  in 2008 (53% of the value) and about 72% of the  $B_{msy}$  level but only 49% of the predicted value had no fishing occurred since 1978 (Table 1.17). This compares with the 21% of  $B_{100\%}$  (based on the SPR expansion from mean recruitment since 1978) and 28% of  $B_0$  (as from the estimated stock-recruitment curve). This range of alternative “reference point” systems serves to illustrate problems associated with developing robust guidelines for management.

### **Abundance and exploitation trends**

The current mid-year biomass estimates (ages 3 and older) derived from the statistical catch-age model suggest that the abundance of Eastern Bering Sea pollock remained at a fairly high level from 1982-88, with estimates ranging from 8 to 10 million t (Table 1.22, Fig. 1.37). Historically, biomass levels have increased from 1979 to the mid-1980's due to the strong 1978 and relatively strong 1982 and 1984 year classes recruiting to the fishable population. The stock is characterized by peaks in the mid 1980s and mid 1990s with a substantial decline to about 4 million t by 1991 and another low point occurring at present with the stock projected to drop to the lowest levels since the late 1970s\*. As predicted in last year's assessment, the stock has continued to decline substantially since 2003 due to apparently poor recruitment between 2000 and 2006.

Compared with past year's assessments, the estimates of age 3+ pollock biomass are significantly lower in the current assessment for recent years but are nearly identical before 2002 (Table 1.22). Overall, compared with the past several assessments, the pattern appears to be balanced between over and under estimates, especially since 1998 when the same statistical model has been used (Fig. 1.38).

The abundance and exploitation pattern estimates show that the spawning exploitation rate (SER, defined as the percent removal of spawning-aged females in any given year) has been below 20% since 1980 until 2006 and 2007 where the rate has averaged more than 25% (Fig. 1.39).

One way to evaluate past management and assessment performance is to plot estimated fishing mortality relative to some reference values. For EBS pollock, we computed the reference fishing mortality as from Tier 1 (unadjusted) and calculated the historical values for  $F_{msy}$  (since selectivity has changed over time). Since 1977 the current estimates of fishing mortality suggest that during the early period, harvest rates were above  $F_{msy}$  until about 1981. Since that time, the levels of fishing mortality have averaged about 44% of  $F_{40\%}$  (Fig. 1.40).

### **Recruitment**

In this year's assessment, the 2000 year class abundance level dropped compared to last year though is still above average. The 2005 year-class, estimated as “about” average last year, also was estimated to be significantly below average. The 2006 year class appears to be well above average and based on survey data (Fig. 1.41, top panel). New data from the BASIS surveys suggests that the abundance of age-0 pollock (near the surface) was low in 2006 and 2007, with high levels observed in 2004 and 2005. This is

---

\* Please refer to Ianelli et al. (2001) for a discussion on the interpretation of age-3+ biomass estimates.



somewhat inconsistent with the relatively abundant 2006 year-class observed in the surveys this year. Further investigations on the water column temperature structure may shed light on the relative availability of pollock young-of-year to the BASIS sampling gear.

The high degree of uncertainty in the magnitude of these year classes can not be overemphasized, particularly as they extend to estimates of future stock size. Management will continue to rely on survey and fishery information to provide reasonable short-term forecasts on stock conditions. The stock-recruitment curve fit within the integrated model shows a fair amount of variability both in the estimated recruitments and in the uncertainty of the curve and also illustrates that the estimate of the 2008 spawning biomass is below the  $B_{msy}$  level (Fig. 1.41, bottom panel).

## Projections and harvest alternatives

### Amendment 56 Reference Points

Amendment 56 to the BSAI Groundfish Fishery Management Plan (FMP) defines “overfishing level” (OFL), the fishing mortality rate used to set OFL ( $F_{OFL}$ ), the maximum permissible ABC, and the fishing mortality rate used to set the maximum permissible ABC. The fishing mortality rate used to set ABC ( $F_{ABC}$ ) may be less than this maximum permissible level, but not greater. Estimates of reference points related to maximum sustainable yield (MSY) are currently available. However, their reliability is questionable. We therefore present both reference points for pollock in the BSAI to retain the option for classification in either Tier 1 or Tier 3 of Amendment 56. These Tiers require reference point estimates for biomass level determinations. For our analyses, the following values from are presented based on recruitment estimates from post-1976 spawning events:

$$\begin{aligned}
 B_{100\%} &= 6,569 \text{ thousand t female spawning biomass}^* \\
 B_{40\%} &= \mathbf{2,627} \text{ thousand t female spawning biomass} \\
 B_{35\%} &= 2,299 \text{ thousand t female spawning biomass} \\
 B_{msy} &= \mathbf{1,876} \text{ thousand t female spawning biomass}
 \end{aligned}$$

### Specification of OFL and Maximum Permissible ABC

The year 2008 spawning biomass is estimated to be **1,380** thousand tons (at the time of spawning, assuming the stock is fished at Tier 1b level). This is below the  $B_{msy}$  value of **1,876**. Under Amendment 56, this stock has qualified under Tier 1 and the harmonic mean value is considered a risk-averse policy since reliable estimates of  $F_{msy}$  and its pdf are available. The exploitation-rate type value that corresponds to the  $F_{msy}$  level was applied to the “fishable” biomass for computing ABC levels. For a future year, the fishable biomass is defined as the sum over ages of predicted begin-year numbers multiplied by age specific fishery selectivity and body weights.

The 2008 estimate of female spawning biomass (at time of spawning assuming 2008 Tier 1b catch levels of 1.17 million t) is **1,380** thousand t. This is below the  $B_{40\%}$  and  $B_{msy}$  values (**2,627** and **1,876** t, respectively). The OFL’s and maximum permissible ABC values by Tier are thus:

| Tier | Year | Max ABC          | OFL              |
|------|------|------------------|------------------|
| 1b   | 2008 | 1,170 thousand t | 1,443 thousand t |
| 1b   | 2009 | 976 thousand t   | 1,204 thousand t |
| 3b   | 2008 | 555 thousand t   | 677 thousand t   |
| 3b   | 2009 | 650 thousand t   | 794 thousand t   |

\* Note that another theoretical “unfished spawning biomass level” (based on stock-recruitment relationship  $\tilde{B}_0$ ) is somewhat lower (**5,013** t).

## ABC Recommendation

This year's population estimates are significantly lower than last year's estimate which reflects a number of issues raised in the data sections above. The stock has dropped to below average levels after a number of years of above-average conditions. The biomass of Eastern Bering Sea pollock is projected to continue declining until the positive signs of incoming year classes recruit to the spawning biomass. The rate of decline since 2003 has been around 20% per year. The spawning exploitation rate has consequently increased by more than 15% from 2003-2007. Under likely catch projections, the spawning stock biomass is expected to be about 72% of  $B_{msy}$  (1,876 thousand t) by 2008 with future status depending on specified catch levels and recruitment (Fig. 1.42).

Given the rapidly declining stock and the recent increases in harvest rates, it could be argued that it would be prudent to consider harvest levels that would 1) provide stability to the fishery; 2) provide added conservation to an important prey species of the endangered stock of Steller sea lions; and 3) provide extra precaution due to uncertain stock removals in Russian waters. There are (as always) reasons to be concerned that specified harvest levels may cause "something bad to happen." Risk aversion is at the core of the NPFMC harvest guidelines. However, all forms of modeling have major weakness when it comes to predicting outcomes of highly variable and uncertain events. The following highlights some points to consider regarding whether the ABC should be set to the Tier 1b maximum permissible level:

- 1) The sum of the survey catchabilities for the accepted model is over 1.5, indicating that there is considerable overlap between the availability of pollock between the two surveys (or some other mechanism such as temporary immigration of pollock into the EBS region from elsewhere).
- 2) The stock-recruitment relationship continues to be constrained within the model which causes the harvest rate to be more conservative.
- 3) 2007 weights-at-age appear to be closer to mean values and are above the lower levels observed in the 2006 fishery.
- 4) In the two surveys conducted in 2007 signs of 1-year old abundance were above average.
- 5) The precautionary MSY harvest rate has been adjusted downward to nearly 72% to account for the stock being below the  $B_{msy}$  level.
- 6) In the 1998 assessment, the outlook for 1999 was fairly pessimistic (although the age 3+ biomass was about 1 million t higher than is presently estimated for 2008). In hindsight, the perception of relatively poor stock conditions at the time changed. For example, at the time, the 1992 year class was estimated to be about average whereas now it appears to be more than twice the average and represents the third highest year class.
- 7) The stock has been at low levels in the past (but this appears to be the lowest since before 1980).
- 8) The Tier 3 ABC levels are substantially lower than the Tier 1 values due to different assumptions about reference biomass levels (hence a larger adjustment).
- 9) Future selectivity patterns are unpredictable given fish distribution.
- 10) If the 2005 year class is in fact below average, then there will have been 5 year classes in sequence that have been below average, an apparently unique event for this stock.
- 11) Spatially equitable catch rates by the fishery may continue to be impacted (lower catch rates overall are to be expected, and this can manifest as spatial differences in pollock availability).
- 12) Absorbing some of the anticipated "adjustment" from the ABC control rule will likely reduce the inter-annual variability ABC recommendations.

Setting the ABC below the maximum permissible value may provide opportunities to reduce the projected declines. For example, under constant catch scenarios of 1.0 and 1.2 million t the stock is projected to rebound sooner (Fig. 1.43). These projections also suggest that the spawning stock exploitation rates will drop if 1.2 million t are caught in 2008. Alternatively, a value of 1.02 million t is projected to be equal to the 1990 exploitation rate (the highest rate observed since 1978). However, if the

stock was managed under the Tier 3b ( $F_{40\%}$ ) harvest level, the decline in exploitation rate would be greater (Fig. 1.44).

The degree to which the ABC should be adjusted downwards is difficult to quantitatively justify. The maximum permissible ABC under Tier 1b seems too high given the continued decline and the lower abundances of older fish seen in the population in recent years. For stability in catches, it may be best to harvest at 1.0 million t in 2008. Since two more surveys will be conducted, next year's assessment should provide clearer guidance on the status of incoming recruits and adult abundance. Catch rates experienced by the fleet are also likely to drop as pollock stock densities reach lower levels.

### **Standard Harvest Scenarios and Projection Methodology**

A standard set of projections is required for each stock managed under Tiers 1, 2, or 3, of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Policy Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA). While EBS pollock is generally considered to fall within Tier 1, the standard projection model requires knowledge of future uncertainty in  $F_{msy}$ . Projections based on Tier 3 are presented along with some considerations for a Tier 1 approach.

For each scenario, the projections begin with the vector of 2007 numbers at age estimated in the assessment. This vector is then projected forward to the beginning of 2008 using the schedules of natural mortality and selectivity described in the assessment and the best available estimate of total (year-end) catch assumed for 2007. In each subsequent year, the fishing mortality rate is prescribed on the basis of the spawning biomass in that year and the respective harvest scenario. In each year, recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments estimated in the assessment. Spawning biomass is computed in each year based on the time of peak spawning and the maturity and weight schedules described in the assessment. Total catch is assumed to equal the catch associated with the respective harvest scenario in all years. This projection scheme is run 1,000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios will be used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2008 and 2009, are as follows (A " $max F_{ABC}$ " refers to the maximum permissible value of  $F_{ABC}$  under Amendment 56):

- Scenario 1:* In all future years,  $F$  is set equal to  $max F_{ABC}$ . (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs).
- Scenario 2:* In all future years,  $F$  is set equal to a value that corresponds to a constant catch level of 1,200,000 t. (Rationale: This catch is close to the mean catch since 1981 and in most years, would satisfy the constraint to be below the maximum permissible under Tier 1 levels).
- Scenario 3:* In all future years,  $F$  is set equal to the 2003-2007 average  $F$ . (Rationale: For some stocks, TAC can be well below ABC, and recent average  $F$  may provide a better indicator of  $F_{TAC}$  than  $F_{ABC}$ .)
- Scenario 4:* In all future years,  $F$  is set equal to  $F_{60\%}$ . (Rationale: This scenario provides a likely lower bound on  $F_{ABC}$  that still allows future harvest rates to be adjusted downward when stocks fall below reference levels. This was requested by public comment for the DSEIS developed in 2006)
- Scenario 5:* In all future years,  $F$  is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

Two other scenarios are needed to satisfy the MSFCMA's requirement to determine whether a stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follow (for Tier 3 stocks, the MSY level is defined as  $B_{35\%}$ ):

- Scenario 6:* In all future years,  $F$  is set equal to  $F_{OFL}$ . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be 1) above its MSY level in 2007 or 2) above  $\frac{1}{2}$  of its MSY level in 2008 and above its MSY level in 2020 under this scenario, then the stock is not overfished.)
- Scenario 7:* In 2008 and 2009,  $F$  is set equal to  $\max F_{ABC}$ , and in all subsequent years,  $F$  is set equal to  $F_{OFL}$ . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be above its MSY level in 2020 under this scenario, then the stock is not approaching an overfished condition.)

### Projections and status determination

For the purposes of these projections, we present results based on selecting the  $F_{40\%}$  harvest rate as the max  $F_{ABC}$  value and use  $F_{35\%}$  as a proxy for  $F_{msy}$ . Scenarios 1 through 7 were projected 14 years from 2006 (Table 1.23). Under Scenarios 1 and 2, the expected spawning biomass will decrease to below the  $B_{35\%}$  then begin increasing after 2008 but not reaching  $B_{40\%}$  (in expectation) until after 2011 (Fig. 1.42).

Any stock that is below its MSST is defined to be overfished. Any stock that is expected to fall below its MSST in the next two years is defined to be approaching an overfished condition. Harvest scenarios 6 and 7 are used in these determinations as follows:

Is the stock overfished? This depends on the stock's estimated spawning biomass in 2008:

- If spawning biomass for 2008 is estimated to be below  $\frac{1}{2} B_{35\%}$  the stock is below its MSST.
- If spawning biomass for 2008 is estimated to be above  $B_{35\%}$ , the stock is above its MSST.
- If spawning biomass for 2008 is estimated to be above  $\frac{1}{2} B_{35\%}$  but below  $B_{35\%}$ , the stock's status relative to MSST is determined by referring to harvest scenario 6 (Table 1.23). If the mean spawning biomass for 2018 is below  $B_{35\%}$ , the stock is below its MSST. Otherwise, the stock is above its MSST.

Is the stock approaching an overfished condition? This is determined by referring to harvest Scenario 7:

- If the mean spawning biomass for 2010 is below  $\frac{1}{2} B_{35\%}$ , the stock is approaching an overfished condition.
- If the mean spawning biomass for 2010 is above  $B_{35\%}$ , the stock is not approaching an overfished condition.
- If the mean spawning biomass for 2010 is above  $\frac{1}{2} B_{35\%}$  but below  $B_{35\%}$ , the determination depends on the mean spawning biomass for 2020. If the mean spawning biomass for 2020 is below  $B_{35\%}$ , the stock is approaching an overfished condition. Otherwise, the stock is not approaching an overfished condition.

For scenarios 6 and 7, we conclude that pollock is not below MSST for the year 2008, nor is it expected to be approaching an overfished condition based on Scenario 7 (the mean spawning biomass in 2010 is above the  $B_{35\%}$  level; Table 1.23). For harvest recommendations, Tier 3 and a proxy for Tier 1 calculations were made that give ABC and OFL values for 2008 and 2009 (assuming catch is 1,170,000 t in 2008 Table 1.24). The Tier 1 projections were approximated by substituting the  $B_{msy}$  values for  $B_{40\%}$  (for the harvest control rule) and setting the  $F_{ABC}$  and  $F_{OFL}$  values to their spawning biomass-per-recruit (SPR) equivalent fishing mortalities. These SPR rates correspond to  $F_{32\%}$  and  $F_{26\%}$ , respectively. Additional projections using alternative future fixed catch levels (and alternative assumptions about the 2006 year class) are shown in Table 1.25.

## Other considerations

### Ecosystem considerations

In general, a number of key issues for ecosystem conservation and management can be highlighted. These include:

- Preventing overfishing;
- Avoiding habitat degradation;
- Minimizing incidental bycatch (via multi-species analyses of technical interactions);
- Controlling the level of discards; and
- Considering multi-species trophic interactions relative to harvest policies.

For the case of pollock in the Eastern Bering Sea, the NPFMC and NMFS continue to manage the fishery on the basis of these issues in addition to the single-species harvest approach. The prevention of overfishing is clearly set out as the main guideline for management. Habitat degradation has been minimized in the pollock fishery by converting the industry to pelagic-gear only. Bycatch in the pollock fleet is closely monitored by the NMFS observer program and managed on that basis. Discard rates of many species have been reduced in this fishery and efforts to minimize bycatch continue.

In comparisons of the Western Bering Sea (WBS) with the Eastern Bering Sea using mass-balance food-web models based on 1980-85 summer diet data, Aydin et al. (2002) found that the production in these two systems is quite different. On a per-unit-area measure, the western Bering Sea has higher productivity than the EBS. Also, the pathways of this productivity are different with much of the energy flowing through epifaunal species (e.g., sea urchins and brittlestars) in the WBS whereas for the EBS, crab and flatfish species play a similar role. In both regions, the keystone species in 1980-85 were pollock and Pacific cod. This study showed that the food web estimated for the EBS ecosystem appears to be relatively mature due to the large number of interconnections among species. In a more recent study based on 1990-93 diet data (see Appendix 1 of Ecosystem Considerations chapter for methods), pollock remain in a central role in the ecosystem (Fig. 1.45). The diet of pollock is similar between adults and juveniles with the exception that adults become more piscivorous (with consumption of pollock by adult pollock representing their third largest prey item). In terms of magnitude, pollock cannibalism may account for 2.5 million t to nearly 5 million t of pollock consumed (based on uncertainties in diet percentage and total consumption rate).

Regarding specific small-scale ecosystems of the EBS, Ciannelli et al. (2004) presented an application of an ecosystem model scaled to data available around the Pribilof Islands region. They applied bioenergetics and foraging theory to characterize the spatial extent of this ecosystem. They compared energy balance, from a food web model relevant to the foraging range of northern fur seals and found that a range of 100 nautical mile radius encloses the area of highest energy balance representing about 50% of the observed foraging range for lactating fur seals. This suggests that fur seals depend on areas outside the energetic balance region. This study develops a method for evaluating the shape and extent of a key ecosystem in the EBS (i.e., the Pribilof Islands). Furthermore, the extent that the pollock fishery extends into northern fur seal foraging habitat (e.g., Robson et al. 2004) will require careful monitoring and evaluation.

### Ecosystem effects on the EBS pollock stock

A brief summary of these two perspectives is given in Table 1.26. Unlike the food-web models discussed above, examining predators and prey in isolation may overly simplify relationships. This table serves to highlight the main connections and the status of our understanding or lack thereof.

In 2006 the EIT survey found an unusually low level of “other” backscatter in the water column based on summaries of the data from acoustic-trawl surveys of the eastern Bering Sea shelf conducted in June-July of 1999, 2000, 2002, 2004, 2006 and 2007 (Fig. 1. 46). These plots represent 38-kHz acoustic backscatter

( $s_A$ ,  $m^2/nmi^2$ ) attributed to an undifferentiated invertebrate-fish species mixture. For the 1999, 2000, 2002 and 2007 surveys, backscatter was measured between 14 m from the surface and 0.5 m off the bottom; in 2004 and 2006 it was measured between 12 m from the surface and 0.5 m off the bottom. These data should be interpreted with care because the exact biological composition of the non-pollock scatterers is unknown. Additionally, classification of non-pollock backscatter was not always performed as rigorously as classification of pollock, so non-pollock backscatter may contain some non-biological scatter. Trawl data suggest that biological components include jellyfish such as *Chrysaora* sp., other macrozooplankton, age 0 pollock, and other fishes. Some animals, such as fish with swimbladders and large medusae, are more easily detected at 38 kHz than small organisms such as copepods and euphausiids. Because these animals all reflect sound at different target strengths, comparison of backscatter both within and between years must be made with extreme caution. However, the data presented indicate that the contribution from non-pollock scatterers in 2006 was quite a bit lower than that in preceding years. In 2007, the contribution (or lack thereof) in the southeastern part of the shelf was similar to that in 2006 and quite different from preceding years. In 2007, most of the non-pollock backscatter was centered just to the north and west of the Pribilof Islands. These data suggest that the ecosystem, particularly in the southeastern region of the EBS, may have been different in the summers of 2006 and 2007, although the nature of the difference cannot be inferred from these data. The impact of this is unknown but should continue to be closely monitored. Work should be encouraged to better characterize the abiotic and biotic components of the non-pollock backscatter.

Furthermore, several other ecosystem indicators may give cause for concern. Zooplankton and non-pollock forage fish have been anomalously low in respective surveys (Figs. 3, 56, 64 in Ecosystem Considerations). While cannibalism still occurs within age-0 pollock (Fig. 61 in Ecosystem Considerations), cannibalism on age 1s by larger pollock has dropped since 1997 and may no longer be a main source of natural mortality for juvenile pollock (Fig. 4B in Ecosystem Considerations).

Moreover, the impact of non-cannibalistic predation may have shifted considerably in recent years. In particular, the increasing population of arrowtooth flounder in the Bering Sea is worth examining, especially considering the large predation caused by these flatfish in the Gulf of Alaska. Overall, the total non-cannibal groundfish predator biomass has gone down in the Bering Sea according to current stock assessments, with the drop of Pacific cod in the 1980s exceeding the rise of arrowtooth in terms of biomass (e.g., see Fig. 4 in Ecosystem Considerations chapter). This also represents a shift in the age of predation, with arrowtooth flounder consuming primarily age-2 pollock, while Pacific cod primarily consume larger pollock. However, the dynamics of this predation interaction may be quite different than in the Gulf of Alaska. A comparison of 1990-94 natural mortality by predator for arrowtooth flounder in the Bering Sea and the Gulf of Alaska shows that they are truly a top predator in the Gulf of Alaska. However, in the Bering Sea, pollock, skates, and sharks all prey on arrowtooth flounder, giving the species a relatively high predation mortality.

The predation on small arrowtooth flounder by large pollock gives rise to a specific concern for the Bering pollock stock. Walters and Kitchell (2001) describe a predator/prey system called “cultivation/depensation” whereby a species such as pollock “cultivates” its young by preying on species that would eat its young (for example, arrowtooth flounder). If these interactions are strong, the removal of the large pollock may lead to an accelerated decline, as the control it exerts on predators of its recruits is removed—this has been cited as a cause for a decline of cod in the Baltic Sea in the presence of herring feeding on cod young (Walters and Kitchell 2001). In situations like this, it is possible that predator culling (e.g., removing arrowtooth) may not have a strong effect towards controlling predation compared to applying additional caution to pollock harvest and thus preserving this natural control. At the moment, this concern for Bering Sea pollock is qualitative; work on extending a detailed, age-structured, multispecies statistical model (e.g., MSM; Jurado-Molina et al. 2005) to more completely model this complex interaction for pollock and arrowtooth flounder is continuing.

## **EBS pollock fishery effects on the ecosystem.**

Since the pollock fishery is primarily pelagic in nature, the bycatch of non-target species is small relative to the magnitude of the fishery (Table 1.27). Jellyfish represent the largest component of the bycatch of non-target species and has been stable at around 5-6 thousand tons per year (except for 2000 when over 9,000 t were caught). The data on non-target species shows a high degree of inter-annual variability which reflects the spatial variability of the fishery and high observation error. This variability may mask any significant trends in bycatch.

The catch of other target species in the pollock fishery represent less than 1% of the total pollock catch. Nonetheless incidental catch of Pacific cod has increased since 1999 but is below the 1997 levels (Table 1.28). The incidental catch of flatfish was variable over time and has increased slightly. Proportionately, the incidental catch has decreased since the overall levels of pollock catch have increased. The catch of prohibited species was also variable but showed noticeable trends (Table 1.29). For example, the level of crab bycatch drops considerably after 1998 when all BSAI pollock fishing was restricted to using only pelagic trawls. Recent levels of salmon bycatch have increased dramatically and current restrictions are under revision to help minimize this problem.

## **Summary**

Summary results are given in Table 1.30.

## **Acknowledgements**

Auke Bay Lab scientists with the BASIS program provided up to date information on their fall surveys. Grant Thompson provided the methodology used for the standard harvest scenarios and the associated text. Terry Hiatt compiled the catch and bycatch records. We thank the staff of the AFSC age-and-growth department for their excellent work in promptly processing the samples used in this assessment. The work of many individuals involved in collecting and processing survey and observer data is greatly appreciated.

## **References**

- Arsenev, V.S. 1967. Currents and water masses in the Bering Sea. Nauka Press, Moscow. English translation by S. Pearson, 1968, U.S. Dept. Commerce, NMFS, Seattle, 147 pp.
- Aydin, K. Y., et al. 2002. A comparison of the Eastern Bering and western Bering Sea shelf and slope ecosystems through the use of mass-balance food web models. U.S. Department of Commerce, Seattle, WA. (NOAA Technical Memorandum NMFS-AFSC-130) 78p.
- Bailey, K.M., T.J. Quinn, P. Bentzen, and W.S. Grant. 1999. Population structure and dynamics of walleye pollock, *Theragra chalcogramma*. *Advances in Mar. Biol.* 37:179-255.
- Barbeaux, S. J., S. Gaichas, J. N. Ianelli, and M. W. Dorn. 2005. Evaluation of biological sampling protocols for at-sea groundfish observers in Alaska. *Alaska Fisheries Research Bulletin* 11(2):82-101.
- Barbeaux, S.J., M. Dorn, J. Ianelli, and J. Horne. 2005. Visualizing Alaska pollock (*Theragra chalcogramma*) aggregation dynamics. ICES CM 2005/U:01.
- Beverton, R. J. H. and S. J. Holt. 1957. On the dynamics of exploited fish populations. *Fish. Invest., Lond., Ser. 2*, 19.
- Boldt, J. 2006. Ecosystem considerations chapter for 2006. <http://access.afsc.noaa.gov/reem/ecoweb>
- Butterworth, D.S., J.N. Ianelli, and R. Hilborn. 2003. A statistical model for stock assessment of southern bluefin tuna with temporal changes in selectivity. *Afr. J. mar. Sci.* 25: 331-361.
- Canino, M.F., and P. Bentzen (2004). Evidence for positive selection at the pantophysin (Pan I) locus in walleye pollock, *Theragra chalcogramma*. *Molecular Biology and Evolution*, Volume 21, No. 7, pp. 1391-1400 (July 2004).

- Canino, M.F., P.T. O'Reilly, L. Hauser, and P. Bentzen. 2005. Genetic differentiation in walleye pollock (*Theragra chalcogramma*) in response to selection at the pantophysin (*Pan I*) locus. *Can. J. Fish. Aquat. Sci.* 62:2519-2529.
- Ciannelli, L., B.W. Robson, R.C. Francis, K. Aydin, and R.D. Brodeur (2004). Boundaries of open marine ecosystems: an application to the Pribilof Archipelago, southeast Bering Sea. *Ecological Applications*, Volume 14, No. 3. pp. 942-953.
- Clark, W.G. 1999. Effects of an erroneous natural mortality rate on a simple age-structured model. *Can. J. Fish. Aquat. Sci.* 56:1721-1731.
- Cotter, A.J.R., L. Burt, C.G.M Paxton, C. Fernandez, S.T. Buckland, and J.X Pan. 2004. Are stock assessment methods too complicated? *Fish and Fisheries*, 5:235-254.
- Deriso, R. B., T. J. Quinn II, and P. R. Neal. 1985. Catch-age analysis with auxiliary information. *Can J. Fish. Aquat. Sci.* 42:815-824.
- Dorn, M.W. 1992. Detecting environmental covariates of Pacific whiting *Merluccius productus* growth using a growth-increment regression model. *Fish. Bull.* 90:260-275.
- Fadeev N.S., Wespestad V. Review of walleye Pollock fishery// *Izv. TINRO.-2001.- Vol.128.- p.75-91.*
- Fair, L.F. 1994. Eastern Bering Sea walleye pollock: revised estimates of population parameters, relation of recruitment to biological and environmental variables, and forecasting. M.S. Thesis, University of Alaska Fairbanks, Fairbanks AK. 131 p.
- Fair, L.F. and T.J. Quinn II, (In prep.). Eastern Bering Sea walleye pollock: a comparison of forecasting methods. Draft MS. Juneau Center, School of Fish. And Ocean Sci. Univ. Alaska Fairbanks. 32 p.
- Fournier, D. 1998. An Introduction to AD model builder for use in nonlinear modeling and statistics. Otter Research Ltd. PO Box 2040, Sidney BC V8L3S3, Canada, 53p.
- Fournier, D.A. and C.P. Archibald. 1982. A general theory for analyzing catch-at-age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.
- Francis, R.C., K. Aydin, R.L. Merrick, and S. Bollens. 1999. Modeling and Management of the Bering Sea Ecosystem. *In "Dynamics of the Bering Sea"*
- Francis, R.I.C.C. 1992. Use of risk analysis to assess fishery management strategies: a case study using orange roughy (*Hoplostethus atlanticus*) on the Chatham Rise, New Zealand. *Can. J. Fish. Aquat. Sci.* 49: 922-930.
- Greiwanck, A., and G.F. Corliss (eds.) 1991. Automatic differentiation of algorithms: theory, implementation and application. Proceedings of the SIAM Workshop on the Automatic Differentiation of Algorithms, held Jan. 6-8, Breckenridge, CO. Soc. Indust. And Applied Mathematics, Philadelphia.
- Harrison, R. C. 1993. Data Report: 1991 bottom trawl survey of the Aleutian Islands area. *Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., NOAA Tech. Memo. NMFS-AFSC-12.*
- Hinckley, S. 1987. The reproductive biology of walleye pollock, *Theragra chalcogramma*, in the Bering Sea, with reference to spawning stock structure. *Fish. Bull.* 85:481-498.
- Hollowed, A. B., J. N. Ianelli, and P. A. Livingston. 2000. Including predation mortality in stock assessments: A case study involving Gulf of Alaska walleye pollock. *ICES Journal of Marine Science*, 57, pp. 279-293.
- Honkalehto, T. N. Williamson, D. Hanson, D. McKelvey, and S. de Blois. 2002b. Results of the Echo Integration-trawl Survey of walleye Pollock (*Theragra chalcogramma*) Conducted on the Southeastern Bering Sea Shelf and in the Southeastern Aleutian Basin Near Bogoslof Island in February and March 2002. AFSC Processed Report 2002-02. 49p.
- Honkalehto, T. N. Williamson, D. McKelvey, and S. Stienessen. 2002a. Results of the Echo Integration-trawl Survey for Walleye Pollock (*Theragra chalcogramma*) on the Bering Sea Shelf and Slope in June and July 2002. AFSC Processed Report 2002-04. 38p.
- Honkalehto, T., D. McKelvey, and N. Williamson. 2005. Results of the echo integration-trawl survey of walleye pollock (*Theragra chalcogramma*) on the U.S. and Russian Bering Sea shelf in June and July 2004. AFSC Processed Rep. 2005-02, 43 p. Alaska Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way NE, Seattle WA 98115.
- Ianelli, J.N. 1996. An alternative stock assessment model of the Eastern Bering Sea pollock fishery. *In: Stock assessment and fishery evaluation report for the groundfish resources*



- of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, Appendix Section 1:1-73.
- Ianelli, J.N. 1997. An alternative stock assessment analysis for Gulf of Maine cod. SARC-24 Working Paper A2. 29p.
- Ianelli, J.N. and D.A. Fournier. 1998. Alternative age-structured analyses of the NRC simulated stock assessment data. *In* Restrepo, V.R. [ed.]. Analyses of simulated data sets in support of the NRC study on stock assessment methods. NOAA Tech. Memo. NMFS-F/SPO-30. 96 p.
- Ianelli, J.N., L. Fritz, T. Honkalehto, N. Williamson and G. Walters 1998. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 1999. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:1-79.
- Ianelli, J.N., L. Fritz, T. Honkalehto, N. Williamson and G. Walters. 2000. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2001. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:1-79.
- Ianelli, J.N., S. Barbeaux, G. Walters, T. Honkalehto, and N. Williamson. 2004. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2005. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:37-126.
- Ianelli, J.N., S. Barbeaux, T. Honkalehto, S. Kotwicki, K. Aydin, and N. Williamson. 2006. Assessment of Alaska Pollock Stock in the Eastern Bering Sea. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:35-138.
- Ianelli, J.N., S. Barbeaux, T. Honkalehto, N. Williamson and G. Walters. 2002. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2003. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:1-101.
- Ianelli, J.N., T. Buckley, T. Honkalehto, G Walters, and N. Williamson 2001. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2002. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/ Aleutian Islands regions. North Pac. Fish. Mgmt. Council Anchorage, AK, Section 1:1-79
- Ianelli, J.N., T. Buckley, T. Honkalehto, N. Williamson and G. Walters. 2001. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2002. *In*: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:1-105.
- Ingraham, W. J., Jr., and Miyahara, R. K. 1988. Ocean surface current simulations in the North Pacific Ocean and Bering Sea (OSCURS -Numerical Model). U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Technical Memorandum, National Marine Fisheries Service F/NWC-130, 155 pp.
- Jurado-Molina J., P. A. Livingston and J. N. Ianelli. 2005. Incorporating predation interactions to a statistical catch-at-age model for a predator-prey system in the eastern Bering Sea. Canadian Journal of Fisheries and Aquatic Sciences. 62(8): 1865-1873.
- Kastelle, C. R., and Kimura, D. K. 2006. Age validation of walleye pollock (*Theragra chalcogramma*) from the Gulf of Alaska using the disequilibrium of Pb-210 and Ra-226. *e* ICES Journal of Marine Science, 63: 1520e1529.
- Kimura, D.K. 1989. Variability in estimating catch-in-numbers-at-age and its impact on cohort analysis. *In* R.J. Beamish and G.A. McFarlane (eds.), Effects on ocean variability on recruitment and an evaluation of parameters used in stock assessment models. Can. Spec. Publ. Fish. Aq. Sci. 108:57-66.
- Kimura, D.K., J.J. Lyons, S.E. MacLellan, and B.J. Goetz. 1992. Effects of year-class strength on age determination. Aust. J. Mar. Freshwater Res. 43:1221-8.

- Kotwicki, S., T.W. Buckley, T. Honkalehto, and G. Walters. 2005. Variation in the distribution of walleye pollock (*Theragra chalcogramma*) with temperature and implications for seasonal migration. *Fish. Bull.* 103:574–587.
- Lauth, R.R., J.N. Ianelli, and W.W. Wakefield. 2004. Estimating the size selectivity and catching efficiency of a survey bottom trawl for thornyheads, *Sebastolobus spp.* using a towed video camera sled. *Fisheries Research.* 70:39-48.
- Livingston, P. A., and Methot, R. D. (1998). "Incorporation of predation into a population assessment model of Eastern Bering Sea walleye pollock. *In* Fishery Stock Assessment Models." NOAA Technical Report 126, NMFS F/NWC-54, Alaska Sea Grant Program, 304 Eielson Building, University of Alaska Fairbanks, Fairbanks, AK 99775. pp. 663-678.
- Low, L.L., and Ikeda. 1980. Average density index of walleye pollock in the Bering Sea. NOAA Tech. Memo. SFRF743.
- Mace, P., L. Botsford, J. Collie, W. Gabriel, P. Goodyear J. Powers, V. Restrepo, A. Rosenberg, M. Sissenwine, G. Thompson, J. Witzig. 1996. Scientific review of definitions of overfishing in U.S. Fishery Management Plans. NOAA Tech. Memo. NMFS-F/SPO-21. 20 p.
- McAllister, M.K. and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling-importance resampling algorithm. *Can. J. Fish. Aquat. Sci.* 54:284-300.
- Methot, R.D. 1990. Synthesis model: an adaptable framework for analysis of diverse stock assessment data. *In* Proceedings of the symposium on applications of stock assessment techniques to Gadids. L. Low [ed.]. *Int. North Pac. Fish. Comm. Bull.* 50: 259-277.
- Miller, T.J. 2005. Estimation of catch parameters from a fishery observer program with multiple objectives. PhD Dissertation. Univ. of Washington. 419p.
- Mueter, F. J., M.C. Palmer, and B.L. Norcross. 2004. Environmental predictors of walleye pollock recruitment on the Eastern Bering Sea shelf. Final Report to the Pollock Conservation Cooperative Research Center. June 2004. 74p.
- O'Reilly, P.T., M.F. Canino, K.M. Bailey and P. Bentzen. 2004. Inverse relationship between  $F_{ST}$  and microsatellite polymorphism in the marine fish, walleye pollock (*Theragra chalcogramma*): implications for resolving weak population structure. *Molecular Ecology* (2004) 13, 1799–1814
- Parma, A.M. 1993. Retrospective catch-at-age analysis of Pacific halibut: implications on assessment of harvesting policies. *In* Proceedings of the International Symposium on Management Strategies of Exploited Fish Populations. Alaska Sea Grant Rep. No. 93-02. Univ. Alaska Fairbanks.
- Pope, J. G. 1972. An investigation of the accuracy of virtual population analysis using cohort analysis. *Res. Bull. Int. Commn. NW Atlant. Fish.* 9: 65-74.
- Press, W.H., S.A. Teukolsky, W.T. Vetterling, B.P. Flannery. 1992. Numerical Recipes in C. Second Ed. Cambridge University Press. 994 p.
- Quinn II, T. J. and J. S. Collie. 1990. Alternative population models for Eastern Bering Sea pollock. INPFC Symposium on application of stock assessment techniques to gadids. *Int. North Pac. Fish. Comm. Bull.* 50:243-258.
- Quinn, T.J. and R.B. Deriso 1999. Quantitative Fish Dynamics. Oxford University Press, New York. 542 p.
- Restrepo, V.R., G.G. Thompson, P.M. Mace, W.L. Gabriel, L.L. Low, A.D. MacCall, R.D. Methot, J.E. Powers, B.L. Taylor, P.R. Wade, and J.F. Witzig. 1998. Technical guidance on the use of precautionary approaches to implementing National Standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. NOAA Tech. Memo. NMFS-F/SPO-31. 54 p.
- Ronholt, L. L., K. Teshima, and D. W. Kessler. 1994. The groundfish resources of the Aleutian Islands region and southern Bering Sea, 1980, 1983, and 1986. *Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., NOAA Tech. Memo. NMFS-AFSC-31.*
- Schnute, J.T. 1994. A general framework for developing sequential fisheries models. *Can. J. Fish. Aquat. Sci.* 51:1676-1688.
- Schnute, J.T. and Richards, L.J. 1995. The influence of error on population estimates from catch-age models. *Can. J. Fish. Aquat. Sci.* 52:2063-2077.

- Shuntov, V. P., A. F. Volkov, O. S. Temnykh, and E. P. Dulepova. 1993. Pollock in the ecosystems of the Far East Seas. TINRO, Vladivostok.
- Smith, G.B. 1981. The biology of walleye pollock. In Hood, D.W. and J.A. Calder, The Eastern Bering Sea Shelf: Oceanography and Resources. Vol. I. U.S. Dep. Comm., NOAA/OMP 527-551.
- Stahl, J. 2004. Maturation of walleye pollock, *Theragra chalcogramma*, in the Eastern Bering Sea in relation to temporal and spatial factors. Masters thesis. School of Fisheries and Ocean Sciences, Univ. Alaska Fairbanks, Juneau. 000p.
- Stepanenko, M.A. 1997. Variations from year to year in the spatial differentiation of the walleye pollock, *Theragra chalcogramma*, and the cod, *Gadus macrocephalus*, in the Bering Sea. Journ. of Ichthyol. 37:14-20.
- Swartzman, G.L., A.G. Winter, K.O. Coyle, R.D. Brodeur, T. Buckley, L. Ciannelli, G.L. Hunt, Jr., J. Ianelli, and S.A. Macklin (2005). Relationship of age-0 pollock abundance and distribution around the Pribilof Islands with other shelf regions of the Eastern Bering Sea. /Fisheries Research/, Vol. 74, pp. 273-287.
- Thompson, G.G. 1996. Risk-averse optimal harvesting in a biomass dynamic model. Unpubl. Manuscr., 54 p. Alaska Fisheries Science Center, 7600 Sand Pt. Way NE, Seattle WA, 98115. Distributed as Appendix B to the Environmental Analysis Regulatory Impact Review of Ammendments 44/44 to the Fishery Management Plans for the Groundfish Fisheries of the Bering Sea and Aleutian Islands Area and the Gulf of Alaska.
- Thompson, G.G. 1996. Spawning exploitation rate: a useful and general measure of relative fishing mortality. Alaska Fisheries Science Center contribution. Unpubl. Manuscr., 7 p.
- Thompson, G.G. and M.W. Dorn. 2004. Chapter 2: assessment of the Pacific cod stock in the Eastern Bering Sea and Aleutian Islands area. In: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 2. p185-302.
- Traynor J. J. and M. O. Nelson. 1985. Results of the U.S. hydroacoustic survey of pollock on the continental shelf and slope. In: R.G. Bakkala and K. Wakabayashi (eds.), Results of cooperative U.S.-Japan groundfish investigations in the Bering Sea during May-August 1979. Int. North Pac. Fish. Comm. Bull. 44: 192-199.
- von Szalay PG, Somerton DA, Kotwicki S. 2007. Correlating trawl and acoustic data in the Eastern Bering Sea: A first step toward improving biomass estimates of walleye pollock (*Theragra chalcogramma*) and Pacific cod (*Gadus macrocephalus*)? Fisheries Research 86(1) 77-83.
- Walters, C. J. 1969. A generalized computer simulation model for fish population studies. Trans. Am. Fish. Soc. 98:505 -512.
- Walters, C. J., and J. F. Kitchell. 2001. Cultivation/depensation effects on juvenile survival and recruitment. Can. J. Fish. Aquat. Sci. 58:39-50.
- Wespestad, V. G. 1990. Walleye pollock. Condition of groundfish resources in the Bering Sea-Aleutian Islands region as assessed in 1989. U.S. Dep. Commer., Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., NOAA Tech. Memo. NMFS F/AKC.
- Wespestad, V. G. and J. M. Terry. 1984. Biological and economic yields for Eastern Bering Sea walleye pollock under differing fishing regimes. N. Amer. J. Fish. Manage., 4:204-215.
- Wespestad, V. G. and J. Traynor. 1989. Walleye pollock. In: L-L. Low and R. Narita (editors), Condition of groundfish resources in the Bering Sea-Aleutian Islands region as assessed in 1988. U.S. Dep. Commer., Natl. Oceanic Atmos. Admin., Natl. Mar. Fish. Serv., NOAA Tech. Memo. NMFS F/AKC-178.
- Wespestad, V. G., J. Ianelli, L. Fritz, T. Honkalehto, G. Walters. 1996. Bering Sea-Aleutian Islands Walleye Pollock Assessment for 1997. In: Stock assessment and fishery evaluation report for the groundfish resources of the Bering Sea/Aleutian Islands regions. North Pac. Fish. Mgmt. Council, Anchorage, AK, section 1:1-73.
- Wespestad, V.G., L.W. Fritz, W.J. Ingraham, and B.A. Megrey. 1997. On Relationships between Cannibalism, climate variability, physical transport and recruitment success of Bering Sea Walleye Pollock, *Theragra chalcogramma*. ICES International Symposium, Recruitment Dynamics of exploited marine populations: physical-biological interactions. Baltimore, MD, Sept 22-24.

Winter, A.G., G.L. Swartzman, and L. Ciannelli (2005). Early- to late-summer population growth and prey consumption by age-0 pollock (*Theragra chalcogramma*), in two years of contrasting pollock abundance near the Pribilof Islands, Bering Sea. /Fisheries Oceanography/, Vol. 14, No. 4, pp. 307-320.

## Tables

Table 1.1 Catch from the Eastern Bering Sea by area, the Aleutian Islands, the Donut Hole, and the Bogoslof Island area, 1979-2007 (2007 values estimated). The southeast area refers to the EBS region east of 170W; the Northwest is west of 170W.

| Year | Eastern Bering Sea |           | Total     | Aleutians | Donut Hole | Bogoslof I. |
|------|--------------------|-----------|-----------|-----------|------------|-------------|
|      | Southeast          | Northwest |           |           |            |             |
| 1979 | 368,848            | 566,866   | 935,714   | 9,446     |            |             |
| 1980 | 437,253            | 521,027   | 958,280   | 58,157    |            |             |
| 1981 | 714,584            | 258,918   | 973,502   | 55,517    |            |             |
| 1982 | 713,912            | 242,052   | 955,964   | 57,753    |            |             |
| 1983 | 687,504            | 293,946   | 981,450   | 59,021    |            |             |
| 1984 | 442,733            | 649,322   | 1,092,055 | 77,595    | 181,200    |             |
| 1985 | 604,465            | 535,211   | 1,139,676 | 58,147    | 363,400    |             |
| 1986 | 594,997            | 546,996   | 1,141,993 | 45,439    | 1,039,800  |             |
| 1987 | 529,461            | 329,955   | 859,416   | 28,471    | 1,326,300  | 377,436     |
| 1988 | 931,812            | 296,909   | 1,228,721 | 41,203    | 1,395,900  | 87,813      |
| 1989 | 904,201            | 325,399   | 1,229,600 | 10,569    | 1,447,600  | 36,073      |
| 1990 | 640,511            | 814,682   | 1,455,193 | 79,025    | 917,400    | 151,672     |
| 1991 | 653,569            | 542,077   | 1,195,646 | 98,604    | 293,400    | 316,038     |
| 1992 | 830,560            | 559,771   | 1,390,331 | 52,352    | 10,000     | 241         |
| 1993 | 1,094,428          | 232,173   | 1,326,601 | 57,132    | 1,957      | 886         |
| 1994 | 1,152,573          | 176,777   | 1,329,350 | 58,659    |            | 556         |
| 1995 | 1,172,304          | 91,941    | 1,264,245 | 64,925    |            | 334         |
| 1996 | 1,086,840          | 105,938   | 1,192,778 | 29,062    |            | 499         |
| 1997 | 819,888            | 304,543   | 1,124,430 | 25,940    |            | 163         |
| 1998 | 965,767            | 135,399   | 1,101,165 | 23,822    |            | 136         |
| 1999 | 783,119            | 206,697   | 989,816   | 1,010     |            | 29          |
| 2000 | 839,175            | 293,532   | 1,132,707 | 1,244     |            | 29          |
| 2001 | 961,975            | 425,219   | 1,387,194 | 824       |            | 258         |
| 2002 | 1,159,730          | 320,465   | 1,480,195 | 1,156     |            | 1,042       |
| 2003 | 932,508            | 557,562   | 1,490,070 | 1,653     |            | 24          |
| 2004 | 1,089,970          | 390,708   | 1,480,678 | 1,150     |            | 0           |
| 2005 | 802,421            | 680,851   | 1,483,271 | 1,621     |            |             |
| 2006 | 826,887            | 659,397   | 1,486,284 | 1,735     |            |             |
| 2007 |                    |           | 1,340,000 |           |            |             |

1979-1989 data are from Pacfin.

1990-2006 data are from NMFS Alaska Regional Office, and includes discards.

2007 EBS catch is estimated

Table 1.2. Time series of ABC, TAC, and catch levels for EBS pollock, 1977-2007 in metric t.  
 Source: compiled from NMFS Regional office web site and various NPFMC reports, catch for 2007 is an estimated projection.

| Year              | ABC       | TAC       | Catch     |
|-------------------|-----------|-----------|-----------|
| 1977              | 950,000   | 950,000   | 978,370   |
| 1978              | 950,000   | 950,000   | 979,431   |
| 1979              | 1,100,000 | 950,000   | 935,714   |
| 1980              | 1,300,000 | 1,000,000 | 958,280   |
| 1981              | 1,300,000 | 1,000,000 | 973,502   |
| 1982              | 1,300,000 | 1,000,000 | 955,964   |
| 1983              | 1,300,000 | 1,000,000 | 981,450   |
| 1984              | 1,300,000 | 1,200,000 | 1,092,055 |
| 1985              | 1,300,000 | 1,200,000 | 1,139,676 |
| 1986              | 1,300,000 | 1,200,000 | 1,141,993 |
| 1987              | 1,300,000 | 1,200,000 | 859,416   |
| 1988              | 1,500,000 | 1,300,000 | 1,228,721 |
| 1989              | 1,340,000 | 1,340,000 | 1,229,600 |
| 1990              | 1,450,000 | 1,280,000 | 1,455,193 |
| 1991              | 1,676,000 | 1,300,000 | 1,195,646 |
| 1992              | 1,490,000 | 1,300,000 | 1,390,331 |
| 1993              | 1,340,000 | 1,300,000 | 1,326,601 |
| 1994              | 1,330,000 | 1,330,000 | 1,329,350 |
| 1995              | 1,250,000 | 1,250,000 | 1,264,245 |
| 1996              | 1,190,000 | 1,190,000 | 1,192,778 |
| 1997              | 1,130,000 | 1,130,000 | 1,124,430 |
| 1998              | 1,110,000 | 1,110,000 | 1,101,165 |
| 1999              | 992,000   | 992,000   | 989,816   |
| 2000              | 1,139,000 | 1,139,000 | 1,132,707 |
| 2001              | 1,842,000 | 1,400,000 | 1,387,194 |
| 2002              | 2,110,000 | 1,485,000 | 1,480,195 |
| 2003              | 2,330,000 | 1,491,760 | 1,490,070 |
| 2004              | 2,560,000 | 1,492,000 | 1,480,678 |
| 2005              | 1,960,000 | 1,478,500 | 1,483,271 |
| 2006              | 1,930,000 | 1,485,000 | 1,486,284 |
| 2007              | 1,394,000 | 1,394,000 | 1,340,000 |
| 1977-2007 average | 1,434,290 | 1,220,557 | 1,197,244 |

Table 1.3. Estimates of discarded pollock (t), percent of total (in parentheses) and total catch for the Aleutians, Bogoslof, Northwest and Southeastern Bering Sea, 1991-2007. Units are in tons, SE represents the EBS east of 170° W, NW is the EBS west of 170° W, source: NMFS Blend and catch-accounting system database.

|      | Discarded pollock |             |              |              |              | Total (retained plus discard) |          |         |           |           |
|------|-------------------|-------------|--------------|--------------|--------------|-------------------------------|----------|---------|-----------|-----------|
|      | Aleutian Is.      | Bogoslof    | NW           | SE           | Total        | Aleutian Is.                  | Bogoslof | NW      | SE        | Total     |
| 1991 | 5,231 (5%)        | 20,327 (6%) | 48,205 (9%)  | 66,789 (10%) | 140,552 (9%) | 98,604                        | 316,038  | 542,056 | 653,552   | 1,610,288 |
| 1992 | 2,982 (6%)        | 240 (100%)  | 57,609 (10%) | 71,195 (9%)  | 132,026 (9%) | 52,352                        | 241      | 559,771 | 830,560   | 1,442,924 |
| 1993 | 1,733 (3%)        | 308 (35%)   | 26,100 (11%) | 83,989 (8%)  | 112,130 (8%) | 57,132                        | 886      | 232,173 | 1,094,431 | 1,384,622 |
| 1994 | 1,373 (2%)        | 11 (2%)     | 16,083 (9%)  | 88,098 (8%)  | 105,565 (8%) | 58,659                        | 556      | 176,777 | 1,152,573 | 1,388,565 |
| 1995 | 1,380 (2%)        | 267 (80%)   | 9,715 (11%)  | 87,491 (7%)  | 98,853 (7%)  | 64,925                        | 334      | 91,941  | 1,172,304 | 1,329,503 |
| 1996 | 994 (3%)          | 7 (1%)      | 4,838 (5%)   | 71,367 (7%)  | 77,206 (6%)  | 29,062                        | 499      | 105,938 | 1,086,840 | 1,222,339 |
| 1997 | 617 (2%)          | 13 (8%)     | 22,557 (7%)  | 71,031 (9%)  | 94,218 (8%)  | 25,940                        | 163      | 304,543 | 819,888   | 1,150,533 |
| 1998 | 164 (1%)          | 3 (2%)      | 1,581 (1%)   | 15,135 (2%)  | 16,883 (2%)  | 23,822                        | 136      | 135,399 | 965,767   | 1,125,123 |
| 1999 | 480 (48%)         | 11 (38%)    | 1,912 (1%)   | 27,089 (3%)  | 29,492 (3%)  | 1,010                         | 29       | 206,697 | 783,119   | 990,855   |
| 2000 | 790 (64%)         | 20 (69%)    | 1,941 (1%)   | 19,678 (2%)  | 22,429 (2%)  | 1,244                         | 29       | 293,532 | 839,175   | 1,133,981 |
| 2001 | 380 (46%)         | 28 (11%)    | 2,450 (1%)   | 14,873 (2%)  | 17,731 (1%)  | 824                           | 258      | 425,219 | 961,889   | 1,388,190 |
| 2002 | 758 (66%)         | 12 (1%)     | 1,439 (0%)   | 19,226 (2%)  | 21,435 (1%)  | 1,156                         | 1,042    | 320,463 | 1,159,730 | 1,482,391 |
| 2003 | 468 (28%)         | NA          | 2,980 (1%)   | 14,063 (2%)  | 17,512 (1%)  | 1,653                         | NA       | 557,552 | 933,459   | 1,492,664 |
| 2004 | 758 (66%)         | NA          | 2,723 (1%)   | 20,302 (2%)  | 23,783 (2%)  | 1,156                         | NA       | 390,414 | 1,089,880 | 1,482,373 |
| 2005 | 324 (20%)         |             | 2,581 (0%)   | 14,838 (2%)  | 17,742 (1%)  | 1,621                         |          | 680,856 | 802,418   | 1,484,895 |
| 2006 | 310 (18%)         |             | 3,672 (1%)   | 11,588 (1%)  | 15,570 (1%)  | 1,735                         |          | 659,397 | 826,887   | 1,488,019 |
| 2007 | 386 (16%)         |             | 4,136 (1%)   | 12,270 (2%)  | 16,792 (1%)  | 2,448                         |          | 592,411 | 657,552   | 1,252,411 |

Table 1.4. Eastern Bering Sea pollock catch at age estimates based on observer data, 1979-2006. Units are in millions of fish.

| Year    | 1     | 2     | 3       | 4       | 5       | 6     | 7     | 8     | 9     | 10    | 11    | 12   | 13   | 14+  | Total   |
|---------|-------|-------|---------|---------|---------|-------|-------|-------|-------|-------|-------|------|------|------|---------|
| 1979    | 101.4 | 543.0 | 719.8   | 420.1   | 392.5   | 215.5 | 56.3  | 25.7  | 35.9  | 27.5  | 17.6  | 7.9  | 3.0  | 1.1  | 2,567.3 |
| 1980    | 9.8   | 462.2 | 822.9   | 443.3   | 252.1   | 210.9 | 83.7  | 37.6  | 21.7  | 23.9  | 25.4  | 15.9 | 7.7  | 3.7  | 2,420.8 |
| 1981    | 0.6   | 72.2  | 1,012.7 | 637.9   | 227.0   | 102.9 | 51.7  | 29.6  | 16.1  | 9.3   | 7.5   | 4.6  | 1.5  | 1.0  | 2,174.6 |
| 1982    | 4.7   | 25.3  | 161.4   | 1,172.2 | 422.3   | 103.7 | 36.0  | 36.0  | 21.5  | 9.1   | 5.4   | 3.2  | 1.9  | 1.0  | 2,003.8 |
| 1983    | 5.1   | 118.6 | 157.8   | 312.9   | 816.8   | 218.2 | 41.4  | 24.7  | 19.8  | 11.1  | 7.6   | 4.9  | 3.5  | 2.1  | 1,744.5 |
| 1984    | 2.1   | 45.8  | 88.6    | 430.4   | 491.4   | 653.6 | 133.7 | 35.5  | 25.1  | 15.6  | 7.1   | 2.5  | 2.9  | 3.7  | 1,938.0 |
| 1985    | 2.6   | 55.2  | 381.2   | 121.7   | 365.7   | 321.5 | 443.2 | 112.5 | 36.6  | 25.8  | 24.8  | 10.7 | 9.4  | 9.1  | 1,920.0 |
| 1986    | 3.1   | 86.0  | 92.3    | 748.6   | 214.1   | 378.1 | 221.9 | 214.3 | 59.7  | 15.2  | 3.3   | 2.6  | 0.3  | 1.2  | 2,040.5 |
| 1987    | 0.0   | 19.8  | 111.5   | 77.6    | 413.4   | 138.8 | 122.4 | 90.6  | 247.2 | 54.1  | 38.7  | 21.4 | 28.9 | 14.1 | 1,378.6 |
| 1988    | 0.0   | 10.7  | 454.0   | 421.6   | 252.1   | 544.3 | 224.8 | 104.9 | 39.2  | 96.8  | 18.2  | 10.2 | 3.8  | 11.7 | 2,192.2 |
| 1989    | 0.0   | 4.8   | 55.1    | 149.0   | 451.1   | 166.7 | 572.2 | 96.3  | 103.8 | 32.4  | 129.0 | 10.9 | 4.0  | 8.5  | 1,783.8 |
| 1990    | 1.3   | 33.0  | 57.0    | 219.5   | 200.7   | 477.7 | 129.2 | 368.4 | 65.7  | 101.9 | 9.0   | 60.1 | 8.5  | 13.9 | 1,745.9 |
| 1991    | 1.0   | 109.1 | 41.0    | 86.8    | 140.0   | 156.0 | 389.1 | 52.5  | 219.4 | 22.2  | 114.8 | 13.9 | 72.0 | 59.3 | 1,477.0 |
| 1992    | 0.0   | 87.8  | 674.4   | 130.7   | 79.2    | 112.6 | 134.9 | 256.0 | 100.6 | 156.3 | 55.6  | 43.3 | 12.7 | 75.3 | 1,919.2 |
| 1993    | 0.1   | 7.5   | 260.9   | 1,153.8 | 102.4   | 64.8  | 62.4  | 52.6  | 91.0  | 21.2  | 33.4  | 12.1 | 12.4 | 23.8 | 1,898.3 |
| 1994    | 0.3   | 35.5  | 54.7    | 357.8   | 1,068.1 | 175.4 | 53.8  | 20.3  | 13.3  | 21.6  | 8.8   | 9.7  | 7.3  | 11.5 | 1,838.1 |
| 1995    | 0.0   | 0.4   | 80.6    | 152.2   | 398.9   | 765.2 | 131.2 | 31.6  | 9.6   | 8.2   | 18.3  | 5.4  | 6.0  | 10.5 | 1,618.2 |
| 1996    | 0.0   | 23.3  | 54.7    | 87.2    | 157.2   | 362.4 | 476.2 | 186.4 | 32.3  | 13.5  | 9.6   | 8.9  | 4.2  | 11.6 | 1,427.7 |
| 1997    | 0.0   | 79.7  | 34.2    | 108.5   | 467.3   | 288.1 | 252.9 | 199.0 | 62.7  | 13.8  | 6.9   | 5.1  | 3.5  | 16.1 | 1,537.6 |
| 1998    | 0.0   | 47.8  | 86.8    | 71.7    | 156.6   | 692.2 | 200.0 | 129.4 | 109.6 | 29.8  | 6.3   | 5.8  | 2.8  | 8.2  | 1,546.8 |
| 1999    | 0.4   | 11.5  | 294.2   | 226.1   | 105.3   | 156.1 | 473.3 | 133.3 | 57.8  | 33.1  | 3.5   | 2.3  | 0.5  | 2.3  | 1,499.6 |
| 2000    | 0.0   | 17.2  | 80.6    | 426.9   | 346.0   | 106.0 | 169.3 | 356.9 | 85.3  | 29.3  | 23.9  | 5.6  | 1.5  | 2.3  | 1,650.7 |
| 2001    | 0.0   | 3.6   | 56.3    | 161.9   | 575.3   | 408.2 | 135.3 | 130.1 | 157.8 | 57.5  | 35.0  | 15.9 | 5.9  | 5.0  | 1,747.8 |
| 2002    | 0.3   | 53.5  | 110.1   | 213.4   | 284.9   | 604.2 | 268.6 | 98.9  | 87.0  | 95.9  | 34.6  | 14.4 | 12.6 | 4.4  | 1,883.0 |
| 2003    | 0.0   | 17.5  | 414.4   | 320.3   | 365.9   | 304.8 | 330.4 | 156.3 | 53.0  | 39.8  | 36.2  | 23.6 | 7.0  | 7.0  | 2,076.3 |
| 2004    | 0.0   | 1.1   | 90.0    | 834.0   | 481.7   | 238.0 | 167.8 | 155.9 | 62.4  | 16.6  | 18.6  | 25.7 | 10.5 | 13.0 | 2,115.4 |
| 2005    | 0.0   | 3.3   | 54.6    | 391.0   | 859.2   | 490.4 | 156.6 | 67.8  | 67.0  | 33.1  | 10.8  | 10.2 | 3.4  | 5.4  | 2,152.7 |
| 2006    | 0.0   | 12.3  | 84.6    | 290.7   | 622.2   | 591.3 | 281.1 | 109.0 | 49.9  | 38.2  | 16.3  | 9.6  | 9.5  | 12.8 | 2,127.5 |
| Average | 4.7   | 71.0  | 235.2   | 363.1   | 382.5   | 323.1 | 207.1 | 118.3 | 69.7  | 37.6  | 25.9  | 13.1 | 8.8  | 12.1 | 1,872.4 |
| Median  | 0.1   | 29.2  | 91.1    | 301.8   | 365.8   | 263.1 | 162.2 | 101.9 | 58.8  | 26.7  | 17.9  | 10.0 | 5.1  | 8.3  | 1,890.7 |

Table 1.5. Numbers of pollock fishery samples measured for lengths and for length-weight by sex and strata, 1977-2007, as sampled by the NMFS observer program.

| Length<br>Frequency     | A Season |         | B Season SE |         | B Season NW |         | Total   |
|-------------------------|----------|---------|-------------|---------|-------------|---------|---------|
|                         | Males    | Females | Males       | Females | Males       | Females |         |
| 1977                    | 26,411   | 25,923  | 4,301       | 4,511   | 29,075      | 31,219  | 121,440 |
| 1978                    | 25,110   | 31,653  | 9,829       | 9,524   | 46,349      | 46,072  | 168,537 |
| 1979                    | 59,782   | 62,512  | 3,461       | 3,113   | 62,298      | 61,402  | 252,568 |
| 1980                    | 42,726   | 42,577  | 3,380       | 3,464   | 47,030      | 49,037  | 188,214 |
| 1981                    | 64,718   | 57,936  | 2,401       | 2,147   | 53,161      | 53,570  | 233,933 |
| 1982                    | 74,172   | 70,073  | 16,265      | 14,885  | 181,606     | 163,272 | 520,273 |
| 1983                    | 94,118   | 90,778  | 16,604      | 16,826  | 193,031     | 174,589 | 585,946 |
| 1984                    | 158,329  | 161,876 | 106,654     | 105,234 | 243,877     | 217,362 | 993,332 |
| 1985                    | 119,384  | 109,230 | 96,684      | 97,841  | 284,850     | 256,091 | 964,080 |
| 1986                    | 186,505  | 189,497 | 135,444     | 123,413 | 164,546     | 131,322 | 930,727 |
| 1987                    | 373,163  | 399,072 | 14,170      | 21,162  | 24,038      | 22,117  | 853,722 |
| 1991                    | 160,491  | 148,236 | 166,117     | 150,261 | 141,085     | 139,852 | 906,042 |
| 1992                    | 158,405  | 153,866 | 163,045     | 164,227 | 101,036     | 102,667 | 843,244 |
| 1993                    | 143,296  | 133,711 | 148,299     | 140,402 | 27,262      | 28,522  | 621,490 |
| 1994                    | 139,332  | 147,204 | 159,341     | 153,526 | 28,015      | 27,953  | 655,370 |
| 1995                    | 131,287  | 128,389 | 179,312     | 154,520 | 16,170      | 16,356  | 626,032 |
| 1996                    | 149,111  | 140,981 | 200,482     | 156,804 | 18,165      | 18,348  | 683,890 |
| 1997                    | 124,953  | 104,115 | 116,448     | 107,630 | 60,192      | 53,191  | 566,527 |
| 1998                    | 136,605  | 110,620 | 208,659     | 178,012 | 32,819      | 40,307  | 707,019 |
| 1999                    | 36,258   | 32,630  | 38,840      | 35,695  | 16,282      | 18,339  | 178,044 |
| 2000                    | 64,575   | 58,162  | 63,832      | 41,120  | 40,868      | 39,134  | 307,689 |
| 2001                    | 79,333   | 75,633  | 54,119      | 51,268  | 44,295      | 45,836  | 350,483 |
| 2002                    | 71,776   | 69,743  | 65,432      | 64,373  | 37,701      | 39,322  | 348,347 |
| 2003                    | 74,995   | 77,612  | 49,469      | 53,053  | 51,799      | 53,463  | 360,390 |
| 2004                    | 75,426   | 76,018  | 63,204      | 62,005  | 47,289      | 44,246  | 368,188 |
| 2005                    | 76,627   | 69,543  | 43,205      | 33,886  | 68,878      | 63,088  | 355,225 |
| 2006                    | 72,353   | 63,108  | 28,799      | 22,363  | 75,180      | 65,209  | 327,010 |
| 2007                    | 66,811   | 64,243  | 14,811      | 11,859  | 30,628      | 27,370  | 215,720 |
| Length – weight samples |          |         |             |         |             |         |         |
| 1977                    | 1,222    | 1,338   | 137         | 166     | 1,461       | 1,664   | 5,988   |
| 1978                    | 1,991    | 2,686   | 409         | 516     | 2,200       | 2,623   | 10,425  |
| 1979                    | 2,709    | 3,151   | 152         | 209     | 1,469       | 1,566   | 9,256   |
| 1980                    | 1,849    | 2,156   | 99          | 144     | 612         | 681     | 5,541   |
| 1981                    | 1,821    | 2,045   | 51          | 52      | 1,623       | 1,810   | 7,402   |
| 1982                    | 2,030    | 2,208   | 181         | 176     | 2,852       | 3,043   | 10,490  |
| 1983                    | 1,199    | 1,200   | 144         | 122     | 3,268       | 3,447   | 9,380   |
| 1984                    | 980      | 1,046   | 117         | 136     | 1,273       | 1,378   | 4,930   |
| 1985                    | 520      | 499     | 46          | 55      | 426         | 488     | 2,034   |
| 1986                    | 689      | 794     | 518         | 501     | 286         | 286     | 3,074   |
| 1987                    | 1,351    | 1,466   | 25          | 33      | 72          | 63      | 3,010   |
| 1991                    | 2,712    | 2,781   | 2,339       | 2,496   | 1,065       | 1,169   | 12,562  |
| 1992                    | 1,517    | 1,582   | 1,911       | 1,970   | 588         | 566     | 8,134   |
| 1993                    | 1,201    | 1,270   | 1,448       | 1,406   | 435         | 450     | 6,210   |
| 1994                    | 1,552    | 1,630   | 1,569       | 1,577   | 162         | 171     | 6,661   |
| 1995                    | 1,215    | 1,259   | 1,320       | 1,343   | 223         | 232     | 5,592   |
| 1996                    | 2,094    | 2,135   | 1,409       | 1,384   | 1           | 1       | 7,024   |
| 1997                    | 628      | 627     | 616         | 665     | 511         | 523     | 3,570   |
| 1998                    | 1,852    | 1,946   | 959         | 923     | 327         | 350     | 6,357   |
| 1999                    | 5,318    | 4,798   | 7,797       | 7,054   | 3,532       | 3,768   | 32,267  |
| 2000                    | 12,421   | 11,318  | 12,374      | 7,809   | 7,977       | 7,738   | 59,637  |
| 2001                    | 14,882   | 14,369  | 10,778      | 10,378  | 8,777       | 9,079   | 68,263  |
| 2002                    | 14,004   | 13,541  | 12,883      | 12,942  | 7,202       | 7,648   | 68,220  |
| 2003                    | 14,780   | 15,495  | 9,401       | 10,092  | 9,994       | 10,261  | 70,023  |
| 2004                    | 7,690    | 7,890   | 6,819       | 6,847   | 4,603       | 4,321   | 38,170  |
| 2005                    | 7,390    | 7,033   | 5,109       | 4,115   | 6,927       | 6,424   | 36,998  |
| 2006                    | 7,324    | 6,989   | 5,085       | 4,068   | 6,842       | 6,356   | 36,664  |
| 2007                    | 6,664    | 6,618   | 1,750       | 1,300   | 3,095       | 2,634   | 22,061  |

Table 1.6. Numbers of pollock fishery samples used for age determination estimates by sex and strata, 1977-2006, as sampled by the NMFS observer program.

|      | <b>Aged</b>     |                |                    |                |                    |                | <b>Total</b> |
|------|-----------------|----------------|--------------------|----------------|--------------------|----------------|--------------|
|      | <b>A Season</b> |                | <b>B Season SE</b> |                | <b>B Season NW</b> |                |              |
|      | <b>Males</b>    | <b>Females</b> | <b>Males</b>       | <b>Females</b> | <b>Males</b>       | <b>Females</b> |              |
| 1977 | 1,229           | 1,344          | 137                | 166            | 1,415              | 1,613          | 5,904        |
| 1978 | 1,992           | 2,686          | 407                | 514            | 2,188              | 2,611          | 10,398       |
| 1979 | 2,647           | 3,088          | 152                | 209            | 1,464              | 1,561          | 9,121        |
| 1980 | 1,854           | 2,158          | 93                 | 138            | 606                | 675            | 5,524        |
| 1981 | 1,819           | 2,042          | 51                 | 52             | 1,620              | 1,807          | 7,391        |
| 1982 | 2,030           | 2,210          | 181                | 176            | 2,865              | 3,062          | 10,524       |
| 1983 | 1,200           | 1,200          | 144                | 122            | 3,249              | 3,420          | 9,335        |
| 1984 | 980             | 1,046          | 117                | 136            | 1,272              | 1,379          | 4,930        |
| 1985 | 520             | 499            | 46                 | 55             | 426                | 488            | 2,034        |
| 1986 | 689             | 794            | 518                | 501            | 286                | 286            | 3,074        |
| 1987 | 1,351           | 1,466          | 25                 | 33             | 72                 | 63             | 3,010        |
| 1991 | 420             | 423            | 272                | 265            | 320                | 341            | 2,041        |
| 1992 | 392             | 392            | 371                | 386            | 178                | 177            | 1,896        |
| 1993 | 444             | 473            | 503                | 493            | 124                | 122            | 2,159        |
| 1994 | 201             | 202            | 570                | 573            | 131                | 141            | 1,818        |
| 1995 | 298             | 316            | 436                | 417            | 123                | 131            | 1,721        |
| 1996 | 468             | 449            | 442                | 433            | 1                  | 1              | 1,794        |
| 1997 | 433             | 436            | 284                | 311            | 326                | 326            | 2,116        |
| 1998 | 592             | 659            | 307                | 307            | 216                | 232            | 2,313        |
| 1999 | 540             | 500            | 730                | 727            | 306                | 298            | 3,100        |
| 2000 | 666             | 626            | 843                | 584            | 253                | 293            | 3,265        |
| 2001 | 598             | 560            | 724                | 688            | 178                | 205            | 2,951        |
| 2002 | 651             | 670            | 834                | 886            | 201                | 247            | 3,489        |
| 2003 | 583             | 644            | 652                | 680            | 260                | 274            | 3,092        |
| 2004 | 560             | 547            | 599                | 697            | 244                | 221            | 2,867        |
| 2005 | 611             | 597            | 613                | 489            | 419                | 421            | 3,149        |
| 2006 | 608             | 599            | 590                | 457            | 397                | 398            | 3,048        |

Table 1.7. NMFS total pollock research catch by year in t, 1963-2007.

|              |      |      |      |      |      |      |      |      |      |      |      |
|--------------|------|------|------|------|------|------|------|------|------|------|------|
| Year         | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 |
| Aleutian Is. | 0    | 0    | 0    | 0    | 0    | 0    | 0    | 0    | 0    | 0    | 0    |
| Bering Sea   | 4    | 0    | 18   | 17   | 21   | 7    | 14   | 9    | 16   | 11   | 69   |
| Year         | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 |
| Aleutian Is. | 0    | 0    | 0    | 0    | 0    | 0    | 193  | 0    | 40   | 454  | 0    |
| Bering Sea   | 83   | 197  | 122  | 35   | 94   | 458  | 139  | 466  | 682  | 508  | 208  |
| Year         | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
| Aleutian Is. | 0    | 292  | 0    | 0    | 0    | 0    | 51   | 0    | 0    | 48   | 0    |
| Bering Sea   | 435  | 163  | 174  | 467  | 393  | 369  | 465  | 156  | 221  | 267  | 249  |
| Year         | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2006 | 2007 |
| Aleutian Is. | 0    | 36   | 0    | 0    | 40   | 0    | 79   | 0    | 51   | 21   | 0    |
| Bering Sea   | 206  | 262  | 121  | 299  | 313  | 241  | 440  | 285  | 363  | 251  | 333  |



Table 1.8. Sampling effort for pollock in the EBS from the NMFS bottom trawl survey 1982-2007.  
 Years where only strata 1-6 were surveyed are shown in italics.

| <b>Year</b> | <b>Number of Hauls</b> | <b>Lengths</b> | <b>Aged</b>  | <b>Year</b> | <b>Number of Hauls</b> | <b>Lengths</b> | <b>Aged</b> |
|-------------|------------------------|----------------|--------------|-------------|------------------------|----------------|-------------|
| <i>1982</i> | <i>329</i>             | <i>40,001</i>  | <i>1,611</i> | 1995        | 376                    | 25,673         | 1,156       |
| <i>1983</i> | <i>354</i>             | <i>78,033</i>  | <i>1,931</i> | 1996        | 375                    | 40,789         | 1,387       |
| <i>1984</i> | <i>355</i>             | <i>40,530</i>  | <i>1,806</i> | 1997        | 376                    | 35,536         | 1,193       |
| 1985        | 434                    | 48,642         | 1,913        | 1998        | 375                    | 37,673         | 1,261       |
| <i>1986</i> | <i>354</i>             | <i>41,101</i>  | <i>1,344</i> | 1999        | 373                    | 32,532         | 1,385       |
| 1987        | 356                    | 40,144         | 1,607        | 2000        | 372                    | 41,762         | 1,545       |
| 1988        | 373                    | 40,408         | 1,173        | 2001        | 375                    | 47,335         | 1,641       |
| 1989        | 373                    | 38,926         | 1,227        | 2002        | 375                    | 43,361         | 1,695       |
| 1990        | 371                    | 34,814         | 1,257        | 2003        | 376                    | 46,480         | 1,638       |
| 1991        | 371                    | 43,406         | 1,083        | 2004        | 375                    | 44,102         | 1,660       |
| 1992        | 356                    | 34,024         | 1,263        | 2005        | 373                    | 35,976         | 1,676       |
| 1993        | 375                    | 43,278         | 1,385        | 2006        | 376                    | 39,211         | 1,573       |
| 1994        | 375                    | 38,901         | 1,141        | 2007        |                        |                | 1,484       |

Table 1.9. Biomass (age 1+) of Eastern Bering Sea pollock as estimated by surveys 1979-2006 (millions of tons). Note that the bottom-trawl survey data only represent biomass from the standard survey strata (1-6) areas in 1982-1984, and 1986. For all other years the estimates include strata 8-9. Also, the 1979 - 1981 bottom trawl survey data were omitted from the model since the survey gear differed.

| Year | Bottom trawl Survey (t) | EIT Survey (t)    | EIT Percent age 3+ | Total* (t) | Near bottom biomass |
|------|-------------------------|-------------------|--------------------|------------|---------------------|
| 1979 | 3.20                    | 7.46              | (22%)              | 10.66      | 30%                 |
| 1980 | 1.00                    |                   |                    |            |                     |
| 1981 | 2.30                    |                   |                    |            |                     |
| 1982 | 2.856                   | 4.90              | (95%)              | 7.76       | 46%                 |
| 1983 | 6.258                   |                   |                    |            |                     |
| 1984 | 4.894                   |                   |                    |            |                     |
| 1985 | 5.928                   | 4.80              | (97%)              | 9.43       | 54%                 |
| 1986 | 4.897                   |                   |                    |            |                     |
| 1987 | 5.515                   |                   |                    |            |                     |
| 1988 | 7.289                   | 4.68              | (97%)              | 11.79      | 63%                 |
| 1989 | 6.519                   |                   |                    |            |                     |
| 1990 | 7.322                   |                   |                    |            |                     |
| 1991 | 5.168                   | 1.45              | (46%)              | 6.56       | 79%                 |
| 1992 | 4.583                   |                   |                    |            |                     |
| 1993 | 5.636                   |                   |                    |            |                     |
| 1994 | 5.027                   | 2.89              | (85%)              | 7.87       | 64%                 |
| 1995 | 5.482                   |                   |                    |            |                     |
| 1996 | 3.371                   | 2.31              | (97%)              | 5.51       | 60%                 |
| 1997 | 3.874                   | 2.59              | (70%)              | 5.62       | 54%                 |
| 1998 | 2.852                   |                   |                    |            |                     |
| 1999 | 3.801                   | 3.29 <sup>†</sup> | (95%)              | 6.86       | 52%                 |
| 2000 | 5.265                   | 3.05              | (95%)              | 8.19       | 63%                 |
| 2001 | 4.200                   |                   |                    |            |                     |
| 2002 | 5.038                   | 3.62              | (82%)              | 8.39       | 57%                 |
| 2003 | 8.458                   |                   |                    |            |                     |
| 2004 | 3.886                   | 3.31              | (99%)              | 7.06       | 53%                 |
| 2005 | 5.318                   |                   |                    |            |                     |
| 2006 | 3.045                   | 1.56              |                    | 4.41       | 65%                 |
| 2007 | 4.338                   | 1.878             |                    | 6.22       | 70%                 |

\* Although the two survey estimates are added in this table, the stock assessment model treats them as separate, independent indices (survey “*q*’s” are estimated).

<sup>†</sup> This figure excludes the zone near the “horseshoe” area of the EBS (southeast) not usually surveyed, the value including this area was 3.35 million tons.

Table 1.10. Survey biomass estimates (age 1+, t) of Eastern Bering Sea pollock based on area-swept expansion methods from NMFS bottom trawl surveys 1982-2006.

| Year | Survey biomass estimates in strata 1-6 | Survey biomass estimates in strata 8 and 9 (NW) | All area Total | NW % Total |
|------|--|---|----------------|------------|
| 1982 | 2,855,539                              |   |                |            |
| 1983 | 6,257,632                              |   |                |            |
| 1984 | 4,893,536                              |   |                |            |
| 1985 | 4,630,111                              | 1,425,625                                       | 6,055,736      | 24%        |
| 1986 | 4,896,780                              |   |                |            |
| 1987 | 5,108,035                              | 416,558   | 5,524,593      | 8%         |
| 1988 | 7,107,258                              | 181,909   | 7,289,168      | 2%         |
| 1989 | 5,927,187                              | 591,622   | 6,518,809      | 9%         |
| 1990 | 7,126,083                              | 195,894   | 7,321,977      | 3%         |
| 1991 | 5,105,224                              | 62,523  | 5,167,748      | 1%         |
| 1992 | 4,367,870                              | 214,676   | 4,582,546      | 5%         |
| 1993 | 5,520,892                              | 114,757   | 5,635,649      | 2%         |
| 1994 | 4,977,019                              | 49,721  | 5,026,740      | 1%         |
| 1995 | 5,413,270                              | 68,983  | 5,482,253      | 1%         |
| 1996 | 3,204,106                              | 167,090   | 3,371,196      | 5%         |
| 1997 | 3,031,557                              | 842,276   | 3,873,833      | 22%        |
| 1998 | 2,212,689                              | 639,715   | 2,852,404      | 22%        |
| 1999 | 3,597,403                              | 203,314   | 3,800,717      | 5%         |
| 2000 | 5,134,616                              | 129,932   | 5,264,548      | 2%         |
| 2001 | 4,145,746                              | 54,162  | 4,199,909      | 1%         |
| 2002 | 4,832,506                              | 205,231   | 5,037,737      | 4%         |
| 2003 | 8,140,573                              | 317,089   | 8,457,662      | 4%         |
| 2004 | 3,756,228                              | 130,227   | 3,886,455      | 3%         |
| 2005 | 5,133,606                              | 160,109   | 5,293,715      | 3%         |
| 2006 | 2,845,507                              | 199,932   | 3,045,438      | 7%         |
| 2007 | 4,156,687                              | 180,856   | 4,337,542      | 4%         |
| Avg. | 4,783,756                              | 297,827   | 5,092,108      | 6%         |

Table 1.11. Bottom-trawl survey estimated numbers (millions) at age used for the stock assessment model, 1982-2007 based on strata 1-8. Shaded cells represent years where only strata 1-6 were surveyed.

| Year | 1     | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    | 11    | 12  | 13  | 14  | 15  | Total  | StdErr | CV  |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----|-----|-----|-----|--------|--------|-----|
| 1982 | 821   | 2,029 | 2,407 | 3,276 | 1,075 | 150   | 103   | 50    | 33    | 18    | 9     | 7   | 2   | 1   | 0   | 9,980  | 1,269  | 13% |
| 1983 | 483   | 670   | 1,638 | 3,060 | 6,663 | 1,979 | 369   | 199   | 78    | 72    | 56    | 19  | 9   | 8   | 3   | 15,306 | 1,198  | 8%  |
| 1984 | 280   | 261   | 348   | 1,196 | 1,400 | 3,551 | 694   | 157   | 68    | 25    | 16    | 6   | 4   | 5   | 2   | 8,012  | 795    | 10% |
| 1985 | 3,053 | 581   | 2,591 | 1,111 | 3,839 | 2,169 | 1,580 | 319   | 81    | 64    | 18    | 6   | 7   | 1   | 0   | 15,420 | 1,967  | 13% |
| 1986 | 1,931 | 278   | 312   | 1,549 | 859   | 1,597 | 1,317 | 1,133 | 389   | 64    | 27    | 12  | 0   | 3   | 0   | 9,473  | 838    | 9%  |
| 1987 | 254   | 474   | 644   | 413   | 3,691 | 815   | 991   | 376   | 1,242 | 181   | 69    | 24  | 4   | 1   | 1   | 9,181  | 1,129  | 12% |
| 1988 | 537   | 449   | 969   | 2,463 | 1,002 | 3,590 | 1,158 | 909   | 522   | 1,180 | 120   | 66  | 12  | 22  | 9   | 13,009 | 1,477  | 11% |
| 1989 | 580   | 223   | 427   | 1,385 | 3,096 | 703   | 2,697 | 400   | 526   | 199   | 517   | 97  | 76  | 43  | 49  | 11,019 | 1,083  | 10% |
| 1990 | 1,333 | 219   | 65    | 590   | 1,038 | 3,810 | 846   | 2,171 | 239   | 406   | 72    | 557 | 42  | 50  | 39  | 11,476 | 1,375  | 12% |
| 1991 | 2,457 | 767   | 98    | 58    | 479   | 445   | 1,445 | 544   | 1,255 | 308   | 429   | 90  | 239 | 34  | 26  | 8,673  | 835    | 10% |
| 1992 | 1,205 | 320   | 1,623 | 363   | 379   | 600   | 475   | 610   | 290   | 657   | 241   | 292 | 127 | 89  | 77  | 7,348  | 812    | 11% |
| 1993 | 1,580 | 290   | 984   | 3,417 | 624   | 475   | 280   | 434   | 580   | 361   | 329   | 235 | 155 | 102 | 122 | 9,968  | 927    | 9%  |
| 1994 | 914   | 448   | 431   | 1,280 | 3,196 | 625   | 155   | 163   | 155   | 286   | 166   | 249 | 87  | 76  | 130 | 8,362  | 973    | 12% |
| 1995 | 1,035 | 65    | 304   | 1,292 | 1,785 | 2,931 | 1,056 | 236   | 183   | 157   | 216   | 101 | 158 | 58  | 80  | 9,658  | 1,803  | 19% |
| 1996 | 1,385 | 331   | 126   | 353   | 829   | 1,087 | 1,075 | 337   | 88    | 82    | 68    | 129 | 35  | 74  | 80  | 6,079  | 498    | 8%  |
| 1997 | 2,254 | 268   | 149   | 172   | 2,317 | 1,026 | 679   | 817   | 130   | 46    | 55    | 68  | 77  | 28  | 101 | 8,186  | 1,111  | 14% |
| 1998 | 582   | 557   | 221   | 170   | 374   | 2,124 | 597   | 392   | 293   | 75    | 25    | 10  | 24  | 22  | 52  | 5,518  | 634    | 11% |
| 1999 | 779   | 649   | 579   | 688   | 402   | 669   | 1,967 | 547   | 312   | 261   | 109   | 37  | 17  | 23  | 71  | 7,111  | 834    | 12% |
| 2000 | 889   | 270   | 258   | 1,160 | 1,168 | 726   | 553   | 2,035 | 741   | 407   | 158   | 124 | 24  | 13  | 71  | 8,596  | 1,052  | 12% |
| 2001 | 1,421 | 777   | 405   | 414   | 1,010 | 1,158 | 447   | 244   | 777   | 574   | 207   | 172 | 60  | 25  | 64  | 7,756  | 695    | 9%  |
| 2002 | 614   | 311   | 541   | 788   | 949   | 1,229 | 681   | 349   | 431   | 872   | 420   | 192 | 114 | 34  | 35  | 7,559  | 763    | 10% |
| 2003 | 298   | 122   | 431   | 1,464 | 1,473 | 1,348 | 1,604 | 897   | 375   | 547   | 1,158 | 485 | 184 | 66  | 47  | 10,501 | 1,887  | 18% |
| 2004 | 293   | 212   | 121   | 946   | 1,103 | 798   | 465   | 512   | 241   | 153   | 151   | 283 | 120 | 28  | 22  | 5,449  | 501    | 9%  |
| 2005 | 336   | 119   | 155   | 837   | 2,227 | 1,643 | 876   | 385   | 296   | 236   | 60    | 121 | 214 | 79  | 79  | 7,663  | 754    | 10% |
| 2006 | 780   | 35    | 36    | 298   | 797   | 1,026 | 681   | 318   | 186   | 162   | 81    | 46  | 71  | 93  | 92  | 4,702  | 427    | 9%  |
| 2007 | 2,113 | 30    | 72    | 321   | 1,052 | 1,252 | 927   | 664   | 287   | 122   | 117   | 105 | 46  | 61  | 109 | 7,279  | 643    | 9%  |
| Avg  | 1,087 | 417   | 615   | 1,120 | 1,648 | 1,443 | 912   | 585   | 377   | 289   | 188   | 136 | 73  | 40  | 52  | 8,984  | 1,025  | 11% |

Table 1.12. Number of (non-YOY) hauls and sample sizes for EBS pollock collected by the EIT surveys.

| <b>Year</b> | <b>Stratum</b>                | <b>No. Hauls</b> | <b>No. lengths</b> | <b>No. otoliths collected</b> | <b>No. aged</b> |
|-------------|-------------------------------|------------------|--------------------|-------------------------------|-----------------|
| 1979        | <b>Total</b>                  | 25               | 7,722              | NA                            | 2,610           |
| 1982        | <b>Total</b>                  | 48               | 8,687              | 3,164                         | 2,741           |
|             | Midwater, east of St Paul     | 13               | 1,725              | 840                           | 783             |
|             | Midwater, west of St Paul     | 31               | 6,689              | 2,324                         | 1,958           |
|             | Bottom                        | 4                | 273                | 0                             | 0               |
| 1985        | <b>Total (Legs 1 &amp; 2)</b> | 73               | 19,872             | 2,739                         | 2,739           |
| 1988        | <b>Total</b>                  | 25               | 6,619              | 1,471                         | 1,471           |
| 1991        | <b>Total</b>                  | 62               | 16,343             | 2,062                         | 1,663           |
| 1994        | <b>Total (US zone)</b>        | 76               | 21,506             | 4,966                         | 1,770           |
|             | East of 170 W                 | 25               |                    | 1,550                         | 612             |
|             | West of 170 W                 | 51               |                    | 3,416                         | 1,158           |
|             | Navarin (Russia)              | 19               |                    | 1,017                         |                 |
| 1996        | <b>Total</b>                  | 57               | 16,824             | 1,949                         | 1,926           |
|             | East of 170 W                 | 15               | 3,551              | 669                           | 815             |
|             | West of 170 W                 | 42               | 13,273             | 1,280                         | 1,111           |
| 1997        | <b>Total</b>                  | 86               | 29,536             | 3,635                         | 2,285           |
|             | East of 170 W                 | 25               | 6,493              | 966                           | 936             |
|             | West of 170 W                 | 61               | 23,043             | 2,669                         | 1,349           |
| 1999        | <b>Total</b>                  | 118              | 42,362             | 4,946                         | 2,446           |
|             | East of 170 W                 | 41               | 13,841             | 1,945                         | 946             |
|             | West of 170 W                 | 77               | 28,521             | 3,001                         | 1,500           |
| 2000        | <b>Total</b>                  | 124              | 43,729             | 3,459                         | 2,253           |
|             | East of 170 W                 | 29               | 7,721              | 850                           | 850             |
|             | West of 170 W                 | 95               | 36,008             | 2,609                         | 1,403           |
| 2002        | <b>Total</b>                  | 126              | 40,234             | 3,307                         | 2,200           |
|             | East of 170 W                 | 47               | 14,601             | 1,424                         | 1,000           |
|             | West of 170 W                 | 79               | 25,633             | 1,883                         | 1,200           |
| 2004        | <b>Total (US zone)</b>        | 90               | 27,158             | 3,169                         | 2,351           |
|             | East of 170 W                 | 33               | 8,896              | 1,167                         | 798             |
|             | West of 170 W                 | 57               | 18,262             | 2,002                         | 1,192           |
|             | Navarin (Russia)              | 15               | 5,893              | 461                           | 461             |
| 2006        | <b>Total</b>                  | 83               | 24,265             | 2,693                         | 2,692           |
|             | East of 170 W                 | 27               | 4,939              | 822                           | 822             |
|             | West of 170 W                 | 56               | 19,326             | 1,871                         | 1,870           |
| 2007        | <b>Total (US zone)</b>        | 69               | 20,355             | 2,832                         | -               |
|             | East of 170 W                 | 23               | 5,492              | 871                           | -               |
|             | West of 170 W                 | 46               | 14,863             | 1,961                         | -               |
|             | Navarin (Russia)              | 4                | 1,407              | 398                           | -               |

Table 1.13. EIT survey estimates of EBS pollock abundance-at-age (millions), 1979-2007. *NOTE: 2007 age specific values are preliminary since they are derived from the bottom-trawl age-length key.*

| Year | Age    |        |       |       |       |       |       |     |     |     | Total   |
|------|--------|--------|-------|-------|-------|-------|-------|-----|-----|-----|---------|
|      | 1      | 2      | 3     | 4     | 5     | 6     | 7     | 8   | 9   | 10+ |         |
| 1979 | 69,110 | 41,132 | 3,884 | 413   | 534   | 128   | 30    | 4   | 28  | 161 | 115,424 |
| 1982 | 108    | 3,401  | 4,108 | 7,637 | 1,790 | 283   | 141   | 178 | 90  | 177 | 17,913  |
| 1985 | 2,076  | 929    | 8,149 | 898   | 2,186 | 1,510 | 1,127 | 130 | 21  | 15  | 17,041  |
| 1988 | 11     | 1,112  | 3,586 | 3,864 | 739   | 1,882 | 403   | 151 | 130 | 414 | 12,291  |
| 1991 | 639    | 5,942  | 967   | 215   | 224   | 133   | 120   | 39  | 37  | 53  | 8,369   |
| 1994 | 453    | 3,906  | 1,127 | 1,670 | 1,908 | 293   | 69    | 67  | 30  | 59  | 9,582   |
| 1996 | 972    | 446    | 520   | 2,686 | 821   | 509   | 434   | 85  | 17  | 34  | 6,525   |
| 1997 | 12,384 | 2,743  | 385   | 491   | 1,918 | 384   | 205   | 143 | 33  | 18  | 18,703  |
| 1999 | 112    | 1,588  | 3,597 | 1,684 | 583   | 274   | 1,169 | 400 | 105 | 90  | 9,601   |
| 2000 | 258    | 1,272  | 1,185 | 2,480 | 900   | 244   | 234   | 725 | 190 | 141 | 7,630   |
| 2002 | 561    | 4,188  | 3,841 | 1,295 | 685   | 593   | 288   | 100 | 132 | 439 | 12,122  |
| 2004 | 16     | 275    | 1,189 | 2,929 | 1,444 | 417   | 202   | 193 | 68  | 101 | 6,834   |
| 2006 | 456    | 209    | 282   | 610   | 695   | 552   | 320   | 110 | 53  | 110 | 3,396   |
| 2007 | 5,699  | 880    | 457   | 337   | 771   | 590   | 331   | 178 | 67  | 74  | 9,385   |

Table 1.14. Pollock biomass estimates by area from summer echo integration-trawl surveys on the U.S. EEZ portion of the Bering Sea shelf from near-surface to 3 m off bottom, 1994-2007. Survey-derived relative sampling errors for the biomass estimates are provided in the last column.

| Year | Survey Dates  | Area (nmi) <sup>2</sup> | Biomass in millions of t<br>(percent of total) |                  |                  |       | Total Biomass (t) | Relative estimation error |
|------|---------------|-------------------------|--|------------------|------------------|-------|-------------------|---------------------------|
|      |               |                         | SCA  | E170-SCA         | W170             |       |                   |                           |
| 1994 | 9 Jul-19 Aug  | 78,251                  | 0.312<br>(10.8%)                               | 0.399<br>(13.8%) | 2.176<br>(75.4%) | 2.886 | 0.047             |                           |
| 1996 | 20 Jul-30 Aug | 93,810                  | 0.215<br>(9.3%)                                | 0.269<br>(11.7%) | 1.826<br>(79.0%) | 2.311 | 0.039             |                           |
| 1997 | 17 Jul-4 Sept | 102,770                 | 0.246<br>(9.5%)                                | 0.527<br>(20.3%) | 1.818<br>(70.2%) | 2.591 | 0.037             |                           |
| 1999 | 7 Jun-5 Aug   | 103,670                 | 0.299<br>(9.1%)                                | 0.579<br>(17.6%) | 2.408<br>(73.2%) | 3.290 | 0.055             |                           |
| 2000 | 7 Jun-2 Aug   | 106,140                 | 0.393<br>(12.9%)                               | 0.498<br>(16.3%) | 2.158<br>(70.8%) | 3.049 | 0.032             |                           |
| 2002 | 4 Jun -30 Jul | 99,526                  | 0.647<br>(17.9%)                               | 0.797<br>(22.0%) | 2.178<br>(60.1%) | 3.622 | 0.031             |                           |
| 2004 | 4 Jun -29 Jul | 99,659                  | 0.498<br>(15.1%)                               | 0.516<br>(15.6%) | 2.293<br>(69.3%) | 3.307 | 0.037             |                           |
| 2006 | 3 Jun -25 Jul | 89,550                  | 0.131<br>(8.4%)                                | 0.254<br>(16.3%) | 1.175<br>(75.3%) | 1.560 | 0.039             |                           |
| 2007 | 2 Jun -30 Jul | 92,944                  | 0.084<br>(4.5%)                                | 0.168<br>(8.9%)  | 1.627<br>(86.6%) | 1.878 |                   |                           |

Key: SCA = Sea lion Conservation Area  
E170 - SCA = East of 170 W minus SCA  
W170 = West of 170 W

Table 1.15. Fishery annual average weights-at-age (kg) as estimated from NMFS observer data. These values are used in the model for computing the predicted fishery catch (in weight) and for computing biomass levels for EBS pollock. *NOTE: 2007 weight-at-age is treated as the three-year average of values from 2004-2006.*

|           | 1     | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    | 11    | 12    | 13    | 14    | 15    |
|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1964-1990 | 0.007 | 0.170 | 0.303 | 0.447 | 0.589 | 0.722 | 0.840 | 0.942 | 1.029 | 1.102 | 1.163 | 1.212 | 1.253 | 1.286 | 1.312 |
| 1991      | 0.007 | 0.170 | 0.288 | 0.476 | 0.604 | 0.723 | 0.844 | 0.883 | 1.001 | 1.126 | 1.121 | 1.240 | 1.229 | 1.272 | 1.227 |
| 1992      | 0.007 | 0.170 | 0.396 | 0.464 | 0.646 | 0.712 | 0.814 | 0.982 | 1.020 | 1.221 | 1.231 | 1.265 | 1.170 | 1.342 | 1.428 |
| 1993      | 0.007 | 0.170 | 0.495 | 0.610 | 0.652 | 0.769 | 0.932 | 1.051 | 1.201 | 1.237 | 1.408 | 1.536 | 1.606 | 1.671 | 1.614 |
| 1994      | 0.007 | 0.170 | 0.393 | 0.647 | 0.729 | 0.746 | 0.705 | 1.012 | 1.382 | 1.334 | 1.341 | 1.434 | 1.383 | 1.314 | 1.401 |
| 1995      | 0.007 | 0.170 | 0.375 | 0.500 | 0.729 | 0.842 | 0.855 | 0.970 | 1.215 | 1.329 | 1.422 | 1.495 | 1.390 | 1.243 | 1.365 |
| 1996      | 0.007 | 0.170 | 0.318 | 0.413 | 0.691 | 0.794 | 0.951 | 1.022 | 1.092 | 1.408 | 1.503 | 1.519 | 1.745 | 1.551 |       |
| 1997      | 0.007 | 0.170 | 0.311 | 0.466 | 0.554 | 0.747 | 0.892 | 1.075 | 1.089 | 1.240 | 1.424 | 1.471 | 1.713 | 1.458 | 1.416 |
| 1998      | 0.007 | 0.170 | 0.371 | 0.588 | 0.624 | 0.622 | 0.775 | 1.033 | 1.177 | 1.241 | 1.295 | 1.413 | 1.546 | 1.546 | 1.620 |
| 1999      | 0.007 | 0.170 | 0.397 | 0.501 | 0.638 | 0.703 | 0.728 | 0.905 | 1.045 | 1.275 | 1.211 | 1.418 | 1.277 | 1.152 | 1.314 |
| 2000      | 0.007 | 0.170 | 0.352 | 0.524 | 0.629 | 0.731 | 0.782 | 0.803 | 0.971 | 1.018 | 1.274 | 1.317 | 1.316 | 1.725 | 1.825 |
| 2001      | 0.007 | 0.170 | 0.322 | 0.497 | 0.669 | 0.786 | 0.964 | 0.994 | 1.059 | 1.134 | 1.327 | 1.451 | 1.581 | 1.463 | 1.660 |
| 2002      | 0.007 | 0.170 | 0.379 | 0.507 | 0.669 | 0.795 | 0.908 | 1.025 | 1.115 | 1.097 | 1.297 | 1.434 | 1.611 | 1.323 | 1.631 |
| 2003      | 0.007 | 0.170 | 0.485 | 0.548 | 0.649 | 0.767 | 0.862 | 0.953 | 1.085 | 1.221 | 1.213 | 1.223 | 1.444 | 1.342 | 1.759 |
| 2004      | 0.007 | 0.170 | 0.404 | 0.581 | 0.640 | 0.770 | 0.891 | 0.929 | 1.027 | 1.208 | 1.167 | 1.188 | 1.373 | 1.303 | 1.254 |
| 2005      | 0.007 | 0.170 | 0.351 | 0.507 | 0.640 | 0.740 | 0.878 | 0.947 | 1.062 | 1.104 | 1.273 | 1.313 | 1.316 | 1.163 | 1.423 |
| 2006      | 0.007 | 0.170 | 0.304 | 0.448 | 0.603 | 0.754 | 0.855 | 0.958 | 1.056 | 1.128 | 1.219 | 1.315 | 1.315 | 1.381 | 1.459 |
| 2007      | 0.007 | 0.170 | 0.353 | 0.512 | 0.627 | 0.754 | 0.875 | 0.945 | 1.048 | 1.147 | 1.220 | 1.272 | 1.334 | 1.282 | 1.379 |

Table 1.16. Pollock sample sizes assumed for the age-composition data likelihoods from the fishery, bottom-trawl survey, and EIT surveys, 1964-2007. The 2007 EIT sample size was down-weighted of the value since the BTS age-length key was used.

| Year      | Fishery | Year          | BTS | EIT       |
|-----------|---------|---------------|-----|-----------|
| 1964-1977 | 10      | 1979 and 1982 | -   | 6         |
| 1978-1990 | 50      |               |     |           |
| 1991      | 179     |               |     |           |
| 1992      | 207     | 1982-2007     | 100 | 55        |
| 1993      | 281     |               |     | (average) |
| 1994      | 111     |               |     |           |
| 1995      | 142     |               |     |           |
| 1996      | 154     |               |     |           |
| 1997      | 265     |               |     |           |
| 1998      | 278     |               |     |           |
| 1999      | 470     |               |     |           |
| 2000      | 467     |               |     |           |
| 2001      | 301     |               |     |           |
| 2002      | 449     |               |     |           |
| 2003      | 402     |               |     |           |
| 2004      | 343     |               |     |           |
| 2005      | 412     |               |     |           |
| 2006      | 339     |               |     |           |

Table 1.17. Summary model results showing the stock condition for EBS pollock. Values in parentheses are coefficients of variation (CV's) of values immediately above.

| <b>Biomass (thousands of t)</b>                                 |              |
|---|--------------|
| Year 2008 spawning biomass *                                    | <b>1,380</b> |
| (CV)  | <b>(20%)</b> |
| 2007 spawning biomass   | <b>1,946</b> |
| $B_{msy}$   | <b>1,876</b> |
| (CV)  | <b>(22%)</b> |
| $B_{40\%}$  | <b>2,627</b> |
| (CV)  | <b>(5%)</b>  |
| $B_{35\%}$  | 2,299        |
| $B_0$ (stock-recruitment curve)                                 | <b>5,013</b> |
| 2008 Percent of $B_{msy}$ spawning biomass                      | <b>72%</b>   |
| 2008 Percent of $B_{40\%}$ spawning biomass                     | <b>57%</b>   |
| Ratio of $B_{2007}$ over $B_{2007}$ under no fishing since 1978 | <b>48%</b>   |
| 2009 Fishable biomass   | <b>3,730</b> |
| Ratio $B_{2009}/B_{2008}$ (fishable biomass)                    | <b>78%</b>   |
| <b>Recruitment (millions of pollock at age 1)</b>               |              |
| Steepness parameter ( $h$ )                                     | 0.68         |
| Average recruitment (all yrs)                                   | 22,032       |
| (CV)  | 64%          |
| Average recruitment (since 1978)                                | 23,690       |
| (CV since 1978)   | 68%          |
| 2000 year class   | 41,060       |
| (CV 2000 year class)  | (8%)         |
| Natural Mortality (age 3 and older)                             | <b>0.3</b>   |

Table 1.18. Summary results of Tier 1 yield projections for EBS pollock.

| <b>Yield projections</b>     |       |
|------------------------------|-------|
| $B_{msy}$ (fishable biomass) | 5,467 |
| 2008 "fishable" biomass (GM) | 4,767 |
| MSYR (HM)                    | 0.341 |
| 2008 MSYR yield (Tier 1 ABC) | 1,170 |
| MSYR (AM)                    | 0.422 |
| 2008 MSYR OFL                | 1,443 |

Notes: MSYR = exploitation rate relative to begin-year age fishable biomass corresponding to  $F_{msy}$ .  $F_{msy}$  yields calculated within the model (i.e., including uncertainty in both the estimate of  $F_{msy}$  and in projected stock size). HM = Harmonic mean, GM = Geometric mean, AM = Arithmetic mean

\*Assuming 2008 catch will be 1,170,00 t



Table 1.19 Estimates of numbers at age for the EBS pollock stock as estimated in 2007 (millions).

|         | <b>1</b> | <b>2</b> | <b>3</b> | <b>4</b> | <b>5</b> | <b>6</b> | <b>7</b> | <b>8</b> | <b>9</b> | <b>10+</b> | <b>Total</b> |
|---------|----------|----------|----------|----------|----------|----------|----------|----------|----------|------------|--------------|
| 1964    | 2,991    | 3,665    | 2,065    | 508      | 233      | 357      | 145      | 61       | 34       | 174        | 10,232       |
| 1965    | 20,529   | 1,214    | 2,306    | 1,436    | 307      | 142      | 222      | 93       | 40       | 142        | 26,431       |
| 1966    | 13,053   | 8,335    | 764      | 1,605    | 870      | 187      | 88       | 143      | 61       | 111        | 25,218       |
| 1967    | 31,519   | 5,301    | 5,226    | 518      | 990      | 545      | 121      | 58       | 94       | 93         | 44,464       |
| 1968    | 28,971   | 12,790   | 3,280    | 3,299    | 276      | 543      | 314      | 71       | 35       | 84         | 49,662       |
| 1969    | 29,855   | 11,757   | 7,891    | 2,081    | 1,803    | 157      | 320      | 188      | 43       | 87         | 54,183       |
| 1970    | 21,139   | 12,117   | 7,263    | 5,034    | 1,149    | 1,038    | 94       | 193      | 115      | 67         | 48,208       |
| 1971    | 8,239    | 8,576    | 7,326    | 4,362    | 2,935    | 647      | 598      | 53       | 110      | 49         | 32,895       |
| 1972    | 10,242   | 3,341    | 5,111    | 4,154    | 2,381    | 1,531    | 348      | 317      | 28       | 57         | 27,509       |
| 1973    | 27,727   | 4,153    | 1,923    | 2,663    | 2,074    | 1,200    | 785      | 178      | 163      | 58         | 40,925       |
| 1974    | 20,578   | 11,235   | 2,317    | 900      | 1,178    | 929      | 549      | 360      | 82       | 32         | 38,160       |
| 1975    | 18,121   | 8,338    | 6,025    | 904      | 367      | 492      | 398      | 236      | 153      | 44         | 35,077       |
| 1976    | 13,429   | 7,348    | 4,632    | 2,676    | 416      | 172      | 236      | 191      | 112      | 36         | 29,247       |
| 1977    | 13,980   | 5,449    | 4,249    | 2,377    | 1,216    | 194      | 83       | 115      | 93       | 52         | 27,808       |
| 1978    | 28,035   | 5,675    | 3,211    | 2,340    | 1,186    | 620      | 101      | 44       | 61       | 56         | 41,328       |
| 1979    | 64,073   | 11,381   | 3,382    | 1,768    | 1,202    | 574      | 309      | 51       | 22       | 52         | 82,813       |
| 1980    | 25,898   | 26,014   | 6,816    | 1,901    | 932      | 599      | 294      | 160      | 26       | 42         | 62,682       |
| 1981    | 29,690   | 10,521   | 16,220   | 4,392    | 1,015    | 451      | 288      | 145      | 79       | 26         | 62,827       |
| 1982    | 15,979   | 12,065   | 6,615    | 11,015   | 2,652    | 576      | 255      | 165      | 83       | 22         | 49,427       |
| 1983    | 54,323   | 6,495    | 7,647    | 4,731    | 7,218    | 1,628    | 341      | 153      | 99       | 40         | 82,673       |
| 1984    | 13,136   | 22,081   | 4,122    | 5,511    | 3,183    | 4,612    | 1,011    | 214      | 96       | 55         | 54,020       |
| 1985    | 35,237   | 5,340    | 14,020   | 2,964    | 3,791    | 2,039    | 2,843    | 620      | 131      | 69         | 67,054       |
| 1986    | 13,056   | 14,324   | 3,391    | 10,088   | 2,042    | 2,435    | 1,261    | 1,750    | 380      | 75         | 48,801       |
| 1987    | 8,338    | 5,307    | 9,089    | 2,447    | 6,944    | 1,357    | 1,517    | 773      | 1,091    | 90         | 36,954       |
| 1988    | 4,783    | 3,390    | 3,373    | 6,609    | 1,720    | 4,760    | 888      | 981      | 506      | 200        | 27,210       |
| 1989    | 10,161   | 1,945    | 2,151    | 2,411    | 4,523    | 1,122    | 3,042    | 529      | 606      | 551        | 27,041       |
| 1990    | 50,836   | 4,131    | 1,234    | 1,538    | 1,649    | 2,947    | 716      | 1,809    | 326      | 526        | 65,713       |
| 1991    | 26,352   | 20,666   | 2,617    | 887      | 983      | 991      | 1,706    | 382      | 1,024    | 511        | 56,118       |
| 1992    | 22,593   | 10,712   | 13,086   | 1,878    | 562      | 584      | 566      | 893      | 213      | 383        | 51,469       |
| 1993    | 51,772   | 9,184    | 6,798    | 9,096    | 1,099    | 342      | 313      | 282      | 430      | 475        | 79,791       |
| 1994    | 14,648   | 21,047   | 5,837    | 4,828    | 5,763    | 715      | 205      | 178      | 157      | 336        | 53,713       |
| 1995    | 9,990    | 5,955    | 13,383   | 4,269    | 3,329    | 3,411    | 364      | 112      | 102      | 350        | 41,265       |
| 1996    | 24,079   | 4,061    | 3,789    | 9,815    | 2,991    | 2,071    | 1,889    | 213      | 68       | 281        | 49,258       |
| 1997    | 32,160   | 9,789    | 2,574    | 2,752    | 6,941    | 2,033    | 1,239    | 991      | 116      | 204        | 58,801       |
| 1998    | 14,840   | 13,074   | 6,207    | 1,873    | 1,953    | 4,749    | 1,237    | 668      | 553      | 146        | 45,300       |
| 1999    | 16,669   | 6,033    | 8,310    | 4,412    | 1,306    | 1,311    | 2,949    | 740      | 392      | 132        | 42,254       |
| 2000    | 26,483   | 6,777    | 3,836    | 5,926    | 3,092    | 884      | 826      | 1,796    | 443      | 289        | 50,351       |
| 2001    | 41,061   | 10,767   | 4,314    | 2,786    | 4,055    | 1,974    | 537      | 479      | 1,028    | 330        | 67,330       |
| 2002    | 21,082   | 16,693   | 6,852    | 3,119    | 1,872    | 2,504    | 1,148    | 295      | 259      | 362        | 54,187       |
| 2003    | 11,623   | 8,571    | 10,613   | 4,872    | 2,135    | 1,137    | 1,321    | 609      | 156      | 574        | 41,611       |
| 2004    | 4,215    | 4,725    | 5,448    | 7,523    | 3,315    | 1,277    | 585      | 683      | 314      | 431        | 28,515       |
| 2005    | 4,446    | 1,714    | 3,007    | 3,949    | 4,888    | 1,988    | 707      | 319      | 376      | 321        | 21,715       |
| 2006    | 11,468   | 1,808    | 1,091    | 2,178    | 2,556    | 2,912    | 1,091    | 383      | 174      | 308        | 23,969       |
| 2007    | 42,147   | 4,662    | 1,150    | 780      | 1,390    | 1,403    | 1,630    | 542      | 192      | 298        | 54,194       |
| Median  | 20,553   | 7,841    | 4,473    | 2,769    | 1,838    | 1,080    | 558      | 259      | 123      | 122        | 46,754       |
| Average | 22,490   | 8,830    | 5,467    | 3,663    | 2,329    | 1,412    | 806      | 437      | 242      | 196        | 45,873       |

Table 1.20. Estimated catch-at-age of EBS pollock (millions).

| Year    | 1    | 2       | 3       | 4       | 5       | 6     | 7     | 8     | 9     | 10+   | Total   |
|---------|------|---------|---------|---------|---------|-------|-------|-------|-------|-------|---------|
| 1964    | 2.6  | 38.9    | 109.3   | 81.4    | 36.1    | 49.2  | 17.1  | 6.2   | 3.0   | 15.7  | 359.6   |
| 1965    | 17.8 | 12.7    | 120.7   | 227.5   | 47.1    | 19.3  | 25.9  | 9.3   | 3.5   | 12.1  | 496.2   |
| 1966    | 9.4  | 112.2   | 56.5    | 233.7   | 116.4   | 21.3  | 9.1   | 13.2  | 5.1   | 10.5  | 587.4   |
| 1967    | 40.9 | 127.1   | 670.0   | 126.5   | 223.4   | 105.7 | 21.2  | 9.2   | 13.8  | 17.8  | 1,355.7 |
| 1968    | 35.2 | 333.3   | 408.4   | 753.7   | 55.2    | 96.2  | 52.2  | 11.2  | 5.3   | 21.0  | 1,771.8 |
| 1969    | 35.0 | 295.7   | 950.4   | 460.8   | 349.5   | 27.0  | 51.6  | 28.7  | 6.3   | 15.5  | 2,220.4 |
| 1970    | 29.4 | 505.9   | 1,193.9 | 932.4   | 240.5   | 200.6 | 18.9  | 38.8  | 22.8  | 23.9  | 3,207.0 |
| 1971    | 14.6 | 453.3   | 1,494.9 | 999.3   | 756.5   | 154.4 | 148.6 | 13.2  | 27.0  | 32.6  | 4,094.4 |
| 1972    | 17.5 | 262.4   | 1,320.7 | 1,182.1 | 663.9   | 411.4 | 93.2  | 83.9  | 7.6   | 32.6  | 4,075.2 |
| 1973    | 61.9 | 420.2   | 619.1   | 939.2   | 717.2   | 400.9 | 261.4 | 58.8  | 54.8  | 25.9  | 3,559.4 |
| 1974    | 46.0 | 1,446.9 | 963.3   | 355.0   | 450.0   | 343.1 | 201.7 | 134.8 | 31.7  | 41.6  | 4,014.1 |
| 1975    | 32.3 | 869.5   | 2,110.9 | 299.3   | 117.4   | 152.1 | 122.1 | 74.1  | 49.3  | 25.7  | 3,852.7 |
| 1976    | 17.8 | 553.1   | 1,240.7 | 905.0   | 134.7   | 52.5  | 70.1  | 56.8  | 33.8  | 32.0  | 3,096.4 |
| 1977    | 15.0 | 334.2   | 948.9   | 676.6   | 330.9   | 49.6  | 20.7  | 28.6  | 23.5  | 26.6  | 2,454.5 |
| 1978    | 27.6 | 299.1   | 717.8   | 625.1   | 359.4   | 176.5 | 28.5  | 12.9  | 17.9  | 30.5  | 2,295.4 |
| 1979    | 58.6 | 558.8   | 709.3   | 443.9   | 343.0   | 153.8 | 81.6  | 14.2  | 6.0   | 22.3  | 2,391.5 |
| 1980    | 14.6 | 463.7   | 768.5   | 462.6   | 282.5   | 183.7 | 86.7  | 46.2  | 7.4   | 15.4  | 2,331.1 |
| 1981    | 10.5 | 117.6   | 1,170.7 | 705.8   | 206.4   | 92.8  | 56.8  | 27.9  | 15.2  | 7.5   | 2,411.2 |
| 1982    | 3.1  | 57.9    | 197.6   | 1,102.3 | 394.9   | 100.2 | 42.4  | 27.6  | 14.1  | 12.0  | 1,952.1 |
| 1983    | 8.3  | 24.5    | 180.2   | 376.8   | 860.8   | 227.5 | 45.6  | 20.5  | 13.4  | 13.0  | 1,770.6 |
| 1984    | 1.7  | 74.5    | 104.2   | 340.8   | 373.0   | 671.1 | 151.2 | 32.7  | 15.5  | 20.9  | 1,785.5 |
| 1985    | 4.4  | 17.7    | 348.0   | 180.1   | 436.8   | 291.8 | 418.1 | 93.2  | 20.8  | 22.5  | 1,833.3 |
| 1986    | 1.3  | 55.0    | 75.4    | 618.6   | 181.9   | 336.0 | 189.2 | 240.7 | 64.1  | 27.6  | 1,789.9 |
| 1987    | 0.6  | 14.6    | 145.1   | 108.5   | 449.1   | 137.1 | 167.0 | 77.9  | 135.5 | 40.6  | 1,275.9 |
| 1988    | 0.4  | 12.5    | 101.8   | 436.6   | 178.2   | 567.5 | 151.1 | 141.2 | 72.2  | 129.3 | 1,790.8 |
| 1989    | 0.8  | 7.2     | 65.4    | 160.5   | 471.9   | 134.7 | 521.1 | 76.7  | 87.1  | 133.3 | 1,658.9 |
| 1990    | 4.4  | 21.9    | 31.4    | 183.2   | 269.9   | 560.5 | 174.8 | 371.0 | 72.3  | 184.1 | 1,873.5 |
| 1991    | 2.4  | 115.6   | 70.2    | 111.3   | 169.1   | 197.9 | 436.3 | 82.2  | 238.0 | 151.2 | 1,574.1 |
| 1992    | 2.1  | 41.4    | 698.2   | 342.8   | 87.3    | 140.4 | 161.7 | 272.5 | 64.8  | 259.9 | 2,071.0 |
| 1993    | 3.2  | 23.6    | 243.2   | 1,142.7 | 116.7   | 57.4  | 63.1  | 60.9  | 92.6  | 107.8 | 1,911.2 |
| 1994    | 0.7  | 46.8    | 64.7    | 288.7   | 1,006.4 | 194.4 | 46.4  | 34.7  | 27.3  | 90.2  | 1,800.2 |
| 1995    | 0.4  | 10.3    | 115.6   | 200.2   | 463.0   | 749.6 | 66.3  | 17.4  | 14.2  | 54.5  | 1,691.6 |
| 1996    | 1.0  | 19.7    | 63.4    | 386.0   | 213.6   | 345.7 | 479.8 | 49.0  | 14.6  | 60.2  | 1,633.1 |
| 1997    | 1.3  | 43.8    | 39.8    | 100.1   | 459.1   | 315.7 | 294.2 | 212.7 | 23.2  | 39.3  | 1,529.2 |
| 1998    | 0.5  | 33.9    | 217.9   | 94.7    | 159.2   | 666.2 | 206.7 | 120.4 | 95.8  | 29.7  | 1,625.1 |
| 1999    | 0.5  | 14.4    | 268.3   | 205.4   | 98.1    | 170.0 | 456.3 | 123.6 | 62.9  | 60.4  | 1,459.8 |
| 2000    | 0.8  | 8.7     | 65.1    | 392.2   | 371.2   | 137.6 | 155.9 | 354.4 | 69.4  | 69.8  | 1,625.2 |
| 2001    | 1.5  | 16.9    | 89.3    | 223.5   | 586.0   | 368.2 | 121.0 | 112.6 | 193.1 | 95.5  | 1,807.9 |
| 2002    | 0.9  | 38.5    | 238.3   | 205.4   | 293.3   | 627.5 | 284.4 | 73.7  | 58.3  | 159.9 | 1,980.1 |
| 2003    | 0.5  | 21.3    | 396.6   | 344.2   | 357.4   | 303.1 | 348.0 | 161.8 | 37.4  | 117.6 | 2,087.7 |
| 2004    | 0.1  | 7.3     | 101.3   | 801.8   | 548.6   | 280.2 | 134.0 | 152.4 | 60.7  | 70.6  | 2,157.1 |
| 2005    | 0.2  | 2.7     | 57.6    | 433.0   | 831.4   | 447.8 | 166.4 | 73.1  | 74.8  | 73.9  | 2,160.8 |
| 2006    | 0.5  | 3.7     | 32.5    | 261.5   | 576.4   | 618.8 | 313.5 | 107.3 | 44.1  | 105.3 | 2,063.6 |
| 2007    | 2.2  | 11.0    | 39.5    | 106.9   | 354.2   | 337.1 | 525.7 | 170.8 | 54.9  | 87.0  | 1,689.3 |
| Median  | 2.9  | 42.6    | 207.7   | 349.6   | 346.2   | 196.2 | 141.3 | 67.0  | 29.5  | 32.6  | 1,853.4 |
| Average | 12.1 | 180.7   | 446.0   | 444.5   | 357.7   | 265.3 | 170.2 | 89.3  | 46.7  | 59.7  | 2,072.1 |

Table 1.21. Estimated EBS pollock age 3+ biomass, female spawning biomass, and age 1 recruitment for 1964-2007. Biomass units are thousands of t, age-1 recruitment is in millions of pollock.

| Year | Age 3+ biomass | Spawning biomass | Age 1 Rec. | Year | Age 3+ biomass | Spawning biomass | Age 1 Rec. |
|------|----------------|------------------|------------|------|----------------|------------------|------------|
| 1964 | 1,717          | 510              | 2,991      | 1986 | 11,773         | 4,016            | 13,056     |
| 1965 | 2,141          | 609              | 20,529     | 1987 | 12,401         | 4,158            | 8,338      |
| 1966 | 2,037          | 680              | 13,053     | 1988 | 11,617         | 4,139            | 4,783      |
| 1967 | 3,206          | 815              | 31,519     | 1989 | 9,875          | 3,718            | 10,161     |
| 1968 | 3,558          | 985              | 28,971     | 1990 | 7,847          | 2,994            | 50,836     |
| 1969 | 5,118          | 1,272            | 29,855     | 1991 | 6,097          | 2,229            | 26,352     |
| 1970 | 6,368          | 1,664            | 21,139     | 1992 | 9,557          | 2,344            | 22,593     |
| 1971 | 7,164          | 1,945            | 8,239      | 1993 | 11,832         | 3,238            | 51,772     |
| 1972 | 6,666          | 1,918            | 10,242     | 1994 | 11,485         | 3,562            | 14,648     |
| 1973 | 4,942          | 1,550            | 27,727     | 1995 | 13,615         | 3,828            | 9,990      |
| 1974 | 3,475          | 1,059            | 20,578     | 1996 | 11,537         | 3,853            | 24,079     |
| 1975 | 3,604          | 856              | 18,121     | 1997 | 10,104         | 3,670            | 32,160     |
| 1976 | 3,584          | 871              | 13,429     | 1998 | 10,178         | 3,383            | 14,840     |
| 1977 | 3,602          | 922              | 13,980     | 1999 | 11,081         | 3,397            | 16,669     |
| 1978 | 3,476          | 950              | 28,035     | 2000 | 10,201         | 3,417            | 26,483     |
| 1979 | 3,363          | 933              | 64,073     | 2001 | 9,898          | 3,427            | 41,061     |
| 1980 | 4,384          | 1,067            | 25,898     | 2002 | 10,224         | 3,226            | 21,082     |
| 1981 | 8,307          | 1,763            | 29,690     | 2003 | 12,865         | 3,472            | 11,623     |
| 1982 | 9,439          | 2,666            | 15,979     | 2004 | 11,784         | 3,613            | 4,215      |
| 1983 | 10,493         | 3,268            | 54,323     | 2005 | 9,598          | 3,264            | 4,446      |
| 1984 | 10,200         | 3,476            | 13,136     | 2006 | 7,178          | 2,617            | 11,468     |
| 1985 | 12,531         | 3,757            | 35,237     | 2007 | 5,363          | 1,946            | 42,147     |
|      |                |                  |            | 2008 | 4,357          | 1,398            |            |
|      |                |                  |            | 2009 | 6,658          | 1,533            |            |



Table 1.23 Projections of catch, fishing mortality, and spawning biomass (thousands of tons) for EBS pollock for the 7 scenarios. Note that the values for  $B_{100\%}$ ,  $B_{40\%}$ , and  $B_{35\%}$  are 6,569; **2,627**; and 2,299 thousand t, respectively.

| <i>Catch (1,000 t)</i>            | <i>Scenario 1</i> | <i>Scenario 2</i> | <i>Scenario 3</i> | <i>Scenario 4</i> | <i>Scenario 5</i> | <i>Scenario 6</i> | <i>Scenario 7</i> |
|-----------------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| 2007                              | 1,340             | 1,340             | 1,340             | 1,340             | 1,340             | 1,340             | 1,340             |
| 2008                              | 555               | 1,000             | 454               | 457               | 0                 | 677               | 555               |
| 2009                              | 650               | 1,200             | 454               | 457               | 0                 | 748               | 650               |
| 2010                              | 1,186             | 1,200             | 637               | 641               | 0                 | 1,353             | 1,445             |
| 2011                              | 1,558             | 1,200             | 854               | 859               | 0                 | 1,752             | 1,786             |
| 2012                              | 1,634             | 1,200             | 976               | 981               | 0                 | 1,785             | 1,797             |
| 2013                              | 1,711             | 1,200             | 1,098             | 1,103             | 0                 | 1,832             | 1,835             |
| 2014                              | 1,687             | 1,200             | 1,137             | 1,142             | 0                 | 1,788             | 1,789             |
| 2015                              | 1,657             | 1,200             | 1,146             | 1,151             | 0                 | 1,747             | 1,748             |
| 2016                              | 1,637             | 1,200             | 1,146             | 1,151             | 0                 | 1,723             | 1,723             |
| 2017                              | 1,634             | 1,200             | 1,149             | 1,153             | 0                 | 1,725             | 1,725             |
| 2018                              | 1,642             | 1,200             | 1,155             | 1,160             | 0                 | 1,738             | 1,738             |
| 2019                              | 1,652             | 1,200             | 1,163             | 1,167             | 0                 | 1,748             | 1,748             |
| 2020                              | 1,648             | 1,200             | 1,165             | 1,169             | 0                 | 1,743             | 1,743             |
| <i>Fishing M.</i>                 | <i>Scenario 1</i> | <i>Scenario 2</i> | <i>Scenario 3</i> | <i>Scenario 4</i> | <i>Scenario 5</i> | <i>Scenario 6</i> | <i>Scenario 7</i> |
| 2007                              | 0.455             | 0.455             | 0.455             | 0.455             | 0.455             | 0.455             | 0.455             |
| 2008                              | 0.258             | 0.510             | 0.207             | 0.209             | 0.000             | 0.323             | 0.258             |
| 2009                              | 0.316             | 0.758             | 0.207             | 0.209             | 0.000             | 0.385             | 0.316             |
| 2010                              | 0.431             | 0.547             | 0.207             | 0.209             | 0.000             | 0.523             | 0.539             |
| 2011                              | 0.455             | 0.415             | 0.207             | 0.209             | 0.000             | 0.554             | 0.558             |
| 2012                              | 0.451             | 0.381             | 0.207             | 0.209             | 0.000             | 0.545             | 0.547             |
| 2013                              | 0.449             | 0.343             | 0.207             | 0.209             | 0.000             | 0.542             | 0.542             |
| 2014                              | 0.445             | 0.328             | 0.207             | 0.209             | 0.000             | 0.535             | 0.535             |
| 2015                              | 0.444             | 0.319             | 0.207             | 0.209             | 0.000             | 0.532             | 0.532             |
| 2016                              | 0.444             | 0.313             | 0.207             | 0.209             | 0.000             | 0.530             | 0.531             |
| 2017                              | 0.444             | 0.306             | 0.207             | 0.209             | 0.000             | 0.531             | 0.531             |
| 2018                              | 0.444             | 0.294             | 0.207             | 0.209             | 0.000             | 0.532             | 0.532             |
| 2019                              | 0.444             | 0.286             | 0.207             | 0.209             | 0.000             | 0.532             | 0.532             |
| 2020                              | 0.444             | 0.283             | 0.207             | 0.209             | 0.000             | 0.532             | 0.532             |
| <i>Spawning Biomass (1,000 t)</i> | <i>Scenario 1</i> | <i>Scenario 2</i> | <i>Scenario 3</i> | <i>Scenario 4</i> | <i>Scenario 5</i> | <i>Scenario 6</i> | <i>Scenario 7</i> |
| 2007                              | 1,945             | 1,945             | 1,945             | 1,945             | 1,945             | 1,945             | 1,945             |
| 2008                              | 1,462             | 1,396             | 1,476             | 1,475             | 1,534             | 1,445             | 1,462             |
| 2009                              | 1,758             | 1,502             | 1,824             | 1,823             | 2,066             | 1,695             | 1,758             |
| 2010                              | 2,374             | 1,999             | 2,552             | 2,550             | 2,969             | 2,272             | 2,342             |
| 2011                              | 2,692             | 2,400             | 3,105             | 3,102             | 3,766             | 2,528             | 2,557             |
| 2012                              | 2,786             | 2,698             | 3,452             | 3,447             | 4,412             | 2,562             | 2,574             |
| 2013                              | 2,818             | 2,953             | 3,685             | 3,679             | 4,955             | 2,561             | 2,565             |
| 2014                              | 2,785             | 3,134             | 3,783             | 3,775             | 5,332             | 2,515             | 2,517             |
| 2015                              | 2,752             | 3,267             | 3,836             | 3,827             | 5,636             | 2,478             | 2,479             |
| 2016                              | 2,740             | 3,383             | 3,880             | 3,871             | 5,890             | 2,467             | 2,468             |
| 2017                              | 2,746             | 3,485             | 3,918             | 3,908             | 6,082             | 2,475             | 2,476             |
| 2018                              | 2,762             | 3,579             | 3,955             | 3,945             | 6,236             | 2,492             | 2,492             |
| 2019                              | 2,769             | 3,652             | 3,980             | 3,970             | 6,354             | 2,497             | 2,497             |
| 2020                              | 2,755             | 3,695             | 3,978             | 3,968             | 6,415             | 2,482             | 2,482             |

Table 1.24 Tier 1b EBS pollock ABC and OFL projections for 2008 and for 2009. **Units are thousands of tons.**

| Year |                    | ABC   | OFL   |
|------|--------------------|-------|-------|
| 2008 |                    | 1,170 | 1,443 |
|      | Assumed 2008 catch | ABC   | OFL   |
| 2009 | 1,170              | 976   | 1,204 |
| 2009 | 1,000              | 1,073 | 1,323 |

Table 1.25 Tier 1 (approximation) mean projections of female spawning biomass for EBS pollock under different assumptions of 2006 year class (as estimated or set to average) and catch scenarios (1.0, 1.2, million t in columns 2-5, and Tier 1 maximum permissible in last two columns with values in parentheses). **Units are thousands of tons.**

| Year | As estimated<br>Catch=1.0 | Average<br>Catch=1.0 | As estimated<br>Catch=1.2 | Average<br>Catch=1.2 | As estimated<br>(catch) | Average<br>(catch) |
|------|---------------------------|----------------------|---------------------------|----------------------|-------------------------|--------------------|
| 2009 | 1,531                     | 1,321                | 1,422                     | 1,209                | 1,529 (976)             | 1,345 (792)        |
| 2010 | 2,118                     | 1,628                | 1,948                     | 1,456                | 2,044 (1,588)           | 1,671 (1,160)      |
| 2011 | 2,603                     | 2,034                | 2,365                     | 1,799                | 2,240 (1,889)           | 1,954 (1,513)      |
| 2012 | 2,919                     | 2,386                | 2,619                     | 2,091                | 2,254 (1,885)           | 2,116 (1,700)      |
| 2013 | 3,134                     | 2,658                | 2,783                     | 2,311                | 2,252 (1,913)           | 2,195 (1,826)      |
| 2014 | 3,269                     | 2,871                | 2,884                     | 2,483                | 2,217 (1,871)           | 2,200 (1,846)      |
| 2015 | 3,412                     | 3,077                | 2,998                     | 2,658                | 2,185 (1,840)           | 2,180 (1,833)      |
| 2016 | 3,578                     | 3,297                | 3,141                     | 2,852                | 2,174 (1,822)           | 2,172 (1,822)      |
| 2017 | 3,721                     | 3,489                | 3,265                     | 3,023                | 2,182 (1,826)           | 2,181 (1,825)      |
| 2018 | 3,826                     | 3,638                | 3,357                     | 3,155                | 2,197 (1,834)           | 2,197 (1,833)      |
| 2019 | 3,905                     | 3,751                | 3,425                     | 3,257                | 2,203 (1,844)           | 2,203 (1,843)      |
| 2020 | 3,979                     | 3,857                | 3,493                     | 3,355                | 2,189 (1,834)           | 2,189 (1,834)      |

Table 1.26. Analysis of ecosystem considerations for BSAI pollock and the pollock fishery.

| Indicator  | Observation   | Interpretation  | Evaluation   |
|--|---|---|--|
| <b>Ecosystem effects on EBS pollock</b>                      |   |   |  |
| <i>Prey availability or abundance trends</i>                 |   |   |  |
| Zooplankton  | Stomach contents, ichthyoplankton surveys, changes mean wt-at-age                     | Data limited, indication of recent declines (especially in summer 2006) | Growing concern<br>Scarcity in inner and middle domain |
| <i>Predator population trends</i>                            |   |   |  |
| Marine mammals   | Fur seals declining, Steller sea lions increasing slightly                            | Possibly lower mortality on pollock                                     | Probably no concern                                    |
| Birds  | Stable, some increasing some decreasing   | Affects young-of-year mortality   | Probably no concern                                    |
| Fish (Pollock, Pacific cod, halibut)                         | Stable to increasing  | Possible increases to pollock mortality                                 |  |
| <i>Changes in habitat quality</i>                            |   |   |  |
| Temperature regime   |   |   | No concern (dealt with in model)                       |
| Winter-spring environmental conditions                       | Cold years pollock distribution towards NW on average<br>Affects pre-recruit survival | Likely to affect surveyed stock   |  |
| Production   | Fairly stable nutrient flow from upwelled BS Basin                                    | Probably a number of factors<br>Inter-annual variability low            | Causes natural variability<br>No concern               |
| <b>Fishery effects on ecosystem</b>                          |   |   |  |
| <i>Fishery contribution to bycatch</i>                       |   |   |  |
| Prohibited species   | Stable, heavily monitored   | Likely to be safe   | No concern   |
| Forage (including herring, Atka mackerel, cod, and pollock)  | Stable, heavily monitored   | Likely to be safe   | No concern   |
| HAPC biota   | Likely minor impact   | Likely to be safe   | No concern   |
| Marine mammals and birds                                     | Very minor direct-take  | Safe  | No concern   |
| Sensitive non-target species                                 | Likely minor impact   |   | No concern   |
|  |   | Data limited, likely to be safe   |  |
| <i>Fishery concentration in space and time</i>               | Generally more diffuse  | Mixed potential impact (fur seals vs Steller sea lions)                 | Possible concern                                       |
| <i>Fishery effects on amount of large size target fish</i>   | Depends on highly variable year-class strength  | Natural fluctuation   | Probably no concern                                    |
| <i>Fishery contribution to discards and offal production</i> | Decreasing  | Improving, but data limited   | Possible concern                                       |
| <i>Fishery effects on age-at-maturity and fecundity</i>      | Maturity study (gonad collection) underway  | NA  | Possible concern                                       |

Table 1.27 Bycatch estimates (t) of non-target species caught in the BSAI directed pollock fishery, 1997-2002 based on observer data, 2003-2006 based on observer data as processed through the catch accounting system (NMFS Regional Office, Juneau, Alaska).

|                  | 1997  | 1998  | 1999  | 2000  | 2001  | 2002  |                | 2003  | 2004  | 2005  | 2006  |
|------------------|-------|-------|-------|-------|-------|-------|----------------|-------|-------|-------|-------|
| Jellyfish        | 6,632 | 6,129 | 6,176 | 9,361 | 3,095 | 1,530 | Jellyfish      | 5,644 | 6,040 | 5,183 | 2,649 |
| Squid            | 1,487 | 1,210 | 474   | 379   | 1,776 | 1,708 | Squid          | 1,151 | 855   | 1,041 | 880   |
| Skates           | 348   | 406   | 376   | 598   | 628   | 870   | Skate          | 452   | 673   | 718   | 1,251 |
| Misc Fish        | 207   | 134   | 156   | 236   | 156   | 134   | Misc fish      | 101   | 77    | 154   | 141   |
| Sculpins         | 109   | 188   | 67    | 185   | 199   | 199   | Large Sculpins | 42.6  | 116   | 137   | 148   |
| Sleeper shark    | 105   | 74    | 77    | 104   | 206   | 149   | Shark          | 81.8  | 107   | 84    | 194   |
| Smelts           | 19.5  | 30.2  | 38.7  | 48.7  | 72.5  | 15.3  | Sea star       | 89.4  | 6.77  | 9.22  | 11.05 |
| Grenadiers       | 19.7  | 34.9  | 79.4  | 33.2  | 11.6  | 6.5   | Other Sculpins | 59.2  | 15.5  | 10.8  | 22.67 |
| Salmon shark     | 6.6   | 15.2  | 24.7  | 19.5  | 22.5  | 27.5  | Grenadier      | 20.4  | 9.40  | 8.99  | 15.73 |
| Starfish         | 6.5   | 57.7  | 6.8   | 6.2   | 12.8  | 17.4  | Eulachon       | 2.49  | 18.8  | 8.98  | 87.37 |
| Shark            | 15.6  | 45.4  | 10.3  | 0.1   | 2.3   | 2.3   | Other osmerids | 7.51  | 1.97  | 3.38  | 4.99  |
| Benthic inverts. | 2.5   | 26.3  | 7.4   | 1.7   | 0.6   | 2.1   | Snails         | 1.26  | 0.94  | 6.91  | 0.16  |
| Sponges          | 0.8   | 21.0  | 2.4   | 0.2   | 2.1   | 0.3   | Eelpouts       | 7.03  | 0.61  | 1.33  | 20.60 |
| Octopus          | 1.0   | 4.7   | 0.4   | 0.8   | 4.8   | 8.1   | Giant Grenad.  | 0.31  | 3.50  | 5.02  | 8.80  |
| Crabs            | 1.0   | 8.2   | 0.8   | 0.5   | 1.8   | 1.5   | Octopus        | 1.10  | 2.58  | 1.16  | 1.67  |
| Anemone          | 2.6   | 1.8   | 0.3   | 5.8   | 0.1   | 0.6   | Sea pens/whips | 0.58  | 0.95  | 1.65  | 1.96  |
| Tunicate         | 0.1   | 1.5   | 1.5   | 0.4   | 3.7   | 3.8   | Birds          | 0.13  | 0.11  | 2.42  | 0.39  |
| Unident. inverts | 0.2   | 2.9   | 0.1   | 4.4   | 0.1   | 0.2   | Anemone        | 0.40  | 0.41  | 0.29  | 0.60  |
| Echinoderms      | 0.8   | 2.6   | 0.1   | 0.0   | 0.2   | 0.1   | Misc crabs     | 0.75  | 0.03  | 0.26  | 0.08  |
| Seapen/whip      | 0.1   | 0.2   | 0.5   | 0.9   | 1.5   | 2.1   | Lanternfish    | 0.29  | 0.07  | 0.63  | 9.59  |
| Birds            | 0.2   | 2.1   | 0.7   | 0.2   | 0.3   | 0.3   | Capelin        | 0.01  | 0.32  | 0.35  | 1.51  |
| Lanternfish      | 0.4   | 0.2   | 0.0   | 0.1   | 0.3   | 2.7   | Urochordate    | 0.00  | 0.01  | 0.49  | 0.01  |
| Coral            | 0.0   | 0.2   | 0.0   | 0.1   | 0.0   | 0.0   | Pandal. shrimp | 0.01  | 0.01  | 0.43  | 0.80  |
| Dogfish          | 0.0   | 0.1   | 0.0   | 0.0   | 0.1   | 0.0   | Corals Bryozo. | 0.01  | 0.04  | 0.35  | 0.01  |
| Sandfish         | 0.0   | 0.0   | 0.1   | 0.4   | 0.1   | 0.3   | Brittle star   | 0.26  | 0.01  | 0.02  | 2.67  |
| Sandlance        | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.2   | Invertebrate   | 0.04  | 0.12  | 0.09  | 0.14  |
| Shrimp           | 0.1   | 0.3   | 0.3   | 0.0   | 0.1   | 0.2   | Stichaeidae    | 0.08  | 0.07  | 0.04  | 0.01  |
| Sticheidae       | 0.1   | 0.0   | 0.0   | 0.0   | 0.3   | 0.0   | Sponge         | 0.10  | 0.05  | 0.03  | 0.01  |
|                  |       |       |       |       |       |       | Other          | 0.09  | 0.08  | 0.07  | 0.22  |

Table 1.28 Bycatch estimates (t) of target species caught in the BSAI directed pollock fishery, 1997-2006 based on then NMFS Alaska Regional Office reports from observers.

|                     | 1997  | 1998  | 1999  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  |
|---------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Pacific Cod         | 8,478 | 6,560 | 3,220 | 3,432 | 3,879 | 5,928 | 5,773 | 6,192 | 6,420 | 6,867 |
| Flathead Sole       | 2,353 | 2,118 | 1,885 | 2,510 | 2,199 | 1,844 | 1,629 | 2,019 | 2,095 | 2,637 |
| Rock Sole           | 1,529 | 779   | 1,058 | 2,688 | 1,673 | 1,885 | 1,345 | 2,301 | 1,041 | 1,189 |
| Yellowfin Sole      | 606   | 1,762 | 350   | 1,466 | 594   | 768   | 150   | 671   | 17    | 148   |
| Arrowtooth Flounder | 1,155 | 1,762 | 273   | 979   | 529   | 607   | 550   | 541   | 551   | 951   |
| Pacific Ocean Perch | 512   | 692   | 121   | 22    | 574   | 545   | 691   | 321   | 503   | 423   |
| Atka Mackerel       | 229   | 91    | 165   | 2     | 41    | 221   | 379   | 369   | 211   | 154   |
| Rex Sole            | 151   | 68    | 34    | 10    | 103   | 169   | 199   | 322   | 307   | 1     |
| Greenland Turbot    | 125   | 178   | 30    | 52    | 68    | 70    | 38    | 18    | 30    | 64    |
| Alaska Plaice       | 1     | 14    | 3     | 147   | 14    | 50    | 7     | 7     | 4     | 5     |
| All other           | 93    | 41    | 31    | 77    | 118   | 103   | 144   | 130   | 137   | 133   |



Table 1.29 Bycatch estimates of prohibited species caught in the BSAI directed pollock fishery, 1997-2006 based on then NMFS Alaska Regional Office reports from observers. Herring and halibut units are in t, all others represent thousands of individuals caught.

|                 | 1997  | 1998  | 1999  | 2000  | 2001  | 2002  | 2003   | 2004   | 2005   | 2006  |
|-----------------|-------|-------|-------|-------|-------|-------|--------|--------|--------|-------|
| Herring         | 1,089 | 821   | 785   | 482   | 224   | 105   | 895    | 963    | 442    | 229   |
| Red king crab   | 0.00  | 5.10  | 0.00  | 0.00  | 0.04  | 0.01  | 0.05   | 0.01   | 0.00   | 0.03  |
| Other king crab | 0.16  | 1.83  | 0.00  | 0.10  | 5.14  | 0.08  | 0.01   | 0.01   | 0.01   | 0.00  |
| Bairdi crab     | 6.53  | 35.59 | 1.08  | 0.17  | 0.09  | 0.65  | 0.78   | 1.20   | 0.59   | 0.08  |
| Opilio crab     | 88.59 | 45.62 | 12.78 | 1.81  | 2.18  | 1.67  | 0.76   | 0.74   | 1.93   | 0.00  |
| Chinook salmon  | 43.34 | 49.37 | 10.19 | 3.97  | 30.11 | 32.22 | 46.04  | 53.34  | 65.34  | 84.4  |
| Other salmon    | 61.50 | 62.28 | 44.59 | 56.71 | 52.84 | 77.00 | 190.15 | 436.18 | 690.32 | 295.5 |
| Halibut         | 127   | 144   | 69    | 80    | 164   | 127   | 97     | 92     | 190    | 102   |

Table 1.30 Bycatch rates (kg / t of pollock) of target species caught in the BSAI directed pollock fishery by season and area for 2007 based on then NMFS Alaska Regional Office reports from observers.

| kg/t of pollock     | Winter (A-season) |        |         | Summer/fall (B-season) |       |         | Total |        |
|---------------------|-------------------|--------|---------|------------------------|-------|---------|-------|--------|
|                     | NW                | SE     | A Total | NW                     | SE    | B Total |       |        |
| Alaska Plaice       | 0.002             | 0.001  | 0.001   | 0.001                  | 0.004 | 0.002   | 0.002 |        |
| Atka mackerel       | 0.001             | 0.178  | 0.159   | 0.009                  | 0.105 | 0.030   | 0.088 |        |
| Arrowtooth flounder | 0.237             | 1.923  | 1.743   | 0.624                  | 2.546 | 1.035   | 1.352 |        |
| Flounder            | 0.018             | 0.242  | 0.218   | 0.094                  | 0.628 | 0.208   | 0.213 |        |
| Flathead sole       | 3.634             | 3.150  | 3.202   | 2.066                  | 3.032 | 2.273   | 2.689 |        |
| Greenland turbot    | 0.004             | 0.060  | 0.054   | 0.135                  | 0.011 | 0.109   | 0.084 |        |
| Northern rockfish   | 0.000             | 0.001  | 0.001   | 0.015                  | 0.052 | 0.023   | 0.013 |        |
| Other               | 1.056             | 1.103  | 1.098   | 1.055                  | 1.858 | 1.227   | 1.169 |        |
| Pacific cod         | 4.075             | 3.754  | 3.789   | 3.786                  | 4.922 | 4.028   | 3.921 |        |
| Pacific ocean perch | 1.413             | 0.126  | 0.264   | 0.304                  | 0.667 | 0.382   | 0.329 |        |
| Rougheye rockfish   | 0.001             | 0.006  | 0.006   | 0.000                  | 0.001 | 0.000   | 0.003 |        |
| Rockfish            | 0.000             | 0.168  | 0.150   | 0.001                  | 0.040 | 0.009   | 0.072 |        |
| Rock sole           | 1.174             | 0.638  | 0.695   | 0.027                  | 0.090 | 0.041   | 0.334 |        |
| Sablefish           | 0.000             | 0.022  | 0.019   | 0.000                  | 0.004 | 0.001   | 0.009 |        |
| Squid               | 0.344             | 1.504  | 1.380   | 0.029                  | 1.263 | 0.293   | 0.780 |        |
| Shortraker          | 0.037             | 0.147  | 0.135   | 0.000                  | 0.000 | 0.000   | 0.061 |        |
| Yellowfin sole      | 0.036             | 0.011  | 0.013   | 0.000                  | 0.097 | 0.021   | 0.017 |        |
| Total               |                   | 12.032 | 13.033  | 12.926                 | 8.148 | 15.321  | 9.681 | 11.135 |

Table 1.31. Summary results for EBS pollock. Tonnage units are thousands of t.

| Age          | 1     | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    | 11    | 12    | 13    | 14    | 15    |
|--------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| M            | 0.900 | 0.450 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 | 0.300 |
| Prop. F.     | 0.000 | 0.004 | 0.145 | 0.321 | 0.421 | 0.451 | 0.474 | 0.482 | 0.485 | 0.500 | 0.500 | 0.500 | 0.500 | 0.500 | 0.500 |
| Mature       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Fish. Select | 0.000 | 0.011 | 0.145 | 0.693 | 1.291 | 1.388 | 1.765 | 1.714 | 1.498 | 1.235 | 1.052 | 1.052 | 1.052 | 1.052 | 1.052 |

|  |                                |                |
|--|--------------------------------|----------------|
|  | Tier (2008)                    | 1b             |
|  | Age 3+ 2008 begin-year biomass | 4,357 t        |
|  | 2008 Spawning biomass          | <b>1,380 t</b> |
|  | $B_{msy}$                      | <b>1,876 t</b> |
|  | $B_{40\%}$                     | <b>2,627 t</b> |
|  | $B_{35\%}$                     | 2,299 t        |
|  | $B_{100\%}$                    | 6,569 t        |
|  | $B_0$                          | <b>5,013 t</b> |

| Yield Considerations                 | 2008       | 2009*   |
|--------------------------------------|------------|---------|
| ABC: Harmonic Mean $F_{msy}$         | 1,170 t    | 976 t   |
| ABC: Yield $F_{40\%}$ (Tier 3)       | 555 t      | 650 t   |
| OFL: Arithmetic Mean $F_{msy}$ Yield | 1,443 t    | 1,204 t |
| OFL: Yield $F_{35\%}$ (Tier 3)       | 677 t      | 794 t   |
| <b>Full Selection F's</b>            |            |         |
|                                      | $F_{msy}$  | 0.919   |
|                                      | $F_{40\%}$ | 0.506   |
|                                      | $F_{35\%}$ | 0.650   |

\* Assuming 2008 catches equal 1,170,000 t

# Figures




Figure 1.1. Alaska pollock catch estimates from the Eastern Bering Sea, Aleutian Islands, Bogoslof Island, and Donut Hole regions, 1964-2007. The 2007 value is based on expected totals for the year.




Figure 1.2. Pollock catch distribution in the fishery 2005-2007, January – May on the EBS shelf. Line delineates catcher-vessel operational area (CVOA). The column height represents relative removal on the same scale in all years.




Figure 1.3. Estimate of EBS pollock catch numbers by sex for the “A season” (January-June) and for the entire annual fishery, 1991-2006.




Figure 1.4. Fishery length frequency for the “A season” (January-May) female EBS pollock, 1991-2007. Data for 2007 are preliminary.




Figure 1.5. Pollock catch distribution during June – December, 2005-2007. Line delineates the catcher-vessel operational area (CVOA). The height of the bars represents relative removal on the same scale over all years.




Figure 1.6. Comparison of foreign reported pollock catch (prior to 1986) with recent fishery data during the summer-fall fishery relative to the northwest zone (W of 170°W) of the EBS.



## B-season females




Figure 1.7. Length frequency of EBS pollock observed in period July-December for 1991-2007. Data for 2007 are preliminary.




Figure 1.8. Relative monthly body mass deviation for EBS pollock standardized by the mean long-term weight at length (from 40 to 50cm), 1991-2006 combined data.





Figure 1.9. Relative monthly body mass deviation for EBS pollock standardized by the mean long-term weight at length (from 40 to 50cm), 1991-2006 combined data.




Figure 1.10. EBS pollock fishery estimated catch-at-age data (in number) for 1991-2006. Age 10 represents pollock age 10 and older.




Figure 1.11. Bottom-trawl survey biomass estimates with approximate 95% confidence bounds (based on sampling error) for EBS pollock, 1982-2007. These estimates **include** the northern strata except for 1982-84, and 1986 (indicated by cross symbols).




Figure 1.12. Maps showing the pollock catch-per-unit effort observed from the 2005 - 2007 NMFS EBS shelf bottom-trawl surveys.




Figure 1.13. Pollock abundance levels by age and year as estimated directly from the NMFS bottom-trawl surveys (1982-2007).




Figure 1.14. Pollock abundance levels by length plotted over time as estimated directly from the NMFS bottom-trawl surveys (1987-2007).







Figure 1.15. Mean EBS pollock body-mass-at-age as observed in the summer bottom trawl surveys, 1982-2007 (top panel) and relative to the mean date of the bottom trawl survey operations (bottom panel) for ages 1-3.




Figure 1.16. Mean EBS pollock body-mass-at-age as observed in the summer bottom trawl surveys, 2004-2007.





Figure 1.17. Evaluation of EBS pollock cohort abundances as observed for age 6 and older in the NMFS summer bottom trawl surveys. The bottom panel shows the raw log-abundances at age while the top panel shows the estimates of total mortality by cohort.






Figure 1.19. Echo-integration trawl survey results for 2006 and 2007. The lower figure is the result from the BTS data in the same years. Vertical lines represent biomass of pollock as observed in the different surveys.




Figure 1.20. EBS pollock population at age estimates from the 2006 echo-integration trawl survey as estimated in 2006 based on one or two age-length keys from the bottom trawl surveys and as revised in 2007 using age-length keys derived from the EIT surveys.

## Hydro-acoustic survey abundance-at-age estimates




Figure 1.21. Time series of estimated abundances at age (numbers) for EBS pollock from the EIT surveys, 1979-2007. Note that the 2007 age compositions were computed using an age-length key derived from the 2007 BTS data and as such, are preliminary. The diagonal lines are placed to follow year classes over time.




Figure 1.22. Length and age relationship for EBS pollock from the 2006 fishery data showing the fit and variability (dashed lines represent  $\pm 2$  standard deviations). This relationship was used to develop age-size conversion to fit the available 2007 fishery length frequency data. Points were randomized slightly for visibility purposes.




Figure 1.23. The impact of the 2006 assessment model with revisions given data and other revisions on Tier 1 ABC values for 2008. NOTE: this does not reflect the addition of any data collected in 2007 and is meant for illustration purposes.






Figure 1.24. Evaluation of how new data affect the model for 2007. The CBE model (right-most) includes all of the new information (**C**atch, fishery **A**ge, **B**ottom-trawl survey data, and **E**cho-integration trawl data). Bars represent predicted EBS pollock begin-year age 3+ biomass and the line indicates the uncertainty in estimating the 2005 year class.




Figure 1.25. Model results of predicted EBS pollock survey proportions-at-age under Model CA where only 2006 fishery age and 2007 length composition data were added.




Figure 1.26. Model results for predicted EBS pollock fishery proportions at age under Model C without the 2006 fishery age and 2007 length data included.




Figure 1.27. Profile likelihood over the combined EIT and bottom-trawl survey selectivity (ages 3-15) relative to the point estimate for EBS pollock. The dashed line traces the change in stock size under different values of catchability and the circle highlights the value estimated (and used) in the assessment.




Figure 1.28. Estimates of 2008 EBS pollock population abundance as estimated in this assessment compared to last year's estimates (top panel) and a similar figure showing cumulative population biomass-at-age differences (bottom panel). Dashed lines in bottom panel show calculations using alternative mean weights-at-age values.




Figure 1.29. Selectivity at age estimates for the EBS pollock fishery, 1978-2007 including the estimates used for the future yield considerations.




Figure 1.30. Fit to the EBS pollock fishery age composition estimates (1979-2006) and to the current-year estimate of fishery length frequency data (bottom most panel). Lines represent model predictions while the vertical columns represent the data. Age data new to this year's assessment are shaded.





Figure 1.31. Estimates of bottom-trawl survey numbers (lower panel) and selectivity-at-age (with maximum value equal to 1.0) over time (upper panel) for EBS pollock, 1982-2007.

## Bottom trawl survey age composition fits




Figure 1.32. Fit to the bottom trawl survey age composition data (proportions) for EBS pollock. Lines represent model predictions while the vertical columns represent the data. Data new to this assessment are shaded (2007).




Figure 1.33. Estimates of EIT survey numbers (lower panel) and selectivity-at-age (with mean value equal to 1.0) over time (upper panel) for EBS pollock age 2 and older. Note that the 1979 observed value (=46,314) is off the scale of the figure.






Figure 1.34. Fit to the EIT survey EBS pollock age composition data (proportions) and age 1 index (bottom panel; log-scale). Lines represent model predictions while the vertical columns and dots represent data. The 2007 age composition data are new to the assessment are shaded and the 2006 data were based on revised values using EIT age-length keys.




Figure 1.35. Bivariate and marginal distributions of key parameters integrated over an MCMC chain (length 4 million with every 500<sup>th</sup> sample selected and a burn-in of 4,000).




Figure 1.36. Distribution of stock-recruitment relationships drawn from posterior distribution for EBS pollock.




Figure 1.37. Estimated age 3+ EBS mid-year pollock biomass, 1978-2008. Approximate upper and lower 95% confidence limits are shown by dashed lines. Superimposed is the estimate of mid-year age 3+ biomass from last year's assessment




Figure 1.38. Comparison of the current assessment results with past assessments of **begin-year** EBS age-3+ pollock biomass, 1978-2006.




Figure 1.39. Estimated spawning exploitation rate (defined as the annual percent removals of spawning females due to the fishery) for EBS pollock. Error bars represent two standard deviations from the estimates.




Figure 1.40. Spawning biomass relative to annually computed  $F_{MSY}$  values and fishing mortality rates for EBS pollock, 1977-2007. Note that as the stock drops below the  $B_{20\%}$  level that the directed pollock fishery would be closed.





Figure 1.41. Year-class strengths by year (as age-1 recruits, upper panel) and relative to female spawning biomass (thousands of tons, lower panel) for EBS pollock. Labels on points correspond to year classes. Solid line in upper panel represents the mean age-1 recruitment for all years since 1964 (1963-2006 year classes). Vertical lines in lower panel indicate  $B_{msy}$  and  $B_{40\%}$  level, curve represents fitted stock-recruitment relationship with dashed lines representing approximate lower and upper 95% confidence limits about the curve.




Figure 1.42. Projected EBS pollock **yield** (top) and **Female spawning biomass** (bottom) relative to the long-term expected values under  $F_{35\%}$  and  $F_{40\%}$  (horizontal lines).  $B_{40\%}$  is computed from average recruitment from 1978-2006. Future harvest rates follow the guidelines specified under Scenario 1, max  $F_{ABC}$  assuming  $F_{ABC} = F_{40\%}$ . Note that this projection method is provided only for reference purposes, the SSC has determined that a Tier 1 approach is recommended for this stock.




Figure 1.43. Estimated EBS pollock female spawning biomass trends, 1990-2012, under different 2008-2012 harvest levels. Note that the  $F_{msy}$  catch levels represent unadjusted arithmetic mean fishing mortality rates. Horizontal solid and dashed lines represent the  $B_{msy}$ , and  $B_{40\%}$  levels, respectively.




Figure 1.44. Estimated EBS pollock spawning exploitation rate (defined as the annual percent removals of spawning females due to the fishery). Error bars represent two standard deviations from the estimate and projections for 2008 show the implications of different harvest levels. Note that the  $F_{40\%}$  level represents the adjusted Tier 3b value.






Figure 1.45. Food web pathways for the EBS region based on data from 1990-1994 emphasizing the position of EBS pollock juveniles (a), adults (b) and the pollock fishery (c). Outlined species and fisheries represent predators of pollock (dark box with light text) and prey of pollock (light boxes with dark text). Labels without boxes indicate no direct connection. Box and text size is proportional to each species' standing stock biomass, while the widths are proportional to the consumption between boxes (tons/year).




Figure 1.46. Geographic distribution of 38 kHz acoustic backscatter ( $s_A$  ( $m^2/nmi^2$ )) from species other than pollock (non-pollock, “other” backscatter) observed along tracklines during June-July eastern Bering Sea shelf acoustic-trawl surveys between 1999 and 2007.

## Model details

### Model structure

We used an explicit age-structured model with the standard catch equation as the operational population dynamics model (e.g., Fournier and Archibald 1982, Hilborn and Walters 1992, Schnute and Richards 1995). Catch in numbers at age in year  $t$  ( $C_{t,a}$ ) and total catch biomass ( $Y_t$ ) were

$$C_{t,a} = \frac{F_{t,a}}{Z_{t,a}} (1 - e^{-Z_{t,a}}) N_{t,a}, \quad 1 \leq t \leq T \quad 1 \leq a \leq A$$

$$N_{t+1,a+1} = N_{t,a} e^{-Z_{t,a}} \quad 1 \leq t \leq T \quad 1 \leq a < A$$

$$N_{t+1,A} = N_{t,A-1} e^{-Z_{t,A-1}} + N_{t,A} e^{-Z_{t,A}} \quad 1 \leq t \leq T$$

$$Z_{t,a} = F_{t,a} + M_{t,a}$$

$$C_t = \sum_{a=1}^A C_{t,a}$$

$$p_{t,a} = C_{t,a} / C_t$$

$$Y_t = \sum_{a=1}^A w_a C_{t,a}, \text{ and}$$

where

$T$  is the number of years,

$A$  is the number of age classes in the population,

$N_{t,a}$  is the number of fish age  $a$  in year  $t$ ,

$C_{t,a}$  is the catch of age class  $a$  in year  $t$ ,

$p_{t,a}$  is the proportion of the total catch in year  $t$ , that is in age class  $a$ ,

$C_t$  is the total catch in year  $t$ ,

$w_a$  is the mean body weight (kg) of fish in age class  $a$ ,

$Y_t$  is the total yield biomass in year  $t$ ,

$F_{t,a}$  is the instantaneous fishing mortality for age class  $a$ , in year  $t$ ,

$M_{t,a}$  is the instantaneous natural mortality in year  $t$  for age class  $a$ , and

$Z_{t,a}$  is the instantaneous total mortality for age class  $a$ , in year  $t$ .

We reduced the freedom of the parameters listed above by restricting the variation in the fishing mortality rates ( $F_{t,a}$ ) following Butterworth et al. (2003) by assuming that

$$F_{t,a} = s_{t,a} \mu^f \exp(\varepsilon_t) \quad \varepsilon_t \sim N(0, \sigma_E^2)$$

$$s_{t+1,a} = s_{t,a} \exp(\gamma_{t,a}), \quad \gamma_{t,a} \sim N(0, \sigma_s^2)$$

where

$s_{t,a}$  is the selectivity for age class  $a$  in year  $t$ , and

$\mu^f$  is the median fishing mortality rate over time.

If the selectivities ( $s_{t,a}$ ) are constant over time then fishing mortality rate decomposes into an age component and a year component. This assumption creates what is known as a separable model. If selectivity in fact changes over time, then the separable model can mask important changes in fish abundance. In our analyses, we constrain the variance term ( $\sigma_s^2$ ) to allow selectivity to change slowly

over time—thus improving our ability to estimate the  $\gamma_{t,a}$ . Also, to provide regularity in the age component, we placed a curvature penalty on the selectivity coefficients using the squared second-differences. We selected a simple random walk as our time-series effect on these quantities. Prior assumptions about the relative variance quantities were made. For example, we assume that the variance of transient effects (e.g.,  $\sigma_E^2$ ) is large to fit the catch biomass precisely. Perhaps the largest difference between the model presented here and those used for other groundfish stocks is in how we model “selectivity” of both the fishery and survey gear types. The approach taken here assumes that large differences between a selectivity coefficient in a given year for a given age should not vary too much from adjacent years and ages (unless the data suggest otherwise, e.g., Lauth et al. 2004). The magnitude of these changes is determined by the prior variances as presented above. For the application here selectivity is allowed to change every two years (previously three years were used). In this application, 2006-2007 were configured to have the same selectivity since the geographical patterns were quite similar compared to other years. The “mean” selectivity going forward for projections and ABC deliberations is the simple mean of the estimates from 2005-2007. Unlike previous years, since 2007 now has age specific information (through length frequency) allowing estimates to extend to such a recent year should help better capture how the fishery is evolving in the short term.

Bottom-trawl survey selectivity was set to be asymptotic yet retain the properties desired for the characteristics of this gear. Namely, that the function should allow flexibility in selecting age 1 pollock over time. The functional form of this selectivity is:

$$\begin{aligned} s_{t,a} &= \left[1 + e^{-\alpha_t(a-\beta_t)}\right]^{-1}, \quad a > 1 \\ s_{t,a} &= \mu_s e^{\delta_t^\mu}, \quad a = 1 \\ \alpha_t &= \bar{\alpha} e^{\delta_t^\alpha} \\ \beta_t &= \bar{\beta} e^{\delta_t^\beta} \end{aligned}$$

where the parameters of the selectivity function follow a random walk process as in Dorn et al. (2000):

$$\begin{aligned} \delta_t^\mu - \delta_{t+1}^\mu &\sim N(0, \sigma_{\delta^\mu}^2) \\ \delta_t^\alpha - \delta_{t+1}^\alpha &\sim N(0, \sigma_{\delta^\alpha}^2) \\ \delta_t^\beta - \delta_{t+1}^\beta &\sim N(0, \sigma_{\delta^\beta}^2) \end{aligned}$$

The parameters to be estimated in this part of the model are thus the  $\bar{\alpha}, \bar{\beta}, \delta_t^\mu, \delta_t^\alpha,$  and  $\delta_t^\beta$  for  $t=1982, 1983, \dots, 2007$ . The variance terms for these process-error parameters were specified to be 0.04.

This year a modification was made to the EIT survey selectivity and how these data are treated. As an option, the age one pollock observed in this trawl can be treated as an index and are not considered part of the age composition (which then ranges from age 2-15). This was done to improve some interaction with the flexible selectivity smoother that is used for this gear and was compared.

### Recruitment

In these analyses, recruitment ( $R_t$ ) represents numbers of age-1 individuals modeled as a stochastic function of spawning stock biomass. A further modification made in Ianelli et al. (1998) was to have an environmental component to account for the differential survival attributed to larval drift (e.g., Weststad et al. 2000). ( $\kappa_t$ ):

$$R_t = f(B_{t-1}) e^{\kappa_t + \tau_t}, \quad \tau_t \sim N(0, \sigma_R^2)$$

with mature spawning biomass during year  $t$  was defined as:

$$B_t = \sum_{a=1}^{15} w_a \phi_a N_{at}$$

and  $\phi_a$ , the proportion of mature females at age, was the same as that presented in Wespestad (1995).

A reparameterized form for the stock-recruitment relationship following Francis (1992) was used. For the Beverton-Holt form we have:

$$R_t = f(B_{t-1}) = \frac{B_{t-1} e^{\varepsilon_t}}{\alpha + \beta B_{t-1}}$$

where

- $R_t$  is recruitment at age 1 in year  $t$ ,
- $B_t$  is the biomass of mature spawning females in year  $t$ ,
- $\varepsilon_t$  is the “recruitment anomaly” for year  $t$ ,
- $\alpha, \beta$  are stock-recruitment function parameters.

Values for the stock-recruitment function parameters  $\alpha$  and  $\beta$  are calculated from the values of  $R_0$  (the number of 0-year-olds in the absence of exploitation and recruitment variability) and the “steepness” of the stock-recruit relationship ( $h$ ). The “steepness” is the fraction of  $R_0$  to be expected (in the absence of recruitment variability) when the mature biomass is reduced to 20% of its pristine level (Francis 1992), so that:

$$\alpha = \tilde{B}_0 \frac{1-h}{4h}$$

$$\beta = \frac{5h-1}{4hR_0}$$

where

- $\tilde{B}_0$  is the total egg production (or proxy, e.g., female spawner biomass) in the absence of exploitation (and recruitment variability) expressed as a fraction of  $R_0$ .

Some interpretation and further explanation follows. For steepness equal 0.2, then recruits are a linear function of spawning biomass (implying no surplus production). For steepness equal to 1.0, then recruitment is constant for all levels of spawning stock size. A value of  $h = 0.9$  implies that at 20% of the unfished spawning stock size will result in an expected value of 90% unfished recruitment level. Steepness of 0.7 is a commonly assumed default value for the Beverton-Holt form (e.g., Kimura 1988). The same prior distribution for steepness based on a beta distribution as in Ianelli et al. (2001) and is shown in Fig. 1.47.

To have the critical value for the stock-recruitment function (steepness,  $h$ ) on the same scale for the Ricker model, we begin with the parameterization of Kimura (1990):

$$R_t = f(B_{t-1}) = \frac{B_{t-1} e^{a\left(1-\frac{B_{t-1}}{\varphi_0 R_0}\right)}}{\varphi_0}$$

It can be shown that the Ricker parameter  $a$  maps to steepness as:

$$h = \frac{e^a}{e^a + 4}$$

so that the prior used on  $h$  can be implemented in both the Ricker and Beverton-Holt stock-recruitment forms. Here the term  $\varphi_0$  represents the equilibrium unfished spawning biomass per-recruit.

### Diagnostics

In 2006 a “replay” feature was added where the time series of recruitment estimates from a particular model is used to compute the subsequent abundance expectation had no fishing occurred. These recruitments are adjusted from the original estimates by the ratio of the expected recruitment given spawning biomass (with and without fishing) and the estimated stock-recruitment curve. I.e., the recruitment under no fishing is modified as:

$$R'_t = \hat{R}_t \frac{f(S'_t)}{f(\hat{S}_t)}$$

where  $\hat{R}_t$  is the original recruitment estimate in year  $t$  with  $f(S'_t)$  and  $f(\hat{S}_t)$  representing the stock-recruitment function given spawning biomass under no fishing and under the fishing scenario, respectively.

The assessment model code allows retrospective analyses (e.g., Parma 1993, and Ianelli and Fournier 1998). This was designed to assist in specifying how recruitment patterns (and uncertainty) have changed relative to Tier 1 and Tier 3 ABC calculations. The retrospective approach simply uses the current model to evaluate how it may change over time with the addition of new data based on the evolution of data collected over the past 14 years.

### Parameter estimation

The objective function was simply the product of the negative log-likelihood function and prior distributions. To fit large numbers of parameters in nonlinear models it is useful to be able to estimate certain parameters in different stages. The ability to estimate stages is also important in using robust likelihood functions since it is often undesirable to use robust objective functions when models are far from a solution. Consequently, in the early stages of estimation we use the following log-likelihood function for the survey and fishery catch at age data (in numbers):

$$f = n \cdot \sum_{a,t} p_{at} \ln(\hat{p}_{at}),$$

$$p_{at} = \frac{O_{at}}{\sum_a O_{at}}, \quad \hat{p}_{at} = \frac{\hat{C}_{at}}{\sum_a \hat{C}_{at}}$$

$$\hat{C} = C \cdot E_{ageing}$$

$$E_{ageing} = \begin{pmatrix} b_{1,1} & b_{1,2} & b_{1,3} & \cdots & b_{1,15} \\ b_{2,1} & b_{2,2} & & & \\ b_{3,1} & & \ddots & & \\ \vdots & & & \ddots & \\ b_{15,2} & & & & b_{15,15} \end{pmatrix},$$

where  $A$ , and  $T$ , represent the number of age classes and years, respectively,  $n$  is the sample size, and  $O_{at}$ ,  $\hat{C}_{at}$  represent the observed and predicted numbers at age in the catch. The elements  $b_{ij}$  represent ageing mis-classification proportions are based on independent agreement rates between otolith age readers. For the models presented this year, the option for including aging errors was omitted as has been recommended in past years.

In 2007 the ability to fit to length frequency data was added. This included 25 “bins” for length categories as follows:

|                  |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|------------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| Bin              | 1  | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9  | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 |
| Lower bound (cm) | 25 | 27 | 29 | 31 | 33 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 48 | 50 | 52 | 54 | 56 | 58 | 60 | 62 |

The growth transition matrix (based on 2006 fishery data) was used to estimate the dispersion of pollock lengths given age and is shown in Fig. 1.22. The mean and standard deviation in length given age was fit as a function of age:

|                  |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
|------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Age              | 1     | 2     | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    | 11    | 12    | 13    | 14    | 15+   |
| Mean length (cm) | 19.07 | 27.90 | 34.74 | 40.02 | 44.11 | 47.27 | 49.71 | 51.61 | 53.07 | 54.20 | 55.08 | 55.76 | 56.28 | 56.68 | 57.99 |
| Std. Dev.        | 2.10  | 2.79  | 3.25  | 3.37  | 3.31  | 3.55  | 3.73  | 3.87  | 4.03  | 4.29  | 4.53  | 4.77  | 4.99  | 5.21  | 5.51  |

The length frequency data were fit in an analogous fashion to that of age-data when ageing errors were assumed—by converting the 2007 assessment model predicted catch-at-age into predicted length frequency using the growth and variability as estimated above. For the length frequency data, a multinomial likelihood was used.

Sample size values were revised and are shown in Table 1.16. Strictly speaking, the amount of data collected for this fishery indicates higher values might be warranted. However, the standard multinomial sampling process is not robust to violations of assumptions (Fournier et al. 1990). Consequently, as the model fit approached a solution, we invoke a robust likelihood function which fit proportions at age as:

$$\prod_{a=1}^A \prod_{t=1}^T \frac{\left( \exp \left\{ -\frac{(p_{t,a} - \hat{p}_{t,a})^2}{2(\eta_{t,a} + 0.1/T) \tau^2} \right\} + 0.01 \right)}{\sqrt{2\pi(\eta_{t,a} + 0.1/T) \tau}}$$

Taking the logarithm we obtain the log-likelihood function for the age composition data:

$$\begin{aligned} & -1/2 \sum_{a=1}^A \sum_{t=1}^T \log_e \left( 2\pi(\eta_{t,a} + 0.1/T) \right) - \sum_{a=1}^A T \log_e(\tau) \\ & + \sum_{a=1}^A \sum_{t=1}^T \log_e \left[ \exp \left\{ -\frac{(p_{t,a} - \hat{p}_{t,a})^2}{2(\eta_{t,a} + 0.1/T) \tau^2} \right\} + 0.01 \right] \end{aligned}$$

where  $\eta_{t,a} = \hat{p}_{t,a} (1 - \hat{p}_{t,a})$

and  $\tau^2 = 1/n$

gives the variance for  $p_{t,a}$

$$(\eta_{t,a} + 0.1/T) \tau^2.$$

Completing the estimation in this fashion reduces the model sensitivity to data that would otherwise be considered “outliers.”

Within the model, predicted survey abundance accounted for within-year mortality since surveys occur during the middle of the year. As in previous years, we assumed that removals by the survey were insignificant (i.e., the mortality of pollock caused by the survey was considered insignificant).

Consequently, a set of analogous catchability and selectivity terms were estimated for fitting the survey observations as:

$$\hat{N}_{t,a}^s = e^{-0.5Z_{t,a}} N_{t,a} q_t^s s_{t,a}^s$$

where the superscript  $s$  indexes the type of survey (EIT or BTS). For these analyses we chose to keep survey catchabilities constant over time (though they are estimated separately for the EIT and bottom trawl surveys). The contribution to the negative log-likelihood function from the surveys is given by

$$\sum_{t^s} \left( \frac{\ln(A_{t^s}^s / \hat{N}_{t^s}^s)^2}{2\sigma_{t^s}^2} \right)$$

where  $A_{t^s}^s$  is the total (numerical) abundance estimate with variance  $\sigma_{t^s}^2$  from survey  $s$  in year  $t$ .

The contribution to the negative log-likelihood function for the observed total catches ( $O_t$ ) by the fishery is given by

$$\lambda_c \sum_t \left( \log(O_t / \hat{C}_t)^2 \right)$$

where  $\lambda_c$  represents prior assumptions about the accuracy of the observed catch data. Similarly, the contribution of prior distributions (in negative log-density) to the log-likelihood function include

$$\lambda_\varepsilon \sum_t \varepsilon_t^2 + \lambda_\gamma \sum_{ta} \gamma_{t,a}^2 + \lambda_\delta \sum_t \delta_t^2$$

where the size of the  $\lambda$ 's represent prior assumptions about the variances of these random variables. Most of these parameters are associated with year-to-year and age specific deviations in selectivity coefficients. For a presentation of this type of Bayesian approach to modeling errors-in-variables, the reader is referred to Schnute (1994). To easily estimate such a large number of parameters in such a non-linear model, automatic differentiation software extended from Greiwanck and Corliss (1991) and developed into C++ class libraries was used. This software provided the derivative calculations needed for finding the posterior mode via a quasi-Newton function minimization routine (e.g., Press et al. 1992). The model implementation language (ADModel Builder) gave simple and rapid access to these routines and provided the ability estimate the variance-covariance matrix for all dependent and independent parameters of interest.

The approach we use to solve for  $F_{msy}$  and related quantities (e.g.,  $B_{msy}$ , MSY) within a general integrated model context was shown in Ianelli et al. (2001). In 2007 this was modified to include uncertainty in weight-at-age as an explicit part of the uncertainty for  $F_{msy}$  calculations. This involved estimating a vector of parameters ( $w_i^{future}$ ) on “future” mean weights for each age  $i$ ,  $i = (1, 2, \dots, 15)$ , given actual observed mean and variances in weight-at-age over the period 1991-2006. The model simply computes the values of  $\bar{w}_i, \sigma_{w_i}^2$  based on available data and (if this option is selected) estimates the parameters subject to the natural constraint:

$$w_i^{future} \sim N(\bar{w}_i, \sigma_{w_i}^2).$$

Note that this converges to the mean values over the time series of data (no other likelihood component within the model is affected by “future” mean weights-at-age) while retaining the natural uncertainty that can propagate through estimates of  $F_{msy}$  uncertainty. This latter point is essentially a requirement of the Tier 1 categorization.






Figure 1.47. Cumulative prior probability distribution of steepness based on the beta distribution with  $\alpha$  and  $\beta$  set to values which assume a mean and CV of 0.45 and 0.15, respectively. .

*(This page intentionally left blank)*