

NIH - CMRR 2003

**Task-Oriented Exercise After Stroke:
Mechanisms of
Neuromuscular Plasticity**

Richard Macko, M.D.

**Baltimore VA Medical Center and
University of Maryland School of Medicine**


Stroke: A Major Public Health Issue

780,000 strokes/yr in U.S. Chronic deficits in 75%.

- **Conventional rehabilitation- front loaded during the sub-acute recovery period (<3 months); emphasizes recovery of ADL function & prevention of complications.**
- **Patients become physically inactive, which compounds disability by physical deconditioning and learned non-use.**

No evidence-based recommendations to promote regular exercise in stroke population.

Percent of stroke patients with stationary gait function and leg motor strength recovery during conventional rehabilitation



N = 804

**95% get
no better
after 11
weeks
with
routine
care.**

What may be deficient in conventional rehabilitation models ?

1. Is the exercise intensity enough?

- Typical 51 minute PT session, < 3 minutes at low aerobic intensity (>40% HRR)

McKay Lyons et al. 2002

2. Is repetition adequate?

- 20-30 minutes practice needed to produce short-term cortical motor adaptations in normal adults.

Classen 1998

3. Do we exploit the full time window for plasticity?

- Extends beyond 3-6 months

Liepert 2001

OUTLINE

- I. What is biological basis for task-oriented exercise after stroke?
 - Peripheral Mechanisms - cardiovascular deconditioning and muscle abnormalities contribute to disability.
 - Central Neural Mechanisms – task repetition is critical for promoting motor learning by neural plasticity.
- II. What is the evidence that task-oriented exercise can improve function and fitness in chronic stroke ?

How unfit are chronic stroke patients?



- We tested fitness levels & energy demands of gait in **53 patients**
- **44 men, 9 women**
- **Mean Age 65 ± 8 years**
- **Mean 3 years post-stroke**
- **Hemiparetic gait, not wheelchair bound.**

ECONOMY OF GAIT



- **Purpose - estimate the energy demands of hemiparetic gait.**
- **Protocol - Treadmill walk at 75% of self-selected floor-walk pace, no incline for 9 minutes.**
- **Rate of V_{O_2} calculated from final 3 minutes at steady state oxygen kinetics.**


Chronic stroke patients have diminished fitness reserve

Low Peak Fitness Levels


- $\dot{V}O_2$ Peak = 14.7 ± 4 ml/kg/min;
- 44% below sedentary controls.

Poor Economy of Gait:


- Mean rate $\dot{V}O_2$ 9.7 ± 2 ml/kg/min



Relationship of peak $\dot{V}O_2$ to thigh muscle mass (DXA)




CT Scan of the Mid-Thigh Cross-Sectional Muscle Area in a Stroke Patient



Muscle area is 20% lower in hemiparetic thigh (N=30, P<.001).

Changes in Paretic Leg Skeletal Muscle Phenotype after Stroke

Myosin Heavy Chain Profile



Increased fast MHC isoforms in paretic leg vastus lateralis.

Paretic leg = *67% Unaffected leg = 51% (*p<.001, N=15)

Fast MHC is prone to fatigue, and insulin resistant.

DeDeyne, submitted

Rationale for exercise after stroke:

- **Physical deconditioning threatens the capacity of stroke patients to meet the high energy demands of hemiparetic gait.**
- **Patients with the most muscular wasting (sarcopenia) in their thighs are at greatest risk for functional aerobic impairment.**
- **Changes in muscle may contribute to motor weakness, fatigue, and increased cardiovascular disease risk by promoting insulin resistance.**

Biomechanical basis for treadmill to promote locomotor re-learning



TM Improves Gait Symmetry

- 20% higher stance:swing ratio on P leg; 20% lower on NP leg.
- 30% improved symmetry of insole forces.
- 40% less cycle-cycle variability.

Initial pilot and feasibility studies of treadmill (TM) training



HYPOTHESIS

6 months TM training will improve fitness and function in chronic hemiparetic stroke?

RESULTS

- Reduces energy demands of HP gait 16%
Macko 1997
- Improves peak fitness 10% & work load 39%
Macko 2001
- Improves floor walking velocity by 21%
Silver 2000
- Increases P leg strength 50% & balance.
Smith 1999, Smith 2000

Study Limitations

Non - controlled. No comparison to conventional care.

RANDOMIZED STUDY

Treadmill AEX vs Active Control

Purpose: Determine whether 6 months TM aerobic training (60 -70% HRR) improves fitness, economy of gait, and floor walking in chronic stroke patients, compared to controls receiving a dose-matched form of conventional care.


- **Active Controls:** 45 minutes stretching exercises; 5 minutes warm-up of low intensity treadmill walking (30-40% HRR).

SUBJECTS:


Clinical & Demographic Features

	TM	Control
• Males : Females	18 : 7	14 : 6
• Age (yrs)	65 ± 10	63 ± 8
• Hemisphere (R : L)	12 : 13	9 : 11
• Time since stroke (M)	32 ± 30	42 ± 65
• Assistive Device		
– None	9 (37.5%)	5 (24%)
– Single point cane	10 (37.5%)	11 (57%)
– Quad cane/walker	6 (25%)	4 (19%)
• Floor Walking Speed (range MPH)	1.4 ± 0.7 (0.25 - 2.7)	1.5 ± 0.7 (0.19 - 2.6)

RESULTS: Effects of Treadmill Training on Fitness after Stroke



Effects of Treadmill Training on Walking Performance



*P < .01 Within groups

CONCLUSIONS - Randomized Study

- I. Treadmill & stretching/low intensity walking both improve economy of gait and 30' floor walk velocity.**
- II. Only TM aerobic training improves cardiovascular metabolic fitness & 6 -minute floor walk performance.**


Biological Mechanisms ?

- Studies investigating the hypothesis that TM exercise mediates central neuroplasticity are ongoing**

Transcranial Magnetic Stimulation (TMS) -Motor evoked potentials from paretic quadriceps before and after a single 20 minute treadmill session:

Before


After



Trained Patients – Increased MEP amplitude paretic VL muscle.

BOLD fMRI

During Knee Movement in a Stroke Patient



Green - activation of motor cortex during movement of unaffected knee.

Red - activation of motor cortex during movement of the paretic knee.

A. Luft,, G Smith, L. Forrester, R Macko, A Goldberg, D. Hanley

Future Research:

Robotics assisted exercise rehabilitation

MIT- ankle robotics module

- Impedance controlled
- 6 degrees of freedom
- Programmable
- Evaluate applications in stroke and other neurological conditions

MIT Newman Lab

Igo Krebs; Neville Hogan, Jason Wheeler.



Thanks to Collaborators

University of Maryland and Baltimore VA Physical Therapy

- Gerald V. Smith, Ph.D., P.T. Larry Forrester, Ph.D.
- Jill Whittall, Ph.D. Sandy Waller, M.S., P.T.
- Michelle Harris-Love, P.T. Patrick DeDeyne, Ph.D.

Baltimore VA & University Maryland Gerontology, Neurology and Rehabilitation

- Andrew P. Goldberg, M.D. Kenneth H. Silver, M.D.
- Alice Ryan, Ph.D.

MIT -Newman Laboratory for Biomechanics and Human Rehabilitation

- H. Igo Krebs, Ph.D. Neville Hogan, Ph.D. Jason Wheeler

Abteilung für Neurologie, Universität Tübingen, Germany, Andi Luft, M.

Johns Hopkins University - Brain Injury Outcomes Division - Dan Hanley, M.D.

Experimental Design

Non-controlled study with measures of...

- **Fitness** - $\dot{V}O_2$ peak, economy of gait.
- **Functional Mobility** - Timed floor walks, Get-Up and Go, translational balance perturbation.
- **Strength**- Isokinetic dynamometry.

...at baseline and after exercise training.