

## Rapid Assessment Reference Condition Model

*The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit [www.landfire.gov](http://www.landfire.gov). Please direct questions to [helpdesk@landfire.gov](mailto:helpdesk@landfire.gov).*

### Potential Natural Vegetation Group (PNVG)

**R2SBMT Mountain Big Sagebrush**

#### General Information

**Contributors** (additional contributors may be listed under "Model Evolution and Comments")

**Modelers**

Louis Provencher    lprovencher@tnc.org

**Reviewers**

Stanley G. Kitchen    skitchen@fs.fed.us  
 Michael Zielinski    mike\_zielinski@nv.blm.gov  
 Gary N. Back    gback@srk.com

**Vegetation Type**

Shrubland

**General Model Sources**

- Literature
- Local Data
- Expert Estimate

**Rapid Assessment Model Zones**

- California
- Great Basin
- Great Lakes
- Northeast
- Northern Plains
- N-Cent. Rockies
- Pacific Northwest
- South Central
- Southeast
- S. Appalachians
- Southwest

**Dominant Species\***

ARTR  
 PUTR2  
 SYOR

**LANDFIRE Mapping Zones**

12    17  
 13    18  
 16

**Geographic Range**

Within the Great Basin, this type occurs in mountain ranges in central and northern Nevada, southern Idaho, Utah, and eastern California. Similar vegetation occurs in Wyoming, Colorado, Oregon, and Washington.

**Biophysical Site Description**

Within the Great Basin modeling zone, elevation ranges from 4500 feet in Idaho to 10,500 feet in the White Mountains of California (Winward and Tisdale 1977, Blaisdell et al. 1982, Cronquist et al. 1994, Miller and Eddleman 2000). However, elevations are predominantly between 5000 and 9000 feet. Mean annual precipitation is typically between 14 and 22 inches, but ranges from 10 to 30 inches (Mueggler and Stewart 1980, Tart 1996).

This type mostly occupies moist, productive rolling upland sites. Soils are typically deep and have well developed dark organic surface horizons (Hironaka et al. 1983, Tart 1996). However, at the high ends of its precipitation and elevation ranges mountain big sagebrush occurs on shallow and/or rocky soils.

**Vegetation Description**

This vegetation type is a mosaic of mountain big sagebrush (*Artemisia tridentata* var. *vaseyana* and *A. tridentata* var. *pauciflora* depending on taxonomy used) and herbaceous communities where tree (conifers) encroachment is unlikely (due to high elevation or soils). Codominant shrubs can include antelope bitterbrush, mountain snowberry, and viscid rabbitbrush. Graminoids are very diverse. Dominant graminoids include Idaho fescue, bluebunch wheatgrass, mountain brome, needlegrasses, slender wheatgrass, bluegrasses, or rough fescue. Among the large number of possible forb species, common forbs may include sulphur buckwheat, pussytoes, lupine, phlox, arrowleaf balsamroot, prairie smoke, and sticky geranium. Mueggler and Stewart (1980), Hironaka et al. (1983), Jensen et al. (1988), and Tart (1996) described several mountain big sagebrush habitat types.

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

## Disturbance Description

Mean fire return intervals in and recovery times of mountain big sagebrush are subjects of lively debate in recent years (Welch and Criddle 2003). Mountain big sagebrush communities were historically subject to stand replacing fires with a mean return interval ranging from 10 years at the Ponderosa pine ecotone, 40+ years at the Wyoming big sagebrush ecotone, and up to 80 years in areas with a higher proportion of low sagebrush in the landscape (Crawford et al. 2004, Johnson 2000, Miller et al. 1994, Burkhardt and Tisdale 1969 and 1976, Houston 1973, Miller and Rose 1995, Miller et al. 2000). Under pre-settlement conditions mosaic burns generally exceeded 75% topkill due to the relatively continuous herbaceous layer. Brown (1982) reported that fire ignition and spread in big sagebrush is largely (90%) a function of herbaceous cover. Mountain big sagebrush communities are also subject to periodic mortality due to insects, diseases, winter kill, rodent outbreaks, and drought (Anderson and Inouye 2001). These disturbances in combination may have significantly reduced the cover of dense stands about every 50 to 100 years.

Recovery rates for shrub canopy cover vary widely in this type, depending post fire weather conditions, abundance of resprouting shrubs, and size and severity of the burn. Mountain big sagebrush typically reaches 5% canopy cover in 8 to 14 years (mean of 12 years). This may take as little as 4 years under favorable conditions and longer than 25 years in unfavorable situations (Pedersen et al. 2003, Miller unpublished data). Mountain big sagebrush typically reaches 25% canopy cover in about 25 years, but an average recovery time of 40 years was used because recovery may take as few as 9 years or as long as 70 years (Winward 1991, Pedersen et al. 2003, Miller unpublished data). Variation in recovery rates is dependent upon burn size and uniformity, survival of residual seed, and upon environmental factors (especially weather). Mountain snowberry and resprouting forms of bitterbrush may return to pre-burn cover values in a few years. Bitterbrush plants less than fifty years old are more likely to resprout than older plants (Simon 1990).

## Adjacency or Identification Concerns

This type may be adjacent to forests dominated by aspen, Ponderosa pine, Douglas-fir or lodgepole pine. It also occurs adjacent to pinyon-juniper woodlands. This type probably served as an ignition source for adjacent aspen stands.

At the lower elevation, dry end of the type, mountain big sagebrush could be confused with Wyoming big sagebrush. At the higher elevation, moist end of the type, mountain big sagebrush, while generally recognized as *A. tridentata* var. *vaseyana*, could be confused with *A. tridentata* var. *pauciflora* and spiked big sagebrush (*Artemisia tridentata* ssp. *Spiciformis*) or mountain shrub communities characterized by *Amelanchier*, *Prunus*, and or *Rosa*.

Uncharacteristic conditions in this type include herbaceous canopy cover less than 40% and dominance of the herbaceous layer by mulesears (*Wyethia amplexcaulis*) on clayey soils.

This PNVG is similar to the PNVG R0SBMT for the Northern and Central Rockies model zone.

## Scale Description

Sources of Scale Data  Literature  Local Data  Expert Estimate

This type occupies areas ranging in size from 10's to 10,000's of acres. Disturbance patch size can also range from 10's to 1,000's of acres. The distribution of past burns was assumed to consist of many small patches in the landscape.

## Issues/Problems

Reviewers and modelers had very different opinions on the range of mean FRIs and mountain big sagebrush recovery times (see Welch and Criddle 2003). It is increasingly agreed upon that a MFI of 20 years, which used to be the accepted norm, is simply too frequent to sustain populations of Greater Sage Grouse and mountain big sagebrush ecosystems whose recovery time varies from 10-70 years. Reviewers consistently suggested longer FRIs and recovery times. The revised model is a compromise with longer recovery times

and FRIs. Modeler and reviewers also disagreed on the choice of FRG: II (modeler) vs. IV (reviewers). In future efforts, this PNVG should be restricted to high elevations where conifer encroachment is unlikely. The PNVG with conifer encroachment is R2SBMTwc.

**Model Evolution and Comments**

The three development classes chosen for this PNVG correspond to the early, mid-, and late seral stages familiar to range ecologists. The depleted or decadent sagebrush condition was considered uncharacteristic of the pre-settlement condition, thus not included.

Resprouting bitterbrush in mountain big sagebrush types is potentially important to wildlife in early stand development.

**Succession Classes\*\***  
*Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).*

**Class A 20 %**

Early1 PostRep

**Description**

Herbaceous cover is variable but typically >50% (50-80%). Shrub cover is 0 to 5%. Replacement fire is uncommon during early recovery (average FRI of 80 yrs) and maintains vegetation in A by causing an ecological setback of 12 yrs. Succession to class B after 12 years.

**Dominant Species\* and Canopy Position**

ARTRV  
 PUTR2  
 SYOR2

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

|                 | Min     | Max     |
|-----------------|---------|---------|
| Cover           | 0 %     | 5 %     |
| Height          | no data | no data |
| Tree Size Class | no data |         |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class B 45 %**

Mid1 Open

**Description**

Shrub cover 6-25%. Mountain big sagebrush cover up to 20%. Herbaceous cover is typically >50%. Replacement fire occurs with a mean FRI of 40 yrs. Succession to class C after 28 years.

**Dominant Species\* and Canopy Position**

ARTRV  
 PUTR2  
 SYMPH

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

|                 | Min     | Max     |
|-----------------|---------|---------|
| Cover           | 6 %     | 25 %    |
| Height          | no data | no data |
| Tree Size Class | no data |         |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Class C 35%**

Late I Closed

**Description**

Shrubs are the dominant lifeform. Shrub cover 26-45+%. Herbaceous cover is typically <50%. Insects and disease every 75 yrs on average will thin the stand and cause a transition to class B. Replacement fire is every 50 yrs on average. Succession keeps vegetation in class C.

**Dominant Species\* and Canopy Position**

ARTRV  
PUTR2  
SYMPH

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

|                 | Min     | Max     |
|-----------------|---------|---------|
| Cover           | 26 %    | 45 %    |
| Height          | no data | no data |
| Tree Size Class | no data |         |

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class D 0%**

**Description**

**Dominant Species\* and Canopy Position**

**Structure Data (for upper layer lifeform)**

|                 | Min     | Max     |
|-----------------|---------|---------|
| Cover           | %       | %       |
| Height          | no data | no data |
| Tree Size Class | no data |         |

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class E 0%**

**Description**

**Dominant Species\* and Canopy Position**

**Structure Data (for upper layer lifeform)**

|                 | Min     | Max     |
|-----------------|---------|---------|
| Cover           | %       | %       |
| Height          | no data | no data |
| Tree Size Class | no data |         |

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Disturbances**

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Disturbances Modeled**

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

**Historical Fire Size (acres)**

Avg: no data  
 Min: no data  
 Max: no data

**Sources of Fire Regime Data**

- Literature
- Local Data
- Expert Estimate

**Fire Regime Group: 4**

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

**Fire Intervals (FI)**

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

|                    | <i>Avg FI</i> | <i>Min FI</i> | <i>Max FI</i> | <i>Probability</i> | <i>Percent of All Fires</i> |
|--------------------|---------------|---------------|---------------|--------------------|-----------------------------|
| <i>Replacement</i> | 48            | 15            | 100           | 0.02083            | 100                         |
| <i>Mixed</i>       |               |               |               |                    |                             |
| <i>Surface</i>     |               |               |               |                    |                             |
| <i>All Fires</i>   | 48            |               |               | 0.02085            |                             |

**References**

Anderson, J.E. and R.S. Inouye 2001. Landscape-scale changes in plant species abundance and biodiversity of a sagebrush steppe over 45 years. *Ecological Monographs* 71:531-556.

Blaisdell, J.P., R.B. Murry, and E.D. McArthur. 1982. Managing Intermountain rangelands-- sagebrush-grass ranges. Gen. Tech. Rep. INT-134. Ogden, UT: U.S. Dept. of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 41p.

Brown J.K. 1982. Fuel and fire behavior prediction in big sagebrush. Res. Pap. INT-197. Ogden, UT: U.S. Dept. of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 10p.

Brown, J. K. and J. K. Smith, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Burkhardt, W.J. and E.W. Tisdale. 1969. Nature and successional status of western juniper vegetation in Idaho. *Journal of Range Management* 22(4):264-270.

Burkhardt, W.J. and E.W. Tisdale. 1976. Causes of juniper invasion in southwestern Idaho. *Ecology* 57:472-484.

Crawford, J.A., R.A. Olson, N.E. West, J.C. Mosley, M.A. Schroeder, T.D. Whitson, R.F. Miller, M.A. Gregg, and C.S. Boyd. 2004. Ecology and management of sage-grouse and sage-grouse habitat. *J. Range Mgmt.* 57:2-19.

Cronquist, A., A.H. Holmgren, N.H. Holmgren, and others. 1994. Intermountain Flora: Vascular Plants of the Intermountain West, U.S.A. Vol. 5. Asterales. New York: The New York Botanical Garden. 496p.

Hironaka, M., M.A. Fosberg, and A.H. Winward. 1983. Sagebrush-Grass Habitat Types of Southern Idaho. University of Idaho Forest, Wildlife and Range Experiment Station, Bulletin Number 35. Moscow, ID. 44p.

Houston, D.B. 1973. Wildfires in northern Yellowstone National Park. *Ecology* 54:1111-1117.

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Jensen, M.E., L.S. Peck, and M.V. Wilson. 1988. A sagebrush community type classification for mountainous northeastern Nevada rangelands. *The Great Basin Naturalist* 48:422-433.

Johnson, Kathleen. 2000. *Artemisia tridentata* ssp. *Vaseyana*. In: Fire Effects Information System [Online], U.S. Dept. of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2004, September 17].

Miller, R.F. And L.L. Eddleman. 2000. Spatial and temporal changes of sage grouse habitat in the sagebrush biome. Technical Bulletin 151. Corvallis, OR: Oregon State University Agricultural Experiment Station. 35p.

Miller, R.F. and J.A. Rose. 1995. Historic expansion of *Juniperus occidentalis* (western juniper) in southeastern Oregon. *The Great Basin Naturalist* 55(1):37-45.

Miller, R.F., T.J. Svejcar, and J.A. Rose. 2000. Impacts of western juniper on plant community composition and structure. *Journal of Range Management* 53(6):574-585.

Miller, R.F.; Svejcar, T.J.; West, N.E. 1994. Implications of livestock grazing in the Intermountain sagebrush region: Plant composition. In: Vavra, M, Laycock, W.A., Pieper, R.D., eds. Ecological implications of livestock herbivory in the West. Denver, CO: Society for Range Management: 101-146.

Mueggler, W.F. and W.L. Stewart. 1980. Grassland and shrubland habitat types of Western Montana. USDA Forest Service GTR INT-66. 154p.

Pedersen, E.K., J.W. Connelly, J.R. Hendrickson, and W.E. Grant. 2003. Effect of sheep grazing and fire on sage grouse populations in southeastern Idaho. *Ecological Modeling* 165:23-47.

Simon, S.A. 1990. Fire effects from prescribed underburning in central Oregon ponderosa pine plant communities: first and second growing season after burning. Pp. 93-109. In *Fire in Pacific Northwest Ecosystems*. Thomas E. Bedell, editor. Department of Rangeland Resources, Oregon State University, Corvallis, OR. 145p.

Tart, D.L. 1996. Big sagebrush plant associations of the Pinedale Ranger district. Pinedale, WY: USDA Forest Service Bridger-Teton National Forest. Jackson, WY. 97 p.

Welch, B. L, C. Criddle. 2003. Countering Misinformation Concerning Big Sagebrush. Research Paper RMRS-RP-40. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 28 p.

Winward, A.H. 1991. A renewed commitment to management in sagebrush grasslands. In: *Management in the Sagebrush Steppes*. Oregon State University Agricultural Experiment Station Special Report 880. Corvallis OR. pp.2-7.

Winward, A.H. and E.W. Tisdale. 1977. Taxonomy of the *Artemisia tridentata* complex in Idaho. Bulletin No. 19. Moscow, ID: University of Idaho, College of Forestry, Wildlife and Range. 15p.

Wyoming Interagency Vegetation Committee. 2002. Wyoming Guidelines for Managing Sagebrush Communities with Emphasis on Fire Management. Wyoming Game and Fish Department and Wyoming BLM. Cheyene, WY 53

pp.