

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R5OASA

Oak Savanna

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

McRee Anderson wanderson@tnc.org

Reviewers

Paul Nelson pwnelson@fs.fed.us

In workshop review

Doug Zollner dzollner@tnc.org

Vegetation Type

Grassland

General Model Sources

- Literature
- Local Data
- Expert Estimate

Rapid Assessment Model Zones

- California
- Great Basin
- Great Lakes
- Northeast
- Northern Plains
- N-Cent.Rockies
- Pacific Northwest
- South Central
- Southeast
- S. Appalachians
- Southwest

Dominant Species*

QUST ANGE
QUMA SCHIZ
QUAL
ANDR

LANDFIRE Mapping Zones

44
43

Geographic Range

Nuzzo(1986) estimated that some 27 to 32 million acres of oak savanna occurred in the Midwest at the time of European settlement extending from southern Texas northward through Missouri into Wisconsin, Minnesota, California and Oregon. Nelson (1987) indicated that perhaps 13 million acres of savanna occurred in Missouri prior to settlement. This number was extrapolated based on interpretations using the extent of prairie cover and descriptions of historic barrens, oak openings and other open woodlands in which grasses dominated the ground cover. The current estimate of six and one half million acres is a relative interpretation excluding presettlement prairie and other natural communities associated with rougher dissected hills. This revised estimate now discounts open woodlands that fall into the woodland natural community descriptions. The estimate is now restricted to the probability of savannas associated with prairie regions and relatively level upland plains.

Biophysical Site Description

Savannas are grasslands interspersed with open-grown scattered trees, groupings of trees of various age, and shrubs. These take on the appearance of widely spaced, orchard-like groves or standing individual trees. They are distinguished from woodlands in that savannas are strongly associated with large prairies on nearly level to dissected plains and are generally dominated by prairie grasses and forbs. The tree canopy cover is generally less than 30 percent. Shrub thickets occur, especially on the northeast-trending lee side of hills or in upland drainages where fire was less frequent or less intense. Savannas are species-rich natural communities, with most diversity found in the understory layer. While no endemic species are presently known to occur in savannas, Packard and Mutel (1997) indicated that oak savanna possesses a distinct herbaceous community characterized by species adapted to frequent large-scale disturbances. Oak Savanna topography is associated with gently rolling plains underlain by Pennsylvanian limestone and sandstone in the unglaciated Osage Plains and the Central Dissected Till Plains sections. However, savannas may occur anywhere upland topography is gently rolling to level, regardless of geologic substrate. Their strongest affinity is to gently rolling plains where prairie occur.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Vegetation Description

In general, three primary vegetation associations dominated savanna natural communities. In the Central Dissected Till Plains Section, bur oak groves (*Quercus Macrocarpa*) once dominated dry to dry-mesic prairie areas underlain by glacial till soils. Chinquapin oak (*Q. muhlenbergii*) co-dominated on the driest, steepest loess hills of Springfield Plain and Central Plateau subsections, especially along the Interstate 44 and Highway 63 (Rolla to Thayer) corridors. In the Springfield Plain Subsection, chinquapin oak and post oak often share dominance where associated with limestone/dolomite bedrock. Rock outcrops on prairies or on rugged, hilly terrain dominated by shrubs such as wild crab (*Malus ioensis*), hawthorn (*Crataegus* species), rough-leaved dogwood (*Cornus drummondii*) and winged sumac (*Rhus copallina*) are often savanna-like in character, but are primarily considered part of the prairie natural community. Moisture modifiers are limited to the primary moisture regime associated with loess and glacial till soils only. Nearly all rock substrate savannas are dry-mesic with inclusions of dry soils, while those found on the deeper soils of glacial till or loess are both mesic and dry-mesic. However, because so little is known about the historic distribution of savanna types developed on bedrock and residuum soils, and distinctions between dry and dry-mesic soils, savannas are named for the primary bedrock substrate only. No wet-mesic or wet savannas are known because either few extant examples remain, or these are too small to function as savannas. Sand savannas are named for the wind or alluvial-deposited sandy soils of terraces or elevated ridges and summits. They are especially characteristic of the Mississippi River Alluvial Basin Section. The typical sand savanna has no moisture modifier because of the difficulty in distinguishing between their dry to dry-mesic soils, and owing to the topographic irregularities of the landscape. Dominant vegetation is listed as big bluestem (*Andropogon gerardii*), little bluestem (*Schizachyrium scoparium*), switchgrass (*Panicum virgatum*), and Indiangrass (*Sorghastrum nutans*).

Six savanna natural communities are described based on differences in soil moisture and rock/parent material substrate:

- Dry-mesic loess/glacial till savanna
- Mesic loess/glacial till savanna
- Limestone/dolomite savanna
- Chert savanna
- Sandstone/shale savanna
- Sand savanna

Disturbance Description

Many oak species are adapted to the frequent, low to moderate intensity fires with the capability of resprouting. Curtis (1959) described brush prairie remnants at Wisconsin savanna sites that burned annually more than 100 years with no observed reduction in the number of oak grubs. Grubs refer to oak (and other species) sprouts killed back by repeated fires and forming large root balls. The presence of these oak grubs account for the rapid degrading of savanna to landscapes overgrown in woody thickets following heavy grazing and the cessation of fire (Schroeder 1981). Savannas, prairies, glades and open woodlands -- all are direct reflections and inextricably linked to natural or aboriginal fires and are relicts of once common grazing sites of American bison (*Bison bison*). In addition, browsing by American elk (*Cervus elaphus*) and white-tailed deer (*Odocoileus virginianus*) influenced the vegetation. Large expanses of level to nearly level landscape coupled with frequent fire and grazing by native herbivores will eventually lead to either prairie or savanna. Though grazing was a natural disturbance questions remain as to the scale that would of altered vegetation changes. Wind and ice storms may have also played a role opening closed communities.

Adjacency or Identification Concerns

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Landscape is adequate in size to contain natural variation in vegetation and disturbance regime. Topographically complex areas can be relatively small (< 1000 acres). Larger landscapes can be up to several thousand acres in size.

Issues/Problems

The causative factors that eventually led to mass degradation, and in some regions total extirpation, of Missouri savannas include suppression of historic natural or anthropogenic fires, replacement of natural herbivory by domestic livestock grazing, logging, conversion to cropland and seeding to cool-season exotic grasses. Because most former savannas (like their associated prairie natural communities) were highly productive in terms of forage (or palatable vegetation), these served as the primary foraging sites for domestic livestock that were allowed to range freely during early settlement. The richest savanna soils, especially in northern Missouri, were rapidly converted to cropland or intensively grazed.

Model Evolution and Comments

Doug Zollner, Tom Foti

Succession Classes**
Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 7%

Early1 All Struct

Description

The early seral open stage is recently burned with a herbaceous species response. Most of the shrubs and oak grubs are top killed by recent replacement fire. However, herbaceous species and oak grubs will resprout and not all are killed by fire.

Dominant Species* and Canopy Position

SCHIZ4 Lower
 ANGE Lower
 ANDR Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 2

Structure Data (for upper layer lifeform)

	Min	Max
Cover	5 %	100 %
Height	no data	Herb Short <0.5m
Tree Size Class	Seedling <4.5ft	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 3%

Mid1 Closed

Description

The mid seral closed stage consists of areas that have not had a recent surface fire. As a result of altered fire regimes, oak grubs have resprouted into medium sized shrubs and pole sized trees. Herbaceous species are present in the ground cover but are limited throughout this stage due to the reduced amount of light reaching the surface.

Dominant Species* and Canopy Position

ANDR Lower
 ANGE Lower
 QUERC Middle
 QUMA Middle

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 9

Structure Data (for upper layer lifeform)

	Min	Max
Cover	30 %	50 %
Height	Shrub Dwarf <0.5m	Shrub Tall >3.0 m
Tree Size Class	Pole 5-9" DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 40%

Mid1 Open
Description

The mid seral open stage consists of areas of the landscape that has recently burned. Due to periodic surface fires some of the oak grub sprouts and shrubs have been top killed resulting in more light reaching the surface propagating the spread of a variety of herbaceous species. Overstory is an intermix of shrubs and pole sized oaks that have not been recently top killed.

Dominant Species* and Canopy Position

QUMA Middle
QUERC Middle
ANGE Lower
ANDR Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 3

Structure Data (for upper layer lifeform)

	Min	Max
Cover	5 %	30 %
Height	Tree Short 5-9m	Tree Medium 10-24m
Tree Size Class	Pole 5-9" DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 45%

Late1 Open
Description

The late open seral stage represents the oak savanna community type. Due to a 3-year surface fire interval the oak grub sprouts and shrubs have been top killed. Tall mature oaks with spreading branches are scattered in a park-like setting with an open canopy allowing light to reach the surface propagating the spread of a variety of herbaceous species.

Dominant Species* and Canopy Position

QUST Upper
QUERC Upper
ANGE Lower
ANDR Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 3

Structure Data (for upper layer lifeform)

	Min	Max
Cover	5 %	30 %
Height	Tree Short 5-9m	Tree Medium 10-24m
Tree Size Class	Very Large >33"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 5%

Late1 Closed
Description

The late closed seral stage represents the oak savanna that has not had recent surface or mixed fire. Tall mature oaks with spreading branches are scattered throughout this type however, oak grubs and shrubs have sprouted into pole size limiting light reaching the surface and therefore reducing the herbaceous species cover. This stage represents a places on the landscape that have an altered fire regime.

Dominant Species* and Canopy Position

QUST Upper
QUERC Upper
QUMA Upper
ANDR Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 9

Structure Data (for upper layer lifeform)

	Min	Max
Cover	30 %	50 %
Height	Tree Short 5-9m	Tree Medium 10-24m
Tree Size Class	Very Large >33"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances

Disturbances Modeled

- Fire
 Insects/Disease
 Wind/Weather/Stress
 Native Grazing
 Competition
 Other:
 Other

Historical Fire Size (acres)

Avg: 10000
 Min: 2000
 Max: 20000

Sources of Fire Regime Data

- Literature
 Local Data
 Expert Estimate

Fire Regime Group: 1

I: 0-35 year frequency, low and mixed severity
 II: 0-35 year frequency, replacement severity
 III: 35-200 year frequency, low and mixed severity
 IV: 35-200 year frequency, replacement severity
 V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
<i>Replacement</i>	100	5	110	0.01	3
<i>Mixed</i>	60	5	250	0.01667	5
<i>Surface</i>	3	1	4	0.33333	93
<i>All Fires</i>	3			0.36	

References

- Adler, P.B., D.A Raff and W.K. Lauenroth. 2001. The effect of grazing on the spatial heterogeneity of vegetation. *Oecologia* 128:465-479.
- Anderson, R.C. 1990. The historic role of fire in North American grasslands. Pp. 8-18 in S.L. Collins and L.L. Wallace (eds.). *Fire in North American tallgrass prairies*. University of Oklahoma Press, Norman.
- Batek, M.J., A.J. Rebertus, W.A. Schroeder, T.L. Haithcoat, E. Compas, and R.P. Guyette. 1999. Reconstruction of early nineteenth-century vegetation and fire regimes in the Missouri Ozarks. *Journal of Biogeography* 26:397-412.
- Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. *Ecological systems of the United States: a working classification of U.S. terrestrial systems*. NatureServe, Arlington, Virginia.
- Fuhlendorf, S.D. and D.M. Engle 2001. Restoring heterogeneity on rangelands; ecosystem management based on evolutionary grazing patterns. *BioScience*, August 2001/Vol. 51 No. 8
- Grazulis, T.P. 2001. *Tornado: nature's ultimate windstorm*. The University of Oklahoma Press, Norman.
- Guyette, R.P. and B.E. Cutter. 1991. Tree-ring analysis of fire history of a post oak savanna in the Missouri Ozarks. *Natural Areas Journal* 11: 93-99.
- Guyette, R.P. and D.C. Dey. 2000. Humans, topography, and wildland fire: the ingredients for long-term patterns in ecosystems. Pp. 28-35 in D.A. Yaussy (ed.). *Proceedings of the workshop on fire, people, and the central hardwoods landscape*. General Technical Report NE-274. U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. Radnor, Pennsylvania.
- Guyette, R.P. and E.A. McGinnes, Jr. 1982. Fire history of an Ozark glade in Missouri. *Transactions of the Missouri Academy of Science* 16:85-93.

- Harlan, J.D., T.A. Nigh and W.A. Schroeder. 2001. The Missouri original General Land Office survey notes project. University of Missouri, Columbia. In progress.
- Kimmel, V.L. and G.E. Probasco. 1980. Change in woody cover on limestone glades between 1938 and 1975. *Transactions of the Missouri Academy of Science* 14:69-74.
- Kline, V. 1997. Orchards of oak and a sea of grass. Pp. 3-22 in S. Packard and C. F. Mutel (eds.). 1997. *The tallgrass restoration handbook for prairies, savannas, and woodlands*. Society for Ecological Restoration. Island Press, Washington, D.C.
- Kuchler, A. W. 1964. Manual to accompany the map of potential natural vegetation of the conterminous United States. American Geographical Society. Spec. Publ. No. 36. Lib. Congress Cat. Card Num. 64-15417. 156 p.
- Ladd, D. 1991. Reexamination of the role of fire in Missouri oak woodlands. Pp. 67-80 in G.V. Burger, J.E. Ebinger and G.S. Wilhelm (eds.). *Proceedings of the oak woods management workshop*. Eastern Illinois University, Charleston, Illinois.
- Ladd, D. and B. Heumann. 1994. Baseline ecological assessment of selected oak woodlands on the Houston-Rolla District, Mark Twain National Forest. U.S. Forest Service challenge cost share agreement.
- McCarty, K. 1998. Landscape-scale restoration in Missouri savannas and woodlands. *Restoration and Management Notes* 16:22-32.
- McCarty, K. and F. Hassien. 1984. Distribution patterns of prairie plant species in a closed-canopy forest situation. Pp. 127-130 in G. Clambey and R. Pemble (eds.). *Proceedings of the ninth North American Prairie Conference. The prairie: past, present and future*. Tri-college University Center for Environmental Studies, North Dakota State University, Fargo.
- Nelson, Paul W. 2005 *The Terrestrial Natural Communities of Missouri*. Missouri Natural Areas Committee (in press).
- Nelson, P.W. and D. Ladd 1983. Preliminary report on the identification, distribution, and classification of Missouri glades. Pp. 59-76 in C.L. Kucera (ed.). *Proceedings of the seventh North American Prairie Conference*. Southwest Missouri State University, Springfield, Missouri.
- Nigh, T.A. 1992. The forests prior to European settlement. Pp. 6-13 in A.R.P. Journet and H.G. Spratt, Jr. (eds.). *Towards a vision for Missouri public forests: proceedings of a conference at Southeast Missouri State University, Cape Girardeau, Missouri*.
- Ozarks Ecoregional Assessment Team. 2003. *Ozarks Ecoregional Conservation Assessment*. The Nature Conservancy, Midwestern Resource Office. Minneapolis, Minnesota.
- Packard, S. and C.F. Mutel (eds.). 1997. *The tallgrass restoration handbook for prairies, savannas, and woodlands*. Society for Ecological Restoration. Island Press, Washington, D.C.
- Rebertus, A.J., S.R. Shifley, R.H. Richards and L.M. Roovers. 1997. Ice storm damage to an old-growth oak-hickory forest in Missouri. *American Midland Naturalist* 137:48-61.
- Rebertus, A.J. and A.J. Meier. 2001. Blowdown dynamics in oak-hickory forests of the Missouri Ozarks.

Journal of the Torrey Botanical Society 128(4):362-369.

Runkle, J.R. 1985. Disturbance regimes in temperate forests. Pp. 17-33 in S.T.A. Pickett and P.S. White (eds.). The ecology of natural disturbance and patch dynamics. Academic Press, New York.

Schoolcraft, H.R. 1821. Journal of a tour into the interior of Missouri and Arkansas from Potosi, or Mine a Burton, in Missouri territory, in a southwest direction, toward the Rocky Mountains: performed in the years 1818 and 1819. Richard Phillips and Company, London.

Taft, J. 1997. Savanna and open woodland communities. Pp. 24-54 in M. Schwartz (ed.). Conservation in highly fragmented landscapes. Chapman and Hall, New York.

U.S. Department of Agriculture, Forest Service. 1999. Ozark-Ouachita highlands assessment: terrestrial vegetation and wildlife. Report 5 of 5. General Technical Report SRS-35. U.S. Department of Agriculture, Forest Service, Southern Research Station. Asheville, North Carolina.