

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R9PCSA Pond Cypress Savanna

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Carl Nordman

carl_nordman@natureserve.org

Reviewers

Vegetation Type

Grassland

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- California
 Great Basin
 Great Lakes
 Northeast
 Northern Plains
 N-Cent. Rockies
- Pacific Northwest
 South Central
 Southeast
 S. Appalachians
 Southwest

Dominant Species*

TAAS
RHMI7
MUFI3
CLMA

LANDFIRE Mapping Zones

56

Geographic Range

Pond cypress savannah occurs in southwest Florida, mainly in the Big Cypress area, Big Cypress National Preserve.

Biophysical Site Description

Pond cypress savannah occurs as often stunted stands of *Taxodium ascendens* growing on shallow sands or marl soils above limestone bedrock (Floherschütz 1978).

Vegetation Description

This PNVG is a wet grassland savannah with scattered pond cypress (*Taxodium ascendens*). The understory is dominated by graminoids including beak rush (*Rhynchospora microcarpa*), sedges (*Cyperus* spp.), muhly grass (*Muhlenbergia filipes*), and sawgrass (*Cladium jamaicense*) (NatureServe, 2005). Vegetation density and diversity are low (Ewel, 1990).

Disturbance Description

The herbaceous graminoid and pond cypress canopy is kept sparse by very low nutrient availability and extreme water level fluctuations. Fires are associated with drought in winter, these droughts occur in association with the El Nino/la nina ENSO, on 7 or 15 year cycles. (David Brownlie, pers com., 2005). In the absence of fire for long periods, hardwood encroachment can occur.

Adjacency or Identification Concerns

This is similar to SW Florida wet prairie, which does not have pond cypress trees. This PNVG is equivalent to CES411.290 South Florida Dwarf Cypress Savanna (NatureServe, 2005).

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Pond cypress savanna occurs primarily in the Big Cypress region of south Florida. Information describing the size of this system was difficult to find. Pond cypress savanna occurs within a matrix of cypress strands, cypress domes, prairie, and pine communities, most contained within Big Cypress National Preserve. Muss

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

et. al. indicated there is approximately 295,100 ha. of cypress within Big Cypress National Preserve, and half of that is open stands of small cypress growing in seasonally flooded grasslands known as cypress prairie.

No information on the scale of disturbances within pond cypress savanna was identified

Issues/Problems

This 3 box model can be brought up to date with the addition of two more boxes to accommodate Melaleuca quinquenervia forests, closed and open.

Model Evolution and Comments

Suggested reviewers Cecil Frost, and somebody from Big Cypress NP.

There was one anonymous reviewer of this model. The reviewer suggested adding a replacement fire disturbance to Class A. No other changes or additions were suggested.

Succession Classes**
Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 15 %

Early1 All Struct

Description

Class A is a pond cypress savanna where medium to high intensity fire in combination with winter ENSO (El Nino/la nina) related drought has killed pond cypress trees.

Dominant Species* and Canopy Position

RHMI7 Lower
 MUFI3 Lower
 CLMAJ Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 2

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	Herb Short <0.5m	Herb Tall > 1m
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 10 %

Mid1 Closed

Description

Class B is characterized by a pond cypress savanna where a lack of fire has led to hardwood encroachment in the understory, and a decline in the herbaceous graminoid groundcover.

Dominant Species* and Canopy Position

TAAS Upper
 ACRU Lower
 SACA5 Lower
 COER2 Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 5

Structure Data (for upper layer lifeform)

	Min	Max
Cover	25 %	100 %
Height	Tree Regen <5m	Tree Medium 10-24m
Tree Size Class	Large 21-33"DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 75%

Mid1 Open
Description

Class C is a pond cypress savanna in which low intensity fire in combination with winter ENSO (El Nino/la nina) related drought has maintained an open pond cypress savanna with a low density, low diversity herbaceous graminoid groundcover.

Dominant Species* and Canopy Position

TAAS Upper
RHMI7 Lower
MUF13 Lower
CLMAJ Lower

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 2

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	25 %
Height	Tree Regen <5m	Tree Medium 10-24m
Tree Size Class	Large 21-33"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 0%

Late1 All Structu
Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	0 %
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 0%

Late1 All Structu
Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: 1200
 Min: 10
 Max: 10000

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Fire Regime Group: 1

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	120			0.00833	17
<i>Mixed</i>	75			0.01333	27
<i>Surface</i>	35			0.02857	57
<i>All Fires</i>	20			0.05024	

References

Anonymous. 1978. Ecological communities-climatic zones Florida. Publisher unknown. Approximately 80 pp.

Brownlie, Dave. U.S. Fish and Wildlife Service. Tallahassee, Fl.

Duever, M.J., et al. 1986. The Big Cypress National Preserve. Res. Rep. No. 8, Nat'l Audubon Society, New York.

Ewel, K. C. 1990b. Swamps. Pages 281-323 in: R. L. Myers and J. J. Ewel, editors. Ecosystems of Florida. University of Central Florida Press, Orlando.

Floherschütz, E. W. 1978. Dwarf cypress in the Big Cypress Swamp of southwestern Florida. Master's thesis, University of Florida, Gainesville. 161 pp.

Muss, Jordan D., Austin, Daniel F. and Snyder, James R. 2003. Plants of the Big Cypress National Preserve. Journal of the Torrey Botanical Society 130(2): 119-142.

Myers, R. L. and John J. Ewel, eds. 1990. Ecosystems of Florida. University of Central Florida Press, Orlando.

NatureServe. 2005. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of February 25, 2005.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.