

**Table 19. Reported AIDS cases for male adults and adolescents, by transmission category and race/ethnicity, 2005 and cumulative—United States and dependent areas**

Transmission category	2005		Cumulative <sup>a</sup>	
	No.	%	No.	%
<b>White, not Hispanic</b>				
Male-to-male sexual contact	6,820	65	248,186	73
Injection drug use	831	8	31,275	9
Male-to-male sexual contact and injection drug use	799	8	30,401	9
Hemophilia/coagulation disorder	50	0	3,971	1
High-risk heterosexual contact <sup>b</sup>	344	3	7,454	2
Sex with injection drug user	68	1	2,249	1
Sex with person with hemophilia	0	0	35	0
Sex with HIV-infected transfusion recipient	1	0	177	0
Sex with HIV-infected person, risk factor not specified	275	3	4,993	1
Receipt of blood transfusion, blood components, or tissue	22	0	3,193	1
Other/risk factor not reported or identified	1,569	15	15,815	5
<b>Total</b>	<b>10,435</b>	<b>100</b>	<b>340,295</b>	<b>100</b>
<b>Black, not Hispanic</b>				
Male-to-male sexual contact	4,731	36	99,269	37
Injection drug use	2,059	16	82,233	30
Male-to-male sexual contact and injection drug use	596	4	21,260	8
Hemophilia/coagulation disorder	11	0	598	0
High-risk heterosexual contact <sup>b</sup>	1,958	15	28,104	10
Sex with injection drug user	224	2	6,675	2
Sex with person with hemophilia	3	0	34	0
Sex with HIV-infected transfusion recipient	8	0	218	0
Sex with HIV-infected person, risk factor not specified	1,723	13	21,177	8
Receipt of blood transfusion, blood components, or tissue	37	0	1,210	0
Other/risk factor not reported or identified	3,868	29	38,097	14
<b>Total</b>	<b>13,260</b>	<b>100</b>	<b>270,771</b>	<b>100</b>
<b>Hispanic</b>				
Male-to-male sexual contact	2,884	44	61,824	43
Injection drug use	1,213	19	45,068	32
Male-to-male sexual contact and injection drug use	293	4	10,390	7
Hemophilia/coagulation disorder	5	0	457	0
High-risk heterosexual contact <sup>b</sup>	740	11	10,301	7
Sex with injection drug user	93	1	2,325	2
Sex with person with hemophilia	0	0	11	0
Sex with HIV-infected transfusion recipient	4	0	119	0
Sex with HIV-infected person, risk factor not specified	643	10	7,846	6
Receipt of blood transfusion, blood components, or tissue	13	0	642	0
Other/risk factor not reported or identified	1,392	21	13,841	10
<b>Total</b>	<b>6,540</b>	<b>100</b>	<b>142,523</b>	<b>100</b>

**Table 19. Reported AIDS cases for male adults and adolescents, by transmission category and race/ethnicity, 2005 and cumulative—United States and dependent areas (cont)**

Transmission category	2005		Cumulative <sup>a</sup>	
	No.	%	No.	%
<b>Asian/Pacific Islander</b>				
Male-to-male sexual contact	214	55	4,317	68
Injection drug use	18	5	312	5
Male-to-male sexual contact and injection drug use	12	3	261	4
Hemophilia/coagulation disorder	2	1	71	1
High-risk heterosexual contact <sup>b</sup>	44	11	356	6
Sex with injection drug user	0	0	55	1
Sex with person with hemophilia	0	0	1	0
Sex with HIV-infected transfusion recipient	0	0	9	0
Sex with HIV-infected person, risk factor not specified	44	11	291	5
Receipt of blood transfusion, blood components, or tissue	1	0	114	2
Other/risk factor not reported or identified	97	25	883	14
<b>Total</b>	<b>388</b>	<b>100</b>	<b>6,314</b>	<b>100</b>
<b>American Indian/Alaska Native</b>				
Male-to-male sexual contact	63	48	1,351	55
Injection drug use	20	15	388	16
Male-to-male sexual contact and injection drug use	19	15	426	17
Hemophilia/coagulation disorder	0	0	31	1
High-risk heterosexual contact <sup>b</sup>	11	8	109	4
Sex with injection drug user	4	3	34	1
Sex with person with hemophilia	0	0	0	0
Sex with HIV-infected transfusion recipient	0	0	3	0
Sex with HIV-infected person, risk factor not specified	7	5	72	3
Receipt of blood transfusion, blood components, or tissue	0	0	9	0
Other/risk factor not reported or identified	18	14	150	6
<b>Total</b>	<b>131</b>	<b>100</b>	<b>2,464</b>	<b>100</b>
<b>Total cases</b>				
Male-to-male sexual contact	14,819	48	416,232	54
Injection drug use	4,168	13	159,676	21
Male-to-male sexual contact and injection drug use	1,742	6	62,940	8
Hemophilia/coagulation disorder	68	0	5,142	1
High-risk heterosexual contact <sup>b</sup>	3,110	10	46,533	6
Sex with injection drug user	389	1	11,371	1
Sex with person with hemophilia	3	0	81	0
Sex with HIV-infected transfusion recipient	13	0	529	0
Sex with HIV-infected person, risk factor not specified	2,705	9	34,552	5
Receipt of blood transfusion, blood components, or tissue	73	0	5,190	1
Other/risk factor not reported or identified	6,976	23	69,095	9
<b>Total</b>	<b>30,956<sup>c</sup></b>	<b>100</b>	<b>764,808<sup>d</sup></b>	<b>100</b>

<sup>a</sup> Reported from the beginning of the epidemic through 2005.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes 202 males of unknown race or multiple races.

<sup>d</sup> Includes 2,441 males of unknown race or multiple races.