

## Complete Summary

---

### **GUIDELINE TITLE**

Preventive health recommendations for adults with mental retardation.

### **BIBLIOGRAPHIC SOURCE(S)**

Massachusetts Department of Mental Retardation, Univ of Massachusetts Medical School's Center for Developmental Disabilities Evaluation and Research. Preventive health recommendations for adults with mental retardation. Boston (MA): Massachusetts Department of Mental Retardation and University of Massachusetts Medical School's Center for Developmental Disabilities Evaluation and Research; 2003 Sep 19. 2 p.

### **GUIDELINE STATUS**

This is the current release of the guideline.

## COMPLETE SUMMARY CONTENT

SCOPE  
 METHODOLOGY - including Rating Scheme and Cost Analysis  
 RECOMMENDATIONS  
 EVIDENCE SUPPORTING THE RECOMMENDATIONS  
 BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS  
 QUALIFYING STATEMENTS  
 IMPLEMENTATION OF THE GUIDELINE  
 INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT  
 CATEGORIES  
 IDENTIFYING INFORMATION AND AVAILABILITY  
 DISCLAIMER

## SCOPE

### **DISEASE/CONDITION(S)**

- Mental retardation
- Cancer (breast, cervical, colorectal, testicular, prostate, and skin)
- Infectious diseases (chlamydia, sexually transmitted diseases, human immunodeficiency virus [HIV], hepatitis B and C, tuberculosis, influenza, pneumonia)
- Hypertension
- Cholesterolemia
- Diabetes
- Osteoporosis
- Hearing and vision impairments

- Glaucoma
- Depression
- Dementia

## **GUIDELINE CATEGORY**

Prevention  
Screening

## **CLINICAL SPECIALTY**

Family Practice  
Internal Medicine  
Preventive Medicine

## **INTENDED USERS**

Health Care Providers  
Physician Assistants  
Physicians

## **GUIDELINE OBJECTIVE(S)**

To present preventive health recommendations for adults with mental retardation

## **TARGET POPULATION**

Adults with mental retardation

## **INTERVENTIONS AND PRACTICES CONSIDERED**

*Specific preventive health interventions and practices for the Department of Mental Retardation-mandated annual physical exam:*

### **Screening**

#### *Cancer Screening*

1. Clinical and self breast examination
2. Mammography
3. Clinical and self testicular examination
4. Pap smear
5. Fecal occult blood testing
6. Skin examination

#### *Infectious Disease Screening*

Chlamydia, sexually transmitted disease, human immunodeficiency virus [HIV], hepatitis B and C, tuberculosis testing

### *Sensory Screening*

Hearing and vision testing

### *Mental and Behavioral Health Screening*

Depression and dementia assessments

### *Other Screening*

1. Thyroid function test
2. Cervical spine x-ray
3. Echocardiogram
4. Hypertension testing
5. Cholesterol testing
6. Diabetes (Type II) testing
7. Liver function testing
8. Osteoporosis testing

### **Prevention**

1. Immunization for influenza, pneumococcus, hepatitis B
2. General counseling and guidance

### **MAJOR OUTCOMES CONSIDERED**

Not stated

## **METHODOLOGY**

### **METHODS USED TO COLLECT/SELECT EVIDENCE**

Searches of Electronic Databases

### **DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE**

Comprehensive review of available Medline/PubMed sources, materials developed by advocacy and state agencies.

### **NUMBER OF SOURCE DOCUMENTS**

16 documents used by the expert panel

### **METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE**

Expert Consensus (Committee)

### **RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE**

Not applicable

## **METHODS USED TO ANALYZE THE EVIDENCE**

Review

## **DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE**

Not stated

## **METHODS USED TO FORMULATE THE RECOMMENDATIONS**

Expert Consensus

## **DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS**

The development of the recommendations required careful consideration of many elements in the lives of adults with mental retardation.

### **Phase I**

Committee members received a package of reference material for review prior to the initial meeting. Members reviewed the general findings from the literature search and discussed their own professional experience in community care for people with mental retardation. Following discussion, committee members agreed to adopt generally accepted *Massachusetts Health Quality Partners (MHQP)* guidelines for the general population as a baseline and develop recommendations for modifications to address areas of greater concern for the population of people with mental retardation.

Based on findings in the literature and the experience of committee members, three focus areas emerged:

1. The need to review and modify generic MHQP standards to ensure they are appropriate to the health needs of the adult population with mental retardation.
2. The need to facilitate effective communication relating to an individual's health, lifestyle, adaptive functioning, and service plan between residential staff or family members and the community clinician.
3. The need to consider additional recommendations for sub-populations such as those with identifiable syndromes or comorbid conditions that are associated with additional health risks.

The committee formed three sub-groups and each group was charged with developing recommendations to address one of the above areas.

### **Phase II**

Subcommittees met to develop recommendations that were shared at a second Advisory Committee meeting. The entire Advisory Committee reviewed and

commented on the work of each subcommittee. The subcommittees then formed their final recommendations.

### Phase III

The final meeting and discussion of the Advisory Committee was held for discussion of the remaining work of each committee. Subcommittee members agreed to complete recommendations for the final report to be forwarded to the Department of Mental of Retardation.

## RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

## COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

## METHOD OF GUIDELINE VALIDATION

Peer Review

## DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Not stated

## RECOMMENDATIONS

### MAJOR RECOMMENDATIONS

**Note:** Items that are indicated in ***bold italic*** are specific recommendations that differ from the Massachusetts Health Quality Partners (MHQP) recommendations in order to reflect particular health concerns of the population with mental retardation.

Health Maintenance Visit, including height and weight measurement

- 19-29 years: ***annually***
- 30-39 years: ***annually***
- 40-49 years: ***annually***
- 50-64 years: annually
- 65+ years: annually

Breast Cancer: Mammography

- 19-39 years: Clinical breast exam and self-exam and self-exam instruction as appropriate. Mammography not routine except for patients at high risk. Accurate and detailed history and family history will identify risk factors.

- 40-49 years: Clinical breast exam and self-exam instruction as appropriate. Mammography every 1 to 2 years, at discretion of physician.
- 50-64 years: Clinical breast exam and self-exam instruction as appropriate. Annual mammography
- 65+ years: Mammography annually through age 69 years. Age 70 years and older, annually at the discretion of the physician.

#### Cervical Cancer: Pap Smear

- 19-64 years: Every 1 to 3 years, at physician's discretion
- 65+ years: May be omitted after age 65 if previous screenings were consistently normal

#### Colorectal Cancer

- 19-49 years: Not routine except for patients at high risk
- 50-65+ years: Fecal Occult Blood testing annually and sigmoidoscopy every 5 years OR colonoscopy every 10 years

#### Testicular and Prostate Cancer

- 19-39 years: Clinical testicular exam and self-exam instruction as appropriate. Prostate cancer screening not routine.
- 40-49 years: Not routine except for patients at high risk. Risk factors include: family history and African-American ancestry.
- 50-65+ years: At physician discretion after discussion of risks and benefits of available screening strategies (prostate specific antigen [PSA], digital rectal examination [DRE]).

#### Skin Cancer

- 19-65+ years: Periodic total skin exams targeting populations at high risk for malignant melanomas. Frequency at physician discretion.

#### Hypertension

- 19-65+ years: ***At least annually***

#### Cholesterol

- 19-64 years: Every five years or at physician discretion
- 65+ years: at physician discretion

#### Diabetes (Type II)

- 19-65+ years: ***At least every 5 years until age 45.*** Every 3 years after age 45. Fasting plasma glucose screen for individuals at high risk. Risk factors include: family history of premature coronary heart disease (CHD), hypertension, diabetes mellitus, peripheral atherosclerosis or carotid artery disease, current cigarette smoking, or high-density lipoprotein (HDL) >35 mg/dL.

***Liver Function***

- 19-65+ years: Annually for hepatitis B carriers. At physician discretion after consideration of risk factors including long term prescription medication

***Osteoporosis***

- 19-64 years: Bone density screening when risk factors are present: long term polypharmacy, mobility impairments, hypothyroid, post-menopausal women. Periodicity of screening at physician discretion. Annually counsel about preventive measures including dietary calcium and vitamin D intake, weight-bearing exercise, and smoking cessation.
- 65+ years: Counsel elderly patients about specific measures to prevent falls.

***Chlamydia and Sexually Transmitted Diseases (STDs)***

- 19-29 years: For all sexually active males and females screen annually <25 years. >25 years screen annually if at risk.
- 30-65+ years: Annually if at risk. Risk factors include inconsistent use of barrier contraceptives, new or multiple sex partners in last 3 months, a new partner since last test, a history of STD, infected with another STD, and partner has had other sexual partner(s).

***Human Immunodeficiency Virus (HIV)***

- 19-65+ years: Periodic testing if at risk and testing of pregnant women at increased risk

***Hepatitis B and C***

- 19-65+ years: Periodic testing if risk factors present

***Tuberculosis***

- 19-65+ years: Tuberculin skin testing every 1 to 2 years when risk factors present. Risk factors include residents or employees of congregate setting, close contact with persons known or suspected to have tuberculosis (TB).

***Hearing assessment***

- 19-65+ years: Screen annually. Re-evaluate if hearing problem is reported or a change in behavior is noted.

***Vision assessment***

- 19-65+ years: Screen annually. Re-evaluate if vision problems are reported or a change in behavior is noted.

***Eye Exam for Glaucoma***

- 19-39 years: Every 3 to 5 years in high risk patients. At least once in patients with no risk factors.
- 40-64 years: every 2 to 4 years
- 65+ years: every 1 to 2 years

### ***Depression***

- 19-65+ years: Screen annually for sleep, appetite disturbance, weight loss, general agitation.

### ***Dementia***

- 19-39 years: Monitor for problems performing daily activities.
- 40-65+ years: In persons with Down Syndrome, annual screen after age 40.

Immunizations (as for the general adult population with the following exceptions)

1. Influenza vaccine  
19-65+ years: Annually
2. Pneumococcal vaccine  
19-65+ years: Once
3. Hepatitis B vaccine  
19-65+ years: Once. Reevaluate antibody status every 5 years.

For persons with Down Syndrome (in addition to the above recommendations)

1. Thyroid function test  
19-65+ years: Every 3 years (sensitive thyroid stimulating hormone [TSH])
2. Cervical spine x-ray to rule out atlanto-axial instability  
19-65+ years: Obtain baseline as adult. Recommend repeat if symptomatic, or 30 years from baseline.
3. Echocardiogram  
19-65+ years: Obtain baseline if no records of cardiac function are available.

General counseling and guidance

1. Prevention counseling  
19-65+ years: Annually counsel regarding prevention of accidents related to falls, fire/burns, choking.
2. Abuse or neglect  
19-65+ years: Annually monitor for behavioral signs of abuse and neglect.
3. Preconception counseling  
19-65+ years: As appropriate, including genetic counseling, folic acid supplementation, discussion of parenting capability.
4. Healthy lifestyle  
19-65+ years: Annually counsel regarding diet/nutrition, incorporating regular physical activity into daily routines, substance abuse.

### **General Counseling and Guidance Recommendations**



**Injury Prevention & Safety:** *Annually* counsel caregivers to be alert to ways to prevent household injuries (fall prevention, choking prevention, fire/burn prevention).

**Family Violence/Abuse:** On *all visits* be alert to physical and behavioral signs and symptoms associated with abuse and neglect. Routinely ask all patients direct, specific questions about abuse including sexual abuse.

**Preconception Counseling:** Preconceptual counseling should include genetic counseling for hereditary conditions, assessment and discussion of parenting capability, as well as folic acid supplementation.

**Menopause Management:** *Ages 40 – 50.* Be alert to signs and symptoms of menopause; consider appropriate therapies to control symptoms.

**Other testing considerations when individual has limited communication capacity.**

For individuals who have significantly limited means of communicating their symptoms, screening labs (blood chemistries, complete blood count [CBC], perhaps TSH) should be considered at each annual health maintenance visit in order to supplement available information. Alternatively, in order to prevent unnecessary follow-up evaluations, it may be preferable to limit testing to the evaluation of specifically identified problems when a thorough and detailed history can be obtained from familiar caregivers.

**Preparation for an examination**

Many adults can be helped to feel more comfortable at a medical visit if they feel adequately prepared for the event. Family or support staff can be encouraged to introduce unfamiliar items such as a stethoscope or a blood pressure cuff at home to allow the instrument to become more familiar and facilitate cooperation during an exam.

During the exam, it is helpful to prepare patients for procedures by explaining them well or allowing patients with sensory impairments to explore the instruments that are about to be used. Performing simple examinations in an office or quiet waiting room may reduce a person's anxiety.

If someone is particularly anxious or an invasive screening procedure is necessary, the clinician might consider sedation prior to the appointment. In some cases, multiple procedures can be performed while the patient is sedated (dental work, or routine blood work, for example) to reduce the number of times a person is exposed to the risks of sedation.

Refer to Attachment D of the companion document "Preventive Health Standards Project FINAL REPORT," Center for Developmental Disabilities Evaluation and Research.

**Communication**

If a patient is unable to communicate regarding his or her own health status, clinicians may have to rely on a family member or support staff to provide information relating to signs or symptoms of health concerns. Questions regarding changes in the individual's behavior and adaptive function can bring underlying physical and mental health issues to light.

### **CLINICAL ALGORITHM(S)**

None provided

## **EVIDENCE SUPPORTING THE RECOMMENDATIONS**

### **TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS**

The recommendations are based on the [Massachusetts Health Quality Partnership \(MHQP\)](#) Adult Preventive Care Recommendations 2003.

## **BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS**

### **POTENTIAL BENEFITS**

Appropriate preventive healthcare for adults with mental retardation

### **POTENTIAL HARMS**

Not stated

## **QUALIFYING STATEMENTS**

### **QUALIFYING STATEMENTS**

With the presentation of these recommendations, the Advisory Committee does not intend to imply that adults with mental retardation all have similar health needs or that clinicians should not continue to use their clinical judgment and seek additional consultation when necessary. The Advisory Committee sees these recommendations as one of a number of steps towards ensuring that every adult with mental retardation receives the most appropriate health care as an individual.

## **IMPLEMENTATION OF THE GUIDELINE**

### **DESCRIPTION OF IMPLEMENTATION STRATEGY**

Massachusetts Department of Mental Retardation had included prompts for preventive health screening as a required element in their residential provider systems. Recommendations pamphlet mailed to all general and family practitioners in the Commonwealth.

## INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

### IOM CARE NEED

Living with Illness  
Staying Healthy

### IOM DOMAIN

Effectiveness  
Patient-centeredness

## IDENTIFYING INFORMATION AND AVAILABILITY

### BIBLIOGRAPHIC SOURCE(S)

Massachusetts Department of Mental Retardation, Univ of Massachusetts Medical School's Center for Developmental Disabilities Evaluation and Research. Preventive health recommendations for adults with mental retardation. Boston (MA): Massachusetts Department of Mental Retardation and University of Massachusetts Medical School's Center for Developmental Disabilities Evaluation and Research; 2003 Sep 19. 2 p.

### ADAPTATION

The recommendations are based on the [Massachusetts Health Quality Partners \(MHQP\)](#) Adult Preventive Care Recommendations 2003.

### DATE RELEASED

2003 Sep 19

### GUIDELINE DEVELOPER(S)

Massachusetts Department of Mental Retardation - State/Local Government Agency [U.S.]  
University of Massachusetts Medical School's Center for Developmental Disabilities Evaluation and Research - Academic Institution

### SOURCE(S) OF FUNDING

Massachusetts Department of Mental Retardation

### GUIDELINE COMMITTEE

Massachusetts Department of Mental Retardation Preventive Health Standards Advisory Committee

## **COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE**

*Committee members:* Victor Acquista, MD; Norberto Alvarez, MD; Bob Baldor, MD; Allen Crocker, MD; Marc Emmerich, MD; Warren Ferguson, MD; Jim Gleason, PT; Adria Hodas, RN NP; Elizabeth King, RN NP; Leo McKenna, PharmD.; Barbara Pilarcik, RN; Julie Lufkin Purtz, RN NP; Van R. Silka, MD; Carol Walsh RNC, MPH; Susan Weiner, OT

## **FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST**

Not stated

## **GUIDELINE STATUS**

This is the current release of the guideline.

## **GUIDELINE AVAILABILITY**

Print copies: Available from the University of Massachusetts Medical School, Center for Developmental Disabilities Evaluation and Research, 200 Trapelo Road, Waltham, MA 02452; Phone: (781) 642-0283; Fax: (781) 642-0162.

## **AVAILABILITY OF COMPANION DOCUMENTS**

The following are available:

- Preventive health recommendations for adults with mental retardation. Guidelines for community practitioners. Boston (MA): Massachusetts Department of Mental Retardation and University of Massachusetts Medical School Center for Development Disabilities Evaluation and Research; 2002 (revised 2003 Sep). 2 p.
- Preventive health standards project. Final report. Center for Developmental Disabilities Evaluation and Research; 2002 (revised 2003 Apr 11). 54 p.

Print copies: Available from the University of Massachusetts Medical School, Center for Developmental Disabilities Evaluation and Research, 200 Trapelo Road, Waltham, MA 02452; Phone: (781) 642-0283; Fax: (781) 642-0162.

## **PATIENT RESOURCES**

None available

## **NGC STATUS**

This summary was prepared by ECRI on January 29, 2004. The information was verified by the guideline developer on February 26, 2004.

## **COPYRIGHT STATEMENT**

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions. Refer to the Center for Developmental Disabilities Education and Research (E.K. Shriver Center for Mental Retardation) for permission to reproduce guidelines.

## DISCLAIMER

### NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 9/22/2008

