

# First Spectra of Neon, Argon, and Xenon 136 in the 1.2–4.0 $\mu\text{m}$ Region

Curtis J. Humphreys\*

Department of Physics, Purdue University, Lafayette, Indiana 47907

Descriptions of the first spectra of neon, argon, and xenon 136, comprising calculated wavelengths, calculated wave numbers, relative intensities, and classifications, are presented. The calculated values are derived from currently best established energy levels, obtained mostly from interferometric observations and adopted as standards by the International Astronomical Union. All listed lines have actually been observed. This paper makes available a compilation of all results previously presented in fragmentary or relatively inaccessible reports with intensities normalized to as nearly a uniform scale as the various observations permit.

Key words: Argon; extraphotographic region; infrared emission spectra; intensities; neon; wavelengths; wavelength standards; wave numbers; xenon.

## 1. Experimental Background

The wavelength limits of the coverage specified in the title represent at the lower end the practical upper limit of photographic response,<sup>1</sup> and at the upper end the long wavelength limit of response of lead sulfide detectors cooled to the temperature of liquid nitrogen. Both of these limits have been set somewhat arbitrarily. Abundant photographic observations at wavelengths short of 1.2  $\mu\text{m}$  have not been significantly augmented by radiometric methods. At the upper end of the scale the dearth of energetic transitions at greater wavelengths than the 4f–5g transitions, which have been reported in a previous publication, [1]<sup>2</sup> do not appear to justify the use of detectors responsive to wavelengths greater than 4  $\mu\text{m}$ .

Very complete descriptions of the noble-gas spectra in the photographic infrared region became available in a series of three NBS publications that appeared about forty years ago. The first by Humphreys and Meggers [2] described Xe I, the second by Meggers and Humphreys [3] covered Ne I, Ar I, and Kr I, and the third by Meggers [4] extended the descriptions of all four of these together with helium to the limit of response of sensitized photographic emulsions, around 13 000 Å.

The investigation of these spectra in the extraphotographic infrared was carried out mostly as a program started in the Radiometry section at the National Bureau of Standards and continued in the Research Department of the former Naval Weapons Center, Corona, California. All lines or unresolved features reported in this article were actually observed in connection with the program. A significant number of these features has been observed independently in other

laboratories. Such work is referenced and notes are introduced into the tables to indicate prior or improved observations.

During 1949, taking advantage of the then newly-available lead sulfide detectors developed and constructed by Cashman [5], Sittner and Peck [6] observed the spectra of argon, krypton, and xenon between about 1.2 and 1.8  $\mu\text{m}$ . They excited the spectra in flash tubes. A short time later Humphreys and Kostkowski [7] at the National Bureau of Standards reobserved these spectra over the same region, making significant extensions and including helium and neon in the program. Geissler tubes were employed as sources.

Additional results, permitted by use of microwave-excited electrodeless tubes were reported by Humphreys and Paul in 1958 [8]. During the same period Hepner reported observations of Ne I and Ar I [9], also a little earlier Xe I [10]. Wavelength measurements on Ne I by Rao are included in the book *Wavelength Standards in the Infrared* [11]. During the period 1960 to 1965 increments to the descriptions resulting from refinements of observing techniques were reported at meetings of the Optical Society of America [12], [13]. In 1965 Séguier [14] reported the observation of several resolved complex features of Ar I between 1.2 and 1.8  $\mu\text{m}$  by use of a grille spectrometer as designed by Girard [15].

Use of liquid-nitrogen cooling of PbS detectors begun by the author about 1961 permitted extension of descriptions of noble-gas spectra to 4.0  $\mu\text{m}$  and interpretation of the 4f–5g transitions that represented the first observed examples in atomic spectra of essentially pure pair coupling. The results were presented at several meetings [16], [17] and reported in final form in an article published jointly with Paul, Cowan, and Andrew in 1967 [1] and are not included in this article.

Reobservation of the 3d–4f transitions in Ne I by Johansson [18] in 1964, and of the 3d–5f transitions by Litzén [19] in 1968, has led to wavelength values considered somewhat more precise than those previously reported.

An essentially complete description of Kr I including the infrared region is provided by the article by Kaufman and Humphreys [20]. Data pertaining to that spectrum

\*Work performed partly at the Naval Weapons Center, Corona, California, under the Foundational Research Program and partly at Purdue with support by the National Aeronautics and Space Administration and the National Science Foundation.

<sup>1</sup>The description of Ne I is started with  $\lambda$  11 143 Å because several very intense lines of the category 3p–4s that are very precisely measured and therefore promising as wavelength standards are located between this point and 12 000 Å.

<sup>2</sup>Figures in brackets indicate the literature references.

Copyright © 1973 by the U.S. Secretary of Commerce on behalf of the United States. This copyright will be assigned to the American Institute of Physics and the American Chemical Society, to whom all requests regarding reproduction should be addressed.

are omitted therefore from this paper. The article on krypton does not include transitions involving  $f$ -levels, since those levels had not been determined precisely by interferometric methods. Essentially all the missing transitions are described in the paper by Humphreys and Kostkowski [7].

The most extensive available descriptions of noble-gas spectra that include coverage of the infrared region as far as  $2.5 \mu\text{m}$  are to be found in the book by Striganov and Sventitskii [21]. Those authors have utilized the data contained in articles referenced here to prepare the compilations. *Wavelength Standards in the Infrared* [11] contains wavelengths, wave numbers, and intensities of the stronger lines of neon, argon, krypton, and xenon. The wavelength entries are in most instances derived from interferometric determinations of the energy levels involved. A list of calculated vacuum wavelengths of Ar I in the infrared was given by Humphreys [22] in an article describing a new interferometric method of wavelength determination. A similar compilation of argon wavelengths is to be found in the report of Commission 14 of the International Astronomical Union [23] covering the meeting of 1964. A compilation of wavelengths of  $^{136}\text{Xe I}$  in the infrared is included in a recent article by Humphreys and Paul [24] reporting interferometric measurements. The list of infrared lines in that article is brief, restricted to only the very intense transitions observable interferometrically.

About 1955 a unique radiometric technique for infrared wavelength determination was developed by Humphreys and Paul and first reported to the OSA [25] and later in the referenced article [22]. These investigations will be discussed in the following sections pertaining specifically to individual spectra only as necessary to explain the origin of the energy-level tables included.

An important consideration prompting the assembly of this material has been the need to achieve a consistent representation of relative intensities. The spectra were reported in small increments covering groups of observed features that had been produced under varying conditions. In each instance relative intensities were estimated within the groups, but there was no direct relationship between the scales applicable to different groups. Following the transfer of the infrared equipment from the Corona Laboratories to Purdue, the amplifying system was modified by the addition of a module to yield a logarithmic output. This modification was designed and built principally by Duane Saufley. Recordings were made of the three spectra under discussion using the modified output. These recordings have provided the principal basis for the estimates of relative intensities reported here. These intensities regarded as reliable over several hundred angstroms are not claimed to be absolutely accurate for the entire interval covered, since the spectral response of the detectors has not been taken into account and the

properties of the gratings for different spectral regions only to a limited extent.

The presentation in the form of calculated wavelengths and wave numbers has been chosen because the use of officially adopted level values achieves a precision better by at least one significant figure over that obtainable by direct scanning except in the instance of features observed interferometrically. Such features are indicated where they appear in the tables.

## 2. Discussion of Individual Spectra

### 2.1. Ne I

Neon has been used for a very long time as a source of wavelength standards mainly because of the very intense red lines arising from  $3s-3p$  transitions whose wavelengths have been known with high precision for half a century. Most of the energy of the infrared spectrum is in the photographic region. The region between  $1.2$  and  $4.0 \mu\text{m}$  is however fairly well populated and many of the lines are particularly useful as internal standards in instances where neon is employed as a carrier gas in the construction of sources.

Most observations of Ne I have utilized the naturally occurring mixture of the isotopes of mass numbers 20 and 22 in the proportions of about 10 to 1. It is now possible to obtain essentially pure isotopes. However a refinement of observing techniques greater than is usually available is necessary to distinguish between the wavelengths of lines originating respectively in natural neon and neon of mass number 20 or to display the isotope effect in the former. This is because the natural line widths preclude the use of very long interference paths.

In common with all noble-gas spectra Ne I exhibits well-developed  $jK$  coupling. This is shown to an increasing extent for levels of large  $l$ -value. The  $f$ -level pairs are unresolvable. Owing to small electrostatic interaction and consequent small values of the  $F$ -parameters the levels of given  $l$ -value tend to form very compact groups. This situation also affects the distribution of the transitions so that the pattern of lines is one of relatively isolated groups. Abundance of overlapping transitions makes it difficult to find single lines among the  $p-d$  and  $d-f$  groups. Many of the  $s-p$  transitions are however sufficiently isolated to offer promise of usefulness as standards.

The description of Ne I appears in table 1. It contains 221 entries of which a considerable number are components of multiple features unresolved on the records. The calculated wavelengths and wave numbers are derived from the set of energy levels recently reported by Kaufman and Minnhagen [26]. No table of neon energy levels is therefore included here. The relative values of most of these levels have been approved by

Commission 14 of the I.A.U.<sup>3</sup> The 3s (not considered in this description), 3p, and 4p levels were adopted by I.A.U. at the 1955 General Assembly [27]. Values of 4s and 3d levels were recommended at the 1958 Assembly [28]. A revised set of 4s levels, determined by interferometric observations in the infrared by Humphreys and Paul, NAVWEPS Report 7190, was reported at the 1964 Assembly. These 4s levels were utilized in combination with the previously adopted 3p and 4p levels to prepare a new set of calculated wavelengths of the 3p-4s and 4s-4p transitions, that are reported along with the 4s levels [23]. These transitions covering the range from 8865 to 33 511 Å appear promising as wavelength standards for the infrared region. Values of the 4d levels have not been adopted by the I.A.U.

Kaufman and Minnhagen [26] have discussed the selection of level values other than I.A.U. adoptions. In particular these selections include the values of 4f levels reported by Johannson [18] and the 5f by Litzén [19]. The absolute scale of the level system has been established by setting the value of the ground level  $s^2p^6\ ^1S_0$  equal to zero. This is the same practice used by Moore-Sitterly in A.E.L. [29]. However a new measurement of the wave number of the transition  $2p^6\ ^1S_0 - 2p^5\ (^3P_{1/2})3s[1\frac{1}{2}]^0_1$  by Kaufman and Minnhagen [26] giving 134 459.28 cm<sup>-1</sup> and superseding the result reported by Petersson [30] has led to a shift of the scale relative to A.E.L. amounting to 1.950 cm<sup>-1</sup>.

The precision of the calculated wave numbers and wavelengths depends upon the uncertainties of the energy-level values entered into the computations. Most of the levels have been adopted by the I.A.U. The rest were selected on the basis of the best experimental evidence available. In some cases these had not been adopted by the I.A.U. because of an insufficient number of independent concordant observations. In all cases the values of the levels were carried to  $10^{-4}$  or  $10^{-3}$  cm<sup>-1</sup>. Accordingly where both levels involved in a transition are carried to four or three decimal places, the same number of places are retained in the wave numbers. For the sake of uniformity and following the practice of the I.A.U. in recent reports the same number of places is retained for the wavelengths. At a little beyond 22 000 Å an uncertainty of 0.001 cm<sup>-1</sup> in the wave number is equivalent to 0.005 Å in the wavelength. It is therefore probably not advantageous to express wavelengths to more than two decimal places beyond that point. The actual location of an unresolved feature involving a transition between a single level and two others that are pair coupled depends on the relative intensities of the two resulting component lines. Such features should not be selected as wavelength standards.

<sup>3</sup> Commission 14 has always designated noble-gas levels in the Paschen notation. In this article the particular category of levels is indicated by the *l*-value of the valence electron preceded by its *n*-value.

## 2.2 Ar I

Naturally occurring argon is a mixture of three isotopes of even mass number, of which 99.6 percent is the isotope of mass number 40. The spectrum is therefore entirely free of hyperfine structure, and for practical purposes, of isotope effect. It has been used extensively as a source material for producing wavelength standards because of the homogeneity of the lines, the abundance of the element, and ease of operation of sources of simple design. Values of the energy levels, 4s, 4p, and 5p were adopted by Commission 14 of the I.A.U. at the 1955 Assembly [27]. These adopted values were based mainly on interferometric measurements by Burns and Adams [31], by Littlefield and Turnbull [32], and by Humphreys [33]. Following the first submission of interferometrically measured infrared wavelengths in the extraphotographic region to the I.A.U. Assembly in 1958, Commission 14 provisionally adopted the values of the 5s and 3d levels. Confirmatory concordant results obtained independently were reported at the 1961 Assembly [34]. Following subsequent new interferometric observations in the 1.0 to 2.0 μm region by Humphreys and Paul, Commission 14 adopted improved values of the 5s and 3d levels at the 1964 Assembly [23]. Further interferometric observations in the same region led to a determination of the 4f levels [35], and in one instance to a direct observation of a resolved f-pair. The foregoing explains the selection of most of the levels used in preparing the description. The 4d and 6s levels have been based mostly on the measurements of Burns and Adams [31] and the entries in A.E.L. [29], where the values have not been discussed specifically in this article.

Table 2, comprising 325 entries, displays the description of the Ar I spectrum in the infrared region. These entries are associated with individual transitions. In many instances recorded features represent unresolved related pairs or juxtaposed lines where there is a probable energy contribution from each. Table 3 contains the energy levels entered in the calculation of the description.

The Ar I results were assembled prior to some new interferometric observations in the photographic region. Norlén [36] has published a set of wavelengths between 5152 and 6953 Å. Observations by Li are now in progress at Purdue. The justification for the recent observations is that although the spectrum has been extensively investigated there have not been a sufficient number of concordant observations to permit final adoption of some of the energy levels. The new determinations are not expected to change the values of many of the levels by as much as 0.001 cm<sup>-1</sup>. It has not been regarded as justifiable to await new energy-level values that might supersede some of those used in the reported calculated wavelengths.

### 2.3. $^{136}\text{Xe}$

Interest in xenon as a source material for the production of wavelength standards was aroused only when the techniques of isotope separation made available the isotope of mass number 136. Owing to small Doppler broadening the lines of xenon isotopes of even mass number should be sharper than those of any of the other noble gases. A further special advantage for infrared intercomparisons is the large number of very intense  $6p-5d$  transitions between 2.0 and 4.0  $\mu\text{m}$  and beyond, the spectrum being unique in this respect. Interferometric observations covering both the photographic and radiometric regions were reported recently by Humphreys and Paul [24]. These latter include only 15 lines. The present description is intended to supplement that of the referenced article by including all observed infrared lines. The background of the observations on  $^{136}\text{Xe}$  I and the determination of level values are to be found in the previous article. No table of levels has therefore been included here. The calculated wavelengths and other descriptive data are compiled in table 4 comprising 148 entries. The same comments regarding multiple features apply here as in the instances of tables 1 and 2.

In addition to bringing together the descriptions of the infrared spectra of neon, argon, and xenon 136, in a single compilation, this article should provide a list of wavelengths of sufficient precision and adequate distribution to be useful as a set of wavelength standards for the region covered.

The assistance of Mr. Duane Saufley in making recordings with the logarithmic output is gratefully acknowledged. We are also indebted to Dr. Hui Li for several computer runs by which energy level differences were converted to wavelengths in air. The contributions of Edward Paul Jr. whose unflagging assistance over a period of many years led to the accumulation of a large part of the experimental data are also acknowledged with thanks and appreciation.

### 3. References

- [1] Humphreys, C. J., Paul, E., Jr., Cowan, R. D., and Andrew, K. L., J. Opt. Soc. Am. **57**, 855 (1967).

- [2] Humphreys, C. J., and Meggers, W. F., J. Research Natl. Bur. Standards **10**, 139 (1933).
- [3] Meggers, W. F., and Humphreys, C. J., J. Research Natl. Bur. Standards **10**, 427 (1933).
- [4] Meggers, W. F., J. Research Natl. Bur. Standards **14**, 487 (1935).
- [5] Cashman, R. J., J. Opt. Soc. Am. **36**, 356A (1946).
- [6] Sittner, W. R., and Peck, E. R., J. Opt. Soc. Am. **39**, 474 (1949).
- [7] Humphreys, C. J., and Kostkowski, H. J., J. Research Natl. Bur. Standards **49**, 73 (1952).
- [8] Paul, E., Jr., and Humphreys, C. J., J. Opt. Soc. Am. **49**, 1186 (1959).
- [9] Hepner, G., Compt. rend. **248**, 8 (1959).
- [10] Hepner, G., Compt. rend. **242**, 1430 (1956).
- [11] Rao, K. N., Humphreys, C. J., and Rank, D. H., *Wavelength Standards in the Infrared*, Academic Press, 1966.
- [12] Humphreys, C. J., and Paul, E., Jr., J. Opt. Soc. Am. **50**, 510A (1960).
- [13] Humphreys, C. J., Adams, K. B. and Weiss, A., J. Opt. Soc. Am. **51**, 1465A (1961).
- [14] Séguier, J., Compt. rend. **261**, 3069 (1965).
- [15] Girard, A., Applied Optics **2**, 79 (1963).
- [16] Humphreys, C. J., *Memoires Soc. R. Liege cinquieme serie, tome IX*, 1016 (1964).
- [17] Humphreys, C. J., J. Opt. Soc. Am. **53**, 506A (1963).
- [18] Johansson, I., Arkiv Fysik **25**, 381 (1963).
- [19] Litzén, U., Arkiv Fysik **38**, 317 (1968).
- [20] Kaufman, V., and Humphreys, C. J., J. Opt. Soc. Am. **59**, 1614 (1969).
- [21] Striganov, A. R., and Sventitskii, N. S., *Tables of Spectral Lines of Neutral and Ionized Atoms*, IFI/Plenum Data Corporation, New York, 1968.
- [22] Humphreys, C. J., Applied Optics **2**, 1155 (1963).
- [23] Trans. Int. Astronom. Union XII B, 173-185 (1966).
- [24] Humphreys, C. J., and Paul, E., Jr., J. Opt. Soc. Am. **60**, 1302 (1970).
- [25] Humphreys, C. J., and Paul, E., Jr., J. Opt. Soc. Am. **45**, 903A (1955).
- [26] Kaufman, V., and Minnhagen, L., J. Opt. Soc. Am. **62**, 92 (1972).
- [27] Trans. Int. Astronom. Union IX, 201-227 (1957).
- [28] Trans. Int. Astronom. Union X, 211-233 (1960).
- [29] Moore-Sitterly, C., *Atomic Energy Levels*, Circular of the Natl. Bur. Standards 467, Vol. I, 1949.
- [30] Petersson, B., Arkiv. Fysik **27**, 317 (1964).
- [31] Burns, K., and Adams, K. B., J. Opt. Soc. Am. **43**, 1020 (1953).
- [32] Littlefield, T. A., and Turnbull, D. T., Proc. Roy. Soc. A, **218**, 577 (1953).
- [33] Humphreys, C. J., J. Research Natl. Bur. Standards **20**, 17 (1938).
- [34] Trans. Int. Astronom. Union XI B, 208-221 (1962).
- [35] Humphreys, C. J., and Paul, E., Jr., J. Opt. Soc. Am. **52**, 591A (1962).
- [36] Norlén, G., Arkiv Fysik **35**, 119 (1968).

TABLE 1. Description of Ne I, region 1.1–4.0 μm

Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm⁻¹)	Classification	Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm⁻¹)	Classification
3000 a	11 143.0201	8971.7709	$3p[2\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	700	13 219.2406	7562.6628	$3p'[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$
15	11 160.219	8957.945	$4s[1\frac{1}{2}]^0-5p[2\frac{1}{2}]_3$	20	14 300.830	6990.690	$4p[\frac{1}{2}]_1-5d[1\frac{1}{2}]^0$
3500 a	11 177.5276	8944.0731	$3p[2\frac{1}{2}]_3-4s[1\frac{1}{2}]^0$	18	14 342.163	6970.543	$4p[\frac{1}{2}]_1-5d[\frac{1}{2}]^0$
5	11 292.964	8852.648	$4s[1\frac{1}{2}]^0-5p[1\frac{1}{2}]_2$	4	14 353.371	6965.100	$4p[\frac{1}{2}]_1-5d[\frac{1}{2}]^0$
2	11 298.450	8848.349	$4s[1\frac{1}{2}]^0-5p[1\frac{1}{2}]_1$	4	14 499.925	6894.703	$4p[\frac{1}{2}]_1-5d[1\frac{1}{2}]^0$
50	11 303.85	8844.12	$4s'[1\frac{1}{2}]^0-5p'[1\frac{1}{2}]_2$	20	14 929.803	6696.182	$4p[2\frac{1}{2}]_3-5d[2\frac{1}{2}]^0$
20	11 304.557	8843.569	$4s[1\frac{1}{2}]^0-5p[1\frac{1}{2}]_1$	100	14 931.179	6695.565	$4p[2\frac{1}{2}]_3-5d[2\frac{1}{2}]^0$
10	11 329.613	8824.010	$4s'[1\frac{1}{2}]^0-5p'[1\frac{1}{2}]_1$	100	14 984.856	6671.581	$4p[2\frac{1}{2}]_3-5d[3\frac{1}{2}]^0$
20	11 333.621	8820.890	$4s'[1\frac{1}{2}]^0-5p'[1\frac{1}{2}]_1$	100	14 986.312	6670.933	$4p[2\frac{1}{2}]_3-5d[3\frac{1}{2}]^0$
10	11 366.673	8795.241	$4s[1\frac{1}{2}]^0-5p[2\frac{1}{2}]_2$	?	15 058.992	6638.736	$4p[1\frac{1}{2}]_1-5d[1\frac{1}{2}]^0$
1600 a	11 390.4340	8776.8937	$3p[2\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	40	15 074.171	6632.052	$4p'[1\frac{1}{2}]_1-5d'[2\frac{1}{2}]^0$
1100 a	11 409.1344	8762.5078	$3p'[1\frac{1}{2}]_1-4s'[1\frac{1}{2}]^0$	40	15 074.990	6631.692	$4p'[1\frac{1}{2}]_1-5d'[1\frac{1}{2}]^0$
3000	11 522.7460	8676.1117	$3p'[1\frac{1}{2}]_2-4s'[1\frac{1}{2}]^0$	20	15 083.904	6627.772	$4p[2\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
1500	11 525.0195	8674.4002	$3p[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$	20	15 085.308	6627.155	$4p[2\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
950	11 536.3445	8665.8847	$3p'[1\frac{1}{2}]_0-3d[1\frac{1}{2}]^0$	50	15 140.101	6603.171	$4p[2\frac{1}{2}]_2-5d[3\frac{1}{2}]^0$
500	11 601.5367	8617.1889	$3p[1\frac{1}{2}]_0-4s[1\frac{1}{2}]^0$	4	15 174.314	6588.283	$4p'[1\frac{1}{2}]_1-5d'[1\frac{1}{2}]^0$
1200 a	11 614.0807	8607.8817	$3p'[1\frac{1}{2}]_1-4s'[1\frac{1}{2}]^0$	15	189.727	6581.598	$4p'[1\frac{1}{2}]_1-5d'[2\frac{1}{2}]^0$
300	11 688.0018	8553.4411	$3p'[1\frac{1}{2}]_0-3d[2\frac{1}{2}]^0$	15	190.558	6581.238	$4p'[1\frac{1}{2}]_1-5d'[1\frac{1}{2}]^0$
2000 a	11 766.7924	8496.1672	$3p'[1\frac{1}{2}]_1-4s'[1\frac{1}{2}]^0$	70	15 190.928	6581.078	$4p'[1\frac{1}{2}]_2-5d'[2\frac{1}{2}]^0$
1500	11 789.0436	8480.1312	$3p'[1\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	15	191.754	6580.720	$4p'[1\frac{1}{2}]_2-5d'[2\frac{1}{2}]^0$
500	11 789.8892	8479.5230	$3p[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$	15	192.585	6580.360	$4p'[1\frac{1}{2}]_2-5d'[1\frac{1}{2}]^0$
1000	11 984.9125	8341.5411	$3p'[1\frac{1}{2}]_0-4s'[1\frac{1}{2}]^0$	800	15 230.7138	6563.8868	$3p'[1\frac{1}{2}]_0-4s'[1\frac{1}{2}]^0$
3000 a	12 066.3339	8285.2540	$3p[1\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	50	15 348.188	6513.647	$4p[1\frac{1}{2}]_1-5d[2\frac{1}{2}]^0$
800	12 459.3888	8023.8806	$3p'[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$	30	15 370.081	6504.369	$4p[1\frac{1}{2}]_1-5d[1\frac{1}{2}]^0$
20 d	12 464.080	8020.860	$3d[1\frac{1}{2}]_0-5f[1\frac{1}{2}]_1$	100	15 407.592	6488.534	$4p[1\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
70 d	12 486.721	8006.317	$3d[1\frac{1}{2}]_1-5f[1\frac{1}{2}]_1$	100	15 409.057	6487.917	$4p[1\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
70 d	12 486.727	8006.313	$3d[1\frac{1}{2}]_1-5f[1\frac{1}{2}]_2$	15	450.890	6470.351	$4p[1\frac{1}{2}]_1-5d[1\frac{1}{2}]^0$
10	12 570.923	7952.690	$3d[3\frac{1}{2}]_2-5f[3\frac{1}{2}]_4$	40	15 451.225	6470.211	$4p[1\frac{1}{2}]_2-5d[1\frac{1}{2}]^0$
10	12 570.934	7952.683	$3d[3\frac{1}{2}]_1-5f[3\frac{1}{2}]_3$	10	15 466.232	6463.933	$4p[1\frac{1}{2}]_2-5d[3\frac{1}{2}]^0$
10	12 573.727	7950.916	$3d[3\frac{1}{2}]_0-5f[3\frac{1}{2}]_4$	20	15 499.487	6450.064	$4p[1\frac{1}{2}]_2-5d[1\frac{1}{2}]^0$
5	12 573.738	7950.909	$3d[3\frac{1}{2}]_0-5f[3\frac{1}{2}]_3$	20	15 500.887	6449.481	$4p[\frac{1}{2}]_1-6s[1\frac{1}{2}]^0$
5	12 577.386	7948.603	$3d[3\frac{1}{2}]_0-5f[2\frac{1}{2}]_3$	30	15 604.203	6406.779	$4p[\frac{1}{2}]_1-6s[1\frac{1}{2}]^0$
250 d	12 584.630	7944.028	$3d[3\frac{1}{2}]_0-5f[4\frac{1}{2}]_5$	50	16 022.732	6239.428	$3d[1\frac{1}{2}]_2-4f'[2\frac{1}{2}]^0$
120 d	12 584.646	7944.018	$3d[3\frac{1}{2}]_0-5f[4\frac{1}{2}]_4$	50	16 022.763	6239.416	$3d[1\frac{1}{2}]_2-4f'[2\frac{1}{2}]^0$
180 d	12 587.456	7942.244	$3d[3\frac{1}{2}]_0-5f[4\frac{1}{2}]_4$	15	16 098.476	6210.072	$3d[1\frac{1}{2}]_1-4f'[2\frac{1}{2}]^0$
300	12 595.0042	7937.4845	$3p'[1\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	15	16 346.923	6115.689	$4p[\frac{1}{2}]_0-5d[1\frac{1}{2}]^0$
100 d	12 604.197	7931.695	$3d[1\frac{1}{2}]_2-5f[2\frac{1}{2}]_2$	20	16 347.729	6115.387	$3d[\frac{1}{2}]^0-5p[1\frac{1}{2}]_1$
100 d	12 604.208	7931.688	$3d[1\frac{1}{2}]_2-5f[2\frac{1}{2}]_3$	80	16 405.242	6093.948	$4p[2\frac{1}{2}]_3-6s[1\frac{1}{2}]^0$
20	12 617.663	7923.230	$3d[1\frac{1}{2}]_2-5f[1\frac{1}{2}]_1$	12	16 468.982	6070.362	$4p'[1\frac{1}{2}]_1-6s[\frac{1}{2}]^0$
120 d	12 640.223	7909.089	$3d'[2\frac{1}{2}]_2-5f'[2\frac{1}{2}]_2$	20	16 528.079	6048.658	$4p'[1\frac{1}{2}]_1-6s[\frac{1}{2}]^0$
120 d	12 640.255	7909.069	$3d'[2\frac{1}{2}]_2-5f'[2\frac{1}{2}]_3$	15	16 607.009	6019.909	$4p'[1\frac{1}{2}]_1-6s[\frac{1}{2}]^0$
150 d	12 642.654	7907.568	$3d'[2\frac{1}{2}]_3-5f'[2\frac{1}{2}]_2$	60	16 609.433	6019.031	$4p'[1\frac{1}{2}]_2-6s[\frac{1}{2}]^0$
60 d	12 642.686	7907.548	$3d'[2\frac{1}{2}]_3-5f'[2\frac{1}{2}]_3$	25	16 634.054	6010.122	$4p'[1\frac{1}{2}]_0-5d'[1\frac{1}{2}]^0$
60 d	12 651.021	7902.338	$3d[1\frac{1}{2}]^0-5f[2\frac{1}{2}]_2$	40	16 788.711	5954.732	$4p[1\frac{1}{2}]_1-6s[1\frac{1}{2}]^0$
120 d	12 658.355	7897.760	$3d'[1\frac{1}{2}]_2-5f'[2\frac{1}{2}]_2$	15	16 789.110	5954.615	$3d[1\frac{1}{2}]^0-5p[2\frac{1}{2}]_2$
120 d	12 658.387	7897.740	$3d'[1\frac{1}{2}]_2-5f'[2\frac{1}{2}]_3$	15	16 834.25	5938.65	$3d[1\frac{1}{2}]_2-5p[\frac{1}{2}]_1$
25	12 664.587	7893.874	$3d[1\frac{1}{2}]^0-5f[1\frac{1}{2}]_1$	20	16 861.640	5929.002	$4p[1\frac{1}{2}]_2-6s[\frac{1}{2}]^0$
12	664.594	7893.870	$3d[1\frac{1}{2}]^0-5f[1\frac{1}{2}]_2$	10	17 144.601	5831.148	$4p'[1\frac{1}{2}]_1-5d[\frac{1}{2}]^0$
40 d	12 683.562	7882.064	$3d'[1\frac{1}{2}]^0-5f'[2\frac{1}{2}]_2$	400	17 161.9296	5825.2596	$3p[\frac{1}{2}]_0-4s[1\frac{1}{2}]^0$
1000	12 689.2008	7878.5617	$3p[1\frac{1}{2}]_0-4s[1\frac{1}{2}]^0$	17	181.727	5818.547	$4p[\frac{1}{2}]_1-5d[2\frac{1}{2}]^0$
120 d	12 746.139	7843.368	$3d[2\frac{1}{2}]_0-5f[3\frac{1}{2}]_3$	150	17 182.499	5818.286	$4p'[1\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
160 d	12 749.036	7841.585	$3d[2\frac{1}{2}]_0-5f[3\frac{1}{2}]_4$	4	17 184.321	5817.669	$4p'[1\frac{1}{2}]_2-5d[2\frac{1}{2}]^0$
160 d	12 749.047	7841.578	$3d[2\frac{1}{2}]_0-5f[3\frac{1}{2}]_5$	15	17 198.578	5812.846	$3d[1\frac{1}{2}]^0-5p[\frac{1}{2}]_1$
10 d	12 752.761	7839.295	$3d[2\frac{1}{2}]_2-5f[2\frac{1}{2}]_2$	400 c	18 029.657	5544.903	$3d[\frac{1}{2}]^0-4f[2\frac{1}{2}]_2$
10 d	12 752.772	7839.288	$3d[2\frac{1}{2}]_2-5f[2\frac{1}{2}]_3$	400 c	18 035.800	5543.015	$3d[\frac{1}{2}]^0-4f[\frac{1}{2}]_1$
13 d	12 755.673	7837.505	$3d[2\frac{1}{2}]_3-5f[2\frac{1}{2}]_2$	1000 c	18 083.206	5528.483	$3d[\frac{1}{2}]^0-4f[\frac{1}{2}]_2$
13 d	12 755.684	7837.498	$3d[2\frac{1}{2}]_3-5f[2\frac{1}{2}]_3$	18	083.245	5528.471	$3d[\frac{1}{2}]^0-4f[\frac{1}{2}]_1$
250	12 769.5248	7829.0034	$3p'[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$	5	18 210.3300	5489.8896	$4s[1\frac{1}{2}]^0-4p[\frac{1}{2}]_0$
?	12 887.1594	7757.5400	$3p'[1\frac{1}{2}]_1-4s[1\frac{1}{2}]^0$	350 c	18 221.100	5486.645	$3d[3\frac{1}{2}]^0-4f[3\frac{1}{2}]_4$
1100	12 912.0141	7742.6073	$3p[1\frac{1}{2}]_2-4s[1\frac{1}{2}]^0$	18	221.129	5486.636	$3d[3\frac{1}{2}]_4-4f[3\frac{1}{2}]_3$

TABLE I. Description of Ne I, region 1.1–4.0  $\mu\text{m}$ —Continued

Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification	Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification
250 <sup>c</sup>	{ 18 226.992 18 227.022	5484.871 5484.862	3d[3½]³-4f[3½]₄ 3d[3½]³-4f[3½]₃	50	{ 24 086.960 24 092.388	4150.492 4149.556	4p'[1½]₁-4d'[1½]² 4p[2½]₂-4d[2½]³
10	18 247.463	5478.718	3d[3½]³-4f[2½]₃	200 <sup>b</sup>	24 098.544	4148.496	4p[2½]₂-4d[2½]²
6	{ 18 253.330 18 253.373	5476.957 5476.944	3d[3½]³-4f[2½]₂	15	{ 24 149.887 24 155.956	4139.676 4138.636	4p'[1½]₁-4d'[1½]² 4p[½]₁-5s'[½]²
2500 <sup>c</sup>	{ 18 276.676 18 276.696	5469.961 5469.955	3d[3½]³-4f[4½]₅	500 <sup>b</sup>	24 161.420	4137.700	4p'[1½]₁-4d'[2½]₂
2000 <sup>c</sup>	18 282.625	5468.181	3d[3½]³-4f[4½]₄	600 <sup>b</sup>	24 218.930	4127.875	4p[2½]₂-4p'[1½]₁
1200 <sup>c</sup>	{ 18 303.930 18 303.974	5461.816 5461.803	3d[1½]₂-4f[2½]₂	1500 <sup>b</sup>	24 249.638	4122.6479	4s'[½]²-4p'[1½]₁
250 <sup>c</sup>	{ 18 359.124 18 359.164	5445.396 5445.384	3d[1½]₂-4f[1½]₂	800	24 365.048	4103.1203	4s[1½]¹-4p[2½]₂
1200 <sup>c</sup>	{ 18 383.972 18 384.013	5438.036 5438.024	3d'[2½]₂-4f'[2½]₂	90	24 371.599	4102.017	4p[2½]₂-4d[3½]³
2000 <sup>c</sup>	{ 18 384.848 18 389.114	5437.777 5436.515	3d'[2½]₂-4f'[3½]₃	400 <sup>b</sup>	24 447.850	4100.038	4p'[1½]₁-4d'[1½]²
1000 <sup>c</sup>	{ 18 389.155 18 389.943	5436.503 5436.270	3d'[2½]₃-4f'[2½]₃	14	24 453.102	4088.345	4p'[1½]₂-4d'[1½]₂
2000 <sup>c</sup>	{ 18 389.943 18 389.990	5436.256 5432.460	3d'[2½]₃-4f'[3½]₃	700 <sup>b</sup>	{ 24 459.366 24 459.670	4087.298	4p'[1½]₂-4d'[2½]₃
1200 <sup>c</sup>	{ 18 402.844 18 422.350	5426.707 5426.695	3d'[1½]₂-4f'[2½]₂	25	24 464.927	4087.247	4p'[1½]₁-4d'[2½]₂
300 <sup>c</sup>	{ 18 422.391 18 423.230	5426.448 5426.448	3d'[1½]₂-4f'[3½]₃	10	24 525.791	4086.369	4p'[1½]₂-4d'[2½]₂
400 <sup>c</sup>	{ 18 458.636 18 458.677	5416.040 5416.028	3d[1½]¹-4f[1½]₂	350 <sup>b</sup>	24 776.460	4034.988	4p[1½]₁-4d[2½]₂
900 <sup>c</sup>	18 475.790	5411.011	3d'[1½]₁-4f'[2½]₂	180	24 903.732	4014.367	4p[1½]₁-4d[1½]²
1600 <sup>c</sup>	{ 18 597.703 18 597.734	5377.321 5375.540	3d[2½]₂-4f[3½]₄	550	24 928.877	4010.318	4p[1½]₂-4d[2½]₃
350 <sup>c</sup>	{ 18 597.734 18 618.917	5375.531 5369.416	3d[2½]₂-4f[3½]₃	5 <sup>b</sup>	24 935.468	4009.258	4p[1½]₂-4d[2½]₂
550 <sup>c</sup>	{ 18 618.917 18 618.962	5369.416 5369.403	3d[2½]₂-4f[2½]₂	30	24 999.792	3998.942	4p[1½]₁-4d[1½]²
20	18 625.124	5367.626	3d[2½]₃-4f[2½]₂	5	25 064.383	3988.637	4p[1½]₁-4d[1½]¹
15	18 625.169	5367.613	3d[2½]₃-4f[2½]₃	250	25 161.689	3973.212	4p[1½]₂-4d[1½]²
50 <sup>b</sup>	18 682.274	5351.206	3d[2½]₃-4f[1½]₂	10	{ 25 227.934 25 228.308	3962.779	4p[1½]₂-4d[3½]³
15	18 687.775	5279.072	4p[½]₁-4d'[1½]₂	50	25 277.246	3962.720	4p[1½]₁-4d[½]²
50 <sup>b</sup>	19 573.7692	5107.4837	4s[1½]₂-4p'[1½]₂	650	25 524.366	3955.048	4p[1½]₁-4d[½]³
170	19 577.1358	5106.6054	4s[1½]₂-4p'[½]₁	50	25 593.188	3936.990	4p[1½]₁-4d[½]¹
120 <sup>b</sup>	20 350.2378	4912.6065	4s[1½]¹-4p'[1½]₂	35	25 854.914	3916.7564	4s[1½]₂-4p[½]₁
20	20 565.1211	4861.2751	4s[1½]¹-4p'[1½]₁	150	25 999.579	3866.6818	4s'[½]¹-4p[½]₀
1200 <sup>b</sup>	21 041.2948	4751.2624	4s'[½]¹-4p'[½]₀	10	26 200.474	3847.0013	4p[½]₁-4d[1½]²
750 <sup>b</sup>	21 708.1449	4605.3090	4s[1½]¹-4p[½]₀	20	26 259.534	3849.910	4s'[½]¹-4p[½]₁
300 <sup>b</sup>	22 247.348	4493.691	4p[½]₁-4d[1½]₂	20	26 594.965	3826.620	4p[½]₁-4d[½]¹
350 <sup>b</sup>	22 428.133	4457.469	4p[½]₁-4d[½]¹	40	26 711.639	3825.207	4p[½]₀-5s'[½]¹
130 <sup>b</sup>	22 466.802	4449.797	4p[½]₁-4d[½]₀	80	26 859.980	3813.7871	4s'[½]₀-4p[½]₁
2250 <sup>b</sup>	22 530.404	4437.2359	4s[1½]₂-4p[1½]₂	450	26 173.094	3803.669	4p[2½]₂-5s[½]¹
400 <sup>b</sup>	22 661.813	4411.5057	4s[1½]₂-4p[1½]₁	30	26 332.683	3799.240	4p'[1½]₁-5s'[½]¹
50	22 687.775	4406.458	4p[½]₀-4d'[1½]₁	1300	26 912.099	3797.472	4p[2½]₂-5s[½]²
600 <sup>b</sup>	23 100.514	4327.7271	4s'[½]₀-4p'[½]₁	40	27 352.352	3798.2520	4s'[½]¹-4p[½]₁
1000 <sup>b</sup>	23 260.302	4297.9975	4s[1½]₂-4p[2½]₂	2200	27 511.327	3794.065	4p'[1½]₁-5s'[½]₀
1050 <sup>b</sup>	23 372.999	4277.2740	4s'[1½]₂-4p'[1½]₁	600	27 849.801	3794.787	4p'[½]₁-5s'[½]¹
850 <sup>b</sup>	23 565.362	4242.3587	4s[1½]¹-4p[1½]₂	100	27 912.099	3794.787	4p'[½]₁-5s'[½]¹
3500 <sup>b</sup>	23 636.515	4229.5880	4s[1½]₂-4p[2½]₃	40	27 947.909	3794.787	4p'[1½]₂-5s'[½]²
300	23 701.643	4217.966	4p[2½]₃-4d[2½]₃	80	27 997.062	3794.787	4p[2½]₂-5s[½]₂
1100 <sup>b</sup>	{ 23 707.601 23 709.160	4216.906 4216.6285	4p[2½]₃-4d[2½]₂	20	28 057.304	3794.787	4p[1½]₁-5s[½]₂
1800 <sup>b</sup>	23 951.417	4173.9793	4s'[½]₀-4p'[1½]₂	120	28 131.310	3794.787	4p[1½]₂-5s[½]₂
600 <sup>b</sup>	23 956.458	4173.1010	4s'[½]₀-4p'[½]₁	18	28 471.442	3794.787	4p[1½]₁-5s[½]₀
10	23 971.820	4170.427	4p[2½]₃-4d[3½]₃	20	28 780.010	3794.787	4p[1½]₂-5s[½]₀
1000	23 978.122	4169.331	4p[2½]₃-4d[3½]₄	15	28 834.784	3794.787	4p[1½]₂-5s[½]₀
				30	28 834.784	3794.787	4p[1½]₂-5s[½]₀
				30	28 736.035	3794.787	4p[1½]₂-5s[½]₀

<sup>a</sup> Observed interferometrically. <sup>b</sup> Reported also by Hepner. <sup>c</sup> Most recent observation by Johansson. <sup>d</sup> Most recent observation by Litzén.

TABLE 2. Description of Ar I, region 1.2–4.0  $\mu\text{m}$ 

Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification	Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification
80	12 026.648	8312.594	$4p[\frac{1}{2}]_0-3d'[\frac{1}{2}]^o$	1500 a, b	13 599.333	7351.292	$4p'[\frac{1}{2}]_2-3d'[\frac{1}{2}]^o$
1300 a, b	12 112.324	8253.795	$4p[\frac{1}{2}]_3-3d[\frac{1}{2}]^o$	7500 a, b	13 622.659	7338.704	$4p[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$
700	12 139.737	8235.157	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$	6	13 652.795	7322.505	$3d[\frac{1}{2}]^o-6p[\frac{1}{2}]_0$
80	12 151.369	8227.274	$3d[\frac{1}{2}]^o-4f'[\frac{3}{2}]^o$	8	{13 658.41}	7319.493	$3d'[\frac{1}{2}]^o-5f[\frac{1}{2}]_2$
30	12 234.397	8171.440	$3d[\frac{3}{2}]_2-6p[\frac{1}{2}]_2$	5000 a, b	{13 658.78}	7319.293	$3d'[\frac{1}{2}]^o-5f[\frac{1}{2}]_1$
900 a, b	12 343.392	8099.285	$4p[\frac{1}{2}]_3-3d[\frac{1}{2}]^o$	10000 a, b	13 678.549	7308.718	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$
450 a, b	12 356.296	8090.826	$3d[\frac{1}{2}]^o-4f'[\frac{3}{2}]_1$	300	13 718.577	7287.393	$4p[\frac{1}{2}]_3-3d[\frac{3}{2}]^o$
22	12 377.194	8077.166	$4p'[\frac{1}{2}]_2-3d'[\frac{1}{2}]^o$	200 b	{13 825.717}	7230.921	$4p[\frac{1}{2}]_2-5s[\frac{1}{2}]^o$
2000	12 402.828	8060.472	$4p[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$	1200 a, b	{13 828.321}	7229.559	$3d[\frac{3}{2}]_3-4f[\frac{3}{2}]_4$
150	12 419.414	8049.707	$3d[\frac{1}{2}]^o-4f'[\frac{1}{2}]_2$	30 b	{13 828.394}	7229.521	$3d[\frac{3}{2}]_3-4f[\frac{3}{2}]_3$
60	12 420.030	8049.308	$3d[\frac{1}{2}]_2-4f'[\frac{1}{2}]_1$	50 b	{13 866.396}	7209.708	$3d[\frac{3}{2}]_3-4f'[\frac{3}{2}]_2$
5000 a, b	12 439.321	8036.825	$4p[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$	100 b	{13 866.998}	7209.395	$3d[\frac{3}{2}]_3-4f'[\frac{1}{2}]_3$
2000 b	12 456.114	8025.990	$4p[\frac{1}{2}]_2-5s[\frac{1}{2}]^o$	1200 a, b	13 907.476	7188.412	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$
2500 a, b	12 487.663	8005.713	$4p[\frac{1}{2}]_3-5s[\frac{1}{2}]^o$	30 b	13 910.556	7186.820	$3d[\frac{3}{2}]_3-4f[\frac{4}{2}]_4$
75	12 554.324	7963.204	$4p[\frac{1}{2}]_3-3d[\frac{1}{2}]^o$	2	13 992.808	7144.575	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$
40 b	{12 596.209}	7936.725	$3d[\frac{1}{2}]^o-4f'[\frac{3}{2}]_1$	2000 b	14 061.991	7109.425	$3d[\frac{1}{2}]^o-6p[\frac{1}{2}]_1$
	{12 596.276}	7936.683	$3d[\frac{1}{2}]^o-4f'[\frac{3}{2}]_3$	1	14 093.640	7093.460	$4p'[\frac{1}{2}]_0-3d[\frac{1}{2}]^o$
90 b	12 621.619	7920.747	$4p[\frac{1}{2}]_0-5s'[\frac{1}{2}]^o$	8	14 174.712	7052.889	$3d[\frac{1}{2}]^o-5p'[\frac{1}{2}]_0$
25 b	12 638.480	7910.180	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$	120 b	14 192.472	7044.063	$3d[\frac{1}{2}]^o-6p[\frac{1}{2}]_2$
20	{12 649.96}	7903.004	$3d'[\frac{1}{2}]_2-5f[\frac{2}{2}]_2$	450	14 249.193	7016.023	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$
	{12 650.34}	7902.764	$3d'[\frac{1}{2}]_2-5f[\frac{2}{2}]_3$	14 254.140	7013.588	$3d'[\frac{1}{2}]_2-4f'[\frac{2}{2}]_2$	
15	12 661.31	7895.920	$3d'[\frac{1}{2}]_3-5f[\frac{3}{2}]_3$	450 b	14 254.844	7013.242	$3d'[\frac{1}{2}]_3-4f'[\frac{2}{2}]_3$
9	{12 684.91}	7881.224	$3d'[\frac{1}{2}]_2-5f[\frac{1}{2}]_2$	14 256.868 a, b	14 266.866	7012.246	$3d'[\frac{1}{2}]_2-4f'[\frac{3}{2}]_3$
8	{12 685.24}	7881.024	$3d'[\frac{1}{2}]_2-5f[\frac{1}{2}]_1$	4	14 444.828	6921.001	$3d[\frac{1}{2}]^o-6p[\frac{1}{2}]_1$
	12 697.00	7873.720	$3d'[\frac{1}{2}]_3-5f[\frac{4}{2}]_4$	15	14 577.458	6858.032	$4p'[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$
1250 b	12 702.280	7870.449	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$	300	{14 595.733}	6849.445	$3d'[\frac{1}{2}]_2-4f'[\frac{2}{2}]_2$
600 b	12 733.418	7851.203	$4p[\frac{1}{2}]_2-5s[\frac{1}{2}]^o$	300 b	{14 596.471 a, b}	6849.099	$3d'[\frac{1}{2}]_3-4f'[\frac{2}{2}]_3$
400 b	12 746.232	7843.310	$4p'[\frac{1}{2}]_1-5s'[\frac{1}{2}]^o$	500	{14 634.414 a, b}	6831.341	$3d'[\frac{1}{2}]_3-4f'[\frac{3}{2}]_4$
12	12 765.954	7831.193	$3d[\frac{1}{2}]_2-6p[\frac{1}{2}]_1$	450 b	{14 634.504}	6831.299	$3d'[\frac{1}{2}]_3-4f'[\frac{3}{2}]_3$
2500 a, b	12 802.737	7808.694	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$	300 b	14 650.346	6823.912	$3d[\frac{1}{2}]_2-4f[\frac{3}{2}]_3$
15	12 813.484	7802.144	$5s[\frac{1}{2}]_2-6p[\frac{1}{2}]_2$	2	14 666.866	6816.226	$5p[\frac{1}{2}]_2-5d'[\frac{1}{2}]^o$
12	12 873.400	7765.831	$3d[\frac{1}{2}]_2-6p[\frac{1}{2}]_2$	2	14 682.028	6809.187	$5p[\frac{1}{2}]_2-6d[\frac{1}{2}]^o$
9	12 903.024	7748.002	$5s'[\frac{1}{2}]_0-6p'[\frac{1}{2}]_1$	2	14 684.646	6807.973	$5p[\frac{1}{2}]_1-6d[\frac{1}{2}]^o$
750 b	12 933.196	7729.926	$4p'[\frac{1}{2}]_1-5s'[\frac{1}{2}]_0$	90 b	14 693.007	6804.099	$3d[\frac{1}{2}]_2-4f'[\frac{2}{2}]_2$
4000 b	12 956.658	7715.929	$4p[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$	6	14 693.683	6803.786	$3d[\frac{1}{2}]_2-4f'[\frac{2}{2}]_3$
25	12 988.701	7696.894	$5s[\frac{1}{2}]_2-6p[\frac{1}{2}]_3$	75 b	14 719.546	6791.832	$3d[\frac{1}{2}]_0-5p'[\frac{1}{2}]_1$
2500 b	13 008.264	7685.319	$4p'[\frac{1}{2}]_2-5s'[\frac{1}{2}]^o$	8	{14 739.139}	6782.803	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$
90 b	13 028.425	7673.426	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$	75 b	{14 783.025 d}	6762.667	$3d[\frac{1}{2}]_2-4f'[\frac{1}{2}]_2$
12	13 051.206	7660.032	$5s'[\frac{1}{2}]_1-6p'[\frac{1}{2}]_2$	8	{14 783.897 d}	6762.268	$3d[\frac{1}{2}]_2-4f'[\frac{1}{2}]_1$
2	13 107.116	7627.357	$5s[\frac{1}{2}]_1-6p[\frac{1}{2}]_2$	40	{14 785.380 d}	6761.590	$5s[\frac{1}{2}]_2-4f'[\frac{2}{2}]_2$
1	13 109.113	7626.195	$5s'[\frac{1}{2}]_1-6p'[\frac{1}{2}]_1$	40	{14 786.064 d, b}	6761.277	$5s[\frac{1}{2}]_2-4f'[\frac{2}{2}]_3$
6	13 130.287	7613.897	$5s[\frac{1}{2}]_0-6p[\frac{1}{2}]_1$	5	14 833.480	6739.664	$3d[\frac{1}{2}]_0-5p'[\frac{1}{2}]_1$
4	13 153.847	7600.260	$5s[\frac{1}{2}]_0-6p[\frac{1}{2}]_1$	11 b	{14 876.537 d}	6720.158	$5s[\frac{1}{2}]_2-4f'[\frac{1}{2}]_2$
3000 b	13 213.991	7565.997	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$	30	{14 877.420 d}	6719.759	$5s[\frac{1}{2}]_2-4f'[\frac{1}{2}]_1$
2500 a, b	13 228.104	7557.595	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$	2	14 898.136	6710.415	$5p[\frac{1}{2}]_3-6d[\frac{3}{2}]^o$
1200	13 230.897	7556.000	$4p[\frac{1}{2}]_1-5s'[\frac{1}{2}]^o$	6	14 966.953	6679.561	$5s'[\frac{1}{2}]^o-4f'[\frac{2}{2}]_2$
12	13 243.981	7548.535	$5s[\frac{1}{2}]_2-6p[\frac{1}{2}]_2$	3	14 974.568	6676.164	$4p[\frac{1}{2}]_3-3d[\frac{1}{2}]_2$
6000 a, b	13 272.635	7532.239	$4p'[\frac{1}{2}]_2-3d'[\frac{1}{2}]^o$	700 b	{15 030.513}	6651.315	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]^o$
225	13 302.312	7515.435	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$	150 b	{15 031.174}	6651.023	$3d[\frac{1}{2}]^o-5p'[\frac{1}{2}]_2$
5500 a, b	13 313.209	7509.283	$4p'[\frac{1}{2}]_1-3d'[\frac{1}{2}]^o$	700 b	{15 046.503}	6644.247	$4p'[\frac{1}{2}]_0-3d'[\frac{1}{2}]^o$
20	13 317.528	7506.848	$3d[\frac{1}{2}]^o-4f'[\frac{2}{2}]_2$	12	{15 052.567}	6641.570	$3d[\frac{1}{2}]^o-5p'[\frac{1}{2}]_1$
25 b	{13 330.112}	7499.761	$3d[\frac{3}{2}]_4-4f'[\frac{3}{2}]_4$	300 b	{15 172.691}	6588.988	$4p[\frac{1}{2}]_0-5s[\frac{1}{2}]^o$
	{13 330.180}	7499.723	$3d[\frac{3}{2}]_4-4f'[\frac{3}{2}]_3$	4	15 177.724	6586.803	$5s[\frac{1}{2}]^o-4f'[\frac{2}{2}]_2$
8500 a, b	13 367.110	7479.003	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$	3	{15 273.799}	6545.371	$5s[\frac{1}{2}]^o-4f'[\frac{1}{2}]_2$
	13 367.827 e	7478.602	$4p'[\frac{1}{2}]_1-5s'[\frac{1}{2}]^o$	3	{15 274.730}	6544.972	$5s[\frac{1}{2}]^o-4f'[\frac{1}{2}]_1$
2500 a, b	{13 406.513}	7457.022	$3d[\frac{3}{2}]_4-4f'[\frac{4}{2}]_4$	500 b	{15 301.881 a}	6533.359	$3d[\frac{2}{2}]_3-4f'[\frac{3}{2}]_4$
	{13 406.586}	7456.981	$3d[\frac{3}{2}]_4-4f'[\frac{4}{2}]_5$	{15 301.970}	6533.321	$3d[\frac{2}{2}]_3-4f'[\frac{3}{2}]_3$	
2	13 421.289	7448.812	$3d[\frac{2}{2}]_3-5s[\frac{1}{2}]^o$	150 b	15 329.344	6521.654	$4p[\frac{2}{2}]_3-3d[\frac{1}{2}]^o$
1200	13 499.406	7405.708	$4p[\frac{1}{2}]_2-5s[\frac{1}{2}]^o$	120	{15 348.516}	6513.508	$3d[\frac{2}{2}]_3-4f'[\frac{2}{2}]_2$
9500 a, b	13 504.190	7403.085	$4p[\frac{1}{2}]_2-3d[\frac{1}{2}]^o$	60 b	{15 349.253 b}	6513.195	$3d[\frac{2}{2}]_3-4f'[\frac{2}{2}]_3$
500	13 544.205	7381.213	$4p[\frac{1}{2}]_1-5s[\frac{1}{2}]^o$	120 b	15 353.128	6511.551	$4p'[\frac{1}{2}]_1-5s[\frac{1}{2}]^o$
750	13 573.618	7365.218	$4p'[\frac{1}{2}]_1-5s[\frac{1}{2}]^o$	120 b	15 402.640	6490.620	$3d[\frac{2}{2}]_3-4f'[\frac{4}{2}]_4$

TABLE 2. Description of Ar I, region 1.2–4.0  $\mu\text{m}$ —Continued

Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification	Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification
10	15 446.772	6472.076	$3d[2\frac{1}{2}]_3-4f[1\frac{1}{2}]_2$	4	19 992.232	5000.578	$5s[1\frac{1}{2}]_2-5p'[1\frac{1}{2}]_2$
6	15 555.460	6426.855	$4p'[1\frac{1}{2}]_2-3d[2\frac{1}{2}]_3$	60	20 025.672	4992.227	$3d[1\frac{1}{2}]_3-5p[\frac{1}{2}]_1$
8	15 734.909	6353.560	$4p'[1\frac{1}{2}]_2-5s[1\frac{1}{2}]_1$	30	20 030.097	4991.125	$5s[1\frac{1}{2}]_2-5p'[\frac{1}{2}]_1$
2	15 776.614	6336.764	$4p'[1\frac{1}{2}]_1-5s[1\frac{1}{2}]_2$	25 <sup>c</sup>	20 068.932	4981.466	$3d[2\frac{1}{2}]_2-5p'[1\frac{1}{2}]_1$
5	15 793.157	6330.127	$3d[1\frac{1}{2}]_2-5p'[1\frac{1}{2}]_2$	160	20 317.011	4920.641	$4p'[\frac{1}{2}]_0-5s[1\frac{1}{2}]_1$
18	15 816.777	6320.674	$3d[1\frac{1}{2}]_2-5p'[1\frac{1}{2}]_1$	1	20 556.994	4863.197	$3d'[\frac{1}{2}]_1-4f[2\frac{1}{2}]_2$
40	15 883.164	6294.255	$4p'[1\frac{1}{2}]_1-3d[2\frac{1}{2}]_2$	75 <sup>c</sup>	20 568.816	4860.402	$3d[1\frac{1}{2}]_2-5p[2\frac{1}{2}]_1$
240 <sup>a, b</sup>	15 899.687	6287.714	$3d'[1\frac{1}{2}]_1-4f'[2\frac{1}{2}]_2$	2500	20 616.229	4849.224	$4p'[1\frac{1}{2}]_2-3d[1\frac{1}{2}]_2$
2	15 913.799	6282.138	$5s[\frac{1}{2}]_1-6p[1\frac{1}{2}]_1$	150	20 647.135	4841.965	$3d[\frac{1}{2}]_1-5p[\frac{1}{2}]_1$
400 <sup>b</sup>	15 989.491	6252.400	$4p'[\frac{1}{2}]_0-5s'[\frac{1}{2}]_1$	25	20 682.40	4833.710	$4d[3\frac{1}{2}]_3-6f[4\frac{1}{2}]_4$
12	16 122.656	6200.758	$4p[2\frac{1}{2}]_2-3d[\frac{1}{2}]_1$	22	20 716.338	4825.791	$5s[\frac{1}{2}]_1-5p'[1\frac{1}{2}]_2$
90 <sup>b</sup>	16 180.023	6178.773	$4p'[1\frac{1}{2}]_2-5s[1\frac{1}{2}]_2$	120	{20 733.634	4821.765	$3d'[\frac{1}{2}]_1-4f[1\frac{1}{2}]_2$
16	16 264.070	6146.843	$4p'[\frac{1}{2}]_1-5s[1\frac{1}{2}]_1$	110	{20 735.350	4821.366	$3d'[\frac{1}{2}]_1-4f[1\frac{1}{2}]_1$
400 <sup>a, b</sup>	16 436.575	6082.331	$3d[1\frac{1}{2}]_1-4f[2\frac{1}{2}]_2$	110	20 811.042	4803.830	$3d[1\frac{1}{2}]_2-5p[2\frac{1}{2}]_3$
500 <sup>b</sup>	16 519.867	6051.664	$4p[1\frac{1}{2}]_2-3d[1\frac{1}{2}]_2$	1200	20 984.286 <sup>e</sup>	4764.170	$5s[\frac{1}{2}]_1-5p'[\frac{1}{2}]_1$
250	{16 549.306 <sup>b, d</sup>	6040.899	$3d[1\frac{1}{2}]_1-4f[1\frac{1}{2}]_2$	44	21 035.834	4763.756	$4p[\frac{1}{2}]_0-3d[\frac{1}{2}]_1$
	{16 550.400 <sup>d</sup>	6040.500	$3d[1\frac{1}{2}]_1-4f[1\frac{1}{2}]_1$	120	21 332.885	4752.496	$3d[2\frac{1}{2}]_3-5p[\frac{1}{2}]_2$
300 <sup>b</sup>	16 740.078	5972.056	$4p'[\frac{1}{2}]_1-5s[1\frac{1}{2}]_2$	750	21 534.207	4642.507	$4p'[\frac{1}{2}]_1-3d[1\frac{1}{2}]_2$
14	16 860.088	5929.547	$4p'[\frac{1}{2}]_1-3d[2\frac{1}{2}]_2$	15	21 669.70	4613.480	$4d[2\frac{1}{2}]_2-6f[3\frac{1}{2}]_3$
5000 <sup>a, b</sup>	16 940.584	5901.372	$4p[1\frac{1}{2}]_2-3d[1\frac{1}{2}]_2$	15	22 039.561	4536.057	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]_2$
8	17 296.104	5780.070	$5p[\frac{1}{2}]_1-7s[1\frac{1}{2}]_2$	15	22 077.181	4528.328	$4p'[\frac{1}{2}]_2-3d[\frac{1}{2}]_2$
22	17 401.908	5744.927	$3d[\frac{1}{2}]_1-5p[\frac{1}{2}]_0$	15	22 112.626	4521.069	$3d[1\frac{1}{2}]_2-5p[\frac{1}{2}]_1$
300	{17 444.903	5730.768	$4p[1\frac{1}{2}]_1-3d[\frac{1}{2}]_1$	15	22 209.669	4501.315	$5p[\frac{1}{2}]_1-6s'[\frac{1}{2}]_1$
	{17 445.248	5730.655	$4p'[1\frac{1}{2}]_2-3d[\frac{1}{2}]_3$	15	22 254.14	4492.320	$4d[2\frac{1}{2}]_3-6f[3\frac{1}{2}]_4$
150	17 823.991	5608.884	$3d'[2\frac{1}{2}]_2-4f[3\frac{1}{2}]_3$	5	22 533.597	4436.607	$5p[\frac{1}{2}]_1-6s'[\frac{1}{2}]_0$
35	{17 887.176 <sup>d</sup>	5589.071	$3d'[2\frac{1}{2}]_2-4f[2\frac{1}{2}]_2$	30	22 938.69	4358.258	$6s[\frac{1}{2}]_2-6f[\frac{1}{2}]_2$
	{17 888.178 <sup>d</sup>	5588.7583	$3d'[2\frac{1}{2}]_2-4f[2\frac{1}{2}]_3$	8	22 939.64	4358.078	$6s[\frac{1}{2}]_2-6f[\frac{1}{2}]_1$
1500 <sup>b</sup>	{17 914.629	5580.506	$4p[1\frac{1}{2}]_1-3d[\frac{1}{2}]_0$	6	23 133.204	4354.921	$4d[1\frac{1}{2}]_2-7p[\frac{1}{2}]_2$
	{17 914.726	5580.476	$4p[1\frac{1}{2}]_2-3d[\frac{1}{2}]_1$	100 <sup>c</sup>	23 845.035	4192.601	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]_1$
6	{18 020.765	5547.639	$3d'[2\frac{1}{2}]_2-4f[1\frac{1}{2}]_2$	40	23 904.766	4182.125	$3d[3\frac{1}{2}]_4-5p[2\frac{1}{2}]_3$
18	{18 022.061	5547.240	$3d'[2\frac{1}{2}]_2-4f[1\frac{1}{2}]_1$	1000	23 956.264	4173.966	$4d[\frac{1}{2}]_1-5f[\frac{1}{2}]_1$
18	18 185.749	5497.310	$5p[1\frac{1}{2}]_3-7s[1\frac{1}{2}]_2$	2000	24 013.230	4171.349	$4p'[\frac{1}{2}]_1-3d[\frac{1}{2}]_0$
15	18 231.349	5483.560	$3d[\frac{1}{2}]_0-5p[\frac{1}{2}]_1$	20	24 776.62	4163.235	$3d[3\frac{1}{2}]_3-5p[1\frac{1}{2}]_2$
14	18 348.006	5448.696	$3d[3\frac{1}{2}]_3-5p'[\frac{1}{2}]_2$	12	24 777.85	4034.962	$4d[\frac{1}{2}]_1-5f[\frac{1}{2}]_2$
9	18 361.332	5444.741	$3d'[1\frac{1}{2}]_2-4f[3\frac{1}{2}]_3$	900	24 778.971	4034.762	$4d[\frac{1}{2}]_1-5f[\frac{1}{2}]_1$
90	{18 418.047	5427.975	$3d'[2\frac{1}{2}]_3-4f[3\frac{1}{2}]_4$	15	25 125.271	3978.971	$3d[3\frac{1}{2}]_3-5p[2\frac{1}{2}]_2$
	{18 418.176	5427.937	$3d'[2\frac{1}{2}]_3-4f[3\frac{1}{2}]_3$	12	25 487.646	3922.399	$3d[3\frac{1}{2}]_3-5p[2\frac{1}{2}]_3$
120	18 427.765	5425.113	$4p'[\frac{1}{2}]_0-3d[1\frac{1}{2}]_1$	1200	25 505.228	3919.695	$5s[\frac{1}{2}]_1-5p[\frac{1}{2}]_0$
200	{18 428.392	5424.928	$3d'[1\frac{1}{2}]_2-4f[2\frac{1}{2}]_2$	900	25 661.022	3895.898	$5s'[\frac{1}{2}]_1-5p'[\frac{1}{2}]_0$
	{18 429.455	5424.615	$3d'[1\frac{1}{2}]_2-4f[2\frac{1}{2}]_3$	120	26 218.593 <sup>f</sup>	3813.047	$5p'[\frac{1}{2}]_2-5d[2\frac{1}{2}]_2$
18	18 485.653	5408.124	$3d'[2\frac{1}{2}]_3-4f[2\frac{1}{2}]_2$	400 <sup>c</sup>	26 234.637	3810.715	$5p[2\frac{1}{2}]_3-4d'[2\frac{1}{2}]_3$
22	18 485.663	5408.121	$5p'[\frac{1}{2}]_1-5d'[\frac{1}{2}]_1$	450 <sup>c</sup>	26 543.041	3766.438	$3d'[2\frac{1}{2}]_2-5p[\frac{1}{2}]_1$
18	18 486.723	5407.811	$3d'[2\frac{1}{2}]_3-4f[2\frac{1}{2}]_3$	50	26 605.288	3757.626	$3d[2\frac{1}{2}]_2-5p[\frac{1}{2}]_0$
26	18 564.219	5385.236	$3d'[2\frac{1}{2}]_3-4f[4\frac{1}{2}]_4$	30	26 835.705	3725.362	$3d[2\frac{1}{2}]_2-5p[\frac{1}{2}]_1$
24	{18 570.219	5383.496	$3d'[1\frac{1}{2}]_2-4f[1\frac{1}{2}]_2$	200	26 909.711	3715.117	$5s[\frac{1}{2}]_2-5p[\frac{1}{2}]_2$
	{18 571.596	5383.097	$3d'[1\frac{1}{2}]_2-4f[1\frac{1}{2}]_1$	75	27 145.454	3682.853	$5s[\frac{1}{2}]_2-5p[\frac{1}{2}]_1$
60	18 632.289	5365.562	$3d[\frac{1}{2}]_1-5p[\frac{1}{2}]_2$	200	27 225.60	3672.012	$4d[\frac{1}{2}]_1-5f[\frac{1}{2}]_{4,5}$
40	18 745.005	5333.298	$3d[\frac{1}{2}]_1-5p[\frac{1}{2}]_1$	1000	27 285.760	3663.916	$3d'[\frac{1}{2}]_2-5p'[\frac{1}{2}]_2$
6	19 024.142	5255.044	$5s'[\frac{1}{2}]_1-4f[2\frac{1}{2}]_2$	100	27 356.342	3654.462	$3d'[\frac{1}{2}]_2-5p'[\frac{1}{2}]_1$
5	19 123.807	5227.657	$5s[\frac{1}{2}]_1-5p'[\frac{1}{2}]_0$	50	27 785.928	3597.962	$5s'[\frac{1}{2}]_1-5p'[\frac{1}{2}]_1$
12	{19 175.325	5213.612	$5s'[\frac{1}{2}]_1-4f[1\frac{1}{2}]_2$	30	27 934.995	3578.763	$6d[\frac{1}{2}]_2-4f[2\frac{1}{2}]_3$
	{19 175.332	5213.610	$4d[2\frac{1}{2}]_2-7p[1\frac{1}{2}]_2$	150	27 937.439	3578.450	$6d[\frac{1}{2}]_2-4f[2\frac{1}{2}]_2$
25	19 294.916	5181.298	$5s'[\frac{1}{2}]_1-4f[1\frac{1}{2}]_1$	75	28 194.726	3545.795	$5s'[\frac{1}{2}]_1-5p'[\frac{1}{2}]_1$
30	19 426.959	5146.081	$5p[1\frac{1}{2}]_2-5d[2\frac{1}{2}]_3$	15	28 238.250	3540.330	$5s[\frac{1}{2}]_1-5p[\frac{1}{2}]_2$
2	{19 469.90	5134.732	$4d[1\frac{1}{2}]_2-6f[1\frac{1}{2}]_2$	150	28 268.69	3536.518	$4d'[\frac{1}{2}]_2-5f'[2\frac{1}{2}]_2$
	{19 470.58	5134.552	$4d[1\frac{1}{2}]_2-6f[1\frac{1}{2}]_1$	11	28 269.97	3536.358	$4d'[\frac{1}{2}]_2-5f'[2\frac{1}{2}]_3$
5	19 628.228	5093.313	$5p[2\frac{1}{2}]_3-5d[3\frac{1}{2}]_4$	300	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[3\frac{1}{2}]_3$
550	19 817.508	5044.666	$3d[1\frac{1}{2}]_2-5p[\frac{1}{2}]_2$	12	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[3\frac{1}{2}]_2$
1	19 860.943	5033.634	$3d[2\frac{1}{2}]_2-5p'[\frac{1}{2}]_1$	400	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[3\frac{1}{2}]_1$
12	19 903.18	5022.952	$4d[3\frac{1}{2}]_4-6f[4\frac{1}{2}]_{4,5}$	12	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[2\frac{1}{2}]_3$
25	19 945.068	5012.402	$3d[1\frac{1}{2}]_2-5p[\frac{1}{2}]_1$	6	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[2\frac{1}{2}]_2$
160	19 965.730	5007.215	$4p'[1\frac{1}{2}]_1-3d[1\frac{1}{2}]_2$	6	28 282.36	3534.808	$4d'[\frac{1}{2}]_2-5f'[2\frac{1}{2}]_1$

TABLE 2. Description of Ar I, region 1.2–4.0 μm—Continued

Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm⁻¹)	Classification	Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm⁻¹)	Classification
300	28 314.045	3530.853	5s[1½]₀–5p[2½]₂	40	32 879.664	3040.564	3d'[1½]₁–5p'[1½]₁
12	{28 414.30	3518.395	4d'[2½]₀–5f'[2½]₂	8	32 930.634	3035.858	3d[1½]₁–5p[1½]₂
	{28 415.59	3518.235	4d'[2½]₀–5f'[2½]₃	90	33 069.750	3023.087	5p[½]₁–6s[1½]₂
45	28 427.265	3516.790	3d[2½]₀–5p[2½]₃	95	33 139.400	3016.733	5s[1½]₁–5p[½]₁
900	28 497.958	3508.066	5s[1½]₁–5p[½]₁	80	33 204.366	3003.594	3d[1½]₁–5p[1½]₁
30	28 523.13	3504.970	4d[3½]₀–5f[3½]₃,₄	8	33 591.86	2976.100	6s'[½]₁–5f[2½]₂
55	28 530.615	3504.051	3d'[1½]₀–5p[½]₀	3	33 912.38	2947.972	6s[1½]₁–5f[2½]₂
1000	28 612.427	3494.032	5s'[½]₁–5p'[½]₂	30	34 950.98	2860.370	4d[1½]₁–5f[2½]₂
300	28 690.049	3484.578	5s'[½]₁–5p'[½]₁	2	35 058.546	2851.594	3d[1½]₁–5p[2½]₂
60	28 704.95	3482.770	4d[3½]₀–5f[4½]₄	2	35 097.327	2848.443	5p[2½]₂–4d[1½]₁
2500 <sup>a</sup>	28 775.083	3474.281	5s[1½]₀–5p[2½]₃	15	{35 219.15	2838.590	4d[1½]₀–5f[1½]₂
450	28 835.223	3467.035	3d[2½]₀–5p[1½]₂		{35 221.64	2838.390	4d[1½]₀–5f[1½]₁
12	28 981.265	3449.564	5p[1½]₁–4d'[1½]₂	10	35 465.914	2818.840	4d[½]₁–6p[½]₀
40	29 100.550	3435.424	5p[1½]₂–4d'[2½]₂	25	36 210.972	2760.841	5p[2½]₂–6s[1½]₁
300	29 126.092	3432.411	5s'[½]₁–5p'[1½]₁	15	36 302.529	2753.878	5p'[1½]₁–6s'[½]₁
60	29 254.880	3417.301	5p[1½]₂–4d'[1½]₂	10	{36 380.538	2747.973	4d[1½]₂–4f'[2½]₂
90	29 272.677	3415.223	3d[1½]₁–5p[½]₀		{36 385.119	2747.627	4d[1½]₂–4f'[2½]₃
40	29 558.23	3382.230	4d'[2½]₀–5f'[3½]₃,₄	5	36 420.419	2744.964	4d[½]₀–6p[1½]₁
1200 <sup>a</sup>	29 788.667	3356.066	5s[1½]₀–5p[2½]₂	30	36 482.046	2740.327	5p[2½]₂–6s[1½]₂
11	30 042.871	3327.669	5d'[2½]₀–4f[2½]₃	9	37 003.491	2701.711	5p'[½]₁–6s'[½]₁
5	30 045.697	3327.356	5d'[2½]₀–4f[2½]₂	14	37 075.797	2696.442	5p[1½]₁–4d[1½]₁
60	30 453.764	3282.771	3d[2½]₀–5p[2½]₂	9	37 133.416	2692.258	5p'[1½]₂–6s'[½]₁
55	30 544.49	3273.020	4d[2½]₀–5f[3½]₃	6	37 176.057	2689.170	5p'[1½]₁–6s'[½]₁
20	{30 635.47	3263.300	4d[2½]₀–5f[2½]₂	8	37 251.067	2683.755	5p[2½]₂–6s[1½]₂
	{30 637.72	3263.060	4d[2½]₀–5f[2½]₃	1	37 911.499	2637.003	5p'[½]₁–6s'[½]₀
80	30 987.774	3226.199	3d[2½]₀–5p[2½]₃	9	38 110.332	2623.245	5p[2½]₂–4d[2½]₃
800 <sup>a</sup>	31 324.485	3191.520	5s[1½]₀–5p[½]₁	2	38 320.762	2608.840	5p[1½]₁–6s[1½]₁
50	31 718.65	3151.060	4d[2½]₀–5f[3½]₃,₄	1	38 384.869	2604.483	5p'[1½]₁–4d'[1½]₁
18	{31 816.76	3142.140	4d[2½]₀–5f[2½]₂	2	38 630.293	2587.936	5s'[½]₁–5p[½]₀
	{31 819.20	3141.900	4d[2½]₀–5f[2½]₃	4	38 800.601	2576.577	5p[1½]₂–6s[1½]₁
20	31 943.64	3129.660	4d[2½]₀–5f[4½]₄	4	38 950.321	2566.673	5p[2½]₂–4d'[2½]₃
25	31 986.21	3125.495	4d'[1½]₁–5f[2½]₂	1	39 169.420	2552.316	5p'[½]₁–4d'[1½]₁
9	32 038.85	3120.360	4d[2½]₀–5f[1½]₂	2	39 319.127	2542.598	3d'[2½]₀–5p[1½]₂
20	32 226.556	3102.185	3d'[1½]₁–5p'[1½]₂	1	39 487.540	2531.754	5p[1½]₁–6s[1½]₁
30	32 247.469	3100.173	5p[½]₁–6s[1½]₁	1	39 824.470	2510.334	3d'[2½]₀–5p[1½]₁
12	32 297.104	3095.409	4p'[½]₀–3d[½]₁	4	39 997.240	2499.491	5p[1½]₂–6s[1½]₀
50	32 325.060	3092.732	3d'[1½]₁–5p'[½]₁	4	40 880.069	2445.513	5p[2½]₂–4d[2½]₂
11	32 650.266	3061.927	5p'[½]₀–5d[½]₁				

<sup>a</sup> Observed interferometrically.<sup>b</sup> First reported by Sittner and Peck.<sup>c</sup> Reported also by Hepner.<sup>d</sup> Observed resolved by Séguier.<sup>e</sup> Unobserved expected transition, probably obscured by adjacent strong line.<sup>f</sup> Does not appear on all records.

TABLE 3. Energy levels of Ar I, partial list

Designation	$E/hc$ (cm $^{-1}$ )	Designation	$E/hc$ (cm $^{-1}$ )	Designation	$E/hc$ (cm $^{-1}$ )	Designation	$E/hc$ (cm $^{-1}$ )
Even levels				Odd levels			
$4p[\frac{1}{2}]_1$	104 102.140	$5p[\frac{1}{2}]_1$	116 660.035	$3d[\frac{1}{2}]_0^o$	111 667.807	$4d[\frac{1}{2}]_0^o$	118 512.234
$4p[2\frac{1}{2}]_3$	105 462.801	$5p[2\frac{1}{2}]_3$	116 942.795	$3d[\frac{1}{2}]_1^o$	111 818.069	$4d[\frac{1}{2}]_1^o$	118 651.438
$4p[2\frac{1}{2}]_2$	105 617.311	$5p[2\frac{1}{2}]_2$	116 999.367	$3d[3\frac{1}{2}]_4^o$	112 750.194	$4d[3\frac{1}{2}]_4^o$	119 023.688
$4p[1\frac{1}{2}]_1$	106 087.301	$5p[1\frac{1}{2}]_1$	117 151.368	$3d[3\frac{1}{2}]_3^o$	113 020.396	$4d[3\frac{1}{2}]_3^o$	119 212.930
$4p[1\frac{1}{2}]_2$	106 237.593	$5p[1\frac{1}{2}]_2$	117 183.631	$3d[1\frac{1}{2}]_2^o$	112 138.965	$4d[1\frac{1}{2}]_2^o$	118 906.648
$4p[\frac{1}{2}]_0$	107 054.313	$5p[\frac{1}{2}]_0$	117 562.996	$3d[\frac{1}{2}]_1^o$	114 147.773	$4d[\frac{1}{2}]_1^o$	119 847.810
$4p'[\frac{1}{2}]_1$	107 131.750	$5p'[1\frac{1}{2}]_1$	118 407.472	$3d[2\frac{1}{2}]_2^o$	113 426.005	$4d[2\frac{1}{2}]_2^o$	119 444.880
$4p'[\frac{1}{2}]_2$	107 289.741	$5p'[1\frac{1}{2}]_2$	118 469.092	$3d[2\frac{1}{2}]_3^o$	113 716.596	$4d[2\frac{1}{2}]_3^o$	119 566.040
$4p'[\frac{1}{2}]_1$	107 496.458	$5p'[\frac{1}{2}]_1$	118 459.639	$3d'[2\frac{1}{2}]_2^o$	114 641.033	$4d'[2\frac{1}{2}]_2^o$	120 619.055
$4p'[\frac{1}{2}]_0$	108 722.661	$5p'[\frac{1}{2}]_0$	118 870.958	$3d'[2\frac{1}{2}]_3^o$	114 821.980	$4d'[2\frac{1}{2}]_3^o$	120 753.510
$4f[1\frac{1}{2}]_1$	120 188.273			$3d'[1\frac{1}{2}]_2^o$	114 805.176	$4d'[1\frac{1}{2}]_2^o$	120 600.932
$4f[1\frac{1}{2}]_2$	120 100.672			$3d'[1\frac{1}{2}]_1^o$	115 366.907	$4d'[1\frac{1}{2}]_1^o$	121 011.955
$4f[4\frac{1}{2}]_5$	120 207.175			$5s[1\frac{1}{2}]_2^o$	113 468.514	$6s[1\frac{1}{2}]_2^o$	119 683.122
$4f[4\frac{1}{2}]_4$	120 207.216			$5s[1\frac{1}{2}]_1^o$	113 643.301	$6s[1\frac{1}{2}]_1^o$	119 760.208
$4f[2\frac{1}{2}]_3$	120 229.791			$5s'[\frac{1}{2}]_0^o$	114 861.676	$6s'[\frac{1}{2}]_0^o$	121 096.642
$4f[2\frac{1}{2}]_2$	120 230.104			$5s'[\frac{1}{2}]_1^o$	114 975.060	$6s'[\frac{1}{2}]_1^o$	121 161.350
$4f[3\frac{1}{2}]_3$	120 249.917			$5d[\frac{1}{2}]_0^o$	121 794.137	$6d[\frac{1}{2}]_0^o$	123 508.953
$4f[3\frac{1}{2}]_4$	120 249.955			$5d[\frac{1}{2}]_1^o$	121 932.885	$6d[\frac{1}{2}]_1^o$	123 468.008
$4f'[2\frac{1}{2}]_3$	121 654.275			$5d[3\frac{1}{2}]_4^o$	122 036.134	$6d[3\frac{1}{2}]_4^o$	123 653.210
$4f'[2\frac{1}{2}]_2$	121 654.621			$5d[3\frac{1}{2}]_3^o$	122 160.184	$6d[3\frac{1}{2}]_3^o$	123 773.892
$4f'[3\frac{1}{2}]_3$	121 653.279			$5d[1\frac{1}{2}]_2^o$	122 086.955	$6d[1\frac{1}{2}]_2^o$	123 808.554
$4f'[3\frac{1}{2}]_4$	121 653.321			$5d[1\frac{1}{2}]_1^o$	122 514.232	$6d[1\frac{1}{2}]_1^o$	
$6p[\frac{1}{2}]_1$	121 068.774	$7p[\frac{1}{2}]_1$	123 171.954	$5d[2\frac{1}{2}]_2^o$	122 282.139	$6d[2\frac{1}{2}]_2^o$	123 826.789
$6p[2\frac{1}{2}]_3$	121 165.408	$7p[2\frac{1}{2}]_3$	123 205.813	$5d[2\frac{1}{2}]_3^o$	122 329.712	$6d[2\frac{1}{2}]_3^o$	123 832.452
$6p[2\frac{1}{2}]_2$	121 191.836	$7p[2\frac{1}{2}]_2$	123 220.730	$5d'[2\frac{1}{2}]_2^o$	123 505.530	$6d'[2\frac{1}{2}]_2^o$	125 113.417
$6p[1\frac{1}{2}]_1$	121 257.198	$7p[1\frac{1}{2}]_1$	123 254.920	$5d'[2\frac{1}{2}]_3^o$	123 557.460	$6d'[2\frac{1}{2}]_3^o$	125 149.968
$6p[1\frac{1}{2}]_2$	121 270.658	$7p[1\frac{1}{2}]_2$	123 261.569	$5d'[1\frac{1}{2}]_2^o$	123 372.959	$6d'[1\frac{1}{2}]_2^o$	125 066.459
$6p[\frac{1}{2}]_0$	121 470.278	$7p[\frac{1}{2}]_0$	123 385.088	$5d'[1\frac{1}{2}]_1^o$	123 815.593	$6d'[1\frac{1}{2}]_1^o$	125 286.280
$6p'[1\frac{1}{2}]_1$	122 609.678	$7p'[1\frac{1}{2}]_1$	124 643.540				
$6p'[1\frac{1}{2}]_2$	122 635.092	$7p'[1\frac{1}{2}]_2$	124 658.490	$7s[1\frac{1}{2}]_2^o$	122 440.105	$8s[1\frac{1}{2}]_2^o$	123 903.255
$6p'[\frac{1}{2}]_1$	122 601.255	$7p'[\frac{1}{2}]_1$	124 650.990	$7s[1\frac{1}{2}]_1^o$	122 479.422	$8s[1\frac{1}{2}]_1^o$	123 935.910
$6p'[\frac{1}{2}]_0$	122 790.586	$7p'[\frac{1}{2}]_0$	124 749.845	$7s'[\frac{1}{2}]_0^o$	123 873.033	$8s'[\frac{1}{2}]_0^o$	125 334.750
				$7s'[\frac{1}{2}]_1^o$	123 882.236	$8s'[\frac{1}{2}]_1^o$	125 353.310

TABLE 4. Description of  $^{136}\text{Xe I}$ , region 1.2–4.0  $\mu\text{m}$ 

Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm $^{-1}$ )	Classification	Relative intensity	Calculated wavelength Air (Å)	Calculated wave number (cm $^{-1}$ )	Classification
50 <sup>b</sup>	12 203.818	8191.915	$5d[3\frac{1}{2}]_3^o-6p'[1\frac{1}{2}]_2$	375 <sup>b</sup>	14 364.987	6959.468	$5d[1\frac{1}{2}]_1^o-4f[1\frac{1}{2}]_2$
375 <sup>b</sup>	12 235.243	8170.875	$6p[\frac{1}{2}]_1-7s[1\frac{1}{2}]_1^o$	35 <sup>b</sup>	14 384.961	6949.805	$5d[1\frac{1}{2}]_1^o-4f[1\frac{1}{2}]_1$
100 <sup>b</sup>	12 257.765	8155.862	$5d[\frac{1}{2}]_0-7p[\frac{1}{2}]_1$	15	14 424.187	6930.905	$7s[1\frac{1}{2}]_1^o-8p[1\frac{1}{2}]_2$
50 <sup>b</sup>	12 271.904	8146.466	$5d[1\frac{1}{2}]_2-7p[2\frac{1}{2}]_3$	10	14 503.404	6893.049	$7s[1\frac{1}{2}]_1^o-8p[1\frac{1}{2}]_1$
20	12 409.131	8056.378	$5d[\frac{1}{2}]_2-6p'[1\frac{1}{2}]_1$	140 <sup>b</sup>	14 660.806	6819.044	$5d[2\frac{1}{2}]_2-7p[1\frac{1}{2}]_1$
75 <sup>b</sup>	12 451.547	8028.934	$5d[\frac{1}{2}]_2-7p[2\frac{1}{2}]_2$	3000 <sup>a, b</sup>	14 732.8055	6785.7189	$6p[2\frac{1}{2}]_3-7s[1\frac{1}{2}]_2$
300 <sup>b</sup>	12 590.203	7940.511	$5d[\frac{1}{2}]_1^o-7p[\frac{1}{2}]_1$	25	14 742.310	6781.344	$7s[1\frac{1}{2}]_1^o-8p[2\frac{1}{2}]_2$
2500 <sup>a, b</sup>	12 623.3912	7919.6350	$6p[\frac{1}{2}]_1-7s[1\frac{1}{2}]_2$	10	14 811.534	6749.650	$6p'[1\frac{1}{2}]_2-10d[3\frac{1}{2}]_3$
75	13 331.868	7498.774	$5d[\frac{1}{2}]_3^o-7p[2\frac{1}{2}]_2$	20	14 850.038	6732.150	$5d[2\frac{1}{2}]_3-6p'[1\frac{1}{2}]_2$
5	13 470.781 <sup>d</sup>	7421.445	$6p[1\frac{1}{2}]_1-7s[\frac{1}{2}]_1^o$	10	15 060.181	6638.212	$6p'[1\frac{1}{2}]_2-7s[\frac{1}{2}]_1^o$
250	13 544.152	7381.242	$5d[3\frac{1}{2}]_3-7p[2\frac{1}{2}]_2$	100	15 099.725	6620.828	$6p[\frac{1}{2}]_1-5d[1\frac{1}{2}]_1^o$
2000 <sup>a, b</sup>	13 657.0548	7320.2214	$6p[2\frac{1}{2}]_2-7s[1\frac{1}{2}]_1^o$	5	15 291.827	6537.654	$6d[3\frac{1}{2}]_3-7f[1\frac{1}{2}]_2$
10	13 814.410	7236.839	$5d[2\frac{1}{2}]_2-6p'[1\frac{1}{2}]_2$	2500 <sup>a, b</sup>	15 418.3935	6483.9883	$6p[1\frac{1}{2}]_1-7s[1\frac{1}{2}]_1^o$
15	13 919.611	7182.145	$7s[1\frac{1}{2}]_2^o-8p[1\frac{1}{2}]_2$	45	15 490.971	6453.610	$5d[2\frac{1}{2}]_2-6p'[1\frac{1}{2}]_1$
5	14 050.741	7115.117	$7s[1\frac{1}{2}]_1^o-8p[\frac{1}{2}]_0$	10	15 518.342 <sup>d</sup>	6442.227	$6p'[\frac{1}{2}]_1-7s[\frac{1}{2}]_0^o$
50	14 128.073	7076.172	$7s[\frac{1}{2}]_2^o-8p[2\frac{1}{2}]_3$	150	15 557.128	6426.166	$5d[2\frac{1}{2}]_2-7p[2\frac{1}{2}]_2$
1250 <sup>a, b</sup>	14 142.4436	7068.9814	$6p[2\frac{1}{2}]_2-7s[\frac{1}{2}]_2^o$	5	15 722.628	6358.523	$7p[\frac{1}{2}]_1-8d[1\frac{1}{2}]_2^o$
?	14 215.639	7032.584	$7s[\frac{1}{2}]_2^o-8p[2\frac{1}{2}]_2$	5	15 866.476	6300.875	$7p[\frac{1}{2}]_1-8d[\frac{1}{2}]_1^o$
800 <sup>b</sup>	14 240.959	7020.080	$5d[\frac{1}{2}]_1^o-4f[2\frac{1}{2}]_2$	250	15 979.536	6256.295	$5d[2\frac{1}{2}]_3-7p[1\frac{1}{2}]_2$

TABLE 4. Description of  $^{136}\text{Xe I}$ , region 1.2–4.0  $\mu\text{m}$ —Continued

Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification	Relative intensity	Calculated wavelength Air ( $\text{\AA}$ )	Calculated wave number ( $\text{cm}^{-1}$ )	Classification
100	16 039.905	6232.748	$6p[1\frac{1}{2}]_1 - 7s[1\frac{1}{2}]_2$	15	28 696.690	3483.772	$6p'[1\frac{1}{2}]_2 - 7d[3\frac{1}{2}]_3$
1000 <sup>b</sup>	16 053.281	6227.555	$6p[1\frac{1}{2}]_2 - 7s[1\frac{1}{2}]_1$	8	29 021.247	3444.812	$6d[1\frac{1}{2}]_2 - 8p[\frac{1}{2}]_1$
125	16 554.489	6039.008	$5d[2\frac{1}{2}]_3 - 7p[2\frac{1}{2}]_3$	75	29 046.734	3441.789	$7p[1\frac{1}{2}]_2 - 7d[\frac{1}{2}]_1$
1500 <sup>b</sup>	16 728.150	5976.315	$6p[1\frac{1}{2}]_2 - 7s[1\frac{1}{2}]_2$	300	29 384.406	3402.238	$7s[1\frac{1}{2}]_1^0 - 7p[\frac{1}{2}]_0$
50	16 745.724	5970.043	$5d[1\frac{1}{2}]_1 - 6p[\frac{1}{2}]_0$	150	29 449.055	3394.884	$7p[2\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_3$
15	16 834.541	5938.546	$7p[2\frac{1}{2}]_2 - 8d[3\frac{1}{2}]_3$	20	29 545.127	3383.730	$7p[1\frac{1}{2}]_1^0 - 7d[\frac{1}{2}]_1$
40	16 883.069	5921.476	$5d[2\frac{1}{2}]_3 - 7p[2\frac{1}{2}]_2$	100	29 649.585	3371.809	$6d[1\frac{1}{2}]_1^0 - 5f[2\frac{1}{2}]_2$
5	[17 139.084	5833.024	$6d[3\frac{1}{2}]_4 - 6f[4\frac{1}{2}]_4$	100	29 813.622	3353.257	$6d[3\frac{1}{2}]_2 - 8p[2\frac{1}{2}]_3$
	[17 140.611	5832.505	$6d[3\frac{1}{2}]_4 - 6f[4\frac{1}{2}]_5$	75	29 985.025	3334.089	$6d[1\frac{1}{2}]_1^0 - 5f[1\frac{1}{2}]_2$
1500 <sup>b</sup>	17 325.767	5770.174	$6p[2\frac{1}{2}]_2 - 5d[1\frac{1}{2}]_1$	12	30 021.886 <sup>d</sup>	3329.995	$8s[1\frac{1}{2}]_2 - 9p[2\frac{1}{2}]_3$
50	17 365.086	5757.109	$7p[2\frac{1}{2}]_3 - 8d[3\frac{1}{2}]_4$	600	30 253.143	3304.540	$7s[1\frac{1}{2}]_1^0 - 7p[1\frac{1}{2}]_1$
350 <sup>b</sup>	18 788.128	5321.057	$6p[\frac{1}{2}]_0 - 7s[1\frac{1}{2}]_1$	60	30 423.535	3286.033	$7p[\frac{1}{2}]_0 - 7d[\frac{1}{2}]_1$
150	20 187.190	4952.285	$5d[1\frac{1}{2}]_1^0 - 7p[\frac{1}{2}]_0$	1500 <sup>a</sup>	30 475.455	3280.4344	$7s[1\frac{1}{2}]_2^0 - 7p[2\frac{1}{2}]_3$
3000 <sup>a, b</sup>	20 262.2419	4933.9412	$6p[1\frac{1}{2}]_1 - 5d[1\frac{1}{2}]_1$	100	30 504.116	3277.352	$7p[2\frac{1}{2}]_3 - 5d[2\frac{1}{2}]_3$
50	21 373.073	4677.508	$6p[1\frac{1}{2}]_2 - 5d[1\frac{1}{2}]_1$	500	30 794.182	3246.481	$7s[1\frac{1}{2}]_1^0 - 7p[1\frac{1}{2}]_2$
250 <sup>c</sup>	21 470.089	4656.372	$6p[\frac{1}{2}]_1 - 5d[2\frac{1}{2}]_2$	15	30 855.221	3240.059	$6d[3\frac{1}{2}]_2 - 8p[2\frac{1}{2}]_3$
60	22 269.836	4489.154	$5d[1\frac{1}{2}]_1^0 - 6p'[\frac{1}{2}]_1$	6000 <sup>a</sup>	31 069.227	3217.7412	$6p[1\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_3$
40	[22 382.762	4466.505	$6d[3\frac{1}{2}]_4 - 5f[4\frac{1}{2}]_4$	80	31 275.972	3196.471	$6d[3\frac{1}{2}]_3 - 8p[2\frac{1}{2}]_2$
	[22 386.390	4465.781	$6d[3\frac{1}{2}]_4 - 5f[4\frac{1}{2}]_5$	125	31 336.011	3190.346	$7s[1\frac{1}{2}]_2^0 - 6p'[\frac{1}{2}]_1$
75	22 406.818	4461.710	$5d[1\frac{1}{2}]_1^0 - 7p[2\frac{1}{2}]_2$	550	31 607.907	3162.903	$7s[1\frac{1}{2}]_2^0 - 7p[2\frac{1}{2}]_2$
90	22 618.283	4419.996	$7s[1\frac{1}{2}]_1^0 - 6p'[\frac{1}{2}]_0$	100	32 293.081	3095.794	$7p[2\frac{1}{2}]_2 - 5d'[\frac{1}{2}]_2$
5	[22 741.699	4396.009	$6d[3\frac{1}{2}]_3 - 5f[3\frac{1}{2}]_4$	70	32 355.650	3089.808	$6d[2\frac{1}{2}]_2 - 8p[\frac{1}{2}]_1$
	[22 742.102	4395.931	$6d[3\frac{1}{2}]_3 - 5f[3\frac{1}{2}]_3$	12	32 581.916	3068.350	$6p'[1\frac{1}{2}]_1 - 5d'[\frac{1}{2}]_2$
40	22 964.776	4353.307	$6d[3\frac{1}{2}]_3 - 5f[4\frac{1}{2}]_4$	1800 <sup>a</sup>	32 739.262	3053.6037	$6p[\frac{1}{2}]_1 - 5d[1\frac{1}{2}]_2$
10	23 022.418	4342.407	$6p'[1\frac{1}{2}]_1 - 7d[1\frac{1}{2}]_2$	75	33 265.433	3005.304	$7p[\frac{1}{2}]_1 - 8s[\frac{1}{2}]_1^0$
45	23 073.456	4332.802	$7p[\frac{1}{2}]_1 - 7d[\frac{1}{2}]_0$	22	33 526.746 <sup>d</sup>	2981.880	$5d'[2\frac{1}{2}]_2 - 8p[2\frac{1}{2}]_3$
8	23 105.265	4326.837	$7p[2\frac{1}{2}]_2 - 7d[2\frac{1}{2}]_2$	50	33 567.470	2978.262	$7p[2\frac{1}{2}]_2 - 5d'[\frac{1}{2}]_2$
1250 <sup>a, c</sup>	23 193.332	4310.4078	$6p[2\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_3$	3500 <sup>a</sup>	33 666.692	2969.4849	$6p[1\frac{1}{2}]_1 - 5d[2\frac{1}{2}]_2$
35	23 252.750	4299.393	$6p'[1\frac{1}{2}]_1 - 7d[2\frac{1}{2}]_2$	150	34 014.669	2939.106	$7s[1\frac{1}{2}]_1^0 - 6p'[\frac{1}{2}]_1$
110	23 279.541	4294.445	$7p[2\frac{1}{2}]_2 - 7d[3\frac{1}{2}]_3$	90	34 074.837	2933.917	$6s'[\frac{1}{2}]_1^0 - 6p[\frac{1}{2}]_0$
35	23 443.639	4264.386	$7p[2\frac{1}{2}]_3 - 7d[2\frac{1}{2}]_3$	450	34 335.274	2911.663	$7s[1\frac{1}{2}]_2^0 - 7p[2\frac{1}{2}]_2$
60	23 796.466	4201.158	$7p[\frac{1}{2}]_1 - 7d[\frac{1}{2}]_1$	170	34 744.002	2877.410	$7p[\frac{1}{2}]_1 - 8s[\frac{1}{2}]_2^0$
30	23 934.491	4176.914	$7p[2\frac{1}{2}]_3 - 7d[3\frac{1}{2}]_3$	75	35 028.676	2854.025	$6p'[\frac{1}{2}]_1 - 7d[\frac{1}{2}]_1^0$
70	24 443.648	4089.926	$7s[1\frac{1}{2}]_2^0 - 6p'[\frac{1}{2}]_1$	5000 <sup>a</sup>	35 070.253	2850.642	$6p[2\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_3$
60	24 702.317	4047.099	$7p[1\frac{1}{2}]_2 - 7d[2\frac{1}{2}]_3$	?	35 083.277	2849.584	$6p'[\frac{1}{2}]_1 - 6d[\frac{1}{2}]_1^0$
30	24 776.187	4035.033	$7p[1\frac{1}{2}]_2 - 7d[1\frac{1}{2}]_2$	110	35 246.924	2836.354	$6d[2\frac{1}{2}]_3 - 8p[\frac{1}{2}]_2$
1800 <sup>a, c</sup>	24 824.712	4027.1453	$6p[2\frac{1}{2}]_3 - 5d[2\frac{1}{2}]_3$	30	35 691.926	2800.990	$7p[2\frac{1}{2}]_2 - 5d'[\frac{1}{2}]_2$
175	25 145.842	3975.716	$7p[2\frac{1}{2}]_3 - 7d[3\frac{1}{2}]_4$	20	36 045.094	2773.546	$6p'[1\frac{1}{2}]_1 - 5d'[\frac{1}{2}]_2$
60	25 159.384	3973.576	$7s[1\frac{1}{2}]_2^0 - 6p'[\frac{1}{2}]_2$	250	36 209.206	2760.976	$7p[1\frac{1}{2}]_2 - 5d'[\frac{1}{2}]_2$
45	25 412.748	3933.960	$7p[\frac{1}{2}]_1 - 7d[2\frac{1}{2}]_2$	150	36 231.741	2759.258	$6s'[\frac{1}{2}]_0 - 6p[\frac{1}{2}]_1$
30	25 820.844	3871.784	$7p[\frac{1}{2}]_0 - 7d[\frac{1}{2}]_1$	450	36 508.360	2738.352	$7s[1\frac{1}{2}]_2^0 - 7p[\frac{1}{2}]_1$
50	26 020.700	3842.046	$6d[\frac{1}{2}]_0 - 8p[\frac{1}{2}]_1$	20	36 614.952	2730.380	$6d[2\frac{1}{2}]_3 - 8p[2\frac{1}{2}]_3$
10	26 043.472	3838.686	$7s[1\frac{1}{2}]_1^0 - 6p'[\frac{1}{2}]_1$	850	36 788.827	2717.475	$6p[\frac{1}{2}]_1 - 5d'[\frac{1}{2}]_1^0$
2000 <sup>a, c</sup>	26 269.084	3805.7180	$6p[2\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_2$	190	36 848.818	2713.051	$6p[1\frac{1}{2}]_2 - 5d[2\frac{1}{2}]_2$
8	26 471.568	3776.608	$7p[2\frac{1}{2}]_2 - 7d[\frac{1}{2}]_1$	25	37 255.184	2683.458	$7p[2\frac{1}{2}]_2 - 5d'[\frac{1}{2}]_2$
2500 <sup>a, c</sup>	26 510.861	3771.0102	$6p[\frac{1}{2}]_0 - 5d[\frac{1}{2}]_1^0$	10	38 496.568	2596.926	$8s[\frac{1}{2}]_2^0 - 5f[\frac{1}{2}]_3$
8	26 601.140 <sup>d</sup>	3758.212	$6p'[\frac{1}{2}]_0 - 5d'[\frac{1}{2}]_1^0$	140	38 685.985	2584.211	$6p'[\frac{1}{2}]_1 - 5d'[\frac{1}{2}]_2$
30	[27 296.636	3662.456	$6d[\frac{1}{2}]_2 - 8p[\frac{1}{2}]_2$	175	38 737.815	2580.753	$7p[2\frac{1}{2}]_2 - 8s[\frac{1}{2}]_1^0$
	[27 298.111	3662.258	$6d[\frac{1}{2}]_0 - 8p[\frac{1}{2}]_1$	270	38 939.602	2567.380	$6p[2\frac{1}{2}]_3 - 5d[3\frac{1}{2}]_3$
15	27 743.212	3603.502	$6d[\frac{1}{2}]_1^0 - 8p[\frac{1}{2}]_1$	20	39 154.184	2553.309	$6p'[\frac{1}{2}]_1 - 8s[\frac{1}{2}]_1^0$
15	28 088.713	3559.178	$6p'[\frac{1}{2}]_1 - 7d[\frac{1}{2}]_2$	8	39 624.876	2522.979	$6d[\frac{1}{2}]_1^0 - 8p[\frac{1}{2}]_0$
50	28 115.553	3555.780	$7s[1\frac{1}{2}]_2^0 - 7p[\frac{1}{2}]_1$	120	39 955.140	2502.125	$6p[\frac{1}{2}]_1 - 5d[\frac{1}{2}]_0$
250	28 381.545	3522.455	$6p[2\frac{1}{2}]_3 - 5d[2\frac{1}{2}]_2$	25	40 196.317	2487.112	$7s[1\frac{1}{2}]_1^0 - 7p[\frac{1}{2}]_1$
8	28 458.790	3512.895	$6d[\frac{1}{2}]_2 - 8p[\frac{1}{2}]_2$	12	40 757.634	2452.859	$7p[2\frac{1}{2}]_2 - 8s[\frac{1}{2}]_2^0$
750	28 582.246	3497.721	$7s[1\frac{1}{2}]_2^0 - 7p[\frac{1}{2}]_2$	15	41 514.978	2408.112	$7p[\frac{1}{2}]_1 - 5d'[\frac{1}{2}]_2$

<sup>a</sup> Observed Interferometrically.<sup>b</sup> First reported by Sittner and Peck.<sup>c</sup> First reported by Hepner.<sup>d</sup> Calculated from AEL levels.