

Table 2. Estimated numbers of diagnoses of HIV/AIDS, by race/ethnicity, sex, and age group, 1999–2002—30 with confidential name-based HIV infection reporting

Age at diagnosis (yrs)	Year of diagnosis			
	1999	2000	2001	2002
White, not Hispanic				
Male				
<13	7	6	9	12
13–19	49	58	60	61
20–24	340	323	372	407
25–34	2,020	1,865	1,804	1,858
35–44	2,584	2,752	2,707	2,885
45–54	1,019	1,196	1,215	1,283
55–64	268	309	381	397
≥65	93	72	102	95
Subtotal	6,379	6,581	6,651	6,999
Female				
<13	9	14	20	11
13–19	67	70	68	65
20–24	157	152	145	142
25–34	415	416	454	409
35–44	442	476	458	439
45–54	190	202	175	210
55–64	40	55	54	52
≥65	19	17	14	20
Subtotal	1,339	1,403	1,389	1,348
Total	7,718	7,984	8,040	8,347
Black, not Hispanic				
Male				
<13	70	53	65	54
13–19	178	208	238	272
20–24	623	727	786	742
25–34	2,488	2,281	2,172	2,206
35–44	3,270	3,145	3,089	3,221
45–54	1,578	1,595	1,706	1,760
55–64	481	502	481	554
≥65	152	168	181	172
Subtotal	8,839	8,680	8,716	8,979
Female				
<13	73	72	76	60
13–19	348	312	309	286
20–24	624	647	623	605
25–34	1,814	1,655	1,625	1,596
35–44	1,730	1,656	1,654	1,649
45–54	695	791	774	877
55–64	199	242	230	255
≥65	76	75	84	91
Subtotal	5,558	5,449	5,374	5,420
Total	14,397	14,129	14,090	14,398
Hispanic				
Male				
<13	10	8	13	7
13–19	20	27	33	61
20–24	132	166	183	173
25–34	754	796	784	906
35–44	765	869	851	919
45–54	265	293	340	378
55–64	79	98	101	122
≥65	25	39	34	41
Subtotal	2,051	2,297	2,339	2,607
Female				
<13	17	9	21	13
13–19	37	18	26	32
20–24	54	68	75	73
25–34	181	205	208	218
35–44	169	197	193	205
45–54	81	90	111	115

Table 2. (continued)

Age at diagnosis (yrs)	Year of diagnosis			
	1999	2000	2001	2002
Hispanic (continued)				
Female (continued)				
55–64	32	28	31	47
≥65	10	8	7	11
Subtotal	580	623	672	715
Total	2,631	2,920	3,012	3,321
Asian/Pacific Islander				
Male				
<13	0	0	0	1
13–19	0	3	0	2
20–24	8	5	7	1
25–34	39	54	39	54
35–44	28	43	36	41
45–54	11	8	20	11
55–64	2	4	4	7
≥65	3	0	0	1
Subtotal	91	118	105	119
Female				
<13	0	0	1	1
13–19	1	1	0	1
20–24	5	3	7	2
25–34	14	16	12	16
35–44	8	10	10	7
45–54	2	0	2	3
55–64	0	0	0	0
≥65	1	2	2	0
Subtotal	32	32	35	30
Total	123	150	140	149
American Indian/Alaska Native				
Male				
<13	0	0	0	1
13–19	3	0	1	2
20–24	11	13	6	8
25–34	39	31	41	38
35–44	35	45	40	47
45–54	17	8	17	9
55–64	3	2	4	9
≥65	1	2	1	1
Subtotal	109	101	110	115
Female				
<13	0	0	0	0
13–19	2	4	1	3
20–24	3	8	5	6
25–34	13	12	10	15
35–44	14	16	6	14
45–54	6	5	8	14
55–64	4	0	2	1
≥65	0	0	0	0
Subtotal	43	46	32	53
Total	153	147	142	168

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV/AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1998, the following 30 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Arkansas, Arizona, Florida, Colorado, Iowa, Idaho, Indiana, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Utah, Virginia, Wisconsin, West Virginia, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.