

**NORA *manufacturing* sector: Proportionate mortality ratio (PMR) adjusted for age, sex, and race for selected work-related respiratory conditions, U.S. residents age 15 and over, selected states and years, 1990–1999**

Condition	Number of Deaths	PMR	95% Confidence Interval	
			LCL	UCL
Byssinosis	50	3.1	2.3	4.1
Silicosis	386	1.7	1.6	1.9
Asbestosis	1,035	1.4	1.3	1.5
Malignant Mesothelioma (all sites)*	134	1.2	1.0	1.4
All Pneumoconioses	1,960	0.7	0.7	0.8
Unspecified and Other Pneumoconioses	236	0.6	0.6	0.7
Hypersensitivity Pneumonitis	13	0.5	0.3	0.8
Coal Workers' Pneumoconiosis	293	0.2	0.2	0.3
All other conditions within this sector	1,034,734			
<b>TOTAL</b>	<b>1,038,841</b>			

NORA - National Occupational Research Agenda

\* data for malignant mesothelioma are for 1999 only.

NOTE: The sum of the individual and all other condition death totals may be greater than the total number of deaths in the selected states and years because some decedents have more than one condition listed on their death certificates. See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.