

PUBLICATIONS EXAMINED

Publications examined is arranged by category, as follows:

Sampling Strategies and Exposure Assessment
Sampling and Analytical Methods
Coal Mine Dust Characteristics and Concentrations
Engineering Controls
Medical Evaluation and Testing
Chronic Obstructive Pulmonary Disease
Lung Function
Coal Workers' Pneumoconiosis
Particle Deposition and Clearance from Lungs
Cellular and Animal Studies
Silica Exposures and Silicosis
Diesel Exposures, Controls, and Health Effects
Health and Safety
Miscellaneous

SAMPLING STRATEGIES AND EXPOSURE ASSESSMENT

Andria GD, Saltsman RD, Kost JA, Zalar JL [1983]. Sampling plan and statistical evaluation for bituminous coal research underground dust-sampling programs. In: Marple VA, Liu BYH, eds. *Aerosols in the mining industrial work environments*. Vol. 1. Ann Arbor, MI: Ann Arbor Science, pp. 235-253.

Armstrong BG, Oakes D [1982]. Effects of approximation in exposure assessments on estimates of exposure-response relationships. *Scand J Work Environ Health* 8(Suppl 1):20-23.

Attfield MD, Hewett P [1992]. Exact expressions for the bias and variance of estimators of the mean of a lognormal distribution. *Am Ind Hyg Assoc J* 53(7):432-435.

Bodsworth PL, Hurley JF, Tran CL, Weston P, Wetherill GZ [1993]. Dust sampling strategy on coalfaces in relation to modern coalmining methods. Edinburgh, Scotland: Institute of Occupational Medicine, Report No. TM/93/06.

Bowman JD, Busch KA, Shulman SA [1983]. Statistical evaluation of sampling strategies for coal mine dust. In: Marple VA, Liu BYH, eds. *Aerosols in the mining and industrial work environments*. Vol. 1. Ann Arbor, MI: Ann Arbor Science Publishers, pp. 283-298.

Breslin AJ, Ong L, Glauberman H, George AC, LeClare P [1967]. The accuracy of dust exposure estimates obtained from occupational air sampling. *Am Ind Hyg Assoc J* 28:56-61.

Checkoway H, Savitz DA, Heyer NJ [1991]. Assessing the effects of nondifferential misclassification of exposures in occupational studies. *Appl Occup Environ Hyg* 6(6):528-533.

Coenen W [1971]. The measurement and evaluation of the concentration of harmful dust, especially silicogenic, at working places of the aboveground industry. *Staub Reinhalt* 31:482-490.

Corn M, Esmen NA [1979]. Workplace exposure zones for classification of employee exposures to physical and chemical agents. *Am Ind Hyg Assoc J* 40(1):47-57.

Esmen NA [1983]. Mathematical basis for an efficient sampling strategy. In: Marple VA, Liu BYH, eds. *Aerosols in the mining and industrial work environments*. Vol. 1. Ann Arbor, MI: Ann Arbor Science, pp. 271-282.

Esmen NA [1992]. A distribution-free double-sampling method for exposure assessment. *Appl Occup Environ Hyg* 7(9):613-621.

Heederik D, Miller BG [1988]. Weak associations in occupational epidemiology: adjustment for exposure estimation error. *Int J Epidemiol* 17(4)(Suppl):970-974.

Heederik D, Boleij JSM, Kromhout H, Smid T [1991a]. Use and analysis of exposure monitoring data in occupational epidemiology: an example of an epidemiological study in the Dutch animal food industry. *Appl Occup Environ Hyg* 6(6):458-463.

Heederik D, Kromhout H, Burema J [1991b]. Assessment of long-term exposures to toxic substances in air [letters]. *Ann Occup Hyg* 35(6):671-673.

Jacobsen M, Dodgson J [1981]. Long-term experience in collecting and using occupational health data in the coal industry. *Ann Occup Hyg* 24(4):391-398.

Lippmann M [1991]. Research needs: retrospective exposure assessment for occupational epidemiology. *Appl Occup Environ Hyg* 6(6):550-554.

Makepeace CE, Horvath FJ, Stocker H [1983]. Application of a stratified random sampling technique to estimation and minimization of respirable quartz exposure to underground miners. In: Marple VA, Liu BYH, eds. *Aerosols in the mining industrial work environments*. Vol. 1. Ann Arbor, MI: Ann Arbor Science, pp. 261-270.

Mutmansky JM, Grayson RL, Rubow KL, Marple VA, Stobbe TL, Meyers WR, et al. [1990]. Formulation, evaluation and verification of improved dust sampling and analytical strategies for

use at surface and underground coal mines. Washington, DC: U.S. Department of the Interior, Bureau of Mines, PB-94-106-713.

Nicas M, Simmons BP, Spear RC [1991]. Environmental versus analytical variability in exposure measurements. *Am Ind Hyg Assoc J* 52(12):533-557.

Rappaport SM [1985]. Smoothing of exposure variability at the receptor: implications for health standards. *Ann Occup Hyg* 29(2):201-214.

Rappaport SM [1991a]. Exposure assessment strategies. In: Rappaport SM, Smith TJ, eds. *Exposure assessment for epidemiology and hazard control*. Chelsea, MI: Lewis Publishers, Inc., pp. 219-249.

Rappaport SM [1991b]. Selection of the measures of exposure for epidemiology studies. *Appl Occup Environ Hyg* 6(6):448-457.

Rappaport SM [1991c]. Review: assessment of long-term exposures to toxic substances in air. *Ann Occup Hyg* 35(1):61-121.

Rappaport SM, Spear RC, Selvin S [1988a]. The influence of exposure variability on dose-response relationships. *Ann Occup Hyg* 32(Suppl 1):529-537.

Rappaport SM, Selvin S, Roach SA [1988b]. A strategy for assessing exposures with reference to multiple limits. *Appl Ind Hyg* 3(11):310-315.

Seixas NS [1994]. Exposure assessment and public health. *New Solutions*, Winter, pp. 34-42.

Seixas N, Robins TG, Moulton LH [1988]. The use of geometric and arithmetic mean exposures in occupational epidemiology. *Am J Ind Med* 14:465-477.

Seixas NS, Moulton LH, Robins TG, Rice CH, Attfield MD, Zellers ET [1991]. Estimation of cumulative exposures for the national study of coal workers' pneumoconiosis. *Appl Occup Environ Hyg* 6(12):1032-1041.

Seixas NS, Robins TG, Becker M [1993]. A novel approach to the characterization of cumulative exposure for the study of chronic occupational disease. *Am J Epidemiol* 137(4):463-471.

Selvin S, Rappaport SM [1989]. A note on the estimation of the mean value from a lognormal distribution. *Am Ind Hyg Assoc J* 50(12):627-630.

Spear RC [1991]. Individual versus group differences in exposure and risk. In: Rappaport SM, Smith TJ, eds. *Exposure assessment for epidemiology and hazard control*. Chelsea, MI: Lewis Publishers, Inc., pp. 283-295.

Smith TJ [1990]. Occupational exposure and dose over time: limitations of cumulative exposure. *Am J Ind Med* 21:35-51.

Stellingwerf J [1983]. Time-dependent aspects in the European communities directives on asbestos and lead. In: Proceedings of the VIth International Pneumoconiosis Conference. Vol. 3. Bochum, Germany: International Labor Office, pp. 1631-1637.

Tuggle RM [1981]. The NIOSH decision scheme. *Am Ind Hyg Assoc J* 42(7):493-498.

Tuggle RM [1982]. Assessment of occupational exposure using one-sided tolerance limits. *Am Ind Hyg Assoc J* 43(5):338-346.

Ulfvarson U [1983]. Limitations to the use of employee exposure data on air contaminants in epidemiologic studies. *Int Arch Occup Environ Health* 52:285-300.

WHO (World Health Organization) [1984]. Evaluation of exposure to airborne particles in the work environment. Geneva, Switzerland: WHO Offset Publication No. 80.

SAMPLING AND ANALYTICAL METHODS

Ashe HB [1955]. Silicosis and dust control. *Public Health Rep* 70(10):983-985.

Bartley DL, Fischback TJ [1993]. Alternative approaches for analyzing sampling and analytical methods. *Appl Occup Environ Hyg* 8(4):381-385.

Bierbaum PJ, Doemeny LJ, Smith JP, Abell MT [1993]. U.S. approach for air sampling of workplace contaminants: current basis and future options. *Appl Occup Environ Hyg* 8(4):247-250.

Busch KA [1977]. Standards completion program (SCP) statistical protocol. In: Taylor DG, Kupel RE, Bryant JM, eds. NIOSH validation tests. Cincinnati, OH: U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health, DHEW (NIOSH) Publication No. 77-185.

Caplan KJ, Doemeny LJ, Sorenson SD [1979]. Performance characteristics of the 10 mm respirable mass sampler: Part II—coal dust studies. *Am Ind Hyg Assoc J* 38:162-173.

Carlson DH, Cornilsen BC, Shan X, Johnson JH [1991]. Measurement and analysis of respirable dust and gaseous pollutant concentrations in three U.S. underground coal mines. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 383-392.

Cecala AB, Volkwein JC, Timko RJ, Williams KL [1983]. Velocity and orientation effects on the 10-mm Dorr-Oliver cyclone. Washington, DC: U.S. Department of the Interior, Bureau of Mines, RI 8764.

CEN (Comite European de Normalisation) [1993]. Workplace atmospheres. Assessment of performance of instruments for measurement of airborne particles. Brussels, Belgium: CEN/TC137/WG2/N125.

Cheng Y, Chen BT, Yeh H [1993]. Performance of an aerodynamic particle sizer. *Appl Occup Environ Hyg* 8(4):307-312.

Corn M [1975]. Remarks on determination of non-compliance with the respirable dust standard, Federal Coal Mine Health and Safety Act of 1969. *Am Ind Hyg Assoc J* 36:404-407.

Corn M [1992]. Historical perspective on approaches to estimation of inhalation risk by air sampling. *Am J Ind Med* 21:113-123.

Corn M, Breysse P, Hall T, Chen G, Risby T, Swift DI [1985]. A critique of MSHA procedures for determination of permissible respirable coal mine dust containing free silica. *Am Ind. Hyg. Assoc J* 46(1):4-8.

Feldman RF [1993]. Degraded instrument performance due to radio interference: criteria and standards. *Appl Occup Environ Hyg* 8(4):351-355.

Gentry SJ [1993]. Instrument performance and standards. *Appl Occup Environ Hyg* 8(4):260-266.

Goldberg SA, Raymond LD, Taylor CD [1976]. Mining enforcement and safety administration procedure for analysis of respirable dust from coal mines. In: *Proceedings of the Symposium on Silica, March 12-13, 1975, Pittsburgh, PA: Industrial Health Foundation, pp. 91-104.*

Goldberg SA, Tomb TF, Kacsmar PM, Baber JJ Jr., Busa MJ [1984]. MSHA's procedure for determining quartz content of respirable coal mine dust. Washington, DC: U.S. Department of Labor, Mine Safety and Health Administration, IR 1152.

Hall TA, Corn M, Zeger S, Law CC [1990]. Respirable dust and free silica variation in mine environments. In: *Proceedings of the VIIth International Pneumoconiosis Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.*

Harris BE, DeSieghardt WC, Riva LA [1976]. Development of an area sampling methodology. Final Report for the U.S. Bureau of Mines on Contract No. HO133108. Washington, DC: U.S. Department of the Interior.

Hearl FJ, Hewett P [1993]. Problems in monitoring dust levels with mines. *Occup Med: State of the Art Rev* 8(1):93-108.

Hinds WC [1982]. *Aerosol technology*. New York, NY: John Wiley and Sons.

HSE [1993]. General methods for the gravimetric determination of respirable and total inhalable dust, MDHS 14 (revised). In *HSE series: Methods for the determination of hazardous substances*. London, England: ISBN 0-11-882046-X.

Hwang CY [1988]. Measurements of respirable dust concentrations by using various samplers in underground coal mines. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 51-59.

ISO [1991]. *Air quality-assessment of performance of instruments used for the health-related sampling of particles at workplaces*. Brussels, Belgium: International Organization for Standardization, Draft Technical Report No. CEN/TC137/WG3/N 100.

John W [1993]. Instrument performance and standards for sampling of aerosols. *Appl Occup Environ Hyg* 8(4):251-259.

Jones, AD, Johnston AM, Vincent JH [1985]. The measurement of electric charge on airborne dusts in quarries and mines. *Staub Reinhalt* 45(10):475-480.

Kinney PL, Thurston GD [1993]. Field evaluation of instrument performance: statistical considerations. *Appl Occup Environ Hyg* 8(4):267-271.

Kissell FN, Ruggieri SK, Jankowski RA [1986]. How to improve the accuracy of coal mine dust sampling. *Am Ind Hyg Assoc J* 47(10):602-606.

Liedel NA, Busch KA [1975]. *Statistical methods for the determination of noncompliance with occupational health standards*. Cincinnati, OH: U.S. Department of Health, Education, and Welfare, Public Health Service, Center for Disease Control, National Institute for Occupational Safety and Health, Division of Laboratories and Criteria Development.

Mark D, Lyons CP, Upton SL [1993]. Performance testing of the respirable dust sampler used in British coal mines. *Appl Occup Environ Hyg* 8(4):370-380.

Mercer TT [1978]. Respirable fraction of airborne dust: quantitative descriptions, formal definitions, and performance characteristics of samplers matched to them. *J Test Eval* 6(1):9-19.

MSHA [1984]. *MSHA's procedure for determining quartz content of respirable coal mine dust*. Washington, DC: U.S. Department of Labor, Mine Safety and Health Administration, IR 1152.

Ondrey R, Stoltz R, Atchison D, Gerbec E [1988]. Dust sampling roof bolting operations. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 264-268.

Tomb TF, Haney RA [1988]. Comparison of number and respirable mass concentration determinations. In: *Advances in air sampling*. Chelsea, MI: Lewis Publishers, pp. 189-202.

Villnave JM, Corn M, Francis M, Hall TA [1991]. Regulatory implications of airborne respirable free silica variability in underground coal mines. *Am Ind Hyg Assoc J* 52(3):107-112.

COAL MINE DUST CHARACTERISTICS AND CONCENTRATIONS

Addison J, Miller BE, Porteous RH, Dodgson J, Robertson A [1990]. Shape, size and composition of coalmine dusts in relation to pneumoconiosis in British coalmines. Edinburgh, Scotland: Institute of Occupational Medicine, NTIS No. PB-91-229-948.

Amandus HE, Piacitelli G [1987]. Dust exposures at U.S. surface coal mines in 1982-1983. *Arch Environ Health* 42(6):374-381.

Andre R, Simonyi T, Grayson R [1988]. Variation in mineral and elemental composition of respirable coal mine dusts by worker location and coal seam. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 124-131.

Babu VS, Pandurangi RS, Seehra MS [1991]. X-ray diffraction and IR spectroscopy of three forms of silica and mine dusts. In: Frantz RL, Ramani RV, eds. *3rd Symposium on Respirable Dust in the Mineral Industries*. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 253-258.

Dieffenbach A [1988]. Trends in coal mine dust exposures for longwell miners: 1981-1987. Cincinnati, OH: U. S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, Division of Respiratory Disease Studies.

Grayson RL [1990]. The potential role of particle characteristics on coal mine respirable dust standards. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., Preprint No. 90-131.

Lee C, Mutmanský JM [1988]. Statistical analysis of the elemental characteristics of airborne coal mine dust. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 111-123.

Parobeck PS [1975]. Effect of the 2.0 Mg/m³ coal mine dust standard on underground environmental dust levels. *Am Ind Hyg Assoc J* 36(8):604-609.

Parobeck PS, Jankowski RA [1979]. Assessment of the respirable dust levels in the nation's underground and surface coal mining operations. *Am Ind Hyg Assoc J* 40:910-915.

Parobeck PS, Tomb TF [1974]. Respirable dust levels—surface work areas of underground coal mines and surface coal mines. *Work Environ Health* 11:43-48.

Ramani RV, Qin J, Jankowski RA [1991]. Characteristics of dust clouds at longwall faces. *Journal of Mine Ventilation Society of South Africa* 44(5):70-79.

Robins TG [1991]. Coal dust particle size and respiratory disease. In: NIOSH Grants Research and Demonstration Projects: Annual Report Fiscal Year 1991. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 92-109.

Shi X, Dalal NS, Hu XN, Vallyathan V [1989]. The chemical properties of silica particle surface in relation to silica-cell interactions. *J Toxicol Environ Health* 27:435-454.

Smith DK, Mutmanský JM, Klimkiewicz M, Marks JA [1991]. Quartz particulate behavior during the mechanical mining of coal seams. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 239-251.

Stobbe TJ, Plummer RW, Kim H, Dower JM [1986]. Characterization of coal mine dust. *Ann Am Conf Gov Ind Hyg* 14:689-696.

Tomb TF, Treafis HN, Gero AJ [1983]. Characteristics of underground coal mine dust aerosols. *Aerosols Mining Ind Work Environ* 2(1):395-405.

Wallace WE, Harrison J, Keane MJ, Bolsaitis P, Eppelsheimer D, Poston J, et al. [1990]. Clay occlusion of respirable quartz particles detected by low voltage scanning electron microscopy—x-ray analysis. *Ann Occup Hyg* 34(2):195-204.

Watts WF, Niewiadomski GE [1990]. Respirable dust trends in coal mines with longwall or continuous miners sections. In: Proceedings of the VIIth International Pneumoconioses Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Watts WF, Parker DR [1987]. Respirable dust levels in coal, metal, and nonmetal mines. Washington, DC: U.S. Department of the Interior, Bureau of Mines, IC 9125.

Weeks JL [1990]. Characteristics of chronically dusty longwall mines in the U.S. In: Proceedings of the VIIth International Pneumoconioses Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Xu L, Mutmanský JM [1991]. Analysis of physical, chemical, and mineralogical characteristics of airborne coal mine dusts. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 289-297.

ENGINEERING CONTROLS

Babbitt C, Bartlett P, Kelly J, Ludlow J, Mangolds A, Rajan S, et al. [1984]. Evaluate fundamental approaches to longwall dust control: Phase III report. BM 8148. Pittsburg, PA: U.S. Bureau of Mines, USBM Contract No. J0318097.

Babbitt C, Muldoon T, Rajan S, Wirch S, Burnett J [1988]. Concepts to control quartz dust exposure of the continuous miner operator and helper. Final Report. Pittsburgh, PA: U.S. Department of the Interior, Bureau of Mines, PB-89-155-402.

Bhaskar R, Gong R [1992]. Effect of foam surfactants on quartz and dust levels in continuous miner sections. *Mining Eng* 44(9):1164-1168.

Colinet JF, McClelland JJ, Jankowski RA [1991]. Fundamental investigation into the interactions and limitations of primary dust controls for continuous miners. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 203-212.

Corn M [1990]. Reflections on progress with mine dust control and dust control technology. In: Proceedings of the VIIth International Pneumoconioses Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Grayson RL, Watts CM, Yuan S, Dean J, Nutter RS [1990]. The mine management support system: a knowledge-based system for better management of underground coal mines. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., Preprint No. 90-131.

Haley WA, Dowd JJ, Turnbull LA [1952]. Modified longwall mining with a German coal planer (plow) in the Pocahontas No. 4 coal bed, Helen, W.VA. Washington, DC: U.S. Department of the Interior, Bureau of Mines, RI 4922.

Kingery DS, Doyle HN, Harris EJ, Jacobson M, Peluso RG, Shutack JB, et al. [1969]. Studies on the control of respirable coal mine dust by ventilation. Washington, DC: U.S. Department of the Interior, Bureau of Mines, Technical Progress Report 19.

Lewis BC [1990]. Longwall mining: future concerns that must be addressed. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., Preprint No. 90-47.

Mundell RL, Taylor CD [1983]. Quantitating respirable-dust levels generated during longwall-mining operations. U.S. Department of the Interior, Mining Enforcement and Safety Administration, IR 1061.

Organiscak JA, Jankowski RA, Kelly JS [1987]. Dust controls to improve quality of longwall intake air. Washington, DC: U.S. Department of the Interior, Bureau of Mines, IC 9114.

Organiscak JA, Doyle-Coombs DM, Cecala AB [1988]. Effect of shearer web depth on dust generation and methane liberation. U.S. Department of the Interior, Bureau of Mines, RI 9180.

Organisak JA, Page SJ, Jankowski RA [1990]. Sources and characteristics of quartz dust in coal mines. Washington, DC: U.S. Department of the Interior, Bureau of Mines, IC 9271.

Potts JD, Jankowski RA [1992]. Dust considerations when using belt entry air to ventilate work areas. Washington, DC: U.S. Department of the Interior, Bureau of Mines, RI 9426.

Ramani RV, Mutmansky JM, Bhaskar R, Qin J [1987]. Fundamental studies on the relationship between quartz levels in the host material and the respirable dust generated during mining. Vol. I. Final Report. Washington, DC: U. S. Department of the Interior, Bureau of Mines, Contract No. H0358031.

Shirley GA, Colinet JF, Kost JA [1985]. Dust control handbook for longwall mining operations. Washington, DC: U.S. Department of the Interior, Bureau of Mines, U.S. Bureau of Mines Contract No. J0348000.

Stein RR, Martin TW [1989]. Coal mining equipment. *Mining Eng* 41(5):338.

Suboleski SC, King RH [1990]. Estimating technology's impact on underground coal productivity. *Mining Eng* 42(10):1159-1164.

Taylor CD, Jankowski RA [1982]. How the six cleanest U.S. longwalls stay in compliance. Proceedings of the 1st Mine Ventilation Symposium, The University of Alabama, University (Tuscaloosa), AL, March 29-31, pp. 67-69.

Thimons ED, Jankowski RA, Finfinger GL [1991]. Health and safety issues related to extended longwalls. In: Hugler E, Bacho A, Karmis M, eds. Proceedings of the XXII Annual Institute on Coal Mining Health, Safety and Research. Blacksburg, VA: Department of Mining and Minerals Engineering, Virginia Polytechnic Institute and State University, pp. 101-111.

Treftis HN, Tomb TF, Carden HF [1981]. Laboratory Evaluation of Racal "Airstream" Helmet. U.S. Department of Labor, Mine Safety and Health Administration, IR 1130.

Xu L, Colinet JF, Mutmansky JM [1990]. Dust control in high-productivity coal mines: observations and strategies in high-quartz sections. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., Preprint No. 90-16.

Zimmer RA, Lueck SR [1986]. Investigation of quartz dust sources and control mechanisms on surface coal mine operations. Vol. I and Vol. II. Final Report. Pittsburgh, PA: U.S. Department of the Interior, Bureau of Mines.

Zimmer RA, Lueck SR [1987]. Optimization of overburden drill dust control systems on surface coal mines. *Int J Surface Mining* 1:155-157.

MEDICAL EVALUATION AND TESTING

ACCP/ATS [1975]. Pulmonary terms and symbols. A report of the ACCP-ATS Joint Committee on pulmonary nomenclature. *Chest* 67:583-593.

ATS (American Thoracic Society) [1969]. Standards for epidemiologic surveys in chronic respiratory disease. New York, NY: National Tuberculosis and Respiratory Disease Association.

ATS (American Thoracic Society) [1982]. Evaluation of impairment/disability secondary to respiratory disease. *Am Rev Respir Dis* 126:945-951.

ATS (American Thoracic Society) [1985]. Guidelines as to what constitutes an adverse respiratory health effect with special reference to epidemiologic studies of air pollution. *Am Rev Respir Dis* 131:666-668.

Balmes JR [1990]. Medical surveillance for pulmonary endpoints. *Occup Med: State of the Art Rev* 5(3):499-513.

Banks DE, Cheng YH, Weber SL, Ma JKH [1993]. Strategies for the treatment of pneumoconiosis. *Occup Med: State of the Art Rev* 8(1):205-232.

Bascom R, Ford E [1992]. Don't just "do spirometry." *Occup Med: State of the Art Rev* 7(2):347-363.

Batra P, Brown K [1991]. Radiology in prevention and surveillance of occupational lung disease. *Occup Med: State of the Art Rev* 6(1):81-100.

Burrows B, [1986]. Pulmonary function testing. In: Merchant JA, ed. Occupational respiratory diseases. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 86-102.

Collins LC, Willing S, Bretz R, Harty M, Lane E, Anderson WH [1993]. High-resolution CT in simple coal workers' pneumoconiosis. *Chest* 104(4):1156-1162.

Conway H, Simmons J, Talbert T [1993a]. Effects of occupational medical surveillance programs as perceived by respondents to the occupational safety and health administration's 1990-1991 survey. *J Occup Med* 35(7):687-697.

Conway H, Simmons J, Talbert T [1993b]. The occupational safety and health administration's 1990-1991 survey of occupational medical surveillance prevalence and type of current practices. *J Occup Med* 35(7):659-669, 693-697.

Conway H, Simmons J, Talbert T [1993c]. The purposes of occupational medical surveillance in US industry and related health findings. *J Occup Med* 35(7):670-686.

Elisburg D [1980]. Respiratory disability in coal miners. *JAMA* 244(19):2158.

Enright, PL [1992]. Surveillance for lung disease: quality assurance using computers and a team approach. *Occup Med: State of the Art Rev* 7(2):209-225.

Epler GR, Saber FA, Gaensler EA [1980]. Determinations of severe impairment (disability) in interstitial lung disease. *Am Rev Respir Dis* 121:647.

Fields CL, Roy TM, Dow FT, Anderson WH [1992]. Impact of arterial blood gas analysis in disability evaluation of the bituminous coal miner with simple pneumoconiosis. *J Med* 34(4):410-413.

Gaensler EA, Wright GW [1966]. Evaluation of respiratory impairment. *AMA Arch Environ Health* 12:146-189.

GAO [1990]. Black lung program: further improvements can be made in claims adjudication. Washington, DC: U.S. General Accounting Office. GAO/HRD-90-75, p. 36.

Goldman RH, Peters JM [1981]. The occupational and environmental health history. *JAMA* 246(24):2831-2836.

Guidotti TL [1979]. Coal workers' pneumoconiosis and medical aspects of coal mining. *South Med J* 72(4):456-466.

Halperin WE [1988]. Screening for occupational disease. In: Levy BS, Wegman DH, eds. *Occupational health: recognizing and preventing work-related disease*. 2nd ed. Boston, MA: Little, Brown and Company, pp. 75-86.

Hansen JE, Wasserman K [1988]. Disability evaluation. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 699-718.

Hathaway JA, [1993]. Medical surveillance: extent and effectiveness? *J Occup Med* 35(7):698-700.

Kibelstis JA [1973]. Diffusing capacity in bituminous coal miners. *Chest* 63:501-504.

Leiderman S, Halbert L, Preefer D, Forwalter D, Silver P [1983]. Review of claims filed under the Black Lung Benefits Amendments of 1981. Washington, DC: U.S. Department of Labor, Employment Standards Administration.

Lyons JP, Campbell H [1976]. Evolution of disability in coalworkers' pneumoconiosis. *Thorax* 31:527-533.

Merchant JA, Schwartz DA [1992]. Chest radiography for assessment of the pneumoconioses. In: Rom WN, ed. *Environmental and occupational medicine*. Boston, MA: Little, Brown and Company, pp. 215-225.

- Nathan SP, Lebowitz MD, Knudson RJ [1979]. Spirometric testing. *Chest* 76:384-388.
- Oboler SK, LaForce MF [1989]. The periodic physical examination in asymptomatic adults. *Ann Intern Med* 110:214-226.
- Osborne R and Fischback T [1985]. Causes of disability in employees of the mining industry: analysis of social security disability benefit awards and allowances to workers. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 85-105.
- Remy-Jardin M, DeGreef JM, Beuscart R, Voisin C, Remy J [1990]. Coal worker's pneumoconiosis: CT assessment in exposed workers and correlation with radiographic findings (1). *Radiology* 177(2):363-371.
- Roy TM, Collins LC, Snider HL, Anderson WH [1989]. Cigarette smoking and federal black lung benefits in bituminous coal miners. *J Occup Med* 31(2):98-101.
- Samet JM [1978]. A historical and epidemiologic perspective on respiratory symptoms questionnaire. *Am J Epidemiol* 108:435-446.
- Schulte PA [1991]. Contribution of biological markers to occupational health. *Am J Ind Med* 20:435-446.
- Sheppard D [1992]. Airway hyperresponsiveness: mechanisms. In: Rom WN, ed. *Environmental and occupational medicine*. Boston, MA: Little, Brown and Company.
- Solu S [1972]. Disability of coal mine workers. *Chest* 61(3):306-307.
- SSA [1986]. Social Security Administration disability evaluation under Social Security. Washington, DC: U.S. Department of Health and Human Services, Social Security Administration, SSA Publication No. 05-10089.
- SSA [1989]. Social Security Administration bulletin—annual statistical supplement, 1989. Washington, DC: U.S. Department of Health and Human Services, Social Security Administration, ISSN 0037-7910.
- Tian J, Whitmoyer B, Sneckenberger JE, Chandler CA, Jerabek MA [1991]. Computation of human upper airway volumes from tracheal acoustic impedance measurements. In: Frantz RL, Ramani RV, eds. *3rd Symposium on Respirable Dust in the Mineral Industries*. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 147-150.
- Treafitis HN, Parobeck PS [1984]. Quality assurance program for field health laboratories. Pittsburgh, PA: U.S. Department of Labor, Mine Safety and Health Administration, IR 1149.

Trout ED, Jacobson G, Moore RT, Shoub EP [1973]. Analysis of the rejection rate of chest radiographs obtained during the coal mine "black lung" program. *Diagn Radiol* 109:25-27.

WHO [1973]. Environmental and health monitoring in occupational health. WHO Technical Report Series No. 535. Geneva, Switzerland: World Health Organization.

WHO [1980]. International classification of impairments, disabilities, and handicaps. Geneva, Switzerland: World Health Organization.

CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Ames RG, Hall DS [1985]. The effects of cigarette smoking cessation on 5-year pulmonary function changes in U.S. underground coal miners. *Occup Med* 35(4):111-113.

Bates DV [1975]. The fate of the chronic bronchitis: a report of the ten-year follow-up in the Canadian department of veteran's affairs coordinated study of chronic bronchitis. *Am Rev Respir Dis* 108:1043-1065.

Cherniak RM, Raber MB [1972]. Normal standards for ventilatory function using an automated wedge spirometer. *Am Rev Respir Dis* 106:38-44.

Heppleston AG [1970]. Emphysema in relation to dust exposure. In: Shapiro A, ed. *Proceedings of the International Conference*. Johannesburg, South Africa: Oxford University Press, pp. 312-314.

Higgins MW, Keller JB, Becker MH, Landis JR, Rotman H, Weg JG, et al. [1982]. An index of risk for obstructive airways disease. *Am Rev Respir Dis* 125:144-151.

Huang X, Zalma R, Pezerat H [1994]. Factors that influence the formation and stability of hydrated ferrous sulfate in coal dusts. Possible relation to the emphysema of coal miners. *Chem Res Toxicol* 7:451-457.

Lamb D [1976]. A survey of emphysema in coal workers and the general population. *Proc R Soc Med* 69(1):14.

Leigh J, Beck R, Cole B [1993]. Emphysema, severity, pneumoconiosis histopathology and lung coal and silica content in coal miners. *J Occup Health Safety—Aust NZ* 9(2):137-146.

Lyons JP, Campbell H [1981]. Relation between progressive massive fibrosis, emphysema, and pulmonary dysfunction in coalworkers' pneumoconiosis. *Br J Ind Med* 38:125-129.

Lyons JP, Ryder RC, Seal RME, Wagner JC [1981]. Emphysema in smoking and nonsmoking coalworkers with pneumoconiosis. *Bull Eur Physiopathol Respir* 17:75-85.

Mannino D, Daniloff E, Townsend M, Petsonk E [1991]. Are atopic symptoms associated with increased airway responsiveness in coal miners? In: Frantz RL, Ramani RV, eds. *3rd Symposium*

on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., p. 145.

Mantel N, Brown C [1973]. A logistic reanalysis of Ashford and Sowden's data on respiratory symptoms in British coal miners. *Biometrics* 29:649-665.

Morgan WKC [1976]. Respiratory problems in coal miners [letters]. *JAMA* 235(13):1324-1325.

Morgan WKC, Lapp NL [1988]. Clinically important respiratory effects of dust exposure and smoking in British coal miners. *Am Rev Respir Dis* 138:1643-1644.

Morgan WKC, Lapp NL, Seaton D [1980]. Respiratory disability in coal miners. *JAMA* 243(23):2401-2404.

Pemberton J [1956]. Chronic bronchitis, emphysema, and bronchial spasm in bituminous coal workers. *Arch Ind Health* 13:529-542.

Petersen M, Castellan RM [1984]. Prevalence of chest symptoms in nonexposed blue-collar workers. *J Occup Med* 26(5):367-374.

PHS [1985]. The health consequences of smoking. Cancer and chronic lung disease in the workplace. A report of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Office on Smoking and Health, DHHS (PHS) Publication No. 85-50207.

Pride NB [1986]. Smoking, allergy and airways obstruction: revival of the "Dutch hypothesis." *Clin Allergy* 16:3-6.

Ruckley VA, Gauld SJ, Chapman JS, Davis JMG, Douglas AN, Fernie JM, et al. [1984]. Emphysema and dust exposure in a group of coal workers. *Am Rev Respir Dis* 129:528-532.

Ryder RC, Dunnill MS, Anderson JA [1971]. A quantitative study of bronchial mucous gland volume, emphysema, and smoking in a necropsy population. *J Pathol* 104:59-71.

Skillrud DM, Offord DP, Miller RD [1986]. Higher risk of lung cancer in chronic obstructive pulmonary disease. *Ann Intern Med* 105:503-507.

Snider GL [1988]. Chronic bronchitis and emphysema. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 1069-1106.

Soutar CA [1987]. Occupational bronchitis. In: Harrington M, ed. *Recent advances in occupational health*. Vol. 3. London, England: Churchill Livingstone, pp. 285-302.

SSA [1992]. Report on chronic bronchitis and emphysema. London, England: Department of Social Security, Social Security Administration, pp. 5-8.

Stenton SC, Hendrick J [1993]. Airflow obstruction and mining. *Occ Med: State of the Art Rev* 8(1):155-170.

Tweeddale PM, Alexander F, McHardy GJR [1987]. Short term variability in FEV₁ and bronchodilator responsiveness in patients with obstructive ventilatory defects. *Thorax* 42:487-490.

Ulmer WT [1975]. Chronic obstructive airway disease in pneumoconiosis in comparison to chronic obstructive airway disease in non-dust exposed workers. *Bull Physiopathol Respir* 11:415-427.

Van de Woestijne CJ [1982]. Rapid decreasing forced expiratory volume in one second or vital capacity and development of chronic airflow obstruction. *Am Rev Respir Dis* 125:553-558.

Worth G [1984]. Emphysema in coal workers (editorial). *Am J Ind Med* 6:401-403.

LUNG FUNCTION

Attfield MD [1992]. Some observations on the variation in height coefficients in prediction equations for forced vital capacity. *J Clin Epidemiol* 45(9):951-957.

Bates DV, Pham QT, Chau N, Pivoteau C, Dechoux J, Sadoul P [1985]. A longitudinal study of pulmonary function in coal miners in Lorraine, France. *Am J Ind Med* 8:21-32.

Buist AS [1988]. Smoking and other risk factors. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 1001-1029.

Burrows B, Lebowitz MD, Camilli AE, Knudson RJ [1986]. Longitudinal changes in forced expiratory volume in one second in adults. *Am Rev Respir Dis* 133:974-980.

Cockcroft A, Berry G, Cotes JE, Lyons JP [1982]. Shape of small opacities and lung function in coalworkers. *Thorax* 37:765-769.

Collins HPR, Dick JA, Bennett JG, Pern PO, Rickards MA, Thomas DJ, et al. [1988]. Irregularly shaped small shadows on chest radiographs, dust exposure, and lung function in coalworkers' pneumoconiosis. *Br J Ind Med* 45:43-55.

Cotes JE, Rossiter CE, Higgins ITT, Gilson JC [1966]. Average normal values for the forced expiratory volume in white caucasian males. *Br Med J* 1:1016-1019.

Cowie H, Lloyd MH, Soutar CA [1985]. Study of lung function data by principal components analysis. *Thorax* 40:438-443.

Crapo RO, Morris AH, Gardner RM [1981]. Reference spirometric values using techniques and equipment that meets ATS recommendations. *Am Rev Respir Dis* 123(6):659-664.

Dockery DW, Ware JH, Ferris BG Jr, Glicksberg DW, Fay ME, Spiro A III, et al. [1985]. Distribution of forced expiratory volume in one second and forced vital capacity in healthy, white, adult never-smokers in six U.S. cities. *Am Rev Respir Dis* 131:511-520.

Dockery DW, Speizer FE, Ferris BG, Ware JH, Louis TA, Spiro A [1988]. Cumulative and reversible effects of lifetime smoking and simple tests of lung function in adults. *Am Rev Respir Dis* 137:286-292.

Friedman GD, Klatshky AL, Siegelau AB [1976]. Lung function and risk of myocardial infarction and sudden cardiac death. *N Engl J Med* 294:1071-1075.

Gauld SJ, Hurley JF, Miller BG [1988]. Differences between long-term participants and non-responders in a study of coalminers' respiratory health and exposure to dust. *Ann Occup Hyg* 32(Suppl 1):545-551.

Ghio AJ, Crapo RO, Elliott CG [1990]. Reference equations used to predict pulmonary function: survey at institutions with respiratory disease training programs in the United States and Canada. *Chest* 97:400-403.

Ghio AJ, Castellan RM, Kinsley KB, Hankinson JL [1991]. Changes in forced expiratory volume in one second and peak expiratory flow rate across a work shift among unexposed blue collar workers. *Am Respir Dis* 143(6):1231-1234.

Glindmeyer HW, Diem JE, Jones RN, Weill H [1982]. Noncomparability of longitudinally and cross-sectionally determined annual change in spirometry. *Am Rev Respir Dis* 125:544-548.

Goldman HI, Becklake MR [1959]. Respiratory function tests: normal values at median altitudes and the prediction of normal results. *Am Rev Tuber* 79:457-463.

Hutchinson J [1846]. On the capacity of the lungs and on the respiratory function, with a view of establishing a precise and easy method of detecting disease by the spirometer. *Med Chir Soc* 29:138-252.

Jaakkola MS, Ernst P, Jaakkola JJK, N'gan'ga LW, Becklake MR [1991]. Effect of cigarette smoking on evolution of ventilatory lung function in young adults: an eight year longitudinal study. *Thorax* 46:907-913.

Jain BL, Patrick JM [1981]. Ventilatory function in Nigerian coal miners. *Br J Ind Med* 38:275-280.

Jones CO, Gauld S, Hurley JF, Rickmann AM [1981]. Personal differences in the breathing patterns and volumes and dust intakes of working miners. Edinburgh, Scotland: Institute of Occupational Medicine, Report No. TM/81/11.

Kannel WB, Hubert H, Lew EA [1983]. Vital capacity as a predictor of cardiovascular disease: the Framingham study. *Am Heart J* 105:311-315.

Knudson RJ, Slatin RC, Lebowitz MD, Burrows B [1976]. The maximal expiratory flow-volume curve. *Am Rev Respir Dis* 113:587.

Kory RC, Callahan R, Boren HG [1961]. The veterans administration—army cooperative study of pulmonary function. *Am J Med* 30:243-258.

Kuller LH, Ockene JK, Townsend M, Browner W, Meilan E, Wentworth DN [1989]. The epidemiology of pulmonary function and COPD mortality in the multiple risk factor intervention trial. *Am Rev Respir Dis* 140:S76-S81.

Kuller LH, Ockene J, Meilahn E, Svendsen KH [1990]. Relation of forced expiratory volume in one second (FEV₁) to lung cancer mortality in the multiple risk factor intervention trial (MRFIT). *Am J Epidemiol* 132(2):265-274.

Lapp NL, Morgan WKC [1975]. Cardio-respiratory function in United States coal workers. *Bull Physiopathol Respir* 11:527-559.

Lewitter FI, Tager IB, McGue M, Tishler PV, Speizer FE [1984]. Genetic and environmental determinants of level of pulmonary function. *Am J Epidemiol* 120(4):518-529.

Love RG [1983]. Lung function studies before and after a work shift. *Br J Ind Med* 40:153-159.

Maclaren WM, Miller BG, Hurley JF, Love RG [1994]. Reference values for lung function decline in working coalminers. Edinburgh, Scotland: Institute of Occupational Medicine, IOM Report TM/94/08.

Morgan WKC, Seaton A, Burgess DB, Lapp NL, Reger R [1972]. Lung volumes in working coal miners. *Ann N Y Acad Sci* 200:478-493.

Morris JF, Koski A, Johnson LC [1971]. Spirometric standards for healthy nonsmoking adults. *Am Rev Respir Dis* 103:57-67.

Morris JF, Temple WP, Koski A [1973]. Normal values for the ratio of one-second forced expiratory volume to forced vital capacity. *Am Rev Respir Dis* 108:1000-1003.

Muir D [1975]. Pulmonary function in miners working in British collieries: epidemiological investigations by the national coal board. *Bull Physiopath Respir* 11:403-414.

Oldham PD, Cole TJ [1983]. Estimation of the FEV. *Thorax* 38:662-667.

Olofson J, Skoogh BE, Bake B, Svardsudd K [1987]. Mortality related to smoking habits, respiratory symptoms, and lung function. *Eur J Respir Dis* 71:69-76.

Pern PO, Love RG, Wightman AJA, Soutar CA [1984]. Characteristics of coalminers who have suffered excessive loss of lung function over 10 years. *Bull Eur Physiopathol Respir* 29:487-493.

Petersen M, Hankinson J [1985]. Spirometry reference values for nonexposed blue-collar workers. *J Occup Med* 27(9):644-650.

Petersen MR, Hodous TK [1988]. Lung volume reference values for blue collar workers not exposed to occupational respiratory hazards. *J Occup Med* 30(8):626-632.

Petersen MR, Lapp NL, Amandus HE [1975]. The relationship of several ventilatory capacities and lung volumes to age, height, and weight. *J Occup Med* 17(6):355-356.

Quanjer PH, ed. [1983]. Standardized lung function testing. *Bull Eur Physiopathol Respir* 19(Suppl 5):1S-9S.

Schoenberg JB, Beck GJ, Bouhuys A [1978]. Growth and decay of pulmonary function in healthy blacks and whites. *Respir Physiol* 33:367-393.

Schwartz J [1989]. Lung function and chronic exposure to air pollution: a cross-sectional analysis of NHANES II. *Environ Res* 50:309-321.

Sorensen JB, Morris AH, Crapo RO, Gardner RM [1980]. Selection of the best spirometric values for interpretation. *Am Rev Respir Dis* 122:802-805.

Susskind H, Acevedo JC, Iwai J, Rasmussen DL, Heydinger DK, Pate HR, et al. [1988]. Heterogeneous ventilation and perfusion: a sensitive indicator of lung impairment in nonsmoking coal miners. *Eur Respir J* 1(3):232-241.

Tockman MS, Anthonisen NR, Wright EC, Donithan MG [1987]. Airways obstruction and the risk for lung cancer. *Ann Intern Med* 106:512-518.

Ullah MI, Cuddihy U, Saunders KB, Addis GJ [1983]. How many blows really make an FEV₁, FVC, or PEF_R? *Thorax* 38:113-118.

Vollmer WM, Johnson LR, McCamant LE, Buist AS [1988]. Longitudinal versus cross-sectional estimation of lung function decline—further insights. *Stat Med* 7:685-696.

Weiss W [1991]. Cigarette smoking and small irregular opacities. *Br J Ind Med* 48:841-844.

COAL WORKERS' PNEUMOCONIOSIS

Amandus HE, Lapp NL, Jacobson G, Reger RB [1976]. Significance of irregular small opacities in radiographs of coal miners in the USA. *Br J Ind Med* 33:13-17.

Attfield MD [1992]. Prevalence and incidence of coalworkers' pneumoconiosis in U.S. underground miners. Proceedings of the 9th International Symposium on Epidemiology in Occupational Health, Cincinnati, OH, September 23-25, 1992. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, pp. 467-473.

Becklake MR [1988]. Pneumoconioses. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 1556-1592.

Blanc PD, Gamsu G [1988]. The effect of cigarette smoking on the detection of small radiographic opacities in inorganic dust diseases. *J Thorac Imag* 3(4):51-56.

Cochrane AL [1976]. An epidemiologist's view of the relationship between simple pneumoconiosis and morbidity and mortality. *Proc R Soc Med* 69(1):12-14.

Cochrane AL, Moore F [1980]. A 20-year follow-up of men aged 55-64 including coal miners and foundry workers in Staveley, Derbyshire. *Br J Ind Med* 37:226-229.

Cochrane AL, Moore F, Moncrieff CB [1982]. Are coalminers, with low "risk factors" for ischaemic heart disease at greater risk of developing progressive massive fibrosis? *Br J Ind Med* 39:265-268.

Cockcroft A, Lyons JP, Andersson N, Saunders MJ [1983]. Prevalence and relation to underground exposure of radiological irregular opacities in South Wales coal workers with pneumoconiosis. *Br J Ind Med* 40(2):169-172.

Dalal NS, Jafari B, Petersen M, Green FHY, Vallyathan V [1991]. Presence of stable coal radicals in autopsied coal miners' lungs and its possible correlation to coal workers' pneumoconiosis. *Arch Environ Health* 46(6):366-372.

Davis JMG, Chapman J, Collings P, Douglas AN, Fernie J, Lamb D, et al. [1983]. Variations in the histological patterns of the lesions of coal workers' pneumoconiosis in Britain and their relationship to lung dust content. *Am Rev Respir Dis* 128:118-124.

Dodgson J, Cowie AJ, Paris I, Whittaker W [1977]. A study of the importance of dust composition in relation to pneumoconiosis in coalminers. Roxburgh place, Edinburgh, Scotland: Institute of Occupational Medicine, Report TM/77/8 (EUR. P22).

Evelo CTA, Bos RP, Borm PJA [1992]. Decreased glutathione content and glutathione S-transferase activity in red blood cells of coal miners with early stages of pneumoconiosis. *Br J Ind Med* 50:633-636.

Gonzalez A, Pedraja DeLa V, Eguidazu J, Alonso JM [1982]. The relation between pneumoconiosis and dust exposure in Spanish coal mines with comparative studies of different gravimetric dust samplers. *Br Occup Hyg Soc* 26(1-4):789-797.

Grayson RL, Andre RA, Simonyi T [1988]. Seeking the "rank factor" in CWP incidence: role of respirable dust particle purity. In: *Proceedings of the VIIth International Pneumoconiosis Conference, August 23-26, 1988, Pittsburgh, PA*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Gross P, Tuma J, deTreville RTP [1971]. Emphysema and pneumoconiosis. *Arch Environ Health* 22(2):194-199.

Heppleston AG [1985]. Silica, pneumoconiosis, and carcinoma of the lung. *Am J Ind Med* 7:285-294.

Heppleston AG [1988]. Prevalence and pathogenesis of pneumoconiosis in coal workers. *Environ Health Perspect* 78:159-170.

Johnson CC [1969]. HEW's eight-point program to combat coal miner's pneumoconiosis. *Ind Med* 38(9):14.

Key MM [1971]. Pulmonary reactions to coal dust. New York, NY: Academic Press, pp. 179-185.

Lamb D [1975]. Physiological/pathological correlations in coal workers' pneumoconiosis. *Bull Physiopathol Respir* 11:471-478.

Lapp NL, Seaton A [1971]. Pulmonary function in coal workers' pneumoconiosis. New York, NY: Academic Press, Inc., pp. 153-177.

Last JA [1985]. Changes in the collagen pathway in fibrosis. *Fundam Appl Toxicol* 5:210-218.

Leibowitz MC, Goldstein B [1987]. Some investigations into the nature and cause of massive fibrosis (MF) in the lungs of South African gold, coal, and asbestos mine workers. *Am J Ind Med* 12:129-143.

Marek K [1974]. Respiratory function impairment in coal workers' pneumoconiosis. In: Zaharski WW, ed. *Coal workers' pneumoconiosis*. Hanover, Germany: The University Press of New England, pp. 31-47.

Marek K [1985]. Lung function in different types of pneumoconiosis. *Z. Erkrank Atm Org* 164(3):270-272.

Martin JC, Daniel-Moussard H, LeBouffant L, Policard A [1972]. The role of quartz in the development of coal workers' pneumoconiosis. In: Selikoff IJ, Key MM, Lee DHK, eds. *Coal workers' pneumoconiosis*, *Ann N Y Acad Sci* 200:127-141.

McConnochie K, Green FHY, Vallyathan V, Wagner JC, Seal RME, Lyons JP [1988]. Interstitial fibrosis in coal workers--experience in Wales and West Virginia. *Ann Occup Hyg* 32(Suppl 1):553-560.

Melville AWT, Parin I, Hurley JF, Soutar CA [1979]. Pneumoconiosis, lung function and exposure to airborne dust: epidemiological research to compare responses of working coalminers with responses of ex-miners. Edinburgh, Scotland: Institute of Occupational Medicine, NTIS No. PB-84-114-081.

Miller BG, Jacobsen MG, Steele RC [1981]. Coalminers' mortality in relation to radiological category, lung function and exposure to airborne dust. Edinburgh, Scotland: Institute of Occupational Medicine, Report No. TM/81/10.

Miller BG, Kinnear AG [1989]. Pneumoconiosis in coalminers and exposure to dust of variable quartz content. Edinburgh, Scotland: Institute of Occupational Medicine, Report No. TM/88/17.

Morgan WKC, Lapp NL, Seaton A [1972]. Respiratory impairment in simple coal workers' pneumoconiosis. *JOM* 14(11):839-844.

Mosquera JA [1988]. Massive melanoptysis: a serious unrecognized complication of coal worker's pneumoconiosis. *Eur Respir J* 1:766-768.

Muir DCF [1976]. Does simple pneumoconiosis of coal-workers cause disability? *Proc Roy Soc Med* 69(1):2-4.

Musk AW, Bevan C, Campbell MJ, Cotes JE [1979]. Factors contributing to the clinical grade of breathlessness in coalworkers with pneumoconiosis. *Bull Europ Physiopath Respir* 15:343-353.

Musk AW, Cotes JE, Bevan C, Campbell MJ [1981]. Relationship between type of simple coalworkers' pneumoconiosis and lung function. A nine-year follow-up study of subjects with small rounded opacities. *Br J Ind Med* 38:313-320.

Naeye RL, Mahon JK, Dellinger WS [1971]. Rank of coal and coal workers pneumoconiosis. *Am Rev Respir Dis* 103:350-355.

Parkes WR, Williamson RGB [1976]. Coronary artery disease and coalworkers' pneumoconiosis. *Br Med J* 2:1319-1320.

Pratt PC [1968]. Role of silica in progressive massive fibrosis. *Arch Environ Health* 16:734-737.

Rasmussen DL [1972]. Patterns of physiological impairment in coal workers' pneumoconiosis. *Ann N Y Acad Sci* 200:455-464.

Rosenman KD, Zhu Z [1995]. Pneumoconiosis and associated medical conditions. *Am J Ind. Med* 27:107-113.

Roy TM, Walker JF, Snider HL, Anderson WH [1989]. Resting gas exchange in nonsmoking bituminous coal-miners with simple pneumoconiosis. *Respiration* 55:28-32.

Scarano D, Fadali MA, Lemole GM [1972]. Carcinoma of the lung and anthracosilicosis. *Chest* 62:251-254.

Seaton A [1991]. Cigarette smoking and small irregular opacities. *Br J Ind Med* 49(6):453-454.

Selikoff I, Key MM, Lee DHK [1972]. Coal workers' pneumoconiosis. *Ann N Y Acad Sci* 200:478-493.

Soutar CA, Coutts I, Parkes WR, Dodi IA, Gauld S, Castro JE, et al. [1983]. Histocompatibility antigens in coal miners with pneumoconiosis. *Br J Ind Med* 40:34-38.

Sweet DV, Crouse WE, Crable JV, Carlberg JR, Lainhart WS [1974]. The relationship of total dust, free silica, and trace metal concentrations to the occupational respiratory disease of bituminous coal miners. *Am Ind Hyg Assoc J* 35:479-488.

Vallyathan V, Althouse R, Green FHY [1990]. Coal workers' pneumoconiosis lesions and their correlation to dust load. In: Proceedings of the VIIth International Pneumoconiosis Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

vanSprundel, MPM [1990]. Pneumoconioses: the situation in developing countries. *Exp Lung Res* 16:5-13.

Viswanathan R, Prasad M, Singh H, Mitra LC, Mandall AS, Abdullah K, et al. [1973] Morbidity survey of Indian coal miners with special reference to pneumoconiosis. *Indian J Med Res* 62(1):150-164.

Walton WH, Dodgson J, Hadden GG, Jacobsen M [1975]. The effect of quartz and other non-coal dusts in coalworkers' pneumoconiosis, part 1: epidemiological studies. In: Walton WH, ed. *Inhaled particles IV*, part 2. New York, NY: Pergamon Press, pp. 669-690.

Young RC Jr, Rachal RE, Carr PG, Press HC [1992]. Patterns of coal workers' pneumoconiosis in Appalachian former coal miners. *J Natl Med Assoc* 84(1):41-48.

PARTICLE DEPOSITION AND CLEARANCE FROM LUNGS

Bartley DL, Doemeny LJ [1986]. Critique of 1985 ACGIH report on particle size-selective sampling in the workplace. *Am Ind Hyg Assoc J* 47(8):443-447.

Bellmann B, Muhle H, Creutzenberg O, Mermelstein R [1990]. Recovery behavior after dust overloading of lungs in rats. *J Aerosol Sci* 21(3):377-380.

Bohning DE, Lippmann M [1992]. Particle deposition and pulmonary defense mechanisms. In: Rom WN, ed. *Environmental and occupational medicine*. Boston, MA: Little, Brown and Company, pp. 171-182.

Brown JH, Cook KM, Ney FB, Hatch T [1950]. Influence of particle size upon the retention of particulate matter in the human being. *Am J Public Health* 40:450-458.

Carlberg JR, Crable JV, Limtiaca LP, Norris HB, Holtz JL, Mauer P, et al. [1971]. Total dust, coal, free silica, and trace metal concentrations in bituminous coal miners' lungs. *Am Ind Hyg Assoc J* 32:432-440.

Clarke SW, Pavia D [1988]. Deposition and clearance. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 313-331.

Freedman AP, Robinson SE [1988]. Noninvasive magnetopneumographic studies of lung dust retention and clearance in coal miners. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 181-186.

Freedman AP, Robinson SE, Johnston RJ [1980]. Non-invasive magnetopneumographic estimation of lung dust loads and distribution in bituminous coal workers. *J Occup Med* 22:613-618.

Freedman AP, Robinson SE, Green FHY [1982]. Magnetopneumography as a tool for the study of dust retention in the lungs. *Ann Occup Hyg* 26:319-335.

Freedman AP, Robinson SE, Street MR [1988]. Magnetopneumographic study of human alveolar clearance in health and disease. *Ann Occup Hyg* 32(Suppl 1):809-820.

Jafari B, Dalal NS, Vallyathan AV, Green FYH [1988]. Detection of organic free radicals in coal-dust exposed lung tissue and correlations with their histopathological parameters. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 223-225.

Kadas I, Szende B, Csikos A, Szendroi M [1984]. Extrapulmonary localisation of carbon and silica particles in anthracosilicosis. *Med Lav* 75(5):404-406.

Kalliomaki K, Kalliomaki P-L, Moilanen M [1988]. Magnetopneumography and its application to occupational hygiene. *Ann Occup Hyg* 32(Suppl 1):821-825.

Knight G [1985]. Definitions of alveolar dust deposition and respirable dust sampling [letters]. *Ann Occup Hyg* 29(4):526-528.

Knight G [1986]. Author's reply [letters]. *Ann Occup Hyg* 30(4):520-521.

Liddell D, Miller K [1983]. Individual susceptibility to inhaled particles. *Scand J Work Environ Health* 9:1-8.

Lippmann M [1970]. "Respirable" dust sampling. *Am Ind Hyg Assoc J* 31:138-159.

Lippmann M, Timbrell V [1990]. Particle loading in the human lung—human experience and implications for exposure limits. *J Aerosol Med* 3(Suppl 1):S155-S168.

Mastin JP, Stettler LE, Shelburne JD [1988]. Quantitative analysis of particulate burden in lung tissue. *Scanning Microsc* 2(3):1613-1629.

McCawley M, Abrons H, Lippmann M [1988]. Modelling an aerosol dispersion test for detecting early airway changes. *Am Occup Hyg* 32(Suppl 1):81-89.

McCawley M, Wu Z, Peng K, Hearl F, Dosemeci M, Chen A, et al. [1992]. Different estimations of dust exposure for mining operations. Presented at the Proceedings of the Society for Mining Engineers (SME), February 24-27, 1992, Phoenix, AR.

McCawley MA [1984]. Performance considerations for size-selective samplers. In: Particle size-selective sampling in the workplace: report of the ACGIH technical committee on air sampling procedures. *Ann Am Conf Ind Hyg* 11:23-100.

Mermelstein R, Dasenbrock C, Takenaka S, Mohr U, Kilpper R, MacKenzie J [1989]. Lung response to test toner upon 2-year inhalation exposure in rats. *Toxicol* 9:548.

Moorman WJ, Lewis TR, Wagner WD [1975]. Maximum expiratory flow-volume studies on monkeys exposed to bituminous coal dust. *J Appl Physiol* 39(3):444-448.

Moorman WJ, Hornung RW, Wagner WD [1977]. Ventilatory functions in germfree and conventional rats exposed to coal dust. *Proc Soc Exp Biol Med* 155:424-428.

Morrow PE [1977]. Clearance kinetics of inhaled particles. In: Brain JD, Proctor D, Reid L, eds. *Respiratory defense mechanisms*. New York, NY: Marcel Dekker, pp. 491-544.

Morrow PE, Gibb FR, Beiter H, Kilpper RW [1979]. Pulmonary retention of neutron-activated coal dust. *Arch Environ Health*, May/June, pp. 178-183.

Muhle H, Bellmann B, Heinrich U [1988]. Overloading of lung clearance during chronic exposure of experimental animals to particles. *Ann Occup Hyg* 32(Suppl 1):141-147.

Oberdoerster G, Green FHY, Freedman AP [1984]. Clearance of $^{59}\text{Fe}_3\text{O}_4$ particles from the lungs of rats during exposure to coal mine dust and diesel exhaust. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, NTIS No. DE-84-000-887.

Oberdorster G, Ferin J, Morse P, Corson NM, Morrow PE [1988]. Volumetric alveolar macrophage (AM) burden as a mechanism of impaired AM mediated particle clearance during chronic dust overloading of the lung. *J Aerosol Med* 1:A207.

Ogden TL [1988]. The ISO and ACGIH standardized size fractions and their relation to human deposition data. *Ann Occup Hyg* 32(Suppl 1):413-421.

Phalen RF, Stuart BO, Liroy PJ [1988]. Rationale for and implications of particle size-selective sampling. In: *Advances in air sampling*. Chelsea, MI: Lewis Publishers, Inc., pp. 3-15.

Raabe OG [1984]. Size-selective sampling criteria for thoracic and respirable mass fractions. In: Particle size-selective sampling in the workplace: report of the ACGIH Technical Committee on Air Sampling Procedures. *Ann Am Conf Ind Hyg* 11:53-65.

Rom WN, Churg A, Peapman R, Fiori C, Swyt C [1990]. Evaluation of alveolar macrophage particle burden in individuals occupationally exposed to inorganic dusts. *J Aerosol Med* 3(Suppl 1):S-43 to S-56.

Smith TJ [1985]. Development and application of a model for estimating alveolar and interstitial dust levels. *Ann Occup Hyg* 29:495-516.

Soderholm SC [1986]. Rationale of ACGIH recommendations for size-selective sampling. *Ann Occup Hyg* 30(4):518-520.

Stober W, Morrow PE, Hoover MD [1989]. Compartmental modeling of the long-term retention of insoluble particles deposited in the alveolar region of the lung. *Fund Appl Toxicol* 13:823-842.

Susskind H, Brill AB, Harold WH [1988]. Clearance of TC-99M DTPA aerosol from coal miners' lungs. *Ann Occup Hyg* 32(Suppl 1):157-169.

Tran CL, Jones AD, Donaldson K [1994]. Development of a dosimetric model for assessing the health risks associated with inhaling coalmine dusts. Edinburgh, Scotland: Institute of Occupational Medicine, Report TM/94/01.

Vincent JH [1993]. Perspectives on international standards for health-related sampling of airborne contaminants. *Appl Occup Environ Hyg* 8(4):233-238.

Vincent JH, Mark D [1988]. On the biologically-relevant sampling of airborne particles. *Ann Occup Hyg* 32(Suppl 1):423-434.

Warheit DB, Hansen JF, Hartsky MA [1991]. Physiological and pathophysiological pulmonary responses to inhaled nuisance-like or fibrogenic dusts. *Anat Rec* 231:107-118.

CELLULAR AND ANIMAL STUDIES

Bartlett GL, Pedersen AB [1988]. Effects of coal dusts and alveolar macrophages on growth of lung fibroblasts. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 142-149.

Bruch J [1987]. Value of in vivo and in vitro tests in regard to the evaluation of mixed mine dusts. In: LeBouffant L, ed. *Silicose et Pneumoconioses 'a Poussierres Mixtes*. *Colloque INSERM* 155:213-226.

Christian RT, Nelson JB, Cody TE, Larson E, Bingham E [1979]. Coal workers' pneumoconiosis: *in vitro* study of the chemical composition and particle size as causes of the toxic effects of coal. *Environ Res* 20:358-365.

Dalal NS, Vallyathan V, Leelarasamee N, Castranova V, Van Dyke K [1990]. Chemiluminescence and biologic reactivity of freshly fractured silica. In: Proceedings of the VIIth International Pneumoconioses Conference, August 23-26, 1988, Pittsburgh, PA. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Dinkelis SS, Kukharensko TA, Derr EA, Chernova TG, Myakishev IA, Abaeva SD, et al. [1981]. Relationship between pathogenicity and physicochemical properties of bituminous coal dusts. *J Hyg Epidemiol Microbiol Immunol* 25(4):393-399.

Fels AOS, Cohn ZA [1986]. The alveolar macrophage. *J Appl Physiol* 60:353-369.

Gernez-Rieux C, Tacquet A, Devulder B, Voisin C, Tonnel A, Aerts C, et al. [1972]. Experimental study of interactions between pneumoconiosis and mycobacterial infections. *Ann N Y Acad Sci* 200:106-126.

Gross P, Braun DC, deTreville RTP [1972]. The pulmonary response to coal dust. *Ann N Y Acad Sci* 200:155-165.

Hahon N, Booth JA, Green F, Lewis TR [1985]. Influenza virus infection in mice after exposure to coal dust and diesel engine emissions. *Environ Res* 37:44-60.

Heppleston AG [1986]. Cell type and response to siliceous minerals. In: Goldsmith DF, Winn DM, Shy CM, eds. *Silica, silicosis, and cancer*. New York, NY: Praeger Publishers, pp. 191-196.

Heppleston AG, Civil GW, Critchlow A [1971]. The effects of duration and intermittency of exposure on the elimination of air-borne dust from high and low rank coal mines. In Walton Wh, ed. *Inhaled particles III*. Vol. I. Surrey, England: The Gresham Press, pp. 261-270.

Heppleston AG, Kulonen E, Potila M [1984]. *In vitro* assessment of the fibrogenicity of mineral dusts. *Am J Ind Med* 6:373-386.

Last JA, Wu J, Chen J, Gelzleichter T, Sun WM, Armstrong LG [1990]. Particle-cell interactions: lung fibrosis. *J Aerosol Med* 3(Suppl 1):S61-S74.

Lehnert BE, Valdez YE, Holland LM [1985]. Pulmonary macrophages: alveolar and interstitial populations. *Exp Lung Res* 9:177-190.

Lehnert BE, Valdez YE, Tietjen GL [1989]. Alveolar macrophage-particle relationships during lung clearance. *Am J Respir Cell Mol Biol* 1:145-154.

Maly ER [1988]. Generation of free oxygen radicals from human polymorphonuclear granulocytes by cytokines from human mononuclear cells, treated with quartz dust DQ12 or coal mine dust TF-1—new aspects in pathogenesis of pneumoconiosis. *Zbl Bakt Hyg B* 187:142-165.

Morgan A, Moores SR, Holmes A, Evans JC, Evans NH, Black A [1980]. The effect of quartz, administered by intratracheal instillation, on the rat lung. The cellular response. *Environ Res* 22:1-12.

Nathan CF [1987]. Secretory products of macrophages. *J Clin Invest* 79:319-326.

Reisner KM, Last JA [1986]. Early cellular events in pulmonary fibrosis. *Exp Lung Res* 10:331-355.

Reisner MTR, Robock K [1977]. Results of epidemiological, mineralogical, and cytotoxicological studies on the pathogenicity of coalmine dusts. In: Walton WH, ed. *Inhaled particles IV*. Oxford, England: Pergamon Press, pp. 703-715.

Reisner MTR, Bruch J, Kriegseis W, Prajsnar D, Robock K, Rosmanith J, et al. [1982]. Specific harmfulness of respirable dusts from West German coal mines, VI: comparison of experimental and epidemiological results. *Ann Occup Hyg* 26(1-4):527-539.

Reisner MTR, Kuhn L, Kriegseis KW, Scharmann A, Strubel G, Rinn G [1988]. Effects of particle size of coalmine dusts in experimental anthraco-silicosis: I. influence of size of mineral composition and surface properties of dusts from same origin. *Ann Occup Hyg* 32(Suppl 1):585-592.

Reist R, Bryner K, Wearden P, Blackford J, Vrma K, Castranova V et al. [1991]. Development of a fibroblast proliferation bioassay to detect mediators of pulmonary fibrosis. In: Frantz RL, Ramani RV, eds. *3rd Symposium on Respirable Dust in the Mineral Industries*. Littleton, CO: Society for Mining, Metallurgy, and Exploration, pp. 179-181.

Reynolds HY [1988]. Bronchoalveolar lavage. In: Murray JF, Nadel JA, eds. *Respiratory medicine*. Philadelphia, PA: W.B. Saunders Company, pp. 597-610.

Richards RJ, Curtis CG [1984]. Biochemical and cellular mechanisms of dust-induced lung fibrosis. *Environ Health Perspect* 55:393-416.

Robertson A, Bolton RE, Chapman JS, Davis JMG, Dodgson J, Gormley IP, et al. [1984]. Animal inhalation experiments to investigate the significance of high and low percentage concentrations of quartz in coalmine dusts in relation to epidemiology and other biological tests. Edinburgh, Scotland: Institute of Occupational Medicine, Report TM/84/5.

Robock K, Bauer HD [1990]. Investigations into the specific fibrogenicity of mine dusts in hardcoal mines of countries in the European community. Part I. In: *Proceedings of the VIIth International Pneumoconiosis Conference, August 23-26, 1988, Pittsburgh, PA*. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease

Control, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 90-108.

Robock K, Reisner MTR [1982]. Specific harmfulness of respirable dusts from West German coal mines. I. Results of cell tests. *Ann Occup Hyg* 26(1-4):473-479.

Rosmanith J, Weller W, Reisner MTR, Kuhn L, Strubel G [1988]. Effects of particle size of coalmine dusts in experimental anthraco-silicosis: II. retention and penetration of the dust. *Ann Occup Hyg* 32(Suppl 1):593-601.

Schreider JP, Culbertson MR, Raabe OG [1985]. Comparative pulmonary fibrogenic potential of selected particles. *Environ Res* 38:256-274.

Shi X, Xianglin S, Dalal NS [1991]. Free radical generation by metal ions adsorbed on mineral dust: a mechanism for chronic lung injury. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 307-311.

Szymczykiwicz KE [1982]. The influence of dust originated in different coal mines on pneumoconiotic changes in white rats. *Medycyna Pracy* 23(4):21-34.

Vallyathan V, Kang JH, Van Dyke K, Dalal NS, Castranova V [1991]. Freshly fractured silica exerts greater toxicity to lung macrophages than aged dust: organosilan (prosil 28) coating markedly reduces dust toxicity to cells. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 117-119.

Vuorio EI, Makela JK, Vuorio TK, Pool A, Wagner JC [1989]. Characterization of excessive collagen production during development of pulmonary fibrosis induced by chronic silica inhalation in rats. *Br J Exp Pathol* 70:305-315.

Wallace WE, Keane MJ [1988]. Characterization of surface properties affecting the activity of the "free silica" fraction of respirable dusts. Final report prepared for the Bureau of Mines, Pittsburgh, PA, by the National Institute for Occupational Safety and Health, Morgantown, WV, pp. 48-70. NTIS No. PB-89-142-988.

Wallace WE, Keane MJ [1991]. Pulmonary surfactant interactions with respirable particles. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 127-130.

Wallace WE, Vallyathan V, Keane MJ, Robinson V [1985]. In vitro biologic toxicity of native and surface-modified silica and kaolin. *J Toxicol Environ Health* 16:415-424.

Wallace WE, Keane MJ, Mike PS, Hill CA, Vallyathan V [1989]. Mineral surface-specific differences in the adsorption and enzymatic removal of surfactant and their correlation with

cytotoxicity. In: Mossman BT, Gegin RO, eds. *Effects of mineral dusts on cells*. NATO ASI Series, Vol. H30. Heidelberg, Germany: Springer-Verlag, pp. 49-56.

Waseem M, Bansal SK, Gupta GSD, Kaw JL [1982]. Cytotoxic and hemolytic action of coal-quartz mixtures. *Arch Environ Health* 37(6):352-357.

Wiedermann CJ, Adamson IYR, Pert CB, Bowden DH [1988]. Enhanced secretion of immunoreactive bombesin by alveolar macrophages exposed to silica. *J Leukocyte Biol* 43:99-103.

SILICA EXPOSURES AND SILICOSIS

Ampian SG, Virta RL [1992]. *Crystalline silica overview: occurrence and analysis*. Washington, DC: U.S. Department of the Interior, Bureau of Mines, IC 9317.

Coffin DL, Palekar LD [1986]. Tumorigenesis and cytotoxicity of silica. In: Goldsmith DF, Winn DM, Shy CM, eds. *Silica, silicosis, and cancer*. New York, NY: Praeger Publishers, pp. 157-165.

Fubini B, Giamello E, Volante M, Bolis V [1990]. Chemical functionalities at the silica surface determining its reactivity when inhaled. Formation and reactivity of surface radicals. *Toxicol Ind Health* 6(6):571-598.

Greskevitch MF, Turk AR, Dieffenbach AL, Roman JM, Groce DW, Hearl FJ [1992]. Quartz analyses of the bulk dust samples collected by the National Occupational Health Survey of Mining. *Appl Occup Environ Hyg* 7(8):527-531.

IARC [1987]. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans: Silica and some silicates. Vol. 42. Lyon, France: World Health Organization, International Agency for Research on Cancer, pp. 33-143.

Lemen RA, Dunnom DD, Wagner WD, Mazzuckelli LF [1986]. Recommended standards for occupational exposure to silica. In: Goldsmith DF, Winn DM, Shy CM, eds. *Silica, Silicosis, and Cancer*. New York, NY: Praeger Publishers, pp. 505-509.

Meloy T [1988]. A working hypothesis on how silica and silica surface may cause silicosis and CWP. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 15-18.

Miller BG, Kinnear AG [1988]. Pneumoconiosis in coalminers and exposure to dust of variable quartz content. Edinburgh, U.K.: Institute of Occupational Medicine, Report No. TM/88/17.

Pandurangi RS, Seehra MS [1991]. Hemolytic ability of different forms of silica and role of the surface silanol species. In: Frantz RL, Ramani RV, eds. *3rd Symposium on Respirable Dust in the Mineral Industries*. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 121-125.

Seaton A, Addison J, Davis JMG, Hurley J, McGovern B, Miller B [1987]. The toxic effects of silica. Edinburgh, Scotland: Institute of Occupational Medicine, Report No. TM/87/13.

Shulman SA, Groff JH, Abell MT [1992]. Performance of laboratories measuring silica in the proficiency analytical testing program. *Am Ind Hyg Assoc J* 53(1):49-56.

Stettler LE, Tucker JH, Riley RD, Platek SF, Mastin JP, Green FHY, et al. [1988]. Silica lung particle size determination in coal miner lungs. Washington, DC: U.S. Department of the Interior, Bureau of Mines, NTIS No. PB-89-142-988, pp. 19-47.

Theriault G, Peters JM, Johnson WM [1974]. Pulmonary function and roentgenographic changes in granite dust exposure. *Arch Environ Health* 28:23-27.

Tomb TF, Peluso RG, Parobeck PS [1986]. Quartz in United States coal mines. *Ann Am Conf Gov Ind Hyg* 14:513-519.

Tomb TF, Parobeck PS, Gero AJ [1988]. MSHA's revised quartz enforcement program. In: Frantz RL, Ramani RV, eds. *Respirable dust in the mineral industries: health effects, characterization and control*. University Park, PA: The Pennsylvania State University, pp. 9-14.

Valiante DJ, Rosenman KD [1989]. Does silicosis still occur? *J Am Med Assoc* 262(21):3003-3007.

Wallace WE Jr., Keane MJ, Vallyathan V, Hathaway P, Regad ED, Castranova V, et al. [1988]. Suppression of inhaled particle cytotoxicity by pulmonary surfactant and retoxification by phospholipase: distinguishing properties of quartz and kaolin. *Ann Occup Hyg* 32(Suppl 1):291-298.

Weber K, Tucker J, Stettler L, Wallace W [1988]. Final report: research on health effects and particle characterization to improve silica dust control in underground coal mines. Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, NTIS No. PB-89-142-988.

DIESEL EXPOSURES, CONTROLS AND HEALTH EFFECTS

Cantrell BK, Rubow KL [1991]. Development of personal diesel aerosol sampler design and performance criteria. *Mining Eng* 43(2):232-236.

Chan LM, Carlson DH, Johnson JH [1992]. Deterioration criteria for servicing mine diesel vehicles based on tailpipe pollutant concentration data. *Mining Eng* 44(12):1472-1476.

Marple VA, Rubow KL, Tao Y [1991]. A personal diesel exhaust aerosol sampler. In: Frantz RL, Ramani RV, eds. *3rd Symposium on Respirable Dust in the Mineral Industries*. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 73-81.

Potts JD, Divers EF [1991]. Powered dust-filtering helmet reduces exposure to diesel-size particulate. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 105-107.

Smith R, Stayner L [1991]. An exploratory assessment of the risk of lung cancer associated with exposure to diesel exhaust based on a study in rats. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 29-49.

Tomb TF, Haney RA, Gero AJ, Parobeck PS [1991]. Evaluation of two diesel particulate sampling devices. In: Frantz RL, Ramani RV, eds. 3rd Symposium on Respirable Dust in the Mineral Industries. Littleton, CO: Society for Mining, Metallurgy, and Exploration, Inc., pp. 83-95.

Vostal JJ [1986]. Factors limiting the evidence for chemical carcinogenicity of diesel emissions in long-term inhalation experiments. In: Ishinishi N, Koizumi A, McClellan RO, Stober W, eds. Carcinogenic and mutagenic effects of diesel engine exhaust. Amsterdam, The Netherlands: Elsevier Science.

Wallace WE, Keane MJ, Hill CA, Xu J, Ong T-M [1987]. Mutagenicity of diesel exhaust particles and oil shale particles dispersed in lecithin surfactant. *J Tox Environ Health* 21:163-171.

Wallace W, Keane M, Xing S, Harrison J, Gautam M, Ong T [1989]. Mutagenicity of diesel exhaust soot dispersed in phospholipid surfactants. *Zbl Hyg* 189:62-101.

HEALTH AND SAFETY

Ary IS [1989]. Keynote address, VPI conference. In: Sutherland WH, Forshey DR, Karmis M, eds. Proceedings of the Twentieth Annual Institute on Coal Mining Health, Safety and Research, Milestones in Health, Safety and Research, Blacksburg, VA, August 29-31, p. 25.

Block DL [1988]. Occupational health and safety programs in the workplace. 2nd ed. In: Levy BS, Wegman DH, eds. Occupational health: recognizing and preventing work-related disease. Boston, MA: Little, Brown, and Co., pp. 75-86.

Breuer H, Reisner MTR [1988]. Criteria for long-term dust standards on the basis of personal dust exposure records. *Ann Occup Hyg* 32(Suppl 1):523-527.

Bridbord K, Costello J, Gamble J, Groce D, Hutchison M, Jones W, et al. [1979]. Occupational safety and health implications of increased coal utilization. *Environ Health Perspect* 33:285-302.

Daniel JH [1993]. Ensuring a healthy environment for miners. *Minerals Today*, August, pp. 14-18.

Jacobsen M [1972]. The basis for the new coal dust standards. *The Mining Engineer* 131:269-278.

Key MM [1972]. Health standards and standard setting in the United States. *Ann N Y Acad Sci* 200:707-716.

Kraus JF [1985]. Fatal and nonfatal injuries in occupational settings: a review. *Annu Rev Public Health* 6:403-418.

Rappaport SM [1984]. The rules of the game: an analysis of OSHA's enforcement strategy. *Am J Ind Med* 6:291-303.

Weeks JL [1991a]. Occupational health and safety regulation in the coal mining industry: public health at the workplace. *Anu Rev Publ Health* 12:195-207.

Weeks JL [1991b]. Tampering with dust samples in coal mines (again). *Am J Ind Med* 20:141-144.

Weeks JL [1993]. From explosions to black lung: a history of efforts to control coal mine dust. *Occup Med: State of the Art Rev* 8(1):1-17.

Weeks JL, Fox M [1983]. Fatality rates and regulatory policies in bituminous coal mining, United States. *Am J Public Health* 73(11):1278-1280.

MISCELLANEOUS

Abraham JL [1992]. Environmental pathology of the lung. In: Rom WN, ed. *Environmental and occupational medicine*. Boston, MA: Little, Brown and Company.

Barth, PS [1987]. *The tragedy of black lung*. Kalamazoo, MI: W.E. Upjohn Institute for Employment Research.

Butani SJ, Bartholomew AM [1988]. *Characterization of the 1986 coal mining workforce*. Washington, DC: U.S. Department of the Interior, Bureau of Mines, IC 9192.

Coggbon D, Barker DJP, Cole RB [1990]. Stomach cancer and work in dusty industries. *B J Ind Med* 47:298-301.

Cohrssen JJ, Covello VT [1989]. Risk assessment. In: *Risk analysis: a guide to principles and methods for analyzing health and environmental risks*. Springfield, VA: U.S. Department of Commerce, The National Technical Information Service, NTIS No. PB-137-772.

Falk HL, Jurgelski W [1979]. Health effects of coal mining and combustion: carcinogens and cofactors. *Environ Health Perspect* 33:203-226.

Fiserova-Bergerova V [1990]. Application of toxicokinetic models to establish biological exposure indicators. *Ann Occup Hyg* 34(6):639-651.

Fox AJ, Collier PF [1976]. Low mortality rates in industrial cohort studies due to selection for work and survival in the industry. *Br J Prev Soc Med* 30(4):225-230.

Hill AB [1965]. The environment and disease: association or causation? *Proc Roy Soc Med* 58:295-300.

Lee HB [1969]. *Bloodletting in Appalachia*. Morgantown, WV: West Virginia University Press.

McFeters JJ [1981]. *Industrial hygiene study of TVA workers in coal-fired power plants*. Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health. Report No. PB-83-139-295.