



# Estimated Number of Adults and Adolescents Living with HIV/AIDS, by Sex, 2003–2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays. Age as of end of year.




# Estimated Number of Persons Living with HIV/AIDS, by Race/Ethnicity, 2003–2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.





# Estimated Number of Asians/Pacific Islanders and of American Indians/Alaska Natives Living with HIV/AIDS 2003–2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.



# Proportion of HIV/AIDS Cases among Adults and Adolescents, by Sex and Transmission Category 2006—33 States




- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU
- High-risk heterosexual contact\*
- Other/not identified †

Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.  
 \*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.  
 † Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




# Proportion of HIV/AIDS Cases among Adults and Adolescents, by Race/Ethnicity, 2003–2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.



# Proportion of HIV/AIDS Cases among Adults and Adolescents, by Transmission Category 2003–2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.  
 \*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.  
 † Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Estimated Number and Proportion of HIV/AIDS Cases among Adults and Adolescents, by Transmission Category 2006—33 States

<u>Transmission category</u>	<u>No.</u>	<u>%</u>
Male-to-male sexual contact	17,465	50
Injection drug use (IDU)	4,728	13
Male-to-male sexual contact and IDU	1,180	3
High-risk heterosexual contact*	11,584	33
<u>Other/not identified</u> <sup>†</sup>	<u>223</u>	1
Total	35,180	

Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\* Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

† Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



# Estimated Number and Proportion of HIV/AIDS Cases Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity 2006—33 States

<u>Race/ethnicity</u>	<u>Cases</u>	<u>%</u>
White, not Hispanic	7,158	41
Black, not Hispanic	6,589	38
Hispanic	3,330	19
Asian/Pacific Islander	220	1
American Indian/ Alaska Native	91	1
<u>Total*</u>	<u>17,465</u>	



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\* Includes 77 persons of unknown race or multiple races.





# Estimated Number and Proportion of HIV/AIDS Cases among Adults and Adolescents Attributed to High-Risk Heterosexual Contact\*, by Race/Ethnicity, 2006—33 States

<u>Race/Ethnicity</u>	<u>Cases</u>	<u>%</u>
White, not Hispanic	1,822	16
Black, not Hispanic	7,606	66
Hispanic	1,953	17
Asian/Pacific Islander	116	1
American Indian/ Alaska Native	36	<1
<u>Total†</u>	<u>11,584</u>	

Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\* Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

† Includes 51 persons of unknown race or multiple races.



# Estimated Number and Proportion of HIV/AIDS Cases Attributed to Male-to-Male Sexual Contact and Injection Drug Use, by Race/Ethnicity, 2006—33 States

<u>Race/Ethnicity</u>	<u>Cases</u>	<u>%</u>
White, not Hispanic	498	42
Black, not Hispanic	440	37
Hispanic	216	18
Asian/Pacific Islander	13	1
American Indian/ Alaska Native	9	1
<u>Total*</u>	<u>1,180</u>	



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\* Total includes 4 persons of unknown race or multiple races.



# Estimated Number and Proportion of HIV/AIDS Cases Attributed to Injection Drug Use, by Race/Ethnicity 2006—33 States

<u>Race/Ethnicity</u>	<u>Cases</u>	<u>%</u>
White, not Hispanic	1,210	26
Black, not Hispanic	2,512	53
Hispanic	912	19
Asian/Pacific Islander	44	1
American Indian/ Alaska Native	27	1
<u>Total*</u>	<u>4,728</u>	



Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

\*Includes 23 persons of unknown race or multiple races.



# Estimated Number of HIV/AIDS Cases and Rates for Male Adults and Adolescents, by Race/Ethnicity 2006—33 States

<u>Race/ethnicity</u>	<u>Cases</u>	<u>Rate (cases per 100,000 population)</u>
White, not Hispanic	9,078	16.7
Black, not Hispanic	11,230	119.1
Hispanic	5,058	50.9
Asian/Pacific Islander	318	13.5
American Indian/ Alaska Native	129	17.7
<u>Total*</u>	<u>25,928</u>	<u>33.8</u>

Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

\* Includes 115 male adults and adolescents of unknown race or multiple races.



# Estimated Number of HIV/AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2006—33 States


<u>Race/ethnicity</u>	<u>Cases</u>	<u>Rate (cases per 100,000 population)</u>
White, not Hispanic	1,664	2.9
Black, not Hispanic	6,033	56.2
Hispanic	1,400	15.1
Asian/Pacific Islander	79	3.2
<u>American Indian/ Alaska Native</u>	<u>35</u>	<u>4.6</u>
Total*	9,252	11.5

Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

\* Includes 41 female adults and adolescents of unknown race or multiple races.




# Estimated Prevalence Rates for Adults and Adolescents Living with HIV Infection (not AIDS), 2006—33 States and 5 U.S. Dependent Areas



Note. Data from 33 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.




# Estimated Prevalence Rates for Children <13 Years of Age Living with HIV Infection (not AIDS), 2006—33 States and 5 U.S. Dependent Areas



Note. Data from 33 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.



# Year of Initiation of Confidential Name-Based HIV Case Surveillance\* as of December 2007




\* As of December 2007 there were 54 areas with confidential name-based HIV infection reporting.





# Reported Cases of HIV Infection (not AIDS), 2006—45 States and 5 U.S. Dependent Areas

N = 52,878\*




Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.  
 \* Includes 132 persons who were residents of areas without HIV infection surveillance but who were reported by areas with HIV infection surveillance. Includes 415 persons whose area of residence is unknown.



# Adolescents 13–19 Years of Age Reported to be Living with HIV Infection (not AIDS), 2006—45 States and 5 U.S. Dependent Areas

N=3,808\*



Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.  
 \* Includes 14 residents of areas without HIV infection surveillance but who were reported from areas with HIV infection surveillance. Includes 24 persons whose state of residence is unknown or missing.



# Reported Cases of HIV Infection (not AIDS), by Age Group at Diagnosis, Cumulative through 2006—45 States and 5 U.S. Dependent Areas

## HIV Infection (not AIDS)

<u>Age (years)</u>	<u>No.</u>	<u>%</u>
<13	5,314	2
13–14	473	<1
15–24	45,514	16
25–34	98,844	34
35–44	88,270	31
45–54	36,916	13
55–64	9,839	3
≥65	2,784	<1
<u>Total</u>	<u>287,954</u>	




Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.



# Reported Cases of HIV Infection (not AIDS) in Children <13 Years of Age at Diagnosis, 2006—45 States and 5 U.S. Dependent Areas


N=566\*



Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.  
 \* Includes 4 persons who were residents of areas without HIV infection surveillance but who were reported by areas with HIV infection surveillance. Includes 8 persons whose area of residence is unknown or missing.



# Reported Cases of HIV Infection (not AIDS), by Age Group and Sex, Cumulative through 2006—45 States and 5 U.S. Dependent Areas




Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.

\*Excludes 6 persons of unknown sex.



# Estimated Diagnosis Rates of HIV/AIDS for Adults and Adolescents, 2006—33 States



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

