

**DAIRY CATTLE: Agricultural Chemical Use
Wisconsin, 2001 1/**

Agricultural chemical	Rate per head per application	Rate per head per year	Total applied
Insecticides:			1,000 lbs.
Coumaphos (Co-Ral)	1.1	10.1	1.4
Cyfluthrin (Countdown)	0.1	0.4	0.1
Dichlorvos	0.5	16.3	3.4
Dipropyl isocinchomeronate (Tox-O-Wik)	0.5	22.3	1.4
Doramectin (Dectomax)	0.4	0.7	*
Eprinomectin (Eprinex)	0.3	0.4	0.3
Ivermectin (Ivomec)	0.3	0.4	0.2
Moxidectin (Cydectin)	0.2	0.3	0.1
N-octy-bicycloheptene dicarbo dicarbo	0.2	8.2	0.7
Permethrin	0.6	5.1	4.4
Piperonyl butoxide	0.3	7.2	4.0
Pyrethrins	0.0	1.2	0.5

* Total applied less than 50 pounds. 1/Insufficient reports to publish data for carbaryl, chlorpyrifos, clorsulon, crotoxyphos, dimethoate, famphur, fenvalerate, malathion, naled, tetrachlorvinphos, toxaphene, and xylene.

**DAIRY CATTLE FACILITIES: Agricultural Chemical Use
Wisconsin, 2001 1/**

Agricultural chemical	Total applied
	1,000 pounds
Insecticides:	
Cyfluthrin (Countdown)	0.5
Dichlorvos	5.1
Dimethoate (Cygon)	3.3
Methomyl (Die Fly)	0.1
N-octy-bicycloheptene dicarbo	0.3
Permethrin	6.3
Piperonyl butoxide	5.4
Pyrethrins	0.9

1/Insufficient reports to publish data for bomyl, chlorpyrifos, dipropyl isocinchomeronate, lambda-cyhalothrin, malathion, moxidectin, petroleum distillate, ronnel, sulfur, and tetrachlorvinphos.

**AGRICULTURAL CHEMICAL USE
Total Amount Applied,
Program States and Total, 2001**

State	Dairy cattle	Dairy cattle facilities
1,000 pounds		
California	13.3	14.0
Colorado	8.0	2.8
Florida	3.7	0.3
Idaho	2.9	2.9
Illinois	2.4	0.8
Indiana	7.7	7.2
Iowa	9.4	4.6
Kentucky	2.0	6.3
Michigan	2.1	2.9
Minnesota	19.6	14.0
Missouri	3.4	0.7
New Mexico	13.2	3.7
New York	11.4	2.8
Ohio	2.2	2.4
Pennsylvania	7.4	5.8
Tennessee	8.0	0.9
Texas	9.4	2.2
Vermont	6.0	2.3
Virginia	2.1	0.2
Washington	5.5	0.4
Wisconsin	22.1	23.2
Total	161.8	100.1