
2003 California Pistachio Objective Measurement Report

Released: July 31, 2003 12:00 NOON PDT

2003 PISTACHIO PRODUCTION FORECAST AT 180 MILLION POUNDS

California pistachio production for 2003 is forecast at 180 million pounds. The 80 percent confidence interval is from 150 to 210 million pounds. This means that the results of our sampling procedures will encompass the true mean 80 percent of the time. This forecast is based on an objective measurement survey conducted by the California Agricultural Statistics Service (CASS) under the sponsorship of the California Pistachio Commission (CPC). The survey collects data such as clusters per tree, nuts per cluster and percent of bearing trees.

Pistachios are an alternate bearing crop with 2003 representing a low year of production. Following the record 2002 crop, the 2003 season got off to a wet start. Rains during key bloom periods resulted in a bad bloom and spotty set. The arrival of warm weather in late May and early June, however, promoted good growth and development. To date, the overall crop is looking better than originally expected.



SURVEY PROCEDURES

Forecasting research on California's pistachio crop began in 1980 by CASS under the sponsorship of the CPC. The Pistachio Objective Measurement (O. M.) Survey uses randomly selected trees throughout the State. These trees are used to gather information on the total number of clusters, nuts within clusters, frequency of blank nuts, as well as weight and size information. This Survey began in 1982 to meet grower and processor needs for accurate production data. An objective measurement survey was not conducted in 1993. CALIFORNIA AGRICULTURAL STATISTICS SERVICE

In 2003 the O.M. survey was modified in order to provide an earlier production forecast to growers and processors. The survey was conducted primarily during July, one month earlier than in previous years, therefore, eliminating much of the sizing data.

The Pistachio O. M. Survey procedures consist of sampling 1,300 randomly selected trees. For each tree, the crosssectional area (CSA) for each primary branch is recorded and a primary branch (path) is randomly selected to obtain additional data. Along this path, CSA measurements are recorded at every branching fork and one branch at each fork is randomly selected until a terminal branch is reached (where only one branch at a fork is greater than 0.9 square inches). Along the path, the number of clusters is recorded. The number of clusters also is recorded for the terminal branch. In addition, randomly selected clusters from the terminal branch are picked so measurements can be obtained. The number of clusters collected from the random path is expanded according to the corresponding branch sizes in order to estimate the total number of clusters on the sample tree. The estimated number of clusters for each sample tree is combined to estimate the number of clusters by rootstock, county, and state. (Starting in 1998, two random paths were performed for each tree.)

Field staff also obtain a "Ten Tree Count" of bearing (female) and pollinator (male) trees. From these counts, the "Estimated Percent Of All Spaces That Contain Bearing Trees" and the "Estimated Percent Of All Spaces That Contain Pollinators" are determined. A tree may be classified as too young or too diseased to be counted as a bearing or pollinator tree.

The clusters are sent to a sizing station where field staff count the nuts on each cluster and obtain in-hull cross-suture width.

In previous years the numbers of filled and blank nuts per cluster were determined, as well as in-hull weight, kernel weight, kernel cross-suture width, kernel suture width, and kernel length measurements for each nut on the cluster.

THE 2003 PISTACHIO OBJECTIVE MEASUREMENT SURVEY

The Pistachio O. M. Survey was completed by July 25. All samplers are employees of the National Association of State Departments of Agriculture and work in cooperation with CASS. Equipment and supplies were furnished, and survey procedures were discussed at training schools prior to the survey. Supervisors also trained enumerators on an individual basis. Quality control checks were made by all field supervisors to assure uniform procedures were followed Statewide.

THE SAMPLE

Data were collected from 636 samples. These samples consist of two trees per sample and two random paths per tree (i.e., 1,272 trees and 2,544 random paths). This year, 219, 382 and 13 samples were obtained from trees with Atlantica, Pioneer Gold I and Pioneer Gold II rootstocks, respectively. Data for some samples could not be obtained due to wet or pulled orchards, or other conditions that prevented the field staff from entering an orchard.


SURVEY DATA


For 2003, the overall average number of clusters per tree decreased 42 percent to 461 from the previous year. The average cluster per tree for Atlantica (616 clusters per tree), Pioneer Gold I (372 clusters per tree) and Pioneer Gold II (606 clusters per tree) all decreased from the previous year by 47 percent, 38 percent and 43 percent, respectively. This is in contrast to 2002 during which all rootstocks increased from the previous year.


The average number of nuts per cluster increased considerably when compared to 2002, from 13.8 to 20.6 nuts per cluster.

The percentage of female trees in California's bearing pistachio orchards (93.4%) decreased slightly from last year (94.0%).

In-hull cross suture measurements decreased from 14.46 millimeters in 2002 to 14.22 in 2003.







	i	i	PISTACHIO OBJECTIVE MEASURE					MENT SURVEY DATA, 1993-2003 1/							1
		Samples	Estimated Average	Of All	Spaces Contain		Count Da	ita	In-Hull Data <u>3</u> /			Kernel Data <u>3</u> /			
Area	Year	Completed <u>2</u> /	Number Of Clusters Per Tree	Bearing Trees	Pollinators	Nuts Per Cluster (Filled and Blank)	Percent Of Nuts Filled	Est. Total Number Of Filled Nuts Per Tree	Weight Per Nut (Includes Blanks)	Weight Per Nut (Filled)	In-Hull Cross Suture	Average Weight Per Kernel	Suture	Cross Suture	Length
Kern	1993	477													
	1994 1995	177 215	984 1,000	87.4 90.5	5.4 5.0	12.1 9.8	79.5 77.7	9,460 7,651	3.04 3.17	 3.29	15.34 15.50	0.986 0.995	10.42 10.22	9.71 9.96	17.10 16.23
	1995	215	702	90.5 89.1	5.0	9.8 11.7	58.8	4,848	2.44	2.60	14.49	0.995	9.51	9.90 9.14	15.73
	1997	236	1,200	89.7	5.1	10.4	76.3	9,563	2.82	3.00	14.83	0.947	10.77	9.62	16.67
	1998	251	1,102	91.5	5.4	13.9	76.2	11,700	2.87	3.07	15.35	0.897	10.35	9.21	16.27
	1999	239	479	92.7	5.2	11.2	66.8	3,589	2.86	3.09	15.16	0.971	10.45	9.88	16.74
	2000	225	1,217	93.5	4.7	13.1	68.3	10,771	2.61	2.85	14.60	0.885	9.96	9.24	16.18
	2001 2002	246 250	751 1,167	93.8 94.9	4.3 4.2	12.7 13.9	68.5 69.1	6,543 11,233	2.97 2.72	3.22 2.94	15.31 14.72	1.045 0.903	10.50 10.27	9.93 9.50	16.58 16.36
	2002	250	379	94.9 94.8	4.2	25.7			2.12	2.94	14.72	0.903		9.50	
Kings	1993														
-	1994	49	776	86.5	3.5	13.4	78.5	8,133	3.08		15.10	1.047	11.08	10.34	17.72
	1995	55	1,165	90.7	3.7	10.8	79.6	10,050	3.09	3.35	15.68	0.983	10.50	10.35	16.86
	1996 1997	39 55	359 1,319	91.0 92.9	5.1 3.8	20.0 11.8	62.5 68.3	4,485 10,659	2.56 2.69	2.76 2.76	14.28 14.62	0.826 0.885	9.75 10.33	9.47 9.76	16.15 16.51
	1997	58	828	92.9	4.1	15.6	76.4	9,899	2.09	3.14	14.02	0.885	10.50	9.96	17.20
	1999	51	626	93.3	2.4	13.9	71.0	6,179	2.79	3.32	16.06	0.912	10.25	9.74	17.00
	2000	53	995	94.9	2.8	11.0	71.5	7,834	2.34	2.78	15.56	0.713	9.59	8.95	15.61
	2001	54	1,159	97.9	1.8	10.0	69.5	8,062	2.87	3.13	16.39	0.993	10.81	10.19	17.00
	2002 2003	60	1,048	95.5	2.6	16.5	71.7	12,375	2.64	2.85	14.47	0.868	10.13	9.47	16.39
Madera	1993	66	580	95.2	3.5	15.1 					14.12				
Maacra	1994	132	673	87.2	7.1	10.8	80.8	5,895	2.70		14.67	0.872	10.36	9.46	16.49
	1995	147	850	88.0	6.2	7.8	81.5	5,385	2.99	3.19	15.55	0.896	10.58	9.77	16.21
	1996	162	932	88.3	5.7	7.9	74.2	5,464	2.54	2.69	15.47	0.751	9.82	8.71	15.11
	1997	162	715	89.3	5.5	8.1	78.4	4,527	2.80	2.92	16.38	0.995	11.15	9.98	17.48
	1998 1999	136 144	634 637	89.7 87.1	3.9 7.7	13.0 11.5	79.1 71.4	6,511 5,232	2.74 2.87	2.93 3.03	14.85 15.22	0.672 0.911	9.97 9.86	9.99 9.86	16.10 16.71
	2000	116	670	92.6	4.7	13.7	71.7	6,567	2.66	2.93	14.73	0.939	10.28	9.86	16.84
	2001	158	570	88.6	8.3	12.6	71.3	5,137	2.89	3.12	15.34	1.046	10.11	9.73	16.30
	2002	132	1,039	91.8	6.6	14.1	73.7	10,861	2.59	2.55	13.88	0.850	9.79	8.90	15.90
Merced	2003 1993	144	555	89.5	7.5	<u>19.0</u>					13.98				
Merceu	1993	21	525	84.3	7.1	14.5	84.4	6,443	2.61		14.35	0.848	10.31	9.39	16.42
	1995	34	753	88.2	5.7	11.2	79.8	6,745	2.91	3.11	15.36	0.852	9.93	9.46	16.44
	1996	29	802	87.2	6.8	10.9	70.6	6,195	2.66	2.86	16.47	0.811	10.23	9.51	16.18
	1997	47	953	85.6	9.3	12.2	80.5	9,380	2.74	2.82	14.23	0.906	9.69	8.89	14.41
	1998 1999	44 35	655 1,087	87.8 86.2	7.7 6.5	14.9 11.0	76.3 71.1	7,434 8,490	2.98 2.68	3.16 2.93	15.19 14.60	0.859 0.944	10.83 10.04	9.70 9.79	16.96 16.67
	2000	36	1,087	86.9	6.4	15.0	83.9	12,890	2.60	2.95	14.00	0.944	10.04	9.79	15.63
	2001	30	722	88.7	6.3	11.2	67.6	5,455	2.70	3.16	15.86	1.032	10.32	10.90	16.76
	2002	31	1,010	90.8	5.8	14.5	74.8	10,937	2.35	2.60	15.92	0.849	10.17	9.05	16.13
Tulore	2003	31	593	90.0	4.9	15.6					14.11				
Tulare	1993 1994	49	 941	 90.4	 5.1	 11.8	 86.5	 9,585	2.74		 14.34	0.866	 10.18	9.63	 16.81
	1995	59	1,002	91.0	4.5	9.9	82.5	8,190	2.97	3.23	15.33	0.950	10.41	10.12	17.20
	1996	48	793	92.5	4.4	11.5	70.6	6,435	2.53	2.76	14.10	0.819	9.86	9.33	16.41
	1997	58	901	90.1	4.7	12.4	74.3	8,322	2.59	2.69	14.12	0.821	9.91	9.26	16.18
	1998	62	859 645	91.1	4.8	12.0	81.3	8,383	2.79	3.00	14.48	0.878	10.12	9.49	16.72
	1999 2000	63 62	645 714	89.5 90.6	4.0 3.7	8.3 12.1	73.4 77.7	3,949 6,695	2.82 2.54	3.32 2.73	16.07 15.78	0.870 0.797	9.83 9.92	9.41 9.16	16.76 16.35
	2000	63	1,284	90.0 93.6	4.3	9.9	71.7	9,112	2.54	2.73	16.21	0.901	10.53	9.65	16.54
	2002	62	1,109	93.1	4.4	11.7	76.0	9,844	2.58	2.76	14.16	0.890	10.35	9.45	16.67
	2003	57	406	93.6	4.2	20.7					14.32				
State	1993 1994	491	 797	 87.4		 11.9	 80 6	7.647	2.92		 15 02		 10 43	 9.68	 16.97
	1994	491 586	797 974	87.4 89.9	6.0 5.4	9.2	80.6 78.9	7,647 7,114	2.92	3.26	15.02 15.51	0.952 0.949	10.43 10.33	9.68 9.94	16.97
	1996	562	739	89.3	5.3	10.3	65.7	5,007	2.52	2.72	14.87	0.775	9.76	9.08	15.70
	1997	642	1,049	89.5	5.4	10.4	76.0	8,326	2.78	2.92	14.92	0.896	10.56	9.60	16.55
	1998	610	895	90.9	5.0	13.8	77.2	9,542	2.86	3.04	15.05	0.828	10.31	9.51	16.48
	1999	603	591 992	90.5	5.6	11.1	70.4 72.2	4,630	2.82 2.57	3.09	15.29	0.928	10.16	9.78	16.72
	2000 2001	555 632	992 805	92.8 92.6	4.5 5.2	13.0 12.0	72.2	9,321 6,737	2.57 2.87	2.84 3.13	14.86 15.59	0.870 1.020	10.01 10.52	9.33 9.99	16.25 16.71
	2001	623	1,108	94.0	4.7	13.8	71.9	11,009	2.65	2.80	14.46	0.889	10.16	9.35	16.34
	2003	636	461	93.4	4.8	20.6					14.22				

1/ 2/ <u>3</u>/

Survey was not conducted in 1993. Sizing data not available in 2003. Number of samples is based on the Pistachio Objective Measurement Survey. There are two trees per sample. All weights are in grams. Suture, cross suture and length measurements are in millimeters.

	Samples Completed <u>b</u> /	Est. Average	Est. Percent Of All Spaces That		Count Data			In-Hull Data			Kernel Data			
Area and Variety			Bearing Trees	Pollinators	Nuts Per Cluster (Filled and Blank)	Percent Of Nuts Filled		Weight Per Nut (Includes Blanks)	Weight Per Nut (Filled)	In-Hull Cross Suture	Average Weight Per Kernel		Cross Suture	Length
KERN														
Kerman/Atlantica	47	461	92.1	5.3	25.6					14.74				
Kerman/Pioneer Gold I	190	372	95.6	3.7	25.3					14.16				
Kerman/Pioneer Gold II	2	205	95.0	5.0	51.9					13.32				
TOTAL	254	379	94.8	4.0	25.7					14.28				
KINGS														
Kerman/Atlantica	13	483	96.9	2.3	21.2					13.88				
Kerman/Pioneer Gold I	49	555	94.7	3.9	15.3					14.22				
Kerman/Pioneer Gold II	2	1,329	100.0	0.0	10.8					13.76				
TOTAL	66	580	95.2	3.5	15.1					14.12				
MADERA														
Kerman/Atlantica	100	664	86.8	9.0	18.3					13.99				
Kerman/Pioneer Gold I	37	312	95.8	3.9	16.0					13.80				
Kerman/Pioneer Gold II	3	531	88.3	10.0	36.0					14.38				
TOTAL	144	555	89.5	7.5	19.0					13.98				
MERCED														
Kerman/Atlantica	30	620	89.7	5.1	15.4					14.11				
Kerman/Pioneer Gold I	1	60	100.0	0.0	20.7					12.77				
Kerman/Pioneer Gold II														
TOTAL	31	593	90.0	4.9	15.6					14.11				
TULARE Kerman/Atlantica	13	559	88.8	3.1	20.0					14.80				
Kerman/Pioneer Gold I	41	371	95.0	4.5	21.6					14.13				
Kerman/Pioneer Gold II	1	351	100.0	0.0	23.7					14.10				
TOTAL	57	406	93.6	4.2	20.7					14.32				
STATE														
Kerman/Atlantica	219	616	89.5	6.8	19.0					14.27				
Kerman/Pioneer Gold I	382	372	95.6	3.8	22.2					14.19				
Kerman/Pioneer Gold II	13	606	95.0	4.6	23.4					14.11				
TOTAL	636	461	93.4	4.8	20.6					14.22				

PISTACHIO OBJECTIVE MEASUREMENT SURVEY DATA, 2003 a/

Sizing data not available in 2003. Number of samples is based on the July Pistachio Objective Measurement Survey. There are two trees per sample. Samples completed may not add to "Total" due to other miscellaneous variety/rootstock which are not listed. <u>a</u>/ b/

		Acreage			Prod	Value of Production			
Year	Bearing <u>a</u> /	Non-Bearing	Total Acres	Marketable In-Hull	Shelling Stock	Total	Yield Per Bearing Acre	Grower Return Per Pound	Total Value
	Acres			1,000	Pounds (In-Hull	Basis)	Pounds	Cents	\$1,000
1980	26,000	9,000	35,000	18,600	8,300	26,900	1,030	205.0	55,145
1981	27,500	13,100	40,600	11,300	3,200	14,500	527	136.0	19,720
1982	29,900	15,600	45,500	39,600	4,400	44,000	1,470	149.0	66,560
1983	31,100	16,000	47,100	20,700	5,700	26,400	849	141.0	37,224
1984	30,800	16,800	47,600	45,200	17,900	63,100	2,050	97.6	61,586
1985	32,300	18,700	51,000	23,100	4,000	27,100	839	137.0	37,127
1986	34,200	20,400	54,600	57,500	17,400	74,900	2,190	112.0	83,888
1987	41,000	16,400	57,400	27,200	5,900	33,100	807	137.0	45,347
1988	47,200	10,300	57,500	76,100	17,900	94,000	1,990	122.0	114,680
1989	50,900	12,000	62,900	33,000	6,000	39,000	766	163.0	63,570
1990	53,700	11,100	64,800	94,600	25,400	120,000	2,230	102.0	122,400
1991	55,700	13,300	69,000	59,000	18,000	77,000	1,280	125.0	96,250
1992	56,500	13,900	70,400	114,500	32,500	147,000	2,600	103.0	151,410
1993	57,000	15,700	72,700	113,000	39,000	152,000	2,670	107.0	162,640
1994	57,500	16,600	74,100	94,600	34,400	129,000	2,235	92.1	118,809
1995	60,300	13,400	73,700	107,500	40,500	148,000	2,454	109.0	161,320
1996	64,300	17,100	81,400	85,000	20,000	105,000	1,630	116.0	121,800
1997	65,400	17,000	82,400	137,000	43,000	180,000	2,750	113.0	203,400
1998	68,000	19,300	87,300	138,000	50,000	188,000	2,760	103.0	193,640
1999	71,000	21,000	92,000	105,000	18,000	123,000	1,730	133.0	163,590
2000	74,600	21,700	96,300	190,000	53,000	243,000	3,260	101.0	245,430
2001	78,000	21,000	99,000	126,500	34,500	161,000	2,060	101.0	162,610
2002	83,000	23,000	106,000	242,000	61,000	303,000	3,650	111.0	336,330
2003	88,000	23,000	111,000	<u>b</u> /	<u>b</u> /	<u>b</u> /	<u>b</u> /	<u>b</u> /	<u>b</u> /
				_	_	_	_	_	_

CALIFORNIA PISTACHIO ACREAGE, PRODUCTION, PRICE AND VALUE, 1980-2003

<u>a</u>/ Bearing acreage for 1988 to date is defined as plantings that are six years old and older. Bearing acreage for 1980 through 1987 is defined as plantings that are seven years old and older.

b/ Pistachio price, total crop value, and production will be available in January 2004.

The California Agriculture Statistics Service would like to thank the California Pistachio Industry for their cooperation and support!

VIC TOLOMEO State Statistician JACK RUTZ Deputy State Statistician

GARY NELSON -- DOUG FLOHR -- SUZANNE ADAMS Agricultural Statisticians -- Estimates Group

California Agricultural Statistic Service publications are available on the Internet at: http://www.nass.usda.gov/ca