REGISTERED AND ACTIVATED PRESCRIBER



Female patients of childbearing potential (FCBP)

Male patients/Female patients not of childbearing potential (FNCBP)

BEFORE TREATMENT

- Educate the patient about isotretinoin and contraception requirements of iPLEDGE
- Screen by obtaining a negative pregnancy test
- Obtain a signed Patient Information/Informed Consent (for all patients) form for treatment
- Obtain a second Patient Information/Informed Consent About Birth Defects (for female patients who can get pregnant) form
- **Register** patient in the program and provide patient ID card
- Counsel patient, or refer to an expert, that she must use 2 effective forms of contraception simultaneously for at least 1 month before starting therapy
- Order pregnancy tests using a CLIA-certified lab during the first 5 days of the menstrual cycle (patients with amenorrhea/irregular cycle, please refer to the PI)
- Access* the system to confirm patient counseling of program and contraception requirements, and to enter pregnancy test result and the patient's forms of contraception
- Provide a prescription for a maximum 30-day supply

- Educate patient about isotretinoin
- Obtain a signed Patient Information/Informed Consent (for all patients) form for treatment
- **Register** patient in the program and provide patient ID card
- Access* the system to confirm patient counseling of program requirements
- **Provide** a prescription for a maximum 30-day supply

EACH MONTH DURING THERAPY

- · Counsel patient on program and contraception requirements
- Order a pregnancy test using a CLIA-certified lab
- Access* the system to confirm patient counseling of program and contraception requirements and to enter pregnancy test result and the patient's forms of contraception
- Provide a prescription for a maximum 30-day supply

- Counsel patient on program requirements
- Access* the system to confirm patient counseling of program requirements
- Provide a prescription for a maximum 30-day supply

AFTER TREATMENT

- Counsel the patient to use 2 forms of contraception every time she has intercourse for at least 1 month after last dose
- Counsel the patient not to donate blood for 1 month after last dose
- Order a pregnancy test using a CLIA-certified lab immediately after the last dose
- Order a pregnancy test 1 month after the last dose
- Access* the system to enter pregnancy test results after every pregnancy test

Counsel patient not to donate blood for 1 month after last dose

*You can access the system via the web site, www.ipledgeprogram.com, or the telephone, 1-866-495-0654.