

Bibliography

1. Burzykowski, T., Molenberghs, G., Abeck, D. et al. 2003. High prevalence of foot diseases in Europe: results of the Achilles Project. *Mycoses.* 46:496-505.
2. Maruyama, R., Hiruma, M., Yamauchi, K. et al. 2003. An epidemiological and clinical study of untreated patients with tinea pedis within a company in Japan. *Mycoses.* 46:208-212.
3. Ogasawara, Y., Hhiruma, M., Muto, M., and Ogawa, H. 2003. Clinical and mycological study of occult tinea pedis and tinea unguium in dermatological patients from Tokyo. *Mycoses.* 46:114-119.
4. Fuchs, A., Fiedler, J., Lebwhol, M., Sapadin, A., et al. 2004. Frequency of Culture-Proven Dermatophyte Infection in Patients with Suspected Tinea Pedis. *Am. J. Med. Sci.* 327(2):77-78.
5. Liu, D., Coloe, S., Baird, R., and Pedersen, J. 2000. Application of PCR to the identification of dermatophyte fungi. *J. Med. Microbiol.* 49:493-497.
6. Miller, M., and Hodgson, Y. 1993. Sensitivity and Specificity of Potassium Hydroxide Smears of Skin Scrapings for the Diagnosis of Tinea Pedis. *Arch. Dermatol.* 129:510-511.
7. Markus, R., Huzaira, M., Anderson, R. R., and Gonzales, S. 2001. A Better Potassium Hydroxide Preparation? In vivo diagnosis of tinea with confocal microscopy. *Arch. Dermatol.* 137(8):1076-8.
8. Semel, J. D. and Goldin, H. 1996. Association of Athlete's Feet with Cellulitis of the Lower Extremities: Diagnostic Value of Bacterial Cultures of Ipsilateral Interdigital Space Samples. *Clin. Infect. Dis.* 23:1162-1164.
9. Bradley, M. C., Leidich, S., Isham, N., Elewski, B. E., and Ghannoum, M. 1999. Antifungal susceptibilities and genetic relatedness of serial *Trichophyton rubrum* isolates from patients with onychomycosis of the toenail. *Mycoses.* 42 (Supp. 2):105-110.
10. Ghannoum, M. A., Espinel-Ingroff, A., Pfaller, M. A., et al. An Intra- and Inter-Laboratory Study of an Antifungal susceptibility Method for the Testing of Dermatophytes.
11. Jessup, C. J., Warner, J., Isham, N., Hasan, I., and Ghannoum, M. 2000. Antifungal Susceptibility Testing of Dermatophytes: Establishing a Medium for Inducing Conidial Growth and Evaluation of Susceptibility of Clinical Isolates. *J. Clin. Microbiol.* 38(1):341-344.

12. Martin, A. G. and Kobayashi, G. S. “Superficial fungal infections: dermatophytosis, tinea nigra, piedra” in Freedberg, I. M., et al. (ed) (1999), *Fitzpatrick’s Dermatology in General Medicine*, New York: McGraw-Hill, Chapter 206, 2349-2351.
13. Mukherjee, P., Leidich, S., and Isham, N., et al. 2003. Clinical *Trichophyton rubrum* Strain Exhibiting Primary Resistance to Terbinafine. *Antimicrob. Agents Chemother.* 47:82-86.
14. Norris, H., Elewski, B., and Ghannoum, M. 1999. Optimal growth conditions for the determination of antifungal susceptibility of three species of dermatophytes with the use of a microdilution method. *J. Am. Acad. Dermatol.* 40(6)(2):S9-S-13.
15. Roberts, D. T. and Bilsland, D. J. “Tinea pedis and skin dermatophytosis” in Lebwohl, M. et al. (ed) (2002), *Treatment of skin disease. Comprehensive therapeutic strategies*, Philadelphia: Mosby, Section 200, 624-626.