

**Assessment
of the Gulf of Maine
and Georges Bank
Witch Flounder Stock
for 2003**

by

**Susan E. Wigley,
Jon K.T. Brodziak,
and Laurel Col**

September 2003

Recent Issues in This Series

- 02-13 **Report of the 35th Northeast Regional Stock Assessment Workshop (35th SAW): Public Review Workshop.** [By Northeast Regional Stock Assessment Workshop No. 35.] September 2002.
- 02-14 **Report of the 35th Northeast Regional Stock Assessment Workshop (35th SAW): Stock Assessment Review Committee (SARC) Consensus Summary of Assessments.** [By Northeast Regional Stock Assessment Workshop No. 35.] September 2002.
- 02-15 **Report of the Workshop on Trawl Warp Effects on Fishing Gear Performance, Marine Biological Laboratory, Woods Hole, Massachusetts, October 2-3, 2002.** [By Workshop on Trawl Warp Effects on Fishing Gear Performance, Marine Biological Laboratory, Woods Hole, Massachusetts, October 2-3, 2002.] October 2002.
- 02-16 **Assessment of 20 Northeast Groundfish Stocks through 2001: A Report of the Groundfish Assessment Review Meeting (GARM), Northeast Fisheries Science Center, Woods Hole, Massachusetts, October 8-11, 2002.** [By Groundfish Assessment Review Meeting, Northeast Fisheries Science Center, Woods Hole, Massachusetts, October 8-11, 2002.] October 2002.
- 03-01 **Manuscript/Abstract/Webpage Preparation, Review, & Dissemination: NEFSC Author's Guide to Policy, Process, and Procedure.** By J.A. Gibson, T.L. Frady, E.L. Kleindinst, and L.S. Garner. January 2003.
- 03-02 **Stock Assessment of Yellowtail Flounder in the Southern New England - Mid-Atlantic Area.** By S.X. Cadrin. [A report of Northeast Regional Stock Assessment Workshop No. 36.] February 2003.
- 03-03 **Stock Assessment of Yellowtail Flounder in the Cape Cod - Gulf of Maine Area.** By S.X. Cadrin and J. King. [A report of Northeast Regional Stock Assessment Workshop No. 36.] February 2003.
- 03-04 **Report of the 36th Northeast Regional Stock Assessment Workshop (36th SAW): Public Review Workshop.** [By Northeast Regional Stock Assessment Workshop No. 36.] February 2003.
- 03-05 **Description of the 2002 Oceanographic Conditions on the Northeast Continental Shelf.** By M.H. Taylor, C. Bascuñán, and J.P. Manning. March 2003.
- 03-06 **Report of the 36th Northeast Regional Stock Assessment Workshop (36th SAW): Stock Assessment Review Committee (SARC) Consensus Summary of Assessments.** [By Northeast Regional Stock Assessment Workshop No. 36.] March 2003.
- 03-07 **Implementation of Electronic Logbook Reporting in a Squid Bottom Trawl Study Fleet during 2002.** By L.C. Hendrickson, D.A. Hiltz, H.M. McBride, B.M. North, and J.E. Palmer. April 2003.
- 03-08 **Northeast Fisheries Science Center Publications, Reports, and Abstracts for Calendar Year 2002.** By L. Garner and J.A. Gibson. July 2003.
- 03-09 **Stock Assessment of Summer Flounder for 2003.** By M. Terceiro. August 2003.
- 03-10 **Comparison of Invertebrate Abundances in Four Bays of the Northeastern United States: Two Bays with Sparse Quahogs and Two Bays with Abundant Quahogs.** By C.L. MacKenzie, Jr. August 2003.
- 03-11 **Accuracy Enhancement of Microscope Enumeration of Picoplankter *Aureococcus anophagefferens*.** By J.B. Mahoney, D. Jeffress, J. Bredemeyer, and K. Wendling. August 2003.
- 03-12 **A Taxonomy of World Whaling: Operations, Eras, and Data Sources.** By R.R. Reeves and T.D. Smith. August 2003.
- 03-13 **Distribution of the Brown Tide Picoplankter *Aureococcus anophagefferens* in Western New York Bight Coastal Waters.** By J.B. Mahoney, D. Jeffress, C. Zetlin, P.S. Olsen, H. Grebe, and J. Brooks. August 2003.

A Report of the 37th Northeast Regional Stock Assessment Workshop

Assessment of the Gulf of Maine and Georges Bank Witch Flounder Stock for 2003

by

Susan E. Wigley^{1,2}, Jon K.T. Brodziak^{1,3}, and Laurel Col^{1,4}

Postal Addresses: ¹*National Marine Fisheries Serv., 166 Water St., Woods Hole, MA 02543-1026*

E-Mail Addresses: ²*Susan.Wigley@noaa.gov*
³*Jon.Brodziak@noaa.gov*
⁴*Laurel.Col@noaa.gov*

**U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts**

September 2003

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review but no technical or copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/series/crdlist.htm>. The electronic version will be available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the desired document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

This document's publication history is as follows: manuscript submitted for review--July 7, 2003; manuscript accepted through technical review--August 26, 2003; manuscript accepted through policy review--September 3, 2003; and camera-ready copy submitted for publication--September 8, 2003. This document may be cited as:

Wigley, S.E.; Brodziak, J.K.T.; Col, L. 2003. Assessment of the Gulf of Maine and Georges Bank witch flounder stock for 2003. *Northeast Fish. Sci. Cent. Ref. Doc.* 03-14; 186 p. Available from: National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543-1026.

TABLE OF CONTENTS

INTRODUCTION	1
THE FISHERY	2
STOCK ABUNDANCE and BIOMASS INDICES	7
MATURITY	10
MORTALITY	11
ESTIMATION OF FISHING MORTALITY RATES AND STOCK SIZE	12
BIOLOGICAL REFERENCE POINTS	16
SHORT-TERM PROJECTIONS	18
CONCLUSIONS	18
ACKNOWLEDGMENTS	18
LITERATURE CITED	18
TABLES	22
FIGURES	84
APPENDIX: Full output of VPA RUN 301-f	155

ABSTRACT

The 2003 analytical assessment for witch flounder in USA waters, covering the period 1982-2002, estimates 2002 fishing mortality and spawning stock biomass for stock status, and provides short-term projections of median landings, discards and spawning stock biomass for various fishing mortality scenarios.

USA commercial landings increased during the 1960's from 1,200 mt to about 3,000 mt, then fluctuated between 2,000 and 3,000 mt until 1983 through 1985 when landings abruptly increased to about 6,000 mt. Landings subsequently declined to 1,500 mt in 1990. Since the early 1990s, landings have fluctuated between 2,000 and 3,000 mt. In 2002, USA commercial landings totaled 3,186 mt, a 5% increase over 2001; and 117% higher than in 1990, the lowest value since 1964. Research survey indices of abundance and biomass remained fairly stable from 1963 until the late 1970s; autumn indices declined during the early and mid 1980s, reaching record low levels in late 1980s and early 1990s. Abundance sharply increased in 1993 and has continued to increase to near record high levels in 2002; however, the age composition still remains truncated.

The VPA indicates that fishing mortality (ages 8-9, unweighted) increased from 0.26 in 1982 to 0.67 in 1985, declined to 0.22 in 1992, increased to 1.13 in 1996, then declined to 0.41 in 2002. Spawning stock biomass declined from 16,897 mt in 1982 to about 3,800 mt in 1996. With recent increases in recruitment and declines in fishing mortality, SSB has increased to 18,296 mt in 2002. Since 1982 recruitment of age 3 has ranged from approximately 3 million fish (1984 year class) to 67.6 million fish (1997 year class). Over the 1982-2002 period, average recruitment of age 3 fish (the 1979 - 2000 year classes) was 19.6 million (the geometric mean equaled 14.4 million fish). The 1995-1999 year classes appear to be above average, and the 1997 year class is the largest in the VPA time series. In addition to the VPA, an alternative model, a statistical catch at age model (SCAA) was also conducted. The SCAA generally confirmed trends in the VPA results.

The biological reference points were updated by applying the approach used to estimate MSY proxies for witch flounder. F_{msy} is approximated as $F_{40\%}$ (0.23), the SSB_{msy} proxy is 25,248 mt, the product of $40\%MSP$ (1.2882 kg spawning biomass) and average long-term recruitment (19.6 million). The MSY proxy is 4,375 mt, the product of yield per recruit at $F_{40\%}$ (0.2232 kg) and average recruitment.

Based on the ADAPT VPA, the witch flounder stock was not overfished, but overfishing was occurring in 2002. Fully recruited fishing mortality in 2002 was 0.41, nearly double F_{msy} (0.23), and spawning stock biomass was estimated to be 18,296 mt in 2002, 72% of SSB_{msy} (25,248 mt). Recent year classes appear to be above average. Although the spawning stock biomass has increased, the age structure still remains truncated. Fishing mortality should be reduced to F_{msy} or below to allow the age structure to rebuild.

INTRODUCTION

The witch flounder (*Glyptocephalus cynoglossus*, L.) or grey sole is a deep water boreal flatfish occurring on both sides of the North Atlantic. In the Northwest Atlantic, witch flounder are distributed from Labrador to Georges Bank and in continental slope waters southward to Cape Hatteras, North Carolina. In U.S. waters, the species is commercially abundant in the Gulf of Maine-Georges Bank region [defined as Northeast Fisheries Science Center (NEFSC) Statistical Reporting Areas (SA) 511-515, 521-522, 525-526, and 561-562; Figure 1], and, in the absence of any stock structure information, is assumed to comprise a single stock unit. Prized as a table fish, witch flounder receives a high ex-vessel price relative to other flounders and represents an important by-catch component in the New England mixed species groundfish fishery. Annual landings during the period 1960-1982 averaged 3,000 metric tons (mt), ranging from 1,000 to 6,000 mt (Lange and Lux 1978, Burnett and Clark 1983). More recently, landings declined from a peak of 6,660 mt in 1984 to a low of 1,490 mt in 1990. Landings for 2002 were 3,186 mt.

Previous witch flounder stock assessments were conducted by Burnett and Clark (1983), Wigley and Mayo (1994) and Wigley et al. (1999). An assessment update was conducted for this stock in 2002 and reviewed at Groundfish Assessment Review Meeting (GARM; NEFSC 2002). The GARM assessment indicated average fishing mortality (ages 7-9, unweighted) increased from 0.21 in 1982 to 0.59 in 1985, declined to 0.24 in 1990, increased to 0.96 in 1996, then declined to 0.45 in 2001. Spawning stock biomass declined from 18,000 tons in 1982 to about 4,000 tons in 1995 and then increased to 11,368 mt in 2001. Since 1982, recruitment at age 3 has ranged from approximately 3 million fish (1984 year class) to 84 million fish (1997 year class) with a mean of 22 million fish. In 2001, the SSB was slightly above $\frac{1}{2}$ SSB_{msy} (9,950 mt), the minimum stock size threshold, and fishing mortality ($F=0.45$) was three times higher than F_{msy} , the maximum fishing mortality threshold; thus, witch flounder was not overfished but overfishing was occurring in 2001.

This assessment of witch flounder in the Gulf of Maine - Georges Bank region and southward (USA Subareas 4, 5 and 6), presents a benchmark analytical assessment for the stock for the 1982-2002 period, estimates 2002 fishing mortality and spawning stock biomass for stock status, and provides short-term projections of median landings, discards and spawning stock biomass for various fishing mortality scenarios. This assessment provides estimates of discards from the shrimp fishery and large-mesh otter trawl fishery based upon analyses of sea sampling, commercial and research vessel survey data through 2002.

Witch flounder is managed under the New England Fishery Management Council's Multi-species Fisheries Management plan since 1987. A brief summary of groundfish management regulations affecting witch flounder is presented in Table 1. Significant changes in regulations include increased minimum size in 1983 and 1987; increases in mesh size in 1982, 1983, 1994, 1999; effort reductions in 1996 and 2002; and implementation of closed areas in 1994 and 1998 (Figure 2). The western Gulf of Maine area closure, Cashes Ledge area closure and the seasonal rolling closures overlap the witch flounder distribution (Figures 3 and 4). Management regulations for the northern shrimp fishery also impact witch flounder (Table 2); significant

changes in the shrimp fishery include a monthly 10% by-catch limit which restricted the possession of groundfish to 10% by weight of shrimp in the mid-1980's to early 1990s; and the implementation of the Nordmore grate to exclude groundfish in 1992.

THE FISHERY

Recreational Catches

There is no recreational fishery for witch flounder.

Commercial Landings

USA commercial landings in 2002 totaled 3,186 mt, a 5% increase over 2001 (Table 3); and 117% higher than in 1990, the lowest value since 1964 (Figure 5). Canadian landings from the stock have been negligible (32 mt in 2001; Table 3). Landings from the Grand Banks (NAFO Divisions 3LNO) during 1985 to 1990 are not included in this assessment. Canadian landings from the western Scotian Shelf (NAFO Division 4X) are not considered due to the fact that, until recently, witch flounder were reported as 'other flounders' by Canada, and cannot be separated from other flounder species. Furthermore, samples from the western Scotian Shelf indicate slower growth of witch flounder than in the Gulf of Maine, suggesting a different phenotypic stock.

The western Gulf of Maine (SA 513 and 514) and the central basin (SA 515) provide nearly a third of the USA witch flounder landings (Table 4); landings from Georges Bank are confined to the deeper waters north of the South Channel (SA 521, 522; Table 4). Otter trawl catches account for about 98% of witch flounder landings, with sink gillnets comprising the remainder (Table 5). Catches are generally highest during March-July (Table 6) when witch flounder form dense pre-spawning aggregations (Burnett et al. 1992). The majority of witch flounder are landed in Maine ports, primarily Portland, with lesser amounts landed in New Bedford and Gloucester, MA (Table 7).

Although culling and grading practices vary by port, witch flounder have historically been landed as either 'small' or 'large'; however, three market categories ('peewee', 'medium', and 'jumbo') were added in some ports beginning in 1982 (Table 8, Figure 6). Since the early 1990s, the proportions of witch flounder landings from the peewee and small market categories have steadily increased. In 2002, witch flounder less than 45 cm ('peewee' and 'small' market categories) constituted 87% of total landings (Table 8, Figure 6). The current regulated minimum landing size for witch flounder is 36 cm (14 inches).

Sampling Intensity

Length frequency and age sampling data for witch flounder landings from the Gulf of Maine-Georges Bank region are summarized by quarter and market category in Table 9 (because some ports do not cull into 'peewee' or 'jumbo' categories, NEFSC sampling protocols incorporate these categories into the 'small' and 'large' categories, respectively). Until 1982, sampling was minimal and sporadic. During 1982-1988, an average of 48 length frequency samples (approximately 100 fish per sample) was obtained annually over all market categories, representing 1 sample per 102 mt landed. In 1990, sampling requirements were adjusted to 1 sample per 50 mt to obtain more samples from the 'large' market category. However, samples for the 'large' market category have been difficult to obtain due to the sharp decrease in the landings of larger fish in recent years (Tables 8 and 9). Sampling intensity during 2001-2002 averaged 39 samples annually, representing 1 sample per 80 mt landed; nonetheless, even with this increased sampling intensity, inadequate numbers of samples were obtained for some market categories and quarter combinations. In 2002, of the 35 samples collected, 15 were small samples (43%), 10 were medium (29%) and 10 were large (29%). Compared with the 2002 market category landings distribution by weight (small 87%; medium: 10%; large: 3%), sampling in 2002 adequately approximated the market category distribution of landings on an annual basis. As in previous years, it was necessary to pool some quarters for some market categories. A summary of pooling procedures by year, market category and quarter is presented in Table 10.

Commercial Landings at Age

Commercial age data for the years 1982 to 2002 were available for this assessment. Quarterly age-length keys (ALKs) were applied to corresponding commercial landings length frequency data by market category. Resulting estimates of annual age compositions (age 0 to 14+) are presented in Table 11 and Figure 7. No discernible changes in growth are evident during the 1982-2002 period; although landings mean weights and mean lengths at ages 6 to 8 declined in 1996-2002, this may be an artifact of poor sampling in recent years.

Discards

The Fisheries Observer Program (FOP), which began in 1989, has generated various levels of coverage for different fisheries. Prior to the FOP, NEFSC conducted sea sampling on an ad-hoc basis. The northern shrimp fishery, the small-mesh otter trawl fishery, and the large-mesh otter trawl fishery are three fisheries in which discarding of witch flounder occurs. In this assessment, discard estimates have been estimated for the shrimp fishery and the large-mesh otter trawl fishery.

Northern shrimp fishery

Since the 'shrimp season' spans a calendar year, in this report, the year in which most of the fishing occurred will be used to identify the entire season. For example, 1990 will refer to the

shrimp season from December 1, 1989 to May 31, 1990. An overview of the procedures used to estimate witch flounder discarded in the shrimp fishery from 1982 to 1998 is presented in Figure 8. These estimation procedures were used in the 1994 assessment (Wigley and Mayo 1994), reviewed by the SAW 18 (NEFSC 1994), and extended through 1997 using the same methodology. The ratio of witch flounder discarded (kg) to days fished was calculated using FOP data for individual shrimp seasons, 1989-1997, by fishing zone. Since depth is an important factor influencing discards (Wigley MS 1994), discard ratios were calculated for each of three fishing zones (zone 1 = 0-3 miles from shore, zone 2 = 3 - 12 miles, and zone 3 = greater than 12 miles) in each season. For the most part, fishing zones are analogous to depth zones. Statistical testing of zonal discard rates indicated differences between fishing zones in most years. The zone-specific discard rates were weighted by the days fished in each zone to calculate a weighted mean discard rate for each season (Table 12). To estimate witch flounder discard rates prior to the FOP, (i.e., 1982-1988), a simple linear regression was employed using 1989-1992 (years in which the Nordmore grate was not required) weighted mean discard rates and annual indices of witch flounder abundance. The NEFSC autumn bottom trawl survey index of age 3 fish was found to be the best predictor of annual discard rates ($r^2 = 0.97$, $p = 0.0127$; Figure 9; Wigley MS 1994).

With no 1998-2002 FOP sampling in the northern shrimp fishery, an alternative method of survey filtering was explored to estimate witch flounder discard rates; however, due to insufficient length frequency data at small sizes, this method did not prove fruitful. As used for the years prior to the FOP, a simple linear regression using 1993-1997 (years in which the Nordmore grate was required) annual shrimp season discard rates and annual survey indices of autumn age 3 fish was employed ($r^2 = 0.87$, $p = 0.0206$). This five-point regression may not be as robust as the r^2 suggests, as four of the points are clustered (Figure 9).

To obtain total weight of witch flounder discarded during a shrimp season, season discard rates (kg per day fished) were multiplied by the total number of days fished by the commercial fleet in each season (Table 13). Estimated discard weight was then translated into discarded numbers at age by applying witch flounder sea-sampled discard length-frequencies (Figure 10) expanded up to the total discard weight and then applying NEFSC spring bottom trawl survey ALKs. Detailed information on this method is given in Wigley (MS 1994). For 1995-2002, days fished were estimated from the Vessel Trip Reports (VTR) using a stratification level of year, ton class, port group, month, and fishing zone. To derive the number of trips by fishing zone, the proportion of VTR trips by fishing zone was applied to the number of trips in the weighout database. Days fished per trip in each fishing zone were derived from the VTR data. Days fished per trip were then multiplied by the estimated number of trips for each fishing zone to derive estimated days fished by fishing zone, and then summed over year and fishing zone.

For the 1982-1997 time period, discard estimates of numbers at age and weight were derived on a shrimp season basis due to the limited number of length frequency samples in December. To adjust the shrimp fishery discard-at-age from a shrimp season basis to calendar year, the ratio of December days fished to the entire shrimp season days fished was used to apportion of the weight and numbers discarded into December and January-May categories. The December

discards-at-age were shifted back one age, and then re-combined with the January-May matrix of the corresponding calendar year. The December discard weight was combined with January-May of the same calendar year. Mean lengths and mean weights at age in the re-combined catch at age were weighted by the numbers at age from each category.

Without 1998-2002 FOP sampling, discard length-frequency data were unavailable to partition the 1998-2002 estimated discard weight into numbers at length; thus discarded numbers at age were derived by apportioning discard weight by the average age composition (calendar year) of discards in 1993-1997 and then dividing by the average 1993-1997 discard mean weights at age (Figure 11). The average 1993-1997 mean weights at age from the FOP were consistent with trends in mean weights from the NEFSC survey during the 1998-2002 time period (Figure 11).

Witch flounder discards in the northern shrimp fishery ranged from a low 0.8 mt in 2002 to a high of 34 mt in 1988 and 1995 (Table 13). Similarly, number of witch flounder discarded ranged from 40,000 fish discarded in 2002 to 1.8 million fish in 1994 (Table 13). Estimates of age compositions of discarded witch flounder in the shrimp fishery are presented in Table 14 and Figure 12. Discarded witch flounder from the shrimp fishery range from age 0 to 6, with ages 1 to 3 most commonly discarded (Table 14 and Figure 12).

Large-mesh otter trawl fishery

Discard estimation from the large-mesh otter trawl fishery is confounded by the lack of FOP coverage prior to 1989, sparse coverage in the beginning of the program, and the recent implementation of year-around and seasonal area closures. As a result, three estimation scenarios were examined (Table 15): 1) utilizing a survey filter method (Table 16); 2) utilizing the at-sea observer data (Table 17); and 3) utilizing the Vessel Trip Report data (Table 18). The estimated discards (in weight and numbers) are presented in Table 19. Each method is described below.

The method used in previous witch flounder assessments to estimate large-mesh otter trawl discards was based upon a method developed by Mayo et al. (1992) which utilizes survey and commercial catch at length data, commercial gear retention ogives, and information on culling practices. An overview of the method is presented in Figure 13. Research vessel length frequency data were filtered through commercial gear retention ogives corresponding to the predominant mesh size employed in the large-mesh fishery (130, 140, and 152 mm) and then through a culling practice ogive. Due to the sparse gear retention studies for witch flounder, mesh selection ogives were taken from Walsh et al. (1992) for American plaice. Given the high value and low abundance of this species, the culling practice of commercial fishermen was assumed to be nearly knife edged at the minimum landing size. A semi-annual ratio estimator of survey filtered 'kept' index to semi-annual numbers landed was used to expand the estimated 'discard' survey index to obtain numbers of fish discarded at length. The method used in this analysis differs from the method described by Mayo et al (1992) which employs an expansion factor derived from a linear regression from the ratios of kept to landed at length. A spreadsheet illustrating the method used is presented in Table 16 for 1993 using the spring survey and

commercial landings from quarters 1 and 2. Semi-annual numbers of discard fish at length were apportioned to age using the corresponding season NEFSC ALK. Estimated numbers of discarded witch flounder in the large-mesh otter trawl fishery are presented by season in Table 19. Results indicate that in recent years, numbers discarded at sea comprised as much as 54% of the witch flounder landed. The general pattern of discarding appears to be consistent with that expected given strong recruitment during 1979-1981 and the mid-1990's.

Given the distribution of juvenile witch flounder in the western Gulf of Maine and the recent implementation of year-around area closures and seasonal rolling area closures in the western Gulf of Maine, there was some concern regarding the application of the survey filter method to estimate discards in recent years. Since the commercial fishery does not have year-around access to the population estimated by the NEFSC survey, it may be inappropriate to use the survey filter method to estimate discards. For the 1989-2002 period, discard weight to kept weight ratios (D/K ratio) were calculated from FOP data on a semi-annual basis (Table 17). Total discard weight was derived by multiplying the D/K ratio by the commercial landings. The number of sea sampled trips varied from no trips in the second half of 1992 to 83 trips in the second half of 2002. The D/K ratios ranged between 0.02 and 0.50. Given the limited number of trips, tows and available discard length frequencies, discards at age were derived only for the 1995-2002 time period (Tables 19).

The Vessel Trip Report data were explored for information on discarding of witch flounder. Reporting of discard information in the logbooks is known to be incomplete. To eliminate problems associated with incomplete reporting, a subset of the VTR data was used. The VTR subset included only logbooks which reported discards of any species (DeLong et al. 1997), assuming that operators who report discards of any species would reliably report witch flounder discards. This subset was used to estimate discard ratios (discard weight/kept weight) semi-annually for large-mesh otter trawl gear from 1994 to 2002. Limitations of this analysis are: 1) the dealer data used to expand discard rates to total discard weight do not contain information on mesh size, precluding partitioning of otter trawl fisheries into small and large mesh trips; 2) there is no area information on dealer data to isolate trips from the Gulf of Maine-Georges Bank region. From this analysis, results suggest that discard rates range between 4% and 9% (Table 18). These estimates should be reviewed cautiously as not all fishermen report discards. Discarded numbers at age were estimated by expanding the FOP length frequencies and applying the survey age/length keys (Table 19).

For estimates of total catch at age, discards from the large-mesh otter trawl fishery were derived using the survey filter method from 1982-1994 and using the FOP method for 1995-2002 (Table 20 and Figure 14). Total discards at age (from the shrimp fishery and large-mesh otter trawl fishery combined) are presented in Figure 15.

Total Catch at Age

Total catch at age compositions (including commercial landings, discards from the northern shrimp fishery and the large-mesh otter trawl fishery) are presented in Table 21 and Figure 16. The age composition data reveal strong 1979-1981 year classes (Table 21). The 1989 and 1993 year classes also appear to have been strong; however, these cohorts were heavily discarded in both the shrimp and large-mesh otter trawl fisheries (Tables 14 and 20; Figures 12 and 14). The poor 1984 year class is also evident as well as the truncated age-structure since the early 1990's.

Since witch flounder landings are highest during March-July, the average weights-at-age in the catch approximate mid-year weights. Mean weights at age at the beginning of the year (January 1; Table 22) were derived from mid-year weights using procedures described by Rivard (1980).

STOCK ABUNDANCE AND BIOMASS INDICES

Commercial LPUE

Commercial catch rates (landings per unit effort, LPUE, expressed as landings in mt per day fished) were derived for vessel tonnage classes 2-4 [Class 2 consists of vessels 5 to 50 gross registered tons (GRT); Class 3, 51 to 150 GRT; and Class 4, 151 to 500 GRT]. These vessel classes account for greater than 95% of annual witch flounder otter trawl landings. LPUE indices for the Georges Bank-Gulf of Maine region were computed for: 1) all trips landing witch flounder, and 2) trips in which 40% or more of the total landings comprised witch flounder (Table 23). These '40% trips' may represent effort that is 'directed' towards witch flounder, a species historically taken as by-catch.

For all trips landing witch flounder, increases in LPUE occurred in 1977-1978 for tonnage classes 2 and 3 and in 1982 for tonnage class 4, and remained high during the early 1980s; however, LPUE indices declined steadily for all tonnage classes from 1986 to 1990. Since the early 1990s, LPUE indices have steadily increased and are among the highest in the time series (Table 23, Figure 19a). Indices for 40% trips peaked in the early 1980's, then declined to a low in 1994, and have increased slightly in recent years (Table 23, Figure 19a). Effort (days fished) associated with all trips and 40% trips increased during the late 1970s and early 1980s, peaked during 1985-1988, and have generally declined since (Figure 19b). While there is some evidence of increased directed effort in the early and mid 1980s [a period in which both witch flounder and American plaice were abundant and a small directed fishery emerged (Burnett and Clark 1983)], it is likely that LPUE indices derived for all trips landing witch flounder provide the best measure of relative abundance. In 1994 the NEFSC commercial data collection system changed from a voluntary to a mandatory system in which fishermen self-report fishing effort. Investigation is still on-going to determine if the time series of LPUE data can be extended (considered one series) or whether the post 1993 LPUE derived under the mandatory system constitutes a separate time series. Effort (days fished) for 1994 to 2002 may be underestimated

in this report since effort is based upon preliminary VTR data, which do not represent 100% of the trips.

Research Vessel Survey Indices

The NEFSC has conducted annual research vessel stratified random bottom trawl surveys during autumn since 1963 and during spring since 1968. Details on survey sampling design and the use of survey data in stock assessments are given in Azarovitz (1981) and Clark (1981), respectively.

In September 2002, an offset in the trawl wraps was detected which may have effected the NEFSC bottom trawl surveys conducted from winter 2000 to the spring 2002. Extensive analyses of existing data sets and experimental studies were conducted to evaluate the offset issue (NEFSC 2002). These analyses were reviewed by a panel of experts and they concluded that no adjustments to the survey time series were justified (Groundfish Science Peer Review, 2003).

The Commonwealth of Massachusetts Division of Marine Fisheries (DMF) began an inshore trawl survey in 1978 which complements the NEFSC survey in coastal Massachusetts waters in that depths less than 27 meters (the lower depth limit sampled by the NEFSC offshore survey) are sampled (for details of this survey, see Howe et al. 1981). Additionally, the Northern Shrimp Technical Committee of the Atlantic States Marine Fisheries Commission (ASMFC) has conducted an annual northern shrimp survey during August in the Gulf of Maine since 1983, with catch data for witch flounder available from 1984 on (for details of the shrimp survey, see Northern Shrimp Technical Committee MS 1984). All three surveys provide useful information relative to trends in abundance, distribution, and recruitment of witch flounder in the Gulf of Maine-Georges Bank region. Strata utilized in the derivation of indices of relative abundance and biomass for witch flounder are as follows: NEFSC, offshore strata 22-30, 36-40 (Figure 18); Massachusetts DMF, regions 4 and 5 (Figure 19); and northern shrimp, strata 1, 3, 6, and 8 (Figure 20).

Witch flounder are generally distributed throughout the Gulf of Maine, along the Northern Edge and southern flank of Georges Bank, and southward along the continental shelf as far south as Cape Hatteras, NC (Figures 3a and 3b). Juvenile witch flounder (< 25 cm) are distributed along the western Gulf of Maine, with a few in the canyon areas in the Mid-Atlantic region (Figure 4a and 4b). Concentrations of witch flounder along the western portion of the Gulf of Maine are observed in the ASMFC shrimp survey. Although this survey has limited spatial coverage (Figure 21), most of the juvenile range is covered.

In response to a research recommendation from SARC 29, analyses were conducted to examine if the use of additional strata in the NEFSC bottom trawl survey might be appropriate. Burnett and Clark (1983) used NEFSC survey strata set 22,24,26-30, 33-40 in the first witch flounder assessment; however, Burnett (MS 1987) suggested that fish from strata 33, 34 and 35 exhibited different growth rates indicating these fish may be from a different stock inhabiting the western Scotian shelf. Based on this information, Wigley and Mayo (1994) revised the witch flounder

survey strata set excluding 33, 34, and 35, and included strata 23 and 25. Following a method developed by Cadrin (2003), witch flounder catches for the entire autumn bottom trawl survey time series were examined by individual stratum. The stratified mean number per tow in each stratum was summed over the time period, and the percentage contribution of each stratum was calculated as well as the percentage of annual stratum sampling which produced no catch (Table 24). Results indicate that the current strata set (22-30, 36-40) accounts for approximately 93% of the survey catch and that only minor differences exist between the strata sets used in previous assessments. This analysis also indicated that stratum 6 contributed to the overall witch flounder catch. The stratified mean weight (kg) per tow was calculated for three strata sets: set 1 (22-30, 36-40); set 2 (22, 24, 26-30, 36-40); and set 3 (6, 22, 24, 26-30, 36-40). The trends of these biomass indices (and their variance) are indistinguishable (Figures 22a and 22b). The inclusion of stratum 6 is not justified due to its geographical discontinuity with the core strata. Since no additional strata were identified as contributing to the total catch, or improved the precision of the estimates of mean weight per tow, the strata set 22-30, 36-40 will continue to be used.

Research vessel survey indices of abundance, biomass, and mean length for NEFSC surveys, Mass. DMF surveys, and ASMFC shrimp surveys are presented in Tables 25-27 and Figures 23-28, respectively. Length frequency data from these surveys are presented in Figures 29-32. A summary of available age data from NEFSC surveys is given in Table 28; survey age samples collected during 1976 to 1979 have not been aged. Too few age samples are collected during DMF surveys to reliably characterize the age composition of witch flounder in the inshore areas, and no age samples are collected on ASMFC surveys. Age-specific relative abundance indices from NEFSC spring and autumn surveys 1980-2002, and preliminary spring 2003 are presented in Table 29, Figures 33 and 34. Mean length and mean weights at age from the NEFSC spring and autumn surveys are given in Tables 30 and 31 and Figures 35 - 37.

While NEFSC spring survey indices tend to be more variable due to the pre-spawning aggregations of witch flounder, spring and autumn indices generally display similar trends. Abundance and biomass remained fairly stable from 1963 until the late 1970s (Table 25, Figures 23 and 24); autumn indices declined during the early and mid 1980s, reaching record low levels in 1987. Abundance sharply increased in 1993, due to a large age 0 index (Table 29, Figure 34) and has continued to increase to near record high levels in 2002. During the same time, mean length declined (Figure 27 and 28). The age structure has been truncated since the late 1980's (Figures 33 and 34).

Length frequency data from the ASMFC shrimp survey suggest that incoming year classes can be identified prior to their appearance in the NEFSC surveys. Thus, the ASMFC survey appears to be more useful in providing a pre-recruit index than in characterizing the population as a whole (Table 27, Figures 26 and 32). The ASMFC survey data indicate improved recruitment in recent years (see length frequency modes at 12 cm, corresponding to age 1 fish, during 1991-1994, 1997, and 1999 (Figure 32). Significant numbers of small fish were also observed in the NEFSC autumn survey during the same years (Figure 30).

Mean lengths at age from NEFSC spring and autumn surveys are presented in Table 30 and for ages 4 to 8 in Figures 35a and 35b. Mean lengths at age for ages 5 to 7 appear to have increased approximately 3-5 cm from 1980 to the late 1980's, and then declined (Figures 35a and 35b); however, Von Bertalanffy growth analyses detected no significant changes in resulting growth parameters over the time period.

NEFSC spring and autumn survey mean weights at age are given in Table 31 and Figures 36 and 37. Survey mean weights are variable, however, similar declines in mean weights for ages 6-9 were observed during the mid-1990s to 2002 in both the commercial landings and spring and autumn surveys.

MATURITY

Witch flounder maturity observations have been collected on the NEFSC research bottom trawl surveys since 1977. The NEFSC spring surveys were used for maturity analyses as these surveys occur closest to and prior to spawning (Halliday 1987). In the previous witch flounder assessment, probit analyses (SAS 1985) of maturity at age data revealed that there have been six maturity stanzas over the assessment period (GARM NEFSC 2002). The proportion at which 50% of the fish are mature at age (A_{50}) was significantly different for the time periods 1980-1982, 1983-1984, 1985-1990, 1991-1993, 1994-1999, and 2000-2002. Due to small sample sizes, it was necessary to pool individual years, however, individual years were examined, and then pooled into time blocks. Trends in female A_{50} and L_{50} were similar, progressively decreasing from 1980-1982 to 1985-1990, then increasing in 1991-1993, then declining in 1994-1999 and increasing in 2000-2002 similar to 1983-1984 levels. The maturity stanzas used revealed sharp changes in proportion mature, uncharacteristic of the assumed gradual biological process. The maturity stanzas also revealed, in a few instances, biologically infeasible outcomes, i.e. over the life span of a cohort, the proportion mature at age would decrease. Given these issues, a method which has been applied to Georges Bank cod (L. O'Brien, NEFSC, pers. comm.) was employed to minimize the abrupt changes yet still capture the changing trends in maturity over time. This method used logistic regression and a five-year moving time block to estimate annual maturity ogives. For example, the proportion mature in 1982 was estimated using NEFSC spring maturity data from 1980, 1981, 1982, 1983 and 1984. Likewise, the 1983 maturity ogive used maturity data from 1981 to 1985. Annual maturity ogives were derived for 1982 to 2001 using 1980 - 2003 data. The annual 2002 maturity ogive was assumed to be equal to the 2001 ogive (Table 32, Figure 38). In addition to the annual maturity ogives, a single ogive using maturity data from the entire time series was also calculated (Table 32). It was concluded that the moving time block method was appropriate for use in the VPA.

Stratified mean weight per tow of mature (spawning stock) witch flounder was calculated for spring NEFSC research vessel surveys (Table 33, Figure 39) using the six maturity stanzas. This analysis will be updated to incorporate the moving time-block maturity estimates in the next assessment update. The spawning stock biomass indices closely track total biomass indices except in most recent years, indicating a larger proportion of immature fish in the population.

MORTALITY

Natural Mortality

Burnett (MS 1987) estimated instantaneous natural mortality (M) to be 0.16 from a regression of survey-derived instantaneous total mortality (Z) estimates on commercial fishing effort.

Halliday (1973) used a value of $M = 0.15$ for females and $M = 0.2$ for males in an assessment of Scotian Shelf witch flounder. In the present study, virtual population analyses, yield per recruit and spawning stock biomass per recruit analyses were performed assuming $M = 0.15$.

Total Mortality

Estimates of instantaneous total mortality (Z) were computed from NEFSC spring and autumn research vessel bottom trawl survey catch per tow at age data by combining cohorts over the following time periods: 1982-1985, 1986-1989, 1990-1993, 1994-1997 and 1997-2001. Given the variability in age at full recruitment to the sampling gear observed during the survey time series (Table 34), estimates were derived for each time period and each season by taking the natural logarithm of the ratio of pooled age 7+ to pooled 8+. For example, the estimate of Z for 1982-1985 was computed as:

$$\begin{array}{ll} \text{Spring:} & \ln (\text{sum age } 7+ \text{ for } 1982-1985 / \text{sum age } 8+ \text{ } 1983-1986) \\ \text{Autumn:} & \ln (\text{sum age } 6+ \text{ for } 1981-1984 / \text{sum age } 7+ \text{ } 1982-1985). \end{array}$$

To evaluate Z over identical year classes within each of the survey series, different age groups were used in the spring and autumn.

Total mortality estimates from the two survey series exhibited similar trends, although autumn estimates were generally lower than those in the spring (Table 34 and Figure 40a). With no objective basis to select one survey series over another, total mortality was calculated by taking the geometric mean of the spring and autumn estimates during each time period. Total mortality ranged between 0.34 and 0.71 over the time series (Table 34). Additionally, annual estimates of total mortality were calculated, and smoothed with a three year moving average (Figure 40b).

ESTIMATION OF FISHING MORTALITY RATES AND STOCK SIZE

Virtual Population Analysis and Calibration

The ADAPT calibration method (Parrack 1986, Gavaris 1988, Conser and Powers 1990) was applied to estimate abundance at age in 2003 using catch-at-age estimates (i.e., landings plus discards from the shrimp and large-mesh otter trawl fishery; Table 21). Estimates of stock sizes, their associated statistics, and F in the terminal year are summarized in the Table 35.

New VPA software is now available in the NOAA Fisheries Toolbox. To bridge the transition between the software used in the last assessment update (FACT 1.5) and the current software, NFTv2.0.11, the accepted 2002 VPA (NEFSC 2002) formulation and input data was re-run using the NFTv2.0.11 software. The summary statistics of the two VPAs (RUN 61-f) reveal only slight changes in stock size estimates and fishing mortality (Table 35), and these minor changes are attributed to the use of the exact catch equation and other improvements in precision.

An initial formulation (RUN 100) based upon the 2002 VPA was performed to estimate 2003 stock sizes for ages 4 to 10 (Table 35) using a catch-at-age matrix including ages 3-11+ and NEFSC spring and autumn abundance indices for ages 3 to 11+ as tuning indices. All indices were given equal weighting. Autumn survey indices were lagged forward one year and one age to calibrate with beginning year population sizes of the subsequent year. A flat-top partial recruitment (PR) pattern was assumed, with full fishing mortality on ages 7 and older. The F on ages 10 and 11+ in the terminal year was estimated as the average of F on ages 7 through 9. The F on ages 10 and 11+ in all years prior to the terminal year was derived from weighted estimates of Z for ages 7 through 9. Instantaneous rate of natural mortality (M) was assumed to be 0.15. Spawning stock biomass (SSB) was calculated at time of spawning (March) and mean weight at age calculated by the Rivard method (Table 22).

The results of the initial run indicated that coefficients of variation (CV) for estimated ages ranged between 29% and 44% and the CVs for survey catchability coefficients (q) were consistent, ranging from 11% to 27%.

Two alternative formulations included: 1) using a total catch at age in which large-mesh otter trawl discards were estimated using the survey filter method for 1982-1994 and Fisheries Observer Program data for 1995 to 2002 [RUN 200]; and 2) estimating age 3 stock size using survey tuning indices [RUN 201]. Results from these alternative formulations provided estimates of stock size, F and spawning stock biomass consistent with the base run [RUN 100]. RUN 201 stock size for age 3 was poorly estimated (CV = 63%). Based on these runs, the partial recruitment pattern indicated that age 7 was not fully recruited. An alternative formulation (RUN 300) was conducted using a partial recruitment vector where the fully recruited age was increased from 7 to 8. Assuming full recruitment at age 8, the F on ages 10 and 11+ in the terminal year was estimated as the average of F on ages 8 and 9. The F on ages 10 and 11+ in all years prior to the terminal year was derived from weighted estimates of Z for ages 8 through 9. This partial recruitment pattern is consistent with recent mesh regulation changes.

The final formulation (RUN 301-f) included a 3 to 11+ catch at age with large-mesh otter trawl discards estimated using both the survey filter method and FOP data; an updated partial recruitment vector reflecting current management regulations was derived from the 1999-2002 F pattern taken from a penultimate calibration run; annual maturity ogives estimated by the five year moving time block with the 2002 maturity vector assumed to be equal to 2001. Ages 1 and 2 were deleted from the catch at age, this allowed recruitment in 2003 to be estimated using the geometric mean; there is no difference between 1-11+ vs 3-11+ on VPA results for fishing

mortality and spawning stock biomass. Based on the final formulation, two sensitivity analyses were conducted to evaluate the selection of tuning indices. The VPA was tuned with only NEFSC spring survey indices and then tuned with only NEFSC autumn survey indices (Table 35). Estimates of F and SSB from analyses using a single tuning series bounded the F and SSB estimated using both spring and autumn tuning indices. Using only the spring tuning series (RUN 301-f-spr), F was slightly higher ($F=0.43$) and SSB is slightly lower (15,798 mt) than the final run (RUN 301-f). Conversely, using only the autumn tuning indices (RUN 301-f-aut), F is slightly lower ($F=0.39$) and SSB is slightly higher (21,569 mt; Table 35) than the final run (RUN 301-f).

VPA Estimates of Fishing Mortality, Spawning Stock Biomass and Recruitment

A full listing of the final ADAPT VPA calibration output and diagnostics is presented in the Appendix and the results, including estimates of F, stock size and spawning stock biomass at age are given in Table 36. The mean residual for the VPA calibration was 0.791 and the CV on age 3-10 stock sizes ranged from 31% to 64% while the CVs on the estimates of survey catchabilities were between 13% and 26% (Appendix). The normalized survey indices and standardized residuals are presented in Figures 41 and 42.

The VPA indicates that fishing mortality (ages 8-9, unweighted) increased from 0.26 in 1982 to 0.67 in 1985, declined to 0.22 in 1992, increased to 1.13 in 1996, then declined to 0.41 in 2002 (Table 37 and Figure 43). Spawning stock biomass declined from 16,897 mt in 1982 to about 3,800 mt in 1996. With recent increases in recruitment and declines in fishing mortality, SSB has increased to 18,296 mt in 2002 (Table 37 and Figure 44). Since 1982 recruitment of age 3 has ranged from approximately 3 million fish (1984 year class) to 67.6 million fish (1997 year class; Table 37 and Figure 44). Over the 1982-2002 period, average recruitment of age 3 fish (the 1979 - 2000 year classes) was 19.6 million (the geometric mean equaled 14.4 million fish). The 1993-2000 year classes appear to be above average, and the 1997 year class is the largest in the VPA time series (Table 37 and Figure 44).

The relationship between spawning stock biomass and recruitment (age 3) is presented in Figure 45. The negative stock-recruitment relationship observed in previous assessments continues with the addition of the 2000 year class.

Precision of F and SSB

The uncertainty associated with the estimates of stock size and fishing mortality from the final VPA was evaluated using a bootstrap procedure (Efron 1982). One thousand bootstrap iterations were performed to derive standard errors, coefficients of variation (CVs) and bias estimates for the stock size estimates at the start of 2003, the catchability estimates (q) of the abundance indices used in calibrating the VPA, and the 2002 fully recruited fishing mortality rate (age 8+). Frequency distributions of the 2002 mean fishing mortality and spawning stock biomass bootstrap estimates were generated and cumulative probability curves produced (Figures 46 and 47).

Bootstrap results suggest that the CVs of the 2003 abundance estimates ranged between 32% to 84%, 24% for 2002 $F_{8,9}$ and 14% for 2002 spawning stock biomass. There is an 80% probability that the 2002 F (0.41) lies between 0.31 and 0.56 (Figure 46), and the 2002 SSB (18,296 mt) lies between 15,603 mt and 22,969 mt (Figure 47).

Retrospective Analyses

A retrospective analysis was conducted on the final VPA (Run 301-f) from 2002 to 1992 by sequentially removing the terminal year of the data to evaluate internal consistency of the current ADAPT formulation with respect to terminal estimates of F , SSB, and recruits at age 3 for the seven years prior to the current assessment. Results indicate that average F was underestimated (Figure 48a) and spawning stock biomass was consistently overestimated (Figure 48b). The retrospective analysis indicated that the number of age 3 recruits were generally overestimated, and the 1995-1997 year classes were considerably overestimated (Figure 48c).

Statistical Catch-at-age model

A statistical catch-at-age analysis was conducted for the witch flounder stock. An age-structured forward-projection model (a.k.a., age-structured production model) was fit to fishery and survey data during 1937-2002. This model provided an alternative long-term perspective on resource dynamics in comparison to VPA-based analyses that were limited to the period 1982-2002. Age-structured population dynamics of witch flounder were described using forward-projection methods for statistical catch-at-age analyses (Fournier and Archibald 1982, Methot 1990, Ianelli and Fournier 1998, Quinn and Deriso 1999). Models were fit to data with the AD Model builder software for nonlinear optimization (Otter Research 2001).

Six alternative statistical catch-at-age models were developed and fit. Brodziak and Wigley (2003 ms) contains a complete description of the basic model and input data. Common features of the six models were:

- Natural mortality was $M=0.15$ for all age classes.
- Catch scenario 2 was used (same as used in VPA).
- Fishery selectivity was estimated for historic (1937-1993) and current (1994-present) time periods.
- NEFSC spring and fall survey biomass and numbers at age data were used.
- Emphasis values for likelihood components were: Recruitment $\lambda_1=10$, Fishery age composition $\lambda_2=1$, NEFSC Fall survey age composition $\lambda_3=1$, NEFSC Fall survey biomass index $\lambda_4=100$, NEFSC Spring survey age composition $\lambda_5=1$, NEFSC Spring survey biomass index $\lambda_6=100$, Catch biomass $\lambda_7=100$, Fishing mortality $\lambda_8=1$, Fishing mortality penalty $\lambda_9=1$

The primary differences among the six alternative models were:

1. Dome-shaped selectivity possible for fishery, spring, and fall surveys; time frame is 1937-2002.

2. Flat-topped selectivity for fishery, spring, and fall surveys; time frame is 1937-2002.
3. Dome-shaped selectivity possible for fishery, spring, and fall surveys; time frame is 1963-2002.
4. Flat-topped selectivity for fishery, spring, and fall surveys; time frame is 1963-2002.
5. Flat-topped selectivity for fishery and spring survey; Dome-shaped selectivity possible for fall survey; time frame is 1963-2002.
6. Flat-topped selectivity for fishery and spring survey; Dome-shaped selectivity possible for fall survey; time frame is 1937-2002.

Models 1, 2, and 6 were considered to be the primary models, while models 3, 4, and 5 provided sensitivity analyses to the choice of time frame. The Northern Demersal Working Group (WG) reviewed the model diagnostics. In general, the selectivity patterns of models that allowed dome-shaped fishery or survey selectivity appeared to be too sharply domed to be biologically plausible. In contrast, models with the assumption of flat-topped selectivity provided a poorer fit to the data, as measured by the root-mean squared errors for the NEFSC fall and spring survey biomass index and the catch biomass fits to the data. The WG chose to reduce the emphasis on the NEFSC fall and spring survey biomass index and the catch biomass likelihood components to 10 down from 100. This choice alleviated the problem of implausible selectivity patterns in the fishery and the survey. As a result, the WG concluded that Model 1 with reduced emphasis values was the best alternative of the statistical catch-at-age analyses (SCAA). Model results are reported to confirm the basic trends of VPA-based results and show the likely effect of extending the assessment time horizon back to 1937.

Model results showed that current fishery selectivity at age was estimated to be lower at ages 1-6 than historic selectivity (Figure 49). This was consistent with increases in fishery mesh size and changes in discarding practices (e.g., shrimp fishery) that occurred around 1994. The resulting catch biomass predictions generally matched observed catch biomass (Figure 50) with some moderate deviations in the early 1980s.

Model results showed that NEFSC fall survey selectivity was dome-shaped with a peak at age-5 (Figure 51). The NEFSC spring survey selectivity was flat-topped with full selection occurring at roughly age-7. The resulting predicted NEFSC fall and spring survey indices generally matched the trends in observed indices (Figures 52 and 53). Both surveys indicate a longterm decline in biomass from the 1970s through the early 1990s. Biomass increases in the late-1990s differed moderately between the fall and spring surveys

There was general agreement between VPA and SCAA results during 1982-2002. Spawning biomass estimates were very similar during 1989-1999 (Figure 54, SCAA estimate of 10.5 kt in 2002). The VPA indicates a smaller decrease in spawning biomass during 1982-1988 and a greater increase during 2000-2002. Fishing mortality estimates were also similar (Figure 55, SCAA estimate of 0.48 in 2002). Both VPA and SCAA estimates increased to roughly the mid-1990s and then declined. Recruitment estimates also exhibited similar patterns (Figure 56, SCAA estimate of 14.1 million age-1 fish in 2002), although the VPA indicated larger increases

in recruitments during the late-1990s. Despite differences in model configuration and estimation approach, the SCAA generally confirmed point estimates and trends in the VPA results.

BIOLOGICAL REFERENCE POINTS

Yield-per-recruit (Y/R) and spawning stock biomass per recruit (SSB/R) analyses were performed using the Thompson and Bell (1934) method for witch flounder ages 3 to 20. Input vectors for partial recruitment, maturation at age and mean weights at age were all updated since the last assessment. Mean weights at age used in the Y/R analyses were computed as an arithmetic average of catch mean weights at age (Table 21) over the 1999-2002 period. Mean weights at age for use in the SSB/R analyses were derived by applying the length-weight relationship for witch flounder to predicted lengths at age from von Bertalanffy growth curve analyses of NEFSC survey data from 1980-2002. The maturation ogive from the entire time series (1980-2003) was also used (Table 32). Given the changes in regulated mesh size in 1999, the exploitation pattern used in the yield and SSB per recruit analyses and short-term projections was computed from the 1999-2002 VPA results. Geometric mean F at age was computed for the 1999-2002 period and divided by the geometric mean of the fully recruited annual Fs to derive the partial recruitment vector. The final exploitation pattern was smoothed, applying full exploitation on ages 8 and older, viz.

Age 3	Age 4	Age 5	Age 6	Age 7	Ages 8+
0.0036	0.0229	0.0703	0.1931	0.5282	1.000

The input data and results for the Y/R and SSB/R analyses are given in Table 38 and Figure 57. The reference points were $F_{0.1} = 0.196$, $F_{max} = 0.545$, and $F_{40\%} = 0.230$.

The biological reference points were updated by applying the approach used to estimate MSY proxies for witch flounder (NEFSC 2002). F_{msy} is approximated as $F_{40\%}$ (0.23), the SSB_{msy} proxy is 25, 248 mt, the product of 40%MSP (1.2882 kg spawning biomass) and average long-term recruitment (19.6 million). The MSY proxy is 4,375 mt, the product of yield per recruit at $F_{40\%}$ (0.2232 kg) and average recruitment.

In 2002, spawning stock biomass was slightly greater than $\frac{1}{2}$ SSB_{msy} (12,624 mt), the minimum stock size threshold, and fishing mortality in 2002 was nearly double F_{msy} , the maximum fishing mortality threshold; therefore, witch flounder was not overfished but overfishing was occurring in 2002 (Figure 58).

To evaluate the effects of simultaneous changes in the three input vectors described above (i.e. partial recruitment, maturation and mean weights) on $F_{40\%}$, Y/R, SSB/R and the SSB_{msy} proxy, a decomposition analysis (P.Rago, NEFSC, pers. comm.) was conducted. This analysis is analogous to decomposing a sum of squares in an analysis of variance (decomposing the total resulting difference into its components).

For F40%, Y/R and SSB/R:

Total effect = effect of vector 1 + effect of vector 2 + effect of vector 3 + interaction terms.

For SSBmsy:

Total effect = SSB/R effect + Recruit effect + interaction term.

The effect is the difference between the former YPR estimate and the current YPR estimate, for F40%, Y/R, SSB/R and for SSBmsy.

To accomplish this, the former YPR analysis (Run 0) was re-run using ages 3-20 to coincide with the ages used in the current YPR analysis (Run 1). Then, YPR analyses were conducted where each former vector was replaced with a current vector (Runs 2 through 7), until all vectors were replaced with current vectors (Run 8). The resultant F, Y/R and SSB/R at 40%MSP from each run (Runs 0 to 8) are reported in Table 39. The total effect of changing all three vectors at once equals Run 1 - Run 8.

Results of the decomposition analysis (Table 39) indicate that changes in F40% were effected most by new partial recruitment vector. Changes in Y/R resulted from the interaction of all three new input vectors while changes in the SSB/R resulted from the interaction between the mean weights and maturity vectors. Changes in SSBmsy were effected most by changes in new mean age 3 recruitment.

SHORT-TERM PROJECTIONS FOR 2004 AND 2005

Short-term stochastic projections were performed to estimate landings, discards and SSB during 2003-2005 under various F scenarios using bootstrapped VPA calibrated stock sizes in 2002. The partial recruitment, maturity ogive, and mean weights at age were the same as described in the yield and SSB per recruit section (Table 40). Recruitment (age 3) in 2003-2005 was derived by re-sampling the cumulative density function based on the empirical observations during 1982-2001 (1979-2000 year classes). Fishing mortality was apportioned among landings and discards based on the proportion observed landed at age during 1999-2002. The proportion of F and M which occurs before spawning equals 0.1667 (March 1); M was assumed to be 0.15. Spawning stock biomass in 2002 was estimated to be 18,296 mt. The F scenarios are: status quo $F_{2003} = 0.41$, $F_{msy} = 0.230$, 75% of $F_{msy} = 0.17$ and $landings_{2003} = landings_{2002}$ ($F = 0.199$). Fishing at the status quo F (0.41) or at the target ($F_{msy} = 0.23$) in 2003 - 2005 is expected to allow biomass to increase above SSBmsy and initiate rebuilding of the age structure (Table 40). Comparison of the current age structure and the age structure under MSY conditions are given in Figure 59.

CONCLUSIONS

Based on the ADAPT VPA, the witch flounder stock was not overfished, but overfishing was occurring in 2002. Fully recruited fishing mortality in 2002 was 0.41, nearly double F_{msy}

(0.23), and spawning stock biomass was estimated to be 18,296 mt in 2002, 72% of SSB_{msy} (25,248 mt) . Recent year classes appear to be above average. Although the spawning stock biomass has increased, the age structure still remains truncated. Fishing mortality should be reduced to F_{msy} or below to allow the age structure to rebuild.

ACKNOWLEDGMENTS

We would like to recognize all those who diligently collected the biological samples from the commercial fisheries and from research vessel surveys. We wish to express our appreciation to Betty Holmes for her technical assistance. We thank Jay Burnett for the age determinations. We thank all members of the Stock Assessment Workshop's Northern Demersal Working Group and the SARC panel for their review and helpful comments.

LITERATURE CITED

- Azarovitz, T.R. 1981. A brief historical review of the Woods Hole Laboratory trawl survey time series. *In*: Doubleday, W.G. and D. Rivard (eds.), Bottom Trawl Surveys, p. 62-67. Can. Spec. Publ. Fish. Aquat. Sci. 58.
- Brodziak, J., and S. Wigley. 2003 manuscript. Statistical catch-at-age analysis of the witch flounder stock, 1937-2002. Northern Demersal Working Group Meeting May 12-16 2003, Working Paper W3, NEFSC, Woods Hole, MA 02543.
- Burnett, J.M. MS 1987. The population biology of the witch flounder, *Glyptocephalus cynoglossus* (L.), in the Gulf of Maine-Georges Bank region. M.Sc. Thesis, Department of Wildlife and Fisheries Biology, University of Massachusetts, Amherst, MA, 116 p.
- Burnett, J. and S.H. Clark. 1983. Status of witch flounder in the Gulf of Maine - 1983. NMFS/NEFC, Woods Hole Laboratory Ref. Doc. No. 83-36, 31 p.
- Burnett, J., M.R. Ross, and S.H. Clark. 1992. Several biological aspects of the witch flounder (*Glyptocephalus cynoglossus* (L.)) in the Gulf of Maine-Georges Bank region. *J. Northw. Atl. Fish. Sci.* 12: 15-25.
- Cadrin, S.X. 2003. Stock assessment of yellowtail flounder in the southern New England - Mid-Atlantic area (SAW 36). NEFSC CRD 03-03.
- Cadrin, S. X. 1999. A precautionary approach to fishery control rules based on Surplus production modeling. NOAA Tech. Memo. NMFS-F/SPO-40. 17-22 pp.
- Clark, S.H. 1981. Use of trawl survey data in assessments. *In*: Doubleday, W.G. and D. Rivard (eds.), Bottom Trawl Surveys, p. 82-92, Can. Spec. Publ. Fish. Aquat. Sci. 58.

- Conser, R.J. and J.E. Powers. 1990. Extensions of the ADAPT VPA tuning method designed to facilitate assessment work on tuna and swordfish stocks. *Int. Comm. Conserv. Atlantic Tunas, Coll. Vol. Sci. Pap.*, 32:461-467.
- Delong, A., K. Sosebee, and S. Cadrin. 1997. Evaluation of vessel logbook data for discard and CPUE estimation. NEFSC Laboratory Reference Document 97-xx.
- Efron, B. 1982. The jackknife, the bootstrap and other resampling plans. *Phila. Soc. for Ind. and Appl. Math.* 38: 92 p.
- Fournier, D. A., and C. P. Archibald. 1982. A general theory for analyzing catch at age data. *Can. J. Fish. Aquat. Sci.* 39:1195-1207.
- Gavaris, S. 1988. An adaptive framework for the estimation of population size. CAFSAC Res. Doc. 88/29, 12 p.
- Halliday, R.G. 1973. The flatfish fisheries of the Scotian Shelf. *Int. Comm. Northw. Atl. Fish., Res. Doc.* 73/103, Ser. No. 3064 (mimeo), 46 p.
- Halliday, R.G. 1987. Size and age at sexual maturity of Atlantic argentine, *Argentina silus*: a critique. *Environ. Biol. Fishes* 19(2):165-261.
- Howe, A.B., F.J. Germano, J.L. Buckley, D. Jimenez, and B.T. Estrella. 1981. Fishery resource assessment, coastal Massachusetts. Completion Report, Massachusetts Division of Marine Fisheries, Commercial Fisheries Review Div. Project 3-287-R-3.
- Ianelli, J. N., and D. A. Fournier. 1998. Alternative age-structured analyses of NRC simulated stock assessment data. NOAA Tech. Memo. NMFS-F/SPO-30. pp. 81-96.
- Lange, A.M.T. and F.E. Lux. 1978. Review of the other flounder stocks (winter flounder, American plaice, witch flounder, and windowpane flounder) off the northeast United States. NMFS, NEFC, Woods Hole Lab. Ref. Doc. No. 78-44, 53 pp.
- Mayo, R.K., L. O'Brien, and N. Buxton. 1992. Discard estimates of American plaice, *Hippoglossoides platessoides*, in the Gulf of Maine northern shrimp fishery and the Gulf of Maine-Georges Bank large-mesh otter trawl fishery. Appendix to NEFSC Ref. Doc. 92-07, Res. Doc. SAW 14/3, 40 p.
- Methot, R. D. 1990. Synthesis model: an adaptive framework for analysis of diverse stock assessment data. *Int. North Pac. Fish. Comm. Bull.* 50:259-277.
- NEFSC [Northeast Fisheries Science Center]. 2002. A Report of the Groundfish Assessment Review Meeting (GARM), October 2002. NEFSC Ref. Doc. 02-16, 511p.

- NEFSC [Northeast Fisheries Science Center]. 2002. Final Report of the Working Group on Re-Evaluation of biological Reference Points for New England Groundfish. NEFSC Ref. Doc. 02-04.
- NEFSC [Northeast Fisheries Science Center]. 1994. Report of the 18th Northeast Regional Stock Assessment Workshop (18th SAW), Stock Assessment Review Committee (SARC) consensus summary of assessments. NEFSC Ref. Doc. 94-22, 281 p.
- Northern Shrimp Technical Committee. MS 1984. Results of the 1983 northern shrimp survey in the western Gulf of Maine, August 1983. Unpublished Report, Woods Hole, MA, 16 p.
- O'Brien, L. and R.W. Brown. 1996. Assessment of the Georges Bank Haddock stock for 1994. NEFSC Ref. Doc. 95-13, 108 p.
- Otter Research. 2001. An introduction to AD Model Builder Version 6.0.2 for use in nonlinear modeling and statistics. Otter Research, Ltd. Sidney, B.C. V8L 3S3, Canada.
- Parrack, M.L. 1986. A method of analyzing catches and abundance indices from a fishery. Int. Comm. Conserv. Atlantic Tunas, Coll. Vol. Sci. Pap., 24:209-211.
- Quinn, T. P., II, and R. B. Deriso. 1999. Quantitative fish dynamics. Oxford University Press, New York, 542 pp.
- Rivard, D. 1980. APL programs for stock assessment. Can. Tech. Rep. Fish. Aquat. Sci. 953.
- SAS. 1985. SAS User's Guide: Statistics, version 5. SAS Institute, Cary, NC. 956 p.
- Thompson, W.F. and F.W. Bell. 1934. Biological statistics of the Pacific halibut fishery. 2. Effect of changes in intensity upon total yield and yield per unit of gear. Rep. Int. Pac. Halibut Comm. 8: 49 p.
- Walsh, S.J., R.B. Millar, C.G. Cooper, and W.M. Hickey. 1992. Codend selection in American plaice: diamond versus square mesh. Fish. Res., 13:235-254.
- Wigley, S.E. MS 1994. Estimation of foregone yield associated with the discarding of witch flounder (*Glyptocephalus cynoglossus*) in the Gulf of Maine northern shrimp (*Pandalus borealis*) fishery, 1982-1992. M.S. thesis. North Carolina State University, Raleigh, NC. 91 p.
- Wigley, S.E. and R.K. Mayo. 1994. Assessment of the Gulf of Maine-Georges Bank witch flounder stock for 1994. Northeast Fisheries Science Center Reference Document 94-17, 86 p.

Wigley, S.E. , J.K.T. Brodziak and S.X. Cadrin. 1999. Assessment of the witch flounder stock in Subareas 5 and 6 for 1999. Northeast Fisheries Science Center Reference Document 99-16, 153 p.

Table 1. Significant changes in management regulations governing the USA commercial fishery for witch flounder (adapted from O'Brien and Brown, 1996), updated by Tom Nies, NEFMC (pers. comm.,2003).

<u>1953-1977</u>		<u>ICNAF era</u>
1953		Minimum mesh in body and codend 4 ½ inches.
1970		Areas 1(A) and 2(B) closed during haddock spawning, from March - April.
1972-1974		Areas 1(A) and 2(B) closure extended to March - May.
1975		Areas 1(A) and 2(B) closures extended to February - May.
<u>1977 - Present</u>		<u>Extended Jurisdiction and National Management</u>
1977		USA Fishery Conservation and Management Act of 1976 (FCMA) in effect.
1977-1982		Fishery Management Plan (FMP) for Atlantic groundfish: Seasonal spawning closures for haddock (Areas 1 and 2), quotas for haddock, etc
1982		Mesh size 5 1/8 inches (130 mm).
1982-1985		The 'Interim Plan' for Atlantic groundfish: Eliminated all catch controls, retained closed area and mesh size regulations, implemented minimum landing sizes.
1983		Mesh size increased to 5 ½ inches (140 mm). Witch flounder minimum size 33 cm.
1984	October	Hague Line separating USA and Canadian fishing zones in the Gulf of Maine and Georges Bank region.
1985		Fishery Management Plan for the Multispecies Fishery.
1987		Witch flounder minimum size increased to 36 cm.
1991		Amendment 4 established overfishing definitions for witch flounder in terms of Fmed (F20%) replacement levels.
1993		Area 2 closure is extended from January 1 - June 30.
1994	January May December	Amendment 5 implemented: expanded Area 2, Area 1 closure not in effect. 6 inch (152 mm) mesh restriction implemented (delayed from March 1). Square or diamond mesh allowed. Area 1, Area 2 and Nantucket Lightship Area closed year-around.
1996	July	Amendment 7 implemented: Days-at-sea (DAS) restrictions. Haddock trip limits Raised to 1000 pounds
1997	May	Additional scheduled DAS restrictions from Amendment 7 accelerated.
1998	May	Western Gulf of Maine Closure Area adopted: Jeffery's Ledge area closed to all groundfishing. Rolling closures in the western Gulf of Maine.
	October	Amendment 9: revised overfishing definitions as required by Sustainable Fisheries Act.
1999	May	Codend mesh regulations changed to 6-inch diamond mesh, 6 1/2-inch square mesh. Additional rolling closures adopted in the western Gulf of Maine Cashes Ledge seasonal closure adopted Roller gear limited to a maximum of 12 inches in an area of the western Gulf of Maine. Gulf of Maine cod trip limit ranged from 30 to 400 lbs. in this fishing year.

continued.

Table 1 continued. Significant changes in management regulations.

2000	May	May closure implementation on northern Georges Bank. Changes to large mesh permit category, granting additional DAS to vessels using larger than 6-inch diamond / 6-inch square mesh.
2002	June 1	Additional restrictions adopted during this fishing year (result of lawsuit over FW33): Vessels limited to 25% of allocated DAS May to July; Increase in minimum mesh size for trawl vessel to 6 ½ inch diamond/ 6 ½ inch square; Reduced number of rolling closures in the western Gulf of Maine (effective in January 2003, with result there were additional rolling closures in calendar year 2002 compared to calendar year 2001); Cashes Ledge seasonal closure expanded to year-around closure; Increase in GOM cod trip limit to 500 lbs. per day/4,000 lbs per trip; Increase in mesh size for large mesh permit category.
2002	August 1	Reduction in allocated DAS based on past history of use for each permit; Front-loading of DAS clock prohibited; Additional restrictions on number and deployment of gillnets.

Table 2. Significant changes in management regulations governing the USA northern shrimp fishery in the Gulf of Maine.

1972	Adoption of mesh regulations, Establishment of count/pound limits, Establishment of closed areas.
1973	Adoption of interim minimum mesh regulations; mesh size not less than 1.5 inches (38 mm).
1975	Minimum mesh regulations increased to 1.75 inches (44 mm). Harvest restricted to 4,200 mt (9.2 million pounds). Fishery closure from July - September.
1976	Harvest restricted to 2,300 mt (5 million pounds) by season closure and quota management. Open season: January 1 - April 15 1976.
1977	Harvest restricted to 1,600 mt (3.5 million pounds). Open season: January 1 - May 15 1977.
1978	Closure of fishery.
1979	Open season: February 1 - March 31, 1979
1980	Open season: February 15 - May 31, 1980
1981	Open season: February 15 - May 31, 1981
1982	Open season: January 1 - April 15, 1982
1983	Open season:
1984	Open season:
1985	Open season:
1986 - 1991	Open season: 183 days, by-catch limit of 10% by weight of groundfish allowed.
1992 April	Nordmore grate regulation (max 25 mm space); no bycatch of groundfish allowed, no Sunday fishing
1993	Open season: Dec 15 - May 15, no Sundays, separator gear Dec 15 - Mar 15; grate Apr-May15
1994	Open season: December 15 - April 15
1995	Open season: December 1 - May 31 with 1 day off per week.
1996	Open season: December 1 - May 31 with 1 week off.
1997	Open season: December 1 - May 27 with 4 or 5 day block off per month.
1998	Open season: December 8 - May 22 with 3 weekends of no fishing
1999	Open season: December 1 - May 30, no weekend fishing.
2000	Open season: (51 day season)
2001	Open season: Jan 9 - March17; April 16-30 and no days off (83 day season)
2002	Open season: February 15 - March 11 (25 day season)
2003	Open season: Jan 15 - Feb 27 no Friday fishing (38 day season)

Table 3. Witch flounder landings, discards and catch (metric tons, live) by country, 1937-2002
 [1937-1959 provisional landings reported in Lange and Lux, 1978; 1960-1963 reported to
 ICNAF/NAFO (Burnett and Clark, 1983)].

Year	LANDINGS						USA Discards	USA Catch
	USA Subarea 4, 5 & 6	USA Subarea 3	USA Total	CAN	Other	Total		
1937			5000			5000		
1938			3600			3600		
1939			3100			3100		
1940			3000			3000		
1941			2000			2000		
1942			1800			1800		
1943			1000			1000		
1944			1000			1000		
1945			1000			1000		
1946			1500			1500		
1947			1500			1500		
1948			1000			1000		
1949			3600			3600		
1950			3000			3000		
1951			2600			2600		
1952			3700			3700		
1953			4200			4200		
1954			4000			4000		
1955			2400			2400		
1956			2000			2000		
1957			1000			1000		
1958			1000			1000		
1959			1000			1000		
1960	1255		1255			1255		
1961	1022		1022	2		1024		
1962	976		976	1		977		
1963	1226		1226	27	121	1374		
1964	1381		1381	37		1418		
1965	2140		2140	22	502	2664		
1966	2935		2935	68	311	3314		
1967	3370		3370	63	249	3682		
1968	2807		2807	56	191	3054		
1969	2542		2542		1310	3852		
1970	3112		3112	19	130	3261		
1971	3220		3220	35	2860	6115		
1972	2934		2934	13	2568	5515		
1973	2523		2523	10	629	3162		
1974	1839		1839	9	292	2140		
1975	2127		2127	13	217	2357		
1976	1871		1871	5	6	1882		
1977	2469		2469	11	13	2493		
1978	3501		3501	18	6	3525		
1979	2878		2878	17		2895		
1980	3128		3128	18	1	3147		
1981	3442		3442	7		3449		

continued.

Table 3 continued. Witch flounder landings, discards and catch (metric tons, live).

Year	LANDINGS							
	USA Subarea 4, 5 & 6	USA Subarea 3	USA Total	CAN	Other	Total	USA Discards	USA Catch
1982	4906		4906	9		4915	48	4954
1983	6000		6000	45		6045	162	6162
1984	6660		6660	15		6675	100	6760
1985	6130	255	6385	46		6431	61	6191
1986	4610	539	5149	67		5216	25	4635
1987	3450	346	3796	23		3819	47	3497
1988	3262	358	3620	45		3665	60	3322
1989	2068	297	2365	13		2378	133	2201
1990	1465	2	1467	12		1479	184	1649
1991	1777		1777	7		1784	95	1872
1992	2227		2227	7		2234	171	2398
1993	2601		2601	10		2611	376	2977
1994	2665		2665	34		2699	422	3087
1995	2209		2209	11		2220	193	2402
1996	2087		2087	10		2097	254	2341
1997	1771		1771	7		1778	300	2071
1998	1848		1848	10		1858	286	2134
1999	2121		2121	19		2140	213	2334
2000	2439		2439	53		2492	115	2554
2001	3019		3019	32		3051	224	3243
2002	3186		3186			3186	279	3465

Table 4. Percentage of USA commercial witch flounder landings (mt) by Statistical Area, 1973 - 2002.

YEAR	Statistical Areas																							TOTAL		
	300	400	464	465	466	500	510	511	512	513	514	515	520	521	522	523	524	525	526	530	537	538	539		540	600
1973	-	1.1	-	0.8	-	-	-	4.0	9.4	18.6	13.8	1.5	-	10.5	16.3	0.8	2.9	7.6	10.7	-	1.0	0.0	0.2	-	0.6	100.0
1974	-	2.7	-	0.1	0.2	-	-	1.0	4.1	17.3	11.6	1.3	-	18.2	16.0	0.9	5.7	7.9	10.4	-	2.2	0.1	0.1	-	0.2	100.0
1975	-	0.7	-	0.8	0.0	-	-	0.8	7.1	16.9	13.6	4.3	-	17.4	11.2	0.5	7.5	13.2	4.9	-	0.6	0.0	0.1	-	0.2	100.0
1976	-	1.2	-	0.3	0.1	-	-	1.3	7.5	25.1	19.5	2.0	-	14.9	11.2	1.3	4.3	7.7	2.7	-	0.6	0.1	0.1	-	0.2	100.0
1977	-	0.2	-	0.2	0.1	-	-	0.6	7.8	30.6	27.6	4.1	-	10.4	10.1	0.8	2.5	2.9	1.2	-	0.5	0.1	0.2	-	0.1	100.0
1978	-	0.3	-	0.1	-	-	-	0.2	9.5	39.1	18.3	4.7	-	10.5	8.7	2.4	2.5	1.1	1.3	-	0.6	0.2	0.1	-	0.3	100.0
1979	-	0.2	-	0.0	-	-	-	2.3	9.4	35.6	14.5	4.2	-	12.8	13.7	3.4	1.2	0.5	1.0	-	0.7	0.0	0.1	-	0.5	100.0
1980	-	0.1	-	0.2	-	-	-	1.4	8.9	42.2	12.3	8.2	-	10.1	7.4	2.1	0.8	1.2	3.5	-	0.6	0.0	0.2	-	0.5	100.0
1981	-	0.2	-	1.0	-	-	-	1.9	9.2	41.0	12.2	9.4	-	11.3	5.3	2.0	1.8	1.4	1.2	-	1.1	0.0	0.3	-	0.8	100.0
1982	-	0.4	-	0.7	-	-	0.0	3.1	15.5	29.2	8.7	15.5	-	11.4	5.9	2.4	1.1	1.4	2.0	-	1.0	0.1	0.2	-	1.3	100.0
1983	-	0.5	-	2.4	-	-	-	4.2	20.6	24.3	8.0	17.4	-	9.3	5.4	2.0	0.8	1.1	1.7	-	1.5	0.0	0.2	-	0.7	100.0
1984	-	0.2	-	2.2	-	-	-	2.4	11.3	23.5	11.8	19.8	-	12.0	6.5	2.3	1.0	1.8	2.7	-	1.5	0.0	0.1	-	0.9	100.0
1985	4.0	0.1	-	1.1	-	-	-	3.7	11.8	23.1	10.3	19.8	-	11.5	7.3	2.0	1.0	1.6	1.7	-	0.5	0.0	0.0	-	0.6	100.0
1986	10.5	0.2	-	1.3	0.0	-	-	4.0	14.9	23.6	9.1	15.3	-	9.3	5.8	1.9	0.4	0.6	1.5	-	0.6	0.0	0.0	-	1.0	100.0
1987	9.1	0.1	-	0.4	-	-	-	2.7	11.6	27.4	9.6	19.0	-	9.1	5.6	1.4	0.5	0.7	1.2	-	0.4	0.0	0.0	-	1.1	100.0
1988	9.9	-	-	0.3	-	-	-	2.6	8.0	26.5	9.7	17.0	-	12.4	5.7	1.5	1.0	2.7	1.3	-	0.4	0.0	0.0	-	1.1	100.0
1989	12.5	0.0	-	0.1	-	-	-	1.3	7.4	21.8	9.4	16.1	-	12.8	5.7	1.6	1.2	2.2	5.4	-	0.9	0.1	0.0	-	1.3	100.0
1990	0.1	0.3	-	0.1	-	-	-	1.6	9.1	29.0	12.4	12.7	-	11.1	5.5	2.4	2.4	3.7	5.2	-	2.6	0.0	0.1	-	1.6	100.0
1991	-	0.1	-	0.1	-	-	-	1.1	9.3	26.1	11.0	15.6	-	8.1	7.7	2.4	3.0	2.0	4.8	-	4.7	0.1	0.1	-	3.7	100.0
1992	-	0.0	-	-	-	-	-	0.6	10.5	23.2	10.1	14.8	-	6.8	8.4	2.0	1.7	2.8	9.8	-	6.4	0.0	0.2	-	2.8	100.0
1993	-	0.5	-	-	-	-	-	0.5	6.7	22.3	16.1	16.2	-	6.9	10.4	3.1	2.5	3.6	5.1	-	3.8	0.0	0.1	-	2.2	100.0
1994*	-	-	0.1	-	-	0.4	0.3	1.7	13.1	15.5	15.5	13.5	0.1	14.3	12.2	2.6	1.5	2.1	1.6	0.1	2.7	0.4	0.1	0.2	1.8	100.0
1995*	-	-	0.5	0.5	-	0.6	0.2	1.1	6.8	14.1	15.2	20.6	0.3	17.3	15.0	1.9	0.7	1.5	0.6	0.2	1.1	0.2	0.0	0.1	1.6	100.0
1996*	-	-	0.1	0.1	-	0.8	1.2	1.7	6.3	18.1	13.8	20.9	1.2	13.7	14.1	2.1	0.4	2.2	0.3	0.0	1.3	0.4	0.1	0.1	1.0	100.0
1997*	-	-	-	0.1	-	1.1	0.7	0.7	9.3	16.5	12.6	21.9	0.6	11.0	16.1	2.7	0.5	3.2	0.7	-	1.2	0.3	0.1	-	0.8	100.0
1998*	-	-	-	0.1	-	1.2	0.1	0.7	8.3	14.5	11.1	21.8	0.2	15.1	16.2	3.5	1.3	2.5	0.5	0.1	1.1	0.6	0.3	0.1	0.5	100.0
1999*	-	-	-	0.1	-	0.4	0.1	0.7	8.2	12.0	11.9	15.8	1.5	17.7	20.9	2.9	1.3	3.0	0.5	-	1.8	-	0.1	-	1.1	100.0
2000*	-	-	-	0.1	-	0.3	0.1	1.0	5.6	12.4	14.5	12.9	0.2	22.8	20.6	2.5	1.0	2.4	0.3	-	0.8	0.2	0.3	0.1	2.0	100.0
2001*	-	-	0.1	0.1	-	-	0.1	1.7	5.2	14.1	15.6	11.2	-	24.8	18.0	4.8	0.5	1.5	0.4	-	0.8	0.1	0.2	-	0.8	100.0
2002*	-	-	-	0.1	-	-	-	1.5	5.5	15.3	23.0	10.5	-	18.4	16.9	3.2	1.6	2.2	0.5	-	0.2	0.3	0.2	0.1	0.5	100.0

¹Note: USA portions of SA 523 and 524 were renamed 561 and 562, respectively, in 1985.

* 1994-2002 spatial distribution based upon Vessel Trip Report data, considered provisional.

Table 5. Percentage of annual USA commercial witch flounder landings by gear type, 1973-2002.

Year	Otter Trawl	Shrimp Trawl	Other	Total
1973	98.7	-	1.3	100.0
1974	99.7	-	0.3	100.0
1975	97.3	2.5	0.2	100.0
1976	98.8	0.9	0.3	100.0
1977	97.4	1.5	1.1	100.0
1978	98.1	-	1.9	100.0
1979	97.9	0.2	1.9	100.0
1980	96.6	0.6	2.8	100.0
1981	97.3	0.8	1.9	100.0
1982	96.8	0.9	2.3	100.0
1984	96.4	0.4	3.2	100.0
1985	95.1	1.0	3.9	100.0
1986	95.9	1.1	3.0	100.0
1987	95.5	1.1	3.4	100.0
1988	96.0	0.8	3.2	100.0
1989	95.3	0.4	4.3	100.0
1990	92.8	0.6	6.6	100.0
1991	95.1	0.5	4.4	100.0
1992	96.2	0.1	3.7	100.0
1993	94.2	0.0	5.8	100.0
1994	96.2	0.0	3.8	100.0
1995	96.1	0.0	3.9	100.0
1996	96.7	0.0	3.3	100.0
1997	96.9	0.0	3.1	100.0
1998	97.5	0.0	2.5	100.0
1999	97.4	0.0	2.6	100.0
2000	97.5	0.0	2.5	100.0
2001	97.5	0.0	2.5	100.0
2002	97.8	0.0	2.2	100.0

Table 6. Percentage of annual USA commercial witch flounder landings by month for all gear types, 1973-2002.

YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	TOTAL
1973	5.0	3.2	8.8	10.9	14.2	14.4	10.8	10.7	5.3	7.6	5.6	3.6	100.0
1974	4.6	3.6	5.6	11.3	16.8	12.1	14.0	10.3	7.3	6.1	4.8	3.4	100.0
1975	3.7	4.3	6.1	10.7	14.3	13.2	13.8	9.1	7.8	6.2	6.9	3.9	100.0
1976	3.2	4.8	7.5	12.2	12.4	16.1	14.0	8.7	7.9	5.2	4.2	3.7	100.0
1977	4.3	6.1	10.2	9.2	9.2	10.9	12.0	10.9	7.8	6.7	7.1	5.4	100.0
1978	5.9	10.6	5.9	10.4	12.9	12.9	10.1	7.8	6.3	7.7	5.0	4.4	100.0
1979	7.2	5.2	8.1	7.2	9.0	15.5	12.1	9.3	7.6	6.9	6.7	5.2	100.0
1980	5.1	8.9	7.2	9.9	10.5	13.6	11.2	9.8	7.9	6.5	4.6	4.9	100.0
1981	7.5	5.8	10.5	8.0	9.0	13.0	9.2	7.8	8.5	6.9	6.6	7.2	100.0
1982	5.9	6.4	10.7	8.8	9.7	9.3	9.2	8.4	7.4	8.2	7.8	8.0	100.0
1983	9.2	8.6	11.4	12.4	10.0	9.6	7.2	7.6	6.3	5.7	5.3	6.8	100.0
1984	9.7	9.1	10.6	11.0	10.8	8.8	6.8	6.9	6.3	7.9	6.2	6.1	100.0
1985	8.0	7.3	8.8	12.6	12.5	8.1	8.3	7.7	7.9	7.4	5.3	6.1	100.0
1986	7.1	7.2	8.1	12.4	13.9	11.9	11.5	6.7	6.2	5.3	4.1	5.7	100.0
1987	6.4	6.8	8.8	9.7	15.1	11.4	8.7	7.7	6.5	5.9	5.7	7.3	100.0
1988	9.4	7.8	11.5	11.4	15.6	12.3	7.7	6.1	6.3	4.3	4.0	3.6	100.0
1989	6.9	7.6	10.4	13.9	17.0	12.7	7.1	6.8	4.7	4.8	4.1	4.0	100.0
1990	9.3	7.5	8.4	8.6	13.5	11.5	8.8	8.7	6.5	5.7	5.9	5.7	100.0
1991	4.8	4.3	3.3	8.4	13.2	10.7	10.4	9.5	10.0	10.2	7.0	8.2	100.0
1992	6.1	6.8	6.7	14.5	14.0	10.6	7.3	7.3	6.4	6.7	8.2	5.3	100.0
1993	7.8	5.9	10.5	8.1	14.7	10.6	7.1	7.7	6.4	7.7	7.1	6.4	100.0
1994	8.5	7.7	7.9	11.0	9.8	10.9	8.1	7.7	7.8	8.7	6.0	5.9	100.0
1995	7.2	5.9	9.7	10.7	11.6	13.3	9.1	7.8	7.1	7.6	5.2	5.0	100.0
1996	6.1	6.3	7.0	8.4	12.9	11.8	9.9	10.0	6.8	7.7	6.0	7.1	100.0
1997	6.2	7.5	7.6	9.4	10.4	14.3	9.7	8.0	9.3	6.2	4.8	6.6	100.0
1998	4.9	7.3	11.1	11.9	12.5	13.4	8.8	8.2	6.8	5.9	4.0	5.2	100.0
1999	6.0	6.7	8.8	8.4	9.9	15.8	9.9	9.7	5.5	5.9	6.0	7.4	100.0
2000	5.5	7.9	9.0	7.6	7.4	13.9	10.5	9.8	7.2	8.5	6.5	6.2	100.0
2001	8.9	5.5	7.9	10.5	7.4	13.6	10.7	8.6	8.0	5.7	5.6	7.6	100.0
2002	8.2	8.4	9.7	13.0	5.1	10.1	13.4	9.7	8.0	5.1	4.7	4.4	100.0

Table 7. Percentage of annual USA commercial witch flounder landings by port, 1982-2002.

Port/State	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
Rockland	22.9	24.4	19.8	18.4	19.6	36.2	35.7	34.8	31.3	32.1	29.3	1.0	0.5	0.7	0.6	0.9	1.2	1.2	0.7	0.1	0.2
Portland	16.3	21.5	12.6	13.6	10.1	11.2	10.5	11.1	1.6	2.4	2.6	35.7	39.3	36.5	40.3	40.8	32.9	29.5	28.2	27.4	23.6
Other Maine	14.0	13.0	11.3	8.8	10.1	11.9	10.6	6.1	8.2	9.0	10.7	10.9	11.0	8.5	8.7	8.7	11.2	11.7	9.5	9.0	9.4
NH	5.2	2.7	4.4	2.3	2.0	3.2	4.2	3.5	2.2	1.6	1.2	1.0	1.0	0.8	0.7	0.9	0.9	0.9	1.9	1.7	3.1
Boston	1.8	2.2	3.0	6.8	10.5	8.3	11.1	10.8	9.3	8.7	8.5	7.1	10.2	9.3	10.7	8.8	9.9	9.5	10.2	9.5	7.2
Gloucester	20.0	17.5	25.0	26.2	26.3	13.7	12.0	13.1	23.5	17.9	14.5	16.5	17.1	15.6	11.4	10.1	12.4	15.8	17.1	16.7	23.1
Chatham	1.3	1.0	1.5	1.4	0.8	1.9	2.1	2.0	1.0	0.8	0.7	1.0	0.6	0.8	0.6	0.7	0.8	1.1	3.0	0.9	1.5
New Bedford	3.3	4.8	8.5	10.4	11.5	6.4	5.5	7.7	8.0	9.1	15.1	13.0	9.6	20.6	19.7	20.2	21.4	18.5	18.7	24.9	23.2
Plymouth	0.7	1.3	1.1	1.1	0.8	0.9	0.8	0.7	0.4	0.8	0.8	0.7	0.2	0.1	0.1	0.0	0.3	0.4	1.7	0.6	1.2
Provincetown	5.8	5.0	4.4	3.3	2.9	1.7	1.4	3.1	4.2	6.0	3.2	4.4	2.6	2.3	1.5	1.5	2.4	1.7	1.8	2.5	1.5
Other Mass.	2.3	2.2	2.6	1.3	1.0	1.2	1.0	0.9	1.4	0.7	0.4	0.7	0.7	1.6	1.5	1.5	1.3	2.5	1.5	2.9	3.6
Newport	3.0	2.2	2.8	3.5	2.0	1.5	2.7	3.5	4.1	4.5	4.7	3.6	2.1	0.7	1.6	2.6	1.5	3.3	2.0	1.3	1.1
Point Judith	1.7	1.4	2.6	2.6	1.6	1.0	1.4	1.7	3.7	4.0	5.8	2.5	1.5	0.5	0.9	2.0	3.0	2.6	1.7	1.3	0.8
Other RI	0.5	0.2	0.0	0.0	0.2	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.5	0.3	0.3	0.1	0.1	0.1	0.1
CT	-	-	-	-	-	-	-	-	-	-	-	-	0.2	0.4	0.3	0.2	-	0.2	0.8	0.3	-
New York	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.2	0.8	1.1	0.7	2.1	0.7	0.5	0.4	0.2	0.6	0.3	0.3	0.1
New Jersey	0.8	0.4	0.4	0.4	0.5	0.4	0.5	0.6	0.8	1.3	1.3	1.0	0.8	0.3	0.3	0.3	0.2	0.3	0.8	0.4	0.3
Maryland	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Virginia	0.3	0.1	0.1	0.0	0.2	0.2	0.1	0.2	0.1	0.2	0.2	0.1	0.4	0.4	0.2	0.0	0.0	0.1	0.0	0.0	0.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 8. Percentage of annual USA commercial witch flounder landings by market category, 1973 - 2002.

Year	Peewee	Small	Medium	Large	Jumbo	Uncl.	Total
1973	0.0	13.5	0.0	45.9	0.0	40.7	100.0
1974	0.0	26.2	0.0	73.8	0.0	0.0	100.0
1975	0.0	26.3	0.0	73.7	0.0	0.0	100.0
1976	0.0	21.5	0.0	78.4	0.0	0.1	100.0
1977	0.0	22.9	0.0	77.1	0.0	0.0	100.0
1978	0.0	30.2	0.0	69.8	0.0	0.0	100.0
1979	0.0	30.8	0.0	69.2	0.0	0.0	100.0
1980	0.0	23.4	0.0	76.0	0.0	0.6	100.0
1981	0.0	30.1	0.0	68.3	0.0	1.6	100.0
1982	0.3	26.3	5.4	64.0	0.0	4.0	100.0
1983	1.4	25.0	14.7	58.4	0.0	0.4	100.0
1984	3.4	25.2	19.1	51.7	0.0	0.6	100.0
1985	7.7	27.8	23.2	40.5	0.1	0.7	100.0
1986	5.1	33.7	25.3	34.6	0.0	1.2	100.0
1987	3.6	37.2	26.0	31.0	0.5	1.7	100.0
1988	2.8	34.3	29.0	30.7	0.6	2.7	100.0
1989	3.3	29.8	31.2	31.5	1.1	3.0	100.0
1990	5.5	26.2	30.6	32.6	0.7	4.4	100.0
1991	6.6	33.1	25.5	31.0	1.3	2.4	100.0
1992	13.2	39.0	20.3	25.0	0.1	2.4	100.0
1993	17.7	39.3	18.5	21.6	0.0	2.9	100.0
1994	19.3	43.7	16.0	16.8	0.0	4.1	100.0
1995	26.0	46.6	11.9	13.0	0.0	2.5	100.0
1996	27.4	53.1	9.9	8.0	0.0	1.7	100.0
1997	18.2	63.7	10.5	6.1	0.0	1.4	100.0
1998	13.2	72.1	9.4	4.6	0.0	0.7	100.0
1999	10.1	74.3	10.1	4.6	0.0	0.9	100.0
2000	8.1	76.6	9.7	3.6	0.0	2.0	100.0
2001	9.0	77.9	9.1	2.9	0.0	1.1	100.0
2002	8.2	78.5	9.7	2.6	0.0	0.9	100.0

Table 9. Summary of USA commercial witch flounder landings (mt), number of length samples (n), number of fish measured (len) and number of age samples (age) by market category and quarter for all gear types, 1981 - 2002. The sampling ratio represents the amount of landings per length sample.

Year	Quarter 1			Quarter 2			Quarter 3			Quarter 4			All	Sampling Ratio
	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large		
1981 mt	260	7	517	269	32	694	242	13	607	230	0	453	3324	
n	1	1	.	1	.	1	.	1	5	
len	101	103	.	89	.	105	.	100	498	
age	26	.	25	.	25	.	25	101	
1982 mt	348	1	726	342	73	886	287	170	739	278	201	669	4720	
n	5	2	6	1	2	2	2	2	6	3	4	2	37	128
len	527	194	626	126	209	216	189	210	514	307	393	189	3700	
age	128	55	150	30	55	50	50	50	150	81	105	50	954	
1983 mt	475	250	910	471	286	1037	298	154	758	257	169	613	5678	
n	5	2	3	5	1	5	8	3	8	6	3	.	49	116
len	680	232	265	685	96	520	1008	123	981	677	344	.	5611	
age	135	30	55	131	16	125	152	0	159	180	75	.	1058	
1984 mt	462	322	1036	513	393	1000	403	248	653	429	286	586	6331	
n	5	9	4	7	1	7	8	1	2	4	2	1	51	124
len	804	1112	400	970	117	775	1045	106	191	615	243	91	6469	
age	154	250	76	186	25	180	210	28	53	105	44	25	1336	
1985 mt	465	377	613	697	453	850	526	291	553	433	310	408	5976	
n	12	1	2	5	4	7	7	7	6	8	2	4	65	92
len	1530	105	229	657	426	698	795	800	684	824	264	349	7361	
age	319	29	50	106	77	153	97	138	113	161	25	29	1297	
1986 mt	384	309	356	654	421	595	375	238	354	312	212	238	4448	
n	6	3	5	5	4	5	4	3	4	5	3	2	49	90
len	662	307	515	558	410	413	302	364	406	416	337	233	4923	
age	123	60	89	106	97	129	63	75	100	87	75	52	1056	

Table 9 continued.

Year	Quarter 1			Quarter 2			Quarter 3			Quarter 4			All	Sampling Ratio
	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large		
1987 mt	349	211	228	432	317	387	296	203	247	298	203	202	3373	69
n	1	1	2	4	2	3	5	5	4	2	3	2	34	
len	85	145	200	323	228	316	354	583	400	204	261	178	3277	
age	25	25	50	77	47	76	78	113	95	48	64	51	749	
1988 mt	424	304	271	436	393	389	184	176	208	140	140	131	3196	65
n	5	4	5	5	5	3	5	4	3	3	4	3	49	
len	335	407	465	344	544	429	396	359	295	229	402	356	4561	
age	70	89	106	71	110	77	70	100	75	61	95	69	993	
1989 mt	230	174	148	255	264	251	98	145	156	85	107	103	2016	112
n	1	2	2	2	2	1	2	2	1	1	2	.	18	
len	94	201	222	230	236	27	150	206	100	125	202	.	1793	
age	25	50	49	50	46	25	40	51	25	25	47	.	433	
1990 mt	113	125	107	147	168	147	100	119	129	84	79	85	1403	40
n	1	2	3	6	3	1	6	2	2	7	2	.	35	
len	134	199	199	335	296	100	349	247	145	381	201	.	2586	
age	15	40	45	81	70	25	69	41	50	103	48	.	587	
1991 mt	71	56	58	219	151	167	192	142	184	168	108	121	1637	40
n	5	2	3	7	2	1	4	2	3	5	4	3	41	
len	262	224	401	537	239	125	212	165	249	300	410	274	3398	
age	53	50	80	93	45	25	49	49	52	66	97	58	717	
1992 mt	180	86	82	466	163	174	205	115	138	212	97	116	2034	68
n	4	2	2	7	1	2	7	1	1	2	.	1	30	
len	259	241	185	501	125	235	477	121	117	129	.	46	2436	
age	42	46	52	78	25	25	86	25	25	27	.	23	454	
1993 mt	350	112	110	442	192	161	263	122	150	331	96	106	2435	76
n	7	1	.	7	1	1	9	1	5	.	.	.	32	
len	830	100	.	741	107	100	728	85	499	.	.	.	3190	
age	55	25	.	56	27	26	74	.	73	.	.	.	336	

Table 9 continued.

Year	Quarter 1			Quarter 2			Quarter 3			Quarter 4			All	Sampling Ratio
	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large		
1994 mt	403	143	98	505	183	154	390	122	117	383	91	80	2670	
n	.	.	.	3	5	6	5	5	1	5	3	4	37	72
len	.	.	.	560	532	749	356	648	105	342	368	407	4067	
age	.	.	.	59	104	134	44	113	26	56	60	82	678	
1995 mt	336	91	77	586	117	100	399	61	70	304	48	40	2212	
n	3	3	3	6	3	5	.	.	.	2	.	1	26	85
len	208	348	347	459	367	517	.	.	.	217	.	94	2557	
age	53	84	89	81	75	135	.	.	.	27	.	25	569	
1996 mt	313	57	36	545	86	60	458	56	44	363	42	28	2088	
n	5	2	3	5	2	1	5	4	4	5	3	3	42	50
len	504	218	292	331	240	127	494	464	468	343	277	348	4106	
age	59	45	78	53	50	26	59	86	101	60	70	69	756	
1997 mt	313	40	25	478	86	41	398	55	27	265	31	16	1775	
n	6	3	3	9	4	3	9	3	1	9	1	1	52	34
len	557	350	351	812	418	309	783	308	107	505	128	50	4678	
age	77	68	70	108	73	77	98	81	20	73	18	23	786	
1998 mt	372	39	19	587	79	31	380	40	20	239	26	14	1849	80
n	5	2	1	4	1	1	5	3	1	.	.	.	23	
len	339	206	128	238	88	135	484	186	100	.	.	.	1904	
age	45	50	19	30	.	29	47	22	242	
1999 mt	386	48	19	616	79	31	436	67	30	353	38	18	2121	51
n	3	.	.	4	.	.	17	2	3	11	1	.	41	
len	282	.	.	308	.	.	1110	201	306	775	109	.	3091	
age	15	.	.	62	.	.	143	.	32	91	16	.	359	
2000 mt	477	53	17	583	93	27	555	89	28	451	50	16	2439	21
n	31	2	.	47	.	.	17	1	.	5	5	2	110	
len	2253	91	.	2445	.	.	994	105	.	308	558	217	6971	
age	390	10	.	460	.	.	224	20	.	67	92	51	1314	

Table 9 continued.

Year	Quarter 1			Quarter 2			Quarter 3			Quarter 4			Sampling	
	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large	Small	Med.	Large	All	Ratio
2001 mt	583	71	17	824	99	30	699	98	28	507	50	13	3019	70
n	8	4	2	3	3	2	8	2	3	5	3	.	43	
len	744	422	134	237	352	159	594	209	213	313	232	.	3609	
age	125	63	42	47	48	64	126	34	46	61	48	.	704	
2002 mt	740	79	18	774	103	26	849	114	29	400	45	9	3186	91
n	4	1	2	3	5	3	5	2	3	3	2	2	35	
len	312	121	107	212	518	209	389	150	194	262	226	115	2815	
age	73	14	44	65	68	63	86	32	62	49	30	49	635	

Table 10. The data pooling to apply age and length frequency samples to landings by market category and quarter to estimate numbers at age of witch flounder from 1982-2002.

Year	Mkt. Cat.	Quarter 1	Quarter 2	Quarter 3	Quarter 4
1982	Small	<=Pooled =>		X	X
	Med.	X	X	X	X
	Large	X	X	X	X
1983	Small	X	X	X	X
	Med.	<=Pooled =>		X	X
	Large	X	X	<=Pooled =>	
1984	Small	X	X	X	X
	Med.	<=Pooled =>		<=Pooled =>	
	Large	X	X	<=Pooled =>	
1985	Small	X	X	X	X
	Med.	X	X	X	X
	Large	X	X	X	X
1986	Small	X	X	X	X
	Med.	X	X	X	X
	Large	X	X	X	X
1987	Small	<=Pooled =>		X	X
	Med.	<=Pooled =>		X	X
	Large	X	X	X	X
1988	Small	X	X	X	X
	Med.	X	X	X	X
	Large	X	X	X	X
1989	Small	<=Pooled =>		<=Pooled =>	
	Med.	X	X	X	X
	Large	<====Pooled====>			
1990	Small	<=Pooled =>		X	X
	Med.	X	X	X	X
	Large	<=Pooled =>		<=Pooled =>	
1991	Small	X	X	X	X
	Med.	X	X	X	X
	Large	<=Pooled =>		X	X
1992	Small	X	X	X	X
	Med.	<====Pooled====>			
	Large	X	X	<=Pooled =>	

Table 10 continued.

Year	Mkt. Cat.	Quarter 1	Quarter 2	Quarter 3	Quarter 4
1993	Small	X	X	<=Pooled =>	
	Med.	<===Pooled ===>			
	Large	<===Pooled ===>			
1994	Small	<=Pooled =>		X	X
	Med.	<=Pooled =>		X	X
	Large	<=Pooled =>		<=Pooled =>	
1995	Small	X	<=== Pooled ===>		
	Med.	X	<===Pooled ===>		
	Large	X	<===Pooled ===>		
1996	Small	X	X	X	X
	Med.	<=Pooled =>		X	X
	Large	<=Pooled =>		X	X
1997	Small	X	X	X	X
	Med.	X	X	<=Pooled =>	
	Large	X	X	<=Pooled =>	
1998	Small	X	X	<=Pooled =>	
	Med.	<=Pooled =>		<=Pooled =>	
	Large	<===Pooled ===>			
1999	Small	<=Pooled =>		X	X
	Med.	<===Pooled ===>			
	Large	<===Pooled ===>			
2000	Small	X	X	X	X
	Med.	<===Pooled ===>			
	Large	<===Pooled ===>			
2001	Small	X	X	X	X
	Med.	<=Pooled =>		<=Pooled =>	
	Large	<===Pooled ===>			
2002	Small	X	X	X	X
	Med.	<=Pooled =>		<=Pooled =>	
	Large	<=Pooled =>		<=Pooled =>	

Table 11. USA commercial landings at age in numbers, weight (thousands of fish; mt) and mean weight (kg) and mean length (cm) at age of witch flounder, 1982 - 2002.

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commercial Landings in Numbers (1000's) at Age																	
1982	0.0	0.0	0.0	117.9	826.6	1119.9	1454.3	665.2	656.0	399.5	239.4	201.0	356.3	183.7	837.4	7057.2	1578.4
1983	0.0	0.0	0.0	219.8	768.6	1033.7	1567.3	1590.2	977.8	737.7	510.4	366.0	287.3	289.1	733.1	9081.0	1675.5
1984	0.0	0.0	0.0	90.6	1012.4	1808.7	1734.3	1486.5	1497.5	696.7	375.1	279.5	356.4	261.3	821.6	10420.0	1718.8
1985	0.0	0.0	0.0	0.0	985.1	2026.8	1933.8	1524.9	1247.9	606.0	400.4	261.2	221.5	170.7	705.8	10084.0	1359.2
1986	0.0	0.0	0.0	6.3	298.5	1441.6	2772.6	1566.9	834.9	412.7	222.8	188.2	157.0	137.0	276.0	8314.5	758.2
1987	0.0	0.0	0.0	0.0	81.5	321.6	1276.0	1574.7	870.9	480.6	252.4	132.4	90.8	62.1	204.1	5347.1	489.4
1988	0.0	0.0	0.0	0.0	50.8	176.0	654.7	1382.7	1154.1	401.5	266.7	124.1	94.0	71.9	307.5	4684.0	597.5
1989	0.0	0.0	0.0	0.0	7.3	49.7	314.3	759.4	882.1	349.7	123.4	73.2	61.1	56.7	157.1	2833.8	349.0
1990	0.0	0.0	0.0	0.0	181.6	574.3	255.6	273.9	471.1	333.9	81.4	43.1	38.5	19.1	76.9	2349.2	179.1
1991	0.0	0.0	0.0	0.0	179.5	732.9	519.4	235.8	244.6	292.1	313.6	51.8	44.0	22.5	139.5	2775.6	260.8
1992	0.0	0.0	0.0	0.0	509.3	839.4	935.5	717.0	201.6	177.9	120.0	217.6	46.3	26.5	86.5	3877.7	380.2
1993	0.0	0.0	0.0	0.0	422.2	1022.8	917.7	597.2	585.6	218.8	278.5	113.9	32.6	103.6	140.4	4433.2	391.1
1994	0.0	0.0	0.0	0.0	201.3	1429.4	1286.2	826.9	196.7	539.2	113.5	71.4	40.2	132.3	80.4	4917.4	324.9
1995	0.0	0.0	0.0	0.0	23.7	763.0	1597.4	848.7	267.5	97.2	269.5	55.0	43.9	8.1	49.9	4023.8	157.1
1996	0.0	0.0	0.0	0.0	45.8	467.7	1263.8	1430.4	263.2	215.5	57.1	78.8	3.6	13.0	18.2	3857.2	113.7
1997	0.0	0.0	0.0	0.0	212.2	527.9	1049.4	1014.0	591.3	83.1	49.8	17.9	36.6	2.2	13.4	3597.8	70.2
1998	0.0	0.0	0.0	0.0	18.1	488.0	1213.5	1583.0	370.5	141.4	15.5	37.2	5.6	19.9	7.7	3900.2	70.3
1999	0.0	0.0	0.0	0.0	185.2	585.7	1391.7	1178.3	763.2	251.3	31.6	40.8	0.0	0.0	13.5	4441.3	54.4
2000	0.0	0.0	0.0	0.0	75.4	266.2	1062.1	1611.1	1027.6	623.7	94.8	174.3	6.2	5.0	27.4	4973.9	212.8
2001	0.0	0.0	0.0	0.0	18.8	382.2	940.5	1669.0	1459.4	634.3	425.4	95.8	163.5	8.6	38.8	5836.3	306.7
2002	0.0	0.0	0.0	0.0	173.2	644.9	1242.7	2098.3	1274.4	632.1	96.4	102.7	11.0	65.6	25.3	6366.6	202.6

Table 11 continued. USA commercial landings.

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commercial Landings Mean Weight (kg) at Age																	
1982	0.000	0.000	0.000	0.216	0.275	0.345	0.424	0.550	0.727	0.886	0.983	1.146	1.255	1.310	1.553	0.695	1.406
1983	0.000	0.000	0.000	0.195	0.257	0.322	0.410	0.518	0.613	0.795	0.977	1.116	1.208	1.321	1.551	0.661	1.357
1984	0.000	0.000	0.000	0.212	0.268	0.346	0.422	0.539	0.664	0.817	0.922	1.004	1.212	1.332	1.511	0.639	1.339
1985	0.000	0.000	0.000	0.000	0.253	0.311	0.429	0.565	0.691	0.842	0.964	1.057	1.193	1.311	1.470	0.608	1.326
1986	0.000	0.000	0.000	0.084	0.227	0.306	0.408	0.533	0.676	0.853	0.975	1.132	1.199	1.317	1.521	0.555	1.321
1987	0.000	0.000	0.000	0.000	0.272	0.342	0.434	0.561	0.686	0.828	0.980	1.067	1.222	1.386	1.467	0.645	1.303
1988	0.000	0.000	0.000	0.000	0.310	0.367	0.435	0.538	0.668	0.819	0.980	1.074	1.190	1.290	1.477	0.696	1.326
1989	0.000	0.000	0.000	0.000	0.260	0.344	0.425	0.574	0.682	0.818	0.968	1.128	1.258	1.315	1.519	0.730	1.358
1990	0.000	0.000	0.000	0.000	0.308	0.323	0.438	0.586	0.688	0.849	1.049	1.213	1.262	1.521	1.669	0.624	1.454
1991	0.000	0.000	0.000	0.000	0.286	0.371	0.443	0.578	0.702	0.836	0.974	1.099	1.369	1.537	1.536	0.640	1.420
1992	0.000	0.000	0.000	0.000	0.328	0.383	0.459	0.614	0.739	0.822	0.882	1.039	1.337	1.459	1.640	0.575	1.243
1993	0.000	0.000	0.000	0.000	0.292	0.364	0.432	0.535	0.666	0.882	1.023	1.118	1.199	1.368	1.519	0.587	1.335
1994	0.000	0.000	0.000	0.000	0.308	0.357	0.430	0.534	0.691	0.832	0.909	1.083	1.172	1.204	1.576	0.542	1.266
1995	0.000	0.000	0.000	0.000	0.284	0.367	0.448	0.561	0.690	0.911	0.974	1.101	1.203	1.411	1.406	0.549	1.243
1996	0.000	0.000	0.000	0.000	0.260	0.355	0.435	0.554	0.708	0.856	0.974	1.114	1.401	1.440	1.558	0.541	1.232
1997	0.000	0.000	0.000	0.000	0.318	0.357	0.407	0.495	0.628	0.871	1.037	1.168	1.196	1.687	1.659	0.492	1.293
1998	0.000	0.000	0.000	0.000	0.235	0.331	0.382	0.492	0.585	0.871	0.978	1.115	1.132	1.261	1.557	0.474	1.206
1999	0.000	0.000	0.000	0.000	0.325	0.355	0.406	0.516	0.584	0.628	0.917	0.683	-	-	1.442	0.477	0.872
2000	0.000	0.000	0.000	0.000	0.319	0.327	0.376	0.450	0.533	0.633	0.677	0.834	1.167	1.298	1.379	0.490	0.925
2001	0.000	0.000	0.000	0.000	0.291	0.325	0.384	0.469	0.550	0.646	0.647	0.718	0.816	1.016	1.206	0.517	0.840
2002	0.000	0.000	0.000	0.000	0.354	0.344	0.416	0.477	0.554	0.651	0.824	0.844	0.716	0.993	1.120	0.501	0.919
mean																	
1982-02	0.000	0.000	0.000	0.034	0.287	0.345	0.421	0.535	0.654	0.807	0.934						1.237
1999-02	0.000	0.000	0.000	0.000	0.322	0.338	0.396	0.478	0.555	0.640	0.766						0.889

Table 11 continued. USA commercial landings.

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commercial Landings Mean Length (cm) at Age																	
1982	0.0	0.0	0.0	32.3	35.0	37.5	39.8	42.9	46.5	49.3	50.9	53.2	54.6	55.2	58.0	44.3	56.3
1983	0.0	0.0	0.0	31.7	34.3	36.8	39.4	42.2	44.2	47.7	50.7	52.8	54.0	56.6	55.8	35.9	55.0
1984	0.0	0.0	0.0	32.6	34.9	37.6	39.8	42.7	45.3	48.2	49.9	51.2	54.1	55.6	57.6	43.6	55.5
1985	0.0	0.0	0.0	0.0	34.2	36.3	40.0	43.3	45.9	48.6	50.6	51.9	53.8	55.3	57.1	42.9	55.3
1986	0.0	0.0	0.0	25.0	33.2	36.2	39.4	42.5	45.6	48.8	50.7	53.0	53.9	55.4	57.7	42.0	55.3
1987	0.0	0.0	0.0	0.0	35.0	37.4	40.1	43.2	45.8	48.4	50.8	52.1	54.2	56.2	57.1	44.3	55.1
1988	0.0	0.0	0.0	0.0	36.4	38.2	40.1	42.7	45.4	48.2	50.8	52.1	53.7	55.0	57.1	45.3	55.3
1989	0.0	0.0	0.0	0.0	34.6	37.5	39.9	43.5	45.6	48.1	50.6	52.9	54.6	55.3	57.6	46.0	55.7
1990	0.0	0.0	0.0	0.0	36.2	36.8	40.2	43.7	45.8	48.7	51.8	54.1	54.6	57.8	59.2	43.5	56.8
1991	0.0	0.0	0.0	0.0	35.4	38.3	40.3	43.3	46.1	48.5	50.6	52.5	56.0	57.9	57.9	43.8	56.5
1992	0.0	0.0	0.0	0.0	37.0	38.7	40.7	44.3	46.8	48.3	49.2	51.7	55.5	57.0	58.9	42.7	54.2
1993	0.0	0.0	0.0	0.0	35.8	38.1	40.0	42.6	45.3	49.3	51.5	52.8	53.9	55.9	57.7	42.8	55.5
1994	0.0	0.0	0.0	0.0	36.0	37.6	39.7	42.3	45.6	48.0	49.1	51.8	53.1	53.4	57.8	41.7	54.1
1995	0.0	0.0	0.0	0.0	35.3	37.9	40.2	42.8	45.4	49.3	50.1	52.0	53.4	56.0	55.8	42.0	53.8
1996	0.0	0.0	0.0	0.0	34.4	37.5	39.8	42.7	45.8	48.4	50.1	52.2	55.8	56.2	57.6	42.0	53.6
1997	0.0	0.0	0.0	0.0	36.4	37.6	39.1	41.3	44.2	48.5	51.1	52.9	53.3	59.0	58.7	40.9	54.4
1998	0.0	0.0	0.0	0.0	33.4	36.8	38.4	41.2	43.3	48.7	50.5	52.3	52.7	54.1	57.6	40.5	53.4
1999	0.0	0.0	0.0	0.0	36.6	37.5	39.0	41.8	43.3	44.3	49.4	45.2	-	-	56.4	40.7	48.0
2000	0.0	0.0	0.0	0.0	36.4	36.7	38.2	40.2	42.2	44.2	45.2	47.7	53.0	54.6	55.3	40.9	49.0
2001	0.0	0.0	0.0	0.0	35.5	36.6	38.4	40.7	42.5	44.6	44.5	45.9	47.7	50.8	53.3	41.6	47.9
2002	0.0	0.0	0.0	0.0	37.4	37.2	39.3	40.9	42.7	44.6	47.9	48.2	45.8	50.5	52.2	41.6	49.3

Table 12. Discard rates (kg/day fished) by fishing zone¹ obtained from a ratio estimator (kg of witch flounder discarded to days fished) using Fisheries Observer Program data collected from the northern shrimp fishery, number of days fished by the shrimp fishery, mean discard rates (kg/df) and estimated discard weight (kg) of witch flounder in the northern shrimp fishery, during the 1989 - 1997 shrimp seasons.

Shrimp Season	Fishing Zone	Sea Sample Data		Commercial days fished	Mean discard rate	Estimated discard weight (kg)	Estimated discard weight (mt)
		Trips	Discard Rate (kg/df)				
1989	1	5	0.0000	398.2	6.0626	17,215	17.2
	2	15	2.2032	1680.2			
	3	16	17.7543	<u>761.1</u>			
				2839.5			
1990	1	4	0.0000	416.9	8.7512	28,044	28.0
	2	23	7.0751	1610.9			
	3	20	14.1459	<u>1176.8</u>			
				3204.6			
1991	1	13	0.9770	528.0	12.6856	32,827	32.8
	2	25	4.4822	1154.8			
	3	24	29.9863	<u>904.9</u>			
				2587.7			
1992	1	30	2.7834	187.3	8.2343	19,048	19.0
	2	60	8.9270	1764.1			
	3	20	7.6787	<u>361.9</u>			
				2313.3			
1993	1	38	1.3559	526.9	4.4485	8,462	8.5
	2	53	3.7619	1094.2			
	3	13	12.9178	<u>281.1</u>			
				1902.2			
1994*	1	37	3.3021	498.7	5.6004	11,102	11.1
	2	56	5.8385	1334			
	3	5	11.1394	<u>149.6</u>			
				1982.3			
1995*	1	24	2.0007	2036.2	11.0492	37,299	37.3
	2	46	27.5162	1109			
	3	18	11.7543	<u>230.5</u>			
				3375.7			
1996*	1	8	0.3532	2079.4	4.3130	13,987	14.0
	2	31	7.6343	958.2			
	3	11	28.919	<u>205.3</u>			
				3242.9			
1997*	1	6	0.4065	1996.1	3.2915	12,051	12.1
	2	19	2.9403	1191.8			
	3	3	16.3461	<u>473.2</u>			
				3661.1			

¹ Fishing zones: 1 = 0-3 miles; 2 = 3 - 12 miles, and 3 = greater than 12 miles from shore.

* Commercial days fished have been estimated from Vessel Trip Report data.

Table 13. Witch flounder discard rates (kg/df), days fished (df), discarded metric tons (mt), numbers of discarded fish (in thousands) in the northern shrimp fishery during 1982-2002.

Year	Shrimp Season Disc. Rate	Calendar Year			mt	Numbers (‘000)
		Days Fished				
		Jan-Apr	Dec.	Total		
1982	5.7025	970.1	35.6	1005.7	5.90	62.14
1983	10.4523	1121.3	141.7	1263	12.56	131.67
1984	5.9234	1612.3	237.6	1849.9	10.93	110.94
1985	5.8129	1843.8	272.8	2116.6	12.12	91.32
1986	5.1502	2122.3	428.9	2551.2	13.14	98.80
1987	5.1502	3279.3	380.4	3659.7	21.79	235.99
1988	12.8824	2434.8	426.9	2861.7	33.95	723.95
1989	6.0626	2412.6	491.9	2904.5	18.93	219.81
1990	8.7512	2712.7	377.6	3090.3	28.53	468.56
1991	12.6856	2210.1	172.3	2382.4	29.46	443.85
1992	8.2343	2141.0	113.2	2254.2	18.13	384.45
1993	4.4485	1789.0	161.7	1950.7	8.86	356.77
1994	5.6004	1820.6	530.8	2351.4	16.06	1891.71
1995	11.0492	2844.9	547.6	3392.5	33.80	1176.37
1996	4.3130	2695.3	645.2	3340.5	13.75	250.46
1997	3.2915	3016.0	361.4	3377.4	13.27	304.52
1998	9.2437	1842.6	96.9	1939.5	18.04	873.52
1999	10.3851	1120.5	0	1120.5	11.64	563.50
2000	9.8775	792.9	0	792.9	7.83	379.26
2001	6.6415	672.8	0	672.8	4.47	216.38
2002	3.5325	238	0	238	0.84	40.68

Note: 1982-1988 discard rates were derived from a linear regression using 1989-1993 discard rates and NEFSC autumn age 3 abundance indices.

1998-2002 discard rates were derived from a linear regression using 1989-1993 discard rates and NEFSC autumn age 3 abundance indices.

Table 14. Witch flounder discards at age in numbers, weight (thousands of fish; mt) and mean weight (kg) and mean length (cm) at age in the shrimp fishery, 1982 - 2002.

Year	Age															TOTAL
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+	
Shrimp Fishery Discards in Numbers (1000's) at Age																
1982	0.00	0.00	1.59	25.24	21.12	11.27	2.91	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	62.14
1983	0.00	0.00	3.62	53.11	44.65	23.81	6.48	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	131.67
1984	0.00	0.33	0.77	46.84	38.55	19.41	5.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	110.94
1985	0.00	0.34	3.37	11.72	47.06	26.39	2.44	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	91.32
1986	0.00	0.53	3.86	15.07	49.83	27.04	2.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	98.80
1987	2.08	18.92	79.51	15.62	74.59	41.46	3.81	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	235.99
1988	0.42	14.62	130.29	495.50	42.57	37.70	2.85	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	723.95
1989	0.74	10.47	47.52	69.23	76.39	15.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	219.81
1990	1.19	5.18	92.78	239.97	97.13	32.31	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	468.56
1991	2.96	17.79	15.98	287.35	102.86	11.59	5.32	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	443.85
1992	2.71	43.41	136.92	118.76	82.06	0.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	384.45
1993	112.06	78.84	107.58	38.69	14.13	5.02	0.46	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	356.77
1994	8.06	1368.46	495.50	19.62	0.00	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1891.71
1995	2.68	49.95	630.10	480.83	12.25	0.20	0.36	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1176.37
1996	5.21	32.68	50.83	99.45	59.21	2.09	0.99	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	250.46
1997	8.68	74.91	102.92	86.49	23.71	7.30	0.52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	304.52
1998	49.78	391.44	264.72	132.04	30.13	4.78	0.63	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	873.52
1999	32.11	252.51	170.76	85.18	19.44	3.09	0.41	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	563.50
2000	21.61	169.95	114.93	57.33	13.08	2.08	0.28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	379.26
2001	12.33	96.96	65.57	32.71	7.46	1.19	0.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	216.38
2002	2.32	18.23	12.33	6.15	1.4	0.22	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	40.68

Note: 1998-2002 estimated using 1993-1997 Fisheries Observer Program data.

Table 14 continued. Discards in the shrimp fishery.

Year	Age														TOTAL	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13		14+
Shrimp fishery Mean Weight (kg) at Age																
1982			0.040	0.043	0.101	0.165	0.256									0.095
1983			0.040	0.044	0.101	0.166	0.256									0.095
1984		0.017	0.044	0.050	0.105	0.165	0.256									0.099
1985		0.017	0.023	0.081	0.123	0.179	0.231									0.133
1986		0.017	0.026	0.089	0.125	0.180	0.231									0.133
1987	0.006	0.015	0.033	0.071	0.126	0.180	0.231									0.092
1988	0.004	0.006	0.017	0.036	0.121	0.206	0.282									0.047
1989	0.010	0.012	0.033	0.058	0.122	0.249										0.086
1990	0.004	0.010	0.029	0.043	0.107	0.155										0.061
1991	0.004	0.014	0.030	0.045	0.117	0.221	0.218									0.066
1992	0.003	0.007	0.021	0.043	0.119	0.225										0.047
1993	0.003	0.009	0.022	0.057	0.136	0.237	0.317									0.025
1994	0.005	0.004	0.019	0.032		0.282										0.009
1995	0.005	0.007	0.023	0.037	0.083	0.289	0.282									0.029
1996	0.004	0.019	0.031	0.056	0.090	0.184	0.289									0.055
1997	0.004	0.023	0.033	0.048	0.115	0.144	0.256									0.044
1998	0.003	0.006	0.023	0.042	0.100	0.184	0.286									0.021
1999	0.003	0.006	0.023	0.042	0.100	0.184	0.286									0.021
2000	0.003	0.006	0.023	0.042	0.100	0.184	0.286									0.021
2001	0.003	0.006	0.023	0.042	0.100	0.184	0.286									0.021
2002	0.003	0.006	0.023	0.042	0.100	0.184	0.286									0.021
Mean																
1982-02	0.004	0.011	0.027	0.050	0.110	0.197	0.267									0.058

Note: 1998-2002 estimated using 1993-1997 Fisheries Observer Program data.

Table 14 continued. Discards in the shrimp fishery.

Year	Age														TOTAL	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13		14+
Shrimp fishery Mean Length (cm) at Age																
1982			20.3	20.6	26.5	30.7	34.9									25.1
1983			20.3	20.6	26.5	30.7	34.9									25.1
1984		15.7	20.7	21.2	26.7	30.7	34.9									25.4
1985		15.7	16.9	24.2	28.1	31.4	33.9									28.3
1986		15.7	17.3	24.9	28.2	31.4	33.9									28.3
1987	10.6	15.3	19.0	23.4	28.2	31.4	33.9									24.3
1988	10.2	10.9	15.6	19.4	27.9	32.8	36.0									19.8
1989	13.6	13.9	18.9	22.2	28.1	34.6										24.0
1990	10.5	13.6	17.9	20.4	27.0	30.2										21.9
1991	9.7	14.2	17.7	20.9	27.6	33.6	33.4									22.5
1992	9.3	10.8	16.6	20.5	27.9	33.7										19.5
1993	9.2	12.0	16.9	22.1	28.9	34.2	37.3									14.7
1994	10.7	9.8	15.9	18.5	-	36.0										11.5
1995	10.9	11.6	17.0	19.6	24.9	36.2	36.0									18.0
1996	10.0	15.3	18.4	22.1	25.6	31.7	36.2									21.2
1997	10.2	16.1	18.9	21.2	27.6	29.5	35.0									19.6
1998	9.4	10.4	16.8	20.3	26.4	31.5	36.1									14.9
1999	9.4	10.4	16.8	20.3	26.4	31.5	36.1									14.9
2000	9.4	10.4	16.8	20.3	26.4	31.5	36.1									14.9
2001	9.4	10.4	16.8	20.3	26.4	31.5	36.1									14.9
2002	9.4	10.4	16.8	20.3	26.4	31.5	36.1									14.9

Note: 1998-2002 estimated using 1993-1997 Fisheries Observer Program data.

Table 15. Three scenarios of large-mesh otter trawl discards were examined: 1) survey filter method utilizing the NEFSC bottom trawl survey; 2) survey filter method and the Fisheries Observer Program (FOP) data; and 3) survey filter method and the Vessel Trip Reports (VTR).

Year	Scenario 1	Scenario 2	Scenario 3
1982	survey filter	survey filter	survey filter
1983		survey filter	survey filter
1984		survey filter	survey filter
1985		survey filter	survey filter
1986		survey filter	survey filter
1987		survey filter	survey filter
1988		survey filter	survey filter
1989		survey filter	survey filter
1990		survey filter	survey filter
1991		survey filter	survey filter
1992		survey filter	survey filter
1993		survey filter	survey filter
1994		survey filter	
1995		FOP	VTR
1996		FOP	VTR
1997		FOP	VTR
1998		FOP	VTR
1999		FOP	VTR
2000		FOP	VTR
2001		FOP	VTR
2002		FOP	VTR

Table 16. Example of the spreadsheet calculations for estimating semi-annual discarded witch flounder in the large-mesh otter trawl fishery. This spreadsheet illustrates 1993, quarters 1 and 2 with the NEFSC spring survey. The **bold** numbers indicate columns and the mathematical operation performed.

1993 Landings from Q1+Q2 and 1993 spring survey

	1	2	3=1*2	4	5=3*4	6=3-5	7	8=6*factor
		140 mm					100's	units
Length	Survey	Prop.	Survey	Prop	Survey	Survey	Numbers	Numbers
(cm)	No/tow	Retained	Retained	Kept	Kept	Discarded	Landed	Discarded
1	0.000	0.00005	0.0000	0.00	0.0000	0.0000	0	0
3	0.000	0.00008	0.0000	0.00	0.0000	0.0000	0	0
5	0.034	0.00015	0.0000	0.00	0.0000	0.0000	0	34
7	0.064	0.00028	0.0000	0.00	0.0000	0.0000	0	120
9	0.051	0.00051	0.0000	0.00	0.0000	0.0000	0	174
11	0.000	0.00092	0.0000	0.00	0.0000	0.0000	0	0
13	0.000	0.00168	0.0000	0.00	0.0000	0.0000	0	0
15	0.011	0.00305	0.0000	0.00	0.0000	0.0000	0	224
17	0.067	0.00554	0.0004	0.00	0.0000	0.0004	0	2483
19	0.042	0.01005	0.0004	0.00	0.0000	0.0004	0	2823
21	0.028	0.01816	0.0005	0.00	0.0000	0.0005	0	3401
23	0.042	0.03261	0.0014	0.00	0.0000	0.0014	0	9161
25	0.061	0.05787	0.0035	0.00	0.0000	0.0035	0	23612
27	0.165	0.10065	0.0166	0.00	0.0000	0.0166	0	111083
29	0.079	0.16938	0.0134	0.00	0.0000	0.0134	0	89503
31	0.205	0.27091	0.0555	0.01	0.0006	0.0550	0	367759
33	0.165	0.40372	0.0666	0.10	0.0067	0.0600	231	401011
35	0.152	0.55231	0.0840	0.99	0.0831	0.0008	2519	5615
37	0.076	0.69211	0.0526	1.00	0.0526	0.0000	4892	0
39	0.042	0.80377	0.0338	1.00	0.0338	0.0000	3984	0
41	0.050	0.88184	0.0441	1.00	0.0441	0.0000	3143	0
43	0.000	0.93150	0.0000	1.00	0.0000	0.0000	1699	0
45	0.000	0.96121	0.0000	1.00	0.0000	0.0000	1658	0
47	0.000	0.97833	0.0000	1.00	0.0000	0.0000	1293	0
49	0.054	0.98799	0.0534	1.00	0.0534	0.0000	1268	0
51	0.046	0.99337	0.0457	1.00	0.0457	0.0000	1225	0
53	0.000	0.99635	0.0000	1.00	0.0000	0.0000	812	0
55	0.018	0.99799	0.0180	1.00	0.0180	0.0000	486	0
57	0.000	1.0000	0.0000	1.00	0.0000	0.0000	354	0
59	0.000	1.0000	0.0000	1.00	0.0000	0.0000	198	0
61	0.000	1.0000	0.0000	1.00	0.0000	0.0000	72	0
63	0.019	1.0000	0.0190	1.00	0.0190	0.0000	27	0
65	0.000	1.0000	0.0000	1.00	0.0000	0.0000	4	0
67	0.000	1.0000	0.0000	1.00	0.0000	0.0000	0	0
69	0.000	1.0000	0.0000	1.00	0.0000	0.0000	0	0
71	0.000	1.0000	0.0000	1.00	0.0000	0.0000	0	0
TOTAL	1.471		0.5088		0.3568	0.1520	2,386,500	1,017,003

Factor = 6,688,621

(col 7/ col 5)

1: From SURVAN, stratified mean number per tow at length.

2: From LOGEST Program using 140 mm mesh in 1993 from adjusted 130-d mm from Walsh et al. (1992) for Am plaice

4: knife-edge at 36 cm

7: From Length BIOSTAT

Table 17 . Summary of number of trips, tows, kept and discard pounds of witch flounder, discard: kept ratio observed in the large-mesh otter trawl fishery in the Georges Bank - Gulf of Maine area (observed tows only; excluding trips targeting loligo) from the Fisheries Observer Program, 1989 - 2002.

Year	January - June								July- December								Annual mt
	Trips	Tows	Fish in		D/K ratio	metric tons		Trips	Tows	fish in disc lf	Pounds		D/K ratio	metric tons			
			disc l-f	kept		discard	Landing				Disc.	kept		Disc	landings	discard	
1989	3	17	68	850	14	0.016	1298.64	20.78	4	19	411	427	142	0.333	663.58	220.97	241.75
1990	2	2	20	50	1	0.020	795.25	15.91	1	12	10	535	48	0.090	573.98	51.66	67.56
1991	4	45	15	1655	49	0.030	757.7	22.73	6	52	148	1882	159	0.084	947.63	79.60	102.33
1992	4	80	31	1896	222	0.117	1273.78	149.03	0	0	0	0	0	0.117	884.62	103.50	252.53
1993	3	37	428	1607	806	0.502	1411.03	708.34	2	24	229	1665	125	0.075	1041.84	78.14	786.48
1994	5	70	63	7298	332	0.045	1427.92	64.26	5	126	125	3794	157	0.041	1139.81	46.73	110.99
1995	31	640	1500	35968	2561	0.071	1244.48	88.36	11	199	571	8240	669	0.081	878.28	71.14	159.50
1996	17	267	272	14016	1387	0.099	1049.86	103.94	1	13	75	638	90	0.141	968.59	136.57	240.51
1997	9	203	593	10907	1831	0.168	945.2	158.79	7	134	77	4518	750	0.166	772.52	128.24	287.03
1998	7	93	200	3786	595	0.157	1095.59	172.01	2	14	3	277	37	0.134	705.15	94.49	266.50
1999	3	46	5	3663	454	0.124	1149.74	142.57	15	148	225	10057	644	0.064	914.79	58.55	201.11
2000	29	211	101	25343	666	0.026	1218.38	31.68	30	266	134	26149	1711	0.065	1159.65	75.38	107.06
2001	34	332	48	36279	1864	0.051	1596.02	81.40	45	383	459	35016	3581	0.102	1356.47	138.36	219.76
2002	27	298	795	36836	2439	0.066	1678.73	110.80	83	500	2361	44451	5268	0.119	1408.75	167.64	278.44
average						0.107								0.115			

Note: in 1993, one 'dirty' trip' during Jan-June; if excluded, d/k ratio = 0.132.; dis mt = 186.3 mt

1999 and 1998 annual discard length frequencies were used due to low numbers of fish in each half year.

Table 18. Number of trips, witch flounder kept and discarded weight (mt), discard:kept ratio from the Vessel Trip Reports, and commercial landings and estimated total discard weight.

Year	Qtr 1+2	Qtr 3+4	Total	Year	Qtr 1+2	Qtr 3+4	Total
1994 VTR trips	1228	2487		1999 VTR trips	1417	1655	
VTR kept (mt)	209.13	631.67		VTR kept (mt)	389.71	334.3	
VTR disc. (mt)	20.07	36.98		VTR disc. (mt)	18.75	17.11	
Ratio (d/k)	0.09597	0.05854		Ratio (d/k)	0.04811	0.05118	
Landings (mt)	1427.29	1139.81	2567.1	Landings (mt)	1149.74	914.79	2064.5
Discards (mt)	136.976	66.7282	203.7	Discards (mt)	55.3171	46.8204	102.1
1995 VTR trips	2674	1801		2000 VTR trips	1421	1608	
VTR kept (mt)	603.17	331.91		VTR kept (mt)	397.84	369.31	
VTR disc. (mt)	32.25	17.41		VTR disc. (mt)	21.84	19.76	
Ratio (d/k)	0.05347	0.05245		Ratio (d/k)	0.0549	0.05351	
Landings (mt)	1244.48	878.28	2122.8	Landings (mt)	1218.38	1159.65	2378.0
Discards (mt)	66.5393	46.0693	112.6	Discards (mt)	66.8847	62.0473	128.9
1996 VTR trips	2216	1662		2001 VTR trips	1327	1165	
VTR kept (mt)	469.79	411.2		VTR kept (mt)	531.92	410.55	
VTR disc. (mt)	21.62	22.93		VTR disc. (mt)	24.28	19.91	
Ratio (d/k)	0.04602	0.05576		Ratio (d/k)	0.04565	0.0485	
Landings (mt)	1049.86	968.59	2018.5	Landings (mt)	1596.02	1356.47	2952.5
Discards (mt)	48.3151	54.0121	102.3	Discards (mt)	72.8519	65.7833	138.6
1997 VTR trips	1906	1360		2002 VTR trips	791	1227	
VTR kept (mt)	355.61	302.67		VTR kept (mt)	494.02	388.52	
VTR disc. (mt)	16.39	17.22		VTR disc. (mt)	23.78	16.15	
Ratio (d/k)	0.04609	0.05689		Ratio (d/k)	0.04814	0.04157	
Landings (mt)	945.2	772.52	1717.7	Landings (mt)	1678.73	1408.75	3087.5
Discards (mt)	43.5641	43.9515	87.5	Discards (mt)	80.8068	58.5589	139.4
1998 VTR trips	1645	1129					
VTR kept (mt)	320.93	223.35					
VTR disc. (mt)	17.06	11.82					
Ratio (d/k)	0.05316	0.05292					
Landings (mt)	1095.59	705.15	1800.7				
Discards (mt)	58.2394	37.3175	95.6				

Table 19. Numbers and weight (mt) of discarded witch flounder in the large-mesh otter trawl fishery derived by three scenarios: 1) Vessel trip reports (VTR), 2) Fisheries Observer Program (FOP), and 3) survey filter method.

Year	VTR	FOP	Survey filter method
Numbers (1000's)			
1982			359.01
1983			1062.39
1984			599.49
1985			320.94
1986			78.75
1987			136.47
1988			202.67
1989			733.52
1990			959.82
1991			455.22
1992			988.11
1993			2171.25
1994	832.65		2155.99
1995	639.95	906.50	1302.62
1996	543.51	1277.79	2432.13
1997	489.12	1604.49	2071.03
1998	549.41	1541.34	1722.55
1999	520.04	1024.35	1932.76
2000	668.38	555.11	3022.89
2001	695.16	1102.21	3820.86
2002	701.19	1400.56	3475.61
Weight (mt)			
1982			42.44
1983			149.04
1984			88.81
1985			48.75
1986			12.00
1987			25.68
1988			26.55
1989		241.8	113.86
1990		67.6	155.75
1991		102.3	65.41
1992		252.5	153.31
1993		786.45	367.77
1994	203.7	110.99	406.30
1995	112.6	159.50	231.23
1996	102.3	240.51	439.88
1997	87.5	287.03	379.83
1998	95.6	266.50	316.89
1999	102.1	201.11	342.33
2000	128.9	107.05	546.34
2001	138.6	219.76	700.93
2002	139.4	278.44	645.23

Table 20. Large-mesh otter trawl discards at age in numbers (thousands of fish), mean weight (kg) and mean length (cm) at age of witch flounder, 1982 - 2002, estimated using a **survey filter method 1982-1994) and FOP data (1995-2002).**

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
Large-mesh Otter Trawl Fishery Discards in Numbers (1000's) at Age																	
1982	0.03	0.06	0.13	47.35	216.75	76.50	18.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	359.01	0.00
1983	0.00	0.02	0.66	64.20	532.92	463.25	1.34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1062.39	0.00
1984	0.00	0.00	0.11	9.17	415.36	174.59	0.26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	599.49	0.00
1985	0.00	0.00	0.10	111.86	143.96	65.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	320.94	0.00
1986	0.00	0.00	0.00	1.58	28.74	48.15	0.28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	78.75	0.00
1987	0.00	0.00	0.42	6.63	25.17	104.00	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	136.47	0.00
1988	0.00	0.04	0.00	104.77	46.54	50.60	0.72	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	202.67	0.00
1989	0.11	0.22	2.80	377.82	352.57	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	733.52	0.00
1990	0.27	1.11	2.52	103.96	355.44	496.52	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	959.82	0.00
1991	0.10	0.11	7.28	154.42	123.36	119.27	50.68	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	455.22	0.00
1992	0.13	0.94	22.51	280.70	664.19	19.17	0.47	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	988.11	0.00
1993	1.70	6.96	22.01	378.54	1371.00	391.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2171.25	0.00
1994	0.00	0.02	0.94	22.35	800.5	1330.43	1.75	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2155.99	0.00
1995	0.00	0.00	25.41	147.96	334.33	278.03	116.71	2.65	1.09	0.31	0.00	0.00	0.00	0.00	0.00	906.50	0.00
1996	0.00	0.00	0.00	22.43	419.44	737.88	98.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1277.79	0.00
1997	0.00	0.00	0.00	48.80	822.57	480.73	243.46	5.05	2.47	1.41	0.00	0.00	0.00	0.00	0.00	1604.49	0.00
1998	0.00	1.45	29.48	199.19	508.67	571.00	201.32	28.61	0.93	0.70	0.00	0.00	0.00	0.00	0.00	1541.34	0.00
1999	0.00	0.43	23.69	75.74	321.49	506.95	69.80	22.80	2.19	1.25	0.00	0.00	0.00	0.00	0.00	1024.35	0.00
2000	0.00	0.00	4.58	46.17	197.98	187.57	95.56	18.26	4.99	0.00	0.00	0.00	0.00	0.00	0.00	555.11	0.00
2001	0.00	0.00	0.85	37.22	316.95	557.06	157.64	32.29	0.09	0.09	0.00	0.00	0.00	0.00	0.00	1102.21	0.00
2002	0.00	2.01	5.09	34.81	574.48	577.81	161.64	33.89	6.61	2.47	1.04	0.00	0.00	0.00	0.00	1400.56	0.00

Table 20 continued. Discards in the large-mesh otter trawl fishery (survey filter method (1982-1994) and FOP data (1995-2002) .

Year	Age														TOTAL	11+	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13			14+
Large-mesh Otter Trawl Fishery Discards Mean Weight (kg) at Age																	
1982	0.000	0.002	0.014	0.050	0.128	0.121	0.169									0.118	-
1983		0.009	0.029	0.081	0.132	0.158	0.209									0.140	-
1984			0.014	0.072	0.144	0.162	0.209									0.148	-
1985			0.031	0.133	0.163	0.160										0.152	-
1986				0.105	0.125	0.170	0.209									0.152	-
1987			0.014	0.105	0.122	0.210	0.256									0.188	-
1988		0.002		0.086	0.161	0.195	0.256									0.131	-
1989	0.001	0.013	0.044	0.134	0.179											0.155	-
1990	0.001	0.018	0.028	0.105	0.146	0.187										0.162	-
1991	0.001	0.010	0.048	0.093	0.140	0.191	0.210									0.144	-
1992	0.001	0.015	0.057	0.129	0.168	0.214	0.256									0.155	-
1993	0.001	0.014	0.050	0.129	0.175	0.199										0.169	-
1994		0.026	0.044	0.103	0.175	0.198	0.256									0.188	-
1995			0.059	0.095	0.152	0.221	0.252	0.473	0.595	0.702						0.176	-
1996				0.077	0.145	0.208	0.251									0.188	-
1997				0.108	0.157	0.185	0.242	0.495	0.471	0.702						0.179	-
1998		0.008	0.041	0.089	0.165	0.200	0.219	0.225	0.355	0.370						0.174	-
1999		0.021	0.037	0.097	0.162	0.218	0.281	0.406	0.447	0.505						0.196	-
2000			0.066	0.096	0.155	0.198	0.280	0.313	0.403							0.193	-
2001			0.070	0.130	0.171	0.208	0.230	0.258	0.309	0.309						0.199	-
2002		0.018	0.054	0.122	0.180	0.205	0.234	0.264	0.414	0.577	0.566					0.199	-
Mean 1982-02	0.001	0.009	0.037	0.102	0.155	0.180	0.236	0.264	0.428	0.528	0.566						

Table 20 continued. Discard in the large-mesh otter trawl fishery (survey filter method (1982-1994) and FOP data (1995-2002) .

Year	Age														TOTAL	11+	
	0	1	2	3	4	5	6	7	8	9	10	11	12	13			14+
Large-mesh Otter Trawl Fishery Discards Mean Length (cm) at Age																	
1982	5.0	7.8	15.0	21.4	28.3	28.1	31.0	-	-	-	-					27.5	-
1983		13.0	18.5	24.7	28.6	30.4	33.0	-	-	-	-					29.2	-
1984		-	15.0	23.6	29.5	30.6	33.0	-	-	-	-					29.7	-
1985		-	19.0	28.8	30.7	30.5	-	-	-	-	-					30.0	-
1986	5.0	-	-	27.0	28.3	31.1	33.0	-	-	-	-					30.0	-
1987		-	15.0	27.0	28.1	33.0	35.0	-	-	-	-					31.8	-
1988		9.0	-	25.4	30.4	32.3	35.0	-	-	-	-					28.3	-
1989	5.9	14.4	20.7	28.8	31.5	-	-	-	-	-	-					30.1	-
1990	6.1	16.0	18.1	26.8	29.6	31.8	-	-	-	-	-					30.4	-
1991	5.5	12.7	21.3	25.8	29.2	32.1	33.0	-	-	-	-					29.1	-
1992	5.7	15.0	22.5	28.4	30.8	33.2	35.0	-	-	-	-					30.0	-
1993	5.5	14.5	21.5	28.5	31.2	32.5	-	-	-	-	-					30.8	-
1994		17.9	20.7	26.5	31.2	32.4	35.0	-	-	-	-					31.9	-
1995			22.7	25.9	29.9	33.5	34.8	41.8	44.7	47.0						30.8	-
1996				24.6	29.5	32.9	34.8									31.7	-
1997				27.1	30.3	31.8	34.4	42.4	41.6	47.0						31.3	-
1998		12.3	20.3	25.6	30.7	32.5	33.3	33.4	38.5	39.0						30.9	-
1999		17.0	19.7	26.1	30.5	33.3	35.9	40.0	41.1	42.7						31.9	-
2000			23.5	26.1	30.0	32.3	35.8	36.7	39.7							31.7	-
2001			23.9	28.5	31.0	32.9	33.9	35.1	37.0	37.0						32.4	-
2002		16.0	21.9	27.7	31.4	32.7	34.0	35.1	40.1	44.2	44.0					32.3	

Table 21. **Total USA commercial catch** [landings + shrimp trawl discards + large-mesh otter trawl discards using survey filter and FOP] in numbers, (thousands of fish), mean weight (kg) and mean length (cm) at age of witch flounder, 1982 - 2002.

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commercial Catch in Numbers (1000's) at Age																	
1982	0.03	0.06	1.72	190.5	1064.5	1207.7	1475.4	665.2	656.0	399.5	239.4	201.0	356.3	183.7	837.4	7478.4	1578.4
1983	0.00	0.02	4.28	337.1	1346.2	1520.8	1575.1	1590.2	977.8	737.7	510.4	366.0	287.3	289.1	733.1	10275.1	1675.5
1984	0.00	0.33	0.88	146.6	1466.3	2002.7	1739.6	1486.5	1497.5	696.7	375.1	279.5	356.4	261.3	821.6	11131.0	1718.8
1985	0.00	0.34	3.47	123.6	1176.1	2118.2	1936.2	1524.9	1247.9	606.0	400.4	261.2	221.5	170.7	705.8	10496.4	1359.2
1986	0.00	0.53	3.86	23.0	377.1	1516.8	2775.4	1566.9	834.9	412.7	222.8	188.2	157.0	137.0	276.0	8492.1	758.2
1987	2.08	18.92	79.93	22.3	181.3	467.1	1280.1	1574.7	870.9	480.6	252.4	132.4	90.8	62.1	204.1	5719.6	489.4
1988	0.42	14.66	130.29	600.3	139.9	264.3	658.3	1382.7	1154.1	401.5	266.7	124.1	94.0	71.9	307.5	5610.6	597.5
1989	0.85	10.69	50.32	447.1	436.3	65.2	314.3	759.4	882.1	349.7	123.4	73.2	61.1	56.7	157.1	3787.2	348.0
1990	1.46	6.29	95.30	343.9	634.1	1103.2	255.6	273.9	471.1	333.9	81.4	43.1	38.5	19.1	76.9	3777.6	177.5
1991	3.06	17.90	23.26	441.8	405.8	863.7	575.4	235.8	244.6	292.1	313.6	51.8	44.0	22.5	139.5	3674.7	257.8
1992	2.84	44.35	159.43	399.5	1255.6	859.2	936.0	717.0	201.6	177.9	120.0	217.6	46.3	26.5	86.5	5250.3	377.0
1993	113.76	85.80	129.59	417.2	1807.3	1419.0	918.1	597.2	585.6	218.8	278.5	113.9	32.6	103.6	140.4	6961.3	390.5
1994	8.06	1368.48	496.44	42.0	1001.8	2759.9	1288.0	826.9	196.7	539.2	113.5	71.4	40.2	132.3	80.4	8965.1	324.3
1995	2.68	49.95	655.51	628.8	370.3	1041.2	1714.5	851.4	268.5	97.5	269.5	55.0	43.9	8.1	49.9	6106.7	156.8
1996	5.21	32.68	50.83	121.9	524.4	1207.7	1362.9	1430.5	263.2	215.5	57.1	78.8	3.6	13.0	18.2	5385.5	113.6
1997	8.68	74.91	102.92	135.3	1058.5	1016.0	1293.4	1019.1	593.8	84.6	49.8	17.9	36.6	2.2	13.4	5506.9	70.1
1998	49.78	392.89	294.20	331.2	556.9	1063.7	1415.5	1611.6	371.4	142.1	15.5	37.2	5.6	19.9	7.7	6315.1	70.3
1999	32.11	252.94	194.45	160.9	526.1	1095.8	1462.0	1201.1	765.3	252.5	31.6	40.8	0.0	0.0	13.5	6029.2	54.4
2000	21.61	169.95	119.51	103.5	286.5	455.9	1158.0	1629.4	1032.6	623.7	94.8	174.3	6.2	5.0	27.4	5908.3	212.8
2001	12.33	96.96	66.42	69.9	343.2	940.4	1098.3	1701.3	1459.6	634.4	425.4	95.8	163.5	8.6	38.8	7154.9	306.7
2002	2.32	20.24	17.42	41.0	749.1	1222.9	1404.4	2132.2	1281.1	634.6	97.5	102.7	11.0	65.6	25.3	7807.9	205.2

Table 21 continued. **Total USA commercial catch** (landings + shrimp trawl discards + LM otter trawl discards) .

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commerical Mean Weight (kg) at Age																	
1982	0.000	0.002	0.038	0.152	0.242	0.329	0.421	0.550	0.727	0.886	0.983	1.146	1.255	1.310	1.553	0.662	1.406
1983		0.009	0.038	0.149	0.202	0.270	0.409	0.518	0.613	0.795	0.977	1.116	1.208	1.321	1.551	0.600	1.357
1984		0.017	0.040	0.151	0.229	0.328	0.421	0.539	0.664	0.817	0.922	1.004	1.212	1.332	1.511	0.607	1.339
1985		0.017	0.023	0.128	0.237	0.305	0.429	0.565	0.691	0.842	0.964	1.057	1.193	1.311	1.470	0.590	1.326
1986		0.017	0.026	0.089	0.206	0.299	0.408	0.533	0.676	0.853	0.975	1.132	1.199	1.317	1.521	0.546	1.321
1987	0.006	0.015	0.033	0.081	0.191	0.298	0.433	0.561	0.686	0.828	0.980	1.067	1.222	1.386	1.467	0.611	1.303
1988	0.004	0.006	0.017	0.045	0.203	0.311	0.434	0.538	0.668	0.819	0.980	1.074	1.190	1.290	1.477	0.592	1.326
1989	0.009	0.012	0.034	0.122	0.170	0.321	0.425	0.574	0.682	0.818	0.968	1.128	1.258	1.315	1.519	0.581	1.358
1990	0.004	0.012	0.029	0.062	0.186	0.257	0.438	0.586	0.688	0.849	1.049	1.213	1.262	1.521	1.669	0.437	1.454
1991	0.004	0.014	0.035	0.062	0.199	0.344	0.420	0.578	0.702	0.836	0.974	1.099	1.369	1.537	1.536	0.509	1.420
1992	0.003	0.007	0.026	0.103	0.230	0.379	0.459	0.614	0.739	0.822	0.882	1.039	1.337	1.459	1.640	0.457	1.243
1993	0.003	0.009	0.027	0.122	0.202	0.318	0.432	0.535	0.666	0.882	1.023	1.118	1.199	1.368	1.519	0.428	1.335
1994	0.005	0.004	0.019	0.070	0.202	0.280	0.430	0.534	0.691	0.832	0.909	1.083	1.172	1.204	1.576	0.344	1.266
1995	0.005	0.007	0.024	0.051	0.158	0.328	0.435	0.561	0.690	0.910	0.974	1.101	1.203	1.411	1.406	0.393	1.243
1996	0.004	0.019	0.031	0.060	0.149	0.265	0.422	0.554	0.708	0.856	0.974	1.114	1.401	1.440	1.558	0.435	1.232
1997	0.004	0.023	0.033	0.070	0.189	0.274	0.376	0.495	0.627	0.868	1.037	1.168	1.196	1.687	1.659	0.376	1.293
1998	0.003	0.006	0.024	0.070	0.163	0.260	0.359	0.487	0.584	0.869	0.978	1.115	1.132	1.261	1.557	0.338	1.206
1999	0.003	0.006	0.024	0.068	0.217	0.291	0.400	0.514	0.584	0.627	0.917	0.683			1.442	0.387	0.872
2000	0.003	0.006	0.024	0.066	0.196	0.273	0.368	0.448	0.532	0.633	0.677	0.834	1.167	1.298	1.379	0.432	0.925
2001	0.003	0.006	0.023	0.089	0.176	0.256	0.362	0.465	0.550	0.646	0.647	0.718	0.816	1.016	1.206	0.453	0.840
2002	0.003	0.007	0.032	0.110	0.220	0.278	0.395	0.474	0.553	0.651	0.821	0.844	0.716	0.993	1.120	0.444	0.918
Mean																	
1982-02	0.0040	0.0106	0.0287	0.0914	0.1984	0.2984	0.4131	0.5344	0.6534	0.8066	0.9339						1.2372
1999-02	0.0031	0.0063	0.0258	0.0830	0.2021	0.2746	0.3813	0.4752	0.5548	0.6393	0.7656						0.8887

Table 21 continued. **Total USA commercial catch** (landings+ shrimp trawl discards + LM otter trawl discards) .

Year	Age															TOTAL	11+
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+		
USA Commerical Mean Length (cm) at Age																	
1982	5.0	7.8	19.9	28.1	33.5	36.8	39.7	42.9	46.5	49.3	50.9	53.2	54.6	55.2	58.0	43.3	56.3
1983		13.0	20.0	28.6	31.8	34.7	39.4	42.2	44.2	47.7	50.7	52.8	54.0	56.6	55.8	35.0	55.0
1984		15.7	20.0	28.4	33.1	36.9	39.7	42.7	45.3	48.2	49.9	51.2	54.1	55.6	57.6	42.7	55.5
1985		15.7	16.9	28.4	33.6	36.1	39.9	43.3	45.9	48.6	50.6	51.9	53.8	55.3	57.1	42.4	55.3
1986		15.7	17.3	25.1	32.2	36.0	39.3	42.5	45.6	48.8	50.7	53.0	53.9	55.4	57.7	41.8	55.3
1987	10.6	15.3	19.0	24.5	31.3	35.9	40.1	43.2	45.8	48.4	50.8	52.1	54.2	56.2	57.1	43.2	55.1
1988	10.2	10.9	15.6	20.4	31.8	36.3	40.1	42.7	45.4	48.2	50.8	52.1	53.7	55.0	57.1	41.4	55.3
1989	12.6	13.9	19.0	27.8	30.9	36.8	39.9	43.5	45.6	48.1	50.6	52.9	54.6	55.3	57.6	41.6	55.7
1990	9.7	14.0	18.0	22.3	31.1	34.3	40.2	43.7	45.8	48.7	51.8	54.1	54.6	57.8	59.2	37.5	56.8
1991	9.6	14.1	18.8	22.6	31.6	37.4	39.6	43.3	46.1	48.5	50.6	52.5	56.0	57.9	57.9	39.4	56.5
1992	9.1	10.9	17.4	26.1	33.1	38.5	40.7	44.3	46.8	48.3	49.2	51.7	55.5	57.0	58.9	38.6	54.2
1993	9.1	12.2	17.7	27.9	32.3	36.6	40.0	42.6	45.3	49.3	51.5	52.8	53.9	55.9	57.7	37.6	55.5
1994	10.7	9.8	15.9	22.8	32.2	35.1	39.7	42.3	45.6	48.0	49.1	51.8	53.1	53.4	57.8	33.0	54.1
1995	10.9	11.6	17.3	21.1	30.1	36.7	39.8	42.8	45.4	49.3	50.1	52.0	53.4	56.0	55.8	35.7	53.8
1996	10.0	15.3	18.4	22.5	29.5	34.7	39.4	42.7	45.8	48.4	50.1	52.2	55.8	56.2	57.6	38.6	53.6
1997	10.2	16.1	18.9	23.3	31.4	34.8	38.2	41.3	44.2	48.5	51.1	52.9	53.3	59.0	58.7	36.9	54.4
1998	9.4	10.4	17.1	23.5	30.5	34.5	37.6	41.1	43.3	48.6	50.5	52.3	52.7	54.1	57.6	34.6	53.4
1999	9.4	10.4	17.2	23.0	32.5	35.6	38.9	41.8	43.3	44.3	49.4	45.2			56.4	36.8	48.0
2000	9.4	10.4	17.1	22.9	31.5	34.8	38.0	40.1	42.1	44.2	45.2	47.7	53.0	54.6	55.3	38.4	49.0
2001	9.4	10.4	16.9	24.7	31.1	34.4	37.7	40.5	42.5	44.6	44.5	45.9	47.7	50.8	53.3	39.3	47.9
2002	9.4	11.0	18.3	26.6	32.8	35.1	38.7	40.8	42.6	44.6	47.9	48.2	45.8	50.5	52.2	39.7	49.3

Table 22. Mean weights at age (kg) at the beginning of the year (January 1) for witch flounder, 1982-2002. Values derived from catch mean weight-at-age data using procedures described by Rivard (1980).

Year	Age										
	1	2	3	4	5	6	7	8	9	10	11+
1982	0.0005	0.0193	0.1316	0.2287	0.2951	0.3789	0.5210	0.6952	0.8437	0.9332	1.4056
1983	0.0042	0.0090	0.0755	0.1753	0.2552	0.3670	0.4667	0.5806	0.7602	0.9304	1.3575
1984	0.0143	0.0191	0.0763	0.1848	0.2577	0.3371	0.4696	0.5865	0.7077	0.8561	1.3393
1985	0.0137	0.0199	0.0719	0.1893	0.2639	0.3751	0.4880	0.6103	0.7477	0.8875	1.3255
1986	0.0121	0.0208	0.0456	0.1623	0.2663	0.3525	0.4781	0.6180	0.7677	0.9061	1.3209
1987	0.0142	0.0236	0.0456	0.1300	0.2477	0.3602	0.4783	0.6047	0.7481	0.9143	1.3031
1988	0.0027	0.0161	0.0384	0.1284	0.2438	0.3598	0.4829	0.6122	0.7496	0.9008	1.3256
1989	0.0076	0.0145	0.0457	0.0874	0.2554	0.3636	0.4992	0.6057	0.7392	0.8904	1.3577
1990	0.0066	0.0187	0.0457	0.1510	0.2092	0.3752	0.4990	0.6284	0.7609	0.9263	1.4542
1991	0.0103	0.0202	0.0426	0.1108	0.2533	0.3286	0.5032	0.6414	0.7584	0.9094	1.4198
1992	0.0033	0.0190	0.0604	0.1192	0.2745	0.3974	0.5081	0.6536	0.7596	0.8587	1.2431
1993	0.0062	0.0135	0.0563	0.1445	0.2703	0.4046	0.4955	0.6395	0.8073	0.9170	1.3353
1994	0.0019	0.0130	0.0434	0.1571	0.2380	0.3698	0.4802	0.6080	0.7444	0.8954	1.2656
1995	0.0036	0.0105	0.0308	0.1050	0.2572	0.3491	0.4909	0.6068	0.7931	0.9002	1.2426
1996	0.0142	0.0151	0.0382	0.0868	0.2048	0.3719	0.4907	0.6301	0.7683	0.9416	1.2316
1997	0.0229	0.0251	0.0465	0.1061	0.2021	0.3154	0.4569	0.5895	0.7840	0.9422	1.2930
1998	0.0030	0.0239	0.0483	0.1069	0.2215	0.3135	0.4279	0.5378	0.7381	0.9215	1.2061
1999	0.0030	0.0120	0.0406	0.1233	0.2180	0.3226	0.4293	0.5333	0.6055	0.8924	0.8721
2000	0.0031	0.0120	0.0399	0.1150	0.2436	0.3272	0.4236	0.5231	0.6078	0.6517	0.9247
2001	0.0026	0.0118	0.0463	0.1076	0.2237	0.3145	0.4137	0.4967	0.5864	0.6400	0.8403
2002	0.0038	0.0138	0.0504	0.1395	0.2213	0.3179	0.4140	0.5072	0.5982	0.7283	0.9179
mean											
1982-2002	0.0073	0.0167	0.0533	0.1361	0.2439	0.3525	0.4722	0.5956	0.7322	0.8735	1.2372
1999-2002	0.0031	0.0124	0.0443	0.1214	0.2267	0.3206	0.4202	0.5151	0.5995	0.7281	0.8888

Table 23. USA commercial witch flounder landings (L), days fished (DF), and landings per day fished (L/DF), by vessel tonnage class, for otter trawl trips which any witch flounder were landed, and for otter trawl trips in which 40% or more of the total catch consisted of witch flounder, in the Gulf of Maine-Georges Bank region (SA 51, 52, 56), 1973 - 2002. *Note: in 1994-2002, Vessel Trip Report data were used.*

YEAR	CLASS 2			CLASS 3			CLASS 4			TOTAL		
	L	DF	L\DF	L	DF	L\DF	L	DF	L\DF	L	DF	L\DF
ALL TRIPS												
1973	802	2620	0.31	1284	6236	0.21	234	859	0.27	2320	9715	0.25
1974	497	2478	0.20	1029	7092	0.15	157	1004	0.16	1683	10574	0.16
1975	679	2354	0.29	1126	7728	0.15	153	1178	0.13	1957	11260	0.19
1976	756	2826	0.27	913	6373	0.14	97	860	0.11	1765	10059	0.19
1977	1074	3183	0.34	1070	6025	0.18	157	872	0.18	2302	10080	0.25
1978	1372	4033	0.34	1658	7053	0.24	277	1225	0.23	3307	12310	0.28
1979	946	4465	0.21	1467	6757	0.22	283	1570	0.18	2696	12792	0.21
1980	1062	4932	0.22	1428	7120	0.20	376	1997	0.19	2866	14049	0.20
1981	1069	3748	0.29	1637	7015	0.23	423	2595	0.16	3129	13358	0.24
1982	1162	4430	0.26	2346	8626	0.27	905	3559	0.25	4413	16615	0.27
1983	1203	3930	0.31	2796	9581	0.29	1308	4544	0.29	5307	18056	0.29
1984	1281	4069	0.31	3245	12157	0.27	1423	4769	0.30	5949	20994	0.28
1985	1195	3794	0.31	2765	12664	0.22	1600	5530	0.29	5560	21988	0.26
1986	806	3289	0.25	2031	10525	0.19	1177	5287	0.22	4015	19101	0.21
1987	647	2833	0.23	1623	9593	0.17	845	5035	0.17	3114	17461	0.18
1988	560	2986	0.19	1463	8948	0.16	951	4871	0.20	2973	16805	0.18
1989	283	2269	0.12	959	8538	0.11	618	4292	0.14	1860	15099	0.12
1990	265	2649	0.10	661	7736	0.09	347	4172	0.08	1274	14557	0.09
1991	316	3135	0.10	830	9076	0.09	383	4681	0.08	1529	16892	0.09
1992	352	3589	0.10	1148	10720	0.11	414	5005	0.08	1914	19314	0.10
1993	380	3321	0.11	1347	10872	0.12	530	4711	0.11	2257	18904	0.12
1994	261	2067	0.13	581	5126	0.11	302	2384	0.13	1143	9578	0.12
1995	291	2784	0.10	852	7328	0.12	462	3911	0.12	1605	14023	0.11
1996	369	2647	0.14	908	6992	0.13	399	3200	0.12	1677	12839	0.13
1997	371	2666	0.14	731	5084	0.14	277	2395	0.12	1379	10145	0.14
1998	393	2641	0.15	694	4697	0.15	255	1867	0.14	1342	9205	0.15
1999	407	2316	0.18	682	4387	0.16	288	1675	0.17	1378	8377	0.17
2000	555	2665	0.21	993	5431	0.18	398	2564	0.16	1946	10660	0.18
2001	321	1349	0.24	771	3750	0.21	396	2115	0.19	1488	7213	0.21
2002	320	875	0.37	446	1781	0.25	185	988	0.19	951	3644	0.28

Table 23 continued.

YEAR	CLASS 2			CLASS 3			CLASS 4			TOTAL		
	L	DF	L\DF	L	DF	L\DF	L	DF	L\DF	L	DF	L\DF
40% TRIPS												
1973	306	208	1.47	392	271	1.45	96	58	1.66	793	536	1.48
1974	134	99	1.34	169	112	1.50	21	16	1.25	323	228	1.42
1975	292	171	1.71	208	168	1.24	4	4	1.09	504	343	1.51
1976	211	144	1.47	137	90	1.54	3	1	3.38	352	234	1.51
1977	151	93	1.62	129	84	1.53	1	4	0.26	281	182	1.57
1978	214	162	1.33	197	82	2.39	7	2	3.58	418	246	1.87
1979	93	79	1.17	103	69	1.49	7	2	3.45	203	151	1.41
1980	93	82	1.14	107	40	2.66	54	25	2.17	254	147	2.00
1981	101	54	1.87	239	108	2.21	22	13	1.69	362	175	2.08
1982	172	112	1.53	289	136	2.13	55	31	1.75	516	279	1.89
1983	183	140	1.30	519	279	1.86	48	30	1.59	750	450	1.70
1984	234	210	1.12	705	595	1.18	176	98	1.80	1115	903	1.27
1985	266	277	0.96	465	580	0.80	177	143	1.24	909	1000	0.93
1986	185	236	0.78	499	785	0.64	127	169	0.75	811	1190	0.69
1987	155	195	0.79	377	569	0.66	86	109	0.78	617	873	0.71
1988	137	176	0.78	517	905	0.57	202	254	0.79	856	1335	0.66
1989	45	67	0.67	128	256	0.50	77	112	0.69	250	435	0.59
1990	36	57	0.63	49	85	0.58	9	16	0.54	94	158	0.60
1991	35	76	0.46	55	106	0.52	1	1	0.83	92	183	0.50
1992	42	65	0.65	181	382	0.48	25	7	3.32	248	454	0.79
1993	76	140	0.54	266	538	0.49	30	42	0.71	372	720	0.52
1994	95	221	0.43	90	225	0.40	17	22	0.77	202	468	0.45
1995	90	237	0.38	155	323	0.48	28	53	0.54	274	613	0.45
1996	139	309	0.45	169	378	0.45	40	60	0.67	348	746	0.47
1997	98	238	0.41	158	311	0.51	17	36	0.48	273	585	0.47
1998	154	327	0.47	172	263	0.65	5	6	0.90	331	596	0.57
1999	164	370	0.44	140	240	0.59	17	14	1.18	321	624	0.54
2000	212	329	0.64	164	200	0.82	21	14	1.49	396	543	0.76
2001	97	117	0.83	138	143	0.96	8	7	1.28	244	267	0.92
2002	132	127	1.05	104	91	1.14	0	0	0.00	236	217	1.09

Table 24. The sum of the 1963-2001 NEFSC autumn stratified mean number per tow by stratum (Stratum sum), the percentage of annual stratum sampling which produced no catch (% Zero Catch), and the percentage of stratum contribution (all years) to the total (% Total).

Strata	Stratum sum	% Zero Catch	% Total
1	0.00	100.0	0.0
2	0.70	84.6	0.1
3	0.22	92.3	0.0
4	2.88	76.9	0.3
5	0.23	97.4	0.0
6	19.30	48.7	1.8
7	0.43	94.9	0.0
8	2.51	64.1	0.2
9	0.53	92.3	0.0
10	8.54	56.4	0.8
11	1.23	92.3	0.1
12	2.01	86.8	0.2
13	7.80	43.6	0.7
14	0.68	89.7	0.1
15	4.26	69.2	0.4
16	0.98	74.4	0.1
17	1.12	92.3	0.1
18	7.40	74.4	0.7
19	0.00	100.0	0.0
20	0.02	97.4	0.0
21	1.43	89.7	0.1
22	73.23	5.1	6.8
23	5.75	66.7	0.5
24	85.01	2.6	7.9
25	0.61	94.9	0.1
26	73.21	5.1	6.8
27	94.59	0.0	8.8
28	22.08	15.4	2.1
29	18.72	7.7	1.7
30	8.93	56.4	0.8
33	6.88	61.5	0.6
34	55.51	2.6	5.2
351	93.13	8.3	8.7
36	48.86	0.0	4.5
37	109.01	0.0	10.2
38	185.36	0.0	17.3
39	38.84	5.1	3.6
40	85.47	10.3	8.0
61	0.00	100.0	0.0
62	0.00	100.0	0.0
63	0.10	97.1	0.0
64	0.60	90.6	0.1
65	0.00	100.0	0.0

Strata	Stratum sum	% Zero Catch	% Total
66	0.00	100.0	0.0
67	0.05	97.1	0.0
68	1.10	75.8	0.1
69	0.02	97.1	0.0
70	0.00	100.0	0.0
71	0.00	100.0	0.0
72	3.35	81.8	0.3
73	0.00	100.0	0.0
74	0.00	100.0	0.0
75	0.05	97.1	0.0
76	1.20	80.0	0.1
Total	1073.89		100.0

Strata set 22-30,36-40	92.5
Strata set 6,22-30,36-40	93.9
Strata set 22,24,26-30,36-40	91.8

Table 25. Stratified mean number, weight (kg), length (cm), and individual weight (kg) per tow of witch flounder in NEFSC offshore spring and autumn bottom trawl surveys in Gulf of Maine-Georges Bank region (strata 22-30,36-40), 1963-2003.

Year	SPRING				AUTUMN			
	Number per tow	Weight per tow	Length per tow	Ave. wt. per tow	Number per tow	Weight per tow	Length per tow	Ave. wt. per tow
1963	-	-	-	-	5.52	3.46	39.7	0.627
1964	-	-	-	-	2.89	2.00	44.2	0.691
1965	-	-	-	-	3.94	2.27	40.6	0.577
1966	-	-	-	-	7.80	4.56	41.2	0.584
1967	-	-	-	-	3.01	2.02	43.7	0.673
1968	4.83	3.35	42.3	0.695	4.82	3.49	44.8	0.724
1969	3.74	2.53	45.3	0.676	5.81	4.40	43.5	0.757
1970	6.39	4.49	44.7	0.702	4.89	3.71	45.0	0.760
1971	2.70	2.04	46.5	0.756	4.32	2.95	42.1	0.683
1972	5.35	4.01	45.8	0.749	3.24	2.42	43.9	0.747
1973	8.20	6.21	44.8	0.758	3.18	2.05	43.6	0.646
1974	6.23	3.62	39.3	0.581	2.38	1.58	41.0	0.666
1975	3.72	2.75	43.9	0.739	1.66	1.03	39.8	0.621
1976	5.50	3.70	42.3	0.673	1.34	0.94	41.9	0.699
1977	4.20	1.96	37.2	0.467	5.06	3.38	42.0	0.669
1978	3.87	2.56	41.7	0.662	4.04	2.94	42.8	0.727
1979	2.91	1.71	38.2	0.587	1.94	1.62	45.2	0.838
1980	8.46	3.89	36.0	0.460	2.62	2.04	43.7	0.777
1981	8.14	4.05	38.0	0.497	3.66	2.19	40.4	0.600
1982	3.64	1.87	37.2	0.513	0.99	0.83	44.7	0.842
1983	6.41	2.74	36.3	0.427	4.72	2.12	36.7	0.448
1984	3.00	1.66	39.9	0.554	4.37	2.34	39.7	0.534
1985	5.18	2.75	40.3	0.531	2.76	1.59	41.9	0.577
1986	2.07	1.35	44.1	0.650	1.59	1.09	43.3	0.683
1987	1.01	0.65	43.4	0.646	0.48	0.37	43.9	0.774
1988	1.43	0.85	42.3	0.590	1.38	0.57	35.2	0.414
1989	1.95	0.74	35.8	0.382	0.89	0.38	31.4	0.423
1990	0.63	0.24	35.2	0.378	2.00	0.40	24.7	0.200
1991	1.68	0.57	31.5	0.341	2.08	0.54	29.2	0.258
1992	1.26	0.48	34.8	0.383	0.94	0.24	29.5	0.254
1993	1.47	0.36	30.3	0.245	5.15	0.54	17.0	0.105
1994	3.13	0.53	27.4	0.170	2.21	0.42	24.9	0.191
1995	1.88	0.47	30.6	0.248	4.74	0.62	25.7	0.132
1996	1.36	0.28	30.5	0.204	5.38	1.02	29.7	0.189
1997	2.22	0.43	31.0	0.195	5.11	0.77	24.9	0.150
1998	4.27	0.77	29.0	0.179	3.70	0.47	24.2	0.127
1999	3.15	0.48	28.1	0.153	5.92	0.88	26.3	0.148
2000	3.45	0.52	27.3	0.151	6.63	1.11	27.1	0.167
2001	4.41	0.75	29.5	0.171	7.94	1.71	32.3	0.216
2002	8.10	1.62	31.4	0.199	4.31	1.06	33.2	0.247
2003	5.20	1.30	34.2	0.250				

Note: During 1963-1984, BMV oval doors were used in the spring and autumn surveys; since 1985, Portuguese polyvalent doors have been used in both surveys. No significant differences in catchability were found for witch flounder, therefore no adjustments have been made (Byrne and Forrester, MS 1991). No significant differences were found between research vessels, and no adjustment have been made (Byrne and Forrester, MS 1991).

Spring surveys during 1973-1981 were accomplished with a 41 Yankee trawl; in all other years, a 36 Yankee trawl was used. No adjustments have been made.

Table 26. Stratified mean number, weight (kg), length (cm) per tow of witch flounder in Massachusetts Division of Marine Fisheries inshore spring and autumn surveys in the Cape Cod Bay and Mass. Bay region (Regions 4 and 5), 1978-2002.

Year	SPRING			AUTUMN		
	Number per tow	Weight per tow	Length per tow	Number per tow	Weight per tow	Length per tow
1978	2.98	2.15	45.3	2.47	2.41	48.2
1979	1.36	1.41	47.9	1.39	1.09	44.4
1980	1.49	1.44	46.0	1.66	1.77	48.2
1981	3.74	3.18	43.7	0.36	0.23	42.3
1982	1.23	0.97	46.1	1.24	0.76	40.3
1983	2.15	1.29	38.9	3.79	2.68	45.9
1984	1.50	1.01	41.6	0.62	0.45	44.7
1985	1.12	0.82	43.9	0.83	0.57	44.2
1986	0.90	0.83	47.1	0.32	0.27	46.3
1987	1.45	1.10	44.7	0.26	0.20	45.7
1988	0.36	0.29	46.1	0.39	0.24	40.8
1989	0.17	0.07	35.9	0.21	0.13	42.2
1990	0.40	0.32	45.0	0.06	0.03	37.5
1991	0.17	0.08	37.4	0.37	0.22	41.5
1992	0.34	0.24	41.3	0.45	0.25	41.3
1993	0.03	0.01	33.0	0.39	0.19	40.2
1994	0.00	0.00	-	0.53	0.21	34.9
1995	0.08	0.05	36.6	2.17	0.46	28.7
1996	0.02	<0.01	21.0	0.05	0.02	40.0
1997	0.04	0.01	31.5	1.20	0.36	36.0
1998	0.00	0.00	-	0.44	0.15	35.7
1999	0.01	<0.01	11.0	1.38	0.38	34.9
2000	0.95	0.11	24.4	1.28	0.43	34.4
2001	0.16	0.04	33.0	0.66	0.20	34.8
2002	0.11	0.03	32.5	2.40	0.79	37.1

Table 27. Stratified mean number, weight (kg), length (cm), and individual weight (kg) per tow of witch flounder in the ASMFC summer shrimp surveys in the Gulf of Maine (Strata set 1,3,6,8), 1984 - 2002.

Year	Number per tow	Weight per tow	Length per tow	Individual Weight per tow
1984	4.68	1.60	33.9	0.341
1985	6.19	2.52	36.2	0.408
1986	2.05	0.74	35.9	0.362
1987	4.87	1.50	26.5	0.307
1988	2.53	0.60	25.8	0.238
1989	2.92	0.31	22.8	0.105
1990	6.66	1.02	24.5	0.154
1991	14.94	1.20	19.6	0.080
1992	24.28	1.91	20.5	0.079
1993	21.42	0.50	12.8	0.023
1994	36.36	2.20	19.1	0.061
1995	17.95	1.48	22.6	0.082
1996	15.45	1.95	25.2	0.126
1997	23.20	1.42	19.1	0.061
1998	7.35	0.52	21.9	0.071
1999	110.07	5.93	18.7	0.054
2000	32.43	3.09	24.2	0.095
2001	41.52	5.57	27.2	0.134
2002	45.25	7.05	28.8	0.156

Table 28. Number of witch flounder caught, aged, percent of fish sampled, and the maximum age observed in the NEFSC spring and autumn bottom trawl surveys (strata 22-30, 36-40), 1980 - 2002.

Year	Spring				Autumn			
	Caught	Aged	% Sampled	Max. Age	Caught	Aged	% Sampled	Max. Age
1980	593	361	60.9	24	189	146	77.2	24
1981	557	209	37.5	23	202	143	70.8	22
1982	245	69	28.2	18	64	53	82.8	24
1983	410	176	42.9	20	359	154	42.9	22
1984	171	145	84.8	26	293	204	69.6	21
1985	269	151	56.1	25	340	232	68.2	30
1986	119	118	99.2	22	258	218	84.5	22
1987	108	108	100.0	24	30	27	90.0	24
1988	74	67	90.5	12	93	82	88.2	20
1989	100	91	91.0	18	59	55	93.2	21
1990	33	27	81.8	16	131	118	90.1	18
1991	93	87	93.5	15	187	107	57.2	11
1992	86	75	87.2	17	79	67	84.8	18
1993	88	81	92.0	19	414	166	40.1	16
1994	196	127	64.8	16	174	102	58.6	21
1995	142	106	74.6	19	352	174	49.4	14
1996	84	72	85.7	13	295	169	57.3	11
1997	129	79	61.2	12	368	243	66.0	12
1998	367	281	76.6	13	359	217	60.4	13
1999	187	138	73.8	10	556	244	43.9	10
2000	231	141	61.0	10	411	224	54.5	12
2001	315	197	62.5	10	475	234	49.3	10
2002	568	289	50.9	21	339	201	59.3	12

Table 29. Stratified mean number per tow at age of witch flounder in NEFSC bottom trawl spring and autumn surveys (Strata 22-30, 36-40), 1980-2002, preliminary spring 2003.

SPRING	Age															Total
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14 +	
1980	0.00	0.06	0.23	0.95	1.52	0.72	1.20	1.02	0.38	0.40	0.31	0.30	0.12	0.16	1.10	8.46
1981	0.00	0.00	0.05	0.82	0.93	2.00	1.02	0.76	0.67	0.42	0.13	0.20	0.24	0.22	0.99	8.40
1982	0.00	0.04	0.04	0.61	0.48	0.38	0.24	0.61	0.36	0.09	0.26	0.17	0.03	0.03	0.29	3.64
1983	0.00	0.00	0.07	0.53	1.26	1.29	0.54	0.72	0.63	0.48	0.21	0.17	0.08	0.05	0.38	6.41
1984	0.00	0.00	0.10	0.01	0.31	0.78	0.40	0.31	0.20	0.20	0.11	0.17	0.12	0.02	0.27	3.00
1985	0.00	0.00	0.00	0.02	0.46	1.06	1.20	0.91	0.41	0.15	0.15	0.04	0.07	0.03	0.69	5.18
1986	0.00	0.00	0.00	0.00	0.04	0.24	0.53	0.41	0.17	0.19	0.08	0.04	0.06	0.05	0.25	2.07
1987	0.00	0.00	0.00	0.00	0.06	0.11	0.13	0.26	0.19	0.01	0.06	0.02	0.00	0.00	0.16	1.01
1988	0.00	0.02	0.02	0.06	0.00	0.07	0.30	0.38	0.24	0.14	0.09	0.08	0.03	0.00	0.00	1.43
1989	0.00	0.02	0.01	0.04	1.00	0.11	0.07	0.08	0.33	0.08	0.02	0.06	0.06	0.02	0.06	1.95
1990	0.00	0.01	0.00	0.04	0.09	0.32	0.00	0.04	0.01	0.05	0.02	0.01	0.01	0.00	0.03	0.63
1991	0.00	0.04	0.00	0.78	0.11	0.09	0.21	0.03	0.10	0.08	0.14	0.02	0.02	0.00	0.06	1.68
1992	0.00	0.05	0.01	0.19	0.37	0.09	0.11	0.15	0.04	0.15	0.02	0.02	0.05	0.00	0.02	1.26
1993	0.00	0.15	0.11	0.14	0.47	0.32	0.06	0.09	0.00	0.02	0.02	0.00	0.07	0.00	0.04	1.47
1994	0.00	0.11	0.70	0.54	0.64	0.81	0.16	0.03	0.03	0.07	0.01	0.00	0.00	0.02	0.02	3.13
1995	0.00	0.04	0.12	0.58	0.32	0.18	0.31	0.12	0.11	0.04	0.00	0.04	0.03	0.00	0.00	1.88
1996	0.00	0.02	0.04	0.24	0.39	0.35	0.22	0.07	0.00	0.00	0.00	0.03	0.00	0.00	0.00	1.36
1997	0.00	0.07	0.07	0.15	0.69	0.62	0.44	0.08	0.08	0.01	0.00	0.00	0.00	0.00	0.00	2.22
1998	0.00	0.11	1.08	0.71	0.39	0.80	0.71	0.21	0.15	0.08	0.00	0.00	0.00	0.03	0.00	4.27
1999	0.00	0.11	0.38	0.97	0.80	0.48	0.16	0.18	0.03	0.01	0.02	0.00	0.00	0.00	0.00	3.15
2000	0.00	0.01	0.25	1.19	0.69	0.66	0.24	0.25	0.12	0.00	0.04	0.00	0.00	0.00	0.00	3.45
2001	0.00	0.11	0.10	0.71	1.48	1.02	0.40	0.29	0.16	0.11	0.03	0.00	0.00	0.00	0.00	4.41
2002	0.00	0.02	0.06	0.90	2.63	2.26	0.82	0.68	0.35	0.19	0.10	0.01	0.00	0.03	0.04	8.10
2003	0.00	0.00	0.00	0.16	0.78	1.57	1.08	0.81	0.43	0.23	0.05	0.07	0.00	0.02	0.01	5.20

Table 29 continued. Stratified mean number per tow at age of witch flounder in NEFSC bottom trawl spring and autumn surveys (Strata 22-30, 36-40), 1980-2002.

AUTUMN	Age															Total
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14 +	
1980	0.04	0.00	0.02	0.00	0.20	0.26	0.28	0.36	0.17	0.15	0.27	0.04	0.16	0.12	0.57	2.62
1981	0.03	0.07	0.03	0.24	0.44	0.61	0.46	0.27	0.26	0.18	0.21	0.17	0.04	0.13	0.48	3.66
1982	0.02	0.00	0.00	0.06	0.01	0.03	0.08	0.24	0.13	0.01	0.03	0.03	0.01	0.04	0.30	0.99
1983	0.00	0.01	0.01	0.51	1.60	0.76	0.55	0.44	0.08	0.14	0.07	0.11	0.02	0.00	0.41	4.72
1984	0.00	0.00	0.00	0.09	0.94	0.99	0.60	0.53	0.31	0.15	0.13	0.07	0.04	0.13	0.38	4.37
1985	0.00	0.00	0.01	0.06	0.08	0.61	0.68	0.48	0.27	0.10	0.12	0.03	0.01	0.09	0.22	2.76
1986	0.01	0.00	0.00	0.00	0.05	0.27	0.35	0.31	0.16	0.11	0.01	0.01	0.02	0.05	0.24	1.59
1987	0.00	0.00	0.02	0.00	0.01	0.02	0.05	0.19	0.07	0.00	0.01	0.00	0.00	0.02	0.08	0.48
1988	0.00	0.01	0.00	0.72	0.05	0.01	0.04	0.21	0.05	0.05	0.05	0.08	0.01	0.04	0.05	1.38
1989	0.17	0.02	0.02	0.08	0.30	0.01	0.02	0.02	0.08	0.08	0.02	0.00	0.03	0.00	0.04	0.89
1990	0.48	0.09	0.14	0.38	0.51	0.22	0.02	0.02	0.02	0.03	0.00	0.00	0.01	0.05	0.03	2.00
1991	0.22	0.02	0.18	0.66	0.33	0.29	0.14	0.07	0.06	0.03	0.05	0.03	0.00	0.00	0.00	2.08
1992	0.10	0.03	0.11	0.26	0.22	0.05	0.06	0.00	0.00	0.02	0.01	0.02	0.00	0.02	0.04	0.94
1993	2.54	0.67	0.15	0.54	0.78	0.22	0.06	0.02	0.08	0.00	0.02	0.04	0.00	0.01	0.01	5.15
1994	0.43	0.16	0.29	0.53	0.17	0.40	0.04	0.11	0.00	0.04	0.01	0.00	0.01	0.00	0.04	2.21
1995	0.51	0.20	0.76	1.62	0.86	0.47	0.23	0.00	0.00	0.01	0.05	0.00	0.00	0.00	0.01	4.74
1996	0.23	0.09	0.26	0.79	1.99	1.39	0.44	0.07	0.06	0.04	0.00	0.03	0.00	0.00	0.00	5.38
1997	0.89	0.34	0.98	0.52	0.87	0.77	0.38	0.33	0.00	0.00	0.00	0.00	0.02	0.00	0.00	5.10
1998	0.64	0.08	0.52	1.36	0.47	0.30	0.17	0.11	0.04	0.01	0.00	0.00	0.00	0.00	0.00	3.70
1999	0.32	0.52	1.18	1.51	1.04	0.60	0.36	0.28	0.05	0.04	0.01	0.00	0.00	0.00	0.00	5.91
2000	0.94	0.10	0.72	1.41	1.75	0.67	0.59	0.23	0.15	0.05	0.00	0.00	0.03	0.00	0.00	6.63
2001	0.00	0.04	0.21	0.95	3.16	1.89	0.81	0.61	0.16	0.06	0.06	0.00	0.00	0.00	0.00	7.94
2002	0.00	0.00	0.27	0.43	1.48	1.00	0.53	0.33	0.15	0.07	0.00	0.05	0.01	0.00	0.00	4.31

Table 30. Witch flounder mean length (cm) at age in spring and autumn NEFSC bottom trawl surveys (Strata 22-30, 36-40), 1980-2002, preliminary 2003.

SPRING	Age														
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14 +
1980	-	9.7	16.4	20.6	26.2	30.6	34.8	38.6	40.6	45.0	48.6	49.2	49.3	52.5	55.2
1981	-	-	13.4	20.2	28.5	32.4	35.4	39.7	44.4	49.4	52.4	49.9	54.5	54.1	57.6
1982	-	7.9	15.0	20.7	27.2	32.8	36.1	41.0	44.1	48.2	50.7	51.3	58.0	53.0	58.1
1983	-	-	17.9	20.9	26.5	31.2	35.5	40.2	43.7	47.4	52.5	54.9	50.0	55.6	56.1
1984	-	-	17.4	19.0	29.4	32.6	37.5	42.2	43.0	46.0	51.3	50.6	54.2	54.3	57.4
1985	-	-	-	19.5	28.8	33.5	36.6	41.2	44.4	46.9	49.3	49.3	48.5	55.0	56.3
1986	-	-	-	-	27.3	35.0	38.5	41.7	45.8	49.1	51.8	52.3	54.2	56.1	57.9
1987	-	-	-	-	28.0	34.4	40.9	40.8	44.1	46.0	51.5	48.0	-	-	56.7
1988	-	9.0	15.0	19.5	-	33.1	39.2	43.0	46.0	50.2	54.2	51.2	58.3	-	-
1989	-	7.0	15.0	20.6	28.5	33.0	39.8	44.0	44.9	50.5	50.2	53.1	58.3	47.0	60.7
1990	-	9.0	-	19.8	28.3	32.4	-	40.9	49.0	49.8	51.5	52.0	53.0	-	54.9
1991	-	7.5	-	20.4	27.4	35.3	37.2	43.1	48.2	48.2	52.7	53.0	54.0	-	52.4
1992	-	8.5	11.0	21.7	29.3	35.0	38.4	42.5	45.7	49.3	46.0	51.0	57.3	-	56.0
1993	-	7.9	17.9	23.5	30.0	34.5	38.1	40.5	-	50.0	50.0	-	50.3	-	60.1
1994	-	10.8	17.9	21.5	29.4	33.7	38.1	41.2	46.0	48.2	48.0	-	-	57.0	57.0
1995	-	9.7	17.3	22.4	27.1	34.3	37.2	43.7	45.7	50.3	-	54.0	58.7	-	-
1996	-	9.4	19.6	22.3	28.1	32.5	37.1	40.6	0.0	-	-	56.3	-	-	-
1997	-	10.8	15.9	22.2	29.5	31.4	36.3	42.3	43.9	48.0	-	-	-	-	-
1998	-	11.0	20.3	24.6	29.9	33.3	35.7	39.1	42.6	45.2	-	-	-	51.7	-
1999	-	10.0	19.5	25.0	28.5	34.0	37.2	40.6	45.5	44.0	50.0	-	-	-	-
2000	-	9.0	18.1	20.9	27.3	31.2	36.5	38.5	41.1	-	50.3	-	-	-	-
2001	-	7.4	15.9	23.5	27.0	32.4	36.0	38.2	40.8	41.5	45.3	-	-	-	-
2002	-	11.0	18.0	20.2	29.0	32.2	34.7	37.8	42.1	44.3	45.1	42.0	-	45.3	54.8
2003	-	-	-	22.4	27.7	31.9	34.9	38.1	40.8	42.7	43.6	45.4	-	47.5	49.5

Table 30 continued. Witch flounder mean length (cm) at age in spring and autumn NEFSC bottom trawl surveys (Strata 22-30, 36-40), 1980-2002.

AUTUMN	Age														
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+
1980	5.5	-	19.5	-	27.3	32.0	34.9	39.1	43.3	47.7	48.8	50.1	51.6	53.7	56.7
1981	5.5	12.6	17.4	23.3	30.6	33.1	38.3	41.4	44.8	47.0	51.4	53.6	52.7	55.0	56.4
1982	5.0	0.0	0.0	22.0	28.5	29.2	36.4	41.9	42.8	47.3	50.2	49.1	51.0	52.0	56.8
1983	0.0	14.0	19.0	24.7	30.1	34.7	39.2	42.2	45.0	48.5	52.0	50.9	51.0	0.0	58.8
1984	0.0	0.0	0.0	25.1	30.7	34.2	38.2	42.8	45.2	46.6	50.2	51.3	54.8	53.5	58.6
1985	0.0	0.0	20.0	26.0	29.7	34.6	38.8	42.7	46.8	49.1	50.8	53.3	55.0	53.1	58.7
1986	6.0	0.0	0.0	0.0	29.7	35.5	38.2	42.4	45.4	49.4	51.0	51.3	49.0	53.9	57.7
1987	0.0	0.0	16.0	0.0	28.0	36.0	39.1	41.6	43.8	0.0	49.0	0.0	0.0	55.0	60.9
1988	0.0	10.0	0.0	25.4	31.5	38.0	42.0	42.9	45.6	48.6	50.0	54.3	56.0	55.1	56.8
1989	5.7	15.0	18.5	24.1	31.1	36.0	44.0	45.0	46.7	49.9	52.0	0.0	54.8	0.0	64.2
1990	6.2	16.3	17.6	26.7	29.8	36.2	40.0	43.0	47.0	50.9	0.0	0.0	58.0	51.2	58.9
1991	5.7	14.9	20.8	26.3	30.5	36.8	41.9	46.4	47.6	46.6	53.5	55.0	0.0	0.0	0.0
1992	5.9	15.8	23.1	27.7	32.1	37.7	37.9	0.0	0.0	46.0	50.0	47.0	0.0	49.0	56.5
1993	5.6	14.2	22.2	28.8	32.2	36.4	42.3	43.6	46.2	0.0	55.0	51.0	0.0	63.0	57.0
1994	5.7	16.0	20.9	23.5	32.7	36.6	43.5	44.0	0.0	54.2	50.0	0.0	51.0	0.0	57.2
1995	6.6	16.7	22.0	26.5	29.9	35.5	39.2	0.0	0.0	54.0	50.6	0.0	0.0	0.0	56.0
1996	5.5	14.2	18.7	25.0	29.6	33.8	39.6	42.1	47.4	50.5	0.0	53.0	0.0	0.0	0.0
1997	6.3	16.3	19.8	25.8	30.6	35.1	38.0	42.0	0.0	0.0	0.0	0.0	47.1	0.0	0.0
1998	5.9	15.7	22.0	25.7	30.7	35.0	39.3	41.9	44.9	43.0	0.0	0.0	0.0	0.0	0.0
1999	5.5	15.6	20.4	26.1	30.2	34.6	38.2	41.6	41.6	45.1	45.0	0.0	0.0	0.0	0.0
2000	6.1	15.6	22.6	26.3	31.0	33.4	38.0	42.3	44.1	46.3	0.0	0.0	51.0	0.0	0.0
2001	0.0	14.6	18.9	28.0	30.6	33.1	36.7	39.9	41.9	41.0	46.4	0.0	0.0	0.0	0.0
2002	0.0	0.0	22.7	26.0	31.3	34.3	37.2	41.7	41.9	44.6	0.0	50.5	51.0	0.0	0.0

Table 31. Mean weight (kg) at age of witch flounder from the Gulf of Maine-Georges Bank region, derived from NEFSC spring and autumn bottom trawl survey, (strata 22-30, 36-40), 1982 - 2002, and preliminary spring 2003.

SPRING	Age															
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14+	14+
1982	0.0000	0.0018	0.0155	0.0479	0.1220	0.2350	0.3174	0.4939	0.6395	0.8643	1.0226	1.0519	1.5952	1.1711	1.6152	
1983	0.0000	0.0000	0.0291	0.0494	0.1069	0.1861	0.2878	0.4433	0.5885	0.7790	1.0889	1.2747	0.9215	1.3281	1.4180	
1984	0.0000	0.0000	0.0261	0.0342	0.1540	0.2234	0.3539	0.5318	0.5647	0.7143	1.0398	0.9979	1.2606	1.2534	1.5434	
1985	0.0000	0.0000	0.0000	0.0368	0.1424	0.2349	0.3247	0.4841	0.6268	0.7516	0.8984	0.8858	0.8318	1.2805	1.4087	
1986	0.0000	0.0000	0.0000	0.0000	0.1115	0.2604	0.3592	0.4737	0.6481	0.8299	0.9880	1.0206	1.1601	1.3053	1.4627	
1987	0.0000	0.0000	0.0000	0.0000	0.1278	0.2542	0.4637	0.4615	0.6012	0.6891	1.0154	0.7974	0.0000	0.0000	1.4347	
1988	0.0000	0.0025	0.0142	0.0364	0.0000	0.2177	0.3875	0.5340	0.6686	0.8996	1.1689	0.9765	1.5001	0.0000	0.0000	
1989	0.0000	0.0010	0.0139	0.0408	0.1255	0.2039	0.3869	0.5537	0.5946	0.8807	0.8600	1.0413	1.4373	0.6851	1.6452	
1990	0.0000	0.0026	0.0000	0.0392	0.1328	0.2150	0.0000	0.4708	0.8717	0.9235	1.0370	1.0687	1.1408	0.0000	1.2886	
1991	0.0000	0.0014	0.0000	0.0429	0.1244	0.2784	0.3348	0.5487	0.8060	0.8278	1.0986	1.1110	1.1845	0.0000	1.0707	
1992	0.0000	0.0024	0.0050	0.0519	0.1449	0.2641	0.3635	0.5224	0.6620	0.8765	0.6694	0.9535	1.4308	0.0000	1.3139	
1993	0.0000	0.0016	0.0256	0.0661	0.1507	0.2395	0.3347	0.4116	0.0000	0.8459	0.8459	0.0000	0.8667	0.0000	1.6168	
1994	0.0000	0.0051	0.0274	0.0524	0.1481	0.2352	0.3566	0.4681	0.6750	0.7940	0.7810	0.0000	0.0000	1.4078	1.4078	
1995	0.0000	0.0032	0.0233	0.0572	0.1089	0.2378	0.3138	0.5452	0.6325	0.8816	0.0000	1.1302	1.5043	0.0000	0.0000	
1996	0.0000	0.0037	0.0351	0.0563	0.1252	0.2024	0.3170	0.4300	0.0000	0.0000	0.0000	1.3229	0.0000	0.0000	0.0000	
1997	0.0000	0.0046	0.0183	0.0532	0.1400	0.1733	0.2840	0.4763	0.5430	0.7288	0.0000	0.0000	0.0000	0.0000	0.0000	
1998	0.0000	0.0053	0.0403	0.0777	0.1507	0.2179	0.2770	0.3836	0.5071	0.6355	0.0000	0.0000	0.0000	0.9754	0.0000	
1999	0.0000	0.0040	0.0347	0.0816	0.1278	0.2311	0.3126	0.4251	0.6275	0.5562	0.8621	0.0000	0.0000	0.0000	0.0000	
2000	0.0000	0.0025	0.0284	0.0464	0.1155	0.1806	0.3075	0.3789	0.4746	0.0000	0.9379	0.0000	0.0000	0.0000	0.0000	
2001	0.0000	0.0015	0.0199	0.0654	0.1052	0.1957	0.2810	0.3412	0.4287	0.4549	0.6114	0.0000	0.0000	0.0000	0.0000	
2002	0.0000	0.0046	0.0255	0.0388	0.1307	0.1867	0.2401	0.3232	0.4634	0.5493	0.5910	0.4552	0.0000	0.5938	1.1331	
2003	0.0000	0.0000	0.0000	0.0588	0.1141	0.1833	0.2461	0.3324	0.4183	0.4860	0.5191	0.5881	0.0000	0.6712	0.7743	

Table 31 continued. Mean weight (kg) at age of witch flounder from the Gulf of Maine-Georges Bank region , derived from NEFSC spring and autumn bottom trawl survey, (strata 22-30, 36-40), 1982 - 2002.

AUTUMN	AGE														
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14 +
1982	0.0004	0.0000	0.0000	0.0604	0.1486	0.1591	0.3462	0.5470	0.5905	0.8193	1.0054	0.9316	1.0589	1.1331	1.5423
1983	0.0000	0.0123	0.0350	0.0883	0.1728	0.2794	0.4271	0.5435	0.6757	0.8757	1.1119	1.0501	1.0340	0.0000	1.7009
1984	0.0000	0.0000	0.0000	0.0905	0.1817	0.2639	0.3836	0.5656	0.6852	0.7623	0.9759	1.0446	1.3091	1.2035	1.6588
1985	0.0000	0.0000	0.0391	0.0969	0.1536	0.2592	0.3875	0.5341	0.7256	0.8551	0.9579	1.1245	1.2535	1.1205	1.5848
1986	0.0007	0.0000	0.0000	0.0000	0.1614	0.3015	0.3832	0.5534	0.6969	0.9241	1.0232	1.0493	0.8921	1.2559	1.5656
1987	0.0000	0.0000	0.0195	0.0000	0.1327	0.3141	0.4168	0.5181	0.6258	0.0000	0.9040	0.0000	0.0000	1.3432	1.9129
1988	0.0000	0.0036	0.0000	0.0900	0.1876	0.3540	0.5152	0.5472	0.6645	0.8253	0.9117	1.2061	1.3376	1.2693	1.4267
1989	0.0007	0.0155	0.0326	0.0791	0.1893	0.3111	0.6190	0.6686	0.7652	0.9710	1.0973	0.0000	1.3101	0.0000	2.2664
1990	0.0008	0.0210	0.0275	0.1135	0.1642	0.3173	0.4439	0.5688	0.7716	1.0181	0.0000	0.0000	1.5867	1.0437	1.7592
1991	0.0006	0.0155	0.0502	0.1083	0.1791	0.3388	0.5257	0.7467	0.8148	0.7657	1.2266	1.3420	0.0000	0.0000	0.0000
1992	0.0006	0.0181	0.0651	0.1236	0.2044	0.3524	0.3850	0.0000	0.0000	0.6915	0.9203	0.7444	0.0000	0.8587	1.3961
1993	0.0006	0.0133	0.0591	0.1416	0.2104	0.3147	0.5233	0.5872	0.7075	0.0000	1.2822	0.9898	0.0000	2.0425	1.4493
1994	0.0006	0.0187	0.0459	0.0707	0.2116	0.3133	0.5605	0.5893	0.0000	1.1912	0.9017	0.0000	0.9650	0.0000	1.4443
1995	0.0008	0.0207	0.0533	0.1000	0.1503	0.2703	0.3794	0.0000	0.0000	1.1259	0.9099	0.0000	0.0000	0.0000	1.2754
1996	0.0005	0.0124	0.0315	0.0828	0.1488	0.2328	0.4005	0.4964	0.7359	0.9408	0.0000	1.0769	0.0000	0.0000	0.0000
1997	0.0008	0.0195	0.0402	0.0950	0.1704	0.2700	0.3529	0.5031	0.0000	0.0000	0.0000	0.0000	0.0000	0.7415	0.0000
1998	0.0006	0.0185	0.0552	0.0928	0.1695	0.2669	0.3936	0.4937	0.6344	0.5333	0.0000	0.0000	0.0000	0.0000	0.0000
1999	0.0005	0.0176	0.0452	0.1021	0.1671	0.2672	0.3707	0.4932	0.4978	0.6581	0.6407	0.0000	0.0000	0.0000	0.0000
2000	0.0007	0.0163	0.0591	0.0993	0.1743	0.2248	0.3510	0.5116	0.5813	0.6765	0.0000	0.0000	0.9435	0.0000	0.0000
2001	0.0000	0.0131	0.0347	0.1227	0.1673	0.2179	0.3122	0.4124	0.4878	0.4468	0.6933	0.0000	0.0000	0.0000	0.0000
2002	0.0000	0.0000	0.0587	0.0959	0.1831	0.2442	0.3225	0.4775	0.4920	0.5955	0.0000	0.9261	0.9391	0.0000	0.0000

Table 32. Proportion mature at age for female witch flounder derived from logistic regression analysis using a 5-year moving average of NEFSC spring bottom trawl surveys, 1980 - 2003.

Age	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
1	0.00	0.00	0.00	0.00	0.00	0.01	0.04	0.01	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01
2	0.00	0.00	0.00	0.00	0.00	0.03	0.13	0.03	0.03	0.03	0.02	0.01	0.01	0.01	0.00	0.00	0.01	0.01	0.02	0.03
3	0.01	0.01	0.02	0.01	0.02	0.12	0.32	0.12	0.08	0.07	0.06	0.04	0.04	0.03	0.02	0.01	0.04	0.05	0.06	0.07
4	0.04	0.06	0.07	0.05	0.09	0.37	0.60	0.36	0.22	0.18	0.18	0.14	0.13	0.13	0.13	0.07	0.14	0.15	0.15	0.16
5	0.14	0.20	0.24	0.24	0.37	0.71	0.83	0.70	0.47	0.39	0.42	0.35	0.34	0.42	0.45	0.37	0.39	0.37	0.33	0.35
6	0.40	0.51	0.59	0.68	0.76	0.91	0.94	0.91	0.74	0.64	0.70	0.65	0.64	0.78	0.82	0.81	0.71	0.67	0.59	0.59
7	0.73	0.82	0.86	0.93	0.95	0.98	0.98	0.98	0.90	0.84	0.88	0.86	0.86	0.95	0.96	0.97	0.91	0.87	0.81	0.79
8	0.92	0.95	0.97	0.99	0.99	0.99	0.99	0.99	0.97	0.94	0.96	0.95	0.95	0.99	0.99	1.00	0.97	0.96	0.93	0.91
9	0.98	0.99	0.99	1.00	1.00	1.00	1.00	1.00	0.99	0.98	0.99	0.99	0.99	1.00	1.00	1.00	0.99	0.99	0.97	0.97
10	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.99	0.99
11+	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

Number of fish

mature	108	153	204	203	186	177	145	111	104	89	80	85	76	81	132	123	127	148	186	191
immature	98	112	115	102	47	24	11	37	60	88	107	139	139	139	190	198	205	247	301	248
total	206	265	319	305	233	201	156	148	164	177	187	224	215	220	322	321	332	395	487	439

Long-term (non-averaged) proportion mature at age for female witch flounder from NEFSC spring bottom trawl surveys, 1980-2003.

Age	1980- 2003
1	0.00
2	0.00
3	0.01
4	0.04
5	0.14
6	0.40
7	0.73
8	0.92
9	0.98
10	0.99
11+	1.00

Table 33. Stratified mean weight (kg) per tow of mature witch flounder (spawning stock biomass) in the NEFSC spring bottom trawl survey in Gulf of Maine-Georges Bank region (strata 22-30, 36-40), 1968-2002.

Year	Mature weight (kg) per tow
1968	2.930
1969	2.300
1970	4.073
1971	1.907
1972	3.772
1973	5.868
1974	3.289
1975	2.499
1976	3.248
1977	1.522
1978	2.278
1979	1.480
1980	2.964
1981	3.104
1982	1.519
1983	2.166
1984	1.383
1985	2.607
1986	1.329
1987	0.638
1988	0.836
1989	0.637
1990	0.200
1991	0.455
1992	0.356
1993	0.186
1994	0.323
1995	0.377
1996	0.174
1997	0.249
1998	0.498
1999	0.251
2000	0.228
2001	0.331
2002	0.782
2003	0.751

Note: 1977-1982, 1983-1984, 1985-1990, 1991-1993, 1994-1999, 2000-2002 ogives were used; No maturity at length data before 1977; the 1977-1982 period was applied to the 1963-1976 period.

Table 34. Estimates of instantaneous total mortality (Z) for witch flounder in the Gulf of Maine-Georges Bank region, 1980-2002, derived from NEFSC spring and autumn bottom trawl survey data.

YEAR	3+	4+	5+	6+	7+	8+				
Spring							Geometric			
							Times period	Spring	Autumn	mean
1980	8.18	7.23	5.71	4.99	3.79	2.77				
1981	8.31	7.49	6.56	4.56	3.54	2.78				
1982	3.56	2.95	2.46	2.09	1.85	1.24	1982-1985	0.48	0.23	0.34
1983	6.34	5.80	4.54	3.25	2.71	1.99	1986-1989	0.77	0.65	0.71
1984	2.90	2.89	2.58	1.80	1.40	1.09	1990-1993	0.59	0.61	0.60
1985	5.18	5.17	4.71	3.65	2.45	1.54	1994-1997	0.53	0.36	0.44
1986	2.07	2.07	2.03	1.79	1.26	0.85	1998-2001	0.52	0.60	0.56
1987	1.01	1.01	0.95	0.83	0.70	0.44				
1988	1.39	1.33	1.33	1.25	0.95	0.57				
1989	1.91	1.87	0.87	0.76	0.69	0.61				
1990	0.62	0.58	0.49	0.17	0.17	0.13				
1991	1.64	0.86	0.75	0.67	0.46	0.42				
1992	1.20	1.01	0.64	0.55	0.44	0.29				
1993	1.21	1.07	0.60	0.28	0.22	0.14				
1994	2.32	1.78	1.14	0.33	0.17	0.14				
1995	1.72	1.14	0.83	0.65	0.33	0.22				
1996	1.31	1.06	0.67	0.32	0.10	0.03				
1997	2.08	1.93	1.24	0.62	0.18	0.10				
1998	3.08	2.37	1.98	1.19	0.47	0.26				
1999	2.67	1.69	0.90	0.41	0.25	0.07				
2000	3.19	2.00	1.30	0.64	0.40	0.15				
2001	4.21	3.49	2.02	1.00	0.60	0.30				
2002	8.02	7.12	4.49	2.23	1.41	0.73				
Autumn										
1980	2.58	2.58	2.38	2.12	1.84	1.48				
1981	3.49	3.25	2.81	2.20	1.74	1.47				
1982	0.97	0.91	0.90	0.87	0.80	0.56				
1983	4.70	4.19	2.60	1.84	1.29	0.85				
1984	4.37	4.28	3.34	2.34	1.74	1.21				
1985	2.75	2.70	2.62	2.01	1.33	0.84				
1986	1.58	1.58	1.53	1.26	0.91	0.60				
1987	0.46	0.46	0.45	0.43	0.38	0.19				
1988	1.37	0.64	0.59	0.58	0.54	0.33				
1989	0.68	0.60	0.30	0.29	0.27	0.25				
1990	1.30	0.92	0.41	0.19	0.17	0.15				
1991	1.66	1.00	0.67	0.38	0.24	0.17				
1992	0.70	0.45	0.22	0.17	0.11	0.11				
1993	1.79	1.24	0.47	0.25	0.19	0.17				
1994	1.33	0.80	0.64	0.24	0.21	0.10				
1995	3.26	1.63	0.78	0.30	0.07	0.07				
1996	4.80	4.02	2.03	0.64	0.20	0.14				
1997	2.89	2.37	1.50	0.73	0.35	0.02				
1998	2.46	1.10	0.63	0.33	0.17	0.05				
1999	3.89	2.38	1.34	0.74	0.37	0.10				
2000	4.87	3.47	1.72	1.05	0.46	0.23				
2001	7.69	6.74	3.58	1.70	0.88	0.27				
2002	4.04	3.61	2.13	1.13	0.60	0.27				

Table 35. Parameter estimates (with associated statistics) and estimates of terminal F from alternative ADAPT VPA formulations; stock sizes in '000s.

	Run 61-f	Run 61-f	Run 100	Run 200	Run 201	Run 300	Run 301-f	Run 301f-spr	Run 301f-aut
Software	FACT 1.5	NFTv2011	NFTv2011	NFTv2011	NFTv2011	NFTv2011	NFTv2011	NFTv2011	NFTv2011
CAA	1982-2001 1-11+	1982-2001 1-11+	1982-2002 1-11+	1982-2002 1-11+	1982-2002 1-11+	1982-2002 1-11+	1982-2002 3-11+	1982-2002 3-11+	1982-2002 3-11+
Est.Ages	4-10	4-10	4 - 10	4 - 10	3-10	3-10	3-10	3-10	3-10
NMFS-s	3-11+	3-11+	3 - 11+	3 - 11+	3-11+	3-11+	3-11+	3-11+	-
NMFS-a	3-11+	3-11+	3 - 11+	3 - 11+	3-11+	3-11+	3-11+	-	3-11+
Notes:	GARM VPA	Re-Run of GARM VPA	LM discards (sf method)	LM discards (sf & FOP)	LM discards (sf & FOP)	LM discards (sf & FOP)	LM discards (sf and FOP)	LM discards (sf and FOP)	LM discards (sf and FOP)
M.S.R.	.7673	.7289	.7469	.7485	.7459	.792	.791	.603	1.00
N3 (cv)					19.621 (.63)	19.703 (.64)	19.759 (.64)	11.429 (.79)	34.449 (1.03)
N4 (cv)	5.76e4 (.45)	57.321 (.44)	25.111 (.44)	24.755 (.44)	25.291 (.44)	25.373 (.45)	25.441 (.45)	36.289 (.56)	17.944 (.73)
N5 (cv)	6.08e4 (.38)	60.440 (.36)	41.902 (.37)	41.897 (.37)	42.456 (.37)	42.628 (.37)	42.739 (.37)	94.857 (.46)	40.928 (.60)
N6 (cv)	2.91e4 (.34)	28.936 (.33)	40.091 (.33)	41.003 (.32)	41.370 (.32)	41.550 (.33)	41.657 (.33)	36.267 (.41)	48.035 (.52)
N7 (cv)	1.67e4 (.32)	16.563 (.31)	19.965 (.31)	20.937 (.30)	21.063 (.30)	21.145 (.31)	21.203 (.31)	16.900 (.38)	26.672 (.48)
N8 (cv)	4.73e3 (.37)	3.844 (.35)	11.861 (.29)	12.420 (.28)	12.261 (.28)	10.122 (.32)	10.370 (.32)	8.067 (.41)	13.403 (.49)
N9 (cv)	1.56e3 (.44)	1.744 (.42)	2.547 (.38)	2.511 (.38)	2.283 (.40)	3.842 (.34)	3.903 (.33)	3.323 (.43)	4.742 (.51)
N10 (cv)	1.06e3 (.44)	1.197 (.42)	1.017 (.43)	972 (.44)	809 (.47)	754 (.47)	791 (.45)	795 (.57)	787 (.72)
Age 3 in T+1	22.643	23.362	6.268	8.026	19.620	19.707	19.760	11.429	34.449
F 1	0.00	0.0004	0.0000	0.0003	0.0004	0.0003	-	-	-
F 2	0.00	0.0026	0.0020	0.0002	0.0008	0.008	-	-	-
F 3	0.00	0.0026	0.0017	0.0015	0.0015	0.0015	0.0015	0.0010	0.0021
F 4	0.02	0.0205	0.0347	0.0164	0.0162	0.0162	0.0161	0.0154	0.0168
F 5	0.07	0.0725	0.0513	0.0273	0.0270	0.0269	0.0268	0.0308	0.0233
F 6	0.07	0.0725	0.0688	0.0603	0.0599	0.0597	0.0595	0.0741	0.0476
F 7	0.28	0.3375	0.1575	0.1474	0.1492	0.1780	0.1741	0.2186	0.1373
F 8	0.62	0.5716	0.3795	0.3857	0.4169	0.2685	0.2648	0.3045	0.2229
F 9	0.44	0.3985	0.4527	0.4706	0.5435	0.5734	0.5526	0.5508	0.5550

	Run 61-f	Run 61-f	Run 100	Run 200	Run 201	Run 300	Run 301-f	Run 301f-spr	Run 301f-aut
F10	0.45	0.4358	0.3299	0.3346	0.3699	0.3339	0.4087	0.4277	0.03890
F11+	0.45	0.4358	0.3299	0.3346	0.3699	0.3339	0.4087	0.4277	0.3890
Ave F 7-9	0.45	0.4358	0.3299	0.3346	0.3699	-	-	-	-
Ave F 8-9						0.3339	0.4087	0.4277	0.3890
SSB ('000 mt)	11,368	11,267	15,331	16,548	16,242	16,212	18,296	15,798	21,569

Table 36. Estimates of beginning year stock size (thousands of fish), instantaneous fishing mortality and spawning stock biomass (mt) for witch flounder estimated from virtual population analysis, 1982-2002.

Stock Numbers (Jan 1) in thousands

Age	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992
3	15404	17700	16340	7650	5414	3001	9740	6214	6670	8831	14342
4	12174	13082	14922	13928	6470	4639	2562	7827	4934	5422	7192
5	9563	9493	10014	11486	10899	5220	3825	2076	6333	3660	4291
6	7830	7114	6765	6768	7929	7978	4060	3047	1726	4431	2353
7	4289	5375	4668	4216	4039	4267	5683	2886	2332	1249	3281
8	2752	3077	3160	2647	2224	2034	2222	3615	1783	1754	857
9	2102	1763	1746	1344	1132	1146	949	853	2297	1100	1283
10	1101	1440	839	862	599	594	544	448	413	1668	677
11+	7260	4728	3844	2926	2039	1151	1218	1263	900	1371	2127
Total	62476	63772	62297	51828	40746	30029	30804	28229	27388	29487	36404

Age	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
3	9670	13610	12671	15878	20226	29659	42904	67651	58704	29603	19760
4	11974	7936	11675	10324	13554	17283	25221	36778	58131	50462	25441
5	5029	8635	5904	9706	8400	10686	14360	21220	31390	49716	42739
6	2900	3019	4887	4119	7237	6290	8213	11345	17842	26146	41658
7	1164	1649	1414	2627	2289	5033	4106	5717	8693	14340	21204
8	2162	453	660	438	949	1033	2846	2426	3417	5910	10370
9	552	1320	209	321	136	274	547	1743	1138	1599	3903
10	940	274	640	91	80	39	105	239	926	398	792
11+	1318	782	373	180	112	179	181	536	667	655	603
Total	35708	37679	38434	43683	52982	70476	98483	147655	180909	178829	166470

Table 36. continued.

Fishing Mortality

Age	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
3	0.013	0.021	0.010	0.018	0.005	0.008	0.069	0.081	0.057	0.055
4	0.099	0.117	0.112	0.095	0.065	0.043	0.061	0.062	0.149	0.084
5	0.146	0.189	0.242	0.221	0.162	0.101	0.077	0.034	0.207	0.292
6	0.226	0.271	0.323	0.366	0.470	0.189	0.191	0.118	0.173	0.150
7	0.182	0.381	0.417	0.490	0.536	0.503	0.303	0.332	0.135	0.227
8	0.295	0.416	0.705	0.700	0.514	0.612	0.807	0.304	0.333	0.162
9	0.228	0.593	0.556	0.657	0.495	0.595	0.602	0.577	0.170	0.335
10	0.266	0.477	0.650	0.685	0.507	0.606	0.741	0.350	0.238	0.226
11+	0.266	0.477	0.650	0.685	0.507	0.606	0.741	0.350	0.238	0.226

Age	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
3	0.030	0.048	0.003	0.055	0.008	0.007	0.012	0.004	0.002	0.001	0.002
4	0.208	0.177	0.146	0.035	0.056	0.088	0.035	0.023	0.008	0.006	0.016
5	0.242	0.360	0.419	0.210	0.144	0.139	0.113	0.086	0.023	0.033	0.027
6	0.554	0.414	0.609	0.471	0.438	0.213	0.276	0.212	0.116	0.069	0.060
7	0.267	0.793	0.766	1.023	0.868	0.646	0.420	0.376	0.365	0.236	0.174
8	0.291	0.343	0.623	0.571	1.021	1.093	0.486	0.340	0.607	0.610	0.265
9	0.161	0.552	0.574	0.688	1.244	1.085	0.805	0.679	0.483	0.900	0.553
10	0.211	0.382	0.586	0.598	1.109	1.092	0.545	0.388	0.553	0.675	0.409
11+	0.211	0.382	0.586	0.598	1.109	1.092	0.545	0.388	0.553	0.675	0.409

Table 36 continued.

Spawning Stock Biomass (mt)

Age	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991
3	20	13	24	5	5	16	115	33	24	25
4	107	132	185	127	91	216	191	238	156	104
5	376	458	580	684	1019	880	745	360	587	336
6	1115	1241	1243	1584	1916	2471	1297	964	454	886
7	1543	1883	1715	1720	1636	1794	2494	1303	999	496
8	1634	1544	1559	1388	1218	1072	1148	2010	1003	1004
9	1632	1172	1088	878	780	757	628	559	1640	754
10	949	1207	628	665	487	479	422	367	358	1411
11+	9521	5781	4505	3374	2414	1323	1392	1577	1227	1829
Total	16897	13431	11528	10425	9567	9008	8433	7410	6447	6844

Age	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
3	50	21	23	11	12	9	56	85	158	186	102
4	145	229	154	155	113	97	251	453	618	975	1096
5	463	437	636	601	852	599	883	1114	1657	2384	3739
6	582	694	630	1200	1139	1740	1304	1671	2095	3192	4736
7	1369	424	585	542	1044	888	1782	1405	1800	2664	4443
8	500	1210	230	352	225	455	485	1343	1040	1361	2545
9	916	392	862	144	196	87	171	286	925	544	825
10	547	789	217	509	69	61	32	86	137	511	262
11+	2490	1610	875	409	180	118	192	145	441	489	548
Total	7063	5806	4212	3922	3828	4053	5156	6587	8871	12305	18296

Table 37. Summary of witch flounder catch (mt), spawning stock biomass (mt), fully recruited fishing mortality, recruitment (age 3, thousands of fish) and the year class.

Year	Catch (mt)	SSB (mt)	Ave F 8-9	Recruits Age 3 (‘000)	Year class
1982	4954	16897	0.262	15.404	1979
1983	6159	13431	0.505	17.700	1980
1984	6759	11527	0.631	16.340	1981
1985	6192	10425	0.679	7.650	1982
1986	4636	9566	0.504	5.414	1983
1987	3494	9008	0.604	3.001	1984
1988	3320	8432	0.704	9.740	1985
1989	2199	7411	0.440	6.214	1986
1990	1645	6448	0.252	6.670	1987
1991	1870	6845	0.249	8.831	1988
1992	2395	7062	0.226	14.342	1989
1993	2973	5806	0.447	9.670	1990
1994	3073	4212	0.598	13.610	1991
1995	2386	3923	0.629	12.671	1992
1996	2338	3830	1.132	15.878	1993
1997	2065	4054	1.089	20.226	1994
1998	2124	5156	0.645	29.659	1995
1999	2327	6588	0.510	42.904	1996
2000	2551	8871	0.545	67.651	1997
2001	3241	12306	0.755	58.704	1998
2002	3466	18296	0.409	29.603	1999
2003				19.760	2000
min	1645	3830	0.226	3.001	
max	6759	18296	1.132	67.651	
mean	3341	8576	0.563	19.613	
geomean				14.448	
median				14.342	

Table 38. Yield and Spawning Stock biomass per recruit results for witch flounder.

Yield per Recruit and Spawning Stock Biomass per Recruit
 ## YPR Version 2.0
 ## Date of Run: 19 May 2003 10:59
 ## Input Data File: H:\WITCHASS\YPR\YPR2003\RUN301-F.DAT

Model Title: Witch Flounder (run 301)
 Start Age = 3
 End Age = 20 (Does Not Include Plus Group)
 Fishing Mortality Upper Bound = 1.0000
 Fishing Mortality Calculation Increment = 0.0001
 Fishing Mortality Printing Increment = 0.05
 Natural Mortality = 0.1500
 Proportion Fishing Mortality Before Spawning = 0.1667
 Proportion Natural Mortality Before Spawning = 0.1667

Age	Selectivity F	Selectivity M	Stock Weight	Catch Weight	Maturity
3	0.0036	1.0000	0.0787	0.0830	0.0200
4	0.0229	1.0000	0.1459	0.2021	0.0800
5	0.0703	1.0000	0.2319	0.2746	0.3000
6	0.1931	1.0000	0.3328	0.3813	0.6200
7	0.5282	1.0000	0.4442	0.4752	0.8700
8	1.0000	1.0000	0.5615	0.5548	0.9700
9	1.0000	1.0000	0.6816	0.6393	1.0000
10	1.0000	1.0000	0.8006	0.7656	1.0000
11	1.0000	1.0000	0.9175	0.9175	1.0000
12	1.0000	1.0000	1.0399	1.0399	1.0000
13	1.0000	1.0000	1.1348	1.1348	1.0000
14	1.0000	1.0000	1.2335	1.2335	1.0000
15	1.0000	1.0000	1.3259	1.3259	1.0000
16	1.0000	1.0000	1.4097	1.4097	1.0000
17	1.0000	1.0000	1.4875	1.4875	1.0000
18	1.0000	1.0000	1.5575	1.5575	1.0000
19	1.0000	1.0000	1.6215	1.6215	1.0000
20	1.0000	1.0000	1.6787	1.6787	1.0000

Reference Point	F	YPR	SSBR	Mean Age	Mean GT	Exp Spawn
F Zero	0.00000	0.00000	3.22009	7.88231	12.35089	3.51035
F-01	0.19560	0.21504	1.42574	6.17333	9.86110	1.89110
F-Max	0.54470	0.23913	0.70920	5.25697	7.88807	1.10782
F at 40 %MSP	0.23030	0.22321	1.28817	6.01053	9.54415	1.75334

FMORT	CTHN	CTHW	STKN	STKW	SPNSTKN	SPNSTKW	MSP	MNAGE	MNGT	EXSP
0.00000	0.00000	0.00000	6.69668	3.73482	3.97773	3.22009	100.00000	7.88231	12.35089	3.51035
0.05000	0.12719	0.10688	6.09150	3.01110	3.36959	2.49680	77.53824	7.27308	11.60967	2.88137
0.10000	0.21103	0.16547	5.65531	2.51453	2.93327	2.00314	62.20754	6.79993	10.93332	2.43786
0.15000	0.26898	0.19799	5.33144	2.16433	2.61075	1.65669	51.44853	6.43444	10.33526	2.11483
0.20000	0.31092	0.21626	5.08406	1.91048	2.36545	1.40670	43.68521	6.15113	9.81886	1.87226
0.25000	0.34258	0.22661	4.89003	1.72147	2.17388	1.22138	37.93009	5.92942	9.38004	1.68494
0.30000	0.36738	0.23250	4.73412	1.57712	2.02062	1.08043	33.55283	5.75361	9.01047	1.53659
0.35000	0.38740	0.23583	4.60609	1.46422	1.89534	0.97064	30.14315	5.61208	8.70022	1.41647
0.40000	0.40401	0.23766	4.49893	1.37397	1.79097	0.88322	27.42858	5.49635	8.43945	1.31728
0.45000	0.41806	0.23861	4.40769	1.30039	1.70256	0.81225	25.22442	5.40025	8.21931	1.23399
0.50000	0.43018	0.23903	4.32886	1.23933	1.62656	0.75359	23.40289	5.31927	8.03229	1.16300
0.55000	0.44078	0.23913	4.25986	1.18785	1.56039	0.70436	21.87379	5.25012	7.87222	1.10172
0.60000	0.45016	0.23902	4.19878	1.14384	1.50214	0.66244	20.57223	5.19034	7.73411	1.04823
0.65000	0.45855	0.23880	4.14416	1.10574	1.45036	0.62633	19.45062	5.13808	7.61399	1.00106
0.70000	0.46613	0.23850	4.09490	1.07240	1.40393	0.59486	18.47345	5.09193	7.50869	0.95912
0.75000	0.47302	0.23815	4.05013	1.04293	1.36197	0.56718	17.61382	5.05081	7.41569	0.92152
0.80000	0.47934	0.23778	4.00915	1.01665	1.32381	0.54262	16.85101	5.01388	7.33296	0.88760
0.85000	0.48516	0.23740	3.97141	0.99303	1.28889	0.52065	16.16884	4.98047	7.25889	0.85680
0.90000	0.49055	0.23701	3.93647	0.97165	1.25675	0.50087	15.55453	4.95004	7.19216	0.82868
0.95000	0.49558	0.23662	3.90396	0.95218	1.22704	0.48294	14.99784	4.92216	7.13170	0.80287
1.00000	0.50028	0.23623	3.87358	0.93434	1.19945	0.46661	14.49050	4.89647	7.07662	0.77908

Table 39. Summary of yield and spawning stock biomass per recruit results, corresponding biological reference points and differences between current and former analyses.

	Results			Age 3		
	F40%	Y/R	SSB/R	Mean Rec	Bmsy	MSY
RUN 0 BRP ages 3-11+	0.1643	0.2405	1.6023	12.42	19,901	2,987
RUN 1 BRP with ages 3-20	0.2033	0.2506	1.3692	12.42	17,006	3,113
RUN 2 with new pr	0.2666	0.2712	1.3694	19.6	26,840	5,316
RUN 3 with new wts	0.1918	0.2151	1.3321	19.6	26,108	4,217
RUN 4 with new maturity	0.1920	0.2473	1.3249	19.6	25,968	4,847
RUN 5 with new pr and wts	0.2478	0.2263	1.3322	19.6	26,111	4,435
RUN 6 with new wts and maturity	0.1807	0.2121	1.2883	19.6	25,251	4,158
RUN 7 with new pr and maturity	0.2485	0.2675	1.3246	19.6	25,963	5,244
RUN 8 with new pr, wts and maturity	0.2303	0.2232	1.2882	19.6	25,248	4,375

Differences between estimates from the current (RUN 8) and former (RUN 1) yield and spawning stock biomass per recruit analyses.

	F40%	Y/R	SSB/R	Bmsy (000's)
Total Effect = RUN 8 - RUN 1	0.07	-0.0173	-0.3142	8.242
P-R effect = RUN 2 - RUN 1	0.06	0.0206	0.0002	
Wt effect = RUN 3 - RUN 1	-0.01	-0.0355	-0.0372	
Mat effect = RUN 4 - RUN 1	-0.01	-0.0033	-0.0443	
2-way interaction				
PR and wt interaction effect	0.04	-0.0244	-0.0371	
Wt and mat interaction effect	-0.02	-0.0385	-0.0809	
PR and mat interaction effect	0.05	0.0169	-0.0446	
3-way interaction				
Total effect - all of the above effects	-0.04	0.0469	-0.0702	
SSB/R effect				-1.006
Recruit effect				9.830
interaction				-0.582

Table 40. Summary of short-term projection results for witch flounder. Projected median estimates of landings (mt), discards (mt), and spawning stock biomass (mt) are provided: 1) status quo fishing mortality ($F_{2003} = F_{2002} = 0.41$); 2) fishing mortality at $F_{msy} = F_{40\%} = 0.23$; 3) fishing mortality at 75% of F_{msy} ; and 4) status quo landings ($landings_{2003} = landings_{2002}$).

Projection input:

Age	Selectivity F	Selectivity M	Stock Weight	Land Weight	Maturity
3	0.0036	1.0000	0.0443	0.0000	0.0700
4	0.0229	1.0000	0.1214	0.3220	0.1600
5	0.0703	1.0000	0.2267	0.3380	0.3500
6	0.1931	1.0000	0.3206	0.3960	0.5900
7	0.5282	1.0000	0.4202	0.4780	0.7900
8	1.0000	1.0000	0.5151	0.5550	0.9100
9	1.0000	1.0000	0.5995	0.6400	0.9700
10	1.0000	1.0000	0.7281	0.7660	0.9900
11+	1.0000	1.0000	0.8888	0.8889	1.0000

Projection results (weight reported in '000 mt)

Scenario	Year	F	Median Landings	Median Discards	Median SSB
F status quo	2003	0.41	6.254	0.251	26.677
F status quo	2004	0.41	8.652	0.191	32.121
F status quo	2005	0.41	10.474	0.132	33.733
F status quo	2003	0.41	6.254	0.251	26.677
Fmsy	2004	0.23	5.174	0.109	32.705
Fmsy	2005	0.23	6.992	0.076	37.600
F status quo	2003	0.41	6.254	0.251	26.677
75% Fmsy	2004	0.17	3.908	0.081	32.902
75% Fmsy	2005	0.17	5.480	0.057	39.080
Landings ₂₀₀₃ = Landings ₂₀₀₂	2003	0.199	3.186	0.121	27.241
Fmsy	2004	0.23	5.781	0.111	35.389
Fmsy	2005	0.23	7.519	0.077	40.160
Landings ₂₀₀₃ = Landings ₂₀₀₂	2003	0.199	3.186	0.121	27.241
75%Fmsy	2004	0.17	4.366	0.083	35.613
75%Fmsy	2005	0.17	5.899	0.058	41.753




Figure 1. Statistical areas included in the witch flounder assessment.






Figure 2. Year-around area closures (shaded polygons) and seasonal rolling closures (open polygons).




Witch Flounder - Spring 1983 - 2002




Witch Flounder - Autumn 1983 - 2002

Figure 3. Number of witch flounder per tow in the NEFSC spring and autumn bottom trawl surveys, 1983-2002; year-around area closures are indicated by shaded polygons.



Witch Flounder - Spring 1983 - 2002 Juveniles < 25 cm



Witch Flounder - Autumn 1983 - 2002 Juveniles < 25 cm

Figure 4. Number of juvenile (< 25 cm) witch flounder per tow in the NEFSC spring and autumn surveys, 1983-2002; year-around area closures are indicated by shaded polygons.




Figure 5. Historical USA witch flounder landings (mt), excluding USA landings from the Grand Banks. Thin line represents provisional landings data taken from Lange and Lux (1978); discards are from the shrimp fishery and large-mesh otter trawl fishery.




Figure 6. Commercial landings of witch flounder by market category, 1973 – 2002.




Figure 7. Witch flounder landings at age (in numbers), 1982 -2002; selected cohorts are labeled.

Overview of Shrimp Fishery discard procedures




Figure 8. Overview of procedures used to estimate winter flounder discarded in the northern shrimp fishery in the Gulf of Maine – Georges Bank region 1982-2002.





Figure 9. Observed witch flounder discard rates (kg per day fished, closed symbols) from the Fisheries Observed Program, and estimated discards rates (kg/df, open symbols) in the northern shrimp fishery estimated from linear regressions (solid lines) of observed discard rates and NEFSC autumn survey age 3 index.

Length frequency of discarded witch flounder from the shrimp fishery




Figure 10. Proportion at length of discarded witch flounder from the Fisheries Observer Program observed tows in the northern shrimp fishery, 1989-1997. Data for 1983, 1985, and 1988 were collected from sea sampled trips prior to the FOP by NEFSC staff.




Figure 11. Mean weights at age of discarded witch flounder in the northern shrimp fishery (lines with symbols); NEFSC spring bottom trawl survey mean weights at age (lines with no symbols).




Figure 12. Witch flounder discards at age (in numbers) from the northern shrimp fishery, 1982 -2002; selected cohorts are labeled.

Large-mesh Otter Trawl Fishery Discards: Survey Filter Method




Figure 13. Overview of survey filter method used to estimate witch flounder discarded in the large-mesh otter trawl fishery in the Gulf of Maine-Georges Bank region, 1982-2002.




Figure 14. Witch flounder discards at age (in numbers) from the large-mesh otter trawl fishery using survey filter method (1982-1994) and Fisheries Observer Program data (1995 – 2002); selected cohorts are labeled.




Figure 15. Witch flounder discards at age (in numbers) from the large-mesh otter trawl fishery and the northern shrimp fishery, 1982 – 2002; selected cohorts are labeled.




Figure 16. Witch flounder catch at age (in numbers), 1982 -2002; selected cohorts are labeled.





Figure 17. Trends in USA landings per day fished (A) and days fished (B) of witch flounder, 1973 - 2002.




Figure 18. NEFSC bottom trawl survey sampling strata.




Figure 19. Massachusetts inshore bottom trawl survey area, delineated by regions.





Figure 20. Atlantic States Marine Fisheries Commission summer northern shrimp survey sampling strata.



Witch Flounder - ASMFC Summer Shrimp Survey 1985 - 2002

Figure 21. Number of witch flounder per tow in the ASMFC northern shrimp survey, 1985-2002; year-around area closures are indicated by shaded polygons.




Figure 22. Stratified mean catch (kg) per tow (A) and variance (B) of witch flounder in the NEFSC autumn bottom trawl survey for three strata sets in the Georges Bank- Gulf of Maine region, 1963 – 2001.





Figure 23. Stratified mean weight (kg) per tow, with 95% confidence limits, of witch flounder in the NEFSC spring (A) and autumn (B) bottom trawl survey, 1963 – 2003.





Figure 24. Stratified mean number per tow, with 95% confidence limits, of witch flounder in the NEFSC spring (A) and autumn (B) bottom trawl survey, 1963 – 2003.





Figure 25. Stratified mean catch per tow, in weight (A) and number (B), of witch flounder in the Massachusetts Division of Marine Fisheries spring and autumn bottom trawl surveys in Cape Cod Bay - Mass. Bay region, 1978 – 2002.




Figure 26. Stratified mean catch per tow, in weight (kg) and numbers, of witch flounder in the Atlantic States Marine Fisheries Commission summer northern shrimp survey, 1984 – 2002.




Figure 27. Stratified mean length (cm) per tow of witch flounder in NEFSC spring and autumn bottom trawl surveys in the Georges Bank- Gulf of Maine region, 1963 – 2003.




Figure 28. Stratified mean length (cm) per tow of witch flounder in Massachusetts Division of Marine Fisheries spring and autumn bottom trawl surveys in the Cape Cod Bay – Mass. Bay region, 1978 – 2002.

NEFSC SPRING SURVEY: stratified mean number per tow




Figure 29. Stratified mean number per tow at length of witch flounder in the NEFSC spring bottom trawl survey, 1968 – 2003.

NEFSC SPRING SURVEY: stratified mean number per tow




Figure 29 continued.

NEFSC SPRING SURVEY: stratified mean number per tow




Figure 29 continued.

NEFSC AUTUMN SURVEY: stratified mean number per tow




Figure 30. Stratified mean number per tow at length of witch flounder in the NEFSC autumn bottom trawl survey, 1968 – 2002.

NEFSC AUTUMN SURVEY: stratified mean number per tow




Figure 30 continued.

NEFSC AUTUMN SURVEY: stratified mean number per tow




Figure 30 continued.

Mass Inshore Survey Length Composition




Figure 31. Stratified mean number per tow at length of witch flounder in the Massachusetts Division of Marine Fisheries inshore spring and autumn bottom trawl survey, 1978 – 2002.

Mass Inshore Survey Length Composition




Figure 31 continued.

Mass Inshore Survey Length Composition




Figure 31 continued.

Mass Inshore Survey Length Composition




Figure 31 continued.




Figure 32. Stratified mean number per tow at length of witch flounder in the Atlantic States Marine Fisheries Commission northern shrimp survey, 1984 – 2002.




Figure 32 continued.

Spring Survey: Stratified mean number per tow at age




Figure 33. Stratified mean number of witch flounder per tow at age from NEFSC spring bottom trawl surveys, 1980-2002, preliminary 2003; selected cohorts are labeled.

Autumn Survey: Stratified mean number per tow at age




Figure 34. Stratified mean number of witch flounder per tow at age from NEFSC autumn bottom trawl surveys, 1980-2002; selected cohorts are labeled.





Figure 35. Mean length (cm) at age of witch flounder, age groups 4 -8, in spring (A) and autumn (B) NEFSC bottom trawl surveys, 1980 – 2002, preliminary spring 2003.




Figure 36. Mean weight at age of witch flounder (age groups 4 – 9) in the NEFSC spring bottom trawl survey, 1980 – 2002, preliminary 2003.




Figure 37. Mean weight at age of witch flounder (age groups 4 to 9) in the NEFSC autumn bottom trawl survey, 1980 – 2002.




Figure 38. Annual estimates of median age (A50) of witch flounder derived from a five-year moving time block of maturity observations collected during NEFSC spring surveys, 1980-2003.




Figure 39. Stratified mean weight (kg) per tow (thick line) and spawning biomass (kg) per tow (thin line) of witch flounder from the NEFSC spring bottom trawl surveys, 1968 – 2002, preliminary 2003.





Figure 40. Estimates of instantaneous total mortality (Z) derived from the NEFSC spring and autumn survey catch per tow at age (log ratio 7+ / 8+) (A) and 3-yr moving average of total mortality (B).





Figure 41. Scaled (Z score) NEFSC spring and autumn survey indices for ages 3 to 11+, 1982-2003.






Figure 41 continued.




Figure 42. Standardized residuals for survey indices (spring solid bar and autumn open bar) at age.




Figure 42. continued.




Figure 43. Trends in total catch and fishing mortality for witch flounder, 1982-2002.




Figure 44. Trends in spawning stock biomass and recruitment (Age 3) for witch flounder, 1982-2002.




Figure 45. Spawning stock biomass and recruits (Age 3) for witch flounder, 1982 – 2002 year classes.




Figure 46. Precision estimates of fishing mortality (F8-9) in 2002 for witch flounder. Vertical bars display both the range of the bootstrap estimates and the probability of individual values in the range. The arrows indicate the 80% confidence intervals.




Figure 47. Precision estimates of spawning stock biomass (mt) in 2002 for witch flounder. Vertical bars display both the range of the bootstrap estimates and the probability of individual values in the range. The arrows indicate the 80% confidence intervals.




Figure 48a. Retrospective analysis results for fishing mortality, (F8-9).




Figure 48b. Retrospective analysis results for spawning stock biomass.




Figure 48c. Retrospective analysis results for age 3 recruitment.

Witch Flounder Fishery Selectivity by Time Period
Model 3: Catch Scenario 2: M=0.15




Figure 49. Selectivity at age.

Witch Flounder Catch Biomass (mt), 1937-2002

Catch Scenario 2: M=0.15




Figure 50. Trends in catch biomass (mt), 1937 – 2002.

Witch Flounder Survey Selectivity
Model 3: Catch Scenario 2: M=0.15




Figure 51. Survey selectivity at age.

Witch Flounder Fall Survey Biomass Index, 1963-2002
Model 3: Catch Scenario 2: M=0.15




Figure 52. Trends in NEFSC autumn survey biomass.

Witch Flounder Spring Survey Biomass Index, 1968-2002

Model 3: Catch Scenario 2: M=0.15




Figure 53. Trends in NEFSC spring survey biomass.

Comparison of ADAPT virtual population analysis (VPA) and statistical catch-at-age analysis (SCAA) estimates of witch flounder spawning biomass, 1982-2002




Figure 54. Comparison of VPA and SCAA estimates of spawning stock biomass.

Comparison of ADAPT virtual population analysis (VPA) and statistical catch-at-age analysis (SCAA) estimates of witch flounder fishing mortality, 1982-2002




Figure 55. Comparison of VPA and SCAA estimates of fishing mortality.

Comparison of ADAPT virtual population analysis (VPA) and statistical catch-at-age analysis (SCAA) estimates of witch flounder recruitment, 1982-2002




Figure 56. Comparison of VPA and SCAA estimates of recruitment.




Figure 57. Yield per recruit (YPR) and spawning stock biomass per recruit (SSB/R) estimates for witch flounder.




Figure 58. Spawning stock biomass and fishing mortality (F 8-9) for witch flounder, 1999-2001, and 2002 with 80% confidence interval.




Figure 59. Age distribution of witch flounder spawning stock biomass, 1982 – 2002 and the expected age distribution of witch flounder at equilibrium.

APPENDIX: Full output of VPA RUN 301-f

VPA Version 2.0.11

Model ID: witch fld. run 301-f new p-r 1980-2002 maturity

Input File: H:\WITCHASS\VPA\VPA2003\SERIES3\RUN301-F.DAT

Date of Run: 15-MAY-2003

Time of Run: 21:15

Levenburg-Marquardt Algorithm Completed 7 Iterations
Residual Sum of Squares = 281.727

Number of Residuals = 364
Number of Parameters = 8
Degrees of Freedom = 356
Mean Squared Residual = 0.791369
Standard Deviation = 0.889589

Number of Years = 21
Number of Ages = 9
First Year = 1982
Youngest Age = 3
Oldest True Age = 10

Number of Survey Indices Available = 42
Number of Survey Indices Used in Estimate = 18

VPA Classic Method - Auto Estimated Q's

Stock Numbers Predicted in Terminal Year Plus One (2003)

Age	Stock Predicted	Std. Error	CV
3	19759.763	0.128070E+05	0.648137E+00
4	25441.190	0.116744E+05	0.458878E+00
5	42739.077	0.161746E+05	0.378451E+00
6	41657.971	0.138230E+05	0.331822E+00
7	21203.570	0.658855E+04	0.310728E+00
8	10370.219	0.335096E+04	0.323133E+00
9	3903.193	0.131462E+04	0.336807E+00
10	791.907	0.361682E+03	0.456723E+00

Catchability Values for Each Survey Used in Estimate

INDEX	Catchability	Std. Error	CV
4	0.134450E-04	0.338182E-05	0.251531E+00
5	0.323264E-04	0.480692E-05	0.148699E+00
6	0.429554E-04	0.514639E-05	0.119808E+00
7	0.430079E-04	0.596770E-05	0.138758E+00
8	0.508706E-04	0.712473E-05	0.140056E+00
9	0.656138E-04	0.131051E-04	0.199731E+00
10	0.788325E-04	0.158102E-04	0.200554E+00
11	0.836507E-04	0.173114E-04	0.206948E+00
18	0.113695E-03	0.150419E-04	0.132300E+00
21	0.834330E-05	0.224168E-05	0.268681E+00
22	0.307342E-04	0.636966E-05	0.207250E+00
23	0.385675E-04	0.103815E-04	0.269179E+00
24	0.410818E-04	0.106353E-04	0.258880E+00
25	0.461380E-04	0.918963E-05	0.199177E+00
26	0.772831E-04	0.152953E-04	0.197913E+00
27	0.796591E-04	0.151773E-04	0.190528E+00
28	0.913192E-04	0.204756E-04	0.224221E+00
35	0.111457E-03	0.183216E-04	0.164383E+00

-- Non-Linear Least Squares Fit --

Default Tolerances Used

Scaled Gradient Tolerance = 6.055454E-06
Scaled Step Tolerance = 3.666853E-11
Relative Function Tolerance = 3.666853E-11
Absolute Function Tolerance = 4.930381E-32

VPA Method Options

- Catch Equation Used in Cohort Solution
- Plus Group Backward Calculation Method Used
- Rivard Weights Used for JAN-1 Biomass
- Rivard Weights Used for SSB Biomass
- Heincke Rule Used in F-Oldest Calculation
- F-Oldest Calculation in Years Prior to Terminal Year
Uses Stock Sizes in Ages 8 to 10
- Calculation of Population of Age 3 In Year 2003
= Stock Estimate

Stock Estimates

Age 3
Age 4
Age 5
Age 6
Age 7
Age 8
Age 9
Age 10

Full F in Terminal Year = 0.4087

F in Oldest True Age in Terminal Year = 0.4087

Full F Calculated Using Classic Method

Age	Input Partial Recruitment	Calc Partial Recruitment	Fishing Mortality	Used In Full F	Comments
3	0.004	0.003	0.0015	NO	Stock Estimate in T+1
4	0.023	0.029	0.0161	NO	Stock Estimate in T+1
5	0.070	0.049	0.0268	NO	Stock Estimate in T+1
6	0.193	0.108	0.0595	NO	Stock Estimate in T+1
7	0.528	0.315	0.1741	NO	Stock Estimate in T+1
8	1.000	0.479	0.2648	YES	Stock Estimate in T+1
9	1.000	1.000	0.5526	YES	Stock Estimate in T+1
10	1.000	0.740	0.4087		Input PR * Full F

JAN-1 Population Numbers

AGE	1982	1983	1984	1985	1986
3	15404.	17700.	16340.	7650.	5414.
4	12174.	13082.	14922.	13928.	6470.
5	9563.	9493.	10014.	11486.	10899.
6	7830.	7114.	6765.	6768.	7929.
7	4289.	5375.	4668.	4216.	4039.
8	2752.	3077.	3160.	2647.	2224.
9	2102.	1763.	1746.	1344.	1132.
10	1101.	1440.	839.	862.	599.
11	7260.	4728.	3844.	2926.	2039.
Total	62476.	63772.	62297.	51828.	40746.
AGE	1987	1988	1989	1990	1991
3	3001.	9740.	6214.	6670.	8831.
4	4639.	2562.	7827.	4934.	5422.
5	5220.	3825.	2076.	6333.	3660.
6	7978.	4060.	3047.	1726.	4431.
7	4267.	5683.	2886.	2332.	1249.
8	2034.	2222.	3615.	1783.	1754.
9	1146.	949.	853.	2297.	1100.
10	594.	544.	448.	413.	1668.
11	1151.	1218.	1263.	900.	1371.
Total	30029.	30804.	28229.	27388.	29487.
AGE	1992	1993	1994	1995	1996
3	14342.	9670.	13610.	12671.	15878.
4	7192.	11974.	7936.	11675.	10324.
5	4291.	5029.	8635.	5904.	9706.
6	2353.	2900.	3019.	4887.	4119.
7	3281.	1164.	1649.	1414.	2627.
8	857.	2162.	453.	660.	438.
9	1283.	552.	1320.	209.	321.
10	677.	940.	274.	640.	91.
11	2127.	1318.	782.	373.	180.
Total	36404.	35708.	37679.	38434.	43683.
AGE	1997	1998	1999	2000	2001
3	20226.	29659.	42904.	67651.	58704.
4	13554.	17283.	25221.	36778.	58131.
5	8400.	10686.	14360.	21220.	31390.
6	7237.	6290.	8213.	11345.	17842.
7	2289.	5033.	4106.	5717.	8693.
8	949.	1033.	2846.	2426.	3417.
9	136.	274.	547.	1743.	1138.
10	80.	39.	105.	239.	926.
11	112.	179.	181.	536.	667.
Total	52982.	70476.	98483.	147655.	180909.

JAN-1 Population Numbers

AGE	2002	2003
3	29603.	19760.
4	50462.	25441.
5	49716.	42739.
6	26146.	41658.
7	14340.	21204.
8	5910.	10370.
9	1599.	3903.
10	398.	792.
11	655.	603.
Total	178829.	166470.

Fishing Mortality Calculated

AGE	1982	1983	1984	1985	1986
3	0.0134	0.0207	0.0097	0.0175	0.0046
4	0.0988	0.1173	0.1117	0.0952	0.0648
5	0.1459	0.1889	0.2417	0.2207	0.1620
6	0.2261	0.2713	0.3227	0.3662	0.4696
7	0.1823	0.3814	0.4173	0.4895	0.5361
8	0.2953	0.4162	0.7051	0.6998	0.5136
9	0.2282	0.5929	0.5562	0.6573	0.4949
10	0.2657	0.4770	0.6495	0.6853	0.5072
11	0.2657	0.4770	0.6495	0.6853	0.5072
AGE	1987	1988	1989	1990	1991
3	0.0080	0.0686	0.0806	0.0571	0.0553
4	0.0430	0.0605	0.0618	0.1487	0.0839
5	0.1012	0.0773	0.0344	0.2072	0.2919
6	0.1892	0.1914	0.1176	0.1733	0.1503
7	0.5025	0.3025	0.3316	0.1350	0.2265
8	0.6118	0.8070	0.3035	0.3332	0.1624
9	0.5951	0.6015	0.5765	0.1699	0.3352
10	0.6058	0.7409	0.3502	0.2380	0.2255
11	0.6058	0.7409	0.3502	0.2380	0.2255
AGE	1992	1993	1994	1995	1996
3	0.0304	0.0475	0.0033	0.0549	0.0083
4	0.2077	0.1769	0.1458	0.0347	0.0562
5	0.2420	0.3602	0.4192	0.2100	0.1436
6	0.5542	0.4143	0.6087	0.4709	0.4376
7	0.2672	0.7926	0.7655	1.0227	0.8680
8	0.2907	0.3431	0.6230	0.5707	1.0205
9	0.1614	0.5517	0.5738	0.6881	1.2443
10	0.2112	0.3821	0.5862	0.5977	1.1092
11	0.2112	0.3821	0.5862	0.5977	1.1092

AGE	1997	1998	1999	2000	2001
3	0.0072	0.0121	0.0040	0.0016	0.0013
4	0.0877	0.0353	0.0227	0.0084	0.0064
5	0.1393	0.1132	0.0857	0.0234	0.0328
6	0.2131	0.2764	0.2122	0.1163	0.0685
7	0.6456	0.4201	0.3762	0.3646	0.2359
8	1.0930	0.4859	0.3402	0.6068	0.6097
9	1.0854	0.8049	0.6793	0.4829	0.9002
10	1.0920	0.5448	0.3876	0.5531	0.6747
11	1.0920	0.5448	0.3876	0.5531	0.6747

AGE	2002
3	0.0015
4	0.0161
5	0.0268
6	0.0595
7	0.1741
8	0.2648
9	0.5526
10	0.4087
11	0.4087

Average Fishing Mortality For Ages 8- 9

Year	Average F	N Weighted	Biomass Wtd	Catch Wtd
1982	0.2618	0.2663	0.2631	0.2699
1983	0.5046	0.4806	0.4920	0.4922
1984	0.6307	0.6521	0.6456	0.6578
1985	0.6786	0.6855	0.6835	0.6859
1986	0.5042	0.5073	0.5063	0.5074
1987	0.6035	0.6058	0.6050	0.6059
1988	0.7043	0.7455	0.7364	0.7540
1989	0.4400	0.3557	0.3646	0.3810
1990	0.2515	0.2412	0.2337	0.2654
1991	0.2488	0.2290	0.2360	0.2565
1992	0.2260	0.2132	0.2086	0.2301
1993	0.4474	0.3855	0.3939	0.3999
1994	0.5984	0.5864	0.5846	0.5870
1995	0.6294	0.5990	0.6051	0.6020
1996	1.1324	1.1152	1.1262	1.1213
1997	1.0892	1.0920	1.0918	1.0920
1998	0.6454	0.5527	0.5710	0.5741
1999	0.5097	0.3948	0.4009	0.4243
2000	0.5448	0.5550	0.5504	0.5601
2001	0.7550	0.6823	0.6917	0.6977
2002	0.4087	0.3261	0.3344	0.3601

Average Fishing Mortality For Ages 8- 9

Year	Average F	N Weighted	Biomass Wtd	Catch Wtd
1982	0.2618	0.2663	0.2631	0.2699
1983	0.5046	0.4806	0.4920	0.4922
1984	0.6307	0.6521	0.6456	0.6578
1985	0.6786	0.6855	0.6835	0.6859
1986	0.5042	0.5073	0.5063	0.5074
1987	0.6035	0.6058	0.6050	0.6059
1988	0.7043	0.7455	0.7364	0.7540
1989	0.4400	0.3557	0.3646	0.3810
1990	0.2515	0.2412	0.2337	0.2654
1991	0.2488	0.2290	0.2360	0.2565
1992	0.2260	0.2132	0.2086	0.2301
1993	0.4474	0.3855	0.3939	0.3999
1994	0.5984	0.5864	0.5846	0.5870
1995	0.6294	0.5990	0.6051	0.6020
1996	1.1324	1.1152	1.1262	1.1213
1997	1.0892	1.0920	1.0918	1.0920
1998	0.6454	0.5527	0.5710	0.5741
1999	0.5097	0.3948	0.4009	0.4243
2000	0.5448	0.5550	0.5504	0.5601
2001	0.7550	0.6823	0.6917	0.6977
2002	0.4087	0.3261	0.3344	0.3601

Back Calculated Partial Recruitment

AGE	1982	1983	1984	1985	1986
3	0.0454	0.0349	0.0138	0.0251	0.0085
4	0.3344	0.1978	0.1584	0.1361	0.1208
5	0.4940	0.3186	0.3428	0.3154	0.3022
6	0.7655	0.4576	0.4577	0.5233	0.8759
7	0.6172	0.6432	0.5918	0.6995	1.0000
8	1.0000	0.7021	1.0000	1.0000	0.9579
9	0.7728	1.0000	0.7889	0.9393	0.9231
10	0.8997	0.8046	0.9212	0.9793	0.9461
11	0.8997	0.8046	0.9212	0.9793	0.9461
AGE	1987	1988	1989	1990	1991
3	0.0131	0.0850	0.1397	0.1713	0.1651
4	0.0702	0.0750	0.1073	0.4462	0.2504
5	0.1654	0.0957	0.0597	0.6218	0.8709
6	0.3092	0.2372	0.2039	0.5201	0.4485
7	0.8212	0.3748	0.5751	0.4052	0.6756
8	1.0000	1.0000	0.5265	1.0000	0.4845
9	0.9726	0.7454	1.0000	0.5099	1.0000
10	0.9901	0.9182	0.6075	0.7143	0.6728
11	0.9901	0.9182	0.6075	0.7143	0.6728
AGE	1992	1993	1994	1995	1996
3	0.0549	0.0600	0.0043	0.0537	0.0067
4	0.3747	0.2232	0.1905	0.0340	0.0452
5	0.4367	0.4545	0.5476	0.2053	0.1154
6	1.0000	0.5227	0.7952	0.4604	0.3516
7	0.4822	1.0000	1.0000	1.0000	0.6976
8	0.5246	0.4329	0.8139	0.5580	0.8202
9	0.2912	0.6961	0.7496	0.6728	1.0000
10	0.3811	0.4821	0.7658	0.5845	0.8914
11	0.3811	0.4821	0.7658	0.5845	0.8914
AGE	1997	1998	1999	2000	2001
3	0.0066	0.0150	0.0060	0.0027	0.0014
4	0.0803	0.0439	0.0334	0.0139	0.0071
5	0.1274	0.1407	0.1261	0.0386	0.0364
6	0.1950	0.3434	0.3124	0.1916	0.0761
7	0.5907	0.5219	0.5538	0.6009	0.2620
8	1.0000	0.6036	0.5008	1.0000	0.6772
9	0.9931	1.0000	1.0000	0.7958	1.0000
10	0.9991	0.6769	0.5707	0.9115	0.7495
11	0.9991	0.6769	0.5707	0.9115	0.7495

Back Calculated Partial Recruitment

AGE	2002
3	0.0027
4	0.0292
5	0.0486
6	0.1077
7	0.3151
8	0.4793
9	1.0000
10	0.7396
11	0.7396

JAN-1 Biomass

AGE	1982	1983	1984	1985	1986
3	2027.	1336.	1247.	550.	247.
4	2784.	2293.	2758.	2637.	1050.
5	2822.	2423.	2581.	3031.	2902.
6	2967.	2611.	2280.	2539.	2795.
7	2235.	2509.	2192.	2058.	1931.
8	1913.	1786.	1853.	1615.	1375.
9	1774.	1340.	1236.	1005.	869.
10	1028.	1340.	718.	765.	543.
11	10204.	6418.	5148.	3878.	2694.
Total	27754.	22056.	20012.	18077.	14406.
AGE	1987	1988	1989	1990	1991
3	137.	374.	284.	305.	376.
4	603.	329.	684.	745.	601.
5	1293.	932.	530.	1325.	927.
6	2874.	1461.	1108.	648.	1456.
7	2041.	2744.	1441.	1164.	629.
8	1230.	1360.	2189.	1120.	1125.
9	857.	712.	631.	1748.	834.
10	543.	490.	399.	382.	1517.
11	1500.	1615.	1714.	1309.	1947.
Total	11077.	10017.	8980.	8745.	9411.
AGE	1992	1993	1994	1995	1996
3	866.	544.	591.	390.	607.
4	857.	1730.	1247.	1226.	896.
5	1178.	1359.	2055.	1518.	1988.
6	935.	1173.	1117.	1706.	1532.
7	1667.	577.	792.	694.	1289.
8	560.	1383.	276.	401.	276.
9	975.	446.	983.	166.	247.
10	581.	862.	245.	576.	85.
11	2644.	1760.	989.	463.	222.
Total	10264.	9833.	8294.	7141.	7141.
AGE	1997	1998	1999	2000	2001
3	941.	1433.	1742.	2699.	2718.
4	1438.	1848.	3110.	4230.	6255.
5	1698.	2367.	3130.	5169.	7022.
6	2282.	1972.	2649.	3712.	5611.
7	1046.	2154.	1763.	2422.	3596.
8	560.	556.	1518.	1269.	1697.
9	106.	202.	331.	1060.	668.
10	75.	36.	94.	156.	592.
11	145.	216.	158.	495.	561.
Total	8290.	10782.	14496.	21212.	28721.

JAN-1 Biomass

AGE	2002	2003
3	1492.	1433.
4	7039.	6106.
5	11002.	14813.
6	8312.	14601.
7	5937.	10411.
8	2998.	5619.
9	956.	2356.
10	290.	561.
11	602.	553.
Total	38628.	56452.

Mean Biomass

AGE	1982	1983	1984	1985	1986
3	2159.	2431.	2286.	902.	444.
4	2604.	2322.	3001.	2925.	1198.
5	2724.	2171.	2719.	2924.	2803.
6	2744.	2375.	2272.	2267.	2410.
7	2007.	2160.	1920.	1760.	1558.
8	1615.	1440.	1410.	1232.	1099.
9	1551.	989.	1023.	776.	711.
10	886.	1045.	532.	563.	428.
11	8349.	4768.	3544.	2629.	1974.
Total	24639.	19702.	18709.	15978.	12626.
AGE	1987	1988	1989	1990	1991
3	226.	392.	679.	372.	494.
4	806.	469.	1202.	796.	961.
5	1376.	1064.	609.	1368.	1018.
6	2932.	1493.	1136.	646.	1609.
7	1758.	2459.	1315.	1189.	602.
8	976.	955.	1982.	973.	1057.
9	669.	547.	496.	1669.	728.
10	408.	353.	341.	359.	1354.
11	1053.	1069.	1349.	1085.	1623.
Total	10205.	8801.	9109.	8455.	9447.
AGE	1992	1993	1994	1995	1996
3	1356.	1073.	878.	580.	877.
4	1389.	2063.	1386.	1689.	1390.
5	1346.	1253.	1846.	1626.	2226.
6	775.	957.	909.	1582.	1314.
7	1647.	403.	577.	467.	913.
8	512.	1137.	218.	324.	183.
9	906.	350.	782.	129.	148.
10	501.	746.	176.	439.	50.
11	2219.	1364.	700.	326.	126.
Total	10652.	9346.	7473.	7162.	7227.
AGE	1997	1998	1999	2000	2001
3	1308.	1917.	2688.	4130.	4821.
4	2274.	2578.	5031.	6656.	9461.
5	1999.	2445.	3722.	5325.	7338.
6	2280.	1837.	2756.	3666.	5800.
7	781.	1869.	1641.	2004.	3354.
8	341.	447.	1313.	906.	1317.
9	68.	153.	233.	818.	455.
10	47.	28.	75.	116.	408.
11	83.	156.	122.	356.	382.
Total	9181.	11429.	17581.	23978.	33336.

Mean Biomass

AGE	2002
3	3022.
4	10224.
5	12691.
6	9320.
7	5801.
8	2677.
9	747.
10	251.
11	461.
=====	
Total	45192.

Spawning Stock Biomass

AGE	1982	1983	1984	1985	1986
3	20.	13.	24.	5.	5.
4	107.	132.	185.	127.	91.
5	376.	458.	580.	684.	1019.
6	1115.	1241.	1243.	1584.	1916.
7	1543.	1883.	1715.	1720.	1636.
8	1634.	1544.	1559.	1388.	1218.
9	1632.	1172.	1088.	878.	780.
10	949.	1207.	628.	665.	487.
11	9521.	5781.	4505.	3374.	2414.
Total	16897.	13431.	11528.	10425.	9567.
AGE	1987	1988	1989	1990	1991
3	16.	115.	33.	24.	25.
4	216.	191.	238.	156.	104.
5	880.	745.	360.	587.	336.
6	2471.	1297.	964.	454.	886.
7	1794.	2494.	1303.	999.	496.
8	1072.	1148.	2010.	1003.	1004.
9	757.	628.	559.	1640.	754.
10	479.	422.	367.	358.	1411.
11	1323.	1392.	1577.	1227.	1829.
Total	9008.	8433.	7410.	6447.	6844.
AGE	1992	1993	1994	1995	1996
3	50.	21.	23.	11.	12.
4	145.	229.	154.	155.	113.
5	463.	437.	636.	601.	852.
6	582.	694.	630.	1200.	1139.
7	1369.	424.	585.	542.	1044.
8	500.	1210.	230.	352.	225.
9	916.	392.	862.	144.	196.
10	547.	789.	217.	509.	69.
11	2490.	1610.	875.	409.	180.
Total	7063.	5806.	4212.	3922.	3828.
AGE	1997	1998	1999	2000	2001
3	9.	56.	85.	158.	186.
4	97.	251.	453.	618.	975.
5	599.	883.	1114.	1657.	2384.
6	1740.	1304.	1671.	2095.	3192.
7	888.	1782.	1405.	1800.	2664.
8	455.	485.	1343.	1040.	1361.
9	87.	171.	286.	925.	544.
10	61.	32.	86.	137.	511.
11	118.	192.	145.	441.	489.
Total	4053.	5156.	6587.	8871.	12305.

Spawning Stock Biomass

AGE	2002
3	102.
4	1096.
5	3739.
6	4736.
7	4443.
8	2545.
9	825.
10	262.
11	548.
=====	
Total	18296.

Catch Biomass

AGE	1982	1983	1984	1985	1986
3	29.	50.	22.	16.	2.
4	257.	272.	335.	279.	78.
5	397.	410.	657.	645.	454.
6	620.	645.	733.	830.	1132.
7	366.	824.	801.	862.	835.
8	477.	599.	994.	862.	564.
9	354.	586.	569.	510.	352.
10	235.	499.	346.	386.	217.
11	2219.	2274.	2302.	1802.	1002.
Total	4955.	6160.	6761.	6192.	4636.
AGE	1987	1988	1989	1990	1991
3	2.	27.	55.	21.	27.
4	35.	28.	74.	118.	81.
5	139.	82.	21.	283.	297.
6	555.	286.	134.	112.	242.
7	883.	744.	436.	161.	136.
8	597.	771.	602.	324.	172.
9	398.	329.	286.	283.	244.
10	247.	261.	119.	85.	305.
11	638.	792.	472.	258.	366.
Total	3494.	3320.	2199.	1646.	1871.
AGE	1992	1993	1994	1995	1996
3	41.	51.	3.	32.	7.
4	288.	365.	202.	59.	78.
5	326.	451.	774.	342.	320.
6	430.	397.	554.	745.	575.
7	440.	320.	442.	477.	792.
8	149.	390.	136.	185.	186.
9	146.	193.	449.	89.	184.
10	106.	285.	103.	262.	56.
11	469.	521.	410.	195.	140.
Total	2395.	2973.	3072.	2386.	2339.
AGE	1997	1998	1999	2000	2001
3	9.	23.	11.	7.	6.
4	200.	91.	114.	56.	60.
5	278.	277.	319.	125.	241.
6	486.	508.	585.	426.	397.
7	504.	785.	617.	731.	791.
8	372.	217.	447.	550.	803.
9	73.	123.	158.	395.	410.
10	52.	15.	29.	64.	275.
11	91.	85.	47.	197.	258.
Total	2066.	2124.	2327.	2550.	3241.

Catch Biomass

AGE	2002
3	5.
4	165.
5	341.
6	555.
7	1010.
8	709.
9	413.
10	80.
11	188.
=====	
Total	3465.

Surplus Production

Average Adjustment Factor (Delta) = 1.0000

Year	Biomass	Delta Biomass	Catch Biomass	Surplus Production
1982	27753.983	-5697.846	4954.638	-743.208
1983	22056.137	-2044.094	6159.976	4115.882
1984	20012.043	-1934.768	6760.517	4825.749
1985	18077.275	-3671.668	6191.704	2520.036
1986	14405.606	-3328.378	4635.851	1307.473
1987	11077.228	-1059.758	3494.396	2434.637
1988	10017.470	-1037.240	3320.343	2283.103
1989	8980.230	-235.331	2199.015	1963.684
1990	8744.899	666.450	1646.456	2312.905
1991	9411.349	852.941	1870.786	2723.727
1992	10264.289	-430.994	2394.876	1963.882
1993	9833.296	-1539.477	2972.811	1433.334
1994	8293.818	-1153.030	3072.149	1919.119
1995	7140.788	-0.006	2385.733	2385.726
1996	7140.782	1149.552	2338.662	3488.214
1997	8290.334	2492.097	2066.078	4558.175
1998	10782.431	3713.086	2124.350	5837.436
1999	14495.517	6716.072	2327.426	9043.498
2000	21211.589	7509.247	2550.036	10059.282
2001	28720.836	9906.770	3241.147	13147.917
2002	38627.606	17824.499	3464.690	21289.189
2003	56452.105			

Summary of Survey Indices Used in the Estimate

INDEX	Survey Tag	Age	Time	Type
4	Spr	3	JAN-1	NUMBER
5	Spr	4	JAN-1	NUMBER
6	Spr	5	JAN-1	NUMBER
7	Spr	6	JAN-1	NUMBER
8	Spr	7	JAN-1	NUMBER
9	Spr	8	JAN-1	NUMBER
10	Spr	9	JAN-1	NUMBER
11	Spr	10	JAN-1	NUMBER
18	Spr	11 - 11	JAN-1	NUMBER
21	Aut	3	JAN-1	NUMBER
22	Aut	4	JAN-1	NUMBER
23	Aut	5	JAN-1	NUMBER
24	Aut	6	JAN-1	NUMBER
25	Aut	7	JAN-1	NUMBER
26	Aut	8	JAN-1	NUMBER
27	Aut	9	JAN-1	NUMBER
28	Aut	10	JAN-1	NUMBER
35	Aut	11 - 11	JAN-1	NUMBER

Survey Index: 4 Tag: Spr AGE = 3
 Time = JAN-1 Type = NUMBER
 Catchability = 0.134450E-04 Variance = 0.126535E+01
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.609600E+00	0.207109E+00	0.107956E+01
1983	0.531000E+00	0.237975E+00	0.802595E+00
1984	0.115000E-01	0.219690E+00	-0.294987E+01
1985	0.167000E-01	0.102859E+00	-0.181795E+01
1986	N/A	0.727929E-01	N/A
1987	N/A	0.403491E-01	N/A
1988	0.620000E-01	0.130952E+00	-0.747699E+00
1989	0.361000E-01	0.835456E-01	-0.839100E+00
1990	0.375000E-01	0.896771E-01	-0.871874E+00
1991	0.780500E+00	0.118731E+00	0.188307E+01
1992	0.187000E+00	0.192833E+00	-0.307154E-01
1993	0.137400E+00	0.130007E+00	0.553119E-01
1994	0.540900E+00	0.182987E+00	0.108382E+01
1995	0.581200E+00	0.170361E+00	0.122717E+01
1996	0.243500E+00	0.213484E+00	0.131556E+00
1997	0.152000E+00	0.271937E+00	-0.581690E+00
1998	0.712200E+00	0.398763E+00	0.579991E+00
1999	0.973800E+00	0.576838E+00	0.523645E+00
2000	0.119360E+01	0.909560E+00	0.271769E+00
2001	0.712600E+00	0.789274E+00	-0.102193E+00
2002	0.897200E+00	0.398006E+00	0.812812E+00
2003	0.159500E+00	0.265669E+00	-0.510209E+00

Survey Index: 5 Tag: Spr AGE = 4
 Time = JAN-1 Type = NUMBER
 Catchability = 0.323264E-04 Variance = 0.464341E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.483800E+00	0.393557E+00	0.206446E+00
1983	0.126160E+01	0.422894E+00	0.109302E+01
1984	0.307200E+00	0.482379E+00	-0.451232E+00
1985	0.459200E+00	0.450245E+00	0.196933E-01
1986	0.439000E-01	0.209159E+00	-0.156118E+01
1987	0.589000E-01	0.149955E+00	-0.934493E+00
1988	N/A	0.828321E-01	N/A
1989	0.100440E+01	0.253028E+00	0.137865E+01
1990	0.913000E-01	0.159510E+00	-0.557958E+00
1991	0.108400E+00	0.175285E+00	-0.480585E+00
1992	0.372800E+00	0.232479E+00	0.472242E+00
1993	0.472300E+00	0.387092E+00	0.198953E+00
1994	0.643500E+00	0.256548E+00	0.919605E+00
1995	0.315600E+00	0.377423E+00	-0.178891E+00
1996	0.394300E+00	0.333725E+00	0.166793E+00
1997	0.692500E+00	0.438141E+00	0.457767E+00
1998	0.388100E+00	0.558705E+00	-0.364359E+00
1999	0.796900E+00	0.815297E+00	-0.228229E-01
2000	0.692300E+00	0.118891E+01	-0.540776E+00
2001	0.147560E+01	0.187918E+01	-0.241772E+00
2002	0.262720E+01	0.163126E+01	0.476563E+00
2003	0.777900E+00	0.822423E+00	-0.556568E-01

Survey Index: 6 Tag: Spr AGE = 5
 Time = JAN-1 Type = NUMBER
 Catchability = 0.429554E-04 Variance = 0.315786E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.376500E+00	0.410784E+00	-0.871507E-01
1983	0.129330E+01	0.407785E+00	0.115421E+01
1984	0.777500E+00	0.430148E+00	0.591955E+00
1985	0.105720E+01	0.493403E+00	0.762052E+00
1986	0.240300E+00	0.468180E+00	-0.666964E+00
1987	0.113800E+00	0.224215E+00	-0.678163E+00
1988	0.724000E-01	0.164290E+00	-0.819428E+00
1989	0.105400E+00	0.891701E-01	0.167217E+00
1990	0.319400E+00	0.272036E+00	0.160509E+00
1991	0.870000E-01	0.157234E+00	-0.591827E+00
1992	0.852000E-01	0.184335E+00	-0.771755E+00
1993	0.319500E+00	0.216027E+00	0.391354E+00
1994	0.810300E+00	0.370923E+00	0.781411E+00
1995	0.178800E+00	0.253603E+00	-0.349503E+00
1996	0.345500E+00	0.416925E+00	-0.187915E+00
1997	0.616900E+00	0.360821E+00	0.536325E+00
1998	0.797800E+00	0.459009E+00	0.552789E+00
1999	0.482400E+00	0.616833E+00	-0.245825E+00
2000	0.659600E+00	0.911522E+00	-0.323482E+00
2001	0.102000E+01	0.134837E+01	-0.279090E+00
2002	0.226340E+01	0.213557E+01	0.581323E-01
2003	0.157250E+01	0.183587E+01	-0.154853E+00

Survey Index: 7 Tag: Spr AGE = 6
 Time = JAN-1 Type = NUMBER
 Catchability = 0.430079E-04 Variance = 0.404332E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.236700E+00	0.336734E+00	-0.352499E+00
1983	0.541000E+00	0.305943E+00	0.570021E+00
1984	0.401200E+00	0.290932E+00	0.321371E+00
1985	0.119940E+01	0.291091E+00	0.141594E+01
1986	0.528500E+00	0.340990E+00	0.438190E+00
1987	0.132800E+00	0.343117E+00	-0.949227E+00
1988	0.299900E+00	0.174623E+00	0.540819E+00
1989	0.727000E-01	0.131053E+00	-0.589262E+00
1990	N/A	0.742453E-01	N/A
1991	0.208600E+00	0.190567E+00	0.904155E-01
1992	0.110500E+00	0.101193E+00	0.879816E-01
1993	0.576000E-01	0.124706E+00	-0.772436E+00
1994	0.164400E+00	0.129855E+00	0.235883E+00
1995	0.311800E+00	0.210201E+00	0.394298E+00
1996	0.218400E+00	0.177149E+00	0.209339E+00
1997	0.437400E+00	0.311234E+00	0.340304E+00
1998	0.713200E+00	0.270515E+00	0.969435E+00
1999	0.164300E+00	0.353213E+00	-0.765378E+00
2000	0.239400E+00	0.487927E+00	-0.712031E+00
2001	0.400700E+00	0.767345E+00	-0.649724E+00
2002	0.822200E+00	0.112450E+01	-0.313107E+00
2003	0.107550E+01	0.179162E+01	-0.510335E+00

Survey Index: 8 Tag: Spr AGE = 7
 Time = JAN-1 Type = NUMBER
 Catchability = 0.508706E-04 Variance = 0.431545E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.608500E+00	0.218195E+00	0.102561E+01
1983	0.716200E+00	0.273446E+00	0.962856E+00
1984	0.309500E+00	0.237453E+00	0.264989E+00
1985	0.907500E+00	0.214493E+00	0.144242E+01
1986	0.411700E+00	0.205471E+00	0.694989E+00
1987	0.259200E+00	0.217053E+00	0.177457E+00
1988	0.379100E+00	0.289099E+00	0.271030E+00
1989	0.811000E-01	0.146812E+00	-0.593470E+00
1990	0.421000E-01	0.118623E+00	-0.103590E+01
1991	0.328000E-01	0.635607E-01	-0.661566E+00
1992	0.151800E+00	0.166929E+00	-0.950066E-01
1993	0.853000E-01	0.591906E-01	0.365411E+00
1994	0.268000E-01	0.838920E-01	-0.114113E+01
1995	0.116200E+00	0.719269E-01	0.479662E+00
1996	0.725000E-01	0.133630E+00	-0.611485E+00
1997	0.841000E-01	0.116435E+00	-0.325330E+00
1998	0.214300E+00	0.256033E+00	-0.177928E+00
1999	0.182300E+00	0.208889E+00	-0.136148E+00
2000	0.252800E+00	0.290841E+00	-0.140178E+00
2001	0.292800E+00	0.442219E+00	-0.412316E+00
2002	0.683000E+00	0.729464E+00	-0.658151E-01
2003	0.808600E+00	0.107864E+01	-0.288150E+00

Survey Index: 9 Tag: Spr AGE = 8
 Time = JAN-1 Type = NUMBER
 Catchability = 0.656138E-04 Variance = 0.797849E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.361900E+00	0.180588E+00	0.695149E+00
1983	0.632200E+00	0.201868E+00	0.114159E+01
1984	0.201700E+00	0.207317E+00	-0.274654E-01
1985	0.411900E+00	0.173675E+00	0.863593E+00
1986	0.171800E+00	0.145945E+00	0.163102E+00
1987	0.185300E+00	0.133445E+00	0.328286E+00
1988	0.238500E+00	0.145792E+00	0.492186E+00
1989	0.327400E+00	0.237172E+00	0.322398E+00
1990	0.870000E-02	0.116991E+00	-0.259878E+01
1991	0.100700E+00	0.115060E+00	-0.133311E+00
1992	0.449000E-01	0.562624E-01	-0.225589E+00
1993	N/A	0.141858E+00	N/A
1994	0.281000E-01	0.297449E-01	-0.568894E-01
1995	0.110200E+00	0.433175E-01	0.933740E+00
1996	N/A	0.287156E-01	N/A
1997	0.828000E-01	0.622769E-01	0.284838E+00
1998	0.153500E+00	0.677777E-01	0.817468E+00
1999	0.305000E-01	0.186736E+00	-0.181197E+01
2000	0.115800E+00	0.159195E+00	-0.318264E+00
2001	0.162700E+00	0.224235E+00	-0.320786E+00
2002	0.350600E+00	0.387771E+00	-0.100769E+00
2003	0.434500E+00	0.680430E+00	-0.448528E+00

Survey Index: 10 Tag: Spr AGE = 9
 Time = JAN-1 Type = NUMBER
 Catchability = 0.788325E-04 Variance = 0.804441E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.932000E-01	0.165726E+00	-0.575587E+00
1983	0.475300E+00	0.138991E+00	0.122954E+01
1984	0.196100E+00	0.137677E+00	0.353717E+00
1985	0.147800E+00	0.105918E+00	0.333194E+00
1986	0.194100E+00	0.892031E-01	0.777457E+00
1987	0.940000E-02	0.903062E-01	-0.226250E+01
1988	0.137200E+00	0.748424E-01	0.606055E+00
1989	0.812000E-01	0.672720E-01	0.188171E+00
1990	0.503000E-01	0.181051E+00	-0.128078E+01
1991	0.828000E-01	0.867019E-01	-0.460472E-01
1992	0.148500E+00	0.101149E+00	0.383994E+00
1993	0.154000E-01	0.435042E-01	-0.103849E+01
1994	0.704000E-01	0.104090E+00	-0.391065E+00
1995	0.420000E-01	0.164975E-01	0.934460E+00
1996	N/A	0.253146E-01	N/A
1997	0.144000E-01	0.107023E-01	0.296772E+00
1998	0.756000E-01	0.215883E-01	0.125331E+01
1999	0.144000E-01	0.431154E-01	-0.109665E+01
2000	N/A	0.137423E+00	N/A
2001	0.112600E+00	0.897399E-01	0.226926E+00
2002	0.191500E+00	0.126036E+00	0.418322E+00
2003	0.225500E+00	0.307698E+00	-0.310800E+00

Survey Index: 11 Tag: Spr AGE = 10
 Time = JAN-1 Type = NUMBER
 Catchability = 0.836507E-04 Variance = 0.770897E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.259300E+00	0.921099E-01	0.103500E+01
1983	0.214100E+00	0.120473E+00	0.575016E+00
1984	0.114600E+00	0.701648E-01	0.490601E+00
1985	0.148700E+00	0.720958E-01	0.723935E+00
1986	0.791000E-01	0.501318E-01	0.456057E+00
1987	0.614000E-01	0.496685E-01	0.212039E+00
1988	0.862000E-01	0.454891E-01	0.639198E+00
1989	0.151000E-01	0.374561E-01	-0.908475E+00
1990	0.175000E-01	0.345202E-01	-0.679343E+00
1991	0.137700E+00	0.139523E+00	-0.131542E-01
1992	0.153000E-01	0.566329E-01	-0.130874E+01
1993	0.154000E-01	0.786150E-01	-0.163020E+01
1994	0.820000E-02	0.228844E-01	-0.102632E+01
1995	N/A	0.535575E-01	N/A
1996	N/A	0.757193E-02	N/A
1997	N/A	0.666199E-02	N/A
1998	N/A	0.330153E-02	N/A
1999	0.228000E-01	0.881578E-02	0.950218E+00
2000	0.354000E-01	0.199644E-01	0.572760E+00
2001	0.281000E-01	0.774422E-01	-0.101376E+01
2002	0.102700E+00	0.333145E-01	0.112582E+01
2003	0.542000E-01	0.662436E-01	-0.200658E+00

Survey Index: 18 Tag: Spr AGE = 11 - 11
 Time = JAN-1 Type = NUMBER
 Catchability = 0.113695E-03 Variance = 0.315058E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.526400E+00	0.825415E+00	-0.449825E+00
1983	0.670900E+00	0.537523E+00	0.221648E+00
1984	0.578700E+00	0.436989E+00	0.280876E+00
1985	0.833300E+00	0.332638E+00	0.918338E+00
1986	0.403000E+00	0.231876E+00	0.552735E+00
1987	0.186200E+00	0.130897E+00	0.352413E+00
1988	0.112500E+00	0.138514E+00	-0.208022E+00
1989	0.185700E+00	0.143569E+00	0.257319E+00
1990	0.511000E-01	0.102310E+00	-0.694227E+00
1991	0.103500E+00	0.155893E+00	-0.409598E+00
1992	0.811000E-01	0.241825E+00	-0.109253E+01
1993	0.105500E+00	0.149822E+00	-0.350734E+00
1994	0.320000E-01	0.888716E-01	-0.102146E+01
1995	0.665000E-01	0.423526E-01	0.451172E+00
1996	0.320000E-01	0.204749E-01	0.446535E+00
1997	N/A	0.127458E-01	N/A
1998	0.284000E-01	0.203523E-01	0.333198E+00
1999	N/A	0.206274E-01	N/A
2000	N/A	0.609106E-01	N/A
2001	N/A	0.758869E-01	N/A
2002	0.803000E-01	0.745089E-01	0.748516E-01
2003	0.960000E-01	0.685140E-01	0.337310E+00

Survey Index: 21 Tag: Aut AGE = 3
 Time = JAN-1 Type = NUMBER
 Catchability = 0.834330E-05 Variance = 0.129941E+01
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.300000E-01	0.128522E+00	-0.145490E+01
1983	N/A	0.147676E+00	N/A
1984	0.109000E-01	0.136329E+00	-0.252631E+01
1985	N/A	0.638295E-01	N/A
1986	0.860000E-02	0.451718E-01	-0.165871E+01
1987	N/A	0.250387E-01	N/A
1988	0.229000E-01	0.812628E-01	-0.126655E+01
1989	N/A	0.518444E-01	N/A
1990	0.182000E-01	0.556493E-01	-0.111765E+01
1991	0.136700E+00	0.736788E-01	0.618073E+00
1992	0.177400E+00	0.119663E+00	0.393729E+00
1993	0.109100E+00	0.806758E-01	0.301826E+00
1994	0.154200E+00	0.113553E+00	0.305981E+00
1995	0.287200E+00	0.105718E+00	0.999404E+00
1996	0.764400E+00	0.132478E+00	0.175268E+01
1997	0.260700E+00	0.168751E+00	0.434945E+00
1998	0.978800E+00	0.247453E+00	0.137510E+01
1999	0.519800E+00	0.357958E+00	0.373029E+00
2000	0.117810E+01	0.564429E+00	0.735843E+00
2001	0.718800E+00	0.489786E+00	0.383615E+00
2002	0.210400E+00	0.246983E+00	-0.160311E+00
2003	0.274600E+00	0.164862E+00	0.510209E+00

Survey Index: 22 Tag: Aut AGE = 4
 Time = JAN-1 Type = NUMBER
 Catchability = 0.307342E-04 Variance = 0.859048E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.240000E+00	0.374173E+00	-0.444079E+00
1983	0.575000E-01	0.402065E+00	-0.194483E+01
1984	0.507300E+00	0.458620E+00	0.100880E+00
1985	0.932000E-01	0.428069E+00	-0.152454E+01
1986	0.588000E-01	0.198857E+00	-0.121844E+01
1987	N/A	0.142569E+00	N/A
1988	N/A	0.787524E-01	N/A
1989	0.724800E+00	0.240565E+00	0.110291E+01
1990	0.817000E-01	0.151654E+00	-0.618547E+00
1991	0.380200E+00	0.166652E+00	0.824792E+00
1992	0.661000E+00	0.221029E+00	0.109546E+01
1993	0.258500E+00	0.368026E+00	-0.353258E+00
1994	0.544200E+00	0.243913E+00	0.802507E+00
1995	0.531500E+00	0.358834E+00	0.392844E+00
1996	0.162390E+01	0.317288E+00	0.163277E+01
1997	0.785100E+00	0.416561E+00	0.633778E+00
1998	0.522300E+00	0.531187E+00	-0.168721E-01
1999	0.136260E+01	0.775141E+00	0.564105E+00
2000	0.151370E+01	0.113036E+01	0.292025E+00
2001	0.140820E+01	0.178663E+01	-0.238017E+00
2002	0.952400E+00	0.155092E+01	-0.487618E+00
2003	0.430900E+00	0.781916E+00	-0.595871E+00

Survey Index: 23 Tag: Aut AGE = 5
 Time = JAN-1 Type = NUMBER
 Catchability = 0.385675E-04 Variance = 0.159406E+01
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.440000E+00	0.368823E+00	0.176459E+00
1983	0.131000E-01	0.366129E+00	-0.333037E+01
1984	0.159570E+01	0.386208E+00	0.141869E+01
1985	0.943400E+00	0.443002E+00	0.755916E+00
1986	0.757000E-01	0.420355E+00	-0.171432E+01
1987	0.510000E-01	0.201311E+00	-0.137303E+01
1988	0.109000E-01	0.147508E+00	-0.260512E+01
1989	0.546000E-01	0.800613E-01	-0.382759E+00
1990	0.300600E+00	0.244248E+00	0.207598E+00
1991	0.506800E+00	0.141172E+00	0.127813E+01
1992	0.329000E+00	0.165505E+00	0.687054E+00
1993	0.223500E+00	0.193960E+00	0.141761E+00
1994	0.777000E+00	0.333033E+00	0.847200E+00
1995	0.165200E+00	0.227698E+00	-0.320862E+00
1996	0.857600E+00	0.374336E+00	0.828983E+00
1997	0.198750E+01	0.323963E+00	0.181400E+01
1998	0.870900E+00	0.412121E+00	0.748211E+00
1999	0.465200E+00	0.553824E+00	-0.174379E+00
2000	0.104380E+01	0.818409E+00	0.243260E+00
2001	0.174610E+01	0.121063E+01	0.366244E+00
2002	0.315550E+01	0.191742E+01	0.498164E+00
2003	0.147540E+01	0.164834E+01	-0.110838E+00

Survey Index: 24 Tag: Aut AGE = 6
 Time = JAN-1 Type = NUMBER
 Catchability = 0.410818E-04 Variance = 0.147441E+01
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.610000E+00	0.321654E+00	0.639984E+00
1983	0.270000E-01	0.292242E+00	-0.238174E+01
1984	0.757500E+00	0.277903E+00	0.100275E+01
1985	0.991400E+00	0.278055E+00	0.127130E+01
1986	0.610200E+00	0.325719E+00	0.627751E+00
1987	0.266400E+00	0.327751E+00	-0.207255E+00
1988	0.229000E-01	0.166803E+00	-0.198568E+01
1989	0.115000E-01	0.125184E+00	-0.238744E+01
1990	0.910000E-02	0.709203E-01	-0.205328E+01
1991	0.219100E+00	0.182033E+00	0.185343E+00
1992	0.289900E+00	0.966616E-01	0.109832E+01
1993	0.537000E-01	0.119121E+00	-0.796728E+00
1994	0.219300E+00	0.124040E+00	0.569839E+00
1995	0.395100E+00	0.200787E+00	0.676893E+00
1996	0.472100E+00	0.169215E+00	0.102602E+01
1997	0.138600E+01	0.297295E+00	0.153945E+01
1998	0.769600E+00	0.258400E+00	0.109136E+01
1999	0.303000E+00	0.337395E+00	-0.107522E+00
2000	0.599800E+00	0.466076E+00	0.252247E+00
2001	0.673800E+00	0.732981E+00	-0.841859E-01
2002	0.188570E+01	0.107414E+01	0.562782E+00
2003	0.997100E+00	0.171139E+01	-0.540208E+00

Survey Index: 25 Tag: Aut AGE = 7
 Time = JAN-1 Type = NUMBER
 Catchability = 0.461380E-04 Variance = 0.872772E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.460000E+00	0.197896E+00	0.843485E+00
1983	0.757000E-01	0.248007E+00	-0.118668E+01
1984	0.547900E+00	0.215362E+00	0.933772E+00
1985	0.604900E+00	0.194538E+00	0.113444E+01
1986	0.683900E+00	0.186356E+00	0.130015E+01
1987	0.353200E+00	0.196860E+00	0.584540E+00
1988	0.457000E-01	0.262204E+00	-0.174702E+01
1989	0.355000E-01	0.133154E+00	-0.132197E+01
1990	0.211000E-01	0.107587E+00	-0.162903E+01
1991	0.240000E-01	0.576475E-01	-0.876293E+00
1992	0.144900E+00	0.151400E+00	-0.438788E-01
1993	0.605000E-01	0.536840E-01	0.119528E+00
1994	0.578000E-01	0.760874E-01	-0.274893E+00
1995	0.370000E-01	0.652354E-01	-0.567085E+00
1996	0.229000E+00	0.121198E+00	0.636298E+00
1997	0.441200E+00	0.105603E+00	0.142981E+01
1998	0.383200E+00	0.232213E+00	0.500900E+00
1999	0.165100E+00	0.189455E+00	-0.137603E+00
2000	0.363500E+00	0.263784E+00	0.320650E+00
2001	0.588900E+00	0.401079E+00	0.384098E+00
2002	0.812900E+00	0.661601E+00	0.205946E+00
2003	0.532000E+00	0.978290E+00	-0.609163E+00

Survey Index: 26 Tag: Aut AGE = 8
 Time = JAN-1 Type = NUMBER
 Catchability = 0.772831E-04 Variance = 0.783392E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.270000E+00	0.212705E+00	0.238515E+00
1983	0.240900E+00	0.237770E+00	0.130787E-01
1984	0.444400E+00	0.244187E+00	0.598789E+00
1985	0.534700E+00	0.204563E+00	0.960829E+00
1986	0.482200E+00	0.171901E+00	0.103144E+01
1987	0.308500E+00	0.157178E+00	0.674343E+00
1988	0.192100E+00	0.171721E+00	0.112144E+00
1989	0.214800E+00	0.279352E+00	-0.262766E+00
1990	0.166000E-01	0.137798E+00	-0.211639E+01
1991	0.229000E-01	0.135524E+00	-0.177801E+01
1992	0.673000E-01	0.662686E-01	0.154447E-01
1993	N/A	0.167087E+00	N/A
1994	0.216000E-01	0.350350E-01	-0.483654E+00
1995	0.106100E+00	0.510214E-01	0.732136E+00
1996	N/A	0.338226E-01	N/A
1997	0.655000E-01	0.733527E-01	-0.113229E+00
1998	0.329400E+00	0.798318E-01	0.141735E+01
1999	0.110300E+00	0.219947E+00	-0.690183E+00
2000	0.275400E+00	0.187507E+00	0.384407E+00
2001	0.229300E+00	0.264115E+00	-0.141352E+00
2002	0.611800E+00	0.456735E+00	0.292302E+00
2003	0.330700E+00	0.801442E+00	-0.885202E+00

Survey Index: 27 Tag: Aut AGE = 9
 Time = JAN-1 Type = NUMBER
 Catchability = 0.796591E-04 Variance = 0.653415E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.260000E+00	0.167464E+00	0.439916E+00
1983	0.132300E+00	0.140448E+00	-0.597687E-01
1984	0.837000E-01	0.139120E+00	-0.508099E+00
1985	0.309600E+00	0.107029E+00	0.106218E+01
1986	0.269700E+00	0.901385E-01	0.109596E+01
1987	0.159700E+00	0.912531E-01	0.559660E+00
1988	0.714000E-01	0.756272E-01	-0.575180E-01
1989	0.484000E-01	0.679774E-01	-0.339676E+00
1990	0.837000E-01	0.182950E+00	-0.781973E+00
1991	0.229000E-01	0.876110E-01	-0.134177E+01
1992	0.591000E-01	0.102209E+00	-0.547791E+00
1993	N/A	0.439604E-01	N/A
1994	0.813000E-01	0.105182E+00	-0.257544E+00
1995	N/A	0.166705E-01	N/A
1996	N/A	0.255801E-01	N/A
1997	0.648000E-01	0.108145E-01	0.179042E+01
1998	N/A	0.218146E-01	N/A
1999	0.425000E-01	0.435674E-01	-0.248060E-01
2000	0.499000E-01	0.138864E+00	-0.102348E+01
2001	0.151600E+00	0.906809E-01	0.513899E+00
2002	0.158700E+00	0.127357E+00	0.220019E+00
2003	0.148400E+00	0.310925E+00	-0.739639E+00

Survey Index: 28 Tag: Aut AGE = 10
 Time = JAN-1 Type = NUMBER
 Catchability = 0.913192E-04 Variance = 0.955223E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.180000E+00	0.100554E+00	0.582264E+00
1983	0.145000E-01	0.131517E+00	-0.220499E+01
1984	0.137200E+00	0.765969E-01	0.582883E+00
1985	0.148800E+00	0.787050E-01	0.636897E+00
1986	0.102600E+00	0.547275E-01	0.628471E+00
1987	0.111800E+00	0.542217E-01	0.723631E+00
1988	N/A	0.496592E-01	N/A
1989	0.462000E-01	0.408898E-01	0.122099E+00
1990	0.783000E-01	0.376847E-01	0.731294E+00
1991	0.250000E-01	0.152314E+00	-0.180707E+01
1992	0.296000E-01	0.618245E-01	-0.736525E+00
1993	0.189000E-01	0.858218E-01	-0.151311E+01
1994	N/A	0.249822E-01	N/A
1995	0.425000E-01	0.584672E-01	-0.318961E+00
1996	0.107000E-01	0.826607E-02	0.258085E+00
1997	0.373000E-01	0.727271E-02	0.163486E+01
1998	N/A	0.360419E-02	N/A
1999	0.123000E-01	0.962394E-02	0.245346E+00
2000	0.370000E-01	0.217946E-01	0.529256E+00
2001	0.494000E-01	0.845414E-01	-0.537291E+00
2002	0.576000E-01	0.363686E-01	0.459818E+00
2003	0.711000E-01	0.723163E-01	-0.169624E-01

Survey Index: 35 Tag: Aut AGE = 11 - 11
 Time = JAN-1 Type = NUMBER
 Catchability = 0.111457E-03 Variance = 0.567455E+00
 Residual = LN(Observed) - LN(Predicted)

Year	Observed	Predicted	Residual
1982	0.103000E+01	0.809166E+00	0.241311E+00
1983	0.409400E+00	0.526941E+00	-0.252396E+00
1984	0.625800E+00	0.428386E+00	0.379005E+00
1985	0.746700E+00	0.326090E+00	0.828490E+00
1986	0.471800E+00	0.227311E+00	0.730237E+00
1987	0.330500E+00	0.128320E+00	0.946081E+00
1988	0.116300E+00	0.135788E+00	-0.154919E+00
1989	0.233300E+00	0.140742E+00	0.505394E+00
1990	0.865000E-01	0.100296E+00	-0.147984E+00
1991	0.980000E-01	0.152824E+00	-0.444320E+00
1992	0.799000E-01	0.237065E+00	-0.108756E+01
1993	0.896000E-01	0.146872E+00	-0.494207E+00
1994	0.867000E-01	0.871221E-01	-0.485638E-02
1995	0.566000E-01	0.415188E-01	0.309862E+00
1996	0.635000E-01	0.200718E-01	0.115172E+01
1997	0.330000E-01	0.124948E-01	0.971192E+00
1998	0.195000E-01	0.199516E-01	-0.228942E-01
1999	N/A	0.202214E-01	N/A
2000	0.940000E-02	0.597115E-01	-0.184882E+01
2001	0.260000E-01	0.743929E-01	-0.105126E+01
2002	0.556000E-01	0.730420E-01	-0.272852E+00
2003	0.507000E-01	0.671652E-01	-0.281229E+00

Bootstrap Summary Report

Number of Bootstrap Repetitions Requested = 1000
 Number of Bootstrap Repetitions Completed = 1000
 Bootstrap Output Variable: Stock Estimates (2003)

	NLLS Estimate	Bootstrap Mean	Bootstrap Std Error	C.V. For NLLS Soln.
N 3	19760.	27028.	22590.	0.8358
N 4	25441.	29186.	14151.	0.4849
N 5	42739.	46607.	18703.	0.4013
N 6	41658.	44306.	15511.	0.3501
N 7	21204.	22124.	7147.	0.3231
N 8	10370.	10947.	3528.	0.3222
N 9	3903.	4039.	1305.	0.3231
N 10	792.	863.	358.	0.4152

	Bias Estimate	Bias Std. Error	Per Cent Bias	NLLS Estimate Corrected For Bias	C.V. For Corrected Estimate
N 3	7268.	750.	36.7840	12491.	1.8085
N 4	3745.	463.	14.7198	21696.	0.6522
N 5	3868.	604.	9.0507	38871.	0.4812
N 6	2648.	498.	6.3573	39010.	0.3976
N 7	920.	228.	4.3405	20283.	0.3524
N 8	577.	113.	5.5662	9793.	0.3602
N 9	136.	41.	3.4768	3767.	0.3464
N 10	71.	12.	8.9993	721.	0.4973

	LOWER 80. % CI	UPPER 80. % CI
N 3	7136.	54576.
N 4	13256.	47904.
N 5	25267.	71594.
N 6	26357.	65268.
N 7	13625.	31371.
N 8	6754.	15709.
N 9	2506.	5701.
N 10	448.	1352.

Bootstrap Output Variable: Catchability Estimates

	NLLS Estimate	Bootstrap Mean	Bootstrap Std Error	C.V. For NLLS Soln.
Q 4	0.134450E-04	0.139774E-04	0.357412E-05	0.2557
Q 5	0.323264E-04	0.324874E-04	0.499383E-05	0.1537
Q 6	0.429554E-04	0.432622E-04	0.531046E-05	0.1228
Q 7	0.430079E-04	0.436599E-04	0.596464E-05	0.1366
Q 8	0.508706E-04	0.514536E-04	0.691624E-05	0.1344
Q 9	0.656138E-04	0.666622E-04	0.132064E-04	0.1981
Q 10	0.788325E-04	0.797692E-04	0.159116E-04	0.1995
Q 11	0.836507E-04	0.852533E-04	0.174415E-04	0.2046
Q 18	0.113695E-03	0.115577E-03	0.152687E-04	0.1321
Q 21	0.834330E-05	0.863377E-05	0.236581E-05	0.2740
Q 22	0.307342E-04	0.311082E-04	0.644467E-05	0.2072
Q 23	0.385675E-04	0.400842E-04	0.104468E-04	0.2606
Q 24	0.410818E-04	0.422332E-04	0.104730E-04	0.2480
Q 25	0.461380E-04	0.468698E-04	0.937549E-05	0.2000
Q 26	0.772831E-04	0.776712E-04	0.151170E-04	0.1946
Q 27	0.796591E-04	0.807409E-04	0.146354E-04	0.1813
Q 28	0.913192E-04	0.930613E-04	0.208653E-04	0.2242
Q 35	0.111457E-03	0.113312E-03	0.181145E-04	0.1599

	Bias Estimate	Bias Std. Error	Per Cent Bias	NLLS Estimate Corrected For Bias	C.V. For Corrected Estimate
Q 4	0.5324E-06	0.1143E-06	3.9602	0.1291E-04	0.2768
Q 5	0.1610E-06	0.1580E-06	0.4979	0.3217E-04	0.1553
Q 6	0.3068E-06	0.1682E-06	0.7142	0.4265E-04	0.1245
Q 7	0.6520E-06	0.1897E-06	1.5160	0.4236E-04	0.1408
Q 8	0.5831E-06	0.2195E-06	1.1461	0.5029E-04	0.1375
Q 9	0.1048E-05	0.4189E-06	1.5978	0.6457E-04	0.2045
Q 10	0.9368E-06	0.5040E-06	1.1883	0.7790E-04	0.2043
Q 11	0.1603E-05	0.5539E-06	1.9158	0.8205E-04	0.2126
Q 18	0.1881E-05	0.4865E-06	1.6548	0.1118E-03	0.1366
Q 21	0.2905E-06	0.7538E-07	3.4815	0.8053E-05	0.2938
Q 22	0.3740E-06	0.2041E-06	1.2168	0.3036E-04	0.2123
Q 23	0.1517E-05	0.3338E-06	3.9327	0.3705E-04	0.2820
Q 24	0.1151E-05	0.3332E-06	2.8027	0.3993E-04	0.2623
Q 25	0.7318E-06	0.2974E-06	1.5862	0.4541E-04	0.2065
Q 26	0.3882E-06	0.4782E-06	0.5023	0.7689E-04	0.1966
Q 27	0.1082E-05	0.4641E-06	1.3581	0.7858E-04	0.1863
Q 28	0.1742E-05	0.6621E-06	1.9078	0.8958E-04	0.2329
Q 35	0.1855E-05	0.5758E-06	1.6642	0.1096E-03	0.1653

	LOWER	UPPER
	80. % CI	80. % CI
Q 4	0.972636E-05	0.189646E-04
Q 5	0.261931E-04	0.392517E-04
Q 6	0.364525E-04	0.502755E-04
Q 7	0.361255E-04	0.513419E-04
Q 8	0.430078E-04	0.606003E-04
Q 9	0.506999E-04	0.842534E-04
Q 10	0.594134E-04	0.998205E-04
Q 11	0.644053E-04	0.107803E-03
Q 18	0.968436E-04	0.134914E-03
Q 21	0.587186E-05	0.118302E-04
Q 22	0.234795E-04	0.397340E-04
Q 23	0.278363E-04	0.546203E-04
Q 24	0.295456E-04	0.568077E-04
Q 25	0.355904E-04	0.591617E-04
Q 26	0.582349E-04	0.965514E-04
Q 27	0.627337E-04	0.997029E-04
Q 28	0.665440E-04	0.120632E-03
Q 35	0.911700E-04	0.136921E-03

Bootstrap Output Variable: Fishing Mortality (2002)

	NLLS Estimate	Bootstrap Mean	Bootstrap Std Error	C.V. For NLLS Soln.
AGE 3	0.0015	0.0016	0.000877	0.5331
AGE 4	0.0161	0.0175	0.007794	0.4466
AGE 5	0.0268	0.0284	0.010337	0.3638
AGE 6	0.0595	0.0629	0.019980	0.3179
AGE 7	0.1741	0.1813	0.057853	0.3192
AGE 8	0.2648	0.2797	0.087777	0.3138
AGE 9	0.5526	0.5753	0.193135	0.3357
AGE 10	0.4087	0.4275	0.101083	0.2365
AGE 11	0.4087	0.4275	0.101083	0.2365

	Bias Estimate	Bias Std. Error	Per Cent Bias	NLLS Estimate Corrected For Bias	C.V. For Corrected Estimate
AGE 3	0.000153	0.000028	10.2505	0.0013	0.6548
AGE 4	0.001341	0.000250	8.3254	0.0148	0.5277
AGE 5	0.001577	0.000331	5.8754	0.0253	0.4092
AGE 6	0.003316	0.000640	5.5688	0.0562	0.3554
AGE 7	0.007176	0.001844	4.1220	0.1669	0.3466
AGE 8	0.014858	0.002815	5.6101	0.2500	0.3511
AGE 9	0.022717	0.006150	4.1112	0.5298	0.3645
AGE 10	0.018787	0.003251	4.5969	0.3899	0.2592
AGE 11	0.018787	0.003251	4.5969	0.3899	0.2592

	LOWER	UPPER
	80. % CI	80. % CI
AGE 3	0.000793	0.002856
AGE 4	0.009644	0.027095
AGE 5	0.017206	0.042028
AGE 6	0.040591	0.090833
AGE 7	0.118230	0.255170
AGE 8	0.188377	0.386366
AGE 9	0.358831	0.831260
AGE 10	0.311489	0.560158
AGE 11	0.311489	0.560158

Bootstrap Output Variable: Average F (2002) AGES 8 - 9

	NLLS Estimate	Bootstrap Mean	Bootstrap Std Error	C.V. For NLLS Soln.
AVG F	0.4087	0.4275	0.101083	0.2365
N WTD	0.3261	0.3329	0.075870	0.2279
B WTD	0.3344	0.3405	0.075636	0.2221
C WTD	0.3601	0.3776	0.081381	0.2155

	Bias Estimate	Bias Std. Error	Per Cent Bias	NLLS Estimate Corrected For Bias	C.V. For Corrected Estimate
AVG F	0.018787	0.003251	4.5969	0.3899	0.2592
N WTD	0.006831	0.002409	2.0948	0.3193	0.2376
B WTD	0.006045	0.002399	1.8075	0.3284	0.2303
C WTD	0.017461	0.002632	4.8483	0.3427	0.2375

	LOWER 80. % CI	UPPER 80. % CI
AVG F	0.311489	0.560158
N WTD	0.243401	0.425261
B WTD	0.251145	0.433677
C WTD	0.286044	0.485475

Bootstrap Output Variable: Biomass

JAN-1 Biomass (2003) Mean Biomass & SSB (2002)

	NLLS Estimate	Bootstrap Mean	Bootstrap Std Error	C.V. For NLLS Soln.
JAN-1	56452.	61054.	11000.	0.1802
MEAN	45192.	48178.	8243.	0.1711
SSB	18296.	19179.	2851.	0.1486

	Bias Estimate	Bias Std. Error	Per Cent Bias	NLLS Estimate Corrected For Bias	C.V. For Corrected Estimate
JAN-1	4602.	377.	8.1521	51850.	0.2121
MEAN	2986.	277.	6.6075	42206.	0.1953
SSB	884.	94.	4.8315	17412.	0.1637

	LOWER 80. % CI	UPPER 80. % CI
JAN-1	46785.	75591.
MEAN	37705.	58929.
SSB	15603.	22969.

Plus Group Diagnostic Report

Calculation Method Selected = Backward

Year	Population Backward	Population Forward	F Forward	F Backward	Ratio
1982	7260.	7260.	0.265727	0.265728	1.000002
1983	4728.	5517.	0.393693	0.477042	1.211710
1984	3844.	3973.	0.620564	0.649539	1.046692
1985	2926.	2215.	1.056192	0.685293	0.648833
1986	2039.	1037.	1.484863	0.507203	0.341583
1987	1151.	513.	4.265155	0.605723	0.142017
1988	1218.	694.	2.328713	0.740857	0.318140
1989	1263.	404.	2.328713	0.350167	0.150369
1990	900.	274.	1.164614	0.237924	0.204294
1991	1372.	354.	1.475319	0.225446	0.152811
1992	2128.	1216.	0.403839	0.211096	0.522722
1993	1318.	1171.	0.441947	0.381925	0.864187
1994	782.	1200.	0.342158	0.585644	1.711616
1995	373.	865.	0.216568	0.596709	2.755296
1996	180.	903.	0.145252	1.105538	7.611176
1997	113.	698.	0.114260	1.076386	9.420518
1998	181.	560.	0.145078	0.536699	3.699394
1999	192.	437.	0.143771	0.362347	2.520317
2000	549.	392.	0.862729	0.535635	0.620861
2001	768.	356.	2.328713	0.557471	0.239390
2002	738.	527.	0.539015	0.353809	0.656399
2003	660.	479.	N/A	N/A	

Warning **** Infeasible Mass Balance in Plus Group

Year = 1991

Year = 2000

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance: All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material lifted from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

Organization: Manuscripts must have an abstract and table of contents, and — if applicable — lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area"/"Experimental Apparatus," "Methods," "Results," "Discussion" and/or "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style: The CRD series is obligated to conform with the style contained in the current edition of the *United States Government Printing Office Style Manual*. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the *CBE Style Manual*. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the International Standardization Organization's guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation: Type a clean/neat, single-spaced version of the document. The document must be paginated continuously from beginning to end and must have a "Table of Contents." Begin the preliminary pages of the document — always the "Table of Contents" — with page "iii." Begin the body of the document — normally the "Introduction" — with page "1," and continuously paginate all pages including tables, figures, appendices, and indices. You can insert blank pages as appropriate throughout the document, but account for them in your pagination (*e.g.*, if your last figure ends on an odd-numbered/right-hand page such as "75," and if your next page is the first page of an appendix, then you would normally insert a blank page after the last figure, and paginate the first page of the appendix as "77" to make it begin on an odd-numbered/right-hand page also). Forward the final version to the Editorial Office as both a paper copy and electronically (*i.e.*, e-mail attachment, 3.5-inch floppy disk, high-density zip disk, or CD). For purposes of publishing the CRD series only, the use of Microsoft Word is preferable to the use of Corel WordPerfect.

Production and Distribution: The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages (pages "i" and "ii") of the document, then combine those covers and preliminary pages with the text that you have supplied. The document will then be issued online.

Paper copies of the four covers and two preliminary pages will be sent to the sole/senior NEFSC author should he/she wish to prepare some paper copies of the overall document as well. The Editorial Office will only produce four paper copies (*i.e.*, three copies for the NEFSC's libraries and one copy for its own archives) of the overall document.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the online version of the document. The sole/senior NEFSC author of the document will receive a list of those so notified.

Research Communications Unit
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "planning, developing, and managing multidisciplinary programs of basic and applied research to: 1) better understand the living marine resources (including marine mammals) of the Northwest Atlantic, and the environmental quality essential for their existence and continued productivity; and 2) describe and provide to management, industry, and the public, options for the utilization and conservation of living marine resources and maintenance of environmental quality which are consistent with national and regional goals and needs, and with international commitments." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Those media are in four categories:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, but no technical or copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. There is no scientific review, nor any technical or copy editing, of this report.

The Shark Tagger -- This newsletter is an annual summary of tagging and recapture data on large pelagic sharks as derived from the NMFS's Cooperative Shark Tagging Program; it also presents information on the biology (movement, growth, reproduction, etc.) of these sharks as subsequently derived from the tagging and recapture data. There is internal scientific review, but no technical or copy editing, of this newsletter.

OBTAINING A COPY: To obtain a copy of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, or to subscribe to the *Resource Survey Report* or the *The Shark Tagger*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2228) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.