

reinforce smoking abstinence (Morgan 1981). Many smokers, particularly women, are concerned about potential weight gain as a result of smoking cessation, and such programs can address these concerns (Ellis 1980).

There are also potential disadvantages of multiple risk factor reduction programs. They may be difficult to implement because staff expertise is required in multiple areas and because some risk factors, such as smoking, may not be relevant for all participants. In addition, multiple risk factor reduction programs must present a large amount of complex information, usually in a limited time, and consequently the amount of attention devoted to a given risk factor such as smoking must often be less than is the case in single modality programs.

Two main types of multiple risk factor reduction programs have involved smoking cessation. The first is large-scale clinical trials for the prevention of coronary heart disease. The Belgian and British WHO studies reported by Kornitzer and Dramaix and colleagues (1980) and by Rose and colleagues (1980) were conducted solely in industrial settings and were discussed in detail in the 1983 Report of the Surgeon General (US DHHS 1983). These studies are well designed and have collected multiple dependent variables, including indices of overall health risk or morbidity and mortality statistics.

The other main type of multiple risk factor reduction program that has been developed is worksite wellness programs conducted by large companies for their employees. Examples include the STAY-WELL program of the Control Data Corporation (Naditch 1984), the Live for Life program of Johnson and Johnson (Nathan 1984), and programs offered by IBM, the Campbell Soup Company, and the Ford Motor Company (Parkinson et al. 1982; Ware and Block 1982). Unfortunately, the outcomes of almost all industry-sponsored programs reported to date are difficult to interpret owing to varying methods of reporting results, difficulties in following subjects, and lack of objective measures of smoking status. Reports of company wellness programs with more than anecdotal data on smoking modification results (e.g., Grove et al. 1979; Sorman 1979) are summarized in Table 1.

Cessation rates in multiple risk factor reduction programs in worksites have ranged from 7 to 33 percent at followup. Many of these rates are lower than those typically reported in other worksite smoking studies and are not consistently better than comparison conditions in controlled studies (Kornitzer, De Backer et al. 1980; Meyer and Henderson 1974). Interpretation of these data is problematic because of the lack of direct comparisons with smoking-cessation-only interventions, because subjects with multiple risk factors may be more recalcitrant than other subjects, and because

TABLE 4.—Organizational characteristics potentially affecting outcome of worksite smoking programs

(1)	Size of worksite	(7)	Union/management relations
(2)	Current worksite smoking policies	(8)	Percent of smokers in the worksite
(3)	Degree of management support for program	(9)	Growth oriented vs. consolidating climate of organization
(4)	History of health promotion efforts in the worksite	(10)	Rank and sociometric standing of primary contact person
(5)	Sex ratio of employees	(11)	Socioeconomic level of employees
(6)	Job stability/turnover		

the risk factor reduction programs reported in this section tend to be ongoing programs rather than one-shot smoking clinics.

Organizational Characteristics and Other Factors

Conducting outcome research in worksite settings involves a number of unique factors that may mediate or interact with program success. The organizational characteristics that may mediate program success are outlined in Table 4. Although this list is certainly not exhaustive, investigators should consider these factors when conducting worksite smoking programs. An example of the potential effects of organizational characteristics is the variability in outcome reported by Glasgow and Klesges and their colleagues (Glasgow et al. 1984, in press; Klesges et al. 1985). The basic treatment programs utilized in these studies were almost identical and were implemented by many of the same therapists. Yet, the 6-month abstinence rates in the different organizations ranged from 14 to 33 percent.

Few of these variables have been addressed in worksite smoking studies. Bishop and Fisher (1984) have conducted similar multilevel smoking cessation programs in a number of different organizations, ranging in size from 200 to 6,000 employees. They reported substantially lower participation rates in large companies, a finding that is consistent with results of studies of worksite weight loss programs (Brownell et al., in press). Also, the studies outlined in Tables 1 through 3 suggest that the highest cessation rates are obtained in smaller worksites. Taken together, these trends suggest that different interventions and different ways to assure participation need to be developed for large corporations. The problem may be one of

implementation, not design. Company policy regarding vesting responsibility in division leadership may be a critical variable.

In terms of the second variable in Table 4, worksite smoking policies, it is important to emphasize that smoking cessation groups are but one way to influence rates of worksite cigarette smoking (Bennett and Levy 1980). Although there have certainly been more reports on cessation programs than on other approaches to occupational smoking control, evaluations of alternative procedures are beginning to appear. In particular, Dawley and colleagues (Dawley and Baldwin 1983; Dawley and Burton, in press; Dawley et al. 1980) and Jason and colleagues (Jason and Liotta 1982; Jason and Clay 1978; Jason and Savio 1978) have studied the effects of no-smoking signs and requests not to smoke. These studies indicate that the posting of nonsmoking signs and the establishment of nonsmoking areas temporarily reduce smoking rates, but that active enforcement of such policies is necessary to produce substantial or lasting decrements in smoking behavior (Dawley et al. 1980; Jason and Liotta 1982; Jason and Savio 1978). One caveat to be kept in mind in evaluating the effects of worksite smoking restrictions is that workers may "compensate" by smoking more during breaks and after work (Meade and Wald 1977). Evaluations of the effectiveness of smoking restrictions should therefore assess smoking rates during both work and nonwork hours and include objective measures of smoking exposure.

Dawley and colleagues subdivided smoking modification efforts into three categories: smoking control (limiting or restricting smoking to designated areas); smoking discouragement (educational efforts to encourage people to stop smoking); and smoking cessation (more formal treatment programs). They also suggested that "worksite smoking cessation programs operate most effectively when offered in conjunction with worksite smoking control and discouragement efforts" (Dawley et al. 1984, p. 329), a highly testable hypothesis that has yet to be experimentally investigated.

The potential to use modifications of the work environment to aid in smoking cessation, including restricting smoking, removing cigarette machines, and altering work rules or situations that promote smoking, make the worksite more than simply a location for cessation interventions. The elimination of environmental supports for smoking, alteration of the smoker's self-image, changing the perception of the smoker among peers, and revising the social norms about smoking in the worksite may all provide a powerful motivation for the smoker to quit and support the successful maintenance of cessation. These changes in the workplace environment and attitudes may be more important than the components of the behavioral intervention used to get workers to quit, and experimental verification of the impact of these changes would provide a useful

guide for the structuring of future comprehensive worksite interventions. Because it would probably be unlikely that researchers would gain access to experimental manipulation of some of the more controversial aspects of guidelines (hiring policies and penalties for smoking), opportunities that may arise to study such changes in noncontrolled research would be worth pursuing.

Few data have been collected on the other variables listed in Table 4. Research on worksite smoking programs should at least provide descriptive information to determine how these variables affect program success. The fit between organizational and program characteristics has been neglected in past occupational smoking control research. It is hoped that future research will be able to identify the types of programs that are most effective in each different worksite setting.

Implementation of Worksite Smoking Programs

This section focuses on two major classes of implementation issues: recruitment procedures and characteristics of intervention programs.

Promotion and Recruitment

The initial contact with a worksite can prove critical to the success of a project. It is generally recommended that the initial meeting be with the chief executive of the organization (Klesges and Glasgow 1985). Although this officer typically does not coordinate the program, support from top-level management appears to be important in program recruitment and implementation (Grove et al. 1979). Another method of enhancing participation and organizational involvement is the formation of a steering committee (Bishop and Fisher 1984; Stachnik and Stoffelmayr 1981) composed of key representatives from both labor and management. Employees should perceive that the program is voluntary and that they have input into its implementation. Steering committees of this kind may be particularly important in large worksites with unionized employees. Management support appears to be quite important to the success of the committee (Bishop and Fisher 1984).

Upon securing permission to offer a program, it is helpful to conduct a brief worksite needs assessment (e.g., Heckler 1980; Kanzler et al. 1976; Klesges and Glasgow 1985). The survey can be used to determine (1) the number and characteristics of smokers in the worksite, (2) the number of smokers potentially interested in participating, and (3) preferences concerning the types of programs that might be offered (e.g., self-help versus group meetings; abstinence versus reduced smoking) and the most convenient times for meetings to be scheduled.

During the recruitment phase, information about the program should come from a variety of sources, such as posters, memos, and brochures. Advertising experts recommend providing multiple exposures to a "product" (in this case, a smoking program) to promote attitude change and to convince participants to take action regarding the product (Sawyer 1981). Promotional materials should include information about the cost of a program, stress that participation is voluntary and individual results are confidential, and counter possible misconceptions (e.g., "I have to quit at the first session"; "I'll lose my job if I don't participate"). It is helpful if at least one memo or announcement comes from top management. At this stage, human resources or personnel directors can be extremely useful in suggesting the best ways to promote the program in their particular setting. Involving the local media may also increase the credibility of the program as well as provide no-cost advertising for both the program and the worksite.

Prior to the actual implementation of a smoking program, some programs prepare worksites for health-behavior change (Andrews 1983; Bennett and Levy 1980; Ellis 1979; Grove et al. 1979; Heckler 1980). These preparatory procedures have ranged from prescreening health exams (Ellis 1979) to the initiation of smoking restrictions (Andrews 1983; Bennett and Levy 1980). Warnings of the impending restrictions with indications of the "target restriction date" allow workers to prepare for changes, such as by joining available programs. Although empirically untested, these recruitment procedures may help to convince employees to join smoking programs.

Program Characteristics

The advantages of occupational smoking control programs discussed earlier do not *automatically* or *necessarily* occur. Programs must be made convenient. Higher participation rates are usually found in programs that offer time off work (e.g., Klesges et al. 1985; Scott et al. 1983). Time off work for participation can be a double-edged sword, however. It may increase the number of smokers who participate primarily to be excused from their work stations, and it may also create demands among nonsmoking employees for time off work to attend other health-related classes. Generally, the benefits of conducting programs during work hours outweigh the potential costs, and if management is not willing to grant time off work, it may at least be possible to negotiate time sharing between employee and employer (e.g., 1/2 hour of work time, 1/2 hour during lunch hour or after work). Investigators should also be aware of the difficulties involved in scheduling group meetings in worksites where employees work rotating shifts, such as hospitals.

In addition to being convenient, programs should be attractive to participants. For example, allowing smokers to choose the type of

program (such as nicotine fading versus aversive smoking), the modality of intervention (self-help manual versus group meetings), the treatment goals (abstinence versus reduced smoking), and the type of group leader (health professional versus peer facilitator) may be helpful in attracting and retaining participants. Different components of a comprehensive program, such as physician advice, no-smoking policies, stop-smoking contests, or group meetings, may mutually reinforce each other. While these suggestions await empirical verification, providing smokers with a number of choices should serve to increase participation rates.

Finally, feedback on progress may serve to increase the magnitude of behavior change. For example, participants can be provided with frequent feedback on carbon monoxide levels as they reduce their smoking (e.g., Rand et al. 1984; Scott et al. 1983). Charts displaying the weekly progress of different groups can be posted in employee lunchrooms or lounges. Periodic progress reports to department supervisors might also be helpful. To avoid stigmatizing particular individuals, public feedback should be provided on progress by the group rather than by individuals.

There are a number of problems in conducting worksite smoking modification groups that should be avoided, or at least anticipated. Group composition is one such sensitive issue. For example, mixing high-ranking executives with production workers can almost eliminate group discussion. However, this may depend on the company's tradition of interaction among workers of different levels, on the skills of the group leaders, and so on. Scheduling difficulties can arise in settings where employees rotate shifts or travel frequently, or where meeting rooms are scarce or distant from work stations. One also needs to be sensitive to negativism or complaining, which can become contagious; the group's focus must be kept positive. A positive perspective is particularly important when conducting competition or incentive interventions in which certain individuals or groups must "lose." A more optimistic perspective that can be used to encourage participants is that everyone can win something by changing their smoking, so there are no losers.

Finally, Marlatt and Gordon's (1985) concept of stopping smoking as a "journey" can be quite helpful. On their journey, people may experience temporary setbacks or detours (relapses), but this should not prevent them from reaching their destination (abstinence). The presence of an ongoing program that makes it easy to try different options or to recycle a procedure can serve to reinforce this concept and to improve long-term results.

Recommendations for Future Research

A number of suggestions for the implementation of worksite smoking modification programs have been outlined. Given the limited nature of the data available, few of these guidelines are experimentally derived. Research is needed to empirically support or refute these recommendations. This section discusses needs for future research in the field of worksite smoking modification. Recommendations are made on both research methodology and substantive issues for further investigation.

Methodological Issues

Greater use should be made of creative experimental and quasi-experimental designs, as discussed by Cook and Campbell (1979). In particular, it should be possible to sequentially introduce an intervention or intervention components in different worksites using time-series or multiple baseline designs or to investigate the incremental effects of adding different strategies, such as physician messages, incentives, and social support procedures, to a basic treatment program.

Greater consistency across studies in the criteria used to define smoking status would substantially aid in the interpretation of results. Berglund and colleagues (1974) and Shipley and colleagues (1982) have provided guidelines for reporting outcomes of smoking cessation studies that should be more widely adopted. For calculating abstinence rates, a standard common denominator representing the number of subjects entering a program should be used across all points in time and any dropouts should be considered conservatively as smokers. In studies in which it is deemed important to evaluate reductions in smoking behavior (e.g., percent reduction in number of cigarettes smoked or nicotine content) in addition to the proportion of abstinent subjects, analyses should be conducted on nonabstinent subjects only. This procedure avoids confounding the results due to cessation with results due to changes in smoking rate or topography. Worksite programs should report cessation success as the fraction of the smokers in the workforce as well as the fraction who agreed to participate in the program.

Objective verification of smoking status is particularly important in programs involving financial incentives, competition between rival organizations, social pressure and support to quit, or controlled smoking instead of abstinence. Each of the biochemical measures of smoking exposure has its own advantages and limitations (Benowitz 1983; Pechacek et al. 1984).

Another methodological problem faced by occupational smoking modification programs concerns the consistency between units of assignment and units of analysis (Biglan and Ary 1985). Typically,

whole companies are assigned to treatment or control conditions, but results are analyzed using individual subjects as the unit. This creates interpretive problems because of the potential dependency among results of smokers within a given worksite (or treatment group). Although there are no easy answers to this dilemma, investigators should consider (1) conducting treatment in a sufficiently large number of companies that the worksite can be used as the unit of analysis; (2) utilizing hierarchical or nesting designs to separate the effects of worksite from intervention condition (Myers 1972); or (3) when feasible, assigning individuals within worksites to different treatment conditions.

Future research should pay greater attention to possible interactions between worksite and treatment variables. For example, interorganizational competition procedures may be highly effective in worksites where employees feel highly committed to the company, but ineffective in settings low in organizational commitment. Organizational and social network factors may also interact with, mediate, or enhance program impact.

More data also need to be collected on the "generalization" effects of worksite smoking modification programs. Employers may be more interested in program effects on employee morale, job satisfaction, and absenteeism than on health outcomes such as smoking status. Similarly, more information should be reported on the costs and health benefits of occupational smoking reduction programs. Progress in this area would be facilitated by a systematic review of and recommendation for procedures to be employed in determining the cost effectiveness and cost benefit of worksite smoking programs.

Substantive Areas

Three primary objectives need to be achieved by future research in worksite smoking modification. First, more research should be conducted on ways to increase participation and followthrough rates in worksite programs. For example, using various incentive procedures (e.g., paycheck bonuses versus team competition versus lotteries) might be expected to enhance participation. Further investigations are needed on the impact on participation rates of interventions such as quitting contests, self-help materials, or hotlines that do not require a large investment of time and effort by participants. The majority of worksite smoking studies to date have focused on group cessation programs, but surveys consistently indicate that most smokers are not interested in participating in such programs (US DHHS 1982; Schneider et al. 1984). For the reasons discussed earlier, renewed emphasis on physician stop-smoking messages is also indicated.

The second main content issue is how to enhance the outcome rates of worksite smoking modification programs. One approach to

this problem is to evaluate the utility of comprehensive intervention programs and environmental changes (no-smoking policies, cigarette machine removal, prominent no-smoking posters) with cessation groups. Other approaches are assessing the impact of multiple risk factor programs versus single modality programs and of ongoing, continuous intervention programs in place for a year or more versus one-time-only program offerings.

The final category of recommendations for future research involves investigating subject and therapist factors that affect treatment outcome (Klesges and Glasgow 1985; Orleans and Shipley 1982). Additional study is needed of the enrollment patterns and success rates of men versus women, white-collar workers versus blue-collar workers, and heavy smokers versus light smokers. Also, little is known about the characteristics of successful program leaders (e.g., ex-smoker coworkers versus professional group leaders).

Summary and Conclusions

1. Smoking modification and maintenance of nonsmoking status among initial quitters has the promise of being more successful in worksite programs than in clinic-based programs. Higher cessation rates in worksite programs are achieved with more intensive programs.
2. Incentives for nonsmoking appear to be associated with higher participation and better success rates. Further research is needed to specify the optimal types of incentive procedures.
3. Success of a worksite smoking program depends upon three primary factors: the characteristics of the intervention program, the characteristics of the organization in which the program is offered, and the interaction between these factors.
4. Research is needed on recruitment strategies and participation rates in worksite smoking programs and on the impact of interventions on the entire workforce of a company.
5. More investigations are needed on worksite characteristics associated with the success of occupational programs and on comprehensive programs including components such as quit-smoking contests, no-smoking policies, physician messages, and self-help materials in addition to smoking cessation clinics.
6. The implementation of broadly based health promotion efforts in the workplace should be encouraged, with smoking interventions representing a major component of the larger effort to improve health through a worksite focus.

References

- ABRAMS, D.B., PINTO, R.P., MONTI, P.M., JACOBUS, S., BROWN, R., ELDER, J.P. *Health Education vs. Intrapersonal Coping vs. Social Network Support for Relapse Prevention in a Worksite Smoking Cessation Program*. Paper presented at the annual meeting of the Society of Behavioral Medicine, New Orleans, March 1985.
- ANDREWS, J.L., Jr. Reducing smoking in the hospital: An effective model program. *Chest* 84(2):206-209, August 1983.
- BAUER, R.B. Bell Laboratories helps employees to quit smoking. *American Lung Association Bulletin* 64(6):11-14, July-August 1978.
- BENNETT, D., LEVY, B.S. Smoking policies and smoking cessation programs of large employers in Massachusetts. *American Journal of Public Health* 70(6):629-631, June 1980.
- BENOWITZ, N.L. The use of biologic fluid samples in assessing tobacco smoke consumption. In: Grabowski, J., Bell, C.S. (eds.). *Measurement in the Analysis and Treatment of Smoking Behavior*. NIDA Monograph No. 48. U.S. Department of Health and Human Services, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, DHHS Pub. No. (ADM)83-1285, 1983, pp. 6-26.
- BERGLUND, E., BERNSTEIN, D.A., EISINGER, R.A., HOCHBAUM, G.M., LICHTENSTEIN, E., SCHWARTZ, J.W., STRAITS, B.G. *Guidelines for Research on the Effectiveness of Smoking Cessation Programs*. Chicago, National Interagency Council on Smoking and Health, American Dental Association, October 1974.
- BIGLAN, A., ARY, D.V. Current methodological issues in research on smoking prevention. In: Bell, C., Battjes, R. (eds.). *Prevention Research: Detering Drug Abuse Among Children and Adolescents*. U.S. Department of Health and Human Services, National Institute on Drug Abuse, 1985.
- BISHOP, D.B., FISHER, E.B. *Employer Assisted Smoking Elimination: The Second Year (9/83-10/84)*. Unpublished, annual report to the American Lung Association of Eastern Missouri, St. Louis, 1984.
- BLISS, R.E., O'CONNELL, K.A. Problems with thiocyanate as an index of smoking status: A critical review with suggestions for improving the usefulness of biochemical measures in smoking cessation research. *Health Psychology* 3(6):563-581, 1984.
- BROWNELL, K.D. Weight control at the workplace: The power of social and behavioral factors. In: Cataldo, M.F., Coates, T.J. (eds.). *Health Promotion in Industry: A Behavioral Medicine Perspective*. New York, John Wiley and Sons, 1985.
- BROWNELL, K.D., COHEN, R.Y., STUNKARD, A.J., FELIX, M.R.J., COOLEY, N.B. Weight loss competitions at the work site: Impact on weight, morale, and cost-effectiveness. *American Journal of Public Health*, in press.
- CAPLAN, R.D., COBB, S., FRENCH, J.R.P., Jr. Relationships of cessation of smoking with job stress, personality, and social support. *Journal of Applied Psychology* 60(2):211-219, April 1975.
- CHESNEY, M.A., FEUERSTEIN, M. Behavioral medicine in the occupational setting. In: McNamara, J.R. (ed.). *Behavioral Approaches to Medicine: Application and Analysis*. New York, Plenum Press, 1979, pp. 267-290.
- COOK, T.D., CAMPBELL, D.T. *Quasi-Experimentation: Design and Analysis Issues for Field Settings*. Chicago, Rand McNally, 1979.
- COOPER, W.A. Johns-Manville says no smoking. *American Lung Association Bulletin* 64(6):7-10, July-August 1978.
- COPPOTELLI, H., ORLEANS, C.S. Spouse support in smoking cessation by women. *Journal of Consulting and Clinical Psychology*, in press.
- CURRY, S.J., GORDON, J.R., MARLATT, G.A. Relapse following smoking cessation: A comparison of aided and unaided quitters. In: Marlatt, G.A., Donovan, D. (eds.). *Assessment of Addictive Behaviors*. New York, Guilford Press, in press.

- DANAHER, B.G. Smoking cessation programs in occupational settings. *Public Health Reports* 95(2):149-157, March-April 1980.
- DARTNELL INSTITUTE OF BUSINESS RESEARCH. Target survey: Smoking in the office—Is it a problem? *Target Survey* 9:1-2, September 1977.
- DAVIS, A.L., FAUST, R., ORDENTLICH, M. Self-help smoking cessation and maintenance programs: A comparative study with 12-month follow-up by the American Lung Association. *American Journal of Public Health* 74(11):1212-1217, November 1984.
- DAWLEY, H.H., Jr., BALDWIN, J. The control of smoking: Smoking rate in designated smoking and no-smoking areas. *International Journal of the Addictions* 18(7):1033-1038, 1983.
- DAWLEY, H.H., Jr., BURTON, M.C. Smoking control in a hospital setting. *Addictive Behaviors*, in press.
- DAWLEY, H.H., Jr., FLEISCHER, B.J., DAWLEY, L.T. Smoking cessation with hospital employees: An example of worksite smoking cessation. *International Journal of the Addictions* 19(3):327-334, 1984.
- DAWLEY, H.H., Jr., MORRISON, J., CARROL, S. Compliance behavior in a hospital setting: Employee and patients' reactions to no-smoking signs. *Addictive Behaviors* 5(4):329-331, 1980.
- ELLIS, B.H., Jr. How to reach and convince asbestos workers to give up smoking. In: Schwartz, J.L. (ed.). *Progress in Smoking Cessation*. Proceedings of the International Conference on Smoking Cessation. New York, American Cancer Society, 1979, pp. 160-182.
- ELLIS, B.H. *Prerequisites for Successful Workplace-Based Smoking Cessation Programs*. Proceedings of Ohio Department of Health Conference "Smoking and the Workplace," Columbus, May 1980.
- EWART, C.K., LI, V.C., COATES, T.J. Increasing physicians' antismoking influence by applying an inexpensive feedback technique. *Journal of Medical Education* 58(6):468-473, June 1983.
- FAGERSTROM, K.-O. Measuring degree of physical dependence to tobacco smoking with reference to individualization of treatment. *Addictive Behaviors* 3(3/4):235-241, 1978.
- FAGERSTROM, K.-O. Effects of nicotine chewing gum and follow-up appointments in physician-based smoking cessation. *Preventive Medicine* 13(5):517-527, September 1984.
- FIELDING, J.E. Health promotion and disease prevention at the worksite. *Annual Review of Public Health* 5:237-265, 1984.
- FREDERIKSEN, L.W., MARTIN, J.E. Carbon monoxide and smoking behavior. *Addictive Behaviors* 4(1):21-30, 1979.
- GLASGOW, R.E., KLESGES, R.C., GODDING, P.R., VASEY, M.W., O'NEILL, H.K. Evaluation of a worksite-controlled smoking program. *Journal of Consulting and Clinical Psychology* 52(1):137-138, February 1984.
- GLASGOW, R.E., KLESGES, R.C., O'NEILL, H.K. Programming social support for smoking modification: An extension and replication. *Addictive Behaviors*, in press.
- GLASGOW, R.E., ROSEN, G.M. Behavioral bibliotherapy: A review of self-help behavior therapy manuals. *Psychological Bulletin* 85(1):1-23, January 1978.
- GROVE, D.A., REED, R.W., MILLER, L.C. A health promotion program in a corporate setting. *Journal of Family Practice* 9(1):83-88, July 1979.
- HECKLER, L.M. Employee education programs: One aspect of a nurse's expanded role in an occupational health program. *Occupational Health Nursing* 28(8): 25-29, August 1980.
- HUGHES, G.H., HYMOWITZ, N., OCKENE, J.K., SIMON, N., VOGT, T.M. The Multiple Risk Factor Intervention Trial (MRFIT): V. Intervention on smoking. *Preventive Medicine* 10(4):476-500, July 1981.

- HUNT, W.A., BESPALC, D.A. An evaluation of current methods of modifying smoking behavior. *Journal of Clinical Psychology* 30(4):431-438, October 1974.
- JANIS, I.L. The role of social support in adherence to stressful decisions. *American Psychologist* 38(2):143-160, February 1983.
- JASON, L.A., CLAY, R. Modifying smoking behaviors in a barber shop. *Man-Environment Systems* 8(1):38-40, January 1978.
- JASON, L.A., LIOTTA, R.F. Reduction of cigarette smoking in a university cafeteria. *Journal of Applied Behavior Analysis* 15(4):573-577, Winter 1982.
- JASON, L.A., SAVIO, D. Reducing cigarette smoke in an office setting. *Health Values: Achieving High Level Wellness* 2(4):180-185, July-August 1978.
- KANZLER, M., ZEIDENBERG, P., JAFFE, J.H. Response of medical personnel to an on-site smoking cessation program. *Journal of Clinical Psychology* 32(3):670-674, July 1976.
- KLESGES, R.C., GLASGOW, R.E. Smoking modification in the worksite. In: Cataldo, M.F., Coates, R.J. (eds.). *Health Promotion in Industry: A Behavioral Medicine Perspective*. New York, John Wiley and Sons, 1985.
- KLESGES, R.C., VASEY, M.W., GLASGOW, R.E. *Evaluation of a Worksite Smoking Competition Program*. Paper presented at the annual meeting of the Society of Behavioral Medicine, New Orleans, March 1985.
- KORNITZER, M., DE BACKER, G., DRAMAIX, M., THILLY, C. The Belgian Heart Disease Prevention Project: Modification of the coronary risk profile in an industrial population. *Circulation* 61(1):18-25, January 1980.
- KORNITZER, M., DRAMAIX, M., KITTEL, F., DE BACKER, G. The Belgian Heart Disease Prevention Project: Changes in smoking habits after two years of intervention. *Preventive Medicine* 9(4):496-503, July 1980.
- KRISTEIN, M. *The Economics of Health Promotion at a Worksite*. (mimeograph). New York, American Health Foundation, 1982.
- LEVENTHAL, H. Findings and theory in the study of fear communications. In: Berkowitz, L. (ed.). *Advances in Experimental Social Psychology*. Vol. 5. New York, Academic Press, 1970, pp. 119-186.
- LI, V.C., KIM, Y.J., EWART, C.K., TERRY, P.B., CUTHIE, J.C., WOOD, J., EMMETT, E.A., PERMUTT, S. Effects of physician counseling on the smoking behavior of asbestos-exposed workers. *Preventive Medicine* 13(5):462-476, September 1984.
- LICHTENSTEIN, E. The smoking problem: A behavioral perspective. *Journal of Consulting and Clinical Psychology* 50(6):804-819, December 1982.
- LICHTENSTEIN, E., BROWN, R.A. Current trends in the modification of cigarette dependence. In: Bellack, A.S., Hersen, M., Kazdin, A.E. (eds.). *International Handbook of Behavior Modification and Therapy*. New York, Plenum Press, 1982, pp. 575-611.
- LICHTENSTEIN, E., DANAHAR, B.G. What can the physician do to assist the patient to stop smoking? In: Brashear, R.E., Rhodes, M.L. (eds.). *Chronic Obstructive Lung Disease: Clinical Treatment and Management*. St. Louis, C. V. Mosby, 1978, pp. 227-241.
- LUCE, B.R., SCHWEITZER, S.O. Smoking and alcohol abuse: A comparison of their economic consequences. *New England Journal of Medicine* 298(10):569-571, March 9, 1978.
- LUEPKER, R.V., PECHACEK, T.F., MURRAY, D.M., JOHNSON, C.A., HUND, F., JACOBS, D.R. Saliva thiocyanate: A chemical indicator of cigarette smoking in adolescents. *American Journal of Public Health* 71(12):1320-1324, December 1981.
- MALOTT, J.M., GLASGOW, R.E., O'NEILL, H.K., KLESGES, R.C. Co-worker social support in a worksite smoking control program. *Journal of Applied Behavior Analysis* 17(4):485-495, Winter 1984.
- MARLATT, G.A., GORDON, J.R. *Relapse Prevention: Maintenance Strategies in the Treatment of Addictive Behaviors*. New York, Guilford Press, 1985, 558 pp.

- McFALL, R.M. Smoking-cessation research. *Journal of Consulting and Clinical Psychology* 46(4):703-712, August 1978.
- McMORROW, M.J., FOX, R.M. Nicotine's role in smoking: An analysis of nicotine regulation. *Psychological Bulletin* 93(2):302-327, March 1983.
- MEADE, T.W., WALD, N.J. Cigarette smoking patterns during the working day. *British Journal of Preventive and Social Medicine* 31:25-29, 1977.
- MERMELSTEIN, R., LICHTENSTEIN, E., McINTYRE, K. Partner support and relapse in smoking-cessation programs. *Journal of Consulting and Clinical Psychology* 51(3):465-466, June 1983.
- MEYER, A.J., HENDERSON, J.B. Multiple risk factor reduction in the prevention of cardiovascular disease. *Preventive Medicine* 3(2):225-236, June 1974.
- MILLER, G.H. SOS stop smoking clinic: A one-year report on the program at the Cummins Engine Company. *Journal of the Indiana State Medical Association* 74(5):292-294, May 1981.
- MORGAN, W.P. Psychological benefits of physical activity. In: Nagle, F.J., Montoye, H.J. (eds.). *Exercise in Health and Disease*. Springfield, Illinois, Charles C Thomas, 1981, pp. 299-314.
- MOSS, R.A., PRUE, D.M. Research on nicotine regulation. *Behavior Therapy* 13(1):31-46, January 1982.
- MYERS, J.L. *Fundamentals of Experimental Design*. 2d ed. Boston, Allyn and Bacon, 1972.
- NADITCH, M.P. The STAYWELL Program. In: Matarazzo, J.D., Weiss, S.M., Herd, J.A., Miller, N.E., Weiss, S.M. (eds.). *Behavioral Health: A Handbook of Health Enhancement and Disease Prevention*. New York, John Wiley and Sons, 1984, pp. 1071-1078.
- NATHAN, P.E. Johnson & Johnson's Live for Life: A comprehensive positive lifestyle change program. In: Matarazzo, J.D., Weiss, S.M., Herd, J.A., Miller, N.E., Weiss, S.M. (eds.). *Behavioral Health: A Handbook of Health Enhancement and Disease Prevention*. New York, John Wiley and Sons, 1984, pp. 1064-1070.
- NATIONAL INTERAGENCY COUNCIL ON SMOKING AND HEALTH. *Smoking and the Workplace: National Interagency Council on Smoking and Health Business Survey*. Unpublished manuscript, 1980.
- NEPPS, M.M. A minimal contact smoking cessation program at the worksite. *Addictive Behaviors* 9(3):291-294, 1984.
- OCKENE, J.K., HYMOWITZ, N., SEXTON, M., BROSTE, S.K. Comparison of patterns of smoking behavior change among smokers in the Multiple Risk Factor Intervention Trial (MRFIT). *Preventive Medicine* 11(6):621-638, November 1982.
- ORLEANS, C.S., SHIPLEY, R.H. Worksite smoking cessation initiatives: Review and recommendations. *Addictive Behaviors* 7(1):1-16, 1982.
- PARKINSON, R.S., BECK, R.N., COLLINGS, G.H., Jr., ERIKSEN, M., MCGILL, A.M., PEARSON, C.E., WARE, B.G. *Managing Health Promotion in the Workplace: Guidelines for Implementation and Evaluation*. Palo Alto, Mayfield Publishing Co., 1982, 314 pp.
- PECHACEK, T.F., FOX, B.H., MURRAY, D.M., LEUPKER, R.V. Review of techniques for measurement of smoking behavior. In: Matarazzo, J.C., Weiss, S.M., Herd, J.A., Miller, N.E., Weiss, S.M. (eds.). *Behavioral Health: A Handbook of Health Enhancement and Disease Prevention*. New York, John Wiley and Sons, 1984.
- RAND, C., STITZER, M., BIGELOW, G., MEAD, A. *Contingent Reinforcement for Smoking Abstinence*. Poster presented at American Psychological Association, Toronto, 1984.
- RAW, M., JARVIS, M.J., FEYERABEND, C., RUSSELL, M.A.H. Comparison of nicotine chewing-gum and psychological treatments for dependent smokers. *British Medical Journal* 281(6238):481-482, August 16, 1980.

- ROSE, G., HAMILTON, P.J.S. A randomised controlled trial of the effect on middle-aged men of advice to stop smoking. *Journal of Epidemiology and Community Health* 32(4):275-281, December 1978.
- ROSE, G., HAMILTON, P.J.S., COLWELL, L., SHIPLEY, M.J. A randomized controlled trial of anti-smoking advice: 10-year results. *Journal of Epidemiology and Community Health* 36(2):102-108, June 1982.
- ROSE, G., HELLER, R.F., PEDOE, H.T., CHRISTIE, D.G.S. Heart disease prevention project: A randomised controlled trial in industry. *British Medical Journal* 280(6216):747-751, March 15, 1980.
- ROSEN, G.M., LICHTENSTEIN, E. An employee incentive program to reduce cigarette smoking. *Journal of Consulting and Clinical Psychology* 45(5):957-959, October 1977.
- RUSSELL, M.A.H., WILSON, C., TAYLOR, C., BAKER, C.D. Effect of general practitioners' advice against smoking. *British Medical Journal* 2(6184):231-235, July 28, 1979.
- SAWYER, A. Repetition, cognitive responses, and persuasion. In: Petty, R., Ostrom, T., Brock, T. (eds.). *Cognitive Responses in Persuasion*. Trenton, Erlbaum, 1981.
- SCHACHTER, S. Recidivism and self-cure of smoking and obesity. *American Psychologist* 37(4):436-444, April 1982.
- SCHLEGEL, R.P., MANSKE, S.R., SHANNON, M.E. *Butt Out: Evaluation of the Canadian Armed Forces smoking cessation program*. In: Forbes, W.F., Frecker, R.C., Nostbakken, D. (eds.). *Proceedings of the Fifth World Conference on Smoking and Health, Winnipeg, Canada, 1983*. Ottawa, Canadian Council on Smoking and Health, 1983, pp. 445-452.
- SCHNEIDER, S.J., BENYA, A., SINGER, H. Computerized direct mail to treat smokers who avoid treatment. *Computers and Biomedical Research* 17(5):409-418, October 1984.
- SCOTT, R.R., DENIER, C.A., PRUE, D.M. *Worksite Smoking Intervention With Health Professionals*. Paper presented at the Association for the Advancement of Behavior Therapy annual convention, Washington, D.C., 1983.
- SHEPARD, D.S. *Incentives for Not Smoking: Experience at the Speedcall Corporation. A Preliminary Report*. Boston, Center for the Analysis Of Health Practices, Harvard School of Public Health, 1980, 14 pp.
- SHEPARD, D.S., PEARLMAN, L.A. Health habits that pay off. *Business and Health*, in press.
- SHIPLEY, R.H., ROSEN, T.J., WILLIAMS, C. Measurement of smoking: Surveys and some recommendations. *Addictive Behaviors* 7(3):299-302, 1982.
- SORMAN, K. This quit-smoking program works. *American Lung Association Bulletin* 65(6):2-6, July-August 1979.
- STACHNIK, T.J., STOFFELMAYR, B.E. Is there a future for smoking cessation programs? *Journal of Community Health* 7(1):47-56, Fall 1981.
- STACHNIK, T., STOFFELMAYR, B. Worksite smoking cessation programs: A potential for national impact. *American Journal of Public Health* 73(12):1395-1396, December 1983.
- STITZER, M.L., BIGELOW, G.E. Contingent reinforcement for reduced carbon monoxide levels in cigarette smokers. *Addictive Behaviors* 7(4):403-412, 1982.
- STITZER, M.L., BIGELOW, G.E. Contingent payment for carbon monoxide reduction: Effects of pay amount. *Behavior Therapy* 14(5):647-656, November 1983.
- STITZER, M.L., BIGELOW, G.E. Contingent reinforcement for reduced breath carbon monoxide levels: Target-specific effects on cigarette smoking. *Addictive Behaviors*, in press.
- TERRY, L.T. The future of an illusion. *American Journal of Public Health* 61(2):233-240, February 1971.

- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Cancer. A Report of the Surgeon General*. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, DHHS Pub. No. (PHS)82-50179, 1982, 322 pp.
- U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES. *The Health Consequences of Smoking: Cardiovascular Disease. A Report of the Surgeon General*. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health. DHHS Pub. No. (PHS)84-50204, 1983, 388 pp.
- U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. *Respiratory Diseases: Task Force Report on Prevention, Control, Education*. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, Heart, Lung, and Blood Institute. DHEW Pub. No. (NIH)77-1248, March 1977.
- U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. *Health: United States*. U.S. Department of Health, Education, and Welfare, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, DHEW Publication No. (PHS)80-1232, 1979.
- WARE, B.G., BLOCK, D.L. Cardiovascular risk intervention at the work site: The Ford Motor Company program. *International Journal of Mental Health* 11(3):68-75, Fall 1982.
- WARNER, K.E. *Economic Incentives and Health Behavior*. Paper presented at the Vermont Conference on the Primary Prevention of Psychopathology, Burlington, June-July, 1983.
- WINDSOR, R.A., BARTLETT, E.E. Employee self-help smoking cessation programs: A review of the literature. *Health Education Quarterly* 11(4):349-359, Winter 1984.