

Preliminary list of species of special status that may be affected by management of resident Canada geese.

Region 1:

Light-footed clapper rail	(<i>Rallus longirostris levipes</i>) [E]
California clapper rail	(<i>Rallus longirostris obsoletus</i>) [E]
Yuma clapper rail	(<i>Rallus longirostris yumanensis</i>) [E]
California least tern	(<i>Sterna antillarum</i>) [E]
Brown pelican	(<i>Pelicanus occidentalis</i>) (Pacific coast population) [E]
Southwestern willow flycatcher	(<i>Empidonax traillii extimus</i>) [E]
California condor	(<i>Gymnogyps californianus</i>) [E]
Least Bell's vireo	(<i>Vireo belli pusillus</i>) [E]
Western snowy plover	(<i>Charadrius alexandrinus nivosus</i>) [T]
Bald eagle	(<i>Haliaeetus leucocephalus</i>) [T]
California gnatcatcher	(<i>Polioptila californica</i>) [T]
Inyo California towhee	(<i>Pipilo crissalis eremophilus</i>) [T]
Marbled murrelet	(<i>Brachyramphus marmoratus</i>) [T]
Northern spotted owl	(<i>Strix occidentalis caurina</i>) [T]
Giant Garter Snake	(<i>Thamnophis gigas</i>) [T]
Fenders blue butterfly	(<i>Icaricia icarioides fenderi</i>) [E]
Bay checkerspot butterfly	(<i>Euphydryas editha bayensis</i>) [T]
Behren's silverspot butterfly	(<i>Speyeria zerene behrensii</i>) [E]
San Joaquin adobe sunburst	(<i>Pseudobahia peirsonii</i>) [T]
Willamette daisy	(<i>Erigeron decumbens</i> var. <i>decumbens</i>) [E]

Region 2

Attwater's greater prairie-chicken	(<i>Tympanuchus cupido attwateri</i>) [E]
Masked bobwhite	(<i>Colinus virginianus ridgewayi</i>) [E]
Red-cockaded woodpecker	(<i>Picoides borealis</i>) [E]
Cactus ferruginous pygmy-owl	(<i>Glaucidium brasilainum cactorum</i>) [E]
Yuma clapper rail	(<i>Rallus longirostris yumanensis</i>) [E]
Least tern	(<i>Sterna antillarum</i>) [E]
Northern aplomado falcon	(<i>Falco femoralis septentrionalis</i>) [E]
Brown pelican	(<i>Pelicanus occidentalis</i>) [E]
Southwestern willow flycatcher	(<i>Empidonax traillii extimus</i>) [E]
Black-capped vireo	(<i>Vireo atricapillus</i>) [E]
Golden-cheeked warbler	(<i>Dendroica chrysoparia</i>) [E]
California condor	(<i>Gymnogyps californianus</i>) [XN]
Mexican spotted owl	(<i>Strix occidentalis lucida</i>) [T]
Bald eagle	(<i>Haliaeetus leucocephalus</i>) [T]
Piping plover	(<i>Charadrius melodus</i>) [T]
Western prairie fringed orchid	(<i>Platanthera praeclara</i>) [T]

Region 3

Piping plover	(<i>Charadrius melodus</i>) [T]
Least tern	(<i>Sterna antillarum</i>) (Interior population) [E]
Bald eagle	(<i>Haliaeetus leucocephalus</i>) [T]
Hine's emerald dragonfly	(<i>Somatochlora hineana</i>) [E]
Karner blue butterfly	(<i>Lycaeides melissa samuelis</i>) [E]
American burying beetle	(<i>Nicrophorus americanus</i>) [E]
Mead's milkweed	(<i>Asclepias meadii</i>) [T]
Virginia sneezeweed	(<i>Helenium virginicum</i>) [T]
Decurrent false aster	(<i>Boltonia decurrents</i>) [T]
Running buffalo clover	(<i>Trifolium stoloniferum</i>) [E]
Prairie bush-clover	(<i>Lespedeza leptostachya</i>) [T]
Leafy prairie-clover	(<i>Dalea foliosa</i>) [E]
Western prairie fringed orchid	(<i>Platanthera praeclara</i>) [T]
Eastern prairie fringed orchid	(<i>Platanthera leucophaea</i>) [T]

Region 4

Red-cockaded woodpecker	(<i>Picoides borealis</i>) [E]
Mississippi sandhill crane	(<i>Grus canadensis pulla</i>) [E]
Piping plover	(<i>Charadrius melodus</i>) [E]
Least tern	(<i>Sterna antillarum</i>) (Interior population) [E]
Everglade snail kite	(<i>Rostrhamus sociabilis plumbeus</i>) [E]
Wood stork	(<i>Mycteria americana</i>) [E]
Brown pelican	(<i>Pelicanus occidentalis</i>) [E]
Cape Sable sparrow	(<i>Ammodramus maritimus mirabilis</i>) [E]
Florida grasshopper sparrow	(<i>Ammodramus savanarum floridanus</i>) [E]
Yellow-shouldered blackbird	(<i>Agelaius xanthomus</i>) [E]
Roseate tern	(<i>Sterna douglalli</i>) [T]
Bald eagle	(<i>Haliaeetus leucocephalus</i>) [T]
Audubon's crested caracara	(<i>Polyborus plancus audubonii</i>) [T]
Florida scrub jay	(<i>Aphelocoma coerulescens</i>) [T]
Bog turtle	(<i>Clemmys muhlenbergii</i>) [T]
Saint Francis' satyr butterfly	(<i>Neonympha mitchellii francisci</i>) []
Schweinitz's sunflower	(<i>Helianthus schweinitzii</i>) [4]
Eggert's sunflower	(<i>Helianthus eggertii</i>) [T]
Spring Creek bladderpod	(<i>Lesquerella perforata</i>) [E]
Running buffalo clover	(<i>Trifolium stoloniferum</i>) [E]
Eastern prairie fringed orchid	(<i>Platanthera leucophaea</i>) [T]

Region 5

Piping plover	(<i>Charadrius melodus</i>) [T]
Roseate tern	(<i>Sterna douglalli</i>) [E]
Bald eagle	(<i>Haliaeetus leucocephalus</i>) [T]

Plymouth redbelly turtle (*Pseudemys rubriventris bangsi*) [E]
Bog turtle (*Clemmys muhlenbergii*) [T]
Karner blue butterfly (*Lycaeides melissa samuelis*) [E]
Virginia sneezeweed (*Helenium virginicum*) [T]
Running buffalo clover (*Trifolium stoloniferum*) [E]
Eastern prairie fringed orchid (*Platanthera leucophaea*) [T]

Region 6

Least tern (*Sterna antillarum*) (Interior population) [E]
Piping plover (*Charadrius melodus*) [T]
Mexican spotted owl (*Strix occidentalis lucida*) [T]
Bald eagle (*Haliaeetus leucocephalus*) [T]
Whooping crane (*Grus americana*) [E]
Mead's milkweed (*Asclepias meadii*) [T]
Western prairie fringed orchid (*Platanthera praeclara*) [T]