

ERRATA

for the FDA Review

of

<p>BL STN 125011/0 Bexxar® (Tositumomab (Anti-B1) and I¹³¹-Tositumomab) Corixa Corporation</p>

Page 5

Table titled “Summary of Efficacy Outcomes by Study”

Correction: **Primary Efficacy Studies, Study RIT-II-004 (N=61)**

Pages 32-33

Correct Subtitles

“Violation of Eligibility Criteria”

“Violations of Informed Consent”

“Violation of Thyroid Protection Protocol”

**“Violation of Timing for Dose Assessment of Administration of
Therapeutic Dose”**

Page 35

Correction: **Time from diagnosis to entry (years)**

Page 41

Correction in Table

Response Rate in Subset without Transformation (N=37)

Corrected Table

Age (Years)		
Median	59	58
Range	(37, 80)	(37, 80)
Gender		
Male (% male)	41 (58%)	23 (58%)
Median time from diagnosis to study entry (years) (range)	6.2 (0.7, 27.8)	5.0 (0.7, 27.8)
Median time from diagnosis to transformation date (years) (range)	1.8 (-0.3, 10.3)	1.9 (0.02, 9.9)
Median time from transformation to study entry (years) (range)	3.4 (0, 24.5)	3.3 (0, 24.5)
Ann Arbor Stage at entry		
1	1 (1%)	1 (2%)
2	7 (10%)	1 (2%)
3	17 (24%)	11 (28%)
4	46 (65%)	27 (68%)
Modified IPI Score	(n = 67)	(n = 38)
0-1	9 (13%)	2 (5%)
2	23 (34%)	14 (37%)
3	23 (34%)	16 (42%)
4-5	12 (18%)	6 (16%)
Number of prior chemotherapies		
Median	4 (3, 5)	4 (3, 5)
IQ	(1, 11)	(1, 9)
Range		
Maximum unidimensional lesion measurement (cm)		
0 to ≤5 cm	24 (34%)	12 (30%)
>5 to ≤10 cm	34 (48%)	20 (50%)
> 10 cm	13 (18%)	8 (20%)
Response to last chemotherapy		
Response (CR+CCR+PR)	35 (49%)	22 (55%)
Complete Response (CR+CCR)	16 (23%)	10 (25%)
Tumor grade at the study entry		
Low	9 (13%)	2 (5%)
Intermediate	59 (83%)	35 (88%)
High	3 (4%)	3 (8%)
Last qualifying chemotherapy end day to study day (yrs)	(n = 66)	(n = 35)
Median	0.5	0.5
Range	(0.1, 5.4)	(0.1, 3.1)