






Galilean Moons of Jupiter



Galilean Moons of Jupiter



The planet Jupiter's four largest moons are called the Galilean satellites, after Italian astronomer Galileo Galilei, who observed them in 1610. The German astronomer Simon Marius claimed to have seen the moons around the same time, but he did not publish his observations and so Galileo is given the credit for their discovery. These large moons, named Io, Europa, Ganymede, and Callisto, are each distinctive worlds.

Io is the most volcanically active body in the solar system. Its surface is covered by sulfur in different colorful forms. As Io travels in its slightly elliptical orbit, Jupiter's immense gravity causes "tides" in the solid surface 100 meters (300 feet) high on Io, generating enough heat to give rise to the volcanic activity and drive off any water. Io's volcanoes are driven by hot silicate magma.

Europa's surface is mostly water ice, and there is evidence that it may be covering an ocean of water or slushy ice. Europa is thought to have twice as much water as does Earth. This moon intrigues astrobiologists because of its potential for having a "habitable zone." Life forms have been found thriving near subterranean volcanoes on Earth and in other extreme locations that may be analogues to what may exist on Europa. Ganymede is the largest moon in the solar system (larger than the planet Mercury), and is the only moon known to have its own internally generated magnetic field. Callisto's surface is extremely heavily cratered and ancient — a record of events from the early history of the solar system. However, the very few small craters on Callisto indicate a small degree of current surface activity.

The interiors of Io, Europa, and Ganymede have a layered structure (as does Earth). Io has a core, and a mantle of at least partially molten rock, topped by a crust of solid rock coated with sulfur compounds. Europa and Ganymede both have a core; a rock envelope around the core; a thick, soft ice layer; and a thin crust of impure water ice. In the case of Europa, a global subsurface water layer probably lies just below the icy crust. Layering at Callisto is less well defined and appears to be mainly a mixture of ice and rock.

Three of the moons influence each other in an interesting way. Io is in a tug-of-war with Ganymede and Europa, and Europa's orbital period (time to go around Jupiter once) is twice Io's period,

and Ganymede's period is twice that of Europa. In other words, every time Ganymede goes around Jupiter once, Europa makes two orbits, and Io makes four orbits. The moons all keep the same face towards Jupiter as they orbit, meaning that each moon turns once on its axis for every orbit around Jupiter.

Pioneers 10 and 11 (1973 to 1974) and Voyagers 1 and 2 (1979) offered striking color views and global perspectives from their flybys of the Jupiter system. From 1995 to 2003, the Galileo spacecraft made observations from repeated elliptical orbits around Jupiter, passing as low as 261 kilometers (162 miles) over the surfaces of the Galilean moons. These close approaches resulted in images with unprecedented detail of selected portions of the surfaces.

Close-up images taken by the Galileo spacecraft of portions of Europa's surface show places where ice has broken up and moved apart, and where liquid may have come from below and frozen smoothly on the surface. The low number of craters on Europa leads scientists to believe that a subsurface ocean has been present in recent geologic history and may still exist today. The heat needed to melt the ice in a place so far from the Sun is thought to come from inside Europa, resulting primarily from the same type of tidal forces that drive Io's volcanoes.

FAST FACTS

Moon	Distance from Jupiter
Io	422,000 km (262,200 mi)
Europa	671,000 km (417,000 mi)
Ganymede	1,070,000 km (665,000 mi)
Callisto	1,883,000 km (1,170,000 mi)

Moon	Mean Radius
Io	1,821.6 km (1,131.9 mi)
Europa	1,560.8 km (969.8 mi)
Ganymede	2,631 km (1,635 mi)
Callisto	2,410 km (1,498 mi)

Moon	Orbital Period (Earth Days)
Io	1.769
Europa	3.551
Ganymede	7.155
Callisto	16.689

Moon	Density (g/cm ³)
Io	3.528
Europa	3.013
Ganymede	1.942
Callisto	1.834

SIGNIFICANT DATES


1610 — Galileo Galilei and Simon Marius independently discover four moons orbiting Jupiter.

1979 — Voyager 1 photographs an erupting volcano on Io: the first ever seen anywhere but on Earth.

1979–2000 — Using data from the Voyager and Galileo spacecraft, scientists gather strong evidence of an ocean beneath the icy crust of Europa; Galileo data indicate possible oceans on Ganymede and Callisto.

2003 — The Galileo mission ends with the spacecraft deliberately descending into Jupiter's atmosphere. In the same year, scientists discover 23 new moons of Jupiter.

ABOUT THE IMAGES



1 A composite "portrait" of Jupiter's four Galilean moons Io, Europa, Ganymede, and Callisto (Jupiter is not at the same scale as the satellites).

2 During one flyby of Io, the Galileo spacecraft photographed Tvashtar Catena, an area of giant erupting volcanoes.

3 This false-color image of Europa shows the icy crust broken up into blocks that "rafted" into new positions.

4 Fresh, bright material was thrown out of an impact crater on Ganymede.

5 Ice on Callisto excavated by younger impact craters contrasts with darker, redder coatings on older surfaces.

FOR MORE INFORMATION

solarsystem.nasa.gov/planets/profile.cfm?Object=Jupiter