

B1. SOUTHERN NEW ENGLAND/MID-ATLANTIC (SNE/MA) WINTER FLOUNDER

TERMS OF REFERENCE

The following terms of reference were addressed for the Southern New England/Mid Atlantic (SNE/MA) stock complex of winter flounder:

- 1) Update the status of SNE/MA winter flounder stock through 2001 providing estimates of fully recruited fishing mortality rate, biomass weighted fishing mortality rate, stock size, mean biomass, spawning stock biomass, and recruitment as appropriate. Characterize uncertainty in SSB and fishing mortality rates.
- 2) Provide short-term (2003) and medium term projections (2009) of catch and biomass (mean biomass, SSB) under status quo F , and ASMFC's $F_{40\%}$ target, and NEFMC's F_{MSY} .
- 3) Develop research recommendations for improving the assessment of SNE/MA winter flounder.
- 4) Comment on and revise, where necessary, the ASMFC and the NEFMC overfishing definitions for this stock. (Note: Currently ASMFC and the NEFMC have different overfishing definitions. The ASMFC Board had recommended that the Winter Flounder Technical Committee develop a single overfishing definition for this stock).

INTRODUCTION

The current assessment of the SNE/MA stock complex of winter flounder is an update of the previous assessment completed in 1998 at SARC 28 (NEFSC 1999). The SARC 28 assessment included catch through 1997, research survey abundance indices through 1998, catch at age analyzed by Virtual Population Analysis (VPA) for 1981-1997, and biological reference points based on a production model conditioned on VPA results. The SARC 28 assessment concluded that the stock complex was fully exploited and at a medium level of biomass. Total biomass in 1997 was estimated to be 17,900 mt, spawning stock biomass was estimated to be 8,600 mt, and the fully recruited fishing mortality rate was estimated to be $F = 0.31$. Subsequent to the SARC 28 assessment, the status of SNE/MA winter flounder has been evaluated annually by projection methods to provide advice to the New England Fishery Management Council (NEFMC). The last such status update was provided in 2001, and projected total biomass to be 25,300 mt, spawning stock biomass to be 13,800 mt, and fully recruited $F = 0.29$, in 1999 (NEFSC 2001). The current assessment updates landings and discard estimates, research survey abundance indices, and assessment models through 2001-2002, as applicable.

Winter flounder (*Pleuronectes americanus*) is a demersal flatfish species commonly found in estuaries and on the continental shelf. The species is distributed between the Gulf of St. Lawrence and North Carolina, although it is not abundant south of Delaware Bay. Within the SNE/MA stock complex, winter flounder undergo migrations from estuaries, where spawning occurs in the late winter and spring, to offshore shelf areas of less than 60 fathoms. Winter flounder reach a maximum size of around 2.25 kg (5 pounds) and 65 cm, with the exception of Georges Bank where growth rate is higher and fish may reach a maximum weight up to 3.6 kg (8 pounds; Bigelow and Schroeder 1953).

Current fishery management is coordinated by the Atlantic States Marine Fisheries Commission (ASMFC) in state waters and the NEFMC in federal waters. Winter flounder fisheries in state waters have been managed by Interstate Agreement under the auspices of the ASMFC Fishery Management Plan (FMP) for Inshore Stocks of Winter Flounder since approval in May, 1992. The plan includes states from Delaware to Maine, with Delaware granted *de minimus* status (habitat regulations applicable but fishery management not required). The Plan's goal is to rebuild spawning stock abundance and achieve a fishing mortality-based management target of $F_{40\%}$ (fishing rate that preserves 40% of the maximum spawning potential of the stock) in three steps: $F_{25\%}$ in 1993-1994, $F_{30\%}$ in 1995-1998, and $F_{40\%}$ in 1999 and later years through implementation of compatible, state-specific regulations. Coastal states from New Jersey to New Hampshire have promulgated a broad suite of indirect catch and effort controls. State agencies have set or increased minimum size limits for recreationally and commercially landed flounder (10-12 inches and 12 inches, respectively); enacted limited recreational closures and bag limits; and instituted seasonal, areal, or state-wide commercial landings/gear restrictions. Minimum codend mesh regulations have been promulgated in directed winter flounder fisheries: 5 inch for NJ and NY, 5.5 inch for CT, 5 inch for RI, and 6 inch for MA.

Winter flounder in the Exclusive Economic Zone (EEZ) are managed under the Northeast Multispecies Fishery FMP developed by the NEFMC. The principle catch of winter flounder in the EEZ has recently occurred as bycatch in directed trawl fisheries for Atlantic cod, haddock, and yellowtail flounder. The management unit encompasses the multispecies finfish fishery that operates from eastern Maine through Southern New England (72°30'). At least one offshore stock, on Georges Bank, has been identified. The FMP extends authority over vessels permitted under the FMP even while fishing in state waters if federal regulations are more restrictive than the state regulations.

The Multispecies FMP was implemented in September, 1986, imposing a codend minimum mesh size of 5.5 inches (previously 5.1 inches) in the large-mesh regulatory area of Georges Bank and the offshore portion of Gulf of Maine. There were closed areas and seasons for haddock and yellowtail flounder. In the western Gulf of Maine, vessels were required to enroll in an Exempted Fisheries Program in order to target small-mesh species such as shrimp, dogfish, or whiting. The bycatch restrictions specified area and season and limited groundfish bycatch to 25% of trip and 10% for the reporting period. In southern New England waters, the groundfish bycatch on vessels fishing with small mesh was not limited in any way. There was a 11 inch

minimum size for winter flounder which corresponded with the length at first capture (near zero percent retention) for 5.5 inch diamond mesh. Although the Multispecies FMP was amended four times by 1991, it was widely recognized that many stocks, including winter flounder, were being overfished.

Time-specific stock rebuilding schedules were a part of Multispecies FMP Amendment 5 which took effect in May, 1994. The rebuilding target for winter flounder, a so-called "large-mesh" species, was $F_{20\%}$ within 10 years. Along with a moratorium on issuance of additional vessel permits, the cornerstone of Amendment 5 was an effort reduction program that required "large-mesh" groundfish vessels to limit days at sea, which would be reduced each year. There was an exemption from effort reduction requirements for groundfishing vessels less than 45 feet in length and for "day boats" (from 2:1 layover day ratio requirement). Dragger retaining more than the "possession limit" of groundfish (10%, by weight, up to 500 lbs) were required to fish with either 5.5 inch diamond or square mesh in Southern New England or 6 inch throughout the net in the regulated mesh area of Georges Bank/ Gulf of Maine, respectively. The possession limit was allowed when using small mesh within the western Gulf of Maine (except Jeffreys Ledge and Stellwagon Bank) and in Southern New England. Vessels fishing in the EEZ west of 72° 30' (the longitude of Shinnecock Inlet, NY) were required to abide by 5.5 inch diamond or 6 inch square codend mesh size restrictions consistent with the Summer Flounder FMP. The minimum landed size of winter flounder increased to 12 inches, appropriate for the increased mesh size in order to reduce discards. There were many additional rules including time/area closures for sink gillnet vessels, seasonal netting closures of prime fishing areas on Georges Bank (Areas I and II), and on Nantucket Shoals to protect juvenile yellowtail flounder.

At the end of 1994, the NEFMC reacted to collapsed stocks of Atlantic cod, haddock, and yellowtail flounder on Georges Bank by recommending a number of emergency actions to tighten existing regulations reducing fishing mortality. Prime fishing areas on Georges Bank (Areas I & II), and the Nantucket Lightship Area were closed. The NEFMC also addressed expected re-direction of fishing effort into Gulf of Maine and Southern New England while, at the same time, developing Amendment 7 to the Multispecies FMP. Under Amendment 7, days-at-sea controls were extended, and any fishing by an EEZ-permitted vessel required use of not less than 6 inch diamond or square mesh in Southern New England east of 72° 30'. Framework 27 in 1999 increased the square mesh minimum size to 6.5 inches in the Gulf of Maine, Georges Bank, and Southern New England mesh areas. Amendment 9 revised the overfishing definitions for New England groundfish, and new overfishing definitions for SNE/MA winter flounder were recommended by SARC 28 (NEFSC 1999).

STOCK STRUCTURE

Although stock groups consist of an assemblage of adjacent estuarine spawning units, the ASMFC FMP originally defined three coastal management units based on similar growth,

maturity and seasonal movement patterns: Gulf of Maine, Southern New England and Mid-Atlantic. Boundaries for a total of four winter flounder stock units as originally defined in the ASMFC management plan (Howell et al., 1992) were:

Gulf of Maine: Coastal Maine, New Hampshire, and Massachusetts north of Cape Cod

Southern New England: Coastal Massachusetts east and south of Cape Cod, including Nantucket Sound, Vineyard Sound, Buzzards Bay, Narragansett Bay, Block Island Sound, Rhode Island Sound, Rhode Island coastal ponds and eastern Long Island Sound to the Connecticut River, including Fishers Island Sound, NY.

Mid-Atlantic: Long Island Sound west of the Connecticut River to Montauk Point, NY, including Gardiners and Peconic Bays, coastal Long Island, NY, coastal New Jersey and Delaware.

Georges Bank

In the current and three previous assessments (e.g., NEFSC 1996, ASMFC 1998, NEFSC 1999) the Southern New England and Mid-Atlantic units have been combined into a single stock complex for assessment purposes. A review of tagging studies for winter flounder (Howell 1996) indicates dispersion (and hence mixing) has occurred between the previously defined Southern New England and Mid-Atlantic units. Howell (1996) noted that differences in growth and maturity among samples from Southern New England to the Mid-Atlantic may reflect discrete sampling along a gradient of changing growth and maturity rates over the range of a stock complex. Differences in growth rates within the Mid-Atlantic unit were observed to be greater than differences between Mid-Atlantic and Southern New England units (Howell, 1996). In offshore waters, the length structure of winter flounder caught in NEFSC research surveys is similar from Southern New England to New Jersey. Most commercial landings are obtained in these offshore regions (greater than 3 miles from shore).

Stock Boundaries and associated Statistical Areas

The Gulf of Maine stock complex extends along the coast of eastern Maine to Provincetown, MA, corresponding to NEFSC commercial fishery statistical division 51. Recreational landings from Maine, New Hampshire and northern Massachusetts (northern half of Barnstable County and north to New Hampshire border) are associated with this stock complex.

The Southern New England/Mid-Atlantic stock complex extends from the coastal shelf east of Provincetown, MA southward along the Great South Channel (separating Nantucket Shoals and Georges Bank) to the southern geographic limits of winter flounder. NEFSC commercial fishery statistical areas within this boundary are 521 and 526, and statistical divisions 53, 61, 62, and 63. The corresponding recreational areas are southern Massachusetts (the southern half of Barnstable County; Dukes, Nantucket and Bristol counties), Rhode Island, Connecticut, New York, New

Jersey, Delaware, Maryland and Virginia. NEFSC survey strata included for this stock extend from the waters of outer Cape Cod to the south and west.

The Georges Bank stock extends eastward of the Great South Channel, including statistical areas 522, 525, and 551-562.

FISHERY DATA

Landings

After reaching an historical peak of 11,977 metric tons (mt) in 1966, then declining through the 1970s, total U.S. commercial landings again peaked at 11,176 mt in 1981, and then steadily declined to a record low of 2,159 mt in 1994. Landings have increased since 1994 to 4,448 mt in 2001 (Table B1.1, Figure B1.1). During 1989-1996, an average of 43% of commercial landings were taken from statistical area 521, 13% from area 526, 13% from area 537, and 11% from area 539, with the remaining landings (20%) obtained from area 538 and divisions 61-62 (Table B1.2). Since 1993, a larger percentage of the commercial landings has been taken from area 521. An unusually high proportion of the commercial landings for the stock complex was reported from NEFSC statistical area 521 in 1997 and 2001, with 62% in 1997 and 56% in 2001. When considered along with the distribution of survey catches, this factor indicates that the commercial fishery is focused on winter flounder along the western side of the Great South Channel. The primary gear in the fishery is the otter trawl which accounts for an average of 95% of landings since 1989. Scallop dredges account for 4%, with handlines, pound nets, fyke nets, and gill nets each accounting for about 1% of total landings.

Recreational landings reached a peak in 1984 of 5,772 mt but declined substantially thereafter (Table B1.3, Figure B1.1). Landings have been less than 1,000 mt since 1991, with the lowest estimated landings in 1998 of 290 mt. Landings in 2001 from the Southern New England/Mid Atlantic stock complex were 552 mt. The principal mode of fishing is private/rental boats, with most recreational landings occurring during January to June.

Sampling Intensity

Length samples of winter flounder are available from both the commercial and recreational landings. In the commercial fishery, annual sampling intensity varied from 63 to 264 mt landed per 100 lengths measured during 1981-1997 (Table B1.4). Overall sampling intensity was 90 mt per 100 lengths in 1998, 75 mt per 100 lengths in 1999, 59 mt per 100 lengths in 2000, and 71 mt per 100 lengths in 2001 (Table B1.5). In the recreational fishery, annual sampling intensity varied from 36 to 231 mt landed per 100 lengths measured during 1981-1997 (Table B1.6). Overall sampling intensity was 47 mt per 100 lengths in 1998, 81 mt per 100 lengths in 1999, 519 mt per 100 lengths in 2000, and 109 mt per 100 lengths in 2001 (Table B1.7).

Landed Age Compositions

Commercial fishery

In the SARC 21 assessment (NEFSC 1996), numbers at age were estimated for 1985-1993 for commercial landings, recreational landings, commercial discards, and recreational discards. Quarterly or half-year commercial age-length samples were applied to corresponding commercial market category landings at length. Unsampled unclassified landings and landings not represented in the weighout database (i.e., state canvas landings) were assumed to have the same age composition as the initial weighout commercial landings at age. Landings at lengths with no associated age data within the quarter were assigned ages based on age at length from adjacent quarters. A comparison was undertaken among age data collected from inshore regions (where the recreational fishery is prosecuted), to determine if all age data were comparable within the stock complex. Data for ages 3-5 from New Jersey, Connecticut, Massachusetts and NEFSC were compared for 1993-1994. Distributions of length at age from New Jersey and Connecticut were similar, while distributions of length at age from Massachusetts lacked smaller fish at age (Howell 1996).

In the ASMFC 1998 assessment (ASMFC 1998), the Technical Committee attempted to update the catch at age matrix for VPA for 1994-1996. Two key market categories of commercial landings were found to lack port samples: medium fish in the second half of 1995 and large fish in the first half of 1996. In addition, several market categories were poorly sampled: medium fish in the first and second half-year of 1996, and large fish in the second half of 1995. The Technical Committee concluded then that the port sampling was insufficient to characterize the length and age frequency of the commercial landings for 1995-1996, and elected to use a non-age dependent model (ASPIC) to assess the stock complex (ASMFC 1998).

In the SARC 28 assessment (NEFSC 1999), commercial fishery port samples for 1995 and 1996 were supplemented with commercial fishery sea sample length data for the second half of 1995 and 1996, to continue the catch at age series. For the second half-year of 1995, 2,979 sea sample lengths (unclassified by market category) were used in place of the available 702 port sample lengths to construct an unclassified length frequency for the second half-year of 1995 landings. For the first half-year of 1996, 55 sea sample lengths were combined with 752 port sample lengths to create an unclassified frequency of 807 lengths for the first-half year of 1996 landings. Also, archived NEFSC research survey and commercial fishery age samples were aged, allowing extension of the NEFSC survey catch at age series back to 1980 and of the fishery catch at age matrix back to 1981 (Table B1.4). Since 1997, port sampling has been adequate to develop the commercial fishery landings at age on a half-year, market category basis across all statistical areas (Tables B1.5 and B1.10).

Recreational fishery

Recreational landings at length were estimated seasonally (January-June and July-December) and geographically. Landings were divided into two geographic regions; 1) Massachusetts and Rhode Island (SNE) and 2) Connecticut and south (MA). For the 1981-1984 period, NEFSC spring age-length keys were used to age both area length frequencies. For 1985-1996, MADMF

survey age-length keys were applied to MA-RI data while CTDEP age-length keys were applied to CT-south data, with the exception of 1993 landings which used a combined NJ/CT age-length key. Since 1997, NEFSC spring and fall keys have been used to age all length frequencies (Tables B1.6, B1.7, and B1.10). For the 1998-1999 recreational catch at age, sample lengths were applied to catch numbers on an annual basis for the two regions, due to low samples size. For the 2000-2001 recreational catch at age, sample lengths were applied to catch numbers on an annual basis for the regions combined, due to low sample sizes in the SNE region (Table B1.7).

Discard estimates and age compositions

Commercial fishery

In the SARC 21 assessment (NEFSC 1996), the Working Group and the SARC concluded that there were too few Fishery Observer sampled trips in which winter flounder were caught to adequately characterize the overall ratio of discards to landings in the commercial fishery. The Fishery Observer sample length frequency data, however, were judged adequate to help characterize the proportion discarded at length. In the SARC 21 assessment, commercial discards for 1985 to 1993 were estimated from length frequency data from NEFSC and the Massachusetts Division of Marine Fisheries (MADMF) bottom trawl surveys, commercial port sampling of landings at length and Fishery Observer sampling of landings and discard at length. The method follows an approach described by Mayo et al. (1992). The year was divided into half year periods. Survey length frequency data (MADMF survey in spring and NEFSC in fall) were smoothed using a three point moving average, then filtered through a mesh selection ogive (Simpson 1989) for 4.5 inch mesh (1984-1989), 5 inch mesh (1990-1992, fall 1993) or 5.5 inch mesh (spring, 1993). The 5.5 inch mesh selection curve was calculated using the 5 inch curve adjusted to an L_{50} for 5.5 inch mesh. The choice of mesh sizes was based on sizes used in the yellowtail assessment for southern New England (Rago et al. 1994) and comparison to length frequencies of commercial landings. The mesh filtering process resulted in a survey length frequency of retained winter flounder. A logistic regression was used to model the percent discarded at length from 1989-1992 sea sampling data, and the resulting percentages at length were applied to the survey numbers at length data to produce the survey-based equivalent of commercial kept and discarded winter flounder. The 1989-1992 average percentage discard at length was applied to 1985-1988. The survey numbers per tow at length "kept" were then regressed against commercial (weighout) numbers landed at length. The linear relationship was calculated for those lengths common to both length frequencies and fitted with an intercept of zero. The slope of the regression provided a conversion factor to re-scale the survey "discard" numbers per tow at length to equivalent commercial numbers at length. The resulting vector of number of fish discarded at length was multiplied by a discard mortality rate of 50% (as averaged in Howell et al., 1992) to produce the vector of fish discarded dead at length per half year. The number of dead discards at length was adjusted by the ratio of weighout landings to total commercial landings and summed across seasons and lengths (and corresponding weight at length) to produce the annual total number and weight of commercial fishery discards for 1985-1993 (Tables B1.10-11, Figure B1.1). In the SARC 28 assessment (NEFSC 1999), this same method using the 4.5 inch mesh ogive and 1989-1992 average discard percentage at length was

used to estimate commercial fishery discards for 1981-1984. NEFSC spring and fall survey age-length keys were applied to convert discard length frequencies to age.

During ASMFC Winter Flounder Technical Committee meetings since 1995, the group has considered the SARC 21 survey length-mesh selection method, NEFSC Fishery Observer data (OB), and NER Vessel Trip Report (VTR) data as sources of information to use in the estimation of commercial fishery discards, with a focus on the latter two sources. The Committee examined the characteristics of both the Fishery Observer and VTR discard data (number of trip samples, frequency distributions of discards to landings ratio per trip, mean and variance of annual half-year discards to landings ratio), and concluded that the VTR mean discard to landed ratio aggregated over all trips in annual half-year season strata (January to June, July to December) provided the most reliable data from which to estimate commercial fishery discards. VTR trawl gear fishery discards to landings ratios on a half-year basis (January to June; July to December) were applied to corresponding commercial fishery landings (all gears) to estimate discards in weight (Table B1.8, Figure B1.1). The Fishery Observer length frequency samples were judged adequate to directly characterize the proportion discarded at length (Table B1.9). The sample proportion at length, converted to weight, was used to convert the discard estimate in weight to numbers at length. As in the SARC 28 assessment (NEFSC 1999), the resulting number of fish discarded at length was multiplied by a discard mortality rate of 50% (as averaged in Howell et al., 1992) to produce the number of fish discarded dead at length. For 1998, discard estimates at length were made by half-year; for 1999-2001, samples length were applied on an annual basis due to low sample sizes (Table B1.9). NEFSC Spring and Fall survey age-length keys were used to convert the discard length frequency to age (Table B1.10).

Recreational fishery

A discard mortality of 15% was assumed for recreational discards (B2 category from MRFSS data), as assumed in Howell et al. (1992). Discard losses peaked in 1984-1985 at 0.7 million fish. Discards have since declined reaching a low in 1999 of 62,000 fish. In 2001, 81,000 fish were estimated to have been discarded (Table B1.3). In the SARC 21 assessment (NEFSC 1996), recreational discards for 1985-1993 were assumed to have the same average weight per fish as spring commercial discards, providing estimates of the total weight of recreational discards ranging from 15 mt in 1992 to 230 mt in 1985. Estimates of recreational discard at age for 1985-1993 were developed using state survey length and age data in a manner similar to that for the commercial discard estimates (Tables B1.10-11; see Gibson (1996) for complete description of computation of 1985-1993 recreational discard numbers at length and age).

The SARC was unable to apply the 1985-1993 method to the 1994-1997 or 1981-1984 periods for the SARC 28 assessment, due to data availability problems (NEFSC 1999). Instead, for 1994-1997, the average proportion at age in the 1991-1993 recreational discard was used to apportion the recreational fishery estimate of discard in numbers to length and age. These discards at age were assumed to have the same mean weight as the landed portion at the same ages, and so this method probably slightly overestimates the discard in weight. For 1981-1984, before implementation of the 12 inch (30 cm) minimum landing size in most states (which

encompasses fish up to age 3), it was assumed that all recreational discard would be age 1 and age 2 fish, and so the discard was allocated to ages 1 and 2 in the same relative proportion as those in the landings, and assumed to have the same mean weight at age. SARC 28 (NEFSC 1999) concluded that since the magnitude of the recreational discard is relatively small compared to the total landings and commercial discards, error in estimation of recreational discard at age due to different methods over the time series and/or error in allocation among ages 1 and 2 would have a minimal effect in terms of estimation of population sizes in the VPA.

Since 1997, irregular sampling of the recreational fisheries by state fisheries agencies has indicated that the discard is usually of fish below the minimum landing size of 12 inches (30 cm). For 1998-2001, the recreational discard has been assumed to have the same length frequency as the landed portion of the catch below 12 inches, and so is still predominantly ages 1, 2, and 3 fish. As with the recreational landings, sample lengths were applied to catch numbers on an annual basis for the two regions for 1998-1999, and on an annual basis for the regions combined for 2000-2001. The recreational discard for 1998-2001 is aged using NEFSC survey spring and fall keys (Table B1.10).

Mean Weights at Age in the Catch

Mean weights at age were determined for the landings and discards in the commercial and recreational fisheries. Length frequencies (cm) for each component were converted to weight (kg) using length-weight equations derived from NEFSC survey samples:

$$\begin{array}{ll} \text{Spring surveys:} & \text{wt} = 0.00000997 * \text{length}^{3.055236} \\ \text{Fall surveys:} & \text{wt} = 0.00000925 * \text{length}^{3.095188} \end{array}$$

The equations from the spring and fall surveys were applied to catches during the corresponding time periods. The annual mean weights at age from the commercial and recreational fisheries were used in the virtual population analysis and yield per recruit calculations.

Total Catch

Estimates of the total catch of winter flounder during 1981-2001 are presented in Table B1.11. These estimates include commercial and recreational landings and discards. The total catch during this period has varied from a high of 15,788 mt (34.6 million fish) in 1984 to a low of 3,095 mt (3.6 million fish) in 1994. The total catch has increased since 1995 to 5,102 mt (9.0 million fish) in 2001 (Table B1.11, Figure B1.1). Total catch and mean weights at age as aggregated for input to the VPA (ages 1-7+) are presented in Tables B1.12-13, and Figures B1.2-3.

RESEARCH SURVEY ABUNDANCE AND BIOMASS INDICES

State and federal surveys were evaluated as fishery independent indices of winter flounder abundance and biomass. Survey methods (with the exception of Rhode Island and the young-of-

year surveys) are reviewed in the proceedings of a 1989 trawl survey workshop sponsored by the ASMFC (Azarovitz et al., 1989).

NEFSC

Mean weight and number per tow abundance indices were determined from fall (1963-2001) and spring (1968-2002) NEFSC bottom trawl surveys. Indices from the spring and fall surveys were based on tows in offshore strata 1-12, 25, and 69-76 and inshore strata 1-29 and 45-56. Spring indices prior to 1973 and fall indices prior to 1972 do not include inshore strata. In addition, offshore surveys from 1963-1966 were not conducted south of Hudson Canyon.

A new series of NEFSC winter trawl surveys was begun in February 1992 specifically to provide improved indices of abundance for flatfish, including winter flounder. A modified 36 Yankee trawl is used in the winter survey that differs from the standard trawl employed during the spring and fall surveys in that 1) long trawl sweeps (wires) are added before the trawl doors, to better herd fish to the mouth of the net, and 2) the large rollers used on the standard gear are absent, and only a chain "tickler" and small spacing "cookies" are present on the footrope. This gear is intended to better target flatfish than the gear used in the spring and fall surveys. The geographical coverage of the winter survey is more limited than the spring and fall surveys, due to time limitations and the use of the flatfish net. Inshore strata and offshore deep strata are irregularly sampled, strata east of the Great South Channel are irregularly sampled, and the Gulf of Maine has never been sampled. For winter flounder, the winter survey indices include offshore strata 1-2, 5-6, 9-10, 69, and 73; generally the offshore between 27 to 110 meters depth (15 to 60 fathoms).

Mean weight per tow and number per tow indices for the spring, fall, and winter time series are presented in Table B1.14. Indices dropped from the beginning of the time series in the 1960s to a low point in the early to mid- 1970s, then rose to a peak by the early 1980s. Following several years of high indices, abundance once again declined to below the low levels of the 1970s. NEFSC survey indices reached near- or record low levels for the time series in the late 1980s-1990s. Indices from the three survey series generally increased during 1993-1998/1999, but have since declined (Figure B1.4).

Massachusetts

The Massachusetts Division of Marine Fisheries (MADMF) spring survey from 1978-2001 was used to characterize the abundance of winter flounder. Survey areas from east and south of Cape Cod were used in the analysis (strata 11-21). The MADMF mean number per tow indices steadily declined from a high value of 53.79 in 1979 to a low of 10.66 in 1991, and then increased to 30-40 fish per tow during 1995-1998, before falling again to 16.00 in 2001. Mean weight per tow indices have varied in a similar manner over the time series, ranging from 15-20 kg/tow in the early 1980s to about 5 kg/tow during 2000-2001 (Tables B1.15-16, Figure B1.4).

The MADMF also conducts an annual juvenile winter flounder seine survey during June. The survey has been conducted since 1975 in coastal ponds and estuaries. The index has shown a

general decline in production, with a high of 0.60 fish per haul in 1977 to a low of 0.07 fish per haul in 1993. The 1997 value was 0.39 fish per haul, and has since declined to 0.10 in 2002 (Table B1.17, Figure B1.5).

Rhode Island

The Rhode Island Division of Fish and Wildlife (RIDFW) conducts a number of research surveys in Narragansett Bay and Rhode Island coastal waters. A seasonal trawl survey was initiated in 1979 to monitor finfish stocks in Narragansett Bay, Rhode Island Sound and Block Island Sound. The survey employs a stratified random design and collects length, weight, and abundance information. Survey results are expressed as un-weighted catch per tow (Tables B1.15-16). Spring survey indices from 1979-2001 showed a steady decline from high values during 1979-1981 (12-13 kg per tow, 63-88 fish per tow) to a low of 0.22 kg per tow and 2.92 fish per tow in 1993. Spring indices increased to 5.83 kg per tow and 31.78 fish per tow in 1995, before declining again to 3.56 kg per tow and 12.49 fish per tow in 2001 (Figure B1. 4). Fall survey indices show similar trends, with peak abundance and biomass during the early and mid 1980s, a decline to low values in the mid-1990s, some rebound during 1995-1997, and a recent decline (Tables B1.15-16).

A juvenile finfish beach seine survey, conducted from June to October since 1986, takes monthly samples at 17 fixed stations in Narragansett Bay. This seine survey provides an index of young-of-year winter flounder. The index shows a great deal of annual variability, although in recent years there have been consistently high levels of recruitment. The index of the 2000 year class is the highest of the time series (Table B1.17, Figure B1.5).

Connecticut

The Connecticut Department of Environmental Protection (CTDEP) Long Island Sound Trawl Survey (1984-present) uses a stratified-random design to sample Connecticut and New York waters of the Sound from Groton to Norwalk. Forty sites are sampled monthly (Apr-June, Sept-Oct) across three sediment (mud, sand, transitional) and four depth intervals (<30 ft, 30-60 ft, 60-90 ft, 90+ft). A 14 m otter trawl with 51 mm codend is towed for 30 min at 3.5 kts from a 15.2 m research vessel.

Winter flounder abundance indices are based on April and May sampling. Winter flounder are counted and measured from each tow. Since 1992 composite biomass (0.1kg) has also been recorded from each tow. Otoliths are collected for aging each spring. Aging samples are stratified by month, area (east/west) and size. Subsamples of 5-7 fish per centimeter are collected from fish up to 36 cm and all fish over 36 cm are retained for aging. Aged fish are measured and weighed in the lab and gonad condition is recorded. Gonad weights were also recorded in some years. In recent years approximately 800 flounder have been aged annually. Otoliths are generally aged whole, however larger fish and difficult bones are sectioned for reading. Indices at age are calculated as a proportion of the overall index. Age length keys are applied by area (east-west)

and year where possible and any remaining unaged fish are aged using a pooled area/year key as necessary.

CTDEP indices exhibited several years of high values between 1988 and 1991, declined to a minimum in 1995, and have since increased to about one-half the time series average during 2000-2002 (Tables B1.15-16, Figure B1.4). A separate young of the year survey index shows above average recruitment during 1994-1996, and below average recruitment since. The 2001 year class index is the smallest of the time series (Table B1.17, Figure B1.5).

New York

The New York Department of Environmental Conservation (NYDEC) has conducted a small-mesh trawl survey in Peconic Bay since 1985. Winter flounder indices for ages 0 and 1 were evaluated for trends in winter flounder abundance (Tables B1.16-17, Figure B1.5). Young of the year indices have increased in recent years from 0.7 in 1985 to the 1993 index of 4.7 and 1996 index of 3.80. The 1992 index indicated the strongest recent year class with an index of 11.4. The corresponding age 1 indices also indicated strong 1992, 1993, and 1996 year classes.

New Jersey

The New Jersey Division of Fish, Game and Wildlife (NJDFW) has conducted a bottom trawl survey in near-shore ocean waters of the state since 1989, and in inshore waters in the Shark and Manasquan Rivers since 1995. Ocean survey samples are collected via a stratified random bottom trawl survey. Surveys are usually conducted in January, April, June, August, and October. Inshore samples are collected via a random station trawl survey in the main channel of the Shark and Manasquan Rivers. Sampling is conducted in March, April and May and results are pooled to calculate mean number per tow indices. Aging of NJDFW samples started in 1993. During both surveys, a sub-sample of fish are aged (fish are aged from April ocean survey only). Age/length keys are constructed, and all lengths are transformed to ages by applying all lengths to the age/length keys. Number at each age are divided by the number of tows to derive catch at age per tow.

Ocean survey indices (mean number per tow in April) tended to decline between 1989 and 1993, and have been quite variable since 1994, with a time series low in 1996, increasing to above the time series mean in 2002 (Tables B1.15-16, Figures B1.4-5). River survey indices exhibit no trend over the short time series (Table B1.16, Figure B1.5).

Delaware

The Delaware Division of Fish and Game (DEDFG) conducts monthly surveys from April to October using a 16 ft. semi-balloon otter trawl with a 0.5 inch stretch mesh liner. An index of young-of-year winter flounder was developed from stations sampled within Indian River and Rehoboth Bays. The re-transformed annual geometric means, presented in Table B1.17, indicate variable annual recruitment with a large year class in 1990. The 1994 index indicates above average recruitment (Table B1.17, Figure B1.5).

ESTIMATES OF MORTALITY AND STOCK SIZE

Natural Mortality and Maturity

Instantaneous natural mortality (M) for winter flounder was assumed to be 0.20 and constant across ages. Commercial catch at age included fish to age 14, under conditions of relatively high fishing mortality. If M = 0.25, less than 5% of the population would reach age 12 under conditions of no fishing mortality. Therefore, the SARC judged that M = 0.20, which represents a maximum age of 15, was representative of the stock complex throughout its range.

In the SARC 28 review of the SNE/MA winter flounder stock assessment (NEFSC 1999), the SARC recommended re-examination of the maturity schedule used in the yield per recruit (YPR) and virtual population analyses (VPA) to incorporate any recent research results. The SARC 28 and previous assessments used the maturity schedule as published in O'Brien et al. (1993) for winter flounder south of Cape Cod, based on data from the MADMF spring trawl survey for strata 11-21 (state waters east of Cape Cod, Nantucket sound, Vineyard Sound, and Buzzards Bay) sampled during 1985-1989 (n = 301 males, n = 398 females). Those data provided estimates of lengths and ages of 50% maturity of 29.0 cm and 3.3 yr for males, and 27.6 cm and 3.0 yr for females, and the following estimated proportions mature at age. The female schedule (with the proportion at age 2 rounded down to 0.00) was used in the SARC 28 assessment YPR and VPA (NEFSC 1999).

Age	1	2	3	4	5	6	7+
Males	0.00	0.04	0.32	0.83	0.98	1.00	1.00
Females	0.00	0.06	0.53	0.95	1.00	1.00	1.00

In response to the SARC 28 recommendation, the SARC has examined NEFSC spring trawl survey data over the 1981-2001 period in an attempt to better characterize the maturity characteristics of the SNE/MA winter flounder stock complex. Data from the NEFSC survey included those judged in the SARC 28 assessment to comprise the SNE/MA complex from Delaware Bay to Nantucket Shoals: NEFSC offshore strata 1-12, 25 and 69-76, and inshore strata 1-29, 45-46. Note that this is a much larger geographic area than that included in the MADMF survey data used in O'Brien et al. (1993). Data were analyzed in 5-6 year blocks (1981-1985, 1986-1990, 1991-1995, and 1996-2001) and for the entire time period (1981-2001), for each sex and combined sexes. Observed proportions mature at age were tabulated, and from those data maturity ogives at length and age were calculated to provide estimated proportions mature at age.

In general, the NEFSC maturity data indicated earlier maturity than the MADMF data, with L50% values ranging from 22-25 cm, rather than from 28-29 cm, and with ~50% maturity for age 2 fish, rather than ~50% maturity for age 3 fish. To investigate the apparent inconsistency between the MADMF and NEFSC maturity data, the SARC compared the two data sets over the same time periods (1985-1989, 1990-1995, 1996-2001) for common/adjacent survey strata (MADMF strata 11-12; NEFSC inshore strata 50-56 and offshore strata 10-12 and 25). Note that

the MAMDF data now have about 160 observations for the 1985-1989 period that were added subsequent to the O'Brien et al. (1993) work. For comparable time periods and geographic areas, the NEFSC maturity data still consistently indicated a smaller size and younger age of 50% maturity than the MADMF data. NEFSC L50% and A50% values range from 22-26 cm and about 2.0 yr, while the MADMF values range from 27-30 cm and about 3.0 yr. The difference in values from this comparison was not as large as for the full NEFSC data set extending southward to Delaware Bay, which incorporates components of the stock complex that mature at smaller sizes and younger ages. However, the difference is still nearly a full age class difference at 50% maturity.

Given that both length and age vary in the same direction, it seems unlikely that the differences could be attributed to aging differences between the two data sets. Since the MADMF and NEFSC geographic areas in this comparison do not match exactly, the difference in maturity rates may be due to the extension of the NEFSC strata to somewhat deeper waters inhabited by fish that mature at a smaller size and younger age (inclusion of fish in offshore strata were necessary for sufficient sample size). Alternatively, for the size range of fish in question (20 to 30 cm length), it may be that immature and mature fish are segregated by area, with mature fish in that size interval tending to occupy inshore areas during the spring, with immature fish tending to remain offshore. Finally, there may be differences in the accuracy and consistency of interpretation of maturity stage between MADMF and NEFSC survey staff.

The SARC considered these data and analyses and the possible causes for the noted inconsistencies, and concluded that more detailed spatial and temporal analyses are needed before revisions to the maturity schedule can be adopted. Therefore, the maturity at age schedule used in the SARC 28 assessment (see above) has been retained for this assessment.

Total Mortality from Mark and Recapture Data

Total mortality in two components of the stock were evaluated using recent tag and recapture data. Northeast Utilities Co. marked and recaptured winter flounder in eastern Long Island Sound from 1983-1998 and the RIDFW has conducted winter flounder tagging programs in Narragansett Bay from 1986-1990 and again from 1996-1998. Mortality estimates were made by maximum likelihood methods using the Brownie class of survivorship models (Brownie et al. 1985). Average estimates of fishing mortality for Long Island Sound averaged 0.59 from 1984-1988 and 0.77 from 1989-1993, and 0.65 from 1993-1996. Fishing mortality in 1996 was estimated to be 0.56. Narragansett Bay estimates of fishing mortality ranged from 0.81 to 1.92 and averaged 1.19 from 1986 to 1989. The most recent tag releases in Narragansett Bay indicate that F had dropped to 0.37 in 1996-1997.

Virtual Population Analysis

Tuning

The Virtual Population Analysis (VPA) was tuned (calibrated) using the NEFSC Woods Hole Fisheries Assessment Compilation Toolbox (FACT) version 1.50 of the ADAPT VPA (Conser and Powers 1990). Abundance indices at age (Tables B1.18-25) were available from several

bottom trawl surveys: NEFSC spring bottom trawl ages 1-7+, NEFSC fall ages 1-5 (advanced to tune January 1 abundance of ages 2-6), NEFSC winter ages 1-5, MADMF spring ages 1-7+, RIDFW fall seine age 0 (advanced to tune age-1), RIDFW spring ages 1-7+, CTDEP spring ages 1-7+, NYDEC age 0 (advanced to tune age-1) and age-1, MADMF summer seine index of age-0 (advanced to tune age-1), DEDFG juvenile trawl survey age-0 (advanced to tune age-1), NJDFW Ocean trawl survey ages 1-7+, and NJDFW River trawl survey ages 1-7+. The indices from the NEFSC winter trawl survey, NYDEC, and NJDFW were included in the VPA tuning for the first time. Survey indices were selected for inclusion in VPA tuning based on consideration of the partial variance in a VPA trial run including all indices, residual error patterns from the various trial runs, and on the significance of the correlation among indices and with VPA abundance estimates from the trial run including all indices.

The SARC considered eight different configurations of tuning indices. In general, tuning indices were excluded if they exhibited high partial variance (indicating a lack of fit within the VPA model) and low correlation with other indices with similar spatial and temporal characteristics and with the VPA estimates of 2002 stock size. Run W36ALL was the initial trial including all indices. Run W36_28 used the same suite of indices as that selected for the SARC 28 VPA (NEFSC 1999), and therefore did not include new indices available from the NEFSC Winter trawl survey, the NYDEC indices, or the two NJDFW index series. Run W36_1 excluded eight indices with high partial variance within the VPA and low correlation with other indices and/or the VPA estimates of stock size, resulting in improvements both in overall fit (Mean Square Residual (MSR) reduced by 14%) and in the precision of the stock size estimates, relative to the W36ALL configuration. Run W36_2 dropped an additional seven indices from the W36_1 configuration, resulting in further improvements in fit (21% improvement over run W36_1) and precision. This was the run adopted as final by the SARC, and is the basis for all further analyses (Table B1.26).

Run W36_3 dropped an additional two indices (from W36_2) to exam the trade-off between overall model fit (MSR) and the precision of the 2002 stock size estimates as degrees of freedom were further reduced. The SARC concluded that the improvement of run W36_3 in overall fit (7%) was balanced by the decrease in precision at ages 6 and 7+, and so retained run W36_2 as final. Two additional runs excluded all state agency indices (W36NEC) and excluded all NEFSC indices (W36STATE). The W36NEC exhibited a better fit than the W36_2 run, but much lower precision of the 2002 stock size estimates, reflecting the fewer degrees of freedom available. The W36STATE run exhibited the poorest fit of the eight considered, along with the lowest precision of the 2002 stock sizes at ages 4 and older. Run W36_28 provided results intermediate to those from the W36_2 and W36STATE runs. Finally, run W36_2IR was the same as run W36_2, but incorporated the iterative re-weighting option of the VPA tuning, which in a second step of tuning gives more influence to indices that fit best within the analysis (tuning weight in inverse proportion to initial fitted variance). The W36_2IR results were very similar to those of runs W36_2 and W36_3 (Figure B1.6).

Stock size estimates for 2002 in the final W36_2 calibration were moderately precise (initial coefficients of variation ranged from 0.21 at age-3 to 0.38 at age-1). Nearly all surveys had years in which all observations deviated from predicted values in the same direction. For example, most surveys exhibited blocks of negative residuals during the late 1980s, and then blocks of positive residuals during the mid to late 1990s, when residuals for all ages are summed within year and survey series. Residuals by age exhibit a similar pattern of blocking, and a tendency for blocks of positive residuals at younger ages during the mid-1990s to move to older ages later in the VPA time series. This pattern of residuals (i.e., overestimation of stock size by the surveys during the mid-to late 1990s) is reflective of the retrospective pattern of VPA estimates evident for terminal years 1995-1999 (see the following Retrospective Analysis section). The correlation analysis of tuning indices also indicated that there are strong year effects in survey indices, due to annual distribution patterns or local recruitment events. However, in concert, the SARC concluded that the surveys appear to provide geographically balanced tuning.

Exploitation Pattern

The exploitation pattern has been variable from year to year, but with the exception of 1996-1997, age-4 fish have been 80%-100% recruited since 1993 (Table B1.26). The SARC noted a recent tendency for partial recruitment at age to decrease substantially at ages 5 and 6 in the terminal year, but further noted that the retrospective analysis indicates that this tendency does not persist, with the expected, flat-topped partial recruitment pattern becoming evident as the VPA converges. For this reason, the average exploitation pattern to be used in yield per recruit analysis and stock projections was calculated as the geometric mean fishing mortality rates for 1998-2000, normalized to age 4. The resulting pattern indicates 1% recruitment at age-1, 27% at age-2 and 75% at age-3. For purposes of yield per recruit and stock projections, full (100%) recruitment was assumed at ages 4 and older. For consistency with the partial recruitment averages, mean weights at age in the landings, discards, and spawning stock biomass were also averaged over 1998-2000.

Fishing Mortality, Spawning Stock Biomass, and Recruitment

During 1981-1993, fishing mortality (fully recruited F , ages 4-5) has varied between 0.4 (1982) and 1.4 (1988), and was as high as 1.2 as recently as 1997. Fishing mortality has been in the range of 0.5-0.6 during 1999-2001 (Table B1.26, Figure B1.7). Accounting for the uncertainty of the 2001 estimate, there is an 80% probability that F in 1997 was between 0.44 and 0.58 (Figure B1.8).

SSB declined from 14,800 mt in 1983 to a record low of 2,700 mt in 1994. SSB has increased since 1994 to 7,600 mt in 2001 (Table B1.26, Figure B1.9). Accounting for the uncertainty of the 2001 estimate, there is an 80% probability that SSB in 2001 was between 6,800 mt and 8,400 mt (Figure B1.8). Recruitment declined from 62.9 million age-1 fish in 1981 to 7.8 million in 1992. Recruitment then averaged 14.7 million fish during 1993-2001, below the VPA time series average of 23.9 million. The 2001 year class is estimated to be the smallest in 22 years, at only 5.7 million fish (Table B1.26, Figure B1.9).

Retrospective analysis

A retrospective analysis of the VPA was conducted back to a terminal catch year of 1997. The VPA exhibits a retrospective pattern of underestimation of F and overestimation of SSB during the late 1990s. The most likely cause of this pattern is a combination of factors including under-reporting of the landings, mis-classification of the landings by stock area, and underestimation of the discards. For 1995-1999, retrospective fishing mortality rates underestimate the current values by an average of 128%, ranging from 232% for 1997 to 44% for 1995. The pattern reversed for 2000 (i.e., F was overestimated), and fishing mortality appears to have been overestimated for 2000 by 7%. The retrospective pattern for spawning stock biomass has been a tendency for overestimation since 1991. The overestimation of SSB averaged 76% from 1995-1999, and was largest for the 1997 and 1998 terminal years (115% and 98% overestimation). The retrospective estimation of age-1 recruits indicated a tendency for overestimation during 1993-2000, with recruitment apparently underestimated for 2001 (2000 year class; Table B1.26, Figure B1.10).

Precision of Stock Size, F , and SSB estimates

The precision of the 2002 stock size, fishing mortality at age in 2001, and SSB estimates from VPA was evaluated using bootstrap techniques (Efron 1982). Five hundred bootstrap iterations were realized in which errors (differences between predicted and observed survey values) were resampled. Estimates of precision and bias are presented in Table B1.27. Bootstrap estimates of stock size at age indicate low bias (<6%) for ages 1-7+ and bootstrap standard errors provide stock size CVs ranging from 18% at age 3 to 34% at age 1.

Bootstrapped estimates of spawning stock biomass indicate a CV of 9%, with low bias (bootstrap mean estimate of spawning stock biomass of 7,705 mt compared with VPA estimate of 7,643 mt; Table B1.27). There is an 80% probability that spawning stock in 2001 was between 6,800 mt and 8,400 mt (Figure B1.8).

The bootstrap estimates of standard error associated with fishing mortality rates at age indicate good precision. Coefficients of variation for F estimates ranged from 16% at age 3 to 21% at ages 1, 6 and 7+ (Table B1.27). There is an 80% probability that fully recruited F for ages 4-5 in 2001 was between 0.44 and 0.58 (Figure B1.8).

BIOLOGICAL REFERENCE POINTS

Yield and Spawning Stock Biomass per Recruit; Stock-recruitment model

NEFSC (2002) re-estimated the biological reference points for SNE/MA winter flounder in 2002 using yield and SSB per recruit (Thompson and Bell 1936) and Beverton-Holt stock-recruitment models (Beverton and Holt 1957, Brodziak et al. 2001, Mace and Doonan 1988) based on the SARC 28 assessment (NEFSC 1999). The yield and SSB per recruit analyses indicate that $F_{40\%} =$

0.21 and $F_{0.1} = 0.25$ (Figure B1.11). The parametric stock-recruitment model indicated that $MSY = 10,600$ mt, $F_{msy} = 0.32$, and $SSB_{msy} = 30,100$ mt (Figure B1.12).

Biological reference points estimated in NEFSC (2002) were updated by the SARC with the partial recruitment pattern and mean weights at age for 1998-2000 (as noted earlier, the 2001 estimates were not included in the averages due to the retrospective variability of the partial recruitment pattern in the terminal year of the VPA). Given the stability of the input data to these analyses and the consistency of the results with the previous work, the SARC elected to retain the NEFSC (2002) estimates of biological reference points for this assessment. The SARC recommends that these parametric stock-recruitment model reference points be adopted as the basis for the ASMFC and NEFMC FMP overfishing definitions.

PROJECTIONS FOR 2002-2013

Stochastic projections were made based on 500 bootstrapped VPA realizations of stock size in numbers at age in 2002. The stochastic forecasts only incorporate uncertainty in 2002 stock sizes due to survey variability, assume current discard to landings proportions, and are not adjusted for the retrospective pattern in VPA stock size estimates. Partial recruitment to the fishery and percentage discarded were estimated as the geometric mean of VPA estimates for 1998-2000. The 2001 estimates were not included in the averages due to the retrospective variability of the partial recruitment pattern in the terminal year of the VPA. For consistency with the partial recruitment averages, mean weights at age in the stock, landings, and discards were similarly estimated as the weighted (by number landed) geometric mean weight at age from 1998-2000. Age-1 recruitment levels in 2003 and later years are estimated from the stochastic, parametric stock-recruitment relationship estimated in NEFSC (2002). Projections were made through 2013 to respond both to the ASMFC terms of reference and more recent NEFMC Plan Development Team requirements.

If F in 2002 is assumed to be 15% less than F in 2001 ($F_{2002} = 0.43$), due to the impact of management measures implemented in response to court orders during 2002, then landings are expected to be about 3,000 mt in 2002. At this reduced F , spawning stock biomass is projected to fall to 5,900 mt in 2002. Given $F = 0.43$ in 2002, a fishing mortality rate of $F_{reb} = 0.24$ will be necessary to rebuild the spawning stock to 30,100 mt by 2013 with 50% probability (Table B1.28, Figure B1.13).

POTENTIAL SENSITIVITY OF VPA ESTIMATES TO HYPOTHETICAL NEFSC SURVEY ADJUSTMENTS

Acting on the advice of industry members, NEFSC staff inspected the trawl cables (warp) on the NOAA Ship Albatross IV's sampling equipment on September 3, 2002. It was determined that the marks on the cable attaching scientific survey gear to the vessel were not at true 50 m length

intervals they are intended to indicate. The marks are used by the vessel crew to determine how much cable is deployed. The cable was most recently replaced in February 2000, and used in eight bottom trawl surveys, beginning with Winter 2000 and ending with Spring 2002.

Therefore, it is likely that at times more cable was deployed on one side of the NEFSC trawl survey net than on the other. This is a matter of inches at shorter lengths, and more pronounced as more cable is deployed. For example, with 100 m (328 ft) of cable deployed, just under 1 inch more cable was out on one side; at 300 m (984 ft) the difference was just under 6 ft. Of all tows made in the surveys, 75% deploy 300 m of cable or less. As a result, the NEFSC trawl survey gear may have fished differently during the Winter 2000 through Spring 2002 survey compared to prior surveys, and the data collected (catch per tow, for example) may have been influenced in a way that should be accounted for.

During September 24-27, 2002, video and net sensor equipment were used in experimental tows to both directly and numerically document net performance. Individuals from the region's commercial industry and the fishery management councils were part of the scientific crew during these observations. During October 2-3, 2002, a workshop was convened to examine the data collected and produce a report. The workshop was open to the public, with invited members to include scientists familiar with fishery survey practices, commercial fishermen and gear providers, the region's fishery management councils. As of this writing, the workshop was in progress, and was expected to produce a report detailing correction factors that can then be used in adjusting the NEFRSC survey indices used in this assessment, if needed.

In the interim, to examine the potential sensitivity of the SNE/MA winter flounder VPA to such corrections, hypothetical adjustments have been applied to the NEFSC winter, spring, and fall survey indices used in the SNE/MA winter flounder VPA. NEFSC indices from the Winter 2000 through Spring 2002 surveys were increased by 10%, 25%, and 100% to explore a range of the potential positive adjustments to the indices that might be necessary to account for reduced catch efficiency of the NEFSC survey gear during those surveys. The effect is nearly linear, with F in 2001 ranging from 0.51 for the baseline, W36_2 VPA to 0.36 for the VPA with all NEFSC survey indices increased by 100% (doubled); SSB in 2001 ranged from 7,600 mt for the baseline to 11,300 mt for NEFSC indices increased 100%. In all cases, the fishing mortality rate remained above F_{msy} , and SSB remained below one-half B_{msy} (Figure B1.14).

CONCLUSIONS

The Southern New England/Mid-Atlantic winter flounder stock complex is overfished and overfishing is occurring (Figure B1.15). Fully recruited fishing mortality in 2001 was 0.51 (exploitation rate = 37%), about 60% above $F_{msy} = 0.32$. The current VPA indicates there is an 80% chance that the 2001 F was between 0.44 and 0.58. Spawning stock biomass was estimated to be 7,600 mt in 2001, about 25% of $SSB_{msy} = 30,100$ mt. There is an 80% chance that the spawning stock biomass was between 6,800 mt and 8,400 mt in 2001.

The current assessment provides a much more pessimistic evaluation of stock status than the SARC 28 assessment in 1998 (NEFSC 1999). This is mainly due to the retrospective pattern of underestimating F and overestimating SSB in the current VPA. However, while the SNE/MA winter flounder VPA provides uncertain estimates of current F and SSB, it provides a better determination of stock status than reliance on survey indices alone. Managers should recognize that given the estimation uncertainty in the assessment, current fishing mortality rates are likely much higher than the 2001 estimate of 0.51, potentially by nearly 100%. Current SSB may in turn be substantially overestimated.

Spawning stock biomass declined substantially from 13,000-14,000 mt during the early 1980s to 2,700 mt during 1994-1996. SSB has increased since the mid 1990s to about 7,600 mt in 2001 due to reduced fishing mortality rates since 1997. Recruitment to the stock has been below average since 1989, and early indications are that the 2001 year class is the smallest in 22 years. Forecasts indicate that it will be necessary to reduce the fishing mortality rate to $F_{reb} = 0.24$ in 2003 and later years to rebuild to spawning stock to the target ($SSB_{msy} = 30,100$ mt) by 2013 with 50% probability.

The SARC elected to retain the NEFSC (2002) estimates of biological reference points for SNE/MA winter flounder for this assessment. The SARC recommends that these parametric stock-recruitment model reference points be adopted as the basis for the ASMFC and NEFMC FMP overfishing definitions. These reference points are a technical improvement over the ASMFC's yield per recruit reference points, as they include the estimates of B_{msy} , MSY , and F_{msy} required by the Sustainable Fisheries Act of 1996.

SARC COMMENTS

The SARC noted that while three of the major research recommendations from the SARC 28 assessment had been addressed, three more dealing mainly with the estimates of fishery discards remain unresolved, and should be addressed before the next assessment. The SARC discussed the use of surveys with different recent trends for tuning the VPA. A VPA run using only NEFSC surveys produced a more optimistic view of stock status than a run using only State indices. It was noted that the different trends among State surveys are likely tracking real trends in different portions of the stock complex. Therefore, combining the indices on a spatial scale or weighting them by survey area before tuning the VPA should be explored. The SARC reviewed a run using iterative re-weighting of the indices, which provided results very similar to the final, accepted VPA.

The SARC discussed the process of selecting indices used to tune the VPA, because the current VPA includes three new tuning series for ages 1-7+ (NEFSC winter, NJDFW river, and NJDFW ocean) and two more recruitment indices (NYDEC) not available for the SARC 28 assessment. The SARC reviewed a VPA run using the same suite of indices as in the last assessment (SARC 28) to determine how the addition of the new series had influenced the VPA results. That run

provided results similar to the final VPA. An examination of the utility of a randomization test on survey indices for determining the influence of the indices on VPA results could be informative.

The SARC noted that the current assessment provides a much more pessimistic evaluation of stock status than the SARC 28 assessment in 1998 (NEFSC 1999). This is mainly due to the retrospective pattern of underestimating F and overestimating SSB in the current VPA. It was noted that an increase in the catchability of the survey could produce the observed retrospective pattern. However, there was no reason to suspect an increase in the catchability of the NEFSC and State research surveys used in the VPA tuning. The mis-classification of landings by stock area could also be a cause of the retrospective pattern. However, the SARC noted a similar retrospective pattern in the Gulf of Maine winter flounder assessment, suggesting that significant SNE-MA landings had not been mis-classified into the Gulf of Maine stock area. An underestimation of the discarded proportion of the catch could also produce the observed retrospective pattern. The use of VTR data in estimating commercial fishery discards is a source of uncertainty. Possible significant discarding in the commercial scallop dredge fishery was noted, but current data provide generally small (less than the trawl fishery) and extremely variable estimates of winter flounder discards in the dredge fishery. Finally, the observed retrospective pattern might be caused by under-reporting or underestimation of the commercial or recreational landings. Given the retrospective pattern, the utility of the current SNE-MA winter flounder VPA was evaluated. The SARC concluded that, while the SNE-MA winter flounder VPA provides uncertain estimates of current F and SSB, it still provides the best available determination of stock status.

As one illustration of the possible magnitude of potential missing catch, the SARC noted that it would take roughly a trebling of the catch during the period 1996-1998 to significantly reduce the magnitude of the retrospective pattern in fishing mortality. The SARC noted that retrospective patterns are evident in several of the New England groundfish stock assessments (e.g. GOM winter flounder, SNE-MA, CC/GOM, and GB yellowtail flounder, GB and GOM cod, and witch flounder). Investigation to determine a common cause for this pattern should be pursued. Alternative assessment methods for dealing with retrospective patterns, such as statistical catch at age models, should be explored.

SOURCES OF UNCERTAINTY

- 1) Stock-specific landings data for 1994 and later are derived by proration from Vessel Trip Report data and are considered provisional.
- 2) Length frequency sampling intensity of the commercial and recreational fishery landings has been low in some recent years, and likely increases the uncertainty of the estimated landings at age.

- 3) Commercial fishery discard estimates are based on rates provided by fishers in the Vessel Trip Reports, owing to inadequate Fishery Observer sampling.
- 4) The SNE-MA winter flounder VPA exhibits a retrospective pattern of underestimating F and overestimating SSB during the late 1990s, increasing the uncertainty of current estimates of F and SSB.

RESEARCH RECOMMENDATIONS

New

- 1) Evaluate the maturity at age of fish sampled in the NEFSC fall and winter surveys.
- 2) Consider fieldwork to record ovary weights along with maturity stage data from 20-30 cm fish in the NEFSC and State agency surveys for 1-2 years to help resolve age/size at maturity differences between State and NEFSC surveys.
- 3) Conduct periodic maturity staging workshops involving State and NEFSC trawl survey staff.
- 4) Examine sources of the differences between NEFSC, MA and CT survey maturity (validity of evidence for smaller size or younger age at 50% maturity in the NEFSC data). Compare NEFSC inshore against offshore strata for differences in maturity. Compare confidence intervals for maturity ogives. Calculate annual ogives and investigate for progression of maturity changes over time. Examine maturity data from NEFSC strata on Nantucket Shoals and near George's Bank separately from more inshore areas. Consider methods for combining maturity data from different survey programs.
- 5) Increase the intensity of commercial fishery discard length sampling.
- 6) Consider post-stratification of NEFSC survey offshore stratum 23, to facilitate inclusion of survey catches from this stratum (east of Cape Cod) in the SNE-MA winter flounder assessment.
- 7) Incorporate State samples (e.g. NY DEC Party Boat Survey and CT DEP Volunteer Angler Survey) in the estimation of recreational fishery landings and discards, if possible.
- 8) Attempt use of a forward projection (statistical catch at age model) in the next assessment.

Old: Pending

- 1) Continue to consider the effects of catch-and-release components of recreational fishery on discard at age (i.e., develop mortality estimates from the American Littoral Society tagging database, if feasible).

- 2) Compare commercial fishery discard estimates from the Mayo survey/mesh algorithm with those from VTR data for comparable time periods.
- 3) Maintain or increase sampling levels (currently supported by individual state funding) and collect age information from MRFSS samples.
- 4) Examine the implications of anthropogenic mortalities caused by pollution and power plant entrainment in estimating yield per recruit, if feasible.
- 5) Examine the implications of stock mixing from data from Great South Channel region.
- 6) Expand sea sampling for estimation of commercial discards.
- 7) Revise the recreational fishery discard estimates by applying a consistent method across all years, if feasible (i.e., the Gibson 1996 method).

Old: Work In Progress

- 1) Re-examine the maturity ogive to incorporate any recent research results.
- 2) Explore the feasibility of stratification of the commercial fishery discard estimation by fishery (e.g., mesh, gear, area).

Old: Completed

- 1) Further examine the comparability of age-length keys from different areas within the stock. Current comparisons are based on two years and three ages. Conduct an age structure exchange between NEFSC, CT DEP, and MADMF, to ensure consistency in ageing protocol.
- 2) Age the archived MA DMF survey age samples for 1978-1989.
- 3) Compile NEFSC Winter Survey abundance indices for winter flounder and evaluate their utility.
- 4) Evaluate the utility of MADMF sea sample data for winter flounder in estimating commercial fishery discards.

LITERATURE CITED

- ASMFC. 1998. Assessment of the Southern New England/Mid-Atlantic and Gulf of Maine Winter Flounder stocks: a report by the ASMFC's Winter Flounder Technical Committee. ASMFC WFTC Document 98-01. 31 p + app.
- Azarovitz, T., J. McGurrin and R. Seagraves, eds. 1989. Proceedings of a workshop on bottom trawl surveys. Atlantic States Marine Fisheries Commission. Special Report No. 17. August 1989.
- Beverton, R.J.H., and S.J. Holt. 1957. On the dynamics of exploited fish populations. Chapman and Hall, London. Facsimile reprint 1993.
- Bigelow, H. and W. Schroeder. 1953. Fishes of the Gulf of Maine. Fishery Bulletin of the Fish and Wildlife Service. V. 53. Fishery Bulletin 74.
- Brodziak, J.T.K., W.J. Overholtz, and P. Rago. 2001. Does spawning stock affect recruitment of New England groundfish? *Can. J. Fish. Aquat. Sci.* 58(2): 306-318.
- Brownie, C., D. Anderson, K. Burnham and D. Robson. 1985. Statistical inference from band recovery data: a handbook. U.S. Fish and Wildlife Service, Res. Publ. No. 156. 305 pp.
- Conser, R. and J. Powers. 1990. Extension of the ADAPT VPA tuning method designed to facilitate assessment work on tuna and swordfish stocks. *Int. Comm. Conserv. Atlantic Tunas. Coll. Vol. Sci. Pap.* 32: 461-467.
- Efron, B. 1982. The jackknife, the bootstrap and other resampling plans. *Phila. Soc. for Ind. and Appl. Math.* 38.
- Gibson, M. 1996. Age structure of winter flounder B2 type recreational discards estimated from inshore trawl survey lengths, age-length keys, and minimum size regulations. NEFSC Res. Doc. 96-05b.
- Howe, A., and P. Coates. 1975. Winter flounder movements, growth, and mortality off Massachusetts. *Trans. Am. Fish. Soc.* 104: 13-29.
- Howell, P. 1996. Identification of stock units. NEFSC Res. Doc. 96-05b.
- Howell, P., A. Howe, M. Gibson and S. Ayvasian. 1992. Fishery management plan for inshore stocks of winter flounder. Atlantic States Marine Fisheries Commission. Fisheries Management Report No. 21. May, 1992.

- Mace, P.M., and I.J. Doonan. 1988. A generalized bioeconomic simulation model for fish population dynamics. N.Z. Fish. Ass. Res. Doc. 88/4.
- Mayo, R. K., L. O'Brien, and N. Buxton. 1992. Discard estimates of American plaice, *Hippoglossoides platessoides*, in the Gulf of Maine northern shrimp fishery and the Gulf of Maine-Georges Bank large-mesh otter trawl fishery. SAW 14 Res. Doc. 14/3, 40 pp.
- NEFSC. 1996. Report of the 21st Northeast Regional Stock Assessment Workshop (21st SAW): Stock Assessment Review Committee (SARC) consensus summary of assessments. Northeast Fish. Sci. Cent. Ref. Doc. 96-05d. 200 p.
- NEFSC. 1999. Report of the 28th Northeast Regional Stock Assessment Workshop (28th SAW): Stock Assessment Review Committee (SARC) consensus summary of assessments. Northeast Fish. Sci. Cent. Ref. Doc. 99-08. 304 p.
- NEFSC. 2001. Assessment of 19 Northeast Groundfish Stocks through 2000: a report to the New England Fishery Management Council's Multi-Species Monitoring Committee. Northeast Fish. Sci. Cent. Ref. Doc. 01-20. 217 p.
- NEFSC. 2002. Final report of the Working Group on re-evaluation of biological reference points for New England groundfish. Northeast Fish. Sci. Cent. Ref. Doc. 02-04. 123 p.
- O'Brien, L., J. Burnett, and R. Mayo. 1993. Maturation of nineteen species of finfish off the northeast coast of the United States, 1985-1990. NOAA Tech. Rep. NMFS 113. 66 pp.
- Rago, P., W. Gabriel, and M. Lambert. 1994. Assessment of yellowtail flounder *Pleuronectes ferrugineus*. NEFSC Ref. Doc. 94-02.
- Simpson, D.G. 1989. Codend selection of winter flounder *Pseudopleuronectes americanus*. NOAA Tech. Rpt. NMFS 75. 10 p.
- Thompson, W. F. and F. H. Bell. 1934. Biological statistics of the Pacific halibut fishery. 2. Effect of changes in intensity upon total yield and yield per recruit of gear. Rep. Int. Fish. (Pacific halibut) Comm. 8: 49 p.

Table B1.1. Winter flounder commercial landings (metric tons) for Southern New England/Mid-Atlantic stock complex area (U.S. statistical reporting areas 521, 526, divisions 53, 61-63) as reported by NEFSC weighout, state bulletin and general canvas data.

Year	Metric Tons
1964	7,474
1965	8,678
1966	11,977
1967	9,478
1968	7,070
1969	8,107
1970	8,603
1971	7,367
1972	5,190
1973	5,573
1974	4,259
1975	3,982
1976	3,265
1977	4,413
1978	6,327
1979	6,543
1980	10,627
1981	11,176
1982	9,438
1983	8,659
1984	8,882
1985	7,052
1986	4,929
1987	5,172
1988	4,312
1989	3,670
1990	4,232
1991	4,823
1992	3,816
1993	3,010
1994	2,159
1995	2,634
1996	2,781
1997	3,441
1998	3,208
1999	3,444
2000	3,783
2001	4,448

Table B1.2. Distribution of commercial landings (percentage of annual total) of winter flounder from Southern New England/Mid-Atlantic stock complex area by U.S. statistical reporting area.

Year	Area								
	521	526	537	538	539	611	612	613	614-622
1989	33.2	10.8	18.9	7.0	12.1	7.1	5.5	4.2	1.2
1990	45.2	16.8	6.1	4.9	9.5	11.1	4.1	2.0	0.1
1991	46.4	14.7	10.8	1.7	13.7	5.7	3.6	2.9	0.4
1992	37.0	12.5	17.4	2.4	9.4	10.1	4.5	3.4	3.4
1993	46.6	10.0	10.8	2.4	8.2	7.7	4.2	8.0	2.1
1994	41.8	13.3	3.3	0.1	17.6	10.3	6.5	3.1	3.3
1995	43.3	9.1	6.7	1.6	15.7	10.8	9.3	2.1	1.4
1996	47.3	12.0	10.8	1.4	12.3	11.0	2.5	2.4	0.3
1997	62.8	3.1	7.5	1.5	12.3	8.5	2.0	2.1	0.2
1998	49.5	12.4	7.6	0.6	15.2	9.9	1.8	2.4	0.6
1999	48.7	12.3	6.9	0.4	13.2	8.2	6.4	2.4	1.5
2000	44.1	7.4	10.7	0.8	15.1	8.5	7.2	4.8	1.4
2001	55.8	7.2	7.4	0.1	9.7	7.7	7.4	3.1	1.6

Table B1.3. Estimated number (N, 000's) and weight (mt) of winter flounder caught, landed, and discarded in the recreational fishery, Southern New England/Mid-Atlantic stock complex.

Year	Catch N (A+B1+B2)	Landed N (A+B1)	Released N (B2)	15% Release Mortality	Landings (A+B1; mt)
1981	11,006	8,089	2,916	437	3,050
1982	10,665	8,392	2,273	341	2,457
1983	11,010	8,365	2,645	397	2,524
1984	17,723	12,756	4,967	745	5,772
1985	18,056	13,297	4,759	714	5,198
1986	9,368	6,995	2,374	356	2,940
1987	9,213	6,900	2,313	347	3,141
1988	10,134	7,358	2,775	416	3,423
1989	5,919	3,682	2,236	335	1,802
1990	3,827	2,486	1,340	201	1,063
1991	4,325	2,795	1,530	230	1,214
1992	1,360	806	555	83	393
1993	2,211	1,180	1,031	155	543
1994	1,829	1,209	620	93	598
1995	1,850	1,390	461	69	661
1996	2,679	1,554	1,125	169	689
1997	1,901	1,207	694	104	621
1998	1,008	584	425	64	290
1999	1,071	658	412	62	320
2000	2,043	1,346	697	105	831
2001	1,441	901	540	81	552

Table B1.4. Winter flounder commercial fishery landed sample lengths (number of fish measured) used for Southern New England/Mid-Atlantic stock complex, 1981-1997. Landings are in metric tons.

Year	Landings	Lengths measured	Metric tons per 100 lengths
1981	11,176	4,230	264
1982	9,438	5,796	163
1983	8,659	5,601	155
1984	8,882	3,697	240
1985	7,052	6,407	110
1986	4,929	5,120	96
1987	5,172	5,271	98
1988	4,312	4,208	102
1989	3,670	3,525	104
1990	4,232	4,088	104
1991	4,823	3,058	158
1992	3,816	4,163	92
1993	3,010	2,354	128
1994	2,159	2,593	83
1995	2,634	4,153	63
1996	2,781	2,019	138
1997	3,441	4,005	86

Table B1.5. Winter flounder commercial fishery landed sample lengths (number of fish measured) used for Southern New England/Mid-Atlantic stock complex, 1998-2001. Landings are in metric tons.

1998		Market Category				
Sample Type	Season	Unclass.	Small	Medium	Large	Total
Port	Jan-Jun	162	105	767	205	1239
Port	Jul-Dec	780	794	558	210	2342
Total lengths used		942	899	1325	415	3581
Landings		644	1453	438	673	3208
Metric tons per 100 lengths		68	162	33	162	90
1999		Market Category				
Sample Type	Season	Unclass.	Small	Medium	Large	Total
Port	Jan-Jun	978	334	502	522	2336
Port	Jul-Dec	1403	464	105	299	2271
Total lengths used		2381	798	607	821	4607
Landings		838	1566	290	750	3444
Metric tons per 100 lengths		35	196	48	91	75
2000		Market Category				
Sample Type	Season	Unclass.	Small	Medium	Large	Total
Port	Jan-Jun	808	377	1868	126	3179
Port	Jul-Dec	845	565	1025	839	3274
Total lengths used		1653	942	2893	965	6453
Landings		848	451	1670	815	3784
Metric tons per 100 lengths		51	48	58	84	59

Table B1.5 continued.

2001		Market Category				
Sample Type	Season	Unclass.	Small	Medium	Large	Total
Port	Jan-Jun	557	510	1067	636	2770
Port	Jul-Dec	203	387	1234	1661	3485
Total lengths used		760	897	2301	2297	6255
Landings		908	1101	1475	962	4446
Metric tons per 100 lengths		119	123	64	42	71

Table B1.6. Winter flounder recreational fishery landed sample lengths (number of fish measured) used for Southern New England/Mid-Atlantic stock complex, 1981-1997. Landings are in metric tons.

Year	Landings	Lengths measured	Metric tons per 100 lengths
1981	3,050	1,725	177
1982	2,457	1,971	125
1983	2,524	2,587	98
1984	5,772	3,123	185
1985	5,198	2,357	221
1986	2,940	2,237	131
1987	3,141	1,360	231
1988	3,423	1,944	176
1989	1,802	2,810	64
1990	1,063	2,548	42
1991	1,214	1,755	69
1992	393	1,083	36
1993	543	1,288	42
1994	598	948	63
1995	661	767	86
1996	689	936	74
1997	621	752	83

Table B1.7. Winter flounder recreational fishery sample lengths (number of fish measured) used for Southern New England/Mid-Atlantic stock complex, 1998-2001. SNE = MA & RI; MA = CT and states south. Landings are in metric tons.

Season/area	1998	1999	2000	2001
Jan-Jun/SNE	105	77	7	80
Jan-Jun/MA	405	256	105	387
Jul-Dec/SNE	85	48	0	3
Jul-Dec/MA	21	14	48	38
Total lengths	616	395	160	508
Landings (A+B1.)	290	320	831	552
Metric tons per 100 Lengths	47	81	519	109

Table B1.8. Winter flounder NEFSC Domestic Fishery Observer Program (OB) and NER Vessel Trip Report (VTR) data: number of OB trips with landed winter flounder (to estimate discards to landings ratio), OB discards to landings ratio, number of VTR trips with winter flounder landings that discarded any species, and VTR discards to landings ratio. VTR data available for 1994 and subsequent years.

Year	Half-year	OB trips	OB ratio	VTR Trips	VTR ratio
1989	Jan-Jun	22	0.235		
	Jul-Dec	28	0.299		
1990	Jan-Jun	21	0.069		
	Jul-Dec	18	0.227		
1991	Jan-Jun	46	0.579		
	Jul-Dec	42	0.283		
1992	Jan-Jun	17	0.021		
	Jul-Dec	21	0.076		
1993	Jan-Jun	11	0.299		
	Jul-Dec	22	0.32		
1994	Jan-Jun	13	0.304	1519	0.241
	Jul-Dec	12	2.84	1488	0.091
1995	Jan-Jun	20	0.044	1484	0.072
	Jul-Dec	36	0.289	764	0.028
1996	Jan-Jun	18	0.358	1002	0.088
	Jul-Dec	38	0.115	576	0.05
1997	Jan-Jun	27	0.175	2138	0.145
	Jul-Dec	18	0.021	1766	0.16

Table B1.8 continued.

Year	Half-year	OB trips	OB ratio	VTR Trips	VTR ratio
1998	Jan-Jun	6	0.306	2114	0.265
	Jul-Dec	18	0.437	1424	0.292
1999	Jan-Jun	13	11.842	2570	0.102
	Jul-Dec	7	0.005	1554	0.238
2000	Jan-Jun	20	0.095	2104	0.16
	Jul-Dec	21	0.042	1586	0.043
2001	Jan-Jun	27	0.04	2508	0.061
	Jul-Dec	22	0.069	2016	0.025

Table B1.9. Winter flounder commercial fishery discard sample lengths (number of fish measured) used for Southern New England/Mid-Atlantic stock complex, 1994-2001. Discard estimates (before impact of 50% mortality rate) are in metric tons.

Season	1994	1995	1996	1997
Jan-Jun	111	73	358	412
Jul-Dec	196	646	245	556
Total lengths	307	719	603	968
Discard Estimate (before mortality)	608	242	346	534
Metric tons per 100 Lengths	198	34	57	55
Season	1998	1999	2000	2001
Jan-Jun	170	354	353	135
Jul-Dec	604	13	128	0
Total lengths	774	367	481	135
Discard Estimate (before mortality)	911	659	296	167
Metric tons per 100 Lengths	118	180	62	124

Table B1.10. Winter flounder catch at age (number in 000s) for the Southern New England/Mid-Atlantic stock complex.

Commercial Landings Year	Age												
	1	2	3	4	5	6	7	8	9	10	11	12	13
1981	194	7154	9740	2750	606	178	42	32	0	0	9	0	0
1982	54	6897	8496	2715	488	187	78	59	21	17	7	7	0
1983	6	2795	7114	3957	1322	584	269	91	34	70	6	29	35
1984	0	4518	6367	3197	1503	768	355	158	67	86	27	33	37
1985	27	3936	5688	3052	1014	326	104	32	17	7	5	2	0
1986	0	2122	4187	2206	551	271	84	27	6	3	1	2	0
1987	0	2488	5465	1895	465	122	40	20	14	12	2	0	0
1988	0	2241	3929	1607	412	122	37	24	3	2	1	0	0
1989	0	1542	4057	1747	431	58	34	13	5	1	0	0	0
1990	0	1003	3977	1757	315	95	37	16	0	3	0	0	0
1991	0	1406	4756	2239	447	143	48	16	5	1	1	0	0
1992	0	484	3416	2127	574	111	32	11	3	0	0	0	0
1993	13	885	2516	1377	361	102	71	7	0	0	2	0	1
1994	0	629	804	401	90	14	10	0	0	0	0	0	0
1995	0	73	1537	587	95	24	5	0	0	0	0	0	0
1996	0	606	1146	470	122	17	11	0	0	0	0	0	0
1997	0	1418	2574	1370	356	70	28	12	5	1	0	0	0
1998	0	1021	3057	1483	450	83	60	63	7	0	0	0	0
1999	0	2009	3347	1538	386	59	11	6	0	0	0	0	0
2000	0	1073	2801	1942	592	135	35	12	0	0	0	0	0
2001	0	1727	3263	1851	620	148	53	23	2	3	0	0	0

Table B1.10 continued

Commercial Discards Year	Age												
	1	2	3	4	5	6	7	8	9	10	11	12	13
1981	322	2514	2186	101	0	0	0	0	0	0	0	0	0
1982	43	2817	1219	192	0	0	0	0	0	0	0	0	0
1983	260	2479	2000	467	45	0	0	0	0	0	0	0	0
1984	159	2102	1502	166	6	1	0	0	0	0	0	0	0
1985	22	1504	2516	442	43	4	0	0	0	0	0	0	0
1986	78	2220	2389	205	10	0	0	0	0	0	0	0	0
1987	11	1600	1755	170	9	0	0	0	0	0	0	0	0
1988	6	887	2540	276	20	0	0	0	0	0	0	0	0
1989	315	2724	2131	555	33	2	1	0	0	0	0	0	0
1990	16	781	1433	322	14	0	1	0	0	0	0	0	0
1991	17	1238	1205	227	12	1	0	0	0	0	0	0	0
1992	15	845	787	150	14	1	0	0	0	0	0	0	0
1993	201	849	467	57	6	0	0	0	0	0	0	0	0
1994	44	204	88	8	0	0	0	0	0	0	0	0	0
1995	15	47	41	4	0	0	0	0	0	0	0	0	0
1996	11	64	66	7	1	0	0	0	0	0	0	0	0
1997	373	580	210	31	6	0	0	0	0	0	0	0	0
1998	43	972	407	78	3	0	0	0	0	0	0	0	0
1999	63	583	314	54	23	22	15	0	0	0	0	0	0
2000	68	218	199	34	8	1	6	0	0	0	0	0	0
2001	11	127	111	33	3	0	0	0	0	0	0	0	0

Table B1.10 continued.

Recreational Landings Year	Age												
	1	2	3	4	5	6	7	8	9	10	11	12	13
1981	776	4054	2426	742	59	4	28	0	0	0	0	0	0
1982	457	4235	2716	823	122	26	13	0	0	0	0	0	0
1983	289	1630	4194	1702	427	112	11	0	0	0	0	0	0
1984	294	4258	6224	1565	267	107	41	0	0	0	0	0	0
1985	219	1585	4270	2558	1895	1513	878	0	335	44	0	0	0
1986	106	1765	2432	1797	491	171	81	77	51	8	17	0	0
1987	16	926	1736	1023	2229	633	82	115	64	77	0	0	0
1988	21	534	2858	2078	775	857	128	51	37	20	0	0	0
1989	99	739	944	1200	385	161	91	36	16	8	3	1	0
1990	7	189	814	851	439	101	52	20	3	3	0	2	5
1991	13	232	1122	879	399	107	38	0	1	0	3	0	0
1992	3	123	235	303	85	50	7	0	0	0	0	0	0
1993	31	233	321	289	218	54	20	10	4	2	0	0	0
1994	5	203	240	303	220	149	89	0	0	0	0	0	0
1995	0	30	268	298	321	267	206	0	0	0	0	0	0
1996	0	106	200	630	220	240	157	0	0	0	0	0	0
1997	1	82	497	410	178	36	0	0	0	0	0	0	0
1998	2	89	191	235	58	7	1	0	0	0	0	0	0
1999	1	101	340	151	49	16	0	0	0	0	0	0	0
2000	0	113	440	472	262	44	14	0	0	0	0	0	0
2001	1	84	267	303	168	62	16	0	0	0	0	0	0

Table B1.10 continued.

Recreational Discards Year	Age												
	1	2	3	4	5	6	7	8	9	10	11	12	13
1981	70	367	0	0	0	0	0	0	0	0	0	0	0
1982	33	308	0	0	0	0	0	0	0	0	0	0	0
1983	62	337	0	0	0	0	0	0	0	0	0	0	0
1984	48	697	0	0	0	0	0	0	0	0	0	0	0
1985	9	340	363	2	0	0	0	0	0	0	0	0	0
1986	32	222	93	9	0	0	0	0	0	0	0	0	0
1987	47	254	43	3	1	0	0	0	0	0	0	0	0
1988	57	279	76	3	0	0	0	0	0	0	0	0	0
1989	49	240	45	1	0	0	0	0	0	0	0	0	0
1990	12	136	51	2	0	0	0	0	0	0	0	0	0
1991	22	151	56	0	0	0	0	0	0	0	0	0	0
1992	7	51	19	1	0	0	0	0	0	0	0	0	0
1993	29	95	26	4	0	0	0	0	0	0	0	0	0
1994	12	60	21	0	0	0	0	0	0	0	0	0	0
1995	9	45	15	0	0	0	0	0	0	0	0	0	0
1996	21	110	37	0	0	0	0	0	0	0	0	0	0
1997	11	55	19	0	0	0	0	0	0	0	0	0	0
1998	5	49	8	1	0	0	0	0	0	0	0	0	0
1999	2	53	6	1	0	0	0	0	0	0	0	0	0
2000	0	38	60	7	0	0	0	0	0	0	0	0	0
2001	1	49	27	5	0	0	0	0	0	0	0	0	0

Table B1.10 continued.

Total Catch Year	Age													Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	
1981	1362	14089	14352	3593	665	182	70	32	0	0	9	0	0	34354
1982	587	14257	12421	3730	610	213	91	59	21	17	7	7	0	32020
1983	617	7241	13308	6126	1794	696	280	91	34	70	6	29	35	30327
1984	501	11575	14093	4928	1776	876	396	158	67	86	27	33	37	34553
1985	277	7366	12836	6054	2953	1843	982	32	352	52	5	2	0	32753
1986	215	6327	9102	4216	1053	442	165	104	57	10	19	2	0	21712
1987	73	5268	8999	3091	2703	755	122	135	78	89	2	0	0	21315
1988	84	3941	9402	3964	1207	979	165	75	39	22	1	0	0	19880
1989	463	5246	7176	3503	849	222	126	49	21	9	3	1	0	17668
1990	36	2109	6275	2931	767	196	89	36	4	5	0	2	5	12455
1991	53	3027	7140	3344	858	251	87	16	6	1	4	0	0	14788
1992	25	1503	4457	2581	674	162	38	11	3	0	0	0	0	9455
1993	274	2062	3329	1728	585	157	91	17	4	2	2	0	1	8251
1994	61	1097	1152	713	311	162	99	0	0	0	0	0	0	3595
1995	24	195	1862	889	415	291	211	0	0	0	0	0	0	3887
1996	32	886	1450	1107	343	258	168	0	0	0	0	0	0	4244
1997	385	2135	3300	1811	540	106	28	12	5	1	0	0	0	8323
1998	50	2132	3663	1797	511	90	61	63	7	0	0	0	0	8374
1999	66	2746	4008	1744	458	97	26	6	0	0	0	0	0	9150
2000	69	1442	3500	2455	862	180	55	12	0	0	0	0	0	8575
2001	13	1987	3668	2191	790	211	69	23	2	3	0	0	0	8957

Table B1.11. Total winter flounder recreational and commercial catch for the Southern New England/Mid-Atlantic stock complex in weight (mt) and numbers (000s).

Year	Commercial Landings		Commercial Discards		Recreational Landings		Recreational Discards		Total Catch		% Discards/Total	
	mt	000s	mt	000s	mt	000s	mt	000s	mt	000s	mt	000s
1981	11,176	20,705	1,343	5,123	3,050	8,089	88	437	15,657	34,354	9.1	16.2
1982	9,438	19,016	1,149	4,271	2,457	8,392	66	341	13,110	32,020	9.3	14.4
1983	8,659	16,312	1,311	5,251	2,524	8,365	125	399	12,619	30,327	11.4	18.6
1984	8,882	17,116	986	3,936	5,772	12,756	148	745	15,788	34,553	7.2	13.5
1985	7,052	14,211	1,534	4,531	5,198	13,297	230	714	14,014	32,753	12.6	16.0
1986	4,929	9,460	1,273	4,902	2,940	6,994	66	356	9,208	21,712	14.5	24.2
1987	5,172	10,524	950	3,545	3,141	6,899	61	347	9,324	21,315	10.8	18.3
1988	4,312	8,377	904	3,728	3,423	7,359	69	416	8,708	19,880	11.2	20.8
1989	3,670	7,888	1,404	5,761	1,802	3,684	49	335	6,925	17,668	21.0	34.5
1990	4,232	7,202	673	2,567	1,063	2,485	31	201	5,999	12,455	11.7	22.2
1991	4,823	9,063	784	2,701	1,214	2,794	51	230	6,872	14,788	12.2	19.8
1992	3,816	6,759	511	1,811	393	802	15	83	4,735	9,455	11.1	20.0
1993	3,010	5,336	457	1,580	543	1,180	31	155	4,041	8,251	12.1	21.0
1994	2,159	1,948	304	344	598	1,210	34	93	3,095	3,595	10.9	12.2
1995	2,634	2,321	121	107	661	1,390	23	69	3,439	3,887	4.2	4.5
1996	2,781	2,372	173	149	689	1,555	64	168	3,707	4,244	6.4	7.5
1997	3,441	5,834	267	1,200	618	1,204	26	85	4,352	8,323	6.7	15.4

Table B1.11 continued.

Year	Commercial Landings		Commercial Discards		Recreational Landings		Recreational Discards		Total Catch		% Discards/Total	
	mt	000s	mt	000s	mt	000s	mt	000s	mt	000s	mt	000s
1998	3,208	6,224	456	1,503	290	584	13	64	3,967	8,375	11.8	18.7
1999	3,444	7,356	329	1,074	320	658	14	62	4,107	9,150	8.4	12.4
2000	3,783	6,590	148	534	831	1,346	30	105	4,792	8,575	3.7	7.5
2001	4,448	7,690	83	285	552	901	19	81	5,102	8,957	2.0	4.1

Table B1.12. Total fishery catch at age used as input to Virtual Population Analysis (VPA) for the Southern New England/Mid-Atlantic winter flounder stock complex.

Year	Age						
	1	2	3	4	5	6	7+
1981	1362	14089	14352	3593	665	182	111
1982	587	14257	12421	3730	610	213	202
1983	617	7241	13308	6126	1794	696	545
1984	501	11575	14093	4928	1776	876	804
1985	277	7366	12836	6054	2953	1843	1424
1986	215	6327	9102	4216	1053	442	357
1987	73	5268	8999	3091	2703	755	426
1988	84	3941	9402	3964	1207	979	303
1989	463	5246	7176	3503	849	222	209
1990	36	2109	6275	2931	767	196	141
1991	53	3027	7140	3344	858	251	115
1992	25	1503	4457	2581	674	162	53
1993	274	2062	3329	1728	585	157	116
1994	61	1097	1152	713	311	162	99
1995	24	195	1862	889	415	291	211
1996	32	886	1450	1107	343	258	168
1997	385	2135	3300	1811	540	106	46
1998	50	2132	3663	1797	511	90	131
1999	66	2746	4008	1744	458	97	32
2000	69	1442	3500	2455	862	180	67
2001	13	1987	3668	2191	790	211	97

Table B1.13. Total fishery mean weights at age used as input to Virtual Population Analysis (VPA) for the Southern New England/Mid-Atlantic winter flounder stock complex.

Year	Age						
	1	2	3	4	5	6	7+
1981	0.130	0.276	0.478	0.802	1.065	1.243	1.202
1982	0.090	0.261	0.438	0.694	1.048	1.253	1.837
1983	0.195	0.237	0.353	0.516	0.774	1.046	1.552
1984	0.146	0.258	0.366	0.542	0.693	0.913	1.282
1985	0.111	0.282	0.364	0.482	0.522	0.467	0.613
1986	0.129	0.292	0.398	0.480	0.685	0.879	0.961
1987	0.046	0.287	0.384	0.551	0.475	0.564	0.853
1988	0.039	0.279	0.351	0.508	0.634	0.517	0.827
1989	0.118	0.258	0.378	0.508	0.660	0.716	1.073
1990	0.082	0.295	0.394	0.525	0.672	0.808	0.990
1991	0.093	0.317	0.420	0.534	0.603	0.823	1.168
1992	0.079	0.287	0.427	0.599	0.802	0.945	1.395
1993	0.169	0.334	0.460	0.592	0.689	0.878	1.167
1994	0.156	0.347	0.448	0.597	0.741	0.692	0.818
1995	0.167	0.323	0.449	0.578	0.714	0.763	0.780
1996	0.193	0.407	0.507	0.569	0.705	0.826	0.853
1997	0.093	0.369	0.510	0.659	0.806	1.071	1.511
1998	0.202	0.332	0.438	0.580	0.665	0.892	1.241
1999	0.079	0.314	0.435	0.562	0.782	0.951	1.317
2000	0.100	0.396	0.484	0.613	0.738	0.915	1.144
2001	0.102	0.419	0.506	0.636	0.796	1.053	1.259

Table B1.14. Winter flounder NEFSC survey index stratified mean number and mean weight (kg) per tow for the Southern New England- Mid-Atlantic stock complex. Spring and fall strata set (offshore 1-12, 25, 69-76 ; inshore 1-29, 45-56); winter strata set (offshore 1-2, 5-6,9-10,69,73).

Year	Spring				Fall			
	Number	N(CV)	Weight	W(CV)	Number	N(CV)	Weight	W(CV)
1963					8.554	33.2	3.284	41.4
1964					13.673	22.1	4.894	19.4
1965					15.537	32.5	4.435	28.7
1966					9.843	31.5	3.275	27.3
1967					9.109	20.6	2.745	18.7
1968	2.444	26.7	0.734	37.2	8.105	21	2.19	18.7
1969	5.64	34.3	3.414	53.7	6.841	34.9	1.939	29.7
1970	2.729	30.9	1.326	35.6	5.11	36.1	2.375	47.8
1971	2.035	32.9	0.756	36.2	3.861	17.5	1.231	19.1
1972	1.865	28.1	0.656	32.1	7.687	39.4	3.053	44.6
1973	7.458	19.9	2.013	20.6	2.691	26.9	0.775	25.8
1974	3.362	21.9	1.043	19.3	2.032	31.1	0.822	29.4
1975	1.135	22.6	0.354	20.8	2.196	20.3	0.688	22.1
1976	3.085	16.3	0.804	17.2	2.376	32.2	1.251	42.9
1977	4.209	17.2	1.189	18.6	4.722	22.5	1.735	25.2
1978	6.695	11.1	1.758	13.3	3.743	17.6	1.43	22.6
1979	2.966	16.8	1.069	25	10.058	18.4	2.606	15.4
1980	15.25	17.5	3.551	13.6	9.964	31	3.216	29.5
1981	18.234	20.9	4.762	16.9	10.206	20.3	3.11	19.9
1982	6.986	20.1	1.918	15.8	4.927	22.8	1.683	25.9
1983	6.262	18.4	2.469	28	8.757	37.6	2.69	31.7
1984	5.524	19	2.072	28.4	2.681	21.1	0.887	21
1985	5.36	17.4	1.983	16.5	2.727	21.5	0.991	21.5
1986	2.266	23.9	0.766	23.4	1.538	21.9	0.487	19.1
1987	1.763	21.3	0.568	17.9	1.167	28.9	0.419	37.8
1988	2.126	19.6	0.73	19.3	1.246	22.4	0.53	27.5
1989	2.485	33.5	0.582	29.6	1.435	40.7	0.341	30.4
1990	1.992	36.8	0.472	33.1	1.979	29.6	0.546	25.8
1991	2.473	15.6	0.692	14.7	1.95	23.6	0.708	25.6

Table B1.14 continued.

Year	Number	Spring			Number	Fall			Number	Winter		
		N(CV)	Weight	W(CV)		N(CV)	Weight	W(CV)		N(CV)	Weight	W(CV)
1992	1.579	23.4	0.435	22.1	2.963	32.4	0.829	31.8	3.68	27.3	0.928	26
1993	0.961	19.1	0.219	14.8	1.382	25	0.392	25.9	2.59	29.4	0.456	21.5
1994	1.51	26.4	0.329	21.9	4.134	24.8	1.482	27.3	3.797	30.8	1.183	35.5
1995	2.097	23.4	0.592	19.1	2.253	20.7	0.626	17.3	2.221	26.1	0.697	29.1
1996	1.517	14.3	0.428	15.2	3.186	39.8	1.063	45.3	3.778	28.4	0.734	25.2
1997	1.436	22.1	0.399	20	7.893	32.6	2.583	26.7	3.906	19.7	1.043	21.6
1998	2.774	20.6	0.845	22.1	6.597	13.6	2.232	9.9	7.169	21.6	1.83	24.1
1999	4.171	16.2	1.245	16.4	3.596	17	1.549	16.5	10.328	31.8	3.1	32.3
2000	3.172	26.6	1.123	31.9	6.168	25.5	2.143	26.2	5.571	32.9	1.525	29.5
2001	1.568	14.3	0.581	13.3	4.877	28.1	2.03	28.5	3.096	31.6	0.873	29
2002	2.043	15.7	0.782	16.3					2.901	27.7	1.188	38.3

NOTE: 1968-1972 spring index does not include inshore strata ; 1963-1971 fall index does not include inshore strata. All indices calculated with trawl door conversion factors where appropriate. Winter trawl survey began in 1992.

Table B1.15. SNE/MA winter flounder mean weight per tow for annual state surveys.

Year	MADMF Spring	RIDFW Spring	RIDFW Fall	CTDEP	NJDFW Ocean (April)
1978	18.12				
1979	18.17	7.72	7.24		
1980	15.18	13.57	4.88		
1981	15.77	12.13	2.12		
1982	14.82	5.23	1.30		
1983	19.67	9.52	2.28		
1984	14.68	8.43	3.38	15.68	
1985	11.60	5.93	3.01	13.82	
1986	10.36	6.47	3.12	10.33	
1987	9.57	8.14	2.25	11.76	
1988	6.64	6.02	1.45	18.29	
1989	8.46	3.09	0.79	22.62	5.86
1990	5.38	3.07	0.71	29.02	4.78
1991	2.91	7.38	0.18	24.59	5.32
1992	7.99	0.95	0.42	12.29	2.48
1993	8.16	0.22	0.50	10.26	3.87
1994	12.59	1.67	0.33	12.20	3.25
1995	7.98	6.04	0.89	7.72	8.06
1996	9.78	4.45	0.91	20.41	3.73
1997	10.02	4.57	0.64	15.53	6.52
1998	7.99	5.00	0.32	14.66	4.17
1999	4.44	3.66	0.57	10.29	6.83
2000	6.52	4.52	0.56	12.63	5.24
2001	3.73	3.56	0.28	14.02	6.36
2002				10.90	8.80
Mean	10.44	5.71	1.66	15.11	5.38

Table B1.16. Winter flounder mean number per tow for annual state surveys.

Year	MADMF Spring	RIDFW Spring	RIDFW Fall	CTDEP	NYDEC (Age-1)	NJDFW Ocean (April)	NJDFW Rivers (March- May)
1978	51.62						
1979	53.78	83.76					
1980	38.94	63.10					
1981	46.12	87.97	25.21				
1982	40.23	31.39	18.55				
1983	56.84	58.97	17.29				
1984	37.36	41.64	19.02	111.96			
1985	38.38	34.97	21.44	83.05	1.96		
1986	36.27	41.02	31.28	63.64			
1987	37.85	56.21	20.90	79.92	1.64		
1988	27.91	34.44	10.64	153.08	1.32		
1989	24.41	20.88	7.17	150.08	3.01	25.60	
1990	25.86	20.33	8.83	226.17	1.79	17.47	
1991	10.66	41.95	1.77	156.06	3.38	22.17	
1992	28.83	4.40	10.60	75.09	1.11	9.88	
1993	46.96	2.92	6.65	69.60	5.42	20.13	
1994	48.55	10.25	2.21	101.60	3.16	14.16	
1995	37.84	32.19	7.00	62.62	1.72	30.04	3.00
1996	30.18	20.67	7.79	129.82	1.32	9.60	3.30
1997	39.31	22.28	5.48	78.79	3.15	36.24	3.60
1998	34.63	19.22	2.02	82.21	3.80	18.05	4.90
1999	25.11	13.45	2.80	50.05	3.25	17.84	3.20
2000	26.23	16.32	2.58	49.74	1.56	10.13	2.60
2001	16.00	12.49	2.10	55.80	5.52	13.83	2.90
2002				43.74		22.72	
Mean	35.83	33.51	11.02	95.95	2.69	19.13	3.36

Table B1.17. State survey indices (stratified mean number per tow or haul) for young-of-year winter flounder in Southern New England/Mid-Atlantic stock complex.

Year	MADMF Seine	RIDFW Seine	CTDEP	NYDEC	DEDFG
1975	0.30				
1976	0.32				
1977	0.60				
1978	0.34				
1979	0.49				
1980	0.40				
1981	0.32				
1982	0.37				
1983	0.23				
1984	0.32				
1985	0.34			0.75	
1986	0.32	29.00			0.17
1987	0.27	11.60		0.97	0.09
1988	0.18	8.90	15.50	0.69	0.02
1989	0.42	18.90	1.90	1.67	0.29
1990	0.33	22.10	3.10	2.71	0.63
1991	0.27	12.00	5.80	2.57	0.03
1992	0.29	33.20	13.70	11.49	0.27
1993	0.07	5.50	6.00	4.73	0.04
1994	0.15	2.60	16.60	2.44	0.31
1995	0.16	5.30	12.50	0.91	0.10
1996	0.22	2.80	19.20	3.80	0.04
1997	0.39	4.40	7.47	4.42	
1998	0.16	2.50	9.38	3.11	
1999	0.19	14.60	8.70	7.49	
2000	0.33	52.90	4.30	0.90	
2001	0.21	12.90	1.30	2.31	
2002	0.10				
Mean	0.27	14.95	8.96	3.19	0.18

Table B1.18. NEFSC Spring survey: stratified mean number per tow at age for winter flounder in the Southern New England/Mid-Atlantic stock complex (strata set: offshore 1-12, 5, 69-76; inshore 1-29, 45-56).

Year	Age									Total	
	1	2	3	4	5	6	7	8	9+		
1980	2.19	8.21	4.15	0.51	0.15	0.04					15.25
1981	2.00	8.08	6.89	0.95	0.26	0.02	0.03				18.23
1982	1.16	3.20	1.56	0.74	0.21	0.09	0.02	0.01			6.99
1983	0.58	0.97	2.14	1.23	0.81	0.37	0.08	0.08			6.26
1984	0.22	1.36	2.18	0.85	0.46	0.29	0.07	0.06	0.03		5.52
1985	0.41	1.21	2.16	0.72	0.51	0.20	0.14	0.01			5.36
1986	0.10	0.49	1.16	0.31	0.15	0.05	0.01				2.27
1987	0.14	0.54	0.70	0.28	0.06	0.02		0.01	0.01		1.76
1988	0.09	0.48	0.99	0.37	0.16	0.02	0.02				2.13
1989	0.14	0.95	0.90	0.34	0.11	0.02	0.02	0.01			2.49
1990	0.23	0.49	0.89	0.28	0.05	0.04	0.01				1.99
1991	0.14	0.60	1.22	0.41	0.05	0.02	0.02	0.01			2.47
1992	0.14	0.39	0.62	0.36	0.05	0.02					1.58
1993	0.14	0.35	0.26	0.12	0.07	0.01	0.01				0.96
1994	0.16	0.74	0.43	0.11	0.04	0.02	0.01				1.51
1995	0.22	0.75	0.87	0.22	0.03		0.01				2.10
1996	0.07	0.54	0.66	0.17	0.06	0.01	0.01				1.52
1997	0.13	0.50	0.56	0.18	0.06	0.01					1.44
1998	0.33	1.21	0.72	0.37	0.13	0.01					2.77
1999	0.41	1.89	1.35	0.36	0.11	0.04	0.01				4.17
2000	0.28	0.70	1.19	0.65	0.27	0.07	0.01				3.17
2001	0.17	0.26	0.47	0.44	0.20	0.02	0.01				1.57
2002	0.11	0.60	0.56	0.38	0.23	0.11	0.04		0.01		2.04
Mean	0.42	1.50	1.42	0.45	0.18	0.07	0.03	0.03	0.02		4.07

Table B1.19. NEFSC Fall survey: stratified mean number per tow at age for winter flounder in the Southern New England/Mid-Atlantic stock complex (strata set: offshore 1-12, 5, 69-76; inshore 1-29, 45-56).

Year	Age									Total
	0	1	2	3	4	5	6	7	8+	
1980	0.40	1.76	4.62	2.74	0.44	0.01	0.01			9.98
1981	0.01	2.06	5.05	2.30	0.31	0.06	0.08	0.03		9.90
1982	0.01	0.76	2.21	1.34	0.47	0.12	0.02			4.93
1983		1.63	3.82	2.06	0.62	0.35	0.11	0.07	0.10	8.76
1984		0.17	1.04	1.17	0.26	0.03	0.01			2.68
1985		0.16	1.18	0.99	0.30	0.09	0.01			2.73
1986		0.23	0.90	0.36	0.03	0.01		0.01		1.54
1987		0.03	0.64	0.36	0.12	0.02				1.17
1988		0.03	0.30	0.64	0.22	0.04	0.01	0.01		1.25
1989		0.28	0.83	0.26	0.05	0.01	0.01			1.44
1990		0.08	0.89	0.85	0.15	0.01				1.98
1991		0.07	1.02	0.73	0.12	0.01				1.95
1992		0.13	1.74	0.79	0.26	0.03	0.01			2.96
1993		0.43	0.52	0.35	0.08					1.38
1994		0.45	2.23	1.08	0.30	0.04	0.03			4.13
1995		0.58	0.93	0.63	0.09	0.01	0.01			2.25
1996		0.61	1.40	0.80	0.31	0.06	0.01			3.19
1997		1.48	3.58	2.20	0.55	0.08				7.89
1998		1.39	2.83	1.91	0.41	0.05	0.01			6.60
1999		0.43	0.95	1.46	0.54	0.18	0.04			3.60
2000		0.90	2.30	2.02	0.71	0.22	0.01	0.01		6.17
2001		0.49	1.79	1.61	0.63	0.30	0.02	0.04		4.88
2002										
Mean		0.64	1.85	1.21	0.32	0.08	0.03	0.03	0.10	4.26

Table B1.20. NEFSC Winter survey: stratified mean number per tow at age for winter flounder in the Southern New England/Mid-Atlantic stock complex (strata set: offshore 1-2, 5-6, 9-10,69, 73).

Year	Age								Total
	1	2	3	4	5	6	7	8+	
1992	0.73	0.86	1.09	0.73	0.24	0.02	0.02		3.68
1993	0.56	1.16	0.54	0.18	0.12	0.02	0.01		2.59
1994	0.36	1.16	1.76	0.25	0.28				3.80
1995	0.04	0.75	1.26	0.17					2.22
1996	1.01	0.87	1.55	0.32	0.02				3.78
1997	0.43	1.49	1.32	0.54	0.13				3.91
1998	0.42	3.52	1.95	0.96	0.32				7.17
1999	0.84	5.94	2.23	0.96	0.20	0.16			10.33
2000	0.23	2.82	2.12	0.24	0.16				5.57
2001	1.04	0.55	0.70	0.54	0.22	0.05			3.10
2002	0.08	1.34	0.74	0.15	0.21	0.06	0.21	0.11	2.90
Mean	0.52	1.86	1.39	0.46	0.19	0.06	0.08	0.11	4.46

Table B1.21. MADMF spring trawl survey mean number per tow at age for winter flounder in the Southern New England/Mid-Atlantic stock complex.

Year	Age									Total
	1	2	3	4	5	6	7	8	9+	
1978	9.93	9.73	15.74	9.33	3.15	1.09	1.33	0.51	0.81	51.62
1979	4.63	12.92	21.14	8.90	2.93	1.00	0.95	0.46	0.85	53.78
1980	1.63	8.21	14.50	9.13	3.01	0.96	0.79	0.28	0.43	38.94
1981	8.35	8.75	13.17	9.38	3.68	1.16	0.75	0.32	0.56	46.12
1982	3.22	11.13	12.36	8.62	2.61	1.05	0.67	0.15	0.42	40.23
1983	1.68	14.84	17.42	13.87	4.08	2.31	1.18	0.56	0.90	56.84
1984	1.17	9.34	11.62	10.06	3.32	1.22	0.48	0.01	0.14	37.36
1985	2.96	9.53	16.09	6.30	2.44	0.73	0.24	0.02	0.07	38.38
1986	3.23	6.81	19.13	5.64	0.82	0.12	0.18	0.16	0.18	36.27
1987	9.29	7.44	11.68	6.46	2.02	0.43	0.35	0.08	0.10	37.85
1988	3.21	7.22	14.45	2.41	0.34	0.08	0.17	0.00	0.03	27.91
1989	2.09	5.41	11.39	4.52	0.96	0.28	0.27	0.12	0.37	25.41
1990	4.22	10.66	7.60	2.90	0.32	0.05	0.10		0.01	25.86
1991	1.64	2.79	4.68	1.15	0.23	0.12	0.02		0.03	10.66
1992	7.93	7.55	6.68	4.16	1.64	0.59	0.07	0.08	0.13	28.83
1993	14.17	17.56	11.70	2.71	0.62	0.14	0.02	0.04		46.96
1994	11.48	16.12	14.65	4.66	0.61	0.58	0.37	0.05	0.03	48.55
1995	13.82	12.05	8.17	1.92	0.60	0.80	0.28	0.14	0.06	37.84
1996	4.81	9.73	7.61	2.84	1.99	1.45	0.84	0.29	0.62	30.18
1997	10.34	10.06	10.38	4.26	1.32	1.01	0.49	0.75	0.70	39.31
1998	8.17	12.59	6.92	3.51	1.46	1.22	0.41	0.31	0.04	34.63
1999	9.23	7.91	5.59	1.79	0.20	0.23	0.13	0.03		25.11
2000	6.62	8.94	6.95	1.69	1.05	0.48	0.22	0.25	0.03	26.23
2001	5.21	5.17	2.46	2.03	0.63	0.19	0.14	0.13	0.04	16.00
Mean	6.21	9.69	11.34	5.34	1.67	0.72	0.44	0.22	0.30	35.87

Table B1.22. CTDEP spring survey for winter flounder in the Southern New England/Mid Atlantic stock complex.

Year	Age													Total
	0	1	2	3	4	5	6	7	8	9	10	11	12	
1984	-	8.21	44.50	31.47	20.83	4.23	1.23	0.67	0.74	0.04	0.01	0.03	0.00	111.96
1985	-	4.10	28.28	32.57	14.13	2.33	0.83	0.45	0.19	0.11	0.04	0.02	0.00	83.05
1986	-	6.69	25.91	15.62	12.27	2.04	0.50	0.25	0.24	0.09	0.01	0.02	0.00	63.64
1987	-	7.32	44.69	14.56	5.05	6.55	1.29	0.11	0.24	0.11	0.00	0.00	0.00	79.92
1988	15.50	14.49	71.87	39.10	8.60	1.82	1.45	0.17	0.04	0.02	0.02	0.00	0.00	153.08
1989	1.90	13.57	78.42	41.23	10.85	2.84	0.98	0.13	0.09	0.06	0.01	0.00	0.00	150.08
1990	3.10	11.31	131.52	64.97	8.97	4.08	1.96	0.19	0.05	0.00	0.02	0.00	0.00	226.17
1991	5.80	8.66	66.88	60.41	9.31	4.05	0.80	0.13	0.01	0.00	0.00	0.01	0.00	156.06
1992	13.70	6.80	31.32	12.78	8.98	1.10	0.36	0.05	0.00	0.00	0.00	0.00	0.00	75.09
1993	6.00	19.11	19.87	15.46	4.81	3.24	0.79	0.15	0.12	0.04	0.01	0.00	0.00	69.60
1994	16.60	9.54	64.06	5.90	3.06	1.15	0.50	0.17	0.06	0.01	0.01	0.00	0.00	101.06
1995	12.50	14.35	23.69	9.77	1.36	0.63	0.20	0.08	0.02	0.02	0.00	0.00	0.00	62.62
1996	19.20	11.46	59.07	24.17	14.41	0.98	0.29	0.13	0.06	0.04	0.01	0.00	0.00	129.82
1997	7.47	12.53	25.53	19.41	9.45	3.76	0.51	0.07	0.03	0.01	0.01	0.01	0.00	78.79
1998	9.28	11.30	32.48	12.18	12.60	3.09	1.05	0.15	0.01	0.07	0.00	0.00	0.00	82.21
1999	8.70	6.53	12.42	11.29	6.09	3.21	1.13	0.61	0.04	0.01	0.02	0.00	0.00	50.05
2000	4.30	7.11	16.66	8.40	7.70	3.44	1.53	0.31	0.26	0.01	0.01	0.00	0.01	49.74
2001	1.30	8.37	19.65	10.87	8.06	5.46	1.26	0.70	0.04	0.09	0.00	0.00	0.00	55.80
2002														0.00
Mean	8.95	10.08	44.27	23.90	9.25	3.00	0.93	0.25	0.12	0.04	0.01	0.01	0.00	100.81

Table B1.23. RIDFW spring survey for winter flounder in the Southern New England-Mid Atlantic stock complex.

Year	Age							Total
	1	2	3	4	5	6	7+	
1981	13.55	32.2	32.99	6.07	1.85	0.79	0.48	87.93
1982	10.59	10.28	6.24	3.21	0.74	0.12	0.14	31.32
1983	16.75	18.51	11.63	7.61	1.9	0.84	0.25	57.49
1984	3.31	21.97	10.46	4.17	1.19	0.3	0.08	41.48
1985	3.77	13.42	14.19	2.44	0.81	0.07	0.04	34.74
1986	9.65	14.16	12.5	3.79	0.57	0.04	0.08	40.79
1987	12.44	20.56	17.09	4.24	0.91	0.14	0.09	55.47
1988	7.33	12.05	10.97	2.94	0.36	0	0.02	33.67
1989	6.67	6.32	5.55	1.58	0.32	0.1	0.03	20.57
1990	5.73	7.63	4.51	2.09	0.19	0.03	0.05	20.23
1991	12.48	14.67	11.29	2.14	0.48	0.22	0.02	41.30
1992	1.19	1.36	1.13	0.51	0.18	0.03	0	4.40
1993	2.35	0.26	0.18	0.05	0.01	0	0	2.85
1994	2.87	4.74	1.9	0.59	0.08	0.02	0.01	10.21
1995	8.33	9.53	11.22	2.03	0.43	0.45	0.2	32.19
1996	2.11	6.45	4.07	1.42	0.53	0.25	0.11	14.94
1997	4.47	7.79	7.42	1.69	0.45	0.25	0.18	22.25
1998	1.5	4.16	8.43	3.87	0.7	0.46	0.11	19.23
1999	1.61	4.07	5.45	1.84	0.16	0.16	0.13	13.42
2000	2.99	4.91	6.09	1.32	0.65	0.20	0.12	16.28
2001	2.11	4.23	2.89	2.53	0.57	0.04	0.08	12.45
Mean	6.28	10.44	8.87	2.67	0.62	0.21	0.11	29.20

Table B1.24. NJDFW Ocean survey (April) for winter flounder in the Southern New England/Mid-Atlantic stock complex. Lengths for 2002 aged with the 2001 age-length key.

Year	Age							Total
	1	2	3	4	5	6	7+	
1993	5.1	6.5	2.5	2.4	1.7	0.4	0.57	19.17
1994	3.7	4.2	3.9	1.4	0.4	0.3	0.16	14.06
1995	8	10.1	8.6	2.4	0.9	0.3	0.11	30.41
1996	0.6	2.9	2.6	1.9	0.9	0.3	0.2	9.40
1997	16.6	5.4	6.1	6	1.5	0.3	0.12	36.02
1998	4.5	3.9	4.8	3.3	1.2	0.4	0.1	18.20
1999	2.40	2.20	5.90	3.10	2.90	0.70	0.59	17.79
2000	0.70	0.30	2.10	3.30	2.00	0.90	0.80	10.10
2001	3.90	0.60	1.30	2.70	3.80	0.70	0.83	13.83
2002	7.56	3.67	3.30	3.00	3.67	0.76	0.77	22.73
Mean	5.06	4.01	4.20	2.94	1.70	0.48	0.39	18.78

Table B1.25. NJDFW Rivers survey (March-May) for winter flounder in the Southern New England/Mid Atlantic stock complex.

Year	Age							Total
	1	2	3	4	5	6	7+	
1995	0.6	0.3	1.4	0.4	0.1	0.01	0.01	2.82
1996	0.3	0.9	0.7	0.7	0.2	0.1	0.15	3.05
1997	1.1	0.4	0.9	0.4	0.4	0.1	0.05	3.35
1998	1.9	0.9	0.4	0.7	0.2	0.1	0.05	4.25
1999	0.20	0.50	1.40	0.50	0.40	0.10	0.13	3.23
2000	0.40	0.20	0.40	0.80	0.20	0.10	0.01	2.11
2001	1.40	0.30	0.20	0.40	0.40	0.10	0.04	2.84
Mean	0.84	0.50	0.77	0.56	0.27	0.09	0.06	3.09

Table B1.26. Virtual Population Analysis for SNE/MA winter flounder, 1981-2001.

Fisheries Assessment Toolbox SNE/MA Winter Flounder Run Number W36_2 9/25/2002 1:11:40 PM
 FACT Version 1.5.0

SNE/MA Winter Flounder 1981 - 2002
 Input Parameters and Options Selected

 Natural mortality is a matrix below
 Oldest age (not in the plus group) is 6
 For all years prior to the terminal year (21), backcalculated
 stock sizes for the following ages used to estimate
 total mortality (Z) for age 6 : 4 5 6
 This method for estimating F on the oldest age is generally used when a
 flat-topped partial recruitment curve is thought to be characteristic of the stock.

Stock size of the 7 + group is then calculated using
 the following method: CATCH EQUATION

Partial recruitment estimate for 2002

1	0.01
2	0.2
3	0.6
4	1
5	1
6	1

The Indices that will be used in this run are:

1	NEC_S1
2	NEC_S2
3	NEC_S3
4	NEC_S4
5	NEC_S5
6	NEC_S6
7	NEC_S7
8	NEC_F2
9	NEC_F3
10	NEC_F4
11	NEC_W1
12	NEC_W2
13	NEC_W3
14	NEC_W4
15	NEC_W5
16	MA_S2
17	MA_S3
18	MA_S4
19	MA_S5
20	RI_S1
21	RI_S2
22	RI_S3
23	RI_S4
24	CT_S1
25	CT_S2
26	CT_S3
27	CT_S4
28	CT_S5
29	CT_S6
30	CT_S7
31	MA_YOY1
32	CT_YOY1
33	NY_PB1.1
34	NJ_O3
35	NJ_O4
36	NJ_O5
37	NJ_O6
38	NJ_O7
39	NJ_R1
40	NJ_R2
41	NJ_R3
42	NJ_R4
43	NJ_R5

Table B1.26 continued.

STOCK NUMBERS (Jan 1) in thousands							
	1981	1982	1983	1984	1985	1986	1987
1	62859	52020	56503	35617	34615	32795	25973
2	52566	50232	42060	45703	28708	28090	26656
3	27768	30289	28226	27884	26945	16839	17273
4	7146	9748	13560	11068	10077	10446	5551
5	1468	2600	4606	5559	4603	2773	4738
6	363	600	1577	2148	2944	1096	1317
7	218	564	1219	1949	2228	876	730
1+	152388	146054	147751	129927	110120	92914	82238
	1988	1989	1990	1991	1992	1993	1994
1	26726	23113	17366	11355	7808	8844	8315
2	21199	21806	18504	14185	9249	6370	6993
3	17057	13790	13106	13242	8875	6212	3350
4	6000	5458	4798	5053	4381	3233	2074
5	1748	1325	1299	1276	1111	1251	1084
6	1433	339	317	369	268	300	495
7	433	312	223	165	86	218	300
1+	74596	66142	55613	45645	31778	26429	22611
	1995	1996	1997	1998	1999	2000	2001
1	12647	17632	21154	18793	13372	12710	19011
2	6753	10333	14407	16971	15341	10889	10343
3	4733	5352	7658	9864	11966	10076	7610
4	1700	2190	3070	3284	4761	6170	5082
5	1053	588	791	875	1063	2320	2830
6	606	487	171	159	254	456	1120
7	433	312	73	228	83	168	512
1+	27925	36893	47324	50174	46840	42788	46509
	2002						
1	5665						
2	15553						
3	6671						
4	2912						
5	2179						
6	1602						
7	1057						
1+	35639						

Table B1.26 continued.

FISHING MORTALITY							
	1981	1982	1983	1984	1985	1986	1987
1	0.02	0.01	0.01	0.02	0.01	0.01	0.00
2	0.35	0.38	0.21	0.33	0.33	0.29	0.25
3	0.85	0.60	0.74	0.82	0.75	0.91	0.86
4	0.81	0.55	0.69	0.68	1.09	0.59	0.96
5	0.69	0.30	0.56	0.44	1.23	0.54	1.00
6	0.81	0.50	0.67	0.60	1.18	0.59	1.00
7	0.81	0.50	0.67	0.60	1.18	0.59	1.00
	1988	1989	1990	1991	1992	1993	1994
1	0.00	0.02	0.00	0.01	0.00	0.03	0.01
2	0.23	0.31	0.13	0.27	0.20	0.44	0.19
3	0.94	0.86	0.75	0.91	0.81	0.90	0.48
4	1.31	1.24	1.12	1.31	1.05	0.89	0.48
5	1.44	1.23	1.06	1.36	1.11	0.73	0.38
6	1.41	1.29	1.15	1.39	1.10	0.86	0.45
7	1.41	1.29	1.15	1.39	1.10	0.86	0.45
	1995	1996	1997	1998	1999	2000	2001
1	0.00	0.00	0.02	0.00	0.01	0.01	0.00
2	0.03	0.10	0.18	0.15	0.22	0.16	0.24
3	0.57	0.36	0.65	0.53	0.46	0.48	0.76
4	0.86	0.82	1.06	0.93	0.52	0.58	0.65
5	0.57	1.04	1.40	1.04	0.65	0.53	0.37
6	0.76	0.88	1.16	0.98	0.55	0.57	0.23
7	0.76	0.88	1.16	0.98	0.55	0.57	0.23
Average F for 4,5							
	1981	1982	1983	1984	1985	1986	1987
4,5	0.75	0.42	0.63	0.56	1.16	0.57	0.98
	1988	1989	1990	1991	1992	1993	1994
4,5	1.38	1.23	1.09	1.34	1.08	0.81	0.43
	1995	1996	1997	1998	1999	2000	2001
4,5	0.72	0.93	1.23	0.98	0.58	0.55	0.51
Biomass Weighted F							
	1981	1982	1983	1984	1985	1986	1987
	0.47	0.42	0.38	0.47	0.61	0.44	0.58
	1988	1989	1990	1991	1992	1993	1994
	0.67	0.56	0.48	0.68	0.64	0.60	0.28
	1995	1996	1997	1998	1999	2000	2001
	0.30	0.23	0.42	0.31	0.36	0.37	0.39

Table B1.26 continued.

BACK-CALCULATED PARTIAL RECRUITMENT							
	1981	1982	1983	1984	1985	1986	1987
1	0.03	0.02	0.02	0.02	0.01	0.01	0.00
2	0.41	0.62	0.29	0.40	0.27	0.31	0.25
3	1.00	1.00	1.00	1.00	0.61	1.00	0.85
4	0.96	0.91	0.94	0.83	0.88	0.65	0.95
5	0.82	0.50	0.76	0.53	1.00	0.60	0.99
6	0.95	0.82	0.91	0.73	0.95	0.65	1.00
7	0.95	0.82	0.91	0.73	0.95	0.65	1.00
	1988	1989	1990	1991	1992	1993	1994
1	0.00	0.02	0.00	0.00	0.00	0.04	0.02
2	0.16	0.24	0.12	0.19	0.18	0.49	0.40
3	0.65	0.66	0.65	0.65	0.73	1.00	1.00
4	0.91	0.96	0.98	0.95	0.95	1.00	1.00
5	1.00	0.95	0.92	0.98	1.00	0.81	0.80
6	0.98	1.00	1.00	1.00	0.99	0.96	0.94
7	0.98	1.00	1.00	1.00	0.99	0.96	0.94
	1995	1996	1997	1998	1999	2000	2001
1	0.00	0.00	0.01	0.00	0.01	0.01	0.00
2	0.04	0.10	0.13	0.14	0.34	0.27	0.31
3	0.66	0.34	0.46	0.51	0.71	0.84	1.00
4	1.00	0.79	0.75	0.89	0.80	1.00	0.85
5	0.66	1.00	1.00	1.00	1.00	0.91	0.48
6	0.88	0.85	0.83	0.94	0.85	0.99	0.31
7	0.88	0.85	0.83	0.94	0.85	0.99	0.31
MEAN BIOMASS (using catch mean weights at age)							
	1981	1982	1983	1984	1985	1986	1987
1	7320	4218	9928	4678	3468	3821	1081
2	11153	9965	8174	9159	6274	6496	6171
3	8228	9117	6470	6403	6338	4048	4094
4	3606	4760	4630	3994	2728	3465	1813
5	1033	2144	2494	2851	1276	1340	1313
6	284	541	1102	1350	747	666	432
7	165	745	1264	1720	742	582	362
1+	31790	31490	34061	30156	21572	20418	15266
	1988	1989	1990	1991	1992	1993	1994
1	943	2445	1289	955	558	1332	1171
2	4807	4409	4639	3590	2190	1570	2009
3	3573	3219	3329	3364	2386	1735	1090
4	1573	1472	1396	1390	1496	1164	899
5	544	465	497	390	497	562	609
6	369	126	140	152	142	162	252
7	178	174	121	97	67	157	180
1+	11987	12310	11412	9937	7335	6682	6210
	1995	1996	1997	1998	1999	2000	2001
1	1912	3081	1766	3436	955	1149	1757
2	1946	3634	4425	4755	3933	3624	3508
3	1482	2082	2635	3068	3806	3532	2475
4	605	782	1152	1142	1908	2627	2180
5	524	238	318	334	561	1216	1718
6	298	246	100	84	170	290	957
7	218	163	60	166	77	134	523
1+	6984	10225	10456	12985	11410	12571	13118

Table B1.26 continued.

SSB AT THE START OF THE SPAWNING SEASON -MALES AND FEMALES (MT) (using SSB mean weights)							
	1981	1982	1983	1984	1985	1986	1987
1	00	00	00	00	00	00	00
2	00	00	00	00	00	00	00
3	4739	4757	3771	3557	3615	2395	2482
4	3893	4592	5119	3855	3106	3541	1958
5	1205	2157	2899	2927	1838	1374	1779
6	341	603	1387	1540	1272	634	644
7	214	900	1590	2129	1037	718	489
1+	10393	13009	14766	14008	10869	8662	7353
	1988	1989	1990	1991	1992	1993	1994
1	00	00	00	00	00	00	00
2	00	00	00	00	00	00	00
3	2282	1923	1831	1980	1414	960	600
4	1863	1642	1556	1627	1626	1242	902
5	744	576	590	526	559	667	639
6	516	169	177	200	156	203	300
7	260	248	169	140	93	206	215
1+	5663	4559	4323	4474	3848	3278	2656
	1995	1996	1997	1998	1999	2000	2001
1	00	00	00	00	00	00	00
2	00	00	00	00	00	00	00
3	849	1028	1563	1817	2128	1756	2579
4	665	857	1311	1354	1990	2548	2103
5	589	293	389	452	563	1251	1692
6	376	301	113	107	170	296	715
7	279	214	84	224	73	169	553
1+	2759	2693	3459	3954	4923	6021	7643

Table B1.26 continued.

Fishing Mortality
Terminal Year

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
1997	0.75	0.42	0.63	0.56	1.16	0.57	0.98	1.37	1.23	1.08	1.31	1.02	0.71	0.34	0.50	0.47	0.37				
1998	0.75	0.42	0.63	0.56	1.16	0.57	0.98	1.38	1.23	1.09	1.33	1.05	0.76	0.38	0.60	0.65	0.54	0.32			
1999	0.75	0.42	0.63	0.56	1.16	0.57	0.98	1.38	1.23	1.09	1.33	1.07	0.79	0.41	0.65	0.76	0.77	0.38	0.36		
2000	0.75	0.42	0.63	0.56	1.16	0.57	0.98	1.38	1.23	1.09	1.34	1.08	0.81	0.42	0.71	0.89	1.10	0.74	0.39	0.59	
2001	0.75	0.42	0.63	0.56	1.16	0.57	0.98	1.38	1.23	1.09	1.34	1.08	0.81	0.43	0.72	0.93	1.23	0.98	0.58	0.55	0.51

Spawning Stock Biomass
Terminal Year

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
1997	10393	13009	14767	14009	10869	8663	7354	5666	4566	4343	4548	4038	3670	3273	3849	4826	7444				
1998	10393	13009	14766	14008	10869	8662	7354	5664	4562	4331	4505	3929	3445	2919	3220	3833	6041	7845			
1999	10393	13009	14766	14008	10869	8662	7354	5664	4561	4327	4488	3887	3355	2783	2969	3357	5233	6245	7280		
2000	10393	13009	14766	14008	10869	8662	7354	5663	4560	4323	4477	3856	3295	2681	2807	2781	3971	4866	5537	6897	
2001	10393	13009	14766	14008	10869	8662	7353	5663	4559	4323	4474	3848	3278	2656	2759	2693	3459	3954	4923	6021	7643

Population Numbers Age: 1
Terminal Year

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
1997	62859	52021	56504	35618	34618	32804	26001	26802	23425	17857	12277	8527	11725	13557	19744	19471	31502	21889			
1998	62859	52021	56504	35617	34617	32799	25985	26759	23243	17580	11733	8140	9992	12293	17810	18933	27084	31936	31205		
1999	62859	52021	56504	35617	34616	32797	25979	26743	23169	17482	11501	8036	9204	11387	16304	17649	22197	21574	17992	15496	
2000	62859	52020	56503	35617	34616	32795	25975	26729	23127	17384	11400	7826	8998	8449	15177	17596	20214	19212	13851	13085	15615
2001	62859	52020	56503	35617	34615	32795	25973	26726	23113	17366	11355	7808	8844	8315	12647	17632	21154	18793	13372	12710	19011

Table B1.27. VPA Bootstrap results: precision of estimates.

The number of bootstraps: 500

Bootstrap Output Variable: N hat

	NLLS ESTIMATE	BOOTSTRAP MEAN	BOOTSTRAP StdError	C.V. FOR NLLS SOLN			
N 1	5665	5960	1905	0.34			
N 2	15553	15895	3191	0.21			
N 3	6671	6691	1176	0.18			
N 4	2912	2938	648	0.22			
N 5	2179	2208	504	0.23			
N 6	1602	1631	369	0.23			
N 7	726	736	159	0.22			

	BIAS ESTIMATE	BIAS STD ERROR	PERCENT BIAS	NLLS EST CORRECTED FOR BIAS	C.V. FOR CORRECTED ESTIMATE	LOWER 80%CI	UPPER 80%CI
N 1	294	85	5.20	5371	0.354683	3746	8342
N 2	342	143	2.20	15211	0.209761	11856	19828
N 3	21	53	0.31	6650	0.176795	5097	8050
N 4	26	29	0.90	2886	0.224561	2114	3811
N 5	29	23	1.35	2149	0.234726	1611	2910
N 6	28	16	1.78	1574	0.234188	1158	2114
N 7	10	07	1.42	715	0.221624	534	937

Bootstrap Output Variable: F t

	NLLS ESTIMATE	BOOTSTRAP MEAN	BOOTSTRAP StdError	C.V. FOR NLLS SOLN			
Age 1	0.0008	0.0008	0.0002	0.21			
Age 2	0.2386	0.2440	0.0395	0.17			
Age 3	0.7607	0.7755	0.1236	0.16			
Age 4	0.6471	0.6599	0.1136	0.18			
Age 5	0.3689	0.3773	0.0754	0.20			
Age 6	0.2336	0.2397	0.0491	0.21			
Age 7	0.2336	0.2397	0.0491	0.21			

	BIAS ESTIMATE	BIAS STD ERROR	PERCENT BIAS	NLLS EST CORRECTED FOR BIAS	C.V. FOR CORRECTED ESTIMATE	LOWER 80%CI	UPPER 80%CI
Age 1	0.0000137	0.0000070	1.815	0.0007423	0.21	0.0006	0.0010
Age 2	0.0053697	0.0017644	2.250	0.2332777	0.17	0.2014	0.3004
Age 3	0.0147454	0.0055258	1.938	0.7459918	0.17	0.6241	0.9413
Age 4	0.0128214	0.0050816	1.981	0.6342562	0.18	0.5193	0.8015
Age 5	0.0084273	0.0033716	2.285	0.3604493	0.21	0.2905	0.4802
Age 6	0.0061558	0.0021967	2.635	0.2274158	0.22	0.1853	0.3045
Age 7	0.0061558	0.0021967	2.635	0.2274158	0.22	0.1853	0.3045

Bootstrap Output Variable: SSB spawn t

	NLLS ESTIMATE	BOOTSTRAP MEAN	BOOTSTRAP StdError	C.V. FOR NLLS SOLN			
	7642.6469	7705.3234	658.0444	0.09			

	BIAS ESTIMATE	BIAS STD ERROR	PERCENT BIAS	NLLS EST CORRECTED FOR BIAS	C.V. FOR CORRECTED ESTIMATE	LOWER 80%CI	UPPER 80%CI
	62.68	29.43	0.82	7579.97	0.09	6777.3392	8444.6451

Table B1.28. Input parameters and stochastic projection results for winter flounder in the Southern New England/Mid-Atlantic stock complex. Starting stock sizes for ages 1 and older on January 1, 2002 are as estimated by SARC 36 VPA, and are not adjusted for the retrospective pattern. Age-1 recruitment levels in 2003 and later years are estimated from a parametric stock-recruitment relationship estimated in NEFSC (2002). Fishing mortality was apportioned among landings and discard based on the proportion landed at age during 1998-2000. Mean weights at age (kg; spawning stock, mean stock biomass, landings, and discards) are weighted (by fishery) geometric means of 1998-2000 values. Proportion of F, M before spawning = 0.20 (spawning peak on 1 March).

Age	Stock Size on 1 Jan 2002 (000s)	Fishing Mortality Pattern	Proportion Landed	Proportion Mature	Mean Weights Spawning Stock	Mean Weights Landings	Mean Weights Discards
1	5688	0.02	0.02	0	0.07	0.325	0.116
2	15592	0.27	0.7	0	0.196	0.383	0.242
3	6712	0.75	0.91	0.53	0.387	0.465	0.317
4	2908	1	0.97	0.95	0.52	0.59	0.417
5	2170	1	0.97	1	0.637	0.725	0.868
6	1612	1	0.97	1	0.793	0.916	0.853
7+	1064	1	0.97	1	1.144	1.125	1.402

F2002 is assumed 0.85*F2001 (15% decrease in F from 2001 to 2002); F during 2003-2013 as indicated; Forecast Medians (50% probability level)

2002				2003				2013				
F	Land	Disc	SSB	'000 Metric tons				F	Land	Disc	SSB	P (%) SSB > 30.1 kmt
0.43	3.0	0.2	5.9	Fsq=0.43	3.3	0.1	7.0	Fsq=0.43	8.0	0.5	16.4	0%
				Fmsy=0.32	2.6	0.2	7.2	Fmsy=0.32	8.3	0.5	23.3	6%
				Freb=0.24	2.0	0.1	7.3	Freb=0.24	8.1	0.4	30.1	50%

SNE/MA Winter Flounder Landings and Discards




Figure B1.1. Commercial landings (1964-2001), commercial discards (1981-2001) recreational landings (1981-2001), recreational discards (1981-2001) and total fishery catch (1981-2001) for the SNE/MA winter flounder stock complex.

SNE/MA winter flounder Total Catch Age Composition




Figure B1.2. Total catch age composition: 1981-2001




Figure B1.3. Trends in mean weight at age in the total catch of SNE/MA winter flounder.

SNE/MA Winter Flounder Survey Biomass Indices




Figure B1.4. Trends in research survey biomass indices for SNE/MA winter flounder.

SNE/MA Winter Flounder Recruitment Indices




Figure B1.5. Trends in survey recruitment indices for SNE/MA winter flounder.
Includes spring survey age-1 indices and fall YOY indices advanced one year

SNE/MA Winter Flounder Recruitment Indices




Figure B1.5 continued.

SNE/MA winter flounder VPA Sensitivity to Tuning Indices




Figure B1.6. Sensitivity of the SARC 36 VPA for SNE/MA winter flounder to alternative combination of survey tuning indices. Run W36_2 was selected as the final run.

SNE/MA Winter Flounder Total Catch and Fishing Mortality




Figure B1.7. Total catch (landings and discards, '000 mt), commercial landings('000 mt), and fishing mortality rate (F, ages 4-5, unweighted) for SNE/MA winter flounder.

SNE/MA Winter Flounder Precision of 2001 Estimates for SSB and F




Figure B1.8. Precision of estimates of spawning stock biomass (ages 3-7+, '000 mt) and fishing mortality rate (F, ages 4-5, unweighted) in 2001 for SNE/MA winter flounder. Vertical bars display the range of the bootstrap estimates and the probability of individual values in the range. The solid curve gives the probability of SSB that is less or fishing mortality that is greater than any value along the X axis.

SNE/MA Winter Flounder SSB and Recruitment




Figure B1.9. Spawning stock biomass (SSB, ages 3-7+, '000 mt) and recruitment (millions of fish at age-1) for SNE/MA winter flounder.

SNE/MA winter flounder retrospective VPAs




Figure B1.10. Retrospective VPAs for SNE/MA winter flounder.

SNE/MA Winter Flounder Yield and SSB per Recruit




Figure B1.11. Yield per recruit (YPR) and spawning stock biomass per recruit (SSB/R) for SNE/MA winter flounder.




Figure B1.12. SNE/MA winter flounder SARC 36 VPA SSB and recruit data for the 1981-2001 year classes. Curved line is the S-R function estimated by NEFSC (2002).

SNE/MA Winter Flounder




Figure B1.13. Median (50% probability) of forecast spawning stock biomass (SSB, mt) for SNE/MA winter flounder under F_{msy} and F_{rebuild} fishing mortality rates during 2003-2013. Assumes $F_{2002} = 0.85 * F_{2001} = 0.43$.

SNE/MA winter flounder sensitivity to hypothetical NEFSC survey index adjustments, 2000-2002




Figure B1.14. SNE/MA winter flounder VPA sensitivity to hypothetical NEFSC winter, spring, and fall survey index adjustments.

SNE/MA Winter Flounder




Figure B1.15. SSB and F for SNE/MA winter flounder. NEFSC (2002) biological reference points ($F_{msy} = 0.32$, $SSB_{msy} = 30,100$ mt) are also shown.