



ENVIRONMENTAL HEALTH PERSPECTIVES

ehponline.org

Methylmercury in Marine Ecosystems: From Sources to Seafood Consumers -- A Work Group Report --

Celia Y. Chen, Nancy Serrell, David C. Evers,
Bethany J. Fleishman, Kathleen F. Lambert,
Jeri Weiss, Robert P. Mason, and Michael S. Bank

doi:10.1289/ehp.11211 (available at <http://dx.doi.org/>)
Online 23 July 2008



NIEHS
National Institute of
Environmental Health Sciences

National Institutes of Health
U.S. Department of Health and Human Services

Methylmercury in Marine Ecosystems: From Sources to Seafood Consumers

-- A Work Group Report --

Celia Y. Chen¹, Nancy Serrell², David C. Evers³, Bethany J. Fleishman², Kathleen F. Lambert⁴,
Jeri Weiss⁵, Robert P. Mason⁶, and Michael S. Bank⁷.

¹Department of Biological Sciences, Dartmouth College, Hanover, NH;

²Dartmouth Toxic Metals Research Program, Dartmouth College, Hanover, NH;

³BioDiversity Research Institute, Gorham, ME;

⁴Ecologic: Analysis & Communications, Woodstock, VT;

⁵United State Environmental Protection Agency, Boston, MA;

⁶Department of Marine Sciences, University of Connecticut, Avery Point, CT;

⁷Department of Environmental Health, Harvard School of Public Health, Boston, MA.

Corresponding Author:

Celia Y. Chen

Research Associate Professor

Department of Biological Sciences

Dartmouth College

HB 6044

(603)646-2376 PH, (603)646-1347 FAX, celia.chen@dartmouth.edu

Acknowledgments

This publication was made possible by NIH Grant Number P42 ESO7373 from the National Institute of Environmental Health Sciences and the NIEHS Workshop Grant Number 3 P42 ES007373-13S1. We also thank all the workshop participants for their valuable input and insights. We acknowledge the guidance and participation of Dr. Claudia Thompson of NIEHS and of Dr. Alan Stern for his help with the editing of the manuscript. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NIH.

Running Title: Methylmercury in Marine Ecosystems

Key Words: bioaccumulation, human health, mercury biomonitoring, mercury exposure, methylmercury

Abbreviations and Definitions:

DGT Diffusive Gel Time-series

MeHg methylmercury

Abstract

Introduction

Theme 1: Biogeochemical Cycling of Mercury in Marine Ecosystems

Research Questions

Monitoring Needs

Theme 2: Methylmercury Trophic Transfer in Marine Food Webs

Research Questions

Monitoring Needs

Theme 3: Human Exposure to Mercury in Fish and Shellfish

Research Questions

Monitoring Needs

Summary

List of References

Tables

Abstract

Mercury and other contaminants in coastal and open ocean ecosystems are an issue of great concern globally and in the United States, where consumption of marine fish and shellfish is a major route of human exposure to methylmercury (MeHg). A recent NIEHS-SBRP workshop, *Fate and Bioavailability of Mercury in Aquatic Ecosystems and Effects on Human Exposure*, convened by the Dartmouth Toxic Metals Research Program on November 15-16, 2006 in Durham NH, brought together human health experts, marine scientists and ecotoxicologists to encourage cross-disciplinary discussion between ecosystem and human health scientists and to articulate research and monitoring priorities to better understand how marine food webs have become contaminated with MeHg. While human health effects of mercury contamination were a major theme of the workshop, effects on marine biota were also explored. The work group focused on three major topics: (1) the biogeochemical cycling of mercury in marine ecosystems, (2) the trophic transfer and bioaccumulation of methylmercury in marine food webs, and (3) human exposure to mercury from marine fish and shellfish consumption. The group concluded that current understanding of mercury in marine ecosystems across a range of habitats, chemical conditions, and ocean basins is severely data-limited. An integrated research and monitoring program is needed to link the processes and mechanisms of MeHg production, bioaccumulation, and transfer with MeHg exposure in humans.

Introduction

Methylmercury (MeHg), a toxic organic form of mercury, is the predominant form found in fish tissue. People who consume large amounts of fish contaminated with MeHg have a higher body burden than non-fish consumers (Mahaffey and Mergler 1998). The developing human nervous system is a sensitive target for MeHg exposure, putting developing fetuses and young children at the highest risk for harm (Clarkson et al. 2003; Grandjean et al. 2005; NAS 2000). Marine apex predators such as sharks, seabirds, and marine mammals are also at risk from elevated MeHg exposure (Braune et al. 2006; Garcia-Hernandez et al. 2007; Kemper et al. 1994).

To date, research on the environmental fate of Hg and MeHg has focused primarily on freshwater systems and upland watersheds. Far less research has been conducted in marine environments where the transport and migration of water, contaminants, and fisheries resources make the identification of sources and receptors of Hg and MeHg contamination particularly challenging (Knap et al. 2002). However, since the main vector for MeHg exposure in humans in the U.S. is fish and shellfish consumption — 60% of which is derived from marine systems (Sunderland 2007) — the links between MeHg bioaccumulation in estuarine, coastal, and open ocean ecosystems and human exposure need to be better understood.

The existing research in marine systems has focused largely on Hg biogeochemistry in the open ocean (Benoit et al. 2003; Fitzgerald et al. 2007; Hammerschmidt and Fitzgerald 2004; Laurier et al. 2004; Lawrence et al. 1999; Mason and Gill 2005; Mason et al. 1999; Mason and Lawrence 1999). A limited number of extensive studies have been conducted in specific coastal systems, such as Long Island Sound, Scheldt River Estuary, and Chesapeake Bay (Baeyens et al. 2003; Hammerschmidt and Fitzgerald 2006; Mason et al. 2006a,b). Early studies on health effects focused on acute exposures (Minimata Japan); later extensive studies have investigated

chronic low-level exposures (Seychelles and Faroe Islands, NAS 2000) resulting from consumption of marine organisms (the Seychelles population consumed marine fish whereas the Faroe Islands population consumed pilot whale). Fetal exposures to Hg in both these populations have been associated with neurological deficits in children, in some cases many years after exposure (Mergler et al. 2007). Although the breadth of Hg research is great, the links between marine sources and ultimate human exposure have not been made across large geographic areas. There is now a need to adopt a systems approach to the study of MeHg in marine ecosystems with more focus on the processes controlling the transfer of MeHg in marine food webs, particularly those that link MeHg sources to seafood consumed by humans.

With this in mind, the Dartmouth Superfund Basic Research Program, with support from NIEHS and New Hampshire Sea Grant, convened a workshop (“Fate and Bioavailability of Mercury in Aquatic Ecosystems and Effects on Human Exposure”) in November 2006. The goal was to bring together ecosystem scientists and human health scientists for cross-disciplinary discussions to identify research and monitoring priorities linking the fate and bioaccumulation of mercury in the marine environment to human exposure. This report summarizes the major research and monitoring needs identified for each of the three workshop themes: (1) the biogeochemical cycling of mercury in marine ecosystems, (2) the trophic transfer and bioaccumulation of MeHg in marine food webs, and (3) human exposure to mercury from marine fish and shellfish consumption. This report is not intended to present an exhaustive review of the state of the science, as several review papers on mercury in marine ecosystems have been published previously (see Table 1).

Theme 1: Biogeochemical Cycling of Mercury in Marine Ecosystems

Research and monitoring on the biogeochemical cycling of mercury in marine ecosystems is critical to expanding current understanding of the MeHg sources that contaminate marine fish. Important insights can be gained from the extensive research in freshwater ecosystems, but biogeochemical processes in freshwater and saltwater likely differ. For example, organic matter decreases from watersheds to the open ocean and sulfate concentrations increase; both constituents are known to strongly influence Hg bioavailability to sulfur-reducing bacteria (Benoit et al. 2003; Sunderland et al. 2006). They also affect the flux of Hg and MeHg from sediments to the water column. Freshwater studies and the existing literature on marine Hg cycling (see Table 1) need to be extended to investigate the factors controlling MeHg production, sediment flux, and biotransfer in marine ecosystems. The work group identified three questions that should be addressed: (1) where is mercury methylation occurring in the ocean? (2) How is methylmercury mobilized from sediments to water in coastal ecosystems? And (3) what is the relative importance of benthic biotransfer of mercury into aquatic food webs?

Research Questions

1. Where is mercury methylation occurring in the ocean?

Methylation is a key process in the transformation of inorganic mercury to the MeHg that bioaccumulates in food webs. Though Hg methylation has been studied in depth in coastal and open ocean environments, studies of tropical and polar regions and deep ocean basin are limited

There are three potential regions of methylation in marine ecosystems – coastal and slope sediments, low-oxygen waters below productive ocean waters, and deep ocean sediments (Kraepiel et al. 2003). Current research suggests that net MeHg production in coastal marine sediments is one of the more important sources and thus a potential source for MeHg in marine

fish (Hammerschmidt and Fitzgerald 2004; Sunderland et al. 2006). Measurements of methylation sources in a wider range of marine conditions are needed to evaluate the relative importance of coastal marine sediments to MeHg concentration in shellfish, fish and other biota.

Methylation in coastal sediments is largely controlled by bacterial activity and the bioavailability of inorganic Hg, which is highly dependent on sediment and porewater concentrations of organic carbon and sulfide. Some, though not all, sulfate-reducing bacteria are thought to be the primary methylators of Hg in sediment (Benoit et al. 2003). However, recent evidence suggests that other bacteria, including iron reducers, methylate Hg as well (Slowey and Brown 2007). Demethylation appears to be influenced by both abiotic (photochemical) and biotic processes, but less is known about these factors. The controls on both methylation and demethylation in marine systems should be a more active area of research.

These patterns would be most effectively examined within the context of total Hg and MeHg concentrations in the water column. However, concentrations of MeHg in most of the oceans, except for coastal oceans and the equatorial Pacific are low and difficult to measure (Table 2; Fitzgerald et al. 2007). For example, recent model results support the idea that the concentrations of total Hg and MeHg differ between ocean basins, as does the rate of change in concentrations (Laurier et al. 2004; Strode et al. 2007; Sunderland and Mason 2007) but more study and data are required to confirm these trends.

2. How is methylmercury mobilized from sediments to water in coastal ecosystems?

Although the main source of MeHg in coastal ecosystems is known to be microbial methylation in sediments, few studies have examined the rates of mobilization or diffusion of that MeHg from the source to the water column where it is available to the food web (e.g. Choe

et al. 2004; Covelli et al. 1999; Mason et al. 2006b). Though biogeochemical factors such as organic carbon, oxygen, and sulfide concentrations appear important, their combined influences on methylation, demethylation, and MeHg mobilization in marine systems are poorly understood. Existing studies suggest that their influence depends on factors that influence partitioning of MeHg to the solid phase, such as organic carbon and solid sulfide content, and sediment redox status (Gill et al. 1999; Mason et al. 2006b), but the relationships between biogeochemical conditions and the mobilization of methylmercury into marine waters needs to be clarified.

Determination of MeHg flux is constrained by the limitations of current methods, which can vary — depending on the method — by an order of magnitude or more. There is a need for *in situ* measurement devices that do not hinder advective processes (e.g. DGT gels (Merritt and Amirbahman 2007)) and for the development of methods and measurements that work across ranges of biogeochemical conditions.

3. What is the relative importance of benthic biotransfer of mercury into aquatic food webs?

The role of benthic biota in transferring Hg to the higher trophic level fish and shellfish species consumed by humans is poorly understood. Existing work suggests that bioturbation of sediments by benthic infauna can affect methylation rates and distribution of MeHg in sediment (Benoit et al. 2006). Benthic fauna in Hg contaminated sediments have also been shown to exhibit higher Hg concentrations than those in more pristine sites, suggesting that biotic transfer from this food pathway may contribute to elevated total Hg levels in high trophic level organisms (Chen et al., unpublished data). However, organic content of sediments diminishes the bioavailability of MeHg to benthic fauna, which may result in lower levels of biotransfer from

highly organic-rich sediments (Lawrence and Mason 2001; Mason and Lawrence 1999). The factors controlling biotransfer of MeHg by benthic fauna need to be identified.

The relative importance of benthic fauna in biotransfer should also be more closely examined. Recent studies in freshwater and marine systems indicate that MeHg concentrations are higher in pelagic than benthic fauna, suggesting that chemical flux into the water column may be more important than biotransfer mechanisms (Chen et al., unpublished data; Gorski 2003; Power et al. 2002) but this has not been extensively investigated.

Monitoring Needs

To identify the important factors controlling methylation, monitoring data should be collected to characterize the spatial and vertical distribution of Hg and MeHg in ocean waters and sediments across a range of marine ecosystems. This range should include coastal margins, where riverine inputs of MeHg may be important, to the open ocean and the deep ocean, where sources of dimethylmercury are present.

Better analytical techniques are needed to improve detection limits of Hg and MeHg in marine waters given that levels in most of the world's oceans are difficult to measure. In addition, measures of ancillary variables in water and sediments are needed to identify the factors controlling Hg methylation and demethylation (e.g. selenium, iron, manganese, sulfide, dissolved organic carbon (DOC), pH, chloride, productivity, and nutrients).

Standardized measurements of methylation rates and MeHg flux from sediments across a range of ecosystem types would aid in validating existing MeHg model results and inform a better understanding of the magnitude of chemical flux of MeHg from sediments. These methylation rates and MeHg fluxes should be linked to measurements of Hg and MeHg in

benthic infauna and epifauna to quantify relative contributions of chemical and biotic flux of MeHg to the water column.

Theme 2: Trophic Transfer and Bioaccumulation of Methylmercury in Marine Food Webs

Trophic transfer and bioaccumulation of MeHg in marine food webs link MeHg production to MeHg exposure in humans and wildlife. Though trophic transfer and effects of MeHg in freshwater food webs have been well characterized in North America (Driscoll et al. 2007; Evers and Clair 2005; Watras et al. 1998), much more attention is needed on marine ecosystems. The work group identified three priority research questions: (1) what is the key entry point for MeHg in the base of the food web in marine ecosystems? (2) What are the factors influencing the transfer of MeHg from the base of the food web to higher trophic level organisms consumed by humans? And (3) what types of MeHg impacts have been measured in marine biota and which organisms could serve as useful indicators for monitoring MeHg spatiotemporal trends in marine ecosystems?

Research Questions

1. What is the key entry point for MeHg into the base of the food web in marine ecosystems?

Studies of inland aquatic ecosystems have found the greatest degree of MeHg bioaccumulation in the food web to occur between concentrations in water and concentrations in phytoplankton. For example the concentration of MeHg has been shown to increase by up to five orders of magnitude, with the percent of total Hg as MeHg increasing an average of 1% in water to 10% in phytoplankton (Driscoll et al. 2007; Fitzgerald et al. 2007).

Although estimates of bioconcentration exist for coastal waters, little is known of the bioconcentration by phytoplankton in the open ocean (Fitzgerald et al. 2007). In freshwater studies, MeHg concentrations in water do not consistently predict concentrations at the base of the food web. Limnological factors such as pH, DOC, and nutrients can have important effects on the bioaccumulation of MeHg by phytoplankton and zooplankton in these ecosystems (Chen et al. 2005; Driscoll et al. 2007; Pickhardt et al. 2002; Watras et al. 1998). In marine ecosystems, the chemical speciation of MeHg and its bioavailability is influenced greatly by the abundance of Cl and S, but less so by DOC and variation in pH (Lawson and Mason 1998). The presence of Hg or MeHg as inorganic or organic complexes determines the passive or active uptake by algal cells (Mason 2002). However, the degree to which quality and quantity of DOC on Hg bioaccumulation is poorly understood. More data are needed to characterize bioaccumulation processes in phytoplankton across a range of marine ecosystems.

2. What are the factors influencing the transfer of MeHg from the base of the food web to higher trophic level organisms consumed by humans?

Existing studies of freshwater and marine food webs show increasing MeHg concentrations with increasing trophic position as measured by stable isotopes (Bank et al. 2007; Driscoll et al. 2007; Hammerschmidt and Fitzgerald 2006). Both freshwater and marine fish also appear to have higher MeHg concentrations with increasing size and age (Hammerschmidt and Fitzgerald 2006; Wiener and Spry 1996). However, other aspects of trophic transfer of MeHg are far more difficult to track in marine food webs. Species-specific life history characteristics, migration patterns, ontogenetic shifts in diet, and differences in lifespan are poorly known for most marine species (Bank et al. 2007; Hammerschmidt and Fitzgerald 2006). In addition to the

need to characterize MeHg concentrations in a range of species across trophic levels, ages, and habitats, a better understanding of the general ecology of marine species is needed to properly interpret differences in MeHg burden between and within species across space and time.

Food web characteristics influence the transfer of MeHg from its sources to higher trophic levels. Humans consume fish from both demersal and pelagic fisheries, but little is known about the relative degree of MeHg bioaccumulation and trophic transfer in these two food webs. Some evidence suggests MeHg burdens in similar trophic level fish are higher in demersal versus pelagic species (Garcia-Hernandez et al. 2007). The influence of different food sources and food webs on MeHg bioaccumulation in fish species needs to be characterized, particularly for species most consumed by humans.

Food sources and food webs also influence the bioaccumulation of MeHg in apex predators such as marine mammals and birds. Marine mammals that have among the highest MeHg body burdens include toothed cetaceans and pinnipeds that feed on fish (Kemper et al. 1994; Thompson 1996; Wagemann et al. 1998). In contrast, baleen cetaceans have lower Hg levels, likely due to their diet of plankton (Hobson et al. 2004). Studies of seabirds suggest that habitat type and functional feeding group may influence MeHg bioaccumulation rates in higher trophic level organisms. For example, MeHg bioaccumulation rates differ between benthic and pelagic feeding birds and between inshore and offshore species (Goodale et al., unpublished data; Thompson et al. 1998). More data are needed to determine whether the influences of habitat type, feeding strategy and diet on MeHg bioaccumulation are consistent across a range of ocean ecosystems and taxonomic groups.

3. What types of MeHg impacts have been measured in marine biota and which organisms could serve as useful indicators for monitoring MeHg spatiotemporal trends in marine ecosystems?

Elevated environmental Hg concentrations have been widely documented in marine biota and extreme levels are regularly reported (Bustamante et al. 2003; Kim et al. 1996). The direct effects of elevated MeHg on marine biota can include impacts on neurological endpoints and memory, locomotion and cognition, as well as changes in brain neurochemical receptor density (Basu et al. 2005; Scheuhammer et al. in press). Adverse effects may further manifest as immunosuppression, which may make individuals more susceptible to disease, as has been measured in cetaceans and pinnipeds (Gauthier et al. 1998; Lalancette et al. 2003). Direct reproductive effects associated with high Hg levels have been documented in bird species in freshwater ecosystems (Burgess and Meyer 2008; Evers et al. 2008) and Braune et al. (2006) documented egg Hg concentrations in the ivory gull that exceed twice the adverse effect threshold for eggs in the common loon (Evers et al. 2003), suggesting that seabirds may be experiencing similar reproductive effects as freshwater birds. More research is needed to understand both the mechanisms and thresholds for adverse neurological, immunosuppressive, and reproductive effects in marine organisms.

The interpretation of Hg effects data in marine biota are complicated by the fact that sensitivity to MeHg toxicity can vary among taxa and foraging guilds. For example, Heinz et al. (in press) found eggs of marine birds dosed with MeHg to be less sensitive than those of terrestrial species. Moreover, information is needed about the interactions of MeHg with other chemicals and contaminants, including the ability of selenium to both reduce and enhance MeHg toxicity (G. Heinz, personal communication). Finally, in addition to the impacts of elevated

MeHg on individuals within a species, more research is needed relating Hg concentrations to population level effects in highly exposed species.

Monitoring Needs

Detailed information on MeHg concentrations in marine food webs across estuary, coastal and open ocean habitats is needed to better understand effects and to monitor changes in environmental MeHg loads over space and time. Emphasis should be placed on capturing a range of productivity from oligotrophic to eutrophic aquatic systems, and MeHg measurements should be conducted across a broad range of indicator taxa. Selected species should represent differing foraging guilds, habitats, and geographic areas and prioritized based on: (1) existing Hg data, (2) commercially harvested species for human consumption, (3) sensitivity to MeHg and (4) degree of conservation concern. Among chosen indicator taxa, emphasis should be placed on measurements of relevant tissue types to best relate MeHg concentrations to specific neurological, behavioral and reproductive effects in marine biota (Wolfe et al. 2007). Monitoring efforts should also include stable isotope measurements, such as delta ^{13}C and delta ^{15}N ratios, in lower and upper trophic level taxa in order to detect shifts in trophic structure and position (Hobson 1993; Hobson et al. 1994).

In order to capture ecologically meaningful changes in MeHg concentrations, monitoring should be conducted in both low and high trophic position organisms. To understand the entry of MeHg at the base of the food web, monitoring should include measurements of MeHg in phytoplankton, zooplankton, and benthic invertebrates. At the top of the food web, five broad groups of apex predators representing some of the highest MeHg concentrations in marine

organisms may be useful indicators: sharks, estuarine birds, seabirds, pinnipeds, and toothed whales.

Studies on these apex predator groups have been conducted in the North Temperate Atlantic Ocean (Spalding et al. 2007) suggesting that monitoring these taxa is feasible. But regional data for Hg bioaccumulation in apex species from some marine waters (e.g. the northeast US) remain limited for sharks, pinnipeds and toothed whales (Gaskin et al. 1973, 1979; Lake et al. 1995), estuarine birds (Cohen et al. 2000; Custer and Mulhern 1983; Rattner et al. 2000; Shriver et al. 2006), and seabirds (Burger 2002; Burger and Gochfeld 1995, 2003, 2004; Gochfeld 1980; Gochfeld and Burger 1998; Gochfeld et al. 1996; Goodale et al., unpublished data). Moreover, the great variability in units of measure, species chosen, age and sex class, and tissue type point to the need for more standardized Hg monitoring protocols for marine biota.

Theme 3. Human Exposure to Mercury from Marine Fish and Shellfish Consumption

Considerable research has been conducted on the human health implications of exposure to MeHg (Clarkson et al. 2003; Clarkson and Magos 2006; Grandjean et al. 2005; Mergler et al. 2007; NAS 2000). In the U.S., most people receive their highest mercury exposure through consumption of seafood (Sunderland 2007). But the relationship between the ecosystem fate of MeHg in freshwater and marine systems is poorly understood, and little is known about the effects of ecosystem variability on human exposure. In addition to Hg, seafood is a potential source of other contaminants, as well as a source of important nutrients such as omega-3 oils. There is a research need for well-defined and meaningful data supporting parameters of risk and benefit. Monitoring needs include increased tracking of the sources of the fish people eat. The work group identified three research questions: (1) what is the cumulative risk of methylmercury

and other co-contaminants in fish (PCBs, dioxin, etc.) and what are the tradeoffs in benefits from eating fish? (2) What are the patterns of mercury exposure and consumption for the most highly exposed human populations? And (3) what are peoples' responses to risk-benefit messages and how can they be improved?

Research Questions

1. What is the cumulative risk of MeHg and other co-contaminants in fish (PCBs, dioxin, etc.) and what are the tradeoffs in benefits from eating fish?

Research on fish contamination and human exposure is often framed in terms of a single contaminant. This approach may not be appropriately holistic for protecting human health. For public policies on fish to be protective of public health, there is a need to define an appropriate baseline for cumulative exposure to MeHg and other contaminants and a means of determining how regulation might alter the combined exposure levels. Developing species-specific pollution matrices in conjunction with fishery sustainability data may assist in identifying those species and fish populations that pose the greatest threat from both human and environmental health standpoints. These matrices will need to be sensitive to variation in species contamination across time and geographies.

In addition, there is a need for investigating the risks and benefits of fish consumption (Budtz-Jorgensen et al. 2007; Stern 2005) in studies that separate and clarify opposite impacts on health outcomes. This has been an area of growing scientific discussion. Studies such as Oken et al (2005) have found higher fish consumption in pregnancy to be associated with better infant cognition, and higher mercury levels to be associated with lower cognition. These findings and others (Domingo et al. 2007) have led many health professionals and organizations such as the

American Heart Association to recommend consumption of fish with high omega-3s and low MeHg. However, there is a need for a better understanding of the cumulative effects of the other nutrients and contaminants in fish. Investigating these effects, independently or interactively, in species most commonly preferred by humans is a priority. Research on human health and fish consumption needs to reflect the variability of individual exposure over time and space.

2. What are the patterns of mercury exposure and consumption for the most highly exposed human populations?

There are numerous factors that affect MeHg exposure levels in individuals or subpopulations, such as ethnic differences in fish consumption preferences, diet, genetic differences, age, and uptake-excretion variation (Canuel et al. 2006). To date, analysis of the NHANES dataset shows Asians, Native Americans, and Pacific Island populations have higher Hg concentrations in their blood (Hightower et al. 2006). In addition, studies show coastal populations having higher Hg concentrations than inland populations (Crépet et al. 2005; Denger et al. 1994; McKelvey et al. 2007). It is important to develop effective tools for capturing these variations and their effects, both for bolus and general exposures. There is also a need for a clinical definition for the subtle symptoms of concern associated with fish consumption resulting in chronic low-level exposures to MeHg that do not fit the current definition for “mercury poisoning” (Hightower and Moore 2003).

Detailed information is needed about the consumption patterns, fish and shellfish species preferences, and regional sources of fish for those who have been shown to carry the highest levels of MeHg in their bodies. In addition, for these highly susceptible individuals, the per-capita exposure should be scaled to the species of fish consumed. Individual variation in per-

capita mercury intake suggests that geography counts where mercury exposure is concerned (Burger et al. 2005). To control exposure, there is a need for research that identifies whether geographic supply regions, fishing methods, and country of origin affect variability in Hg concentrations in the fish available to consumers in different areas of the US.

3. What are peoples' responses to risk-benefit messages and how can they be improved?

Challenges still remain with regard to balancing risk-benefit messages regarding fish consumption and mercury, and there is a great need to study how people respond to different messages on fish consumption advisories, especially those intended for pregnant women (Knuth et al. 2003; Oken et al. 2003). Some investigators argue that fish consumption advisories are unbalanced in their focus on the health risks of fish consumption without informing the public about the health benefits of consuming appropriate amounts of low mercury fish (Arnold et al. 2005). However, some advocacy groups contend that consumption advisories do not go far enough in warning vulnerable populations about the risks of eating the most popular forms of fish. High-risk populations, who subsist heavily on fish and foods originating from the ocean, may require more intensive education efforts to ensure species-specific and local water body consumption advisory information is available (Arnold et al. 2005).

Monitoring Needs

Mercury monitoring efforts for human exposure should focus on highly exposed populations, commonly consumed fish and areas of high commercial seafood production. Commonly consumed fish may require a tracking system, similar to beef, to identify catch locations and sources of highly contaminated fish. Such a tracking system should include

information on catch location, distribution chain from source areas to consumers, type of fish and brand of the fish product, and assessments of MeHg levels in fish from different regions. There appear to be significant regional differences in mercury concentration by species that are not captured in the FDA's national database (Burger et al. 2005; Sunderland 2007). Commercial and non-commercial sources (i.e., recreational fish) will likely need to be evaluated separately.

Information on species-specific MeHg concentrations, frequency data, and amount of consumption will be required for human exposure studies to be informative (Burger et al. 2005; Mergler et al. 2007; Sunderland 2007). In addition, special monitoring may be required to evaluate MeHg levels in uncommon fish species that are regularly consumed by certain ethnic sectors of the public. In order to address the problem of multiple exposures to contaminants via fish consumption, measurement of other contaminants such organic (PCB's, dioxin, PAH's) could also be monitored in commercially harvested species as has been done in some existing monitoring programs (e.g. U.S. EPA National Coastal Assessment and Gulfwatch Programs).

Consumption data suggest that a common route of MeHg exposure for most Americans is canned tuna (Sunderland 2007). There is known variation between white (albacore tuna) and light (skipjack tuna) varieties, the former having MeHg concentrations more than 3 times higher than the latter (Burger and Gochfield 2004; Sunderland 2007). Better monitoring of the national distribution system for canned tuna is needed (type, brand, variation, region, etc.). Data on the sources, species, MeHg concentrations, and markets for canned tuna are necessary to evaluate an important source of MeHg exposure to humans.

Summary

Mercury research in marine ecosystems is a growing field that holds great promise for

improving our understanding of the critical linkages among the sources of methylmercury in marine systems, the processes that govern bioaccumulation in higher trophic level organisms, and human exposure to methylmercury through seafood consumption. In order to realize this potential, however, research and monitoring efforts must be coordinated in a way that helps answer targeted scientific and policy-relevant questions. Based on the deliberations of the 43 participants in the Dartmouth workshop, this report identifies several key research questions and recommended monitoring approaches that should guide future interdisciplinary work. We suggest that the study of mercury in marine ecosystems where humans have the greatest potential to be exposed is severely data-limited and that research and monitoring efforts should be expanded considerably. Moreover, we suggest that research and monitoring initiatives should take an integrated approach which addresses the poorly understood linkages among marine sources, biotransfer processes and bioaccumulation mechanisms that put humans at risk of exposure to MeHg. Within this integrated approach, there is an over-arching need to collect data and information across a range of habitats, chemical conditions, and ocean basins and to relate the resulting spatial patterns of MeHg bioaccumulation to the food sources of at-risk human and wildlife populations. Finally, we suggest that to advance these recommendations, an organized Hg monitoring effort in marine systems should be developed to characterize the spatial and temporal variability of MeHg in various compartments much like the proposed Hg monitoring network for freshwater and upland systems (Mason et al. 2005). Together, these recommendations will help elucidate the patterns and processes influencing the transfer of MeHg from marine sources to human exposure.

List of References

- Arnold SM, Lynn TV, Verbrugge LA, Middaugh JP. 2005. Human biomonitoring to optimize fish consumption advice: reducing uncertainty when evaluating benefits and risks. *American J of Public Health* 95:393–397.
- Baeyens W, Leermakers M, Papina T, Saprykin A, Brion N, Noyen J, et al. 2003. Bioconcentration and biomagnification of mercury and methylmercury in North Sea and Scheldt Estuary fish. *Arch. Environ Contam Toxicol* 45:498-508.
- Bank MS, Chesney E, Shine J, Maage A, Senn D. 2007. Mercury bioaccumulation and trophic transfer in sympatric snapper species from the Gulf of Mexico. *Ecol Applic* 17: 2100-2110.
- Basu N, Scheuhammer A, Grochowina N, Kelnavic K, Evans D, O'Brien M, et al. 2005. Effects of mercury on neurochemical receptors in wild river otters (*Lontra canadensis*). *Environ Sci Technol* 39:3585-3591.
- Benoit JM, Gilmour CC, Heyes A, Mason RP, and Miller CL. 2003. Geochemical and biological controls over methylmercury production and degradation in aquatic ecosystems. In: *Biogeochemistry of environmentally important trace elements*. American Chemical Society Publ, ACS symposium series 835: 262–297 pp.
- Benoit JM, Shull DH, Robinson P, Ucran LR. 2006. Infaunal burrow densities and sediment monomethyl mercury distributions in Boston Harbor, Massachusetts. *Mar Chem* doi:10.1016/j.marchem.2005.09.017.
- Braune BM, Mallory ML, Gilchrist HG. 2006. Elevated mercury levels in declining population of ivory gulls in the Canadian Arctic. *Marine Poll. Bull.* 52:969-87.
- Budtz-Jørgensen E, Grandjean P, Weihe P. 2007. Separation of risks and benefits of seafood intake. *Environ. Health Perspect* 115:323-327.
- Burger J. 2002. Food chain differences affect heavy metals in bird eggs in Barnegat Bay, New Jersey. *Environ Res* 90:33-39
- Burger J, Gochfeld M. 1995. Heavy metal and selenium concentrations in eggs of herring gulls (*Larus argentatus*): temporal differences from 1989 to 1994. *Arch Environ Contam Toxicol* 29:192-197.
- Burger J, Gochfeld M. 2003. Spatial and temporal patterns in metal levels in eggs of common terns (*Sterna hirundo*) in New Jersey. *Sci Total Environ* 311:91-100
- Burger J, Gochfeld M. 2004. Mercury in canned tuna: white vs. light and temporal variation. *Environ Res* 96:239-249.

- Burger J, Stern AH, Gochfeld M. 2005. Mercury in commercial fish: optimizing individual choices to reduce risk. *Environ Health Perspect* 113:266-271.
- Burgess NM, Meyer MW. 2008. Methylmercury exposure associated with reduced productivity in common loons. *Ecotoxicol* 17:83-91.
- Bustamante P, Garrigue C, Breau L, Caurant F, Dabin W, Greaves J, et al. 2003. Trace elements in two odontocete species (*Kogia breviceps* and *Globicephala macrorhynchus*) stranded in New Caledonia (South Pacific). *Environ Pollut* 24:263-271.
- Canuel R, de Grosbois SB, Atikesse L, Lucotte M, Arp P, Ritchie C, et al. 2006. New evidence on variations of human body burden of methylmercury from fish consumption. *Environ Health Perspect* 114:302-306.
- Chen CY, Stemberger RS, Kamman NC, Mayes BM, Folt CL. 2005. Patterns of Hg bioaccumulation and transfer in aquatic food webs across multi-lake studies in the Northeast US. *Ecotoxicol* 14: 135–147.
- Choe K-Y, Gill GA, Lehman RD, Han S, Heim WA, Coale KH. 2004. Sediment–water exchange of total mercury and monomethyl mercury in the San Francisco Bay–Delta. *Limnol Oceanogr* 49:1512–1527.
- Clarkson TW, Magos L, Myers GJ. 2003. The toxicology of mercury—current exposures and clinical manifestations. *New England J Medicine* 349:1731–1737.
- Clarkson TW, Magos L. 2006. The toxicology of mercury and its chemical compounds. *Crit Rev Toxicol* 36:609-62.
- Cohen JB, Barclay JS, Major AR, Fisher JP. 2000. Wintering greater scaup as bioindicators of metal concentrations in federal refuges in the Long Island region. *Arch Environ Contam Toxicol* 38:83-92.
- Cossa D, Coquery M. 2005. The Mediterranean mercury anomaly: A geochemical or a biochemical issue? In: *Handbook of Environmental Chemistry*, (Salot, A, ed.) Review Series in Chemistry, Vol 5. Springer-Verlag.
- Cossa D, Martin JM, Takayanagi K, Sanjuan J. 1997. The distribution and cycling of mercury species in the western Mediterranean. *Deep-Sea Res II* 44:721-740.
- Covelli S, Faganeli J, Horvat M, Brambati A. 1999. Porewater distribution and benthic flux measurements of mercury and methylmercury in the Gulf of Trieste (Northern Adriatic Sea). *Est Coast Shelf Sci* 48:415-428.
- Crépet A, Tressou J, Verger P, Leblanc JC. 2005. Management Options to Reduce Exposure to Methylmercury through the Consumption of Fish and Fishery Products by the French Population. *Regulatory Toxicol Pharmacol* 42:179-189.

- Custer TW, Mulhern BL. 1983. Heavy metal residues in prefledging black-crowned night-heron from three Atlantic coast colonies. *Bull Environ Contam Toxicol* 30:178-185.
- Denger R, Adams C, Moss S, Mack S. 1994. *Per capita fish and shellfish consumption in Florida*. Gainesville, FL: Florida Agriculture Market Research Center. University of Florida.
- Domingo JL, Bocio A, Falco G, Llobet JM. 2007. Benefits and risks of fish consumption Part I. A quantitative analysis of the intake of omega-3 fatty acids and chemical contaminants. *Toxicol* 230:219-226.
- Driscoll CT, Han YJ, Chen CY, Evers DC, Lambert KF, Holsen TM, et al. 2007. Mercury contamination in remote forest and aquatic ecosystems in the northeastern U.S.: Sources, transformations and management options. *Bioscience* 57:17-28.
- Evers DC, Clair TA. 2005. Mercury in northeastern North America: A synthesis of existing databases. *Ecotoxicol* 14:7-14.
- Evers DC, Taylor KM, Major A, Taylor RJ, Poppenga RH, Scheuhammer AM. 2003. Common Loon eggs as indicators of methylmercury availability in North America. *Ecotoxicol* 12:69-81.
- Evers DC, Savoy L, DeSorbo CR, Yates D, Hanson W, Taylor KM, et al. 2008. Adverse effects from environmental mercury loads on breeding common loons. *Ecotoxicol* 17:69-81.
- Fitzgerald WF, Clarkson TW. 1991. Mercury and monomethylmercury: Present and future concerns. *Environ Health Perspect* 96:159-166.
- Fitzgerald WF, Engstrom DR, Mason RP, Nater EA. 1998. The case for atmospheric mercury contamination in remote areas. *Environ Science Technol* 32:1-7.
- Fitzgerald WF, Lamborg CH, Hammerschmidt CR. 2007. Marine biogeochemical cycling of mercury. *Chem Rev* 107:641-662.
- Garcia-Hernandez J, Cadena-Cardenas L, Betancourt-Lozano M, Garcia-de-la-Parra LM, Garcia-Rico L, Marquez-Farias F. 2007. Total mercury content found in edible tissues of top predator fish from the Gulf of California, Mexico. *Toxicol Environ Chem* 89:507-522.
- Gaskin DE, Frank R, Holdrinet M, Ishida K, Walton CJ, Smith M. 1973. Mercury, DDT, and PCB in harbour seals (*Phoca vitulina*) from the Bay of Fundy and Gulf of Maine. *J Fish Res Board Can* 30:471-475.
- Gaskin, DE, Stonefield KI, Suda P, Franck R. 1979. Changes in mercury levels in harbor porpoises from the Bay of Fundy, Canada, and adjacent waters during 1969-1977. *Arch Environ Contam Toxicol* 8:733-762.

- Gauthier JM, Dubeau H, Rassart E. 1998. Mercury-induced micronuclei in skin fibroblasts of beluga whales. *Environ Toxicol Chem* 17:2487-2493.
- Gill GA, Bloom NS, Cappellino S, Driscoll CT, Mason R, Rudd JWM. 1999. Sediment–water fluxes of mercury in Lavaca Bay, Texas. *Environ Sci Technol* 33:663–669.
- Gochfeld M. 1980. Tissue distribution of mercury in normal and abnormal young common terns. *Mar Pollut Bull* 11:362-377.
- Gochfeld, M, Burger J. 1998. Temporal trends in metal levels in eggs of endangered roseate tern (*Sterna dougallii*) in New York. *Environ Res* 77:36-42.
- Gochfeld M, Belant JL, Shukla T, Benson T, Burger J. 1996. Heavy metals in laughing gulls: gender, age and tissue differences. *Environ Toxicol Chem* 15:2275-2283.
- Gorski P R, Cleckner LB, Hurley JP, Sierszen ME, Armstrong DE. 2003. Factors affecting enhanced mercury bioaccumulation in inland lakes of Isle Royale National Park, USA. *Sci Total Environ* 304:327-348.
- Grandjean P, Cordier S, Kjellström T, Weihe P, Budtz-Jørgensen E. 2005. Health effects and risk assessments. In: *Dynamics of Mercury Pollution on Regional and Global Scales: Atmospheric Processes and Human Exposures around the World* (Pirrone N, Mahaffey KR, eds). Norwell, MA:Springer, 499–523.
- Hammerschmidt CR, Fitzgerald WF. 2004. Geochemical controls on the production and distribution of methylmercury in near-shore marine sediments. *Environ Sci Technol* 38:1487–1495.
- Hammerschmidt CR, Fitzgerald WF. 2006. Bioaccumulation and trophic transfer of methylmercury in Long Island Sound. *Arch Environ Contam Toxicol* 51:416-424.
- Harris R, Krabbenhoft DP, Mason RP, Murray MW, Reash R and Saltman T, eds. 2007. *Ecosystem Responses to Mercury Contamination: Indicators of Change*. SETAC, CRC Press, Boca Raton.
- Heinz GH, Hoffman DJ, Kilmstra, JD, Stebbins KR, Kondrad SL, Erwin CA. In press. Species differences in the sensitivity of avian embryos to methylmercury. *Arch Environ Contam Toxicol*.
- Hightower, JM, Moore. 2003. Mercury levels in high-end consumers of fish. *Environ Health Perspect* 111:604-608.
- Hightower JM, O’Hare A, Hernandez GT. 2006. Blood mercury reporting in NHANES: identifying Asian, Pacific Islander, Native American, and multiracial groups. *Environ Health Perspect* 114:173-175.

- Hobson KA. 1993. Trophic relationships among high Arctic seabirds: insights from tissue-dependent stable-isotope models. *Mar Ecol Prog Ser* 95:7-16.
- Hobson KA, Piatt JF, Pitocchelli J. 1994. Using stable isotopes to determine seabird trophic relationships. *J Anim Ecol* 63:786-798.
- Hobson KA, Riget FF, Outridge PM, Dietz R, Born E. 2004. Baleen as a biomonitor of mercury content and dietary history of North Atlantic minke whales (*Balaenoptera acutorostrata*): combining elemental and stable isotope approaches. *Sci Total Environ* 331:69-82.
- Horvat M, Kotnik J, Logar M, Fajon V, Zvonaric T, Pirrone N. 2003. Speciation of mercury in surface and deep-sea waters in the Mediterranean Sea. *Atmos Environ* 37:S93-S108 Suppl. 1.
- Kemper C, Gibbs P, Obendorf D, Marvanek S, Lenghaus C. 1994. A review of heavy metal and organochlorines levels in marine mammals in Australia. *Sci Total Environ* 154:129-139.
- Kim EY, Murakami T, Saeki K, Tatsukawa R. 1996. Mercury levels and its chemical form in tissues and organs of seabirds. *Arch Environ Contam Toxicol* 30:259-266.
- Knap A, Dewailly E, Furgal C, Galvin J, Baden D, Bowen R, et al. 2002. Indicators of ocean health and human health: Developing a research and monitoring framework. *Environ Health Perspect* 110:839-845.
- Knuth BA, Connelly NA, Sheeshka J, Patterson J. 2003. Weighing health benefit and health risk information when consuming sport-caught fish. *Risk Analysis* 23:1185-1197.
- Kraepiel AML, Keller K, Chin HB, Malcolm EG, Morel FMM. 2003. Sources and variations of mercury in tuna. *Environ Sci Technol* 37:5551-5558.
- Lake CA, Lake JL, Haebler R, McKinney R, Boothman WS, Sadove SS. 1995. Contaminant levels in harbor seals from the northeastern United States. *Environ Contam Toxicol* 29:128-134.
- Lalancette A, Morin Y, Measures L, Fournier M. 2003. Contrasting changes of sensitivity by lymphocytes and neutrophils to mercury in developing grey seals. *Develop Comp Immunol* 27:735-747.
- Laurier FJG, Mason RP, Gill GA, Whalin L. 2004. Mercury distributions in the North Pacific Ocean – 20 years of observations. *Mar Chem* 90:3-19.
- Lawrence AL, Mason RP. 2001. Factors controlling the bioaccumulation of mercury and methylmercury by the estuarine amphipod *Leptocheirus plumulosus*. *Environ Poll* 111:217-231.

- Lawrence AL, Mcaloon KM, Mason RP, Mayer LM. 1999. Intestinal solubilization of particle-associated organic and inorganic mercury as a measure of bioavailability in benthic invertebrates. *Environ Sci Technol* 33:1871-1876.
- Lawson NM, Mason RP. 1998. Accumulation of mercury in estuarine food chains. *Biogeochem.*40:235-247.
- Mahaffey K R, Mergler D. 1998. Blood levels of total and organic mercury in residents of the Upper St. Lawrence River Basin, Quebec: Association with age, gender, and fish consumption. *Environ Res* 77:104-114.
- Mason RP. 2002. The bioaccumulation of mercury, methylmercury and other toxic elements into pelagic and benthic organisms. In: *Coastal and Estuarine Risk Assessment* (Newman MC, Roberts MH, Hale RC, eds). Boca Raton:Lewis Publishers, 127-149.
- Mason RP, Abbott ML, Bodaly RA, Bullock OR, Driscoll CT, Evers D, et al. 2005. Monitoring the environmental response to changing atmospheric mercury deposition. *Environ Sci Technol* 39:14A-22A.
- Mason RP, Benoit JM. 2003. Organo-mercury compounds in the environment. In: *Organometallics in the Environment* (Craig P, ed). New York:John Wiley and Sons, 57-99.
- Mason RP, Fitzgerald WF. 1993. The distribution and biogeochemical cycling of mercury in the equatorial Pacific Ocean. *Deep-Sea Res.* 40:1897-1924.
- Mason RP, Gill GA. 2005. Mercury in the Marine Environment. In: *Mercury: Sources, Measurements, Cycles and Effects* (Parsons MB, Percival JB, eds). Mineralogical Association of Canada Short Course Series 34:179-216.
- Mason RP, Heyes D, Sveinsdottir A. 2006a. Methylmercury concentrations in fish from tidal waters of the Chesapeake Bay. *Arch Environ Contam Toxicol* 51:425-437.
- Mason RP, Kim E-H, Cornwell J, Heyes D. 2006b. The influence of sediment redox status on the flux of mercury, methylmercury and other constituents from estuarine sediment. *Mar Chem* 102:96-110
- Mason RP, Lawrence AL. 1999. Concentration, distribution, and bioavailability of mercury and methylmercury in sediments of Baltimore Harbor and Chesapeake Bay, Maryland, USA. *Environ Toxicol Chem* 18:2438-2447.
- Mason RP, Lawson NM, Lawrence AL, Leaner JJ, Lee JG, Sheu GR. 1999. Mercury in the Chesapeake Bay. *Mar Chem* 65:77-96.
- Mason RP, Rolfhus KR, Fitzgerald WF. 1995. Bioaccumulation of mercury and methylmercury. *Water Air Soil Poll.* 80:915-919.

- Mason RP, Rolfhus KR, Fitzgerald WF. 1998. Mercury in the North Atlantic. *Mar Chem* 61:37-53.
- Mason RP, Sullivan KA. 1999. The distribution and speciation of mercury in the South and equatorial Atlantic. *Deep-Sea Res II*, 46:937-956.
- McKelvey W, Gwynn RC, Jeffery N, Kass D, Thorpe LE, Garg RK, et al. 2007. *Environ Health Perspect* 115: 1435-1441.
- Mergler D, Anderson HA, Chan HM, Mahaffey KR, Murray M, Sakamoto M, et al. 2007. Methylmercury exposure and health effects in humans: A worldwide concern. *Ambio* 36:3-11.
- Merritt KA, Amirbahman A. 2007. Mercury mobilization in estuarine sediment porewaters: A diffusive gel time-series study. *Environ Sci Technol* 41:717-722.
- Morel FMM, Kraepiel AML, Amyot M. 1998. The chemical cycle and bioaccumulation of mercury. *Ann Rev Ecology System* 29:543-566.
- Myers RA, Baum JK, Shepherd TD, Powers SP, Peterson CH. 2007. Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science* 315:1846-1849
- National Academy of Sciences Committee on the Toxicological Effects of Methylmercury, Board on Environmental Studies and Toxicology, National Research Council. 2000. *Toxicological Effects of Methylmercury*. The National Academies Press.
- Oken E, Kleinman KP, Berland WE, Simon SR, Rich-Edwards JW, Gillman MW. 2003. Decline in fish consumption among pregnant women after a national mercury advisory. *Obstet Gynecol* 102:346-351.
- Oken E, Wright RO, Kleinman KP, Bellinger D, Amarasiriwardena CJ, Hu H, et al. 2005. Maternal fish consumption, hair mercury, and infant Cognition in a U.S. cohort. *Environ Health Perspect* 113:1376-1380.
- Pickhardt PC, Folt CL, Chen, CY, Klaue B, Blum JD. 2002. Algal blooms reduce the uptake of methylmercury in freshwater food webs. *Proc Nation Acad Sciences* 99:4419-4423.
- Power M, Klein GM, Guiguer K, Kwan MKH. 2002. Mercury accumulation in the fish community of a sub-Arctic lake in relation to trophic position and carbon sources. *J Applied Ecol* 39:819-830.
- Rattner BA, Hoffman DJ, Melancon MJ, Olsen GH, Schmidt SR, Parsons KC. 2000. Organochlorine and metal Contaminant exposure and effects in hatching Black-Crowned Night Herons (*Nycticorax nycticorax*) in Delaware Bay. *Arch Environ Contam Toxicol* 38:38-45.

- Riget F, Muir D, Kwan M, Savinova T, Nyman M, Woshner V, et al. 2004. Circumpolar pattern of mercury and cadmium in ringed seals. *Sci Total Environ* 351/352:312-322.
- Scheuhammer AM, Basu AM, Burgess N, Elliot NM, Campbell JE, Wayland, GD, et al. In press. Relationships among mercury, selenium, and neurochemical parameters in common loons (*Gavia immer*) and bald eagles (*Haliaeetus leucocephalus*). *Ecotoxicol.*
- Shriver G, Evers DC, Hodgman TP, MacCulloch BJ, Taylor RJ. 2006. Mercury in sharp-tailed sparrows breeding in coastal wetlands. *Environ. Bioindicators* 1:129-135.
- Slowey AJ, Brown GE. 2007. Transformations of mercury, iron, and sulfur during the reductive dissolution of iron oxyhydroxide by sulfide. *Geochimica et Cosmochimica Acta* 71:877-894.
- Spalding MD, Fox HE, Allen GR, Davidson N, Ferdana ZA, Finlayson M, et al. 2007. Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *BioScience* 57:573-583.
- Storelli MM, Zizzo N, Marcotrigiano GO. 1999. Heavy metals and methylmercury in tissues of Risso's dolphin (*Grampus griseus*) and Cuvier's beaked whale (*Ziphius cavirostris*) stranded in Italy (South Adriatic Sea). *Bull Environ Contam Toxicol* 63:703-710.
- Stern AH. 2005. A review of the studies of the cardiovascular health effects of methylmercury with consideration of their suitability for risk assessment. *Environ Res* 98:133-142.
- Strode SA, Jaegle L, Selin NE, Jacob DJ, Park RJ, Yantosca RM, et al. 2007. Air-sea exchange in global mercury cycle. *Global Biogeochem Cycles* 21:GB1017-
- Sunderland EM. 2007. Mercury exposure from domestic and imported estuarine and marine fish in the U.S. seafood market. *Environ Health Perspect* 115:235-242.
- Sunderland EM, Gobas FAPC, Branfireun BA, Heyes A. 2006. Environmental controls on the speciation and distribution of mercury in coastal sediments. *Mar Chem* 102:111-123.
- Sunderland EM, Mason RP. 2007. Human impacts on open ocean mercury concentrations, *Global Biogeochem. Cycles*, doi:10.1029/2006GB002876 [accepted 30 August 2007].
- Thompson, D. R. 1996. Mercury in Birds and Terrestrial Animals. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. (Beyer WN, Heinz GH, Redmon-Norwood AW, eds). Clemson, SC: Lewis Publisher 341-355.
- Thompson DR, Furness RW, Monteiro LR. 1998. Seabirds as biomonitors of mercury inputs to epipelagic and mesopelagic marine food chains. *Sci. Total Environ.* 213:299-305.
- Ullrich SM, Tanton TW, Abdrashitova SA. 2001. Mercury in the aquatic environment: A review of factors affecting methylation. *Critical Rev. Environ Science Technol* 31:241-293.

- United States Commission on Ocean Policy. 2004. Preliminary report available at <http://www.oceancommission.gov/documents/prelimreport/welcome.html#full>.
- Wagemann R, Trebacz E, Boila G, Lockhart WL. 1998. Methylmercury and total mercury in tissues of Arctic marine mammals. *Sci Total Environ* 218:19-31.
- Watras CJ, Back RC, Halvorsen S, Hudson RJ, Morrison KA, Wentz SP. 1998. Bioaccumulation of mercury in pelagic freshwater food webs, *Sci Total Environ* 219:183-208.
- Wiener J.G, Krabbenhoff D P, Heinz GH, Scheuhammer AM. 2003. Ecotoxicology of Mercury. In: *Handbook of Ecotoxicology*, 2nd ed. (Hoffman DJ, Rattner BA, Burton Jr GA, Cairns Jr J, eds). Boca Raton: CRC Press, 407–461.
- Wiener JG, Spry DJ. 1996. Toxicological significance of mercury in freshwater fish. In: *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. (Beyer WN, Heinz GH, Redmon-Norwood AW, eds) Boca Raton: Lewis, 297–340.
- Wolfe, M.F., Atkeson, T., Bowerman, W., Burger, K., Evers, D.C., Murray, et al. 2007. Wildlife Indicators. In: R. Harris, D.P. Krabbenhoft, R. Mason, M.W. Murray, R. Reash and T. Saltman, eds), *Ecosystem Response to Mercury Contamination: Indicators of Change* SETAC, Webster, NY: CRC Press, 123-189.

Table 1. Selected reviews of mercury biogeochemistry, trophic transfer, biomonitoring and human health.

<u>Topic</u>	<u>Key Reviews</u>
Mercury Biogeochemistry	Fitzgerald and Clarkson 1991 Fitzgerald et al. 1998 Morel et al. 1998 Ullrich et al. 2001 Mason and Benoit 2003 Fitzgerald et al. 2007
Mercury Trophic Transfer	Mason et al. 1995 Mason 2002 Wiener et al. 2003 Mason and Benoit 2003
Biomonitoring	Harris et al. 2007 Mason et al. 2005
Human Health	Clarkson et al. 2003 Grandjean et al. 2005 Stern 2005 Clarkson and Magos 2006 Mergler et al. 2007

Table 2. Range or mean mercury concentrations in subsurface water at different stations of the world ocean. HgT: total mercury; MeHg: methylated mercury including MeHg and Me₂Hg; pM: pica-moles. Table from: Mason, RP and Gill, GA. 2005. Mercury in the Marine Environment. In: Mercury: Sources, Measurements, Cycles and Effects (Parsons MB, Percival JB) Mineralogical Association of Canada Short Course Series 34:179-216.

Location	HgT (pM)	MeHg (pM)	%MeHg	Reference
South and Equatorial Atlantic	0.8 - 2.4	0.025 - .200	5-10	a
North Atlantic	2.4 ± 1.6	0.029 – 0.160	2-7	b
North Pacific	1.2 ± 0.9	<0.050	<4	c
Equatorial Pacific	0.5-4.0	0.035 – 0.670	2-15	d
Mediterranean	0.5 – 4.0	0.020 – 0.460	1 – 35	e-g

^aMason and Sullivan 1999; ^bMason et al. 1998; ^cLaurier et al. 2004; ^dMason and Fitzgerald 1993; ^eCossa et al. 1997; ^fHorvat et al. 2003; ^gCossa and Coquery 2005.