

Index

A

- AARP *See* American Association of Retired Persons
- Abdomen, 310
- Abdominal trauma, 281
- Abdominal aortic aneurysm, 14
- Abortion, spontaneous, 14, 272, 281–283, 307, 370
 abruptio placentae, 281
 environmental tobacco smoke, 16, 364, 368, 372
 infertility, 273, 276
 risk, 304
- Abruptio placentae, 14, 279–281, 307, 370
- Abstinence, 581, 606; *See also* Smoking cessation
 lung cancer risk, 200
 menstrual cycle, 559–560
- Academic performance, 66, 472
- Acculturation, 581–582
- ACS *See* American Cancer Society
- Action on Smoking and Health, 603
- Addiction *See* Nicotine dependence
- Adenocarcinoma, 201, 635
 cervical cancer, 220
 diet and lung cancer, 208
 esophageal cancer, 226
 incidence, 202
 oropharyngeal cancers, 225
 urinary tract cancers, 228
- Adenoma, colorectal, 226
- Adipose tissue, nicotine metabolism, 339
- Adolescents
 body weight, 308
 depression, 134, 333–334
 dysmenorrhea, 561
 lung growth, 370
 nicotine dependence, 334, 557
 promotions and smoking initiation, 504
 restrictive tobacco policies, 593, 606
 smoking cessation, 572–573, 606
 smoking initiation, 63, 82, 453–477
 smoking and lung development, 252, 261
 smoking prevalence, 142, 635
 tobacco advertisements, 511
 tobacco marketing campaigns, 500–505
- Adult Use of Tobacco Surveys, 23, 28, 29, 553
 cigar use, 116–117
 cigarette brand preference, 46
 environmental tobacco smoke, 122
 heavy smoking, 42
 nicotine replacement therapy, 98, 99
 physician advice about smoking, 112
 pipe use, 118–119
 smoking cessation, 96, 100, 106, 551
 smoking continuum, 107, 108
- Adults, older, smoking cessation, 486–487, 562
- Advertising, 490–511; *See* Marketing
 awareness, 504
 counteractivities, 603–605
 direct, 521–522
 gays and lesbians, 509
 indirect, 521
 power of, 497
 restrictions, 515, 596–597
- Advocacy, anti-tobacco, 634
 programs, 600–606
- Advocacy Institute, 603
- Affect, 569
 negative, 331, 482, 483
- AFL-CIO, 602
- Africa, smoking prevalence, 136
- African American men *See* Black men
- African American women *See* Black women
- Age
 assisted reproduction, 273
 atherosclerosis, 241
 breast cancer risk, 212
 chronic obstructive pulmonary disease prevalence, 257
 cigar use, 116
 cigarette brand preference, 47
 coronary heart disease, 232, 233, 369
 gestational *See* Gestational age
 maternal *See* Maternal age
 mortality risk, 184–186
 pregnancy and smoking cessation, 484
 smoking after pregnancy, 484
 smoking cessation, 485–487, 386–387
 smoking initiation, 74
 stages of smoking cessation, 110
- Age factors
 older smokers, 554
 smoking cessation, 553, 562
 smoking cessation during pregnancy, 579
- Ageing, smoking cessation, 486–487, 562, 583–584
- Air pollution, 208, 261
- Airflow, expiratory, 249
- Airflow obstruction, 249
- Airline flights, 602
- Alaska Native girls
 physician advice about smoking, 115
 smokeless tobacco, 121
 smoking prevalence, 61
- Alaska Native women, 142
 birth weight, 302
 cigar use, 117
 current smoking, 39
 environmental tobacco smoke, 123
 ever smoking, 33
 heavy smokers, 43–44, 45, 46
 smokeless tobacco, 119
 smoking cessation, 100, 101, 578, 582
 smoking during pregnancy, 73
 smoking prevalence, 11, 37, 40–41, 48, 635
 smoking prevention, 636

- Alaska Native young women, current smoking, 49
- Alcohol consumption, 129, 130, 484, 486
- binge drinking, 130
 - combined with tobacco, 224–225
 - esophageal cancer, 226
 - laryngeal cancer, 225–226
 - liver cancer, 227
 - oropharyngeal cancers, 224–225
 - ovarian cancer, 220
 - pancreatic cancer, 228
 - smoking cessation, 485
 - smoking during pregnancy, 484
 - spontaneous abortion, 281
- Alcohol dependence, 16, 335, 336, 371
- Alpha-tocopherol, 208
- Altitude, 281
- Alzheimer's disease, 16, 336–337, 372
- Amenorrhea, 266, 370
- American Association of Retired Persons study, smoking cessation attempts, 96
- American Cancer Society, 26, 184; *See* Cancer Prevention Study I; Cancer Prevention Study II
- community intervention programs, 589
- American Cancer Society survey
- ever smoking, 59
 - physician advice about smoking, 112, 115
 - smoking cessation, 95
- American College of Chest Physicians, 601
- American College of Obstetricians and Gynecologists, 601
- American Indian girls
- current smoking, 59
 - physician advice about smoking, 115
 - smokeless tobacco, 121
 - smoking intensity, 62
 - smoking prevalence, 61
- American Indian infants, sudden infant death syndrome, 306
- American Indian women *See Also* Canadian Indian women; Cherokee tribe; Hopi women; Lumbee tribe
- cigar use, 117
 - current smoking, 39
 - environmental tobacco smoke, 123
 - ever smoking, 33
 - heavy smokers, 43–44, 45, 46
 - smokeless tobacco, 119
 - smoking cessation, 100, 101, 578, 582
 - smoking cessation during pregnancy, 111
 - smoking during pregnancy, 73
 - smoking prevalence, 11, 37, 40–41, 48, 142, 635
 - smoking prevention, 636
- American Indian Women's Health Education Resource Center, 601
- American Indian young women, current smoking, 49
- American Lung Association, nicotine replacement therapy, 98, 99
- American Medical Women's Association, 601, 605
- American Nurses Association, 601
- American Public Health Association, 603, 605
- American Stop Smoking Intervention Study, 589
- American Tobacco Company, 493, 494, 495, 510
- Amniotic fluid, 278
- Ampulla of Vater, 228
- Anal atresia, 306
- Anaplastic thyroid cancer *See* Thyroid cancer
- Androgen, 217, 265, 311
- Androstenedione, 265
- Anencephaly *See* Neural tube defects
- Anger, 473
- Ankle brachial pressure index, 245
- Antidepressants, 556, 557, 569–570
- Antioxidants, cervical cancer, 223
- Antismoking campaign, effect on prevalence, 29
- Antismoking materials, 523
- Anxiety, 473
- Anxiety disorders, 16, 334–335, 336, 371
- Anxiolytics, 556, 557
- Aortic aneurysm, abdominal, 247, 248, 370
- Aortic atherosclerosis, 247
- Appetite, suppression, 475–476
- Arthritis, 15, 327–330, 371
- Asbestos, 207
- Ascorbic acid, maternal levels, 278
- Asia
- oral cancers, 225
 - smokeless tobacco use, 138
 - smoking prevalence, 136
- Asian or Pacific Islander girls
- physician advice about smoking, 115
 - smoking intensity, 62
 - smoking prevalence, 61
- Asian or Pacific Islander women, 44, 46
- current smoking, 39
 - environmental tobacco smoke, 123
 - ever smoking, 33
 - lung cancer incidence, 195
 - smokeless tobacco, 119
 - smoking cessation, 100, 583
 - smoking during pregnancy, 73
 - smoking prevalence, 11, 37, 41, 48, 142, 635
- Asian or Pacific Islander young women
- current smoking, 49
 - smoking prevalence, 50
- Association of Flight Attendants, 602
- Association of Professional Flight Attendants, 602
- Asthma, 249, 252, 258
- Athens, Greece, air pollution and lung cancer risk, 209
- Atherosclerosis *See* Carotid atherosclerosis
- Atherosclerosis Risk in Communities Study, 243
- Athletics, 600
- Atomic bomb survivors, lung cancer, 209
- Attention, 341, 560
- Attention deficit disorder, 15, 334–335, 336, 371
- Attitudes, 592, 594, 595
- Attorneys General settlement *See* Master Settlement Agreement

Australia
 chronic obstructive pulmonary disease mortality, 261
 community-based antismoking campaigns, 588
 lung cancer, 210–211
 smoking trends, 138
 Australian National Women's Magazine Project, 604
 Austria, lung cancer mortality, 211
 Auto racing, sponsorship, 514
 Autopsy, 350
 AUTS *See* Adult Use of Tobacco Surveys

B

Badge products, 492
 Bailancho Saad, 526
 Basic brand, advertising, 510
 Beaver Dam (Wisconsin) Eye Study, 330
 Behavioral control, smoking initiation among adolescents, 471
 Behavioral programs, 566–567, 570
 Behavioral Risk Factor Surveillance System, 23, 40–41, 42, 72
 Behavioral training, smoking cessation, 584
 Benign breast disease, 217
 Benson & Hedges brand, 46, 47
 advertising, 510, 521–522
 international sponsorship, 523
 Benzo[a]pyrene, 207
 Benzodiazepines, 556
 Bernays, Edward, 494, 495
 Beta-blockers, 556
 Beta-carotene, 208
 cervical cancer, 223
 Betelquid, 138
 Bidis, 121, 147
 Bile ducts, 228
 Biliary tract cancer, 228
 Billboards, 521
 Binge drinking, 130, 472; *See also* Alcohol consumption
 Biological factors, smoking initiation, 476–477
 Biopsychosocial factors, 552, 606
 smoking cessation, 17
 Birth cohort studies, 28–29
 Birth cohorts, 74–77
 and ever smoking by age, 77–81
 forced expiratory volume, 253
 initiation, 142
 lung cancer trends, 203
 smoking cessation, 105, 553
 smoking initiation, 74–77, 81, 499
 smoking initiation and coronary heart disease, 233
 Birth defects, 290, 303
 Birth outcomes, 290–306
 smoking cessation, 564
 Birth weight 15, 296, 302–303, 371; *See also* Intrauterine growth retardation
 environmental tobacco smoke, 357, 368, 372
 low *See* Low birth weight

preterm delivery, 291
 Bisexual women, current smoking, 42
 Bisexual youth, smoking prevalence, 59
 Black adolescents
 emotional distress, 473
 risk taking behavior, 472
 smoking initiation, 470, 472
 smoking prevalence, 635
 Black boys, smoking prevalence, 61
 Black children, maternal smoking and lung function, 251
 Black girls
 cigarette brand preference, 70
 current smoking, 57, 58
 ever smoking, 55, 59
 physician advice about smoking, 115
 smoking and body weight, 126
 smoking intensity, 62–63
 smoking prevalence, 7, 11–12, 57, 60, 61, 70, 142
 weight control, 476
 Black infants, sudden infant death syndrome, 306
 Black men
 chronic obstructive pulmonary disease mortality, 260
 ever smoking, 49
 lung cancer incidence, 194
 smoking cessation, 106
 smoking prevalence, 29
 Black women
 age at smoking initiation, 75, 76, 77, 80
 birth weight, 302
 body weight, 127
 bone fractures, 311
 breastfeeding, 484
 chronic obstructive pulmonary disease mortality, 260
 cigarette brand preference, 46
 coronary heart disease risk, 232
 current smoking, 39
 environmental tobacco smoke, 123
 ever smoking, 33, 78
 first cigarette of the day, 83
 forced expiratory volume impairment, 259
 heavy smokers, 43, 45, 46
 lung cancer, 194, 197, 200
 neonatal mortality, 294
 nicotine dependence, 88
 physician advice about smoking, 113
 preeclampsia risk, 288
 pregnancy and smoking cessation, 483
 relapse to smoking, 487
 smokeless tobacco, 119
 smoking cessation, 97, 100, 101, 106, 578, 579–580
 smoking cessation during pregnancy, 111, 563, 577, 579
 smoking cessation programs, 580
 smoking continuum, 107
 smoking during pregnancy, 71, 72, 73
 smoking prevalence, 11, 27, 28–29, 35, 48, 142
 stages of smoking cessation, 110

stillbirth, 291
Black young women
 current smoking, 49
 ever smoking, 49
 smoking cessation, 110
 smoking prevalence, 50
Blacks, television and smoking cessation, 577
Bladder cancer, 13, 228, 231, 369
Blood pressure *See* Hypertension
Blood vessel diseases, 232
Body fat distribution, 15, 310–311, 371
Body image, 125, 134
Body mass index, 329, 565; *See also* Body weight
 maternal, 291
Body weight, 15, 125–128, 134, 307–311, 371; *See also* Body mass index
 bone density, 315, 317
 endometrial cancer, 217
 maternal, 303
 nicotine effects, 341
 osteoarthritis, 329
 perceived, 126–128
 smoking initiation, 125
Bone density, 15, 311–318, 321, 371
Bone fractures, 318–321
 risk, 15, 321, 371
Boredom, and smoking, 134
Bowling, James, 498
Boys
 cigar use, 117
 current smoking prevalence, 60, 61
 depression, 341
 maternal smoking and lung function, 251
 religion and smoking, 67
 restrictions to access, 595
 senior class, 59
 smokeless tobacco, 121
 smoking initiation, 500
 smoking prevalence, 142
Brain, nicotine metabolism, 339
Brand stretching, 522
BRCA1 genes, 216
BRCA2 genes, 216
Breast cancer, 13, 212–217
 correlations with male lung cancer, 216
 environmental tobacco smoke, 216, 369
 risk, 213, 224, 369
 susceptibility, 634
Breast disease, benign, 217
Breastfeeding, 15, 306, 307, 371, 484, 488
 relapse to smoking, 487
 smoking cessation, 484, 577, 579
BRFSS *See* Behavioral Risk Factor Surveillance System
Brief interventions *See* Clinical interventions, minimal
Brill, A.A., 494
Bronchial hyperresponsiveness, 253
Bronchiectasis, 249
Bronchioles, 249
Bronchioloalveolar regions, carcinogens, 203

Bronchitis, chronic, 249, 250
Brown & Williamson, 506, 510, 511
Buccal mucosa, 225
Bulgaria, lung cancer mortality, 210–211
Bulimia nervosa, 16, 334–335, 336, 371
Bupropion, 556
 pharmacology, 557
 smoking cessation, 565, 570
 weight control, 569
Bureau of Indian Affairs, 56, 59
Bus stops, advertising, 521

C

Cadmium, 318
 neonatal mortality, 365
Caffeine, preterm delivery, 291
Calcaneus, 312
Calcitonin, 318
Calcium, 315, 318
California, tobacco control programs, 636
California Department of Health Services, 603, 604
California Environmental Protection Agency, lung cancer and environmental tobacco smoke, 344
California Tobacco Survey, 587
 promotions and smoking initiation, 504
 susceptibility to smoking, 475, 504
 tobacco product advertising, 455
California Youth Tobacco Survey, smoking initiation, 472
Camel brand, 46, 504
 advertising, 494, 496–497, 509, 510
 brand stretching, 522
 preference among girls, 69, 70
 promotions, 510
Canada
 chronic obstructive pulmonary disease mortality, 261
 lung cancer, 209–211
 smoking trends, 138
Canadian Association for the Advancement of Women and Sport and Physical Activity, 600
Canadian Indian women, smoking prevalence, 136
Canadian Network of Women Against Tobacco, 605
Cancer, 12–13, 193–231; *See also under* individual type
 predisposition to, 205–207
Cancer Prevention Study I, 26, 28, 37, 42
 age at smoking initiation, 75, 76
 aortic aneurysm mortality risk, 247
 chronic obstructive pulmonary disease mortality, 260
 coronary heart disease, 353
 death rates, 186–187
 lung cancer risk, 197, 200
 multiple myeloma, 231
 stroke mortality risk in former smokers, 239
Cancer Prevention Study II, 42
 age at smoking initiation, 75

- biliary tract cancer, 228
- bladder cancer mortality, 229
- breast cancer and environmental tobacco smoke, 216
- chronic obstructive pulmonary disease mortality, 260, 261
- coronary heart disease, 223, 233, 353
- death rates, 185–187
- esophageal cancer, 226
- laryngeal cancer, 226
- liver cancer mortality, 227
- low-tar cigarette preference, 48
- lung cancer, 200, 201
- lymphoid leukemias, 231
- multiple myeloma, 231
- non-Hodgkin's lymphoma, 231
- pancreatic cancer mortality, 228
- relative risk, 189–190
- risk and smoking cessation, 197–198
- smoking and breast cancer mortality, 215
- smoking risk in lung cancer, 197, 198
- spouse exposure to environmental tobacco smoke, 344
- stomach cancer, 226
- stroke risk, 240
- Cancer and Steroid Hormone Study, 213
 - ovarian study, 219–220
- Capri brand, 504, 508, 509
 - advertising restrictions, 525
 - appeal to women, 522
 - promotions, 510
- Carbon monoxide, 303, 340, 341, 573
 - congenital malformations, 304
 - fetal growth retardation, 364
 - neonatal mortality, 364
- Carboxyhemoglobin, 303
- Carcinogens, lung cancer, 203
- Carcinoma, adenosquamous, cervical cancer, 220
- Carcinoma, bronchioloalveolar, 201
- Carcinoma, large cell, 201, 208
- Carcinoma in situ, cervical, 220
- Carcinoma, small cell, 201, 202
 - diet and lung cancer, 208
- Carcinoma, squamous cell, 201, 202, 203
 - cervical cancer, 220
 - diet and lung cancer, 208
 - esophageal cancer, 226
- Cardiac birth defects, 304
- Cardiovascular defects, environmental tobacco smoke, 366
- Cardiovascular disease, 13–14, 232–238, 369
 - environmental tobacco smoke, 122, 123
 - mortality, 187
- Cardiovascular function markers, 356
- Caribbean Islands, smoking prevalence, 136
- Carlton brand, advertising, 509
- Carotenoids, 208, 223
- Carotid artery stenosis, 241–242
- Carotid atheromatous plaques, 242
- Carotid atherosclerosis, 14, 241–244, 248, 370
- Case-control studies
 - breast cancer, 212
 - coronary heart disease and environmental tobacco smoke, 353
 - environmental tobacco smoke, 345–350
 - ethnicity and lung cancer, 200
 - lung cancer, 198–200
- CASH *See* Cancer and Steroid Hormone Study
- Cataract, 15, 330, 331, 371
- Catecholamines, 265
- Cavalier brand, advertising, 497
- Censorship, 524
- Centers for Disease Control and Prevention
 - Office on Smoking and Health, 602
 - public health partnerships, 600
- Central American women, smoking during pregnancy, 73
- Central nervous system, congenital malformations, 303, 304
- Cerebral infarction *See* Stroke
- Cerebrovascular disease *See* Stroke
- Cervical cancer, 13, 220–224
 - risk, 224, 369
 - smoking and disease progression, 220
- Cervical dysplasia, 220
- Cervical epithelium, effects of smoking, 223
- Cervical incompetence, 278
- Cervical intraepithelial neoplasia, 220
- Cesarean section, placenta previa, 281
- Cessation *See* Smoking cessation
- Change, transtheoretical model, 474, 478
- Chelsea brand, 508
- Cherokee tribe, *See also* American Indian women
 - smokeless tobacco, 119
- Chesterfield brand, 46, 499, 500
 - advertising, 493, 494, 496, 498
- Children
 - cigarette advertising, 504
 - coronary heart disease and environmental tobacco smoke, 353, 355
 - environmental tobacco smoke, 345
 - forced expiratory volume, 251
 - nicotine dependence, 334
 - type 1 diabetes, 264
- China, 136, 138
 - air pollution and lung cancer risk, 208–209
 - environmental tobacco smoke, 345
- Chinese women, 41–42; *See also* Asian or Pacific Islander women
 - ethnicity and lung cancer, 200
 - lung cancer, 195, 211–212
 - smoking during pregnancy, 73
- Chinese young women, smoking prevalence, 50
- Cholecystectomy, 321
- Cholecystitis, 15, 323, 325, 371
- Cholelithiasis, 321, 323
- Cholesterol, 232, 247
 - lung cancer risk, 208

- Chromosomal abnormalities, 281, 283, 303
- Chronic obstructive lung disease *See* Chronic obstructive pulmonary disease
- Chronic obstructive pulmonary disease, 14, 249–262, 370
 - death rate, 250
 - diagnosis, 249
 - mortality, 259–261
 - prevalence, 257
 - risk, 250
 - risk and smoking intensity, 261
- Cigar use, 116
- Cigarette addiction *See* Nicotine dependence
- Cigarette advertisements, 490–511
- Cigarette brands
 - choices and advertising, 455
 - images, 490
 - loyalty, 491
 - preference, 46–48, 69–71
- Cigarette marketing *See* Marketing, 490–526
- Cigarette promotions *See* Promotions
- Cigarettes
 - brand stretching, 522
 - craving, 342
 - feminized, 522
 - filtered, 47, 99, 203, 554
 - first of the day, 82–85
 - generic, 47, 70
 - low-tar, 47, 116, 203
 - low-yield, 302
 - number smoked, 42, 482, 483; *See also* Smoking intensity
 - packaging, 492–493, 522
 - price, 454–455, 510, 594
 - reasons for use, 85–86
- Cigars, 12, 116–117, 142
 - marketing, 516–517
- Cimetidine, 324
- Circulatory anomalies, 303
- Civic improvement, tobacco sponsorship, 511
 - Clear Horizons*, 583, 584
 - Clearing the Air*, 584
- Cleft lip *See* Oral clefts
- Cleft palate *See* Oral clefts
- Clinical interventions, intensive, 555–556
 - pregnancy, 578
- Clinical interventions, minimal, 554–555, 556
 - pregnancy, 578
- Cliques, smoking initiation among adolescents, 469
- Clomiphene citrate challenge, 273
- Clonidine, 556, 557
- Cocaine use, 129, 130
- Cognitive readiness, 488
- Cohabitant *See* Partner
- Cohort studies
 - breast cancer, 212
 - coronary heart disease and environmental tobacco smoke, 351, 353
 - lung cancer, 196–198
 - lung cancer and environmental tobacco smoke, 344–345
 - mortality risk, 368
- Colitis, ulcerative, 15, 325, 371
- College students, 67, 565
- Colon cancer, 225, 226–227
- Colorectal cancer, 13, 226–227, 231, 369
- Commission for a Healthy New York, 602
- COMMIT *See* Community Intervention Trial for Smoking Cessation
- Community, Hypertension, Atherosclerosis, and Diabetes Program, 588
- Community Intervention Trial for Smoking Cessation, 558
 - brand preference, 47, 70
 - international programs, 588, 589
 - smoking after awakening, 83
 - smoking cessation, 482–483, 574
 - workplace policies, 587
- Community-based interventions, 581, 587–590, 598–599
- Compliance checks, 454
- Conception, delayed, 14, 15, 272–273, 276, 307, 371
 - environmental tobacco smoke, 16, 366, 368, 372
- Condylomata, 224
- Confidence, smoking cessation, 527
- Congenital malformations, 15, 303–306, 307, 371
 - environmental tobacco smoke, 366
- Conotruncal malformations, 304
- Consolidated Cigar Corporation, marketing, 516
- Contraception, spontaneous abortion, 281
- Copenhagen City Heart Study, 241
 - chronic obstructive pulmonary disease mortality, 261
- Coping skills, 474, 483
- Coronary Artery Surgery Study, 235
- Coronary heart disease, 232–238, 369
 - environmental tobacco smoke, 16, 343, 350–356, 372
 - hormone replacement therapy, 237–238
 - mortality, 233, 356
 - risk, 13, 232–235, 248, 353
- Coronary Risk Factor Study, 588
- Corpora lutea, 273
- Cortisol, 217, 265, 318
- Cotinine, 16, 83, 123, 372
 - birth weight, 302, 357, 362
 - blood pressure, 248
 - in cervical mucus, 223
 - fetal growth retardation, 364
 - infant lung development, 251
 - levels during pregnancy, 563
 - nicotine metabolism, 339
 - preterm delivery, 291
 - smoking intensity, 335
 - weight gain, 568
- Counteradvertising, 603

Covariate studies, 315
 CPS-I *See* Cancer Prevention Study I
 CPS-II *See* Cancer Prevention Study II
 Craniosynostosis, 304
 Craven "A" brand, 519
 Craving, cigarette, 342
 Crohn's disease, 15, 325, 371
 Cross-sectional studies
 bone density, 313–315
 smoking initiation among adolescents, 454
 Crow's foot area, 332
 Cuban American women, 39, 45
 cigarette brand preference, 47
 smoking during pregnancy, 73
 Cultural influences, 453
 Current Population Survey, 23, 26, 46
 age at smoking initiation, 75
 cigar use, 117
 smoking cessation, 101
 smoking initiation, 74, 80, 81, 82, 499
 smoking intensity, 44, 74
 smoking prevalence, 26, 42
 trends in smoking cessation, 100
 workplace tobacco smoke exposure, 123
 CYP1A1 genes, 216
 Cystic fibrosis, 249
 Czech Republic, smoking prevalence, 136

D

D₂ dopamine receptors, 476
d-Fenfluramine, weight control, 568–569
 Dakota brand
 advertising, 511
 counteradvertising, 603
 Dakota Papers, 603
 Death, premature, 12
 Death rate *See also* Mortality, 184–187
 lung cancer, 197
 smoking-specific, 184–186
 temporal trends, 186–189
 Deaths, smoking attributable, 190
 Deaths, premature, 7, 194, 468
 Decision making, 478
 Dehydroepiandrosterone, 262
 Delinquency, smoking initiation among adolescents, 472
 Delivery, preterm, 15, 278, 290–291, 307, 371
 abruptio placentae, 281
 Demand for cigarettes, pricing, 455
 Denmark
 cigar use, 138
 human papillomavirus infection, 223
 lung cancer, 210–211
 smoking prevalence, 136
 smoking trends, 138
 Dentists, advice about smoking, 555
 Department of Defense Survey of Health Related Behaviors Among Military Personnel, 39
 Depression, 15, 333–334, 336, 371, 473, 557
 adolescents, 476
 major depressive disorder, 569, 570
 premenstrual dysphoric disorder, 561
 smoking cessation, 486, 569–570, 571
 smoking risk, 334
 social support, 571
 Depression and smoking, 15, 130, 134
 Desogestrel, dose, 237
 Detroit, Michigan, environmental tobacco smoke exposure, 347
 Developed countries, 519
 deaths from smoking, 190
 lung cancer trends, 209–211
 smoking behavior, 135
 Developing countries
 lung cancer mortality, 211–212
 marketing, 520
 smoking behavior, 135
 smoking prevention, 637
 DHEAS, pregnant smokers, 262
 Diabetes mellitus
 14, 264, 370; *See also* Gestational diabetes
 preeclampsia, 288
 smoking intensity, 264
Diagnostic and Statistical Manual of Mental Disorders, 82, 86, 334–335, 561
 Diet, lung cancer, 208
 Dieting, 125, 565
 Diffusion of innovations model, 471
 Digestive malformations, 305–306
 Disease outcomes, 634, 636
 Displays, point-of-sale, 521
 Distal influences, 453
 District of Columbia Interagency Council on Smoking and Health, 604
 DNAdamage, human papillomavirus infection, 223
 Doctors Ought to Care, 604
 Dopamine transporter gene, 476
 Doral brand, advertising, 510
 Dose-response relationship *See also* Smoking intensity
 carcinoma, 202
 carotid atherosclerosis, 241
 coronary heart disease and environmental tobacco smoke, 353
 depression, 333
 emphysema, 250
 limb reductions, 305
 low birth weight, 302
 lung cancer, 197
 menopause, 269
 neonatal mortality, 295
 paternal smoking and congenital malformations, 366
 preterm delivery, 290–291
 smoking cessation and weight gain, 341–342

smoking intensity and stroke, 240
spontaneous abortion, 281–283
stomach cancer, 226
sudden infant death syndrome, 307
tar and lung cancer, 200
urinary tract cancers, 229
urinary tract malformations, 305
Down syndrome, 305
DRD4 gene, 334
Drug use
 intravenous, abruptio placentae, 281
 smoking, 129–130, 134
 smoking initiation, 130
DSM *See Diagnostic and Statistical Manual of Mental Disorders*
Dual photon absorptiometry, 312
Dual X-ray absorptiometry, 312
Duodenal ulcer *See Peptic ulcer*
Duodenum, 324
Dysmenorrhea, 14, 266, 272, 370, 561

E

Eastern Europe, smoking prevalence, 136
Eclampsia, 283, 288
Ectopic pregnancy, 14
Edema, 283
Edinburgh Artery Study, 245
Education
 plans after graduation, 66, 71
 tobacco control advocacy programs, 601–602
Education level, 575–576
 current smoking, 48
 environmental tobacco smoke, 123
 ever smoking, 33
 heavy smoking, 46
 maternal smoking, 73
 parental, 64
 physician advice about smoking, 113
 pregnancy and smoking cessation, 484, 577, 579
 smoking after pregnancy, 484
 smoking bans, 123
 smoking cessation, 101, 110, 483, 486, 488, 527
 smoking cessation during pregnancy, 111
 smoking prevalence, 11, 37, 42, 142
 smoking prevalence among young women, 50, 50–51
 stages of smoking cessation, 110
 television and smoking cessation, 577
 workplace exposure, 124
Embassy brand, 519
Emotions, 134, 473
Emphysema, 249; *See also* Chronic obstructive pulmonary disease
Emphysema Slims tennis tournaments, 604
Employment
 pregnancy, 288

 smoking cessation, 527
Employment status, smoking cessation, 101, 102
Encephalocele *See Neural tube defects*
Endometrial cancer, 13, 217–218, 369
 risk, 224
Endometrial hyperplasia, 218
Endometriosis, 269–271
England, lung cancer, 210
Entertainment, sponsorship, 492
Environmental exposure *See Environmental tobacco smoke*
Environmental factors, smoking initiation, 454–470
Environmental Protection Agency, lung carcinogens, 344
Environmental tobacco smoke, 7, 14, 122–125, 343–368, 487, 592–593, 636
 breast cancer, 13, 216, 224, 369
 cervical cancer, 224
 childhood lung function, 14
 children, 251–252, 345
 conception delay, 368
 coronary heart disease, 16, 350–356, 372
 developing countries, 637
 exposure indicators, 345
 exposure levels, 345
 home exposure, 122–123
 infant pulmonary function, 250
 lung cancer, 16, 372
 lung development, 250, 370
 maternal exposure and neonatal mortality, 364–365
 perinatal effects, 357–366
 reproductive outcomes, 16, 356–358, 372
 smoking cessation, 484
 sudden infant death syndrome, 306
 in utero exposure, 217, 261
 workplace, 123–125, 350
Esophageal cancer, 13, 226, 231, 369
Essence (magazine), 604
Established Populations for Epidemiologic Studies of the Elderly, 223, 486
Estradiol, 262
 breast cancer risk, 212
 cancer, 217
 infertility, 273
Estriol, 217
Estrogen, 212, 230; *See also* Hormone replacement therapy
 bone density, 315
 dose and stroke risk, 241
 endometrial cancer, 218
 excretion, 262
 exogenous *See* Hormone replacement therapy
 metabolism, 262, 370
 oral, 237
Estrogen replacement therapy *See* Hormone replacement therapy
Estrogen-deficiency disorders, 14, 265, 370
Estrogen-dependent disorders, 14, 265, 270, 370
Estrogen-receptor tumors, 213–214
Estrogens, 14
Estrone, 212, 262

- Ethinyl estradiol, 562
 dose, 236, 237
- Ethnicity
 age at smoking initiation, 76
 cigarette brand preference, 70
 lung cancer, 200
 smoking prevalence during pregnancy, 73
- Europe
 environmental tobacco smoke exposure, 348
 esophageal cancer incidence, 226
 lung cancer rates, 210
 smoking prevalence, 136
- European Union 1998 Directive, advertising restrictions, 525
- Eve brand, 506
 product placement, 523
- Ever smoked, 142
 by age, 77–81
 breast cancer risk, 216
 trends, 29–34
 young women, 48
- Exercise, 315
 smoking cessation, 584
- Exercise programs, 567
- Experimental smoking, 474
- Expiratory flow rates, 251
- Extroversion, smoking cessation, 482
- Eye disease, 15, 330–331, 371
- F**
-
- Facial wrinkling, 15, 332, 371
- Fagerström Tolerance Questionnaire, 574
- Fallopian tubes, 273, 276
- Family conflict, smoking initiation, 455, 468
- Family history, 205–207
- Fashion, cigarette advertising, 497
- Fat, dietary, lung cancer risk, 208
- Fat distribution *See* Body fat distribution
- Fatalism, smoking initiation among adolescents, 471–472
- Fecundity, environmental tobacco smoke, 366–368
- Federation Internationale de Football Association (FIFA), 600
- Female cancers, 13, 212–224, 369; *See also* Breast cancer; Cervical cancer; Endometrial cancer; Ovarian cancer; Vulvar cancer
- Fertility
 environmental tobacco smoke, 366–368
 smoking cessation, 552
- Fertilization rates, 273
- Fertilization, in vitro, 273
- Fetal growth, 303, 357
- Fetal hypoxia, 303
 congenital malformations, 304
 nicotine, 364
- Fetal hypoxia-ischemia, 294
- Fetal loss, 291
 environmental tobacco smoke, 357, 364–365
- Fetal malpresentation, placenta previa, 281
- Fetal morbidity and mortality *See* Neonatal morbidity; Neonatal mortality
- Fetal nutrition, 303
- Fetuses, multiple, 288
- FEV *See* Forced expiratory volume
- Fibroadenoma, 217
- Fibrocystic disease, 217
- Fighting, 472
- Filipino women *See also* Asian or Pacific Islander women
 lung cancer, 195, 200
 smoking during pregnancy, 73
- Filipino young women, smoking prevalence, 50
- Film, 521
- Finland
 lung cancer, 210–211
 smoking trends, 138
- Flair brand, 506
- Flight attendant associations, 602
- Flight Attendant Non-Smokers, 602
- Florida, environmental tobacco smoke, 345
- Fluoxetine hydrochloride, weight control, 568
- Folic acid, 303
- Follicular atresia, 269
- Follicular fluid, 273
- Follicular phase
 smoking cessation, 560
 withdrawal symptoms, 559
- Follicular thyroid cancer *See* Thyroid cancer
- Forced expiratory volume, 249, 252, 253
 smoking cessation, 253, 257
- Forced vital capacity, 249, 252
- Forearm, 312
- Fortuna brand, brand stretching, 522
- Fortune Magazine* survey, 23, 26
- Four A's protocol, 579–580
- Framework Convention for Tobacco Control, 637
- Framingham Study, 26
 bone density, 315
 carotid atherosclerosis, 241
 gallbladder disease, 323
 menopause, 268
 osteoarthritis, 329
 peripheral vascular disease risk, 245
 smoking cessation, 103
 type 2 diabetes, 264
- France
 lung cancer, 209, 210–211
 smoking prevalence, 136
 smoking trends, 138
- Freedom from Smoking for You and Your Family*, 566
- Friends *See* Peers
- Frontier Menthol Slims brand, advertising restrictions, 525
- FVC *See* Forced vital capacity

G

- Gallbladder, 228
- Gallbladder disease, 321–324, 325, 371
- Gallstones, 15, 321, 325, 371
- Gallup Poll, 26
- Gamete interaction, 273
- Gastric mucosa, 324
- Gastric ulcer *See* Peptic ulcer
- Gastrointestinal abnormalities, 303, 305–306
- Gastrointestinal diseases, 15, 321–327, 371
- Gastroschisis, 306
- Gauloises Blondes brand, direct advertising, 521
- Gay men
- HIV disease, 331
 - targeted advertising, 509
- Gender
- acculturation, 581–582
 - adolescent smoking, 66
 - Alzheimer's disease, 337
 - attitudes towards smoking, 475
 - chronic obstructive pulmonary disease, 249, 250
 - cigarette advertising, 509
 - depression, 371
 - depression and adolescents, 333, 334
 - dieting among college students, 565
 - esophageal cancer incidence, 226
 - first cigarette of the day, 83
 - inflammatory bowel disease, 325
 - intensive clinical interventions, 555–556
 - laryngeal cancer, 225
 - lung cancer, 201, 203, 209
 - lung function, 253
 - nicotine dependence, 88, 91, 142, 342
 - nicotine intake, 372
 - nicotine pharmacology, 339, 476
 - nicotine replacement therapy, 556
 - physician advice about smoking, 112
 - reasons to quit smoking, 572
 - relapse to smoking, 107
 - restrictions on youth access, 595
 - risk for urinary tract cancers, 228
 - schizophrenia, 335
 - smokeless tobacco, 120
 - smoking cessation, 102–106, 482, 483, 550–552
 - smoking cessation among adolescents, 111
 - smoking cessation among older smokers, 583–584
 - smoking cessation methods, 553, 554, 588, 606
 - smoking and disease outcomes, 634
 - smoking initiation among adolescents, 454
 - smoking prevalence, 11–12, 142
 - social support, 571–572
 - stages of smoking cessation, 110
 - taxation policies, 594
 - television and smoking cessation, 576
 - tobacco-related policies, 591–592
 - in utero lung development, 251
 - weight gain, 565
- Gender susceptibility, lung cancer, 201
- Genetic polymorphisms, breast cancer risk, 216
- Genetic predisposition, lung cancer, 205
- Genetics, 476
- Mendelian inheritance, 206
 - psychiatric disorders, 336
- Genital warts, 224
- Genitourinary defects, environmental tobacco smoke, 366
- Germany
- environmental tobacco smoke exposure, 349
 - lung cancer mortality, 210–211
- Gestational age, 290
- birth weight and environmental tobacco smoke, 357
 - preterm delivery, 291
- Gestational age, small, 15, 290, 296, 303, 307, 370
- Gestational diabetes, 14, 265, 370; *See also* Diabetes mellitus
- Gestational hypertension, 283
- Gestodene, dose, 237
- Gingival recession, 116
- Girl Scouts of the USA, 601
- Girls, *See also under* entries at Alaska Native, American Indian, Asian or Pacific Islander, Black, Hispanic, Latin American, Mexican American, Navajo, and White
- attitudes about smoking, 67–69
 - cigar use, 117
 - cigarette brand preference, 69–71
 - cigarette smoking, 52–70
 - current or ever smoking, 59
 - current smoking, 56–61, 59–61
 - depression, 333
 - emotions and smoking, 134
 - ever smoking, 54, 55
 - first cigarette of the day, 83–85
 - lung function, 261, 370
 - lung growth, 14
 - nicotine dependence, 82, 88
 - other drug use, 129–131, 134
 - physician advice about smoking, 115
 - pregnant, 71–73, 111
 - psychosocial problems and smoking, 134
 - reasons for cigarette use, 86
 - restrictions to access, 595
 - senior class, 59–61
 - smokeless tobacco, 121
 - smoking cessation, 17, 111, 115
 - smoking initiation, 16, 500, 527
 - smoking intensity, 62–63, 70, 82
 - smoking and lung function, 14
 - smoking prevalence, 7, 11–12, 70, 142
 - weight, 125–126, 527
- Girls Incorporated of Alameda County, 601
- Girls organizations, tobacco control advocacy programs, 601
- Glaucoma, 15, 331, 371
- Glucose regulation, 14, 265, 370

Glucose tolerance, 265
 Goiter, 230, 263
 Golden Handcuffs Challenge, 604
 Golden Tobacco Company, targeting women, 526
 Gonadotropins, 269, 273
 Graves' disease, 263
 with ophthalmopathy, 263, 265, 370
 Graves' ophthalmopathy, 14
 Great Britain *See* United Kingdom
 Greece
 air pollution and lung cancer risk, 209
 lung cancer mortality, 210–211
 smoking prevalence, 136
 smoking trends, 138
 GSTM1 genes, 216
 Gudakhu, 138
 Gynecologic conditions, benign, 14, 266–272, 370

H

Harbin, China, environmental tobacco smoke exposure, 345
 Harvard Six Cities Study, air pollution and chronic obstructive pulmonary disease, 261
 Hawaiian women *See also* Asian or Pacific Islander women
 lung cancer, 195, 200
 smoking during pregnancy, 73
 Health consequences of smoking, 12, 475, 633, 635
 Health consequences of smoking for women, 549
 Health motives, smoking cessation, 550
 Health Promotion and Disease Prevention Supplement, 553
 Health promotion strategies, 17
 Health Resources and Services Administration, Maternal and Child Health Bureau, 602
 Healthy Worker Project, 586
 Heart, nicotine metabolism, 339
 Heart defects, 304
 conotruncal, environmental tobacco smoke, 366
 Heart disease *See* Cardiovascular disease
 Heel, 312
Helicobacter pylori, 325
 Hematologic cancers, 13, 231, 369
 Hemoglobin, fetal growth retardation, 364
 Hemoglobin A_{1c}, 265
 Hemorrhage
 intracerebral, 240
 postpartum, 281
 Hepatitis B infection, liver cancer, 227
 Hepatocellular carcinoma, 227
 Herstories, 602
 HHANES *See* Hispanic Health and Nutrition Examination Survey
 High school drop outs, 66
 Hill, George Washington, 493
 Hip, 312
 osteoarthritis, 329
 Hip fracture, 15, 371
 mortality, 311
 risk, 318–319, 321
 Hispanic adolescents, risk taking behavior, 472
 Hispanic girls
 cigarette brand preference, 70
 current smoking, 56
 ever try smoking, 55, 56
 physician advice about smoking, 115
 smoking and body weight, 126
 smoking intensity, 62–63
 smoking prevalence, 61
 Hispanic Health and Nutrition Examination Survey, 229
 cigarette preference, 47
 environmental tobacco smoke, 122
 smoking prevalence, 39
 Hispanic men
 smoking cessation, 106
 smoking prevalence, 29
 Hispanic women
 age at smoking initiation, 77, 80
 body weight, 127
 cigarette preference, 47
 current smoking, 39
 environmental tobacco smoke, 122, 123
 ever smoking, 33
 heavy smokers, 45, 46
 lung cancer, 195, 200
 nicotine dependence, 88
 physician advice about smoking, 113
 smokeless tobacco, 119
 smoking cessation, 97, 100, 101, 106, 578, 581–582
 smoking continuum, 107
 smoking during pregnancy, 71, 73
 smoking prevalence, 11, 27, 28–29, 37, 48, 71, 142, 635
 Hispanic young men, ever smoking, 49
 Hispanic young women
 current smoking, 49
 ever smoking, 49
 smoking cessation, 110
 smoking prevalence, 50
 HIV-1 infection, 15
 HMO study, lung cancer risk, 201
 Hodgkin's disease, mortality, 231
 Home exposure
 environmental tobacco smoke, 122–123, 353
 smoking bans, 123
 Homosexual men
 HIV-1 disease, 331
 targeted advertising, 509
 Hopelessness, 473
 Hopi women *See also* American Indian women
 smoking prevalence, 40
 Hormone replacement therapy, 15, 217, 218; *See also* Estrogen
 breast cancer risk, 213
 coronary heart disease, 237–238, 369
 ovarian cancer, 220

sex hormones, 262
smoking cessation, 562
stroke, 241, 248, 370
stroke risk, 14
thyroid cancer risk, 230
Hormones, smoking cessation, 558
Hostility, 473
Household structure, 71
 and smoking, 63
HPV *See* Human papillomavirus infection
Human chorionic gonadotropin hormone, 281
Human immunodeficiency virus, 15, 331, 371
Human papillomavirus infection, 13, 223, 369
 cervical cancer, 220
 vulvar cancer, 224
Humerus, fractures, 319
Hungary, lung cancer, 210–211
25-Hydroxy vitamin D₃, 318
2-Hydroxyestradiol, 262
2-Hydroxyestrones, 217
Hyperinsulinemia, 265
Hypertension, 232, 240, 247–248
 abruptio placentae, 281
 pregnancy, 283, 288–290
Hyperthyroidism, 263
Hypothalamic-pituitary axis, effects of smoking, 561
Hypothyroidism, 263
Hysterectomy, 269

I

Image, 475
Imipramine hydrochloride, 569
Immunosuppression, human papillomavirus, 220
IMPACT, 589
Implantation rates, 273
Income
 adolescent personal, 66
 family, 64
 smoking prevalence, 42
Income level, 71; *See also* Poverty
 self-help programs, 554
 smoking cessation, 101, 563, 579
 smoking prevalence, 38
 television and smoking cessation, 577
Income status, ever smoking, 33
India
 community-based interventions, 589
 oral cancers, 225
 smokeless tobacco use, 139
Indian Health Clinics, smoking cessation project, 582
Indoor air laws, 592
 cigarette demand, 455
Industrialized countries *See* Developed countries
Infection, during pregnancy, 278
Infertility, 14, 15, 273–276, 307, 370, 371
 endometrial cancer, 217
Inflammatory bowel disease, 325

Inhalant use, 130
Initiation *See* Smoking initiation
Initiatives to Mobilize for the Prevention and Control of Tobacco Use, 589
Insulin, 265
Integumental anomalies, 303
Intelligence, smoking initiation among adolescents, 472
Intention to smoke, 474
Intermittent claudication, 244
International Classification of Diseases, ninth revision, 249
International Conference on Tobacco and Health, 605
International Conference on Women and Smoking, 605
International Network of Women Against Tobacco, 526, 602, 605
International Union Against Cancer, 605
Internet marketing, 516
Interventions
 brief *See* Clinical interventions, minimal
 population-based, 558
Intracerebral hemorrhage, 369
Intrauterine devices, 277
Intrauterine growth retardation, 16, 290, 362–364, 372, 564
 environmental tobacco smoke, 368
INWAT *See* International Network of Women Against Tobacco
Iowa Women's Health Study, pancreatic cancer mortality, 228
Ireland, *See also* United Kingdom
 lung cancer, 210–211
 pancreatic cancer, 228
 smoking trends, 138
Irritability, 473, 551
Ischemic heart disease, fatal, 240
Ischemic stroke *See* Stroke
Italy
 esophageal cancer, 226
 lung cancer, 211
 smoking prevalence, 136

J

Japan
 biliary tract cancer, 228
 bladder cancer, 229
 lung cancer, 211
 pancreatic cancer, 228
 smoking prevalence, 136
 smoking trends, 138
Japan Tobacco International, 523
Japanese men, arsenic in drinking water, 208
Japanese women, *See also* Asian or Pacific Islander women
 arsenic in drinking water, 208
 lung cancer, 195
 pregnancy, 73
 smoking prevalence, 41–42
Japanese young women, smoking prevalence, 50

Jasira Warriors, 601
 JCAHO *See* Joint Commission on the Accreditation of Health Care Organizations
 Job stress, 486
 smoking after pregnancy, 484
 Joe Camel, 500, 510
 recognition among children, 504
 Joint Commission on the Accreditation of Health Care Organizations, 592
 Josephine Camel, 510

K

Kaiser Permanente Medical Care Program cohort study, 188–189, 261
 Kaiser Permanente Medical Health Care Program Study, lung cancer risk, 197, 198
 Kent brand
 advertising, 498
 brand stretching, 522
 Khaini, 138
 KICKIT!, 580
 Kidney, nicotine metabolism, 339
 Kidney cancer, 13, 231, 369
 mortality, 229
 Kidney malformations, 305
 Kim brand
 appeal to women, 522
 sponsorship, 511
 Knee, osteoarthritis, 329
 Kobe Declaration, 637
 Kool brand, 46, 47, 504
 sponsorships, 511
 Korea
 smoking prevalence, 136
 smoking trends, 138
 Korean women *See also* Asian or Pacific Islander women
 lung cancer, 195
 smoking prevalence, 41
 Korean young women, smoking prevalence, 50
 Kreteks, 116, 122, 148
 Kreyberg type II carcinoma *See* Carcinoma, bronchioloalveolar

L

L&M brand
 advertising, 497, 498
 appeal to women, 522
 Lactation, 306
 Langerhans' cells, 223
 Lark brand
 advertising, 498
 product placement, 523
 Laryngeal cancer, 13, 225–226, 231, 369
 Lasker, A.D., 494
 Latin America
 smoking and cervical cancer, 223

 smoking prevalence, 136
 Latin American girls
 smoking intensity, 62
 smoking prevalence, 61
 Latin American Women Association on Smoking Control, 605
 Learning, nicotine effects, 341
 Leiomyomas *See* Uterine fibroids, 271
 Leisure World Cohort Study, 233
 Lesbian women
 current smoking, 42, 51
 smoking prevalence, 59, 635, 636
 targeted advertising, 509
 Leukemia, acute myeloid, 13, 231, 369
 Leukoplakia, 116
 Life expectancy, 368
 U.S. women and smoking, 193
 Life Skills Training Project, 598
 Liggett & Myers, 494, 498, 506
 product placement, 523
 Lilith Fair concerts, 603
 Limb-reduction defects, 305
 environmental tobacco smoke, 366
 Linked Telephone Survey, smoking cessation during pregnancy, 111
 Lipid Research Clinics Follow-up Study, coronary heart disease, 232
 Lipoprotein lipase, 311
 Liver, nicotine metabolism, 339
 Liver cancer, 13, 227–228, 231, 369
 Locus of control, smoking initiation among adolescents, 471–472
 Loneliness, and smoking, 134
 Longitudinal studies, 315
 Lorillard, 506, 508
 sponsorship, 511
 Low birth weight, 15, 290, 296, 362–364, 371, 564
 environmental tobacco smoke, 362–364
 smoking cessation, 564
 Lucky Strike brand, 46, 499, 500
 advertising, 493, 494, 495
 Lumbee tribe, *See also* American Indian women
 smokeless tobacco, 119
 smoking cessation, 582
 Lung, nicotine pharmacology, 339
 Lung cancer, 7, 13, 193–212, 369; *See also* Adenocarcinoma; Carcinoma, large cell; Carcinoma, small cell; Carcinoma, squamous cell
 differences by gender, 200–201
 environmental tobacco smoke, 16, 122, 123, 207, 343–350, 372
 epidemiologic studies, 344–350
 gender, 194, 209
 genetic predisposition, 205–207
 international trends, 13, 209–212, 369
 mortality *See* Lung cancer mortality
 survival rates, 195
 Lung cancer histology, dose-response rate relative risk, 202

Lung cancer mortality, 197, 209–211, 369
 age-standardized rates, 211
 developing countries, 211–212
 international, 212
 in men, 211
Lung cancer risk, 196–205, 209
 case-control studies, 198–200
 duration of smoking, 198
 nonsmokers and environmental tobacco smoke, 343
 risk factors, 207–209
Lung function, 14, 250–257, 370
 decline in, 252
 smoking cessation, 14
 in utero exposure to maternal smoking, 261
Lung Health Study, 253, 555, 575, 583
 smoking cessation among heavy smokers, 575
 weight gain, 309–310
Lung parenchyma, 249
Lupus *See* Systemic lupus erythematosus
Luteal phase
 smoking cessation, 560
 withdrawal symptoms, 559
Lymphoma, non-Hodgkin's, 231
Lymphoproliferative cancers, 13, 231, 369

M

Macroalbuminuria, 264
Macular degeneration, 15, 330–331, 371
Magazines
 advertising, 505–510, 517–519, 521
 counteradvertising, 603–604
Major depressive disorder *See* Depression
Mania, 131
Manitoba Health Services Commission, 258
Maori women *See also* Asian or Pacific Islander women
 smoking prevalence, 136
Marijuana, 129, 130, 472
Marital status, smoking after pregnancy, 485
Marketing, 490–521; *See also* Advertising
 international, 519–526
 restrictions, 515, 524–525, 596–597
 targeting of women, 526, 527
 strategies, 520–524
Marlboro brand, 46, 47, 500, 504
 advertising, 493, 510–511
 brand stretching, 522
 international advertising, 520
 international sponsorship, 523
 nitrosamine content, 205
 preference among girls, 69, 70
 product placement, 523
 promotions, 510
 sponsorship, 511, 514
Mass media
 counteradvertising, 603
 smoking cessation, 576–577
Massachusetts Tobacco Survey, 587, 594
Master Settlement Agreement, 515–516, 597
Maternal age
 abruptio placentae, 281
 birth weight, 302
 congenital malformations, 303
 placenta previa, 281
 preterm delivery, 290
 stillbirth, 291
Maternal and Child Health Block Grant programs, 602
Maternal and Child Health Bureau, Health Resources and Services Administration, 602
Maternal smoking
 birth outcomes, 290–306
 childhood lung function, 251–252
 diabetes in children, 264
 infant lung function, 14, 261
 smoking cessation, 551
 smoking initiation among adolescents, 468–469
 sudden infant death syndrome, 306
 in utero lung development, 250–251, 370
Maudsley Personality Inventory, 482
Mecamylamine, 556, 557
Media
 advertising, 521
 self-censorship, 517–519
 tobacco advertising, 527
Media censorship, 524
Medullary thyroid cancer *See* Thyroid cancer
Membrane rupture, premature, 20
Membranes, preterm premature rupture, 278–279, 307, 370, 371
Memory, nicotine effects, 341
Memphis Health Project, 454
Men, *See also under* Asian Pacific Islander, Black, Hispanic, and White
 body weight, 127
 bone density, 315
 cigar use, 117
 diabetes risk, 265
 gallstone, 321
 HIV-1 infection, 331
 international lung cancer mortality, 211
 lung cancer, 201, 216, 343, 369
 nicotine, 339
 nicotine nasal spray, 340
 physician advice about smoking, 112–113
 smoking cessation, 102–106, 486
 vertebral fractures, 319
Menopausal hormones *See* Hormone replacement therapy
Menopause, 14, 268–269, 272, 370, 371
 bone density, 315
 hormone replacement therapy, 262
 perimenopausal symptoms, 561
 smoking cessation, 552, 562
Menopause status
 benign breast disease, 217

- bone density, 313–314
- breast cancer, 213, 216
- endometrial cancer, 217, 224
- hormone levels, 262
- ovarian cancer, 220
- sex hormones, 262
- vertebral fractures, 319
- Menses, 560
- Menstrual cycle, 552
 - irregularity, 266, 269, 370
 - nicotine, 558, 559
 - smoking cessation, 558–561
 - withdrawal and abstinence, 559–560
- Menstrual disorders, smoking and endometrial cancer, 217
- Menstrual function, 14, 266–268, 272, 370
- Mental health, and smoking, 131–135
- Mental illness, smoking prevalence, 333
- Merit brand, advertising, 509
- Meta-analyses
 - coronary heart disease studies, 356
 - lung cancer and environmental tobacco smoke, 344
- Methacholine, 253
- Methodologic issues, smoking studies, 52
- 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone, 204
- Mexican American girls, smoking intensity, 62
- Mexican American women
 - age at smoking initiation, 77
 - birth weight, 302
 - body mass index, 128
 - cigarette brand preference, 47
 - environmental tobacco smoke, 122
 - ever smoking, 33
 - heavy smokers, 45
 - pregnancy, 73
 - smoking prevalence, 29, 39
- Mexican Americans, smoking cessation, 106
- Mexican girls, smoking prevalence, 61
- Microalbuminuria, 264
- Middle East, smoking prevalence, 136
- Midwestern Prevention Project, 598, 599
- Mila Schön brand, promotions, 522
- Military women, 39
 - smokeless tobacco, 120
 - smoking and body weight, 128
- Milwaukee Journal* surveys, 26, 42
- Mineral supplements, 291
- Minimal contact *See* Clinical interventions, minimal
- Minnesota Heart Health Program, 599
 - coronary heart disease, 588–589
 - gender and smoking cessation, 554
 - health behavior, 473
 - smoking prevention, 599
 - smoking reduction, 99
- Minnesota Heart Survey, black women, 579
- Minnesota Nursing Network for Tobacco Control, 601
- Minnesota Smoking Prevention Program, 598
- Minority women, smoking cessation, 578–584, 606
- Miscarriage *See* Abortion, spontaneous
- Mishri, 138
- Misty brand
 - advertising, 509
 - promotions, 510
- Mobilization, tobacco control advocacy programs, 601–602
- Molecular genetics, depression, 334
- Monitoring the Future Surveys, 23, 52, 54, 57
 - academic performance, 66
 - attitudes about smoking, 67, 68
 - cigarette brand preference, 69, 70
 - current smoking among girls, 58
 - current smoking prevalence, 60, 61
 - daily smoking, 60
 - household structure, 63
 - other drug use, 130
 - parental education and smoking initiation, 471
 - personal income, 66
 - religion, 66
 - smokeless tobacco, 121
 - smoking cessation, 95, 111
- Mood, 135
 - adolescents, 476
 - dysphoric, 333
 - menstrual cycle, 560
 - negative, 473
 - nicotine, 341
- Mood disorders, behavioral interventions, 570
- Morbidity
 - coronary heart disease, 356
 - fetal, 290
 - neonatal, 290
- More brand, 506
 - appeal to women, 522
 - sponsorship, 511
- Mormon women, cervical cancer, 223
- Mortality, 12, 13, 183–193, 368; *See also* Death rate; Neonatal mortality; Perinatal mortality
 - Alzheimer's disease, 338
 - chronic obstructive pulmonary disease, 14, 250
 - coronary heart disease, 233, 351, 356
 - fetal, 290
 - Parkinson's disease, 336
 - peptic ulcers, 324
 - perinatal, 15, 370
 - smoking intensity, 188–189
 - smoking-related, 193
 - temporal trends, 186–189
- Mothers, self-help programs, 554
- Motor performance
 - menstrual cycle, 560
 - nicotine effects, 341
- Mouth cancer, 224–225
- Ms. brand
 - appeal to women, 522
 - targeting women, 526
- MTF *See* Monitoring the Future Surveys
- Multiparity *See* Parity

Multiple births, stillbirth risk, 291
Muscle, skeletal, nicotine metabolism, 339
Musculoskeletal anomalies, 303
Myeloid leukemia, 13
Myeloma, multiple, 231
Myocardial infarction, 232
 oral contraceptives, 237

N

N-acetyltransferase 2 genes, 216
N^o-nitrosoanabasine, 204–205
N^o-nitrosoanatabine, 204–205
N^o-nitrosonornicotine, 203–205
Nasal spray, nicotine, 340
Nass, 138
Naswar, 138
NAT1 genes, 216
NAT2 genes, 216
National Association of African Americans for Positive Imagery, 602
National Association of Flight Attendants, 602
National Association of Public Health Policy, 601
National Cancer Institute, 193
 community intervention programs, 589
 Four A's protocol, 579–580
 public health partnerships, 600
National Comorbidity Survey, smoking prevalence, 333
National Health Interview Surveys, 23, 257
 age at smoking initiation, 223
 Cancer Control Supplement, 550
 risk taking behaviors, 472
 smoking initiation, 499
 tobacco product pricing, 455
National Health and Nutrition Examination Survey Epidemiologic Followup Study, 324, 486
National Health and Nutrition Examination Survey I, weight gain, 309, 310
National Health and Nutrition Examination Survey III, 23
 body mass index, 128
 environmental tobacco smoke, 123
 smoking and body weight, 128
 workplace exposure, 124
National Hospital Ambulatory Medical Care Survey, 257
National Hospital Discharge Survey, 257, 258
National Household Surveys on Drug Abuse, 23, 46, 52
 binge drinking, 130
 cigar use, 117
 current smoking, 39
 current smoking among girls, 56–63, 57, 58
 ever smoked, 33, 53, 54, 595
 ever smoking by age, 78
 frequent smoking, 62
 high school drop outs, 66
 nicotine dependence, 86, 88, 89, 91, 338, 556–557
 other drug use, 129–130, 130
 smokeless tobacco use, 119
 smoking among high school senior girls, 59
 smoking cessation, 101
 smoking cessation among young women, 110
 smoking during pregnancy, 73
 smoking prevalence among young women, 50
 young women, 49
National Household Surveys on Drug Abuse, cigarette brand preference, 69
National Institute for Occupational Safety and Health, 123
 environmental tobacco smoke and lung cancer, 343
National Longitudinal Study of Adolescent Health, 334
National Maternal and Infant Health Survey, 71, 111
National Natality Survey, 71
National Organization of School Nurses, 601
National Organization for Women, 601, 602, 603, 605
National Park Service, 604
National Pregnancy and Health Survey, 71
National Smoking Cessation Campaign for African American Women, 602
National Survey of Family Growth, 71, 272
National Survey of Health and Development, 482
National Teenage Tobacco Surveys, 23, 52, 54, 59
National Vital Statistics Systems, 257
National Youth Risk Behavior Survey *See* Youth Risk Behavior Survey
National Youth Tobacco Survey, 23, 52, 54–56, 67, 68, 69
 bidis, 121
 cigars, 117–118
 physician advice about smoking, 115
 pipes, 119
 smokeless tobacco, 121
 smoking cessation, 95
Native American *See under* entries at American Indian
Nauru, *See also* Asian or Pacific Islander women type 2 diabetes, 264
Navajo girls, *See also* American Indian girls
 current smoking, 59
 smokeless tobacco, 121
Navajo students, smoking prevention, 598
Navajo women, *See also* American Indian women, 40
Neonatal morbidity, 290
Neonatal mortality, 290, 294–296; *See also* Perinatal mortality
 environmental tobacco smoke, 364–365, 372
Neoplasia, cervical intraepithelial, 220
 environmental tobacco smoke, 224
Nervousness, 473
Netherlands
 lung cancer mortality, 210–211
 smoking trends, 138
Neural tube defects, 303, 304
 environmental tobacco smoke, 366
Neurologic diseases, 15, 336–338, 372
Neuroticism, 482
Neurotransmitters, 570
Never active smokers, breast cancer risk, 213, 216
New Mexico, air pollution and lung cancer risk, 208
New Zealand
 lung cancer trends, 210–211

- smoking prevalence, 136
 - smoking trends, 138
 - New Zealand, Toxic Substances Board, 525
 - Newport brand, 46, 504
 - advertising, 498
 - preference among girls, 69, 70
 - sponsorship, 511
 - NHANES *See* National Health and Nutrition Examination Survey
 - NHIS *See* National Health Interview Surveys
 - NHSDA *See* National Household Surveys on Drug Abuse
 - Nicotine, 83, 273, 303
 - antidepressive effects, 333
 - birth weight, 364
 - breast fluid, 212
 - cervical mucus, 223
 - clearance, 339
 - congenital malformations, 304
 - effect on sex hormones, 269
 - hair samples, 364
 - hormones, 558
 - inflammatory bowel disease, 325
 - low-nicotine content, 506
 - osteoblasts, 318
 - pharmacology, 338–342, 372
 - physical dependence, 342
 - psychoactive effects, 340–342
 - reduced content, 97
 - self-medication, 333
 - stillbirth, 294
 - tolerance, 342
 - transdermal system, 342
 - Nicotine dependence, 12, 16, 17, 82–93, 142, 334, 372
 - anxiety disorders, 334
 - gender, 91, 606
 - heavy smoking, 574
 - indicators, 86, 89
 - pharmacologic processes, 16, 478
 - pharmacotherapy, 556–558
 - smoking cessation, 552
 - teens, 74
 - treatment methods, 552–558
 - Nicotine gum, 342, 556, 584
 - weight control, 566, 567–568
 - Nicotine inhalers, 556
 - weight control, 568
 - Nicotine intake, 342
 - Nicotine metabolism, 552
 - gender differences, 476
 - Nicotine nasal spray, 340, 556
 - Nicotine patch, 556, 557
 - pregnancy, 557
 - weight control, 568
 - Nicotine replacement therapy, 98, 555, 556–557
 - depression, 333
 - gender, 342
 - weight control, 567–568
 - Nicotine withdrawal, 335, 338, 342
 - symptoms in girls, 93
 - NIOSH *See* National Institute for Occupational Safety and Health
 - Nonsmokers
 - chronic obstructive pulmonary disease, 259
 - congenital malformations, 366
 - coronary heart disease, 351, 356, 372
 - environmental tobacco smoke, 123
 - hormone replacement therapy, 237
 - lung cancer, 197, 343, 372
 - lung function, 252
 - maintaining status, 469
 - risk of death, 12
 - Norms, social, 469
 - North Africa, smoking prevalence, 136
 - North Carolina, oral cancers, 225
 - North Karelia Youth Project, 588, 598
 - Northwest Portland Area Indian Health Board, 602
 - Nortriptyline, 334, 556, 557, 570
 - Norway
 - bladder cancer, 229
 - lung cancer, 210–211
 - pancreatic cancer, 228
 - smoking prevalence, 136
 - NTTS *See* National Teenage Tobacco Surveys
 - Nulliparity *See* Parity
 - Nurses, *See also* U.S. Nurses' Health Study
 - self-help programs, 554
 - weight gain and smoking cessation, 565
 - Nurses' Health Study *See* U.S. Nurses' Health Study
-
- O**
- Obesity *See* Body weight
 - Obstructive airways disease *See* Chronic obstructive pulmonary disease
 - Occupation
 - lung cancer risk, 208
 - smoking status, 486
 - Occupational exposure, environmental tobacco smoke, 123
 - Occupational settings *See* Worksite smoking cessation programs
 - Occupational status, 39, 576
 - Office on Smoking and Health, Centers for Disease Prevention and Control, 602
 - Old Gold brand, advertising, 494
 - Older smokers, cessation, 554
 - Ontario, Canada, air pollution and lung cancer risk, 208
 - Oocytes, 273
 - Oophorectomy, 269
 - Ophthalmopathy, in Graves' disease, 263
 - Oral cancer, 13, 116, 224–225, 369
 - smokeless tobacco, 231
 - Oral clefts, 305
 - environmental tobacco smoke, 366
 - Oral contraceptives
 - breast cancer, 213
 - coronary heart disease, 236–237, 369

coronary heart disease risk, 248
ectopic pregnancy, 277
heart disease risk, 13–14
nicotine, 561–562
smoking cessation, 552
stroke, 241, 248, 370
stroke risk, 14
thyroid cancer, 230
Organisation for Economic Co-operation and Development, 597
Oropharyngeal cancer, 231, 369
Oslo Youth Study Smoking Prevention Program, 598
Osteoarthritis, 15, 329, 330, 371
Osteoblasts, 318
Osteoporosis, 311, 317, 583
Ovarian atresia, 219
Ovarian cancer, 13, 219–220
 risk, 223, 369
Ovarian cysts, 272
Ovarian function, 269
Ovarian pathology, 273
Ovulation frequency, cancer risk, 219
Oxford Family Planning Contraceptive Study, 321

P

p53-Tumor-suppressor gene, 207
p53 Positive tumors, 216
Pacific Islands, smoking prevalence, 136
Pack-years, 198
 carotid atherosclerosis, 241
 chronic obstructive pulmonary disease, 261
 coronary heart disease risk, 353
 environmental tobacco smoke, 344, 353
 lung cancer, 201
 menopause, 269
 oocytes, 273
 rheumatoid arthritis, 328
Pall Mall brand, advertising, 498
Pancreatic cancer, 13, 228, 231, 369
Panic disorders, 131
Papillary thyroid cancer *See* Thyroid cancer
Parathyroid hormone, 318
Parental, smoking, 468–469, 470
Parental education, smoking initiation, 471
Parental smoking, 252, 468–469, 470; *See also*
 Maternal smoking
 diabetes in children, 264
Parents, smoking initiation, 455, 468–469
Parity, 483
 low birth weight, 296
 placenta previa, 281
 preeclampsia, 288
 smoking cessation, 579
 stillbirth, 291
Parkinson's disease, 16, 338, 372
Parliament brand, advertising, 497, 509
Partner *See also* Spouse
 smoking status and smoking after pregnancy, 483–485
Passive smoking *See* Environmental tobacco smoke
Paternal smoking
 birth weight, 364
 congenital malformations, 366
 neonatal mortality, 365
Pawtucket Heart Health Program, 589
Peers, 67
 attitudes toward smoking, 470
 smoking initiation among adolescents, 469
Pelvic inflammatory disease, 276, 278
Peptic ulcers, 15, 324–325, 371
Peribronchiolar alveoli, inflammation, 249
Perinatal mortality, 15, 290, 303, 307, 368, 370, 372; *See also*
Neonatal mortality
 environmental tobacco smoke, 16
Periodontitis, 116
Peripheral vascular disease, 244
 atherosclerosis, 14, 370
 smoking cessation, 248
 subclinical, 247
Personal characteristics, 478
 smoking initiation, 471
Personal influences, 453
Personality disorders, 131
Peter Stuyvesant brand, brand stretching, 522
Pharmacologic approaches
 smoking cessation, 556–558
 weight control, 567–569
Pharmacologic processes, 486–487
Pharyngeal cancer, 13, 224–225
Phenylpropanolamine gum, 569
Philip Morris, 490, 498
 advertising, 510, 520, 521, 524
 cigarette brand, 46
 international sponsorship, 523
 product placement, 523
 promotions, 510
 sponsorship, 513–514
Physical activity, 486
Physician intervention, 554
Physicians, advice about smoking, 112–115, 116
Pima Indian women, *See also* American Indian women
 type 2 diabetes, 264
Pipes, 12, 118–119, 132
Placental abnormalities, 291
Placenta previa, 14–15, 281, 307, 370
Plaque thickness, 243
Players Navy Cut brand, 519
Pneumonia, 252
Polacrilex gum, 332, 556; *See also* Nicotine gum
Poland
 air pollution and lung cancer risk, 209
 lung cancer, 209, 211
Policy, smoking cessation, 584–597
Polycyclic aromatic hydrocarbons, 219, 269, 273
Polymorphism, 476

- Population attributable risk, 190
- Portugal
 lung cancer, 209, 211
 smoking prevalence, 136
 smoking trends, 138
- Postpartum period, 564
 relapse to smoking, 487, 555
- Poverty, 575; *See also* Income level
 smoking cessation, 101
 smoking prevalence, 42
- Predictor variables, 479
- Predictors, smoking cessation, 483
- Preeclampsia, 15, 283, 288, 307, 370
- Pregnancy, 8, 12, 71–73, 142
 adverse outcomes, 288
 bleeding during, 278
 diabetes in children, 264
 ectopic, 276, 277–278, 307, 370
 estriol levels, 262
 estrogen levels, 262
 gallstones, 321
 maternal smoking and infant lung development, 251
 nausea and vomiting, 561
 nicotine replacement therapy, 557
 prenatal smoking, 484
 self-help programs, 554
 smoking, 71–73
 smoking cessation, 17, 111–112, 483–485, 552, 555, 562–564, 606
 smoking cessation among black women, 579–580
 smoking cessation among low income women, 575
 smoking prevalence, 73
 smoking reduction, 602
 socioeconomic status and smoking cessation, 577–578
- Pregnancy rates, 273
- Pregnancy Risk Assessment Monitoring System, 73
- Premature rupture of membranes, 290
- Premenstrual distress, smoking cessation, 560
- Premenstrual dysphoric disorder, 561
- Premenstruum
 nicotine intake, 559
 withdrawal symptoms during, 560
- Prenatal smoking, 484
- Press *See* Media
- Preterm delivery *See* Delivery, preterm
- Preterm premature rupture of membranes, 14, 15, 278–279, 307, 370, 371
- Prevalence *See* Smoking prevalence
- Price, tobacco, 510
- Print media, 521
- Proctitis, ulcerative, 325
- Product packaging, 492–493
- Product placement, 523
- Progestins, 236, 262
- Progestogen, 217
- Project ALERT, 598
- Project SHOUT, 598
- Project SMART, 598
- Project Towards No Tobacco Use, 598
- Prolactin, 306
- Promotions, 490, 491–492, 509–511
 restrictions, 524–526
- Proteinuria, 283, 288
- Proximal duodenum, 324
- Proximal humerus, fractures, 319
- Proximal influences, 453
- Prozac *See* Fluoxetine hydrochloride
- Psychiatric disorders, 15–16, 131, 333–336, 371
- Psychological disturbances, 473
- Psychosocial problems and smoking, 134
- Psychosocial resources, smoking after pregnancy, 484
- Psychosocial variables, smoking cessation, 552
- Public attitudes, 592
- Public health initiatives, 599–600
- Public Health Service Update (2000), 555
- Public health settings, smoking cessation, 576
- Public transportation, 521
- Pueblo students, *See also* American Indian girls
 smoking prevention, 598
- Puerto Rican American girls
 smoking intensity, 62
 smoking prevalence, 61
- Puerto Rican American women, 39
 cigarette brand preference, 47
 environmental tobacco smoke, 122
 heavy smokers, 45
 smoking during pregnancy, 73
 smoking prevalence, 29
- Puff volume, 330
-
- Q**
-
- Quangzhou, China, environmental tobacco smoke exposure, 345
- Quit ratio, 100
- Quit Victoria, 604
- Quitting smoking *See* Smoking cessation
-
- R**
-
- Race
 age at smoking initiation, 76
 birth weight, 302
 cigarette brand preference, 70
 lung cancer, 200
 preeclampsia, 288
 relapse to smoking, 487
 smoking among girls, 60, 61
 smoking prevalence, 57, 73
 spontaneous abortion, 281
 stillbirth, 291
- Radiation
 congenital malformations, 303
 ionizing, 209

thyroid cancer, 230
Radio, 521
Radius, 312
Radon, 207
 lung cancer, 209
Rancho Bernardo Heart and Chronic Disease Study, 253
Ranitidine, 324
Rate difference, smoking status and mortality, 186
Rebelliousness, smoking initiation among adolescents, 472
Receptors, D₂ dopamine, 476
Reducing Tobacco Use, 556
Reemtsma R1 Minima brand, appeal to women, 522
Refusal skills, 474
Regent brand, advertising, 496
Reinforcers, conditioned, 342
Relapse to smoking *See* Smoking, relapse to
Relative risk, defined, 184
Relaxation, smoking, 473
Religion, 66, 67, 71
Religiousness, 473
Renal cell cancer, 228
Renal parenchyma, 228
Renal pelvis cancer, 228
Report on Carcinogens, 344
Reproduction
 assisted, 273
 environmental tobacco smoke, 343
Reproductive age, smoking prevalence, 35–36
Reproductive history, smoking, 219
Reproductive outcomes, 14, 272–307, 370–371
 and environmental tobacco smoke, 16
 environmental tobacco smoke, 356–368, 372
Research, 636
 smoking prevention, 597–600
Respiratory anomalies, 303
Respiratory tract, lower, 252
Restaurants, smoking restrictions, 593
Restlessness, and smoking, 134
Retinol, 208
Rheumatoid arthritis, 15, 327–329, 330
Risk behavior syndrome, 472
Risk factors
 other than smoking, 189–190
 postpartum relapse, 564
Ritz brand, 506
R.J. Reynolds, 490, 506, 508
 advertising, 510
 counteradvertising, 603
 international sponsorship, 523
 promotions, 510
 sponsorship, 511, 513–514
RJR Nabisco *See* R.J. Reynolds
Robert Wood Johnson Foundation
 cigar use, 118
 community intervention programs, 589
Robert Wood Johnson Youth Access Survey, 595

Romania, lung cancer mortality, 211
Rompa con el Vicio: Una Guía Para Dejar De Fumar, 582
Rural population, smoking cessation among blacks, 581

S

Safe@Work coalition, 513
Saint Laurent, Yves, 506; *See also* Yves Saint Laurent brand
Salem brand, 46, 47
 advertising, 506
 international sponsorship, 523
Salivary gland tumors, 225
SAMMEC *See* Smoking Attributable Mortality, Morbidity, and Economic Costs
Satin brand, 506
Schizophrenia, 16, 131, 335, 336, 371
School grade, 56
 current smoking, 58, 59
School-based interventions, 597–598
Schools
 smoking cessation programs, 572
 smoking initiation, 472
 smoking restrictions, 593
Scotland, lung cancer in women, 210
Seizures, 288
Selenium, 208
Self-efficacy, 470, 483, 488
Self-esteem, 473
Self-help programs, 553–554, 564, 581
 black women, 579
Self-image, 490
 tobacco advertising, 504
Self-medication, 15, 333, 336, 340
 smoking and depression, 471
Self-report, smoking during pregnancy, 563
Sensation seeking, 472
Sensory cues, 342
Sensory motivation, 482
Sensory performance, nicotine effects, 341
Sex hormone-binding globulin, 218
 body fat distribution, 311
Sex hormones, 14, 230, 262–263, 370
 effects of nicotine, 269
 oxidase systems, 217
Sexual behavior, 223
Sexual intercourse, smoking and cervical cancer, 220
Sexually transmitted diseases, 223, 277, 331
Shenyang, China, air pollution and lung cancer risk, 208
Sibling smoking, smoking initiation among adolescents, 469
Silk Cut brand, brand stretching, 522
Silver acetate, 556
Singapore, smoking prevalence, 136
Single photon absorptiometry, 312

- Skeleton, axial, 312
- SLC6A3-9*, 476
- Sleeping position, infant, 306
- Smoke *See* Environmental tobacco smoke; Tobacco smoke
- Smoke inhalation
 - aortic aneurysm, 247
 - cancer risk, 202
 - cervical cancer, 220
- Smoke-free policies, 123–124
- SmokeFree Kids & Soccer Campaign, 600
- SmokeLess States program, 589
- Smokeless tobacco, 12, 116, 142
 - cessation programs, 549–550
 - international use, 138–139
 - marketing, 516–517
 - oral cancer, 225, 231
 - prevalence, 122
 - use by women, 119–121
- Smokers, current
 - benign breast disease, 217
 - breast cancer, 212, 215
 - cervical cancer, 223
 - chronic obstructive pulmonary disease, 259
 - dysmenorrhea, 266
 - endometrial cancer, 369
 - gestational diabetes, 265
 - hip fractures, 318
 - inflammatory bowel disease, 325
 - intermittent smoking, 100
 - lung function, 250, 252
 - mental illness, 333
 - peripheral vascular disease, 247
 - risk of death, 12
 - stroke risk, 240
 - ulcerative colitis, 371
 - weight gain, 565
- Smokers, former
 - benign breast disease, 217
 - body weight, 311
 - breast cancer risk, 212
 - chronic obstructive pulmonary disease, 250, 261, 370
 - diabetes risk, 264
 - dysmenorrhea, 266
 - endometrial cancer, 369
 - facial wrinkling, 332
 - gestational diabetes, 265
 - hip fracture, 318
 - infertility, 273–276
 - inflammatory bowel disease, 325
 - lung cancer risk, 13, 200, 209, 369
 - lung function, 252
 - spontaneous abortion, 282
 - stroke risk, 230
 - ulcerative colitis, 325
- Smoking
 - acute myeloid leukemia, 231
 - age at initiation, 12
 - antiestrogenic effects, 217, 230, 262
 - aortic atherosclerosis, 247, 248
 - birth cohorts, 27
 - blood pressure, 247, 248
 - body weight, 125–128
 - breast cancer, 212–217, 224
 - calming effects, 85
 - cervical cancer, 220–224
 - combined with alcohol, 224–225, 226
 - coronary heart disease, 232–238
 - current, 34–42
 - daily *See* Smoking, current
 - depression risk, 334
 - disease outcomes, 634
 - endometrial cancer risk, 224
 - escalation to, 454
 - esophageal cancer, 226
 - experimental, 67, 474; *See also* Smoking initiation
 - frequent, 62
 - goiter, 263
 - heavy *See* Smoking intensity
 - impact on women's health, 633
 - insulin resistance, 265
 - laryngeal cancer, 225–226
 - lung cancer, 209
 - oral cancer risk, 225
 - oral contraceptives, 236–237
 - oropharyngeal cancers, 224–225
 - ovarian cancer, 219–220, 224
 - perceived norms, 469
 - peripheral vascular disease, 244–247
 - pharyngeal cancer risk, 224–225
 - relapse to, 485, 487–488, 551–552
 - relaxing effects, 85
 - stomach cancer, 226
 - transition from, 479–487
 - trends, 34–39
 - type 2 diabetes risk, 264
 - uterine fibroids, 271–272
 - vulvar cancer risk, 224
 - young women, 49–51
- Smoking abstinence
 - chronic obstructive pulmonary disease, 250
 - coronary heart disease risk, 233
 - lung function, 252
 - stroke risk, 248
- Smoking, age at initiation
 - breast cancer risk, 213
 - coronary heart disease, 233, 233
- Smoking Attributable Mortality, Morbidity, and Economic Costs, 190
- Smoking behavior, research, 635
- Smoking cessation 8, 12, 17, 115, 142, 477–488; *See also*
- Smoking reduction
 - abruptio placentae, 281
 - adolescent girls, 572–573
 - adults, 485–486
 - age groups, 482–487

- alcohol consumption, 485
 - antidepressive drug treatment, 334, 336
 - attempts, 16, 93–99, 96, 479, 485
 - birth weight, 303
 - body weight, 125, 309–310
 - bone density, 315
 - carotid atherosclerosis, 248, 370
 - chronic obstructive pulmonary disease, 250, 253, 261
 - community studies, 588–590
 - conception delay, 307
 - coronary heart disease, 233, 248, 369
 - demographic characteristics, 101–102
 - depression, 333, 372
 - duration of abstinence, 488
 - economic benefits, 564
 - gender, 102–106, 551
 - girls, 86, 95–96
 - infertility, 307
 - innovations, 550
 - issues for women, 558–564
 - liver cancer risk, 228
 - low birth weight, 307
 - lung cancer risk, 369
 - lung function, 262, 370
 - mass media approach, 576–577
 - menstrual cycle, 560–561
 - methods, 17, 96–99, 116, 549–558
 - mood disorders, 372
 - neonatal mortality, 295
 - nicotine dependence, 88
 - nicotine dependence treatment methods, 552–558
 - older adults, 486–487, 562
 - older women, 583
 - overweight women, 127
 - peptic ulcers, 324
 - peripheral vascular disease, 244, 370
 - pharmacological treatments, 636
 - predictive variables, 86, 487
 - predictors, 485, 488, 527, 553
 - pregnancy, 15, 111–112, 471, 483–485, 562–564
 - premature death risk, 368
 - preterm delivery, 291, 307
 - preterm premature rupture of membranes, 279, 307
 - programmatic and policy approaches, 584–597
 - psychiatric disorders, 16
 - psychosocial predictors, 482
 - public health settings, 576
 - reasons, 106
 - research, 635
 - stages, 109–110
 - stroke, 248
 - stroke risk, 240, 370
 - study population, 479
 - trends, 100–101
 - unassisted, 552–553, 574
 - weight gain, 309, 311, 431, 371
 - worksite activities, 584–585
 - young persons, 482–483
- Smoking Cessation, Clinical Practice Guideline, Agency for Health Care Policy and Research, 555*
 - Smoking continuum, 115
 - ever smoking, 107–109
 - Smoking duration
 - aortic aneurysm risk, 247
 - chronic obstructive pulmonary disease, 370
 - coronary heart disease, 232
 - lung cancer, 202
 - lung cancer risk, 198
 - mortality, 193, 368
 - Smoking initiation, 74–82
 - age at, 198
 - among adolescents, 63
 - biological factors, 476–477
 - birth cohorts, 77–81
 - body weight, 308
 - cigarette price, 454–455
 - decision to smoke, 455
 - environmental factors, 453–470
 - influencing factors, 454–477
 - lung cancer, 198
 - mood, 476
 - other drug use, 130, 134
 - perceived approval, 470
 - personal factors, 475–476
 - predictors, 454, 527
 - prevention, 549
 - rate, 81–82
 - smoking duration risk, 198
 - social factors, 454–470
 - stages, 474
 - tobacco marketing, 498–505, 527
 - Smoking intensity 12, 13, 62–63, 193, 372; *See also* Cigarettes, number smoked; Dose-response relationship
 - abruptio placentae, 281
 - all-cause death rates, 186
 - among girls, 70
 - birth weight, 303
 - body weight, 308
 - breastfeeding, 306
 - chronic bronchitis, 250
 - chronic obstructive pulmonary disease, 13, 261, 370
 - coronary heart disease, 233, 233
 - diabetes mellitus, 264
 - dysmenorrhea, 266
 - environmental tobacco smoke, 344, 353
 - fertility, 276
 - gallstones, 321
 - heavy, 42–46, 48, 574
 - hip fracture, 318
 - human immunodeficiency virus, 331
 - human papillomavirus, 223
 - intermittent, 42, 100
 - liver cancer, 227–228
 - low birth weight, 302
 - lung cancer, 197, 198, 200, 368

- macular degeneration, 330
- mortality risk, 188–189, 368
- nicotine dependence, 88, 89, 142, 372
- nicotine exposure, 340
- osteoarthritis, 329
- ovarian cancer, 219
- pancreatic cancer, 228
- pregnancy, 563
- pregnant women, 73
- preterm delivery, 290
- smoking cessation, 86, 100, 482, 485, 486, 488, 527
- stroke risk, 240
- trends, 42–46
- women and smoking cessation, 573
- Smoking intention, 470
- Smoking maintenance, 477–488
 - study populations, 479
- Smoking policies, 17
 - restrictive, 585–586, 587, 592
- Smoking, pregnancy, 564
- Smoking prevalence, 7, 11, 28–29, 39–42, 48
 - among high school senior girls, 60
 - among men, 29
 - during pregnancy, 73
 - ethnicity, 635
 - international patterns, 135–139
 - international trends, 12, 138, 142
 - lung cancer, 203
 - mental illness, 333
 - perceived norms, 469
 - race, 635
 - sexual orientation, 635
 - socioeconomic status, 635
- Smoking prevention, 597–600, 606
 - research, 635
- Smoking reduction, 8, 12, 11
- Smoking reduction, 17, 86, 88, 97, 99, 636; *See also* Smoking cessation
- Smoking, relapse to, 107, 551, 552
 - body weight, 125
 - menstrual cycle, 560–561
 - mental health, 131
 - postpartum period, 564
 - pregnancy, 563–564
- Smoking status
 - attitudes, 67
 - benign breast disease, 217
- Smoking trends, 138
 - historical, 23–28
- Sociability, smoking initiation among adolescents, 471
- Social acceptability, 521
- Social bonding, relationship to smoking, 63
- Social class, spontaneous abortion, 281
- Social desirability of smoking, 67
- Social factors, 453, 478
 - smoking cessation, 551
 - smoking initiation, 454–470
- Social network, pregnancy and smoking cessation, 484
- Social reality, 491
- Social strata, 486
- Social support, 570–572
 - smoking cessation, 488, 527
- Sociodemographic factors, 65–66, 71
- Socioeconomic status, 575–576
 - cessation activities at work, 578
 - lung development, 251
 - relationship to smoking, 63–67
 - smoking cessation, 575
 - smoking initiation among adolescents, 471
- Somatostatin, 265
- South Africa
 - smokeless tobacco use, 139
 - smoking prevalence, 136
- South African women, smoking prevalence, 136
- South American women, smoking during pregnancy, 73
- Southwest Cardiovascular Curriculum Project, 598
- Spain
 - lung cancer, 211
 - smoking prevalence, 136
 - smoking trends, 138
- Special Lights brand, international sponsorship, 523
- Sperm, environmental tobacco smoke, 366
- Spina bifida. *See* Neural tube defects
- Spine, 312
- Spirometry, 249, 253
- Sponsorship, 490, 492, 511–515, 633
 - counteractivities, 604–605
 - international, 522–523
 - restrictions, 515
 - women's tennis, 604–605
- Sporting events, sponsorship, 492
- Spousal smoking, 350
 - coronary heart disease, 353, 372
 - delayed conception, 366
 - environmental tobacco smoke, 356
- Spouse
 - environmental tobacco smoke, 343, 344
 - smoking after pregnancy, 485
- Spud brand, advertising, 494
- Squamous intraepithelial neoplasms, 220
- Stages-of-change model, 109–110, 478
- Stanford Five-City Project, 36, 588–589
 - smoking intensity, 45
- Stanford Three-Community Study, 588
- Stillbirth, 15, 291–294, 304, 307, 370
 - environmental tobacco smoke, 364
- Stimulation, nicotine, 341
- Stomach cancer, 13, 226, 231, 369
- Strategic Coalition of Girls and Women United Against Tobacco, 601, 605
- Stress, 473, 557
 - nicotine, 341
- Stress reduction, 569
 - smoking, 473
- Stroke, 14, 232, 238–241, 369
 - hemorrhagic, 240

ischemic, 240
risk, 248
Students, restrictive smoking policies, 593
Study of Osteoporotic Fractures, 315
Style brand, 508
Subarachnoid hemorrhage, 14, 369
 oral contraceptives, 241
 risk, 240, 248
Substance abuse, adolescents, 573
Sudden infant death syndrome, 15, 290, 306–307, 370
Sun exposure, 332
Survey of American Indians and Alaska Natives of the
 National Medical Expenditure Survey (1987), 40
Surveys, public tobacco-related attitudes, 592
Survival rates, lung cancer, 195
Susceptibility to smoking, 474
Sweden
 beauty pageant, 606
 bladder cancer, 229
 lung cancer mortality, 210–211
 pancreatic cancer, 228
 smokeless tobacco use, 138
 smoking trends, 138
Swedish National Institute of Public Health, 606
Swedish Nurses Against Tobacco, 601
Switzerland, lung cancer mortality, 210–211
Systemic lupus erythematosus, 15, 329–330, 371

T

Take Heart project, 586
TAPS I *See* Teenage Attitudes and Practices Survey I
TAPS II *See* Teenage Attitudes and Practices Survey II
Tar content
 adenocarcinoma, 203
 low, 48, 97, 506
 lung cancer risk, 200
Tareyton brand, advertising, 494
Task Force on Women & Girls, Tobacco & Lung
 Cancer, 601
Taxation policies, 594–595
Tecumseh Community Health Study
 coronary heart disease, 232
 type 2 diabetes, 264
Teenage Attitudes and Practices Survey I, 23, 52
 body weight, 126
 cigarette brand preference, 70, 500
 depression in girls, 134
 emotions, 134
 ever smoked, 55
 smokeless tobacco, 121
 smoking cessation among girls, 111
 smoking cessation attempts, 95
 weekly income, 66
Teenage Attitudes and Practices Survey II, 23, 52
 cigarette brand preference, 69, 500
 depression in girls, 134
 emotions, 134
 ever smoked, 54
 first cigarette of the day, 84
 nicotine dependence indicators, 91
 nicotine withdrawal, 88–89, 91, 93
 reasons for smoking, 85
 smokeless tobacco, 121
 smoking cessation among girls, 111
 weekly income, 66
Television, 521
Temporal trends, smoking-associated risks, 202–203
Tennis, tobacco company sponsorship, 511, 604–605
Tennis Legends Tour, 604
Testosterone
 body fat distribution, 311
 maternal levels, 477
 metabolism, 262–263
Tetralogy of Fallot, environmental tobacco smoke, 366
Thailand, smoking prevalence, 136
Thiocyanate, 263
 levels during pregnancy, 563
Throat irritation, 494, 495, 498
Thyroid cancer, 13, 230–231, 369
Thyroid disorders, 74, 263–264, 370
Thyroid hormones, 74, 263, 265, 370
Thyroid nodules, 230
Thyroid volume, 263
Thyroid-stimulating hormone, 230, 263
Thyroiditis, autoimmune, 263
Thyrototoxicosis, 230
Thyroxine, 263
Tobacco company sponsorship *See* Sponsorship
Tobacco control, women's leadership, 605–606
Tobacco control advocacy programs, 600–606
Tobacco control programs, 636
Tobacco industry, 8
 marketing, 16–17
 promotion, 524
 targeted advertising, 633–634
Tobacco Institute, sponsorship, 514
Tobacco Institute in Japan, advertising restrictions, 525
Tobacco prevention strategies, 600
Tobacco products
 accessibility, 454
 advertising and promotion, 455
 price, 510
 sales to minors, 454
Tobacco settlement *See* Master Settlement Agreement
Tobacco smoke, 273; *See also* Environmental tobacco
 smoke
 exposure and gender, 202
 inhaled, 198
 lung cancer histology, 201–202
Tobacco Society for Voice Culture, 494
Tobacco use, 16
 patterns, 11–12
 twin studies, 476
Tobacco-related policies, 591–597
Tobacco-specific nitrosamines, 203–205, 635

Toluene, neonatal mortality, 365
 Tongue cancer, 225
 Tooth loss, 110
 Transdermal nicotine patch *See* Nicotine patch
 Transforming growth factor alpha, 305, 366
 Transient ischemic attack, hormone replacement therapy, 241
 Transtheoretical model of change, 474, 478
 Treatment approaches, older women, 583–584
 Trends *See* Smoking trends
 Triiodothyronine, 263
 TSNAs *See* Tobacco-specific nitrosamines
 Tucson Epidemiologic Study of Obstructive Lung Diseases, 249, 252
 Tucson Epidemiological Study of Airways Obstructive Disease, 253
 Tumor-suppressor genes, 216
 Twin studies

- bone density, 315
- gallstones, 322
- peptic ulcers, 324
- rheumatoid arthritis, 328
- tobacco use, 476

 Type 1 diabetes *See* Diabetes mellitus
 Type 2 diabetes *See* Diabetes mellitus

U

Ultimate influences, 453
 Underreporting, 53, 58

- smoking during pregnancy, 73

 Unemployment, 486
 United Kingdom

- advertising, 519
- counteradvertising, 603
- lung cancer, 209, 210–211
- smoking trends, 138

 United States

- lung cancer in developed countries, 209–211
- smoking trends, 138

 United States Tobacco Company, 490
 Uranium miners, lung cancer, 209
 Urinary tract cancers, 228–230
 Urinary tract malformations, 305–306
 Urogenital malformations, 303, 305–306
 U.S. Department of Defense survey, smokeless tobacco, 120
 U.S. Environmental Protection Agency, 122
 U.S. Food and Drug Administration, Drug Abuse Advisory Committee, 99
 U.S. Nurses' Health Study

- body weight, 308
- cataract, 330
- cerebral hemorrhage, 240
- coronary heart disease, 232, 233, 234, 353
- duration of smoking, 188–189
- environmental tobacco smoke, 353
- ever smoking by age, 78

- gallstones, 322
- menopause, 269
- mortality risk, 189
- pancreatic cancer, 228
- polymorphism and breast cancer, 216
- rheumatoid arthritis, 327
- smoking cessation, 485, 487
- systemic lupus erythematosus, 330
- type 2 diabetes, 264
- weight gain, 310

U.S. Women's National Soccer team, 600
 Uterine fibroids, 14, 271–272, 370
 Uteroplacental circulation, 303

V

Variables, dependent, 454
 Vascular disruption, congenital malformations, 304
 Vasoconstriction, fetal growth retardation, 364
 Vegetable intake, 486
 Vertebral fractures, 319
 Victorian Smoking and Health Program, 604
 Video, 521
 Vietnamese women, *See also* Asian or Pacific Islander women

- lung cancer incidence, 195

 Virginia Slims brand, 504–506, 508

- advertising, 510, 521, 525
- appeal to women, 522
- nitrosamine content, 203, 205
- promotions, 509
- sponsorship, 511
- tennis tournament, 604

 Vitamin A, 208
 Vitamin C, 208

- cervical cancer, 223

 Vitamin E, 208
 Vitamin supplements, 291
 Vlagtwedde-Vlaardingen study, 253
 Vogue brand, appeal to women, 521
 Vulvar cancer, 13, 224, 369

W

WAGAT *See* Women And Girls Against Tobacco
 Waist-to-hip ratio, 310, 315
 Wales, lung cancer, 210
 Walnut Creek Study, coronary heart disease, 232
 Waterloo Smoking Prevention Project, 598
 Weight *See* Body weight
 Weight control, 475, 488, 527, 565–569

- cigarette advertising, 494
- smoking initiation, 16

 Weight gain, 15, 309, 311, 551, 565, 606

- pregnancy, 288
- smoking cessation, 17, 486

 Weight loss, 15, 134, 311, 371
 Well-being, 333

- Western countries *See* Developed countries
- Wheezing, 251, 252
- White adolescents
- emotional distress, 473
 - risk taking behavior, 472
 - smoking prevalence, 635
 - weight control, 475
- White boys, smoking prevalence, 61
- White girls
- cigarette brand preference, 70
 - current smoking, 58
 - ever smoked, 55, 56, 59
 - physician advice about smoking, 115
 - smoking intensity, 62–63
 - smoking prevalence, 7, 11–12, 57, 60, 67, 70, 142
 - weight control, 476
- White men
- chronic obstructive pulmonary disease mortality, 260
 - lung cancer incidence, 194
 - smoking cessation, 106
 - smoking prevalence, 29
- White women
- age at smoking initiation, 74, 76, 77, 80
 - birth weight, 302
 - body mass index, 128
 - body weight, 127
 - bone fractures, 311
 - chronic obstructive pulmonary disease mortality, 260
 - cigarette brand preference, 46
 - coronary heart disease risk, 232
 - current smoking, 39
 - environmental tobacco smoke, 123
 - ever smoking, 33, 78
 - facial wrinkling, 332
 - first cigarette of the day, 83
 - heavy smokers, 43, 45, 46
 - hip fracture risk, 318
 - lung cancer, 193, 197, 200
 - neonatal mortality, 294
 - nicotine dependence, 88
 - pregnancy and smoking cessation, 483, 577
 - relapse to smoking, 487
 - smokeless tobacco, 119
 - smoking cessation, 97, 100, 101, 105, 110
 - smoking cessation during pregnancy, 111, 484
 - smoking continuum, 107
 - smoking during pregnancy, 71, 73
 - smoking initiation rate, 82
 - smoking prevalence, 11, 28–29, 35, 48, 142
 - stillbirth, 297
- White young men, ever smoking, 49
- White young women
- current smoking, 49
 - ever smoking, 49
 - smoking cessation, 110
 - smoking prevalence, 50
- WHO Collaborative Study, oral contraceptives and
- myocardial infarction, 237
- WHO MONICA survey, smoking prevalence, 136
- Winston brand, 46, 47, 504
- sponsorship, 514
- Withdrawal symptoms, 86, 88, 557
- gender differences, 476
 - menstrual cycle, 559–560
 - pharmacology, 557
- Women, *See under* entries at Alaska Native, American Indian, Asian or Pacific Islander, Black, Canadian Indian, Chinese, Cuban American, Filipino, Hispanic, Hopi, Japanese; entries at Korean, Maori, Military, Navajo, Puerto Rican American, South African, and White
- Women And Girls Against Tobacco, 603, 604, 605
- Women Talk to Women About Smoking, 602
- Women United Against Tobacco Workshop, 605
- Women vs. Smoking Network, 603
- Women's Action on Smoking, 526
- Women's Health Equity Act, 514
- Women's Health Initiative, 237, 562
- Women's magazine projects, 603–604
- Women's magazines *See* Magazines
- Women's organizations
- tobacco control advocacy programs, 601–602
 - tobacco sponsorship, 511, 513
- Working Well Trial, 586–587, 593
- Workplace
- environmental tobacco smoke, 123–125, 350, 353
 - pregnancy and smoking cessation, 484
- Workplace exposure
- coronary heart disease, 353
 - low birth weight, 362
- Worksite
- smoking cessation, 578
 - smoking restriction policies, 593
- Worksite smoking cessation programs, 575, 584–587
- efficacy, 586–587
- World Conference on Smoking and Health, 605
- World Conference on Tobacco OR Health, 600, 605
- World Health Organization, 605, 637
- Collaborative Study, 237
 - MONICA survey, 136
 - Regional Office for the Western Pacific, 525
 - SmokeFree Soccer campaign, 600
- World Smoking and Health*, 602
- Wrinkling *See* Facial wrinkling
-
- ## Y
- Years of potential life lost, 190–191
- U.S. women, 193
- Young adults
- attitudes about smoking, 67
 - smoking cessation, 485–486
 - smoking initiation rates, 82
- Young men
- ever smoking, 49

sexual orientation, 59
 smoking initiation, 82, 500
 smoking prevalence, 51
 Young women, *See also under* entries at Alaska Native, American Indian, Asian or Pacific Islander, Black, Chinese, Filipino, Hispanic, Japanese, Korean, and White
 cigarette smoking, 48–51
 current smoking, 49–51
 drug use, 129–130
 first cigarette of the day, 83–85
 nicotine dependence, 89, 91
 other drug use, 130
 physician advice about smoking, 115
 reasons for cigarette use, 86
 sexual orientation, 59
 smoking cessation, 110–111
 smoking initiation by age, 80
 Young Women’s Christian Association, 601
 Youth *See* Adolescents
 Youth, bisexual, 59
 Youth Access Survey, 595
 Youth Risk Behavior Survey, 23, 52, 129–130, 472
 binge drinking, 130
 body weight, 125–126

cigar use, 118
 current smoking, 57
 ever smoked, 54, 56
 frequent smoking, 63
 sexual orientation, 59
 smokeless tobacco, 121
 smoking cessation attempts, 95–96
 smoking initiation rates, 82
 YPLL *See* Years of potential life lost
 YRBS *See* Youth Risk Behavior Survey
 Yves Saint Laurent brand *See also* Saint Laurent, Yves
 appeal to women, 522
 international advertising, 520

Z

Zinc, maternal levels, 278
 Zyban *See* Bupropion