

Drug Facts (continued)

■ taking any other drug

right away.

Directions

Weight (lb)

under 24 lb

24-35 lb

36-47 lb

48-59 lb

60-71 lb

72-95 lb

Ask a doctor or pharmacist before use if the child is Ask a doctor or pnarmacist before use if the child i
under a doctor's care for any serious condition
taking any other product that contains ibuprofen,
or any other pain reliever/fever reducer
taking a prescription product for anticoagulation
(blood thinning)

When using this product give with food or milk if

Stop use and ask a doctor if
an allergic reaction occurs. Seek medical help right away.

from the first term of the first days from the first day (24 hours) of treatment

■ stomach pain or upset gets worse or lasts
■ redness or swelling is present in the painful area
■ any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center

■ do not give more than directed
■ shake well before using
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
■ repeat dose every 6-8 hours, if needed
■ do not use more than 4 times a day

measure only with the blue dosing cup provided.
 Blue dosing cup to be used with Children's Advil
 Suspension only. Do not use with other products
 Dose lines account for product remaining in cup
 due to thickness of suspension.

Dosing Chart

Age (yr)

under 2 yr

2-3 yr

4-5 yr

6-8 yr

9-10 yr

11 yr

Dose (tsp)

ask a doctor

1 tsp

1 1/2 tsp

2 tsp

2 1/2 tsp

3 tsp

Drug Facts (continued)

TQuestions or comments?

Wyeth®

Wyeth Consumer Healthcare Madison, NJ 07940 Made in USA U.S. Patent No 4,788,220

F.P.O.

UK24326-1

or health care professional

Alcohol free

Other information


expiration date Inactive ingredients

venuoi luentintation	I IIIIIGI NOIGS	
Production Packaging Dept. Linda Hill 6.27.3	PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIG PRINTER TO SUPPLY COLOR MATCHPRINT FOR CO TO PRESS RUN.	
DEDARTMENT	DEVIEWED BY	DATE

DEPARTMENT	REVIEWED BY		DATE	Wi
	Print Name	Signature		Chang
Graphic Artist				
Packaging Engineer				
Labeling Compliance				
Medical				
Regulatory Affairs				
Marketing				
Legal				
Trademark				一一

DRUG FACTS TEXT DEFINED	TYPE SIZE
DRUG FACTS Title	9 pt
DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
SPACE AFTER BULLETED SECTION	2 ems
BARLINES, HAIRLINES	1.5 pt, .5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

RELEASED		
(1)	Date	
(2)	Date	





Children's

mportant: Read all product information before using. Keep this box for important information. This product is intended for use in children ages years - 11 years.

Drug Facts Active ingredients (in each 5 mL)

.Fever reducer/Pain reliever uprofen 100 mg.

Purposes

Uses temporarily:
■ reduces fever
■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe
allergic reaction which may include:

allergic reaction which may include:
■ hives
■ asthma (wheezing)
■ facial swelling
■ shock
Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever Ask a doctor before use if the child has

■ not been drinking fluids ■ ulcers


- bleeding problems
 lost a lot of fluid due to continued vomiting or diarrhea

 problems or serious side effects from taking fever reducers or pain relievers

 stomach problems that last or come back,

- such as heartburn, upset stomach or pain

 high blood pressure, heart or kidney disease or is taking a diuretic



Weight (lb) Age (yr) Dose (tsp) under 24 lb under 2 yr ask a doctor 24-35 lb 2-3 yr 1 tsp 36-47 lb 4-5 yr 1 1/2 tsp 48-59 lb 6-8 yr 2 tsp 60-71 lb 9-10 yr 2 1/2 tsp 72-95 lb 11 yr 3 tsp

Drug Facts (continued)

■ taking any other drug

stomach upset occurs

right away.

Directions

Ask a doctor or pharmacist before use if the child is

Ask a ductor or priarriacts a benefit use if the clint is under a doctor's care for any serious condition taking any other product that contains ibuprofen, or any other pain reliever/fever reducer taking a prescription product for anticoagulation (blood thinning)

When using this product give with food or milk if

Stop use and ask a doctor if
an allergic reaction occurs. Seek medical help right away.

from the first term of the first days from the first day (24 hours) of treatment

■ stomach pain or upset gets worse or lasts
■ redness or swelling is present in the painful area
■ any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center

■ do not give more than directed
■ shake well before using
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
■ repeat dose every 6-8 hours, if needed
■ do not use more than 4 times a day

measure only with the blue dosing cup provided.
 Blue dosing cup to be used with Children's Advil
 Suspension only. Do not use with other products
 Dose lines account for product remaining in cup
 due to thickness of suspension.

Dosing Chart



Drug Facts (continued)

one dose lasts 6-8 hours
 store at 20-25°C (68-77°F)
 see bottom of box for lot number and

artificial flavors, carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C red no. 40,

glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Call weekdays from 9am to 5pm EST at 1-800-88-ADVIL or ask your pharmacist, doctor

Great Convenience: Long duration means fewer doses needed per day.

Long Lasting: Up to 8 hours of fever relief.

TQuestions or comments?

or health care professional

Other information

expiration date Inactive ingredients

UK24323-1



4 FL OZ (120 mL)


Job Information	C	olor Lege	nd	
Date 6/27/03 PASS #3 Component # UK24323-1 Drawing # 3919 Rev 3				
Identification	DANTONE	PANTONE	PDOOFOO	PDOOFOO
Wyeth	PANTONE 3005 C	Rhodamine Red C	PROCESS YELLOW	PROCESS BLACK
Consumer Healthcare				
Production Packaging Dept. Five / Farms				
Madison, NJ 07940 Phone: (973) 660-5595/5153	PANTONE 485 C			
Fax: (973) 660-6048	h			
Vendor Identification	P	<u> Printer Not</u>	es	
Production Packaging Dept. Linda Hill SC: 6.27.3		ACE FPO UPC'S WITH Ly Color Matchpr		

DRUG FACTS Title
DRUG FACTS CONTIN
• HEADINGS
SUBHEADS/BODY TE
• LEADING
• # OF CHARACTERS P
• BULLETS
SPACE AFTER BULLE
• BARLINES, HAIRLINE
SPACE BETWEEN HAI

DRUG FACTS TEXT DEFINED	TYPE SIZE
DRUG FACTS Title	9 pt
DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
SPACE AFTER BULLETED SECTION	2 ems
BARLINES, HAIRLINES	1.5 pt, .5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

DEPARTMENT	REVIEWED BY		DATE	
	Print Name	Signature		Chan
Graphic Artist				
Packaging Engineer				
Labeling Compliance				
Medical				
Regulatory Affairs				
Marketing				
Legal				
Trademark				

RELEASED			
(1)	Date		
(2)	Date		



Suspension

Ibuprofen 100 mg >> Fever Reducer/Pain Reliever

per 5 mL

Fever

Aches & Pains

Lasts Up To

Hours

BLUE

Alcohol Free

Drug Facts (continued)

- Ask a doctor or pharmacist before use if the child is
- Ask a ductor or priarriacts a benefit use if the clint is under a doctor's care for any serious condition taking any other product that contains ibuprofen, or any other pain reliever/fever reducer taking a prescription product for anticoagulation (blood thinning)
- taking any other drug

When using this product give with food or milk if

stomach upset occurs

- Stop use and ask a doctor if
 an allergic reaction occurs. Seek medical help right away.
- figure away.

 fever or pain gets worse or lasts more than 3 days

 the child does not get any relief within first day
 (24 hours) of treatment
- stomach pain or upset gets worse or lasts
 redness or swelling is present in the painful area
 any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not give more than directed
 shake well before using
 find right dose on chart below. If possible, use weight to dose; otherwise use age.
 repeat dose every 6-8 hours, if needed
 do not use more than 4 times a day

- measure only with the blue dosing cup provided.
 Blue dosing cup to be used with Children's Advil
 Suspension only. Do not use with other products
 Dose lines account for product remaining in cup
 due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (tsp)	
under 24 lb	under 2 yr	ask a doctor	
24-35 lb	2-3 yr	1 tsp	
36-47 lb	4-5 yr	1 1/2 tsp	
48-59 lb	6-8 yr	2 tsp	
60-71 lb	9-10 yr	2 1/2 tsp	
72-95 lb	11 yr	3 tsp	

Drug Facts (continued)

Other information

- one dose lasts 6-8 hours
 store at 20-25°C (68-77°F)
 see bottom of box for lot number and expiration date

Inactive ingredients

carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no. 1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucrose, xanthan gum

TQuestions or comments? Call weekdays from 9am to 5pm EST at 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever relief. **Great Convenience:** Long duration means fewer doses needed per day. Alcohol free

Wyeth®

Wyeth Consumer Healthcare Madison, NJ 07940 Made in USA U.S. Patent No 4,788,220





UK24321-1



mportant: Read all product information before using. Keep this box for important information. This product is intended for use in children ages years - 11 years.

Drug Facts

Active ingredients (in each 5 mL)

Purposes uprofen 100 mg.

Uses temporarily:
■ reduces fever
■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

allergic reaction which may include:

■ hives
■ asthma (wheezing)
■ facial swelling
■ shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

Sore throat warning: Severe or persistent sore threat a cere threat becare the threat thre throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever Ask a doctor before use if the child has

■ not been drinking fluids ■ ulcers

- bleeding problems
 lost a lot of fluid due to continued vomiting
- or diarrhea

 problems or serious side effects from taking fever reducers or pain relievers

 stomach problems that last or come back,
- such as heartburn, upset stomach or pain


 high blood pressure, heart or kidney disease
 or is taking a diuretic

Job Information	Color Legend	
Date 7/14/03 PASS #3 Component # UK24321-1 Drawing # 3919 Rev 3	Color Esgena	
Identification	PANTONE PANTONE PROCESS PROCESS	
Wyeth	3005 C 300 C YELLOW BLACK	
Consumer Healthcare		
Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048	PANTONE 485 C	
Vendor Identification	Printer Notes	
LARS + ASSOCIATES INC.	PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN.	

DRUG FACTS TEXT DEFINED	TYPE SIZE
DRUG FACTS Title	9 pt
DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
SPACE AFTER BULLETED SECTION	2 ems
BARLINES, HAIRLINES	1.5 pt, .5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

DEPARTMENT	REVIEWED BY		DATE	
	Print Name	Signature		Cha
Graphic Artist				
Packaging Engineer				
Labeling Compliance				
Medical				
Regulatory Affairs				
Marketing				
Legal				
Trademark				1

	RELEASED
(1)	Date
(2)	Date



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
SUBHEADS/BODY TEXT	4.5 pt
• LEADING	5 pt
• BULLETS	3.5
SPACE AFTER BULLETED SECTION	2 ems
WARNING BOX LINE, HAIRLINES	.5
Job Information	Color L

Job Information		С	olor Lege	end	
Date 12/17/03 PASS #3 Component # U0171-30-N2 Drawing # 5691 Rev.2					
Identification					
Wyeth Consumer Healthcare	PROCE YELLO		PROCESS BLACK	PANTONE 3005 C	PANTONE 266 C
Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048					
Vendor Identification		Р	rinter No	tes	
PPD: Cheng Lin		SUPPL		H HI-RES DIGITAL A Rint for Color Ai	

DEPARTMENT	REVIEWED BY				
	Print Name	Signature			
Graphic Artist					
Packaging Engineer					
Labeling Compliance					
Medical					
Regulatory Affairs					
Marketing					
Legal					
Trademark					



DIGITAL MECHANICAL COLOR REPRESENTATION MAY BE OFF FOLLOW COLOR LEGEND FOR ACTUAL COLOR.

Warnings	Do Not Use if plastic wrap imprinted v
Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:	"CHILDREN'S ADVIL SAFETY SEALE
■ hives ■ asthma (wheezing) ■ facial swelling ■ shock	is broken or missing

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding. Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever Ask a doctor before use if the child has ■ not been drinking fluids ■ ulcers ■ bleeding problems

■ lost a lot of fluid due to continued vomiting or diarrhea

problems or serious side effects from taking fever reducers or pain relievers stomach problems that last or come back, such as heartburn, upset stomach or pain

■ high blood pressure, heart or kidney disease or is taking a diuretic

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition

■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

■ taking a prescription product for anticoagulation (blood thinning) ■ taking any other drug

When using this product give with food or milk if upset stomach occurs

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

fever or pain gets worse or lasts more than 3 days

■ the child does not get any relief within first day (24 hours) of treatment

stomach pain or upset gets worse or lasts

redness or swelling is present in the painful area

■ any new symptoms appear

with ED"

is broken or missing. Children's

Suspension Fever Reducer/Pain Reliever

Active ingredient (in each 5 mL)......lbuprofen 100 mg Uses temporarily

reduces fever

relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Grape Flavored Liquid 4 FL OZ (120 mL)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

■ do not give more than directed ■ shake well before using

■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day

measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines

account for product remaining in cup due to thickness of suspension. Dosing Chart

Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-95 lb	11 yr	3 tsp

One Dose Lasts 6-8 Hours

Store at 20-25°C (68-77°F)


Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist doctor or health care professional

Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA

U.S. Patent No. 4,788,220

U0171-30-N2

150%



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
SUBHEADS/BODY TEXT	4.5 pt
• LEADING	5 pt
• BULLETS	3.5
SPACE AFTER BULLETED SECTION	2 ems
WARNING BOX LINE, HAIRLINES	.5


Job Information	Color Legend				
Date 12/16/03 PASS #4 Component # U0170-30-N2 Drawing # 5691 Rev2					
Identification					
Wyeth	PROCESS YELLOW	PROCESS BLACK	PANTONE 3005 C	PANTONE 199 C	
Consumer Healthcare					
Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048					
Vendor Identification	Р	rinter No	tes		
Production Packaging Dept. Cheng Lin SC: 7.1.3	PRINTER TO REPLI PRINTER TO SUPPI TO PRESS RUN.				

DEPARTMENT	REVIE	DATE	with	
	Print Name	Signature		Change
Graphic Artist				
Packaging Engineer				
Labeling Compliance				
Medical				1
Regulatory Affairs				1
Marketing				1 🗂
Legal				1
Trademark				┧╒╗

	Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction which may include: ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.	Do Not Use if plastic wrap imprinted with "CHILDREN'S ADVIL SAFETY SEALED" is broken or missing.	rigl Dir	ht away. rections do not give more than dire	ected ■ shake well bef			ontrol Center
	Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vorniting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever Ask a doctor before use if the child has not been drinking fluids uccers belong problems	Children's		 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ repeat dose every 6-8 hours, if needed ■ do not use more than 4 times a day ■ measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension. ■ Dosing Chart 				
	 ■ lost a lot of fluid due to continued vomiting or diarrhea ■ problems or serious side effects from taking fever reducers or pain relievers 			Weight (lb)	Age (yr)	Dose (tsp)]	
	■ stomach problems that last or come back, such as heartburn, upset stomach or pain	Suspension		under 24 lb	under 2 yr	ask a doctor	'	
Ш	■ high blood pressure, heart or kidney disease or is taking a diuretic	Suspension		24-35 lb	2-3 yr	1 tsp	1 i	
ľ	Ask a doctor or pharmacist before use if the child is ■ under a doctor's care for any serious condition	Fever Reducer/Pain Reliever		36-47 lb	4-5 yr	1 1/2 tsp	1 -├ ──	
lı	■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer		ı	48-59 lb	6-8 yr	2 tsp	1. —	
<u>'</u> '	■ taking a prescription product for anticoagulation (blood thinning)	Active ingredient (in each 5 mL)Ibuprofen 100 mg	ı	60-71 lb	9-10 yr	2 1/2 tsp	11	
h	■ taking any other drug When using this product give with food or milk if upset stomach occurs	Uses temporarily:	ı	72-95 lb	11 yr	3 tsp	1	
 	Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. fever or pain gets worse or lasts more than 3 days	 ■ reduces fever ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches 		ore at 20-25°C (68-77°F)	One Dose Lasts 6-8 Hou	rs VIL or ask your pharmacis	<u> </u>	
ľ	 the child does not get any relief within first day (24 hours) of treatment stomach pain or upset gets worse or lasts 	Fruit Flavored Liquid	dod	ctor or health care profest reth Consumer Healthcare	sional	, ,) ^{1,}	
//	■ redness or swelling is present in the painful area ■ any new symptoms appear ■ — — — — — — — — — — — — — — — — — — —	4 FL OZ (120 mL)		S. Patent No. 4,788,220		U0 <u>17</u> 0-30-N2	2	

150%

	RELEASED
(1)	Date
(2)	Date



Job Information Color Legend		
Date 12/18/03 PASS #1 Component # U0174-30-N2 Drawing # 5691 Rev.2		
Identification		
Wyeth Consumer Healthcare	PROCESS PROCESS PANTONE 3005 C	
Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048		
Vendor Identification	Vendor Identification Printer Notes	
Production Packaging Dept. Cheng Lin SC: 7.15.3	PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN.	

5 pt

3.5

2 ems

LEADING

• BULLETS

• SPACE AFTER BULLETED SECTION

WARNING BOX LINE HAIRLINES.

DEPARTMENT	REVIEWED BY		DATE	Wi
	Print Name	Signature		Chang
Graphic Artist				
Packaging Engineer				
Labeling Compliance				
Medical				
Regulatory Affairs				1
Marketing				1
Legal				7 🗂
Trademark				

	RELEASED
(1)	Date
(2)	Date

DIGITAL MECHANICAL COLOR REPRESENTATION MAY BE OFF FOLLOW COLOR LEGEND FOR ACTUAL COLOR.

Do Not Use if plastic wrap imprint	ed wit
"CHILDREN'S ADVIL SAFETY SE	ALED'
is broken or missing.	



Active ingredient (in each 5 mL)......lbuprofen 100 mg

reduces fever

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

problems or serious side effects from taking fever reducers or pain relievers stomach problems that last or come back, such as heartburn, upset stomach or pain

■ high blood pressure, heart or kidney disease or is taking a diuretic Ask a doctor or pharmacist before use if the child is

■ taking a prescription product for anticoagulation (blood thinning)

■ an allergic reaction occurs. Seek medical help right away.

When using this product give with food or milk if upset stomach occurs

■ fever or pain gets worse or lasts more than 3 days
 ■ the child does not get any relief within first day (24 hours) of treatment

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use

more than 2 days or administer to children under 3 years of age unless directed by doctor.

■ taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high

■ hives ■ asthma (wheezing) ■ facial swelling ■ shock

■ not been drinking fluids ■ ulcers ■ bleeding problems

■ lost a lot of fluid due to continued vomiting or diarrhea

under a doctor's care for any serious condition

stomach pain or upset gets worse or lasts

redness or swelling is present in the painful area

taking any other drug

Stop use and ask a doctor if

any new symptoms appear

Ask a doctor before use if the child has

relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Blue Raspberry Flavored Liquid 4 FL OZ (120 mL)

th Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions

- do not give more than directed shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed do not use more than 4 times a day measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-95 lb	11 yr	3 tsp

Store at 20-25°C (68-77°F)

Questions or comments? Call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

One Dose Lasts 6-8 Hours

Wyeth Consumer Healthcare, Madison, NJ 07940 Made in USA

U.S. Patent No. 4,788,220

U0174-30-N2

150%