

Table 35. Tuberculosis Cases and Percentages by Initial Drug Regimen: Reporting Areas, 2005

Reporting Area	Total Cases	Cases in Persons Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percentage of Cases in Persons with Initial Drug Regimen ^{1,2}			
			No.	(%)	IR	IRZ	IRZ,E/S	IRZE ³
United States	14,097	13,802	13,772	(99.8)	(1.2)	(5.5)	(83.5)	(83.4)
Alabama	216	213	213	(100.0)	(0.5)	(4.7)	(87.8)	(86.9)
Alaska	59	59	59	(100.0)	(3.4)	(10.2)	(81.4)	(81.4)
Arizona	281	276	270	(97.8)	(0.4)	(5.2)	(84.8)	(84.8)
Arkansas	114	109	109	(100.0)	(15.6)	(39.4)	(29.4)	(29.4)
California	2,904	2,860	2,859	(100.0)	(1.1)	(2.8)	(87.4)	(87.2)
Colorado	101	98	98	(100.0)	(0.0)	(5.1)	(79.6)	(79.6)
Connecticut	95	93	93	(100.0)	(1.1)	(8.6)	(81.7)	(81.7)
Delaware	26	26	26	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
District of Columbia	56	55	55	(100.0)	(0.0)	(3.6)	(85.5)	(85.5)
Florida	1,094	1,067	1,067	(100.0)	(0.6)	(6.5)	(83.4)	(83.3)
Georgia	505	503	503	(100.0)	(0.4)	(4.8)	(77.3)	(77.3)
Hawaii	112	111	111	(100.0)	(4.5)	(8.1)	(74.8)	(74.8)
Idaho	23	23	22	(95.7)	(0.0)	(18.2)	(81.8)	(81.8)
Illinois	596	588	585	(99.5)	(0.3)	(6.2)	(83.8)	(83.8)
Indiana	146	145	145	(100.0)	(1.4)	(11.0)	(80.7)	(80.7)
Iowa	55	55	54	(98.2)	(0.0)	(7.4)	(88.9)	(88.9)
Kansas	60	59	59	(100.0)	(5.1)	(1.7)	(83.1)	(83.1)
Kentucky	124	120	120	(100.0)	(0.8)	(9.2)	(82.5)	(82.5)
Louisiana	257	246	245	(99.6)	(0.4)	(5.7)	(88.6)	(88.6)
Maine	17	17	17	(100.0)	(0.0)	(5.9)	(88.2)	(88.2)
Maryland	283	278	278	(100.0)	(2.2)	(1.8)	(91.7)	(91.7)
Massachusetts	265	260	260	(100.0)	(0.4)	(2.3)	(86.9)	(86.9)
Michigan	246	240	240	(100.0)	(5.0)	(19.6)	(65.0)	(65.0)
Minnesota	199	197	197	(100.0)	(0.0)	(6.6)	(87.3)	(87.3)
Mississippi	103	95	95	(100.0)	(1.1)	(8.4)	(77.9)	(77.9)
Missouri	108	103	103	(100.0)	(1.9)	(1.9)	(75.7)	(75.7)
Montana	10	9	9	(100.0)	(0.0)	(0.0)	(77.8)	(77.8)
Nebraska	35	33	33	(100.0)	(0.0)	(6.1)	(81.8)	(81.8)
Nevada	112	110	110	(100.0)	(0.0)	(0.0)	(94.5)	(94.5)
New Hampshire	4	4	4	(100.0)	(0.0)	(25.0)	(75.0)	(75.0)
New Jersey	485	471	470	(99.8)	(0.6)	(7.2)	(82.3)	(82.3)
New Mexico	39	37	36	(97.3)	(0.0)	(11.1)	(83.3)	(83.3)
New York State ³	305	298	298	(100.0)	(1.0)	(6.7)	(82.6)	(82.6)
New York City	984	968	968	(100.0)	(0.4)	(2.8)	(86.7)	(86.7)
North Carolina	329	322	322	(100.0)	(0.3)	(2.2)	(90.1)	(89.8)
North Dakota	6	6	6	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Ohio	260	251	251	(100.0)	(0.8)	(9.6)	(81.3)	(81.3)
Oklahoma	144	139	138	(99.3)	(10.9)	(15.9)	(61.6)	(61.6)
Oregon	103	103	103	(100.0)	(0.0)	(2.9)	(93.2)	(93.2)
Pennsylvania	325	319	319	(100.0)	(0.9)	(4.1)	(62.1)	(62.1)
Rhode Island	47	46	46	(100.0)	(0.0)	(2.2)	(76.1)	(76.1)
South Carolina	261	252	252	(100.0)	(0.8)	(12.7)	(77.8)	(77.8)
South Dakota	16	16	16	(100.0)	(0.0)	(18.8)	(75.0)	(75.0)
Tennessee	298	291	291	(100.0)	(0.3)	(5.5)	(82.8)	(82.8)
Texas	1,535	1,491	1,479	(99.2)	(1.8)	(5.3)	(84.5)	(84.5)
Utah	29	29	29	(100.0)	(0.0)	(0.0)	(93.1)	(93.1)
Vermont	8	8	8	(100.0)	(0.0)	(75.0)	(25.0)	(25.0)
Virginia	355	349	348	(99.7)	(0.9)	(5.5)	(88.2)	(88.2)
Washington	256	250	249	(99.6)	(0.0)	(2.8)	(90.8)	(90.8)
West Virginia	28	28	28	(100.0)	(10.7)	(3.6)	(67.9)	(67.9)
Wisconsin	78	76	76	(100.0)	(2.6)	(6.6)	(80.3)	(80.3)
Wyoming	0	0	0	—	—	—	—	—
American Samoa ⁴	5	5	5	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Fed. States of Micronesia ⁴	74	74	74	(100.0)	(1.4)	(16.2)	(73.0)	(73.0)
Guam ⁴	64	63	63	(100.0)	(0.0)	(3.2)	(87.3)	(87.3)
Marshall Islands ⁴	66	66	66	(100.0)	(0.0)	(1.5)	(93.9)	(93.9)
N. Mariana Islands ⁴	56	56	56	(100.0)	(0.0)	(0.0)	(98.2)	(98.2)
Puerto Rico ⁴	113	101	99	(98.0)	(0.0)	(3.0)	(91.9)	(91.9)
Republic of Palau ⁴	10	10	10	(100.0)	(0.0)	(20.0)	(80.0)	(80.0)
U.S. Virgin Islands ⁴

¹Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

³Streptomycin is no longer considered a first-line drug. CDC. Treatment of Tuberculosis. American Thoracic Society, CDC, and Infectious Diseases Society of America. MMWR 2003;52(No. RR-11): 1–77.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Excluding cases with no information on drug regimen, 406 (2.9%) persons were not started on any drugs, 7 (<0.1%) were started on one drug, and 1,189 (8.4%) had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.

Ellipses indicate data not available.