

Carex arcta. (A) Inflorescences, (B) pistillate scales, (C) perigynia, (D) achenes. B through D: Left and center—dorsal views; right—ventral view.

HABIT: Plants densely tufted from very short rhizomes. **Culms:** Sharply triangular, 1-8 dm tall, shorter than the leaves, slightly **aphyllopodic**.

LEAVES: Clustered toward the base. **Blades:** Flat, light green, 1.5-4 mm wide. **Sheaths:** Thin and hyaline ventrally, brownish-tinged at the truncate mouth; lower sheaths at least sparsely red-dotted ventrally.

BRACTS: Lowest sheathless, setaceous-prolonged to 5 cm; upper short and inconspicuous.

SPIKES: 4-15, **gynaecandrous**, sessile, 5-10 mm long, crowded in a dense, pale green to greenish brown, oblong to elongate head 1.5-4 cm long; upper spikes crowded; lower spikes well-differentiated; **staminate** flowers inconspicuous; perigynia spreading-ascending to widely spreading at maturity.

PISTILLATE SCALES: Ovate with an acute tip, shorter than the perigynia, largely hyaline-scarious, brownish to nearly colorless with a firm, light brownish green midvein.

PERIGYNIA: Ovate or lance-ovate, rounded and somewhat spongy at the stipitate base, plano-convex, thin-edged, not winged, distally serrulate from above the middle, greenish-stramineous with greener margins, becoming greenish brown at maturity, 2-3.4 mm long, 1-1.2 mm wide. **Nerves:** Several dorsally, few or none ventrally. **Beaks:** Tapered, flat, serrulate, bidentate, 0.6-1.2 mm long.

ACHENES: Obovate to ovate, lenticular, slightly tapered to a substipitate base, stramineous to brownish, 1.2-1.6 mm long, 0.8-1.1 mm wide. **Stigmas:** 2.

HABITAT AND DISTRIBUTION: Occasional. Streambanks, wet meadows, swampy woods, thickets, and other wet places. Lowlands to near timberline. Southern Yukon to Quebec, south to Washington, Oregon, northern California (coast ranges), northern and central Idaho, and northwestern Montana and east to Minnesota and New York. June-August.