

 $\label{eq:Carex} \textit{Carex echinata}. \ (A) \ Pistillate \ scales, \ (B) \ perigynia, \ (C) \ achenes, \ (D) \ inflorescences. \\ A \ through \ C: \ Left-dorsal \ view; \ right-ventral \ view.$ 

## Carex echinata Murray Spiny star sedge

SYNONYM: C. muricata L.

**HABIT:** Densely tufted. **Culms:** 1-6 dm tall, equaling to exceeding the leaves, somewhat **aphyllopodic.** 

**LEAVES:** All borne on the lower part of culm. **Blades:** Elongate, slender, light green, 1-3.3 mm wide. **Sheaths:** Thin, hyaline ventrally, slightly thickened, and shallowly concave at the mouth.

**BRACTS:** Lowest small and inconspicuous to setaceous-prolonged, sheathless; upper reduced and scale-like.

**SPIKES:** 3-6, green, turning to brown, clustered to remote, forming an elongated to oblong head. **Terminal: Gynaecandrous. Lateral: Gynaecandrous** or wholly **pistillate**; perigynia 5-15, crowded, widely spreading to deflexed at maturity.

**PISTILLATE SCALES:** Broadly ovate, acute or cuspidate at the tip, shorter and narrower than the perigynia, hyaline or scarious, brownish tinged, greenish along the firm midvein.

**PERIGYNIA:** Lance-triangular to lance-ovate, plano-convex, thin-walled or subcoriaceous with abundant spongy tissue at the base, sharp edged, serrulate distally, green or tan, 1.6-3.5(4.0) mm long, 0.8-2 mm wide. **Nerves:** Few to many dorsally, few or absent ventrally, prominent on the margins. **Beaks:** Slender, abruptly tapered, greener than the body, bidentate, 1.1-1.6 mm long.

**ACHENE:** Ovate, lenticular, substipitate, golden yellow at maturity, 1.1-1.6 mm long, 0.8-1.3 mm wide. **Stigmas:** 2.

**HABITAT AND DISTRIBUTION:** Swamps, sphagnum bogs, and other wet places from **sea level to moderate elevations**. Circumboreal. In North America extending south to central Oregon and California and east to Utah, Colorado, Wisconsin, and North Carolina. **May-August.** 

**SIMILAR SPECIES:** *C. echinata*—Perigynium beak slender; longer (1.1-1.6 mm long), equaling the body, thin-walled, bidentate; style slender. *C. interior*—Perigynium beak broader and shorter (0.5-1 mm long), thick-walled, very shallowly bidentate; style stout.