

TABLE 1
 THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
 FROM HUMAN SOURCES REPORTED TO CDC IN 1997 AND FROM
 NONHUMAN SOURCES REPORTED TO CDC AND USDA IN 1997

HUMAN 1997				NONHUMAN 1997			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	TYPHIMURIUM *	9116	26.3	1	TYPHIMURIUM *	3717	20.8
2	ENTERITIDIS	7924	22.9	2	HEIDELBERG	1916	10.7
3	HEIDELBERG	2104	6.1	3	KENTUCKY	939	5.3
4	NEWPORT	1584	4.6	4	ANATUM	756	4.2
5	AGONA	740	2.1	5	DERBY	670	3.8
6	MONTEVIDEO	718	2.1	6	AGONA	652	3.7
7	THOMPSON	695	2.0	7	SENFENBERG	648	3.6
8	JAVIANA	675	2.0	8	HADAR	631	3.5
9	INFANTIS	651	1.9	9	DUBLIN	615	3.4
10	HADAR	643	1.9	10	MBANDAKA	599	3.4
11	ORANIENBURG	623	1.8	11	ENTERITIDIS	564	3.2
12	BRAENDERUP	559	1.6	12	BREDENEY	463	2.6
13	MUENCHEN	543	1.6	13	MONTEVIDEO	450	2.5
14	SAINTPAUL	436	1.3	14	CHOLERAESUIS **	396	2.2
15	TYPHI	349	1.0	15	INFANTIS	389	2.2
16	POONA	293	0.8	16	MUENSTER	334	1.9
17	ANATUM	208	0.6	17	SCHWARZENGRUND	272	1.5
18	MISSISSIPPI	205	0.6	18	WORTHINGTON	249	1.4
19	MBANDAKA	189	0.5	19	SAINTPAUL	219	1.2
20	JAVA	184	0.5	20	OHIO	197	1.1
SUB TOTAL		28439	82.2			14676	82.3
TOTAL		34608				17829	

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN

** CHOLERAESUIS INCLUDES VAR. KUNZENDORF