

Appendix A.

Table A. Listing of all benthic macrofauna collected by Van Veen grab in Newark Bay from June 1993 to March 1994, abundance and weight (mg) in each sample. Scientific names are as given by Cove Corporation.

Sample month	Station	Abundance	Weight, mg	Taxon
1	B01	1	1.3	<i>Leitoscoloplos</i> sp.
1	B01	5	23.5	<i>Spio setosa</i>
1	B01	4	53.4	<i>Marenzelleria viridis</i>
1	B01	0	3.6	Annelida fragments
1	B01	28	2.4	Oligochaeta
1	B01	1	139.3	<i>Mulinia lateralis</i>
1	B01	1	0	<i>Tellina agilis</i>
1	B01	8	218.4	<i>Mya arenaria</i>
1	B01	1	1.6	<i>Neomysis americana</i>
1	B01	1	2	<i>Monoculodes</i> sp. 1
1	B01	1	447.6	<i>Molgula manhattensis</i>
1	B02	20	8.4	<i>Hypereteone heteropoda</i>
1	B02	1	0	<i>Proceraea cornuta</i>
1	B02	1	0	<i>Streptosyllis pettiboneae</i>
1	B02	1	35.5	<i>Neanthes succinea</i>
1	B02	22	112.6	<i>Leitoscoloplos robustus</i>
1	B02	102	12.6	<i>Polydora cornuta</i>
1	B02	4	3.5	<i>Spio setosa</i>
1	B02	1152	146.2	<i>Streblospio benedicti</i>
1	B02	166	71.5	<i>Tharyx</i> sp. A
1	B02	1	0	<i>Heteromastus filiformis</i>
1	B02	61	5.1	<i>Mediomastus ambiseta</i>
1	B02	40	184.6	<i>Sabellaria vulgaris</i>
1	B02	1	6.2	<i>Pectinaria gouldii</i>
1	B02	0	47.4	Annelida fragments
1	B02	259	23.8	Oligochaeta
1	B02	2	1.9	<i>Odostomia</i> sp.
1	B02	8	206.2	<i>Mytilus edulis</i>
1	B02	1	10.4	<i>Mulinia lateralis</i>
1	B02	29	7161.4	<i>Mya arenaria</i>
1	B02	2	0	<i>Balanus improvisus</i>
1	B02	1	0	<i>Cyathura polita</i>
1	B02	21	16	<i>Edotea triloba</i>
1	B02	2	0	<i>Microdeutopus</i> spp.
1	B02	4	3.6	<i>Unciola serrata</i>
1	B02	6	0	<i>Parapleustes aestuarius</i>
1	B02	1	0	<i>Paracaprella tenuis</i>
1	B02	7	117.9	<i>Rhithropanopeus harrisi</i>
1	B02	66	2422	<i>Molgula manhattensis</i>
1	B03	3	1.2	<i>Hypereteone heteropoda</i>
1	B03	9	18.9	<i>Leitoscoloplos</i> sp.

Sample month	Station	Abundance	Weight, mg	Taxon
1	B03	8	0	<i>Polydora cornuta</i>
1	B03	1	2	<i>Spio setosa</i>
1	B03	168	72.3	<i>Streblospio benedicti</i>
1	B03	4	1.5	<i>Tharyx</i> sp. A
1	B03	16	1.8	<i>Mediomastus ambiseta</i>
1	B03	1	193.2	<i>Pectinaria gouldii</i>
1	B03	4	30	<i>Asabellides oculata</i>
1	B03	0	1.5	Annelida fragments
1	B03	7	0	Oligochaeta
1	B03	5	288.4	<i>Mulinia lateralis</i>
1	B03	2	0	<i>Macoma balthica</i>
1	B03	2	2387.1	<i>Mya arenaria</i>
1	B03	3	2.4	<i>Neomysis americana</i>
1	B03	2	0	<i>Leucon americanus</i>
1	B03	4	0	<i>Edotea triloba</i>
1	B03	1	3.4	<i>Ampelisca abdita</i>
1	B03	2	0	<i>Monoculodes</i> sp. 1
1	B04	13	8.9	<i>Hypereteone heteropoda</i>
1	B04	1	12.9	<i>Glycera americana</i>
1	B04	14	38.1	<i>Leitoscoloplos robustus</i>
1	B04	18	1.5	<i>Polydora cornuta</i>
1	B04	300	86.1	<i>Streblospio benedicti</i>
1	B04	17	3.4	<i>Tharyx</i> sp. A
1	B04	43	4.8	<i>Mediomastus ambiseta</i>
1	B04	2	252.9	<i>Pectinaria gouldii</i>
1	B04	12	43.5	<i>Asabellides oculata</i>
1	B04	0	14.4	Annelida fragments
1	B04	13	1	Oligochaeta
1	B04	7	4.5	<i>Odostomia</i> sp.
1	B04	1	12	<i>Mytilus edulis</i>
1	B04	5	280.5	<i>Mulinia lateralis</i>
1	B04	1	1.1	<i>Ensis directus</i>
1	B04	1	0	<i>Macoma balthica</i>
1	B04	1	0	<i>Tellina agilis</i>
1	B04	31	3382.9	<i>Mya arenaria</i>
1	B04	3	5.4	<i>Balanus improvisus</i>
1	B04	5	0	<i>Leucon americanus</i>
1	B04	2	0	<i>Oxyurostylis smithi</i>
1	B04	6	0	<i>Edotea triloba</i>
1	B04	5	2.2	<i>Molgula manhattensis</i>
1	B05	6	3.5	<i>Hypereteone heteropoda</i>
1	B05	4	168.2	<i>Neanthes succinea</i>
1	B05	31	199.1	<i>Leitoscoloplos robustus</i>
1	B05	35	4.9	<i>Polydora cornuta</i>
1	B05	4	6.6	<i>Spio setosa</i>
1	B05	71	35.9	<i>Streblospio benedicti</i>
1	B05	22	6.2	<i>Tharyx</i> sp. A

Sample month	Station	Abundance	Weight, mg	Taxon
1	B05	56	4.7	<i>Mediomastus ambiseta</i>
1	B05	2	12.3	<i>Sabellaria vulgaris</i>
1	B05	22	81.8	<i>Asabellides oculata</i>
1	B05	0	15.2	Annelida fragments
1	B05	20	1.4	Oligochaeta
1	B05	19	11.2	<i>Odostomia</i> sp.
1	B05	22	762.6	<i>Mulinia lateralis</i>
1	B05	1	1.6	<i>Ensis directus</i>
1	B05	39	38551	<i>Mya arenaria</i>
1	B05	1	8.7	<i>Balanus improvisus</i>
1	B05	6	1.6	<i>Neomysis americana</i>
1	B05	7	1.4	<i>Edotea triloba</i>
1	B05	2	0	<i>Microdeutopus</i> spp.
1	B05	10	16.2	<i>Unciola serrata</i>
1	B05	3	625.3	<i>Crangon septemspinosa</i>
1	B05	6	2.4	<i>Rhithropanopeus harrisi</i>
1	B06	1	0	Nereididae
1	B06	13	50.6	<i>Leitoscoloplos robustus</i>
1	B06	6	5.5	<i>Polydora cornuta</i>
1	B06	66	42.3	<i>Streblospio benedicti</i>
1	B06	3	0.9	<i>Tharyx</i> sp. A
1	B06	6	0	<i>Mediomastus ambiseta</i>
1	B06	7	17.7	<i>Asabellides oculata</i>
1	B06	0	6.6	Annelida fragments
1	B06	1	0	Oligochaeta
1	B06	4	1.5	<i>Odostomia</i> sp.
1	B06	9	332.3	<i>Mulinia lateralis</i>
1	B06	1	655	<i>Mya arenaria</i>
1	B06	1	2.3	<i>Neomysis americana</i>
1	B06	1	0	<i>Oxyurostylis smithi</i>
1	B06	1	0	<i>Unciola serrata</i>
1	B06	1	0	<i>Monoculodes</i> sp. 1
1	B06	1	0	<i>Crangon septemspinosa</i>
1	B06	1	0	Xanthidae
1	B06	1	0	<i>Molgula manhattensis</i>
1	B07	1	1.2	<i>Hypereteone heteropoda</i>
1	B07	9	43.4	<i>Leitoscoloplos robustus</i>
1	B07	2	0	<i>Polydora cornuta</i>
1	B07	50	12.3	<i>Streblospio benedicti</i>
1	B07	1	2	<i>Tharyx</i> sp. A
1	B07	9	1	<i>Mediomastus ambiseta</i>
1	B07	1	110.5	<i>Pectinaria gouldii</i>
1	B07	5	1.2	<i>Asabellides oculata</i>
1	B07	0	0.1	Annelida fragments
1	B07	1	0	Oligochaeta
1	B07	2	0	<i>Leucon americanus</i>
1	B07	1	0	<i>Edotea triloba</i>

Sample month	Station	Abundance	Weight, mg	Taxon
1	B07	1	1.5	<i>Crangon septemspinosa</i>
1	B08	1	1	<i>Paranaitis speciosa</i>
1	B08	1	1.7	<i>Leitoscoloplos</i> sp.
1	B08	3	0.8	<i>Polydora cornuta</i>
1	B08	1	0	<i>Streblospio benedicti</i>
1	B08	1	1.2	<i>Sabellaria vulgaris</i>
1	B08	1	1	<i>Mulinia lateralis</i>
1	B08	1	0.5	<i>Tellina agilis</i>
1	B08	1	2.8	<i>Mya arenaria</i>
1	B08	1	0	<i>Leucon americanus</i>
1	B08	1	6.6	<i>Cancer irroratus</i>
1	B09	36	231.3	<i>Leitoscoloplos robustus</i>
1	B09	10	1.4	<i>Streblospio benedicti</i>
1	B09	1	24.1	<i>Pectinaria gouldii</i>
1	B09	0	0.1	Annelida fragments
1	B09	1	88.6	<i>Mulinia lateralis</i>
1	B09	30	5.9	<i>Leucon americanus</i>
1	B10	12	88.3	<i>Leitoscoloplos robustus</i>
1	B10	11	3.2	<i>Streblospio benedicti</i>
1	B10	1	0	<i>Mediomastus ambiseta</i>
1	B10	1	204.2	<i>Pectinaria gouldii</i>
1	B10	3	5.5	<i>Asabellides oculata</i>
1	B10	1	1.9	<i>Macoma balthica</i>
1	B10	1	892.8	<i>Mya arenaria</i>
1	B10	9	2.9	<i>Leucon americanus</i>
1	B10	5	1.3	<i>Edotea triloba</i>
1	B11	1	31.6	<i>Neanthes succinea</i>
1	B11	47	3.4	<i>Streblospio benedicti</i>
1	B11	2	0	Oligochaeta
1	B11	1	4.8	<i>Mya arenaria</i>
1	B11	2	0	<i>Neomysis americana</i>
1	B12	1	0	<i>Leitoscoloplos</i> sp.
1	B12	54	21.8	<i>Streblospio benedicti</i>
1	B12	11	3.5	<i>Tharyx</i> sp. A
1	B12	2	0	Oligochaeta
1	B12	2	2.2	<i>Odostomia</i> sp.
1	B12	3	8	<i>Mulinia lateralis</i>
1	B12	3	1.9	<i>Macoma balthica</i>
1	B12	14	1.2	<i>Leucon americanus</i>
1	B13	1	0	<i>Hypereteone heteropoda</i>
1	B13	18	179.8	<i>Leitoscoloplos robustus</i>
1	B13	88	27.5	<i>Streblospio benedicti</i>
1	B13	47	17.7	<i>Tharyx</i> sp. A
1	B13	12	1.3	<i>Mediomastus ambiseta</i>
1	B13	8	0	Oligochaeta
1	B13	1	1.2	<i>Macoma balthica</i>
1	B13	1	941.3	<i>Mya arenaria</i>

Sample month	Station	Abundance	Weight, mg	Taxon
1	B13	6	0	<i>Neomysis americana</i>
1	B13	19	2.9	<i>Leucon americanus</i>
1	B13	2	0	<i>Edotea triloba</i>
1	B13	2	0	<i>Monoculodes</i> sp. 1
1	B13	1	669.3	<i>Molgula manhattensis</i>
1	B14	2	25.8	<i>Leitoscoloplos robustus</i>
1	B14	10	2.2	<i>Streblospio benedicti</i>
1	B14	1	193.2	<i>Pectinaria gouldii</i>
1	B14	2	0	Oligochaeta
1	B14	1	0	<i>Macoma balthica</i>
1	B14	3	0	<i>Neomysis americana</i>
1	B14	5	0	<i>Leucon americanus</i>
1	B15	1	4.1	<i>Hypereteone heteropoda</i>
1	B15	1	3.9	<i>Nephtys incisa</i>
1	B15	5	31.9	<i>Leitoscoloplos robustus</i>
1	B15	10	1.4	<i>Streblospio benedicti</i>
1	B15	1	1.6	<i>Macoma balthica</i>
1	B15	1	0.5	<i>Tellina agilis</i>
1	B15	3	1405.9	<i>Mya arenaria</i>
1	B15	16	3.1	<i>Leucon americanus</i>
1	B16	2	0	<i>Neomysis americana</i>
1	B16	5	2.2	<i>Leucon americanus</i>
1	B17	1	1.3	<i>Leitoscoloplos robustus</i>
1	B17	1	9.9	<i>Polydora cornuta</i>
1	B17	17	4.1	<i>Streblospio benedicti</i>
1	B17	2	0	Ampharetidae
1	B17	2	0	<i>Neomysis americana</i>
1	B17	13	1.8	<i>Leucon americanus</i>
1	B18	1	2.5	<i>Hypereteone heteropoda</i>
1	B18	1	35.2	<i>Neanthes succinea</i>
1	B18	8	19.2	<i>Leitoscoloplos robustus</i>
1	B18	1	0	<i>Polydora cornuta</i>
1	B18	16	2	<i>Streblospio benedicti</i>
1	B18	1	9.3	<i>Asabellides oculata</i>
1	B18	1	0	<i>Macoma balthica</i>
1	B18	1	0	<i>Mya arenaria</i>
1	B18	21	3.7	<i>Leucon americanus</i>
1	B18	1	0	<i>Edotea triloba</i>
1	B18	1	0	<i>Monoculodes</i> sp. 1
1	V1	2	0	<i>Hypereteone heteropoda</i>
1	V1	4	0	<i>Microphthalmus szcelkowi</i>
1	V1	1	0	<i>Streptosyllis pettiboneae</i>
1	V1	2	0	Nereididae
1	V1	24	392.4	<i>Leitoscoloplos robustus</i>
1	V1	27	10.6	<i>Polydora cornuta</i>
1	V1	15	24	<i>Spio setosa</i>
1	V1	487	63.5	<i>Streblospio benedicti</i>

Sample month	Station	Abundance	Weight, mg	Taxon
1	V1	4	6.6	<i>Marenzelleria viridis</i>
1	V1	16	3.8	<i>Tharyx</i> sp. A
1	V1	34	2.4	<i>Mediomastus ambiseta</i>
1	V1	0	44.2	Annelida fragments
1	V1	8	18.3	Oligochaeta
1	V1	1	0	<i>Odostomia</i> sp.
1	V1	1	0	<i>Mya arenaria</i>
1	V1	18	39.4	<i>Balanus improvisus</i>
1	V1	1	0	<i>Neomysis americana</i>
1	V2	2	0	<i>Hypereteone heteropoda</i>
1	V2	2	111.4	<i>Neanthes succinea</i>
1	V2	12	30.6	<i>Leitoscoloplos</i> sp.
1	V2	18	5.4	<i>Polydora cornuta</i>
1	V2	2	0	<i>Spio setosa</i>
1	V2	403	96	<i>Streblospio benedicti</i>
1	V2	2	9.5	<i>Marenzelleria viridis</i>
1	V2	2	26.2	<i>Heteromastus filiformis</i>
1	V2	58	5	<i>Mediomastus ambiseta</i>
1	V2	1	138	<i>Pectinaria gouldii</i>
1	V2	2	0	Oligochaeta
1	V2	4	1.9	<i>Odostomia</i> sp.
1	V2	1	0	<i>Doradella obscura</i>
1	V2	1	0	<i>Macoma balthica</i>
1	V2	1	0	<i>Tellina agilis</i>
1	V2	2	1605.4	<i>Mya arenaria</i>
1	V2	94	355.9	<i>Balanus improvisus</i>
1	V2	1	0	<i>Edotea triloba</i>
1	V2	3	5.4	<i>Melita nitida</i>
1	V3	1	0	<i>Hypereteone heteropoda</i>
1	V3	1	9	<i>Neanthes succinea</i>
1	V3	36	615.4	<i>Leitoscoloplos robustus</i>
1	V3	2	3	<i>Polydora cornuta</i>
1	V3	22	6.7	<i>Streblospio benedicti</i>
1	V3	1	1.1	<i>Tharyx</i> sp. A
1	V3	18	2.1	<i>Mediomastus ambiseta</i>
1	V3	0	1.1	Annelida fragments
1	V3	1	0	Oligochaeta
1	V3	9	5.6	<i>Odostomia</i> sp.
1	V3	1	136.7	<i>Mulinia lateralis</i>
1	V3	1	2.2	<i>Macoma balthica</i>
1	V3	1	0	<i>Monoculodes</i> sp. 1
1	V4	1	0	<i>Hypereteone heteropoda</i>
1	V4	23	244	<i>Leitoscoloplos robustus</i>
1	V4	9	2.1	<i>Streblospio benedicti</i>
1	V4	10	0	<i>Mediomastus ambiseta</i>
1	V4	1	0	Oligochaeta
1	V4	2	1.8	<i>Odostomia</i> sp.

Sample month	Station	Abundance	Weight, mg	Taxon
1	V4	5	977.5	<i>Mulinia lateralis</i>
1	V4	1	0	<i>Macoma balthica</i>
1	V4	1	1263.7	<i>Mya arenaria</i>
1	V4	1	0	<i>Neomysis americana</i>
1	V5	1	0	<i>Hypereteone heteropoda</i>
1	V5	46	888.8	<i>Leitoscoloplos robustus</i>
1	V5	7	3	<i>Streblospio benedicti</i>
1	V5	1	0	<i>Tharyx</i> sp. A
1	V5	2	0	<i>Odostomia</i> sp.
1	V5	20	3397.3	<i>Mulinia lateralis</i>
1	V5	1	2250.2	<i>Mya arenaria</i>
1	V5	4	0	<i>Leucon americanus</i>
1	V5	1	0	<i>Oxyurostylis smithi</i>
1	V6	95	1394.4	<i>Leitoscoloplos robustus</i>
1	V6	2	5.4	<i>Polydora cornuta</i>
1	V6	1	4.5	<i>Marenzelleria viridis</i>
1	V6	1	0	<i>Tharyx</i> sp. A
1	V6	1	0	Oligochaeta
1	V6	1	1.2	<i>Odostomia</i> sp.
1	V6	22	5449.8	<i>Mulinia lateralis</i>
1	V6	2	53.8	<i>Macoma balthica</i>
1	V6	1	3358.3	<i>Mya arenaria</i>
1	V6	1	0	<i>Leucon americanus</i>
1	V7	101	1461.8	<i>Leitoscoloplos robustus</i>
1	V7	3	0.5	<i>Streblospio benedicti</i>
1	V7	24	3708.4	<i>Mulinia lateralis</i>
1	V7	1	4.3	<i>Macoma balthica</i>
1	V7	1	0	<i>Leucon americanus</i>
1	V7	2	0.6	<i>Ampelisca abdita</i>
2	B01	2	0	<i>Hypereteone heteropoda</i>
2	B01	9	74.4	<i>Leitoscoloplos robustus</i>
2	B01	1	1.2	<i>Spio setosa</i>
2	B01	197	43.6	<i>Streblospio benedicti</i>
2	B01	194	143.5	<i>Tharyx</i> sp. A
2	B01	18	2	<i>Mediomastus ambiseta</i>
2	B01	0	2.9	Annelida fragments
2	B01	29	1.6	Oligochaeta
2	B01	5	11.4	<i>Mulinia lateralis</i>
2	B01	1	33.9	<i>Macoma balthica</i>
2	B01	2	4	<i>Mya arenaria</i>
2	B01	2	0.7	<i>Leucon americanus</i>
2	B01	1	13.9	<i>Cyathura polita</i>
2	B01	2	0.6	<i>Edotea triloba</i>
2	B02	2	0	<i>Hypereteone heteropoda</i>
2	B02	1	0	Nereididae
2	B02	4	15.1	<i>Leitoscoloplos</i> sp.
2	B02	1	0	<i>Polydora cornuta</i>

Sample month	Station	Abundance	Weight, mg	Taxon
2	B02	1	3.7	<i>Spio setosa</i>
2	B02	360	93.4	<i>Streblospio benedicti</i>
2	B02	89	48.4	<i>Tharyx</i> sp. A
2	B02	1	7.1	<i>Heteromastus filiformis</i>
2	B02	2	0.5	<i>Mediomastus ambiseta</i>
2	B02	1	0	<i>Sabellaria vulgaris</i>
2	B02	0	5.2	Annelida fragments
2	B02	43	2.6	Oligochaeta
2	B02	1	0	<i>Odostomia</i> sp.
2	B02	1	443.3	<i>Mya arenaria</i>
2	B02	1	0.6	<i>Balanus improvisus</i>
2	B02	1	0	<i>Melita nitida</i>
2	B03	2	97.7	Anthozoa
2	B03	4	0.5	<i>Hypereteone heteropoda</i>
2	B03	1	499.1	<i>Glycera americana</i>
2	B03	7	17.1	<i>Leitoscoloplos robustus</i>
2	B03	6	0	<i>Polydora cornuta</i>
2	B03	9	42.6	<i>Spio setosa</i>
2	B03	487	188.5	<i>Streblospio benedicti</i>
2	B03	506	360.7	<i>Tharyx</i> sp. A
2	B03	85	5.9	<i>Mediomastus ambiseta</i>
2	B03	2	0	<i>Sabellaria vulgaris</i>
2	B03	4	18.3	<i>Asabellides oculata</i>
2	B03	0	46.3	Annelida fragments
2	B03	25	1	Oligochaeta
2	B03	5	1.6	<i>Odostomia</i> sp.
2	B03	3	0	<i>Mulinia lateralis</i>
2	B03	4	352.8	<i>Mya arenaria</i>
2	B03	3	59.1	<i>Balanus improvisus</i>
2	B03	4	1.1	<i>Edotea triloba</i>
2	B03	1	0	<i>Unciola serrata</i>
2	B03	1	0	<i>Melita nitida</i>
2	B03	1	118.2	<i>Rhithropanopeus harrisi</i>
2	B04	1	0.7	Rhynchozoela
2	B04	2	0.8	<i>Hypereteone heteropoda</i>
2	B04	23	180.7	<i>Leitoscoloplos robustus</i>
2	B04	207	48.6	<i>Streblospio benedicti</i>
2	B04	5	1.3	<i>Tharyx</i> sp. A
2	B04	1	0	<i>Sabellaria vulgaris</i>
2	B04	2	0	<i>Asabellides oculata</i>
2	B04	0	1.2	Annelida fragments
2	B04	3	0.9	Oligochaeta
2	B04	3	0.6	<i>Odostomia</i> sp.
2	B04	9	1.8	<i>Mulinia lateralis</i>
2	B04	2	338	<i>Mya arenaria</i>
2	B04	1	0	<i>Leucon americanus</i>
2	B05	12	485.3	Anthozoa

Sample month	Station	Abundance	Weight, mg	Taxon
2	B05	9	1.5	<i>Hypereteone heteropoda</i>
2	B05	1	0	Nereididae
2	B05	1	40.6	Glycera spp.
2	B05	1	0	<i>Glycera americana</i>
2	B05	22	43.4	<i>Leitoscoloplos robustus</i>
2	B05	13	0	<i>Polydora cornuta</i>
2	B05	684	207.4	<i>Streblospio benedicti</i>
2	B05	20	8.9	<i>Tharyx</i> sp. A
2	B05	325	54.7	<i>Mediomastus ambiseta</i>
2	B05	31	41.2	<i>Sabellaria vulgaris</i>
2	B05	2	187	<i>Pectinaria gouldii</i>
2	B05	2	5.1	<i>Asabellides oculata</i>
2	B05	0	11.5	Annelida fragments
2	B05	175	10.6	Oligochaeta
2	B05	3	489.1	<i>Mytilus edulis</i>
2	B05	1	20.3	<i>Mulinia lateralis</i>
2	B05	1	5.1	<i>Macoma balthica</i>
2	B05	8	2268.5	<i>Mya arenaria</i>
2	B05	3	1	<i>Edotea triloba</i>
2	B05	1	0	<i>Microdeutopus gryllotalpa</i>
2	B05	79	40.9	<i>Unciola serrata</i>
2	B05	1	0	<i>Paracaprella tenuis</i>
2	B05	6	248.8	<i>Rhithropanopeus harrisi</i>
2	B05	1	885	<i>Molgula manhattensis</i>
2	B06	19	62.5	<i>Leitoscoloplos robustus</i>
2	B06	5	3.8	<i>Polydora cornuta</i>
2	B06	91	27.3	<i>Streblospio benedicti</i>
2	B06	2	0.7	<i>Tharyx</i> sp. A
2	B06	10	0.4	<i>Mediomastus ambiseta</i>
2	B06	2	1.6	<i>Sabellaria vulgaris</i>
2	B06	1	5.9	<i>Asabellides oculata</i>
2	B06	0	0.9	Annelida fragments
2	B06	6	0.6	Oligochaeta
2	B06	4	0.6	<i>Odostomia</i> sp.
2	B06	3	274	<i>Mulinia lateralis</i>
2	B06	4	4619.3	<i>Mya arenaria</i>
2	B06	5	0.8	<i>Edotea triloba</i>
2	B06	1	96.7	<i>Crangon septemspinosa</i>
2	B07	6	19.4	<i>Leitoscoloplos robustus</i>
2	B07	7	1.4	<i>Streblospio benedicti</i>
2	B07	1	0	<i>Tharyx</i> sp. A
2	B07	2	149.1	<i>Mulinia lateralis</i>
2	B07	1	0	<i>Leucon americanus</i>
2	B07	1	0	<i>Microdeutopus gryllotalpa</i>
2	B08	31	166.2	<i>Leitoscoloplos robustus</i>
2	B08	35	6.1	<i>Streblospio benedicti</i>
2	B08	1	0	<i>Tharyx</i> sp. A

Sample month	Station	Abundance	Weight, mg	Taxon
2	B08	2	0	<i>Mediomastus ambiseta</i>
2	B08	1	0	<i>Pectinaria gouldii</i>
2	B08	1	0.5	Oligochaeta
2	B08	10	375.5	<i>Mulinia lateralis</i>
2	B08	1	0	<i>Leucon americanus</i>
2	B09	2	4.2	<i>Leitoscoloplos robustus</i>
2	B09	15	1.5	<i>Streblospio benedicti</i>
2	B09	1	0	<i>Odostomia</i> sp.
2	B09	3	0.8	<i>Rictaxis punctostriatus</i>
2	B09	2	0.7	<i>Mulinia lateralis</i>
2	B10	2	12.8	<i>Leitoscoloplos</i> sp.
2	B10	1	0	<i>Odostomia</i> sp.
2	B10	1	0	<i>Rictaxis punctostriatus</i>
2	B11	1	0.5	<i>Hypereteone heteropoda</i>
2	B11	1	1.8	<i>Leitoscoloplos</i> sp.
2	B11	86	20.2	<i>Streblospio benedicti</i>
2	B11	2	0.5	<i>Tharyx</i> sp. A
2	B11	1	0	Oligochaeta
2	B11	1	0	<i>Mulinia lateralis</i>
2	B11	1	0	<i>Leucon americanus</i>
2	B11	1	0	<i>Edotea triloba</i>
2	B11	1	281.9	<i>Rhithropanopeus harrisii</i>
2	B12	5	67.5	<i>Leitoscoloplos robustus</i>
2	B12	28	6	<i>Streblospio benedicti</i>
2	B12	24	19.5	<i>Tharyx</i> sp. A
2	B12	1	0	<i>Pectinaria gouldii</i>
2	B12	1	0	<i>Asabellides oculata</i>
2	B12	3	0	Oligochaeta
2	B12	3	0	<i>Mulinia lateralis</i>
2	B12	8	1.5	<i>Leucon americanus</i>
2	B12	1	0	<i>Edotea triloba</i>
2	B12	1	0	<i>Corophium acutum</i>
2	B13	2	76.1	<i>Leitoscoloplos robustus</i>
2	B13	80	20.2	<i>Streblospio benedicti</i>
2	B13	16	8	<i>Tharyx</i> sp. A
2	B13	6	1.6	<i>Mediomastus ambiseta</i>
2	B13	1	0	<i>Pectinaria gouldii</i>
2	B13	2	0	<i>Odostomia</i> sp.
2	B13	12	2.7	<i>Mulinia lateralis</i>
2	B13	1	0	<i>Mya arenaria</i>
2	B13	14	3.4	<i>Leucon americanus</i>
2	B14	1	802.3	<i>Glycera americana</i>
2	B14	2	59	<i>Leitoscoloplos robustus</i>
2	B14	2	0	<i>Streblospio benedicti</i>
2	B14	7	1.2	<i>Leucon americanus</i>
2	B14	1	0	<i>Edotea triloba</i>
2	B15	1	77	<i>Nephtys incisa</i>

Sample month	Station	Abundance	Weight, mg	Taxon
2	B15	7	72.6	<i>Leitoscoloplos robustus</i>
2	B15	6	0.6	<i>Streblospio benedicti</i>
2	B15	1	0	<i>Tharyx</i> sp. A
2	B15	1	42.4	<i>Mulinia lateralis</i>
2	B15	1	0	<i>Neomysis americana</i>
2	B15	6	0.8	<i>Leucon americanus</i>
2	B16	1	0	Anthozoa
2	B16	3	29.5	<i>Leitoscoloplos robustus</i>
2	B16	5	0.6	<i>Streblospio benedicti</i>
2	B16	4	0	<i>Mulinia lateralis</i>
2	B16	1	0	<i>Tellina agilis</i>
2	B16	2	0.8	<i>Neomysis americana</i>
2	B16	1	0	<i>Leucon americanus</i>
2	B17	14	88.1	<i>Leitoscoloplos robustus</i>
2	B17	4	1.9	<i>Polydora cornuta</i>
2	B17	29	4	<i>Streblospio benedicti</i>
2	B17	1	0	<i>Tharyx</i> sp. A
2	B17	1	0	<i>Mediomastus ambiseta</i>
2	B17	1	0	<i>Rictaxis punctostriatus</i>
2	B17	2	197.8	<i>Mulinia lateralis</i>
2	B17	1	370.7	<i>Mya arenaria</i>
2	B17	7	0.8	<i>Leucon americanus</i>
2	B17	2	0.5	<i>Edotea triloba</i>
2	B18	11	113.6	<i>Leitoscoloplos robustus</i>
2	B18	5	0.7	<i>Streblospio benedicti</i>
2	B18	1	0	<i>Mediomastus ambiseta</i>
2	B18	1	0	<i>Pectinaria gouldii</i>
2	B18	1	0	<i>Rictaxis punctostriatus</i>
2	B18	1	4	<i>Mulinia lateralis</i>
2	B18	8	2	<i>Leucon americanus</i>
2	V1	1	8.1	Rhynchocoela
2	V1	16	217.7	<i>Leitoscoloplos robustus</i>
2	V1	4	0.8	<i>Polydora cornuta</i>
2	V1	1	0.8	<i>Spio setosa</i>
2	V1	335	116.3	<i>Streblospio benedicti</i>
2	V1	5	58.2	<i>Marenzelleria viridis</i>
2	V1	1	1.7	<i>Heteromastus filiformis</i>
2	V1	50	4.4	<i>Mediomastus ambiseta</i>
2	V1	0	11.4	Annelida fragments
2	V1	3	1.1	<i>Odostomia</i> sp.
2	V1	17	138.7	<i>Mulinia lateralis</i>
2	V1	2	60.9	<i>Mya arenaria</i>
2	V1	1	2.6	<i>Neomysis americana</i>
2	V1	1	17.2	<i>Crangon septemspinosa</i>
2	V2	1	0	<i>Hypereteone heteropoda</i>
2	V2	47	384.5	<i>Leitoscoloplos robustus</i>
2	V2	4	7.3	<i>Polydora cornuta</i>

Sample month	Station	Abundance	Weight, mg	Taxon
2	V2	2	0	<i>Pygospio elegans</i>
2	V2	166	30.4	<i>Streblospio benedicti</i>
2	V2	1	13.8	<i>Marenzelleria viridis</i>
2	V2	2	0	<i>Tharyx</i> sp. A
2	V2	7	2.4	<i>Heteromastus filiformis</i>
2	V2	13	1.1	<i>Mediomastus ambiseta</i>
2	V2	0	2.5	Annelida fragments
2	V2	15	3.2	<i>Rictaxis punctostriatus</i>
2	V2	11	86.1	<i>Mulinia lateralis</i>
2	V2	1	22.1	<i>Macoma balthica</i>
2	V2	2	2915.1	<i>Mya arenaria</i>
2	V2	2	0	<i>Balanus improvisus</i>
2	V2	1	0	<i>Molgula manhattensis</i>
2	V3	1	88.5	<i>Neanthes succinea</i>
2	V3	32	363.8	<i>Leitoscoloplos robustus</i>
2	V3	1	1.2	<i>Polydora cornuta</i>
2	V3	187	60.4	<i>Streblospio benedicti</i>
2	V3	1	4.6	<i>Marenzelleria viridis</i>
2	V3	1	0	<i>Tharyx</i> sp. A
2	V3	3	0.9	<i>Heteromastus filiformis</i>
2	V3	28	1.6	<i>Mediomastus ambiseta</i>
2	V3	0	4.8	Annelida fragments
2	V3	1	0	<i>Odostomia</i> sp.
2	V3	1	0	<i>Rictaxis punctostriatus</i>
2	V3	1	0	<i>Mytilus edulis</i>
2	V3	13	854.8	<i>Mulinia lateralis</i>
2	V3	1	0	<i>Leucon americanus</i>
2	V4	47	610.8	<i>Leitoscoloplos robustus</i>
2	V4	1	1.1	<i>Polydora cornuta</i>
2	V4	37	8.3	<i>Streblospio benedicti</i>
2	V4	12	5.5	<i>Rictaxis punctostriatus</i>
2	V4	9	6.5	<i>Mulinia lateralis</i>
2	V4	1	41.7	<i>Macoma balthica</i>
2	V4	3	7210.8	<i>Mya arenaria</i>
2	V4	3	0	<i>Leucon americanus</i>
2	V4	1	0	<i>Ampelisca</i> sp.
2	V5	1	0.8	<i>Rhynchocoela</i>
2	V5	78	708.5	<i>Leitoscoloplos robustus</i>
2	V5	6	4	<i>Polydora cornuta</i>
2	V5	76	21.1	<i>Streblospio benedicti</i>
2	V5	3	8.8	<i>Heteromastus filiformis</i>
2	V5	3	0.5	<i>Mediomastus ambiseta</i>
2	V5	0	0.1	Annelida fragments
2	V5	1	0.5	<i>Oligochaeta</i>
2	V5	4	0.9	<i>Rictaxis punctostriatus</i>
2	V5	34	5895.5	<i>Mulinia lateralis</i>
2	V5	1	0.7	<i>Leucon americanus</i>

Sample month	Station	Abundance	Weight, mg	Taxon
2	V6	68	1267.5	<i>Leitoscoloplos robustus</i>
2	V6	1	0.6	<i>Polydora cornuta</i>
2	V6	44	11.8	<i>Streblospio benedicti</i>
2	V6	6	0	<i>Mediomastus ambiseta</i>
2	V6	1	0	<i>Asabellides oculata</i>
2	V6	3	0.6	<i>Oligochaeta</i>
2	V6	1	0	<i>Odostomia sp.</i>
2	V6	9	3	<i>Rictaxis punctostriatus</i>
2	V6	25	4932.8	<i>Mulinia lateralis</i>
2	V6	3	0	<i>Leucon americanus</i>
2	V7	1	936.5	<i>Glycera americana</i>
2	V7	58	719	<i>Leitoscoloplos robustus</i>
2	V7	34	7.2	<i>Streblospio benedicti</i>
2	V7	3	0.9	<i>Tharyx sp. A</i>
2	V7	0	11.6	<i>Annelida fragments</i>
2	V7	10	2.2	<i>Rictaxis punctostriatus</i>
2	V7	9	899.6	<i>Mulinia lateralis</i>
2	V7	2	0.5	<i>Neomysis americana</i>
3	B01	8	298.6	<i>Anthozoa</i>
3	B01	11	3.1	<i>Hypereteone heteropoda</i>
3	B01	3	0	<i>Eumida sanguinea</i>
3	B01	2	0	<i>Streptosyllis pettiboneae</i>
3	B01	2	16.8	<i>Neanthes succinea</i>
3	B01	4	9.3	<i>Glycera americana</i>
3	B01	29	94.1	<i>Leitoscoloplos robustus</i>
3	B01	9	6.1	<i>Polydora cornuta</i>
3	B01	180	65.1	<i>Streblospio benedicti</i>
3	B01	311	294.4	<i>Tharyx sp. A</i>
3	B01	21	1.3	<i>Mediomastus ambiseta</i>
3	B01	22	52.3	<i>Sabellaria vulgaris</i>
3	B01	2	253.9	<i>Pectinaria gouldii</i>
3	B01	1	9.5	<i>Asabellides oculata</i>
3	B01	0	30.4	<i>Annelida fragments</i>
3	B01	91	9.1	<i>Oligochaeta</i>
3	B01	33	13.2	<i>Odostomia sp.</i>
3	B01	1	1968.8	<i>Mytilus edulis</i>
3	B01	3	146.4	<i>Mulinia lateralis</i>
3	B01	17	1685.3	<i>Mya arenaria</i>
3	B01	1	1.4	<i>Balanus improvisus</i>
3	B01	4	5	<i>Edotea triloba</i>
3	B01	26	22.8	<i>Unciola serrata</i>
3	B01	1	38.4	<i>Rhithropanopeus harrisi</i>
3	B01	12	107.1	<i>Molgula manhattensis</i>
3	B02	1	0	<i>Hypereteone heteropoda</i>
3	B02	6	18.4	<i>Glycera americana</i>
3	B02	45	221.5	<i>Leitoscoloplos robustus</i>
3	B02	1	9.3	<i>Spio setosa</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B02	863	1078.7	<i>Tharyx sp. A</i>
3	B02	1	0.7	<i>Heteromastus filiformis</i>
3	B02	19	1.9	<i>Mediomastus ambiseta</i>
3	B02	1	0.9	<i>Sabellaria vulgaris</i>
3	B02	4	206.6	<i>Pectinaria gouldii</i>
3	B02	0	5	<i>Annelida fragments</i>
3	B02	62	4.5	<i>Oligochaeta</i>
3	B02	74	37.6	<i>Odostomia sp.</i>
3	B02	11	3.8	<i>Rictaxis punctostriatus</i>
3	B02	3	183.7	<i>Mulinia lateralis</i>
3	B02	1	293.4	<i>Macoma balthica</i>
3	B02	9	3.4	<i>Mya arenaria</i>
3	B02	2	0.6	<i>Leucon americanus</i>
3	B02	1	0	<i>Edotea triloba</i>
3	B02	1	0	<i>Monoculodes sp. 1</i>
3	B03	1	0	<i>Hypereteone heteropoda</i>
3	B03	1	0	<i>Podarkeopsis levifuscina</i>
3	B03	4	9.4	<i>Glycera americana</i>
3	B03	90	145.4	<i>Leitoscoloplos robustus</i>
3	B03	7	13.6	<i>Polydora cornuta</i>
3	B03	9	2.1	<i>Streblospio benedicti</i>
3	B03	6	1.3	<i>Tharyx sp. A</i>
3	B03	1	0	<i>Heteromastus filiformis</i>
3	B03	148	15.7	<i>Mediomastus ambiseta</i>
3	B03	13	1233	<i>Pectinaria gouldii</i>
3	B03	2	12.6	<i>Asabellides oculata</i>
3	B03	0	5.3	<i>Annelida fragments</i>
3	B03	23	1.6	<i>Oligochaeta</i>
3	B03	27	8.5	<i>Odostomia sp.</i>
3	B03	9	33.5	<i>Rictaxis punctostriatus</i>
3	B03	20	331.5	<i>Mulinia lateralis</i>
3	B03	1	0	<i>Ensis directus</i>
3	B03	1	125.9	<i>Macoma balthica</i>
3	B03	27	23	<i>Mya arenaria</i>
3	B03	9	1.2	<i>Leucon americanus</i>
3	B03	9	5.1	<i>Edotea triloba</i>
3	B03	2	0	<i>Unciola serrata</i>
3	B03	2	0	<i>Monoculodes sp. 1</i>
3	B04	1	5.4	<i>Lepidonotus squamatus</i>
3	B04	1	0	<i>Eumida sanguinea</i>
3	B04	1	0.9	<i>Neanthes succinea</i>
3	B04	1	14	<i>Glycera americana</i>
3	B04	22	4.5	<i>Polydora cornuta</i>
3	B04	1	0	<i>Heteromastus filiformis</i>
3	B04	66	457.1	<i>Sabellaria vulgaris</i>
3	B04	2	1.2	<i>Asabellides oculata</i>
3	B04	0	1.9	<i>Annelida fragments</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B04	10	5	<i>Odostomia sp.</i>
3	B04	1	5.8	<i>Rictaxis punctostriatus</i>
3	B04	1	859.2	<i>Mytilus edulis</i>
3	B04	19	4347.7	<i>Mya arenaria</i>
3	B04	14	1020.6	<i>Bivalve siphons</i>
3	B04	6	1.5	<i>Edotea triloba</i>
3	B04	18	12	<i>Unciola serrata</i>
3	B04	2	0.8	<i>Melita nitida</i>
3	B04	1	146.2	<i>Rhithropanopeus harrisi</i>
3	B04	6	8.9	<i>Molgula manhattensis</i>
3	B05	1	29.1	<i>Lepidonotus sublevis</i>
3	B05	5	1.3	<i>Hypereteone heteropoda</i>
3	B05	13	2.5	<i>Eumida sanguinea</i>
3	B05	1	18.3	<i>Neanthes succinea</i>
3	B05	2	1.1	<i>Glycera americana</i>
3	B05	11	8.1	<i>Leitoscoloplos robustus</i>
3	B05	12	2.2	<i>Polydora cornuta</i>
3	B05	447	288.2	<i>Streblospio benedicti</i>
3	B05	3	0.8	<i>Tharyx sp. A</i>
3	B05	20	1.1	<i>Mediomastus ambiseta</i>
3	B05	183	725.6	<i>Sabellaria vulgaris</i>
3	B05	2	25.4	<i>Pectinaria gouldii</i>
3	B05	3	13	<i>Asabellides oculata</i>
3	B05	1	0	<i>Sabellidae</i>
3	B05	0	7.8	<i>Annelida fragments</i>
3	B05	20	1.2	<i>Oligochaeta</i>
3	B05	38	10.4	<i>Odostomia sp.</i>
3	B05	5	0	<i>Ensis directus</i>
3	B05	42	2919.5	<i>Mya arenaria</i>
3	B05	1	0	<i>Neomysis americana</i>
3	B05	19	3.6	<i>Leucon americanus</i>
3	B05	1	0	<i>Sphaeroma quadridentata</i>
3	B05	5	2.2	<i>Edotea triloba</i>
3	B05	222	172.2	<i>Unciola serrata</i>
3	B05	1	0	<i>Elasmopus laevis</i>
3	B05	5	2.8	<i>Melita nitida</i>
3	B05	3	4.8	<i>Parapleustes aestuarius</i>
3	B05	1	127.5	<i>Crangon septemspinosa</i>
3	B05	2	66.5	<i>Rhithropanopeus harrisi</i>
3	B05	23	115.9	<i>Molgula manhattensis</i>
3	B06	2	0.9	<i>Hypereteone heteropoda</i>
3	B06	1	5.7	<i>Neanthes succinea</i>
3	B06	44	67.1	<i>Polydora cornuta</i>
3	B06	221	112.8	<i>Streblospio benedicti</i>
3	B06	1	0	<i>Tharyx sp. A</i>
3	B06	1	0.5	<i>Mediomastus ambiseta</i>
3	B06	1	0	<i>Oligochaeta</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B06	25	5.7	<i>Odostomia sp.</i>
3	B06	1	0	<i>Rictaxis punctostriatus</i>
3	B06	1	3.5	<i>Mulinia lateralis</i>
3	B06	1	0	<i>Ensis directus</i>
3	B06	14	3.8	<i>Mya arenaria</i>
3	B06	1	1.5	<i>Neomysis americana</i>
3	B06	1	0	<i>Leucon americanus</i>
3	B06	3	1.9	<i>Edotea triloba</i>
3	B07	3	1.7	<i>Rhynchocoela</i>
3	B07	2	10.7	<i>Glycera americana</i>
3	B07	93	92.1	<i>Leitoscoloplos sp.</i>
3	B07	4	0.7	<i>Streblospio benedicti</i>
3	B07	1	0	<i>Tharyx sp. A</i>
3	B07	19	2.5	<i>Mediomastus ambiseta</i>
3	B07	2	64	<i>Pectinaria gouldii</i>
3	B07	3	0.6	<i>Asabellides oculata</i>
3	B07	4	1.1	<i>Odostomia sp.</i>
3	B07	1	0	<i>Rictaxis punctostriatus</i>
3	B07	9	240.8	<i>Mulinia lateralis</i>
3	B07	2	1	<i>Mya arenaria</i>
3	B07	20	4.5	<i>Leucon americanus</i>
3	B07	1	0	<i>Edotea triloba</i>
3	B07	1	0	<i>Microdeutopus gryllotalpa</i>
3	B07	1	0	<i>Unciola serrata</i>
3	B08	5	20.9	<i>Glycera americana</i>
3	B08	136	83.6	<i>Leitoscoloplos sp.</i>
3	B08	8	1.2	<i>Streblospio benedicti</i>
3	B08	7	1	<i>Tharyx sp. A</i>
3	B08	37	230.7	<i>Mediomastus ambiseta</i>
3	B08	14	1141	<i>Pectinaria gouldii</i>
3	B08	6	0	<i>Oligochaeta</i>
3	B08	5	1	<i>Odostomia sp.</i>
3	B08	15	38.8	<i>Rictaxis punctostriatus</i>
3	B08	9	578.7	<i>Mulinia lateralis</i>
3	B08	1	20.5	<i>Tellina agilis</i>
3	B08	1	0	<i>Mya arenaria</i>
3	B08	6	0.9	<i>Leucon americanus</i>
3	B09	1	0	<i>Hypereteone heteropoda</i>
3	B09	1	0	<i>Podarkeopsis levifuscina</i>
3	B09	1	74.2	<i>Nephtys incisa</i>
3	B09	3	31.7	<i>Glycera americana</i>
3	B09	36	41.4	<i>Leitoscoloplos robustus</i>
3	B09	2	0	<i>Tharyx sp. A</i>
3	B09	14	1.4	<i>Mediomastus ambiseta</i>
3	B09	2	55.1	<i>Pectinaria gouldii</i>
3	B09	1	0	<i>Asabellides oculata</i>
3	B09	0	0.6	<i>Annelida fragments</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B09	2	0	<i>Oligochaeta</i>
3	B09	2	0.6	<i>Odostomia sp.</i>
3	B09	4	1.1	<i>Rictaxis punctostriatus</i>
3	B09	1	112.6	<i>Mulinia lateralis</i>
3	B09	2	0	<i>Ensis directus</i>
3	B09	1	0	<i>Tellina agilis</i>
3	B09	1	0.5	<i>Mya arenaria</i>
3	B09	26	5.6	<i>Leucon americanus</i>
3	B09	5	2.5	<i>Edotea triloba</i>
3	B09	5	5.4	<i>Ampelisca abdita</i>
3	B09	1	0	<i>Unciola serrata</i>
3	B10	1	0	<i>Leitoscoloplos robustus</i>
3	B10	1	0	<i>Polydora cornuta</i>
3	B10	2	0	<i>Mediomastus ambiseta</i>
3	B10	3	4.2	<i>Sabellaria vulgaris</i>
3	B10	3	295.2	<i>Pectinaria gouldii</i>
3	B10	1	0	<i>Oligochaeta</i>
3	B10	4	0	<i>Rictaxis punctostriatus</i>
3	B10	42	9.4	<i>Leucon americanus</i>
3	B10	1	0	<i>Oxyurostylis smithi</i>
3	B10	1	0	<i>Unciola serrata</i>
3	B11	2	0	<i>Leitoscoloplos sp.</i>
3	B11	3	0	<i>Streblospio benedicti</i>
3	B11	1	0	<i>Tharyx sp. A</i>
3	B11	0	0.6	<i>Annelida fragments</i>
3	B11	1	0	<i>Oligochaeta</i>
3	B11	3	5.4	<i>Mulinia lateralis</i>
3	B11	1	0	<i>Leucon americanus</i>
3	B11	1	0	<i>Chironomidae</i>
3	B12	1	0	<i>Rhynchocoela</i>
3	B12	1	0	<i>Hypereteone heteropoda</i>
3	B12	1	1.4	<i>Glycera americana</i>
3	B12	51	261.8	<i>Leitoscoloplos robustus</i>
3	B12	1	0	<i>Streblospio benedicti</i>
3	B12	37	39	<i>Tharyx sp. A</i>
3	B12	1	0	<i>Mediomastus ambiseta</i>
3	B12	1	44.8	<i>Pectinaria gouldii</i>
3	B12	1	0	<i>Asabellides oculata</i>
3	B12	2	0	<i>Oligochaeta</i>
3	B12	3	1.3	<i>Odostomia sp.</i>
3	B12	2	13.1	<i>Rictaxis punctostriatus</i>
3	B12	7	9.7	<i>Mulinia lateralis</i>
3	B12	2	0.9	<i>Mya arenaria</i>
3	B12	31	8.3	<i>Leucon americanus</i>
3	B13	8	1.6	<i>Hypereteone heteropoda</i>
3	B13	1	13.2	<i>Hypereteone foliosa</i>
3	B13	4	21.8	<i>Glycera americana</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B13	31	16.6	<i>Leitoscoloplos robustus</i>
3	B13	3	0	<i>Polydora cornuta</i>
3	B13	170	78.1	<i>Streblospio benedicti</i>
3	B13	27	3.3	<i>Tharyx sp. A</i>
3	B13	11	1.9	<i>Mediomastus ambiseta</i>
3	B13	4	14	<i>Sabellaria vulgaris</i>
3	B13	3	63.2	<i>Pectinaria gouldii</i>
3	B13	3	0.7	<i>Asabellides oculata</i>
3	B13	0	14	<i>Annelida fragments</i>
3	B13	10	1.8	<i>Oligochaeta</i>
3	B13	77	30	<i>Odostomia sp.</i>
3	B13	13	46.1	<i>Rictaxis punctostriatus</i>
3	B13	9	43.8	<i>Mulinia lateralis</i>
3	B13	2	401	<i>Ensis directus</i>
3	B13	40	68.6	<i>Mya arenaria</i>
3	B13	1	0.8	<i>Balanus improvisus</i>
3	B13	28	4.8	<i>Leucon americanus</i>
3	B13	1	12.4	<i>Cyathura polita</i>
3	B13	3	0.9	<i>Edotea triloba</i>
3	B13	1	0.8	<i>Ampelisca sp.</i>
3	B13	1	53.8	<i>Rhithropanopeus harrisii</i>
3	B13	3	0	<i>Molgula manhattensis</i>
3	B14	2	13.8	<i>Glycera americana</i>
3	B14	26	51.4	<i>Leitoscoloplos sp.</i>
3	B14	1	0	<i>Mediomastus ambiseta</i>
3	B14	1	24.3	<i>Pectinaria gouldii</i>
3	B14	40	15.1	<i>Leucon americanus</i>
3	B15	1	0	<i>Rhynchocoela</i>
3	B15	2	0	<i>Hypereteone heteropoda</i>
3	B15	39	51	<i>Leitoscoloplos robustus</i>
3	B15	1	0	<i>Tharyx sp. A</i>
3	B15	1	0	<i>Mediomastus ambiseta</i>
3	B15	0	7.2	<i>Annelida fragments</i>
3	B15	1	1	<i>Neomysis americana</i>
3	B15	85	20.5	<i>Leucon americanus</i>
3	B15	1	0	<i>Unciola serrata</i>
3	B16	1	0	<i>Hypereteone heteropoda</i>
3	B16	2	40.4	<i>Glycera americana</i>
3	B16	109	168	<i>Leitoscoloplos robustus</i>
3	B16	2	0	<i>Mediomastus ambiseta</i>
3	B16	2	43.6	<i>Pectinaria gouldii</i>
3	B16	0	0.1	<i>Annelida fragments</i>
3	B16	3	0	<i>Oligochaeta</i>
3	B16	1	0	<i>Odostomia sp.</i>
3	B16	5	11.9	<i>Rictaxis punctostriatus</i>
3	B16	54	14	<i>Leucon americanus</i>
3	B17	1	0.8	<i>Rhynchocoela</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	B17	1	1.1	<i>Hypereteone heteropoda</i>
3	B17	1	26.7	<i>Glycera americana</i>
3	B17	25	25.4	<i>Leitoscoloplos robustus</i>
3	B17	1	72.9	<i>Pectinaria gouldii</i>
3	B17	1	6.9	<i>Mya arenaria</i>
3	B17	64	16.7	<i>Leucon americanus</i>
3	B18	1	0	<i>Hypereteone heteropoda</i>
3	B18	7	5.8	<i>Leitoscoloplos robustus</i>
3	B18	1	12	<i>Pectinaria gouldii</i>
3	B18	1	0	<i>Rictaxis punctostriatus</i>
3	V1	3	0	<i>Hypereteone heteropoda</i>
3	V1	1	0	<i>Neanthes succinea</i>
3	V1	3	13	<i>Glycera americana</i>
3	V1	22	39.8	<i>Leitoscoloplos robustus</i>
3	V1	1	0	<i>Polydora spp.</i>
3	V1	1	0	<i>Polydora cornuta</i>
3	V1	2	2.5	<i>Spio setosa</i>
3	V1	294	213.2	<i>Streblospio benedicti</i>
3	V1	2	0	<i>Tharyx sp. A</i>
3	V1	1	0.6	<i>Heteromastus filiformis</i>
3	V1	65	3.4	<i>Mediomastus ambiseta</i>
3	V1	1	19.1	<i>Pectinaria gouldii</i>
3	V1	0	3.4	<i>Annelida fragments</i>
3	V1	1	0	<i>Oligochaeta</i>
3	V1	19	5	<i>Odostomia sp.</i>
3	V1	1	0.6	<i>Rictaxis punctostriatus</i>
3	V1	3	0.7	<i>Mulinia lateralis</i>
3	V1	1	0	<i>Ensis directus</i>
3	V1	8	9.5	<i>Mya arenaria</i>
3	V1	1	0	<i>Balanus improvisus</i>
3	V1	1	0	<i>Leucon americanus</i>
3	V1	1	0	<i>Corophium sp.</i>
3	V2	3	14	<i>Rhynchocoela</i>
3	V2	2	0	<i>Hypereteone heteropoda</i>
3	V2	1	0	<i>Podarkeopsis levifuscina</i>
3	V2	2	228.1	<i>Glycera americana</i>
3	V2	44	313.9	<i>Leitoscoloplos robustus</i>
3	V2	10	16.9	<i>Polydora cornuta</i>
3	V2	1	0	<i>Tharyx sp. A</i>
3	V2	2	1.4	<i>Heteromastus filiformis</i>
3	V2	20	1.5	<i>Mediomastus ambiseta</i>
3	V2	0	1.6	<i>Annelida fragments</i>
3	V2	4	4.7	<i>Rictaxis punctostriatus</i>
3	V2	12	380	<i>Mulinia lateralis</i>
3	V2	13	13.9	<i>Mya arenaria</i>
3	V2	20	7.1	<i>Leucon americanus</i>
3	V2	1	1.5	<i>Ampelisca abdita</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	V2	1	0	<i>Monoculodes sp. 1</i>
3	V3	3	68.2	<i>Glycera americana</i>
3	V3	28	102.3	<i>Leitoscoloplos robustus</i>
3	V3	4	10.6	<i>Polydora cornuta</i>
3	V3	1	0	<i>Tharyx sp. A</i>
3	V3	4	6.8	<i>Heteromastus filiformis</i>
3	V3	1	0	<i>Mediomastus ambiseta</i>
3	V3	0	0.9	<i>Annelida fragments</i>
3	V3	1	0	<i>Gastropoda</i>
3	V3	5	2.9	<i>Rictaxis punctostriatus</i>
3	V3	24	881.9	<i>Mulinia lateralis</i>
3	V3	2	0.6	<i>Ensis directus</i>
3	V3	14	2284.1	<i>Mya arenaria</i>
3	V3	7	1	<i>Leucon americanus</i>
3	V3	3	3.6	<i>Ampelisca abdita</i>
3	V4	1	1	<i>Hypereteone heteropoda</i>
3	V4	2	45.5	<i>Glycera americana</i>
3	V4	53	415	<i>Leitoscoloplos robustus</i>
3	V4	5	19.6	<i>Polydora cornuta</i>
3	V4	1	0	<i>Tharyx sp. A</i>
3	V4	1	0.5	<i>Heteromastus filiformis</i>
3	V4	0	1	<i>Annelida fragments</i>
3	V4	5	1	<i>Rictaxis punctostriatus</i>
3	V4	35	348	<i>Mulinia lateralis</i>
3	V4	3	7.4	<i>Mya arenaria</i>
3	V4	54	10.2	<i>Leucon americanus</i>
3	V4	1	0	<i>Ampelisca abdita</i>
3	V5	1	34.5	<i>Glycera americana</i>
3	V5	61	599.6	<i>Leitoscoloplos robustus</i>
3	V5	3	19	<i>Polydora cornuta</i>
3	V5	1	0	<i>Tharyx sp. A</i>
3	V5	3	1.3	<i>Rictaxis punctostriatus</i>
3	V5	19	269.7	<i>Mulinia lateralis</i>
3	V5	7	1313.8	<i>Mya arenaria</i>
3	V5	19	5.5	<i>Leucon americanus</i>
3	V5	1	5.4	<i>Ampelisca abdita</i>
3	V6	1	0	<i>Podarkeopsis levifuscina</i>
3	V6	2	138.4	<i>Glycera americana</i>
3	V6	53	457.8	<i>Leitoscoloplos robustus</i>
3	V6	2	10.7	<i>Polydora cornuta</i>
3	V6	2	0.5	<i>Tharyx sp. A</i>
3	V6	1	234.8	<i>Pectinaria gouldii</i>
3	V6	1	0	<i>Odostomia sp.</i>
3	V6	15	7	<i>Rictaxis punctostriatus</i>
3	V6	1	0	<i>Cylichnella bidentata</i>
3	V6	23	667.3	<i>Mulinia lateralis</i>
3	V6	6	3.5	<i>Mya arenaria</i>

Sample month	Station	Abundance	Weight, mg	Taxon
3	V6	47	17.8	<i>Leucon americanus</i>
3	V6	1	0	<i>Oxyurostylis smithi</i>
3	V6	12	16.7	<i>Ampelisca abdita</i>
3	V7	1	0	<i>Hypereteone heteropoda</i>
3	V7	1	7.6	<i>Hypereteone foliosa</i>
3	V7	1	1.8	<i>Neanthes succinea</i>
3	V7	2	125.8	<i>Glycera americana</i>
3	V7	49	510.2	<i>Leitoscoloplos robustus</i>
3	V7	1	3.3	<i>Polydora cornuta</i>
3	V7	1	0.7	<i>Sabellaria vulgaris</i>
3	V7	7	4.5	<i>Rictaxis punctostriatus</i>
3	V7	9	42.9	<i>Mulinia lateralis</i>
3	V7	2	0.5	<i>Mya arenaria</i>
3	V7	22	6.7	<i>Leucon americanus</i>
4	B01	1	0.1	<i>Hypereteone heteropoda</i>
4	B01	1	0.1	<i>Glycera americana</i>
4	B01	8	17.1	<i>Leitoscoloplos robustus</i>
4	B01	1	0.1	<i>Polydora cornuta</i>
4	B01	2	14.8	<i>Spio setosa</i>
4	B01	47	20.9	<i>Streblospio benedicti</i>
4	B01	281	387.8	<i>Tharyx sp. A</i>
4	B01	3	1.9	<i>Heteromastus filiformis</i>
4	B01	3	1.4	<i>Sabellaria vulgaris</i>
4	B01	35	1.3	<i>Oligochaeta</i>
4	B01	25	11.2	<i>Odostomia sp.</i>
4	B01	2	0.1	<i>Macoma balthica</i>
4	B01	3	49.9	<i>Mya arenaria</i>
4	B01	9	1	<i>Leucon americanus</i>
4	B01	1	0.1	<i>Edotea triloba</i>
4	B02	1	30.1	<i>Anthozoa</i>
4	B02	5	1.9	<i>Hypereteone heteropoda</i>
4	B02	1	0.1	<i>Glycera americana</i>
4	B02	30	73.7	<i>Leitoscoloplos robustus</i>
4	B02	4	5	<i>Polydora cornuta</i>
4	B02	139	52.3	<i>Streblospio benedicti</i>
4	B02	502	590.7	<i>Tharyx sp. A</i>
4	B02	1	0.1	<i>Heteromastus filiformis</i>
4	B02	17	1	<i>Mediomastus ambiseta</i>
4	B02	3	12	<i>Sabellaria vulgaris</i>
4	B02	3	151.3	<i>Pectinaria gouldii</i>
4	B02	0	16.4	<i>Annelida fragments</i>
4	B02	56	4.9	<i>Oligochaeta</i>
4	B02	1	0.1	<i>Epitonium sp.</i>
4	B02	28	15.8	<i>Odostomia sp.</i>
4	B02	11	33	<i>Rictaxis punctostriatus</i>
4	B02	2	99.3	<i>Mulinia lateralis</i>
4	B02	5	0.1	<i>Macoma balthica</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	B02	12	153.6	<i>Mya arenaria</i>
4	B02	3	574.9	<i>Bivalve siphons</i>
4	B02	1	6.8	<i>Balanus improvisus</i>
4	B02	4	0.1	<i>Leucon americanus</i>
4	B02	1	1.3	<i>Unciola serrata</i>
4	B02	1	93.9	<i>Crangon septemspinosa</i>
4	B02	1	0.1	<i>Echinoidea</i>
4	B03	3	0.1	<i>Hypereteone heteropoda</i>
4	B03	5	15.6	<i>Glycera americana</i>
4	B03	68	73.3	<i>Leitoscoloplos robustus</i>
4	B03	4	17.8	<i>Polydora cornuta</i>
4	B03	7	1.8	<i>Streblospio benedicti</i>
4	B03	348	318.7	<i>Tharyx sp. A</i>
4	B03	190	12.2	<i>Mediomastus ambiseta</i>
4	B03	12	742.6	<i>Pectinaria gouldii</i>
4	B03	0	19.7	<i>Annelida fragments</i>
4	B03	34	1.3	<i>Oligochaeta</i>
4	B03	41	18.7	<i>Odostomia sp.</i>
4	B03	2	6.5	<i>Rictaxis punctostriatus</i>
4	B03	4	128.3	<i>Mulinia lateralis</i>
4	B03	2	0.1	<i>Macoma balthica</i>
4	B03	22	427.1	<i>Mya arenaria</i>
4	B03	21	4.7	<i>Leucon americanus</i>
4	B04	6	0.1	<i>Hypereteone heteropoda</i>
4	B04	4	24.1	<i>Glycera americana</i>
4	B04	54	150.7	<i>Leitoscoloplos robustus</i>
4	B04	9	7.1	<i>Polydora cornuta</i>
4	B04	4	0.1	<i>Streblospio benedicti</i>
4	B04	17	2.9	<i>Tharyx sp. A</i>
4	B04	6	50.4	<i>Heteromastus filiformis</i>
4	B04	8	0.9	<i>Mediomastus ambiseta</i>
4	B04	9	728.9	<i>Pectinaria gouldii</i>
4	B04	0	5.9	<i>Annelida fragments</i>
4	B04	11	0.1	<i>Oligochaeta</i>
4	B04	13	5	<i>Odostomia sp.</i>
4	B04	7	67	<i>Mulinia lateralis</i>
4	B04	19	62.2	<i>Mya arenaria</i>
4	B04	60	13.2	<i>Leucon americanus</i>
4	B04	6	3.5	<i>Edotea triloba</i>
4	B04	1	37	<i>Crangon septemspinosa</i>
4	B05	4	15.9	<i>Anthozoa</i>
4	B05	4	1.7	<i>Hypereteone heteropoda</i>
4	B05	3	0.1	<i>Eumida sanguinea</i>
4	B05	1	0.1	<i>Proceraea cornuta</i>
4	B05	1	2.2	<i>Neanthes succinea</i>
4	B05	8	5	<i>Leitoscoloplos robustus</i>
4	B05	14	11.6	<i>Polydora cornuta</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	B05	285	157.5	<i>Streblospio benedicti</i>
4	B05	4	0.1	<i>Tharyx sp. A</i>
4	B05	12	0.1	<i>Mediomastus ambiseta</i>
4	B05	35	127.6	<i>Sabellaria vulgaris</i>
4	B05	2	0.1	<i>Asabellides oculata</i>
4	B05	0	3.9	<i>Annelida fragments</i>
4	B05	26	2.3	<i>Oligochaeta</i>
4	B05	29	6.7	<i>Odostomia sp.</i>
4	B05	12	9221.3	<i>Mya arenaria</i>
4	B05	36	4.5	<i>Leucon americanus</i>
4	B05	1	0.5	<i>Edotea triloba</i>
4	B05	58	55	<i>Unciola serrata</i>
4	B05	1	193.8	<i>Crangon septemspinosa</i>
4	B05	9	7.2	<i>Molgula manhattensis</i>
4	B06	2	0.1	<i>Leitoscoloplos robustus</i>
4	B06	3	2.7	<i>Polydora cornuta</i>
4	B06	6	1.1	<i>Streblospio benedicti</i>
4	B06	2	0.1	<i>Tharyx sp. A</i>
4	B06	1	0.1	<i>Mediomastus ambiseta</i>
4	B06	4	237.9	<i>Pectinaria gouldii</i>
4	B06	1	0.1	<i>Oligochaeta</i>
4	B06	1	0.1	<i>Gastropoda</i>
4	B06	3	194	<i>Mulinia lateralis</i>
4	B06	9	9.5	<i>Mya arenaria</i>
4	B06	53	16	<i>Leucon americanus</i>
4	B06	1	51.7	<i>Crangon septemspinosa</i>
4	B07	7	1	<i>Leitoscoloplos robustus</i>
4	B07	1	0.5	<i>Polydora cornuta</i>
4	B07	3	142.1	<i>Pectinaria gouldii</i>
4	B07	3	293.6	<i>Mulinia lateralis</i>
4	B07	26	6.3	<i>Leucon americanus</i>
4	B08	7	0.5	<i>Polydora cornuta</i>
4	B09	1	250.5	<i>Nephtys incisa</i>
4	B09	22	8.7	<i>Leitoscoloplos robustus</i>
4	B09	2	0.5	<i>Tharyx sp. A</i>
4	B09	2	5.9	<i>Heteromastus filiformis</i>
4	B09	5	341.5	<i>Pectinaria gouldii</i>
4	B09	2	0.1	<i>Oligochaeta</i>
4	B09	4	118.5	<i>Mulinia lateralis</i>
4	B09	23	4.8	<i>Leucon americanus</i>
4	B10	1	1.1	<i>Hypereteone heteropoda</i>
4	B10	4	244	<i>Glycera americana</i>
4	B10	65	51.7	<i>Leitoscoloplos robustus</i>
4	B10	58	40.5	<i>Polydora cornuta</i>
4	B10	1	0.1	<i>Tharyx sp. A</i>
4	B10	2	4.3	<i>Heteromastus filiformis</i>
4	B10	6	0.1	<i>Mediomastus ambiseta</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	B10	6	2.4	<i>Sabellaria vulgaris</i>
4	B10	12	419.8	<i>Pectinaria gouldii</i>
4	B10	0	10.8	<i>Annelida fragments</i>
4	B10	2	0.1	<i>Oligochaeta</i>
4	B10	2	0.1	<i>Rictaxis punctostriatus</i>
4	B10	5	468.2	<i>Mulinia lateralis</i>
4	B10	2	4.3	<i>Barnea truncata</i>
4	B10	1	22	<i>Balanus improvisus</i>
4	B10	7	0.8	<i>Leucon americanus</i>
4	B10	4	2.4	<i>Unciola serrata</i>
4	B10	1	0.1	<i>Parapleustes aestuarius</i>
4	B10	1	484.6	<i>Rhithropanopeus harrisii</i>
4	B10	1	4	<i>Molgula manhattensis</i>
4	B11	1	3.6	<i>Hypereteone foliosa</i>
4	B11	5	2.4	<i>Leitoscoloplos robustus</i>
4	B11	2	0.8	<i>Streblospio benedicti</i>
4	B11	1	0.1	<i>Tharyx sp. A</i>
4	B11	2	33.5	<i>Mya arenaria</i>
4	B11	11	1.6	<i>Leucon americanus</i>
4	B12	2	0.1	<i>Hypereteone heteropoda</i>
4	B12	29	180.6	<i>Leitoscoloplos robustus</i>
4	B12	21	11.5	<i>Tharyx sp. A</i>
4	B12	3	0.1	<i>Mediomastus ambiseta</i>
4	B12	12	5.6	<i>Oligochaeta</i>
4	B12	1	0.5	<i>Odostomia sp.</i>
4	B12	2	6.5	<i>Rictaxis punctostriatus</i>
4	B12	3	6.2	<i>Mulinia lateralis</i>
4	B12	1	0.1	<i>Tellina agilis</i>
4	B12	3	17.2	<i>Mya arenaria</i>
4	B12	60	17.1	<i>Leucon americanus</i>
4	B13	2	0.1	<i>Hypereteone heteropoda</i>
4	B13	1	21.8	<i>Glycera americana</i>
4	B13	17	103	<i>Leitoscoloplos robustus</i>
4	B13	1	1.6	<i>Polydora cornuta</i>
4	B13	61	73	<i>Streblospio benedicti</i>
4	B13	11	3.3	<i>Tharyx sp. A</i>
4	B13	1	3.3	<i>Heteromastus filiformis</i>
4	B13	25	1.2	<i>Mediomastus ambiseta</i>
4	B13	1	46.4	<i>Pectinaria gouldii</i>
4	B13	9	0.1	<i>Oligochaeta</i>
4	B13	6	1.5	<i>Odostomia sp.</i>
4	B13	1	5	<i>Rictaxis punctostriatus</i>
4	B13	7	26.8	<i>Mulinia lateralis</i>
4	B13	4	60.8	<i>Mya arenaria</i>
4	B13	116	43.8	<i>Leucon americanus</i>
4	B13	1	4	<i>Edotea triloba</i>
4	B14	37	191.6	<i>Leitoscoloplos robustus</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	B14	3	1.2	<i>Streblospio benedicti</i>
4	B14	1	0.1	<i>Tharyx sp. A</i>
4	B14	2	2.2	<i>Heteromastus filiformis</i>
4	B14	3	153	<i>Pectinaria gouldii</i>
4	B14	1	0.1	<i>Oligochaeta</i>
4	B14	24	4.7	<i>Leucon americanus</i>
4	B15	7	5.4	<i>Leitoscoloplos robustus</i>
4	B15	7	1	<i>Leucon americanus</i>
4	B16	127	155.9	<i>Leitoscoloplos robustus</i>
4	B16	1	0.1	<i>Streblospio benedicti</i>
4	B16	1	0.6	<i>Heteromastus filiformis</i>
4	B16	2	132.8	<i>Pectinaria gouldii</i>
4	B16	1	0.1	<i>Oligochaeta</i>
4	B16	15	3.1	<i>Leucon americanus</i>
4	B17	1	34.2	<i>Glycera americana</i>
4	B17	8	78.8	<i>Leitoscoloplos robustus</i>
4	B17	1	0.1	<i>Streblospio benedicti</i>
4	B17	1	0.1	<i>Tharyx sp. A</i>
4	B17	6	479.7	<i>Pectinaria gouldii</i>
4	B17	2	0.1	<i>Oligochaeta</i>
4	B17	1	1464.9	<i>Macoma balthica</i>
4	B17	115	35.5	<i>Leucon americanus</i>
4	B18	2	0.1	<i>Leitoscoloplos robustus</i>
4	B18	1	0.1	<i>Mediomastus ambiseta</i>
4	B18	2	150.7	<i>Pectinaria gouldii</i>
4	B18	30	5.3	<i>Leucon americanus</i>
4	V1	2	0.1	<i>Hypereteone heteropoda</i>
4	V1	1	2.6	<i>Glycera americana</i>
4	V1	33	175.8	<i>Leitoscoloplos robustus</i>
4	V1	4	5.7	<i>Polydora cornuta</i>
4	V1	4	1.8	<i>Spio setosa</i>
4	V1	117	46.4	<i>Streblospio benedicti</i>
4	V1	9	1.8	<i>Tharyx sp. A</i>
4	V1	4	4.4	<i>Heteromastus filiformis</i>
4	V1	21	0.9	<i>Mediomastus ambiseta</i>
4	V1	1	0.1	<i>Oligochaeta</i>
4	V1	32	9.6	<i>Odostomia sp.</i>
4	V1	2	12.8	<i>Mulinia lateralis</i>
4	V1	3	10.4	<i>Mya arenaria</i>
4	V1	2	0.1	<i>Leucon americanus</i>
4	V2	3	183.2	<i>Anthozoa</i>
4	V2	1	0.1	<i>Hypereteone heteropoda</i>
4	V2	8	33.3	<i>Neanthes succinea</i>
4	V2	2	5	<i>Glycera americana</i>
4	V2	16	14.7	<i>Leitoscoloplos robustus</i>
4	V2	9	1.8	<i>Polydora cornuta</i>
4	V2	277	111.3	<i>Streblospio benedicti</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	V2	4	0.1	<i>Marenzelleria viridis</i>
4	V2	5	1.2	<i>Tharyx sp. A</i>
4	V2	163	7.7	<i>Mediomastus ambiseta</i>
4	V2	1	0.1	<i>Asabellides oculata</i>
4	V2	1	0.8	<i>Sabellidae</i>
4	V2	0	1.8	<i>Annelida fragments</i>
4	V2	1	0.1	<i>Oligochaeta</i>
4	V2	1	2.9	<i>Epitonium sp.</i>
4	V2	3	0.7	<i>Odostomia sp.</i>
4	V2	4	2.4	<i>Doradella obscura</i>
4	V2	11	401.5	<i>Mya arenaria</i>
4	V2	32	2565.1	<i>Balanus improvisus</i>
4	V2	10	2.2	<i>Leucon americanus</i>
4	V2	2	0.7	<i>Unciola serrata</i>
4	V2	8	6.2	<i>Melita nitida</i>
4	V2	2	80.1	<i>Rhithropanopeus harrisi</i>
4	V2	1	0.1	<i>Asteroidea</i>
4	V2	3	258.9	<i>Molgula manhattensis</i>
4	V3	2	56.9	<i>Glycera americana</i>
4	V3	52	292.4	<i>Leitoscoloplos robustus</i>
4	V3	4	6.8	<i>Polydora cornuta</i>
4	V3	9	1.8	<i>Tharyx sp. A</i>
4	V3	42	1.9	<i>Mediomastus ambiseta</i>
4	V3	1	0.1	<i>Oligochaeta</i>
4	V3	1	0.1	<i>Odostomia sp.</i>
4	V3	4	0.9	<i>Rictaxis punctostriatus</i>
4	V3	41	740.6	<i>Mulinia lateralis</i>
4	V3	17	104.2	<i>Mya arenaria</i>
4	V3	205	64.1	<i>Leucon americanus</i>
4	V3	1	0.1	<i>Edotea triloba</i>
4	V3	1	0.1	<i>Ampelisca abdita</i>
4	V3	1	8	<i>Crangon septemspinosa</i>
4	V4	1	0.1	<i>Rhynchocoela</i>
4	V4	1	84.8	<i>Neanthes succinea</i>
4	V4	1	25.3	<i>Glycera americana</i>
4	V4	49	385.1	<i>Leitoscoloplos robustus</i>
4	V4	6	28.3	<i>Polydora cornuta</i>
4	V4	1	0.1	<i>Streblospio benedicti</i>
4	V4	3	1.2	<i>Tharyx sp. A</i>
4	V4	1	9.3	<i>Heteromastus filiformis</i>
4	V4	4	0.1	<i>Mediomastus ambiseta</i>
4	V4	1	0.1	<i>Odostomia sp.</i>
4	V4	3	0.8	<i>Rictaxis punctostriatus</i>
4	V4	29	551.7	<i>Mulinia lateralis</i>
4	V4	18	3549.6	<i>Mya arenaria</i>
4	V4	78	26.1	<i>Leucon americanus</i>
4	V4	1	0.1	<i>Ampelisca abdita</i>

Sample month	Station	Abundance	Weight, mg	Taxon
4	V5	2	19.2	<i>Glycera americana</i>
4	V5	55	382.1	<i>Leitoscoloplos robustus</i>
4	V5	5	21.4	<i>Polydora cornuta</i>
4	V5	2	0.1	<i>Tharyx sp. A</i>
4	V5	1	0.1	<i>Mediomastus ambiseta</i>
4	V5	1	0.5	<i>Rictaxis punctostriatus</i>
4	V5	27	278.8	<i>Mulinia lateralis</i>
4	V5	2	1.5	<i>Mya arenaria</i>
4	V5	7	2.2	<i>Leucon americanus</i>
4	V6	2	68.1	<i>Glycera americana</i>
4	V6	28	223.5	<i>Leitoscoloplos robustus</i>
4	V6	2	2	<i>Polydora cornuta</i>
4	V6	1	0.1	<i>Streblospio benedicti</i>
4	V6	1	0.1	<i>Tharyx sp. A</i>
4	V6	1	0.1	<i>Mediomastus ambiseta</i>
4	V6	1	80.8	<i>Pectinaria gouldii</i>
4	V6	1	0.1	<i>Odostomia sp.</i>
4	V6	3	1.6	<i>Rictaxis punctostriatus</i>
4	V6	14	175.5	<i>Mulinia lateralis</i>
4	V6	13	59.9	<i>Mya arenaria</i>
4	V6	36	10.7	<i>Leucon americanus</i>
4	V7	3	2.5	<i>Hypereteone heteropoda</i>
4	V7	1	4	<i>Hypereteone foliosa</i>
4	V7	1	124	<i>Glycera americana</i>
4	V7	62	429.7	<i>Leitoscoloplos robustus</i>
4	V7	14	83.7	<i>Polydora cornuta</i>
4	V7	2	0.1	<i>Streblospio benedicti</i>
4	V7	2	1.5	<i>Tharyx sp. A</i>
4	V7	1	0.8	<i>Heteromastus filiformis</i>
4	V7	7	0.1	<i>Mediomastus ambiseta</i>
4	V7	2	0.1	<i>Odostomia sp.</i>
4	V7	4	35.7	<i>Mulinia lateralis</i>
4	V7	17	4.4	<i>Leucon americanus</i>