

TABLE 20a. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Assured immediate access to medication*	Had full-time registered nurse**	Identified and tracked students with asthma	Obtained and used an Asthma Action Plan <sup>§</sup>	Provided intensive case management for students with asthma <sup>§§</sup>
<b>Weighted Data</b>					
Alabama	90.7	21.7	72.4	52.5	29.8
Alaska	66.6	17.9	50.5	27.9	12.7
Arizona	81.9	46.8	79.9	46.4	36.9
Arkansas	91.9	54.3	79.7	56.3	33.5
Connecticut	88.9	89.7	94.1	69.1	46.8
Delaware	96.1	98.0	92.2	74.0	46.1
Georgia	86.5	41.8	63.0	40.2	35.7
Hawaii	95.5	15.0	79.5	62.6	30.9
Idaho	90.4	13.5	54.7	29.3	35.6
Illinois <sup>+</sup>	94.5	42.5	88.5	57.1	29.6
Iowa	94.3	40.6	85.7	57.2	30.9
Kentucky	92.6	28.0	66.4	34.4	23.1
Maine	90.2	45.0	90.2	57.9	39.5
Massachusetts	88.2	95.4	93.9	66.8	45.4
Michigan	95.2	10.4	73.6	38.5	21.4
Minnesota	92.4	35.6	86.3	62.2	34.7
Missouri	91.3	71.7	93.0	64.1	38.1
Montana	88.7	10.9	65.4	31.6	22.4
Nebraska	92.1	25.0	79.9	43.5	26.4
New Hampshire	89.9	89.6	95.3	56.6	48.0
New Jersey	96.6	97.5	98.1	79.7	48.6
New Mexico	92.5	49.4	81.9	57.7	38.8
New York	91.1	90.6	94.4	62.9	33.3
North Carolina	96.6	14.7	76.1	61.2	41.9
North Dakota	88.0	2.2	52.7	26.8	27.1
Oklahoma	88.3	28.1	59.8	31.0	31.3
Tennessee	92.9	18.7	71.9	54.4	24.4
Texas <sup>+</sup>	91.7	71.8	82.4	55.9	36.0
Utah	92.9	1.4	43.9	33.4	40.9
Vermont	90.9	63.1	96.3	64.5	53.7
Virginia	92.4	59.8	87.8	68.5	43.3
Wisconsin	96.2	17.2	83.3	46.1	32.7
<b>Unweighted Data</b>					
California	88.0	21.4	84.6	44.2	28.5
Oregon	92.2	10.4	80.8	50.8	25.8
Pennsylvania <sup>+</sup>	88.0	79.4	93.0	54.9	26.6
Rhode Island	100.0	84.6	90.6	70.3	43.5
South Carolina	90.7	57.0	86.1	58.7	45.3
Washington	94.1	15.2	86.4	62.3	31.2
<b>State Median</b>	<b>91.8</b>	<b>41.2</b>	<b>82.8</b>	<b>56.4</b>	<b>34.1</b>

(continued)

TABLE 20a. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Assured immediate access to medication*	Had full-time registered nurse**	Identified and tracked students with asthma	Obtained and used an Asthma Action Plan <sup>§</sup>	Provided intensive case management for students with asthma <sup>§§</sup>
<b>Weighted Data</b>					
Chicago	94.5	8.9	79.2	61.3	49.5
Dallas	91.9	70.4	82.0	75.9	61.2
Fort Lauderdale	90.2	24.1	70.2	40.6	29.0
Houston	88.9	100.0	95.6	95.6	79.4
Los Angeles	84.4	92.4	95.3	69.5	49.4
Miami	83.5	13.2	24.7	19.5	29.3
Orange County	89.8	47.9	64.6	49.1	32.8
Palm Beach	100.0	91.9	97.3	91.9	46.0
San Bernardino	78.6	14.3	71.4	35.7	35.7
San Diego	93.9	79.6	98.0	80.9	64.4
San Francisco	91.3	6.1	52.5	41.0	37.7
<b>Unweighted Data</b>					
District of Columbia	86.7	63.3	80.0	60.0	43.3
Philadelphia	85.1	88.5	96.5	84.5	60.5
<b>Local Median</b>	<b>89.8</b>	<b>63.3</b>	<b>80.0</b>	<b>61.3</b>	<b>46.0</b>

\* As prescribed by a physician and approved by parents.

\*\* All day every day.

§ For all students with asthma.

§§ For students with asthma who are absent 10 days or more per year.

+ Survey did not include students from one of the state's large school districts.

TABLE 20b. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Educated school staff about asthma	Educated students with asthma about asthma management	Taught asthma awareness to all students*	Encouraged full participation in physical education and physical activity**	Modified physical education and physical activities <sup>§</sup>
<b>Weighted Data</b>					
Alabama	53.7	41.1	21.8	94.4	80.9
Alaska	34.3	26.4	19.2	77.5	57.1
Arizona	46.6	46.5	15.1	84.3	73.7
Arkansas	52.0	42.9	28.5	97.0	81.4
Connecticut	58.5	68.6	19.8	97.4	86.9
Delaware	57.6	60.8	28.5	98.0	77.7
Georgia	45.2	37.0	23.1	93.5	81.7
Hawaii	25.0	38.4	24.9	92.3	83.7
Idaho	38.2	28.4	26.1	94.1	83.7
Illinois <sup>§§</sup>	65.2	44.9	31.2	97.4	75.9
Iowa	59.9	51.8	22.8	98.9	85.1
Kentucky	35.3	24.0	13.5	92.4	71.7
Maine	54.2	63.8	24.6	97.9	82.1
Massachusetts	52.0	71.6	27.8	98.5	85.4
Michigan	52.4	27.6	22.7	95.3	79.7
Minnesota	54.4	54.8	28.6	95.0	84.8
Missouri	60.0	55.9	31.7	97.3	84.5
Montana	49.0	29.3	28.4	94.3	76.1
Nebraska	57.0	39.6	22.8	97.5	76.3
New Hampshire	59.2	73.5	21.0	99.4	89.1
New Jersey	67.4	77.4	47.1	98.7	85.3
New Mexico	54.6	58.6	26.5	95.1	80.7
New York	49.9	59.9	34.8	99.1	80.9
North Carolina	57.3	54.4	22.0	94.6	85.4
North Dakota	38.9	22.2	27.1	92.8	76.7
Oklahoma	51.9	31.8	14.3	94.4	67.1
Tennessee	50.0	37.1	17.1	93.2	80.3
Texas <sup>§§</sup>	50.8	47.1	20.8	96.8	78.7
Utah	33.7	19.6	27.3	93.3	83.3
Vermont	65.5	75.0	23.7	99.0	84.7
Virginia	58.6	60.0	32.9	95.6	84.2
Wisconsin	50.6	42.8	30.3	98.8	77.6
<b>Unweighted Data</b>					
California	49.6	41.2	18.4	94.1	78.4
Oregon	38.5	39.1	27.8	96.6	83.1
Pennsylvania <sup>§§</sup>	53.7	61.8	31.7	98.7	81.8
Rhode Island	50.8	70.3	47.6	98.5	92.3
South Carolina	50.4	53.9	20.2	95.4	82.3
Washington	46.4	48.3	16.7	92.5	83.8
<b>State Median</b>	<b>51.9</b>	<b>46.8</b>	<b>24.8</b>	<b>95.5</b>	<b>81.8</b>

(continued)

TABLE 20b. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites:  
School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Educated school staff about asthma	Educated students with asthma about asthma management	Taught asthma awareness to all students*	Encouraged full participation in physical education and physical activity**	Modified physical education and physical activities§
<b>Weighted Data</b>					
Chicago	72.0	64.2	38.6	97.2	85.5
Dallas	44.9	61.2	31.7	92.2	88.2
Fort Lauderdale	53.5	38.0	32.0	95.3	85.5
Houston	70.7	77.9	46.0	100.0	91.2
Los Angeles	57.1	69.5	40.5	97.1	95.1
Miami	27.9	23.1	25.2	81.3	65.6
Orange County	34.6	40.5	29.8	88.8	70.4
Palm Beach	63.5	77.3	36.9	97.3	94.5
San Bernardino	35.7	50.0	15.4	92.9	78.6
San Diego	75.5	81.3	8.7	91.7	87.2
San Francisco	64.8	50.5	34.7	94.4	80.1
<b>Unweighted Data</b>					
District of Columbia	53.3	70.0	40.0	90.0	66.7
Philadelphia	67.8	84.7	50.0	95.4	91.9
<b>Local Median</b>	<b>57.1</b>	<b>64.2</b>	<b>34.7</b>	<b>94.4</b>	<b>85.5</b>

\* In at least one grade.

\*\* When students with asthma were doing well.

§ As specified by the student's Asthma Action Plan.

§§ Survey did not include students from one of the state's large school districts.

TABLE 21. Percentage of Schools\* in Which Students Could Purchase Selected Items from Vending Machines or at the School Store, Canteen, or Snack Bar, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Chocolate candy	Other kinds of candy	Salty snacks not low in fat**	Salty snacks low in fat <sup>S</sup>	Fruits or vegetables	Low-fat baked goods	Soft drinks, sports drinks, or fruit drinks <sup>SS</sup>	100% fruit juice	Bottled water
<b>Weighted Data</b>									
Alabama	74.2	79.8	91.8	84.9	20.5	70.5	95.0	72.4	86.9
Alaska	76.1	77.8	75.6	67.1	21.7	47.4	96.3	66.4	62.5
Arizona	68.2	72.2	80.8	76.9	41.5	57.5	96.9	64.5	87.0
Arkansas	69.5	70.7	73.2	66.3	22.8	49.8	95.2	63.1	83.0
Connecticut	41.3	45.4	79.7	83.8	53.5	63.2	92.1	76.8	85.0
Delaware	51.1	58.7	74.0	79.3	40.8	68.4	89.7	84.3	86.8
Georgia	66.4	65.1	70.8	69.1	19.6	59.3	94.4	75.1	96.0
Hawaii	31.2	31.8	31.2	30.7	18.9	25.6	96.4	53.4	90.3
Idaho	81.3	81.2	76.1	76.2	36.3	59.2	97.1	77.4	92.2
Illinois <sup>+</sup>	61.6	63.4	75.9	74.0	49.5	59.1	91.0	79.3	89.5
Iowa	69.6	70.3	73.8	74.2	37.3	62.2	94.2	87.7	89.4
Kentucky	82.7	85.0	85.8	84.9	20.4	68.7	96.9	67.0	94.5
Maine	50.1	55.7	71.3	78.0	41.9	60.8	89.0	91.2	97.7
Massachusetts	40.9	46.8	78.6	80.2	49.7	62.5	88.9	87.1	92.3
Michigan	71.4	74.4	82.9	85.3	61.4	67.1	95.0	85.1	94.3
Minnesota	83.2	84.3	87.8	87.6	49.7	71.3	98.2	90.5	92.7
Missouri	69.8	69.7	77.2	73.9	30.0	57.4	97.0	78.8	86.1
Montana	64.9	67.9	60.2	60.6	19.8	41.1	96.8	80.2	82.8
Nebraska	58.8	60.3	57.5	57.0	23.2	49.3	96.5	79.3	84.8
New Hampshire	49.7	52.7	75.5	79.6	49.0	68.7	93.9	89.8	96.9
New Jersey	54.5	58.8	83.3	82.3	52.7	56.3	89.8	74.1	84.4
New Mexico	78.0	81.0	78.6	74.8	34.9	58.3	95.3	68.6	94.3
New York	47.3	52.7	72.3	82.6	46.1	59.6	90.5	85.2	89.2
North Carolina	54.5	60.6	83.7	82.3	37.7	64.3	96.3	79.8	91.7
North Dakota	60.4	63.2	52.8	50.9	15.5	37.7	95.0	83.5	94.6
Oklahoma	85.8	88.0	83.1	80.5	19.3	58.6	99.0	51.5	82.9
Tennessee	81.0	82.9	82.3	81.8	29.0	64.9	97.8	73.6	93.3
Texas <sup>+</sup>	83.3	85.9	88.9	87.4	44.2	70.1	98.0	74.5	92.3
Utah	94.6	94.1	92.1	91.2	51.7	75.8	96.1	83.8	83.0
Vermont	23.7	28.8	63.5	68.9	50.0	53.9	88.3	83.4	81.3
Virginia	68.5	71.8	85.8	85.1	36.9	68.0	93.2	80.0	90.2
Wisconsin	64.2	69.3	72.4	76.1	49.4	66.3	94.5	84.1	85.9
<b>Unweighted Data</b>									
California	59.9	67.7	83.7	78.4	57.1	60.7	94.1	76.4	89.8
Oregon	68.7	76.4	81.3	81.3	47.0	60.6	97.6	75.5	92.1
Pennsylvania <sup>+</sup>	66.0	72.8	86.6	86.5	48.5	67.4	93.5	84.0	93.9
Rhode Island	62.9	69.4	91.9	91.8	64.5	67.7	96.8	91.9	98.4
South Carolina	75.6	80.5	89.7	83.8	30.2	64.6	95.5	68.0	90.9
Washington	75.0	77.9	80.1	77.8	39.1	62.4	96.8	80.1	92.2
<b>State Median</b>	<b>67.3</b>	<b>70.0</b>	<b>79.1</b>	<b>79.4</b>	<b>39.9</b>	<b>62.3</b>	<b>95.3</b>	<b>79.3</b>	<b>90.3</b>

(continued)

TABLE 21. Percentage of Schools\* in Which Students Could Purchase Selected Items from Vending Machines or at the School Store, Canteen, or Snack Bar, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

## LOCAL SURVEYS

Site	Chocolate candy	Other kinds of candy	Salty snacks not low in fat**	Salty snacks low in fat <sup>§</sup>	Fruits or vegetables	Low-fat baked goods	Soft drinks, sports drinks, or fruit drinks <sup>§§</sup>	100% fruit juice	Bottled water
<b>Weighted Data</b>									
Chicago	27.6	31.1	46.7	50.9	30.6	47.7	78.6	79.1	80.1
Dallas	82.1	86.0	96.1	79.9	61.0	76.4	100.0	88.2	96.0
Fort Lauderdale	58.7	61.9	71.8	78.1	16.3	65.5	96.4	72.8	82.3
Houston	84.6	82.4	93.3	76.0	80.3	58.2	100.0	91.2	95.5
Los Angeles	91.2	90.1	91.1	89.3	62.7	66.4	95.0	82.4	97.0
Miami	66.0	65.9	79.7	74.2	42.4	68.0	94.7	84.2	87.8
Orange County	53.2	59.0	82.2	80.4	65.2	63.1	97.7	90.3	100.0
Palm Beach	34.9	37.9	60.6	63.6	54.6	54.1	94.7	91.3	91.5
San Bernardino	27.3	27.3	72.7	72.7	72.7	54.5	100.0	81.8	90.9
San Diego	72.5	70.0	92.5	90.0	55.0	66.7	97.5	59.0	95.0
San Francisco	60.0	60.0	88.3	87.7	55.3	68.2	97.1	68.8	97.1
<b>Unweighted Data</b>									
District of Columbia	72.7	77.3	63.6	59.1	23.8	54.5	100.0	81.8	72.7
Philadelphia	36.9	56.1	80.3	75.8	53.8	57.8	89.4	84.6	75.4
<b>Local Median</b>	<b>60.0</b>	<b>61.9</b>	<b>80.3</b>	<b>76.0</b>	<b>55.0</b>	<b>63.1</b>	<b>97.1</b>	<b>82.4</b>	<b>91.5</b>

\* Among schools which allowed students to purchase snack foods or beverages from vending machines or at the school store, canteen, or snack bar.

\*\* Such as regular potato chips.

§ Such as pretzels, baked chips, or other low-fat chips.

§§ Not 100% fruit juice.

+ Survey did not include students from one of the state's large school districts.

TABLE 22. Percentage of Schools with a Policy That Prohibited Tobacco Use and, Among Those Schools, the Percentage That Designated an Individual to Enforce the Policy and the Percentage That Had an "Ideal" Tobacco-Use Prevention Policy,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Had a policy prohibiting tobacco use	Had an individual to enforce the policy	Had an "ideal" tobacco-use prevention policy
<b>Weighted Data</b>			
Alabama	99.6	63.3	59.0
Alaska	99.4	60.1	36.7
Arizona	97.8	64.7	63.0
Arkansas	100.0	64.3	54.3
Connecticut	97.8	50.9	41.5
Delaware	96.1	38.4	46.2
Georgia	97.9	56.1	45.5
Hawaii	100.0	64.2	65.4
Idaho	100.0	64.5	37.7
Illinois**	99.2	62.6	43.6
Iowa	98.5	62.5	21.0
Kentucky	97.5	60.9	12.6
Maine	99.5	59.2	53.3
Massachusetts	99.3	65.4	58.1
Michigan	98.1	56.1	34.2
Minnesota	98.4	60.8	47.7
Missouri	99.7	64.1	24.7
Montana	100.0	70.0	28.7
Nebraska	99.0	65.5	19.0
New Hampshire	98.9	59.6	40.4
New Jersey	97.5	59.7	46.8
New Mexico	96.9	62.4	48.6
New York	97.1	50.8	54.0
North Carolina	98.7	59.1	23.1
North Dakota	98.3	68.4	20.4
Oklahoma	100.0	64.3	24.6
Tennessee	99.3	67.5	27.1
Texas**	100.0	60.4	64.7
Utah	99.5	79.5	50.0
Vermont	100.0	73.9	55.7
Virginia	99.2	60.5	34.4
Wisconsin	98.1	61.2	54.5
<b>Unweighted Data</b>			
California	99.4	66.7	65.4
Oregon	98.9	67.6	44.7
Pennsylvania**	99.7	61.4	52.4
Rhode Island	96.9	58.1	59.3
South Carolina	98.5	57.9	40.4
Washington	99.2	68.2	56.8
<b>State Median</b>	<b>99.1</b>	<b>62.4</b>	<b>45.9</b>

(continued)

TABLE 22. Percentage of Schools with a Policy That Prohibited Tobacco Use and, Among Those Schools, the Percentage That Designated an Individual to Enforce the Policy and the Percentage That Had an "Ideal" Tobacco-Use Prevention Policy,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Had a policy prohibiting tobacco use	Had an individual to enforce the policy	Had an "ideal" tobacco-use prevention policy
<b>Weighted Data</b>			
Chicago	87.3	48.4	38.5
Dallas	100.0	54.3	69.8
Fort Lauderdale	97.9	35.5	39.4
Houston	100.0	62.4	55.7
Los Angeles	99.1	94.1	62.9
Miami	96.7	60.9	52.1
Orange County	100.0	62.4	51.7
Palm Beach	97.5	36.8	44.8
San Bernardino	100.0	64.3	64.3
San Diego	100.0	85.4	71.1
San Francisco	97.6	86.7	66.0
<b>Unweighted Data</b>			
District of Columbia	96.7	69.0	62.1
Philadelphia	95.3	48.1	34.6
<b>Local Median</b>	<b>97.9</b>	<b>62.4</b>	<b>55.7</b>

\* An "ideal" tobacco-use policy, as described in CDC's *Guidelines for School Health Programs to Prevent Tobacco Use and Addiction*, prohibits all tobacco use by students, all school staff, and visitors on school property, in school vehicles, and at school-sponsored functions away from school property.

\*\* Survey did not include students from one of the state's large school districts.



TABLE 23a. Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Referred to school counselor	Referred to school administrator	Encouraged to participate in a program**	Required to participate in a program**	Referred to legal authorities
<b>Weighted Data</b>					
Alabama	61.0	99.7	35.7	14.3	17.6
Alaska	61.7	95.5	57.3	26.3	45.9
Arizona	69.1	97.7	69.0	33.9	49.0
Arkansas	60.5	99.7	40.5	15.3	27.7
Connecticut	79.7	98.7	67.3	31.3	24.8
Delaware	80.8	97.9	71.9	42.4	15.0
Georgia	62.0	100.0	37.1	17.2	15.1
Hawaii	79.9	100.0	72.7	21.2	15.4
Idaho	88.4	99.4	72.8	58.6	96.3
Illinois <sup>§</sup>	70.5	98.3	51.2	24.4	37.6
Iowa	79.3	100.0	68.3	37.8	73.5
Kentucky	64.1	99.6	60.0	31.3	17.4
Maine	88.1	100.0	76.6	45.6	81.9
Massachusetts	77.4	99.5	69.9	39.8	20.0
Michigan	76.8	99.7	66.1	37.1	68.5
Minnesota	68.7	100.0	71.0	40.6	78.3
Missouri	56.5	99.7	35.6	14.3	38.5
Montana	79.8	100.0	60.9	44.8	82.2
Nebraska	70.5	99.0	57.7	24.6	46.3
New Hampshire	81.7	99.4	70.2	41.0	82.1
New Jersey	88.6	99.4	63.7	34.7	35.4
New Mexico	78.0	99.4	61.0	29.3	13.3
New York	83.2	97.4	66.4	34.2	17.9
North Carolina	69.3	100.0	51.4	40.7	15.2
North Dakota	77.7	98.6	55.7	25.3	65.4
Oklahoma	68.8	99.4	36.5	15.2	39.2
Tennessee	66.6	97.8	50.5	33.6	75.7
Texas <sup>§</sup>	68.1	98.9	38.5	20.9	67.5
Utah	76.1	99.0	77.1	57.2	96.1
Vermont	94.2	100.0	82.0	57.5	71.3
Virginia	71.6	98.8	61.5	45.0	63.8
Wisconsin	77.9	97.0	66.8	35.9	83.9
<b>Unweighted Data</b>					
California	78.0	98.2	68.7	50.1	49.1
Oregon	80.9	99.4	74.4	50.8	71.3
Pennsylvania <sup>§</sup>	71.9	100.0	62.2	35.4	87.1
Rhode Island	83.6	95.1	78.3	48.3	23.0
South Carolina	70.8	100.0	52.9	24.3	17.6
Washington	82.1	96.6	72.3	60.8	44.6
<b>State Median</b>	<b>77.1</b>	<b>99.4</b>	<b>64.9</b>	<b>35.0</b>	<b>46.1</b>

(continued)

TABLE 23a. Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Referred to school counselor	Referred to school administrator	Encouraged to participate in a program**	Required to participate in a program**	Referred to legal authorities
<b>Weighted Data</b>					
Chicago	75.6	94.3	37.8	20.3	26.0
Dallas	65.4	100.0	49.0	44.4	53.2
Fort Lauderdale	82.2	100.0	71.7	49.9	61.0
Houston	84.6	100.0	76.3	42.9	52.3
Los Angeles	89.2	92.9	75.6	82.2	47.1
Miami	93.5	96.7	80.5	48.4	24.7
Orange County	88.3	97.0	76.8	54.0	62.5
Palm Beach	91.3	100.0	85.7	68.2	44.1
San Bernardino	92.9	100.0	69.2	85.7	78.6
San Diego	91.8	98.0	80.9	89.4	64.6
San Francisco	90.8	90.6	82.6	76.2	29.6
<b>Unweighted Data</b>					
District of Columbia	85.7	100.0	55.2	41.4	17.9
Philadelphia	81.3	93.9	54.9	23.1	17.7
<b>Local Median</b>	<b>88.3</b>	<b>98.0</b>	<b>75.6</b>	<b>49.9</b>	<b>47.1</b>

\* Among schools with a policy prohibiting tobacco use.

\*\* An assistance, education, or cessation program.

§ Survey did not include students from one of the state's large school districts.

TABLE 23b. Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Placed in detention	Given in-school suspension	Suspended from school	Informed parents or guardians
<b>Weighted Data</b>				
Alabama	56.3	69.1	81.2	98.9
Alaska	45.5	52.2	74.5	97.7
Arizona	58.5	71.6	77.3	97.4
Arkansas	52.0	77.3	79.8	100.0
Connecticut	53.9	71.9	77.6	98.7
Delaware	68.1	74.0	74.5	100.0
Georgia	41.2	84.5	75.6	100.0
Hawaii	64.5	62.2	89.4	100.0
Idaho	56.1	70.3	78.1	100.0
Illinois**	53.4	68.3	81.3	98.6
Iowa	43.1	68.8	65.3	100.0
Kentucky	63.7	78.2	66.8	98.6
Maine	35.8	50.0	89.2	100.0
Massachusetts	62.7	65.0	78.3	99.0
Michigan	42.5	57.0	86.3	100.0
Minnesota	49.3	66.4	74.7	98.9
Missouri	62.2	82.7	70.9	99.2
Montana	59.0	72.4	72.2	100.0
Nebraska	59.3	78.7	75.3	100.0
New Hampshire	46.0	62.5	83.8	99.3
New Jersey	66.7	66.2	78.7	98.7
New Mexico	67.9	70.2	64.7	95.7
New York	64.9	78.5	59.6	100.0
North Carolina	62.0	80.8	78.4	99.1
North Dakota	60.6	70.3	66.3	98.2
Oklahoma	66.4	72.3	73.7	99.0
Tennessee	47.1	58.5	70.9	97.5
Texas**	63.7	88.1	53.6	99.1
Utah	45.3	54.0	79.6	99.0
Vermont	44.3	72.9	74.7	100.0
Virginia	45.3	63.4	89.6	99.2
Wisconsin	53.1	68.9	75.4	99.0
<b>Unweighted Data</b>				
California	59.3	70.7	84.8	100.0
Oregon	57.8	65.7	86.4	99.4
Pennsylvania**	37.9	62.6	70.0	99.7
Rhode Island	81.4	74.6	70.5	96.7
South Carolina	48.8	63.2	93.1	98.4
Washington	50.2	62.9	83.8	97.5
<b>State Median</b>	<b>56.2</b>	<b>69.7</b>	<b>76.5</b>	<b>99.1</b>

(continued)

TABLE 23b. Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Placed in detention	Given in-school suspension	Suspended from school	Informed parents or guardians
<b>Weighted Data</b>				
Chicago	71.9	67.2	82.0	96.6
Dallas	79.1	85.1	74.4	100.0
Fort Lauderdale	63.5	80.8	85.8	100.0
Houston	79.9	86.8	81.6	100.0
Los Angeles	72.7	65.3	82.1	100.0
Miami	71.4	84.2	70.1	96.7
Orange County	59.8	93.1	81.1	100.0
Palm Beach	41.2	41.9	100.0	100.0
San Bernardino	71.4	78.6	78.6	100.0
San Diego	68.8	68.8	77.6	98.0
San Francisco	51.5	65.3	56.1	96.9
<b>Unweighted Data</b>				
District of Columbia	71.4	79.3	86.2	96.6
Philadelphia	72.0	64.6	84.0	93.9
<b>Local Median</b>	<b>71.4</b>	<b>78.6</b>	<b>81.6</b>	<b>100.0</b>

\* Among schools with a policy prohibiting tobacco use.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 24. Percentage of Schools That Prohibited Tobacco Advertising in Specific Places, Through Sponsorship of School Events, and on Student Apparel, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Prohibited tobacco advertising in specific locations*	Prohibited tobacco sponsorship of school events	Prohibited students from wearing tobacco brand-name apparel
<b>Weighted Data</b>			
Alabama	91.1	92.2	95.6
Alaska	92.1	92.0	78.6
Arizona	93.8	92.2	94.7
Arkansas	89.2	91.5	94.5
Connecticut	90.5	91.8	78.4
Delaware	92.2	80.3	77.1
Georgia	91.7	92.1	94.3
Hawaii	94.9	94.8	92.4
Idaho	91.0	94.5	96.7
Illinois**	94.2	92.3	92.3
Iowa	88.6	92.8	98.2
Kentucky	91.3	87.9	84.9
Maine	92.7	94.4	91.9
Massachusetts	93.2	93.2	64.7
Michigan	93.7	95.4	94.1
Minnesota	92.9	95.6	92.2
Missouri	90.6	90.2	93.1
Montana	92.2	91.2	95.7
Nebraska	90.8	94.7	97.8
New Hampshire	94.5	94.6	90.4
New Jersey	96.0	92.1	69.7
New Mexico	93.4	95.5	95.5
New York	93.0	91.9	65.5
North Carolina	90.0	90.3	81.7
North Dakota	90.8	93.1	97.1
Oklahoma	91.4	89.8	96.9
Tennessee	93.6	91.9	94.1
Texas**	93.7	94.2	98.1
Utah	94.5	95.0	98.6
Vermont	94.7	91.6	93.1
Virginia	94.6	93.5	85.5
Wisconsin	92.8	96.1	88.9
<b>Unweighted Data</b>			
California	96.3	96.9	95.7
Oregon	95.6	94.5	92.3
Pennsylvania**	91.1	89.3	85.1
Rhode Island	95.4	92.2	64.6
South Carolina	95.4	90.8	87.8
Washington	95.9	91.9	91.9
<b>State Median</b>	<b>93.0</b>	<b>92.2</b>	<b>92.4</b>

(continued)

TABLE 24. Percentage of Schools That Prohibited Tobacco Advertising in Specific Places, Through Sponsorship of School Events, and on Student Apparel, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (*continued*)**LOCAL SURVEYS**

Site	Prohibited tobacco advertising in specific locations*	Prohibited tobacco sponsorship of school events	Prohibited students from wearing tobacco brand-name apparel
<b>Weighted Data</b>			
Chicago	88.8	87.6	84.7
Dallas	98.1	84.4	90.3
Fort Lauderdale	92.8	84.5	95.6
Houston	95.7	95.7	97.9
Los Angeles	92.2	91.5	93.3
Miami	94.4	94.7	80.3
Orange County	95.0	92.3	94.2
Palm Beach	97.5	94.7	94.5
San Bernardino	78.6	71.4	85.7
San Diego	93.9	100.0	100.0
San Francisco	94.5	92.0	82.6
<b>Unweighted Data</b>			
District of Columbia	93.3	90.0	70.0
Philadelphia	90.6	91.8	75.3
<b>Local Median</b>	<b>93.9</b>	<b>91.8</b>	<b>90.3</b>

\* In school buildings, on school grounds, on school buses, and in school publications.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 25. Percentage of Schools That Implemented Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Required visitors to report to main office	Maintained a closed campus	Used staff to monitor halls	Checked bags, desks, and lockers	Prohibited backpacks	Required school uniforms	Required student identification badges	Used metal detectors	Had uniformed police
<b>Weighted Data</b>									
Alabama	100.0	96.1	94.2	70.9	32.9	21.7	7.0	31.5	42.8
Alaska	86.6	35.3	70.6	25.5	9.8	1.3	0.8	1.2	9.0
Arizona	99.6	84.8	82.8	35.4	13.9	21.0	6.7	5.6	49.6
Arkansas	100.0	95.3	93.2	67.0	13.5	5.5	11.5	25.1	43.7
Connecticut	99.1	90.9	82.6	18.6	39.1	2.3	4.6	3.5	50.5
Delaware	100.0	88.4	88.1	31.6	38.0	9.9	11.7	0.0	51.6
Georgia	100.0	95.4	94.5	65.5	29.8	4.2	18.6	26.5	72.7
Hawaii	100.0	98.7	93.9	2.7	0.0	23.7	33.9	0.0	50.1
Idaho	98.9	54.5	87.6	54.5	21.5	0.0	0.5	0.0	53.4
Illinois*	98.7	76.6	82.4	51.9	44.8	5.9	7.7	8.6	26.3
Iowa	98.1	73.4	85.0	39.8	33.5	0.3	0.4	0.7	14.2
Kentucky	99.5	96.7	93.2	59.5	34.0	11.3	4.9	22.0	51.3
Maine	98.0	85.2	76.0	20.0	30.5	0.0	1.0	0.5	25.2
Massachusetts	98.9	91.7	81.6	28.5	28.3	5.4	7.7	3.9	40.5
Michigan	99.7	85.1	84.6	48.5	38.8	9.1	7.2	4.6	37.4
Minnesota	99.1	69.5	79.9	41.5	34.5	1.4	2.4	2.4	39.9
Missouri	100.0	95.6	90.5	53.5	29.0	2.6	10.9	5.6	40.8
Montana	98.8	39.5	86.3	41.2	20.3	0.0	1.3	2.2	17.9
Nebraska	98.0	67.7	87.6	46.3	26.4	1.0	1.0	0.0	15.8
New Hampshire	100.0	79.7	76.9	20.8	20.2	0.0	1.8	0.6	31.9
New Jersey	100.0	84.0	83.6	30.6	31.3	4.7	13.6	4.9	46.9
New Mexico	100.0	75.5	91.4	46.4	6.2	10.9	5.9	2.2	48.5
New York	99.0	74.2	84.1	35.4	32.5	5.0	7.7	6.8	50.1
North Carolina	100.0	92.2	94.9	65.7	25.2	4.1	8.2	38.5	81.0
North Dakota	92.8	56.1	79.6	34.8	18.4	0.5	0.5	0.6	10.1
Oklahoma	98.8	74.6	92.5	67.9	14.6	2.7	2.2	8.1	26.6
Tennessee	99.6	95.3	92.5	65.7	22.0	5.6	5.2	23.2	55.5
Texas*	99.4	79.5	90.0	64.1	17.0	6.9	10.2	11.3	52.8
Utah	97.6	57.1	87.8	38.9	18.7	3.5	2.4	0.0	52.1
Vermont	94.0	75.7	63.6	16.4	9.5	0.0	0.0	0.0	22.5
Virginia	100.0	94.8	92.0	53.6	25.1	1.9	6.4	20.7	82.9
Wisconsin	98.7	74.8	78.4	54.3	46.3	1.1	3.4	5.8	30.4
<b>Unweighted Data</b>									
California	99.1	84.2	85.3	27.9	2.6	11.2	2.9	8.3	56.4
Oregon	100.0	66.1	88.5	33.7	28.0	1.1	0.5	1.6	48.9
Pennsylvania*	100.0	94.0	85.7	45.8	50.7	5.3	7.3	14.0	39.5
Rhode Island	100.0	92.3	87.7	24.2	35.4	0.0	3.1	1.5	32.8
South Carolina	100.0	95.4	92.8	70.6	14.0	3.8	36.6	28.1	90.4
Washington	99.6	73.9	89.0	28.7	16.0	2.0	7.3	3.3	45.7
<b>State Median</b>	<b>99.6</b>	<b>84.1</b>	<b>87.6</b>	<b>41.4</b>	<b>25.8</b>	<b>3.7</b>	<b>5.5</b>	<b>4.3</b>	<b>44.7</b>

(continued)

TABLE 25. Percentage of Schools That Implemented Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Required visitors to report to main office	Maintained a closed campus	Used staff to monitor halls	Checked bags, desks, and lockers	Prohibited backpacks	Required school uniforms	Required student identification badges	Used metal detectors	Had uniformed police
<b>Weighted Data</b>									
Chicago	100.0	96.9	94.0	61.5	28.3	75.7	20.7	67.0	90.3
Dallas	100.0	100.0	98.0	86.3	3.9	7.9	55.7	90.2	88.0
Fort Lauderdale	100.0	93.6	97.9	28.7	0.0	32.1	43.0	19.4	100.0
Houston	100.0	95.7	95.6	69.2	30.0	80.7	30.8	18.3	97.9
Los Angeles	100.0	98.1	95.1	91.1	3.9	43.4	8.7	90.3	86.6
Miami	100.0	90.1	91.6	64.7	4.3	36.3	38.8	49.4	93.6
Orange County	100.0	97.8	97.1	48.1	9.7	5.2	49.9	5.2	97.8
Palm Beach	100.0	97.3	95.1	51.3	5.2	15.0	38.8	10.6	100.0
San Bernardino	100.0	100.0	85.7	71.4	0.0	0.0	0.0	28.6	92.9
San Diego	100.0	100.0	93.9	6.3	4.1	24.5	0.0	0.0	77.6
San Francisco	100.0	73.1	91.5	21.4	2.5	26.4	0.0	0.0	75.6
<b>Unweighted Data</b>									
District of Columbia	100.0	96.7	100.0	96.7	30.0	40.0	76.7	96.7	93.3
Philadelphia	100.0	97.7	96.6	47.7	21.8	96.6	34.5	28.7	88.5
<b>Local Median</b>	<b>100.0</b>	<b>97.3</b>	<b>95.1</b>	<b>61.5</b>	<b>4.3</b>	<b>32.1</b>	<b>34.5</b>	<b>28.6</b>	<b>92.9</b>

\* Survey did not include students from one of the state's large school districts.



TABLE 26. Percentage of Schools with a Written Policy on HIV\*-Infected Students or School Staff and, Among Those Schools, Percentage That Addressed Specific Topics, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Had a written policy	Topic addressed by a written policy							
		Attendance of students**	Protection from discrimination <sup>§</sup>	Maintenance of confidentiality <sup>§</sup>	Worksite safety	Confidential counseling <sup>§§</sup>	Communication of policy <sup>+</sup>	Training for school staff about HIV infection	Procedure for implementing the policy
<b>Weighted Data</b>									
Alabama	69.8	96.3	97.8	98.4	99.0	88.2	90.3	87.5	95.4
Alaska	48.1	92.5	97.3	99.1	96.8	78.6	90.0	82.8	87.8
Arizona	68.5	90.9	93.7	96.7	96.3	74.2	82.8	83.0	91.7
Arkansas	49.7	84.8	94.1	97.2	92.3	83.2	87.2	69.2	84.9
Connecticut	73.8	88.4	97.8	100.0	100.0	79.1	84.8	87.8	89.9
Delaware	58.7	89.4	92.9	96.3	96.3	81.9	85.0	81.9	89.2
Georgia	66.6	90.3	95.0	99.4	98.8	87.9	91.3	83.3	87.4
Hawaii	60.4	82.6	100.0	100.0	100.0	72.5	85.1	86.0	86.0
Idaho	70.3	96.3	96.7	94.5	95.3	81.6	82.4	65.5	86.7
Illinois <sup>++</sup>	58.2	92.9	96.7	99.0	96.9	77.3	86.5	85.0	90.6
Iowa	65.3	89.5	92.2	95.3	98.1	68.0	85.7	89.1	92.4
Kentucky	32.5	86.4	92.9	97.4	98.3	77.0	82.6	92.1	90.9
Maine	76.8	91.9	96.7	98.7	97.3	80.7	82.7	90.6	92.0
Massachusetts	74.3	92.6	97.9	98.6	99.6	79.1	85.8	77.4	89.8
Michigan	60.7	84.5	90.8	94.9	96.1	66.7	78.4	83.8	85.1
Minnesota	54.0	88.8	97.8	98.6	97.9	69.6	87.0	85.7	89.6
Missouri	69.2	92.0	97.4	98.7	98.3	77.6	86.8	84.0	91.7
Montana	70.9	90.5	94.7	97.7	95.1	79.4	86.6	81.5	87.9
Nebraska	70.7	89.7	95.8	97.2	98.1	58.9	85.9	74.6	86.7
New Hampshire	83.8	88.1	94.7	98.5	96.2	70.8	80.3	79.2	86.3
New Jersey	78.0	95.1	98.0	99.6	99.2	81.1	87.5	91.7	95.8
New Mexico	61.6	88.6	94.9	98.7	91.0	81.0	80.3	82.0	89.3
New York	72.3	83.1	96.7	99.0	97.1	76.5	90.1	84.1	91.6
North Carolina	56.6	88.1	96.6	96.4	100.0	80.9	84.5	90.9	92.9
North Dakota	61.3	86.8	92.6	94.3	96.4	78.4	88.9	86.0	83.9
Oklahoma	71.3	91.0	97.0	99.4	97.1	77.9	90.6	93.4	94.2
Tennessee	72.9	89.5	95.6	97.8	95.4	77.1	88.0	88.4	92.3
Texas <sup>++</sup>	60.2	85.4	95.5	96.3	94.5	77.3	84.7	78.8	88.1
Utah	68.8	90.6	97.1	97.1	97.8	71.3	78.5	80.9	86.9
Vermont	90.3	96.1	99.1	100.0	97.0	77.3	89.9	87.7	92.7
Virginia	67.1	90.1	94.7	98.9	98.3	72.9	78.8	85.5	89.5
Wisconsin	56.7	86.8	94.4	95.9	97.8	73.4	80.8	84.6	88.1
<b>Unweighted Data</b>									
California	56.7	92.0	96.0	97.7	97.2	74.0	80.6	80.4	86.9
Oregon	85.0	92.4	98.6	99.3	98.6	79.5	83.0	87.1	89.7
Pennsylvania <sup>++</sup>	66.8	90.7	96.9	98.4	99.0	75.5	82.2	78.5	90.1
Rhode Island	84.6	88.7	94.3	100.0	98.1	86.5	90.2	84.3	92.2
South Carolina	73.3	92.0	96.1	98.3	99.4	83.3	84.6	93.2	92.1
Washington	63.8	91.1	95.0	97.2	97.9	77.5	81.6	92.3	89.4
<b>State Median</b>	<b>67.8</b>	<b>90.2</b>	<b>96.0</b>	<b>98.4</b>	<b>97.8</b>	<b>77.5</b>	<b>85.1</b>	<b>84.4</b>	<b>89.6</b>

(continued)

TABLE 26. Percentage of Schools with a Written Policy on HIV\*-Infected Students or School Staff and, Among Those Schools, Percentage That Addressed Specific Topics, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002  
(continued)

## LOCAL SURVEYS

Site	Had a written policy	Topic addressed by a written policy							
		Attendance of students**	Protection from discrimination <sup>§</sup>	Maintenance of confidentiality <sup>§</sup>	Worksite safety	Confidential counseling <sup>§§</sup>	Communication of policy <sup>+</sup>	Training for school staff about HIV infection	Procedure for implementing the policy
<b>Weighted Data</b>									
Chicago	65.5	93.3	97.0	97.7	94.2	86.0	94.0	82.3	90.4
Dallas	61.9	90.4	100.0	100.0	100.0	89.9	86.2	79.2	86.5
Fort Lauderdale	90.2	74.7	97.0	100.0	100.0	85.2	91.3	91.3	91.3
Houston	71.5	93.6	93.6	96.9	93.4	93.6	96.9	87.0	96.9
Los Angeles	72.8	90.3	97.2	98.6	100.0	82.0	86.2	79.5	90.1
Miami	85.5	87.7	98.6	100.0	98.6	93.3	96.1	93.4	97.3
Orange County	56.5	90.7	100.0	100.0	100.0	100.0	95.6	100.0	100.0
Palm Beach	66.1	96.2	95.8	100.0	100.0	87.7	87.0	100.0	91.3
San Bernardino	92.9	92.3	92.3	92.3	92.3	92.3	84.6	84.6	92.3
San Diego	87.5	97.6	100.0	100.0	100.0	75.6	95.1	100.0	100.0
San Francisco	80.1	92.9	100.0	100.0	96.1	87.9	92.9	74.8	85.2
<b>Unweighted Data</b>									
District of Columbia	90.0	96.3	96.3	96.3	96.3	59.3	92.6	85.2	92.6
Philadelphia	62.8	92.2	96.1	100.0	98.0	90.2	86.3	72.5	90.2
<b>Local Median</b>	<b>72.8</b>	<b>92.3</b>	<b>97.0</b>	<b>100.0</b>	<b>98.6</b>	<b>87.9</b>	<b>92.6</b>	<b>85.2</b>	<b>91.3</b>

\* HIV = human immunodeficiency virus.

\*\* With HIV infection.

§ For HIV-infected students and staff members.

§§ For HIV-infected students.

+ To students, school staff, and parents.

++ Survey did not include students from one of the state's large school districts.

TABLE 27. Percentage of Schools That Involved Parents and Families in Health Education Activities, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Provided families with information*	Met with PTA/PTO**	Invited family members to attend a health education class
<b>Weighted Data</b>			
Alabama	59.6	23.3	37.7
Alaska	51.3	24.8	42.5
Arizona	58.2	18.3	34.9
Connecticut	79.3	23.9	31.8
Delaware	84.1	29.4	42.5
Georgia	75.0	27.8	47.5
Hawaii	71.3	7.9	29.1
Idaho	59.5	8.7	44.3
Illinois <sup>§</sup>	60.9	10.4	30.4
Iowa	62.6	12.2	29.4
Kentucky	60.3	14.3	34.7
Maine	73.6	15.8	33.8
Massachusetts	86.5	43.0	35.5
Michigan	72.5	27.7	41.8
Minnesota	63.4	16.3	37.9
Missouri	73.4	20.6	29.7
Montana	56.9	14.5	33.5
Nebraska	49.6	9.3	28.9
New Hampshire	80.7	21.4	29.5
New Mexico	59.5	24.5	48.8
New York	82.6	32.7	33.9
North Dakota	51.9	11.6	24.1
Tennessee	57.3	28.6	36.7
Utah	76.4	20.3	49.6
Vermont	83.0	19.0	45.9
Virginia	71.8	25.0	39.4
Wisconsin	72.9	11.5	39.0
<b>Unweighted Data</b>			
Arkansas	47.2	11.8	27.8
California	69.5	21.1	36.4
North Carolina	70.2	22.8	43.9
Oklahoma	46.4	15.3	25.7
Pennsylvania <sup>§</sup>	72.8	26.0	37.0
Rhode Island	74.6	21.9	28.1
South Carolina	68.5	20.1	51.5
Texas <sup>§</sup>	53.7	20.2	31.3
<b>State Median</b>	<b>69.5</b>	<b>20.3</b>	<b>35.5</b>

(continued)

TABLE 27. Percentage of Schools That Involved Parents and Families in Health Education Activities, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (*continued*)

### LOCAL SURVEYS

Site	Provided families with information*	Met with PTA/PTO**	Invited family members to attend a health education class
<b>Weighted Data</b>			
Chicago	71.5	28.4	29.8
Dallas	52.7	26.3	47.6
Fort Lauderdale	81.8	25.0	48.5
Houston	73.0	29.4	61.5
Los Angeles	86.7	28.5	55.5
Miami	74.8	32.4	29.5
Orange County	88.3	14.4	56.6
Palm Beach	75.6	10.5	40.4
San Diego	89.8	64.6	56.3
San Francisco	91.0	62.0	47.1
<b>Unweighted Data</b>			
District of Columbia	71.0	38.7	64.5
Philadelphia	64.7	18.8	34.1
San Bernardino	87.5	37.5	57.1
<b>Local Median</b>	<b>75.6</b>	<b>28.5</b>	<b>48.5</b>

\* On the health education program.

\*\* To discuss the health education program.

§ Survey did not include students from one of the state's large school districts.

TABLE 28. Percentage of Schools in Which Students Participated in Health-Related Community Activities,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Performed volunteer work**	Attended a health fair	Gathered information <sup>§</sup>	Compared prices <sup>§§</sup>	Identified potential injury sites <sup>+</sup>	Analyzed advertising <sup>++</sup>	Advocated for health-related issues	Completed homework with family members
<b>Weighted Data</b>								
Alabama	23.8	33.6	59.7	34.0	72.7	66.9	51.1	82.5
Alaska	12.3	27.9	49.3	15.0	41.0	43.3	40.9	66.4
Arizona	23.6	32.2	50.2	18.9	48.1	52.8	49.1	69.6
Connecticut	23.0	25.2	55.0	17.8	50.5	71.1	54.9	76.8
Delaware	13.6	30.4	63.6	28.2	56.1	83.6	68.6	81.4
Georgia	20.8	36.3	64.3	26.2	62.4	72.1	51.7	77.8
Hawaii	23.4	34.9	61.3	27.6	42.8	82.4	67.1	78.5
Idaho	14.6	25.0	53.6	28.6	48.5	64.4	43.6	82.7
Illinois <sup>¶</sup>	10.6	16.6	42.9	22.2	51.4	60.3	30.4	73.8
Iowa	12.6	15.0	50.8	25.0	45.8	62.1	41.7	77.4
Kentucky	19.3	35.6	65.1	25.4	68.6	69.6	50.8	82.9
Maine	12.2	18.4	69.0	24.8	50.8	67.4	50.0	79.1
Massachusetts	16.6	26.8	55.7	23.3	51.4	73.4	56.1	78.3
Michigan	12.3	12.4	42.8	15.1	39.7	65.1	46.9	86.8
Minnesota	14.6	22.8	58.8	21.3	51.8	73.0	50.3	81.2
Missouri	11.7	35.4	52.7	25.8	62.0	67.5	39.8	74.2
Montana	10.2	25.5	41.2	16.7	44.4	60.3	49.3	72.0
Nebraska	11.2	14.0	43.0	19.1	45.8	52.0	39.5	75.5
New Hampshire	13.9	19.7	54.2	19.6	42.6	61.6	47.4	79.1
New Mexico	16.4	35.5	63.6	30.5	52.5	73.8	54.2	84.4
New York	24.2	32.8	69.9	29.4	59.7	71.7	55.2	84.3
North Dakota	11.9	32.5	48.8	17.0	49.4	61.1	35.6	76.0
Tennessee	20.8	36.3	62.1	33.0	68.4	64.7	46.5	75.5
Utah	23.2	18.4	56.2	31.4	52.4	69.4	49.3	90.2
Vermont	13.6	29.8	54.3	18.6	43.7	70.2	50.3	82.8
Virginia	20.1	33.2	60.7	30.6	70.7	73.2	46.8	80.3
Wisconsin	14.1	13.7	54.4	26.0	44.4	65.2	49.0	82.9
<b>Unweighted Data</b>								
Arkansas	11.8	18.2	51.9	29.3	56.7	52.1	35.1	68.6
California	20.5	22.1	49.0	22.9	44.6	68.5	44.8	81.4
North Carolina	13.1	31.9	54.4	24.6	64.9	64.6	44.0	75.7
Oklahoma	23.4	32.5	53.2	27.6	63.2	47.4	36.4	71.6
Pennsylvania <sup>¶</sup>	19.0	29.3	55.0	22.9	54.8	70.5	40.6	78.3
Rhode Island	21.7	28.3	71.7	35.6	52.5	73.3	68.3	71.7
South Carolina	18.9	28.3	50.0	28.9	57.0	61.7	44.6	71.1
Texas <sup>¶</sup>	13.7	20.9	47.9	18.7	50.7	57.1	42.1	70.9
<b>State Median</b>	<b>14.6</b>	<b>28.3</b>	<b>54.4</b>	<b>25.0</b>	<b>51.4</b>	<b>66.9</b>	<b>47.4</b>	<b>78.3</b>

(continued)

TABLE 28. Percentage of Schools in Which Students Participated in Health-Related Community Activities,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Performed volunteer work**	Attended a health fair	Gathered information <sup>§</sup>	Compared prices <sup>§§</sup>	Identified potential injury sites <sup>+</sup>	Analyzed advertising <sup>++</sup>	Advocated for health-related issues	Completed homework with family members
<b>Weighted Data</b>								
Chicago	19.1	38.9	51.7	27.4	62.2	55.6	49.7	82.9
Dallas	25.7	58.4	67.6	39.2	55.6	71.3	67.8	69.0
Fort Lauderdale	39.1	32.9	63.0	47.3	74.2	78.5	60.0	97.0
Houston	16.2	68.0	65.7	35.6	57.2	65.7	69.8	84.1
Los Angeles	27.4	41.7	74.0	55.9	59.5	88.1	62.9	92.7
Miami	39.1	46.0	65.4	37.9	69.7	79.1	61.9	85.0
Orange County	19.7	31.3	51.8	37.9	47.2	93.4	74.9	96.7
Palm Beach	56.4	46.3	71.3	52.0	56.4	69.0	70.0	91.8
San Diego	2.4	19.0	21.4	0.0	19.0	69.0	35.7	45.2
San Francisco	33.8	42.9	56.5	27.9	48.7	85.7	56.5	80.5
<b>Unweighted Data</b>								
District of Columbia	53.6	71.4	75.0	35.7	53.6	60.7	50.0	75.0
Philadelphia	38.6	45.1	62.9	42.3	70.0	70.8	55.7	84.7
San Bernardino	16.7	16.7	66.7	16.7	50.0	83.3	50.0	50.0
<b>Local Median</b>	<b>27.4</b>	<b>42.9</b>	<b>65.4</b>	<b>37.9</b>	<b>56.4</b>	<b>71.3</b>	<b>60.0</b>	<b>84.1</b>

\* In a required health education course.

\*\* At a hospital, a local health department, or other community organization that addresses health issues.

§ About health services available in the community.

§§ Visited a store to compare prices of health products.

+ At school, home, or in the community.

++ Advertising in the community that was designed to influence health behaviors or health risk behaviors.

¶ Survey did not include students from one of the state's large school districts.