



Coeur d'Alene, Idaho
31 March – 4 April, 2003

Florida's Bioassessment Program Applications

Ellen McCarron

Russ Frydenborg

Florida Department of Environmental Protection



Mission Statement

- Provide cost-effective and accurate ecological information to enable legally defensible environmental decisions



Bioassessment Program Steps

1. **Classify environments (e.g., regionalization)**
2. **Standardize sampling methods**
3. **Develop assessment approach (IBI)**
4. **Perform biological surveys**
5. **Select metrics (positive biological signals)**
6. **Incorporate Quality Assurance activities**
7. **Incorporate training and testing (certification)**
8. **Integrate into programs**
9. **Report results (Ecosummaries)**
10. **Revise biocriteria**

1. Classify environments (regionalization)

DRAFT ECOREGIONS/SUBREGIONS OF FLORIDA

Glenn E. Griffith¹, James M. Omernik², and Suzanne M. Pierson¹

¹ManTech Environmental Technology, Inc.
Corvallis, Oregon 97333

²U.S. Environmental Protection Agency
Corvallis, Oregon 97333

SOUTHEASTERN PLAINS ECOREGION (65)

- Southern Pine Plains and Hills (65f)
- Dougherty/Marianna Plains (65g)
- Tifton Upland/Tallahassee Hills (65h)

SOUTHERN COASTAL PLAIN ECOREGION (75)


- Gulf Coast Flatwoods (75a)
- Southwestern Florida Flatwoods (75b)
- Central Florida Ridges and Uplands (75c)
- Eastern Florida Flatwoods (75d)
- Okefenokee Swamps and Plains (75e)
- Sea Island Flatwoods (75f)

SOUTHERN FLORIDA COASTAL PLAIN ECOREGION (76)


- Everglades (76a)
- Big Cypress (76b)
- Miami Ridge/Atlantic Coastal Strip (76c)
- Southern Coast and Islands (76d)

● sampled reference site

Albers equal area projection



Bioregions of Florida



Panhandle

Peninsula

“Northeast”
(Peninsula)

Ecoregion
76 not
included

2. Standardize methods

Dipnet
Sampling





Stream Bioassessments (SCI + BioRecon)

- Habitat Assessment Procedures
- Physical/Chemical Characterization




3. Develop assessment approach (IBI)

- Stream Condition Index (20 dip net sweeps - macroinverts)
- BioRecon (4 dip net sweeps - macroinverts)
- Lake Condition Index (ponar grabs)
- Floristic Quality Index (macrophytes)
- Wetlands Condition Index (vegetation, macroinvertebrates, algae)



4. Perform bioassessments

5. Select metrics





Components of the Stream Condition Index (SCI)

Response to disturbance

- | | |
|---------------------------------|----------|
| ■ Taxa Richness | Decrease |
| ■ EPT Index | Decrease |
| ■ % Contribution Dominant Taxon | Increase |
| ■ Florida Index | Decrease |
| ■ # Chironomidae | Decrease |
| ■ % Filter-feeders | Decrease |
| ■ % Diptera | Increase |



Recalibrating SCI tool:

- *Leska Fore* – Statistical Design, Inc.
- First draft due spring '03
- 10 metrics for SCI

6. Incorporate QA






Bioassessment Program Quality Assurance Activities at FDEP

- Habitat Assessment Testing (Certification)
- SCI/BioRecon field audits
- Taxonomic Round Robins
- Intra-DEP Variability Studies
- Ongoing taxonomic manual updates
- Expert taxonomic verifications

Taxonomic RR Results





7. Training & Testing Program

- DEP - ongoing, continuous training
- Consultants
- Water Management Districts
- County and City Governments
- Regulated Industries



8. DEP Programs Using Bioassessments

App 1 - TMDL

App 2 - Springs Initiative

App 3 - Point Source Studies

App 4 - Ambient Monitoring

App 5 - RCRA (Hazardous Waste)

App 6 - Forestry BMP Effectiveness

App 7 - Mitigation Studies



Application 1: TMDL Program

- Impaired Waters Rule - Ch. 62-303, F.A.C.
- Collect biological, habitat, and water quality data to support FDEP's Impaired Waters Rule and TMDL Program
 - De-listing tool
 - Listing tool
 - Verification tool
 - Tool to evaluate watershed remediation
- 2 SCIs or Biorecons to list or de-list



**Application 2:
Florida Springs Program**


Pictured: Ichetucknee Springs



What's Different about Springs?

- More constant flow
- More alkaline (higher pH) water
- Better transparency
- More submerged aquatic vegetation
- Lower dissolved oxygen at boil
- Nitrate-nitrite levels increasing

Health of Spring-Dominated Streams Compared with Reference Sites



Springs Protection Strategy

Classification: Regional Model Springs

← Represent regional geologic/biological functions and forms basis for understanding range of human threats

Threat Identification

- Groundwater withdrawals
- Ditching/drainage to sinks
- Water quality alterations: toxics, nutrients, other
- Habitat alterations: recreational impacts, siltation, plant overgrowth, exotics

Corrective Action

- BMPs, education
- Land use/activity restrictions
- Purchase of critical areas
- *In situ* restoration

Application 3: Point Source Program

Fifth Year Inspections (NPDES)



Biological Assessment of
Bonifay Wastewater Treatment Facility
Holmes County, Florida
NPDES #FL0027731
Sampled April 2002

January 2003

Biology Section Bureau of Laboratories
Division of Resource Assessment & Management

Comprehensive Quality Assurance Plan #870346G

Florida Department of Environmental Protection

Fifth Year Inspection Summary

Discharger: Bonifay Waste Water Treatment Facility
County: Holmes
NPDES Number: FL0027731
Permit Expiration: 31 July 2002
Date Sampled: 15 and 16 April 2002

Toxics

Toxics Sampling Inspection (XSI)

Date Sampled: 15 April 2002

Results: Aluminum, cadmium, copper, iron, lead and zinc were found in the effluent at levels that complied with Class III Water Quality Standards. Silver was found in the effluent (0.25 µg/L) at a level that exceeded Class III Water Quality Standards (0.07 µg/L) (62-302.530(60) FAC). Diazinon (0.067 µg/L) was detected in the effluent. Total residual chlorine measured in the bioassay sample in the laboratory (0.54 mg/L) exceeded permit limits and Class III Water Quality Standards (FAC 62-302.530(19)) by a factor of more than 50.

Bioassay

Compliance Biomonitoring Inspection (CBI)

Date Sampled: 15 April 2002

Results: The effluent sample was toxic to both the fish, *Cyprinella leedsi*, and the water flea, *Ceriodaphnia dubia*, a violation of Class III Water Quality Standards (FAC 62-302.530 (62)). A dechlorinated aliquot of the effluent sample was not toxic to the fish, *Cyprinella leedsi*, or to the water flea, *Ceriodaphnia dubia*, during 48-hour acute screening bioassays.

Chemistry

Water Quality Inspection (WQI)

Date Sampled: 15 April 2002

Results: Effluent concentrations of ortho-phosphate (0.68 mg/L), total phosphorus (0.87 mg/L), ammonia (0.53 mg/L), nitrate+nitrite (0.28 mg/L), and total Kjeldahl nitrogen (1.5 mg/L) contributed to enrichment of nitrogen and phosphorus at the Test Site. The Test Site values were found to be greater than 40-80% of other Florida waters. In contrast, the nutrients at the Control Site were only greater than 20-30% of typical of Florida streams, except ammonia (0.072 mg/L), which was greater than about 40% of typical Florida streams. The Control Site AGP value (6.0 mg dry weight/L) was just over the "problem threshold" of 5.0 mg dry weight/L, while the Test Site value (18.5 mg dry weight/L) was more than three times the problem threshold. The effluent AGP was 30.8 mg dry weight/L. These results are a further indication of enrichment downstream of the facility due to the effluent.

Bioassessment



Impact Bioassessment Inspection (IBI)

Date Sampled: 15 and 16 April 2002

Results: Macroinvertebrate community data suggest degradation at the Test Site related to the facility's effluent. The Shannon-Weaver Diversity Index was 63% lower at the Test Site (0.9) compared to the Control Site (2.4), a violation of the Class III Biological Integrity Criterion (62-302.530 (11) FAC). Quantitative macroinvertebrate data showed 95.6% of the total community consisted of dipterans and the number of individuals rose from 87 at the Control Site to 1,811 at the Test Site. These results may indicate an increase in productivity in response to nutrient enrichment from the facility. In qualitative dipnet data, the dominant taxon comprised 81.6% of the community compared with 25.5%, and taxa richness was 68% lower at the Test Site compared to the Control Site. The Control Site received a SCI score of 21, which placed it in the "Good" category, while the Test Site received the lowest possible score of 7, placing it in the "Very Poor" category. Periphyton cell density was twice as high at the Test Site compared to the Control Site, reflecting a ten-fold increase in percentage of green algae at the Test Site. This change in algal community structure may also be related to the facility's effluent. The SCI and algal community composition data indicate an imbalance of flora or fauna, a violation of FAC 62-302.530 (48)(b).


Control (upstream) site "A"



Test (downstream) site "B"



**WWTP
pipe discharge**



Control Site A

Table 6. Stream Condition Index

Bonifay WWTF Control Site	Value	5	3	1	Score
Total Number of Taxa	22	≥31	30-16	<16	3
Number of EPT Taxa	0	≥7	6-4	<4	1
Number of Chironomid Taxa	9	≥9	8-5	<5	5
Percent Contribution of Dominant Taxon	25.5	≤22	23-61	>61	3
Percent Diptera	16.4	-	≤50	>50	3
Florida Index	2	≥16	15-8	<8	1
Percent Suspension Feeders and Filterers	23.2	≥12	11-6	<6	5
Total Score		Panhandle			21
Interpretation of Scores by Region		Excellent			27-33
		Good			21-26
		Poor			14-20
		Very Poor			7-13

Summer Index Period: April 1-October 31. Stream Condition Index (SCI) for Florida Panhandle. Values calculated from benthic macroinvertebrates collected with 20 standardized dipnet sweeps (Barbour *et al.* 1996a, b).

Test Site B


Bonifay WWTF Test Site	Value	5	3	1	Score
Total Number of Taxa	7	≥31	30-16	<16	1
Number of EPT Taxa	0	≥7	6-4	<4	1
Number of Chironomid Taxa	4	≥9	8-5	<5	1
Percent Contribution of Dominant Taxon	81.6	≤22	23-61	>61	1
Percent Diptera	88.6	-	≤50	>50	1
Florida Index	1	≥16	15-8	<8	1
Percent Suspension Feeders and Filterers	0.9	≥12	11-6	<6	1
Total Score		Panhandle			7
Interpretation of Scores by Region		Excellent			27-33
		Good			21-26
		Poor			14-20
		Very Poor			7-13

Summer Index Period: April 1-October 31. Stream Condition Index (SCI) for Florida Panhandle. Values calculated from benthic macroinvertebrates collected with 20 standardized dipnet sweeps (Barbour *et al.* 1996a, b).



Application 4: Ambient Monitoring Program

- Probabilistic network
- 5-year rotating basin program



Basin Rotation Groups and Schedule

Map prepared April 23, 2002 by the Bureau of Watershed Management, Division of Water Resource Management. This map is a representation of ground conditions and is not intended for delineations or analysis of the features shown.
 For more information or copies, contact Hollie Brandt at (850) 921-9469, or hollie.brandt@dep.state.fl.us
 Location: \\GERRAID\GIS2sub\atlaz\basin_map

Legend

Water Management Districts

Basin Rotation Groups and Schedule

- Group 1
- Group 2
- Group 3
- Group 4
- Group 5

County Boundaries




20 0 20 40 Miles





Ambient Monitoring Program – cont.

- Selected tools = Stream Condition Index, and Floristic Quality Index
- To describe condition of individual rotating basins (yearly)
- To describe statewide conditions (5 years)
-- 305(b) report
- To report on effectiveness of all water programs going on both statewide and in each basin



Application 5: Resource Conservation and Recovery Act (RCRA)

- Joint endeavor by DEP Waste Division and Water Division
- SCI tool requested by Waste Mgt. Division

RCRA Site





Hazardous Waste (RCRA) Studies

- Concern: leachate from waste sites impacting aquatic systems
- Designed ecological assessments with Waste Management staff
- Answers the question: Is leachate affecting nearby stream communities?



Application 6: Effectiveness of Forestry Best Management Practices

- Joint project between Florida DEP, Florida Department of Agriculture and the silviculture industry.
- Purpose: to determine if forestry BMPs, when properly applied, protect aquatic biota in adjacent streams




Examples of forestry BMPs


- Undisturbed buffer zone (SMZ)
- Site preparation to prevent erosion
- Control fertilizers and pesticides
- Design roads/drainage easements for minimum erosion/deposition

Experimental Design


**Before
Silviculture
Activities**



**After
Silviculture
Activities**



San Julia SCI Results





Application 7: Wetland Mitigation Program

- Reclaimed phosphate streams
- Wetland restoration sites

9. Reporting



EcoSummary

BioRecon Report



White Oak Creek below County Road 191 Santa Rosa County March 16, 2000

BioRecon: A rapid, cost-effective screening mechanism for identification of biological impairment

Purpose

A bioassessment was performed on White Oak Creek in an effort to document the environmental health of streams flowing into the Yellow River Aquatic Preserve (YRAP). The BioRecon was conducted in partnership with the YRAP staff.

Background

White Oak Creek at the bioassessment site is a second order stream originating above I-10 about 5 miles south of Milton in Santa Rosa County (Lat. 30° 32' 00.5" Long. 87° 03' 08.5"). This stream flows to White Oak Bayou in Blackwater Bay, at Eagle Point, then into Pensacola Bay. This site drains the Gulf Coast Flatwoods subcoregion (75a).



Results

The BioRecon indicated an impaired biological community. All 3 biological indicators failed thresholds established for a healthy aquatic wildlife community:

Biometrics	Value	Thresholds
Taxa Richness	17	≥24
Florida Index	4	≥22
EPT	3	≥17

The biota was dominated by pollution tolerant aquatic wildlife. The impoundment created by the CR-191 road culvert caused water quality problems in White Oak Creek. The dissolved oxygen concentrations (3-4.7 mg/l in March, 2 mg/l in May) did not meet State Water Quality Standards. The May biochemical demand was very high (5.1 mg/l) with elevated nutrients (ammonia 89 ug/l, nitrogen 1000 ug/l, phosphorus 61 ug/l) present. The elevated nutrients led to an algae bloom with a very high chlorophyll A concentration of 56 ug/l. Sediments were anaerobic with a hydrogen sulfide odor. Silt smothering of fish and wildlife habitats was severe.



Significance

This White Oak Creek site did not meet Class III State Water Quality Standards 62-302 for recreation and the propagation and maintenance of a healthy, well-balanced population of fish and wildlife. The dam effects of the CR191 culvert crossing contributed to White Oak Creek not meeting State designated use. Organic sediment and nutrient loading from the impoundment created an elevated biochemical oxygen demand (5.1mg/l) that caused oxygen depletion. Reduced stream flow from the damming negatively affect the system's fish and wildlife nursery function. Altered habitats from culverts and clearing streambank riparian zones negatively affects the Yellow River Aquatic Preserve's fish and wildlife in White Oak Bayou and the lower portion of Blackwater Bay. Reduced flows, nutrient, and organic sediment loading could affect the federally endangered Gulf of Mexico sturgeon.

Suggestions


Restoration of the stream's natural hydrology (i.e. flow, quantity) and preserving riparian wetland forests buffer zones would benefit the Aquatic Preserve's fish and wildlife community including the sturgeon. Replacing culverts with bridges that span the streamside wetlands and restoration of the watershed's natural hydrological patterns could enhance Live Oak Bayou as a nursery area for aquatic wildlife in the Pensacola Bay basin. Nutrient sources that created the algae bloom were unknown. Background nutrient concentrations in area watersheds are naturally very low. Possibly fallout from 2 major industrial manufacturing air pollution dischargers within 3 and 5 miles of the watershed is a nutrient source. For more information, contact Donald Ray, FDEP Northwest District, 160 Government Center, Pensacola, FL 32501 (850) 595-8300 x1126 or SC 695-8300




10. Develop or revise biocriteria

- Integrate into Tiered Aquatic Life Use System (TALUS)

Tiered Aquatic Life Uses: Draft Conceptual Framework



Tiered Aquatic Life Use Conceptual Model: Draft Biological Tiers





Conclusions

- FDEP Bioassessment Program provides practical support for a variety of FDEP programs
- QA and training are critical for demonstrating legal defensibility

It's all about clean water for future generations!

