

Wisconsin, 2006

Primary and secondary syphilis

- **CASES:** Wisconsin reported 68 cases in 2006.
- **RATES:** Wisconsin ranked 38 among 50 states, DC, and 3 territories with 1.2 cases per 100,000 population compared to the total rate of 3.3 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Wisconsin, the rate among males was 2.4 per 100,000 population compared to the total male rate of 5.7 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the total female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.8 among whites, 5.3 among blacks, 3.6 among Hispanics, 0.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 6.6 times that of whites.
- The rate among Hispanics was 4.5 times that of whites.
- The rate among Asian/Pacific Islanders was 1.1 times that of whites.

Congenital syphilis

- **CASES:** Wisconsin reported 0 cases in 2006.
- **RATES:** Wisconsin had a rate of 0.0 cases per 100,000 live births compared to the total rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 1997-2006



Total P&S Syphilis Rates by County, Wisconsin, 2006

