

Vaccination Schedule for Children



Hepatitis B¹

DTP
(Diphtheria,
Tetanus,
Pertussis)

Polio

Hib
(Haemophilus
influenza type

MMR
(Measles,
Mumps,
Rubella)

Td
(Tetanus,
Diphtheri

Chickenpox

	Hepatitis B ¹	DTP (Diphtheria, Tetanus, Pertussis)	Polio	Hib (Haemophilus influenza type	MMR (Measles, Mumps, Rubella)	Td (Tetanus, Diphtheri	Chickenpox
Birth	✓						
1-2 months	✓						
2 months		✓	✓	✓			
4 months		✓	✓	✓			
6 months	✓	✓		2			
12-18 months				3			5
15 months					✓		
15-18 months		✓	✓				
4-6 years		✓	✓		4		
14-16 years						✓	

1 — This schedule is for high-risk infants (those whose mothers have tested positive for hepatitis B). These infants should receive hepatitis B immune globulin at or shortly after the first vaccine dose. Infants not at high risk also should be vaccinated against hepatitis B. Check with your doctor for the appropriate schedule.

2 — May not be required, depending on type of vaccine used. Check with your doctor.

3 — Given at 12, 15, or 18 months, depending on type of vaccine. Check with your doctor.

4 — Where required by public health authorities for school entry. Otherwise: 11-12 years. Check with your doctor.

5 — Can be given along with MMR or anytime between 12 to 18 months. Older children who have not had chickenpox and have not been vaccinated previously should get the vaccine by their 13th birthday.

(Source: U.S. Food and Drug Administration)