

## 2. HEALTH EFFECTS

### 2.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of methylene chloride. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

### 2.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals and others address the needs of persons living or working near hazardous waste sites, the information in this section is organized first by route of exposure (inhalation, oral, and dermal) and then by health effect (death, systemic, immunological, neurological, reproductive, developmental, genotoxic, and carcinogenic effects). These data are discussed in terms of three exposure periods: acute (14 days or less), intermediate (15–364 days), and chronic (365 days or more).

Levels of significant exposure for each route and duration are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. "Serious" effects are those that evoke failure in a biological system and can lead to morbidity or mortality (e.g., acute respiratory distress or death). "Less serious" effects are those that are not expected to cause significant dysfunction or death, or those whose significance to the organism is not entirely clear. ATSDR acknowledges that a considerable amount of judgment may be required in establishing whether an end point should be classified as a NOAEL, "less serious" LOAEL, or "serious" LOAEL, and that in some cases, there will be insufficient data to decide whether the effect is indicative of significant dysfunction. However, the Agency has established guidelines and policies that are used to classify these end points. ATSDR believes that there is sufficient merit in this approach to warrant an attempt at distinguishing between "less serious" and "serious" effects. The distinction between "less serious" effects and "serious" effects is

## 2. HEALTH EFFECTS

considered to be important because it helps the users of the profiles to identify levels of exposure at which major health effects start to appear. LOAELs or NOAELs should also help in determining whether or not the effects vary with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown in the Levels of Significant Exposure (LSE) tables and figures may differ depending on the user's perspective. Public health officials and others concerned with appropriate actions to take at hazardous waste sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAEL) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans (Minimal Risk Levels or MRLs) may be of interest to health professionals and citizens alike.

Levels of exposure associated with carcinogenic effects (Cancer Effect Levels, CELs) of methylene chloride are indicated in Tables 2-1 and 2-2 and Figures 2-1 and 2-2. Because cancer effects could occur at lower exposure levels, Figures 2-1 and 2-2 also show a range for the upper bound of estimated excess risks, ranging from a risk of 1 in 10,000 to 1 in 10,000,000 ( $10^{-4}$  to  $10^{-7}$ ), as developed by EPA.

Estimates of exposure levels posing minimal risk to humans (Minimal Risk Levels or MRLs) have been made for methylene chloride. An MRL is defined as an estimate of daily human exposure to a substance that is likely to be without an appreciable risk of adverse effects (noncarcinogenic) over a specified duration of exposure. MRLs are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration within a given route of exposure. MRLs are based on noncancerous health effects only and do not consider carcinogenic effects. MRLs can be derived for acute, intermediate, and chronic duration exposures for inhalation and oral routes. Appropriate methodology does not exist to develop MRLs for dermal exposure.

Although methods have been established to derive these levels (Barnes and Dourson 1988; EPA 1990d), uncertainties are associated with these techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of the procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

## 2. HEALTH EFFECTS

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

### 2.2.1 Inhalation Exposure

#### 2.2.1.1 Death

Case studies of methylene chloride poisoning during paint stripping operations have demonstrated that inhalation exposure can be fatal to humans (Bonventre et al. 1977; Hall and Rumack 1990; Stewart and Hake 1976). Although quantitative estimates of exposure levels were not reported for these cases, levels of methylene chloride in various tissues were reported: liver (14.4 mg/dL), blood (51 mg/dL), serum (29 µg/mL), and brain (24.8 mg/100 g) (Bonventre et al. 1977; Hall and Rumack 1990). The cause of death in these cases was uncertain; however, myocardial infarction was reported in one case (Stewart and Hake 1976). Death also occurred in two workers involved in oleoresin extraction processes and liquid cleaning operations (Moskowitz and Shapiro 1952; Winek et al. 1981). Exposure reportedly occurred from less than 1 hour up to 3 hours, but the concentration of methylene chloride was not reported. The compound was detected in the lung (0.1 mL/500 g wet tissue), brain (0.27 g/L), and blood (29.8 mg%) (Moskowitz and Shapiro 1952; Winek et al. 1981). Two cases of lethal poisoning following acute inhalation of extremely high concentrations of methylene chloride in air (estimated as up to 168,000 ppm) occurred in two workers burying barrels containing mixed solvents and solid chemical waste in a well about 2 meters below ground level (Manno et al. 1992). Methylene chloride concentrations in blood of the two workers were 572 and 601 mg/L, respectively. Blood carboxyhemoglobin (COHb) concentrations were about 30% higher than normal. Death appears to have been caused by narcosis and respiratory depression due to the acute effects of high concentration methylene chloride on the central nervous system. One death in the United Kingdom resulting from acute inhalation occupational exposure to methylene chloride (concentration not provided) was attributed to acute narcosis resulting in respiratory depression (Bakinson and Jones 1985); necropsy revealed evidence of liver and spleen congestion. One case of fatal gassing occurred during paint-stripping of a chemical tank (Tay et al. 1995). The paint stripper contained 74% w/w (weight by weight) of methylene chloride; the concentration of methylene chloride vapor within the tank was later estimated to have been well above 100,000 ppm. The worker did not wear a respirator and did not engage a forced ventilation system into the confined space of the tank. He was found unconscious and died 4 days later. His methylene chloride blood concentration was 281 mg/L.

## 2. HEALTH EFFECTS

Another case of fatal poisoning was reported in Korea: the body of the chief executive of a painting and coating factory was discovered in an underground trichloroethylene tank in which methylene chloride was being used to removed rust from iron sheets (Kim et al. 1996a). Autopsy revealed pronounced organ congestion in the brain, kidneys, liver, and lungs; blood tissue samples contained 3% COHb and 252 mg/L of methylene chloride. This methylene chloride blood concentration is similar to that reported for other fatalities (Manno et al. 1992; Tay et al. 1995). Concentrations of methylene chloride in other tissues ranged from 26 mg/kg in the lungs to 75 mg/kg in the brain (Kim et al. 1996a). The data were sufficient to confirm that the death was due to methylene chloride poisoning.

In another study, no increase in deaths in methylene chloride workers, assessed by life table analysis, was found after exposure to 30–120 ppm (time weighted averages) for over 30 years (Friedlander et al. 1978). Fiber production workers exposed to methylene chloride (140–475 ppm) for at least 3 months did not have a significant increase in mortality (Lanes et al. 1993; Ott et al. 1983b).

Studies in animals confirm that methylene chloride may be lethal after inhalation exposure at high concentrations. Acute exposure to 16,000–19,000 ppm of methylene chloride for 4–8 hours caused death in rats and mice (NTP 1986; Svirbely et al. 1947). Also, one of four female monkeys died after 10 days of continuous exposure to 5,000 ppm methylene chloride (MacEwen et al. 1972). Data suggest there is a narrow margin between concentrations causing anesthesia and death. An  $LC_{50}$  of 16,189 ppm was reported in mice acutely exposed to methylene chloride (Svirbely et al. 1947). No deaths were found in mice exposed for 4 hours to 16,800 ppm, but 70% of the mice exposed to 17,250 ppm died (NTP 1986). Repeated exposure from intermediate to lifetime duration at levels ranging from 1,000 to 16,000 ppm can cause increased deaths in rats, mice, guinea pigs, rabbits, and dogs (Burek et al. 1984; Heppel et al. 1944; NTP 1986). Results of the different inhalation studies described in the NTP (1986) report illustrate that with increasing duration, the lethal exposure level decreases. A 19-day intermittent exposure to 6,500 ppm of methylene chloride or a 13-week exposure to 4,200 ppm were not lethal to rats or mice, but exposure for 2 years at 4,000 ppm reduced survival in female rats and in mice of both sexes (NTP 1986). Although the same target organs (central nervous system, lungs, liver) are affected in mammals, the available mortality data suggest differences in sensitivity among species, with dogs being more sensitive than mice and rats.

An  $LC_{50}$  value and all reliable LOAEL values for lethality in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation

Key to figure	Species (strain)	Exposure/ duration/ frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
<b>ACUTE EXPOSURE</b>							
<b>Death</b>							
1	Monkey (Rhesus)	10 d 24 hr/d				5000 F (1/4 died)	MacEwen et al. 1972
2	Rat (Fischer- 344)	4 hr				17250 death in 2/15	NTP 1986
3	Mouse (B6C3F1)	4 hr				17250 death in 7/10	NTP 1986
4	Mouse (Swiss-Webster)	8 hr				16189 (LC <sub>50</sub> )	Svirbely et al. 1947
<b>Systemic</b>							
5	Human	8 hr/d 3 d	Cardio	100 M			Cherry et al. 1981
6	Human	8 hr	Cardio	475			Ott et al. 1983c
7	Rat (Sprague-Dawley)	Gd 6-15 7 hr/d	Hepatic	1250 F			Schwetz et al. 1975
8	Mouse (Swiss-Webster)	Gd 6-15 7 hr/d	Hepatic		1250 F (increased absolute liver weight)		Schwetz et al. 1975
9	Gn Pig (Hartley)	6 hr	Hepatic		5200 M (increased hepatic triglycerides)		Morris et al. 1979

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
10	Dog (NS)	6 d 4 hr/d	Hepatic		10000 F (moderate centrilobular fatty degeneration)		Heppel et al. 1944
11	Rabbit (New Zealand)	10 min	Ocular		490M (increased corneal thickness and intraocular tension)		Ballantyne et al. 1976
<b>Neurological</b>							
12	Human	24 hr			300 F (decreased critical flicker frequency)		Fodor and Winneke 1971
13	Human	4 hr			200 (decreased eye-hand coordination and peripheral visual response and auditory function)		Putz et al. 1979
14	Human	1-2 hr			515M (altered visual evoked response)		Stewart et al. 1972
15	Human	3-4 hr			300 <sup>b</sup> F (decreased critical flicker frequency and auditory vigilance)		Winneke 1974
16	Monkey (NS)	1 d 4 hr/d				10000 F (reduced activity; incoordination)	Heppel et al. 1944
17	Rat (NS)	5 d/wk 4 hr/d				10000 (gait disturbance, somnolence, prostration, respiratory depression)	Heppel et al. 1944
18	Rat (Fischer- 344)	60 min			5000M (increased latency of auditory evoked potential)		Rebert et al. 1989

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
19	Rat (Wistar)	2 wk 5 d/wk 6 hr/d			500 M (decreased succinate dehydrogenase activity in cerebellum)		Savolainen et al. 1981
20	Gn Pig (NS)	5 d/wk 7 hr/d			10000 M (somnolence)		Heppel et al. 1944
21	Dog (NS)	6 d 4 hr/d				10000 (incoordination, excitability, hyperactivity)	Heppel et al. 1944
22	Rabbit (NS)	5 d/wk 4 hr/d				10000 M (excitement, inactivity, postural disturbance)	Heppel et al. 1944
<b>Developmental</b>							
23	Rat (Sprague-Dawley)	Gd 6-15 7 hr/d			1250 (increased incidence of dilated renal pelvis and delayed ossification of sternebra)		Schwetz et al. 1975
24	Mouse (Swiss-Webster)	Gd 6-15 7 hr/d			1250 (increased incidence of extra sternal ossification center)		Schwetz et al. 1975
<b>INTERMEDIATE EXPOSURE</b>							
<b>Death</b>							
25	Rat (NS)	8 wk 5 d/wk 4 hr/d				10000 (death in 2/16)	Heppel et al. 1944
26	Rat (Fischer-344)	19 d 5 d/wk 6 hr/d				16000 (death in 9/10)	NTP 1986

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
27	Mouse (ICR)	4 wk 7 d/wk 24 hr/d				5000 F (LC <sub>30</sub> )	MacEwen et al. 1972
28	Mouse (B6C3F1)	19 d 5 d/wk 6 hr/d				13000 (death in 7/10)	NTP 1986
29	Mouse (B6C3F1)	13 wk 5 d/wk 6 hr/d				8400 (LC <sub>50</sub> )	NTP 1986
30	Gn Pig (NS)	6 mo 5 d/wk 7 hr/d				5000 M (death in 3/8)	Heppel et al. 1944
31	Dog (Beagle)	14 wk 7 d/wk 24 hr/d				1000 F (death in 6/8)	MacEwen et al. 1972
32	Rabbit (NS)	8 wk 5 d/wk 4 hr/d				10000 (death in 3/5)	Heppel et al. 1944
<b>Systemic</b>							
33	Monkey (NS)	100 d 24 hr/d	Hemato	100			Haun et al. 1972
			Hepatic	100			
			Renal	100			
34	Monkey (Rhesus)	14 wk 7 d/wk 24 hr/d	Hepatic		1000 F (fat accumulation)		MacEwen et al. 1972
			Bd Wt		1000 F (decreased weight gain)		



TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
35	Monkey (Rhesus)	4 wk 7 d/wk 24 hr/d	Hepatic		5000 F (atrophy; fatty change)		MacEwen et al. 1972
36	Rat (NS)	100 d 24 hr/d	Hepatic		25 <sup>c</sup>	(cytoplasmic vacuolization, fatty infiltration)	Haun et al. 1972
			Renal		25	(fatty infiltration, degenerative changes)	
37	Rat (NS)	8 wk 5 d/wk 4 hr/d	Resp			10000 (pulmonary congestion, edema, focal extravasion of blood)	Heppel et al. 1944
38	Rat (Sprague-Dawley)	14 wk 7 d/wk 24 hr/d	Hepatic		5000 M (iron pigmentation)		MacEwen et al. 1972
			Renal		5000 M (cortical tubular degeneration)		
39	Rat (Sprague-Dawley)	4 wk 7 d/wk 24 hr/d	Hepatic		5000 M (iron pigmentation, cellular vacuolization)		MacEwen et al. 1972
40	Rat (Wistar)	28 d 5 hr/d	Hepatic	250 M			Norpoth et al. 1974
41	Rat (Fischer-344)	13 wk 5 d/wk 6 hr/d	Resp			8400 (pneumonia)	NTP 1986
42	Mouse (NS)	100 d 24 h/d	Hepatic	25			Haun et al. 1972

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/ duration/ frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
43	Mouse (NMRI)	90 d 24 hr/d	Hepatic		75 (fatty infiltration; increased liver weight)		Kjellstrand et al. 1986
44	Mouse (ICR)	14 wk 7 d/wk 24 hr/d	Hepatic		1000 F (iron pigmentation nuclear degeneration; pyknotic cells)		MacEwen et al. 1972
45	Mouse (B6C3F1)	13 wk 5 d/wk 6 hr/d	Hepatic		4200 (centrilobular hydropic degeneration)		NTP 1986
46	Gn Pig (NS)	6 mo 5 d/wk 7 hr/d	Resp			5000 M (pneumonia)	Heppel et al. 1944
			Hepatic			5000 M (centrilobular fatty degeneration)	
47	Gn Pig (NS)	8 wk 5 d/wk 4 hr/d	Hepatic		10000 (slight to moderate fatty degeneration)		Heppel et al. 1944
48	Dog (NS)	100 d 24 hr/d	Hemato	100			Haun et al. 1972
			Hepatic	100			
			Renal	100			
49	Dog	14 wk 7 d/wk 24 hr/d	Hepatic		1000 F (fatty changes, increased enzyme)		MacEwen et al. 1972
			Renal		1000 F (renal tubular vacuolization)		
50	Rabbit (NS)	8 wks 5 d/wk 4 hr/d	Resp			10000 M (pulmonary congestion, edema, focal necrosis)	Heppel et al. 1944

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
<b>Immunological/Lymphoreticular</b>							
51	Rat (Sprague-Dawley)	28 d 5 d/wk 6 hr/d		5187			Halogenated Solvent Industry Alliance, Inc 2000
<b>Neurological</b>							
52	Rat (Long-Evans)	Gd 1-17		4500 F			Bornschein et al. 1980
53	Rat (Fischer-344)	13 wk 5 d/wk 6 hr/d		2000			Mattsson et al. 1990
54	Gerbil (Mongolian)	3 mo			210	(alterations in brain amino acids)	Briving et al. 1986
55	Gerbil (Mongolian)	7-16 wk 24 hr/d			210	(decreased hippocampal DNA concentrations)	Rosengren et al. 1986
<b>Reproductive</b>							
56	Rat (Fischer-344)	2 gen		1500			Nitschke et al. 1988b
57	Mouse (Swiss-Webster)	6 wk 5 d/wk 2 hr/d		200 M			Raje et al. 1988
<b>CHRONIC EXPOSURE</b>							
<b>Death</b>							
58	Rat (Sprague-Dawley)	2 yr 5 d/wk 6 hr/d				3500 (90% mortality)	Burek et al. 1984

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
<b>Systemic</b>							
59	Human	>1-5 yr 8 hr/d	Hemato	475			Ott et al. 1983a
60	Human	>10 yr	Cardio Hemato Hepatic	475 475 475			Soden 1993
61	Rat (Sprague-Dawley)	2 yr 5 d/wk 6 hr/d	Hemato  Hepatic	3500	500	(hepatocellular vacuolization, multinucleated hepatocytes)	Burek et al. 1984
62	Rat (Fischer-344)	102 wk 5 d/wk 6 hr/d	Resp  Renal		1000	(nasal cavity squamous metaplasia)	Mennear et al. 1988; NTP 1986
					2000 F	(tubular cell degeneration)	
63	Rat (Sprague-Dawley)	2 yr 5 d/wk 6 hr/d	Hemato  Hepatic	50  50 <sup>d</sup>	200	(significant increase in carboxyhemoglobin)	Nitschke et al. 1988a
					200 F	(multinucleated hepatocytes)	
64	Mouse (B6C3F1)	102 wk 5 d/wk 6 hr/d	Renal		4000	(increased incidence of kidney/tubule casts)	Mennear et al. 1988; NTP 1986

TABLE 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Key to figure <sup>a</sup>	Species (strain)	Exposure/duration/frequency	System	NOAEL (ppm)	LOAEL		Reference
					Less serious (ppm)	Serious (ppm)	
<b>Cancer</b>							
65	Rat (Fischer- 344)	102 wk 5 d/wk 6 hr/d				1000	(CEL: mammary fibroadenoma or adenocarcinoma) Mennear et al 1988; NTP 1986
66	Rat (Fischer- 344)	2 yr 5 d/wk 6 hr/d				500	(CEL mammary gland tumors) Nitschke et al. 1988a
67	Mouse (B6C3F1)	102 wk 5 d/wk 6 hr/d				2000	(CEL: liver and lung adenoma and carcinomas) Mennear et al. 1988; NTP 1986

<sup>a</sup>The number corresponds to entries in Figure 2-1.

<sup>b</sup>Used to derive an acute inhalation MRL of 0.6 ppm; the LOAEL was adjusted for a 24-hour exposure by Rietz et al. (1997), yielding a duration-adjusted LOAEL of 60 ppm which was divided by an uncertainty factor of 100 (10 for use of a LOAEL and 10 for human variability).

<sup>c</sup>Used to derive an intermediate inhalation MRL of 0.3 ppm, human equivalent concentration divided by an uncertainty factor of 90 (3 for the use of a minimal LOAEL, 3 for extrapolation from animals to humans, and 10 for human variability).

<sup>d</sup>Used to derive a chronic inhalation MRL of 0.3 ppm; human equivalent concentration adjusted for intermittent exposure and divided by an uncertainty factor of 30 (3 for extrapolation from animals to humans and 10 for human variability).

Cardio = cardiovascular; CEL = cancer effect level; d = day(s); Derm/oc = dermal/ocular; DNA = deoxyribonucleic acid; Gd = gestation day; gen = generation; Gn pig = guinea pig; Hemato = hematological; hr = hour(s); LC50 = lethal concentration, 50% kill; LOAEL = lowest-observed-adverse-effect level; min = minute(s); mo = month(s); NOAEL = no-observed-adverse-effect level; wk = week(s); yr = year(s)

Figure 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation  
Acute ( $\leq 14$  days)




Figure 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)  
Intermediate (15-364 days)





Figure 2-1. Levels of Significant Exposure to Methylene Chloride - Inhalation (continued)

Chronic (≥365 days)





## 2. HEALTH EFFECTS

**2.2.1.2 Systemic Effects**

No studies were located regarding musculoskeletal or dermal effects in humans or animals after inhalation exposure to methylene chloride. Effects of methylene chloride on the respiratory, cardiovascular, gastrointestinal, hepatic, renal, and ocular systems are discussed below.

The highest NOAEL values and all reliable LOAEL values for systemic effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

**Respiratory Effects.** Asphyxia was determined to be the cause of death in the case of a male worker who was subjected to acute inhalation exposure (concentration unknown) for 1 hour (Winek et al. 1981); the autopsy revealed bilateral pulmonary congestion with focal hemorrhage. Respiratory symptoms (cough, breathlessness, chest tightness) were reported in only 4 of 33 cases of acute inhalation exposure to methylene chloride that were reported to occupational health authorities in the United Kingdom between 1961 and 1980 (Bakinson and Jones 1985); no exposure levels were provided in this study. No pulmonary function abnormalities were found in humans exposed to methylene chloride vapors (50–500 ppm) for 6 weeks (NIOSH 1974). Irritative symptoms of the respiratory tract were more prevalent among 12 Swedish male graffiti removers, employed to clean underground stations by using methylene chloride-based solvent, than those of the general population (Anundi et al. 1993). The 8-hour time-weighted average (TWA) to which these workers were exposed ranged from 18–1,200 mg/m<sup>3</sup>.

Two clinical case studies (Snyder et al. 1992a, 1992b) were reported in which two men who had been working in confined spaces with a nationally advertised brand of paint remover (consisting of >80% w/w methylene chloride) presented to the hospital emergency department complaining of dyspnea, cough, and discomfort in the midchest. In chest x-rays, each of the patients showed alveolar and interstitial infiltrates. One patient was treated with oxygen and albuterol and his symptoms improved over 48 hours; a repeat chest x-ray showed complete clearing of the infiltrates. During the next year, the patient continued to have episodic cough with wheeze and breathlessness which improved with albuterol therapy. The patient had no prior history of asthma or cough. A methacholine challenge test verified that he had hyperactive airways. The second patient was treated with oxygen and his symptoms improved during the next 48 to 72 hours; a repeat chest x-ray taken 3 days later revealed marked, but not complete, resolution of previously-noted lung infiltrates. Ten days later he was asymptomatic and his chest x-ray was normal.

## 2. HEALTH EFFECTS

Pulmonary effects were observed in animals that died following exposure to high concentrations of methylene chloride (Heppel et al. 1944). Extreme pneumonia was found in 3/14 guinea pigs exposed to 5,000 ppm for up to 6 months, and pulmonary congestion and edema with focal necrosis was found in 3/5 rabbits and 2/16 rats exposed to 10,000 ppm for up to 8 weeks (Heppel et al. 1944). A high incidence of foreign body pneumonia, involving focal accumulation of mononuclear and multinucleate inflammatory cells, was observed in 10/20 rats exposed to methylene chloride at 8,400 ppm for 13 weeks (NTP 1986). The significance of this finding is uncertain since the effect was observed only at the highest concentration tested. Male B6C3F<sub>1</sub> mice exposed to 4,000 ppm methylene chloride for 6 hours/day, 5 days/week for 13 weeks showed acute Clara cell damage in the lung after a 1-day exposure to methylene chloride, which appeared to resolve after 5 consecutive daily exposures (Foster et al. 1992). The appearance and disappearance of the lesion in Clara cells correlated well with the activity of cytochrome P-450 monooxygenase in Clara cells, as assessed immunocytochemically in the whole lung, and biochemically in freshly isolated Clara cells. Nasal cavity squamous metaplasia was observed in rats exposed intermittently to 1,000 ppm methylene chloride in the NTP (1986) bioassay.

**Cardiovascular Effects.** Studies in humans exposed to methylene chloride vapors between 50 and 500 ppm have not reported significant electrocardiographic abnormalities (Cherry et al. 1981; Ott et al. 1983c; NIOSH 1974). In cohort studies of methylene chloride workers, no increased ischemic heart disease mortality was observed with chronic time-weighted average exposures from 26 to 1,700 ppm (Hearne et al. 1990; Lanes et al. 1993; Ott et al. 1983b). There were no differences in cardiac effects, as measured by a health history questionnaire relating to heart problems (e.g., chest discomfort with exercise; racing, skipping, or irregular heartbeat), between 150 workers occupationally exposed for more than 10 years to relatively high levels of methylene chloride (8-hour TWA of 475 ppm), and a similar, nonexposed group of employees at a polyester staple plant (Soden 1993). The exposed cohort were also exposed to mean 8-hour TWA concentrations of 900 and 100 ppm of acetone and methanol, respectively.

Data in animals are limited to one study evaluating cardiac arrhythmia in the mouse (Aviado and Belej 1974). Atrioventricular block was observed following acute exposure to methylene chloride at concentrations greater than 200,000 ppm. Exposure to high concentrations of this sort is not likely to occur in the environment under normal conditions.

## 2. HEALTH EFFECTS

**Gastrointestinal Effects.** Nausea and vomiting were reported in 13 out of 33 cases of acute inhalation exposure to methylene chloride that were registered with occupational health authorities in the United Kingdom between 1961 and 1980 (Bakinson and Jones 1985); concentration levels were not provided in this study. Dilatation of the stomach was reported in mice after inhalation exposure to 4,000 ppm methylene chloride for 2 years (NTP 1986).

**Hematological Effects.** In humans, average blood COHb levels measure less than 1% in an atmosphere free of carbon monoxide, and less than 4% in a normal atmosphere. Blood COHb concentrations were about 30% higher than normal in two cases of lethal poisoning following acute inhalation of extremely high concentrations of methylene chloride in air (estimated ~168,000 ppm) in workers who were burying barrels containing mixed solvents and solid chemical waste in a well about 2 meters below ground level (Manno et al. 1992). Employees monitored at the end of 1 work day following exposure to methylene chloride at 7–90 ppm (8-hour TWA) had average COHb concentrations between 1.7 and 4.0% for nonsmokers, and between 4.95 and 6.35% for smokers (Soden et al. 1996). Additional daily cumulative exposure to methylene chloride did not produce increased levels of COHb. In volunteers who were exposed to methylene chloride at 200 ppm for 4 hours, blood COHb levels rose to approximately 5% (Putz et al. 1979); this was equivalent to the levels seen in volunteers after inhaling 70 ppm of carbon monoxide for 4 hours. In nonsmoking volunteers exposed to 50, 100, 150, or 200 ppm of methylene chloride for 7.5 hours, blood COHb levels rose to 1.9, 3.4, 5.3, and 6.8%, respectively, and blood COHb levels declined immediately following exposure (DiVincenzo and Kaplan 1981).

Other studies in humans reported increases in the red cell count, hemoglobin, and hematocrit in women occupationally exposed to concentrations up to 475 ppm during an 8-hour workday, but no effects were found in men. These effects were judged by the authors to be suggestive of compensatory hematopoiesis (Ott et al. 1983d). It may be anticipated that stress polycythemia will occur in the majority of individuals, especially cigarette smokers, who are chronically exposed to methylene chloride vapor concentrations in the 500 ppm range.

In animals, no significant hematologic or clinical chemistry alterations were reported in dogs and monkeys exposed continuously to up to 100 ppm methylene chloride for 100 days (Haun et al. 1972). In the dogs, COHb increased from 0.5 to about 2% during exposure to 100 ppm methylene chloride, but no significant increase was seen at 25 ppm. In the monkeys, COHb levels were approximately 0.5, 1.7, and 4.5% in controls, 25 ppm, and 100 ppm exposed groups, respectively. No treatment-related effects on

## 2. HEALTH EFFECTS

common hematologic parameters (cell counts, hemoglobin concentration differentials, white cell counts, etc.) were observed among rats chronically exposed to methylene chloride at concentrations up to 3,500 ppm (Burek et al. 1984; Nitschke et al. 1988a).

**Musculoskeletal Effects.** No musculoskeletal effects have been reported in either animals or humans after inhalation exposure to methylene chloride.

**Hepatic Effects.** There is very little published information on the hepatic effects of methylene chloride in humans. One autoworker, who was exposed to methylene chloride by inhalation and dermally for 1.5 years, was reported to have an enlarged liver in addition to adverse neurological and reproductive effects (Kelly 1988). NIOSH found workplace levels of methylene chloride to average 68 ppm (range of 3.3–154.4 ppm), which may be an underestimate given evaporation of the volatile liquid from the applicator pads and cotton gloves that were used; the worker was also exposed to low levels of styrene (7.2 ppm, range of 1.5–10.4 ppm). Exposure to methylene chloride was verified by a blood COHb level of 6.4% in a sample taken more than 24 hours after work. The relative contributions of the inhalation and dermal exposures to the hepatic effect was not determined. There were no alterations in serum enzyme activity (alkaline phosphatase, alanine aminotransferase [ALT], or lactic dehydrogenase) or in serum bilirubin, calcium, and phosphorus in humans exposed to methylene chloride vapors (50–500 ppm) for 6 weeks (NIOSH 1974). In a clinical epidemiologic assessment of methylene chloride workers, an exposure-related increase (not clinically significant) in serum bilirubin was observed in workers exposed to methylene chloride (up to 475 ppm) and methanol, but there were no concentration-related changes in serum enzyme levels that could indicate liver injury (Ott et al. 1983a). In Swedish graffiti removers employed to clean underground stations using methylene chloride solvent, no exposure-related deviations in serum concentrations of creatinine, aspartate transaminase (AST), ALT, or gamma-glutamyl transpeptidase were observed (Anundi et al. 1993). Based on these data, the liver appears to be a less sensitive target organ in humans than it is in rodents (see below).

In animals, the effects of methylene chloride have been studied more extensively. For the most part, exposure to methylene chloride has resulted in fatty changes in the liver and elevated plasma enzymes. These effects were reversible when exposure ceased. No histopathological changes were observed in guinea pigs following acute exposure to 5,200 ppm; however, there was a 2.5-fold increase in hepatic triglycerides (Morris et al. 1979). When male guinea pigs were exposed to 5,000 ppm of methylene chloride for up to 6 months, 3/8 died and exhibited moderate centrilobular fatty degeneration of the liver

## 2. HEALTH EFFECTS

(Heppel et al. 1944); no deaths, but similar liver histopathology was observed after exposure to 10,000 ppm for 8 weeks (guinea pigs) or 1 week (dogs). Fatty changes in the liver were noted in monkeys, mice, and dogs continuously exposed to 5,000 ppm for 4 weeks (MacEwen et al. 1972). In addition, mice exposed to 1,000 ppm exhibited iron pigmentation, nuclear degeneration, and pyknotic cells (MacEwen et al. 1972). Hepatic microsomal enzymes were elevated at 500 ppm ( $p < 0.01$ ) following 10 days of exposure, but were not increased significantly over control levels in rats exposed to methylene chloride at 250 ppm for 28 days (Norpoth et al. 1974). Continuous exposure of mice and rats for 100 days to 25 or 100 ppm caused fatty changes in the liver (Haun et al. 1972; Kjellstrand et al. 1986; Weinstein and Diamond 1972). No effects were seen in mice continuously exposed at 25 ppm, but cytoplasmic vacuolization was reported in rats at this exposure level (Haun et al. 1972). No adverse liver effects were reported in dogs or monkeys exposed to up to 100 ppm methylene chloride in the Haun et al. (1972) study. Using results from the Haun et al. (1972) study, an intermediate inhalation MRL of 0.3 ppm was derived based on the LOAEL of 25 ppm for liver effects in rats. In 13-week studies, centrilobular hydropic degeneration was observed in female mice exposed to 4,200 ppm, and in both sexes at 8,400 ppm (NTP 1986). Repeated exposure of rats to 200–500 ppm or greater for 2 years resulted in increased incidences of hepatocellular vacuolization and multinucleate hepatocytes (Burek et al. 1984; Nitschke et al. 1988a; NTP 1986), but not at 50 ppm (Nitschke et al. 1988a). In the 2-year NTP (1986) study, other liver effects in rats included hemosiderosis, focal necrosis of hepatocytes, basophilic change (females only), hepatocytomegaly, bile duct fibrosis in males, and granulomatous inflammation in females. The NOAEL of 50 ppm identified in the Nitschke et al. (1988a) study was used as the basis for derivation of a chronic inhalation MRL of 0.3 ppm.

**Renal Effects.** Daily exposures to concentrations up to 500 ppm methylene chloride for 6 weeks did not alter blood urea nitrogen or urine urobilinogen levels in humans (NIOSH 1974). In Swedish graffiti removers employed to clean underground stations using methylene chloride solvent, no exposure-related deviations in urinary concentrations of microglobulins or N-acetyl-beta-glucosaminidase were observed (Anundi et al. 1993).

Renal tubular vacuolization was observed in dogs following continuous inhalation exposure to 1,000 ppm for 4 weeks and in rats following exposure at 5,000 ppm for 14 weeks (MacEwen et al. 1972). Nonspecific renal tubular degenerative and regenerative changes were observed after continuous exposure in rats at 25 and 100 ppm for 100 days (Haun et al. 1972). No significant gross or histopathologic alterations in the kidneys were reported in dogs or monkeys exposed continuously to up to 100 ppm

## 2. HEALTH EFFECTS

methylene chloride for 100 days (Haun et al. 1972). Inhalation exposure to 2,000 ppm for 2 years resulted in statistically significant increases in the incidences of kidney degeneration in female rats and of kidney/tubule casts in mice of both sexes (NTP 1986).

**Ocular Effects.** One human study reported mild eye irritation in males exposed to 500 ppm methylene chloride vapors after 1 hour (NIOSH 1974). This was most likely due to direct contact of methylene chloride vapor with the eyes. Irritative symptoms of the eyes were more prevalent among 12 Swedish male graffiti removers (employed to clean underground stations using a methylene chloride-based solvent) compared to the general population (Anundi et al. 1993). The 8-hour TWA to which these workers were exposed ranged from 5 to 340 ppm.

One study reported transient increases in the thickness of the cornea in rabbits that were acutely exposed to vapors of methylene chloride at 490 ppm (Ballantyne et al. 1976). It is likely that the effects observed were due to direct effect of vapors on the cornea (see Section 2.2.3.2).

### 2.2.1.3 Immunological and Lymphoreticular Effects

No studies were located regarding immunological effects in humans after inhalation exposure to methylene chloride.

Some animal studies have reported alterations in secondary lymphoid organs following exposure to methylene chloride. Splenic fibrosis was observed in a chronic rat study (Mennear et al. 1988) and splenic atrophy in an intermediate dog study (MacEwen et al. 1972) at exposure concentrations of 1,000 ppm or greater. The significance of these effects on the spleen as an indicator of chemical insult to the immune system is not clear since no information was obtained on alterations in germinal cell proliferation, the lymphocyte subpopulation, or other immunological parameters. For these reasons, these studies are not presented in Table 2-1. However, a recent study by the Halogenated Solvent Industry Alliance, Inc. (2000) in which male and female rats were exposed whole body to 5,187 ppm methylene chloride 6 hours/day, 5 days/week, for 28 days found no evidence of immunotoxicity as judged by gross and microscopical examination of lymphoid tissues, hematology, or IgM antibody response to sheep red blood cells (SRBC). This NOAEL is listed in Table 2-1 and plotted in Figure 2-1.

## 2. HEALTH EFFECTS

**2.2.1.4 Neurological Effects**

A number of human studies reveal that the nervous system is perhaps the most important target of acute methylene chloride toxicity. All 33 cases of acute inhalation exposure to methylene chloride that were reported to occupational health authorities in the United Kingdom between 1961 and 1980 involved depression of the central nervous system (Bakinson and Jones 1985). Unconsciousness occurred in 13 of these cases and other common effects included headache and dizziness; a few instances of confusion, intoxication, incoordination, and paresthesia were also reported. Acute inhalation exposure to methylene-chloride-based paint strippers in rooms with inadequate ventilation led to unconsciousness in four cases and to generalized seizures in one of these (Hall and Rumack 1990); 10/21 respondents to an occupational health questionnaire reported experiencing dizziness and headache while working in these conditions, but the symptoms abated when they moved to fresh air. In volunteers, a single 4-hour exposure to 200 ppm methylene chloride significantly decreased visual and psychomotor performance and auditory function (Putz et al. 1979). Auditory monitoring, eye-hand coordination, and high-difficulty peripheral brightness test performances were not degraded until the final hour of exposure, by which time, the level of carbon monoxide in exhaled breath had risen to 50 ppm and the level of COHb in blood had risen to 5%. A single 3- to 4-hour exposure to methylene chloride at 300 ppm caused decreased visual and auditory functions in volunteers, but the adverse effects were reversible once exposure ceased (Fodor and Winneke 1971; Winneke 1974). Winneke (1974) attributed these effects to methylene chloride rather than its metabolite COHb, since exposure to carbon monoxide at concentrations up to 100 ppm did not cause similar effects. At the lowest exposure level (300 ppm of methylene chloride), critical flicker fusion frequency (visual) and auditory vigilance tasks were impaired. These higher-order functions involved complex visual and central nervous system processes that are assumed to be influenced by the degree of “cortical alertness” mediated by subcortical structures, especially the reticular formation (Fodor and Winneke 1971). Similarly, psychomotor performance (reaction time, hand precision, steadiness) was impaired, but this occurred at higher exposure levels (800 ppm for 4 hours) (Winneke 1974). Since these parameters are sensitive indicators of overt central nervous system-related depression, drowsiness, or narcosis, the Winneke (1974) study was selected as an appropriate basis for deriving an MRL for acute inhalation effects of methylene chloride. Alterations in visual evoked response were observed in humans exposed to methylene chloride at 515–986 ppm for 1–2 hours (Stewart et al. 1972). In another study, there were no effects on spontaneous electroencephalogram, visual evoked response, or a battery of cognitive effects in humans exposed to concentrations of methylene chloride up to 500 ppm (NIOSH

## 2. HEALTH EFFECTS

1974). While some changes in tests related to mood have been reported in humans after acute combined exposure to methylene chloride (28–173 ppm) and methanol (Cherry et al. 1983), no evidence of neurological or behavioral impairment was observed at exposure levels of 75–100 ppm (Cherry et al. 1981). Dementia and gait impairment were reported in one case of a person exposed to methylene chloride (500–1,000 ppm) for 3 years (Barrowcliff and Knell 1979). Based on a LOAEL of 300 ppm for neurological effects (Winneke 1974), an acute inhalation MRL of 0.6 ppm was calculated as described in Table 2-1 and Section 2.5.

No acute central nervous system effects were observed among 12 Swedish male graffiti removers employed to clean underground stations using methylene-chloride-based solvent compared to the general population (Anundi et al. 1993). The 8-hour TWA to which these workers were exposed ranged from 5 to 340 ppm. Two cases of men using a paint remover (>80% methylene chloride by weight) in small confined spaces were studied by Snyder et al. (1992a, 1992b) in a hospital emergency room. One reported symptom was severe headache, which disappeared within 24 hours after cessation of exposure. The authors considered this symptom to be associated with methylene chloride neurotoxicity. No neurologic effects, as measured by responses to questions relating to neurotoxicity (e.g., recurring severe headaches, numbness/tingling in hands or feet, loss of memory, dizziness) were reported in a group of 150 employees in a fiber plant occupationally exposed to methylene chloride (mean 8-hour TWA=475 ppm) for more than 10 years, when compared to a similar, nonexposed cohort (Soden 1993). In a retrospective epidemiology study, there were no significant associations between potential solvent exposure and self-reported neurological symptoms (based on a standard battery of medical surveillance questions) among workers exposed to a variety of solvents, including methylene chloride, at a pharmaceutical company (Bukowski et al. 1992). However, Bukowski et al. (1992) concluded that questionnaires were not the most appropriate tool to investigate potential neurobehavioral changes caused by low-level exposure to solvents, and recommended the use of neurological test batteries. This caveat would also apply to the study of Soden (1993).

In a group of 34 autoworkers, which included a subgroup of 26 ‘bonders’, complaints of central nervous system dysfunction were common following occupational exposure to methylene chloride for up to 3 years (Kelly 1988); the precise number of individuals with neurological complaints was not provided, since the report focused on reproductive effects. The bonding job involved soaking pads from open buckets of methylene chloride using ungloved hands, so exposures were both by inhalation and by the dermal route; exposure to methylene chloride was confirmed by analysis of blood COHb levels. In workplace air samples, NIOSH measured 3.3–154.4 ppm (average=68 ppm) of methylene chloride, but



## 2. HEALTH EFFECTS

worker exposure is likely to have been sometimes higher from the evaporation of liquid spilled onto clothing; bonders were also exposed to styrene at 1.5–10.4 ppm (average=7.2 ppm), which is less than the threshold limit value (TLV) for that chemical (ACGIH 1999). The neurological complaints reported in this group included dizziness, lightheadedness, memory loss, personality changes, and depression. Affected individuals were recommended for further neurological testing, but no test results were reported.

The neurotoxicity of occupational exposure to methylene chloride was examined in a cohort study of retired airline mechanics who had been chronically exposed to methylene chloride at concentrations ranging from a mean 8-hour TWA of 105 to 336 ppm, with short-term high exposures ranging from 395 to 660 ppm (Lash et al. 1991). Five categories of variables were assessed: demographic and some potential confounders, health symptoms, and physiological, psychophysical, and psychological variables. There were three tests of physiological characteristics (each measuring olfactory, visual, and auditory parameters) and four tests of psychophysical variables (finger tapping, simple reaction time, choice reaction time, and complex choice reaction time). Six psychological variables were assessed, most by more than one test: short-term visual memory, retention measure of visual memory, short-term verbal memory, and retention measures of verbal memory, attention, and spatial ability. None of the measured variables were statistically different between the exposed and control groups. However, trends in the effect sizes appeared within clusters of some variables. In the group of psychological variables showing effects on memory and attention, the exposed group scored higher than the unexposed group on the verbal memory tasks but lower on the attention tasks. The major potential confounder was exposure misclassification. Lack of precision, sampling biases, and random measurement errors might also have affected the results. However, the authors concluded that overall no effects on the central nervous system were attributable to chronic, low-level exposures to methylene chloride, a finding they reported as being consistent with that of Cherry et al. (1981).

White et al. (1995) conducted a 2-year prospective study among workers in the screen printing industry to investigate the association between exposure to mixed solvents with known neurotoxic properties and neurobehavioral deficits. Thirty subjects participated in the study which involved medical, demographic, occupational, and neurological screening, completion of a questionnaire on medical history, life style, and occupational history, and neurobehavioral assessment using a standard battery of neuropsychological tests. Subjects were evaluated and tested twice. Air monitoring studies identified workplaces within the plant that were most exposed to the following chemicals: toluene, methyl ethyl ketone (MEK), mineral

## 2. HEALTH EFFECTS

spirits,  $\beta$ -ether, methylene chloride, diacetone alcohol, acetic acid, and lead. A measure of total hydrocarbon content (THC) was also assessed to account for unidentified solvents in the workplaces. Generally, exposure to each of the major airborne chemical constituents was below the TLV. Among the 12 subjects classified as having high-acute exposure, performance, adjusted for age and education, was poorer for tasks involving visual memory and manual dexterity. Whether or to what degree exposure to methylene chloride contributed to these neurological effects cannot be determined.

Acute studies in animals are consistent with findings in humans that methylene chloride affects the central nervous system. Narcotic effects of methylene chloride (incoordination, reduced activity, somnolence) were observed in monkeys, rabbits, rats, and guinea pigs exposed to 10,000 ppm for up to 4 hours (Heppel et al. 1944); reduced activity was measured in rats exposed to 5,000 ppm (Heppel and Neal 1944). Dogs exposed to 10,000 ppm for 4 hours, first became uncoordinated, then excited and hyperactive to the extent of bruising themselves, but rapidly recovered afterwards (Heppel et al. 1944). Somatosensory-evoked potentials were altered in rats after 1 hour of exposure to methylene chloride at concentration levels of 5,000 ppm or greater (Rebert et al. 1989). Decreased levels of succinate dehydrogenase were measured in the cerebellum of rats exposed to 500 ppm of methylene chloride for 2 weeks (Savolainen et al. 1981).

Changes in neurotransmitter amino acids and brain enzymes were observed in gerbils after continuous exposure to 210 ppm for 3 months (Briving et al. 1986; Karlsson et al. 1987; Rosengren et al. 1986). The DNA concentration decreased in the hippocampus and cerebellum in gerbils exposed to 210 ppm of methylene chloride, indicating decreased cell density in these brain regions, probably due to cell loss (Karlsson et al. 1987; Rosengren et al. 1986). Methylene chloride (4,500 ppm) did not affect wheel running activity and avoidance learning in rats born to dams exposed prior to and/or during gestation (Bornschein et al. 1980). No treatment-related alterations in sensory evoked potentials, reflexes, posture, or locomotion were observed in rats exposed at 2,000 ppm (Mattsson et al. 1990).

The highest NOAEL values and all reliable LOAEL values for neurological effects for each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

## 2. HEALTH EFFECTS

**2.2.1.5 Reproductive Effects**

Exposure to methylene chloride has been reported to result in adverse reproductive effects in humans and animals.

One group of case studies reported reproductive effects in 8 out of 34 men who complained of central nervous system dysfunction following occupational exposure to methylene chloride for 0.4–2.9 years (Kelly 1988). The eight men (ages 20–47) were part of a subgroup of 26 ‘bonders’ who applied methylene chloride to automobile parts during assembly; this involved soaking pads from open buckets of methylene chloride using ungloved hands, so exposures were both by inhalation and by the dermal route. In workplace air samples, NIOSH measured 3.3–154.4 ppm (average=68 ppm) of methylene chloride, but worker exposure is likely to have been sometimes higher from the evaporation of liquid spilled onto clothing; bonders were also exposed to styrene at 1.5–10.4 ppm (average=7.2 ppm), which is less than the TLV for that chemical (ACGIH 1999). Exposure to methylene chloride within the group was suggested by blood COHb levels of 1.2–11% for six nonsmokers and 7.3 and 17.3% for two smokers. All eight men had recent histories of infertility and complained of genital pain (testicular, epididymal, and/or prostatic); the testes were atrophied in two workers (ages 27 and 47). Infectious disease was eliminated as a cause of these reproductive effects. Four men who submitted to testing had reduced sperm counts and the proportion of sperm with abnormal morphology ranged from 38 to 50%. Uncertainty regarding this study involves the small number of subjects, the multiple exposure to other organic chemicals, the lack of blood or urine samples quantifying exposure to methylene chloride, and the lack of a control group. No other studies were located that reported similar male reproductive effects from exposure to methylene chloride. Contrary to Kelly’s (1988) findings, Wells et al. (1989) found no evidence of oligospermia in four workers who had been exposed to levels of methylene chloride that were twice as high as in the Kelly study, while involved in furniture stripping for at least 3 months. It is not certain whether the longer duration of exposure (minimum occupational exposure 1.4 years) or exposure to other chemicals in the Kelly (1988) study contributed to the different outcomes in the two studies.

In a retrospective study of pregnancy outcomes among Finnish pharmaceutical workers during the late 1970s, female workers at eight factories had a higher rate of spontaneous abortions compared to the general population (Taskinen et al. 1986). It is likely that the women were exposed to multiple solvents, including methylene chloride, prior to conception, as well as during pregnancy. In the case-control study

## 2. HEALTH EFFECTS

of 44 pharmaceutical workers who had spontaneous abortions, exposure to various solvents, including methylene chloride, was associated with a slightly higher abortion rate (Taskinen et al. 1986). Exposure to methylene chloride was associated with an odds ratio (OR) for spontaneous abortion of 2.3, but the increase was of borderline significance ( $p=0.06$ ; 95% confidence interval [CI]=1.0–5.7). In a logistic regression model, the odds ratio of spontaneous abortion increased with increasing frequency of exposure to methylene chloride, but the sample size was too small for statistical significance. In addition, smoking and alcohol consumption were not considered. The authors mentioned that improved industrial hygiene procedures that eliminated solvent vapors in the workplace may have contributed to an overall decline in rates of spontaneous abortion among pharmaceutical workers during the period in question.

No adverse effects on reproduction were observed in rats exposed to concentrations up to 1,500 ppm of methylene chloride for two generations (Nitschke et al. 1988b). In dominant lethal tests involving male mice exposed to #200 ppm methylene chloride for up to 6 weeks, no microscopic lesions were found in the testes (Raje et al. 1988). Uterine, ovarian, and testicular atrophy was observed in rats and mice exposed to vapors of methylene chloride (4,000 ppm) for 2 years (NTP 1986), but the authors considered this effect to be secondary to malignant hepatic and alveolar neoplasms, as described in Section 2.2.1.8 Cancer. Existing data suggest reproductive toxicity may occur following chronic exposure to relatively high concentrations of methylene chloride.

The highest NOAEL values and all reliable LOAEL values for each study for reproductive effects are recorded in Table 2-1 and plotted in Figure 2-1.

### **2.2.1.6 Developmental Effects**

The only information reported on potential developmental effects of methylene chloride in humans was part of retrospective study of pregnancy outcomes among Finnish pharmaceutical workers (Taskinen et al. 1986). In the case-control study of these workers, exposure to methylene chloride (probably both before and during pregnancy) was associated through an odds-ratio prediction with a higher risk of spontaneous abortion (Taskinen et al. 1986). (See Section 2.2.1.5 Reproductive Effects for additional details).

In a study examining the relationship between birth weights and environmental exposures to methylene chloride from Kodak manufacturing processes in Monroe County, New York, no significant adverse

## 2. HEALTH EFFECTS

effects on birth weight were found among 91,302 single births from 1976 through 1987 (Bell et al. 1991). The highest predicted environmental concentration of methylene chloride was 0.01 ppm, which is substantially lower than levels found in occupational settings, but significantly higher than the average ambient background level of 50 parts per trillion.

A study in rats demonstrated that methylene chloride crosses the placental barrier (see Section 2.3.2.1; Anders and Sunram 1982). No treatment-related visceral abnormalities were reported in fetuses of mice and rats exposed to 1,250 ppm of methylene chloride during gestation, but an increase in the incidence of minor skeletal variants (e.g., delayed ossification of sternebra or extra sternebrae) was observed in both species; rats also exhibit an increased incidence of dilated renal pelvis. A maternal effect of increased liver weight was observed (Schwetz et al. 1975). When rats were exposed to 4,500 ppm, maternal liver weights increased and fetal body weights decreased, but teratogenic effects were not observed and viability and growth were not affected (Bornschein et al. 1980; Hardin and Manson 1980). Wheel running activity and avoidance learning were not affected in rats born to dams exposed prior to and/or during gestation to methylene chloride at 4,500 ppm (Bornschein et al. 1980). Longer-term exposure (for two generations) to concentrations of 1,500 ppm of methylene chloride did not affect neonatal survival or neonatal growth in rats (Nitschke et al. 1988b). Although fetal body weights were decreased, the absence of other fetotoxic effects, major skeletal variants, or significant embryoletality suggests that developmental toxicity is not a major area of concern following exposure to methylene chloride.

The highest NOAEL values and all reliable LOAEL values for developmental effects for each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

### 2.2.1.7 Genotoxic Effects

No studies were located regarding genotoxic effects in humans after inhalation exposure to methylene chloride.

Results of *in vivo* assays following inhalation exposure in animals were mixed. Inhalation exposure of mice to methylene chloride for 10 days at concentrations of 4,000 ppm or higher resulted in significant increases in frequencies of sister chromatid exchanges in lung cells and peripheral blood lymphocytes, chromosomal aberrations in lung and bone marrow cells, and micronuclei in peripheral blood erythrocytes (Allen et al. 1990). However, no evidence of chromosomal abnormalities was seen in bone marrow cells

## 2. HEALTH EFFECTS

of rats exposed by inhalation to concentrations up to 3,500 ppm for 6 months (Burek et al. 1984). The difference in responses observed may be due, in part, to species differences or test methodology. The relevance of these two studies to clastogenic mechanisms in humans is not certain.

More recent *in vivo* genotoxicity studies have attempted to elucidate the mechanism(s) of genotoxic action of methylene chloride. Casanova et al. (1992) pre-exposed male B6C3F<sub>1</sub> mice and Syrian Golden hamsters for 2 days (6 hours/day) to 4,000 ppm of methylene chloride. On the third day, animals were exposed for 6 hours to a decaying concentration of [<sup>14</sup>C] methylene chloride and then examined for the presence of DNA-protein crosslinks. DNA-protein crosslinks were detected in mouse liver, but not in mouse lung, hamster liver, or hamster lung. Similar results were observed in a second experiment (Casanova et al. 1996). B6C3F<sub>1</sub> mice, exposed to methylene chloride for 6 hours/day for 3 days at concentrations ranging from approximately 500 to 4,000 ppm formed DNA-protein crosslinks in the liver. The formation of DNA-protein crosslinks was a nonlinear function of airborne concentrations of methylene chloride. In addition, mice exposed for 6 hours/day for 3 days to concentrations ranging from approximately 1,500 to 4,000 ppm showed an increased rate of DNA synthesis in the lung, indicating cell proliferation, but increased cell turnover was not detected in mouse lung at exposure concentrations of 150 or 500 ppm. Hamsters showed no evidence of cell proliferation in the lung at any concentration, nor did cell proliferation occur in the livers of either species.

Devereux et al. (1993) analyzed liver and lung tumors induced in female B6C3F<sub>1</sub> female mice by inhalation of 2,000 ppm of methylene chloride for 6 hours/day, 5 days/week exposure for up to 104 weeks for the presence of activated *ras* proto-oncogenes. In methylene chloride-induced liver tumors, mutations, mainly transversions or transitions in base 1 or base 2, were detected, and were similar to those observed for the H-*ras* gene in spontaneous liver tumors. Mutations were also identified in the lung. The K-*ras* activation profiles in the methylene chloride-induced tumors were not significantly different from those in spontaneously occurring tumors. No other transforming genes were found in the nude mouse tumorigenicity assay. The authors were unable to identify any transforming genes other than *ras* genes in either mouse liver or lung tumors. Based on liver tumor data, they suggested that methylene chloride may affect the liver by promoting cells with spontaneous lesions.

Hegi et al. (1994) generated allelotypes of 38 methylene chloride-induced lung carcinomas from female B6C3F<sub>1</sub> mice exposed 6 hours/day, 5 days/week for 2 years to 2,000 ppm. The allelotypes were examined for various genotoxic endpoints, and the results compared to genotoxicity findings in two other

## 2. HEALTH EFFECTS

reciprocal-cross mouse strains. Throughout the genome, allelic losses occurred infrequently, except for markers on chromosome 4, which were lost in approximately half of the carcinomas. In lung adenomas, chromosome 4 losses were associated with malignant conversion. Methylene chloride-induced liver tumors did not demonstrate chromosome 4 loss, which indicated that this finding was specific for lung carcinomas. Preferential loss of the maternal chromosome 4 was also observed in carcinomas in B6C3F<sub>1</sub> mice. On chromosome 6, an association between *K-ras* gene activation and allelic imbalances was also found in B6C3F<sub>1</sub> mouse lung tumors. When allelotypes of tumors in mice from two reciprocal cross strains, AC3F<sub>1</sub> and C3AF<sub>1</sub>, were examined and compared with the findings in B6C3F<sub>1</sub> mice, one allele of the putative chromosome 4 tumor suppressor gene was shown to be inactivated. Whereas the results in B6C3F<sub>1</sub> mice suggested that nondisjunction events were responsible for the chromosome 4 losses, tumors from both reciprocal-cross mouse strains appeared to show small interstitial deletions in a chromosomal region that is homologous with a region in human chromosomes which is often lost in a variety of tumors, including lung cancers. In human chromosomes, a candidate tumor suppressor gene, MTS1, is located in this region.

In another genotoxic analysis with the same cohort (Hegi et al. 1994), loss of heterozygosity at markers near the p53 gene on chromosome 11 and within the retinoblastoma tumor suppressor gene were examined in methylene chloride-induced liver and lung tumors and compared to spontaneous tumors in control mice. The authors concluded that inactivations of p53 and the retinoblastoma tumor suppressor gene were infrequent events in lung and liver tumorigenesis in mice exposed to methylene chloride.

Replicative DNA synthesis was examined by Kanno et al. (1993) to evaluate the potential role of treatment-induced lung cell proliferation on pulmonary carcinogenicity in female B6C3F<sub>1</sub> mice exposed to 2,000 or 8,000 ppm of methylene chloride for 6 hours/day, 5 days/week for 2 years. By the end of the study, there was a statistically significant increase in lung tumors in exposed animals when compared to controls. Cell proliferation was assessed in the lung after 1, 2, 3, or 4 weeks of inhalation exposure to 2,000 or 8,000 ppm, and after 13- and 26-week exposures to 2,000 ppm, as measured by changes in labeling indices (LI). The LI of both bronchiolar epithelium and terminal bronchioles were substantially decreased in mice exposed to 2,000 ppm of methylene chloride for 2–26 weeks. Similar findings, but not as severe, were observed in mice exposed to 8,000 ppm. The decreases in LI were not accompanied by cytotoxicity. The authors concluded that high-concentration exposure to methylene chloride for up to 26 weeks reduces cell proliferation in lung epithelial cells in female B6C3F<sub>1</sub> mice.

## 2. HEALTH EFFECTS

Maronpot et al. (1995b) assessed replicative DNA synthesis after 13, 26, 52, and 78 weeks of inhalation exposure of female B6C3F<sub>1</sub> mice to 2,000 ppm of methylene chloride for 6 hours/day, 5 days/week. A statistically significant decrease in the hepatocyte LI was only observed at 13 weeks. In lung epithelial cells, the results were similar to those observed by Kanno et al. (1993). No increases in replicative DNA synthesis were found in liver foci cells or lung parenchymal cells. *K-ras* gene activation in liver tumors and *H-ras* gene activation in lung tumors did not differ among methylene chloride-induced tumors and those observed in control animals. The authors concluded that these oncogenes were not involved in mouse tumorigenesis.

Other genotoxicity studies are discussed in Section 2.5.

**2.2.1.8 Cancer**

No excess risk of death from malignant neoplasms has been detected in workers exposed to methylene chloride at levels up to 475 ppm (Friedlander et al. 1978; Hearne et al. 1987, 1990; Lanes et al. 1993; Ott et al. 1983a). Lanes et al. (1990) reported excess mortality associated with cancer of the buccal cavity and pharynx (combined), and liver and biliary passages (combined) in workers occupationally exposed to methylene chloride (#1,700 ppm) in the cellulose fiber production industry for more than 20 years. Although the actual number of cases was small, the excess mortality for the combined liver/biliary cancer cases was statistically significant (standard mortality rate [SMR]=5.75; 95% CI=1.82–13.78); since three of the four deaths were biliary cancer, the SMR was 20 (95% CI=5.2–56.0) for that site alone (Lanes et al. 1990). In a follow-up study of the same cohort (Lanes et al. 1993), no new cancer cases were found, but there was still an excess mortality for the cohort (SMR=2.98; 95% CI=0.81–7.63). Lanes et al. (1993) concluded that the excess death in this cohort from liver/biliary cancer was “statistically unstable”, but warranted further monitoring.

Tomenson et al. (1997) studied mortality due to cancer in a group of workers occupationally exposed to methylene chloride vapors at a mean exposure concentration of 19 ppm (8-hour TWA); the average length of employment was 9 years. Compared to national and local rates, the occupationally-exposed group had lower rates for all cancers, including those of the liver, lung, pancreas, and biliary tract. The authors suggested that the significant reduction in mortality due to lung cancer was likely associated with restrictions on smoking in the workplace.



## 2. HEALTH EFFECTS

Gibbs et al. (1996) examined causes of mortality in an occupationally exposed group of workers similar to that studied by Lanes et al. (1990, 1993). Exposure was classified categorically, with the airborne concentrations in the high-exposure group ranging from 350 to 700 ppm, and in the low exposure group from 50 to 100 ppm. Because there were no jobs in this cohort that did not involve methylene chloride exposure, a “0” exposure category was created as an internal control. There was no increase in mortality from cancers of the lung, liver, pancreas, or biliary tract. An unexpected finding was a concentration-related increase in mortality from prostate cancer among men with 20 or more years of employment. Another unexpected finding was an increase in mortality from cervical cancer among women with 20 or more years of employment, although the increase in women was not concentration-related. These results are not consistent with those from other studies. It should be noted that the confidence intervals were very large and no potentially confounding variables were measured. The authors concluded that the results were difficult to interpret biologically and required further investigation.

In a case control study, Heineman et al. (1994) evaluated exposures of men in the petroleum refining and chemical manufacturing industries to chlorinated aliphatic hydrocarbons (CAHs), including methylene chloride, as potential risk factors for astrocytic brain tumors. Job-exposure matrices for six individual CAHs, including methylene chloride, and for total organic solvents, were developed by estimating the probability of exposure and the frequency and magnitude of exposure to CAH solvents by industry and by job classification, based on likely solvent usage over 6 decades (1920–1980). An increase in the incidence of mortality due to astrocytic brain cancer was observed for exposure to four CAHs (carbon tetrachloride, methylene chloride, tetrachloroethylene, and trichloroethylene); the strongest association was with methylene chloride. In occupations judged to be associated with methylene chloride exposure, risk of astrocytic brain cancer increased with increasing exposure, as measured by the job exposure matrix in conjunction with duration of employment. The authors stated that these trends could not be explained by exposures to the other solvents.

As first evidence of such an association, these results should be interpreted very cautiously. The principal limitation of the study was the lack of direct information on exposure to solvents; no quantitative measurements were made, nor were specific-use records available. Instead, qualitative estimates of exposure were made by industrial hygienists based on work histories provided by next-of-kin; there were no workplace records. A lack of quantitative information on the use of specific solvents in various occupations, the ability of solvents to be used interchangeably for many industrial applications, and the use of multiple, or mixtures of, solvents also contributed to a high potential for exposure

## 2. HEALTH EFFECTS

misclassification. The authors stated that “few individual risks were statistically significant and most confidence intervals were broad”; interpretation of the results was based on “patterns of trends” (Heineman et al. 1994). The authors added that “...the trends and consistency of the methylene chloride and brain cancer association suggest that chance seems unlikely to entirely explain the results.”

In another case-control study, Cocco et al. (1999) examined the occupational risk of central nervous system cancer among women, using a study design similar to that described for Heineman et al. (1994). From death certificates in a U.S. 24-state database for the period 1984–1992, the authors identified 12,980 cases of cancers of the brain and other parts of the central nervous system. For each case, four controls were selected among women who died from nonmalignant diseases (excluding neurological disorders), frequency-matched by state, race, and 5-year age-group. Job exposure matrices were developed for 11 occupational hazards, including methylene chloride. An estimate of intensity of exposure was developed for each occupation and industry listed in the U.S. Census code. A final intensity level score and a probability of exposure score were then developed for each occupation/industry combination appearing in death certificates of the study subjects. The ORs were calculated with logistic regression for each workplace exposure, adjusting for marital status, socioeconomic status, and age at death. The authors found that potential exposure to methylene chloride was associated with a modest, but statistically significant, 20–30% increase in risk of mortality from central nervous system cancer (OR=1.2; 95% CI=1.1–1.2). However, the authors characterized the association as “equivocal”, since risk did not show a clear increase by probability or intensity of exposure. Admitted weaknesses of the study include poor occupational information in the death certificates, possible diagnostic bias among lower socioeconomic status cases, and the absence of more detailed information to supplement that provided in the death certificates.

Cantor et al. (1995) conducted a case control study with 33,509 cases and 117,794 controls, matched for age, gender, and race, to investigate the association between occupational exposure to workplace chemicals and the incidence of breast cancer in women. Exposure was indirectly estimated using a job exposure matrix that ranked the probability and level of 31 workplace exposures, among them methylene chloride. All workplace chemical exposures were evaluated separately. After adjusting for socioeconomic status imputed from occupation, the OR in the highest exposure category of methylene chloride (probability and level of exposure combined) was slightly greater than 1.0, and statistically significant (OR=1.46, CI=1.2–1.7). The authors caution that this analysis is crude and should be considered a first-level “hypothesis-generating” evaluation rather than one which is “hypothesis testing.”

## 2. HEALTH EFFECTS

They also noted that the study had numerous methodologic limitations, including the use of death certificates as the primary source of individual information and the use of a job-exposure matrix rather than quantitative workplace measurements to estimate exposure. Additionally, no adjustments were made for other risk factors such as smoking, obesity, family history, duration of employment, or other confounding variables.

In rats exposed to low levels of methylene chloride (100 ppm) for 2 years, there was a nonsignificant increase in the total incidence of malignant tumors (Maltoni et al. 1988). In mice and rats, inhalation of very high levels of methylene chloride significantly increased the incidence of liver and lung cancer (Mennear et al. 1988; NTP 1986) and benign mammary gland tumors (fibroadenomas or adenomas) (Mennear et al. 1988; Nitschke et al. 1988a; NTP 1986).

In the NTP (1986) study, groups of 50 animals of each sex were exposed to methylene chloride by inhalation 6 hours/day, 5 days/week for 102 weeks. F344/N rats were exposed to 0, 1,000, 2,000, or 4,000 ppm of methylene chloride and B6C3F<sub>1</sub> mice were exposed to 0, 2,000, or 4,000 ppm. At or above 2,000 ppm, the incidence of liver tumors (mostly hepatocellular adenomas or carcinomas) in mice was significantly higher than in chamber and historical control groups (NTP 1986); at 4,000 ppm, the incidence of liver tumors was highly significant ( $p \leq 0.001$ ). The incidence of combined benign and malignant liver tumors was high (67–83%) in the treated animals. There was also a statistically significant increase in the incidence of lung tumors in mice ( $p < 0.001$ ) exposed at 2,000 ppm or above; these tumors were primarily alveolar/bronchiolar adenomas or carcinomas. The incidence of combined benign and malignant lung tumors was 54–85% in the treated animals. The NTP (1986) report concluded that there was “some evidence of carcinogenicity” in male rats and “clear evidence of carcinogenicity” in female rats, based on the increased incidence of benign mammary neoplasms following 2 years of inhalation exposure to methylene chloride. The report concluded that there was “clear evidence for carcinogenicity” for methylene chloride chronic inhalation exposure, based on the increased incidence of alveolar/bronchiolar neoplasms and of hepatocellular neoplasms.

In two related studies, Kari et al. (1993) and Maronpot et al. (1995b) examined the progressive development of lung and liver tumors in B6C3F<sub>1</sub> mice exposed via chamber inhalation to 2,000 ppm methylene chloride for 6 hours/day, 5 days/week, for 104 weeks. In addition, a series of stop exposure experiments were performed to evaluate the effects of differing exposure durations on tumor development. Kari et al. (1993) examined histology and histopathology of lung and liver tumors, whereas

## 2. HEALTH EFFECTS

Maronpot et al. (1995b) evaluated DNA synthesis and oncogene expression during tumor development. (mechanistic aspects of Maronpot et al. (1995b) are discussed in Sections 2.2.1.7 and 2.4.2). Chronic high-concentration exposure to methylene chloride resulted in: (1) an 8-fold increase in the incidence of animals having lung adenomas or carcinomas as compared to controls; (2) a 13-fold increase in the total number of lung tumors in each animal at risk; (3) a 2.5-fold increase in the incidence of mice having liver adenomas or carcinomas compared to controls; and (4) a 3-fold increase in the number of liver tumors in each animal at risk. The development of the first lung tumors in methylene chloride exposed mice occurred 1 year earlier than in control animals. In contrast, there was no difference in the latency to first liver tumor period between exposed and control animals. The incidences of tumors in lungs, but not liver, continued to increase after cessation of exposure. Maronpot et al. (1995b) found that 26 weeks of exposure was sufficient to significantly and irreversibly increase the incidence of lung tumors at 2 years, whereas the incidence of hepatic tumors increased with 78 weeks of exposure, but not with 25 or 52 weeks of exposure. Furthermore, vulnerability to methylene chloride may have been age-related, since no lung tumor increase was observed in mice that were kept under control conditions for 52 weeks prior to methylene chloride exposure for 52 weeks. Based on these results, Kari et al. (1993) and Maronpot et al. (1995b) concluded that methylene chloride is a more potent lung than liver carcinogen in female B6C3F<sub>1</sub> mice; the differing incidence of lung and liver tumors under various exposure regimes suggests that the mechanisms of tumorigenesis in these target organs may be different.

The EPA (1985b) reviewed the NTP data on the carcinogenic effects of methylene chloride and calculated a human potency estimate. The potency factor ( $q_1^*$ ) which represents a 95% upper confidence limit of the extra lifetime human risk, is  $1.4 \times 10^{-2} (\text{mg/kg/day})^{-1}$ . The unit risk estimate (the excess cancer risk associated with lifetime exposure to  $1 \mu\text{g}/\text{m}^3$ ) for inhalation exposures is  $4.1 \times 10^{-6}$ . The EPA (1987a, 1987b) lowered this risk estimate to  $4.7 \times 10^{-7} \mu\text{g}/\text{m}^3$  on the basis of pharmacokinetics data reported by Andersen et al. (1987). Based on this value, cancer risk levels of  $10^{-4}$ ,  $10^{-5}$ ,  $10^{-6}$ , and  $10^{-7}$  correspond to 70 years of continuous exposure to 0.06, 0.006, 0.0006, and 0.00006 ppm, respectively. The predicted cancer risks are considered conservative upper estimates. The actual risk of cancer is unlikely to be higher and may be substantially lower. These values are recorded in Figure 2-1. EPA is planning to re-evaluate potential human risks associated with inhalation exposure to methylene chloride based on new mechanistic data and more recent pharmacokinetic modeling using tissue dosimetry (see Sections 2.3.5 and 2.4).

## 2. HEALTH EFFECTS

**2.2.2 Oral Exposure****2.2.2.1 Death**

Hughes and Tracey (1993) reported a case in which a woman ingested 300 mL of Nitromors, a paint remover solvent containing 75–80% methylene chloride, and died 25 days later. Ingestion of this paint remover is known to cause severe corrosion of the gastrointestinal tract, and the autopsy revealed that death was due to the corrosive effects of the paint remover rather than to the metabolic consequences of methylene chloride ingestion.

Acute oral LD<sub>50</sub> values of 2,100 (Kimura et al. 1971) and 2,300 mg/kg (Marzotko and Pankow 1987) were reported for methylene chloride in rats. Ninety-five percent lethality was reported in rats dosed with 4,382 mg/kg of methylene chloride (Ugazio et al. 1973). The cause of death appeared to be respiratory failure as a result of depression of the central nervous system. Statistically significant increases in mortality occurred among male rats gavaged with 320 mg/kg/day of methylene chloride and among male and female mice receiving 64 mg/kg/day for more than 36 weeks (Maltoni et al. 1988).

LD<sub>50</sub> values and LOAEL values for lethality in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**2.2.2.2 Systemic Effects**

No studies were located regarding respiratory, cardiovascular, musculoskeletal, or dermal effects in humans or animals following oral exposure to methylene chloride. Gastrointestinal, hematological, hepatic, renal, endocrine, and metabolic effects after oral exposure to methylene chloride are discussed below.

The highest NOAEL values and all reliable LOAEL values for systemic effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Gastrointestinal Effects.** A fatal single oral dose of Nitromors, a paint solvent containing 75–80% methylene chloride resulted in severe corrosion of the gastrointestinal tract, perforation, peritonitis, septicemia, and death (Hughes and Tracey 1993). A man who ingested 1–2 pints of Nitromors in a

TABLE 2-2. Levels of Significant Exposure to Methylene Chloride - Oral

Key to figure <sup>a</sup>	Species (Strain)	Exposure/Duration/Frequency (Specific Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference
					Less Serious (mg/kg/day)	Serious (mg/kg/day)	
<b>ACUTE EXPOSURE</b>							
<b>Death</b>							
1	Rat	once (GW)				2100 (LD <sub>50</sub> )	Kimura et al. 1971
2	Rat (albino)	once (G)				2300 M (LD <sub>50</sub> )	Marzotko and Pankow 1987
3	Rat (Wistar)	once (GO)				4382 M (95% mortality)	Ugazio et al. 1973
<b>Systemic</b>							
4	Rat (albino)	once (G)	Hemato	798 M		1325 M (hemolysis)	Marzotko and Pankow 1987
			Renal	798 M	1325 M (inhibited diuresis)		
			Endocr	399 M	526 M (increased catecholamine secretion)		
5	Rat (Wistar)	once (GO)	Hepatic			1095 M (liver necrosis)	Ugazio et al. 1973
<b>Neurological</b>							
6	Human	4 hr (W)			16 <sup>b</sup>	(decreased critical flicker frequency and auditory vigilance function)	Reitz et al. 1997
<b>INTERMEDIATE EXPOSURE</b>							
<b>Systemic</b>							
7	Rat (Fischer- 344)	90 d (W)	Hepatic		166 M (hepatocellular vacuolization; increased serum ALT)	1200 M (centrilobular necrosis)	Kirschman et al. 1986
			Renal	607 F	1469 F (increased kidney weight)		

TABLE 2-2. Levels of Significant Exposure to Methylene Chloride - Oral (continued)

Key to figure	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference
					Less Serious (mg/kg/day)	Serious (mg/kg/day)	
8	Mouse (B6C3F1)	90 d (W)	Hepatic	226 M	587 M (centrilobular fatty changes)		Kirschman et al. 1986
<b>CHRONIC EXPOSURE</b>							
<b>Death</b>							
9	Rat (Sprague-Dawley)	64 wk 4-5 d/wk 1x/d (GO)				320 M (increased mortality after 36 weeks)	Maltoni et al. 1988
10	Mouse (Swiss-Webster)	64 wk 4-5 d/wk 1x/d (GO)				64 (increased mortality after 36 weeks)	Maltoni et al. 1988
<b>Systemic</b>							
11	Rat (Fischer- 344)	78-104 wk (W)	Hemato	6			Serota et al. 1986a.
			Hepatic	6 <sup>c</sup>	55	(increased foci of cellular alteration and fatty changes)	
			Ocular	249			
			Bd Wt	55	131	(unquantified reduction in body weight gain)	
			Other	55	131	(decreased food and water consumption)	
12	Mouse (B6C3F1)	104 wk (W)	Hemato	236			Serota et al. 1986b
			Hepatic	175	236	(histochemical evidence of increased liver fat)	
			Bd Wt	236			

TABLE 2-2. Levels of Significant Exposure to Methylene Chloride - Oral

Key to figure <sup>a</sup>	Species (Strain)	Exposure/ Duration/ Frequency (Specific Route)	System	LOAEL			Reference
				NOAEL (mg/kg)	Less Serious (mg/kg)	Serious (mg/kg)	
13	Mouse (Swiss- Webster)	64 wk 4-5 d/wk 1x/d (GO)				320 M CEL (pulmonary tumors)	Maltoni et al. 1988

<sup>a</sup>The number corresponds to entries in Figure 2-2.

<sup>b</sup>LOAEL based on PBPK modeling of inhalation-to-oral route. Used to derive an acute oral minimal risk level (MRL) of 0.2 mg/kg/day; equivalent oral dose divided by an uncertainty factor of 100 (10 for the use of a LOAEL and 10 for human variability).

<sup>c</sup>Used to derive a chronic oral MRL of 0.06 mg/kg/day; dose divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability).

d = day(s); (GO) = gavage - oil; (BW) = gavage - water; Hemato = hematological; LD50 = lethal dose, 50% kill; LOAEL = lowest-observed-adverse-effect level;  
NOAEL = no-observed-adverse-effect level; (W) = water; wk = week(s); x = time(s)



Figure 2-2. Levels of Significant Exposure to Methylene Chloride - Oral  
Acute (≤14 days)




Figure 2-2. Levels of Significant Exposure to Methylene Chloride - Oral (Continued)  
Intermediate (15-364 days)





Figure 2-2. Levels of Significant Exposure to Methylene Chloride - Oral (continued)

Chronic (≥365 days)



## 2. HEALTH EFFECTS

suicide attempt, and started treatment with diuresis and hydrocortisone 1.5 hours later, had episodic gastrointestinal hemorrhage and duodenojejunal ulceration that developed into diverticula 6 months later (Roberts and Marshall 1976). No studies were located regarding gastrointestinal effects in animals after oral exposure to methylene chloride.

**Hematological Effects.** COHb levels were elevated to 9% in a woman following lethal ingestion of 300 mL of Nitromors, a paint remover solvent containing 75–80% methylene chloride (Hughes and Tracey 1993). The authors reported that this case study was the first to reveal that ingestion of methylene chloride results in the formation of COHb, as occurs with methylene chloride inhalation. A man who ingested 1–2 pints of Nitromors in a suicide attempt exhibited gross hemoglobinuria, a symptom of intravascular hemolysis (Roberts and Marshall 1976).

In an acute rat study, hemolysis developed within 2 days following a single gavage dose of 1,325 mg/kg of methylene chloride (Marzotko and Pankow 1987); no such effect was seen at 798 mg/kg. In rats that were given 420 mg/kg/day of methylene chloride in the drinking water for 3 months, mean hemoglobin concentrations were elevated in males, and erythrocyte counts were elevated, although mean corpuscular hemoglobin was reduced, in females (Kirschman et al. 1986); because no quantitative data were presented, this information is not presented in Table 2-2. In rats of both sexes that were given 55–249 mg/kg/day of methylene chloride in drinking water for 2 years, red blood cell counts and hematocrit and hemoglobin levels were increased over concurrent control levels (Serota et al. 1986a); however, Serota et al. (1986a) indicate that half of these increases were statistically significant without specifying which ones. A dose level of 6 mg/kg/day was a NOAEL. In mice exposed similarly to 60–236 mg/kg/day no significant hematological effects were observed after 104 weeks of treatment (Serota et al. 1986b).

**Hepatic Effects.** No studies were located regarding hepatic effects in humans after oral exposure to methylene chloride.

In rats given two doses (39–1,275 mg/kg each) of methylene chloride by gavage, there were no significant effects (within 24 hours) in the liver on levels of glutathione, cytochrome P-450, or serum ALT (Kitchin and Brown 1989). However, at the highest dose level, the activity of ornithine decarboxylase, an enzyme involved in cell growth, was significantly increased, and DNA damage was detected in the livers of rats. In another rat study, acute exposure to doses of 1,095 mg/kg/day of

## 2. HEALTH EFFECTS

methylene chloride by gavage resulted in liver necrosis (Ugazio et al. 1973). After ingestion of methylene chloride in drinking water at or above doses of 1,200 mg/kg/day for 3 months, hepatic effects in rats included centrilobular necrosis, granulomatous foci, accumulation of ceroid or lipofuscin, and dose-related hepatocytic vacuolation (Kirschman et al. 1986). Liver damage was indicated by increases in serum ALT in males (166 mg/kg/day) and females (1,469 mg/kg/day), and by an increase in serum AST in the latter group. In a parallel 3-month study in mice, there were dose-related hepatic centrilobular fatty changes over the dose range between 226 and 2,030 mg/kg/day (Kirschman et al. 1986). Chronic ingestion of methylene chloride in drinking water has been associated with histological alterations of the liver (cellular foci and areas of cellular alterations) of rats exposed to dose levels of 55 mg/kg/day or greater (Serota et al. 1986a) and fatty changes in mice exposed to levels of 236 mg/kg/day (Serota et al. 1986b); the NOAELs were 6 and 175 mg/kg/day in rats and mice, respectively. F344 rats and B6C3F<sub>1</sub> mice were exposed for 104 weeks to methylene chloride in deionized drinking water at target doses of 0, 5, 50, 125, or 250 mg/kg/day (for rats) and 0, 60, 125, 185, or 250 mg/kg/day (for mice) (Serota et al. 1986a, 1986b). Based on the calculated intake of 6 mg/kg/day for rats (Serota et al. 1986a), a chronic oral MRL of 0.06 mg/kg/day was calculated as described in the footnote of Table 2-2.

**Renal Effects.** Hemoglobinuria as a result of hemolysis was noted in the case of a suicide attempt by ingestion of the paint remover Nitromors (Roberts and Marshall 1976); the authors indicated that the renal damage that could have been a consequence of hemolysis was averted by treatment with diuresis and hydrocortisone. In an acute animal study, administration of a single oral dose of 1,325 mg/kg of methylene chloride inhibited diuresis in rats (Marzotko and Pankow 1987). In a 3-month study, the pH of the urine was lowered in all rats that ingested methylene chloride at levels  $\geq$  166 mg/kg/day, and increased kidney weights were observed in female rats receiving 1,469 mg/kg/day of methylene chloride, but not 607 mg/kg/day (Kirschman et al. 1986).

**Endocrine Effects.** No studies were located regarding endocrine effects in humans following oral exposure to methylene chloride. In the only animal study that mentions endocrine effects, administration of a single oral dose of methylene chloride (526 mg/kg) to male rats resulted in angiectasis (dilatation of capillaries) of the adrenal medulla and statistically significant increases in secretion of catecholamines (epinephrine and norepinephrine) (Marzotko and Pankow 1987). The authors suggested that these findings appeared to be stress-related.

## 2. HEALTH EFFECTS

**Ocular Effects.** No studies were located regarding ocular effects in humans after oral exposure to methylene chloride. No treatment-related ophthalmologic findings in rats administered up to 249 mg methylene chloride/kg/day in the drinking water for 104 weeks (Serota et al.1986a). No further relevant information was located.

**Metabolic Effects.** Metabolic acidosis was detected in a man who attempted suicide by drinking 1–2 pints of Nitromors paint remover that contained methylene chloride as the active ingredient (Roberts and Marshall 1976); he recovered following treatment with diuresis and hydrocortisone.

### 2.2.2.3 Immunological and Lymphoreticular Effects

No studies were located regarding immunological or lymphoreticular effects in humans or animals after oral exposure to methylene chloride.

### 2.2.2.4 Neurological Effects

One and a half hours after drinking 1–2 pints of paint remover (9,000–18,000 mg/kg) that contained methylene chloride as the active ingredient, a man was deeply unconscious and unresponsive to painful stimuli; his pupils were reactive, but tendon jerks were depressed and plantar response was absent (Roberts and Marshall 1976). He was treated by diuresis and hydrocortisone and regained consciousness by 14 hours after the initial event; at this time, no apparent cerebral damage was detected. The authors suggested that recovery would have been unlikely without medical intervention.

### 2.2.2.5 Reproductive Effects

No studies were located regarding reproductive effects in humans or animals following oral exposure to methylene chloride.

### 2.2.2.6 Developmental Effects

No studies were located regarding developmental effects in humans or animals following oral exposure to methylene chloride.

## 2. HEALTH EFFECTS

**2.2.2.7 Genotoxic Effects**

No studies were located regarding genotoxic effects in humans after oral exposure to methylene chloride.

In rats given two doses of 1,275 mg/kg of methylene chloride by gavage (17 hours apart), significant DNA breakage was detected in the liver 4 hours after the last dose (Kitchin and Brown 1989). Methylene chloride did not induce a statistically significant increase of micronuclei of polychromatic erythrocytes in mice when administered at doses up to 4,000 mg/kg (Sheldon et al. 1987). In mice given a single dose of 1,720 mg/kg of methylene chloride, DNA breaks were detected in nuclei from the liver and lung, but not from stomach, kidney, urinary bladder, brain, or bone marrow; the authors indicated that there was no evidence of cytotoxicity that might have caused the genetic damage (Sasaki et al. 1998). The variability in genotoxicity may reflect tissue-specific variation in the metabolism of methylene chloride.

Other genotoxicity studies are discussed in Section 2.5.

**2.2.2.8 Cancer**

No studies were located regarding carcinogenic effects in humans after oral exposure to methylene chloride.

Studies in animals provide suggestive evidence that ingestion of methylene chloride may increase the incidence of liver cancer. In an acute rat study, two doses of 1,275 mg/kg of methylene chloride (given 17 hours apart) caused an increase in the liver activity of ornithine decarboxylase, an enzyme that may contribute to the promotion of hepatic cancer (Kitchin and Brown 1989). Liver tumors were observed in female, but not in male rats that ingested methylene chloride (up to 250 mg/kg/day) for 2 years, but the cancer incidence rates were within historical control ranges (Serota et al. 1986a). Although liver cancer was observed in male mice, the incidence was not significantly elevated. Female mice did not have increased liver tumor incidence (Serota et al. 1986b). An increased incidence of mammary tumors was found in female rats that received methylene chloride by gavage at 500 mg/kg/day for 64 weeks but the results were not statistically significant (Maltoni et al. 1988). Under the same exposure conditions, an increased incidence of pulmonary tumors in male mice was statistically significant ( $p < 0.05$ ) when the increased mortality rate was taken into account (Maltoni et al. 1988). The EPA (1985b, 1987a) reviewed

## 2. HEALTH EFFECTS

the available data on the carcinogenic effects of methylene chloride and concluded that there was borderline evidence for carcinogenicity. The EPA estimated that the upper bound incremental unit carcinogenic risk for drinking water containing 1 µg/L methylene chloride for a lifetime was  $2.1 \times 10^{-7}$  (µg/L)<sup>-1</sup>. This risk estimate derivation was based on the mean of the carcinogenic risk estimates from the finding of liver tumors in the NTP (1986) inhalation study in female mice; the lung tumor data from the NTP study were not included in EPA's analysis. Since the extrapolation model is linear at low doses, additional lifetime cancer risk is directly proportional to the water concentration of methylene chloride. Thus, risk levels of  $10^{-4}$ ,  $10^{-5}$ ,  $10^{-6}$ , and  $10^{-7}$  are associated with 0.5, 0.05, 0.005, and 0.0005 mg/L, respectively (0.1, 0.01, 0.001, and 0.0001 mg/kg/day). Because these values are based on upper bound estimates, the true risk could be lower. These values are shown in Figure 2-2.

### 2.2.3 Dermal Exposure

#### 2.2.3.1 Death

No studies were located regarding death in humans or animals after dermal exposure to methylene chloride.

#### 2.2.3.2 Systemic Effects

**Respiratory Effects.** Shortness of breath was reported among some autoworkers who were exposed to methylene chloride both dermally and by inhalation for periods of up to 3 years (Kelly 1988). However, it is likely that inhalation exposure is largely responsible for this effect. No studies were located regarding respiratory effects in animals following dermal exposure to methylene chloride.

No studies were located regarding the following systemic effects in humans or animals after dermal exposure to methylene chloride:

**Cardiovascular Effects.**

**Gastrointestinal Effects.**

**Hematological Effects.**



## 2. HEALTH EFFECTS

**Musculoskeletal Effects.****Renal Effects.**

**Hepatic Effects.** One autoworker who was exposed to methylene chloride by dermal contact and by inhalation (3.3–154.4 ppm, average of 68 ppm) for 1.5 years was reported to have an enlarged liver in addition to adverse neurological and reproductive effects (Kelly 1988). This study is described in more detail under “Inhalation Exposure” in Section 2.1.3.2. The relative contributions of the two exposure routes to hepatotoxicity were not determined.

No studies were located regarding hepatic effects in animals after dermal exposure to methylene chloride.

**Dermal Effects.** There are few studies of dermal effects of methylene chloride in humans. In several cases, workers who were rendered unconscious while stripping furniture using a methylene chloride-based compound in an open tank became partially immersed in the liquid (Hall and Rumack 1990); first or second degree chemical burns developed on areas of the body having direct contact with the liquid. In another case, a worker who was cleaning the interior of a tank with methylene chloride became unconscious and fell into the solvent when the bucket overturned (Wells and Waldron 1984); during the 30 minutes before he was removed, second and third degree burns developed on the areas of contact. In a similar workplace accident, in which a man was found dead after 1 hour of exposure to methylene chloride, chemical burns with excoriation had developed on the areas of contact (Winek et al. 1981).

**Ocular Effects.** Data are limited regarding ocular effects in humans after dermal exposure to methylene chloride. Severe corneal burns developed in a worker who was found unconscious and slumped over with his face partially submerged in an open tank of paint stripper containing methylene chloride (Hall and Rumack 1990). The duration of exposure was not indicated.

In animals, methylene chloride (0.01–0.1 mL) caused eye irritation and inflammation, increased corneal thickness, and increased intraocular tension in rabbits following instillation of the liquid solvent into the conjunctival sac (Ballantyne et al. 1976). Effects were reversible within 3–9 days after treatment. In the same study, rabbits exposed to vapors of methylene chloride at concentrations of 490 ppm or greater for 10 minutes also showed effects on the eyes. There were small increases in corneal thickness and intraocular tension. Effects were reversible within 2 days. It is likely that effects observed were due to

## 2. HEALTH EFFECTS

direct effect of vapors on the cornea rather than to inhaled methylene chloride or its metabolites acting on the eyes.

### **2.2.3.3 Immunological and Lymphoreticular Effects**

No studies were located regarding immunological or lymphoreticular effects in humans or animals after dermal exposure to methylene chloride.

### **2.2.3.4 Neurological Effects**

Neurological effects (loss of memory, loss of concentration, sensory deficit) were reported in a group of 34 male autoworkers, who were exposed to methylene chloride by dermal contact and by inhalation (3.3–154.4 ppm, average of 68 ppm) for up to 3 years (Kelly 1988). This study is described in more detail under “Inhalation Exposure” in Section 2.1.3.4. The relative contributions of the two exposure routes to neurotoxicity were not determined.

No studies were located regarding neurological effects in animals after dermal exposure to methylene chloride.

### **2.2.3.5 Reproductive Effects**

One group of case studies reported reproductive effects in 8 out of 34 men who complained of central nervous system dysfunction following occupational exposure to methylene chloride for 0.4–2.9 years (Kelly 1988). The eight men were part of a subgroup of 26 ‘bonders’ who applied methylene chloride to automobile parts during assembly; this involved soaking pads from open buckets of methylene chloride using ungloved hands, so exposures were both by dermal contact and by inhalation (3.3–154.4 ppm, average of 68 ppm). This study is described in more detail under “Inhalation Exposure” in Section 2.1.3.5. Infectious disease was eliminated as a cause of the adverse reproductive effects (genital pain, testicular atrophy, and oligospermia). The relative contributions of the two exposure routes to reproductive toxicity were not determined. Uncertainty regarding this study involves the small number of subjects, the multiple exposure to other organic chemicals, the lack of blood or urine samples quantifying exposure to methylene chloride, and the lack of a control group. No other studies were located that reported similar male reproductive effects from exposure to methylene chloride. Contrary to Kelly’s

## 2. HEALTH EFFECTS

(1988) findings, Wells et al. (1989) found no evidence of oligospermia among four workers who had been exposed to levels of methylene chloride that were twice as high as in the Kelly study while involved in furniture stripping for at least 3 months; dermal exposure may have occurred, but this was not specified in the report. It is not certain whether the longer duration of exposure (minimum occupational exposure 1.4 years) or exposure to other chemicals in the Kelly (1988) study contributed to the different outcomes in the two studies.

### 2.2.3.6 Developmental Effects

No studies were located regarding developmental effects in humans or animals after dermal exposure to methylene chloride.

### 2.2.3.7 Genotoxic Effects

No studies were located regarding genotoxic effects in humans or animals after dermal exposure to methylene chloride.

Genotoxicity studies are discussed in Section 2.5.

### 2.2.3.8 Cancer

No studies were located regarding cancer effects in humans or animals after dermal exposure to methylene chloride.

## 2.3 TOXICOKINETICS

Inhalation is the main route of exposure to methylene chloride for humans. Within the first few minutes of exposure, approximately 70–75% of inhaled vapor is absorbed. However, as the concentration of methylene chloride in the blood increases, the net uptake is greatly reduced until at steady-state, it is equal to metabolic clearance, which has a maximum (determined by the fraction of blood flowing to the liver) of 25% (EPA 1994). Under conditions of continuous exposure to air concentrations of up to approximately 300 ppm, blood steady state concentrations of methylene chloride are reached in about 4 hours. Pulmonary absorption is influenced by exercise and body fat. In animals, pulmonary absorption

## 2. HEALTH EFFECTS

is proportional to magnitude and duration of exposure over a concentration range of 100–8,000 ppm. An increase of the steady state blood/air concentration ratio at high exposure levels reflects saturation of metabolic pathways rather than an increased absorption coefficient. There is only qualitative evidence of oral absorption in humans. In animals, methylene chloride is easily absorbed from the gastrointestinal tract, particularly from aqueous media. Seventy-five to 98% of an administered dose may be absorbed in 10–20 minutes. There are no quantitative data on dermal absorption of methylene chloride, although it is known to occur.

Distribution data in humans are lacking, but methylene chloride has been found in human breast milk and blood. Methylene chloride is widely distributed in animal tissues after inhalation exposure. The highest concentrations are found in adipose tissue and liver. Methylene chloride has been found in blood from rats' fetuses. After acute exposure, methylene chloride disappears rapidly from fat. Distribution of methylene chloride does not seem to be route-dependent and it does not bioaccumulate in tissues.

There are two main competing metabolic pathways for methylene chloride; one initially catalyzed by cytochrome P-450 enzymes (CYP2E1) and the other by a theta glutathione-S-transferase (GSSTI-1). The P-450 pathway (MFO) produces carbon monoxide and carbon dioxide via formyl chloride and the glutathione pathway (GST) produces carbon dioxide via a postulated glutathione conjugate (S-chloromethyl glutathione) and formaldehyde. Both pathways can give rise to toxic metabolites. The oxidative pathway is preferred at lower exposure concentrations and becomes saturated as exposure levels increase. Oxidative biotransformation of methylene chloride is similar in rats and humans. In rats, the MFO pathway is high-affinity low capacity, whereas the GST pathway has lower affinity, but higher capacity. The GST pathway is more active in mice than in rats and less active in hamsters and humans than in rats.

After inhalation exposure, humans eliminate methylene chloride mainly in expired air, but also in the urine. In rats, following a single exposure to radioactive methylene chloride, exhaled air had the most radioactivity, but radioactivity was also found in urine and feces. In exhaled air, the radiolabel was mostly as carbon monoxide and carbon dioxide. Physiologically based pharmacokinetic (PBPK) models have been developed to describe disposition of methylene chloride in humans and animals. These models were designed to distinguish contributions of the two metabolic pathways in lung and liver tissue, to look for correlations between tumor incidence and various measures of target tissue dose predicted by the models, and to extrapolate cancer risks from mice to humans.

## 2. HEALTH EFFECTS

**2.3.1 Absorption**

There is no information to determine whether absorption of methylene chloride in children is different than in adults.

**2.3.1.1 Inhalation Exposure**

The principal route of human exposure to methylene chloride is inhalation. During absorption through the lungs, the concentration of methylene chloride in alveolar air in equilibrium with pulmonary venous blood content approaches the concentration in inspired air until a steady state is achieved. After tissue and total body steady states are reached through the lungs and other routes, uptake is balanced by metabolism and elimination. Steady state blood methylene chloride concentrations appear to be reached after 2–4 hours of exposure (DiVincenzo and Kaplan 1981; McKenna et al. 1980).

Evaluation of pulmonary uptake in humans indicated that 70–75% of inhaled methylene chloride vapor was absorbed initially (DiVincenzo and Kaplan 1981). Initial absorption of methylene chloride was rapid as indicated by an uptake of methylene chloride into the blood of approximately 0.6 mg/L in the first hour of exposure to levels of 100–200 ppm. At a concentration of 50 ppm, the increase in blood methylene chloride concentration was 0.2 mg/L for the first hour (DiVincenzo and Kaplan 1981). There was a direct correlation between the steady state blood methylene chloride values and the exposure concentration, with a proportionality constant of approximately 0.008 ppm in blood per ppm in air (DiVincenzo and Kaplan 1981). The blood concentrations reached steady state values during the 4th through 8th hour of continuous exposure to the vapor. Once exposure ceased, methylene chloride was rapidly cleared from the blood. Six hours after the end of exposure, only traces of methylene chloride were present in the blood in the highest-concentration group, and pre-exposure baseline blood levels were detected in the other exposure groups.

Similar to other lipophilic organic vapors, methylene chloride absorption appears to be influenced by factors other than the vapor concentration. Increased physical activity increases the amount of methylene chloride absorbed by the body due to an increase in ventilation rate and cardiac output (Astrand et al. 1975; DiVincenzo et al. 1972). Uptake also increases with the percent body fat since methylene chloride dissolves in fat to a greater extent than it dissolves in aqueous media (Engstrom and Bjurstrom 1977). Therefore, obese subjects will absorb and retain more methylene chloride than lean subjects exposed to

## 2. HEALTH EFFECTS

the same vapor concentration. This does not mean obese subjects are more sensitive to toxicity. Under controlled conditions, there was a 30% greater absorption and retention of methylene chloride by obese subjects exposed to 75 ppm for 1 hour as compared to lean subjects (Engstrom and Bjurstrom 1977).

Studies of the relationship of inhalation exposures of animals to their blood methylene chloride concentrations indicate that absorption is proportional to the magnitude and duration of the exposure over a methylene chloride concentration range of 100–8,000 ppm. This conclusion is based on the monitoring of blood methylene chloride concentrations following inhalation exposure in dogs and rats (DiVincenzo et al. 1972; MacEwen et al. 1972; McKenna et al. 1982). As was the case with humans, blood methylene chloride levels reached a steady state value as the duration of exposure increased (McKenna et al. 1982).

Studies of blood methylene chloride values during 6-hour exposures of rats to between 50 and 1,500 ppm of methylene chloride suggest that the steady state blood/air concentration ratio increases as the exposure concentration increases. The ratio of the steady state methylene chloride concentration in the blood to the exposure concentration increased from 0.001 to 0.005 and 0.007 as the exposure increased from 50 to 500 to 1,500 ppm, respectively (McKenna et al. 1982). It is postulated that the increased ratio at steady state results from saturation of metabolic pathways as exposure increases rather than from an increased absorption coefficient.

### 2.3.1.2 Oral Exposure

No quantitative studies were located regarding absorption in humans after oral exposure to methylene chloride. There is qualitative evidence that the compound is absorbed when ingested. A male became deeply unconscious within 1.5 hours after ingestion of 1–2 pints of a paint remover (9,000–18,000 mg/kg) (Roberts and Marshall 1976).

In animals, the limited available data suggest that methylene chloride is easily absorbed from the gastrointestinal tract, particularly if exposed via aqueous media. Ten minutes after treatment, 24% of the administered dose (50 mg/kg in aqueous solution) was recovered from the upper gastrointestinal tract of mice when the stomach and small intestinal tissues and contents were analyzed (Angelo et al. 1986a). Only 2.2% of the methylene chloride was in the stomach and small intestines 20 minutes after compound administration and less than 1% remained after 40 minutes. At 10 minutes the large intestines and

## 2. HEALTH EFFECTS

caecum contained 0.08% of the dose; these values were less than those detected after 20 minutes. Thus, 75% of the dose was absorbed within 10 minutes and about 98% of the dose was absorbed within 20 minutes.

On the other hand, after treatment of mice with 10–1,000 mg methylene chloride in corn oil, approximately 55% of the administered dose was detected in the stomach and small intestines at 20 minutes and remained there at 2 hours (Angelo et al. 1986a). The large intestines and caecum tissues and contents contained about 5–8% of the dose at 10 minutes. This value declined to 1–2% at the end of 2 hours. This suggests that the absorption processes are slowed when methylene chloride is presented in a hydrophobic vehicle. During the 2 hours of observation, approximately 40–45% of the 50 mg dose was absorbed from the oil vehicle as compared to essentially all of a comparable dose in water. These data are consistent with the slower emptying of chyme from the stomach when lipids are present, the partitioning of methylene chloride between the corn oil and the aqueous digestive fluids, the necessity that fats be emulsified by bile prior to digestion and absorption, and delayed mobilization of lipophilic substances from the gastrointestinal tract.

Staats et al. (1991) developed a two-compartment model of oral absorption, dependent in part on the vehicle (aqueous or lipid) and on the lipophilic characteristics of the ingested compound. Absorption along the gastrointestinal tract was predicted to increase in the first compartment (likely the stomach), when fat-soluble toxicants are administered in water. When administered in corn oil, the toxicants are likely to adhere to the lipid vehicle in the first compartment and absorption in the stomach is likely to decrease. The model also predicts that passage of the compound from the first compartment (stomach) to the second compartment (small intestine) is enhanced when the vehicle is oil rather than water. However, existing experimental data suggest that absorption from the second compartment (small intestine) is generally slow for all lipophilic compounds, irrespective of vehicle. Using previously-determined absorption and transfer constant values, the model developed by Staats et al. (1991) was able to predict reasonably experimental data for trichloroethylene. The ability of the model to predict the gastrointestinal absorption of methylene chloride was not determined. However, in general, the model is consistent with methylene chloride experimental data (Angelo et al. 1986a).

## 2. HEALTH EFFECTS

**2.3.1.3 Dermal Exposure**

No studies were located regarding absorption in humans following dermal exposure to methylene chloride.

Dermal permeability constants for rats were obtained for 3 concentrations of methylene chloride (30,000, 60,000, and 100,000 ppm) in air for use in developing a pharmacokinetics model for dermal absorption of vapors (McDougal et al. 1986). The mean permeability constant was 0.28 cm/hr. The total amount absorbed was determined to be 34.4, 57.5, and 99.4 mg, respectively, for the three concentrations tested.

**2.3.2 Distribution**

There is no information to ascertain whether distribution of methylene chloride would be different in children than in adults.

**2.3.2.1 Inhalation Exposure**

When methylene chloride is absorbed by the lungs it is expected that it will dissolve in the lipoprotein components of the blood and enter the systemic circulation after passage through the heart. It is distributed from the systemic circulation to the body organs. However, no quantitative data were located which showed the distribution of methylene chloride following human inhalation exposure. Some data are available which relate to the uptake of methylene chloride by human adipose tissues. These data indicate that the methylene chloride concentrations in the adipose deposits of lean subjects are greater than those in obese subjects. However, the total methylene chloride adipose tissue load is greater for the obese subjects due to their greater adipose mass (Engstrom and Bjurstrom 1977).

Distribution studies in rats demonstrate that methylene chloride and/or its metabolites are present in the liver, kidney, lungs, brain, muscle, and adipose tissues after inhalation exposures (Carlsson and Hultengren 1975; McKenna et al. 1982). One hour after exposure, the highest concentration of radioactive material was found in the white adipose tissue, followed by the liver. The concentration in the kidney, adrenals, and brain were less than half that in the liver (Carlsson and Hultengren 1975). Radioactivity in the fat deposits declined rapidly during the first 2 hours after exposure (Carlsson and Hultengren 1975). Concentrations in the other tissues declined more slowly. On the other hand, after



## 2. HEALTH EFFECTS

5 days of exposure to 200 ppm, 6 hours per day, the concentration of methylene chloride in the perirenal fat was 6- to 7-fold greater than that in the blood and liver (Savolainen et al. 1977).

The animal data are accordingly consistent with the human adipose tissue data discussed above. When pregnant rats were exposed to 500 ppm of methylene chloride by inhalation for 1 hour, the chemical was found in fetal blood, but at a lower level than in maternal blood (Anders and Sunram 1982).

### 2.3.2.2 Oral Exposure

No studies were located regarding distribution in humans following oral exposure to methylene chloride.

In animals, radioactivity from labeled methylene chloride was detected in the liver, kidney, lung, brain, epididymal fat, muscle, and testes after exposure of rats to a single oral gavage dose of 1 or 50 mg/kg methylene chloride (McKenna and Zempel 1981). The tissue samples were taken 48 hours after dosing. At that time, the lowest concentration of radioactivity was found in the fat. The highest concentrations were in the liver and kidney; this was true for both doses. Radioactivity was found in the blood, liver, and carcass of mice. Similar results were observed in rats administered doses of 50–1,000 mg/kg methylene chloride for 14 days (Angelo et al. 1986a, 1986b). At each dose tested, and in each tissue, the label was rapidly cleared during the 240 minutes after each exposure. These data suggest that methylene chloride and/or its metabolites do not bioaccumulate in any tissues.

### 2.3.2.3 Dermal Exposure

In humans, direct dermal contact with pure methylene chloride causes an intense burning, mild erythema and paresthesia (Stewart and Dodd 1964). Absorption is relatively rapid (McDougal et al. 1986).

### 2.3.3 Metabolism

Methylene chloride metabolism is not known to be qualitatively different in children than adults. Information regarding the developmental expression of enzymes that metabolize methylene chloride is discussed in Section 2.7, Children's Susceptibility.

## 2. HEALTH EFFECTS

***Inhalation Exposure***


Available data suggest that there are two pathways by which methylene chloride is metabolized. One pathway utilizes the mixed function oxidase (MFO) enzymes and produces carbon monoxide (CO) (Figure 2-3, Pathway 1). The other pathway involves the glutathione transferase (GST) and produces carbon dioxide (CO<sub>2</sub>) (Figure 2-3, Pathway 2). It has been postulated that CO<sub>2</sub> can also be produced by the MFO pathway if the reactive intermediate in this pathway (postulated to be formyl chloride) reacts with a nucleophile prior to elimination of the chloride ion and formation of CO (Figure 2-3, Pathway 3) (Gargas et al. 1986).

The MFO pathway seems to be the preferred pathway for methylene chloride metabolism following inhalation exposures. Human subjects exposed by inhalation to 500 ppm or greater for 1 or 2 hours experienced elevated COHb concentrations indicating that methylene chloride was metabolized to CO by the MFO pathway (Stewart et al. 1972). The COHb concentrations rose to an average of 10.1% saturation 1 hour after the exposure of 3 subjects to 986 ppm of methylene chloride for 2 hours. The mean COHb concentration at 17 hours-post exposure remained elevated (3.9% saturation) above the pre-exposure baseline value (1–1.5% saturation). The exposure of 8 subjects to 515 ppm of methylene chloride for 1 hour increased the COHb level, which remained elevated above baseline for more than 21 hours.

In human subjects exposed by inhalation to 50–500 ppm of methylene chloride for up to 5 weeks, COHb concentrations could be predicted from methylene chloride exposure parameters (Peterson 1978). However, the exhaled breath concentrations of methylene chloride correlated better with exposure parameters than did COHb concentrations. No differences in methylene chloride metabolism between male and female subjects were detectable and there was no induction of metabolism to CO during 5-weeks exposure to concentrations ranging from 100 to 500 ppm of methylene chloride (Peterson 1978).

Metabolism of methylene chloride in animals has been shown to be similar to that in humans in an experiment by Fodor et al. (1973). Albino rats exposed to methylene chloride showed COHb formation, confirming the observation that methylene chloride is metabolized to CO following inhalation exposures. Among rats exposed to methylene chloride for 4 hours, concentrations of 500 or 2,500 ppm resulted in similar maximal blood levels of COHb (Wirkner et al. 1997). A concentration-response relationship between inhaled methylene chloride and the maximum changes in COHb values was observed when

Figure 2-3. Proposed Pathways for Methylene Chloride Metabolism



Source: Gargas et al. 1986

- 1 Mixed Function Oxidase Pathway
- 2 Glutathione Transferase Pathway
- 3 Nucleophile Pathway

## 2. HEALTH EFFECTS

4 rabbits were exposed to 1,270–11,520 ppm of methylene chloride for a 4-week period (Roth et al. 1975). The larger the exposure concentration, the longer it took for the changes in blood COHb values to reach their maximum values. The maximum COHb concentration was observed 1.5 hours after exposure to 1,270 ppm and 3.5 hours after exposure to 11,520 ppm.

In rats, inhalation of 50 ppm for 6 hours resulted in 26 and 27% of the body burden being recovered as expired CO and CO<sub>2</sub>, respectively, during the first 48 hours after the exposure period (McKenna et al. 1982). At 1,500 ppm, 14 and 10% of the body burden were recovered as CO and CO<sub>2</sub>, respectively. These values are consistent with the concept that the enzymes responsible for the metabolic conversion of methylene chloride to CO and CO<sub>2</sub> become saturated at high-exposure concentrations.

Anders and Sunram (1982) monitored the fetal and maternal blood concentrations of methylene chloride and carbon monoxide following inhalation exposure of pregnant rats to methylene chloride at 500 ppm for 1 hour. Whereas the level of methylene chloride in fetal blood was significantly lower than in maternal blood, the levels of carbon monoxide were about the same. The authors suggested that the maternal liver has higher biotransforming activity than the fetal liver and that the metabolically generated carbon monoxide equilibrates between the fetal and maternal circulations.

Thier et al. (1991) investigated whether the human metabolism of methylene chloride was similar to monohalogenated methanes; these compounds are metabolized via a glutathione-dependent pathway in human erythrocytes in a subgroup of the human population (called “conjugators”), whereas another subgroup does not exhibit erythrocyte glutathione-mediated metabolism (called “nonconjugators”). Blood samples were taken from 10 volunteers who had previously been determined to be either “conjugators” (subgroup B) or “nonconjugators” (subgroup A) of monohalogenated methanes. The samples were exposed to radiolabeled methylene chloride, incubated, centrifuged to separate blood plasma from cellular fraction, and the distribution of radioactivity between the different blood compartments measured. For individuals from subgroup B, radioactivity in blood plasma increased over time, reaching 30% in the low-molecular weight fraction and 5% in the high-molecular weight fraction after 9 hours. For individuals in subgroup A, almost no radioactivity was found in either blood plasma molecular fraction. In all samples from both groups, no radioactivity was found in either erythrocyte cytoplasm or membranes. Thus, although dihalogenated methylene chloride does not appear to undergo metabolism in erythrocytes, some metabolic transformation of methylene chloride appears to have occurred in the plasma of all individuals classified as “conjugators” (subgroup B), whereas this

## 2. HEALTH EFFECTS

metabolism did not occur in “nonconjugators” (subgroup A). The authors concluded that these data provide evidence for enzyme polymorphism in humans with respect to methylene chloride metabolism.

The contributions of the MFO and GST pathways to the metabolism of methylene chloride were studied in male rats by Gargas et al. (1986). Based on the data from these studies, the MFO pathway is a high affinity-low capacity pathway with a metabolic rate of 47  $\mu\text{mol/kg/hour}$ . The GST pathway, on the other hand, has a lower affinity than the MFO pathway but a higher capacity. As part of this study, rats were pretreated with an inhibitor of the MFO pathway or a glutathione (GSH) depleting agent. Concentrations of COHb were measured to assess the relative contribution of each pathway to the total metabolism of methylene chloride. Pretreatment with the MFO inhibitor essentially abolished CO production by the high-affinity saturable MFO pathway and limited metabolism to the GSH-transferase pathway. Concentrations of COHb following treatment with the GSH-depleting agent were increased 20–30% compared to untreated controls, indicating increased activity of the MFO pathway. Some  $\text{CO}_2$  was produced, indicating that despite some depletion of GSH, some methylene chloride was still metabolized by the GST pathway or by conversion of CO to  $\text{CO}_2$  via the MFO pathway.

There appear to be species differences in pathway preference. The GST pathway is more active in the mouse than the rat, based on studies of metabolic end products in both species during and immediately after 6 hours of exposure to 500, 1,000, 2,000, and 4,000 ppm of methylene chloride (Green et al. 1986c). The cytochrome MFO pathway leading to CO and COHb was shown to be saturated at the 500 ppm-exposure level. After saturation of this pathway had occurred, the blood levels of methylene chloride in the rat increased almost linearly with concentration, indicating little further metabolism in this species. In contrast, there was evidence for significant metabolism of methylene chloride in the mouse at high-concentration levels by the GST pathway, which leads to  $\text{CO}_2$ . Comparison of expired  $\text{CO}_2$  levels after 4,000-ppm exposure for 6 hours showed almost an order of magnitude more  $\text{CO}_2$  produced per kilogram of body weight in the mouse than in the rat. A marked difference was seen in the rate of clearance of methylene chloride from tissues, as measured by its elimination in expired air. Although cleared from blood rapidly in both species, the rate of clearance from rat tissues was markedly slower than in the mouse. This slow release in the rat sustained metabolism for up to 8 hours after the end of exposure and, consequently, had a marked effect on the overall body burden of metabolites. Methylene chloride was cleared from tissues in the mouse in less than 2 hours. Overall, saturation of the cytochrome P-450 MFO pathway occurred at similar levels in both species, but significantly more methylene chloride was

## 2. HEALTH EFFECTS

metabolized by the GST pathway in the mouse when assessed either from the blood levels of methylene chloride or by CO<sub>2</sub> formation at high-concentration levels.

The isoenzymes involved in the metabolism and biotransformation of methylene chloride have recently been identified for each major pathway. The MFO pathway involves cytochrome P-450 2E1 and the glutathione-mediated pathway involves a  $\Theta$  (theta) class glutathione S-transferase, GSTT1-1 (Blocki et al. 1994; Mainwaring et al. 1996b; Meyer et al. 1991). GSTT1-1 is expressed in a number of human organs in a tissue-specific manner which is different from the pattern of expression of other glutathione S-transferases, specifically those from the  $\alpha$ ,  $\mu$ , and  $\pi$  classes (Sheratt et al. 1997). A recent *in vitro* study by Sheratt et al. (1997) detected very low levels of GSTT1-1 in human lung cells, suggesting that the human lung is likely to have a low capacity for activating methylene chloride into reactive metabolites. Although Mainwaring et al. (1996b) found that the overall distribution of mRNA and protein for GSTT1-1 and GSTT2-2 in the liver and lungs of humans was lower than in mice and rats, they found localized high concentrations of GSTT2-2 enzyme in human bile-duct epithelial cells. However, since the enzyme did not localize to the nucleus in these cells, the risk of genotoxic effects from methylene chloride metabolism would appear to be small. Mainwaring et al. (1996b) also found locally high concentrations of GSTT1-1 mRNA in a small number of Clara cells and ciliated cells of the alveolar/bronchiolar junction in one human lung sample out of four. Mainwaring et al. (1996b) also found that rates of metabolism of methylene chloride were low in human tissue, and lower in the lung than in the liver. Thus, it is possible that, in some individuals, specific cell types may be vulnerable to genotoxic effects from reactive intermediates of methylene chloride metabolism, although the overall risk is likely to be low.

The oxidative cytochrome P-450 2E1 is presumed to metabolize methylene chloride to carbon dioxide via a reactive intermediate, formyl chloride; this metabolic pathway also produces carbon monoxide (Green 1997; Reitz 1990). The glutathione pathway metabolizes methylene chloride to carbon dioxide following the formation of both formaldehyde and a glutathione conjugate, putatively chloromethyl glutathione. No carbon monoxide is produced during glutathione-mediated metabolism (Green 1997; Reitz 1990). Neither the formyl chloride, nor the chloromethyl glutathione, nor any other glutathione conjugate of methylene chloride, has been isolated and characterized. However, according to Green (1997), the formation of these reactive intermediates is consistent with the end products formed, and with the enzymes and cofactors available.

## 2. HEALTH EFFECTS

An extremely detailed analysis of these two metabolic pathways was conducted in mice, rats, hamsters, and humans, both *in vivo* and *in vitro*. These studies provide evidence for the concentration-dependent behaviors of the two pathways and compare the metabolic rates by each pathway in the four species (Bogaards et al. 1993; Green 1991, 1997; Reitz et al. 1989). *In vivo*, the cytochrome P-450 MFO pathway was the major route of metabolism at low-concentration inhalation exposures to methylene chloride. In both rats and mice, saturation of the MFO pathway occurred at concentration levels of 500 ppm and above, with maximum COHb levels reported to be 12–15%. *In vitro* studies verified that this pathway was similar in all four species. In contrast, the glutathione S-transferase pathway was the major metabolic pathway at exposure concentrations used in the rodent cancer bioassays, and showed a linear concentration response. Furthermore, metabolic activity in mouse tissue was more than 10-fold greater than metabolic activity in rat tissue. The glutathione-mediated metabolic rates in hamster and human tissues were even lower than those observed in the rat (Green 1997). In human tissues, metabolic rates in the lung were about 10-fold lower than those in the liver (Green 1997). Although the  $\Theta$  (theta) class glutathione S-transferase (enzyme 5-5 in rat) has a high specific activity for metabolizing methylene chloride (Meyer et al. 1991; Sheratt et al. 1997), the  $\mu$ -class GSTs (enzymes 3-3, 3-4, and 4-4) are, as a group, 800-fold more abundant in rat liver (Blocki et al. 1994); in rat liver cytosol, the  $\Theta$ -class enzyme (5-5) and the  $\mu$ -class group of enzymes (3-3, 3-4, and 4-4) are each responsible for half of the metabolism of methylene chloride to formaldehyde (Blocki et al. 1994). In addition, another  $\Theta$ -class enzyme (12-12) is present in rat liver, but its lability during isolation has prevented analysis of its specific activity (Meyer et al. 1991). Although Schroder et al. (1996) demonstrated that the human erythrocyte GSTT1-1 is polymorphic, from N-terminal modification, and differs from liver and lung GSTT1-1, it is not yet known whether liver and lung human GSTT1-1 are polymorphic. Thus, the enzymatic basis for methylene chloride metabolism in different tissues in different species is not completely elucidated.

Nelson et al. (1995) examined the distribution of erythrocyte GSTT1-1 polymorphisms among five different ethnic groups: North American Caucasians, African-Americans, Mexican-Americans, Chinese, and Koreans. Polymerase chain reaction (PCR)-based genotyping of erythrocyte GSTT1-1 demonstrated that significant ethnic variations occur. The prevalence of the nonfunctional genotype (i.e., the one lacking the ability to metabolize methylene chloride) was highest among Chinese (64%), followed by Koreans (60%), African-Americans (22%), Caucasians (20%), and Mexican-Americans (10%). These data suggest that there are ethnic differences in metabolizing capacity; additionally, substantial variations in GSTT1-1 polymorphisms also occur within ethnic groups (Nelson et al. 1995)

## 2. HEALTH EFFECTS

Kim et al. (1996b) conducted a series of experiments using the vial equilibration technique on freshly isolated liver tissue of male Sprague Dawley rats in order to characterize the metabolism of methylene chloride. Several different hepatic microsomal preparations were used to compare and contrast the MFO- and GSH-mediated pathways. To produce glutathione depletion in one study, rats were injected intraperitoneally with 250 mg phorone/kg body weight in corn oil prior to sacrifice and preparation of liver homogenate for *in vitro* testing; controls received corn oil only. A glycerol buffer preparation significantly inhibited methylene chloride metabolism while a sucrose buffer containing EDTA and KCl did not. The use of substrates for P-450 2E1 (i.e., ethanol, pyrazole) completely inhibited methylene chloride metabolism, indicating that the methylene chloride was metabolized preferentially by the MFO pathway under the conditions of this study. Pretreatment with phorone to produce hepatic glutathione depletion had little effect on the metabolic rate of methylene chloride, demonstrating that the glutathione-mediated metabolism was not a major pathway under the conditions of the study. The authors concluded that these results demonstrate that at low exposure levels, little methylene chloride is metabolized by the GSH pathway, and thus these results confirm the results of other investigators.

Several PBPK models have been developed that can be used to predict tissue-specific exposures to methylene chloride, taking into account absorption, distribution, and metabolism. A PBPK model was used to provide quantitative estimates of the levels of methylene chloride in various organs of four mammalian species (rats, mice, hamsters, and humans) following inhalation exposure (Andersen et al. 1987). The model, which incorporates a variety of variables representing the blood and tissue concentrations of methylene chloride, exhaled methylene chloride, and instantaneous rates of metabolism by each pathway, was validated by comparing predictions of concentrations of methylene chloride in blood with time-course data obtained with Fischer-344 rats, Syrian Golden hamsters, B6C3F<sub>1</sub> mice, and human volunteers. The predicted values for each of the four species were in agreement with the experimental data. The model was also shown to predict the appearance and elimination of methylene chloride metabolites reasonably well.

Other PBPK models are discussed in Section 2.3.5.

**Oral Exposure** In a lethal poisoning case, a 56 year-old woman ingested approximately 300 mL of Nitromors, a paint remover solvent whose major ingredient is 75–80% methylene chloride (Hughes and Tracey 1993). COHb levels in the patient were increased to about 9% in blood samples taken several hours later. According to the authors, this case demonstrated that conversion of ingested methylene



## 2. HEALTH EFFECTS

chloride to COHb occurs in humans; previously, the conversion of ingested methylene chloride to COHb had only been reported in rats.

No other studies were located regarding metabolism in humans after oral exposure to methylene chloride.

Animal data on metabolism indicate that the process appears to be similar for inhalation and oral exposures. When rats were gavaged with a single oral dose of 526 mg/kg of methylene chloride, COHb levels in blood rose to nearly 10% (Wirkner et al. 1997). When methylene chloride was administered to rats and mice by gavage at daily doses of 50 and 200 mg/kg in the mice and 50 mg/kg in the rats, there was a dose-dependent biotransformation of methylene chloride to CO<sub>2</sub> and CO (Angelo et al. 1986a, 1986b). Pulmonary excretion of methylene chloride, CO, and CO<sub>2</sub> could be detected within 30 minutes of administration. Initially, exhaled CO<sub>2</sub> levels from the methylene chloride subjects exceeded CO levels in the rats. However, the amount of exhalant did increase with time. A similar profile of metabolism was apparent in mice based on exhaled methylene chloride, CO, and CO<sub>2</sub> values except that the exhaled CO<sub>2</sub> values support studies which indicated that the GST pathway is more important in mice than in rats (Green et al. 1986c, 1988).

Immediately following administration of doses of 500 or 1,000 mg/kg of methylene chloride in corn oil, the values for pulmonary excretion of methylene chloride, CO<sub>2</sub>, and CO were all lower than when methylene chloride was administered in aqueous solution, reflecting the slower absorption of methylene chloride from the corn oil vehicle (Angelo et al. 1986a). Three hours after administration the corn oil, pulmonary excretory patterns were similar, but still had lower values than those for the methylene chloride in aqueous solution.

Pankow and Jagielki (1993) studied the *in vivo* metabolism of methylene chloride to carbon monoxide as measured by the COHb level in blood under the following conditions: pretreatment with methanol; simultaneous administration of methanol; and pretreatment with glutathione-depleting chemicals. Male Wistar rats were administered 6.2 mmol (=0.4 mL)/kg body mass methylene chloride via gavage and >148 mmol/kg of methanol. Six hours following methylene chloride administration, blood samples were taken for COHb determination. The animals were then sacrificed and glutathione levels were measured in the liver. The authors concluded that the cytochrome P-450 2E1 oxidative pathway is responsible for the formation of COHb and the metabolism of methylene chloride to carbon monoxide, and suggested that the two metabolic pathways of methylene chloride (i.e, oxidation cytochrome P-450 2E1 and conjugation

## 2. HEALTH EFFECTS

via glutathione/glutathione-S-transferase) may be independent at study doses in rats. However, the rationale for this suggestion was unclear.

A comparison of the rates of pulmonary elimination of methylene chloride, CO<sub>2</sub>, and CO for oral and intravenous exposures of rats indicated that biotransformation rather than absorption was the rate-limiting factor that controlled the CO and CO<sub>2</sub> production (Angelo et al. 1986b). The same situation was true in mice when the pulmonary elimination patterns of the methylene chloride in water were compared to those from intravenous injection (Angelo et al. 1986a). This was not the case when the methylene chloride was given in corn oil.

These data suggest that both the MFO and GST pathways can participate in the metabolism of methylene chloride. Factors which influence the metabolism of methylene chloride by the oral exposure route are the rate of absorption and the distribution of the MFO and GST enzyme systems in the various tissues and the transportation of methylene chloride via the aqueous blood components to the liver or via the chylomicrons to systemic circulation.

***Dermal Exposure*** No studies were located on the metabolism of methylene chloride after dermal exposure to humans or animals.

### 2.3.4 Elimination and Excretion

Elimination and excretion of methylene chloride in children are not expected to be different from adults; however, this has not been investigated.

#### 2.3.4.1 Inhalation Exposure

Methylene chloride is removed from the body primarily in expired air and urine. In 4 human subjects exposed to 100 ppm of methylene chloride for 2 hours, an average of 22.6 µg (0.003%) methylene chloride was excreted in the urine within 24 hours after the exposure; in 7 subjects exposed to 200 ppm of methylene chloride for 2 hours, the corresponding value was 81.5 µg (0.006%) (DiVincenzo et al. 1972). The percentage values are based on a respiration rate of 1 mg/m<sup>3</sup> and the assumption that methylene chloride is completely absorbed. No data were found on amounts recovered in feces. Methylene chloride

## 2. HEALTH EFFECTS

excretion in the expired air was most evident in the first 30 minutes after exposure. Initial postexposure methylene chloride concentrations in expired breath following 2- and 4-hour exposure periods were about 20 ppm and dropped to about 5 ppm at the end of 30 minutes. Small amounts of methylene chloride remained in the expired air at 2.5 hours.

In rats, methylene chloride was excreted in the expired air, urine, and feces following a single 6-hour exposure to 50, 500, or 1,500 ppm of methylene chloride (McKenna et al. 1982). Exhaled air accounted for 58–79% of the radioactive concentration. At the 50 ppm exposure only 5% of the exhaled label was found as methylene chloride. The remainder was exhaled as CO and CO<sub>2</sub>. As the exposures increased, so did the amount of unmetabolized methylene chloride exhaled. Methylene chloride accounted for 30% of the label from the 500 ppm concentration and 55% of the label for the 1,500 ppm concentration. Exhaled methylene chloride, CO<sub>2</sub>, and CO accounted for 58, 71, and 79% of the inhaled methylene chloride for the 50, 500, and 1,500 concentrations, respectively. Urinary excretions accounted for 7.2–8.9% of the absorbed dose and 1.9–2.3% of the absorbed dose was in the feces.

### 2.3.4.2 Oral Exposure

No studies were located regarding excretion in humans after oral exposure to methylene chloride.

Expired air accounted for 78–90% of the excreted dose in rats in the 48-hour period following a 1 or 50 mg/kg methylene chloride dose in aqueous solution (McKenna and Zemple 1981). The radiolabel was present in the exhaled air as CO and CO<sub>2</sub>, as well as in expired methylene chloride. The amount of methylene chloride in the expired air increased from 12 to 72% as the dose was increased from 1 to 50 mg/kg. Radiolabel in the urine accounted for 2–5% of the dose under the above exposure conditions, while 1% or less of the dose was found in the feces. These data indicate that the lungs are the major organ of methylene chloride excretion even under oral exposure conditions.

### 2.3.4.3 Dermal Exposure

No studies were located regarding excretion in humans or animals after dermal exposure to methylene chloride.

## 2. HEALTH EFFECTS

**2.3.5 Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models**

Physiologically based pharmacokinetic (PBPK) models use mathematical descriptions of the uptake and disposition of chemical substances to quantitatively describe the relationships among critical biological processes (Krishnan et al. 1994). PBPK models are also called biologically based tissue dosimetry models. PBPK models are increasingly used in risk assessments, primarily to predict the concentration of potentially toxic moieties of a chemical that will be delivered to any given target tissue following various combinations of route, dose level, and test species (Clewell and Andersen 1985). Physiologically based pharmacodynamic (PBPD) models use mathematical descriptions of the dose-response function to quantitatively describe the relationship between target tissue dose and toxic end points.

PBPK/PD models refine our understanding of complex quantitative dose behaviors by helping to delineate and characterize the relationships between: (1) the external/exposure concentration and target tissue dose of the toxic moiety, and (2) the target tissue dose and observed responses (Andersen et al. 1987; Andersen and Krishnan 1994). These models are biologically and mechanistically based and can be used to extrapolate the pharmacokinetic behavior of chemical substances from high to low dose, from route to route, between species, and between subpopulations within a species. The biological basis of PBPK models results in more meaningful extrapolations than those generated with the more conventional use of uncertainty factors.

The PBPK model for a chemical substance is developed in four interconnected steps: (1) model representation, (2) model parametrization, (3) model simulation, and (4) model validation (Krishnan and Andersen 1994). In the early 1990s, validated PBPK models were developed for a number of toxicologically important chemical substances, both volatile and nonvolatile (Krishnan and Andersen 1994; Leung 1993). PBPK models for a particular substance require estimates of the chemical substance-specific physicochemical parameters, and species-specific physiological and biological parameters. The numerical estimates of these model parameters are incorporated within a set of differential and algebraic equations that describe the pharmacokinetic processes. Solving these differential and algebraic equations provides the predictions of tissue dose. Computers then provide process simulations based on these solutions.

The structure and mathematical expressions used in PBPK models significantly simplify the true complexities of biological systems. If the uptake and disposition of the chemical substance(s) is

## 2. HEALTH EFFECTS

adequately described, however, this simplification is desirable because data are often unavailable for many biological processes. A simplified scheme reduces the magnitude of cumulative uncertainty. The adequacy of the model is, therefore, of great importance, and model validation is essential to the use of PBPK models in risk assessment.

PBPK models improve the pharmacokinetic extrapolations used in risk assessments that identify the maximal (i.e., the safe) levels for human exposure to chemical substances (Andersen and Krishnan 1994). PBPK models provide a scientifically sound means to predict the target tissue dose of chemicals in humans who are exposed to environmental levels (for example, levels that might occur at hazardous waste sites) based on the results of studies where doses were higher or were administered in different species. Figure 2-4 shows a conceptualized representation of a PBPK model.


PBPK models for methylene chloride are discussed in this section in terms of their use in risk assessment, tissue dosimetry, and dose, route, and species extrapolations.

### **2.3.5.1 Methylene Chloride PBPK Model Comparison**

The first methylene chloride-specific pharmacokinetic model was developed by Andersen et al. (1987) for use in methylene chloride human risk assessment, formulated by integrating information on mouse physiology, methylene chloride solubility characteristics, and metabolic rate constants to describe the disposition of methylene chloride in target tissues. This model was actually a modification of earlier models developed by Ramsey and Andersen (1984) for generic PBPK analysis of volatile chemicals and by Gargas et al. (1986) for study of the kinetics and metabolism of dihalomethanes. The model could predict the time-course of the parent chemical and the production of metabolites by both GSH and MFO pathways. A more comprehensive model developed by Andersen et al. (1991) is also capable of describing the production of carbon monoxide during oxidative metabolism by the MFO pathway and the production of COHb by CO binding to blood hemoglobin. This model (Andersen et al. 1991) provided a coherent description of experimental data from both rodents and humans. In an earlier draft of this toxicological profile, the Andersen et al. (1991) model was used to conduct route-to-route extrapolation of the inhalation data in humans from the Putz et al. (1979) study to develop an equivalent oral dose that was evaluated as the basis for an acute oral MRL. (However, a different critical study has been selected and has been modeled as described below). The model calculated the peak level (6.37%) of carbon

## 2. HEALTH EFFECTS

**Figure 2-4. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance**



Source: adapted from Krishnan et al. 1994

Note: This is a conceptual representation of a physiologically based pharmacokinetic (PBPK) model for a hypothetical chemical substance. The chemical substance is shown to be absorbed via the skin, by inhalation, or by ingestion, metabolized in the liver, and excreted in the urine or by exhalation.

## 2. HEALTH EFFECTS

monoxide in the blood that produced a neurological effect (decreased visual and psychomotor performance and auditory function) following exposure to 200 ppm of methylene chloride (Putz et al. 1979). The equivalent concentration of methylene chloride in drinking water that will produce the same neurological effect was 865 mg/L. Using a daily drinking water consumption value of 2 L and an average human body weight of 70 kg, the LOAEL was calculated to be 25 mg/kg/day. Reitz et al. (1988, 1989), Reitz (1990, 1991), and Andersen and Krishnan (1994) used the Andersen et al. (1987) model to make predictions about GSH-mediated target tissue doses associated with tumorigenesis, or the lack thereof, in the long-term inhalation and drinking water bioassays. Using target tissue dosimetry as a concentration surrogate for atmospheric concentrations of methylene chloride, these investigators calculated a unit inhalation risk for methylene chloride that was two orders of magnitude lower than EPA's, using the same linearized multistage methodology.

Dankovic and Bailer (1994) used the models developed by Andersen et al. (1987) and Reitz et al. (1988, 1989; Reitz 1990, 1991) to examine the effects of exercise and intersubject variability on estimation of human dose levels of methylene chloride. Casanova et al. (1992) developed an extended PBPK model for DNA-protein crosslink (DPX) formation in mouse liver, based on the model developed by Andersen et al. (1987).

Reitz et al. (1997) and DeJongh et al. (1998) have expanded earlier models (Andersen et al. 1987, DeJongh and Blaauboer 1996, Ramsey and Andersen 1984) to include simulations of brain concentrations of methylene chloride. Reitz et al. (1997) used their model to make route-to-route dose extrapolations based on brain methylene chloride concentrations. Winneke (1974) had concluded that neurological effects (decreased visual and auditory functions) of acute exposure to methylene chloride were based primarily on the properties of the parent compound, and not on the accumulation of COHb, which had been the conclusion of Putz et al. (1979). Based on the study of Winneke (1974) in humans, an inhalation exposure to 300 ppm methylene chloride for 4 hours was predicted to result in a similar peak brain methylene chloride concentration as an exposure to 565 mg methylene chloride/L drinking water, both exposures correspond to approximately 4 mg/L of brain tissue. Using a daily drinking water consumption value of 2 L and an average human body weight of 70 kg, the LOAEL was calculated to be 16 mg/kg/day. This LOAEL was used as the basis for the acute oral MRL (see Section 2.5). DeJongh et al. (1998) used a rat PBPK model to estimate brain methylene chloride concentrations in rats corresponding to inhalation exposures to the rat 15-minute or 6-hour LC<sub>50</sub>.

## 2. HEALTH EFFECTS

Earlier models have also been extended to simulate humans during pregnancy and lactation. The Reitz et al. (1997) model is based on an earlier model of trichloroethylene pharmacokinetics in the pregnant rat (Fisher et al. 1989) and includes compartments representing mammary tissue, placenta, and the fetus. The Reitz et al. (1997) model has been used to simulate fetal doses of methylene chloride in rats that were exposed to methylene chloride by the inhalation route. Exposure to 1,250 ppm methylene chloride ( $4,342 \text{ mg/m}^3$ ) for 7 hours/day during gestation days 6–15 was predicted to result in a dose to the fetus of 3.67 mg/L of fetus. The corresponding drinking water intake in the human that would result in the same fetal dose was estimated from the human PBPK model to be 5,000 mg/L drinking water. Fisher et al. (1997) described an extension of the PBPK model of Ramsey and Andersen (1984) which simulates amounts and concentrations of methylene chloride in human breast milk that would result from inhalation exposures. The human lactation model includes a breast milk compartment and parameters for simulating breast milk production and intake of breast milk by nursing infants. Simulations of a workday maternal exposure to methylene chloride at an air concentration of 50 ppm ( $174 \text{ mg/m}^3$ ) yielded a predicted intake in the infant of 0.213 mg/24 hours.

Advances in modeling methylene chloride pharmacokinetics have also been achieved by linking methylene chloride models to other PBPK models or to other types of simulation or prediction models. For example, OSHA (1997) developed a PBPK model that utilizes a Monte Carlo approach to simulate variability in methylene chloride pharmacokinetics. The model outputs a probability distribution of the amount of methylene chloride metabolized in the lung through the glutathione-S-transferase pathway. Poulin and Krishnan (1999) developed a structure-based PBPK model that uses information on chemical structure to estimate tissue:blood and tissue:air partition coefficients needed to run the model. This approach can be used to model volatile organic chemicals for which empirically-based estimates are not available (Poulin and Krishnan 1999). A PBPK model of methylene chloride has been linked to a PBPK model of toluene (Pelekis and Krishnan 1997). The composite model has been used to simulate the pharmacokinetic and toxicodynamic outcomes (e.g., COHb concentrations) that might result from interactions between methylene chloride and toluene at the level of the mixed function oxidase pathway.



## 2. HEALTH EFFECTS

**2.3.5.2 Discussion of Methylene Chloride PBPK Models****The Reitz (1990) Model (Also the Andersen et al. [1987], the Reitz et al. [1988, 1989, 1997], and Andersen and Krishnan [1994] Models)**

**Risk assessment.** An inhalation to oral route extrapolation was pharmacokinetically modeled using data from the Haun et al. (1972) study. Reitz (1990) incorporates PBPK principles into estimating excess lifetime cancer risk for humans continuously exposed to  $1 \mu\text{g}/\text{m}^3$ . Using the linearized multistage model, the upper-bound estimate of excess lifetime human cancer risk is  $2.8 \times 10^{-4}$  ppm which differs from that of EPA ( $4.1 \times 10^{-6}$ ) by 2 orders of magnitude. It should be noted that EPA used default assumptions of low-dose linear extrapolation and of surface area adjustment to account for interspecies differences.

**Description of the model.** Reitz (1990) used the mathematical model developed by Andersen et al. (1987), which can quantitatively describe the production of metabolites in target tissues by either the MFO or glutathione (GSH) pathway, to test whether model predictions are consistent with the results obtained in the rodent inhalation and drinking water cancer bioassays of methylene chloride. The MFO pathway is oxidative and appears to yield carbon monoxide as well as considerable amounts of carbon dioxide. The glutathione-dependent pathway produces formaldehyde and carbon dioxide, but no carbon monoxide. Potentially reactive intermediates are formed in each of the metabolic pathways for methylene chloride. Distribution of methylene chloride metabolism between these pathways is dose dependent. The MFO pathway is a high-affinity, limited-capacity pathway which saturates at relatively low airborne concentrations (about 200–500 ppm). In contrast, the GSH pathway has a lower affinity for methylene chloride, and does not appear to saturate at experimental concentrations ( $<5,000$  ppm). Thus, at low concentrations, most of the methylene chloride is metabolized by the MFO pathway. As exposure concentrations increase and the MFO pathway saturates, metabolism by the secondary GSH pathway is observed.

Predicted amounts of methylene chloride metabolism produced by each pathway (mg equivalents of methylene chloride metabolized per L of liver tissue per day) are presented in Reitz's (1990) analysis. The predicted amounts of MFO metabolites formed in lung and liver tissue are nearly identical for the 2 concentration levels in the mouse inhalation cancer bioassay (2,000 and 4,000 ppm) because saturation of the MFO pathway is reached between 200 and 500 ppm of administered methylene chloride. However, mouse liver and lung tumors are consistently higher in the 4,000 ppm exposure group than in the

## 2. HEALTH EFFECTS

2,000 ppm group in this study. Therefore, there is no concentration-response relationship between rodent tumor incidence and MFO-metabolite levels. The PBPK model also predicts that levels of MFO metabolites produced by drinking water administration of 250 mg/kg/day methylene chloride to B6C3F<sub>1</sub> mice should be similar to the amounts of MFO metabolites produced in the inhalation study. Because no statistically significant increases in tumors in the drinking water study were observed, MFO metabolites would not be predicted to be involved in rodent tumorigenicity. Predicted rates of metabolism of methylene chloride by the GSH pathway correlate more clearly with the induction of lung and liver tumors, or lack thereof, in the two chronic studies. In the inhalation study, predicted levels of GSH metabolites at 4,000 ppm are higher than predicted levels of GSH metabolites at 2,000 ppm. In contrast, predicted levels of GSH metabolites formed in target tissues during drinking water administration of methylene chloride are very low. Thus, the pattern of tumor induction, or lack thereof, in both studies shows a good correlation with the rates of metabolism of methylene chloride by the GSH pathway. Because the concentration-dependency of GSH-metabolite formation is nonlinear at low concentrations, the delivered dose arriving at the target sites cannot be directly extrapolated from very high inhalation concentrations (4,000 ppm) to very low concentrations (<1 ppm) typical of human exposure. Additionally, with regard to interspecies sensitivity, humans would only be more sensitive than mice (based on the default surface area interspecies adjustment) if the parent chemical were directly responsible for observed toxicity. In the case of methylene chloride, metabolism mediated by the GSH pathway is necessary to activate methylene chloride to reactive intermediates, so this assumption is not applicable. Reitz et al. (1990) incorporates these PBPK findings and principles into estimating excess lifetime risk for humans continuously exposed to 1 µg/m<sup>3</sup>, using the linearized multistage model; his upper-bound estimate of excess lifetime human cancer risk is 2.8x10<sup>-4</sup> ppm, which differs from that of EPA (4.1x10<sup>-6</sup>) by two orders of magnitude (EPA used default assumptions of low-dose linear extrapolation and surface area adjustment for interspecies differences).

**Validation of the model.** The model was validated by comparing predicted responses with experimental data.

**Target tissues.** The target tissues were the liver and the lung.

**Species extrapolation.** Human physiological parameters and assumptions about interspecies sensitivity were derived from PBPK modeling and use in extrapolation from mice to humans.

## 2. HEALTH EFFECTS

**Interroute extrapolation.** The results and conclusions from this model application are limited to exposure via inhalation.

**The Andersen et al. (1991) Model**

**Risk assessment.** A modification of the Andersen et al. (1991) model by Reitz et al. (1997) was used to derive an acute oral MRL from inhalation data in humans from Winneke (1974). For acute neurological effects, the associated dose measure was defined as the peak concentration of methylene chloride in brain tissue of humans exposed to 300 ppm of methylene chloride for 4 hours by inhalation. The modified PBPK model calculated that the administered inhalation dose was equivalent to 3.95 mg of methylene chloride per liter of brain tissue. The equivalent administered human concentration in drinking water that will produce the same neurological effects was 565 mg of methylene chloride/liter. Using a daily water consumption of 70 kg, the LOAEL was calculated to be 16 mg/kg/day.

**Description of the model.** Andersen et al. (1991) previously developed a PBPK model expanding the original PBPK model of Ramsey and Andersen (1984) for the disposition of inhaled volatiles. This model expands the previously developed Andersen et al. (1987) model by both describing the kinetics of carbon monoxide (CO), COHb, and parent compound methylene chloride, and by comparing the inhalation kinetics of CO and methylene chloride in rats and humans. The description of CO and COHb kinetics was adapted from the Coburn-Forster-Kane equation, which assumes that most heme is bound with oxygen and that endogenous CO production is constant. Methylene chloride kinetics and metabolism were originally described by the generic PBPK model for volatile chemicals (Ramsey and Andersen (1984). Predictions in humans from the model were compared to several experimental data sets in the literature from human volunteers exposed to CO or to methylene chloride, and to experimental data collected by Andersen et al. (1991) from six male volunteers exposed to methylene chloride vapors at concentrations of 100 or 350 ppm for a period of 6 hours.

Physiological and biochemical constants for CO were first estimated by exposing rats to 200 ppm of CO for 2 hours and examining the time course of COHb after cessation of CO exposure. The CO inhalation studies provided estimates of CO diffusing capacity under free breathing and for the Haldane coefficient, i.e., the relative equilibrium distribution ratio for hemoglobin between CO and oxygen. The CO model was then coupled with the PBPK model for methylene chloride both to predict COHb time-course concentrations during and after methylene chloride exposures in rats, and to estimate the yield of CO

## 2. HEALTH EFFECTS

produced from oxidation of methylene chloride. In rats, only about 0.7 mol of CO was produced from 1 mol of methylene chloride during oxidation. The combined model predicted COHb and methylene chloride behavior following 4-hour exposures to 200 or 1,000 ppm of methylene chloride, and COHb behavior following 30-minute exposure to 5,160 ppm of methylene chloride. The rat PBPK model was scaled to predict methylene chloride, COHb, and CO kinetics in humans exposed to either methylene chloride or CO. Three human data sets from the literature were examined: inhalation of CO at 50, 100, 250, and 500 ppm; seven 30-minute inhalation exposures to 50, 100, 250, and 500 ppm of methylene chloride; and 2-hour inhalation exposures to 986 ppm of methylene chloride. Additional experimental data from volunteers exposed to 100 or 350 ppm of methylene chloride were also reported. Endogenous CO production rates and the initial amount of CO in the blood compartment varied in each study, as necessary, to provide a baseline value of COHb. The combined PBPK model accurately predicted the experimental findings in all four human studies.

**Validation of the model.** In rats, the combined model adequately represented COHb and methylene chloride behavior following 4-hour exposures to 200 or 1,000 ppm of methylene chloride, and COHb behavior following ½-hour exposure to 5,160 ppm of methylene chloride. In addition, short-duration exposures conducted with 5,000 ppm of bromochloromethane, with adjustment of metabolic parameters and partition coefficients for this different chemical, demonstrated that the PBPK model gave a good description of COHb levels for up to 6 hours postexposure. In humans, the combined PBPK model provided a good representation of the experimental data in all four studies examined.

**Target tissues.** Blood was identified as the target. Both concentrations of COHb and methylene chloride were considered in the model.

**Species extrapolation.** The application of the model was validated in both rats and humans. The authors concluded that this model could be useful for developing biological monitoring strategies for CO and methylene chloride, based on observed COHb blood concentrations following exposure.

**Interroute extrapolation.** The model was used to convert inhalation data in humans from Winneke (1974) into equivalent oral doses that were used as the basis for the acute oral MRL (Reitz 1997).

## 2. HEALTH EFFECTS

**The Dankovic and Bailer (1994) Model**

**Risk assessment.** Risk assessment was not addressed directly in this model application. However, the results indicated that some occupationally-exposed individuals may receive glutathione S-transferase-metabolized methylene chloride doses several-fold greater than human doses previously estimated by Reitz et al. (1989), based on differing levels of physical activity, and inter-individual variability in methylene chloride metabolism. Therefore, the assessment of excess lifetime human risk associated with methylene chloride exposure would be affected.

**Description of the model.** Dankovic and Bailer (1994) examined the impact of exercise and inter-individual variability on human dose estimates to methylene chloride, using the models developed by Andersen et al. (1987) and used by Reitz et al. (1989; Reitz 1990, 1991). Earlier models used physiological parameters appropriate for humans at rest and metabolic parameters based on average rates of methylene chloride metabolism. Dankovic and Bailer (1994) increased model parameters describing cardiac output, alveolar ventilation, and blood flows to tissues to account for exercise, assuming an 8-hour workday exposed to mean methylene chloride concentrations of 25 ppm. The GSH-mediated metabolized doses for human liver and lung were increased by a factor of 2.9 and 2.4, respectively, as compared with the metabolized GSH-mediated dose estimates of Reitz et al. (1989). The model was also modified to account for inter-individual variability in methylene chloride metabolism. Modeled metabolized GSH-mediated dose estimates for human liver ranged from 0 to 5.4-fold greater than the doses estimated by Reitz et al. (1989); for human lung, estimates were 0–3.6-fold greater than those of Reitz et al. (1989). The authors concluded that their results indicated that some occupationally-exposed individuals may receive GST-metabolized doses several-fold greater than human doses previously estimated.

**Validation of the model.** The model used by Dankovic and Bailer (1994) has been previously validated (Andersen et al. 1987; Reitz 1990, 1991; Reitz et al. 1988, 1989). The authors merely modified the model parameters to be consistent with light work conditions, as opposed to the resting condition parameters used in the earlier models, and to reflect inter-individual variability in methylene chloride metabolism.

## 2. HEALTH EFFECTS

**Target tissues.** Target tissues were the human liver and the human lung.

**Species extrapolation.** The model has been previously validated for use with rodents and humans.

**Interroute extrapolation.** The results and conclusions from this model application are limited to exposure via inhalation.

### **The Casanova et al. (1996) Model**

**Risk assessment.** The use of DNA-protein crosslinks as a tissue dosimeter of methylene chloride exposure markedly reduced the upper-bound estimate and improved the precision of the low-dose excess lifetime liver cancer risk estimates (as defined by the ratio of the upper-bound estimate to the maximum likelihood estimate), while having only a minor effect on the maximum likelihood estimate. The reduction in excess lifetime cancer risk was 2 orders of magnitude less than those estimated by using airborne methylene chloride vapor concentrations of 10, 30, and 100 ppm.

**Description of the model.** Casanova et al. (1996) developed an extended PBPK model for DNA-protein crosslink (DPX) formation in mouse liver associated with chronic methylene chloride exposure by modifying the model originally developed by Andersen et al. (1987). This extended PBPK model estimated area under the curve for methylene chloride in mouse liver as the independent variable. Formaldehyde, one of at least two reactive intermediates formed during glutathione-mediated metabolism, was considered to be the proximate metabolite associated with tumorigenicity. Parameter estimates for formaldehyde disposition in methylene chloride-exposed mouse liver were derived from the published literature. The amount of DPX formed in the mouse liver was estimated for methylene chloride concentrations used in rodent cancer inhalation bioassay (i.e., 2,000 and 4,000 ppm). Using the linearized multistage model, the tumor incidence data in mice were fitted to two alternative measures of exposure: DPX yields and airborne concentration of methylene chloride. The 2 dose measures gave similar maximum likelihood estimates for the excess lifetime cancer risk at administered concentrations ranging from 10 to 100 ppm, but the upper 95% confidence limit on this risk estimate was reduced by 2 orders of magnitude when DPX yield, as compared with airborne concentrations, was used as the measure of exposure. These results demonstrate that in the case of methylene chloride, the use of DNA-protein crosslinks as a dose surrogate markedly reduced the upper-bound estimate and improved the precision of the low-dose risk estimates (as defined by the ratio of the upper-bound estimate to the maximum-

## 2. HEALTH EFFECTS

likelihood estimate), while having only a slight effect on the maximum-likelihood estimate. However, Casanova et al. (1996) point out that DNA-protein crosslinks cannot be used directly as a surrogate for the internal dose in humans because human hepatocytes, unlike mouse hepatocytes, do not appear to form DNA-protein crosslinks in measurable amounts (Casanova et al. 1996). A surrogate for the internal dose, RNA adducts of formaldehyde, has been developed and can be detected in human hepatocytes exposed to methylene chloride (Casanova et al. 1996, 1997). The utility of this measure is under study.

**Validation of the model.** The model has been previously validated in other applications and for other endpoints. For this application, predicted concentrations of DPX were compared with actual concentrations induced by experimental concentrations.

**Target tissues.** The target tissue effect was the formation of DNA-protein crosslinks in mouse liver.

**Species extrapolation.** The results and conclusions from this model application are limited to DPX formation in mouse liver.

**Interroute extrapolation.** Casanova et al. (1996) point out that DNA-protein crosslinks cannot be used directly as a surrogate for the internal dose in humans, because human hepatocytes, unlike mouse hepatocytes, do not appear to form DNA-protein crosslinks in measurable amounts (Casanova et al. 1996). As a surrogate for the internal dose, RNA adducts of formaldehyde have been developed and can be detected in human hepatocytes exposed to methylene chloride (Casanova et al. 1996). The formation of this dose surrogate in the hepatocytes of different species is currently being examined.

### **The Andersen and Krishnan (1994) Model**

**Risk assessment.** The PBPK model-based risk assessment estimated an excess lifetime cancer risk to humans of  $3.7 \times 10^{-8}$  for a lifetime inhalation exposure of  $2.8 \times 10^{-4}$ , which is lower by more than two orders of magnitude than that calculated by the EPA using the linearized multistage model for low-dose extrapolation and a default body surface correction factor for interspecies scaling.

## 2. HEALTH EFFECTS

**Description of the model.** The mouse PBPK model developed earlier by Andersen et al. (1987) was validated by comparing model predictions with observed pharmacokinetic data. To predict the tissue dosimetry of methylene chloride and its metabolites in humans, the physiological parameters of the model were scaled and methylene chloride-specific parameters for humans were determined. The metabolic rate constants for humans were estimated from volunteer human exposure studies (Andersen et al. 1991). The high-dose to low-dose extrapolation and interspecies extrapolation of methylene chloride induced cancer risk were conducted with the tissue doses of the glutathione pathway metabolite predicted by the PBPK model. This was validated by demonstration of a concentration-response association between this dose metric and the degree of methylene chloride-induced rodent cancers rodent bioassay concentrations between 2,000 and 4,000 ppm. The cancer risk assessment was conducted using the linearized multistage model to relate tissue dose of methylene chloride glutathione metabolite to the increases in tumor incidence in mice exposed to high concentrations of methylene chloride by inhalation. In assessing the excess lifetime cancer risks associated with human exposure to methylene chloride, the authors assumed that humans are as sensitive (i.e., not more sensitive) as the most sensitive target species; this assumption is consistent with what is known about methylene chloride chemistry and metabolism, and about interspecies variability in carcinogenic response to xenobiotics. The PBPK model-based risk assessment estimated an excess lifetime human cancer inhalation unit risk  $3.7 \times 10^{-8}$  per inhalation exposure of  $2.8 \times 10^{-4}$ . This human risk estimate is lower by two orders of magnitude than that calculated by the EPA using the linearized multistage model for low-dose extrapolation and a default body surface adjustment factor for interspecies scaling.

**Validation of the model.** The model has been previously validated by comparing predicted with actual pharmacokinetic data.

**Target tissues.** The target tissues were the liver and the lung.

**Species extrapolation.** The animal PBPK model was used for interspecies extrapolation of pharmacokinetic behavior of methylene chloride by scaling the physiological parameters and determining chemical-specific parameters in the species of interest (i.e., humans). Metabolic rates for humans were developed from experimental literature on human volunteers. It was also assumed that humans are as sensitive (i.e., not more sensitive than) as the most sensitive target species; this assumption is consistent with what is known about methylene chloride chemistry and metabolism, and about interspecies variability in carcinogenic response to xenobiotics.



## 2. HEALTH EFFECTS

**Interroute extrapolation.** The results and conclusions from this model application are limited to exposure via inhalation.

### The Reitz et al. (1997) Model

#### 1. Inhalation Route-to-Oral Route Extrapolation in Volunteers

**Risk assessment.** In this toxicological profile, ATSDR (2000) establishes a minimal risk level (MRL) for acute inhalation using the study reported by Winneke (1974). Volunteers were exposed to 300–800 ppm of methylene chloride for approximately 4 hours and tested for neurobehavioral effects. Concentration-dependent adverse effects included decreased performance in auditory vigilance-performance tasks; and three small decrements in the visual critical flicker fusion frequency. Because similar exposures to 50–100 ppm of carbon monoxide alone did not produce these effects, the author concluded that they were mediated by methylene chloride directly and not by its oxidative metabolite, carbon monoxide. A LOAEL of 300 ppm from this study was used to determine an acute oral drinking water equivalent. Using pharmacokinetic modeling, Reitz et al. (1997) was able to estimate that an inhalation concentration of 300 ppm of methylene chloride would produce a target organ-specific (brain-specific) dose equivalent to that produced by a drinking water concentration of 565 mg/L of methylene chloride. Multiplying the drinking water concentration by the default daily water consumption rate (2 L) and dividing by the default human body weight yields an acute oral dose of 16 mg/kg/day.

**Description of the model.** Reitz et al. (1997) modified the basic PBPK methylene chloride model developed by Andersen et al. (1987), Reitz et al. (1988), and Andersen et al. (1991) in the following manner: (1) liver weights for rodents were based on the actual organ weights of control animals which were sacrificed during chronic toxicity studies at 6–18 months of age; (2) partition coefficients for methylene chloride derived from *in vitro* experiments performed by Gargas et al. (1986) were used for liver, fat, muscle, and blood, and (3) a brain compartment was added to the methylene chloride model so that central nervous system effects could be assessed in female and male rodents and humans. Size of rodent brains, blood flow rates to the brain, and partition coefficients for brain tissue were obtained from either published literature (e.g., Stott et al. 1983; Thomas 1975) or personal communication by the authors. The modified methylene chloride model thus contained six tissue compartments: fat, muscle (slowly perfused tissue), rapidly perfused tissue, liver tissue, mammary tissue, and brain tissue.

## 2. HEALTH EFFECTS

For acute neurological effects, the associated dose measure was defined as the peak concentration of methylene chloride in brain tissue (mg/L of brain tissue) of humans exposed to 300 ppm of methylene chloride for 4 hours, by inhalation. The modified PBPK model calculated that the administered inhalation concentration was equivalent to 3.95 mg methylene chloride/L of brain tissue. Human exposure patterns in the PBPK model simulated realistic human drinking water consumption patterns, (i.e., consisting of bouts of drinking during the day, with and between meals, and little-to-no drinking during the night). PBPK modeling predicted that peak concentrations of methylene chloride in the brain would increase rapidly after each episode of drinking water consumption, and then drop sharply, to near-zero, between bouts of drinking. Additionally, there would be no cumulative effects of repeated exposure. The equivalent administered human concentration in drinking water was calculated to be 565 mg methylene chloride/L. Using a daily drinking water consumption value of 2 L and an average human body weight of 70 kg, the LOAEL was calculated to be 16 mg/kg/day.

**Validation of the model.** Model predictions of brain methylene chloride concentrations in humans or rats have not been evaluated for comparability with empirical observations. However, this model has been previously validated with human and animal data.

**Target tissues.** The target organ of the model was the central nervous system.

**Species extrapolation.** There was no species extrapolation because the experiment that was pharmacokinetically-modeled (Winneke 1974) was conducted in human volunteers.

**Interroute extrapolation.** The PBPK modeling was conducted to extrapolate from the inhalation route of exposure to an oral route, specifically drinking water concentration of methylene chloride.

## **2. Inhalation Route-to-Oral Route Extrapolation and Rodent-to-Human Species Extrapolation Using PBPK Modeling of Subchronic Toxicity Data**

**Risk assessment.** ATSDR established an intermediate inhalation minimal risk level (MRL) using the study reported by Haun et al. (1972). Rats were exposed continuously to either 25 or 100 ppm of methylene chloride for 100 days. Data on liver histopathology of rats exposed to 25 or 100 ppm of methylene chloride were selected as the critical effect. The authors report that liver cytoplasmic vacuolization and Oil-Red-O staining associated with fatty deposits were observed in rats at both exposure concentrations. In mice, fatty deposits (but no liver vacuolization) were reported at the higher

## 2. HEALTH EFFECTS

exposure concentration, 100 ppm. Additionally, renal changes were also reported in this study. ATSDR considered the liver effects at 25 ppm to be adverse and established that concentration as a LOAEL. Using pharmacokinetic modeling and data on human drinking water consumption patterns, Reitz et al. (1997) estimated that a drinking water concentration of 6,170 mg/L of methylene chloride produces a target organ-specific dose equivalent to that produced by an inhalation concentration of 25 ppm. Multiplying the drinking water concentration by the default daily water consumption rate (2 L) and dividing by the default human body weight yields an intermediate oral dose of 176 mg/kg/day.

**Description of the model.** For inhalation-to-oral and rat-to-human extrapolations, Reitz et al. (1997) modified the basic PBPK methylene chloride model developed by Andersen et al. (1987), Reitz et al. (1988), and Andersen et al. (1991) in the following manner: (1) liver weights for rodents were based on the actual organ weights of control animals which were sacrificed during chronic toxicity studies at 6–18 months of age; (2) partition coefficients for methylene chloride derived from *in vitro* experiments performed by Gargas et al. (1986), were used for liver, fat, muscle, and blood; and (3) a brain compartment was added to the methylene chloride model so that central nervous system effects could be assessed in female and male rodents and humans. Size of rodent brains, blood flow rates to the brain, and partition coefficients for brain tissue were obtained from either published literature (Stott et al. 1983; Thomas 1975) or personal communication by the authors. The modified methylene chloride model thus contained six tissue compartments: fat, muscle (slowly perfused tissue), rapidly perfused tissue, liver tissue, mammary tissue, and brain tissue.

The PBPK model was utilized to compare the average daily production of methylene chloride metabolites per L of liver in rats exposed to methylene chloride via inhalation for 24 hours/day with the mean daily production of methylene chloride metabolites per L of liver in humans drinking water that contained specific concentrations of methylene chloride. The dose measure was defined as the average daily concentration of metabolites per L of liver tissue. No distinctions were made between metabolites produced via the MFO pathway and those produced during glutathione conjugation by the GSH pathway. Total metabolite production was obtained by integrating the rate of metabolite production during simulation; the result was then divided by the number of 24-hour exposure periods simulated and the volume of the liver in each species (rodent and human). At inhalation concentrations by rats of 25 ppm for 24 hours/day for 100 days, the metabolized tissue-specific dose calculated by the model was 1,259 mg

## 2. HEALTH EFFECTS

metabolites methylene chloride/L liver/day. In humans, the drinking water concentration yielding an equivalent concentration of metabolized tissue-specific dose was found to be approximately 6,170 mg/L.

Multiplying this value by 2 L/day (human drinking water consumption rate) and dividing by the default human body weight of 70 kg yields a daily dose (NOAEL) of 142 mg/kg/day.

**Validation of the model.** Model predictions of the concentrations of methylene chloride or its metabolites in liver of humans or rats have not been evaluated for comparability with empirical observations. However, this model has been previously validated with human and animal data.

**Target tissue.** The target tissue was the liver.

**Species extrapolation.** Rat-to-human extrapolation was conducted using appropriate partition coefficients, metabolic rates, and other species-specific PBPK variables.

**Interroute extrapolation.** The PBPK modeling was conducted to extrapolate from the inhalation route of exposure to an oral route, described in terms of drinking water concentration of methylene chloride.

### 3. Inhalation Route-to-Oral Route Extrapolation and Rodent-to-Human Species Extrapolation Using PBPK Modeling of Developmental Toxicity Data

**Risk assessment.** The developmental toxicity of methylene chloride in rodents was studied by Schwetz et al. (1975) who exposed rats to 1,250 ppm of methylene chloride vapors for 7 hours/day from day 6–15 of gestation. There were slight, statistically significant increases in the incidence of minor skeletal variants in the offspring of females exposed to 1,250 ppm of methylene chloride during gestation. Therefore, 1,250 ppm was considered to be a LOAEL. The PBPK model was used to extrapolate from pregnant rodents to pregnant humans and from inhalation route to oral route of exposure. Using pharmacokinetic modeling and data on human drinking water consumption patterns, Reitz et al. (1997) estimated that a maternal drinking water concentration of 5,000 mg/L of methylene chloride produces a fetal dose equivalent to that produced by a maternal inhalation concentration of 1,250 ppm. Multiplying

## 2. HEALTH EFFECTS

the drinking water concentration by the default daily water consumption rate (2 L) and dividing by the default human body weight of 70 kg yields an intermediate oral dose of 142 mg/kg/day.

**Description of the model.** For determination of developmental effects, modifications to the basic PBPK methylene chloride model (Andersen et al. 1987) were based on procedures reported by Fisher et al. (1989). The model consisted of the addition of three compartments: mammary tissue; placental tissue; and fetal tissue. The fetal dose measure was of the parent compound in fetal tissue and was estimated by integrating the fetal concentration of methylene chloride (expressed as mg/L of fetal tissue) during the exposure period in order to calculate the total area under the curve (AUC) for the exposure period. The AUC was then divided by the total number of hours during which gestational exposure occurred to give the average concentration of methylene chloride during the exposure period. Because it was assumed that human fetuses would be exposed continuously throughout gestation, the total AUC for the human fetus was divided by the total gestation period to calculate the fetal concentration of parent compound methylene chloride (not metabolites), expressed in mg methylene chloride/L of fetal tissue.

Simulation of the exposure paradigm used by Schwetz et al. (1975) with the PBPK model yielded 3.67 mg methylene chloride/L of fetal tissue in the rodent. PBPK simulation of human drinking water exposures gave equivalent values of mg methylene chloride/L of fetal tissue at methylene chloride concentrations approximately 5,000 mg/L of water. Multiplying by 2 L/day and dividing by 70 kg body weight yielded a human LOAEL of 142 mg/kg/day.

**Validation of the model.** Model predictions of the concentrations of methylene chloride in rat or human fetal tissues have not been evaluated for comparability with empirical observations. However, this model has been previously validated with human and animal data.

**Target tissues.** The target tissue was the developing fetus.

**Species extrapolation.** Rat-to-human extrapolation was connected using appropriate partition coefficients, metabolic rates, and other species-specific PBPK variables.

## 2. HEALTH EFFECTS

**Interroute extrapolation.** The PBPK modeling was conducted to extrapolate from the inhalation route of exposure to an oral route, described in terms of drinking water concentration of methylene chloride.

**The Fisher et al. (1997) Model**

**Risk assessment.** The model provides an approach to estimating rates of intake of methylene chloride and other volatile chemicals by the nursing infant for a given temporal pattern of maternal exposure and infant nursing. Simulations of a workday maternal exposure to methylene chloride at an air concentration of 50 ppm (174 mg/m<sup>3</sup>) yielded a predicted intake in the infant of 0.213 mg/24 hours (Fisher et al. 1997).

**Description of the model.** Fisher et al. (1997) described a PBPK model for estimating amounts and concentrations of methylene chloride in human breast milk that would result from inhalation exposures to methylene chloride (other air borne volatile chemicals are also simulated in the model). The human lactation model is an adaptation of the PBPK model of Ramsey and Andersen (1984), with the addition of a breast milk compartment and parameters for simulating breast milk production and intake of breast milk by nursing infants. The model simulates seven tissue compartments: blood, lung, fat, liver, richly perfused tissues, poorly perfused tissues, and breast milk. Two metabolic pathways for methylene chloride are assumed to occur exclusively in the liver. A glutathione-S-transferase pathway is represented by an allometrically scaled first order rate constant, and a mixed function oxidase pathway is represented by a Michaelis-Menton function with constants  $V_{\max}$  and  $K_m$ .

The major innovation in this model is simulation of a breast milk compartment, which allows calculations of the rates of transfer of chemicals from blood into breast milk and rates of transfer to the infant during breast feeding. The volume of the breast milk compartment is assumed to decrease from an initial volume of 0.125 L at the beginning of each nursing session to a residual volume of 0.010 L at the end of each session. The rate of change in the milk volume is simulated as the difference between a zero order production rate of 0.06 L/hour and the loss rate from nursing, defined as the product the current milk volume and a first order loss constant of 20/hour. The amount of methylene chloride in breast milk is calculated using standard PBPK algorithms for flow-limited transfer from blood, assuming that methylene chloride partitions from blood directly into breast milk according to empirically derived milk/blood

## 2. HEALTH EFFECTS

partition coefficients (Fisher et al. 1997). The resulting concentrations in breast milk are used to calculate rates of intake by the nursing infant for a given temporal pattern of maternal exposure and infant nursing.

**Validation of the model.** The human lactation model, as described in Fisher et al. (1997), has not been calibrated against an empirical data set or validated for any specific use of the model.

**Target tissues.** Output from the model described are estimates of the amount and concentration of methylene chloride in breast milk and the rate of intake of methylene chloride by a nursing infant.

**Species extrapolation.** The model is designed to predict the transfer of inhaled methylene chloride to breast milk in humans. Extrapolation to other species would require modification of the model to account for different tissue masses, blood flows, and possibly other kinetic variables.

**Interroute extrapolation.** The model is designed to simulate the pharmacokinetics of methylene chloride when exposure is by inhalation to airborne methylene chloride. The pharmacokinetics of methylene chloride would be expected to be different for other routes of exposure; therefore, the output of the model cannot be extrapolated to other exposure pathways (e.g., dietary, drinking water) or routes (e.g., oral, dermal) without modification of the model.

### The DeJongh et al. (1998) Model

**Risk assessment.** The model provides an approach to estimating the brain concentrations of methylene chloride associated with acute inhalation exposures in rats. DeJongh et al. (1998) reported the results of simulations of exposures at the 15-minute and 6-hour  $LC_{50}$  in the rat. The predicted methylene chloride concentrations in brain were as follows: 15-minute  $LC_{50}$  exposure, 95,781 ppm (331,770  $mg/m^3$ , brain 24.3 mM, brain lipid 97.6 mM; and 6-hour  $LC_{50}$  exposure, 25,181 ppm (87,469  $mg/m^3$ ), brain 17.3 mM (1,594 mg/L), brain lipid 69.5 mM (6,403 mg/L).

**Description of the model.** DeJongh et al. (1998) developed a PBPK model for estimating brain concentrations of methylene chloride and other airborne volatile chemicals in rats. The model is an adaptation of PBPK models of toluene (DeJongh and Blaauboer 1996) and styrene (Ramsey and Andersen 1984), with the addition of a brain compartment. The model simulates seven tissue compartments: blood, lung, fat, liver, richly perfused tissues, slowly perfused tissues, and brain. Two

## 2. HEALTH EFFECTS

metabolic pathways for methylene chloride are assumed to occur exclusively in the liver. A glutathione-S-transferase pathway is represented by an allometrically scaled first order rate constant, and a mixed function oxidase pathway is represented by a Michaelis-Menten function with constants  $V_{\max}$  and  $K_m$ .

The major innovation in this model is the simulation of a brain compartment, which allows calculations of the rates of transfer of methylene chloride from blood into water and lipid compartments of brain. The amount of methylene chloride in the brain is calculated using standard PBPK algorithms for flow-limited transfer from blood, assuming that methylene chloride partitions from blood directly into the brain according to an empirically-derived blood-brain partition coefficient (Gargas et al. 1989). Partitioning of methylene chloride between aqueous and lipid compartments in the brain is calculated from an octanol-water partition coefficient (Meylan and Howard 1995).

**Validation of the model.** The model, as described in DeJongh et al. (1998), has not been calibrated against an empirical data set or validated for predicting brain methylene chloride concentrations.

**Target tissues.** Outputs from the model described in DeJongh et al. (1998) are estimates of the amount and concentration of methylene chloride in whole brain or brain lipid.

**Species extrapolation.** The model is designed to predict the distribution of inhaled methylene chloride to the brain in rats. Extrapolation to other species would require modification of the model to account for different tissue masses, blood flows, and possibly other kinetic variables.

**Interroute extrapolation.** The model is designed to simulate the pharmacokinetics of methylene chloride when exposure is by inhalation to airborne methylene chloride. The kinetics would be expected to be different for other routes of exposure; therefore, the output of the model cannot be extrapolated to other exposure pathways (e.g., dietary, drinking water) or routes (e.g., oral, dermal) without modification of the model.

### **The Poulin and Krishnan (1999) Model**

**Risk assessment.** The model provides an approach to estimating the concentrations of methylene chloride in venous blood, and possibly other tissues, associated with acute inhalation exposures. Poulin and Krishnan (1998) reported the results of simulations of a 6-hour inhalation exposure of an adult human



## 2. HEALTH EFFECTS

to 100 ppm methylene chloride (347 mg/m<sup>3</sup>). The predicted methylene chloride concentrations in venous blood after 6 hours of exposure were approximately 0.5 and 1.5 mg/L for the bounding assumptions, liver extraction ratio of one or zero, respectively.

**Description of the model.** Poulin and Krishnan (1999) described a quantitative structure-toxicokinetic relationship (QST<sub>k</sub>R) model for estimating venous blood concentrations, and tissue:air and tissue:blood partition coefficients of methylene chloride (and other airborne volatile chemicals) in humans. The QST<sub>k</sub>R model is an adaptation of the PBPK model of Andersen et al. (1991) with the addition of a model for estimating values of the partition coefficients based on molecular structure fragment and tissue composition information. The model simulates blood, lung, fat, liver, richly perfused tissues, and slowly perfused tissues. Two innovations in this model are the approach to the simulation of methylene chloride metabolism in the liver and the inclusion of the structure-based estimation of partition coefficients.

The rate of metabolism of methylene chloride in the liver is represented as the products of the liver blood flow rate, the arterial concentration of methylene chloride, and the liver extraction ratio (ratio of hepatic clearance to hepatic blood flow). This approach does not require specification of values for the kinetic constants of metabolism (e.g., V<sub>max</sub>, K<sub>m</sub>), but does require specification of a value, or range of values, for the liver extraction ratio. In the absence of an empirical basis for any given value for the extraction ratio, bounding estimates for venous blood concentrations, as affected by hepatic metabolism, can be made by running simulations in which the extraction ratio is assumed to be either zero or one (Poulin and Krishnan 1999).

The octanol:water partition coefficient, the water:air partition coefficient, and the boiling point are estimated using molecular structure fragment information (Poulin and Krishnan 1996, 1998). The estimated values for the above parameters are used with tissue composition information (e.g., water and lipid content) to estimate values for blood:air and tissue:blood partition coefficients.

**Validation of the model.** Estimates of blood:air and tissue:blood concentration predicted with the QST<sub>k</sub>R compared well with empirical determinations for methylene chloride (and a variety of other volatile organic compounds). Measured venous blood concentrations of methylene chloride in humans exposed to 100 ppm methylene chloride (347 mg/m<sup>3</sup>) for 6 hours (Andersen et al. 1991) were within the bounding estimates (liver extraction ratio assumed to be zero or one) predicted with the QST<sub>k</sub>R.

## 2. HEALTH EFFECTS

**Target tissues.** Output from the model described in Poulin and Krishnan (1998) are estimates of the concentration of methylene chloride in venous blood.

**Species extrapolation.** The model is designed to predict venous blood concentration of methylene chloride in humans. Extrapolation to other species would require modification of the model to account for different tissue masses, blood flows, and possibly other kinetic variables.

**Interroute extrapolation.** The model is designed to simulate the pharmacokinetics of methylene chloride when exposure is by inhalation to airborne methylene chloride. The kinetics would be expected to be different for other routes of exposure; therefore, the output of the model cannot be extrapolated to other exposure pathways (e.g., dietary, drinking water) or routes (e.g., oral, dermal) without modification of the model.

### **The Pelekis and Krishnan (1997) Model**

**Risk assessment.** The model provides an approach to estimating the COHb levels in blood that would result from oral or inhalation exposures to mixtures of methylene chloride and toluene if certain assumptions are made regarding the mechanism of interaction between the two chemicals: (1) the interaction occurs at the level of the mixed function oxidase-mediated metabolism of methylene chloride to carbon monoxide in the liver; and (2) the mechanism of inhibition can be simulated by Michaelis-Menten type models of competitive, noncompetitive or uncompetitive inhibition of the pathway. Pelekis and Krishnan (1997) extrapolated the rat model to humans by replacing the rat values with human values for physiological variables and by assuming that partition coefficients and the metabolism kinetic constants are the same in the rat and human. The resulting human model was used to predict the area under the COHb-time curve for a human exposure to the ACGIH 8-hour TLV for methylene chloride (50 ppm, 174 mg/m<sup>3</sup>), or to a simultaneous exposure to the methylene chloride and toluene at their respective TLVs (50 ppm, 188 mg/m<sup>3</sup>). The model predicted a 2–9% decrease in COHb levels in the mixed exposure compared to the exposure to methylene chloride alone, depending on which mechanism of inhibition of the mixed function oxidase pathway was assumed.

**Description of the model.** Pelekis and Krishnan (1997) linked a PBPK model for methylene chloride (Andersen et al. 1991) with a PBPK model of toluene (Tardif et al. 1993). The composite model was used to simulate the kinetics of COHb production resulting from exposures to mixtures of the two

## 2. HEALTH EFFECTS

chemicals in rats or humans. The model simulates seven tissue compartments: blood, gastrointestinal tract, lung, fat, liver, richly perfused tissues, and slowly perfused tissues. Metabolic pathways for methylene chloride are assumed to occur exclusively in the liver. A glutathione-S-transferase pathway is represented by a first order rate constant, and a mixed function oxidase pathway, which produces carbon monoxide as a product, is represented by a Michaelis-Menton function with constants  $V_{max}$  and  $K_m$ . The metabolism of toluene is assumed to occur in the liver through a mixed function oxidase pathway. An innovation in this model is the simulation of interactions between methylene chloride and toluene at the level of the mixed function oxidase pathway. This is achieved with Michaelis-Menton equations that simulate four possible interaction possibilities: (1) no interaction between methylene chloride and toluene; (2) competitive inhibition of the mixed function oxidase pathway (increase in apparent  $K_m$ ); (3) noncompetitive inhibition of the pathway (decrease in  $V_{max}$ ); or (4) uncompetitive inhibition (increase in  $K_m$  and decrease in  $V_{max}$ ). The relationship between carbon monoxide production (including endogenous production) and percent COHb in blood is simulated using the Coburn-Foster-Kane model, as implemented in the model described by Andersen et al. (1991).

**Validation of the model.** The results of simulations were compared with observations made in studies of single or mixed exposures to methylene chloride and toluene in rats (Ciuchta et al. 1979; Pankow et al. 1991a, 1991b). Pankow et al. (1991a, 1991b) reported the levels of COHb in rats exposed to either a single oral dose of methylene chloride (6.2 mmol/kg, 527 mg/kg), or combined single oral doses of 6.2 mmol/kg methylene chloride (527 mg/kg) and 18.8 mmol/kg toluene (1,732 mg/kg) (Pankow et al. 1991a, 1991b). The observed COHb level 6 hours after dosing with methylene chloride was 9.3% compared to 1.7% after combined dosing with methylene chloride and toluene. The corresponding COHb levels simulated with the PBPK model were 8.7% for dosing with methylene chloride alone, and 2.1% for the combined dosing, assuming competitive inhibition of the mixed function oxidase pathway, or 0.6%, assuming either noncompetitive or uncompetitive inhibition (Pelekis and Krishnan 1997). The PBPK model also predicted COHb levels 12 hours after dosing with methylene chloride alone and the corresponding 12-hour area under the COHb-time curve that agreed well with observations from Pankow et al. (1991). Observed peak COHb levels following exposure of rats for 1 hour to 5,000 ppm (17,368 mg/m<sup>3</sup>) methylene chloride were 10–12% compared to less than 1% when the methylene chloride exposure occurred 30 minutes after a single intraperitoneal dose of 0.005 mmol/kg (0.46 mg/kg) of toluene (Ciuchta et al. 1979). Corresponding model simulations agreed well with these observations, only when uncompetitive or noncompetitive inhibition of the mixed function oxidase pathway was assumed. The competitive inhibition model severely overestimated the peak COHb concentrations observed in the

## 2. HEALTH EFFECTS

combined exposure experiment. These results suggest that, of the three interactions mechanisms simulated, the noncompetitive and uncompetitive mechanisms more accurately predicted the *in vivo* observations.

**Target tissues.** Output from the model described in Pelekis and Krishnan (1997) are estimates of the COHb level in arterial blood (expressed in units of percent saturation of hemoglobin).

**Species extrapolation.** The model was calibrated against observed arterial blood COHb levels in rats. Although Pelekis and Krishnan (1997) have extrapolated the rat model to simulate human exposures to methylene chloride and toluene, model outputs have not been compared to empirical observations in humans; therefore, the accuracy of the extrapolation has not been evaluated. Extrapolation to other species (other than human) would require additional modifications to the model to account for different tissue masses, blood flows, and possibly other kinetic variables.

**Interroute extrapolation.** The model is designed to simulate the pharmacokinetics of methylene chloride and toluene when exposures are by the inhalation or oral routes. COHb levels predicted by the model are highly sensitive to the assumed value of the gastrointestinal absorption rate constant of methylene chloride, and less sensitive to the value of the rate constant for absorption of toluene (Pelekis and Krishnan 1997). This suggests that potential dose level or exposure medium effects (e.g., diet, drinking water) or interaction effects on the absorption rate constants should be considered in simulations of the oral route. The kinetics would be expected to be different for other routes of exposure; therefore, the output of the models cannot be extrapolated to other exposure pathways (e.g., dermal) without modification of the model.

### The OSHA (1997) Model

**Risk assessment.** The OSHA model provides an approach to estimating various internal dose surrogates corresponding to inhalation exposures to methylene chloride. OSHA (1997) used the mouse PBPK model to translate methylene chloride inhalation exposure levels used in an NTP (1986) mouse bioassay to equivalent amounts of methylene chloride metabolized by the mouse lung glutathione-S-transferase pathway. The resulting internal dose estimates were used to establish an internal dose-response relationship for lung tumors observed in the mouse bioassay. The mouse internal dose-response relationship was then translated into an equivalent human internal dose-response relationship using the

## 2. HEALTH EFFECTS

human PBPK model. The mean and 95th percentile cancer risks (based on the maximum likelihood of the dose-response parameters) associated with a human exposure to 25 ppm methylene chloride ( $87 \text{ mg/m}^3$ ), 8 hours/day, 5 days/week for 45 years were  $1.24 \times 10^{-3}$  and  $3.63 \times 10^{-3}$ , respectively.

**Description of the model.** OSHA (1997) proposed a PBPK model in support of its Final Rule on limits for occupational exposure to methylene chloride. The OSHA model is based on several more recent extensions of the Andersen et al. (1991) model as described by Clewell (unpublished report available as an exhibit in OSHA (1997) and Reitz et al. (1997)). The model simulates eight tissue compartments: blood, gastrointestinal tract, lung, fat, liver, well perfused tissues, poorly perfused tissues, and bone marrow.

Innovations in the OSHA model include the following. (1) Bone marrow is simulated as a distinct tissue rather than including it in the well or poorly perfused tissue compartment. (2) Metabolism of methylene chloride is assumed to occur in the liver and lung tissues. The liver and lung  $K_m$  values for the mixed function oxidase pathway are assumed to be identical. The values for the  $V_{\max}$  for the mixed function oxidase pathway in the lung and the rate constant for the glutathione-S-transferase pathway in the lung are set as fixed fractions of their respective values in liver. These estimates take into account relative tissue volumes, *in vitro* estimates of metabolism rates in the two tissues, and the relative abundance of microsomal or soluble protein in the two tissues. (3) Alveolar ventilation rates are dependent on cardiac output. The relationship between the two variables is assumed to be a direct proportionality with a ventilation-perfusion ratio as the proportionality term. (4) Cardiac output, tissue distribution of cardiac output, and the ventilation-perfusion ratio are related to work intensity allowing various work-related exposure scenarios to be simulated. (5) Tissue blood flows, as a fraction of the cardiac output, are constrained so that either the fractional flow to the well-perfused tissues (mouse model) or poorly perfused tissues (human model) is set as 1 minus the sum of the fractional flows to all other tissues. This constraint was imposed as an approach to ensure mass balance of flows when a Monte Carlo sampling approach was used to select parameter values.

Probability distributions for input parameters in the mouse and human versions of the OSHA model were developed using a Bayesian analysis of empirical data from gas uptake studies in mice and human open chamber inhalation studies, respectively. The resulting probability distributions were used in a Monte Carlo approach to implement the PBPK models. This approach results in probability distributions for model outputs, rather than single point estimates.

## 2. HEALTH EFFECTS

**Validation of the model.** The mouse and human models were calibrated using data from gas uptake studies in mice and human open chamber inhalation studies (OSHA 1997). Comparisons of model outputs with empirical observations other than those used in the Bayesian analysis were not reported in OSHA (1997).

**Target tissues.** The mouse and human models have been used to estimate the amounts of methylene chloride metabolized through the glutathione-S-transferase pathway in liver and lung (OSHA 1997).

**Species extrapolation.** Separate models for the mouse and human have been developed, enabling extrapolations of internal dose surrogates between these two species (OSHA 1997). Extrapolation to other species would require additional modifications to the models to account for different tissue masses, blood flows, and possibly other kinetic variables.

**Interroute extrapolation.** The mouse and human models have been used to simulate the pharmacokinetics of methylene chloride when exposures are by the inhalation route (OSHA 1997). The models include gastrointestinal tissue compartment and, therefore, could be applied to oral exposures, provided that validation studies support such uses of the model. The kinetics would be expected to be different for other routes of exposure; therefore, the output of the models cannot be extrapolated to other exposure pathways (e.g., dermal) without modification of the models.

## 2.4 MECHANISMS OF ACTION

### 2.4.1 Pharmacokinetic Mechanisms

The physical properties of methylene chloride, particularly its lipophilic nature, high vapor pressure, and high serum/air partition coefficient, suggest that it is likely to be absorbed across the alveolar membranes of the lung, mucosal membranes of the gastrointestinal tract, and the skin by passive diffusion. Once in the body, it is widely distributed, with the greatest amounts accumulating in the more lipophilic tissues; this probably also occurs by passive diffusion.

Andersen et al. (1987), Reitz (1990, 1991), Reitz et al. (1988), and Andersen and Krishnan (1994) have used a PBPK model to predict amounts of methylene chloride metabolism produced (mg equivalents of methylene chloride metabolized per L of liver tissue per day) produced by the two major pathways of

## 2. HEALTH EFFECTS

methylene chloride metabolism: the MFO pathway and the GSH-mediated pathway. This model identifies the pathway which appears to activate methylene chloride to reactive intermediates but neither characterizes the putative metabolite nor describes the mechanisms of action. In essence, the MFO pathway is oxidative and appears to yield carbon monoxide as well as considerable amounts of carbon dioxide. The glutathione-dependent pathway produces formaldehyde and carbon dioxide, but no carbon monoxide. Potentially reactive intermediates are formed in each of the metabolic pathways for methylene chloride: formyl chloride via the MFO pathway and methylchlorogluthathione via the GSH pathway. Distribution of methylene chloride metabolism between these pathways is dose dependent. The MFO pathway is a high-affinity, limited-capacity pathway which saturates at relatively low airborne concentrations (about 200–500 ppm). In contrast, the GSH pathway has a lower affinity for methylene chloride, and does not appear to saturate at experimental concentrations (<5,000 ppm). Thus, at low concentrations, most of the methylene chloride is metabolized by the MFO pathway. As exposure concentrations increase and the MFO pathway saturates, metabolism by the secondary GSH pathway is observed.

Predicted amounts of methylene chloride metabolism produced by each pathway (mg equivalents of methylene chloride metabolized per L of liver tissue per day) are presented in Reitz's (1990, 1991) analyses. The predicted amounts of MFO metabolites formed in lung and liver tissue are nearly identical for the 2 concentration levels in the mouse inhalation cancer bioassay (2,000 and 4,000 ppm) because saturation of the MFO pathway occurs at administered airborne concentrations of approximately 200–500 ppm. However, mouse liver and lung tumors are consistently higher in the 4,000 ppm exposure group than in the 2,000 ppm group in this bioassay. Therefore, there is no concentration-response relationship between rodent tumor incidence and MFO metabolite levels. The PBPK model also predicts that levels of MFO metabolites produced by long-term drinking water administration of 250 mg/kg/day methylene chloride to B6C3F<sub>1</sub> mice should be similar to the levels of MFO metabolites produced in the inhalation bioassay. However, no statistically significant increases in tumors in the drinking water bioassay were observed; therefore, MFO metabolites would not be predicted to be involved in rodent tumorigenicity by either route of administration. Predicted rates of metabolism of methylene chloride by the GSH pathway correlate more clearly with the induction of lung and liver tumors, or lack thereof, in the two chronic studies. In the inhalation study, predicted levels of GSH metabolites at 4,000 ppm are higher than predicted levels of GSH metabolites at 2,000 ppm. In contrast, predicted levels of GSH metabolites formed in target tissues during drinking water administration of methylene chloride are very low. Thus, the pattern of tumor induction, or lack thereof, in both studies shows a good correlation with

## 2. HEALTH EFFECTS

the rates of metabolism of methylene chloride by the GSH pathway. Because the dose dependency of GSH metabolite formation is nonlinear at low concentrations, the delivered dose arriving at the target sites cannot be directly extrapolated from very high inhalation concentrations (4,000 ppm) to the very low concentrations (<1 ppm) typical of human exposure.

Casanova et al. (1996) developed an extended PBPK model for DNA-protein crosslink (DPX) formation in mouse liver, based on the model originally developed by Andersen et al. (1987). The extended PBPK model estimated area under the curve for methylene chloride in mouse liver as the independent variable. Tissue-specific yields of DPX were used as the dose surrogate. Estimates were made of the amount of DPX formed in the mouse liver at methylene chloride inhalation concentrations used in the bioassay (i.e., 2,000 and 4,000 ppm) and plotted against liver tumor yields in the mouse. DPX thus served as a concentration surrogate for airborne methylene chloride concentrations. The model assumes that DPX formation is associated with methylene chloride mouse liver tumorigenicity. Because formaldehyde produces DPX, and GSH-mediated metabolism of methylene chloride produces formaldehyde as a reactive intermediate, the authors suggest that formaldehyde is involved in the genotoxic mechanism(s) of action associated with mouse liver tumorigenicity. When excess lifetime cancer risk was estimated using the mouse liver tumor data and two alternate dose measures, DPX and airborne vapor concentrations, the maximum likelihood estimates were similar, but the upper-bound estimate using DPX was two orders of magnitude lower than that using airborne concentrations. Thus, DPX appears to be a reasonable dose surrogate for methylene chloride inhalation exposure. However, most other investigators do not consider DPX formed by the weakly mutagenic activity of formaldehyde to be the putative mechanism of action of methylene chloride-induced liver tumorigenicity (see Section 2.4.2, Mechanisms of Toxicity). Furthermore, human hepatocytes do not appear to form DPX in measurable amounts, as do mouse hepatocytes (Casanova et al. 1996).

### 2.4.2 Mechanisms of Toxicity

The lung, the blood system, and the nervous system are the major target organs of toxicity associated with exposure to methylene chloride.

***Non-neoplastic Mechanisms.*** In humans, Snyder et al. (1992a, 1992b) have reported headache, chest discomfort, cough, and the presence of alveolar and interstitial infiltrates in the lung as a result of short-term high-concentration vapor exposure to methylene chloride in confined, unventilated rooms or



## 2. HEALTH EFFECTS

basements. In B6C3F<sub>1</sub> mice exposed to 4,000 ppm of methylene chloride vapors for 6 hours (Foster et al. 1992), the major initial morphological effect observed in mouse lung was acute Clara cell damage. However, the damage appeared to resolve after five consecutive daily exposures to methylene chloride. The appearance and disappearance of the lesion in the Clara cell correlated well with the activity of cytochrome P-450 monooxygenase in the Clara cell, as assessed immunocytochemically (CYP2B1 and CYP2B2) in the whole lung and biochemically in the freshly isolated Clara cell (as determined by ethoxycoumarin O-dealkylation and aldrin epoxidation).

Over 13 weeks (5 days/week) of exposure, the acute Clara cell damage, which developed after a 1-day exposure but resolved after 5 consecutive exposures, reappeared on reexposure after a 2-day weekly break. The severity of the lesion diminished as the study progressed. The authors suggest that the reason for the decrease or disappearance of the lesion was due to an adaptation/tolerance in the Clara cell to methylene chloride that was linked to a marked decrease of methylene chloride metabolism by cytochrome P-450 pathways. Glutathione (GST) activity in the Clara cell either remained unchanged or increased following methylene chloride exposure.

Inhalation and ingestion exposures to methylene chloride result in the production of carbon monoxide associated mainly with metabolism via the MFO pathway. CO binds to hemoglobin, and can cause carboxyhemoglobinemia. In two fatal human cases of methylene chloride poisoning, COHb was elevated to approximately 30% (Manno et al. 1992). Other reports on human and animals show that COHb increases from baselines of 0–2 to 4–15%, under varying regimes of methylene chloride inhalation exposure.

Neurotoxicity resulting from exposure to methylene chloride is believed to be associated with the lipophilic properties of methylene chloride; however, the precise mechanisms of neurotoxicity are not known. Presumably, the methylene chloride enters cell membranes, which in the case of neurons, interferes with signal transmission, in a manner similar to general anesthetics (De Jongh et al. 1998; Sikkema et al. 1995). Neurotoxicity is also assumed to be caused by the hypoxia that results from the formation of COHb.

**Cancer.** With regard to tumor induction in the rodent lung and liver, methylene chloride is postulated to be activated to an unknown reactive intermediate via metabolism. There are two major metabolic pathways: the MFO pathway, specifically cytochrome P-450 2E1 and glutathione-glutathione

## 2. HEALTH EFFECTS

S-transferase-mediated (GSH-GST) pathway. The isoenzyme involved in the GSH-GST pathway has been identified as a  $\Theta$  (theta) class glutathione S-transferase, GSTT1-1, which is present in moderate quantities in the mouse lung, but has been detected only at very low levels in human lung and liver tissue samples (Mainwaring et al. 1996b; Sheratt et al. 1997). These findings suggest that in humans the lung and liver are likely to have little capacity to activate methylene chloride into its reactive metabolites. However, higher than background levels of GSTT1-1 mRNA were detected in a small number of Clara cells and alveolar/bronchiolar ciliated epithelial cells of one human lung sample (out of four) and of GSTT2-2 enzyme in the biliary epithelium of the human liver (Mainwaring et al. 1996b). Thus, it is possible that, in some individuals, these specific cell types may be vulnerable to genotoxic effects from reactive intermediates of methylene chloride metabolism, although the overall risk is likely to be low.

The MFO pathway is oxidative and appears to yield carbon monoxide as well as considerable amounts of carbon dioxide. The glutathione-dependent pathway produces formaldehyde and carbon dioxide, but no carbon monoxide. Potentially reactive intermediates are formed in each of the metabolic pathways for methylene chloride: formyl chloride in the oxidative pathway, and formaldehyde and chloromethyl glutathione in the conjugative pathway. Neither formyl chloride nor the glutathione conjugate of methylene chloride has been isolated or characterized, although Green (1997) reports that their formation is entirely consistent with available information on glutathione-mediated metabolism. Distribution of methylene chloride metabolism between these pathways is dose dependent. The MFO pathway is a high-affinity, limited-capacity pathway which saturates at relatively low atmospheric concentrations (approximately 200–500 ppm). The GSH pathway, in contrast, has a lower affinity for methylene chloride, but does not appear to saturate at experimentally produced concentrations (<5,000 ppm). Thus, the MFO pathway accounts for most of the metabolized methylene chloride at concentrations less than 500 ppm, but as exposure concentrations increase above the MFO saturation level, increases in the amount of methylene chloride metabolized by the secondary GSH pathway are seen (Reitz 1990). The concentration dependency of these two metabolic pathways is consistent with the tumor results obtained in long-term rodent inhalation and drinking water cancer bioassays of methylene chloride and supports the assertion that GSH-mediated metabolism is responsible for methylene chloride-induced tumorigenicity in B6CF<sub>1</sub> mice.

There is no evidence to suggest that methylene chloride is a direct acting carcinogen; the marked species differences in carcinogenicity induced by methylene chloride are not typical behavior of direct-acting compounds. Methylene chloride also does not exhibit the chemical reactivity towards nucleophiles

## 2. HEALTH EFFECTS

normally associated with direct action (Green 1997). Therefore, metabolic activation is required which interacts in some way with mouse tissues to cause tumors.

A series of bacterial mutagenicity tests has demonstrated that: methylene chloride induction of bacterial mutagenicity is expressed more strongly in *Salmonella typhimurium* TA 1535 modified to express a mammalian GST  $\Theta$  class enzyme (NM5004 strain) than in the original strain (Oda et al. 1996); methylene chloride induction of bacterial mutagenicity *S. typhimurium* strain TA 100 is unaffected by the presence of GST  $\alpha$  or  $\pi$  classes (Simula et al. 1993); methylene chloride is less mutagenic in a *S. typhimurium* GSH-deficient strain (TA100/NG11) as compared to TA 100 (Graves et al. 1994a); and bacterial testing with 3 K12 strains of *Escherichia coli* showed that methylene chloride (activated by S9 mouse liver fraction) and formaldehyde were mutagenic only in the wild-type *E. coli*, a characteristic shared with crosslinking agents; these data initially suggested a mutagenic role for metabolically-derived formaldehyde in *E. coli* (Graves et al. 1994a).

These bacterial assays demonstrated that in *in vitro* tests, methylene chloride was activated by a  $\Theta$  class GST enzyme to a bacterial mutagen in *S. typhimurium* and behaved similarly to formaldehyde in *E. coli* tester strains. However, in the Chinese Hamster ovary (CHO) assay involving the hypoxanthine-guanine phosphoribosyl transferase (HPRT) gene assay, studies of DNA single strand breaks and DNA-protein crosslinks at mutagenic concentrations of methylene chloride and formaldehyde showed that both these compounds induced DNA single-strand breaks; only formaldehyde induced significant DNA-protein crosslinking (Graves et al. 1996). Similar findings were observed in cultured, freshly isolated mouse hepatocytes (Graves and Green 1996), but not in rat hepatocytes (Graves et al. 1994b, 1995). The authors concluded that, although formaldehyde might play a role in methylene chloride genotoxicity, its weak mutagenicity and the absence of methylene chloride-induced DNA-protein crosslinking in the CHO/HPRT assay suggested that methylene chloride-induced DNA damage and resulting mutations are likely produced by its glutathione conjugate, putatively chloromethylglutathione. Graves and Green (1996) also concluded that these results suggested that the mechanism for methylene chloride tumorigenicity in the mouse liver was likely to be genotoxic and mediated by the GSH pathway. Observed species differences in liver tumorigenicity between the mouse and the rat might result from species differences in the amount of GSH-mediated metabolism induced by methylene chloride exposure.

A series of studies have been conducted to elucidate the precise genetic mechanisms of methylene chloride carcinogenicity. Female B6C3F<sub>1</sub> mice were exposed to vapor concentrations of

## 2. HEALTH EFFECTS

2,000–8,000 ppm for 2 years and sacrificed at various intervals to evaluate a number of genotoxic endpoints.

Replicative DNA synthesis in the bronchiolar epithelium, examined by the use of the labeling index (LI), indicated that mice exposed to 2,000 ppm of methylene chloride for 2–26 weeks decreased to 40–60% of controls (Kanno et al. 1993). Mice exposed to 8,000 ppm of methylene chloride have a less dramatic decrease in LI. No pathological changes were found in the exposed lungs. Thus, high-concentration exposure to methylene chloride for up to 26 weeks reduces the cell turnover of bronchiolar cells in these mice; therefore, increased cell proliferation does not appear to be involved in mouse lung tumorigenesis. DNA single-strand breaks were detected in the livers of B6C3F<sub>1</sub> male mice, but not Syrian Golden male hamsters, immediately following inhalation exposure to 2,000–8,000 ppm for 6 hours, but not 2 hours after exposure, suggesting active DNA repair (Graves et al. 1995). The DNA of mouse Clara cells incubated *in vitro* with methylene chloride was also damaged at high concentrations. Pretreatment of mice with a glutathione depletor prior to inhalation exposure caused a decrease in the amount of DNA damage detected, suggesting a GST-mediated mechanism; similar findings were observed in Clara cells incubated *in vitro* with methylene chloride and a glutathione depletor.

Devereux et al. (1993) analyzed liver and lung tumor, induced in female B6C3F<sub>1</sub> female mice by inhalation of 2,000 ppm of methylene chloride for 6 hours/day, 5 days/week exposure for up to 104 weeks, for the presence of activated *ras* proto-oncogenes. In methylene chloride-induced liver tumors, mutations, mainly transversions or transitions in base 1 or base 2, were detected and were similar to those observed for the H-*ras* gene in spontaneous liver tumors. Mutations were also identified in the lung. The K-*ras* activation profile in the methylene chloride-induced tumors was not significantly different from the profile in spontaneously-occurring tumors. No other transforming genes were found in the nude mouse tumorigenicity assay. The authors concluded that at present, no transforming genes other than *ras* genes could be identified in either mouse liver or lung tumors. Based on liver tumor data, they also suggested that methylene chloride may affect the liver by promoting cells with spontaneous lesions.

Hegi et al. (1993) studied allelotypes of 38 methylene chloride-induced lung carcinomas from female B6C3F<sub>1</sub> mice exposed 6 hours/day, 5 days/week for 2 years to 2,000 ppm. The allelotypes were examined for various genotoxic endpoints, and the results were compared with genotoxicity findings in two other reciprocal-cross mouse strains. Throughout the genome, allelic losses occurred infrequently,

## 2. HEALTH EFFECTS

except for markers on chromosome 4, which were lost in approximately half of the carcinomas. In lung adenomas, chromosome 4 losses were associated with malignant conversion. Methylene chloride-induced liver tumors did not demonstrate chromosome 4 loss, which indicated that this finding was specific for lung carcinomas. Preferential loss of the maternal chromosome 4 was also observed in carcinomas in B6C3F<sub>1</sub> mice. On chromosome 6, an association between *K-ras* gene activation and allelic imbalances was also found in B6C3F<sub>1</sub> mouse lung tumors. When allelotypes of tumors in mice from two reciprocal cross strains, AC3F<sub>1</sub> and C3AF<sub>1</sub>, were examined and compared to the findings in B6C3F<sub>1</sub> mice, one allele of the putative chromosome 4 tumor suppressor gene was shown to be inactivated. Whereas the results in B6C3F<sub>1</sub> mice suggested that nondisjunction events were responsible for the chromosome 4 losses, tumors from both reciprocal-cross mouse strains appeared to show small interstitial deletions in a chromosomal region homologous with a region in human chromosome which is often lost in a variety of tumors, including lung cancers. In human chromosomes, a candidate tumor suppressor gene, *MTS1*, is located in this region.

In another genotoxic analysis with the same cohort (Hegi et al. 1994), loss of heterozygosity at markers near the *p53* gene on chromosome 11 and within the retinoblastoma tumor suppressor gene were examined in methylene chloride-induced liver and lung tumors and compared to spontaneous tumors in control mice. The authors concluded that inactivations of *p53* and the retinoblastoma tumor suppressor gene were infrequent events in lung and liver tumorigenesis in mice exposed to methylene chloride.

Replicative DNA synthesis was examined by Kanno et al. (1993) to evaluate the potential role of treatment-induced lung cell proliferation on pulmonary carcinogenicity in female B6C3F<sub>1</sub> mice exposed to 2,000 or 8,000 ppm of methylene chloride for 6 hours/day, 5 days/week for 2 years. By the end of the study, there was a statistically significant increase in lung tumors in exposed animals when compared to controls. Cell proliferation was assessed in the lung after 1, 2, 3, or 4 weeks of inhalation exposure to 2,000 or 8,000 ppm, and after 13 and 26 weeks exposure to 2,000 ppm, as measured by changes in labeling indices (LI). The LI of both bronchiolar epithelium and terminal bronchioles were substantially decreased in mice exposed to 2,000 ppm of methylene chloride for 2–26 weeks. Similar findings, but not as severe, were observed in mice exposed to 8,000 ppm. The decreases in LI were not accompanied by cytotoxicity. The authors concluded that high-concentration exposure to methylene chloride for up to 26 weeks reduces cell proliferation in lung epithelial cells in female B6C3F<sub>1</sub> mice.

## 2. HEALTH EFFECTS

Maronpot et al. (1995b) assessed replicative DNA synthesis after 13, 26, 52, and 78 weeks of inhalation exposure by female B6C3F<sub>1</sub> mice to 2,000 ppm of methylene chloride for 6 hours/day, 5 days/week. A statistically significant decrease in the hepatocyte LI was only observed at 13 weeks. In lung epithelial cells, the results were similar to those observed by Kanno et al. (1993). No increases in replicative DNA synthesis were found in liver foci cells or lung parenchymal cells. *K-ras* gene activation in liver tumors and *H-ras* gene activation in lung tumors did not differ between methylene chloride-induced tumors and those observed in control animals. The authors concluded that these oncogenes were not involved in mouse tumorigenesis.

DNA-protein crosslinks (DPX) in lung and liver were examined by Casanova et al. (1992, 1996) in male B6C3F<sub>1</sub> mice exposed to 2,000 and 4,000 ppm of methylene chloride for 6 hours/day for 2 days and in male Syrian Golden hamsters exposed to 3,500 ppm. The authors suggested that formaldehyde derived from GSH-mediated methylene chloride metabolism might be forming DPX in mouse liver. Although DPX were detected in mouse liver, there was no evidence of DPX formation in mouse lung, hamster liver, or hamster lung. Additionally, DPX are not formed in measurable amounts in human liver tissue (Casanova et al. 1996). Therefore, the induction of DPX by formaldehyde in mouse liver might be a species-specific, tissue-specific response. A subsequent *in vitro* study by this laboratory confirmed the absence of DPX formation in response to methylene chloride in human, rat, and hamster hepatocytes, whereas a dose-response was observed in mouse hepatocytes (Casanova et al. 1997). In a different experiment, a dose-response in the formation of RNA-formaldehyde adducts was observed in hepatocytes of all four species (mouse>human>rat>hamster). RNA adduct production was related to the expression of the GSTT1-1 enzyme; the human liver sample lacking GSTT1-1 did not produce RNA adducts.

*In vitro* tests using two different strains of *Salmonella*, one with and one without GST expression, revealed that methylene chloride may be mutagenic by at least two pathways (DeMarini et al. 1997). The bacterial strains employed were TA100 and RSJ100; the latter is a derivative of the strain TA1535 (not normally mutagenized by methylene chloride) that contains recombinant rat GSTT1-1. In RSJ100, methylene chloride was mutagenic at moderate doses, and produced a single class of mutation (GC ÷ AT transversions) only at the middle C of the target CCC. Although methylene chloride was mutagenic in TA100, it required a much higher dose to match the mutation frequency in RSJ100. Furthermore, it produced a variety of lesions and mutations (predominantly GC ÷ TA transversions) at the first and second positions of the CCC target. The implication of these results is that genotoxic effects of methylene chloride will be different, depending on the GSTT1 phenotype. Those who lack the functional

## 2. HEALTH EFFECTS

gene would only be susceptible to genetic damage from methylene chloride under high exposure levels, whereas carriers might be vulnerable to genetic damage at much lower exposure levels.

According to Maronpot et al. (1995b), the precise mode of action of methylene chloride-induced mouse tumorigenicity appears to be elusive and has not yet been confirmed.

### 2.4.3 Animal-to-Human Extrapolations

The two major difficulties in applying the results of rodent cancer bioassays to humans involve extrapolation from the high-concentration rodent exposures to the lower concentrations typical of human exposure conditions; and the interspecies extrapolation. The predominant tumors of interest with regard to methylene chloride-induced tumorigenicity are mouse lung and mouse liver. However, species differences among rodent tumorigenic, genotoxic, and morphological responses to methylene chloride, as well as differences between mice and humans, appear to limit the applicability of mouse tumor data to humans, according to some authors. Foster et al. (1992) have suggested that the Clara cell may have a role in mouse lung tumor induction; there is a substantially higher number of Clara cells in the mouse than in other rodent species or in humans. GSH-mediated activation of the proximate carcinogenic agent of methylene chloride in mice has been associated with a specific isoenzyme, the  $\Theta$  (theta) class GSTs; Sheratt et al. (1997) have shown that this isozyme is expressed at very low levels in human pulmonary cells *in vitro*, suggesting that in humans, the lung has little capacity to activate methylene chloride into biologically reactive intermediates. *In vitro* studies (e.g., Graves et al. 1995) have shown species differences in the ability to induce DNA single-strand breaks in mouse, rat, hamster, and human cells that are compatible with the known rodent carcinogenicity, or lack thereof, in chronic cancer bioassays. These *in vitro* studies suggested that humans are unlikely to be more susceptible than rodents to methylene chloride-induced liver cancer.

In a comparison of rates of methylene chloride metabolism by each of the two major pathways in four species, the mouse, rat, hamster, and human, Green (1997) presented *in vivo* and *in vitro* evidence that the MFO pathway metabolic rates are similar among all four species and saturated at concentrations of 500 ppm or above. In contrast, the GSH-mediated metabolism is linear over the concentration range studied; it is also the major metabolic pathway for methylene chloride in mice at concentration levels used in cancer inhalation bioassays. Furthermore, the activity in mouse tissues is more than an order of magnitude greater than the activity in rat tissues. Hamster and human tissues show metabolic rates for this pathway that are even lower than those found in the rat. These findings are corroborated by the localization study of Mainwaring et al. (1996b), which found higher levels of GSTT1-1 mRNA and protein in the liver and lung of mice than in rats or humans. In mechanistic terms, Green (1997)

## 2. HEALTH EFFECTS

concludes that these studies demonstrate that differences in glutathione-mediated metabolism among mice, rats, hamsters, and humans are correlated with differences in carcinogenicity of methylene chloride. This is supported by the levels of glutathione that have been detected in the livers of the different species. The hepatic concentrations of GSH (in nmol/10<sup>6</sup> cells) were approximately 129 in B6C3F<sub>1</sub> mice (Ruch et al. 1989), 50 in Sprague-Dawley rats (Jones et al. 1978), 22 in Syrian Golden hamsters, and 21 in humans (Steinmetz et al. 1988). Thus, the mouse is the most sensitive species to metabolic activation of methylene chloride by glutathione metabolism, whereas humans appear to be the least sensitive.

However, other evidence suggests that methylene chloride may be potentially carcinogenic in humans. Mainwaring et al. (1996b) analysis of the distribution of mRNA and protein for GSTT1-1 and GSTT2-2 in the analysis of the liver and lungs of mice, rats, and humans corroborated the finding of Sheratt et al. (1997) that the overall levels in human tissues are much lower than in those of mice. However, the immunodetection of localized high concentrations of GSTT2-2 enzyme in human bile-duct epithelial cells is potentially significant, considering the increased incidence of biliary cancer following chronic exposure to methylene chloride as reported by Lanes et al. (1990). On the other hand, the GSTT2-2 antibody did not localize to the nucleus of human biliary epithelial cells, which would tend to reduce the potential genotoxic effect in humans. Although Mainwaring et al. (1996b) found that rates of metabolism of methylene chloride were very low in human lung, they also detected higher than background amounts of GSTT1-1 mRNA in a few Clara cells and ciliated cells of the alveolar/ bronchiolar junction of the lung in one human sample out of four. Therefore, despite the general low level of GSTT1-1 and GSTT2-2 in human tissue, it is possible that in some individuals, specific cell types within the human liver (bile duct) and lung might produce genotoxic reactive intermediates as a result of methylene chloride metabolism. Based on GSTT enzyme distributions and concentrations, the carcinogenic risk from methylene chloride in humans appears to be low as in rats rather than high as in mice.

The mouse model has been employed in assessment of excess lifetime cancer risks in humans from inhalation exposure to methylene chloride by EPA and others. Recent data, both *in vitro* and *in vivo*,



## 2. HEALTH EFFECTS

strongly indicate that the higher sensitivity of the mouse, relative to humans, must be taken into consideration if mouse data are to be used to estimate potential human health risks.

## 2.5 RELEVANCE TO PUBLIC HEALTH

Issues relevant to children are explicitly discussed in 2.7, Children's Susceptibility, and 5.6, Exposures of Children.

**Overview.** Methylene chloride has been widely used in industrial processes, food preparation, agriculture, and consumer products; consequently, there have been numerous studies describing its effects in a variety of experimental animal species. Humans have not been clinically studied as extensively. Although methylene chloride uses in agricultural goods and some consumer products have declined in recent years, there is still potential public health concern due to its continued use in industrial processes, and continued releases into the environment.

The central nervous system is affected adversely in humans and animals at inhalation exposure levels of 200 ppm or higher (Putz et al. 1979). Effects in animals were also reported on the liver and kidney following continuous exposure at concentrations of 25 ppm or greater (Haun et al. 1972), and on the cardiovascular system, but at extremely high exposures (Aviado and Belej 1974). Long-term inhalation exposure to methylene chloride (500 ppm or greater) increased tumors in some animals (Nitschke et al. 1988a), but did not cause teratogenic or reproductive effects in a two-generation study (Nitschke et al. 1988b). Since inhalation is the principal route of exposure to methylene chloride, most of these effects have been tested for or observed by this route. Data on effects observed after oral and dermal exposure are somewhat more limited. Further details are presented below.

### Minimal Risk Levels for Methylene Chloride.

#### *Inhalation MRLs.*

- C An MRL of 0.6 ppm has been derived for acute inhalation exposure (0–14 days) to methylene chloride. *This MRL supersedes the previous MRL of 3 ppm derived in the 1998 draft for public comment profile. Refer to chapter 7 for additional information.*

## 2. HEALTH EFFECTS

The acute inhalation MRL was derived from the behavioral toxicity study by Winneke (1974) in which a randomized blind clinical chamber experiment was used to expose 6–20 volunteers to vapors of methylene chloride (300, 500, or 800 ppm) or filtered air for 3–4 hours. Subjects were tested at 45-minute intervals with standard neurobehavioral tests measuring: (1) critical flicker fusion frequency (visual); (2) auditory vigilance performance; and (3) performance on psychomotor tasks. The tested parameters were considered to reflect the status of ‘cortical alertness’ (Fodor and Winneke 1971). A statistically significant depression in critical flicker fusion (CFF) frequency was observed at all concentrations. The magnitude of CFF frequency depression was similar at exposure concentrations of 300 and 500 ppm and was larger at 800 ppm. Thus, there was no dose-response at the two lowest concentrations, and a dose-response was evident at the highest concentration. A decrease in auditory vigilance performance was observed at 500 ppm and psychomotor task performance was impaired at 800 ppm. Thus, reduced CFF frequency is the most sensitive neurological response to acute inhalation exposure to methylene chloride. Based on this endpoint, the LOAEL is 300 ppm. A PBPK model for this experiment was used to adjust the dosage to a 24 hour exposure period, thus resulting in a LOAEL of 60 ppm for the same endpoint (Reitz et al. 1997). The MRL was derived by dividing the LOAEL of 60 ppm by an uncertainty factor of 100 (10 for the use of a LOAEL and 10 for human variability).

- C An MRL of 0.3 ppm has been derived for intermediate inhalation exposure (15–364 days) to methylene chloride.

The intermediate inhalation MRL was derived from a study by Haun et al. (1972) in which groups of mice, rats, dogs, and monkeys were continuously exposed to methylene chloride for 14 weeks at chamber concentrations of either 0, 25, or 100 ppm. Body weights and clinical signs were monitored throughout the study. Necropsy was performed and tissues were examined histopathologically and organ-to-body weight were determined at the end of the exposure. Data on liver histopathology of rats exposed to 25 or 100 ppm of methylene chloride were selected as the critical effect. Liver cytoplasmic vacuolization and staining associated with fatty deposits were observed in rats at both exposure concentrations; Haun et al. (1972) did not mention whether there were quantitative differences in the effect observed at the two exposure levels. The MRL was derived based on a LOAEL of 25 ppm for hepatic effects. Because the critical effect observed is an extrarspiratory effect (rat liver), a human equivalent concentration (HEC) was calculated. Since the ratio of the blood:air partition coefficient in the rat to the blood:air partition coefficient in the human was  $> 1$ , the value of 1.0 was used to calculate the  $LOAEL_{[HEC]}$  (EPA 1994). This resulted in an MRL of 0.3 ppm by dividing the LOAEL of 25 ppm by an uncertainty factor of 90 (3 for use of a minimal LOAEL, 3 for extrapolation from animals to humans, and 10 for human variability).

## 2. HEALTH EFFECTS

CAn MRL of 0.3 ppm has been derived for chronic inhalation exposure (365 days) to methylene chloride.

The chronic inhalation MRL was derived from a study by Nitschke et al. (1988a) in which groups of 90 male and 108 female Sprague-Dawley rats were exposed to methylene chloride at 0 (controls), 50, 200, or 500 ppm for 6 hours/day, 5 days/week for 2 years. A number of satellite groups were also exposed to assess the temporal relationship between methylene chloride exposure and evidence of toxicity. Subgroups of females in the main study were sacrificed after 6, 12, 15, and 18 months of exposure. The following end points were evaluated: body weight, food consumption rates, organ weights, hematology, clinical chemistry, urinalysis, pathology, histopathology, and blood COHb levels. Blood COHb levels were consistently higher than 10% in animals exposed to 200 ppm. No pathologic or histopathologic nontumor findings were reported except in the liver. Hepatocellular cytoplasmic vacuolization consistent with fatty changes, and multinucleate hepatocytes were elevated in female rats exposed to methylene chloride at 200 and 500 ppm; a slight increase in the incidence of hepatocellular vacuolization was also observed in male rats exposed to 500 ppm.

The NOAEL of 50 ppm was adjusted for continuous exposure (6 hour/day, 5 day/week) resulting in a NOAEL<sub>[ADJ]</sub> of 8.92 ppm. Whereas the MRL was derived based on hepatic effects (extrarespiratory), a human equivalent concentration (HEC) was calculated. Since the ratio of the blood:air partition coefficient in the rat to the blood:air partition coefficient in the human was > 1, the value of 1.0 was used to calculate the LOAEL<sub>[HEC]</sub> (EPA 1994). This resulted in an MRL of 0.3 ppm by dividing the LOAEL of 8.92 ppm by an uncertainty factor of 30 (3 for extrapolation from animals to humans, and 10 for human variability).

**Oral MRLs.**

CAn MRL of 0.2 mg/kg/day has been derived for acute oral exposure (0–14 days) to methylene chloride. *This MRL supersedes the previous MRL of 0.5 mg/kg/day derived in the 1998 draft for public comment profile. Refer to chapter 7 for additional information.*

The acute oral MRL was derived by route-to-route extrapolation of the data from Winneke (1974) in which a randomized blind clinical chamber experiment was used to expose 6–12 volunteers to vapors of methylene chloride (300, 500, or 800 ppm) or filtered air for 3–4 hours. Subjects were tested at 45-minute intervals with standard neurobehavioral tests measuring: (1) critical flicker fusion frequency

## 2. HEALTH EFFECTS

(visual); (2) auditory vigilance performance; and (3) performance on psychomotor tasks. A statistically significant depression in critical flicker fusion (CFF) frequency was observed at all concentrations. The magnitude of CFF frequency depression was similar at exposure concentrations of 300 and 500 ppm and was larger at 800 ppm. Thus, there was no dose-response at the two lowest concentrations, and a dose-response was evident at the highest concentration. A decrease in auditory vigilance performance was observed at 500 ppm and psychomotor task performance was impaired at 800 ppm. Thus, reduced CFF frequency is the most sensitive neurological response to acute inhalation exposure to methylene chloride. Based on this end point, the LOAEL is 300 ppm.

Reitz et al. (1997) modified the basic PBPK model for methylene chloride that was developed by Andersen et al. (1987), Reitz et al. (1988), and Andersen et al. (1991) as described in Section 2.3.5.2. The major modification of the model was the inclusion of a brain compartment so that central nervous system effects could be assessed. Reitz et al. (1997) modeled the Winneke (1974) data to obtain the target organ (brain) concentrations of methylene chloride associated with administered inhalation concentrations, and then calculate the human drinking water concentrations (mg/L) that would result in the equivalent target organ-specific doses. Human exposure patterns in the PBPK model simulated realistic human drinking water consumption patterns, (i.e., consisting of bouts of drinking during the day, with and between meals, and little-to-no drinking during the night). PBPK modeling predicted that peak concentrations of methylene chloride in the brain would increase rapidly after each episode of drinking water consumption, and then drop sharply, to near-zero, between bouts of drinking. Additionally, there would be no cumulative effects from repeated exposure.

For acute neurological effects, the associated dose measure was defined as the peak concentration of methylene chloride in brain tissue (mg/L of brain tissue) of humans exposed to 300 ppm of methylene chloride for 4 hours by inhalation. The modified PBPK model calculated that the administered inhalation dose was equivalent to 3.95 mg of methylene chloride per L of brain tissue. The equivalent administered human concentration in drinking water that will produce the same neurological effects was 565 mg of methylene chloride/L. Using a daily drinking water consumption value of 2L and an average human body weight of 70 kg, the LOAEL was calculated to be 16 mg/kg/day. An acute oral MRL was calculated by dividing the LOAEL (16 mg/kg/day) by an uncertainty factor of 100 (10 for the use of a LOAEL and 10 for human variability), to yield 0.2 mg/kg/day.

## 2. HEALTH EFFECTS

CAn MRL of 0.06 mg/kg/day has been derived for chronic oral exposure (365 days) to methylene chloride. *This MRL supersedes the previous MRL of 0.2 mg/kg/day derived in the 1998 draft for public comment profile. Refer to chapter 7 for additional information.*

The chronic oral MRL was derived from a study by Serota et al. (1986a) in which F344 rats (85/sex/dose) were exposed to methylene chloride in deionized drinking water at concentrations to provide target doses of 0, 5, 50, 125, or 250 mg/kg/day for 104 weeks. The nominal mean doses were 0, 6, 55, 131, and 249 mg/kg/day. There were no treatment-related effects on survival or on the incidence of adverse clinical signs. Organ weights were not significantly affected by treatment. Histopathology was only observed in the liver, which therefore is the critical target organ. Statistically significant cellular changes (hepatic foci/areas of cellular alterations and fatty changes) were observed at dose levels  $\geq$  50 mg/kg/day. Reduced weight gain was observed at 131 and 249 mg/kg/day, but quantitative data were not provided. Hematological effects (increased mean hematocrit, hemoglobin, and erythrocyte count) were observed at all dose levels except 6 mg/kg/day. Therefore, the lowest dose in rats was identified as the NOAEL. Based on measured, mean drinking water consumption rates, this dose was calculated to be 6 mg/kg/day. The chronic oral MRL was calculated by dividing the NOAEL (6 mg/kg/day) by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability).

No intermediate oral MRL was derived because of an inadequate database.

**Death.** Acute inhalation exposure to methylene chloride has caused death in humans (Bakinson and Jones 1985; Bonventre et al. 1977; Hall and Rumack 1990; Stewart and Hake 1976). Although exposure levels were not measured, estimates suggest that a combination of high exposure levels and/or inadequate ventilation has contributed to these lethal accidents. Measurements of methylene chloride in tissues or of COHb in blood have corroborated high exposures in some cases (Manno et al. 1992; Moskowitz and Shapiro 1952; Winek et al. 1981; Tay et al. 1995). The biologic cause of death was not verified in all cases, but is thought to have been respiratory depression secondary to narcosis. Asphyxia accompanied by bilateral pulmonary congestion and focal hemorrhage was reported in one case (Winek et al. 1981) and myocardial infarction was reported in another (Stewart and Hake 1976). Mortality risk was not increased in humans exposed occupationally to 30–120 ppm of methylene chloride for over 30 years (Friedlander et al. 1978) and no excess mortality was found in workers exposed to 140–475 ppm for at least 3 months (Lanes et al. 1993; Ott et al. 1983b). Inhalation exposure to concentrations of 3,500 ppm for longer

## 2. HEALTH EFFECTS

durations (14 weeks to 2 years) was lethal in some animals (Burek et al. 1984; MacEwen et al. 1972). No mortality increase was noted in chronic inhalation studies at 500 ppm in the rat (Nitschke et al. 1988a).

In one suicide case, ingestion of paint remover containing 75–80% methylene chloride resulted in death from corrosion of the gastrointestinal tract (Hughes and Tracey 1993). In animals, acute exposure to high doses (2,100 mg/kg or greater) by gavage caused death (Kimura et al. 1971; Ugazio et al. 1973); intermediate-duration exposures to 64 or 320 mg/kg/day by gavage significantly increased mortality in female mice and male rats, respectively (Maltoni et al. 1988). However, exposure to methylene chloride in drinking water (up to 250 mg/kg/day) did not significantly affect survival (Serota et al. 1986a, 1986b). Gavage administration may result in more severe effects than *ad libitum* water ingestion since the administered dose may temporarily saturate normal metabolic processes, and result in a different metabolic profile. No data were found on death in humans or animals from dermal exposure.

### Systemic Effects

**Respiratory Effects.** In one workplace accident, acute inhalation exposure to methylene chloride (probably at a high concentration) resulted in bilateral pulmonary congestion with focal hemorrhage (Winek et al. 1981). Less severe respiratory symptoms (cough, breathlessness, chest tightness) were reported in occupational exposure incidents (concentrations unknown; Bakinson and Jones 1985). Exposure to 18–1,200 mg/m<sup>3</sup> (5–340 ppm; 8-hour TWA) resulted in irritation of the respiratory tract in one occupational study (Anundi et al. 1993). However, a study in humans found no effect on pulmonary function following repeated exposures to methylene chloride vapors (up to 500 ppm) (Stewart et al. 1972). Studies in animals corroborated the severe pulmonary effects (congestion, edema, inflammation) of acute or intermediate exposures at high concentrations (Heppel et al. 1944; NTP 1986). Resolution of acute Clara cell damage in mice exposed to 4,000 ppm of methylene chloride was correlated with cytochrome P-450 activity in the lung (Foster et al. 1992). No information was found on the respiratory effects of low levels of methylene chloride in humans near hazardous waste sites or industrial urban areas or in animals.

**Cardiovascular Effects.** Myocardial infarction occurred in one case of acute inhalation occupational exposure (Stewart and Hake 1976). However, occupational studies in humans did not find any association between exposure to methylene chloride at 75–475 ppm and cardiac abnormalities (Cherry et al. 1981) or excess mortality due to ischemic heart disease (Hearne et al. 1990; Ott et al. 1983b). Further,

## 2. HEALTH EFFECTS

a cross-sectional study of workers showed no excess of electrocardiographic abnormalities among those exposed to methylene chloride (Ott et al. 1983c). Another study in humans did not reveal effects on cardiovascular functions at concentrations up to 500 ppm (NIOSH 1974). These findings suggest the cardiovascular system is not a sensitive target for exposure to methylene chloride in humans. One study in mice reported atrioventricular block following acute inhalation exposure to very high levels of methylene chloride (>200,000 ppm) (Aviado and Belej 1974). The relevance of this finding is limited since the exposure level was so high.

In an experiment with rats, using an ischemia-reperfusion model, intravenous methylene chloride infusion (leading to blood concentrations of <0.1 mg/mL) markedly increased the atrioventricular block during the reperfusion phase (Scholz et al. 1991). From these results, the authors concluded that the initial coma resulting from methylene chloride-induced poisoning is likely to result not only from anesthetic effects, but also from sudden onset of cardiac arrhythmias.

***Gastrointestinal Effects.*** Nausea and vomiting were reported in 13 out of 33 occupational cases of acute inhalation exposure to methylene chloride in the United Kingdom (Bakinson and Jones 1985); exposure levels were not reported in these cases. In mice exposed to 4,000 ppm of methylene chloride for 2 years, dilatation of the stomach was reported (NTP 1986). In humans attempting suicide, ingestion of a single oral dose of Nitromors, a paint remover solvent containing 75–80% methylene chloride, resulted in severe corrosion and ulceration of the gastrointestinal tract (Hughes and Tracey 1993), peritonitis and septicemia occurred in the fatal case, whereas intestinal diverticuli developed during recovery in the another case (Roberts and Marshall 1976). No other studies were located regarding gastrointestinal effects in humans or animals after exposure to methylene chloride.

***Hematological Effects.*** Metabolism of methylene chloride results in excess carbon monoxide and increases in COHb, which contributes to hypoxia (Tomaszewski 1998). Blood COHb concentrations were about 30% higher than normal in 2 lethal cases in which workers were estimated to be exposed to extremely high concentrations (up to 168,000 ppm) of methylene chloride in a confined work space (Manno et al. 1992). Several hours after an adult woman ingested a fatal oral dose of Nitromors, a paint remover containing 75–80% methylene chloride, her COHb level was 9%. In all three fatal exposures, cause of death was *not* associated with elevated COHb. In autoworkers who were exposed to methylene chloride dermally and by inhalation (3–154 ppm), blood COHb measurements taken within 24 hours of exposure were 1.2–11% for nonsmokers and 7.3 and 17.3% for two smokers (Kelly 1988). One-day

## 2. HEALTH EFFECTS

occupational exposures to methylene chloride at levels below ACGIH standard (50 ppm, 8-hour TWA) produced small increases in COHb levels in both nonsmoking and smoking adults (Soden et al. 1996); additional daily cumulative exposure to methylene chloride did not further increase COHb levels. Increases in blood COHb levels between 5 and 6.8% were measured in nonsmoking volunteers exposed to methylene chloride at concentrations up to 200 ppm for 4 to 7.5 hours (DiVincenzo and Kaplan 1981; Putz et al. 1979).

Other studies have reported increases in red cell count, hemoglobin, and hematocrit in women, but not in men, occupationally exposed to concentrations of up to 475 ppm during an 8-hour workday (Ott et al. 1983d); these effects were judged by the authors to be suggestive of compensatory hematopoietic effects. Similar findings were not observed in rodents chronically exposed to methylene chloride by inhalation at concentrations up to 3,500 ppm (Burek et al. 1984), but were observed in rodents exposed orally for 3 months at 480 mg/kg/day (Kirschman et al. 1986), or for 2 years at 55–249 mg/kg/day (Serota et al. 1986a, 1986b). Intravascular hemolysis was reported in the case of a man who attempted suicide by ingesting the paint remover, Nitromors (Roberts and Marshall 1976) and in a high-dose acute gavage study in rats (Marzotko and Pankow 1987).

**Hepatic Effects.** Human data are limited on the effects of methylene chloride on the liver. A slight exposure-related increase in serum bilirubin (but not at levels of clinical significance) was observed in workers with exposure up to an average of 475 ppm of methylene chloride, but serum levels of hepatic enzymes (e.g., aspartate aminotransferase, alanine aminotransferase, lactate dehydrogenase, and alkaline phosphatase) were not elevated (Ott et al. 1983a); another occupational study found no exposure-related changes in hepatic enzymes (Anundi et al. 1993). In another study, methylene chloride vapors (up to 500 ppm) did not affect comparable serum enzyme activity in volunteers (Stewart et al. 1972). However, the liver appears to be a major target organ following methylene chloride exposure in animals, particularly at high exposure levels (>5,000 ppm; Heppel et al. 1944). Histomorphological and biochemical changes of the liver occur following acute inhalation (6 hours to 7 days) at high concentration levels (5,200 ppm) (Morris et al. 1979), while fatty changes and biochemical alterations (altered cytochrome P-450 levels) were also observed at lower concentrations (100 ppm) for continuous, 24-hour intermediate-duration exposure (100 days) (Haun et al. 1972; Kjellstrand et al. 1986; Weinstein and Diamond 1972). Cytoplasmic vacuolization was observed in rats at 25 ppm (Haun et al. 1972). Using these data, ATSDR derived an intermediate inhalation MRL of 0.3 ppm, as calculated in Table 2-1. Exposure to 1,000–4,000 ppm for 2 years resulted in an increased incidence of hemosiderosis and focal



## 2. HEALTH EFFECTS

hepatic necrosis in rats (NTP 1986). An increased incidence of fatty changes occurred following chronic exposure at 500 ppm, but not at 200 ppm (Nitschke et al. 1988a); fatty changes were reversible when exposure ceased. Using data from the Nitschke et al. (1988a) study, ATSDR derived a chronic inhalation MRL of 0.3 ppm, as shown in Table 2-1. Necrosis or fatty changes in the liver have been observed in rats given high oral doses (>1,000 mg/kg/day) of methylene chloride (Kirschmann et al. 1986; Ugazio et al. 1973). Chronic ingestion of methylene chloride in drinking water has been associated with fatty changes in rats at 55 mg/kg/day or greater and in mice at 175 mg/kg/day or greater, but not at 6 mg/kg/day (Serota et al. 1986a, 1986b). Based on this value (6 mg/kg/day), a chronic oral MRL of 0.06 mg/kg/day was calculated as described in Table 2-2.

**Endocrine Effects.** No relevant information was located regarding endocrine effects in human or animals associated with exposure to methylene chloride.

**Renal Effects.** Kidney function was not altered in humans repeatedly exposed to methylene chloride vapors (up to 500 ppm) for 6 weeks (Stewart et al. 1972) and no alterations in urinary microglobulins or N-acetyl-beta-glucosaminidase were detected in workers chronically exposed to methylene chloride (Anundi et al. 1993). In rats, nonspecific renal tubular and degenerative changes occurred after continuous intermediate-duration exposure to methylene chloride vapors (100–5,000 ppm) (Haun et al. 1972; MacEwen et al. 1972) or chronic exposure to 4,000 ppm (NTP 1986). Similar renal changes occurred in dogs exposed to 1,000 ppm for 4 weeks (MacEwen et al. 1972). There were no studies reporting renal effects following oral exposure to methylene chloride in humans. In rat oral studies using methylene chloride at doses >1,300 mg/kg/day, a single dose inhibited diuresis (Marzotko and Pankow 1987) and treatment for 3 months increased kidney weights in females (Kirschman et al. 1986). At 166 mg/kg/day, methylene chloride lowered the pH of urine in rats of both sexes (Kirschman et al. 1986). There are no data that would enable an assessment of the potential for renal effects in humans living near hazardous waste sites.

**Dermal Effects.** No studies were located regarding dermal effects in human or animals associated with inhalation or oral exposure to methylene chloride. However, in some occupational accidents, direct contact with methylene chloride has resulted in second or third degree chemical burns within 30 minutes (Hall and Rumack 1990; Wells and Waldron 1984; Winek et al. 1981).

## 2. HEALTH EFFECTS

**Ocular Effects.** No studies were located on ocular effects in humans by oral or dermal exposure. However, repeated direct exposure to methylene chloride vapors (up to 500 ppm) caused mild irritation to the eyes of volunteers (Stewart et al. 1972). In one occupational accident, direct contact with the liquid (duration unspecified) resulted in severe corneal burns (Hall and Rumack 1990). In animals, small increases in corneal thickness and intraocular tension were reported after exposure to vapors of 490 ppm of methylene chloride or greater, but effects were reversible within 2 days after exposure ceased (Ballantyne et al. 1976). Inflammation of the conjunctivas and eyelids as well as increases in corneal thickness and intraocular tension were observed following direct contact of methylene chloride (0.1 mL) with the eyes of rabbits. Effects were reversible within 3–9 days (Ballantyne et al. 1976).

**Immunological and Lymphoreticular Effects.** No studies were located regarding immunological effects in humans after inhalation, oral, or dermal exposure. Splenic atrophy was evident in dogs that died following continuous intermediate-duration exposure to vapors of methylene chloride (1,000 ppm) (MacEwen et al. 1972), but the incidence did not increase over control levels in rats and mice chronically, but discontinuously, exposed at a level of 4,000 ppm or less (NTP 1986). Splenic fibrosis was observed in rats after chronic inhalation exposure at 1,000 ppm methylene chloride (Mennear et al. 1988), but was not observed in a recent intermediate-duration study in rats exposed to 5,187 ppm methylene chloride (Halogenated Solvent Industry Alliance, Inc. (2000)). This latter study also found that the IgM antibody response to SRBC was not significantly altered in exposed rats relative to controls. Due to the very limited and inconsistent nature of the database, further studies are necessary before conclusions can be drawn about the relevance of these findings to human health.

**Neurological Effects.** Studies in humans and animals indicate the central nervous system is an important target for methylene chloride; in the case of acute exposures, anesthetic responses, which subsided once exposure ceased, have been reported in humans and animals (Bakinson and Jones 1985; Hall and Rumack 1990; Heppel et al. 1944; Snyder et al. 1992a, 1992b). Neurological effects in humans have included headache, dizziness, confusion, memory loss, intoxication, incoordination, paresthesia, and in severe cases, unconsciousness and seizures (Bakinson and Jones 1985; Hall and Rumack 1990; Kelly 1988). Degraded performance in various psychomotor tasks was reported in humans acutely exposed to methylene chloride (200 ppm or greater) in experimental studies (Fodor and Winneke 1971; Putz et al. 1979; Stewart et al. 1972; Winneke 1974). Impaired performance during exposure to methylene chloride was associated with a rise in blood levels COHb to about 5%, which occurred after 3 hours of exposure at 200 ppm (Putz et al. 1979). In more specific measures of cortical function, Winneke (1974) attributed

## 2. HEALTH EFFECTS

impaired performance in volunteers following 3–4 hours of exposure to 300 ppm of methylene chloride to the properties of the parent compound; exposure to 50–100 ppm of carbon monoxide did not generate the same adverse effects. Based on this value (300 ppm), an acute inhalation MRL of 0.6 ppm was calculated as described in Table 2-1 (Winneke 1974). Studies in factory workers chronically exposed to methylene chloride revealed no evidence of neurological or behavioral impairment at exposure levels of 75–100 ppm (Cherry et al. 1981). There was a reduction in test scores pertaining to mood changes in workers in 1 of 3 rapid rotation shifts exposed to vapors of methylene chloride (28–173 ppm), but no effects were observed on performance as determined by digit symbol substitution scores (Cherry et al. 1983). Mood changes are very subjective and these results may reflect other causes.

Neurological effects have been noted in animals following exposure to methylene chloride independent of the route. Narcotic effects of methylene chloride (incoordination, gait disturbances, reduced activity, somnolence) were observed in monkeys, rabbits, rats and guinea pigs following acute exposure at 10,000 ppm (Heppel and Neal 1944; Heppel et al. 1944); dogs exposed at this level became uncoordinated, then excited and hyperactive (Heppel et al. 1944). These effects wore off when exposure ceased. Studies in rats and gerbils suggest that methylene chloride induces regional alterations in DNA concentrations, amino acids levels, and enzyme activities in the brain. There was a decrease in succinic dehydrogenase activity in the cerebellum in rats exposed to vapors of methylene chloride at concentrations of 500 ppm or greater and signs of increased protein breakdown in the cerebrum at 1,000 ppm (Savolainen et al. 1981). The DNA concentration decreased in the hippocampus and cerebellum in gerbils exposed to 210 ppm of methylene chloride, indicating decreased cell density in these brain regions, probably due to cell loss (Karlsson et al. 1987; Rosengren et al. 1986). Levels of aminobutyric acid increased in the posterior cerebellar vermis of gerbils exposed to 210 ppm of methylene chloride; however, the significance of this finding is uncertain (Briving et al. 1986). On the other hand, studies in rats exposed to methylene chloride (2,000 ppm) for 13 weeks revealed that the compound did not cause clinical, postural, sensory, locomotor, evoked potential, or pathological effects (Mattsson et al. 1990). However, when rats were given an intraperitoneal injection of methylene chloride, 115 mg/kg was sufficient to alter the pattern of flash evoked potentials (Herr and Boyes 1997); in this study, the responses generated in methylene chloride-treated rats were different from those generated by other solvents, leading the authors to conclude that lipid solubility alone was insufficient to predict the neurotoxic effects. The mechanism by which methylene chloride exerts its effects on the central nervous system is not clear. Anaesthetic effects of the parent compound and hypoxic effects of its metabolite carbon monoxide may both contribute to the observed symptoms.

## 2. HEALTH EFFECTS

**Reproductive Effects.** Data on reproductive toxicity in humans are limited; there are no studies involving oral exposure. One group of cases reported genital pain, testicular atrophy, recent infertility, and low/abnormal sperm counts in workers who inhaled vapors of methylene chloride and who had direct contact with the liquid on the job for more than a year (Kelly 1988). Exposure to methylene chloride in this group was confirmed by blood COHb analysis and the presence of typical neurological deficits. A retrospective study of pregnancy outcomes among Finnish pharmaceutical workers reported a slightly increased risk of spontaneous abortion (not quite statistically significant) associated with inhalation exposure to methylene chloride (Taskinen et al. 1986). No animal reproductive studies were conducted for oral or dermal exposures to methylene chloride, and the inhalation studies were negative. Methylene chloride did not adversely affect fertility and litter size in rats that inhaled methylene chloride vapors at concentrations up to 1,500 ppm or less for two generations (Nitschke et al. 1988b). After intermediate-duration exposure (6 weeks) to vapors of methylene chloride (200 ppm or less), there were no microscopic lesions in testes of rats (Raje et al. 1988). Uterine, ovarian, and testicular atrophy has been noted in rats and mice chronically exposed for 2 years to 4,000 ppm of methylene chloride, but this was reported to be secondary to malignant neoplasms (NTP 1986). Based on these data, methylene chloride does not appear to pose a hazard to human reproduction, except at very high exposure levels.

**Developmental Effects.** There are few studies addressing developmental effects in humans. A retrospective study of pregnancy outcomes among Finnish pharmaceutical workers reported a slightly increased risk of spontaneous abortion associated with inhalation exposure to methylene chloride (Taskinen et al. 1986). A study examining the effect of low environmental concentrations of methylene chloride (<0.01 ppm) on over 90,000 births found no significant effect on birth weight (Bell et al. 1991). Studies in rats demonstrated that methylene chloride crosses the placenta and that metabolism of methylene chloride by the maternal liver elevates the blood levels of carbon monoxide in the fetus (Anders and Sunram 1982). Animal studies demonstrated that inhalation of methylene chloride vapors at concentrations of 1,250 ppm produced minor skeletal variants: delayed ossification of sternbrae in rats and extra center of ossification in the sternum of mice and rats (Schwetz et al. 1975). Fetal weight was reduced and behavioral changes occurred in rat pups following exposure of dams to 4,500 ppm of methylene chloride (Bornschein et al. 1980; Hardin and Manson 1980). The significance of these observations is uncertain since each of the three studies used only one concentration level and the observed effects occurred at maternally toxic concentrations. Growth (yolk sac blood vessel growth, body length, protein concentration) and development (somite addition) were significantly retarded in rat embryos cultured in the presence of methylene chloride at 0.5 mg/mL, but not at 0.2 mg/mL.

## 2. HEALTH EFFECTS

(Brown-Woodman et al. 1998); the effect on the yolk sac blood vessels may have interfered with the ability of the embryos to take up nutrients. Methylene chloride had no effect on heart function in these embryos. Although fetal body weights were decreased in these animal studies, the absence of other fetotoxic effects, embryo lethality, or major malformations, suggests that methylene chloride is not likely to cause developmental effects or behavioral changes at levels normally encountered in the environment; with current standards and procedures, workplace exposure of pregnant women is unlikely to be hazardous to the fetus. The reduction in the rate of spontaneous abortions observed during the later years of the Finnish study was attributed partly to improved industrial hygiene (Taskinen et al. 1986).

**Genotoxic Effects.** No studies were located regarding the genotoxic effects of methylene chloride in humans after inhalation, oral, or dermal exposure. *In vitro* results were mixed in bacterial assays and in tests employing mammalian cells (Table 2-3). Methylene chloride has caused chromosomal aberrations in some studies, but not in others. Given the evidence of *in vitro* clastogenicity and its negative results in unscheduled DNA-synthesis and DNA-binding studies, methylene chloride may be a weak mutagen in mammalian systems. The chemical has been evaluated in several *in vivo* assay systems in animals to assess its potential to induce gene mutation and cause chromosomal aberrations or DNA damage and repair. Many studies were negative and some were positive (Table 2-4). Tissue-specific genetic damage in mice following exposure to methylene chloride, suggests that the ability of tissues to metabolize methylene chloride may determine its genotoxicity (Sasaki et al. 1998). Specific polymorphisms in the metabolizing enzymes are associated with genotoxicity. *In vitro*, low concentrations of methylene chloride were more mutagenic when a functional GSTT1-1 gene was present (DeMarini et al. 1997).

Human erythrocytes expressing the GSTT1-null phenotype had a higher incidence of sister chromatid exchange than normal following exposure to methylene chloride (Hallier et al. 1994). Similarly, expression of the DraI DD mutant of CYP2E1 was associated with an increase in bleomycin-induced single-strand breaks in human lymphocyte DNA (El-Zein et al. 1997a); possibly methylene chloride exposure would induce similar genetic damage in individuals with that genotype. Additional details are presented in Section 2.10, Populations that are Unusually Susceptible.

**Table 2-3. Genotoxicity of Methylene Chloride *In Vitro***

Species (test system)	End point	Results		Reference
		With activation	Without activation	
Mammalian cells:				
Peripheral lymphocytes	Chromosomal aberrations	+	0	Thilagar et al. 1984a
Mouse/lymphoma L5178Y	Chromosomal aberrations	+	+	Thilagar et al. 1984a
Chinese hamster	Chromosomal aberrations	+	+	Thilagar et al. 1984a
Human/primary fibroblasts	Unscheduled DNA synthesis	Not tested	–	Jongen et al. 1981
Chinese hamster (V79)	Unscheduled DNA synthesis	Not tested	–	Jongen et al. 1981
Chinese hamster (V79)	Sister chromatid exchanges	(+)	(+)	Jongen et al. 1981
Human/peripheral lymphocytes	Unscheduled DNA synthesis	–	–	Perocco and Prodi 1981
Prokaryotic organisms:				
<i>Salmonella typhimurium</i> (TA98, TA100)	Gene mutation	+	+	Gocke et al. 1981
<i>S. typhimurium</i> (TA1535, TA1538, TA1537)	Gene mutation	–	–	Gocke et al. 1981

DNA = deoxyribonucleic acid; + = positive result; – = negative result; (+) = weakly positive result

**Table 2-4. Genotoxicity of Methylene Chloride *In Vivo***

Species (test system)	End point	Results	Reference
Mammalian cells:			
Mouse (bone marrow and lung cells)	Chromosomal aberrations	+	Allen et al. 1990
Mouse (peripheral erythrocytes)	Micronuclei	+	Allen et al. 1990
Mouse (peripheral lymphocytes and lung cells)	Sister chromatid exchanges	+	Allen et al. 1990
Mouse (bone marrow cells)	Micronuclei	–	Sheldon et al. 1987
Mouse	Dominant lethality	–	Raje et al. 1988
Mouse (liver and lung)	DNA breakage	+	Sasaki et al. 1998
Mouse (stomach, urinary bladder, kidney, brain and bone marrow)	DNA breakage	–	Sasaki et al. 1998
Rat (liver)	DNA breakage	+	Kitchin and Brown 1989
Rat	Unscheduled DNA synthesis	–	Trueman and Ashby 1987
Mouse	Unscheduled DNA synthesis	–	Trueman and Ashby 1987
Rat (bone marrow cells)	Chromosomal aberrations	–	Burek et al. 1984
Mouse (bone marrow cells)	Chromosomal aberrations	–	Gocke et al. 1981
Rat (liver and lung cells)	DNA alkylation	–	Green et al. 1988
Mouse (liver and lung cells)	DNA alkylation	–	Green et al. 1988
Eukaryotic organisms:			
Insect:			
<i>Drosophila</i>	Gene mutation (sex-linked recessive lethal)	(+)	Gocke et al. 1981

+ = positive result; – = negative result; (+) = weakly positive result; DNA = deoxyribonucleic acid

## 2. HEALTH EFFECTS

**Cancer.** A significant increase in bile-duct cancer was observed within a cohort of workers who had been exposed to methylene chloride (at #1,700 ppm, 8-hour TWA) for up to 28 years (Lanes et al. 1990). Epidemiology studies have not revealed a causal relationship between deaths due to cancer and occupational exposure to methylene chloride at lower levels (475 ppm or less) (Friedlander et al. 1978; Hearne et al. 1987, 1990; Ott et al. 1983b). It should be noted that these latter studies had limited power to detect very small increases in cancer and are not sufficient to rule out a carcinogenic potential of methylene chloride. Studies in animals exposed via inhalation have demonstrated that methylene chloride can increase the incidence of naturally-occurring tumors. When administered by inhalation, methylene chloride (2,000 ppm or greater) increased the incidence of alveolar/bronchiolar neoplasms in mice of both sexes (NTP 1986). Concentrations of 500 ppm or greater of methylene chloride increased the incidence of benign mammary gland tumors per animal in females and male rats (Burek et al. 1984; Nitschke et al. 1988a; NTP 1986). The incidence of liver tumors increased over concurrent control levels in male mice and female rats administered methylene chloride (50–250 mg/kg/day) in drinking water; however, the incidence of lesions in treated groups were within the historical range of control values and showed no dose response (Serota et al. 1986a, 1986b). The results of recent toxicokinetics studies suggest that the parent compound and/or reactive metabolites produced by the GST pathway are the source of methylene chloride-induced tumor increases. Based on these findings, the EPA has ranked methylene chloride as a Group B2 carcinogen (probable human carcinogen). The EPA has calculated that an upper limit  $10^{-6}$  risk level corresponds to 0.0006 ppm and 0.001 mg/kg/day for inhalation and oral 70-year continuous exposures, respectively. OSHA (1997) determined that methylene chloride is a potential occupational carcinogen, based on rodent inhalation studies and PBPK modeling, and established an inhalation unit risk of  $3.62 \times 10^{-3} (\mu\text{g}/\text{m}^3)^{-1}$  for occupational exposures. The Department of Health and Human Services (NTP 1999) has determined that methylene chloride may reasonably be anticipated to be a human carcinogen. IARC (1987) has classified methylene chloride in Group 2B (possibly carcinogenic to humans). EPA (IRIS 1999) has determined that methylene chloride is a probable human carcinogen. Section 2.10 discusses polymorphisms in genes that metabolize methylene chloride that may be associated with an increased risk of cancer.



## 2. HEALTH EFFECTS

**2.6 ENDOCRINE DISRUPTION**

Recently, attention has focused on the potential hazardous effects of certain chemicals on the endocrine system because of the ability of these chemicals to mimic or block endogenous hormones, or otherwise interfere with the normal function of the endocrine system. Chemicals with this type of activity are most commonly referred to as endocrine disruptors. Some scientists believe that chemicals with the ability to disrupt the endocrine system are a potential threat to the health of humans, aquatic animals, and wildlife. Others believe that endocrine disrupting chemicals do not pose a significant health risk, particularly in light of the fact that hormone mimics exist in the natural environment. Examples of natural hormone mimics are the isoflavonoid phytoestrogens (Adlercreutz 1995; Livingston 1978; Mayr et al. 1992). These compounds are derived from plants and are similar in structure and action as endogenous estrogen. While there is some controversy over the public health significance of endocrine disrupting chemicals, it is agreed that the potential exists for these compounds to affect the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body that are responsible for the maintenance of homeostasis, reproduction, development, and/or behavior (EPA 1997). As a result, endocrine disruptors may play a role in the disruption of sexual function, immune suppression, and neurobehavioral function. Endocrine disruption is also thought to be involved in the induction of breast, testicular, and prostate cancers, as well as endometriosis (Berger 1994; Giwercman et al. 1993; Hoel et al. 1992).

There is no evidence suggesting that methylene chloride is an endocrine disruptor.

**2.7 CHILDREN'S SUSCEPTIBILITY**

This section discusses potential health effects from exposures during the period from conception to maturity at 18 years of age in humans, when all biological systems will have fully developed. Potential effects on offspring resulting from exposures of parental germ cells are considered, as well as any indirect effects on the fetus and neonate resulting from maternal exposure during gestation and lactation. Relevant animal and *in vitro* models are also discussed.

Children are not small adults. They differ from adults in their exposures and may differ in their susceptibility to hazardous chemicals. Children's unique physiology and behavior can influence the extent of their exposure. Exposures of children are discussed in Section 5.6 Exposures of Children.

## 2. HEALTH EFFECTS

Children sometimes differ from adults in their susceptibility to hazardous chemicals, but whether there is a difference depends on the chemical (Guzelian et al. 1992; NRC 1993). Children may be more or less susceptible than adults to health effects, and the relationship may change with developmental age (Guzelian et al. 1992; NRC 1993). Vulnerability often depends on developmental stage. There are critical periods of structural and functional development during both prenatal and postnatal life and a particular structure or function will be most sensitive to disruption during its critical period(s). Damage may not be evident until a later stage of development. There are often differences in pharmacokinetics and metabolism between children and adults. For example, absorption may be different in neonates because of the immaturity of their gastrointestinal tract and their larger skin surface area in proportion to body weight (Morselli et al. 1980; NRC 1993); the gastrointestinal absorption of lead is greatest in infants and young children (Ziegler et al. 1978). Distribution of xenobiotics may be different; for example, infants have a larger proportion of their bodies as extracellular water and their brains and livers are proportionately larger (Altman and Dittmer 1974; Fomon 1966; Fomon et al. 1982; Owen and Brozek 1966; Widdowson and Dickerson 1964). The infant also has an immature blood-brain barrier (Adinolfi 1985; Johanson 1980) and probably an immature blood-testis barrier (Setchell and Waites 1975). Many xenobiotic metabolizing enzymes have distinctive developmental patterns. At various stages of growth and development, levels of particular enzymes may be higher or lower than those of adults, and sometimes unique enzymes may exist at particular developmental stages (Komori et al. 1990; Leeder and Kearns 1997; NRC 1993; Vieira et al. 1996). Whether differences in xenobiotic metabolism make the child more or less susceptible also depends on whether the relevant enzymes are involved in activation of the parent compound to its toxic form or in detoxification. There may also be differences in excretion, particularly in newborns who all have a low glomerular filtration rate and have not developed efficient tubular secretion and resorption capacities (Altman and Dittmer 1974; NRC 1993; West et al. 1948). Children and adults may differ in their capacity to repair damage from chemical insults. Children also have a longer remaining lifetime in which to express damage from chemicals; this potential is particularly relevant to cancer.

Certain characteristics of the developing human may increase exposure or susceptibility while others may decrease susceptibility to the same chemical. For example, although infants breathe more air per kilogram of body weight than adults breathe, this difference might be somewhat counterbalanced by their alveoli being less developed, which results in a disproportionately smaller surface area for alveolar absorption (NRC 1993).

## 2. HEALTH EFFECTS

There is no evidence from human toxicity studies that children are, or are likely to be, more susceptible to health effects from inhalation, oral, or dermal exposure to methylene chloride than adults. No cases of accidental poisoning in children due to methylene chloride exposure have been reported. The nervous system is a sensitive acute target of methylene chloride exposure in adults and the response in children is likely to be similar. Methylene chloride is neurotoxic at high concentrations, in part because of its lipophilic characteristics. There are no data in the literature that suggest that children are more susceptible to the acute or chronic neurotoxic effects of methylene chloride than adults. However, there is little information available in the literature on the chronic low-concentration neurotoxic effects of methylene chloride exposure in either children or adults. Methylene chloride did not produce any toxicologically significant developmental effects in animals in a two-generation reproductive study with rats (Nitschke et al. 1988b) or in mouse and rat developmental studies, except at very high, maternally toxic doses (Bornschein et al. 1980; Hardin and Manson 1980; Schwetz et al. 1975).

The only animal study that is suggestive of potentially age-related vulnerability to methylene chloride is that of Maronpot et al. (1995b), in which B6C3F<sub>1</sub> mice were exposed to 2,000 ppm of methylene chloride by inhalation for 26, 52, or 78 weeks. The results of stop-exposure experiments showed that early exposure to methylene chloride was more effective than late exposure in inducing pulmonary neoplasms. However, the authors mentioned the possibility that the late-exposed animals were sacrificed too early for tumors to have developed.

There is no information regarding the pharmacokinetics of methylene chloride in children or regarding the nutritional factors that may influence the absorption of methylene chloride. A PBPK model has been developed for estimating the amounts and concentrations of methylene chloride in human breast milk that would result from inhalation exposures to methylene chloride (Fisher et al. 1997). Animal studies demonstrate that methylene chloride and/or its metabolites distribute in the liver, kidney, lungs, brain, muscle, and adipose tissues after inhalation exposures (Carlsson and Hultengren 1975; McKenna et al. 1982) and has also been shown to cross the placenta in rats (Anders and Sunram 1982). Toxicokinetic data indicate that methylene chloride is rapidly cleared after cessation of exposure. Small amounts of methylene chloride have been found in breast milk (EPA 1980d; Pellizzari et al. 1982).

It is not known whether the metabolism of methylene chloride in children is different than in adults, but there are theoretical reasons to suspect it might be. Available data suggest that there are two pathways by which methylene chloride is metabolized. One pathway utilizes the mixed function oxidase (MFO)

## 2. HEALTH EFFECTS

enzymes and produces carbon monoxide (CO). The other pathway involves the glutathione transferase (GST) and produces carbon dioxide (CO<sub>2</sub>). No information was located regarding the possibility that the metabolism of methylene chloride via the GST metabolic pathway is developmentally regulated.

CYP2E1 (the specific MFO pathway for methylene chloride) expression is low in fetal liver but increases several hours after birth in humans and continues to increase during the first week of life (Carpenter et al. 1996, 1997; Jones et al. 1992; Viera et al. 1996). In the human fetal liver, CYP2E1 transcripts were not detected at 10 weeks of gestation, but were detected as early as 19 weeks (Carpenter et al. 1996). The CYP2E1 gene in the liver is modified by cytosine methylation at the 3' region during fetal development, which contributes to the low expression of the gene during fetal life (Jones et al. 1992; Viera et al. 1996). CYP2E1 expression in the fetal brain, however, has been detected near baseline levels at 46 days of gestation, but at significant levels beginning at 58 days (Brzezinski et al. 1999). The early expression of CYP2E1 in the fetal brain suggests that the brain may be particularly vulnerable to oxidative stress as a result of xenobiotic metabolism.

In cases of heavy ethanol consumption by the mother, CYP2E1 is induced in the placenta (Rasheed et al. 1997) and in the liver (Carpenter et al. 1997). *In vitro* tests demonstrated that ethanol treatment of human fetal hepatocytes upregulated expression of CYP2E1 (Carpenter et al. 1996). This presumably would increase the rate of metabolism of methylene chloride in the fetus.

Evidence from pregnant rats exposed to methylene chloride indicates that the maternal liver metabolizes methylene chloride at a higher rate than the fetus, but that the metabolite carbon monoxide equilibrates between the dam and fetus (Anders and Sunram 1982). The high affinity of fetal hemoglobin for both carbon monoxide and oxygen suggests that fetuses may be at risk from hypoxia following maternal exposures to high levels of methylene chloride (Anders and Sunram 1982; Longo 1977). The greater risk of hypoxia and the potential for greater neurological damage would last until the expression of adult hemoglobin during the first year of life.

As mentioned above, methylene chloride has been isolated from human breast milk (EPA 1980d; Pellizzari et al. 1982), so it is possible that maternal exposures could transmit the compound to infants. However, PBPK modeling suggests that lactating females who breast feed their infants will not deliver methylene chloride in significant quantities (Fisher et al. 1997). Simulation of a workday maternal inhalation exposure to methylene chloride at 50 ppm yielded a predicted daily intake in the infant of only 0.213 mg, which is significantly less than the 2.0 mg/day equivalent EPA Health Advisory Intake.

## 2. HEALTH EFFECTS

There are no biomarkers of exposure or effect for methylene chloride that have been validated in children or in adults exposed as children. There are no biomarkers in adults that identify previous childhood exposure. No information was located regarding pediatric-specific methods for reducing peak absorption following exposure to methylene chloride, reducing body burden or interfering with the mechanism of action for toxic effects. In addition, no data were located regarding whether methods for reducing toxic effects might be contraindicated in children.

### 2.8 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility (NAS/NRC 1989).

Due to a nascent understanding of the use and interpretation of biomarkers, implementation of biomarkers as tools of exposure in the general population is very limited. A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule(s) or cell(s) that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself or substance-specific metabolites in readily obtainable body fluid(s), or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic substance (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time samples can be taken. It may be difficult to identify individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc, and selenium). Biomarkers of exposure to methylene chloride are discussed in Section 2.8.1.

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction (e.g., increased liver enzyme activity or pathologic changes in female genital epithelial

## 2. HEALTH EFFECTS

cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are not often substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by methylene chloride are discussed in Section 2.8.2.

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic substance. It can be an intrinsic genetic or other characteristic or a preexisting disease that results in an increase in absorbed dose, a decrease in the biologically effective dose, or a target tissue response. If biomarkers of susceptibility exist, they are discussed in Section 2.10 "Populations That Are Unusually Susceptible".

### 2.8.1 Biomarkers Used to Identify or Quantify Exposure to Methylene Chloride

Measurements of parent methylene chloride and its metabolites in expired air, blood, and urine have been used as indicators of exposure. Elimination of methylene chloride from the body occurs primarily through pulmonary excretion; approximately 70–75% is excreted unchanged at concentrations from 50 to 200 ppm (DiVincenzo and Kaplan 1981). Pulmonary excretion was rapid during the first hour, then began to decline as steady state was approached. By 7 hours postexposure, expired air contained less than 1 ppm of methylene chloride for exposures between 50 and 150 ppm. The 7-hour values for an exposure of 200 ppm were twice that for the other concentrations. At 16 hours, negligible levels of methylene chloride were detected in all concentration groups. Therefore, measurements of methylene chloride in expired air as a biomarker are useful only if they occur within 6–8 hours of the most recent exposure (DiVincenzo and Kaplan 1981).

Methylene chloride can be detected in blood. Because it is cleared from blood very rapidly, this method is only useful for monitoring recent exposures. A plasma half-life of inhaled methylene chloride in humans is estimated to be 40 minutes (DiVincenzo et al. 1972).

Methylene chloride has been detected in adipose tissue in humans, but animal studies suggest that methylene chloride is cleared so rapidly (90% decrease in 2 hours), that it is useful for measuring only recent exposures (Carlsson and Hultengren 1975). Methylene chloride was detected in the breast milk of mothers living in urban industrial areas, but the route of exposure was not analyzed (EPA 1980e; Pellizzari et al. 1982).

## 2. HEALTH EFFECTS

Methylene chloride is also excreted in urine. Humans exposed to 100 ppm of methylene chloride vapor for 2 hours averaged 26.6  $\mu\text{g}$  methylene chloride in the urine within 24 hours after exposure and 85.5  $\mu\text{g}$  at exposure levels of 200 ppm (DiVincenzo et al. 1972).

Levels of COHb in the blood may also be used as an indicator of exposure to methylene chloride. Levels of COHb are concentration- and time-dependent (Stewart et al. 1972). Human subjects exposed to concentrations of 500 ppm or less for 1 hour experienced elevation in COHb levels (1–4%). These levels rose to an average of 10% saturation within 1 hour after exposure to higher concentrations (1,000 ppm for 2 hours) (Stewart et al. 1972). Exposure to concentrations as low as 100 ppm for 7.5 hours or 200 ppm for 4 hours resulted in COHb elevation above 5% in nonsmokers (Putz et al. 1979; Stewart et al. 1972). Levels of COHb can remain elevated above preexposure levels for more than 21 hours post exposure (Stewart et al. 1972). DiVincenzo and Kaplan (1981) showed recovery to baseline each day for 5 consecutive days. COHb saturation levels were lower than in Stewart et al. (1972). Different methods of COHb detection were used by the authors. Although COHb levels can be used as an indicator of exposure to methylene chloride, this biomarker is not specific. Exposure to other sources of carbon monoxide, such as tobacco smoke, incomplete organic fuel combustion, and automobile exhaust, will also increase COHb levels.

Ghittori et al. (1993) examined several methods for conducting biological monitoring of workers exposed occupationally to methylene chloride. Twenty males (12 smokers and 8 nonsmokers), employed at different jobs in a pharmaceutical factory where methylene chloride was used to wash gelatin capsules, wore a personal passive dosimeter in the respiratory zone during half of work days (4 hours) in order to measure the weighted mean-inspired environmental concentration of methylene chloride.

Tetrachloroethylene was also present in the work environment. Immediately after the end of the exposure, a urine sample was collected using gas-tight samplers. Carbon monoxide (CO) was determined at the end of the shift using a portable instrument.

Ambient exposures ranged from 49 to 168 ppm. No significant correlation was observed among the CO of all subjects and the concentration of methylene chloride in ambient air. When those workers who smoked were removed from analysis, a correlation between the methylene chloride concentration in air and the CO concentration in alveolar air was found ( $R = 0.87$ ). Significant linear correlation was found between the environmental concentration of methylene chloride in the breathing zone and methylene chloride concentration in urine ( $R = 0.9$  uncorrected,  $0.72$  corrected for creatinine). The authors suggest

## 2. HEALTH EFFECTS

that measuring methylene chloride in the urine was a more useful measure of exposure, since unlike COHb in the blood or CO in exhaled breath, it was unaffected by smoking.

Concentrations of CO in exhaled air studied in other groups of volunteers were shown to reach a maximum at 1–2 hours after exposure and were also directly proportional to the magnitude of exposure both during and after exposure. These findings are consistent with those studies.

The authors concluded that the results indicated that methylene chloride urinary concentration could be used as a possible biological index to evaluate methylene chloride air exposure, according to what has been observed for other solvents. However, at high ambient exposures (i.e., those exceeding 5–7 times the threshold limit value-time weighted average [TLV-TWA] of 49 ppm), the use of methylene chloride as a biological index might be problematic because exposure to very high concentrations gives disproportionately less COHb in the blood and more unmetabolized methylene chloride is produced.

### **2.8.2 Biomarkers Used to Characterize Effects Caused by Methylene Chloride**

As discussed in Section 2.2, the effects that are most often observed in humans exposed to methylene chloride vapors are central nervous system depression and behavioral effects. Clinical signs and symptoms which may be monitored include irritability, narcosis, and fatigue. Impairment of visual, auditory, and psychomotor functions can also be evaluated to detect early effects on the central nervous system. Since these effects also occur following exposure to numerous other chemicals, they are not specific for methylene chloride exposure and evaluation is often subjective.

Honma and Suda (1997) have investigated the effects of a single intraperitoneal administration of several short-chain chlorinated compounds, including methylene chloride, on lipoproteins in plasma and liver in male Fischer F344 rats. Changes in lipoproteins caused by these solvents were compared with hepatotoxicity markers such as GPT (ALT). Following administration of methylene chloride in olive oil, concentrations of lipoproteins (VLDL, LDL, HDL), triglyceride, cholesterol, and GPT in plasma were determined, as were changes in liver weight and amounts of triglyceride and glutathione in liver tissue. For methylene chloride, peaks of changes in these endpoints were observed at 8 or 19 hours following compound administration. The dose dependency of these changes were investigated after intraperitoneal dosing of 300 and 1,000 mg/kg methylene chloride. HDL decreased significantly at a dose of 300 mg/kg, whereas a marked increase in LDL occurred at 1,000 mg/kg. GPT also increased significantly at a dose



## 2. HEALTH EFFECTS

of 1,000 mg/kg. The authors conclude that changes in some lipoproteins in plasma occurred at lower doses than those causing elevation of GPT activity; therefore, these lipoproteins may be able to serve as sensitive and simple markers of adverse liver effects (i.e., biomarkers of effects). However, further investigation is necessary to determine whether these endpoints can serve as biomarkers of effects in humans when exposure is via inhalation and confounding variables are be numerous.

No other biomarkers have been identified to characterize effects associated with exposure to methylene chloride.

For more information on biomarkers for renal and hepatic effects of chemicals see ATSDR/CDC Subcommittee Report on Biological Indicators of Organ Damage (1990) and for information on biomarkers for neurological effects see OTA (1990).

### 2.9 INTERACTIONS WITH OTHER CHEMICALS

Limited studies were found in the available literature on the interactions of methylene chloride with other chemicals. Exposure of adult human subjects to 500 ppm of methylene chloride resulted in levels of COHb in the blood comparable with those produced by the TLV for CO (50 ppm) (Fodor and Roscovana 1976). A 4-hour exposure of human subjects to 200 ppm of methylene chloride or 70 ppm of CO resulted in similar blood COHb levels (Putz et al. 1979). This suggests that simultaneous exposure to methylene chloride and CO has an additive effect on blood COHb levels.

A possible additive effect of methylene chloride and toluene (as well as mineral spirits and methyl ethyl ketone) may have been operative in an occupational mixed-solvent exposure study in which manual dexterity, and visual memory task performances were impaired, even though the concentrations of the specific chemicals were below the recommended their threshold limit values (White et al. 1995); the reported adverse neurobehavioral effects are similar to those reported for individual exposures of methylene chloride at higher concentrations (Putz et al. 1979). In a rat study, methylene chloride and toluene interacted in an additive manner to cause retardation of embryonic growth and development *in vitro* (Brown-Woodman et al. 1998). In rats, combined CO and methylene chloride exposure yielded additive increases in the COHb levels (ACGIH 1986).

## 2. HEALTH EFFECTS

Methylene chloride decreases the nerve conduction velocity in animals when used alone but when the compound is administered with ethanol there is a more pronounced decrease of nerve conduction velocity (Glatzel et al. 1987). The additive or synergist effect of ethanol coadministration decreases at higher doses. Rats maintained on drinking water containing 10% (v/v) ethanol show increased levels of COHb after a single oral dose or during inhalation exposure to methylene chloride compared to rats that have not been exposed to ethanol (Wirkner et al. 1997). This interaction has been attributed to the induction of CYP2E1 by ethanol and a resulting increased production of carbon monoxide from methylene chloride. Wirkner et al. (1997) indicate that ethanol has no synergistic effect when methylene chloride is administered at high doses because the cytochrome P-450-dependent monooxygenase pathway becomes saturated at exposures >500 ppm.

Interactions resulting from the inhibition of metabolism of methylene chloride have been described in rats. Blood COHb levels in rats, after an oral dose of methylene chloride or during inhalation exposure to methylene chloride, are decreased by a concurrently administered oral dose of toluene (Ciuchta et al. 1979; Pankow et al. 1991a, 1991b). This interaction has been attributed to inhibition of the metabolism of methylene chloride to carbon monoxide, possibly as a result of both chemicals being a substrate for the 2E1 isoenzyme of cytochrome P-450 (Pelekis and Krishnan 1997).

In a rat study, methylene chloride (coinjected intraperitoneally) potentiated the hepatotoxic effect of carbon tetrachloride (Kim 1997). Coadministration reduced the methylene-chloride-induced rise in blood COHb, but augmented the carbon tetrachloride-induced increase in liver microsomal enzyme activities 3- to 8-fold. Methylene chloride increased the covalent binding of carbon tetrachloride metabolites to lipids, which the author suggests may be a cause of enhanced hepatotoxicity of the mixture.

### **2.10 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE**

A susceptible population will exhibit a different or enhanced response to methylene chloride than will most persons exposed to the same level of methylene chloride in the environment. Reasons may include genetic makeup, age, health and nutritional status, and exposure to other toxic substances (e.g., cigarette smoke). These parameters result in reduced detoxification or excretion of methylene chloride, or compromised function of organs affected by methylene chloride. Populations who are at greater risk due to their unusually high exposure to methylene chloride are discussed in Section 5.7, Populations With Potentially High Exposures.

## 2. HEALTH EFFECTS

There are certain subgroups of the general population that may be more susceptible to methylene chloride than others. One basis for this concern is the potential effect of COHb, produced from CO, a metabolite of methylene chloride. The COHb generated from methylene chloride is expected to be additive to COHb from other sources. Thus, methylene chloride exposure at high concentrations may pose an additional human health burden. Of particular concern are smokers (who maintain significant constant levels of COHb), and persons with existing cardiovascular disease. In addition, higher than normal levels of CO may result when alcoholics are exposed to methylene chloride, since ethanol increases the expression and activity of CYP2E1 (Carpenter et al. 1996). Similarly, enhanced expression of CYP2E1 occurs in the condition of diabetes, although insulin erases that effect (Thomas et al. 1987).

Varying susceptibility to methylene chloride may be correlated with polymorphism in its metabolizing enzymes. Genetic polymorphisms have been identified for both GSTT1 and CYP2E1 (Garte and Crosti 1999), and the health consequences of these variants are being explored (d'Errico et al. 1999; Lang and Pelkonen 1999; Pelkonen et al. 1999; Strange and Fryer 1999; Stubbins and Wolf 1999). In the case of GSTT1, which is the major metabolic pathway for methylene chloride at concentrations >500 ppm, there is, in addition to the wild type gene, a nonfunctioning allele with a deletion, so that three phenotypes exist: "high conjugators" homozygous for the wild type allele; "nonconjugators" homozygous for the deletion ("null) allele; and "low conjugators", heterozygotes with one of each (Thier et al. 1998). Thier et al. (1998) demonstrated that tissue samples representing the three phenotypes varied in the ability to metabolize methylene chloride. The health implications of the homozygous GSTT1-null genotype for people exposed to methylene chloride exposure are not clear. A potentially positive effect is that no toxic reactive GST-intermediates of methylene chloride would be produced, as demonstrated by the absence of RNA-formaldehyde adduct formation in GSTT1-null human hepatocytes treated with methylene chloride *in vitro* (Casanova et al. 1997). However, the overall rate of methylene chloride metabolism and elimination would be reduced in GSTT1-null individuals, so that narcotic effects of the parent compound would last longer. Furthermore, increased genotoxic damage (sister chromatid exchange) occurs following exposure to methylene chloride, in null-phenotype, compared to wild type lymphocytes (Hallier et al. 1994). Variations in the incidence of GSTT1 polymorphisms have been identified in different ethnic groups (Strange and Fryer 1999). In one study, the proportion of individuals carrying functional GSTT1 alleles was 36% among Chinese, 80% among Caucasians, and 90% among Mexican-Americans (Nelson et al. 1995). In a study of 416 subjects, the GSTT1-null genotype occurred among 24.1% of the blacks and 15% of the whites (Chen et al. 1996). The GSTT1-null phenotype was detected in 21.1% of

## 2. HEALTH EFFECTS

lung cancer patients with either squamous cell carcinoma or adenocarcinoma (El-Zein et al. 1997a).

Polymorphisms have also been detected in CYP2E1 (Garte and Crosti 1999; Lang and Pelkonen 1999). One polymorphism involves a mutation in the 5' flanking region of the CYP2E1 gene that leads to a higher rate of transcription, a higher level of protein, and higher enzyme activity compared to cells containing the wild type allele (Wan et al. 1998). The frequency of the mutant allele was 16% in a group of 203 Mexican-Americans (Wan et al. 1998), which is considerably higher than the 1–5% frequency measured in Caucasians (Stephens et al. 1994). Presumably, this allele would result in higher rates of metabolism of methylene chloride, and possibly more severe toxic effects, but this has not been assayed. A number of studies have tried to associate specific polymorphisms with cancer incidence (d'Errico et al. 1999). The CYP2E1 DraI DD genotype was found to be associated with a significantly higher risk of lung cancer among Mexican-Americans and African-Americans (Wu et al. 1998); expression of this genotype was also associated with an increase in bleomycin-induced single-strand breaks in lymphocytes tested *in vitro*. In another study, a rare CYP2E1 PstI variant (mutated in the transcription regulation region) was found in 7/57 lung cancer patients and 2/48 controls (El-Zein et al. 1997a); all seven patients developed adenocarcinoma. In a case control study, combined homozygosity for the CYP2E1 PstI variant and the GSTT1-null genotype was associated with an increased risk of developing lung cancer (El-Zein et al. 1997b). Presumably, individuals expressing these mutant genotypes would be susceptible to adverse effects following exposure to methylene chloride.

### 2.11 METHODS FOR REDUCING TOXIC EFFECTS

This section will describe clinical practice and research concerning methods for reducing toxic effects of exposure to methylene chloride. However, because some of the treatments discussed may be experimental and unproven, this section should not be used as a guide for treatment of exposures to methylene chloride. When specific exposures have occurred, poison control centers and medical toxicologists should be consulted for medical advice. The following texts provide specific information about treatment following exposures to methylene chloride: Ellenhorn 1997; Stewart and Dodd 1964.

## 2. HEALTH EFFECTS

**2.11.1 Reducing Peak Absorption Following Exposure**

Human exposure to methylene chloride may occur by inhalation, ingestion, or by dermal contact. Mitigation approaches to reduce absorption of methylene chloride have included general recommendations of separating contaminated food, water, air, or clothing from the exposed individual. Externally, exposed eyes and skin are flushed with a clean neutral solution such as water or normal saline. If the victim is alert and oriented and has an intact gag reflex, water or milk may be administered after ingestion of small amount of methylene chloride to wash residual chemical through the esophagus and dilute the contents in the stomach. In cases where persons have ingested more than several swallows, emesis can be induced with ipecac syrup after it has been determined that the victim is alert and oriented and precautions have been taken to protect the respiratory tract from aspiration of gastric contents. Once the individual is placed in the care of a health professional, gastric lavage can be performed within 1 hour of the exposure. Activated charcoal and cathartics are frequently recommended; however, no data exist to support their efficacy (Ellenhorn 1997).

Once methylene chloride has been inhaled or ingested, it is readily absorbed through the lungs or gastrointestinal tract. Dermal exposure also results in absorption, although at a slower rate than the other exposure routes and some references question whether the amount absorbed would be sufficient to cause systemic toxicity (Ellenhorn 1997; Stewart and Dodd 1964). Once absorbed, methylene chloride is rapidly metabolized in the liver in part to carbon monoxide. Because of the affinity of hemoglobin for carbon monoxide, COHb levels in the blood will increase, but free circulating carbon monoxide will not increase.

**2.11.2 Reducing Body Burden**

Inhalation data demonstrate that a portion of the absorbed methylene chloride is stored in fat tissue as evidenced by continued increase in COHb levels. It is likely to occur only with long-term exposure and at high concentrations (DiVincenzo et al. 1972). Investigations in humans following oral exposure to methylene chloride have failed to detect significant retention in fat or other tissue stores (Angelo et al. 1986a, 1986b; Ellenhorn 1997).

Following absorption, methylene chloride is distributed mainly to the liver, brain and subcutaneous adipose tissue (Carlsson and Hultengren 1975). The liver is the primary site of metabolism, although

## 2. HEALTH EFFECTS

additional transformation occurs in the lungs and kidneys. In the liver, methylene chloride may undergo metabolism by two pathways. The first pathway produces CO and CO<sub>2</sub> and is saturable at a few hundred ppm. The second pathway yields formaldehyde and formic acid and shows no indication of saturation at inhaled concentrations up to 10,000 ppm. Acute toxic effects (central nervous system depression) may persist for hours after removal from the source of exposure because of continued metabolism of methylene chloride released from tissue storage (ATSDR 1990). COHb levels can continue to rise, peaking 5–6 hours after exposure. Peak levels of 12 (NIOSH 1974) and 50% (Ellenhorn 1997) have been reported. Most of the effects seen following acute high-concentration exposure are due to the anesthetic properties of the parent compound. Fatalities have occurred with COHb levels less than 10%; thus, it is not likely that carboxyhemoglobine was the cause of death. A study by Scholz et al. (1991) suggests that, in addition to anesthetic effects, the sudden onset of cardiac arrhythmia induced by acute exposure to methylene chloride may contribute to its lethality.

The metabolic contribution of each pathway appears to vary in humans, particularly with the exposure level, and therefore toxicity extrapolation between high and low doses is complex (Gargas et al. 1986). Furthermore, recent studies suggest that the second pathway is considerably more active in certain animal species, particularly mice, a finding that complicates interspecies comparisons (ATSDR 1990; Green et al. 1986b, 1986c).

The body eliminates methylene chloride primarily through the lungs. A small amount of unchanged methylene chloride is also eliminated in the urine and feces. At low doses, a large percentage of methylene chloride is metabolized to form COHb and eliminated as carbon monoxide, while at higher doses more of the unchanged parent compound is exhaled (ATSDR 1990). The administration of 100% oxygen is efficacious in reducing the half-life of carbon monoxide and should be continued until COHb is less than 5%.

The prompt application of hemoperfusion and diuretic therapy eliminated metabolic acidosis and hemoglobinuria in the case of an acute oral exposure to paint remover containing methylene chloride (Roberts and Marshall 1976). This treatment was thought to have prevented renal damage, but did not address the problem of gastrointestinal ulceration.

## 2. HEALTH EFFECTS

**2.11.3 Interfering with the Mechanism of Action for Toxic Effects**

The primary effects of exposure to methylene chloride appear to be hepatotoxicity and neurotoxicity. However, mechanisms of action for these toxic effects are not well characterized and it is difficult to speculate concerning specific methods for preventing these effects.

Because one consequence of metabolism of methylene chloride is production of COHb, methylene chloride exposure victims are sometimes given supplemental oxygen. The administration of oxygen increases the dissociation of carbon monoxide from hemoglobin and hastens the reduction of COHb. The half-life of COHb is normally approximately 5.3 hours but this can be reduced to 60–90 minutes with inhalation of 100% oxygen. Hyperbaric oxygen as used in carbon monoxide poisoning from direct carbon monoxide inhalation may be useful when high COHb levels are present (Haddad and Winchester 1990). At high COHb levels, hyperbaric oxygen will reduce the half-life to 20–40 minutes (ATSDR 1990). Because carbon monoxide is generated metabolically, this often necessitates a longer duration of oxygen therapy after methylene chloride poisoning than with carbon monoxide poisoning.

Specific mechanisms of action relating to neurological effects are not well understood. Steroids and mannitol have been used to decrease cerebral edema caused by methylene chloride toxicity, but their value in preventing later neurologic sequelae remains unproven (ATSDR 1990).

Victims with cardiopulmonary disease or workers who are exposed to carbon monoxide sources could have an elevated risk from methylene chloride metabolism and should be monitored closely. One reference suggests that physical exercise and smoking be avoided because these activities may have an additive effect on the COHb level (Ellenhorn 1997).

**2.12 ADEQUACY OF THE DATABASE**

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of methylene chloride is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of methylene chloride.

## 2. HEALTH EFFECTS

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

**2.12.1 Existing Information on Health Effects of Methylene Chloride**

The existing data on health effects of inhalation, oral, and dermal exposure of humans and animals to methylene chloride are summarized in Figure 2-5. The purpose of this figure is to illustrate the existing information concerning the health effects of methylene chloride. Each dot in the figure indicates that one or more studies provide information associated with that particular effect. The dot does not necessarily imply anything about the quality of the study or studies, nor should missing information in this figure be interpreted as a “data need”. A data need, as defined in ATSDR’s *Decision Guide for Identifying Substance-Specific Data Needs Related to Toxicological Profiles* (ATSDR 1989), is substance-specific information necessary to conduct comprehensive public health assessments. Generally, ATSDR defines a data gap more broadly as any substance-specific information missing from the scientific literature.

As shown in Figure 2-5, studies of humans exposed to methylene chloride by the inhalation route are available. These have focused mainly on neurological, cancer, and systemic effects (e.g., cardiovascular). One report focused on possible reproductive effects following inhalation (and possible dermal) exposure. Other end points (immunological, developmental, and genotoxic effects) have not been evaluated. There are no studies on the effects of methylene chloride in humans after ingestion. Other than one study evaluating potential reproductive effects following inhalation and possibly dermal exposure, no other end points have been evaluated following dermal exposure.

Studies in animals have also focused mainly on inhalation exposure and several end points have been evaluated. Effects of oral exposure have focused on death, systemic effects after intermediate and chronic exposure, genotoxicity, and cancer. Other end points have not been evaluated. There are limited reports on the effects of methylene chloride after direct ocular exposure of animals.



2. HEALTH EFFECTS

**Figure 2-5. Existing Information of Health Effects of Methylene Chloride**

		Systemic									
		Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation		●	●		●		●	●			●
Oral							●		●		
Dermal		●	●					●			

**Human**

		Systemic									
		Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation		●	●	●	●	●	●	●	●	●	●
Oral		●	●	●	●				●		●
Dermal			●								

**Animal**

● Existing Studies

## 2. HEALTH EFFECTS

**2.12.2 Identification of Data Needs**

**Acute-Duration Exposure.** Available data indicate that the central nervous system is the primary target of inhaled methylene chloride in humans (Bakinson and Jones 1985; Fodor and Winneke 1971; Hall and Rumack 1990; Putz et al. 1979; Stewart et al. 1972; Winneke 1974), rats (Heppel and Neal 1944; Rebert et al. 1989; Savolainen et al. 1981), guinea pigs, rabbits, dogs, and monkeys (Heppel et al. 1944). An acute inhalation MRL was derived for methylene chloride based on a LOAEL of 300 ppm for neurological effects in humans (Winneke 1974). Respiratory effects (cough, breathlessness, and in one fatal case, pulmonary congestion with focal hemorrhage) were observed in several cases of acute occupational exposure to methylene chloride in humans (Bakinson and Jones 1985; Snyder et al. 1992a, 1992b; Winek et al. 1981). No cardiovascular effects were seen in humans acutely exposed to concentrations between 100 and 475 ppm (Cherry et al. 1981; Ott et al. 1983c). Gastrointestinal effects (nausea and vomiting) were reported in several cases of acute occupational exposure to methylene chloride (Bakinson and Jones 1985). Acute inhalation exposure to methylene chloride increased the blood COHb level in humans (DiVincenzo and Kaplan 1981; Manno et al. 1992; Putz et al. 1979; Soden et al. 1996). An acute inhalation study in mice (Aviado and Belej 1974) suggests that the cardiovascular system may be a target for methylene chloride toxicity; however, the effects were observed at lethal concentrations. An acute study in guinea pigs demonstrated an increase in hepatic triglycerides, but no histopathological effects on the liver following exposure to methylene chloride at 5,200 ppm (Morris et al. 1979). Since the most sensitive acute effects in humans are neurological, additional animal studies that evaluate changes in analogous functions (visual, auditory discrimination tasks) at low levels of exposure might be needed. These experiments should monitor CO/COHb levels in the animals to correlate exposure and effect. Furthermore, the animals should be genotyped with respect to for GSTT1 and CYP2E1 to evaluate the mechanism of toxicity.

The only studies in humans on the effects of acute oral exposure to methylene chloride are reports of suicide attempts by ingestion of paint removers. One of these studies reported suppression of the central nervous system and metabolic acidosis (Roberts and Marshall 1976). Corrosion of the gastrointestinal tract was also reported (Hughes and Tracey 1993; Roberts and Marshall 1976). Hematological effects included elevated COHb (Hughes and Tracey 1993) and intravascular hemolysis leading to hemoglobinuria (Roberts and Marshall 1976). No acute respiratory, cardiovascular, hepatic, endocrine, or immunological effects have been reported in humans following ingestion of methylene chloride, and there are no acute oral studies on the effect of low doses in humans. Limited studies in animals involved

## 2. HEALTH EFFECTS

exposure of rats to high doses of methylene chloride via gavage, resulting in increased mortality (Kimura et al. 1971; Ugazio et al. 1973); as in suicide cases, respiratory failure as a result of suppression of the central nervous system was the cause of death in these animal studies. Hemolysis occurred and diuresis was inhibited in rats gavaged with 1,325 mg/kg of methylene chloride (Marzotko and Pankow 1987); in rats given single oral doses 526 mg/kg, the only endocrine effects noted were dilatation of capillaries of the adrenal medulla and an increase in the secretion of catecholamines (Marzotko and Pankow 1987). In rats gavaged twice with 1,275 mg/kg, DNA damage and an increase in the activity of ornithine decarboxylase were detected in the liver (Kitchin and Brown 1989). Genotoxicity (DNA breakage) was detected in nuclei from selected rat organs following a single dose of methylene chloride (1,720 mg/kg) (Sasaki et al. 1998). Since there is a potential for oral exposure for humans near hazardous waste sites (through contaminated groundwater sources where volatilization is restricted), additional animal data on the effects of acute oral exposure to lower doses of methylene chloride would strengthen the database. An acute oral MRL was derived by using a PBPK model to extrapolate human inhalation data from Winneke (1974) to estimate the concentration of methylene chloride in drinking water needed to produce a tissue specific concentration equivalent to that produced by exposure to 300 ppm of methylene chloride (Reitz et al. 1997).

Case reports indicate that methylene chloride can cause eye and skin damage following direct contact with the liquid (Hall and Rumack 1990; Wells and Waldron 1984; Winek et al. 1981). An acute study in rabbits demonstrated adverse ocular effects (Ballantyne et al. 1976). Dermal absorption of methylene chloride has been demonstrated in animals (McDougal et al. 1986), and therefore, is possible in humans. Additional dermal studies in animals are needed to provide further insight into the potential risks of living near hazardous waste sites.

**Intermediate-Duration Exposure.** Suggestive evidence for liver effects in humans exists, as workers exposed to methylene chloride (up to 475 ppm) via inhalation have shown increased serum bilirubin levels, but not at clinically significant levels (Ott et al. 1983a). Other liver injury parameters were normal. Repeated exposures to methylene chloride vapors (up to 500 ppm) did not alter serum enzyme activity, pulmonary function, or urine chemistry significantly in exposed and control groups (NIOSH 1974). However, irritation of the respiratory tract, but no changes in urinary chemistry were noted after exposure to a TWA of 5–340 ppm in an occupational study (Anundi et al. 1993).

## 2. HEALTH EFFECTS

Available data for rats, dogs, monkeys, rabbits, guinea pigs, gerbils, and mice indicate that repeated exposure to methylene chloride via inhalation produces central nervous system effects (Briving et al. 1986; Karlsson et al. 1987; Rosengren et al. 1986), respiratory effects (Heppel et al. 1944; NTP 1986) hepatic effects (Haun et al. 1972; Heppel et al. 1994; Kjellstrand et al. 1986; MacEwen et al. 1972; Norpoth et al. 1974; NTP 1986; Weinstein and Diamond 1972), and kidney effects (Haun et al. 1972; MacEwen et al. 1972). An intermediate inhalation MRL was based on a LOAEL of 25 ppm for hepatic and renal effects in rats (Haun et al. 1972).

No quantitative data are available for oral exposures of intermediate duration in humans. Hepatic damage and lowered urinary pH in rats was reported following oral exposure \$166 mg/kg/day for 3 months (Kirschman et al. 1986), but the reporting of results was incomplete in this study. Furthermore, since the lowest dose was a LOAEL, additional intermediate oral studies are needed in animals to evaluate specific tissue and organ effects at lower doses and to determine threshold levels and dose-response relationships. Reitz et al. (ATSDR 1997) developed a PBPK model to extrapolate inhalation data for an oral MRL, based on the inhalation study of Haun et al. (1972), but the reporting of results in this study was incomplete; a new intermediate oral animal study would be preferable. No intermediate oral MRL was derived because of lack of adequate data.

There are no quantitative intermediate dermal exposure data available for humans or animals. Given the potential for human exposure to methylene chloride through paint strippers, adhesives, glues, paint thinners, wood stain, varnishes, spray paint, and automobile spray primers, information on the effects of dermal exposures are needed to estimate more accurately the risks to human health from methylene chloride.

**Chronic-Duration Exposure and Cancer.** No data are available on the non-neoplastic effects in humans after chronic exposure to methylene chloride via the oral or dermal routes. Neurological and reproductive effects have been reported in workers following combined inhalation/dermal exposures for up to 3 years (Kelly 1988). Studies in animals suggest that the liver is a target organ following chronic inhalation (Burek et al. 1984; Nitschke et al. 1988a; NTP 1986) and oral (Serota 1986a, 1986b) exposure. Both a chronic inhalation MRL and a chronic oral MRL were derived using hepatotoxicity as the critical endpoint (Nitschke et al. 1988a; Serota et al. 1986a). Renal effects were also observed following chronic inhalation of methylene chloride in rodents (NTP 1986). Dermal data in animals after chronic exposure

## 2. HEALTH EFFECTS

to methylene chloride are not available. Additional dermal studies in animals are needed to enhance our understanding of potential risk to people living near hazardous waste sites.

Epidemiological studies of human occupational cohorts show no increase in cancer of the lung, liver, or any other organs from occupational inhalation exposures (Friedlander et al. 1978; Hearne et al. 1987, 1990; Ott et al. 1983a). There are no human oral or dermal exposure data for the cancer endpoint. Inhalation studies in animals show a concentration-dependent, statistically significant increase in liver and lung adenomas and carcinomas in mice exposed to high concentration of methylene chloride (Mennear et al. 1988; NTP 1986) and benign mammary gland tumors in rats (Mennear et al. 1988; NTP 1986) following 2 years of exposure to methylene chloride. The evidence for carcinogenicity in animals from oral exposures (Serota et al. 1986a, 1986b) is inconclusive, and there are no dermal data available. Therefore, additional chronic oral and dermal studies are needed to clarify the cancer risk of ingested methylene chloride. The carcinogenic mechanism of inhaled methylene chloride in animals is not yet understood despite an extensive database on toxicokinetics and potential mode(s) of action. Additional information on the mechanisms of carcinogenesis is needed to provide further insight regarding current findings that (1) the concentration-response may be nonlinear at low exposure concentrations and there is a concentration range below which carcinogenicity is unlikely to occur and (2) the mouse is not an appropriate model for investigation of potential human carcinogenicity because it is much more sensitive than any other species.

**Genotoxicity.** Genotoxicity data show mixed results. Generally, mutagenesis assays are negative in mammalian *in vivo* studies (Burek et al. 1984; Sheldon et al. 1987). Methylene chloride is mutagenic in bacteria (Gocke et al. 1981), but results for *Drosophila* (Gocke et al. 1981) are contradictory. This compound has shown a dose-response relationship for clastogenesis in mammalian cells *in vitro* (Thilagar et al. 1984a) but rats exposed *in vivo* (Burek et al. 1984) have shown no effects. In mice exposed *in vivo*, tissue-specific DNA breaks were observed in mice (Sasaki et al. 1998), which suggests that genotoxicity of methylene chloride may be dependent on the expression of metabolizing enzymes. In view of the unknown carcinogenic mechanisms, additional *in vivo* studies of clastogenesis are needed to provide useful markers, both for dosimetry and specific mechanisms. A recent *in vitro* test using strains of *Salmonella* with or without recombinant GSTT1 has demonstrated that methylene chloride may have more than one mechanism of genotoxicity (De Marini et al. 1997). In bacteria expressing mammalian GSTT1, methylene chloride, at moderate doses, caused a single type of genetic lesion. In bacteria without GSTT1, higher doses were required to induce genetic lesions and the results were more varied. These

## 2. HEALTH EFFECTS

results may explain the variability observed in previous studies. Therefore, additional genotoxicity studies should be carried out on mammalian cells for which the expression of GSTT1 and/or CYP2E1 isoenzymes is defined.

**Reproductive Toxicity.** Data on reproductive toxicity in humans are limited. One case study reported genital pain, and low sperm counts (testicular atrophy in some cases) in workers who inhaled vapors of methylene chloride and had direct contact with the liquid on the job for up to 3 years (Kelly 1988); they were also exposed to styrene at low levels. Exposure to methylene chloride was confirmed by blood COHb levels. In another study, workers exposed to methylene chloride for a shorter time showed no reproductive effects (Wells et al. 1989b). A case-control occupational study reported a nearly significant association of methylene chloride exposure and the incidence of spontaneous abortion (Taskinen et al. 1986). There are no oral or dermal exposure data that pertain to reproductive effects in humans. A two-generation inhalation study in rats reported no effects on fertility and litter size (Nitschke et al. 1988b). In addition, no effects on the testes were observed in dominant lethal tests involving male mice that inhaled concentrations of methylene chloride up to 200 ppm for up to 6 weeks (Raje et al. 1988). Uterine, ovarian, and testicular atrophy were observed in rodents following chronic exposure to methylene chloride at 4,000 ppm (NTP 1986). There are no data on reproductive effects in animals following oral or dermal exposure. Intermediate-duration oral and dermal studies that incorporate histopathological analysis of the reproductive organs are needed to address this data need. Part of the analysis should include a determination of whether the reproductive organs express GSTT1 and/or CYP2E1, as a first step in evaluating their possible role in the reproductive toxicity of methylene chloride.

**Developmental Toxicity.** No data are available on developmental toxicity of methylene chloride following inhalation, oral, or dermal exposure in humans, aside from the case-control study relating exposure and the rate of spontaneous abortion mentioned above (Taskinen et al. 1986). Exposure to 0.01 ppm of methylene chloride had no effect on birth outcome in a study of over 90,000 pregnancies (Bell et al. 1991). Several inhalation studies in animals indicate that methylene chloride can cross the placenta (Anders and Sunram 1982). Some of these studies showed statistically nonsignificant malformations in rats and mice or decreased fetal weight at maternally toxic concentrations (Bornschein et al. 1980; Hardin and Manson 1980; Schwetz et al. 1975). However, there was a statistically significant increase in the incidence of delayed ossification of sternbrae (Schwetz et al. 1975). The use of only one concentration in these studies precludes any evaluation of concentration-response relationships.

## 2. HEALTH EFFECTS

However, Reitz et al. (1997) developed an inhalation route-to-oral route extrapolation and rodent-to-human species extrapolation using PBPK modeling of the developmental toxicity data in Schwetz et al. (1975). The resulting LOAEL was an intermediate oral dose of 142 mg/kg/day. Additional studies for inhalation and dermal exposures in two species would be useful in clarifying the developmental toxicity potential of this chemical. Conducting studies on animals with known genotypes with respect to metabolizing enzymes GSTT1 and CYP2E1 are needed to evaluate the risk of exposure to methylene chloride.

**Immunotoxicity.** No human data are available on immunotoxicity of methylene chloride. Animal data are limited on the immunotoxicity of methylene chloride. One study was located that indicated that methylene chloride caused splenic atrophy in dogs following continuous intermediate-duration inhalation exposure (MacEwen et al. 1972). Splenic fibrosis was observed in a chronic rat study (Mennear et al. 1988), but occurred only at the highest concentration tested (1,000 ppm) and only in one sex. A recent intermediate-duration study, also in rats, found no evidence of gross or microscopical alterations in the spleen at a concentration of 5,187 ppm methylene chloride (Halogenated Solvent Industry Alliance, Inc. (2000). This study also found no effect on IgM response to SRBC. Additional immunotoxicity studies including evaluation of humoral- and cell-mediated immunity are needed to determine whether this system is susceptible to methylene chloride, as some chlorinated hydrocarbons do affect the immune system.

**Neurotoxicity.** The central nervous system is a target for both short- and long-term inhalation exposures in humans. These data are derived from experimental (Fodor and Winneke 1971; Putz et al. 1979; Stewart et al. 1972; Winneke 1974;) and occupational studies (Lash et al. 1991; White et al. 1995) that reported alterations in behavioral performance and various psychomotor tasks following exposure to methylene chloride. Other neurotoxic effects noted in occupational studies included dizziness, headaches, nausea, memory loss, paresthesia, tingling in hands and feet, and loss of consciousness (Bakinson and Jones 1985; Hall and Rumack 1990; Kelly 1988). It should be noted that some of these workers were exposed to other unspecified solvents at the same time. Winneke (1974) attributed the neurological effects in volunteers following a 3–4 hour inhalation exposure to 300–800 ppm of methylene chloride to the anaesthetic properties of the parent compound since exposures to 50–100 ppm of carbon monoxide alone did not produce these effects. The specific parameters in the Winneke (1974) study, critical flicker fusion frequency, auditory vigilance, and other psychomotor tasks, were considered to be specific measures of ‘cortical alertness’ or central nervous system depression. Putz et al. (1979) correlated

## 2. HEALTH EFFECTS

neurological deficits following inhalation exposure to methylene chloride (200 ppm) with the accumulation of COHb in the blood and carbon monoxide in the exhaled breath; performance did not deteriorate until 3 hours of exposure had elapsed and blood COHb levels exceeded 5%. Although the Putz et al. (1979) study seems to implicate the metabolite CO as the neurotoxic agent, there is a need for studies to determine the relative contributions of parent compound and metabolite to the neurotoxic effects of methylene chloride. The Winneke (1974) study, which was deemed to have used more specific measures of central nervous system depression than the Putz et al. (1979) study, was used to derive an acute inhalation MRL and, by route-to-route extrapolation, also to derive an acute oral MRL (Reitz et al. 1997).

Subtle neurological effects were reported in rats after acute exposure to methylene chloride. Changes in somatosensory-evoked potentials were observed after 1 hour exposure to 5,000 ppm of methylene chloride (Rebert et al. 1989), and changes in cerebellar enzymes were detected following 2 weeks of exposure at 500 ppm (Savolainen et al. 1981). Inhalation studies in rats did not reveal neurobehavioral effects after intermediate-duration exposure at 4,500 ppm of methylene chloride vapors (Bornschein et al. 1980). In other studies, neurochemical changes were reported in gerbils at 210 ppm (Briving et al. 1986; Karlsson et al. 1987; Rosengren et al. 1986). There were no treatment-related neurophysiological or neuropathological effects in rats exposed to concentrations of 2,000 ppm for 13 weeks (Mattsson et al. 1990).

There are no oral or dermal studies in either humans or animals with regard to neurotoxicity of methylene chloride. Data on electrophysiology, functional observational batteries, and neuropathology from 90-day oral or dermal exposures in at least two different species would be useful to further evaluate neurotoxic effects from these routes of exposure, since human exposure can occur from all three routes at hazardous waste sites.

**Epidemiological and Human Dosimetry Studies.** Information on the effects of methylene chloride in humans comes from occupational studies, case reports of acute high exposure, and studies with volunteers. Asphyxia and eventually death occurred in a subject acutely exposure to a high but undetermined concentration of methylene chloride in the air (Winek et al. 1981). Exposure to lower concentrations affects primarily the central nervous system; signs and symptoms observed include dizziness, incoordination, loss of balance, unconsciousness, and decreased performance in tests of sensory and motor functions (Bakinson and Jones 1985; Fodor and Winneke 1971; Hall and Rumack 1990; Putz



## 2. HEALTH EFFECTS

et al. 1979; Stewart et al. 1972; Winneke 1974). These effects are likely to be caused by a combination of the anaesthetic properties of the parent compound and accumulation of COHb which forms as a result of methylene chloride metabolism. There is a potential for low-level general population exposure from ambient methylene chloride emissions from paint removal, aerosol use, metal degreasing, electronics and pharmaceutical manufacturing, and food processing (Callahan 1981; CPSC 1987, 1990; NAS 1978). Such exposure levels are unlikely to produce adverse neurological effects. Hazardous waste sites release methylene chloride into the air, groundwater, surface water, and soil. There are very few monitoring data for methylene chloride outside the occupational setting (Singh et al. 1981). Because methylene chloride evaporates readily from water and soil, inhalation is the main route of potential exposure. In the unlikely event that long-term exposure of the general population (in the past or present) to low levels of primarily methylene chloride is identified, individuals should be monitored for hematological and hepatic effects, as these have been observed in studies in animals.

**Biomarkers of Exposure and Effect.**

**Exposure.** The presence of methylene chloride in the postexposure expired breath is the most commonly used biomarker of exposure. Methylene chloride is easily detected in expired air for 24 hours following a vapor exposure (NIOSH 1974). Methylene chloride can be detected in blood, but because clearance is so rapid, this method is only useful for monitoring recent exposures (DiVincenzo and Kaplan 1980). The half-life of COHb following methylene chloride exposure is twice that following exposure to carbon monoxide (NIOSH 1974; Stewart and Hake 1976). COHb in blood may be monitored; however, COHb is cleared so rapidly (plasma half-life of 40 minutes) that it is unlikely to be useful for monitoring environmental exposures (DiVincenzo et al. 1972). Urinary excretion of methylene chloride is measurable for several hours postexposure (DiVincenzo et al. 1972).

**Effect.** Changes in the nervous system are sensitive, but not specific, biomarkers of methylene chloride effects. Clinical signs and symptoms which may be monitored include narcosis, fatigue, and analgesia (Bakinson and Jones 1985; Hall and Rumack 1990; Putz et al. 1979; Winneke 1974). Impairment of neurobehavioral functions and electromyographic measurements of nerve conduction velocity and amplitude can be monitored to detect early signs of neurotoxicity in people exposed to methylene chloride (Glatzel et al. 1987; White et al. 1995). Additional studies that couple measurement of methylene chloride with tests for determining nervous system effects are needed to correlate exposure with adverse health effects of this chemical.

## 2. HEALTH EFFECTS

While neurobehavioral tests are not specific for methylene chloride-induced toxicity, they do identify potential health impairment. Studies to develop more specific biomarkers of methylene chloride-induced effects are needed to assess the potential health risk of methylene chloride exposure near hazardous waste sites.

**Absorption, Distribution, Metabolism, and Excretion.** Methylene chloride is a volatile liquid with high lipid solubility and modest solubility in water. In humans, it is rapidly absorbed by the inhalation (DiVincenzo and Kaplan 1981; McKenna et al. 1980) and ingestion routes of exposure (Roberts and Marshall 1976). Dermal absorption is suspected to occur in humans, but quantitative data are lacking. Methylene chloride is also readily absorbed by animals following inhalation (DeVincenzo et al. 1972; MacEwen et al. 1972; McKenna et al. 1982) and oral exposure (Angelo et al. 1986a); dermal absorption data in animals are also lacking. Once absorbed, methylene chloride is quickly distributed to a wide range of tissues and body fluids. Absorption and distribution of methylene chloride can be affected by a number of factors, including dose level, vehicle, physical activity, duration of exposure, and amount of body fat (Astrand et al. 1975; DiVincenzo et al. 1972; Engstrom and Bjorstrom 1977).

Distribution data in humans are lacking, but it has been found in human breast milk (EPA 1980e; Pellizzari et al. 1982). Methylene chloride is widely distributed in animal tissues after inhalation exposure (Carlsson and Hultengren 1975; McKenna et al. 1982). The highest concentrations are found in adipose tissue and liver. Methylene chloride has been found in blood from rats' fetuses (Anders and Sunram 1982). After acute exposure, methylene chloride disappears rapidly from fat (Carlsson and Hultengren 1975). No studies were located regarding distribution of methylene chloride following dermal exposure in animals. Distribution of methylene chloride does not seem to be route-dependent and it does not bioaccumulate in tissues.

Methylene chloride is metabolized via two pathways, the MFO pathway, which produces CO and CO<sub>2</sub>, and the GST pathway which yields only CO<sub>2</sub> (Gargas et al. 1986). Methylene chloride is metabolized almost exclusively by the MFO pathway at low exposures. The only data available for humans on the toxicokinetics parameters are for inhalation exposures. These data show that the GST activity is low in humans relative to other animal species (Andersen et al. 1987). It has been postulated that the activity of the GST pathway in rats is less than that of mice and might only become significant when the P-450 pathway (MFO) has been saturated (Green et al. 1986b, 1986c). Toxicokinetic data for oral and dermal exposures would be useful because some exposures to humans are expected to be from contaminated

## 2. HEALTH EFFECTS

drinking water and soils. Toxicokinetic models (Andersen and Krishnan 1994; Andersen et al. 1987, 1991; Dankovic and Bailer 1994; Ramsey and Andersen 1984; Reitz 1990) have been developed that account for the nonlinearities in the internal dose across exposure levels that arise from concentration-dependent changes in absorption, distribution, excretion, and saturation of metabolism following inhalation exposure to methylene chloride. Reitz et al. (1997) have developed an inhalation route-to-oral route extrapolation for methylene chloride. The metabolism of methylene chloride in humans appears to be qualitatively similar to that in animals (Fodor et al. 1973; Wirkner et al. 1997).

Methylene chloride is excreted via expired air primarily as CO and CO<sub>2</sub> at low concentrations (McKenna et al. 1982). As concentration increases, more unchanged parent compound is exhaled. A small fraction of absorbed methylene chloride or metabolites has been detected in urine of humans occupationally exposed (DiVincenzo et al. 1972) and in the urine and feces of experimental animals (McKenna et al. 1982). Additional data on excretion parameters as functions of concentration are needed to understand the toxicokinetics of methylene chloride.

**Comparative Toxicokinetics.** Generally, animal data on toxicokinetics parameters substantiate those from humans. There is rapid absorption through the lung (DiVincenzo and Kaplan 1981; DiVincenzo et al. 1972; MacEwen et al. 1972; McKenna et al. 1982). The metabolites of methylene chloride in different species indicate that they are metabolized by the same pathways to CO or to CO<sub>2</sub> (Fodor et al. 1973; Gargas et al. 1986; Wirkner et al. 1997). The data from both human and animal studies indicate that the target organs of methylene chloride toxicity are the same across species. Interesting differences do exist, however, in some metabolic parameters. The rate of clearance from rat tissues was markedly slower than in the mouse following inhalation exposures. In contrast to findings of a direct proportional relationship between inspired air concentration of methylene chloride and blood level in man and other animals, there is a report that the ratio at steady state between blood levels of methylene chloride and exposure increases as the concentration increases in rats (Green et al. 1986b, 1986c). Distribution of methylene chloride across tissues in rats and humans has been shown to be lipophilic except for one rat study that showed less of the chemical in adipose tissue (McKenna and Zempel 1981). Species differences in placental anatomy and physiology may contribute to variations in fetal exposure to methylene chloride or its metabolites following maternal exposure. Additional data on the impact of differences in species sensitivity to specific internal doses by different routes of exposure are needed to resolve questions about extrapolation of predicted effects from different routes in different species.

## 2. HEALTH EFFECTS

**Methods for Reducing Toxic Effects.** There is no specific treatment or antidote for methylene chloride intoxication. Available methods and treatments (e.g., gastric lavage with the airway protected, or the use of oxygen as in carbon monoxide poisoning) for reducing peak absorption have been shown to be beneficial (Shih 1998; Tomaszewski 1998). There is no evidence that activated charcoal or cathartics are effective (Shih 1998). Because methylene chloride is metabolized to carbon monoxide, mitigation strategies (e.g., hyperbaric oxygen) used for carbon monoxide intoxication have been effective in increasing the oxygen carrying capacity of the blood (Tomaszewski 1998).

**Children's Susceptibility.** Data needs relating to both prenatal and childhood exposures, and developmental effects expressed either prenatally or during childhood, are discussed in detail in the Developmental Toxicity subsection above.

There are no studies that specifically addressed effects of exposure to methylene chloride in children. There is no evidence from human toxicity studies that children are, or are likely to be, more susceptible to health effects of exposure to methylene chloride than adults. No cases of accidental poisoning in children due to methylene chloride exposure have been reported. The nervous system is a sensitive target for acute exposure to methylene chloride in adults, and the response in children is likely to be similar.

Although developmental studies in animals indicate that methylene chloride is not a teratogen (Bornschein et al. 1980; Hardin and Manson 1980), there is a need to evaluate neurological/neuro-behavioral effects in animals exposed *in utero*. Subtle neurological effects could result from hypoxia (CO-mediated) or from reactive intermediates of metabolism that would only be revealed by appropriate behavioral testing of the offspring. It is not clear that the "wheel running activity" and "avoidance learning" tests that Bornschein et al. (1980) employed in rats exposed *in utero* were adequate to reveal neurological deficits. Acute effects observed in an adult human study involved degraded performance on visual and auditory discrimination tasks (Putz et al. 1979). If neurological effects were detected in mice, it would be useful to conduct additional developmental studies using mice in which functional genes for GSTT1 and/or CYP2E1 have been knocked-out, to discover which metabolic pathway is implicated in developmental neurological effects.

There is no direct evidence regarding pharmacokinetics or the mechanism of toxicity of methylene chloride in children. However, considering what is known about the expression of metabolizing enzymes (see Section 2.7 Children's Susceptibility), there is no reason to expect that children over the age of

## 2. HEALTH EFFECTS

6 months will be qualitatively different from adults; by 6 months, the last of the fetal hemoglobin has been replaced by the adult isoforms. At fetal and early postnatal stages, however, children will be more vulnerable to methylene-chloride-generated CO because of its effect on fetal hemoglobin (Longo 1977). CO binding to Hb causes oxygen to be held more tightly, so that tissues receive less oxygen; the net effect is that the CO clears more slowly from the circulation. It is not known to what extent methylene chloride (or metabolites other than CO or CO<sub>2</sub>) can cross the placenta in humans or accumulate in breast milk. A study in animals showed that methylene chloride crosses the placental barrier (Anders and Sunram 1982), but quantitative data are lacking. There are no animal studies testing whether methylene chloride can pass into breast milk. There are no studies quantifying maternal exposure and output into breast milk. The Fisher et al. (1997) PBPK model predicts a relatively low transfer into breast milk following maternal inhalation exposure.

In adults, the biomarker of exposure to methylene chloride is the parent compound, and the biomarkers of effect are CO in the exhaled breath or COHb in the blood. There is no reason to expect that biomarkers in children would be any different.

It is expected that any chemical interactions involving methylene chloride would have the same effects in children as in adults. The exception would be the fetal period, during which concentrations of metabolizing enzymes are significantly lower than in the adult. A fetus exposed to methylene chloride and another chemical that requires CYP2E1 for metabolism, might be at higher risk than an adult.

Treatment with 100% oxygen has been used to enhance the rate of clearance of CO from the bloodstream after acute methylene chloride exposure. Although there is no specific information regarding pediatric treatments for methylene chloride exposure, 100% oxygen has been used for treating pregnant women exposed to CO (Longo 1977). In these cases, it is recommended that the oxygen treatment be extended 3.5–5 times longer than the time required to bring the maternal COHb level down to 3%, to ensure that the fetus is adequately treated. If an exposed mother has neurological symptoms or a COHb level above 15%, a single hyperbaric oxygen treatment is indicated (Tomaszewski 1998). It does not appear that any pediatric-specific methods need to be developed.

Since methylene chloride may be genotoxic, depending on the genotype with respect to GSTT1 and CYP2E1 (see Section 2.10), it is possible that parental exposure could affect the fetus. At issue would be whether the gonads express alleles that are associated with genotoxicity, since the effect is intracellular.

## 2. HEALTH EFFECTS

Child health data needs relating to exposure are discussed in 5.8.1 Identification of Data Needs: Exposures of Children.

**2.12.3 Ongoing Studies**

A number of research projects are in progress investigating the health effects and mechanism of action of methylene chloride. These projects have been identified from FEDRIP (1999).

Dr. J.B. Wheeler, at Vanderbilt University, Nashville, Tennessee, is comparing the ability of rat and human glutathione S-transferases (GST) to form mutagenic DNA adducts with glutathione in the presence of different haloalkanes. He will examine the mechanism of activation by GST, the stability of the glutathione conjugates, and the DNA adducts produced. This research is sponsored by the National Cancer Institute.

Dr. T.R. Devereux, at the NIEHS, is identifying critical target genes and genetic alterations that may be important in chemical carcinogenesis. He is characterizing genetic alterations in oncogenes (e.g., ras) and tumor suppressor genes (e.g., p53 and p16) from rodent tumors and human cancers generated by exposure to chemical agents. He is also examining polymorphisms in the human p16 gene that may predispose to lung cancer.

Dr. D.A. Bell, of the Genetic Risk Group at the NIEHS, is conducting case-control studies of 5,000 genotyped individuals to test the impact of cancer susceptibility genes on cancer of the bladder, lung, liver, colon, stomach, prostate, and breast. He and his colleagues have identified ethnic differences in the frequency of at-risk genotypes for glutathione transferase (M1, theta 1, and Pi), and N-acetyltransferase (1 and 2).