

§ 24.180

wine; however, wine spirits may not be added to specially sweetened natural wine. Specially sweetened natural wines may be blended with each other, or with natural wine or heavy bodied blending wine (including juice or concentrated fruit juice to which wine spirits have been added), in the further production of specially sweetened natural wine only if the wines (or juice) so blended are made from the same kind of fruit. (Sec. 201, Pub. L. 85-859, 72 Stat. 1383, as amended, 1384, as amended, 1385, as amended, 1386, as amended (26 U.S.C. 5382, 5383, 5384, 5385))

[T.D. ATF-299, 55 FR 24989, June 19, 1990, as amended by T.D. ATF-312, 56 FR 31078, July 9, 1991]

§ 24.180 Use of concentrated and unconcentrated fruit juice.

Concentrated fruit juice reduced with water to its original density, or to 22 degrees Brix, or to any degree of Brix between its original density and 22 degrees Brix, and unconcentrated fruit juice reduced with water to not less than 22 degrees Brix, is considered juice for the purpose of standard wine production. Concentrated fruit juice reduced with water to any degree of Brix greater than 22 degrees Brix may be further reduced with water to any degree of Brix between its original density and 22 degrees Brix. The proprietor, prior to using concentrated fruit juice in wine production, shall obtain a statement in which the producer certifies the kind of fruit from which it was produced and the total solids content of the juice before and after concentration. Concentrated or unconcentrated fruit juice may be used in juice or wine made from the same kind of fruit for the purposes of chaptalizing or sweetening, as provided in this part. Concentrated fruit juice, or juice which has been concentrated and reconstituted, may not be used in standard wine production if at any time it was concentrated to more than 80 degrees Brix. (Sec. 201, Pub. L. 85-859, 72 Stat. 1383, as amended (26 U.S.C. 5382))

(Approved by the Office of Management and Budget under control number 1512-0298)

[T.D. ATF-299, 55 FR 24989, June 19, 1990, as amended by T.D. ATF-413, 64 FR 46845, Aug. 27, 1999]

27 CFR Ch. I (4-1-08 Edition)

§ 24.181 Use of sugar.

Only sugar, as defined in § 24.10, may be used in the production of standard wine. The quantity of sugar used will be determined either by measuring the increase in volume or by considering that each 13.5 pounds of pure dry sugar results in a volumetric increase of one gallon. (Sec. 201, Pub. L. 85-859, 72 Stat. 1383, as amended, 1384, as amended, 1385, as amended, 1387, as amended (26 U.S.C. 5382, 5383, 5384, 5392))

[T.D. ATF-299, 55 FR 24989, June 19, 1990, as amended by T.D. ATF-312, 56 FR 31078, July 9, 1991]

§ 24.182 Use of acid to correct natural deficiencies.

(a) *General.* Acids of the kinds occurring in grapes or other fruit (including berries) may be added within the limitations of § 24.246 to juice or wine in order to correct natural deficiencies; however, no acid may be added to juice or wine which is ameliorated to correct natural deficiencies except that in the production of grape wine, tartaric acid may be used to reduce the pH of the juice or wine. If tartaric acid is used to correct the pH of grape juice or wine, the fixed acid level of the juice shall be measured prior to the addition of any tartaric acid to determine the maximum quantity of ameliorating material allowed. In addition, when using tartaric acid to reduce the pH of ameliorated grape juice or wine, the pH cannot be reduced below 3.0.

(b) *Grape wine.* Tartaric acid or malic acid, or a combination of tartaric acid and malic acid, may be added prior to or during fermentation, to grapes or juice from grapes. In addition, after fermentation is completed, citric acid, fumaric acid, malic acid, lactic acid or tartaric acid, or a combination of two or more of these acids, may be added to correct natural deficiencies. However, the use of these acids, either prior to, during or after fermentation, may not increase the fixed acid level of the finished wine (calculated as tartaric acid) above 9.0 grams per liter. In cases where the wine contains 8.0 or more grams of total solids per 100 milliliters of wine, acids may be added to the extent that the finished wine does not contain more than 11.0 grams per liter