SHORELINE CHANGES IN THE ISLES DERNIERES BARRIER ISLAND ARC: 1887 - 1996

Terrebonne Parish, Louisiana

Jim Penland, Chris Kindinger, Karen Westphal, Paul Connell, Jeff Lee, and S. Jeffress Williams

INTRODUCTION

The U.S. Geological Survey (USGS), in cooperation with the Coastal Research Laboratory at the University of Southwestern Louisiana and the University of New Orleans, initiated the Coastal Geology Investigations Program in 1979 (Penland and Boyd, 1981). Coastal changes are a common theme in many of the studies undertaken by this program. Previous studies have focused on barrier and headland shorelines in Louisiana (Gulledge and others, 1985; Gulledge and Penland, 1986; Penland and others, 1986; Penland and Ramsey, 1990; Penland and others, 1988; Stone Street, 1988; Stone and others, 1992) and on the Chandeleur Islands in Louisiana (Penland and others, 1988), the Mississippi Sound (Penland and others, 1989), the Mississippi-Alabama barrier islands (Williams and others, 1992), and the Florida Panhandle coast (Stone and others, 1992). This report is the first in a series of publications that will document coastal changes in the Louisiana barrier islands.

The Isles Dernieres are located about 100 km southwest of New Orleans (Figure 1). The barrier island arc is 35 km long and extends from Caillou Boca on the west to Whiskey Pass. The Isles Dernieres consist of a narrow island arc, a wide island arc, and a remnant island arc (Figure 2). The narrow island arc is the youngest of the three arcs and is the only arc that has been subaerially exposed since 1887. The wide island arc was subaerially exposed from 1887 to 1988 and is now mostly submerged. The remnant island arc is the oldest of the three arcs and was subaerially exposed from 1887 to 1932. The Isles Dernieres are separated by Caillou Boca, Whiskey Pass, and Trinity Shoals. The Isles Dernieres are bounded on the east by Timbalier Island and on the west by Raccoon Island. The Isles Dernieres are the subject of this report.

The Isles Dernieres have undergone extensive subaerial and subaqueous changes since the first survey in 1887. Subaerial changes include island segmentation, island migration, island drowning, and island overwash. Subaqueous changes include the development of flood-tidal deltas and the migration of ebb-tidal deltas. Island segmentation has occurred at Raccoon Island, Whiskey Island, and Trinity Island. Island migration has occurred at Raccoon Island, Whiskey Island, Trinity Island, and Wine Island. Island drowning has occurred at Trinity Island, East Island, and Wine Island. Overwash has occurred at Raccoon Island, Whiskey Island, Trinity Island, and Wine Island. Flood-tidal deltas have developed at Caillou Boca, Whiskey Pass, and Trinity Shoals. Ebb-tidal deltas have developed at Caillou Boca and Whiskey Pass.

This report presents historical changes in the Isles Dernieres from 1887 to 1996. Changes in island area, island width, island length, island segmentation, island migration, island drowning, island overwash, flood-tidal delta development, and ebb-tidal delta development are described. Changes in island area, island width, and island length are presented in tables. Changes in island segmentation, island migration, island drowning, island overwash, flood-tidal delta development, and ebb-tidal delta development are presented in figures.

METHODS

Historical changes in the Isles Dernieres were determined from T-sheets, topographic maps, and aerial photographs. T-sheets are nautical charts published by the U.S. Coast and Geodetic Survey (USC&GS) in the late 1800's and early 1900's. T-sheets were published at a scale of 1:20,000 and were based on surveys conducted in the late 1870's and early 1880's. Topographic maps were published by the USC&GS and the U.S. Geological Survey (USGS) from the early 1900's to the present. Topographic maps were published at a scale of 1:24,000 from 1934 to the present. Aerial photographs were taken by the USGS, the U.S. Department of Agriculture (USDA), and the National Aeronautics and Space Administration (NASA) from 1932 to the present. Aerial photographs were published at a scale of 1:20,000 from 1932 to 1956 and at a scale of 1:40,000 from 1956 to the present.

Historical changes in the Isles Dernieres were determined by digitizing shorelines from T-sheets, topographic maps, and aerial photographs. Shorelines were digitized using the Intergraph Interactive Graphics Design System (IGDS) software package. Shorelines were digitized from T-sheets at a scale of 1:20,000, from topographic maps at a scale of 1:24,000, and from aerial photographs at a scale of 1:20,000. Shorelines were digitized from T-sheets, topographic maps, and aerial photographs at a scale of 1:20,000 to minimize digitizing errors. Digitized shorelines were then plotted at a scale of 1:20,000 for analysis.

Historical changes in island area, island width, and island length were determined from digitized shorelines. Island area was calculated using the Intergraph Interactive Graphics Design System (IGDS) software package. Island width was measured perpendicular to the island axis. Island length was measured parallel to the island axis.

Historical changes in island segmentation, island migration, island drowning, island overwash, flood-tidal delta development, and ebb-tidal delta development were determined from digitized shorelines. Island segmentation was determined by comparing the number of islands in each arc in 1887 with the number of islands in each arc in 1996. Island migration was determined by comparing the position of the 1887 shoreline with the position of the 1996 shoreline. Island drowning was determined by comparing the position of the 1887 shoreline with the position of the 1996 shoreline. Island overwash was determined by comparing the position of the 1887 shoreline with the position of the 1996 shoreline. Flood-tidal delta development was determined by comparing the position of the 1887 shoreline with the position of the 1996 shoreline. Ebb-tidal delta development was determined by comparing the position of the 1887 shoreline with the position of the 1996 shoreline.

Table 1

Isles Dernieres gulfward migration (-) or accretion (+)				Isles Dernieres bayward migration (+) or loss (-)			
1887 - Jan 88	Jan 88 - Dec 96	1887 - Dec 96		1887 - Jan 88	Jan 88 - Dec 96	1887 - Dec 96	
m	m	m		m	m	m	
0	0	0		0	0	0	
-48	0	-48		48	0	48	
-96	0	-96		96	0	96	
-144	0	-144	