

Table G3.A4. Metabolic Syndrome: Physical Activity Longitudinal Studies

Author, Journal, Year	N	Finding
Laaksonen et al./ <i>Diabetes Care</i> / 2002 (1)	612 M	Inverse association
Carnethon et al./ <i>Diabetes Care</i> / 2004 (2)	N=4,192	None
Palaniappan et al./ <i>Diabetes Care</i> / 2004 (3)	N=714 318 M 396 W	None
Ferreira et al./ <i>Arch. Intern. Med.</i> / 2005 (4)	N=364	Inverse association
Ekelund et al./ <i>Diabetes Care</i> / 2005 (5)	N=605 246 M 359 W	Inverse association
Ekelund et al./ <i>Diabetes Care</i> / 2007 (6)	N=393 176 M 217 W	Inverse association
Holme et al./ <i>BMC Public Health</i> / 2007 (7)	16,209 M	Inverse association

M, men; N, number; W, women

Reference List

1. Laaksonen DE, Lakka HM, Salonen JT, Niskanen LK, Rauramaa R, Lakka TA. Low levels of leisure-time physical activity and cardiorespiratory fitness predict development of the metabolic syndrome. *Diabetes Care* 2002 Sep;25(9):1612-8.
2. Carnethon MR, Loria CM, Hill JO, Sidney S, Savage PJ, Liu K. Risk factors for the metabolic syndrome: the Coronary Artery Risk Development in Young Adults (CARDIA) study, 1985-2001. *Diabetes Care* 2004 Nov;27(11):2707-15.
3. Palaniappan L, Carnethon MR, Wang Y, Hanley AJ, Fortmann SP, Haffner SM, Wagenknecht L. Predictors of the incident metabolic syndrome in adults: the Insulin Resistance Atherosclerosis Study. *Diabetes Care* 2004 Mar;27(3):788-93.
4. Ferreira I, Henry RM, Twisk JW, van MW, Kemper HC, Stehouwer CD. The metabolic syndrome, cardiopulmonary fitness, and subcutaneous trunk fat as independent determinants of arterial stiffness: the Amsterdam Growth and Health Longitudinal Study. *Arch.Intern.Med.* 2005 Apr 25;165(8):875-82.
5. Ekelund U, Brage S, Franks PW, Hennings S, Emms S, Wareham NJ. Physical activity energy expenditure predicts progression toward the metabolic syndrome independently of aerobic fitness in middle-aged healthy Caucasians: the Medical Research Council Ely Study. *Diabetes Care* 2005 May;28(5):1195-200.
6. Ekelund U, Franks PW, Sharp S, Brage S, Wareham NJ. Increase in physical activity energy expenditure is associated with reduced metabolic risk independent of change in fatness and fitness. *Diabetes Care* 2007 Aug;30(8):2101-6.

7. Holme I, Tonstad S, Sogaard AJ, Larsen PG, Haheim LL. Leisure time physical activity in middle age predicts the metabolic syndrome in old age: results of a 28-year follow-up of men in the Oslo study. *BMC.Public Health* 2007;7(147):154.