

Table G3.A5. Children: Exercise and Metabolic Syndrome

Author, Journal, Year	N	Random/Control	Finding
Brage et al./ <i>Diabetes Care</i> / 2004 (1)	N=589 310 F 279 M	Cross-sectional	Inverse association
Eisenmann et al./ <i>Intl. J. Obesity</i> / 2005 (2)	N=761 345 F 416 M	Inverse association	Inverse association
Eisenmann et al./ <i>Am. Heart J.</i> / 2005 (3)	N=48	Prospective	Inverse association
Ferreira et al./ <i>Arch. Intern. Med</i> / 2005 (4)	N=364 189 F 175 M	Prospective	Inverse association
Platat et al./ <i>Diabetologia</i> / 2006 (5)	N=640 325 F 315 M	Cross-sectional	Inverse association
DuBose et al./ <i>Pediatrics</i> / 2007 (6)	N=375 193 F 182 M	Cross-sectional	Inverse association
Kelishadi et al./ <i>Horm. Res.</i> / 2007 (7)	N=4,811 2,563 F 2,248 M	Cross-sectional	Inverse association
Eisenmann et al./ <i>Med. Sci. Sports Exerc.</i> / 2007 (8)	N=1,615 755 F 860 M	Cross-sectional	Inverse association

F, females; M, males; N, number

Reference List

1. Brage S, Wedderkopp N, Ekelund U, Franks PW, Wareham NJ, Andersen LB, Froberg K. Features of the metabolic syndrome are associated with objectively measured physical activity and fitness in Danish children: the European Youth Heart Study (EYHS). *Diabetes Care* 2004 Sep;27(9):2141-8.
2. Eisenmann JC, Katzmarzyk PT, Perusse L, Tremblay A, Despres JP, Bouchard C. Aerobic fitness, body mass index, and CVD risk factors among adolescents: the Quebec family study. *Int.J.Obes.(Lond)* 2005 Sep;29(9):1077-83.
3. Eisenmann JC, Wickel EE, Welk GJ, Blair SN. Relationship between adolescent fitness and fatness and cardiovascular disease risk factors in adulthood: the Aerobics Center Longitudinal Study (ACLS). *Am.Heart J.* 2005 Jan;149(1):46-53.
4. Ferreira I, Henry RM, Twisk JW, van MW, Kemper HC, Stehouwer CD. The metabolic syndrome, cardiopulmonary fitness, and subcutaneous trunk fat as independent determinants of arterial stiffness: the Amsterdam Growth and Health Longitudinal Study. *Arch.Intern.Med.* 2005 Apr 25;165(8):875-82.
5. Platat C, Wagner A, Klumpp T, Schweitzer B, Simon C. Relationships of physical activity with metabolic syndrome features and low-grade inflammation in adolescents. *Diabetologia* 2006 Sep;49(9):2078-85.

6. DuBose KD, Eisenmann JC, Donnelly JE. Aerobic fitness attenuates the metabolic syndrome score in normal-weight, at-risk-for-overweight, and overweight children. *Pediatrics* 2007 Nov;120(5):e1262-e1268.
7. Kelishadi R, Razaghi EM, Gouya MM, Ardalan G, Gheiratmand R, Delavari A, Motaghian M, Ziaee V, Siadat ZD, Majdzadeh R, et al. Association of physical activity and the metabolic syndrome in children and adolescents: CASPIAN Study. *Horm.Res.* 2007;67(1):46-52.
8. Eisenmann JC, Welk GJ, Ihmels M, Dollman J. Fatness, fitness, and cardiovascular disease risk factors in children and adolescents. *Med.Sci.Sports Exerc.* 2007 Aug;39(8):1251-6.