#### HIV/AIDS in Urban and Nonurban Areas

- Many ways to characterize urban and nonurban areas and populations
- CDC uses metropolitan statistical areas (MSAs) as defined by the Office of Management and Budget (OMB)
- OMB subdivides MSAs as:
  - >500,000 population
  - 50,000–500,000 population
  - Nonmetropolitan (<50,000)</li>






# Reported AIDS Cases and Rates among Adults and Adolescents, by Population of Area of Residence, 2006 and Cumulative, 50 States and DC

MSA Population	Number	Rate/100,000	Cumulative through 2006
>500,000	30,607	19.3	807,912
50,000–500,000	4,239	9.2	88,041
Nonmetropolitan	2,696	6.4	51,146





# Proportion of Reported AIDS Cases among Adults and Adolescents, by Population of Area of Residence 1994–2006—50 States and DC





Note. Data based on residence at time of AIDS diagnosis.



#### **Regions of the United States**






# Reported AIDS Cases and Rates for Adults and Adolescents by Region and Population of Area of Residence 2006—50 States and DC

		MSA of >500,000		MSA of 50,000–500,000		Nonmetropolitan area	
Region	No.	Rate/ 100,000	No.	Rate/ 100,000	No.	Rate/ 100,000	
Northeast	8,628	23.8	598	11.6	324	7.1	
Midwest	3,480	11.5	555	4.8	366	2.8	
South	13,189	25.4	2,517	12.9	1,805	9.8	
West	5,310	13.0	569	5.8	201	3.4	





### Proportion of Reported AIDS Cases among Adults and Adolescents by Region and Population of Area of Residence 2006—50 States and DC








## Reported AIDS Cases among Adults and Adolescents by Race/Ethnicity and Population of Area of Residence 2006—50 States and DC

Population	White, not Hispanic	Black, not Hispanic	Hispanic	Asian/ Pacific Islander	American Indian/ Alaska Native
MSA of >500,000	8,499	15,162	6,135	456	94
MSA of 50,000 -500,000	1,607	1,938	566	48	33
Non- metropolitan	1,051	1,292	264	13	34





Proportion of Estimated AIDS Cases among Male Adults and Adolescents, by Transmission Category and Population of Area of Residence, 2002–2006—50 States and DC






<sup>\*</sup>Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.



#### Proportion of Estimated AIDS Cases among Female Adults and Adolescents, by Transmission Category and Population of Area of Residence, 2002–2006—50 States and DC






Note. Data based on residence at time of AIDS diagnosis. Data have been adjusted for reporting delays and cases

without risk factor information were proportionally redistributed.

Includes heterosexual contact with a bisexual, a person with hemophilia, an HIV-infected transfusion recipient, or an HIV infected person whose risk factor is unknown or not identified.




### Reported AIDS Cases among Adults and Adolescents by Age at Diagnosis and Population of Area of Residence 2006—50 States and DC







#### Reported AIDS Cases among Adults and Adolescents in Nonmetropolitan Areas, 2006—50 States and DC






Note. Data based on residence at time of AIDS diagnosis.
According to the U.S. Office of Management and Budget, the District of Columbia and New Jersey do not have any nonmetropolitan areas.

\* (nonmetropolitan cases/total cases) x100




Reported Cases of HIV Infection (not AIDS) among Adults and Adolescents in Nonmetropolitan Areas







### AIDS Rates for Adults and Adolescents, by Population of Area of Residence and Region Reported in 2006—50 States and DC



Note. Data based on residence at time of AIDS diagnosis.



# Reported AIDS Cases and Rates for Adults and Adolescents, by Population of Area of Residence and Sex, 2006—50 States and DC

	MSA of	MSA of	Nonmetropolitan	
	>500,000	50,000–500,000	area	
Sex	Rate/	Rate/	Rate/	
	No. 100,000	No. 100,000	No. 100,000	
Male	22,327 28.8	3,078 13.7	2,001 9.6	
Female	8,279 10.2	1,161 4.9	695 3.3	



