

Table 14. Tuberculosis Cases and Percentages by Reason TB Therapy Stopped: United States, 1993-2001

Year	Total Cases ¹	Completed Therapy		Moved		Lost		Refused		Died ²		Other ³	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	23,741	18,037	(76.0)	1,121	(4.7)	1,087	(4.6)	223	(0.9)	3,051	(12.9)	222	(0.9)
1994	23,052	17,761	(77.1)	1,195	(5.2)	737	(3.2)	182	(0.8)	2,743	(11.9)	434	(1.9)
1995	21,712	17,284	(79.6)	971	(4.5)	562	(2.6)	155	(0.7)	2,389	(11.0)	351	(1.6)
1996	20,298	16,507	(81.3)	783	(3.9)	521	(2.6)	156	(0.8)	1,990	(9.8)	341	(1.7)
1997	18,930	15,650	(82.7)	668	(3.5)	434	(2.3)	119	(0.6)	1,755	(9.3)	304	(1.6)
1998	17,584	14,766	(84.0)	535	(3.0)	399	(2.3)	104	(0.6)	1,578	(9.0)	202	(1.2)
1999	16,862	14,231	(84.4)	459	(2.7)	355	(2.1)	105	(0.6)	1,438	(8.5)	274	(1.6)
2000	15,784	13,664	(84.7)	416	(2.6)	387	(2.5)	112	(0.7)	1,291	(8.2)	214	(1.4)
2001	15,402	13,038	(84.7)	401	(2.6)	353	(2.3)	94	(0.6)	1,103	(7.2)	413	(2.7)

¹Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

²Died = died of any cause (not only TB).

³Includes cases in persons reporting reason therapy stopped = Other, Missing, or Unknown.

Note: Data for all years are updated through March 30, 2004.

Data complete through 2001 only. See Technical Notes (page 9) for details.