

Table 41. Tuberculosis Cases and Percentages by Completion of Tuberculosis Therapy (COT): 59 Reporting Areas, 2001

Reporting Area	Total Cases	Therapy \leq 1 Year Indicated ¹		Therapy $>$ 1 Year Indicated ²		All Drug Therapy		
		No. ³	COT \leq 1 Year(%)	No. ³	COT(%)	No. ³	COT(%)	
United States	15,945	14067	(80.5)	(91.4)	232	(79.7)	14299	(91.2)
Alabama	264	224	(78.6)	(91.5)	3	(100.0)	227	(91.6)
Alaska	54	50	(88.0)	(92.0)	0	...	50	(92.0)
Arizona	289	254	(81.9)	(93.3)	9	(66.7)	263	(92.4)
Arkansas	162	138	(92.0)	(95.7)	0	...	138	(95.7)
California	3,329	2969	(79.2)	(90.7)	44	(65.9)	3013	(90.3)
Colorado	138	128	(93.0)	(96.1)	3	(66.7)	131	(95.4)
Connecticut	121	113	(70.8)	(92.0)	1	(100.0)	114	(92.1)
Delaware	33	30	(93.3)	(93.3)	0	...	30	(93.3)
District of Columbia	74	63	(85.7)	(92.1)	0	...	63	(92.1)
Florida	1,139	1018	(80.6)	(92.5)	12	(100.0)	1030	(92.6)
Georgia	577	507	(80.3)	(91.1)	6	(83.3)	513	(91.0)
Hawaii	150	138	(72.5)	(92.8)	1	(0.0)	139	(92.1)
Idaho	9	7	—	—	0	...	7	—
Illinois	703	643	(74.3)	(85.4)	4	(100.0)	647	(85.5)
Indiana	115	97	(92.8)	(97.9)	1	(100.0)	98	(98.0)
Iowa	43	36	(77.8)	(100.0)	1	(0.0)	37	(97.3)
Kansas	81	70	(72.9)	(80.0)	3	(100.0)	73	(80.8)
Kentucky	152	131	(88.5)	(96.9)	1	(100.0)	132	(97.0)
Louisiana	294	259	(77.2)	(89.2)	0	...	259	(89.2)
Maine	20	19	(94.7)	(100.0)	0	...	19	(100.0)
Maryland	262	235	(87.7)	(93.6)	5	(100.0)	240	(93.8)
Massachusetts	270	250	(78.0)	(90.8)	6	(100.0)	256	(91.0)
Michigan	329	275	(80.4)	(90.2)	6	(50.0)	281	(89.3)
Minnesota	239	222	(83.8)	(94.1)	6	(100.0)	228	(94.3)
Mississippi	154	137	(81.8)	(91.2)	2	(50.0)	139	(90.6)
Missouri	156	138	(84.8)	(94.2)	1	(100.0)	139	(94.2)
Montana	20	17	(94.1)	(100.0)	0	...	17	(100.0)
Nebraska	40	31	(83.9)	(93.5)	1	(100.0)	32	(93.8)
Nevada	96	87	(80.5)	(95.4)	1	(0.0)	88	(94.3)
New Hampshire	20	19	(89.5)	(100.0)	0	...	19	(100.0)
New Jersey	530	459	(75.4)	(91.7)	7	(57.1)	466	(91.2)
New Mexico	54	46	(84.8)	(95.7)	0	...	46	(95.7)
New York State ⁴	413	375	(80.3)	(91.2)	8	(87.5)	383	(91.1)
New York City	1,231	1065	(86.7)	(94.7)	30	(83.3)	1095	(94.4)
North Carolina	397	340	(88.5)	(95.6)	3	(100.0)	343	(95.6)
North Dakota	6	5	(100.0)	(100.0)	0	...	5	(100.0)
Ohio	306	258	(79.1)	(91.5)	3	(33.3)	261	(90.8)
Oklahoma	193	159	(79.9)	(91.8)	4	(100.0)	163	(92.0)
Oregon	123	113	(87.6)	(96.5)	2	(100.0)	115	(96.5)
Pennsylvania	349	281	(74.0)	(86.1)	8	(100.0)	289	(86.5)
Rhode Island	60	60	(73.3)	(91.7)	0	...	60	(91.7)
South Carolina	263	227	(85.0)	(95.2)	3	(100.0)	230	(95.2)
South Dakota	13	12	(66.7)	(83.3)	0	...	12	(83.3)
Tennessee	313	265	(84.2)	(95.1)	1	(100.0)	266	(95.1)
Texas	1,631	1442	(75.8)	(86.9)	31	(87.1)	1473	(86.9)
Utah	35	33	(81.8)	(93.9)	0	...	33	(93.9)
Vermont	7	7	(100.0)	(100.0)	0	...	7	(100.0)
Virginia	306	271	(80.8)	(90.4)	5	(60.0)	276	(89.9)
Washington	261	241	(84.6)	(94.6)	8	(62.5)	249	(93.6)
West Virginia	32	25	(64.0)	(80.0)	0	...	25	(80.0)
Wisconsin	86	75	(81.3)	(96.0)	2	(100.0)	77	(96.1)
Wyoming	3	3	(100.0)	(100.0)	0	...	3	(100.0)
American Samoa ⁵
Fed. States of Micronesia ⁵
Guam ⁵	63	56	—	—	3	—	59	—
N. Mariana Islands ⁵
Puerto Rico ⁵	121	92	(92.4)	(97.8)	2	(50.0)	94	(96.8)
Republic of Palau ⁵
U.S. Virgin Islands ⁵

¹Initial isolate susceptible to rifampin (n=10,571) or susceptibility unknown (n=434); culture-negative (n=2,349); culture status unknown (n=713); age unknown (n=2).

²Initial isolate rifampin resistant, or pediatric (aged <15) patients with meningeal, bone, joint, or military disease.

³Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed, who did not die during therapy. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for \geq 90% of cases.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Ellipses indicate data not available. See Technical Notes for description of COT calculation (page 9).