

**Table 34. Tuberculosis Cases and Percentages by Excess Alcohol Use,<sup>1</sup> Age ≥15: Reporting Areas, 2006**

Reporting Area	Total Cases	Cases with Information on Excess Alcohol Use		Cases Reporting Excess Alcohol Use	
		No.	(%)	No.	(%)
<b>United States</b>	<b>12,971</b>	<b>12,674</b>	<b>(97.7)</b>	<b>1,795</b>	<b>(14.2)</b>
Alabama	191	191	(100.0)	21	(11.0)
Alaska	63	63	(100.0)	30	(47.6)
Arizona	274	232	(84.7)	38	(16.4)
Arkansas	95	95	(100.0)	6	(6.3)
California	2,646	2,576	(97.4)	222	(8.6)
Colorado	117	117	(100.0)	11	(9.4)
Connecticut	85	85	(100.0)	11	(12.9)
Delaware	27	27	(100.0)	3	(11.1)
District of Columbia	67	67	(100.0)	4	(6.0)
Florida	981	974	(99.3)	199	(20.4)
Georgia	465	452	(97.2)	83	(18.4)
Hawaii	113	74	(65.5)	--	--
Idaho	19	16	(84.2)	3	(18.8)
Illinois	525	492	(93.7)	74	(15.0)
Indiana	109	109	(100.0)	22	(20.2)
Iowa	40	40	(100.0)	3	(7.5)
Kansas	81	81	(100.0)	17	(21.0)
Kentucky	84	84	(100.0)	11	(13.1)
Louisiana	194	179	(92.3)	36	(20.1)
Maine	15	15	(100.0)	3	(20.0)
Maryland	245	242	(98.8)	20	(8.3)
Massachusetts	251	245	(97.6)	20	(8.2)
Michigan	211	210	(99.5)	23	(11.0)
Minnesota	193	192	(99.5)	10	(5.2)
Mississippi	107	107	(100.0)	28	(26.2)
Missouri	92	91	(98.9)	13	(14.3)
Montana	13	13	(100.0)	4	(30.8)
Nebraska	22	22	(100.0)	1	(4.5)
Nevada	97	97	(100.0)	8	(8.2)
New Hampshire	15	15	(100.0)	2	(13.3)
New Jersey	492	491	(99.8)	31	(6.3)
New Mexico	46	44	(95.7)	11	(25.0)
New York State <sup>2</sup>	301	280	(93.0)	21	(7.5)
New York City	916	901	(98.4)	133	(14.8)
North Carolina	357	355	(99.4)	64	(18.0)
North Dakota	9	8	(88.9)	3	(37.5)
Ohio	223	221	(99.1)	27	(12.2)
Oklahoma	133	133	(100.0)	31	(23.3)
Oregon	77	74	(96.1)	6	(8.1)
Pennsylvania	307	302	(98.4)	19	(6.3)
Rhode Island	23	23	(100.0)	0	(0.0)
South Carolina	215	214	(99.5)	65	(30.4)
South Dakota	13	13	(100.0)	2	(15.4)
Tennessee	261	261	(100.0)	37	(14.2)
Texas	1,479	1,479	(100.0)	314	(21.2)
Utah	28	28	(100.0)	2	(7.1)
Vermont	8	8	(100.0)	0	(0.0)
Virginia	306	306	(100.0)	27	(8.8)
Washington	250	244	(97.6)	35	(14.3)
West Virginia	22	19	(86.4)	8	(42.1)
Wisconsin	64	63	(98.4)	10	(15.9)
Wyoming	4	4	(100.0)	1	(25.0)
American Samoa <sup>3</sup>	...	...	...	...	...
Fed. States of Micronesia <sup>3</sup>	54	48	(88.9)	1	(2.1)
Guam <sup>3</sup>	44	43	(97.7)	0	(0.0)
Marshall Islands <sup>3</sup>	24	23	(95.8)	1	(4.3)
N. Mariana Islands <sup>3</sup>	34	32	(94.1)	0	(0.0)
Puerto Rico <sup>3</sup>	104	104	(100.0)	19	(18.3)
Republic of Palau <sup>3</sup>	3	3	(100.0)	0	(0.0)
U.S. Virgin Islands <sup>3</sup>	...	...	...	...	...

<sup>1</sup>Excess alcohol use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

<sup>2</sup>Excludes New York City.

<sup>3</sup>Not included in U.S. totals. Data for 2006 are interim as of April 6, 2007.

**Note:** Ellipses indicate data not available.