

**Table 37. Completion of Tuberculosis Therapy (COT): 59 Reporting Areas, 2000**

Reporting Area	Total Cases	Therapy ≤1 Year Indicated <sup>1</sup>			Therapy >1 Year Indicated <sup>2</sup>		Overall	
		No. <sup>3</sup>	COT ≤1 Year(%)	COT(%)	No. <sup>3</sup>	COT(%)	No. <sup>3</sup>	COT(%)
<b>United States</b>	<b>16,313</b>	<b>14,263</b>	<b>80.2</b>	<b>91.5</b>	<b>230</b>	<b>78.7</b>	<b>14,493</b>	<b>91.3</b>
Alabama	310	263	87.8	97.3	3	100.0	266	97.4
Alaska	108	104	89.4	97.1	3	100.0	107	97.2
Arizona	261	228	77.6	88.6	2	50.0	230	88.3
Arkansas	199	176	84.1	91.5	0	...	176	91.5
California	3,292	2,936	77.9	90.1	54	75.9	2,990	89.9
Colorado	97	83	94.0	95.2	4	100.0	87	95.4
Connecticut	105	89	69.7	84.3	3	100.0	92	84.8
Delaware	28	23	82.6	87.0	2	100.0	25	88.0
District of Columbia	85	71	80.3	88.7	1	0.0	72	87.5
Florida	1,163	1,002	82.9	94.7	9	100.0	1,011	94.8
Georgia	696	600	80.5	89.8	10	90.0	610	89.8
Hawaii	136	120	72.5	90.8	3	100.0	123	91.1
Idaho	16	13	--	--	1	--	14	--
Illinois	738	636	83.5	91.4	10	70.0	646	91.0
Indiana	145	119	89.9	95.8	1	100.0	120	95.8
Iowa	40	38	89.5	94.7	0	...	38	94.7
Kansas	77	70	82.9	88.6	0	...	70	88.6
Kentucky	147	117	88.0	96.6	0	...	117	96.6
Louisiana	331	285	--	--	2	--	287	--
Maine	24	22	72.7	90.9	0	...	22	90.9
Maryland	283	255	77.6	87.8	1	100.0	256	87.9
Massachusetts	285	261	84.3	92.7	3	66.7	264	92.4
Michigan	287	248	83.5	91.9	5	80.0	253	91.7
Minnesota	178	172	83.7	91.9	1	100.0	173	91.9
Mississippi	173	153	83.0	92.8	2	0.0	155	91.6
Missouri	210	184	81.0	94.0	3	100.0	187	94.1
Montana	21	19	89.5	94.7	0	...	19	94.7
Nebraska	24	22	--	--	0	...	22	--
Nevada	97	89	--	--	0	...	89	--
New Hampshire	22	20	75.0	90.0	2	50.0	22	86.4
New Jersey	565	480	74.2	90.8	9	88.9	489	90.8
New Mexico	46	38	71.1	78.9	0	...	38	78.9
New York State <sup>4</sup>	409	358	77.7	91.1	4	75.0	362	90.9
New York City	1,309	1,122	84.0	93.7	27	66.7	1,149	93.0
North Carolina	447	391	90.8	95.4	3	66.7	394	95.2
North Dakota	5	5	100.0	100.0	0	...	5	100.0
Ohio	338	298	74.2	86.9	1	100.0	299	87.0
Oklahoma	153	128	82.8	92.2	1	100.0	129	92.2
Oregon	119	110	80.9	92.7	3	66.7	113	92.0
Pennsylvania	382	336	74.1	87.5	3	66.7	339	87.3
Rhode Island	49	43	74.4	88.4	0	...	43	88.4
South Carolina	286	245	76.7	90.6	2	100.0	247	90.7
South Dakota	16	12	75.0	91.7	0	...	12	91.7
Tennessee	383	323	80.5	95.7	2	100.0	325	95.7
Texas	1,496	1,324	78.8	92.4	32	81.3	1,356	92.2
Utah	49	41	70.7	87.8	1	100.0	42	88.1
Vermont	4	2	100.0	100.0	0	...	2	100.0
Virginia	292	254	85.8	92.5	11	90.9	265	92.5
Washington	258	233	82.0	95.3	4	75.0	237	94.9
West Virginia	33	24	50.0	83.3	0	...	24	83.3
Wisconsin	92	75	78.7	100.0	2	50.0	77	98.7
Wyoming	4	3	100.0	100.0	0	...	3	100.0
American Samoa <sup>5</sup>	...	...	...	...	...	...	...	...
Fed. States of Micronesia <sup>5</sup>	...	...	...	...	...	...	...	...
Guam <sup>5</sup>	54	48	--	--	1	--	49	--
N. Mariana Islands <sup>5</sup>	75	72	83.3	84.7	3	33.3	75	82.7
Puerto Rico <sup>5</sup>	174	128	87.5	95.3	1	0.0	129	94.6
Republic of Palau <sup>5</sup>	...	...	...	...	...	...	...	...
U.S. Virgin Islands <sup>5</sup>	...	...	...	...	...	...	...	...

<sup>1</sup>Initial isolate susceptible to rifampin (n=10,785) or susceptibility unknown (n=327); culture-negative (n=2,394); culture status unknown (n=756); age unknown (n=1).

<sup>2</sup>Initial isolate rifampin resistant, or pediatric (aged <15) case with meningeal, bone or joint, or miliary disease.

<sup>3</sup>Number of cases in persons alive at diagnosis, with initial drug regimen of one or more drugs prescribed, who did not die during therapy. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information on reason therapy stopped reported for ≥90% of cases.

<sup>4</sup>Excludes New York City.

<sup>5</sup>Not included in U.S. totals.

Ellipses indicate data not available.

**Note:** See Technical Notes (Appendix A) for description of COT calculation.