


**Figure 13.1. Percent of adults aged 18 years and over who experienced serious psychological distress during the past 30 days: United States, 1997–2003**




NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0-4) of the six items for each person are summed to yield a scale with a 0-to-24 range. A value of 13 or more for this scale is used here to define serious psychological distress (8). CI is confidence interval. Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-02 were recalculated using weights derived from the 2000 census. See "About This Release" and appendix tables in this release for more details.

DATA SOURCE: Sample Adult Core component of the 1997-2003 National Health Interview Surveys.

- In 2003, 3.1% (95% CI = 2.9%-3.4%) of adults aged 18 years and over experienced serious psychological distress during the past 30 days.
- The annual percent of adults who experienced serious psychological distress during the past 30 days declined significantly from 3.3% in 1997 to 2.4% in 1999, then increased from 2.4% in 1999 to 3.2% in 2001, and shows no trend in more recent years.

**Figure 13.2. Percent of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by age group and sex: United States, 2003**




NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0-4) of the six items for each person are summed to yield a scale with a 0-to-24 range. A value of 13 or more for this scale is used here to define serious psychological distress (8).

DATA SOURCE: Sample Adult Core component of the 2003 National Health Interview Survey.

- For both sexes combined, persons aged 45-64 years (3.9%) were more likely to have experienced serious psychological distress during the past 30 days than persons aged 18-44 years (2.9%) and 65 years and over (2.4%).
- For all age groups, women were more likely than men to have experienced serious psychological distress during the past 30 days.

**Figure 13.3. Age-sex-adjusted percent of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by race/ethnicity: United States, 2003**



NOTES: Six psychological distress questions are included in the Sample Adult Core component. These questions ask how often a respondent experienced certain symptoms of psychological distress during the past 30 days. The response codes (0-4) of the six items for each person are summed to yield a scale with a 0-to-24 range. A value of 13 or more for this scale is used here to define serious psychological distress (8). Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: 18-44 years, 45-64 years and 65 years and over.

DATA SOURCE: Sample Adult Core component of the 2003 National Health Interview Survey.

- The age-sex-adjusted prevalence of serious psychological distress was 3.9% for Hispanic persons, 3.0% for non-Hispanic white persons, and 3.1% for non-Hispanic black persons.
- Of the three race/ethnicity groups, Hispanic persons were most likely to have experienced serious psychological distress during the past 30 days.

## Data tables for figures 13.1-13.3:

**Data table for figure 13.1. Percent of adults aged 18 years and over who experienced serious psychological distress during the past 30 days: United States, 1997-2003**

Year	Percent	95% confidence interval
1997	3.3	3.1-3.5
1998	3.0	2.7-3.2
1999	2.4	2.2-2.6
2000	2.7	2.5-2.9
2001	3.2	2.9-3.4
2002	3.0	2.8-3.2
2003	3.1	2.9-3.4

**Data table for figure 13.2. Percent of adults aged 18 and over who experienced serious psychological distress during the past 30 days, by age group and sex: United States, 2003**

Age and sex	Percent	95% confidence interval
<b>18-44 years</b>		
Total	2.9	2.6-3.2
Men	2.0	1.7-2.4
Women	3.8	3.3-4.2
<b>45-64 years</b>		
Total	3.9	3.4-4.3
Men	2.7	2.2-3.3
Women	5.0	4.3-5.6
<b>65 years and over</b>		
Total	2.4	2.0-2.9
Men	1.7	1.1-2.4
Women	2.9	2.3-3.6

**Data table for figure 13.3. Age-sex-adjusted percent of adults aged 18 years and over who experienced serious psychological distress during the past 30 days, by race/ethnicity: United States, 2003**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
Hispanic or Latino	3.9	3.3-4.5
Not Hispanic or Latino		
White, single race	3.0	2.7-3.3
Black, single race	3.1	2.4-3.8

<sup>1</sup>Estimates are age-sex-adjusted to the 2000 projected U.S. standard population using three age groups: 18-44 years, 45-64 years, and 65 years and over.