

Wisconsin, 2005

Primary and secondary syphilis

- **CASES:** Wisconsin reported 41 cases in 2005.
- **RATES:** Wisconsin ranked 44 among 50 states + DC + 3 territories: 0.7 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Wisconsin, the rate among males was 1.4 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In Wisconsin, the race/ethnicity adjusted rates per 100,000 population were 0.6 among whites, 3.0 among blacks, 1.3 among Hispanics, 0.0 among Asian/Pacific Islanders, and 2.0 among American Indians/Alaska Natives.

- The rate among blacks was 5 times that of whites.
- The rate among Hispanics was 2.2 times that of whites.
- The rate among American Indians/Alaska Natives was 3.3 times that of whites.


Congenital syphilis

- **CASES:** Wisconsin reported 2 cases in 2005.
- **RATES:** Wisconsin ranked 22 among 28 states + 2 territories reporting at least one congenital syphilis case: 2.9 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Wisconsin, 1996-2005



Total P&S Syphilis Rates by County, Wisconsin, 2005



Rate (per 100,000 population)

