

TABLE 1. A summary of Atlantic and Gulf of Mexico marine mammal stock assessment reports for stocks of marine mammals under NMFS authority that occupy waters under U.S. jurisdiction. (A "Y" under the heading "SAR revised" indicates which 1999 stock assessment reports have been revised relative to the 1998 reports.)

| Species | Stock Area | SRG Region | NMFS Center | Nmin | Rmax | Fr | PBR | Total Annual Mort. | Annual Fish. Mort. | Strategic Status | SAR Revised |
|------------------------------|----------------------------|------------|-------------|--------------------|------|------|-------|--------------------|--------------------|------------------|-------------|
| Harbor seal | Western North Atlantic | ATL | NEC | 30,990 | 0.12 | 1.0 | 1,859 | 943 | 943 | N | Y |
| Gray seal | Western North Atlantic | ATL | NEC | NA | NA | NA | NA | 67 | 67 | N | Y |
| Harp seal | Western North Atlantic | ATL | NEC | N/A | N/A | N/A | N/A | 383 | 383 | N | Y |
| Hooded seal | Western North Atlantic | ATL | NEC | N/A | N/A | N/A | N/A | 5.6 | 5.6 | N | Y |
| Harbor porpoise | Gulf of Maine/Bay of Fundy | ATL | NEC | 48,289 | 0.04 | 0.5 | 483 | 1,850 | 1,850 | Y | Y |
| Risso's dolphin | Western North Atlantic | ATL | NEC | 11,140 | 0.04 | 0.48 | 107 | 7.4 | 7.4 | N | Y |
| Atlantic white-sided dolphin | Western North Atlantic | ATL | NEC | 19,196 | 0.04 | 0.48 | 184 | 287 | 287 | Y | Y |
| White-beaked dolphin | Western North Atlantic | ATL | NEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | N |
| Common dolphin | Western North Atlantic | ATL | NEC | 16,060 | 0.04 | 0.48 | 154 | 780 | 780 | Y | Y |
| Atlantic spotted dolphin | Western North Atlantic | ATL | NEC | 1,617 ¹ | 0.04 | 0.5 | 16 | 9.9 ² | 9.9 ² | N | Y |
| Pantropical spotted dolphin | Western North Atlantic | ATL | NEC | 1,617 ¹ | 0.04 | 0.5 | 16 | 9.9 ² | 9.9 ² | N | Y |
| Striped dolphin | Western North Atlantic | ATL | NEC | 18,220 | 0.04 | 0.5 | 182 | 11 | 11 | N | N |

| Species | Stock Area | SRG Region | NMFS Center | Nmin | Rmax | Fr | PBR | Total Annual Mort. | Annual Fish. Mort. | Strategic Status | SAR Revised |
|---|----------------------------------|------------|-------------|--------------------|-------|------|-----|--------------------|--------------------|------------------|-------------|
| Spinner dolphin | Western North Atlantic | ATL | NEC | N/A | N/A | N/A | N/A | 0.31 | 0.31 | N | N |
| Bottlenose dolphin | Western North Atlantic, offshore | ATL | NEC | 8,794 ³ | 0.04 | 0.5 | 88 | 10 | 10 | N | Y |
| Bottlenose dolphin | Western North Atlantic, coastal | ATL | SEC | 2,482 | 0.04 | 0.5 | 25 | 29 | 29 | Y | Y |
| Dwarf sperm whale | Western North Atlantic | ATL | SEC | N/A | 0.04 | N/A | N/A | 0.20 | 0.20 | N | Y |
| Pygmy sperm whale | Western North Atlantic | ATL | SEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | Y |
| Killer whale | Western North Atlantic | ATL | NEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | N |
| Pygmy killer whale | Western North Atlantic | ATL | SEC | 6 | 0.04 | 0.5 | 0.1 | 0.00 | 0.00 | N | N |
| Northern bottlenose whale | Western North Atlantic | ATL | NEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | N |
| Cuvier's beaked whale | Western North Atlantic | ATL | NEC | 895 ⁴ | 0.04 | 0.5 | 8.9 | 9.7 | 9.7 ⁵ | Y | Y |
| Mesoplodon beaked whale | Western North Atlantic | ATL | NEC | 895 ⁴ | 0.04 | 0.5 | 8.9 | 9.7 | 9.7 ⁵ | Y | Y |
| Pilot whale, long-finned (<i>Globicephala</i> sp.) | Western North Atlantic | ATL | NEC | 4,968 ⁶ | 0.04 | 0.45 | 45 | 40 | 40 ⁷ | N | Y |
| Pilot whale, short-finned | Western North Atlantic | ATL | NEC | 457 | 0.04 | 0.45 | 4.4 | 40 | 40 ⁷ | Y | Y |
| Sperm whale | North Atlantic | ATL | NEC | 1,617 | 0.04 | 0.1 | 3.2 | 0.00 | 0.00 | Y | Y |
| North Atlantic right whale | Western North Atlantic | ATL | NEC | 295 | 0.025 | 0.1 | 0.4 | 2.0 | 1.0 ⁸ | Y | Y |

| Species | Stock Area | SRG Region | NMFS Center | Nmin | Rmax | Fr | PBR | Total Annual Mort. | Annual Fish. Mort. | Strategic Status | SAR Revised |
|--------------------|--|------------|-------------|--------|-------|-----|------|--------------------|--------------------|------------------|-------------|
| Humpback whale | Western North Atlantic | ATL | NEC | 10,019 | 0.065 | 0.1 | 32.6 | 4.4 | 3.2 ⁹ | Y | Y |
| Fin whale | Western North Atlantic | ATL | NEC | 1,803 | 0.04 | 0.1 | 3.6 | 0.6 | 0.20 | Y | Y |
| Sei whale | Nova Scotia | ATL | NEC | N/A | 0.04 | 0.1 | N/A | 0.00 | 0.00 | Y | Y |
| Minke whale | Canadian east coast | ATL | NEC | 2,145 | 0.04 | 0.5 | 21 | 5.8 | 4.8 | N | Y |
| Blue whale | Western North Atlantic | ATL | NEC | 308 | 0.04 | 0.1 | 0.6 | 0.00 | 0.00 | Y | Y |
| Bottlenose dolphin | Gulf of Mexico bay, sound, and estuarine | ATL | SEC | 3,933 | 0.04 | 0.5 | 39.7 | N/A | N/A | Y | Y |
| Dwarf sperm whale | Northern Gulf of Mexico | ATL | SEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | Y |
| Pygmy sperm whale | Northern Gulf of Mexico | ATL | SEC | N/A | 0.04 | N/A | N/A | 0.00 | 0.00 | N | Y |

1. This value includes either or both of *Stenella frontalis* or *Stenella attenuata*.
2. Mortality data are not separated by species; therefore, species-specific estimates are not available. The mortality estimate represents both Atlantic and Pantropical spotted dolphins.
3. Estimates may include sightings of the coastal form.
4. This estimate includes Cuvier's beaked whales and *Mesoplodon* spp. beaked whales.
5. This is the average mortality of beaked whales (*Mesoplodon* spp.) based on 5 years of observer data. This annual mortality rate includes an unknown number of Cuvier's beaked whales.
6. This estimate may include both long-finned and short-finned pilot whales.
7. Mortality data are not separated by species; therefore, species-specific estimates are not available. This mortality estimate represents both long-finned and short-finned pilot whales.
8. This is the average mortality of right whales based on 5 years of observer data (0.0) and additional fishery impact records (1.0).
9. This is the average mortality of humpback whales based on 5 years of observer data (0.6) and additional fishery impact records (2.6).