

Table 33. Tuberculosis Cases and Percentages by Noninjecting Drug Use,¹ Age ≥15: Reporting Areas, 2005

Reporting Area	Total Cases	Cases with Information on Noninjecting Drug Use		Cases Reporting Noninjecting Drug Use	
		No.	(%)	No.	(%)
United States	13,234	12,915	(97.6)	1,004	(7.8)
Alabama	201	201	(100.0)	8	(4.0)
Alaska	52	52	(100.0)	2	(3.8)
Arizona	244	225	(92.2)	20	(8.9)
Arkansas	105	105	(100.0)	0	(0.0)
California	2,753	2,683	(97.5)	165	(6.1)
Colorado	84	81	(96.4)	5	(6.2)
Connecticut	93	93	(100.0)	5	(5.4)
Delaware	25	25	(100.0)	0	(0.0)
District of Columbia	55	55	(100.0)	4	(7.3)
Florida	1,037	1,027	(99.0)	145	(14.1)
Georgia	463	446	(96.3)	54	(12.1)
Hawaii	109	70	(64.2)	—	—
Idaho	21	17	(81.0)	1	(5.9)
Illinois	550	498	(90.5)	50	(10.0)
Indiana	137	137	(100.0)	21	(15.3)
Iowa	51	51	(100.0)	0	(0.0)
Kansas	57	57	(100.0)	4	(7.0)
Kentucky	118	118	(100.0)	8	(6.8)
Louisiana	241	224	(92.9)	44	(19.6)
Maine	16	16	(100.0)	0	(0.0)
Maryland	266	266	(100.0)	11	(4.1)
Massachusetts	252	246	(97.6)	7	(2.8)
Michigan	238	236	(99.2)	13	(5.5)
Minnesota	179	178	(99.4)	3	(1.7)
Mississippi	100	100	(100.0)	9	(9.0)
Missouri	105	103	(98.1)	4	(3.9)
Montana	10	10	(100.0)	1	(10.0)
Nebraska	35	35	(100.0)	2	(5.7)
Nevada	107	107	(100.0)	6	(5.6)
New Hampshire	4	4	(100.0)	0	(0.0)
New Jersey	461	460	(99.8)	32	(7.0)
New Mexico	39	33	(84.6)	1	(3.0)
New York State ²	287	286	(99.7)	11	(3.8)
New York City	943	897	(95.1)	85	(9.5)
North Carolina	309	307	(99.4)	42	(13.7)
North Dakota	6	4	(66.7)	—	—
Ohio	241	239	(99.2)	25	(10.5)
Oklahoma	125	125	(100.0)	14	(11.2)
Oregon	99	94	(94.9)	8	(8.5)
Pennsylvania	312	312	(100.0)	7	(2.2)
Rhode Island	44	44	(100.0)	0	(0.0)
South Carolina	235	233	(99.1)	28	(12.0)
South Dakota	16	16	(100.0)	1	(6.3)
Tennessee	277	277	(100.0)	37	(13.4)
Texas	1,424	1,423	(99.9)	96	(6.7)
Utah	25	25	(100.0)	0	(0.0)
Vermont	7	7	(100.0)	1	(14.3)
Virginia	330	330	(100.0)	5	(1.5)
Washington	247	240	(97.2)	11	(4.6)
West Virginia	27	27	(100.0)	3	(11.1)
Wisconsin	72	70	(97.2)	5	(7.1)
Wyoming	0	0	—	—	—
American Samoa ³	5	5	(100.0)	0	(0.0)
Fed. States of Micronesia ³	63	27	(42.9)	—	—
Guam ³	53	51	(96.2)	1	(2.0)
Marshall Islands ³	60	60	(100.0)	2	(3.3)
N. Mariana Islands ³	55	55	(100.0)	0	(0.0)
Puerto Rico ³	108	108	(100.0)	21	(19.4)
Republic of Palau ³	9	9	(100.0)	0	(0.0)
U.S. Virgin Islands ³

¹Noninjecting drug use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.