

Table HIST002_2. Deaths for 113 selected causes by 5-year age groups, race and sex: United States, 1979-98

Plate 2 of 2 Age groups 45-49 years to 85+ years

The mortality trend data in this table are classified by the Ninth Revision, International Classification of Diseases (ICD-9) and sorted into the list of 113 selected causes of death developed for the Tenth Revision (ICD-10). ICD-10 was introduced with 1999 mortality data. Data in these tables were constructed to be most comparable with the ICD-9 cause-of-death titles used in the list of 113 selected causes of death. These data are the most appropriate ICD-9 data to use with [published comparability ratios](#) describing the discontinuities between ICD-9 and ICD-10. Used in conjunction with the comparability ratios, this table provides trend data comparable to that published for data year 1999 and beyond. Although in some cases categories from the list of 113 selected causes are identical to those in the old list of 72 selected causes of death used with ICD-9, it is important to note that many of these categories are not comparable with categories in the list of 72 selected causes even though the cause-of-death titles may be the same.

Description of symbols used in this table:

- Quantity zero**
- Category not applicable**

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
All races, both sexes											
1998	2,337,256	65,410	81,069	100,645	133,079	189,780	269,202	329,463	352,200	612,575	411
1997	2,314,245	65,090	79,792	98,130	133,863	194,776	269,498	325,799	344,731	594,068	401
1996	2,314,690	67,032	77,297	96,726	136,999	200,045	273,849	321,223	342,067	576,541	523
1995	2,312,132	65,623	77,377	96,641	138,871	204,347	276,543	315,516	336,661	561,259	463
1994	2,278,994	62,517	75,380	95,897	141,222	208,394	275,275	310,348	329,866	538,690	414
1993	2,268,553	59,270	72,545	96,333	145,248	213,366	274,453	312,774	325,264	528,437	507
1992	2,175,613	56,533	68,497	94,582	146,409	211,071	266,845	301,736	308,116	487,446	474
1991	2,169,518	53,627	67,049	96,553	151,525	214,468	264,168	301,822	305,668	477,401	576
1990	2,148,463	51,856	66,752	97,865	154,831	217,333	260,616	301,114	300,325	463,105	562
1989	2,150,466	50,857	67,409	101,474	158,584	219,097	262,127	301,225	297,981	457,358	562
1988	2,167,999	49,684	67,788	105,363	164,386	221,504	267,041	302,580	299,334	459,710	587
1987	2,123,323	47,613	68,307	107,583	165,741	220,110	265,993	296,714	287,621	439,248	582
1986	2,105,361	46,415	68,746	110,323	168,706	218,649	266,890	293,520	281,629	427,473	999
1985	2,086,440	46,254	70,380	115,192	171,288	216,749	265,897	290,149	278,699	419,051	877
1984	2,039,369	45,723	71,490	116,975	170,380	214,732	261,838	281,892	269,020	399,466	524
1983	2,019,201	45,536	74,357	119,779	169,161	214,916	259,464	277,591	263,627	389,363	623
1982	1,974,797	46,412	76,677	119,833	167,181	213,426	252,394	268,121	253,838	366,727	666
1981	1,977,981	48,546	80,687	123,693	166,316	213,890	250,758	261,631	253,018	362,499	614
1980	1,989,841	49,787	83,370	126,016	166,165	216,277	250,344	263,611	253,646	357,970	568
1979	1,913,841	50,779	84,486	123,622	163,344	209,007	240,248	251,822	241,854	328,725	652
All causes											
All races, male											
1998	1,157,260	41,707	50,020	60,895	79,239	110,193	149,150	171,426	162,786	199,057	324
1997	1,154,039	41,447	49,140	59,184	79,692	113,790	150,085	169,810	159,581	194,161	317
1996	1,163,569	43,056	47,815	58,974	82,196	116,660	152,513	167,377	157,882	187,719	373
1995	1,172,959	42,609	48,313	58,848	83,442	119,415	154,586	164,372	155,036	182,823	331
1994	1,162,747	40,224	47,235	58,691	85,497	122,008	154,231	161,437	151,492	176,102	311
1993	1,161,797	38,084	45,364	59,129	87,777	125,445	154,161	163,677	149,882	172,778	354
1992	1,122,336	36,622	42,649	58,083	88,797	124,228	149,937	158,257	141,640	161,236	356
1991	1,121,665	34,363	41,665	59,342	92,094	126,381	149,475	158,268	140,682	156,823	437
1990	1,113,417	33,097	41,557	60,527	94,119	128,356	147,711	157,873	137,533	151,897	403
1989	1,114,190	32,718	42,105	62,981	96,628	129,847	148,559	157,090	135,580	149,735	405
1988	1,125,540	31,597	42,318	65,194	100,174	131,931	151,970	158,662	136,059	151,559	417
1987	1,107,958	30,122	42,735	67,093	101,309	131,122	151,935	155,809	131,142	145,331	409
1986	1,104,005	29,361	42,978	69,198	103,133	130,613	152,433	154,284	128,547	142,953	533
1985	1,097,758	29,181	44,139	72,223	105,488	130,265	152,752	152,649	127,223	141,653	491
1984	1,076,514	28,805	45,026	73,197	105,419	129,211	150,167	148,140	122,297	136,799	353
1983	1,071,923	28,707	46,691	75,130	104,678	130,217	149,457	146,886	119,875	134,651	412
1982	1,056,440	29,342	48,665	75,332	103,990	129,819	145,539	141,221	114,716	128,564	415
1981	1,063,772	30,732	51,138	78,042	103,843	130,290	145,260	138,021	114,500	127,874	425
1980	1,075,078	31,469	53,003	80,075	104,187	132,459	144,912	138,208	114,557	128,134	394
1979	1,044,959	32,366	53,837	78,814	103,474	129,066	140,105	132,212	110,104	119,006	458

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
All races, female											
1998	1,179,996	23,703	31,049	39,750	53,840	79,587	120,052	158,037	189,414	413,518	87
1997	1,160,206	23,643	30,652	38,946	54,171	80,986	119,413	155,989	185,150	399,907	84
1996	1,151,121	23,976	29,482	37,752	54,803	83,385	121,336	153,846	184,185	388,822	150
1995	1,139,173	23,014	29,064	37,793	55,429	84,932	121,957	151,144	181,625	378,436	132
1994	1,116,247	22,293	28,145	37,206	55,725	86,386	121,044	148,911	178,374	362,588	103
1993	1,106,756	21,186	27,181	37,204	57,471	87,921	120,292	149,097	175,382	355,659	153
1992	1,053,277	19,911	25,848	36,499	57,612	86,843	116,908	143,479	166,476	326,210	118
1991	1,047,853	19,264	25,384	37,211	59,431	88,087	114,693	143,554	164,986	320,578	139
1990	1,035,046	18,759	25,195	37,338	60,712	88,977	112,905	143,241	162,792	311,208	159
1989	1,036,276	18,139	25,304	38,493	61,956	89,250	113,568	144,135	162,401	307,623	157
1988	1,042,459	18,087	25,470	40,169	64,212	89,573	115,071	143,918	163,275	308,151	170
1987	1,015,365	17,491	25,572	40,490	64,432	88,988	114,058	140,905	156,479	293,917	173
1986	1,001,356	17,054	25,768	41,125	65,573	88,036	114,457	139,236	153,082	284,520	466
1985	988,682	17,073	26,241	42,969	65,800	86,484	113,145	137,500	151,476	277,398	386
1984	962,855	16,918	26,464	43,778	64,961	85,521	111,671	133,752	146,723	262,667	171
1983	947,278	16,829	27,666	44,649	64,483	84,699	110,007	130,705	143,752	254,712	211
1982	918,357	17,070	28,012	44,501	63,191	83,607	106,855	126,900	139,122	238,163	251
1981	914,209	17,814	29,549	45,651	62,473	83,600	105,498	123,610	138,518	234,625	189
1980	914,763	18,318	30,367	45,941	61,978	83,818	105,432	125,403	139,089	229,836	174
1979	868,882	18,413	30,649	44,808	59,870	79,941	100,143	119,610	131,750	209,719	194
All causes											
White, both sexes											
1998	2,015,984	48,731	63,425	80,555	108,757	160,753	233,911	292,762	319,419	561,836	310
1997	1,996,393	48,511	62,893	78,397	109,450	165,581	234,217	290,591	313,012	545,264	301
1996	1,992,966	50,239	60,640	77,494	112,647	170,389	238,485	286,315	310,103	529,736	384
1995	1,987,437	49,275	60,509	77,317	114,299	174,015	241,309	281,834	304,570	515,856	352
1994	1,959,875	47,185	59,042	76,885	117,431	178,047	240,705	277,518	298,482	496,083	296
1993	1,951,437	45,103	56,767	77,318	120,918	182,643	240,176	279,742	294,280	486,211	339
1992	1,873,781	43,451	53,689	75,750	122,213	180,788	234,117	270,238	279,507	448,984	351
1991	1,868,904	41,199	52,454	78,133	127,160	183,809	232,010	269,816	276,797	439,797	433
1990	1,853,254	39,788	52,364	79,196	129,901	186,667	229,536	269,085	272,644	426,982	384
1989	1,853,841	38,963	52,670	82,414	133,021	187,834	230,704	268,221	270,882	421,669	384
1988	1,876,906	38,143	53,296	86,289	138,357	190,770	235,866	269,922	272,232	425,306	426
1987	1,843,067	36,709	53,975	88,913	140,201	190,472	234,828	265,687	261,500	406,652	409
1986	1,831,083	35,906	54,471	91,491	142,976	189,952	235,871	262,791	256,545	395,825	812
1985	1,819,054	35,864	56,340	95,543	145,694	188,473	234,483	260,372	253,582	389,513	702
1984	1,781,897	35,526	57,651	97,365	145,262	187,563	230,931	253,910	244,706	372,354	372
1983	1,765,582	35,482	60,214	99,854	144,273	187,966	228,799	250,004	240,532	363,410	419
1982	1,729,085	36,257	62,428	100,325	143,338	186,977	223,610	242,120	232,390	342,651	474
1981	1,731,233	37,980	65,786	103,698	142,992	187,095	222,350	236,476	231,851	338,658	407
1980	1,738,607	38,886	67,748	105,852	142,793	188,394	221,932	236,997	233,370	334,240	393
1979	1,676,145	40,024	68,970	104,028	141,277	182,308	213,718	226,584	224,321	307,964	477

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
White, male											
1998	990,190	31,547	39,365	49,142	65,269	94,345	130,959	153,165	148,428	181,696	251
1997	986,884	31,226	38,974	47,548	65,755	97,752	131,749	152,257	145,618	177,227	249
1996	991,984	32,574	37,785	47,550	68,138	100,389	133,921	149,924	143,738	171,449	277
1995	997,277	32,221	37,939	47,443	69,377	102,592	135,885	147,548	140,862	166,772	256
1994	988,823	30,564	37,138	47,535	71,793	105,210	135,694	145,252	137,367	161,104	225
1993	988,329	29,159	35,704	47,806	73,750	108,341	135,772	147,381	135,805	157,645	254
1992	956,957	28,343	33,681	47,042	74,994	107,427	132,273	142,422	128,484	147,419	272
1991	956,497	26,506	32,815	48,337	78,173	109,220	132,362	142,329	127,340	143,266	325
1990	950,812	25,485	32,805	49,456	79,857	111,237	130,930	141,699	124,811	138,807	277
1989	950,852	25,166	32,966	51,499	81,919	112,194	131,599	140,663	123,249	136,813	281
1988	965,419	24,324	33,489	53,910	85,141	114,474	135,187	142,069	123,668	139,042	312
1987	953,382	23,333	34,078	56,006	86,542	114,258	135,122	139,916	119,096	133,284	289
1986	952,554	22,869	34,352	57,922	88,223	114,255	135,671	138,505	116,918	131,147	420
1985	950,455	22,707	35,640	60,582	90,456	114,175	135,744	137,460	115,596	130,583	377
1984	934,529	22,517	36,608	61,493	90,573	113,733	133,421	133,916	111,144	126,344	254
1983	931,779	22,534	38,248	63,255	89,997	114,930	132,775	132,738	109,081	124,677	282
1982	919,239	23,059	40,051	63,681	89,933	114,655	129,707	127,905	104,832	119,069	292
1981	925,490	24,153	42,071	66,011	90,093	114,982	129,666	125,170	104,633	118,475	285
1980	933,878	24,816	43,490	67,809	90,363	116,502	129,502	124,611	105,008	118,549	285
1979	910,137	25,752	44,334	66,724	90,438	113,753	125,620	119,296	101,651	110,591	333
All causes											
White, female											
1998	1,025,794	17,184	24,060	31,413	43,488	66,408	102,952	139,597	170,991	380,140	59
1997	1,009,509	17,285	23,919	30,849	43,695	67,829	102,468	138,334	167,394	368,037	52
1996	1,000,982	17,665	22,855	29,944	44,509	70,000	104,564	136,391	166,365	358,287	107
1995	990,160	17,054	22,570	29,874	44,922	71,423	105,424	134,286	163,708	349,084	96
1994	971,052	16,621	21,904	29,350	45,638	72,837	105,011	132,266	161,115	334,979	71
1993	963,108	15,944	21,063	29,512	47,168	74,302	104,404	132,361	158,475	328,566	85
1992	916,824	15,108	20,008	28,708	47,219	73,361	101,844	127,816	151,023	301,565	79
1991	912,407	14,693	19,639	29,796	48,987	74,589	99,648	127,487	149,457	296,531	108
1990	902,442	14,303	19,559	29,740	50,044	75,430	98,606	127,386	147,833	288,175	107
1989	902,989	13,797	19,704	30,915	51,102	75,640	99,105	127,558	147,633	284,856	103
1988	911,487	13,819	19,807	32,379	53,216	76,296	100,679	127,853	148,564	286,264	114
1987	889,685	13,376	19,897	32,907	53,659	76,214	99,706	125,771	142,404	273,368	120
1986	878,529	13,037	20,119	33,569	54,753	75,697	100,200	124,286	139,627	264,678	392
1985	868,599	13,157	20,700	34,961	55,238	74,298	98,739	122,912	137,986	258,930	325
1984	847,368	13,009	21,043	35,872	54,689	73,830	97,510	119,994	133,562	246,010	118
1983	833,803	12,948	21,966	36,599	54,276	73,036	96,024	117,266	131,451	238,733	137
1982	809,846	13,198	22,377	36,644	53,405	72,322	93,903	114,215	127,558	223,582	182
1981	805,743	13,827	23,715	37,687	52,899	72,113	92,684	111,306	127,218	220,183	122
1980	804,729	14,070	24,258	38,043	52,430	71,892	92,430	112,386	128,362	215,691	108
1979	766,008	14,272	24,636	37,304	50,839	68,555	88,098	107,288	122,670	197,373	144

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
All other, both sexes											
1998	321,272	16,679	17,644	20,090	24,322	29,027	35,291	36,701	32,781	50,739	101
1997	317,852	16,579	16,899	19,733	24,413	29,195	35,281	35,208	31,719	48,804	100
1996	321,724	16,793	16,657	19,232	24,352	29,656	35,364	34,908	31,964	46,805	139
1995	324,695	16,348	16,868	19,324	24,572	30,332	35,234	33,682	32,091	45,403	111
1994	319,119	15,332	16,338	19,012	23,791	30,347	34,570	32,830	31,384	42,607	118
1993	317,116	14,167	15,778	19,015	24,330	30,723	34,277	33,032	30,984	42,226	168
1992	301,832	13,082	14,808	18,832	24,196	30,283	32,728	31,498	28,609	38,462	123
1991	300,614	12,428	14,595	18,420	24,365	30,659	32,158	32,006	28,871	37,604	143
1990	295,209	12,068	14,388	18,669	24,930	30,666	31,080	32,029	27,681	36,123	178
1989	296,625	11,894	14,739	19,060	25,563	31,263	31,423	33,004	27,099	35,689	178
1988	291,093	11,541	14,492	19,074	26,029	30,734	31,175	32,658	27,102	34,404	161
1987	280,256	10,904	14,332	18,670	25,540	29,638	31,165	31,027	26,121	32,596	173
1986	274,278	10,509	14,275	18,832	25,730	28,697	31,019	30,729	25,084	31,648	187
1985	267,386	10,390	14,040	19,649	25,594	28,276	31,414	29,777	25,117	29,538	175
1984	257,472	10,197	13,839	19,610	25,118	27,169	30,907	27,982	24,314	27,112	152
1983	253,619	10,054	14,143	19,925	24,888	26,950	30,665	27,587	23,095	25,953	204
1982	245,712	10,155	14,249	19,508	23,843	26,449	28,784	26,001	21,448	24,076	192
1981	246,748	10,566	14,901	19,995	23,324	26,795	28,408	25,155	21,167	23,841	207
1980	251,234	10,901	15,622	20,164	23,372	27,883	28,412	26,614	20,276	23,730	175
1979	237,696	10,755	15,516	19,594	22,067	26,699	26,530	25,238	17,533	20,761	175
All causes											
All other, male											
1998	167,070	10,160	10,655	11,753	13,970	15,848	18,191	18,261	14,358	17,361	73
1997	167,155	10,221	10,166	11,636	13,937	16,038	18,336	17,553	13,963	16,934	68
1996	171,585	10,482	10,030	11,424	14,058	16,271	18,592	17,453	14,144	16,270	96
1995	175,682	10,388	10,374	11,405	14,065	16,823	18,701	16,824	14,174	16,051	75
1994	173,924	9,660	10,097	11,156	13,704	16,798	18,537	16,185	14,125	14,998	86
1993	173,468	8,925	9,660	11,323	14,027	17,104	18,389	16,296	14,077	15,133	100
1992	165,379	8,279	8,968	11,041	13,803	16,801	17,664	15,835	13,156	13,817	84
1991	165,168	7,857	8,850	11,005	13,921	17,161	17,113	15,939	13,342	13,557	112
1990	162,605	7,612	8,752	11,071	14,262	17,119	16,781	16,174	12,722	13,090	126
1989	163,338	7,552	9,139	11,482	14,709	17,653	16,960	16,427	12,331	12,922	124
1988	160,121	7,273	8,829	11,284	15,033	17,457	16,783	16,593	12,391	12,517	105
1987	154,576	6,789	8,657	11,087	14,767	16,864	16,813	15,893	12,046	12,047	120
1986	151,451	6,492	8,626	11,276	14,910	16,358	16,762	15,779	11,629	11,806	113
1985	147,303	6,474	8,499	11,641	15,032	16,090	17,008	15,189	11,627	11,070	114
1984	141,985	6,288	8,418	11,704	14,846	15,478	16,746	14,224	11,153	10,455	99
1983	140,144	6,173	8,443	11,875	14,681	15,287	16,682	14,148	10,794	9,974	130
1982	137,201	6,283	8,614	11,651	14,057	15,164	15,832	13,316	9,884	9,495	123
1981	138,282	6,579	9,067	12,031	13,750	15,308	15,594	12,851	9,867	9,399	140
1980	141,200	6,653	9,513	12,266	13,824	15,957	15,410	13,597	9,549	9,585	109
1979	134,822	6,614	9,503	12,090	13,036	15,313	14,485	12,916	8,453	8,415	125

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
All other, female											
1998	154,202	6,519	6,989	8,337	10,352	13,179	17,100	18,440	18,423	33,378	28
1997	150,697	6,358	6,733	8,097	10,476	13,157	16,945	17,655	17,756	31,870	32
1996	150,139	6,311	6,627	7,808	10,294	13,385	16,772	17,455	17,820	30,535	43
1995	149,013	5,960	6,494	7,919	10,507	13,509	16,533	16,858	17,917	29,352	36
1994	145,195	5,672	6,241	7,856	10,087	13,549	16,033	16,645	17,259	27,609	32
1993	143,648	5,242	6,118	7,692	10,303	13,619	15,888	16,736	16,907	27,093	68
1992	136,453	4,803	5,840	7,791	10,393	13,482	15,064	15,663	15,453	24,645	39
1991	135,446	4,571	5,745	7,415	10,444	13,498	15,045	16,067	15,529	24,047	31
1990	132,604	4,456	5,636	7,598	10,668	13,547	14,299	15,855	14,959	23,033	52
1989	133,287	4,342	5,600	7,578	10,854	13,610	14,463	16,577	14,768	22,767	54
1988	130,972	4,268	5,663	7,790	10,996	13,277	14,392	16,065	14,711	21,887	56
1987	125,680	4,115	5,675	7,583	10,773	12,774	14,352	15,134	14,075	20,549	53
1986	122,827	4,017	5,649	7,556	10,820	12,339	14,257	14,950	13,455	19,842	74
1985	120,083	3,916	5,541	8,008	10,562	12,186	14,406	14,588	13,490	18,468	61
1984	115,487	3,909	5,421	7,906	10,272	11,691	14,161	13,758	13,161	16,657	53
1983	113,475	3,881	5,700	8,050	10,207	11,663	13,983	13,439	12,301	15,979	74
1982	108,511	3,872	5,635	7,857	9,786	11,285	12,952	12,685	11,564	14,581	69
1981	108,466	3,987	5,834	7,964	9,574	11,487	12,814	12,304	11,300	14,442	67
1980	110,034	4,248	6,109	7,898	9,548	11,926	13,002	13,017	10,727	14,145	66
1979	102,874	4,141	6,013	7,504	9,031	11,386	12,045	12,322	9,080	12,346	50
All causes											
Black, both sexes											
1998	278,440	14,846	15,419	17,596	21,217	25,173	30,434	31,448	27,967	43,341	93
1997	276,520	14,737	14,877	17,253	21,391	25,288	30,613	30,212	27,115	42,115	90
1996	282,089	15,009	14,762	16,962	21,497	25,960	30,789	30,236	27,462	40,534	121
1995	286,401	14,705	14,914	17,077	21,691	26,788	30,917	29,435	27,798	39,502	101
1994	282,379	13,752	14,519	16,849	21,101	26,818	30,437	28,746	27,360	37,202	95
1993	282,151	12,726	14,192	16,932	21,663	27,401	30,471	29,122	27,132	37,011	141
1992	269,219	11,753	13,252	16,727	21,669	27,011	29,124	27,875	25,260	33,856	114
1991	269,525	11,229	13,135	16,536	21,912	27,578	28,860	28,475	25,707	33,110	135
1990	265,498	10,920	12,951	16,852	22,621	27,665	28,022	28,681	24,756	32,047	166
1989	267,642	10,781	13,329	17,258	23,262	28,452	28,507	29,815	24,205	31,725	167
1988	264,019	10,481	13,151	17,384	23,769	28,084	28,377	29,622	24,399	30,875	147
1987	254,814	9,933	13,068	16,966	23,540	27,268	28,421	28,242	23,687	29,222	150
1986	250,326	9,615	13,033	17,395	23,789	26,604	28,532	28,086	22,696	28,517	165
1985	244,207	9,530	12,956	18,142	23,694	26,166	28,866	27,178	22,822	26,675	158
1984	235,884	9,352	12,746	18,144	23,346	25,207	28,649	25,681	22,198	24,439	137
1983	233,124	9,230	13,098	18,538	23,273	25,095	28,452	25,448	21,166	23,533	156
1982	226,513	9,320	13,211	18,256	22,308	24,681	26,844	23,978	19,710	21,875	167
1981	228,560	9,766	13,909	18,803	21,974	25,141	26,608	23,300	19,579	21,833	177
1980	233,135	10,171	14,645	18,970	22,100	26,232	26,591	24,698	18,719	21,649	165
1979	220,818	10,082	14,564	18,483	20,819	25,168	24,889	23,457	16,117	18,944	160

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All causes											
Black, male											
1998	143,417	9,045	9,315	10,260	12,130	13,714	15,560	15,365	11,951	13,964	67
1997	144,110	9,114	8,953	10,086	12,179	13,893	15,767	14,874	11,597	13,767	62
1996	149,472	9,434	8,963	10,056	12,382	14,226	16,015	14,927	11,866	13,340	81
1995	154,175	9,410	9,254	10,082	12,463	14,850	16,217	14,533	11,958	13,250	69
1994	153,019	8,683	9,004	9,899	12,169	14,806	16,129	14,017	11,994	12,424	69
1993	153,502	8,050	8,746	10,152	12,531	15,233	16,188	14,101	11,891	12,671	82
1992	146,630	7,493	8,021	9,824	12,380	14,946	15,580	13,782	11,253	11,646	76
1991	147,331	7,149	7,996	9,915	12,535	15,362	15,246	13,964	11,453	11,498	106
1990	145,359	6,931	7,888	10,055	12,938	15,350	15,000	14,235	11,058	11,236	118
1989	146,393	6,857	8,317	10,446	13,340	16,031	15,209	14,555	10,644	11,056	115
1988	144,228	6,641	8,040	10,280	13,727	15,850	15,139	14,811	10,811	10,818	96
1987	139,551	6,189	7,925	10,096	13,590	15,475	15,177	14,165	10,600	10,506	101
1986	137,214	5,948	7,882	10,460	13,748	15,074	15,246	14,147	10,195	10,377	104
1985	133,610	5,931	7,890	10,759	13,898	14,842	15,468	13,533	10,271	9,780	106
1984	129,147	5,781	7,756	10,857	13,785	14,315	15,322	12,739	9,909	9,225	90
1983	127,911	5,710	7,819	11,066	13,650	14,138	15,261	12,804	9,673	8,918	102
1982	125,610	5,784	8,007	10,941	13,082	14,088	14,590	11,973	8,881	8,466	106
1981	127,296	6,109	8,474	11,287	12,912	14,290	14,373	11,713	8,925	8,468	116
1980	130,138	6,219	8,937	11,498	13,036	14,904	14,191	12,377	8,669	8,534	103
1979	124,433	6,220	8,928	11,361	12,253	14,327	13,404	11,776	7,658	7,522	115
All causes											
Black, female											
1998	135,023	5,801	6,104	7,336	9,087	11,459	14,874	16,083	16,016	29,377	26
1997	132,410	5,623	5,924	7,167	9,212	11,395	14,846	15,338	15,518	28,348	28
1996	132,617	5,575	5,799	6,906	9,115	11,734	14,774	15,309	15,596	27,194	40
1995	132,226	5,295	5,660	6,995	9,228	11,938	14,700	14,902	15,840	26,252	32
1994	129,360	5,069	5,515	6,950	8,932	12,012	14,308	14,729	15,366	24,778	26
1993	128,649	4,676	5,446	6,780	9,132	12,168	14,283	15,021	15,241	24,340	59
1992	122,589	4,260	5,231	6,903	9,289	12,065	13,544	14,093	14,007	22,210	38
1991	122,194	4,080	5,139	6,621	9,377	12,216	13,614	14,511	14,254	21,612	29
1990	120,139	3,989	5,063	6,797	9,683	12,315	13,022	14,446	13,698	20,811	48
1989	121,249	3,924	5,012	6,812	9,922	12,421	13,298	15,260	13,561	20,669	52
1988	119,791	3,840	5,111	7,104	10,042	12,234	13,238	14,811	13,588	20,057	51
1987	115,263	3,744	5,143	6,870	9,950	11,793	13,244	14,077	13,087	18,716	49
1986	113,112	3,667	5,151	6,935	10,041	11,530	13,286	13,939	12,501	18,140	61
1985	110,597	3,599	5,066	7,383	9,796	11,324	13,398	13,645	12,551	16,895	52
1984	106,737	3,571	4,990	7,287	9,561	10,892	13,327	12,942	12,289	15,214	47
1983	105,213	3,520	5,279	7,472	9,623	10,957	13,191	12,644	11,493	14,615	54
1982	100,903	3,536	5,204	7,315	9,226	10,593	12,254	12,005	10,829	13,409	61
1981	101,264	3,657	5,435	7,516	9,062	10,851	12,235	11,587	10,654	13,365	61
1980	102,997	3,952	5,708	7,472	9,064	11,328	12,400	12,321	10,050	13,115	62
1979	96,385	3,862	5,636	7,122	8,566	10,841	11,485	11,681	8,459	11,422	45

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
All races, both sexes											
1998	37	-	2	3	1	4	4	5	7	7	-
1997	51	2	4	-	2	2	1	6	12	8	-
1996	59	1	1	3	5	3	8	5	4	12	-
1995	66	4	2	1	4	5	9	11	10	8	-
1994	50	2	2	2	3	5	4	3	10	6	-
1993	52	6	1	3	1	4	4	8	7	6	-
1992	47	1	-	1	2	4	9	3	8	8	-
1991	54	2	2	3	6	4	4	3	5	15	-
1990	81	2	4	4	8	7	7	15	10	11	-
1989	99	4	2	2	6	12	11	10	14	18	-
1988	66	1	5	2	6	1	6	13	6	13	-
1987	107	5	1	3	7	8	14	16	10	27	-
1986	104	1	3	4	8	11	10	16	14	14	-
1985	117	1	6	5	12	7	18	17	13	17	-
1984	90	7	3	3	5	3	16	13	7	19	-
1983	85	3	5	4	3	7	14	6	14	11	-
1982	91	2	1	6	7	14	11	10	7	17	-
1981	107	4	5	6	3	13	17	12	7	23	-
1980	91	1	3	4	12	10	11	10	14	7	-
1979	73	1	5	3	9	6	12	8	7	10	-
Salmonella infections (002-003)											
All races, male											
1998	14	-	1	1	1	3	1	2	2	2	-
1997	34	1	2	-	1	2	1	5	5	6	-
1996	27	-	1	2	3	2	5	1	1	5	-
1995	35	2	1	1	2	4	6	7	3	5	-
1994	30	-	1	1	2	4	3	3	6	3	-
1993	27	3	1	3	-	2	1	3	3	2	-
1992	24	1	-	-	1	3	3	-	6	3	-
1991	33	2	2	3	6	2	2	2	1	6	-
1990	49	1	2	3	7	5	5	10	5	3	-
1989	48	3	-	1	4	10	5	4	4	5	-
1988	34	1	4	2	2	-	3	7	2	4	-
1987	58	5	1	2	4	5	6	9	5	9	-
1986	71	1	3	3	7	8	3	8	11	7	-
1985	63	-	4	5	3	3	10	8	6	8	-
1984	49	3	3	1	2	2	10	5	3	10	-
1983	40	1	3	-	1	2	4	3	5	6	-
1982	49	2	1	2	2	7	5	7	3	8	-
1981	58	1	2	3	1	9	10	9	3	9	-
1980	48	1	2	3	6	8	8	4	5	3	-
1979	37	1	2	1	8	5	5	4	2	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
All races, female											
1998	23	-	1	2	-	1	3	3	5	5	-
1997	17	1	2	-	1	-	-	1	7	2	-
1996	32	1	-	1	2	1	3	4	3	7	-
1995	31	2	1	-	2	1	3	4	7	3	-
1994	20	2	1	1	1	1	1	-	4	3	-
1993	25	3	-	-	1	2	3	5	4	4	-
1992	23	-	-	1	1	1	6	3	2	5	-
1991	21	-	-	-	-	2	2	1	4	9	-
1990	32	1	2	1	1	2	2	5	5	8	-
1989	51	1	2	1	2	2	6	6	10	13	-
1988	32	-	1	-	4	1	3	6	4	9	-
1987	49	-	-	1	3	3	8	7	5	18	-
1986	33	-	-	1	1	3	7	8	3	7	-
1985	54	1	2	-	9	4	8	9	7	9	-
1984	41	4	-	2	3	1	6	8	4	9	-
1983	45	2	2	4	2	5	10	3	9	5	-
1982	42	-	-	4	5	7	6	3	4	9	-
1981	49	3	3	3	2	4	7	3	4	14	-
1980	43	-	1	1	6	2	3	6	9	4	-
1979	36	-	3	2	1	1	7	4	5	6	-
Salmonella infections (002-003)											
White, both sexes											
1998	32	-	2	3	-	3	4	5	6	6	-
1997	40	2	3	-	-	2	1	6	12	6	-
1996	41	1	-	2	3	2	7	4	3	9	-
1995	50	2	1	-	3	5	8	9	10	6	-
1994	33	2	1	1	2	4	3	2	7	4	-
1993	37	4	-	2	1	3	3	4	6	5	-
1992	32	-	-	1	1	2	8	2	8	7	-
1991	34	1	1	1	3	3	3	1	3	11	-
1990	54	2	4	2	7	4	3	11	8	9	-
1989	78	2	1	1	6	10	7	10	13	16	-
1988	50	-	5	1	4	1	6	9	6	12	-
1987	81	3	-	2	7	6	10	14	8	27	-
1986	86	-	3	4	6	11	9	15	12	10	-
1985	89	-	2	4	7	6	14	16	12	15	-
1984	67	6	2	2	2	1	11	13	6	17	-
1983	60	2	4	3	2	4	12	5	9	8	-
1982	72	1	1	6	6	9	9	9	6	15	-
1981	86	2	4	4	3	10	17	12	7	18	-
1980	70	1	3	2	11	7	9	8	12	7	-
1979	57	-	4	2	8	4	11	7	6	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
White, male											
1998	12	-	1	1	-	2	1	2	2	2	-
1997	28	1	2	-	-	2	1	5	5	5	-
1996	18	-	-	2	2	1	4	1	-	4	-
1995	28	1	1	-	2	4	6	5	3	4	-
1994	16	-	-	-	1	3	3	2	3	2	-
1993	17	2	-	2	-	2	1	-	2	2	-
1992	14	-	-	-	-	2	3	-	6	2	-
1991	20	1	1	1	3	2	1	1	1	4	-
1990	31	1	2	1	6	2	2	7	4	3	-
1989	41	1	-	1	4	9	4	4	4	5	-
1988	24	-	4	1	2	-	3	5	2	3	-
1987	40	3	-	1	4	4	5	8	4	9	-
1986	59	-	3	3	5	8	2	8	9	6	-
1985	50	-	2	4	3	3	6	7	6	8	-
1984	34	3	2	1	1	1	7	5	3	8	-
1983	27	1	2	-	1	-	2	3	4	5	-
1982	40	1	1	2	2	6	5	6	2	7	-
1981	46	1	2	2	1	7	10	9	3	6	-
1980	35	1	2	1	5	6	7	3	4	3	-
1979	29	-	2	1	8	3	4	4	2	2	-
Salmonella infections (002-003)											
White, female											
1998	20	-	1	2	-	1	3	3	4	4	-
1997	12	1	1	-	-	-	-	1	7	1	-
1996	23	1	-	-	1	1	3	3	3	5	-
1995	22	1	-	-	1	1	2	4	7	2	-
1994	17	2	1	1	1	1	-	-	4	2	-
1993	20	2	-	-	1	1	2	4	4	3	-
1992	18	-	-	1	1	-	5	2	2	5	-
1991	14	-	-	-	-	1	2	-	2	7	-
1990	23	1	2	1	1	2	1	4	4	6	-
1989	37	1	1	-	2	1	3	6	9	11	-
1988	26	-	1	-	2	1	3	4	4	9	-
1987	41	-	-	1	3	2	5	6	4	18	-
1986	27	-	-	1	1	3	7	7	3	4	-
1985	39	-	-	-	4	3	8	9	6	7	-
1984	33	3	-	1	1	-	4	8	3	9	-
1983	33	1	2	3	1	4	10	2	5	3	-
1982	32	-	-	4	4	3	4	3	4	8	-
1981	40	1	2	2	2	3	7	3	4	12	-
1980	35	-	1	1	6	1	2	5	8	4	-
1979	28	-	2	1	-	1	7	3	4	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
All other, both sexes											
1998	5	-	-	-	1	1	-	-	1	1	-
1997	11	-	1	-	2	-	-	-	-	2	-
1996	18	-	1	1	2	1	1	1	1	3	-
1995	16	2	1	1	1	-	1	2	-	2	-
1994	17	-	1	1	1	1	1	1	3	2	-
1993	15	2	1	1	-	1	1	4	1	1	-
1992	15	1	-	-	1	2	1	1	-	1	-
1991	20	1	1	2	3	1	1	2	2	4	-
1990	27	-	-	2	1	3	4	4	2	2	-
1989	21	2	1	1	-	2	4	-	1	2	-
1988	16	1	-	1	2	-	-	4	-	1	-
1987	26	2	1	1	-	2	4	2	2	-	-
1986	18	1	-	-	2	-	1	1	2	4	-
1985	28	1	4	1	5	1	4	1	1	2	-
1984	23	1	1	1	3	2	5	-	1	2	-
1983	25	1	1	1	1	3	2	1	5	3	-
1982	19	1	-	-	1	5	2	1	1	2	-
1981	21	2	1	2	-	3	-	-	-	5	-
1980	21	-	-	2	1	3	2	2	2	-	-
1979	16	1	1	1	1	2	1	1	1	2	-
Salmonella infections (002-003)											
All other, male											
1998	2	-	-	-	1	1	-	-	-	-	-
1997	6	-	-	-	1	-	-	-	-	1	-
1996	9	-	1	-	1	1	1	-	1	1	-
1995	7	1	-	1	-	-	-	2	-	1	-
1994	14	-	1	1	1	1	-	1	3	1	-
1993	10	1	1	1	-	-	-	3	1	-	-
1992	10	1	-	-	1	1	-	-	-	1	-
1991	13	1	1	2	3	-	1	1	-	2	-
1990	18	-	-	2	1	3	3	3	1	-	-
1989	7	2	-	-	-	1	1	-	-	-	-
1988	10	1	-	1	-	-	-	2	-	1	-
1987	18	2	1	1	-	1	1	1	1	-	-
1986	12	1	-	-	2	-	1	-	2	1	-
1985	13	-	2	1	-	-	4	1	-	-	-
1984	15	-	1	-	1	1	3	-	-	2	-
1983	13	-	1	-	-	2	2	-	1	1	-
1982	9	1	-	-	-	1	-	1	1	1	-
1981	12	-	-	1	-	2	-	-	-	3	-
1980	13	-	-	2	1	2	1	1	1	-	-
1979	8	1	-	-	-	2	1	-	-	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
All other, female											
1998	3	-	-	-	-	-	-	-	1	1	-
1997	5	-	1	-	1	-	-	-	-	1	-
1996	9	-	-	1	1	-	-	1	-	2	-
1995	9	1	1	-	1	-	1	-	-	1	-
1994	3	-	-	-	-	-	1	-	-	1	-
1993	5	1	-	-	-	1	1	1	-	1	-
1992	5	-	-	-	-	1	1	1	-	-	-
1991	7	-	-	-	-	1	-	1	2	2	-
1990	9	-	-	-	-	-	1	1	1	2	-
1989	14	-	1	1	-	1	3	-	1	2	-
1988	6	-	-	-	2	-	-	2	-	-	-
1987	8	-	-	-	-	1	3	1	1	-	-
1986	6	-	-	-	-	-	-	1	-	3	-
1985	15	1	2	-	5	1	-	-	1	2	-
1984	8	1	-	1	2	1	2	-	1	-	-
1983	12	1	-	1	1	1	-	1	4	2	-
1982	10	-	-	-	1	4	2	-	-	1	-
1981	9	2	1	1	-	1	-	-	-	2	-
1980	8	-	-	-	-	1	1	1	1	-	-
1979	8	-	1	1	1	-	-	1	1	-	-
Salmonella infections (002-003)											
Black, both sexes											
1998	3	-	-	-	-	1	-	-	1	-	-
1997	9	-	-	-	2	-	-	-	-	1	-
1996	15	-	1	1	2	1	1	1	-	3	-
1995	13	2	1	1	1	-	1	1	-	1	-
1994	16	-	1	1	1	1	-	1	3	2	-
1993	10	1	1	-	-	-	1	3	1	1	-
1992	11	-	-	-	1	2	-	1	-	-	-
1991	18	1	1	2	2	1	1	2	2	4	-
1990	24	-	-	2	1	2	4	4	2	1	-
1989	19	2	1	1	-	2	3	-	1	2	-
1988	16	1	-	1	2	-	-	4	-	1	-
1987	25	2	1	1	-	2	4	2	2	-	-
1986	16	1	-	-	2	-	1	-	1	4	-
1985	26	1	4	1	5	1	3	1	1	1	-
1984	21	1	1	-	3	2	5	-	1	1	-
1983	24	1	1	1	1	3	2	1	5	3	-
1982	18	1	-	-	1	4	2	1	1	2	-
1981	21	2	1	2	-	3	-	-	-	5	-
1980	21	-	-	2	1	3	2	2	2	-	-
1979	13	1	1	1	1	1	-	1	1	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Salmonella infections (002-003)											
Black, male											
1998	1	-	-	-	-	1	-	-	-	-	-
1997	5	-	-	-	1	-	-	-	-	-	-
1996	7	-	1	-	1	1	1	-	-	1	-
1995	6	1	-	1	-	-	-	1	-	1	-
1994	14	-	1	1	1	1	-	1	3	1	-
1993	7	1	1	-	-	-	-	2	1	-	-
1992	7	-	-	-	1	1	-	-	-	-	-
1991	11	1	1	2	2	-	1	1	-	2	-
1990	16	-	-	2	1	2	3	3	1	-	-
1989	5	2	-	-	-	1	-	-	-	-	-
1988	10	1	-	1	-	-	-	2	-	1	-
1987	17	2	1	1	-	1	1	1	1	-	-
1986	11	1	-	-	2	-	1	-	1	1	-
1985	12	-	2	1	-	-	3	1	-	-	-
1984	14	-	1	-	1	1	3	-	-	1	-
1983	12	-	1	-	-	2	2	-	1	1	-
1982	9	1	-	-	-	1	-	1	1	1	-
1981	12	-	-	1	-	2	-	-	-	3	-
1980	13	-	-	2	1	2	1	1	1	-	-
1979	5	1	-	-	-	1	-	-	-	2	-
Salmonella infections (002-003)											
Black, female											
1998	2	-	-	-	-	-	-	-	1	-	-
1997	4	-	-	-	1	-	-	-	-	1	-
1996	8	-	-	1	1	-	-	1	-	2	-
1995	7	1	1	-	1	-	1	-	-	-	-
1994	2	-	-	-	-	-	-	-	-	1	-
1993	3	-	-	-	-	-	1	1	-	1	-
1992	4	-	-	-	-	1	-	1	-	-	-
1991	7	-	-	-	-	1	-	1	2	2	-
1990	8	-	-	-	-	-	1	1	1	1	-
1989	14	-	1	1	-	1	3	-	1	2	-
1988	6	-	-	-	2	-	-	2	-	-	-
1987	8	-	-	-	-	1	3	1	1	-	-
1986	5	-	-	-	-	-	-	-	-	3	-
1985	14	1	2	-	5	1	-	-	1	1	-
1984	7	1	-	-	2	1	2	-	1	-	-
1983	12	1	-	1	1	1	-	1	4	2	-
1982	9	-	-	-	1	3	2	-	-	1	-
1981	9	2	1	1	-	1	-	-	-	2	-
1980	8	-	-	-	-	1	1	1	1	-	-
1979	8	-	1	1	1	-	-	1	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Shigellosis and amebiasis (004,006)											
All races, both sexes											
1998	8	-	-	-	-	-	-	-	1	4	-
1997	6	-	-	-	-	1	-	1	-	2	-
1996	9	2	-	2	-	1	-	-	2	1	-
1995	12	1	-	-	2	1	1	1	2	3	-
1994	15	-	1	2	-	1	1	1	-	4	-
1993	11	-	-	-	-	1	3	2	-	-	-
1992	14	1	3	-	-	1	1	1	-	4	-
1991	15	-	-	-	1	-	1	2	2	1	-
1990	15	-	1	1	1	-	1	2	2	1	-
1989	20	-	-	-	-	1	-	5	1	3	-
1988	15	1	2	-	-	-	-	2	1	3	-
1987	22	-	1	3	2	1	2	-	-	3	-
1986	12	1	-	-	-	1	-	1	2	-	-
1985	27	2	-	1	1	4	4	1	1	3	-
1984	18	-	-	2	1	2	3	-	1	4	-
1983	30	2	2	2	3	3	3	2	3	3	-
1982	16	-	-	3	1	-	1	4	-	1	-
1981	27	2	3	-	1	2	3	1	4	1	-
1980	37	2	-	1	4	1	3	3	3	1	-
1979	38	3	3	1	3	3	3	2	-	4	-
Shigellosis and amebiasis (004,006)											
All races, male											
1998	3	-	-	-	-	-	-	-	1	-	-
1997	1	-	-	-	-	-	-	-	-	1	-
1996	3	2	-	1	-	-	-	-	-	-	-
1995	7	1	-	-	1	-	1	1	1	1	-
1994	7	-	1	2	-	1	1	1	-	-	-
1993	5	-	-	-	-	-	2	-	-	-	-
1992	12	1	3	-	-	1	-	1	-	3	-
1991	11	-	-	-	1	-	1	2	2	-	-
1990	5	-	1	-	1	-	-	1	-	-	-
1989	9	-	-	-	-	-	-	4	-	-	-
1988	5	-	1	-	-	-	-	-	-	-	-
1987	11	-	1	1	1	1	-	-	-	-	-
1986	9	1	-	-	-	1	-	1	1	-	-
1985	16	1	-	1	-	2	4	-	1	-	-
1984	11	-	-	2	1	2	2	-	1	1	-
1983	17	1	2	2	-	2	2	2	1	2	-
1982	8	-	-	1	1	-	1	1	-	1	-
1981	16	2	2	-	1	1	3	1	1	1	-
1980	21	1	-	-	2	1	1	1	1	1	-
1979	26	3	3	-	2	1	2	-	-	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Shigellosis and amebiasis (004,006)											
All races, female											
1998	5	-	-	-	-	-	-	-	-	4	-
1997	5	-	-	-	-	1	-	1	-	1	-
1996	6	-	-	1	-	1	-	-	2	1	-
1995	5	-	-	-	1	1	-	-	1	2	-
1994	8	-	-	-	-	-	-	-	-	4	-
1993	6	-	-	-	-	1	1	2	-	-	-
1992	2	-	-	-	-	-	1	-	-	1	-
1991	4	-	-	-	-	-	-	-	-	1	-
1990	10	-	-	1	-	-	1	1	2	1	-
1989	11	-	-	-	-	1	-	1	1	3	-
1988	10	1	1	-	-	-	-	2	1	3	-
1987	11	-	-	2	1	-	2	-	-	3	-
1986	3	-	-	-	-	-	-	-	1	-	-
1985	11	1	-	-	1	2	-	1	-	3	-
1984	7	-	-	-	-	-	1	-	-	3	-
1983	13	1	-	-	3	1	1	-	2	1	-
1982	8	-	-	2	-	-	-	3	-	-	-
1981	11	-	1	-	-	1	-	-	3	-	-
1980	16	1	-	1	2	-	2	2	2	-	-
1979	12	-	-	1	1	2	1	2	-	2	-
Shigellosis and amebiasis (004,006)											
White, both sexes											
1998	6	-	-	-	-	-	-	-	1	4	-
1997	6	-	-	-	-	1	-	1	-	2	-
1996	9	2	-	2	-	1	-	-	2	1	-
1995	10	1	-	-	2	1	1	1	2	2	-
1994	12	-	1	1	-	1	1	1	-	4	-
1993	9	-	-	-	-	1	2	2	-	-	-
1992	10	1	2	-	-	1	1	1	-	3	-
1991	13	-	-	-	1	-	1	2	2	1	-
1990	11	-	1	-	1	-	1	2	2	-	-
1989	12	-	-	-	-	-	-	4	1	3	-
1988	8	-	-	-	-	-	-	1	1	3	-
1987	16	-	1	3	1	1	1	-	-	3	-
1986	11	1	-	-	-	1	-	1	2	-	-
1985	20	-	-	1	1	4	3	1	1	3	-
1984	16	-	-	2	1	2	2	-	1	3	-
1983	23	-	2	2	2	2	3	2	2	3	-
1982	12	-	-	2	-	-	-	4	-	1	-
1981	22	2	2	-	1	1	2	1	4	1	-
1980	28	1	-	1	4	1	2	3	3	-	-
1979	29	3	2	-	2	2	3	2	-	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Shigellosis and amebiasis (004,006)											
White, male											
1998	2	-	-	-	-	-	-	-	1	-	-
1997	1	-	-	-	-	-	-	-	-	1	-
1996	3	2	-	1	-	-	-	-	-	-	-
1995	5	1	-	-	1	-	1	1	1	-	-
1994	6	-	1	1	-	1	1	1	-	-	-
1993	4	-	-	-	-	-	1	-	-	-	-
1992	8	1	2	-	-	1	-	1	-	2	-
1991	10	-	-	-	1	-	1	2	2	-	-
1990	5	-	1	-	1	-	-	1	-	-	-
1989	6	-	-	-	-	-	-	3	-	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	7	-	1	1	1	1	-	-	-	-	-
1986	8	1	-	-	-	1	-	1	1	-	-
1985	10	-	-	1	-	2	3	-	1	-	-
1984	11	-	-	2	1	2	2	-	1	1	-
1983	15	-	2	2	-	2	2	2	1	2	-
1982	5	-	-	1	-	-	-	1	-	1	-
1981	13	2	2	-	1	1	2	1	1	1	-
1980	14	-	-	-	2	1	1	1	1	-	-
1979	22	3	2	-	2	1	2	-	-	2	-
Shigellosis and amebiasis (004,006)											
White, female											
1998	4	-	-	-	-	-	-	-	-	4	-
1997	5	-	-	-	-	1	-	1	-	1	-
1996	6	-	-	1	-	1	-	-	2	1	-
1995	5	-	-	-	1	1	-	-	1	2	-
1994	6	-	-	-	-	-	-	-	-	4	-
1993	5	-	-	-	-	1	1	2	-	-	-
1992	2	-	-	-	-	-	1	-	-	1	-
1991	3	-	-	-	-	-	-	-	-	1	-
1990	6	-	-	-	-	-	1	1	2	-	-
1989	6	-	-	-	-	-	-	1	1	3	-
1988	6	-	-	-	-	-	-	1	1	3	-
1987	9	-	-	2	-	-	1	-	-	3	-
1986	3	-	-	-	-	-	-	-	1	-	-
1985	10	-	-	-	1	2	-	1	-	3	-
1984	5	-	-	-	-	-	-	-	-	2	-
1983	8	-	-	-	2	-	1	-	1	1	-
1982	7	-	-	1	-	-	-	3	-	-	-
1981	9	-	-	-	-	-	-	-	3	-	-
1980	14	1	-	1	2	-	1	2	2	-	-
1979	7	-	-	-	-	1	1	2	-	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Shigellosis and amebiasis (004,006)											
All other, both sexes											
1998	2	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	1	-
1994	3	-	-	1	-	-	-	-	-	-	-
1993	2	-	-	-	-	-	1	-	-	-	-
1992	4	-	1	-	-	-	-	-	-	1	-
1991	2	-	-	-	-	-	-	-	-	-	-
1990	4	-	-	1	-	-	-	-	-	1	-
1989	8	-	-	-	-	1	-	1	-	-	-
1988	7	1	2	-	-	-	-	1	-	-	-
1987	6	-	-	-	1	-	1	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	7	2	-	-	-	-	1	-	-	-	-
1984	2	-	-	-	-	-	1	-	-	1	-
1983	7	2	-	-	1	1	-	-	1	-	-
1982	4	-	-	1	1	-	1	-	-	-	-
1981	5	-	1	-	-	1	1	-	-	-	-
1980	9	1	-	-	-	-	1	-	-	1	-
1979	9	-	1	1	1	1	-	-	-	-	-
Shigellosis and amebiasis (004,006)											
All other, male											
1998	1	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	1	-
1994	1	-	-	1	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	1	-	-	-	-
1992	4	-	1	-	-	-	-	-	-	1	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	3	-	-	-	-	-	-	1	-	-	-
1988	3	-	1	-	-	-	-	-	-	-	-
1987	4	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	6	1	-	-	-	-	1	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	2	1	-	-	-	-	-	-	-	-	-
1982	3	-	-	-	1	-	1	-	-	-	-
1981	3	-	-	-	-	-	1	-	-	-	-
1980	7	1	-	-	-	-	-	-	-	1	-
1979	4	-	1	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Shigellosis and amebiasis (004,006)											
All other, female											
1998	1	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	2	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	4	-	-	1	-	-	-	-	-	1	-
1989	5	-	-	-	-	1	-	-	-	-	-
1988	4	1	1	-	-	-	-	1	-	-	-
1987	2	-	-	-	1	-	1	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	1	-	-	-	-	-	-	-	-	-
1984	2	-	-	-	-	-	1	-	-	1	-
1983	5	1	-	-	1	1	-	-	1	-	-
1982	1	-	-	1	-	-	-	-	-	-	-
1981	2	-	1	-	-	1	-	-	-	-	-
1980	2	-	-	-	-	-	1	-	-	-	-
1979	5	-	-	1	1	1	-	-	-	-	-
Shigellosis and amebiasis (004,006)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	1	-
1994	2	-	-	1	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	2	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	1	-	-	-	-	-	-	-
1989	6	-	-	-	-	1	-	-	-	-	-
1988	2	-	-	-	-	-	-	1	-	-	-
1987	5	-	-	-	1	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	7	2	-	-	-	-	1	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	1	-
1983	5	1	-	-	1	1	-	-	1	-	-
1982	3	-	-	-	1	-	1	-	-	-	-
1981	3	-	-	-	-	1	-	-	-	-	-
1980	7	1	-	-	-	-	1	-	-	-	-
1979	6	-	1	-	-	1	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Shigellosis and amebiasis (004,006)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	1	-
1994	1	-	-	1	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	2	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	6	1	-	-	-	-	1	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	1	-	-	-	-	-	-	-	-	-
1982	3	-	-	-	1	-	1	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	5	1	-	-	-	-	-	-	-	-	-
1979	3	-	1	-	-	-	-	-	-	-	-
Shigellosis and amebiasis (004,006)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	1	-	-	-	-	-	-	-
1989	4	-	-	-	-	1	-	-	-	-	-
1988	2	-	-	-	-	-	-	1	-	-	-
1987	1	-	-	-	1	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	1	-
1983	4	-	-	-	1	1	-	-	1	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	1	-	-	-	-	-
1980	2	-	-	-	-	-	1	-	-	-	-
1979	3	-	-	-	-	1	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
All races, both sexes											
1998	1,059	11	11	21	25	52	75	129	164	302	-
1997	1,023	9	17	16	24	54	84	133	168	259	-
1996	824	6	10	9	43	43	55	82	109	224	-
1995	793	8	9	9	14	33	60	83	103	225	-
1994	746	8	12	8	27	39	53	83	76	179	-
1993	631	8	9	10	22	33	58	76	89	156	-
1992	529	4	6	5	17	28	36	56	65	138	-
1991	469	8	7	11	12	28	33	29	55	122	-
1990	415	3	7	5	15	17	34	45	49	101	-
1989	339	4	4	7	15	25	23	36	31	70	-
1988	305	4	3	10	6	17	23	31	34	69	-
1987	307	4	2	4	13	18	25	30	23	53	-
1986	347	10	6	9	10	14	19	26	17	70	1
1985	301	5	4	6	3	18	16	23	30	54	-
1984	331	3	5	6	15	9	19	23	26	51	1
1983	354	2	8	9	9	11	21	33	32	62	-
1982	329	4	1	5	14	12	15	27	23	44	-
1981	371	8	6	8	14	14	19	28	41	59	-
1980	364	5	8	10	17	15	25	26	33	68	-
1979	298	6	5	9	9	10	17	24	32	46	-
Certain other intestinal infections (007-009)											
All races, male											
1998	464	7	7	11	9	27	32	54	67	94	-
1997	435	5	8	7	13	22	41	60	62	62	-
1996	356	3	6	5	17	21	26	41	33	52	-
1995	347	6	7	7	8	18	27	38	34	55	-
1994	348	6	5	5	19	18	27	35	32	45	-
1993	292	3	6	7	13	12	33	28	31	57	-
1992	255	2	2	2	9	16	20	25	25	44	-
1991	215	3	5	6	8	12	18	16	17	36	-
1990	180	3	3	2	7	6	14	16	16	34	-
1989	172	2	4	6	7	12	11	18	9	25	-
1988	133	3	1	4	3	8	8	14	13	17	-
1987	137	3	1	2	5	7	14	8	13	16	-
1986	192	5	3	4	6	6	10	11	4	24	-
1985	146	3	3	4	-	7	10	9	8	16	-
1984	156	2	2	1	5	5	4	7	12	15	-
1983	172	1	4	3	3	8	11	12	14	12	-
1982	148	2	1	2	6	6	5	12	5	14	-
1981	170	4	3	5	5	5	12	18	12	9	-
1980	162	3	6	4	7	8	6	12	11	19	-
1979	143	1	3	4	6	7	7	8	15	14	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
All races, female											
1998	595	4	4	10	16	25	43	75	97	208	-
1997	588	4	9	9	11	32	43	73	106	197	-
1996	468	3	4	4	26	22	29	41	76	172	-
1995	446	2	2	2	6	15	33	45	69	170	-
1994	398	2	7	3	8	21	26	48	44	134	-
1993	339	5	3	3	9	21	25	48	58	99	-
1992	274	2	4	3	8	12	16	31	40	94	-
1991	254	5	2	5	4	16	15	13	38	86	-
1990	235	-	4	3	8	11	20	29	33	67	-
1989	167	2	-	1	8	13	12	18	22	45	-
1988	172	1	2	6	3	9	15	17	21	52	-
1987	170	1	1	2	8	11	11	22	10	37	-
1986	155	5	3	5	4	8	9	15	13	46	1
1985	155	2	1	2	3	11	6	14	22	38	-
1984	175	1	3	5	10	4	15	16	14	36	1
1983	182	1	4	6	6	3	10	21	18	50	-
1982	181	2	-	3	8	6	10	15	18	30	-
1981	201	4	3	3	9	9	7	10	29	50	-
1980	202	2	2	6	10	7	19	14	22	49	-
1979	155	5	2	5	3	3	10	16	17	32	-
Certain other intestinal infections (007-009)											
White, both sexes											
1998	882	8	10	19	22	44	66	122	152	283	-
1997	850	6	11	12	20	47	72	125	159	247	-
1996	672	5	9	8	41	40	50	76	106	210	-
1995	635	6	7	9	11	26	52	79	93	218	-
1994	574	6	11	6	17	31	47	78	75	175	-
1993	510	7	6	9	16	31	50	67	84	151	-
1992	430	4	4	4	15	24	35	53	62	132	-
1991	363	6	5	10	11	23	29	27	54	113	-
1990	342	3	7	5	11	14	33	43	46	96	-
1989	252	3	3	6	11	23	23	31	30	64	-
1988	239	4	3	9	6	13	20	25	32	63	-
1987	246	3	1	4	13	15	22	27	23	53	-
1986	257	5	2	7	7	11	18	26	16	66	1
1985	231	3	3	5	2	17	15	22	29	50	-
1984	245	3	3	5	13	9	17	21	22	51	-
1983	262	2	5	8	9	10	17	32	29	59	-
1982	233	2	-	3	13	9	12	26	21	42	-
1981	290	8	6	7	14	11	18	25	38	57	-
1980	286	2	7	9	15	12	21	22	32	68	-
1979	228	5	5	7	8	9	15	19	31	44	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
White, male											
1998	379	6	6	10	9	22	29	50	65	89	-
1997	343	3	7	5	11	21	37	57	57	57	-
1996	268	2	5	4	16	19	22	39	32	50	-
1995	264	4	6	7	7	14	24	36	32	53	-
1994	244	5	5	3	12	13	21	33	31	43	-
1993	221	2	5	6	11	10	27	24	30	56	-
1992	205	2	2	1	8	14	19	23	24	43	-
1991	157	2	4	6	7	11	16	14	17	32	-
1990	141	3	3	2	5	5	13	15	13	32	-
1989	120	1	3	5	5	11	11	15	9	23	-
1988	99	3	1	4	3	8	7	12	12	14	-
1987	111	2	-	2	5	6	12	7	13	16	-
1986	134	3	1	2	4	4	10	11	4	21	-
1985	107	2	2	3	-	7	9	9	7	15	-
1984	102	2	1	1	3	5	4	5	11	15	-
1983	117	1	3	2	3	7	9	12	13	11	-
1982	103	2	-	2	6	3	4	12	5	14	-
1981	124	4	3	5	5	4	11	16	11	8	-
1980	122	1	6	3	6	7	5	10	11	19	-
1979	107	1	3	2	5	6	6	7	14	13	-
Certain other intestinal infections (007-009)											
White, female											
1998	503	2	4	9	13	22	37	72	87	194	-
1997	507	3	4	7	9	26	35	68	102	190	-
1996	404	3	4	4	25	21	28	37	74	160	-
1995	371	2	1	2	4	12	28	43	61	165	-
1994	330	1	6	3	5	18	26	45	44	132	-
1993	289	5	1	3	5	21	23	43	54	95	-
1992	225	2	2	3	7	10	16	30	38	89	-
1991	206	4	1	4	4	12	13	13	37	81	-
1990	201	-	4	3	6	9	20	28	33	64	-
1989	132	2	-	1	6	12	12	16	21	41	-
1988	140	1	2	5	3	5	13	13	20	49	-
1987	135	1	1	2	8	9	10	20	10	37	-
1986	123	2	1	5	3	7	8	15	12	45	1
1985	124	1	1	2	2	10	6	13	22	35	-
1984	143	1	2	4	10	4	13	16	11	36	-
1983	145	1	2	6	6	3	8	20	16	48	-
1982	130	-	-	1	7	6	8	14	16	28	-
1981	166	4	3	2	9	7	7	9	27	49	-
1980	164	1	1	6	9	5	16	12	21	49	-
1979	121	4	2	5	3	3	9	12	17	31	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
All other, both sexes											
1998	177	3	1	2	3	8	9	7	12	19	-
1997	173	3	6	4	4	7	12	8	9	12	-
1996	152	1	1	1	2	3	5	6	3	14	-
1995	158	2	2	-	3	7	8	4	10	7	-
1994	172	2	1	2	10	8	6	5	1	4	-
1993	121	1	3	1	6	2	8	9	5	5	-
1992	99	-	2	1	2	4	1	3	3	6	-
1991	106	2	2	1	1	5	4	2	1	9	-
1990	73	-	-	-	4	3	1	2	3	5	-
1989	87	1	1	1	4	2	-	5	1	6	-
1988	66	-	-	1	-	4	3	6	2	6	-
1987	61	1	1	-	-	3	3	3	-	-	-
1986	90	5	4	2	3	3	1	-	1	4	-
1985	70	2	1	1	1	1	1	1	1	4	-
1984	86	-	2	1	2	-	2	2	4	-	1
1983	92	-	3	1	-	1	4	1	3	3	-
1982	96	2	1	2	1	3	3	1	2	2	-
1981	81	-	-	1	-	3	1	3	3	2	-
1980	78	3	1	1	2	3	4	4	1	-	-
1979	70	1	-	2	1	1	2	5	1	2	-
Certain other intestinal infections (007-009)											
All other, male											
1998	85	1	1	1	-	5	3	4	2	5	-
1997	92	2	1	2	2	1	4	3	5	5	-
1996	88	1	1	1	1	2	4	2	1	2	-
1995	83	2	1	-	1	4	3	2	2	2	-
1994	104	1	-	2	7	5	6	2	1	2	-
1993	71	1	1	1	2	2	6	4	1	1	-
1992	50	-	-	1	1	2	1	2	1	1	-
1991	58	1	1	-	1	1	2	2	-	4	-
1990	39	-	-	-	2	1	1	1	3	2	-
1989	52	1	1	1	2	1	-	3	-	2	-
1988	34	-	-	-	-	-	1	2	1	3	-
1987	26	1	1	-	-	1	2	1	-	-	-
1986	58	2	2	2	2	2	-	-	-	3	-
1985	39	1	1	1	-	-	1	-	1	1	-
1984	54	-	1	-	2	-	-	2	1	-	-
1983	55	-	1	1	-	1	2	-	1	1	-
1982	45	-	1	-	-	3	1	-	-	-	-
1981	46	-	-	-	-	1	1	2	1	1	-
1980	40	2	-	1	1	1	1	2	-	-	-
1979	36	-	-	2	1	1	1	1	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
All other, female											
1998	92	2	-	1	3	3	6	3	10	14	-
1997	81	1	5	2	2	6	8	5	4	7	-
1996	64	-	-	-	1	1	1	4	2	12	-
1995	75	-	1	-	2	3	5	2	8	5	-
1994	68	1	1	-	3	3	-	3	-	2	-
1993	50	-	2	-	4	-	2	5	4	4	-
1992	49	-	2	-	1	2	-	1	2	5	-
1991	48	1	1	1	-	4	2	-	1	5	-
1990	34	-	-	-	2	2	-	1	-	3	-
1989	35	-	-	-	2	1	-	2	1	4	-
1988	32	-	-	1	-	4	2	4	1	3	-
1987	35	-	-	-	-	2	1	2	-	-	-
1986	32	3	2	-	1	1	1	-	1	1	-
1985	31	1	-	-	1	1	-	1	-	3	-
1984	32	-	1	1	-	-	2	-	3	-	1
1983	37	-	2	-	-	-	2	1	2	2	-
1982	51	2	-	2	1	-	2	1	2	2	-
1981	35	-	-	1	-	2	-	1	2	1	-
1980	38	1	1	-	1	2	3	2	1	-	-
1979	34	1	-	-	-	-	1	4	-	1	-
Certain other intestinal infections (007-009)											
Black, both sexes											
1998	158	3	1	2	2	5	7	6	9	16	-
1997	154	3	4	4	2	5	10	6	9	12	-
1996	140	1	1	1	2	2	4	5	2	11	-
1995	145	2	2	-	3	5	6	4	8	6	-
1994	154	1	-	1	10	8	5	5	1	2	-
1993	110	1	2	1	4	2	7	8	5	4	-
1992	94	-	1	1	2	4	-	3	3	5	-
1991	96	2	2	1	1	3	4	2	1	7	-
1990	67	-	-	-	2	3	1	1	3	5	-
1989	78	1	1	1	3	2	-	5	1	6	-
1988	57	-	-	1	-	3	2	5	-	4	-
1987	56	-	1	-	-	3	3	3	-	-	-
1986	82	5	3	2	3	3	1	-	1	3	-
1985	65	2	1	1	1	1	1	1	-	4	-
1984	80	-	2	-	2	-	1	2	4	-	1
1983	84	-	2	1	-	1	4	1	2	1	-
1982	87	2	1	2	1	3	3	1	2	1	-
1981	75	-	-	1	-	3	1	2	3	2	-
1980	72	3	-	1	2	3	3	3	1	-	-
1979	65	1	-	2	1	1	2	3	1	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain other intestinal infections (007-009)											
Black, male											
1998	77	1	1	1	-	3	2	4	1	3	-
1997	81	2	-	2	-	1	4	2	5	5	-
1996	82	1	1	1	1	1	3	2	1	2	-
1995	75	2	1	-	1	4	1	2	1	2	-
1994	94	1	-	1	7	5	5	2	1	1	-
1993	64	1	1	1	-	2	5	3	1	1	-
1992	47	-	-	1	1	2	-	2	1	-	-
1991	53	1	1	-	1	1	2	2	-	3	-
1990	34	-	-	-	1	1	1	-	3	2	-
1989	46	1	1	1	1	1	-	3	-	2	-
1988	29	-	-	-	-	-	1	1	-	2	-
1987	23	-	1	-	-	1	2	1	-	-	-
1986	53	2	2	2	2	2	-	-	-	2	-
1985	36	1	1	1	-	-	1	-	-	1	-
1984	51	-	1	-	2	-	-	2	1	-	-
1983	51	-	-	1	-	1	2	-	-	-	-
1982	39	-	1	-	-	3	1	-	-	-	-
1981	42	-	-	-	-	1	1	1	1	1	-
1980	36	2	-	1	1	1	-	2	-	-	-
1979	35	-	-	2	1	1	1	1	1	1	-
Certain other intestinal infections (007-009)											
Black, female											
1998	81	2	-	1	2	2	5	2	8	13	-
1997	73	1	4	2	2	4	6	4	4	7	-
1996	58	-	-	-	1	1	1	3	1	9	-
1995	70	-	1	-	2	1	5	2	7	4	-
1994	60	-	-	-	3	3	-	3	-	1	-
1993	46	-	1	-	4	-	2	5	4	3	-
1992	47	-	1	-	1	2	-	1	2	5	-
1991	43	1	1	1	-	2	2	-	1	4	-
1990	33	-	-	-	1	2	-	1	-	3	-
1989	32	-	-	-	2	1	-	2	1	4	-
1988	28	-	-	1	-	3	1	4	-	2	-
1987	33	-	-	-	-	2	1	2	-	-	-
1986	29	3	1	-	1	1	1	-	1	1	-
1985	29	1	-	-	1	1	-	1	-	3	-
1984	29	-	1	-	-	-	1	-	3	-	1
1983	33	-	2	-	-	-	2	1	2	1	-
1982	48	2	-	2	1	-	2	1	2	1	-
1981	33	-	-	1	-	2	-	1	2	1	-
1980	36	1	-	-	1	2	3	1	1	-	-
1979	30	1	-	-	-	-	1	2	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
All races, both sexes											
1998	1,112	58	56	60	91	95	134	134	146	177	-
1997	1,166	64	61	73	84	90	144	163	190	161	-
1996	1,202	68	61	71	96	99	146	149	160	192	-
1995	1,336	76	67	70	87	142	158	172	187	187	1
1994	1,478	85	67	94	110	147	166	188	169	221	2
1993	1,631	85	107	100	109	168	182	210	208	197	1
1992	1,705	86	74	103	139	152	226	198	184	215	-
1991	1,713	80	94	110	144	159	170	199	212	198	-
1990	1,810	96	92	117	152	183	199	210	196	209	-
1989	1,970	109	92	125	145	208	218	237	229	242	2
1988	1,921	101	94	124	154	182	222	234	219	256	2
1987	1,755	87	89	114	170	179	210	211	202	195	3
1986	1,782	75	108	114	165	185	188	230	186	211	3
1985	1,752	81	102	134	165	186	203	223	207	199	1
1984	1,729	78	99	147	155	192	214	205	200	176	1
1983	1,779	88	110	152	154	198	238	221	196	212	3
1982	1,807	62	115	158	167	207	226	265	175	231	1
1981	1,937	92	130	171	200	233	243	214	232	203	-
1980	1,978	89	154	172	192	249	293	236	200	214	2
1979	2,007	115	148	200	177	251	243	249	189	217	-
Tuberculosis (010-018)											
All races, male											
1998	658	40	38	40	60	58	76	83	85	84	-
1997	662	39	41	50	56	52	74	77	98	86	-
1996	739	48	44	55	64	67	90	87	95	87	-
1995	814	57	47	48	67	93	91	96	105	93	1
1994	884	54	46	63	76	100	96	107	88	110	2
1993	1,018	68	79	68	79	111	113	120	114	90	1
1992	1,059	72	59	79	101	91	147	120	99	94	-
1991	1,048	62	66	87	100	94	94	113	117	100	-
1990	1,147	73	66	81	100	115	122	131	117	106	-
1989	1,284	85	64	96	95	143	138	141	120	131	2
1988	1,221	80	75	90	111	124	145	143	110	123	2
1987	1,169	64	68	87	123	112	138	138	122	109	3
1986	1,157	57	90	83	100	119	119	136	110	107	3
1985	1,135	67	79	102	120	116	109	126	126	116	1
1984	1,149	59	79	110	108	128	142	124	113	101	1
1983	1,176	69	84	116	107	139	159	129	117	119	2
1982	1,202	51	90	124	120	136	152	169	105	113	-
1981	1,299	72	89	123	145	164	172	140	142	109	-
1980	1,331	59	113	130	144	180	194	141	132	131	2
1979	1,371	81	105	133	129	186	180	179	114	122	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
All races, female											
1998	454	18	18	20	31	37	58	51	61	93	-
1997	504	25	20	23	28	38	70	86	92	75	-
1996	463	20	17	16	32	32	56	62	65	105	-
1995	522	19	20	22	20	49	67	76	82	94	-
1994	594	31	21	31	34	47	70	81	81	111	-
1993	613	17	28	32	30	57	69	90	94	107	-
1992	646	14	15	24	38	61	79	78	85	121	-
1991	665	18	28	23	44	65	76	86	95	98	-
1990	663	23	26	36	52	68	77	79	79	103	-
1989	686	24	28	29	50	65	80	96	109	111	-
1988	700	21	19	34	43	58	77	91	109	133	-
1987	586	23	21	27	47	67	72	73	80	86	-
1986	625	18	18	31	65	66	69	94	76	104	-
1985	617	14	23	32	45	70	94	97	81	83	-
1984	580	19	20	37	47	64	72	81	87	75	-
1983	603	19	26	36	47	59	79	92	79	93	1
1982	605	11	25	34	47	71	74	96	70	118	1
1981	638	20	41	48	55	69	71	74	90	94	-
1980	647	30	41	42	48	69	99	95	68	83	-
1979	636	34	43	67	48	65	63	70	75	95	-
Tuberculosis (010-018)											
White, both sexes											
1998	667	21	29	40	55	55	82	95	108	124	-
1997	695	21	26	39	47	48	94	112	138	111	-
1996	724	31	26	43	42	62	100	98	112	145	-
1995	804	29	32	38	54	83	100	123	140	128	-
1994	863	36	24	39	59	91	107	131	119	168	2
1993	969	34	56	50	58	103	126	145	163	136	1
1992	994	34	31	57	81	79	154	138	136	161	-
1991	978	29	44	53	74	101	115	142	155	141	-
1990	1,033	33	50	54	90	116	138	138	141	151	-
1989	1,152	37	35	58	96	129	144	156	177	185	1
1988	1,172	40	44	73	99	113	162	171	164	187	1
1987	1,082	32	54	68	115	121	143	142	140	147	2
1986	1,065	36	45	72	100	113	132	167	132	160	1
1985	1,101	32	53	75	110	129	143	167	156	144	1
1984	1,047	33	45	84	93	124	145	154	150	135	-
1983	1,119	37	56	90	97	145	159	164	147	156	-
1982	1,135	22	69	82	99	140	155	194	130	176	1
1981	1,211	36	66	98	130	147	175	153	177	158	-
1980	1,253	34	77	98	129	177	195	178	142	173	1
1979	1,294	57	64	114	110	181	178	186	150	174	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
White, male											
1998	381	15	21	29	39	33	43	55	62	48	-
1997	380	15	18	25	34	28	44	52	67	58	-
1996	440	21	21	36	28	43	62	59	56	63	-
1995	474	20	21	23	44	53	56	68	78	62	-
1994	506	22	15	23	40	63	66	70	56	87	2
1993	591	26	40	36	46	73	84	79	85	56	1
1992	602	28	26	47	59	45	101	75	70	73	-
1991	593	26	27	43	59	61	62	77	82	71	-
1990	630	26	37	39	57	70	85	80	81	72	-
1989	724	28	27	43	62	89	92	91	88	99	1
1988	730	34	35	56	76	77	100	101	82	86	1
1987	724	24	40	52	82	76	100	92	88	86	2
1986	679	26	36	54	63	74	90	96	75	84	1
1985	703	27	43	56	85	78	73	93	98	81	1
1984	689	27	34	62	69	83	98	91	86	78	-
1983	718	29	42	68	67	101	105	94	82	88	-
1982	745	18	54	65	72	99	100	125	78	87	-
1981	797	30	52	72	96	100	122	97	107	80	-
1980	856	24	56	80	96	129	132	106	94	108	1
1979	909	44	51	79	78	138	134	138	95	99	-
Tuberculosis (010-018)											
White, female											
1998	286	6	8	11	16	22	39	40	46	76	-
1997	315	6	8	14	13	20	50	60	71	53	-
1996	284	10	5	7	14	19	38	39	56	82	-
1995	330	9	11	15	10	30	44	55	62	66	-
1994	357	14	9	16	19	28	41	61	63	81	-
1993	378	8	16	14	12	30	42	66	78	80	-
1992	392	6	5	10	22	34	53	63	66	88	-
1991	385	3	17	10	15	40	53	65	73	70	-
1990	403	7	13	15	33	46	53	58	60	79	-
1989	428	9	8	15	34	40	52	65	89	86	-
1988	442	6	9	17	23	36	62	70	82	101	-
1987	358	8	14	16	33	45	43	50	52	61	-
1986	386	10	9	18	37	39	42	71	57	76	-
1985	398	5	10	19	25	51	70	74	58	63	-
1984	358	6	11	22	24	41	47	63	64	57	-
1983	401	8	14	22	30	44	54	70	65	68	-
1982	390	4	15	17	27	41	55	69	52	89	1
1981	414	6	14	26	34	47	53	56	70	78	-
1980	397	10	21	18	33	48	63	72	48	65	-
1979	385	13	13	35	32	43	44	48	55	75	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
All other, both sexes											
1998	445	37	27	20	36	40	52	39	38	53	-
1997	471	43	35	34	37	42	50	51	52	50	-
1996	478	37	35	28	54	37	46	51	48	47	-
1995	532	47	35	32	33	59	58	49	47	59	1
1994	615	49	43	55	51	56	59	57	50	53	-
1993	662	51	51	50	51	65	56	65	45	61	-
1992	711	52	43	46	58	73	72	60	48	54	-
1991	735	51	50	57	70	58	55	57	57	57	-
1990	777	63	42	63	62	67	61	72	55	58	-
1989	818	72	57	67	49	79	74	81	52	57	1
1988	749	61	50	51	55	69	60	63	55	69	1
1987	673	55	35	46	55	58	67	69	62	48	1
1986	717	39	63	42	65	72	56	63	54	51	2
1985	651	49	49	59	55	57	60	56	51	55	-
1984	682	45	54	63	62	68	69	51	50	41	1
1983	660	51	54	62	57	53	79	57	49	56	3
1982	672	40	46	76	68	67	71	71	45	55	-
1981	726	56	64	73	70	86	68	61	55	45	-
1980	725	55	77	74	63	72	98	58	58	41	1
1979	713	58	84	86	67	70	65	63	39	43	-
Tuberculosis (010-018)											
All other, male											
1998	277	25	17	11	21	25	33	28	23	36	-
1997	282	24	23	25	22	24	30	25	31	28	-
1996	299	27	23	19	36	24	28	28	39	24	-
1995	340	37	26	25	23	40	35	28	27	31	1
1994	378	32	31	40	36	37	30	37	32	23	-
1993	427	42	39	32	33	38	29	41	29	34	-
1992	457	44	33	32	42	46	46	45	29	21	-
1991	455	36	39	44	41	33	32	36	35	29	-
1990	517	47	29	42	43	45	37	51	36	34	-
1989	560	57	37	53	33	54	46	50	32	32	1
1988	491	46	40	34	35	47	45	42	28	37	1
1987	445	40	28	35	41	36	38	46	34	23	1
1986	478	31	54	29	37	45	29	40	35	23	2
1985	432	40	36	46	35	38	36	33	28	35	-
1984	460	32	45	48	39	45	44	33	27	23	1
1983	458	40	42	48	40	38	54	35	35	31	2
1982	457	33	36	59	48	37	52	44	27	26	-
1981	502	42	37	51	49	64	50	43	35	29	-
1980	475	35	57	50	48	51	62	35	38	23	1
1979	462	37	54	54	51	48	46	41	19	23	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
All other, female											
1998	168	12	10	9	15	15	19	11	15	17	-
1997	189	19	12	9	15	18	20	26	21	22	-
1996	179	10	12	9	18	13	18	23	9	23	-
1995	192	10	9	7	10	19	23	21	20	28	-
1994	237	17	12	15	15	19	29	20	18	30	-
1993	235	9	12	18	18	27	27	24	16	27	-
1992	254	8	10	14	16	27	26	15	19	33	-
1991	280	15	11	13	29	25	23	21	22	28	-
1990	260	16	13	21	19	22	24	21	19	24	-
1989	258	15	20	14	16	25	28	31	20	25	-
1988	258	15	10	17	20	22	15	21	27	32	-
1987	228	15	7	11	14	22	29	23	28	25	-
1986	239	8	9	13	28	27	27	23	19	28	-
1985	219	9	13	13	20	19	24	23	23	20	-
1984	222	13	9	15	23	23	25	18	23	18	-
1983	202	11	12	14	17	15	25	22	14	25	1
1982	215	7	10	17	20	30	19	27	18	29	-
1981	224	14	27	22	21	22	18	18	20	16	-
1980	250	20	20	24	15	21	36	23	20	18	-
1979	251	21	30	32	16	22	19	22	20	20	-
Tuberculosis (010-018)											
Black, both sexes											
1998	346	33	20	18	32	30	40	26	28	40	-
1997	342	32	32	25	27	29	36	31	33	32	-
1996	369	32	30	22	41	26	33	36	31	34	-
1995	418	44	32	22	29	46	40	33	34	42	1
1994	499	38	36	44	41	44	46	42	36	39	-
1993	549	46	44	39	41	55	42	51	30	49	-
1992	612	45	39	40	50	54	58	46	41	45	-
1991	619	50	46	45	58	47	45	40	36	38	-
1990	668	59	32	55	50	54	47	56	46	48	-
1989	707	62	54	54	38	68	62	63	40	49	1
1988	641	56	42	42	45	61	51	45	43	54	1
1987	582	51	29	43	49	47	50	58	43	39	1
1986	628	38	58	34	54	68	49	49	41	42	2
1985	579	45	49	57	51	51	51	45	39	46	-
1984	619	42	52	59	55	62	62	43	42	32	1
1983	577	50	51	53	52	45	65	46	38	49	2
1982	587	35	39	71	65	58	63	57	31	46	-
1981	657	54	61	68	64	72	61	53	46	40	-
1980	651	50	72	67	60	66	87	48	52	33	1
1979	643	55	76	81	62	61	54	54	32	37	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Tuberculosis (010-018)											
Black, male											
1998	213	23	13	11	19	18	26	18	14	26	-
1997	204	19	21	19	17	18	23	15	18	14	-
1996	236	25	20	15	26	19	21	23	24	15	-
1995	271	35	25	18	20	33	24	19	22	18	1
1994	315	27	28	31	30	30	24	28	24	18	-
1993	361	38	37	28	28	31	23	31	19	27	-
1992	400	39	31	28	38	36	39	35	26	17	-
1991	390	35	37	36	38	28	24	27	21	18	-
1990	446	44	22	38	37	34	27	40	29	28	-
1989	485	49	35	45	23	47	39	36	24	29	1
1988	422	43	34	29	31	43	40	29	20	26	1
1987	395	38	24	32	39	31	32	37	22	19	1
1986	424	30	52	26	30	44	24	30	25	20	2
1985	393	39	36	45	32	35	29	27	21	31	-
1984	425	30	44	45	36	42	40	27	23	17	1
1983	409	40	39	41	37	34	44	31	27	26	2
1982	412	29	33	56	46	31	48	36	20	22	-
1981	455	41	35	47	45	57	45	37	29	25	-
1980	443	35	57	48	46	47	57	27	36	19	1
1979	423	37	51	52	49	44	37	35	14	17	-
Tuberculosis (010-018)											
Black, female											
1998	133	10	7	7	13	12	14	8	14	14	-
1997	138	13	11	6	10	11	13	16	15	18	-
1996	133	7	10	7	15	7	12	13	7	19	-
1995	147	9	7	4	9	13	16	14	12	24	-
1994	184	11	8	13	11	14	22	14	12	21	-
1993	188	8	7	11	13	24	19	20	11	22	-
1992	212	6	8	12	12	18	19	11	15	28	-
1991	229	15	9	9	20	19	21	13	15	20	-
1990	222	15	10	17	13	20	20	16	17	20	-
1989	222	13	19	9	15	21	23	27	16	20	-
1988	219	13	8	13	14	18	11	16	23	28	-
1987	187	13	5	11	10	16	18	21	21	20	-
1986	204	8	6	8	24	24	25	19	16	22	-
1985	186	6	13	12	19	16	22	18	18	15	-
1984	194	12	8	14	19	20	22	16	19	15	-
1983	168	10	12	12	15	11	21	15	11	23	-
1982	175	6	6	15	19	27	15	21	11	24	-
1981	202	13	26	21	19	15	16	16	17	15	-
1980	208	15	15	19	14	19	30	21	16	14	-
1979	220	18	25	29	13	17	17	19	18	20	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
All races, both sexes											
1998	815	48	36	35	68	69	100	104	114	145	-
1997	875	44	44	49	63	63	113	124	151	136	-
1996	912	46	47	58	66	75	105	121	125	161	-
1995	1,044	56	47	54	73	116	123	138	150	162	1
1994	1,129	62	44	57	85	109	124	159	137	194	2
1993	1,276	70	84	78	84	140	153	157	165	172	1
1992	1,315	65	56	80	118	109	187	162	158	171	-
1991	1,310	62	76	83	119	118	137	149	171	161	-
1990	1,378	68	61	79	112	146	161	176	159	174	-
1989	1,536	80	67	100	110	167	174	184	192	210	1
1988	1,488	83	71	97	119	144	179	177	190	215	1
1987	1,398	64	73	89	145	147	173	172	168	167	2
1986	1,407	57	85	91	135	151	152	191	152	181	3
1985	1,368	60	82	110	134	149	158	176	173	169	1
1984	1,342	65	77	119	124	157	164	168	158	148	-
1983	1,418	66	102	113	131	168	186	178	161	176	1
1982	1,468	51	93	129	139	178	184	219	140	198	-
1981	1,586	69	106	146	168	205	197	179	192	169	-
1980	1,638	70	129	148	165	209	243	201	167	185	2
1979	1,657	92	123	172	149	208	207	215	152	177	-
Respiratory tuberculosis (010-012)											
All races, male											
1998	528	35	30	27	50	44	59	67	75	73	-
1997	537	28	31	38	46	39	65	62	85	78	-
1996	600	31	37	48	51	54	69	76	79	79	-
1995	684	44	37	41	60	81	78	81	92	85	1
1994	722	44	32	44	65	77	81	94	73	100	2
1993	837	57	67	56	68	94	98	99	95	78	1
1992	853	61	51	65	86	71	127	99	86	77	-
1991	841	50	57	64	86	78	80	88	99	85	-
1990	907	55	43	57	79	94	102	114	98	92	-
1989	1,042	67	51	81	77	120	112	117	103	117	1
1988	986	68	59	80	94	106	117	114	99	102	1
1987	971	49	56	74	108	95	121	118	108	98	2
1986	948	44	75	70	89	101	102	121	92	89	3
1985	923	51	65	87	105	95	90	104	110	99	1
1984	932	52	65	90	89	110	120	107	98	83	-
1983	982	57	80	92	96	119	130	112	98	103	1
1982	1,014	43	76	110	105	120	128	142	83	102	-
1981	1,118	54	78	108	125	152	151	119	126	94	-
1980	1,148	54	96	112	127	158	162	126	122	114	2
1979	1,205	70	91	122	118	161	161	161	100	106	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
All races, female											
1998	287	13	6	8	18	25	41	37	39	72	-
1997	338	16	13	11	17	24	48	62	66	58	-
1996	312	15	10	10	15	21	36	45	46	82	-
1995	360	12	10	13	13	35	45	57	58	77	-
1994	407	18	12	13	20	32	43	65	64	94	-
1993	439	13	17	22	16	46	55	58	70	94	-
1992	462	4	5	15	32	38	60	63	72	94	-
1991	469	12	19	19	33	40	57	61	72	76	-
1990	471	13	18	22	33	52	59	62	61	82	-
1989	494	13	16	19	33	47	62	67	89	93	-
1988	502	15	12	17	25	38	62	63	91	113	-
1987	427	15	17	15	37	52	52	54	60	69	-
1986	459	13	10	21	46	50	50	70	60	92	-
1985	445	9	17	23	29	54	68	72	63	70	-
1984	410	13	12	29	35	47	44	61	60	65	-
1983	436	9	22	21	35	49	56	66	63	73	-
1982	454	8	17	19	34	58	56	77	57	96	-
1981	468	15	28	38	43	53	46	60	66	75	-
1980	490	16	33	36	38	51	81	75	45	71	-
1979	452	22	32	50	31	47	46	54	52	71	-
Respiratory tuberculosis (010-012)											
White, both sexes											
1998	495	16	20	21	42	44	58	75	83	103	-
1997	532	13	19	26	34	33	75	86	112	99	-
1996	551	21	18	35	27	47	72	82	85	120	-
1995	641	19	20	28	48	64	82	104	119	109	-
1994	678	25	12	27	43	74	81	108	97	148	2
1993	769	28	45	41	42	84	104	112	130	120	1
1992	780	26	24	43	69	56	128	114	118	128	-
1991	754	20	30	41	62	78	90	110	126	116	-
1990	812	24	37	38	68	93	112	116	118	129	-
1989	927	27	27	46	75	108	120	126	149	162	1
1988	957	35	33	62	82	92	132	137	147	159	1
1987	900	25	47	57	105	100	119	118	126	125	2
1986	859	27	36	55	83	96	111	137	108	135	1
1985	882	27	44	63	90	106	113	134	125	126	1
1984	828	28	31	66	76	99	114	129	115	117	-
1983	925	27	53	72	84	126	130	135	122	133	-
1982	962	20	57	69	87	124	134	162	105	153	-
1981	1,012	29	56	83	116	134	142	131	147	134	-
1980	1,067	27	65	86	113	151	167	154	123	149	1
1979	1,106	48	57	102	95	159	157	164	121	144	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
White, male											
1998	306	13	17	17	33	27	32	44	53	43	-
1997	309	10	13	20	26	20	39	41	60	54	-
1996	361	14	16	31	21	35	50	52	45	58	-
1995	400	15	15	20	40	45	49	59	69	55	-
1994	423	18	7	19	33	53	56	62	46	79	2
1993	490	22	34	30	38	60	71	67	73	47	1
1992	488	23	22	38	51	36	86	61	62	60	-
1991	474	18	20	33	50	51	51	64	70	60	-
1990	518	19	27	29	46	61	72	70	70	65	-
1989	600	23	22	35	52	77	78	77	77	87	1
1988	611	30	28	52	64	66	80	84	76	72	1
1987	623	18	35	46	75	66	88	80	84	76	2
1986	564	21	30	46	56	64	77	85	64	68	1
1985	585	24	35	48	73	64	62	81	83	70	1
1984	575	24	26	47	58	71	84	81	74	66	-
1983	616	25	40	56	61	88	91	80	68	76	-
1982	647	16	46	56	65	89	89	105	62	80	-
1981	702	23	48	65	85	95	106	85	96	72	-
1980	752	22	46	70	85	113	115	97	90	93	1
1979	812	40	46	73	71	121	124	125	82	86	-
Respiratory tuberculosis (010-012)											
White, female											
1998	189	3	3	4	9	17	26	31	30	60	-
1997	223	3	6	6	8	13	36	45	52	45	-
1996	190	7	2	4	6	12	22	30	40	62	-
1995	241	4	5	8	8	19	33	45	50	54	-
1994	255	7	5	8	10	21	25	46	51	69	-
1993	279	6	11	11	4	24	33	45	57	73	-
1992	292	3	2	5	18	20	42	53	56	68	-
1991	280	2	10	8	12	27	39	46	56	56	-
1990	294	5	10	9	22	32	40	46	48	64	-
1989	327	4	5	11	23	31	42	49	72	75	-
1988	346	5	5	10	18	26	52	53	71	87	-
1987	277	7	12	11	30	34	31	38	42	49	-
1986	295	6	6	9	27	32	34	52	44	67	-
1985	297	3	9	15	17	42	51	53	42	56	-
1984	253	4	5	19	18	28	30	48	41	51	-
1983	309	2	13	16	23	38	39	55	54	57	-
1982	315	4	11	13	22	35	45	57	43	73	-
1981	310	6	8	18	31	39	36	46	51	62	-
1980	315	5	19	16	28	38	52	57	33	56	-
1979	294	8	11	29	24	38	33	39	39	58	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
All other, both sexes											
1998	320	32	16	14	26	25	42	29	31	42	-
1997	343	31	25	23	29	30	38	38	39	37	-
1996	361	25	29	23	39	28	33	39	40	41	-
1995	403	37	27	26	25	52	41	34	31	53	1
1994	451	37	32	30	42	35	43	51	40	46	-
1993	507	42	39	37	42	56	49	45	35	52	-
1992	535	39	32	37	49	53	59	48	40	43	-
1991	556	42	46	42	57	40	47	39	45	45	-
1990	566	44	24	41	44	53	49	60	41	45	-
1989	609	53	40	54	35	59	54	58	43	48	-
1988	531	48	38	35	37	52	47	40	43	56	-
1987	498	39	26	32	40	47	54	54	42	42	-
1986	548	30	49	36	52	55	41	54	44	46	2
1985	486	33	38	47	44	43	45	42	48	43	-
1984	514	37	46	53	48	58	50	39	43	31	-
1983	493	39	49	41	47	42	56	43	39	43	1
1982	506	31	36	60	52	54	50	57	35	45	-
1981	574	40	50	63	52	71	55	48	45	35	-
1980	571	43	64	62	52	58	76	47	44	36	1
1979	551	44	66	70	54	49	50	51	31	33	-
Respiratory tuberculosis (010-012)											
All other, male											
1998	222	22	13	10	17	17	27	23	22	30	-
1997	228	18	18	18	20	19	26	21	25	24	-
1996	239	17	21	17	30	19	19	24	34	21	-
1995	284	29	22	21	20	36	29	22	23	30	1
1994	299	26	25	25	32	24	25	32	27	21	-
1993	347	35	33	26	30	34	27	32	22	31	-
1992	365	38	29	27	35	35	41	38	24	17	-
1991	367	32	37	31	36	27	29	24	29	25	-
1990	389	36	16	28	33	33	30	44	28	27	-
1989	442	44	29	46	25	43	34	40	26	30	-
1988	375	38	31	28	30	40	37	30	23	30	-
1987	348	31	21	28	33	29	33	38	24	22	-
1986	384	23	45	24	33	37	25	36	28	21	2
1985	338	27	30	39	32	31	28	23	27	29	-
1984	357	28	39	43	31	39	36	26	24	17	-
1983	366	32	40	36	35	31	39	32	30	27	1
1982	367	27	30	54	40	31	39	37	21	22	-
1981	416	31	30	43	40	57	45	34	30	22	-
1980	396	32	50	42	42	45	47	29	32	21	1
1979	393	30	45	49	47	40	37	36	18	20	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
All other, female											
1998	98	10	3	4	9	8	15	6	9	12	-
1997	115	13	7	5	9	11	12	17	14	13	-
1996	122	8	8	6	9	9	14	15	6	20	-
1995	119	8	5	5	5	16	12	12	8	23	-
1994	152	11	7	5	10	11	18	19	13	25	-
1993	160	7	6	11	12	22	22	13	13	21	-
1992	170	1	3	10	14	18	18	10	16	26	-
1991	189	10	9	11	21	13	18	15	16	20	-
1990	177	8	8	13	11	20	19	16	13	18	-
1989	167	9	11	8	10	16	20	18	17	18	-
1988	156	10	7	7	7	12	10	10	20	26	-
1987	150	8	5	4	7	18	21	16	18	20	-
1986	164	7	4	12	19	18	16	18	16	25	-
1985	148	6	8	8	12	12	17	19	21	14	-
1984	157	9	7	10	17	19	14	13	19	14	-
1983	127	7	9	5	12	11	17	11	9	16	-
1982	139	4	6	6	12	23	11	20	14	23	-
1981	158	9	20	20	12	14	10	14	15	13	-
1980	175	11	14	20	10	13	29	18	12	15	-
1979	158	14	21	21	7	9	13	15	13	13	-
Respiratory tuberculosis (010-012)											
Black, both sexes											
1998	251	28	12	12	23	20	30	21	22	32	-
1997	250	23	23	17	21	22	26	25	25	22	-
1996	284	23	25	19	30	20	23	30	24	31	-
1995	314	36	25	17	21	40	27	21	23	36	1
1994	368	29	28	24	35	29	32	38	28	33	-
1993	416	38	36	32	33	47	35	32	20	40	-
1992	463	37	29	31	42	43	47	37	33	35	-
1991	467	41	43	33	50	32	39	26	27	28	-
1990	491	41	20	37	36	43	39	48	33	38	-
1989	533	48	40	44	27	53	45	42	33	44	-
1988	454	45	33	27	31	46	41	31	32	42	-
1987	432	37	22	29	37	40	40	46	27	34	-
1986	478	29	47	28	44	52	36	41	35	38	2
1985	432	31	38	45	40	41	36	36	37	35	-
1984	463	34	45	49	41	52	45	33	35	24	-
1983	431	39	46	34	43	34	46	36	32	36	1
1982	444	26	33	55	50	48	45	44	23	40	-
1981	522	40	47	59	47	60	50	41	39	31	-
1980	518	40	61	56	50	54	67	39	39	29	1
1979	504	41	63	67	50	44	43	44	25	28	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Respiratory tuberculosis (010-012)											
Black, male											
1998	176	20	10	10	16	13	20	17	14	22	-
1997	165	14	16	13	16	15	20	13	15	11	-
1996	186	17	19	14	22	14	14	19	20	13	-
1995	225	28	21	15	17	30	20	14	18	17	1
1994	252	22	24	20	26	20	20	25	20	16	-
1993	294	32	32	23	26	28	21	23	12	24	-
1992	323	36	27	23	32	30	34	30	21	13	-
1991	319	31	36	26	35	23	22	18	17	16	-
1990	341	33	12	27	29	25	23	36	22	23	-
1989	386	40	29	41	18	38	27	27	20	27	-
1988	319	35	27	23	26	36	33	22	15	20	-
1987	310	30	18	25	32	26	28	31	15	18	-
1986	339	22	44	21	28	36	21	26	21	18	2
1985	306	27	30	38	29	29	21	20	21	25	-
1984	328	26	38	40	28	36	32	22	20	12	-
1983	327	32	37	30	32	27	31	29	25	22	1
1982	328	23	28	51	39	27	35	29	15	19	-
1981	376	31	28	39	36	51	41	29	25	18	-
1980	373	32	50	41	41	42	43	22	30	18	1
1979	363	30	44	48	45	36	31	31	13	15	-
Respiratory tuberculosis (010-012)											
Black, female											
1998	75	8	2	2	7	7	10	4	8	10	-
1997	85	9	7	4	5	7	6	12	10	11	-
1996	98	6	6	5	8	6	9	11	4	18	-
1995	89	8	4	2	4	10	7	7	5	19	-
1994	116	7	4	4	9	9	12	13	8	17	-
1993	122	6	4	9	7	19	14	9	8	16	-
1992	140	1	2	8	10	13	13	7	12	22	-
1991	148	10	7	7	15	9	17	8	10	12	-
1990	150	8	8	10	7	18	16	12	11	15	-
1989	147	8	11	3	9	15	18	15	13	17	-
1988	135	10	6	4	5	10	8	9	17	22	-
1987	122	7	4	4	5	14	12	15	12	16	-
1986	139	7	3	7	16	16	15	15	14	20	-
1985	126	4	8	7	11	12	15	16	16	10	-
1984	135	8	7	9	13	16	13	11	15	12	-
1983	104	7	9	4	11	7	15	7	7	14	-
1982	116	3	5	4	11	21	10	15	8	21	-
1981	146	9	19	20	11	9	9	12	14	13	-
1980	145	8	11	15	9	12	24	17	9	11	-
1979	141	11	19	19	5	8	12	13	12	13	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
All races, both sexes											
1998	297	10	20	25	23	26	34	30	32	32	-
1997	291	20	17	24	21	27	31	39	39	25	-
1996	290	22	14	13	30	24	41	28	35	31	-
1995	292	20	20	16	14	26	35	34	37	25	-
1994	349	23	23	37	25	38	42	29	32	27	-
1993	355	15	23	22	25	28	29	53	43	25	-
1992	390	21	18	23	21	43	39	36	26	44	-
1991	403	18	18	27	25	41	33	50	41	37	-
1990	432	28	31	38	40	37	38	34	37	35	-
1989	434	29	25	25	35	41	44	53	37	32	1
1988	433	18	23	27	35	38	43	57	29	41	1
1987	357	23	16	25	25	32	37	39	34	28	1
1986	375	18	23	23	30	34	36	39	34	30	-
1985	384	21	20	24	31	37	45	47	34	30	-
1984	387	13	22	28	31	35	50	37	42	28	1
1983	361	22	8	39	23	30	52	43	35	36	2
1982	339	11	22	29	28	29	42	46	35	33	1
1981	351	23	24	25	32	28	46	35	40	34	-
1980	340	19	25	24	27	40	50	35	33	29	-
1979	350	23	25	28	28	43	36	34	37	40	-
Other tuberculosis (013-018)											
All races, male											
1998	130	5	8	13	10	14	17	16	10	11	-
1997	125	11	10	12	10	13	9	15	13	8	-
1996	139	17	7	7	13	13	21	11	16	8	-
1995	130	13	10	7	7	12	13	15	13	8	-
1994	162	10	14	19	11	23	15	13	15	10	-
1993	181	11	12	12	11	17	15	21	19	12	-
1992	206	11	8	14	15	20	20	21	13	17	-
1991	207	12	9	23	14	16	14	25	18	15	-
1990	240	18	23	24	21	21	20	17	19	14	-
1989	242	18	13	15	18	23	26	24	17	14	1
1988	235	12	16	10	17	18	28	29	11	21	1
1987	198	15	12	13	15	17	17	20	14	11	1
1986	209	13	15	13	11	18	17	15	18	18	-
1985	212	16	14	15	15	21	19	22	16	17	-
1984	217	7	14	20	19	18	22	17	15	18	1
1983	194	12	4	24	11	20	29	17	19	16	1
1982	188	8	14	14	15	16	24	27	22	11	-
1981	181	18	11	15	20	12	21	21	16	15	-
1980	183	5	17	18	17	22	32	15	10	17	-
1979	166	11	14	11	11	25	19	18	14	16	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
All races, female											
1998	167	5	12	12	13	12	17	14	22	21	-
1997	166	9	7	12	11	14	22	24	26	17	-
1996	151	5	7	6	17	11	20	17	19	23	-
1995	162	7	10	9	7	14	22	19	24	17	-
1994	187	13	9	18	14	15	27	16	17	17	-
1993	174	4	11	10	14	11	14	32	24	13	-
1992	184	10	10	9	6	23	19	15	13	27	-
1991	196	6	9	4	11	25	19	25	23	22	-
1990	192	10	8	14	19	16	18	17	18	21	-
1989	192	11	12	10	17	18	18	29	20	18	-
1988	198	6	7	17	18	20	15	28	18	20	-
1987	159	8	4	12	10	15	20	19	20	17	-
1986	166	5	8	10	19	16	19	24	16	12	-
1985	172	5	6	9	16	16	26	25	18	13	-
1984	170	6	8	8	12	17	28	20	27	10	-
1983	167	10	4	15	12	10	23	26	16	20	1
1982	151	3	8	15	13	13	18	19	13	22	1
1981	170	5	13	10	12	16	25	14	24	19	-
1980	157	14	8	6	10	18	18	20	23	12	-
1979	184	12	11	17	17	18	17	16	23	24	-
Other tuberculosis (013-018)											
White, both sexes											
1998	172	5	9	19	13	11	24	20	25	21	-
1997	163	8	7	13	13	15	19	26	26	12	-
1996	173	10	8	8	15	15	28	16	27	25	-
1995	163	10	12	10	6	19	18	19	21	19	-
1994	185	11	12	12	16	17	26	23	22	20	-
1993	200	6	11	9	16	19	22	33	33	16	-
1992	214	8	7	14	12	23	26	24	18	33	-
1991	224	9	14	12	12	23	25	32	29	25	-
1990	221	9	13	16	22	23	26	22	23	22	-
1989	225	10	8	12	21	21	24	30	28	23	-
1988	215	5	11	11	17	21	30	34	17	28	-
1987	182	7	7	11	10	21	24	24	14	22	-
1986	206	9	9	17	17	17	21	30	24	25	-
1985	219	5	9	12	20	23	30	33	31	18	-
1984	219	5	14	18	17	25	31	25	35	18	-
1983	194	10	3	18	13	19	29	29	25	23	-
1982	173	2	12	13	12	16	21	32	25	23	1
1981	199	7	10	15	14	13	33	22	30	24	-
1980	186	7	12	12	16	26	28	24	19	24	-
1979	188	9	7	12	15	22	21	22	29	30	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
White, male											
1998	75	2	4	12	6	6	11	11	9	5	-
1997	71	5	5	5	8	8	5	11	7	4	-
1996	79	7	5	5	7	8	12	7	11	5	-
1995	74	5	6	3	4	8	7	9	9	7	-
1994	83	4	8	4	7	10	10	8	10	8	-
1993	101	4	6	6	8	13	13	12	12	9	-
1992	114	5	4	9	8	9	15	14	8	13	-
1991	119	8	7	10	9	10	11	13	12	11	-
1990	112	7	10	10	11	9	13	10	11	7	-
1989	124	5	5	8	10	12	14	14	11	12	-
1988	119	4	7	4	12	11	20	17	6	14	-
1987	101	6	5	6	7	10	12	12	4	10	-
1986	115	5	6	8	7	10	13	11	11	16	-
1985	118	3	8	8	12	14	11	12	15	11	-
1984	114	3	8	15	11	12	14	10	12	12	-
1983	102	4	2	12	6	13	14	14	14	12	-
1982	98	2	8	9	7	10	11	20	16	7	-
1981	95	7	4	7	11	5	16	12	11	8	-
1980	104	2	10	10	11	16	17	9	4	15	-
1979	97	4	5	6	7	17	10	13	13	13	-
Other tuberculosis (013-018)											
White, female											
1998	97	3	5	7	7	5	13	9	16	16	-
1997	92	3	2	8	5	7	14	15	19	8	-
1996	94	3	3	3	8	7	16	9	16	20	-
1995	89	5	6	7	2	11	11	10	12	12	-
1994	102	7	4	8	9	7	16	15	12	12	-
1993	99	2	5	3	8	6	9	21	21	7	-
1992	100	3	3	5	4	14	11	10	10	20	-
1991	105	1	7	2	3	13	14	19	17	14	-
1990	109	2	3	6	11	14	13	12	12	15	-
1989	101	5	3	4	11	9	10	16	17	11	-
1988	96	1	4	7	5	10	10	17	11	14	-
1987	81	1	2	5	3	11	12	12	10	12	-
1986	91	4	3	9	10	7	8	19	13	9	-
1985	101	2	1	4	8	9	19	21	16	7	-
1984	105	2	6	3	6	13	17	15	23	6	-
1983	92	6	1	6	7	6	15	15	11	11	-
1982	75	-	4	4	5	6	10	12	9	16	1
1981	104	-	6	8	3	8	17	10	19	16	-
1980	82	5	2	2	5	10	11	15	15	9	-
1979	91	5	2	6	8	5	11	9	16	17	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
All other, both sexes											
1998	125	5	11	6	10	15	10	10	7	11	-
1997	128	12	10	11	8	12	12	13	13	13	-
1996	117	12	6	5	15	9	13	12	8	6	-
1995	129	10	8	6	8	7	17	15	16	6	-
1994	164	12	11	25	9	21	16	6	10	7	-
1993	155	9	12	13	9	9	7	20	10	9	-
1992	176	13	11	9	9	20	13	12	8	11	-
1991	179	9	4	15	13	18	8	18	12	12	-
1990	211	19	18	22	18	14	12	12	14	13	-
1989	209	19	17	13	14	20	20	23	9	9	1
1988	218	13	12	16	18	17	13	23	12	13	1
1987	175	16	9	14	15	11	13	15	20	6	1
1986	169	9	14	6	13	17	15	9	10	5	-
1985	165	16	11	12	11	14	15	14	3	12	-
1984	168	8	8	10	14	10	19	12	7	10	1
1983	167	12	5	21	10	11	23	14	10	13	2
1982	166	9	10	16	16	13	21	14	10	10	-
1981	152	16	14	10	18	15	13	13	10	10	-
1980	154	12	13	12	11	14	22	11	14	5	-
1979	162	14	18	16	13	21	15	12	8	10	-
Other tuberculosis (013-018)											
All other, male											
1998	55	3	4	1	4	8	6	5	1	6	-
1997	54	6	5	7	2	5	4	4	6	4	-
1996	60	10	2	2	6	5	9	4	5	3	-
1995	56	8	4	4	3	4	6	6	4	1	-
1994	79	6	6	15	4	13	5	5	5	2	-
1993	80	7	6	6	3	4	2	9	7	3	-
1992	92	6	4	5	7	11	5	7	5	4	-
1991	88	4	2	13	5	6	3	12	6	4	-
1990	128	11	13	14	10	12	7	7	8	7	-
1989	118	13	8	7	8	11	12	10	6	2	1
1988	116	8	9	6	5	7	8	12	5	7	1
1987	97	9	7	7	8	7	5	8	10	1	1
1986	94	8	9	5	4	8	4	4	7	2	-
1985	94	13	6	7	3	7	8	10	1	6	-
1984	103	4	6	5	8	6	8	7	3	6	1
1983	92	8	2	12	5	7	15	3	5	4	1
1982	90	6	6	5	8	6	13	7	6	4	-
1981	86	11	7	8	9	7	5	9	5	7	-
1980	79	3	7	8	6	6	15	6	6	2	-
1979	69	7	9	5	4	8	9	5	1	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
All other, female											
1998	70	2	7	5	6	7	4	5	6	5	-
1997	74	6	5	4	6	7	8	9	7	9	-
1996	57	2	4	3	9	4	4	8	3	3	-
1995	73	2	4	2	5	3	11	9	12	5	-
1994	85	6	5	10	5	8	11	1	5	5	-
1993	75	2	6	7	6	5	5	11	3	6	-
1992	84	7	7	4	2	9	8	5	3	7	-
1991	91	5	2	2	8	12	5	6	6	8	-
1990	83	8	5	8	8	2	5	5	6	6	-
1989	91	6	9	6	6	9	8	13	3	7	-
1988	102	5	3	10	13	10	5	11	7	6	-
1987	78	7	2	7	7	4	8	7	10	5	-
1986	75	1	5	1	9	9	11	5	3	3	-
1985	71	3	5	5	8	7	7	4	2	6	-
1984	65	4	2	5	6	4	11	5	4	4	-
1983	75	4	3	9	5	4	8	11	5	9	1
1982	76	3	4	11	8	7	8	7	4	6	-
1981	66	5	7	2	9	8	8	4	5	3	-
1980	75	9	6	4	5	8	7	5	8	3	-
1979	93	7	9	11	9	13	6	7	7	7	-
Other tuberculosis (013-018)											
Black, both sexes											
1998	95	5	8	6	9	10	10	5	6	8	-
1997	92	9	9	8	6	7	10	6	8	10	-
1996	85	9	5	3	11	6	10	6	7	3	-
1995	104	8	7	5	8	6	13	12	11	6	-
1994	131	9	8	20	6	15	14	4	8	6	-
1993	133	8	8	7	8	8	7	19	10	9	-
1992	149	8	10	9	8	11	11	9	8	10	-
1991	152	9	3	12	8	15	6	14	9	10	-
1990	177	18	12	18	14	11	8	8	13	10	-
1989	174	14	14	10	11	15	17	21	7	5	1
1988	187	11	9	15	14	15	10	14	11	12	1
1987	150	14	7	14	12	7	10	12	16	5	1
1986	150	9	11	6	10	16	13	8	6	4	-
1985	147	14	11	12	11	10	15	9	2	11	-
1984	156	8	7	10	14	10	17	10	7	8	1
1983	146	11	5	19	9	11	19	10	6	13	1
1982	143	9	6	16	15	10	18	13	8	6	-
1981	135	14	14	9	17	12	11	12	7	9	-
1980	133	10	11	11	10	12	20	9	13	4	-
1979	139	14	13	14	12	17	11	10	7	9	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other tuberculosis (013-018)											
Black, male											
1998	37	3	3	1	3	5	6	1	-	4	-
1997	39	5	5	6	1	3	3	2	3	3	-
1996	50	8	1	1	4	5	7	4	4	2	-
1995	46	7	4	3	3	3	4	5	4	1	-
1994	63	5	4	11	4	10	4	3	4	2	-
1993	67	6	5	5	2	3	2	8	7	3	-
1992	77	3	4	5	6	6	5	5	5	4	-
1991	71	4	1	10	3	5	2	9	4	2	-
1990	105	11	10	11	8	9	4	4	7	5	-
1989	99	9	6	4	5	9	12	9	4	2	1
1988	103	8	7	6	5	7	7	7	5	6	1
1987	85	8	6	7	7	5	4	6	7	1	1
1986	85	8	8	5	2	8	3	4	4	2	-
1985	87	12	6	7	3	6	8	7	-	6	-
1984	97	4	6	5	8	6	8	5	3	5	1
1983	82	8	2	11	5	7	13	2	2	4	1
1982	84	6	5	5	7	4	13	7	5	3	-
1981	79	10	7	8	9	6	4	8	4	7	-
1980	70	3	7	7	5	5	14	5	6	1	-
1979	60	7	7	4	4	8	6	4	1	2	-
Other tuberculosis (013-018)											
Black, female											
1998	58	2	5	5	6	5	4	4	6	4	-
1997	53	4	4	2	5	4	7	4	5	7	-
1996	35	1	4	2	7	1	3	2	3	1	-
1995	58	1	3	2	5	3	9	7	7	5	-
1994	68	4	4	9	2	5	10	1	4	4	-
1993	66	2	3	2	6	5	5	11	3	6	-
1992	72	5	6	4	2	5	6	4	3	6	-
1991	81	5	2	2	5	10	4	5	5	8	-
1990	72	7	2	7	6	2	4	4	6	5	-
1989	75	5	8	6	6	6	5	12	3	3	-
1988	84	3	2	9	9	8	3	7	6	6	-
1987	65	6	1	7	5	2	6	6	9	4	-
1986	65	1	3	1	8	8	10	4	2	2	-
1985	60	2	5	5	8	4	7	2	2	5	-
1984	59	4	1	5	6	4	9	5	4	3	-
1983	64	3	3	8	4	4	6	8	4	9	-
1982	59	3	1	11	8	6	5	6	3	3	-
1981	56	4	7	1	8	6	7	4	3	2	-
1980	63	7	4	4	5	7	6	4	7	3	-
1979	79	7	6	10	8	9	5	6	6	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
All races, both sexes											
1998	5	-	-	-	-	-	-	-	-	-	-
1997	6	-	-	-	-	-	-	-	-	-	-
1996	4	-	-	-	-	-	-	-	-	-	-
1995	6	-	-	-	-	-	-	-	-	-	-
1994	8	-	-	-	-	-	-	-	-	-	-
1993	7	-	-	-	-	-	-	-	-	-	-
1992	5	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	12	-	-	-	-	-	-	-	-	-	-
1989	12	-	-	-	-	-	-	-	-	-	-
1988	4	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	6	-	-	-	-	-	1	-	-	-	-
1985	4	-	-	-	-	-	-	-	-	-	-
1984	7	-	-	-	-	-	-	-	-	-	-
1983	5	-	-	-	-	-	-	-	-	-	-
1982	4	-	-	-	-	-	-	-	-	-	-
1981	6	-	-	-	-	-	-	-	-	-	-
1980	11	-	-	-	-	-	-	-	-	-	-
1979	6	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
All races, male											
1998	5	-	-	-	-	-	-	-	-	-	-
1997	3	-	-	-	-	-	-	-	-	-	-
1996	2	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	2	-	-	-	-	-	-	-	-	-	-
1993	3	-	-	-	-	-	-	-	-	-	-
1992	3	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	4	-	-	-	-	-	-	-	-	-	-
1989	4	-	-	-	-	-	-	-	-	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	4	-	-	-	-	-	1	-	-	-	-
1985	3	-	-	-	-	-	-	-	-	-	-
1984	2	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	4	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
All races, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	3	-	-	-	-	-	-	-	-	-	-
1996	2	-	-	-	-	-	-	-	-	-	-
1995	4	-	-	-	-	-	-	-	-	-	-
1994	6	-	-	-	-	-	-	-	-	-	-
1993	4	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	8	-	-	-	-	-	-	-	-	-	-
1989	8	-	-	-	-	-	-	-	-	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	5	-	-	-	-	-	-	-	-	-	-
1983	4	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	4	-	-	-	-	-	-	-	-	-	-
1980	7	-	-	-	-	-	-	-	-	-	-
1979	6	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
White, both sexes											
1998	4	-	-	-	-	-	-	-	-	-	-
1997	4	-	-	-	-	-	-	-	-	-	-
1996	4	-	-	-	-	-	-	-	-	-	-
1995	6	-	-	-	-	-	-	-	-	-	-
1994	7	-	-	-	-	-	-	-	-	-	-
1993	5	-	-	-	-	-	-	-	-	-	-
1992	4	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	9	-	-	-	-	-	-	-	-	-	-
1989	10	-	-	-	-	-	-	-	-	-	-
1988	3	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	3	-	-	-	-	-	1	-	-	-	-
1985	2	-	-	-	-	-	-	-	-	-	-
1984	6	-	-	-	-	-	-	-	-	-	-
1983	5	-	-	-	-	-	-	-	-	-	-
1982	4	-	-	-	-	-	-	-	-	-	-
1981	5	-	-	-	-	-	-	-	-	-	-
1980	8	-	-	-	-	-	-	-	-	-	-
1979	4	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
White, male											
1998	4	-	-	-	-	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	2	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	2	-	-	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	-	-	-	-	-	-
1992	3	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	3	-	-	-	-	-	-	-	-	-	-
1989	3	-	-	-	-	-	-	-	-	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	3	-	-	-	-	-	1	-	-	-	-
1985	2	-	-	-	-	-	-	-	-	-	-
1984	2	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	4	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
White, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	2	-	-	-	-	-	-	-	-	-	-
1995	4	-	-	-	-	-	-	-	-	-	-
1994	5	-	-	-	-	-	-	-	-	-	-
1993	3	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	6	-	-	-	-	-	-	-	-	-	-
1989	7	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	4	-	-	-	-	-	-	-	-	-	-
1983	4	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	3	-	-	-	-	-	-	-	-	-	-
1980	4	-	-	-	-	-	-	-	-	-	-
1979	4	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
All other, both sexes											
1998	1	-	-	-	-	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	3	-	-	-	-	-	-	-	-	-	-
1989	2	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	3	-	-	-	-	-	-	-	-	-	-
1985	2	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
All other, male											
1998	1	-	-	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	1	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	3	-	-	-	-	-	-	-	-	-	-
1989	2	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	3	-	-	-	-	-	-	-	-	-	-
1985	2	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Whooping cough (033)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	1	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Whooping cough (033)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
All races, both sexes											
1998	3	-	-	-	1	1	-	1	-	-	-
1997	5	-	1	-	-	-	-	-	1	3	-
1996	3	-	-	-	1	-	1	-	1	-	-
1995	3	-	-	-	-	-	-	-	1	2	-
1994	3	-	-	-	-	1	-	1	-	-	-
1993	8	-	-	1	-	-	-	1	-	6	-
1992	1	-	-	-	-	-	-	-	-	1	-
1991	1	-	-	-	-	-	1	-	-	-	-
1990	5	-	-	-	-	-	2	-	1	2	-
1989	8	-	-	1	1	1	-	1	-	4	-
1988	4	-	-	1	-	1	2	-	-	-	-
1987	3	-	-	-	2	-	-	-	-	1	-
1986	6	-	2	-	2	-	-	-	1	1	-
1985	6	-	-	-	1	-	1	-	-	3	-
1984	6	1	1	-	-	-	1	-	1	-	-
1983	1	-	-	-	-	-	-	1	-	-	-
1982	10	1	1	-	1	1	-	-	1	4	-
1981	9	-	-	-	1	-	1	2	2	1	-
1980	10	-	1	-	2	1	-	1	3	2	-
1979	12	1	-	-	-	1	2	-	3	3	-
Scarlet fever and erysipelas (034.1-035)											
All races, male											
1998	2	-	-	-	1	1	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	1	-
1996	1	-	-	-	1	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	1	-	-	-
1993	2	-	-	1	-	-	-	-	-	1	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	1	-	-	-	-	-	1	-	-	-	-
1989	2	-	-	1	-	-	-	-	-	1	-
1988	1	-	-	-	-	-	1	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	3	-	2	-	1	-	-	-	-	-	-
1985	2	-	-	-	-	-	1	-	-	1	-
1984	2	-	-	-	-	-	-	-	1	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	5	1	1	-	-	-	-	-	-	2	-
1981	5	-	-	-	1	-	-	1	-	1	-
1980	3	-	-	-	-	1	-	-	1	1	-
1979	6	1	-	-	-	-	1	-	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
All races, female											
1998	1	-	-	-	-	-	-	1	-	-	-
1997	4	-	1	-	-	-	-	-	1	2	-
1996	2	-	-	-	-	-	1	-	1	-	-
1995	3	-	-	-	-	-	-	-	1	2	-
1994	2	-	-	-	-	1	-	-	-	-	-
1993	6	-	-	-	-	-	-	1	-	5	-
1992	1	-	-	-	-	-	-	-	-	1	-
1991	1	-	-	-	-	-	1	-	-	-	-
1990	4	-	-	-	-	-	1	-	1	2	-
1989	6	-	-	-	1	1	-	1	-	3	-
1988	3	-	-	1	-	1	1	-	-	-	-
1987	3	-	-	-	2	-	-	-	-	1	-
1986	3	-	-	-	1	-	-	-	1	1	-
1985	4	-	-	-	1	-	-	-	-	2	-
1984	4	1	1	-	-	-	1	-	-	-	-
1983	1	-	-	-	-	-	-	1	-	-	-
1982	5	-	-	-	1	1	-	-	1	2	-
1981	4	-	-	-	-	-	1	1	2	-	-
1980	7	-	1	-	2	-	-	1	2	1	-
1979	6	-	-	-	-	1	1	-	2	2	-
Scarlet fever and erysipelas (034.1-035)											
White, both sexes											
1998	3	-	-	-	1	1	-	1	-	-	-
1997	5	-	1	-	-	-	-	-	1	3	-
1996	3	-	-	-	1	-	1	-	1	-	-
1995	3	-	-	-	-	-	-	-	1	2	-
1994	3	-	-	-	-	1	-	1	-	-	-
1993	8	-	-	1	-	-	-	1	-	6	-
1992	1	-	-	-	-	-	-	-	-	1	-
1991	1	-	-	-	-	-	1	-	-	-	-
1990	5	-	-	-	-	-	2	-	1	2	-
1989	8	-	-	1	1	1	-	1	-	4	-
1988	3	-	-	-	-	1	2	-	-	-	-
1987	3	-	-	-	2	-	-	-	-	1	-
1986	6	-	2	-	2	-	-	-	1	1	-
1985	5	-	-	-	1	-	1	-	-	3	-
1984	5	1	1	-	-	-	1	-	1	-	-
1983	1	-	-	-	-	-	-	1	-	-	-
1982	8	-	-	-	1	1	-	-	1	4	-
1981	9	-	-	-	1	-	1	2	2	1	-
1980	10	-	1	-	2	1	-	1	3	2	-
1979	11	1	-	-	-	1	2	-	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
White, male											
1998	2	-	-	-	1	1	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	1	-
1996	1	-	-	-	1	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	-	-	-	-	-	-	1	-	-	-
1993	2	-	-	1	-	-	-	-	-	1	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	1	-	-	-	-	-	1	-	-	-	-
1989	2	-	-	1	-	-	-	-	-	1	-
1988	1	-	-	-	-	-	1	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	3	-	2	-	1	-	-	-	-	-	-
1985	2	-	-	-	-	-	1	-	-	1	-
1984	2	-	-	-	-	-	-	-	1	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	3	-	-	-	-	-	-	-	-	2	-
1981	5	-	-	-	1	-	-	1	-	1	-
1980	3	-	-	-	-	1	-	-	1	1	-
1979	5	1	-	-	-	-	1	-	1	1	-
Scarlet fever and erysipelas (034.1-035)											
White, female											
1998	1	-	-	-	-	-	-	1	-	-	-
1997	4	-	1	-	-	-	-	-	1	2	-
1996	2	-	-	-	-	-	1	-	1	-	-
1995	3	-	-	-	-	-	-	-	1	2	-
1994	2	-	-	-	-	1	-	-	-	-	-
1993	6	-	-	-	-	-	-	1	-	5	-
1992	1	-	-	-	-	-	-	-	-	1	-
1991	1	-	-	-	-	-	1	-	-	-	-
1990	4	-	-	-	-	-	1	-	1	2	-
1989	6	-	-	-	1	1	-	1	-	3	-
1988	2	-	-	-	-	1	1	-	-	-	-
1987	3	-	-	-	2	-	-	-	-	1	-
1986	3	-	-	-	1	-	-	-	1	1	-
1985	3	-	-	-	1	-	-	-	-	2	-
1984	3	1	1	-	-	-	1	-	-	-	-
1983	1	-	-	-	-	-	-	1	-	-	-
1982	5	-	-	-	1	1	-	-	1	2	-
1981	4	-	-	-	-	-	1	1	2	-	-
1980	7	-	1	-	2	-	-	1	2	1	-
1979	6	-	-	-	-	1	1	-	2	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
All other, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	1	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	2	1	1	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (034.1-035)											
All other, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	2	1	1	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	1	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (034.1-035)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	1	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	1	-	1	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Scarlet fever and erysipelas (034.1-035)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	1	-	1	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (034.1-035)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	1	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
All races, both sexes											
1998	234	12	11	5	-	8	2	5	4	10	-
1997	309	8	12	6	9	7	7	6	6	14	-
1996	290	12	11	10	6	7	8	10	3	7	-
1995	273	16	8	5	5	2	6	6	8	14	-
1994	276	11	12	5	7	5	7	5	8	9	-
1993	260	9	6	9	7	4	9	9	5	7	-
1992	201	6	7	6	4	3	6	6	5	8	-
1991	198	2	4	3	5	7	4	4	1	8	-
1990	215	3	5	5	8	5	6	7	3	6	-
1989	273	10	5	8	10	11	8	7	7	7	-
1988	278	12	6	1	11	4	6	7	6	5	-
1987	258	5	7	8	8	4	7	5	2	3	-
1986	286	9	3	5	12	8	8	6	6	5	-
1985	257	7	6	6	8	7	10	4	6	7	-
1984	300	8	8	12	12	7	9	4	6	6	-
1983	299	6	9	10	10	10	7	11	5	6	-
1982	364	5	6	7	15	13	9	9	5	4	-
1981	459	17	14	17	11	13	12	10	12	7	-
1980	387	12	9	13	10	16	7	8	7	6	-
1979	404	11	9	22	16	10	14	5	5	6	-
Meningococcal infection (036)											
All races, male											
1998	132	4	6	4	-	3	1	1	2	2	-
1997	152	4	7	3	6	2	1	1	1	1	-
1996	141	5	4	2	2	3	2	3	1	-	-
1995	137	6	4	-	3	1	3	-	1	1	-
1994	132	5	6	3	1	2	5	2	1	1	-
1993	124	2	2	4	1	1	3	2	-	2	-
1992	98	3	1	2	1	1	4	1	1	-	-
1991	103	1	2	-	2	3	1	2	1	1	-
1990	113	1	2	4	3	2	2	2	2	3	-
1989	127	5	4	2	3	5	5	1	1	2	-
1988	143	4	2	-	7	2	-	3	1	2	-
1987	137	2	5	4	1	-	2	2	-	1	-
1986	146	5	2	2	5	6	2	-	2	2	-
1985	127	2	3	2	4	1	2	3	-	-	-
1984	146	3	1	2	4	5	2	1	1	1	-
1983	157	3	1	4	6	5	2	3	2	-	-
1982	177	-	1	3	2	3	5	2	1	1	-
1981	233	10	5	6	4	6	2	5	2	2	-
1980	204	7	3	5	3	6	2	4	2	-	-
1979	234	7	5	8	7	4	4	4	2	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
All races, female											
1998	102	8	5	1	-	5	1	4	2	8	-
1997	157	4	5	3	3	5	6	5	5	13	-
1996	149	7	7	8	4	4	6	7	2	7	-
1995	136	10	4	5	2	1	3	6	7	13	-
1994	144	6	6	2	6	3	2	3	7	8	-
1993	136	7	4	5	6	3	6	7	5	5	-
1992	103	3	6	4	3	2	2	5	4	8	-
1991	95	1	2	3	3	4	3	2	-	7	-
1990	102	2	3	1	5	3	4	5	1	3	-
1989	146	5	1	6	7	6	3	6	6	5	-
1988	135	8	4	1	4	2	6	4	5	3	-
1987	121	3	2	4	7	4	5	3	2	2	-
1986	140	4	1	3	7	2	6	6	4	3	-
1985	130	5	3	4	4	6	8	1	6	7	-
1984	154	5	7	10	8	2	7	3	5	5	-
1983	142	3	8	6	4	5	5	8	3	6	-
1982	187	5	5	4	13	10	4	7	4	3	-
1981	226	7	9	11	7	7	10	5	10	5	-
1980	183	5	6	8	7	10	5	4	5	6	-
1979	170	4	4	14	9	6	10	1	3	1	-
Meningococcal infection (036)											
White, both sexes											
1998	180	10	10	4	-	8	1	4	4	9	-
1997	236	6	10	4	9	7	5	5	6	13	-
1996	223	9	10	8	3	7	8	10	3	7	-
1995	207	15	4	4	5	2	5	6	8	14	-
1994	232	10	11	4	6	5	6	4	8	9	-
1993	201	7	5	6	5	4	8	8	5	6	-
1992	167	6	7	5	4	3	4	6	5	6	-
1991	165	2	3	3	4	6	3	3	1	8	-
1990	166	3	4	5	5	4	6	6	3	6	-
1989	214	9	4	6	10	11	8	7	4	7	-
1988	198	6	4	-	10	4	5	6	6	5	-
1987	192	4	5	8	7	3	6	4	2	3	-
1986	216	7	3	4	11	7	7	4	6	5	-
1985	210	4	5	6	7	4	10	4	6	6	-
1984	226	6	5	11	11	7	7	4	5	4	-
1983	229	4	7	9	8	9	6	11	5	5	-
1982	282	4	5	7	12	12	7	8	4	4	-
1981	380	12	13	14	10	13	12	9	10	6	-
1980	306	8	8	10	10	11	7	8	7	5	-
1979	310	8	7	18	13	8	11	4	5	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
White, male											
1998	98	3	6	4	-	3	-	-	2	1	-
1997	111	3	6	2	6	2	1	1	1	1	-
1996	103	4	3	1	1	3	2	3	1	-	-
1995	97	6	1	-	3	1	2	-	1	1	-
1994	107	5	5	2	1	2	5	1	1	1	-
1993	89	2	2	3	1	1	2	1	-	2	-
1992	79	3	1	2	1	1	2	1	1	-	-
1991	87	1	2	-	2	3	1	2	1	1	-
1990	80	1	1	4	1	1	2	1	2	3	-
1989	103	4	4	1	3	5	5	1	1	2	-
1988	102	2	2	-	7	2	-	2	1	2	-
1987	100	1	3	4	-	-	2	1	-	1	-
1986	113	4	2	2	5	5	2	-	2	2	-
1985	107	2	3	2	4	1	2	3	-	-	-
1984	107	2	-	2	3	5	1	1	1	-	-
1983	122	2	1	4	4	4	1	3	2	-	-
1982	132	-	1	3	1	2	3	2	1	1	-
1981	191	8	4	5	3	6	2	4	1	2	-
1980	158	4	3	4	3	3	2	4	2	-	-
1979	179	7	5	6	6	3	3	3	2	4	-
Meningococcal infection (036)											
White, female											
1998	82	7	4	-	-	5	1	4	2	8	-
1997	125	3	4	2	3	5	4	4	5	12	-
1996	120	5	7	7	2	4	6	7	2	7	-
1995	110	9	3	4	2	1	3	6	7	13	-
1994	125	5	6	2	5	3	1	3	7	8	-
1993	112	5	3	3	4	3	6	7	5	4	-
1992	88	3	6	3	3	2	2	5	4	6	-
1991	78	1	1	3	2	3	2	1	-	7	-
1990	86	2	3	1	4	3	4	5	1	3	-
1989	111	5	-	5	7	6	3	6	3	5	-
1988	96	4	2	-	3	2	5	4	5	3	-
1987	92	3	2	4	7	3	4	3	2	2	-
1986	103	3	1	2	6	2	5	4	4	3	-
1985	103	2	2	4	3	3	8	1	6	6	-
1984	119	4	5	9	8	2	6	3	4	4	-
1983	107	2	6	5	4	5	5	8	3	5	-
1982	150	4	4	4	11	10	4	6	3	3	-
1981	189	4	9	9	7	7	10	5	9	4	-
1980	148	4	5	6	7	8	5	4	5	5	-
1979	131	1	2	12	7	5	8	1	3	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
All other, both sexes											
1998	54	2	1	1	-	-	1	1	-	1	-
1997	73	2	2	2	-	-	2	1	-	1	-
1996	67	3	1	2	3	-	-	-	-	-	-
1995	66	1	4	1	-	-	1	-	-	-	-
1994	44	1	1	1	1	-	1	1	-	-	-
1993	59	2	1	3	2	-	1	1	-	1	-
1992	34	-	-	1	-	-	2	-	-	2	-
1991	33	-	1	-	1	1	1	1	-	-	-
1990	49	-	1	-	3	1	-	1	-	-	-
1989	59	1	1	2	-	-	-	-	3	-	-
1988	80	6	2	1	1	-	1	1	-	-	-
1987	66	1	2	-	1	1	1	1	-	-	-
1986	70	2	-	1	1	1	1	2	-	-	-
1985	47	3	1	-	1	3	-	-	-	1	-
1984	74	2	3	1	1	-	2	-	1	2	-
1983	70	2	2	1	2	1	1	-	-	1	-
1982	82	1	1	-	3	1	2	1	1	-	-
1981	79	5	1	3	1	-	-	1	2	1	-
1980	81	4	1	3	-	5	-	-	-	1	-
1979	94	3	2	4	3	2	3	1	-	1	-
Meningococcal infection (036)											
All other, male											
1998	34	1	-	-	-	-	1	1	-	1	-
1997	41	1	1	1	-	-	-	-	-	-	-
1996	38	1	1	1	1	-	-	-	-	-	-
1995	40	-	3	-	-	-	1	-	-	-	-
1994	25	-	1	1	-	-	-	1	-	-	-
1993	35	-	-	1	-	-	1	1	-	-	-
1992	19	-	-	-	-	-	2	-	-	-	-
1991	16	-	-	-	-	-	-	-	-	-	-
1990	33	-	1	-	2	1	-	1	-	-	-
1989	24	1	-	1	-	-	-	-	-	-	-
1988	41	2	-	-	-	-	-	1	-	-	-
1987	37	1	2	-	1	-	-	1	-	-	-
1986	33	1	-	-	-	1	-	-	-	-	-
1985	20	-	-	-	-	-	-	-	-	-	-
1984	39	1	1	-	1	-	1	-	-	1	-
1983	35	1	-	-	2	1	1	-	-	-	-
1982	45	-	-	-	1	1	2	-	-	-	-
1981	42	2	1	1	1	-	-	1	1	-	-
1980	46	3	-	1	-	3	-	-	-	-	-
1979	55	-	-	2	1	1	1	1	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
All other, female											
1998	20	1	1	1	-	-	-	-	-	-	-
1997	32	1	1	1	-	-	2	1	-	1	-
1996	29	2	-	1	2	-	-	-	-	-	-
1995	26	1	1	1	-	-	-	-	-	-	-
1994	19	1	-	-	1	-	1	-	-	-	-
1993	24	2	1	2	2	-	-	-	-	1	-
1992	15	-	-	1	-	-	-	-	-	2	-
1991	17	-	1	-	1	1	1	1	-	-	-
1990	16	-	-	-	1	-	-	-	-	-	-
1989	35	-	1	1	-	-	-	-	3	-	-
1988	39	4	2	1	1	-	1	-	-	-	-
1987	29	-	-	-	-	1	1	-	-	-	-
1986	37	1	-	1	1	-	1	2	-	-	-
1985	27	3	1	-	1	3	-	-	-	1	-
1984	35	1	2	1	-	-	1	-	1	1	-
1983	35	1	2	1	-	-	-	-	-	1	-
1982	37	1	1	-	2	-	-	1	1	-	-
1981	37	3	-	2	-	-	-	-	1	1	-
1980	35	1	1	2	-	2	-	-	-	1	-
1979	39	3	2	2	2	1	2	-	-	-	-
Meningococcal infection (036)											
Black, both sexes											
1998	51	2	1	1	-	-	1	1	-	1	-
1997	63	2	2	2	-	-	2	-	-	1	-
1996	58	2	1	2	1	-	-	-	-	-	-
1995	57	1	4	1	-	-	1	-	-	-	-
1994	36	1	1	1	1	-	-	1	-	-	-
1993	49	2	1	1	2	-	-	1	-	1	-
1992	31	-	-	1	-	-	2	-	-	2	-
1991	29	-	1	-	1	-	1	1	-	-	-
1990	43	-	-	-	3	-	-	1	-	-	-
1989	50	1	-	1	-	-	-	-	3	-	-
1988	71	6	2	1	1	-	1	1	-	-	-
1987	65	1	2	-	1	1	1	1	-	-	-
1986	66	2	-	1	1	1	1	2	-	-	-
1985	44	3	-	-	1	3	-	-	-	1	-
1984	70	2	3	1	1	-	2	-	1	2	-
1983	67	2	2	1	2	1	1	-	-	1	-
1982	75	1	1	-	2	1	2	1	1	-	-
1981	75	5	1	3	1	-	-	1	1	1	-
1980	72	4	1	2	-	5	-	-	-	1	-
1979	83	2	-	4	3	1	3	-	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningococcal infection (036)											
Black, male											
1998	32	1	-	-	-	-	1	1	-	1	-
1997	35	1	1	1	-	-	-	-	-	-	-
1996	35	1	1	1	1	-	-	-	-	-	-
1995	34	-	3	-	-	-	1	-	-	-	-
1994	20	-	1	1	-	-	-	1	-	-	-
1993	28	-	-	-	-	-	-	1	-	-	-
1992	18	-	-	-	-	-	2	-	-	-	-
1991	15	-	-	-	-	-	-	-	-	-	-
1990	28	-	-	-	2	-	-	1	-	-	-
1989	22	1	-	-	-	-	-	-	-	-	-
1988	37	2	-	-	-	-	-	1	-	-	-
1987	36	1	2	-	1	-	-	1	-	-	-
1986	31	1	-	-	-	1	-	-	-	-	-
1985	19	-	-	-	-	-	-	-	-	-	-
1984	37	1	1	-	1	-	1	-	-	1	-
1983	33	1	-	-	2	1	1	-	-	-	-
1982	44	-	-	-	1	1	2	-	-	-	-
1981	40	2	1	1	1	-	-	1	1	-	-
1980	41	3	-	1	-	3	-	-	-	-	-
1979	49	-	-	2	1	1	1	-	-	1	-
Meningococcal infection (036)											
Black, female											
1998	19	1	1	1	-	-	-	-	-	-	-
1997	28	1	1	1	-	-	2	-	-	1	-
1996	23	1	-	1	-	-	-	-	-	-	-
1995	23	1	1	1	-	-	-	-	-	-	-
1994	16	1	-	-	1	-	-	-	-	-	-
1993	21	2	1	1	2	-	-	-	-	1	-
1992	13	-	-	1	-	-	-	-	-	2	-
1991	14	-	1	-	1	-	1	1	-	-	-
1990	15	-	-	-	1	-	-	-	-	-	-
1989	28	-	-	1	-	-	-	-	3	-	-
1988	34	4	2	1	1	-	1	-	-	-	-
1987	29	-	-	-	-	1	1	-	-	-	-
1986	35	1	-	1	1	-	1	2	-	-	-
1985	25	3	-	-	1	3	-	-	-	1	-
1984	33	1	2	1	-	-	1	-	1	1	-
1983	34	1	2	1	-	-	-	-	-	1	-
1982	31	1	1	-	1	-	-	1	1	-	-
1981	35	3	-	2	-	-	-	-	-	1	-
1980	31	1	1	1	-	2	-	-	-	1	-
1979	34	2	-	2	2	-	2	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
All races, both sexes											
1998	23,731	525	706	898	1,195	1,820	2,654	3,461	3,970	7,107	2
1997	22,396	540	648	787	1,065	1,685	2,494	3,294	3,732	6,874	1
1996	21,423	471	571	687	1,072	1,640	2,418	3,190	3,591	6,498	1
1995	20,965	476	497	671	1,068	1,556	2,429	3,079	3,554	6,281	1
1994	20,360	421	496	672	1,016	1,562	2,285	2,954	3,453	6,185	1
1993	20,634	387	454	642	967	1,650	2,347	2,990	3,499	6,360	2
1992	19,667	403	448	604	1,027	1,600	2,295	2,858	3,321	5,810	4
1991	19,691	323	440	635	1,033	1,589	2,198	2,953	3,355	5,793	3
1990	19,169	329	438	632	1,058	1,582	2,194	2,843	3,181	5,551	4
1989	19,333	351	430	622	1,073	1,547	2,117	2,875	3,239	5,650	2
1988	20,925	327	454	706	1,248	1,822	2,460	3,131	3,601	5,862	3
1987	19,916	268	486	741	1,261	1,846	2,424	2,974	3,210	5,414	3
1986	18,795	301	443	764	1,220	1,721	2,194	2,837	3,020	5,049	6
1985	17,182	255	412	755	1,214	1,584	2,068	2,633	2,796	4,321	3
1984	15,028	268	398	673	1,117	1,377	1,930	2,120	2,327	3,730	1
1983	13,394	216	388	657	1,018	1,279	1,651	2,013	2,032	3,196	4
1982	11,493	195	363	622	882	1,085	1,468	1,654	1,736	2,606	4
1981	10,443	233	333	574	803	1,041	1,243	1,441	1,576	2,241	4
1980	9,438	173	357	571	690	961	1,174	1,286	1,455	1,967	3
1979	8,024	179	323	449	600	861	1,036	1,137	1,118	1,545	-
Septicemia (038)											
All races, male											
1998	10,225	301	383	485	621	914	1,324	1,649	1,651	2,149	-
1997	9,655	306	353	376	576	891	1,299	1,621	1,514	2,047	1
1996	9,246	289	321	389	589	864	1,198	1,492	1,492	1,868	1
1995	8,991	271	273	383	574	807	1,210	1,435	1,464	1,848	1
1994	8,810	252	294	367	553	840	1,179	1,402	1,368	1,812	-
1993	8,916	232	251	356	498	883	1,188	1,437	1,503	1,820	-
1992	8,527	238	268	315	549	853	1,181	1,366	1,313	1,712	2
1991	8,610	184	266	342	580	863	1,110	1,423	1,314	1,742	2
1990	8,542	188	253	342	588	878	1,159	1,348	1,313	1,641	2
1989	8,489	202	250	337	597	840	1,116	1,338	1,309	1,678	1
1988	9,132	189	248	379	690	961	1,303	1,486	1,433	1,715	2
1987	8,969	153	259	417	696	1,004	1,301	1,396	1,295	1,715	3
1986	8,786	175	261	441	714	958	1,177	1,402	1,299	1,661	2
1985	8,017	155	232	428	710	863	1,179	1,256	1,163	1,396	1
1984	7,068	146	223	384	636	774	1,007	1,033	953	1,302	1
1983	6,417	120	225	384	594	740	854	994	887	1,083	2
1982	5,469	102	209	347	523	578	780	771	745	964	2
1981	5,047	137	196	338	451	575	642	684	684	836	4
1980	4,684	99	203	333	415	537	665	618	637	728	3
1979	3,972	109	183	237	343	472	536	565	517	606	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
All races, female											
1998	13,506	224	323	413	574	906	1,330	1,812	2,319	4,958	2
1997	12,741	234	295	411	489	794	1,195	1,673	2,218	4,827	-
1996	12,177	182	250	298	483	776	1,220	1,698	2,099	4,630	-
1995	11,974	205	224	288	494	749	1,219	1,644	2,090	4,433	-
1994	11,550	169	202	305	463	722	1,106	1,552	2,085	4,373	1
1993	11,718	155	203	286	469	767	1,159	1,553	1,996	4,540	2
1992	11,140	165	180	289	478	747	1,114	1,492	2,008	4,098	2
1991	11,081	139	174	293	453	726	1,088	1,530	2,041	4,051	1
1990	10,627	141	185	290	470	704	1,035	1,495	1,868	3,910	2
1989	10,844	149	180	285	476	707	1,001	1,537	1,930	3,972	1
1988	11,793	138	206	327	558	861	1,157	1,645	2,168	4,147	1
1987	10,947	115	227	324	565	842	1,123	1,578	1,915	3,699	-
1986	10,009	126	182	323	506	763	1,017	1,435	1,721	3,388	4
1985	9,165	100	180	327	504	721	889	1,377	1,633	2,925	2
1984	7,960	122	175	289	481	603	923	1,087	1,374	2,428	-
1983	6,977	96	163	273	424	539	797	1,019	1,145	2,113	2
1982	6,024	93	154	275	359	507	688	883	991	1,642	2
1981	5,396	96	137	236	352	466	601	757	892	1,405	-
1980	4,754	74	154	238	275	424	509	668	818	1,239	-
1979	4,052	70	140	212	257	389	500	572	601	939	-
Septicemia (038)											
White, both sexes											
1998	18,855	318	482	610	842	1,380	2,118	2,820	3,334	6,067	-
1997	17,832	343	439	526	748	1,277	1,944	2,702	3,188	5,876	1
1996	17,130	285	375	446	777	1,282	1,912	2,645	3,068	5,575	-
1995	16,784	294	348	452	779	1,203	1,916	2,537	3,029	5,410	1
1994	16,452	268	337	503	724	1,224	1,815	2,479	2,899	5,384	1
1993	16,696	243	312	459	714	1,269	1,926	2,489	2,927	5,524	-
1992	15,792	263	295	406	762	1,203	1,841	2,376	2,819	5,049	1
1991	15,884	205	290	441	780	1,222	1,764	2,455	2,863	5,066	3
1990	15,456	204	288	442	794	1,235	1,758	2,365	2,695	4,848	3
1989	15,557	223	271	441	795	1,185	1,713	2,378	2,773	4,912	1
1988	16,943	202	295	496	924	1,397	2,004	2,585	3,091	5,122	2
1987	15,936	159	310	537	943	1,420	1,967	2,475	2,703	4,650	1
1986	15,156	196	280	545	887	1,375	1,784	2,350	2,570	4,383	5
1985	13,740	152	283	541	902	1,223	1,672	2,150	2,326	3,761	3
1984	11,976	162	266	480	852	1,074	1,520	1,752	1,950	3,238	1
1983	10,696	125	266	480	752	993	1,288	1,676	1,714	2,798	2
1982	9,192	122	254	428	677	876	1,169	1,364	1,483	2,267	3
1981	8,311	152	240	426	612	823	983	1,216	1,352	1,916	2
1980	7,461	116	235	402	533	717	933	1,048	1,245	1,704	1
1979	6,376	129	211	327	456	668	828	922	973	1,352	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
White, male											
1998	8,114	188	258	325	444	717	1,088	1,356	1,405	1,837	-
1997	7,633	197	232	257	418	670	1,031	1,350	1,300	1,749	1
1996	7,247	170	208	249	420	674	957	1,268	1,259	1,604	-
1995	7,040	164	189	250	423	631	952	1,185	1,258	1,558	1
1994	7,018	159	198	285	391	659	945	1,169	1,165	1,589	-
1993	7,087	141	168	247	377	697	982	1,200	1,274	1,531	-
1992	6,735	155	167	215	415	658	955	1,152	1,097	1,471	-
1991	6,812	109	178	233	442	673	880	1,197	1,117	1,522	2
1990	6,786	113	162	238	453	700	941	1,133	1,107	1,431	1
1989	6,728	134	160	231	452	645	910	1,135	1,118	1,432	1
1988	7,270	115	155	265	511	733	1,068	1,246	1,219	1,485	2
1987	7,096	89	163	303	533	783	1,061	1,174	1,100	1,459	1
1986	6,962	113	165	308	530	771	947	1,155	1,118	1,424	2
1985	6,321	91	158	295	529	680	976	1,014	961	1,210	1
1984	5,602	87	153	281	510	620	800	855	803	1,111	1
1983	5,119	70	157	299	452	586	691	819	745	940	-
1982	4,357	61	152	249	404	475	624	641	641	827	2
1981	4,003	89	146	255	347	469	508	576	599	699	2
1980	3,672	65	139	236	326	401	541	500	552	620	1
1979	3,121	79	125	169	254	372	420	462	446	531	-
Septicemia (038)											
White, female											
1998	10,741	130	224	285	398	663	1,030	1,464	1,929	4,230	-
1997	10,199	146	207	269	330	607	913	1,352	1,888	4,127	-
1996	9,883	115	167	197	357	608	955	1,377	1,809	3,971	-
1995	9,744	130	159	202	356	572	964	1,352	1,771	3,852	-
1994	9,434	109	139	218	333	565	870	1,310	1,734	3,795	1
1993	9,609	102	144	212	337	572	944	1,289	1,653	3,993	-
1992	9,057	108	128	191	347	545	886	1,224	1,722	3,578	1
1991	9,072	96	112	208	338	549	884	1,258	1,746	3,544	1
1990	8,670	91	126	204	341	535	817	1,232	1,588	3,417	2
1989	8,829	89	111	210	343	540	803	1,243	1,655	3,480	-
1988	9,673	87	140	231	413	664	936	1,339	1,872	3,637	-
1987	8,840	70	147	234	410	637	906	1,301	1,603	3,191	-
1986	8,194	83	115	237	357	604	837	1,195	1,452	2,959	3
1985	7,419	61	125	246	373	543	696	1,136	1,365	2,551	2
1984	6,374	75	113	199	342	454	720	897	1,147	2,127	-
1983	5,577	55	109	181	300	407	597	857	969	1,858	2
1982	4,835	61	102	179	273	401	545	723	842	1,440	1
1981	4,308	63	94	171	265	354	475	640	753	1,217	-
1980	3,789	51	96	166	207	316	392	548	693	1,084	-
1979	3,255	50	86	158	202	296	408	460	527	821	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
All other, both sexes											
1998	4,876	207	224	288	353	440	536	641	636	1,040	2
1997	4,564	197	209	261	317	408	550	592	544	998	-
1996	4,293	186	196	241	295	358	506	545	523	923	1
1995	4,181	182	149	219	289	353	513	542	525	871	-
1994	3,908	153	159	169	292	338	470	475	554	801	-
1993	3,938	144	142	183	253	381	421	501	572	836	2
1992	3,875	140	153	198	265	397	454	482	502	761	3
1991	3,807	118	150	194	253	367	434	498	492	727	-
1990	3,713	125	150	190	264	347	436	478	486	703	1
1989	3,776	128	159	181	278	362	404	497	466	738	1
1988	3,982	125	159	210	324	425	456	546	510	740	1
1987	3,980	109	176	204	318	426	457	499	507	764	2
1986	3,639	105	163	219	333	346	410	487	450	666	1
1985	3,442	103	129	214	312	361	396	483	470	560	-
1984	3,052	106	132	193	265	303	410	368	377	492	-
1983	2,698	91	122	177	266	286	363	337	318	398	2
1982	2,301	73	109	194	205	209	299	290	253	339	1
1981	2,132	81	93	148	191	218	260	225	224	325	2
1980	1,977	57	122	169	157	244	241	238	210	263	2
1979	1,648	50	112	122	144	193	208	215	145	193	-
Septicemia (038)											
All other, male											
1998	2,111	113	125	160	177	197	236	293	246	312	-
1997	2,022	109	121	119	158	221	268	271	214	298	-
1996	1,999	119	113	140	169	190	241	224	233	264	1
1995	1,951	107	84	133	151	176	258	250	206	290	-
1994	1,792	93	96	82	162	181	234	233	203	223	-
1993	1,829	91	83	109	121	186	206	237	229	289	-
1992	1,792	83	101	100	134	195	226	214	216	241	2
1991	1,798	75	88	109	138	190	230	226	197	220	-
1990	1,756	75	91	104	135	178	218	215	206	210	1
1989	1,761	68	90	106	145	195	206	203	191	246	-
1988	1,862	74	93	114	179	228	235	240	214	230	-
1987	1,873	64	96	114	163	221	240	222	195	256	2
1986	1,824	62	96	133	184	187	230	247	181	237	-
1985	1,696	64	74	133	181	183	203	242	202	186	-
1984	1,466	59	70	103	126	154	207	178	150	191	-
1983	1,298	50	68	85	142	154	163	175	142	143	2
1982	1,112	41	57	98	119	103	156	130	104	137	-
1981	1,044	48	50	83	104	106	134	108	85	137	2
1980	1,012	34	64	97	89	136	124	118	85	108	2
1979	851	30	58	68	89	100	116	103	71	75	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
All other, female											
1998	2,765	94	99	128	176	243	300	348	390	728	2
1997	2,542	88	88	142	159	187	282	321	330	700	-
1996	2,294	67	83	101	126	168	265	321	290	659	-
1995	2,230	75	65	86	138	177	255	292	319	581	-
1994	2,116	60	63	87	130	157	236	242	351	578	-
1993	2,109	53	59	74	132	195	215	264	343	547	2
1992	2,083	57	52	98	131	202	228	268	286	520	1
1991	2,009	43	62	85	115	177	204	272	295	507	-
1990	1,957	50	59	86	129	169	218	263	280	493	-
1989	2,015	60	69	75	133	167	198	294	275	492	1
1988	2,120	51	66	96	145	197	221	306	296	510	1
1987	2,107	45	80	90	155	205	217	277	312	508	-
1986	1,815	43	67	86	149	159	180	240	269	429	1
1985	1,746	39	55	81	131	178	193	241	268	374	-
1984	1,586	47	62	90	139	149	203	190	227	301	-
1983	1,400	41	54	92	124	132	200	162	176	255	-
1982	1,189	32	52	96	86	106	143	160	149	202	1
1981	1,088	33	43	65	87	112	126	117	139	188	-
1980	965	23	58	72	68	108	117	120	125	155	-
1979	797	20	54	54	55	93	92	112	74	118	-
Septicemia (038)											
Black, both sexes											
1998	4,487	194	206	261	325	412	483	594	582	963	2
1997	4,241	182	200	238	289	383	508	543	512	926	-
1996	3,996	174	184	220	272	333	472	507	489	864	1
1995	3,845	166	136	206	263	322	472	507	472	809	-
1994	3,639	142	143	153	277	309	430	443	513	759	-
1993	3,632	132	128	165	232	352	392	457	530	779	2
1992	3,607	127	144	180	241	372	410	453	470	714	3
1991	3,535	109	140	182	239	342	411	450	454	671	-
1990	3,465	116	137	172	247	328	413	452	453	650	1
1989	3,555	120	151	167	264	343	386	462	442	688	1
1988	3,740	121	145	199	312	395	434	510	475	690	1
1987	3,748	103	159	194	300	411	429	468	475	720	2
1986	3,417	94	157	202	325	325	384	461	419	614	1
1985	3,257	91	124	204	293	340	374	465	435	533	-
1984	2,897	102	124	186	257	285	391	357	353	473	-
1983	2,550	84	112	170	253	268	347	318	301	376	2
1982	2,162	68	105	178	194	198	276	276	240	319	1
1981	2,028	74	89	139	185	209	244	215	215	311	2
1980	1,877	53	118	163	149	237	230	231	195	253	2
1979	1,561	42	107	116	137	186	198	204	141	183	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Septicemia (038)											
Black, male											
1998	1,923	105	116	145	163	185	208	268	222	280	-
1997	1,871	102	115	108	148	208	248	250	195	268	-
1996	1,844	111	110	127	154	174	221	208	213	240	1
1995	1,779	101	79	129	136	159	240	233	177	255	-
1994	1,659	89	87	74	153	169	214	217	184	206	-
1993	1,673	83	74	102	111	172	191	217	203	262	-
1992	1,668	78	96	92	122	185	202	203	197	224	2
1991	1,656	69	83	103	128	177	216	199	174	200	-
1990	1,624	69	84	92	127	167	204	198	191	191	1
1989	1,643	66	87	100	139	183	192	183	178	223	-
1988	1,729	73	86	107	172	208	224	216	199	203	-
1987	1,760	59	89	110	154	214	222	204	183	237	2
1986	1,697	57	91	125	180	172	214	230	164	212	-
1985	1,595	55	70	130	170	170	191	233	180	175	-
1984	1,383	57	64	98	124	144	198	171	135	179	-
1983	1,221	45	64	83	133	144	152	163	132	135	2
1982	1,046	40	55	93	112	97	145	124	97	128	-
1981	987	44	48	78	101	103	123	101	81	129	2
1980	958	31	63	94	85	131	115	115	78	105	2
1979	797	24	55	67	83	96	109	94	68	72	-
Septicemia (038)											
Black, female											
1998	2,564	89	90	116	162	227	275	326	360	683	2
1997	2,370	80	85	130	141	175	260	293	317	658	-
1996	2,152	63	74	93	118	159	251	299	276	624	-
1995	2,066	65	57	77	127	163	232	274	295	554	-
1994	1,980	53	56	79	124	140	216	226	329	553	-
1993	1,959	49	54	63	121	180	201	240	327	517	2
1992	1,939	49	48	88	119	187	208	250	273	490	1
1991	1,879	40	57	79	111	165	195	251	280	471	-
1990	1,841	47	53	80	120	161	209	254	262	459	-
1989	1,912	54	64	67	125	160	194	279	264	465	1
1988	2,011	48	59	92	140	187	210	294	276	487	1
1987	1,988	44	70	84	146	197	207	264	292	483	-
1986	1,720	37	66	77	145	153	170	231	255	402	1
1985	1,662	36	54	74	123	170	183	232	255	358	-
1984	1,514	45	60	88	133	141	193	186	218	294	-
1983	1,329	39	48	87	120	124	195	155	169	241	-
1982	1,116	28	50	85	82	101	131	152	143	191	1
1981	1,041	30	41	61	84	106	121	114	134	182	-
1980	919	22	55	69	64	106	115	116	117	148	-
1979	764	18	52	49	54	90	89	110	73	111	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
All races, both sexes											
1998	45	2	1	3	1	4	3	6	4	11	-
1997	62	3	2	3	2	6	8	8	11	7	-
1996	73	1	3	2	4	5	9	11	5	15	-
1995	65	4	5	2	2	4	7	5	9	9	-
1994	79	4	3	2	3	4	14	9	13	8	-
1993	80	2	1	4	3	8	5	10	9	7	-
1992	91	5	1	6	6	4	11	9	8	15	-
1991	93	5	-	3	4	5	8	6	10	16	-
1990	106	5	4	3	4	3	6	13	12	13	-
1989	105	4	3	6	4	7	8	12	11	15	-
1988	85	2	3	1	6	4	6	17	7	12	-
1987	98	1	2	2	10	7	13	13	16	8	-
1986	80	-	3	3	2	5	12	16	15	13	-
1985	80	1	3	6	6	6	9	12	10	14	-
1984	105	1	1	5	7	14	17	19	15	15	-
1983	121	1	5	6	5	16	20	25	14	14	-
1982	126	2	3	10	9	10	19	19	19	16	-
1981	136	3	6	7	8	13	27	28	14	14	-
1980	154	2	3	4	8	22	29	34	18	22	-
1979	180	2	5	7	22	24	33	29	18	25	-
Syphilis (090-097)											
All races, male											
1998	25	1	-	2	1	2	1	5	3	4	-
1997	39	3	2	3	2	5	4	4	6	2	-
1996	50	1	3	2	4	4	4	9	3	9	-
1995	41	3	3	1	2	4	5	3	3	2	-
1994	46	4	1	2	1	2	9	4	8	2	-
1993	52	2	1	4	3	6	2	5	7	3	-
1992	60	4	1	3	6	3	7	5	5	5	-
1991	52	4	-	2	4	4	4	2	8	5	-
1990	63	5	4	2	2	-	2	6	7	6	-
1989	65	2	2	6	1	5	2	9	5	8	-
1988	51	2	3	-	3	3	3	10	6	6	-
1987	57	1	2	1	6	3	5	9	9	3	-
1986	42	-	1	1	1	3	7	10	7	5	-
1985	49	-	2	6	5	4	5	7	5	8	-
1984	67	1	-	3	6	12	6	15	9	9	-
1983	68	1	4	3	4	9	10	11	6	10	-
1982	76	2	1	8	7	6	8	14	9	8	-
1981	76	1	4	3	4	8	15	15	9	6	-
1980	92	-	2	3	4	11	14	23	13	14	-
1979	107	2	1	4	16	14	25	14	13	9	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
All races, female											
1998	20	1	1	1	-	2	2	1	1	7	-
1997	23	-	-	-	-	1	4	4	5	5	-
1996	23	-	-	-	-	1	5	2	2	6	-
1995	24	1	2	1	-	-	2	2	6	7	-
1994	33	-	2	-	2	2	5	5	5	6	-
1993	28	-	-	-	-	2	3	5	2	4	-
1992	31	1	-	3	-	1	4	4	3	10	-
1991	41	1	-	1	-	1	4	4	2	11	-
1990	43	-	-	1	2	3	4	7	5	7	-
1989	40	2	1	-	3	2	6	3	6	7	-
1988	34	-	-	1	3	1	3	7	1	6	-
1987	41	-	-	1	4	4	8	4	7	5	-
1986	38	-	2	2	1	2	5	6	8	8	-
1985	31	1	1	-	1	2	4	5	5	6	-
1984	38	-	1	2	1	2	11	4	6	6	-
1983	53	-	1	3	1	7	10	14	8	4	-
1982	50	-	2	2	2	4	11	5	10	8	-
1981	60	2	2	4	4	5	12	13	5	8	-
1980	62	2	1	1	4	11	15	11	5	8	-
1979	73	-	4	3	6	10	8	15	5	16	-
Syphilis (090-097)											
White, both sexes											
1998	21	1	1	1	-	2	1	1	3	8	-
1997	37	2	2	2	-	3	3	3	7	6	-
1996	39	1	2	2	2	3	4	4	3	8	-
1995	36	2	2	-	1	3	3	2	7	7	-
1994	41	3	2	1	1	-	9	6	8	5	-
1993	32	-	-	1	2	4	3	2	7	2	-
1992	43	1	-	2	1	2	8	6	4	10	-
1991	44	3	-	2	1	3	2	5	6	10	-
1990	41	-	3	-	3	-	2	8	8	7	-
1989	47	3	-	3	3	2	4	7	5	10	-
1988	40	1	-	1	2	4	3	8	5	8	-
1987	51	-	-	1	6	5	6	8	9	4	-
1986	43	-	3	2	1	3	4	8	9	6	-
1985	49	-	1	4	2	2	7	9	8	9	-
1984	55	1	1	4	2	8	7	5	12	7	-
1983	70	1	1	5	3	8	14	15	8	9	-
1982	84	-	3	5	5	7	14	13	13	12	-
1981	88	2	4	5	6	10	18	18	6	9	-
1980	93	2	1	2	5	11	21	16	11	15	-
1979	101	1	2	3	8	16	18	19	9	15	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
White, male											
1998	12	1	-	1	-	1	-	1	2	4	-
1997	20	2	2	2	-	2	1	1	4	1	-
1996	26	1	2	2	2	2	1	4	1	5	-
1995	22	1	2	-	1	3	1	1	3	2	-
1994	24	3	1	1	1	-	5	3	6	1	-
1993	22	-	-	1	2	3	1	1	5	1	-
1992	23	1	-	2	1	1	5	4	1	2	-
1991	27	2	-	1	1	3	1	2	6	2	-
1990	21	-	3	-	1	-	-	3	3	3	-
1989	27	1	-	3	-	2	1	6	1	4	-
1988	25	1	-	-	2	3	1	5	4	4	-
1987	32	-	-	1	5	2	3	6	5	2	-
1986	22	-	1	1	-	2	1	5	4	3	-
1985	29	-	1	4	2	-	3	4	5	6	-
1984	33	1	-	2	2	6	3	4	7	3	-
1983	35	1	1	2	3	3	5	9	2	6	-
1982	47	-	1	5	4	4	4	10	5	5	-
1981	47	1	3	2	3	5	9	9	4	4	-
1980	50	-	-	2	2	5	8	9	11	7	-
1979	60	1	1	2	7	8	14	10	7	4	-
Syphilis (090-097)											
White, female											
1998	9	-	1	-	-	1	1	-	1	4	-
1997	17	-	-	-	-	1	2	2	3	5	-
1996	13	-	-	-	-	1	3	-	2	3	-
1995	14	1	-	-	-	-	2	1	4	5	-
1994	17	-	1	-	-	-	4	3	2	4	-
1993	10	-	-	-	-	1	2	1	2	1	-
1992	20	-	-	-	-	1	3	2	3	8	-
1991	17	1	-	1	-	-	1	3	-	8	-
1990	20	-	-	-	2	-	2	5	5	4	-
1989	20	2	-	-	3	-	3	1	4	6	-
1988	15	-	-	1	-	1	2	3	1	4	-
1987	19	-	-	-	1	3	3	2	4	2	-
1986	21	-	2	1	1	1	3	3	5	3	-
1985	20	-	-	-	-	2	4	5	3	3	-
1984	22	-	1	2	-	2	4	1	5	4	-
1983	35	-	-	3	-	5	9	6	6	3	-
1982	37	-	2	-	1	3	10	3	8	7	-
1981	41	1	1	3	3	5	9	9	2	5	-
1980	43	2	1	-	3	6	13	7	-	8	-
1979	41	-	1	1	1	8	4	9	2	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
All other, both sexes											
1998	24	1	-	2	1	2	2	5	1	3	-
1997	25	1	-	1	2	3	5	5	4	1	-
1996	34	-	1	-	2	2	5	7	2	7	-
1995	29	2	3	2	1	1	4	3	2	2	-
1994	38	1	1	1	2	4	5	3	5	3	-
1993	48	2	1	3	1	4	2	8	2	5	-
1992	48	4	1	4	5	2	3	3	4	5	-
1991	49	2	-	1	3	2	6	1	4	6	-
1990	65	5	1	3	1	3	4	5	4	6	-
1989	58	1	3	3	1	5	4	5	6	5	-
1988	45	1	3	-	4	-	3	9	2	4	-
1987	47	1	2	1	4	2	7	5	7	4	-
1986	37	-	-	1	1	2	8	8	6	7	-
1985	31	1	2	2	4	4	2	3	2	5	-
1984	50	-	-	1	5	6	10	14	3	8	-
1983	51	-	4	1	2	8	6	10	6	5	-
1982	42	2	-	5	4	3	5	6	6	4	-
1981	48	1	2	2	2	3	9	10	8	5	-
1980	61	-	2	2	3	11	8	18	7	7	-
1979	79	1	3	4	14	8	15	10	9	10	-
Syphilis (090-097)											
All other, male											
1998	13	-	-	1	1	1	1	4	1	-	-
1997	19	1	-	1	2	3	3	3	2	1	-
1996	24	-	1	-	2	2	3	5	2	4	-
1995	19	2	1	1	1	1	4	2	-	-	-
1994	22	1	-	1	-	2	4	1	2	1	-
1993	30	2	1	3	1	3	1	4	2	2	-
1992	37	3	1	1	5	2	2	1	4	3	-
1991	25	2	-	1	3	1	3	-	2	3	-
1990	42	5	1	2	1	-	2	3	4	3	-
1989	38	1	2	3	1	3	1	3	4	4	-
1988	26	1	3	-	1	-	2	5	2	2	-
1987	25	1	2	-	1	1	2	3	4	1	-
1986	20	-	-	-	1	1	6	5	3	2	-
1985	20	-	1	2	3	4	2	3	-	2	-
1984	34	-	-	1	4	6	3	11	2	6	-
1983	33	-	3	1	1	6	5	2	4	4	-
1982	29	2	-	3	3	2	4	4	4	3	-
1981	29	-	1	1	1	3	6	6	5	2	-
1980	42	-	2	1	2	6	6	14	2	7	-
1979	47	1	-	2	9	6	11	4	6	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
All other, female											
1998	11	1	-	1	-	1	1	1	-	3	-
1997	6	-	-	-	-	-	2	2	2	-	-
1996	10	-	-	-	-	-	2	2	-	3	-
1995	10	-	2	1	-	-	-	1	2	2	-
1994	16	-	1	-	2	2	1	2	3	2	-
1993	18	-	-	-	-	1	1	4	-	3	-
1992	11	1	-	3	-	-	1	2	-	2	-
1991	24	-	-	-	-	1	3	1	2	3	-
1990	23	-	-	1	-	3	2	2	-	3	-
1989	20	-	1	-	-	2	3	2	2	1	-
1988	19	-	-	-	3	-	1	4	-	2	-
1987	22	-	-	1	3	1	5	2	3	3	-
1986	17	-	-	1	-	1	2	3	3	5	-
1985	11	1	1	-	1	-	-	-	2	3	-
1984	16	-	-	-	1	-	7	3	1	2	-
1983	18	-	1	-	1	2	1	8	2	1	-
1982	13	-	-	2	1	1	1	2	2	1	-
1981	19	1	1	1	1	-	3	4	3	3	-
1980	19	-	-	1	1	5	2	4	5	-	-
1979	32	-	3	2	5	2	4	6	3	5	-
Syphilis (090-097)											
Black, both sexes											
1998	21	1	-	2	-	2	1	4	1	3	-
1997	24	1	-	-	2	3	5	5	4	1	-
1996	33	-	1	-	2	2	5	6	2	7	-
1995	26	2	2	2	1	1	4	3	1	2	-
1994	36	1	1	1	1	4	4	3	5	3	-
1993	47	2	1	3	1	4	2	8	2	5	-
1992	46	3	1	3	5	2	3	3	4	5	-
1991	48	1	-	1	3	2	6	1	4	6	-
1990	64	5	1	3	1	3	4	5	4	5	-
1989	56	1	3	2	1	5	4	5	5	5	-
1988	45	1	3	-	4	-	3	9	2	4	-
1987	45	1	2	1	4	2	7	4	6	4	-
1986	37	-	-	1	1	2	8	8	6	7	-
1985	29	1	2	2	4	3	2	3	2	5	-
1984	46	-	-	1	5	4	10	13	2	8	-
1983	50	-	4	1	2	8	6	10	6	5	-
1982	40	2	-	5	4	3	5	6	5	4	-
1981	47	1	2	1	2	3	9	10	8	5	-
1980	59	-	2	2	3	11	8	18	5	7	-
1979	76	1	2	4	14	8	15	10	9	9	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Syphilis (090-097)											
Black, male											
1998	11	-	-	1	-	1	1	3	1	-	-
1997	18	1	-	-	2	3	3	3	2	1	-
1996	24	-	1	-	2	2	3	5	2	4	-
1995	18	2	1	1	1	1	4	2	-	-	-
1994	21	1	-	1	-	2	3	1	2	1	-
1993	30	2	1	3	1	3	1	4	2	2	-
1992	36	2	1	1	5	2	2	1	4	3	-
1991	24	1	-	1	3	1	3	-	2	3	-
1990	41	5	1	2	1	-	2	3	4	2	-
1989	36	1	2	2	1	3	1	3	3	4	-
1988	26	1	3	-	1	-	2	5	2	2	-
1987	23	1	2	-	1	1	2	2	3	1	-
1986	20	-	-	-	1	1	6	5	3	2	-
1985	19	-	1	2	3	3	2	3	-	2	-
1984	31	-	-	1	4	4	3	11	1	6	-
1983	32	-	3	1	1	6	5	2	4	4	-
1982	28	2	-	3	3	2	4	4	3	3	-
1981	28	-	1	-	1	3	6	6	5	2	-
1980	41	-	2	1	2	6	6	14	1	7	-
1979	45	1	-	2	9	6	11	4	6	4	-
Syphilis (090-097)											
Black, female											
1998	10	1	-	1	-	1	-	1	-	3	-
1997	6	-	-	-	-	-	2	2	2	-	-
1996	9	-	-	-	-	-	2	1	-	3	-
1995	8	-	1	1	-	-	-	1	1	2	-
1994	15	-	1	-	1	2	1	2	3	2	-
1993	17	-	-	-	-	1	1	4	-	3	-
1992	10	1	-	2	-	-	1	2	-	2	-
1991	24	-	-	-	-	1	3	1	2	3	-
1990	23	-	-	1	-	3	2	2	-	3	-
1989	20	-	1	-	-	2	3	2	2	1	-
1988	19	-	-	-	3	-	1	4	-	2	-
1987	22	-	-	1	3	1	5	2	3	3	-
1986	17	-	-	1	-	1	2	3	3	5	-
1985	10	1	1	-	1	-	-	-	2	3	-
1984	15	-	-	-	1	-	7	2	1	2	-
1983	18	-	1	-	1	2	1	8	2	1	-
1982	12	-	-	2	1	1	1	2	2	1	-
1981	19	1	1	1	1	-	3	4	3	3	-
1980	18	-	-	1	1	5	2	4	4	-	-
1979	31	-	2	2	5	2	4	6	3	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
All races, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	1	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	1	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	3	-	-	-	-	2	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	6	1	-	-	1	2	-	1	-	-	-
1979	4	-	1	-	2	-	-	-	-	-	-
Acute poliomyelitis (045)											
All races, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	1	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	2	-	-	-	-	1	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	4	1	-	-	1	1	-	1	-	-	-
1979	4	-	1	-	2	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
All races, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	1	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	1	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (045)											
White, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	1	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	1	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	3	-	-	-	-	2	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	4	1	-	-	1	1	-	1	-	-	-
1979	4	-	1	-	2	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
White, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	1	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	2	-	-	-	-	1	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	3	1	-	-	1	-	-	1	-	-	-
1979	4	-	1	-	2	-	-	-	-	-	-
Acute poliomyelitis (045)											
White, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	1	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	1	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
All other, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (045)											
All other, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (045)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute poliomyelitis (045)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	1	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (045)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
All races, both sexes											
1998	5	-	-	-	-	2	-	-	-	1	-
1997	4	-	1	-	-	-	-	-	-	-	-
1996	3	-	-	2	-	-	-	-	-	-	-
1995	7	-	1	-	-	1	1	1	1	-	-
1994	3	1	-	-	1	1	-	-	-	-	-
1993	3	-	-	-	1	-	1	1	-	-	-
1992	3	-	-	-	-	1	-	-	-	-	-
1991	10	-	-	2	1	1	2	1	-	1	-
1990	15	-	-	2	2	1	1	1	5	3	-
1989	1	-	-	-	-	-	-	1	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	5	-	-	-	1	1	1	-	-	-	-
1986	4	-	-	-	1	-	-	-	-	-	-
1985	3	-	-	2	1	-	-	-	-	-	-
1984	3	-	-	-	1	-	-	-	-	-	-
1983	6	-	-	1	-	2	1	1	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	7	-	1	-	1	-	1	1	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
All races, male											
1998	2	-	-	-	-	1	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	2	-	-	2	-	-	-	-	-	-	-
1995	6	-	1	-	-	1	-	1	1	-	-
1994	2	1	-	-	-	1	-	-	-	-	-
1993	3	-	-	-	1	-	1	1	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	5	-	-	1	1	1	1	-	-	-	-
1990	8	-	-	-	1	1	-	1	3	2	-
1989	1	-	-	-	-	-	-	1	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	1	1	1	-	-	-	-
1986	2	-	-	-	1	-	-	-	-	-	-
1985	2	-	-	2	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	5	-	-	1	-	1	1	1	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	6	-	1	-	1	-	1	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
All races, female											
1998	3	-	-	-	-	1	-	-	-	1	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	1	-	-	-	-
1994	1	-	-	-	1	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	1	-	-	-	-	-
1991	5	-	-	1	-	-	1	1	-	1	-
1990	7	-	-	2	1	-	1	-	2	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	1	-	-	-	-	-	-
1984	2	-	-	-	1	-	-	-	-	-	-
1983	1	-	-	-	-	1	-	-	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	1	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
White, both sexes											
1998	5	-	-	-	-	2	-	-	-	1	-
1997	4	-	1	-	-	-	-	-	-	-	-
1996	2	-	-	1	-	-	-	-	-	-	-
1995	3	-	-	-	-	-	-	1	-	-	-
1994	1	-	-	-	-	1	-	-	-	-	-
1993	2	-	-	-	1	-	1	-	-	-	-
1992	2	-	-	-	-	1	-	-	-	-	-
1991	6	-	-	1	1	-	1	-	-	1	-
1990	9	-	-	2	1	-	-	1	3	2	-
1989	1	-	-	-	-	-	-	1	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	1	1	1	-	-	-	-
1986	2	-	-	-	1	-	-	-	-	-	-
1985	3	-	-	2	1	-	-	-	-	-	-
1984	3	-	-	-	1	-	-	-	-	-	-
1983	6	-	-	1	-	2	1	1	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	7	-	1	-	1	-	1	1	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
White, male											
1998	2	-	-	-	-	1	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	1	-	-	1	-	-	-	-	-	-	-
1995	3	-	-	-	-	-	-	1	-	-	-
1994	1	-	-	-	-	1	-	-	-	-	-
1993	2	-	-	-	1	-	1	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	4	-	-	1	1	-	1	-	-	-	-
1990	5	-	-	-	1	-	-	1	2	1	-
1989	1	-	-	-	-	-	-	1	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	1	1	1	-	-	-	-
1986	2	-	-	-	1	-	-	-	-	-	-
1985	2	-	-	2	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	5	-	-	1	-	1	1	1	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	6	-	1	-	1	-	1	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
White, female											
1998	3	-	-	-	-	1	-	-	-	1	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	1	-	-	-	-	-
1991	2	-	-	-	-	-	-	-	-	1	-
1990	4	-	-	2	-	-	-	-	1	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	1	-	-	-	-	-	-
1984	2	-	-	-	1	-	-	-	-	-	-
1983	1	-	-	-	-	1	-	-	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	1	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
All other, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	1	-	-	-	-	-	-	-
1995	4	-	1	-	-	1	1	-	1	-	-
1994	2	1	-	-	1	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	1	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	4	-	-	1	-	1	1	1	-	-	-
1990	6	-	-	-	1	1	1	-	2	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
All other, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	1	-	-	-	-	-	-	-
1995	3	-	1	-	-	1	-	-	1	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	1	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	1	-	-	-	-	-
1990	3	-	-	-	-	1	-	-	1	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	1	-	-	-	-
1994	1	-	-	-	1	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	3	-	-	1	-	-	1	1	-	-	-
1990	3	-	-	-	1	-	1	-	1	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	1	-	-	-	-	-	-	-
1995	4	-	1	-	-	1	1	-	1	-	-
1994	2	1	-	-	1	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	1	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	4	-	-	1	-	1	1	1	-	-	-
1990	6	-	-	-	1	1	1	-	2	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Arthropod-borne viral encephalitis (062-064)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	1	-	-	-	-	-	-	-
1995	3	-	1	-	-	1	-	-	1	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	1	-	-	-	-	-	-	1	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	1	-	-	-	-	-
1990	3	-	-	-	-	1	-	-	1	1	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (062-064)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	1	-	-	-	-
1994	1	-	-	-	1	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	3	-	-	1	-	-	1	1	-	-	-
1990	3	-	-	-	1	-	1	-	1	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
All races, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	4	-	-	-	-	-	-	-	-	-	-
1991	27	-	1	-	-	-	-	-	-	-	-
1990	64	-	-	-	-	-	-	-	-	-	-
1989	32	-	-	-	-	-	-	-	1	-	-
1988	3	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	1	-	-	-	-	-	-	-
1986	2	-	-	-	1	-	-	-	-	-	-
1985	4	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	4	-	-	-	-	-	-	-	-	-	-
1982	2	-	1	-	-	-	-	-	-	-	-
1981	2	1	-	-	-	-	-	-	-	-	-
1980	11	-	-	-	-	-	-	-	-	-	-
1979	6	-	1	1	-	-	-	-	-	-	-
Measles (055)											
All races, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	10	-	-	-	-	-	-	-	-	-	-
1990	34	-	-	-	-	-	-	-	-	-	-
1989	16	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	1	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	3	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	6	-	-	-	-	-	-	-	-	-	-
1979	2	-	1	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
All races, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	17	-	1	-	-	-	-	-	-	-	-
1990	30	-	-	-	-	-	-	-	-	-	-
1989	16	-	-	-	-	-	-	-	1	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	1	-	-	-	-	-	-
1985	3	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	1	-	1	-	-	-	-	-	-	-	-
1981	2	1	-	-	-	-	-	-	-	-	-
1980	5	-	-	-	-	-	-	-	-	-	-
1979	4	-	-	1	-	-	-	-	-	-	-
Measles (055)											
White, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	4	-	-	-	-	-	-	-	-	-	-
1991	19	-	1	-	-	-	-	-	-	-	-
1990	43	-	-	-	-	-	-	-	-	-	-
1989	18	-	-	-	-	-	-	-	1	-	-
1988	3	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	1	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	4	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	3	-	-	-	-	-	-	-	-	-	-
1982	2	-	1	-	-	-	-	-	-	-	-
1981	2	1	-	-	-	-	-	-	-	-	-
1980	8	-	-	-	-	-	-	-	-	-	-
1979	5	-	1	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
White, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	9	-	-	-	-	-	-	-	-	-	-
1990	23	-	-	-	-	-	-	-	-	-	-
1989	10	-	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	1	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	3	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	4	-	-	-	-	-	-	-	-	-	-
1979	2	-	1	-	-	-	-	-	-	-	-
Measles (055)											
White, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	2	-	1	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	10	-	1	-	-	-	-	-	-	-	-
1990	20	-	-	-	-	-	-	-	-	-	-
1989	8	-	-	-	-	-	-	-	1	-	-
1988	2	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	3	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	1	-	1	-	-	-	-	-	-	-	-
1981	2	1	-	-	-	-	-	-	-	-	-
1980	4	-	-	-	-	-	-	-	-	-	-
1979	3	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
All other, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	8	-	-	-	-	-	-	-	-	-	-
1990	21	-	-	-	-	-	-	-	-	-	-
1989	14	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	1	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	1	-	-	-	-	-	-	-
Measles (055)											
All other, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	11	-	-	-	-	-	-	-	-	-	-
1989	6	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	7	-	-	-	-	-	-	-	-	-	-
1990	10	-	-	-	-	-	-	-	-	-	-
1989	8	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	1	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	1	-	-	-	-	-	-	-
Measles (055)											
Black, both sexes											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	7	-	-	-	-	-	-	-	-	-	-
1990	10	-	-	-	-	-	-	-	-	-	-
1989	11	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	1	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	1	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Measles (055)											
Black, male											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	5	-	-	-	-	-	-	-	-	-	-
1989	6	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Measles (055)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	6	-	-	-	-	-	-	-	-	-	-
1990	5	-	-	-	-	-	-	-	-	-	-
1989	5	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	1	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	1	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	1	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
All races, both sexes											
1998	4,796	794	539	374	366	433	421	412	248	153	-
1997	4,057	580	372	341	298	388	422	355	262	160	-
1996	3,780	460	322	294	298	401	406	379	216	139	1
1995	3,400	374	274	277	294	355	396	321	225	121	-
1994	3,061	280	205	223	251	342	344	322	204	105	-
1993	2,489	208	192	195	237	270	284	263	151	79	-
1992	2,001	151	136	137	184	229	224	216	140	80	-
1991	1,840	141	114	160	180	210	229	177	116	66	-
1990	1,578	96	93	101	147	206	191	178	115	65	-
1989	1,516	90	93	108	136	191	171	138	115	72	-
1988	1,290	65	91	102	151	147	156	125	87	57	-
1987	1,182	66	64	112	97	121	146	115	70	39	-
1986	1,006	50	60	82	92	117	115	96	55	40	-
1985	942	64	46	72	95	117	104	76	42	33	-
1984	869	42	55	75	90	92	82	53	40	42	-
1983	863	42	61	78	83	91	95	62	39	32	-
1982	814	44	63	82	78	91	74	75	30	36	1
1981	862	58	68	68	95	85	67	54	41	32	1
1980	809	54	54	71	74	67	82	74	40	35	-
1979	753	45	41	70	79	74	64	49	39	31	1
Viral hepatitis (070)											
All races, male											
1998	2,982	603	380	229	208	223	192	187	119	73	-
1997	2,442	435	256	216	162	189	189	165	117	62	-
1996	2,273	354	220	188	159	197	196	184	90	57	1
1995	2,076	286	193	176	170	181	194	141	108	50	-
1994	1,869	208	136	140	129	183	180	160	97	46	-
1993	1,554	159	140	128	128	145	149	132	73	30	-
1992	1,249	111	102	88	112	131	125	112	64	31	-
1991	1,132	110	75	100	96	109	131	74	64	25	-
1990	927	76	66	62	82	110	93	92	47	20	-
1989	906	64	63	71	89	104	99	71	51	24	-
1988	768	49	59	59	86	86	91	61	45	22	-
1987	725	46	46	71	64	72	87	54	32	18	-
1986	619	37	38	60	65	66	71	50	24	21	-
1985	531	43	29	37	59	69	56	39	17	13	-
1984	484	28	38	48	52	49	44	25	17	14	-
1983	494	24	37	40	58	57	50	30	17	13	-
1982	444	23	38	41	41	48	39	44	12	17	1
1981	466	34	41	42	50	40	32	30	19	9	-
1980	415	34	24	42	38	28	40	40	20	14	-
1979	399	27	23	37	45	46	25	21	24	11	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
All races, female											
1998	1,814	191	159	145	158	210	229	225	129	80	-
1997	1,615	145	116	125	136	199	233	190	145	98	-
1996	1,507	106	102	106	139	204	210	195	126	82	-
1995	1,324	88	81	101	124	174	202	180	117	71	-
1994	1,192	72	69	83	122	159	164	162	107	59	-
1993	935	49	52	67	109	125	135	131	78	49	-
1992	752	40	34	49	72	98	99	104	76	49	-
1991	708	31	39	60	84	101	98	103	52	41	-
1990	651	20	27	39	65	96	98	86	68	45	-
1989	610	26	30	37	47	87	72	67	64	48	-
1988	522	16	32	43	65	61	65	64	42	35	-
1987	457	20	18	41	33	49	59	61	38	21	-
1986	387	13	22	22	27	51	44	46	31	19	-
1985	411	21	17	35	36	48	48	37	25	20	-
1984	385	14	17	27	38	43	38	28	23	28	-
1983	369	18	24	38	25	34	45	32	22	19	-
1982	370	21	25	41	37	43	35	31	18	19	-
1981	396	24	27	26	45	45	35	24	22	23	1
1980	394	20	30	29	36	39	42	34	20	21	-
1979	354	18	18	33	34	28	39	28	15	20	-
Viral hepatitis (070)											
White, both sexes											
1998	3,829	613	421	270	283	351	360	357	217	139	-
1997	3,250	439	289	252	224	306	362	304	239	138	-
1996	3,065	364	243	226	240	317	343	343	188	128	-
1995	2,769	284	218	209	226	295	361	287	209	107	-
1994	2,441	217	159	155	186	274	295	288	183	99	-
1993	2,029	152	150	145	191	227	254	234	140	70	-
1992	1,614	112	107	97	148	192	193	198	128	73	-
1991	1,487	106	76	126	150	181	206	155	106	62	-
1990	1,265	69	66	76	124	173	165	162	103	58	-
1989	1,232	71	71	84	107	162	151	124	108	66	-
1988	1,056	49	68	83	130	128	136	111	81	51	-
1987	935	47	47	97	77	102	127	96	63	38	-
1986	830	37	49	69	76	101	96	87	51	36	-
1985	765	47	31	63	82	101	92	73	38	30	-
1984	699	29	45	61	77	86	71	46	37	40	-
1983	683	34	42	70	72	76	75	57	36	30	-
1982	646	33	52	68	66	74	64	70	26	32	1
1981	694	44	50	55	85	75	63	51	37	27	1
1980	667	37	39	64	65	62	78	69	37	32	-
1979	594	33	28	59	72	64	59	43	35	29	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
White, male											
1998	2,382	470	299	163	163	180	168	162	107	66	-
1997	1,988	335	199	169	128	152	165	149	113	52	-
1996	1,868	282	166	145	135	162	170	175	83	53	-
1995	1,714	226	155	132	137	159	180	127	104	46	-
1994	1,543	167	110	103	100	158	158	154	89	42	-
1993	1,270	122	114	98	100	127	135	115	70	27	-
1992	1,015	83	80	63	91	111	113	103	58	26	-
1991	930	87	56	81	84	93	122	70	60	23	-
1990	755	55	48	50	70	96	80	82	45	19	-
1989	744	50	46	55	68	94	91	65	47	23	-
1988	647	38	48	48	77	77	85	54	41	20	-
1987	580	34	34	63	53	63	74	47	28	17	-
1986	517	27	31	50	56	58	59	47	21	18	-
1985	447	35	23	33	50	62	50	39	15	12	-
1984	396	20	31	40	43	48	39	21	17	13	-
1983	396	20	25	38	51	46	40	29	17	12	-
1982	360	16	31	38	35	41	36	41	11	13	1
1981	388	30	31	35	47	36	31	30	17	7	-
1980	350	25	17	41	36	26	38	37	17	14	-
1979	310	18	19	31	42	41	22	18	20	10	1
Viral hepatitis (070)											
White, female											
1998	1,447	143	122	107	120	171	192	195	110	73	-
1997	1,262	104	90	83	96	154	197	155	126	86	-
1996	1,197	82	77	81	105	155	173	168	105	75	-
1995	1,055	58	63	77	89	136	181	160	105	61	-
1994	898	50	49	52	86	116	137	134	94	57	-
1993	759	30	36	47	91	100	119	119	70	43	-
1992	599	29	27	34	57	81	80	95	70	47	-
1991	557	19	20	45	66	88	84	85	46	39	-
1990	510	14	18	26	54	77	85	80	58	39	-
1989	488	21	25	29	39	68	60	59	61	43	-
1988	409	11	20	35	53	51	51	57	40	31	-
1987	355	13	13	34	24	39	53	49	35	21	-
1986	313	10	18	19	20	43	37	40	30	18	-
1985	318	12	8	30	32	39	42	34	23	18	-
1984	303	9	14	21	34	38	32	25	20	27	-
1983	287	14	17	32	21	30	35	28	19	18	-
1982	286	17	21	30	31	33	28	29	15	19	-
1981	306	14	19	20	38	39	32	21	20	20	1
1980	317	12	22	23	29	36	40	32	20	18	-
1979	284	15	9	28	30	23	37	25	15	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
All other, both sexes											
1998	967	181	118	104	83	82	61	55	31	14	-
1997	807	141	83	89	74	82	60	51	23	22	-
1996	715	96	79	68	58	84	63	36	28	11	1
1995	631	90	56	68	68	60	35	34	16	14	-
1994	620	63	46	68	65	68	49	34	21	6	-
1993	460	56	42	50	46	43	30	29	11	9	-
1992	387	39	29	40	36	37	31	18	12	7	-
1991	353	35	38	34	30	29	23	22	10	4	-
1990	313	27	27	25	23	33	26	16	12	7	-
1989	284	19	22	24	29	29	20	14	7	6	-
1988	234	16	23	19	21	19	20	14	6	6	-
1987	247	19	17	15	20	19	19	19	7	1	-
1986	176	13	11	13	16	16	19	9	4	4	-
1985	177	17	15	9	13	16	12	3	4	3	-
1984	170	13	10	14	13	6	11	7	3	2	-
1983	180	8	19	8	11	15	20	5	3	2	-
1982	168	11	11	14	12	17	10	5	4	4	-
1981	168	14	18	13	10	10	4	3	4	5	-
1980	142	17	15	7	9	5	4	5	3	3	-
1979	159	12	13	11	7	10	5	6	4	2	-
Viral hepatitis (070)											
All other, male											
1998	600	133	81	66	45	43	24	25	12	7	-
1997	454	100	57	47	34	37	24	16	4	10	-
1996	405	72	54	43	24	35	26	9	7	4	1
1995	362	60	38	44	33	22	14	14	4	4	-
1994	326	41	26	37	29	25	22	6	8	4	-
1993	284	37	26	30	28	18	14	17	3	3	-
1992	234	28	22	25	21	20	12	9	6	5	-
1991	202	23	19	19	12	16	9	4	4	2	-
1990	172	21	18	12	12	14	13	10	2	1	-
1989	162	14	17	16	21	10	8	6	4	1	-
1988	121	11	11	11	9	9	6	7	4	2	-
1987	145	12	12	8	11	9	13	7	4	1	-
1986	102	10	7	10	9	8	12	3	3	3	-
1985	84	8	6	4	9	7	6	-	2	1	-
1984	88	8	7	8	9	1	5	4	-	1	-
1983	98	4	12	2	7	11	10	1	-	1	-
1982	84	7	7	3	6	7	3	3	1	4	-
1981	78	4	10	7	3	4	1	-	2	2	-
1980	65	9	7	1	2	2	2	3	3	-	-
1979	89	9	4	6	3	5	3	3	4	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
All other, female											
1998	367	48	37	38	38	39	37	30	19	7	-
1997	353	41	26	42	40	45	36	35	19	12	-
1996	310	24	25	25	34	49	37	27	21	7	-
1995	269	30	18	24	35	38	21	20	12	10	-
1994	294	22	20	31	36	43	27	28	13	2	-
1993	176	19	16	20	18	25	16	12	8	6	-
1992	153	11	7	15	15	17	19	9	6	2	-
1991	151	12	19	15	18	13	14	18	6	2	-
1990	141	6	9	13	11	19	13	6	10	6	-
1989	122	5	5	8	8	19	12	8	3	5	-
1988	113	5	12	8	12	10	14	7	2	4	-
1987	102	7	5	7	9	10	6	12	3	-	-
1986	74	3	4	3	7	8	7	6	1	1	-
1985	93	9	9	5	4	9	6	3	2	2	-
1984	82	5	3	6	4	5	6	3	3	1	-
1983	82	4	7	6	4	4	10	4	3	1	-
1982	84	4	4	11	6	10	7	2	3	-	-
1981	90	10	8	6	7	6	3	3	2	3	-
1980	77	8	8	6	7	3	2	2	-	3	-
1979	70	3	9	5	4	5	2	3	-	1	-
Viral hepatitis (070)											
Black, both sexes											
1998	694	159	88	66	43	52	34	36	15	11	-
1997	535	105	63	54	40	37	31	32	16	18	-
1996	480	81	59	37	26	41	39	15	21	8	1
1995	429	64	37	39	32	39	21	17	12	10	-
1994	413	48	26	37	31	37	28	24	13	5	-
1993	294	40	22	25	21	22	19	18	8	6	-
1992	256	28	15	20	20	22	17	8	9	4	-
1991	221	26	19	16	14	13	14	12	6	3	-
1990	212	18	13	10	11	21	18	9	9	6	-
1989	206	13	10	14	15	22	15	9	6	6	-
1988	162	10	17	10	12	10	11	10	4	4	-
1987	165	12	5	8	8	15	11	15	5	-	-
1986	126	9	6	9	5	13	13	3	3	2	-
1985	118	9	9	5	7	13	5	1	4	2	-
1984	119	8	3	10	5	4	7	4	2	2	-
1983	135	7	14	7	8	10	12	4	2	2	-
1982	125	6	5	8	9	14	8	4	4	3	-
1981	139	8	12	13	8	9	4	3	3	3	-
1980	113	12	11	7	7	4	2	3	2	2	-
1979	127	9	9	6	4	8	4	6	3	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Viral hepatitis (070)											
Black, male											
1998	429	118	59	38	28	27	12	14	3	5	-
1997	311	76	44	29	22	15	14	10	1	7	-
1996	273	62	42	21	12	14	16	3	3	2	1
1995	243	41	25	23	15	12	8	8	3	1	-
1994	216	30	16	19	11	12	11	5	7	3	-
1993	185	27	16	16	14	9	9	10	3	2	-
1992	155	21	12	13	13	11	5	4	5	2	-
1991	124	16	8	9	4	8	3	4	2	1	-
1990	105	12	7	4	5	10	8	5	1	-	-
1989	106	9	7	8	10	8	5	2	3	1	-
1988	79	6	7	5	6	3	2	5	3	2	-
1987	92	6	5	4	4	7	5	6	3	-	-
1986	66	7	4	6	1	6	8	1	2	1	-
1985	54	5	2	2	4	5	3	-	2	1	-
1984	55	3	2	6	3	-	3	2	-	1	-
1983	72	3	8	2	6	8	5	1	-	1	-
1982	56	3	3	1	4	6	2	2	1	3	-
1981	58	2	4	7	2	3	1	-	1	1	-
1980	50	6	4	1	2	2	-	1	2	-	-
1979	60	6	1	1	-	3	2	3	3	1	-
Viral hepatitis (070)											
Black, female											
1998	265	41	29	28	15	25	22	22	12	6	-
1997	224	29	19	25	18	22	17	22	15	11	-
1996	207	19	17	16	14	27	23	12	18	6	-
1995	186	23	12	16	17	27	13	9	9	9	-
1994	197	18	10	18	20	25	17	19	6	2	-
1993	109	13	6	9	7	13	10	8	5	4	-
1992	101	7	3	7	7	11	12	4	4	2	-
1991	97	10	11	7	10	5	11	8	4	2	-
1990	107	6	6	6	6	11	10	4	8	6	-
1989	100	4	3	6	5	14	10	7	3	5	-
1988	83	4	10	5	6	7	9	5	1	2	-
1987	73	6	-	4	4	8	6	9	2	-	-
1986	60	2	2	3	4	7	5	2	1	1	-
1985	64	4	7	3	3	8	2	1	2	1	-
1984	64	5	1	4	2	4	4	2	2	1	-
1983	63	4	6	5	2	2	7	3	2	1	-
1982	69	3	2	7	5	8	6	2	3	-	-
1981	81	6	8	6	6	6	3	3	2	2	-
1980	63	6	7	6	5	2	2	2	-	2	-
1979	67	3	8	5	4	5	2	3	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
All races, both sexes											
1998	13,426	2,022	1,098	642	337	181	120	40	24	7	3
1997	16,516	2,340	1,173	648	417	219	117	57	15	7	6
1996	31,130	4,168	2,091	1,138	656	328	175	59	28	18	7
1995	43,115	5,421	2,758	1,477	843	468	213	60	19	10	8
1994	42,114	4,949	2,687	1,426	760	403	184	77	18	11	6
1993	37,267	4,216	2,271	1,161	682	385	160	63	19	17	10
1992	33,566	3,666	1,909	1,087	698	360	159	61	27	14	13
1991	29,555	3,022	1,706	936	622	310	137	71	21	10	10
1990	25,188	2,476	1,336	851	463	262	107	47	19	12	13
1989	22,082	2,092	1,193	723	421	224	103	48	22	12	13
1988	16,602	1,427	925	543	324	194	92	64	15	11	3
1987	13,468	1,211	627	468	293	149	81	50	23	13	3
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
All races, male											
1998	10,291	1,648	915	512	277	147	96	28	13	3	3
1997	12,892	1,892	970	538	344	194	86	40	6	4	5
1996	25,277	3,516	1,805	968	546	273	148	46	21	7	7
1995	35,950	4,674	2,442	1,279	724	381	156	46	11	5	8
1994	35,641	4,372	2,383	1,239	665	340	144	46	12	6	6
1993	32,093	3,789	2,047	1,029	607	321	125	45	11	11	10
1992	29,325	3,333	1,771	962	616	317	114	47	16	6	13
1991	26,046	2,768	1,550	836	557	261	103	43	15	7	10
1990	22,386	2,320	1,245	782	409	218	71	32	10	4	11
1989	19,758	1,969	1,114	665	389	185	75	28	16	8	13
1988	14,811	1,347	867	501	283	157	66	44	9	8	2
1987	12,088	1,153	586	436	262	122	57	27	14	6	2
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
All races, female											
1998	3,135	374	183	130	60	34	24	12	11	4	-
1997	3,624	448	203	110	73	25	31	17	9	3	1
1996	5,853	652	286	170	110	55	27	13	7	11	-
1995	7,165	747	316	198	119	87	57	14	8	5	-
1994	6,473	577	304	187	95	63	40	31	6	5	-
1993	5,174	427	224	132	75	64	35	18	8	6	-
1992	4,241	333	138	125	82	43	45	14	11	8	-
1991	3,509	254	156	100	65	49	34	28	6	3	-
1990	2,802	156	91	69	54	44	36	15	9	8	2
1989	2,324	123	79	58	32	39	28	20	6	4	-
1988	1,791	80	58	42	41	37	26	20	6	3	1
1987	1,380	58	41	32	31	27	24	23	9	7	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
White, both sexes											
1998	6,106	897	496	287	158	80	54	22	10	6	1
1997	7,840	1,081	554	310	179	101	43	23	7	4	4
1996	16,804	2,264	1,167	606	345	181	75	27	13	10	2
1995	25,509	3,242	1,709	899	471	271	122	31	11	6	5
1994	25,578	3,086	1,692	864	458	228	98	47	10	9	4
1993	23,586	2,786	1,519	771	458	240	100	41	13	16	7
1992	21,921	2,562	1,344	708	476	236	107	46	19	11	7
1991	19,850	2,129	1,181	672	441	210	107	54	14	8	6
1990	17,255	1,846	1,002	620	330	194	80	38	14	8	8
1989	15,095	1,563	883	542	318	167	82	39	21	10	5
1988	11,267	1,063	685	393	253	153	79	53	15	10	2
1987	9,328	899	482	364	224	126	74	47	22	10	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
White, male											
1998	5,184	792	440	246	137	68	44	14	4	3	1
1997	6,688	957	484	270	154	96	36	16	2	3	4
1996	14,662	2,030	1,064	533	299	155	66	21	11	2	2
1995	22,670	2,951	1,563	823	426	240	94	21	7	2	5
1994	22,993	2,862	1,567	795	413	202	79	27	8	5	4
1993	21,455	2,609	1,412	702	421	206	78	29	7	10	7
1992	20,161	2,405	1,272	650	437	213	73	39	11	5	7
1991	18,366	2,018	1,116	624	413	177	78	31	9	6	6
1990	16,106	1,781	961	580	300	167	53	27	8	1	8
1989	14,114	1,510	844	510	297	136	61	24	16	6	5
1988	10,479	1,025	657	370	225	124	55	37	9	7	2
1987	8,700	875	453	340	205	106	54	26	13	4	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
White, female											
1998	922	105	56	41	21	12	10	8	6	3	-
1997	1,152	124	70	40	25	5	7	7	5	1	-
1996	2,142	234	103	73	46	26	9	6	2	8	-
1995	2,839	291	146	76	45	31	28	10	4	4	-
1994	2,585	224	125	69	45	26	19	20	2	4	-
1993	2,131	177	107	69	37	34	22	12	6	6	-
1992	1,760	157	72	58	39	23	34	7	8	6	-
1991	1,484	111	65	48	28	33	29	23	5	2	-
1990	1,149	65	41	40	30	27	27	11	6	7	-
1989	981	53	39	32	21	31	21	15	5	4	-
1988	788	38	28	23	28	29	24	16	6	3	-
1987	628	24	29	24	19	20	20	21	9	6	-
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
All other, both sexes											
1998	7,320	1,125	602	355	179	101	66	18	14	1	2
1997	8,676	1,259	619	338	238	118	74	34	8	3	2
1996	14,326	1,904	924	532	311	147	100	32	15	8	5
1995	17,606	2,179	1,049	578	372	197	91	29	8	4	3
1994	16,536	1,863	995	562	302	175	86	30	8	2	2
1993	13,681	1,430	752	390	224	145	60	22	6	1	3
1992	11,645	1,104	565	379	222	124	52	15	8	3	6
1991	9,705	893	525	264	181	100	30	17	7	2	4
1990	7,933	630	334	231	133	68	27	9	5	4	5
1989	6,987	529	310	181	103	57	21	9	1	2	8
1988	5,335	364	240	150	71	41	13	11	-	1	1
1987	4,140	312	145	104	69	23	7	3	1	3	2
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
All other, male											
1998	5,107	856	475	266	140	79	52	14	9	-	2
1997	6,204	935	486	268	190	98	50	24	4	1	1
1996	10,615	1,486	741	435	247	118	82	25	10	5	5
1995	13,280	1,723	879	456	298	141	62	25	4	3	3
1994	12,648	1,510	816	444	252	138	65	19	4	1	2
1993	10,638	1,180	635	327	186	115	47	16	4	1	3
1992	9,164	928	499	312	179	104	41	8	5	1	6
1991	7,680	750	434	212	144	84	25	12	6	1	4
1990	6,280	539	284	202	109	51	18	5	2	3	3
1989	5,644	459	270	155	92	49	14	4	-	2	8
1988	4,332	322	210	131	58	33	11	7	-	1	-
1987	3,388	278	133	96	57	16	3	1	1	2	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
All other, female											
1998	2,213	269	127	89	39	22	14	4	5	1	-
1997	2,472	324	133	70	48	20	24	10	4	2	1
1996	3,711	418	183	97	64	29	18	7	5	3	-
1995	4,326	456	170	122	74	56	29	4	4	1	-
1994	3,888	353	179	118	50	37	21	11	4	1	-
1993	3,043	250	117	63	38	30	13	6	2	-	-
1992	2,481	176	66	67	43	20	11	7	3	2	-
1991	2,025	143	91	52	37	16	5	5	1	1	-
1990	1,653	91	50	29	24	17	9	4	3	1	2
1989	1,343	70	40	26	11	8	7	5	1	-	-
1988	1,003	42	30	19	13	8	2	4	-	-	1
1987	752	34	12	8	12	7	4	2	-	1	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
Black, both sexes											
1998	7,180	1,110	590	343	177	99	66	17	14	1	2
1997	8,525	1,247	606	332	234	117	72	34	8	3	2
1996	13,997	1,855	906	522	302	145	96	32	15	8	5
1995	17,139	2,124	1,029	559	363	194	89	27	8	4	3
1994	16,079	1,816	963	547	294	170	84	29	7	2	2
1993	13,319	1,382	728	379	222	142	59	22	6	1	3
1992	11,378	1,077	547	371	218	121	50	15	8	3	6
1991	9,437	866	512	256	175	97	28	15	6	2	4
1990	7,730	607	323	224	129	65	25	9	5	4	5
1989	6,795	507	302	173	99	56	20	9	1	1	7
1988	5,197	354	233	141	67	39	13	10	-	1	1
1987	4,040	300	140	103	65	18	6	2	1	3	2
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Human immunodeficiency virus (HIV) disease (042-044)											
Black, male											
1998	4,994	843	464	256	140	77	52	14	9	-	2
1997	6,078	928	475	263	188	97	50	24	4	1	1
1996	10,329	1,441	725	426	239	116	78	25	10	5	5
1995	12,875	1,675	861	440	291	139	61	24	4	3	3
1994	12,245	1,466	789	429	246	134	63	18	3	1	2
1993	10,324	1,138	614	317	184	114	47	16	4	1	3
1992	8,925	901	484	306	175	103	39	8	5	1	6
1991	7,440	725	422	205	140	81	23	11	6	1	4
1990	6,097	516	273	196	107	49	16	5	2	3	3
1989	5,475	438	263	150	89	49	13	4	-	1	7
1988	4,202	313	204	123	55	31	11	7	-	1	-
1987	3,301	267	128	95	55	12	3	-	1	2	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---
Human immunodeficiency virus (HIV) disease (042-044)											
Black, female											
1998	2,186	267	126	87	37	22	14	3	5	1	-
1997	2,447	319	131	69	46	20	22	10	4	2	1
1996	3,668	414	181	96	63	29	18	7	5	3	-
1995	4,264	449	168	119	72	55	28	3	4	1	-
1994	3,834	350	174	118	48	36	21	11	4	1	-
1993	2,995	244	114	62	38	28	12	6	2	-	-
1992	2,453	176	63	65	43	18	11	7	3	2	-
1991	1,997	141	90	51	35	16	5	4	-	1	-
1990	1,633	91	50	28	22	16	9	4	3	1	2
1989	1,320	69	39	23	10	7	7	5	1	-	-
1988	995	41	29	18	12	8	2	3	-	-	1
1987	739	33	12	8	10	6	3	2	-	1	1
1986	---	---	---	---	---	---	---	---	---	---	---
1985	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---
1981	---	---	---	---	---	---	---	---	---	---	---
1980	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
All races, both sexes											
1998	6	-	2	-	1	1	-	1	-	-	-
1997	7	1	-	2	-	-	-	-	1	-	-
1996	4	-	1	1	1	-	-	-	-	-	-
1995	8	-	-	1	-	3	1	-	-	-	-
1994	3	1	-	-	-	-	1	-	-	-	-
1993	12	-	3	-	1	1	-	-	-	-	-
1992	8	-	1	-	1	1	1	-	1	-	-
1991	4	-	2	-	-	1	-	-	-	-	-
1990	3	-	-	-	-	1	-	-	-	-	-
1989	11	1	-	1	1	1	-	2	-	-	-
1988	7	-	1	1	1	2	-	-	-	-	-
1987	5	-	-	-	1	-	1	-	-	-	-
1986	5	-	1	-	-	2	-	-	-	-	-
1985	13	1	-	1	1	2	1	3	1	-	-
1984	7	1	1	1	-	-	-	-	-	-	-
1983	3	-	-	-	-	-	1	-	-	-	-
1982	2	1	-	-	-	-	-	-	-	-	-
1981	7	1	-	-	2	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	3	-	-	1	-	-	-	1	-	-	-
Malaria (084)											
All races, male											
1998	4	-	1	-	1	-	-	1	-	-	-
1997	6	1	-	2	-	-	-	-	1	-	-
1996	3	-	-	1	1	-	-	-	-	-	-
1995	5	-	-	-	-	3	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	7	-	1	-	1	-	-	-	-	-	-
1992	5	-	1	-	1	1	-	-	1	-	-
1991	3	-	2	-	-	1	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	7	1	-	1	-	1	-	2	-	-	-
1988	3	-	1	1	-	1	-	-	-	-	-
1987	3	-	-	-	-	-	1	-	-	-	-
1986	3	-	1	-	-	2	-	-	-	-	-
1985	7	1	-	-	1	1	1	1	-	-	-
1984	6	1	1	1	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	1	-	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	6	1	-	-	2	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
All races, female											
1998	2	-	1	-	-	1	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	1	-	1	-	-	-	-	-	-	-	-
1995	3	-	-	1	-	-	1	-	-	-	-
1994	2	-	-	-	-	-	1	-	-	-	-
1993	5	-	2	-	-	1	-	-	-	-	-
1992	3	-	-	-	-	-	1	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	3	-	-	-	-	1	-	-	-	-	-
1989	4	-	-	-	1	-	-	-	-	-	-
1988	4	-	-	-	1	1	-	-	-	-	-
1987	2	-	-	-	1	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	6	-	-	1	-	1	-	2	1	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	1	-	-	-	1	-	-	-
Malaria (084)											
White, both sexes											
1998	3	-	1	-	-	-	-	1	-	-	-
1997	5	1	-	2	-	-	-	-	1	-	-
1996	2	-	-	1	1	-	-	-	-	-	-
1995	6	-	-	1	-	3	1	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	5	-	2	-	-	-	-	-	-	-	-
1992	3	-	-	-	1	1	-	-	-	-	-
1991	2	-	-	-	-	1	-	-	-	-	-
1990	1	-	-	-	-	1	-	-	-	-	-
1989	8	1	-	1	-	1	-	2	-	-	-
1988	6	-	1	1	-	2	-	-	-	-	-
1987	2	-	-	-	1	-	1	-	-	-	-
1986	5	-	1	-	-	2	-	-	-	-	-
1985	10	-	-	1	1	2	1	3	1	-	-
1984	5	1	1	1	-	-	-	-	-	-	-
1983	3	-	-	-	-	-	1	-	-	-	-
1982	2	1	-	-	-	-	-	-	-	-	-
1981	2	1	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	3	-	-	1	-	-	-	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
White, male											
1998	3	-	1	-	-	-	-	1	-	-	-
1997	4	1	-	2	-	-	-	-	1	-	-
1996	2	-	-	1	1	-	-	-	-	-	-
1995	4	-	-	-	-	3	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	3	-	-	-	-	-	-	-	-	-	-
1992	2	-	-	-	1	1	-	-	-	-	-
1991	1	-	-	-	-	1	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	6	1	-	1	-	1	-	2	-	-	-
1988	3	-	1	1	-	1	-	-	-	-	-
1987	1	-	-	-	-	-	1	-	-	-	-
1986	3	-	1	-	-	2	-	-	-	-	-
1985	4	-	-	-	1	1	1	1	-	-	-
1984	5	1	1	1	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	1	-	-	-	-
1982	1	1	-	-	-	-	-	-	-	-	-
1981	2	1	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	1	-	-	-	-	-	-	-	-	-	-
Malaria (084)											
White, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	2	-	-	1	-	-	1	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	2	-	2	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	1	-	-	-	-	1	-	-	-	-	-
1989	2	-	-	-	-	-	-	-	-	-	-
1988	3	-	-	-	-	1	-	-	-	-	-
1987	1	-	-	-	1	-	-	-	-	-	-
1986	2	-	-	-	-	-	-	-	-	-	-
1985	6	-	-	1	-	1	-	2	1	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	1	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	2	-	-	1	-	-	-	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
All other, both sexes											
1998	3	-	1	-	1	1	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	2	-	1	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	3	1	-	-	-	-	1	-	-	-	-
1993	7	-	1	-	1	1	-	-	-	-	-
1992	5	-	1	-	-	-	1	-	1	-	-
1991	2	-	2	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	3	-	-	-	1	-	-	-	-	-	-
1988	1	-	-	-	1	-	-	-	-	-	-
1987	3	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	3	1	-	-	-	-	-	-	-	-	-
1984	2	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	5	-	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Malaria (084)											
All other, male											
1998	1	-	-	-	1	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	4	-	1	-	1	-	-	-	-	-	-
1992	3	-	1	-	-	-	-	-	1	-	-
1991	2	-	2	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	3	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	4	-	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
All other, female											
1998	2	-	1	-	-	1	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	1	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	2	-	-	-	-	-	1	-	-	-	-
1993	3	-	-	-	-	1	-	-	-	-	-
1992	2	-	-	-	-	-	1	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	2	-	-	-	1	-	-	-	-	-	-
1988	1	-	-	-	1	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Malaria (084)											
Black, both sexes											
1998	2	-	1	-	1	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	2	-	1	-	-	-	-	-	-	-	-
1995	2	-	-	-	-	-	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	5	-	-	-	-	1	-	-	-	-	-
1992	2	-	-	-	-	-	-	-	-	-	-
1991	1	-	1	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	2	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	2	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	3	-	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malaria (084)											
Black, male											
1998	1	-	-	-	1	-	-	-	-	-	-
1997	2	-	-	-	-	-	-	-	-	-	-
1996	1	-	-	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	1	-	1	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	3	-	-	-	1	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Malaria (084)											
Black, female											
1998	1	-	1	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	1	-	1	-	-	-	-	-	-	-	-
1995	1	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	3	-	-	-	-	1	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	-	-	-	-	-	-	-	-
1989	1	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All races, both sexes											
1998	6,052	293	332	378	428	553	689	705	611	792	-
1997	6,102	304	348	339	456	541	681	731	618	693	-
1996	6,334	369	294	356	406	616	711	696	571	681	-
1995	6,388	319	341	331	404	546	666	698	529	680	-
1994	6,496	368	307	347	478	575	675	632	539	670	-
1993	6,468	309	294	329	455	573	670	656	538	728	1
1992	6,184	318	298	330	446	545	618	649	502	653	-
1991	6,234	295	274	331	484	592	653	593	505	631	1
1990	6,442	307	248	337	492	636	643	651	548	629	1
1989	6,576	260	280	348	517	663	668	713	530	654	-
1988	6,122	228	279	330	494	588	678	633	534	670	1
1987	5,569	202	228	359	465	569	620	545	453	565	-
1986	8,785	497	399	503	546	637	624	583	452	590	1
1985	6,931	342	338	391	490	568	586	535	389	553	2
1984	5,627	283	247	364	431	504	510	492	392	482	1
1983	4,528	169	236	315	405	451	479	466	340	398	-
1982	4,071	134	210	267	373	456	469	402	289	399	-
1981	3,820	128	229	283	380	412	388	369	315	397	-
1980	3,693	120	205	303	310	406	385	343	298	411	1
1979	3,234	136	162	268	303	331	310	294	256	263	1
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All races, male											
1998	3,134	181	199	203	234	299	374	360	285	242	-
1997	3,179	190	194	184	252	295	346	382	286	240	-
1996	3,409	237	166	191	236	325	366	339	261	256	-
1995	3,555	217	188	181	212	302	353	368	252	257	-
1994	3,685	256	190	219	266	311	354	310	216	243	-
1993	3,641	202	186	194	269	317	358	309	267	252	1
1992	3,498	207	190	194	247	310	329	321	231	240	-
1991	3,642	208	168	203	283	336	358	327	266	222	1
1990	3,678	213	151	193	279	363	354	325	265	222	-
1989	3,744	180	174	209	300	384	351	377	250	220	-
1988	3,420	142	182	205	269	342	376	312	239	216	-
1987	3,105	146	146	229	270	308	324	286	203	192	-
1986	6,090	410	296	351	342	368	338	311	205	190	1
1985	4,508	266	239	255	284	343	309	273	185	185	-
1984	3,444	211	170	239	278	272	278	242	174	167	-
1983	2,552	112	125	185	223	244	260	252	166	153	-
1982	2,188	82	131	150	225	248	245	206	139	150	-
1981	2,019	75	138	161	219	245	197	182	148	156	-
1980	1,937	69	121	177	171	247	216	155	138	149	1
1979	1,736	77	97	152	169	190	181	156	130	98	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All races, female											
1998	2,918	112	133	175	194	254	315	345	326	550	-
1997	2,923	114	154	155	204	246	335	349	332	453	-
1996	2,925	132	128	165	170	291	345	357	310	425	-
1995	2,833	102	153	150	192	244	313	330	277	423	-
1994	2,811	112	117	128	212	264	321	322	323	427	-
1993	2,827	107	108	135	186	256	312	347	271	476	-
1992	2,686	111	108	136	199	235	289	328	271	413	-
1991	2,592	87	106	128	201	256	295	266	239	409	-
1990	2,764	94	97	144	213	273	289	326	283	407	1
1989	2,832	80	106	139	217	279	317	336	280	434	-
1988	2,702	86	97	125	225	246	302	321	295	454	1
1987	2,464	56	82	130	195	261	296	259	250	373	-
1986	2,695	87	103	152	204	269	286	272	247	400	-
1985	2,423	76	99	136	206	225	277	262	204	368	2
1984	2,183	72	77	125	153	232	232	250	218	315	1
1983	1,976	57	111	130	182	207	219	214	174	245	-
1982	1,883	52	79	117	148	208	224	196	150	249	-
1981	1,801	53	91	122	161	167	191	187	167	241	-
1980	1,756	51	84	126	139	159	169	188	160	262	-
1979	1,498	59	65	116	134	141	129	138	126	165	-
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
White, both sexes											
1998	4,944	205	261	293	350	484	599	625	555	721	-
1997	4,910	211	278	276	376	459	585	645	550	623	-
1996	5,063	265	223	273	330	524	626	600	521	602	-
1995	5,178	247	257	269	331	475	594	630	466	628	-
1994	5,236	281	240	285	385	507	576	565	473	614	-
1993	5,184	224	222	261	378	498	582	577	484	653	-
1992	4,953	223	229	259	358	471	542	587	444	587	-
1991	5,000	227	214	266	399	501	570	537	453	571	-
1990	5,168	225	177	268	403	554	561	569	497	569	-
1989	5,277	192	217	289	449	569	584	622	475	600	-
1988	5,023	169	218	269	418	524	587	560	479	617	1
1987	4,546	148	155	279	400	484	553	486	406	512	-
1986	7,079	393	322	413	458	557	554	516	412	543	-
1985	5,643	262	273	317	423	477	517	480	355	513	2
1984	4,600	219	188	304	368	440	449	442	348	451	1
1983	3,768	132	184	257	359	401	431	411	308	368	-
1982	3,420	104	171	228	323	395	419	349	260	372	-
1981	3,253	104	187	244	323	358	348	331	285	364	-
1980	3,102	84	166	258	261	353	341	298	268	378	1
1979	2,742	116	131	231	257	294	270	266	233	249	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
White, male											
1998	2,546	128	151	152	198	265	333	320	262	219	-
1997	2,534	130	153	148	201	252	298	336	263	215	-
1996	2,679	169	120	143	195	273	331	295	238	220	-
1995	2,874	173	149	151	180	261	324	324	225	231	-
1994	2,945	194	155	183	224	269	307	279	177	216	-
1993	2,875	146	141	148	223	279	307	269	240	228	-
1992	2,772	144	147	149	202	269	286	298	204	213	-
1991	2,875	160	130	159	224	287	311	294	234	196	-
1990	2,920	156	100	156	233	316	301	286	244	197	-
1989	2,960	134	134	177	261	327	312	326	216	200	-
1988	2,757	105	140	165	227	302	326	280	219	188	-
1987	2,513	104	101	176	235	267	287	260	179	172	-
1986	4,866	331	237	289	281	328	294	275	182	168	-
1985	3,629	204	191	213	248	290	277	243	164	168	-
1984	2,770	160	131	205	240	238	244	218	146	152	-
1983	2,085	87	103	144	199	215	232	221	150	135	-
1982	1,795	65	105	128	195	212	216	177	122	134	-
1981	1,710	60	114	142	188	211	177	166	135	138	-
1980	1,618	48	97	149	143	210	192	135	125	134	1
1979	1,472	66	79	131	145	174	160	137	116	93	1
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
White, female											
1998	2,398	77	110	141	152	219	266	305	293	502	-
1997	2,376	81	125	128	175	207	287	309	287	408	-
1996	2,384	96	103	130	135	251	295	305	283	382	-
1995	2,304	74	108	118	151	214	270	306	241	397	-
1994	2,291	87	85	102	161	238	269	286	296	398	-
1993	2,309	78	81	113	155	219	275	308	244	425	-
1992	2,181	79	82	110	156	202	256	289	240	374	-
1991	2,125	67	84	107	175	214	259	243	219	375	-
1990	2,248	69	77	112	170	238	260	283	253	372	-
1989	2,317	58	83	112	188	242	272	296	259	400	-
1988	2,266	64	78	104	191	222	261	280	260	429	1
1987	2,033	44	54	103	165	217	266	226	227	340	-
1986	2,213	62	85	124	177	229	260	241	230	375	-
1985	2,014	58	82	104	175	187	240	237	191	345	2
1984	1,830	59	57	99	128	202	205	224	202	299	1
1983	1,683	45	81	113	160	186	199	190	158	233	-
1982	1,625	39	66	100	128	183	203	172	138	238	-
1981	1,543	44	73	102	135	147	171	165	150	226	-
1980	1,484	36	69	109	118	143	149	163	143	244	-
1979	1,270	50	52	100	112	120	110	129	117	156	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All other, both sexes											
1998	1,108	88	71	85	78	69	90	80	56	71	-
1997	1,192	93	70	63	80	82	96	86	68	70	-
1996	1,271	104	71	83	76	92	85	96	50	79	-
1995	1,210	72	84	62	73	71	72	68	63	52	-
1994	1,260	87	67	62	93	68	99	67	66	56	-
1993	1,284	85	72	68	77	75	88	79	54	75	1
1992	1,231	95	69	71	88	74	76	62	58	66	-
1991	1,234	68	60	65	85	91	83	56	52	60	1
1990	1,274	82	71	69	89	82	82	82	51	60	1
1989	1,299	68	63	59	68	94	84	91	55	54	-
1988	1,099	59	61	61	76	64	91	73	55	53	-
1987	1,023	54	73	80	65	85	67	59	47	53	-
1986	1,706	104	77	90	88	80	70	67	40	47	1
1985	1,288	80	65	74	67	91	69	55	34	40	-
1984	1,027	64	59	60	63	64	61	50	44	31	-
1983	760	37	52	58	46	50	48	55	32	30	-
1982	651	30	39	39	50	61	50	53	29	27	-
1981	567	24	42	39	57	54	40	38	30	33	-
1980	591	36	39	45	49	53	44	45	30	33	-
1979	492	20	31	37	46	37	40	28	23	14	-
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All other, male											
1998	588	53	48	51	36	34	41	40	23	23	-
1997	645	60	41	36	51	43	48	46	23	25	-
1996	730	68	46	48	41	52	35	44	23	36	-
1995	681	44	39	30	32	41	29	44	27	26	-
1994	740	62	35	36	42	42	47	31	39	27	-
1993	766	56	45	46	46	38	51	40	27	24	1
1992	726	63	43	45	45	41	43	23	27	27	-
1991	767	48	38	44	59	49	47	33	32	26	1
1990	758	57	51	37	46	47	53	39	21	25	-
1989	784	46	40	32	39	57	39	51	34	20	-
1988	663	37	42	40	42	40	50	32	20	28	-
1987	592	42	45	53	35	41	37	26	24	20	-
1986	1,224	79	59	62	61	40	44	36	23	22	1
1985	879	62	48	42	36	53	32	30	21	17	-
1984	674	51	39	34	38	34	34	24	28	15	-
1983	467	25	22	41	24	29	28	31	16	18	-
1982	393	17	26	22	30	36	29	29	17	16	-
1981	309	15	24	19	31	34	20	16	13	18	-
1980	319	21	24	28	28	37	24	20	13	15	-
1979	264	11	18	21	24	16	21	19	14	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
All other, female											
1998	520	35	23	34	42	35	49	40	33	48	-
1997	547	33	29	27	29	39	48	40	45	45	-
1996	541	36	25	35	35	40	50	52	27	43	-
1995	529	28	45	32	41	30	43	24	36	26	-
1994	520	25	32	26	51	26	52	36	27	29	-
1993	518	29	27	22	31	37	37	39	27	51	-
1992	505	32	26	26	43	33	33	39	31	39	-
1991	467	20	22	21	26	42	36	23	20	34	-
1990	516	25	20	32	43	35	29	43	30	35	1
1989	515	22	23	27	29	37	45	40	21	34	-
1988	436	22	19	21	34	24	41	41	35	25	-
1987	431	12	28	27	30	44	30	33	23	33	-
1986	482	25	18	28	27	40	26	31	17	25	-
1985	409	18	17	32	31	38	37	25	13	23	-
1984	353	13	20	26	25	30	27	26	16	16	-
1983	293	12	30	17	22	21	20	24	16	12	-
1982	258	13	13	17	20	25	21	24	12	11	-
1981	258	9	18	20	26	20	20	22	17	15	-
1980	272	15	15	17	21	16	20	25	17	18	-
1979	228	9	13	16	22	21	19	9	9	9	-
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
Black, both sexes											
1998	928	79	61	67	59	58	76	58	44	56	-
1997	1,002	78	60	50	60	66	79	68	55	61	-
1996	1,078	94	56	72	60	75	69	82	38	62	-
1995	1,039	63	70	49	60	58	59	59	51	41	-
1994	1,080	78	59	57	70	54	79	51	51	48	-
1993	1,105	74	66	51	67	57	69	64	47	67	1
1992	1,072	86	63	60	67	57	67	52	47	55	-
1991	1,096	63	54	57	77	67	70	48	45	48	1
1990	1,122	75	62	57	78	67	67	71	43	47	1
1989	1,154	62	52	50	60	80	74	79	42	49	-
1988	972	50	55	53	63	54	74	70	48	46	-
1987	901	46	65	66	57	73	58	52	41	47	-
1986	1,552	97	69	79	69	65	57	54	36	36	1
1985	1,156	71	59	63	59	71	55	46	27	35	-
1984	906	58	48	49	54	57	50	39	36	26	-
1983	676	35	47	47	40	44	43	47	28	22	-
1982	565	29	31	30	45	57	44	38	25	21	-
1981	477	22	36	36	48	44	33	31	23	24	-
1980	523	33	32	35	45	47	41	38	26	30	-
1979	420	17	27	30	42	32	35	20	20	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
Black, male											
1998	505	47	42	45	29	32	38	27	18	17	-
1997	535	52	36	28	38	34	38	33	17	19	-
1996	619	61	37	42	33	41	28	37	16	25	-
1995	596	41	33	25	28	33	24	39	21	21	-
1994	642	54	33	36	30	33	38	24	28	22	-
1993	653	50	41	37	40	28	37	30	23	20	1
1992	635	57	40	40	37	34	35	22	18	20	-
1991	687	46	37	38	55	36	40	26	28	18	1
1990	663	51	44	28	40	34	41	33	19	15	-
1989	698	42	33	28	34	48	36	46	24	17	-
1988	597	34	40	36	36	32	39	31	19	24	-
1987	523	36	41	45	32	36	31	24	20	16	-
1986	1,121	73	53	57	48	30	34	25	19	17	1
1985	798	56	44	39	31	41	25	24	16	15	-
1984	593	48	33	26	32	29	25	18	21	12	-
1983	414	24	20	32	22	26	25	25	12	12	-
1982	339	16	21	17	30	32	25	18	14	11	-
1981	256	14	22	17	25	28	14	14	9	12	-
1980	281	20	20	21	27	31	23	15	12	12	-
1979	226	10	15	17	23	13	17	15	11	3	-
Other and unspecified infectious and parasitic diseases and their sequelae (001,005,020-032,037,039-041,046-054,056-061,065-066,071-083,085-088,098-134,136-139,771.3)											
Black, female											
1998	423	32	19	22	30	26	38	31	26	39	-
1997	467	26	24	22	22	32	41	35	38	42	-
1996	459	33	19	30	27	34	41	45	22	37	-
1995	443	22	37	24	32	25	35	20	30	20	-
1994	438	24	26	21	40	21	41	27	23	26	-
1993	452	24	25	14	27	29	32	34	24	47	-
1992	437	29	23	20	30	23	32	30	29	35	-
1991	409	17	17	19	22	31	30	22	17	30	-
1990	459	24	18	29	38	33	26	38	24	32	1
1989	456	20	19	22	26	32	38	33	18	32	-
1988	375	16	15	17	27	22	35	39	29	22	-
1987	378	10	24	21	25	37	27	28	21	31	-
1986	431	24	16	22	21	35	23	29	17	19	-
1985	358	15	15	24	28	30	30	22	11	20	-
1984	313	10	15	23	22	28	25	21	15	14	-
1983	262	11	27	15	18	18	18	22	16	10	-
1982	226	13	10	13	15	25	19	20	11	10	-
1981	221	8	14	19	23	16	19	17	14	12	-
1980	242	13	12	14	18	16	18	23	14	18	-
1979	194	7	12	13	19	19	18	5	9	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
All races, both sexes											
1998	541,532	18,167	27,580	37,365	49,659	67,917	86,836	87,486	71,031	70,916	13
1997	539,577	18,099	27,330	36,447	49,867	70,044	86,702	86,242	70,049	69,876	11
1996	539,533	18,405	26,222	35,858	50,970	72,347	88,517	84,798	69,682	67,644	25
1995	538,455	18,003	26,183	36,057	51,841	73,634	89,230	83,677	68,431	66,170	18
1994	534,310	17,533	26,055	35,894	53,357	75,492	88,303	82,187	67,201	63,003	16
1993	529,904	17,097	25,275	35,998	54,687	76,201	87,108	81,765	64,772	61,703	27
1992	520,578	16,752	24,454	35,807	55,802	76,434	84,791	79,813	62,835	58,187	30
1991	514,657	16,013	23,909	36,901	57,294	76,507	82,806	78,439	60,962	56,054	21
1990	505,322	15,798	24,025	36,960	57,959	76,599	80,814	76,775	58,236	52,963	27
1989	496,152	15,401	23,737	37,633	58,478	75,222	79,806	74,387	56,225	50,566	22
1988	485,048	15,031	23,735	38,577	59,079	73,260	77,564	71,201	53,848	48,314	26
1987	476,927	14,555	23,685	39,386	59,038	72,614	76,432	69,502	51,263	46,388	33
1986	469,376	14,241	23,559	39,755	59,050	71,134	75,669	67,004	49,628	44,749	46
1985	461,563	14,215	23,988	40,984	59,642	68,941	73,601	65,179	47,880	43,082	43
1984	453,492	13,859	24,499	41,870	58,185	68,105	71,636	63,064	46,468	42,089	42
1983	442,986	13,767	24,768	41,811	56,684	66,644	70,219	61,094	44,324	40,645	63
1982	433,795	13,831	25,584	41,357	55,880	66,057	67,126	59,241	42,900	38,958	64
1981	422,094	13,896	26,244	41,618	53,752	64,300	65,246	56,364	41,425	37,129	52
1980	416,509	14,230	26,800	41,600	53,045	63,469	63,961	54,773	40,466	35,720	28
1979	403,395	14,474	27,127	40,394	51,694	61,555	61,147	52,447	39,315	33,458	39
Malignant neoplasms (140-208)											
All races, male											
1998	282,065	8,920	14,151	20,157	27,508	38,290	47,957	47,126	35,991	30,430	8
1997	281,110	8,748	13,961	19,461	27,649	39,607	47,908	46,417	35,524	30,161	7
1996	281,898	8,889	13,389	19,342	28,345	40,749	49,230	45,489	35,390	29,238	15
1995	281,611	8,518	13,294	19,302	28,963	41,444	49,474	44,913	34,853	28,852	7
1994	280,465	8,288	13,428	19,359	30,090	42,369	49,025	44,129	34,204	27,493	11
1993	279,375	8,162	12,936	19,632	30,732	42,998	48,695	44,155	33,093	26,789	12
1992	274,838	8,105	12,498	19,467	31,431	43,234	47,039	43,427	32,031	25,473	17
1991	272,380	7,655	12,326	19,963	32,249	43,309	46,568	42,419	31,391	24,257	14
1990	268,283	7,511	12,369	20,290	32,763	43,267	45,467	41,855	29,835	23,049	15
1989	263,309	7,504	12,211	20,636	32,947	42,388	44,962	40,426	28,866	21,788	11
1988	258,088	7,219	12,326	21,047	33,154	41,587	43,629	39,183	27,517	20,855	16
1987	254,653	6,940	12,187	21,650	33,125	41,481	43,334	38,136	26,416	19,947	19
1986	250,559	6,827	12,110	21,799	33,009	40,486	42,633	36,936	25,557	19,332	22
1985	246,914	6,802	12,325	22,411	33,345	39,380	41,733	36,021	24,541	18,658	21
1984	242,790	6,647	12,671	22,967	32,621	38,779	40,942	34,798	23,691	18,325	27
1983	238,383	6,736	12,779	22,771	31,853	38,246	40,106	34,185	22,775	17,870	35
1982	233,864	6,694	13,311	22,416	31,546	38,044	38,568	32,854	21,974	17,422	37
1981	227,885	6,810	13,586	22,624	30,473	37,154	37,785	31,062	21,105	16,574	34
1980	225,948	6,904	13,869	22,739	30,128	36,980	36,881	30,601	20,730	16,149	16
1979	220,015	7,056	14,060	22,000	29,632	35,953	35,858	29,265	20,253	15,334	19

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
All races, female											
1998	259,467	9,247	13,429	17,208	22,151	29,627	38,879	40,360	35,040	40,486	5
1997	258,467	9,351	13,369	16,986	22,218	30,437	38,794	39,825	34,525	39,715	4
1996	257,635	9,516	12,833	16,516	22,625	31,598	39,287	39,309	34,292	38,406	10
1995	256,844	9,485	12,889	16,755	22,878	32,190	39,756	38,764	33,578	37,318	11
1994	253,845	9,245	12,627	16,535	23,267	33,123	39,278	38,058	32,997	35,510	5
1993	250,529	8,935	12,339	16,366	23,955	33,203	38,413	37,610	31,679	34,914	15
1992	245,740	8,647	11,956	16,340	24,371	33,200	37,752	36,386	30,804	32,714	13
1991	242,277	8,358	11,583	16,938	25,045	33,198	36,238	36,020	29,571	31,797	7
1990	237,039	8,287	11,656	16,670	25,196	33,332	35,347	34,920	28,401	29,914	12
1989	232,843	7,897	11,526	16,997	25,531	32,834	34,844	33,961	27,359	28,778	11
1988	226,960	7,812	11,409	17,530	25,925	31,673	33,935	32,018	26,331	27,459	10
1987	222,274	7,615	11,498	17,736	25,913	31,133	33,098	31,366	24,847	26,441	14
1986	218,817	7,414	11,449	17,956	26,041	30,648	33,036	30,068	24,071	25,417	24
1985	214,649	7,413	11,663	18,573	26,297	29,561	31,868	29,158	23,339	24,424	22
1984	210,702	7,212	11,828	18,903	25,564	29,326	30,694	28,266	22,777	23,764	15
1983	204,603	7,031	11,989	19,040	24,831	28,398	30,113	26,909	21,549	22,775	28
1982	199,931	7,137	12,273	18,941	24,334	28,013	28,558	26,387	20,926	21,536	27
1981	194,209	7,086	12,658	18,994	23,279	27,146	27,461	25,302	20,320	20,555	18
1980	190,561	7,326	12,931	18,861	22,917	26,489	27,080	24,172	19,736	19,571	12
1979	183,380	7,418	13,067	18,394	22,062	25,602	25,289	23,182	19,062	18,124	20
Malignant neoplasms (140-208)											
White, both sexes											
1998	470,139	14,219	22,386	31,026	41,744	58,742	76,372	78,155	64,289	64,016	9
1997	468,521	14,320	22,505	30,197	41,950	60,652	76,137	77,089	63,365	62,880	4
1996	469,406	14,671	21,589	29,789	43,176	62,672	78,019	75,770	62,891	61,126	16
1995	468,897	14,508	21,438	30,089	43,742	63,962	78,774	75,063	61,674	59,828	10
1994	465,797	14,093	21,420	29,793	45,462	65,671	78,015	73,817	60,534	57,045	7
1993	461,904	13,766	20,611	29,995	46,722	66,357	76,956	73,323	58,358	55,784	9
1992	454,531	13,648	19,934	29,631	47,779	66,652	75,124	71,772	56,834	52,779	22
1991	449,506	13,023	19,501	30,825	49,348	66,777	73,429	70,361	54,866	50,814	12
1990	441,585	12,793	19,654	30,786	49,745	66,951	71,867	68,841	52,681	48,092	15
1989	434,156	12,495	19,450	31,610	50,517	65,649	70,976	66,587	51,016	45,969	14
1988	425,140	12,288	19,459	32,563	51,151	64,075	69,277	63,790	48,868	43,940	18
1987	418,473	11,853	19,452	33,469	51,094	63,884	68,166	62,421	46,372	42,259	16
1986	412,352	11,533	19,407	33,998	51,086	62,690	67,444	60,279	45,071	40,810	33
1985	405,727	11,637	19,825	34,831	51,787	60,960	65,440	58,725	43,345	39,422	30
1984	398,631	11,233	20,394	35,586	50,361	60,267	63,787	56,843	42,078	38,672	24
1983	390,009	11,166	20,671	35,733	49,272	59,037	62,390	55,030	40,287	37,496	31
1982	382,652	11,301	21,424	35,362	48,667	58,573	59,839	53,586	39,023	35,970	43
1981	372,674	11,426	22,097	35,750	46,980	56,887	58,275	51,006	37,764	34,316	27
1980	368,162	11,713	22,544	35,780	46,442	56,150	57,237	49,567	37,068	32,987	17
1979	357,336	11,917	22,944	34,820	45,391	54,436	54,920	47,411	36,261	31,089	29

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
White, male											
1998	244,109	6,957	11,368	16,714	23,041	33,143	42,149	41,896	32,434	27,223	4
1997	243,192	6,885	11,382	15,937	23,179	34,341	41,996	41,321	32,079	26,888	2
1996	243,952	7,039	10,948	15,905	23,885	35,204	43,233	40,409	31,768	26,133	9
1995	244,000	6,858	10,766	16,027	24,353	35,890	43,477	40,061	31,281	25,772	2
1994	242,995	6,568	10,870	15,943	25,573	36,668	43,051	39,459	30,613	24,616	5
1993	241,908	6,465	10,427	16,224	26,128	37,289	42,757	39,426	29,545	23,885	3
1992	238,499	6,552	10,056	15,991	26,839	37,574	41,382	38,820	28,692	22,853	12
1991	236,409	6,108	9,868	16,480	27,774	37,558	41,162	37,836	28,029	21,727	8
1990	232,608	5,958	9,967	16,819	27,973	37,557	40,179	37,223	26,754	20,630	9
1989	228,301	5,990	9,794	17,073	28,328	36,779	39,656	35,896	25,920	19,528	5
1988	224,514	5,785	9,929	17,677	28,544	36,112	38,721	34,804	24,803	18,719	12
1987	221,757	5,587	9,889	18,288	28,516	36,167	38,410	33,944	23,666	17,898	10
1986	218,381	5,448	9,867	18,530	28,317	35,388	37,724	32,933	23,010	17,384	18
1985	215,079	5,465	9,986	18,853	28,716	34,528	36,824	32,213	21,972	16,850	14
1984	211,369	5,275	10,416	19,278	27,922	34,048	36,170	31,092	21,254	16,607	17
1983	208,053	5,358	10,563	19,249	27,348	33,642	35,400	30,570	20,447	16,334	12
1982	204,477	5,372	10,972	18,950	27,216	33,487	34,115	29,530	19,802	15,856	22
1981	199,370	5,466	11,302	19,259	26,352	32,562	33,493	27,896	19,043	15,143	16
1980	198,188	5,632	11,512	19,318	26,201	32,414	32,855	27,505	18,775	14,747	9
1979	193,444	5,747	11,736	18,705	25,858	31,596	32,042	26,306	18,460	14,099	13
Malignant neoplasms (140-208)											
White, female											
1998	226,030	7,262	11,018	14,312	18,703	25,599	34,223	36,259	31,855	36,793	5
1997	225,329	7,435	11,123	14,260	18,771	26,311	34,141	35,768	31,286	35,992	2
1996	225,454	7,632	10,641	13,884	19,291	27,468	34,786	35,361	31,123	34,993	7
1995	224,897	7,650	10,672	14,062	19,389	28,072	35,297	35,002	30,393	34,056	8
1994	222,802	7,525	10,550	13,850	19,889	29,003	34,964	34,358	29,921	32,429	2
1993	219,996	7,301	10,184	13,771	20,594	29,068	34,199	33,897	28,813	31,899	6
1992	216,032	7,096	9,878	13,640	20,940	29,078	33,742	32,952	28,142	29,926	10
1991	213,097	6,915	9,633	14,345	21,574	29,219	32,267	32,525	26,837	29,087	4
1990	208,977	6,835	9,687	13,967	21,772	29,394	31,688	31,618	25,927	27,462	6
1989	205,855	6,505	9,656	14,537	22,189	28,870	31,320	30,691	25,096	26,441	9
1988	200,626	6,503	9,530	14,886	22,607	27,963	30,556	28,986	24,065	25,221	6
1987	196,716	6,266	9,563	15,181	22,578	27,717	29,756	28,477	22,706	24,361	6
1986	193,971	6,085	9,540	15,468	22,769	27,302	29,720	27,346	22,061	23,426	15
1985	190,648	6,172	9,839	15,978	23,071	26,432	28,616	26,512	21,373	22,572	16
1984	187,262	5,958	9,978	16,308	22,439	26,219	27,617	25,751	20,824	22,065	7
1983	181,956	5,808	10,108	16,484	21,924	25,395	26,990	24,460	19,840	21,162	19
1982	178,175	5,929	10,452	16,412	21,451	25,086	25,724	24,056	19,221	20,114	21
1981	173,304	5,960	10,795	16,491	20,628	24,325	24,782	23,110	18,721	19,173	11
1980	169,974	6,081	11,032	16,462	20,241	23,736	24,382	22,062	18,293	18,240	8
1979	163,892	6,170	11,208	16,115	19,533	22,840	22,878	21,105	17,801	16,990	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
All other, both sexes											
1998	71,393	3,948	5,194	6,339	7,915	9,175	10,464	9,331	6,742	6,900	4
1997	71,056	3,779	4,825	6,250	7,917	9,392	10,565	9,153	6,684	6,996	7
1996	70,127	3,734	4,633	6,069	7,794	9,675	10,498	9,028	6,791	6,518	9
1995	69,558	3,495	4,745	5,968	8,099	9,672	10,456	8,614	6,757	6,342	8
1994	68,513	3,440	4,635	6,101	7,895	9,821	10,288	8,370	6,667	5,958	9
1993	68,000	3,331	4,664	6,003	7,965	9,844	10,152	8,442	6,414	5,919	18
1992	66,047	3,104	4,520	6,176	8,023	9,782	9,667	8,041	6,001	5,408	8
1991	65,151	2,990	4,408	6,076	7,946	9,730	9,377	8,078	6,096	5,240	9
1990	63,737	3,005	4,371	6,174	8,214	9,648	8,947	7,934	5,555	4,871	12
1989	61,996	2,906	4,287	6,023	7,961	9,573	8,830	7,800	5,209	4,597	8
1988	59,908	2,743	4,276	6,014	7,928	9,185	8,287	7,411	4,980	4,374	8
1987	58,454	2,702	4,233	5,917	7,944	8,730	8,266	7,081	4,891	4,129	17
1986	57,024	2,708	4,152	5,757	7,964	8,444	8,225	6,725	4,557	3,939	13
1985	55,836	2,578	4,163	6,153	7,855	7,981	8,161	6,454	4,535	3,660	13
1984	54,861	2,626	4,105	6,284	7,824	7,838	7,849	6,221	4,390	3,417	18
1983	52,977	2,601	4,097	6,078	7,412	7,607	7,829	6,064	4,037	3,149	32
1982	51,143	2,530	4,160	5,995	7,213	7,484	7,287	5,655	3,877	2,988	21
1981	49,420	2,470	4,147	5,868	6,772	7,413	6,971	5,358	3,661	2,813	25
1980	48,347	2,517	4,256	5,820	6,603	7,319	6,724	5,206	3,398	2,733	11
1979	46,059	2,557	4,183	5,574	6,303	7,119	6,227	5,036	3,054	2,369	10
Malignant neoplasms (140-208)											
All other, male											
1998	37,956	1,963	2,783	3,443	4,467	5,147	5,808	5,230	3,557	3,207	4
1997	37,918	1,863	2,579	3,524	4,470	5,266	5,912	5,096	3,445	3,273	5
1996	37,946	1,850	2,441	3,437	4,460	5,545	5,997	5,080	3,622	3,105	6
1995	37,611	1,660	2,528	3,275	4,610	5,554	5,997	4,852	3,572	3,080	5
1994	37,470	1,720	2,558	3,416	4,517	5,701	5,974	4,670	3,591	2,877	6
1993	37,467	1,697	2,509	3,408	4,604	5,709	5,938	4,729	3,548	2,904	9
1992	36,339	1,553	2,442	3,476	4,592	5,660	5,657	4,607	3,339	2,620	5
1991	35,971	1,547	2,458	3,483	4,475	5,751	5,406	4,583	3,362	2,530	6
1990	35,675	1,553	2,402	3,471	4,790	5,710	5,288	4,632	3,081	2,419	6
1989	35,008	1,514	2,417	3,563	4,619	5,609	5,306	4,530	2,946	2,260	6
1988	33,574	1,434	2,397	3,370	4,610	5,475	4,908	4,379	2,714	2,136	4
1987	32,896	1,353	2,298	3,362	4,609	5,314	4,924	4,192	2,750	2,049	9
1986	32,178	1,379	2,243	3,269	4,692	5,098	4,909	4,003	2,547	1,948	4
1985	31,835	1,337	2,339	3,558	4,629	4,852	4,909	3,808	2,569	1,808	7
1984	31,421	1,372	2,255	3,689	4,699	4,731	4,772	3,706	2,437	1,718	10
1983	30,330	1,378	2,216	3,522	4,505	4,604	4,706	3,615	2,328	1,536	23
1982	29,387	1,322	2,339	3,466	4,330	4,557	4,453	3,324	2,172	1,566	15
1981	28,515	1,344	2,284	3,365	4,121	4,592	4,292	3,166	2,062	1,431	18
1980	27,760	1,272	2,357	3,421	3,927	4,566	4,026	3,096	1,955	1,402	7
1979	26,571	1,309	2,324	3,295	3,774	4,357	3,816	2,959	1,793	1,235	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
All other, female											
1998	33,437	1,985	2,411	2,896	3,448	4,028	4,656	4,101	3,185	3,693	-
1997	33,138	1,916	2,246	2,726	3,447	4,126	4,653	4,057	3,239	3,723	2
1996	32,181	1,884	2,192	2,632	3,334	4,130	4,501	3,948	3,169	3,413	3
1995	31,947	1,835	2,217	2,693	3,489	4,118	4,459	3,762	3,185	3,262	3
1994	31,043	1,720	2,077	2,685	3,378	4,120	4,314	3,700	3,076	3,081	3
1993	30,533	1,634	2,155	2,595	3,361	4,135	4,214	3,713	2,866	3,015	9
1992	29,708	1,551	2,078	2,700	3,431	4,122	4,010	3,434	2,662	2,788	3
1991	29,180	1,443	1,950	2,593	3,471	3,979	3,971	3,495	2,734	2,710	3
1990	28,062	1,452	1,969	2,703	3,424	3,938	3,659	3,302	2,474	2,452	6
1989	26,988	1,392	1,870	2,460	3,342	3,964	3,524	3,270	2,263	2,337	2
1988	26,334	1,309	1,879	2,644	3,318	3,710	3,379	3,032	2,266	2,238	4
1987	25,558	1,349	1,935	2,555	3,335	3,416	3,342	2,889	2,141	2,080	8
1986	24,846	1,329	1,909	2,488	3,272	3,346	3,316	2,722	2,010	1,991	9
1985	24,001	1,241	1,824	2,595	3,226	3,129	3,252	2,646	1,966	1,852	6
1984	23,440	1,254	1,850	2,595	3,125	3,107	3,077	2,515	1,953	1,699	8
1983	22,647	1,223	1,881	2,556	2,907	3,003	3,123	2,449	1,709	1,613	9
1982	21,756	1,208	1,821	2,529	2,883	2,927	2,834	2,331	1,705	1,422	6
1981	20,905	1,126	1,863	2,503	2,651	2,821	2,679	2,192	1,599	1,382	7
1980	20,587	1,245	1,899	2,399	2,676	2,753	2,698	2,110	1,443	1,331	4
1979	19,488	1,248	1,859	2,279	2,529	2,762	2,411	2,077	1,261	1,134	4
Malignant neoplasms (140-208)											
Black, both sexes											
1998	61,193	3,353	4,434	5,460	6,842	7,851	8,997	7,983	5,810	6,015	4
1997	61,333	3,194	4,155	5,415	6,869	8,121	9,189	7,935	5,747	6,145	7
1996	60,766	3,157	4,003	5,290	6,811	8,471	9,102	7,849	5,841	5,734	9
1995	60,603	2,991	4,095	5,227	7,089	8,520	9,195	7,525	5,873	5,548	7
1994	59,939	2,962	4,032	5,374	6,941	8,621	9,095	7,294	5,875	5,266	7
1993	59,873	2,888	4,137	5,249	7,054	8,744	9,010	7,452	5,666	5,240	13
1992	58,401	2,699	3,970	5,469	7,106	8,709	8,622	7,127	5,342	4,809	5
1991	57,921	2,605	3,886	5,395	7,112	8,743	8,388	7,193	5,496	4,676	9
1990	57,077	2,649	3,878	5,569	7,449	8,688	8,051	7,108	5,002	4,384	11
1989	55,564	2,581	3,847	5,429	7,173	8,680	7,934	7,034	4,672	4,123	8
1988	53,968	2,410	3,821	5,381	7,198	8,398	7,523	6,738	4,463	3,939	8
1987	53,027	2,404	3,860	5,357	7,340	7,987	7,545	6,443	4,421	3,745	15
1986	51,979	2,446	3,769	5,278	7,318	7,809	7,558	6,153	4,117	3,575	13
1985	50,906	2,330	3,804	5,659	7,216	7,375	7,473	5,874	4,107	3,338	13
1984	50,266	2,407	3,743	5,801	7,230	7,260	7,238	5,720	4,004	3,093	17
1983	48,754	2,356	3,755	5,638	6,919	7,095	7,282	5,608	3,688	2,859	26
1982	47,316	2,304	3,826	5,607	6,722	6,964	6,828	5,249	3,567	2,736	19
1981	45,827	2,269	3,844	5,532	6,343	6,952	6,524	4,943	3,362	2,581	15
1980	45,037	2,332	3,973	5,480	6,229	6,860	6,317	4,828	3,126	2,499	11
1979	42,981	2,390	3,907	5,252	5,938	6,726	5,840	4,663	2,807	2,183	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms (140-208)											
Black, male											
1998	32,523	1,682	2,405	3,004	3,877	4,410	4,994	4,460	3,048	2,741	4
1997	32,719	1,578	2,238	3,043	3,899	4,591	5,155	4,403	2,930	2,830	5
1996	32,973	1,593	2,158	3,024	3,920	4,872	5,181	4,423	3,109	2,675	6
1995	32,880	1,448	2,227	2,898	4,088	4,917	5,255	4,230	3,093	2,644	5
1994	32,861	1,502	2,258	3,036	4,002	5,026	5,263	4,092	3,144	2,501	5
1993	33,071	1,483	2,250	3,023	4,130	5,111	5,266	4,164	3,093	2,530	5
1992	32,155	1,381	2,160	3,102	4,100	5,055	5,042	4,053	2,939	2,294	2
1991	32,094	1,379	2,201	3,146	4,047	5,170	4,845	4,063	2,989	2,231	6
1990	31,995	1,404	2,152	3,182	4,363	5,135	4,768	4,113	2,753	2,146	6
1989	31,452	1,362	2,202	3,253	4,152	5,113	4,761	4,078	2,623	2,007	6
1988	30,321	1,287	2,173	3,051	4,215	5,008	4,460	3,960	2,417	1,904	4
1987	29,928	1,203	2,111	3,087	4,287	4,895	4,510	3,790	2,453	1,835	9
1986	29,363	1,268	2,047	3,042	4,338	4,712	4,499	3,639	2,254	1,756	4
1985	29,028	1,208	2,169	3,279	4,272	4,497	4,485	3,425	2,305	1,630	7
1984	28,851	1,266	2,073	3,431	4,372	4,408	4,377	3,373	2,215	1,557	9
1983	27,946	1,269	2,047	3,291	4,194	4,308	4,349	3,310	2,116	1,398	19
1982	27,226	1,225	2,173	3,267	4,034	4,249	4,157	3,059	1,980	1,437	13
1981	26,477	1,246	2,136	3,171	3,861	4,307	3,998	2,923	1,882	1,311	10
1980	25,861	1,189	2,221	3,223	3,715	4,280	3,756	2,849	1,778	1,269	7
1979	24,831	1,242	2,188	3,116	3,566	4,113	3,559	2,728	1,635	1,137	6
Malignant neoplasms (140-208)											
Black, female											
1998	28,670	1,671	2,029	2,456	2,965	3,441	4,003	3,523	2,762	3,274	-
1997	28,614	1,616	1,917	2,372	2,970	3,530	4,034	3,532	2,817	3,315	2
1996	27,793	1,564	1,845	2,266	2,891	3,599	3,921	3,426	2,732	3,059	3
1995	27,723	1,543	1,868	2,329	3,001	3,603	3,940	3,295	2,780	2,904	2
1994	27,078	1,460	1,774	2,338	2,939	3,595	3,832	3,202	2,731	2,765	2
1993	26,802	1,405	1,887	2,226	2,924	3,633	3,744	3,288	2,573	2,710	8
1992	26,246	1,318	1,810	2,367	3,006	3,654	3,580	3,074	2,403	2,515	3
1991	25,827	1,226	1,685	2,249	3,065	3,573	3,543	3,130	2,507	2,445	3
1990	25,082	1,245	1,726	2,387	3,086	3,553	3,283	2,995	2,249	2,238	5
1989	24,112	1,219	1,645	2,176	3,021	3,567	3,173	2,956	2,049	2,116	2
1988	23,647	1,123	1,648	2,330	2,983	3,390	3,063	2,778	2,046	2,035	4
1987	23,099	1,201	1,749	2,270	3,053	3,092	3,035	2,653	1,968	1,910	6
1986	22,616	1,178	1,722	2,236	2,980	3,097	3,059	2,514	1,863	1,819	9
1985	21,878	1,122	1,635	2,380	2,944	2,878	2,988	2,449	1,802	1,708	6
1984	21,415	1,141	1,670	2,370	2,858	2,852	2,861	2,347	1,789	1,536	8
1983	20,808	1,087	1,708	2,347	2,725	2,787	2,933	2,298	1,572	1,461	7
1982	20,090	1,079	1,653	2,340	2,688	2,715	2,671	2,190	1,587	1,299	6
1981	19,350	1,023	1,708	2,361	2,482	2,645	2,526	2,020	1,480	1,270	5
1980	19,176	1,143	1,752	2,257	2,514	2,580	2,561	1,979	1,348	1,230	4
1979	18,150	1,148	1,719	2,136	2,372	2,613	2,281	1,935	1,172	1,046	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All races, both sexes											
1998	7,965	447	591	865	884	1,034	1,111	964	802	878	-
1997	7,894	443	619	714	921	1,144	1,158	931	758	784	1
1996	7,854	420	546	795	965	1,141	1,126	950	728	757	1
1995	8,060	379	638	788	957	1,208	1,179	952	750	810	-
1994	7,915	418	604	781	1,005	1,231	1,089	993	707	701	1
1993	8,241	438	649	788	1,100	1,283	1,155	966	732	709	-
1992	8,107	388	572	819	1,130	1,235	1,218	978	673	695	-
1991	8,278	387	592	847	1,200	1,260	1,186	1,035	690	670	1
1990	8,405	363	565	878	1,260	1,308	1,229	996	707	697	-
1989	8,104	378	566	968	1,233	1,270	1,168	909	621	598	3
1988	8,199	365	646	922	1,232	1,289	1,198	943	642	577	1
1987	7,968	367	571	957	1,218	1,296	1,153	866	591	587	-
1986	8,292	387	582	956	1,326	1,333	1,195	883	601	634	-
1985	8,290	386	588	1,008	1,290	1,371	1,220	895	565	554	-
1984	8,462	357	688	1,076	1,397	1,347	1,188	848	602	560	1
1983	8,486	386	642	1,124	1,369	1,369	1,162	897	533	578	4
1982	8,528	366	727	1,113	1,371	1,401	1,106	873	620	563	4
1981	8,506	364	764	1,169	1,373	1,392	1,229	803	514	556	-
1980	8,553	390	817	1,198	1,320	1,383	1,112	810	619	523	1
1979	8,589	404	869	1,155	1,386	1,363	1,132	823	550	520	3
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All races, male											
1998	5,281	363	449	659	652	736	753	605	431	355	-
1997	5,201	339	487	545	672	798	790	560	391	324	1
1996	5,215	343	428	596	707	825	761	570	378	317	1
1995	5,327	296	509	582	690	839	769	589	414	347	-
1994	5,227	328	490	595	751	819	677	598	391	294	1
1993	5,515	342	499	605	806	886	768	591	416	294	-
1992	5,366	291	441	639	828	842	806	578	356	298	-
1991	5,464	310	456	635	826	880	771	635	363	281	1
1990	5,636	295	429	660	951	915	782	582	395	315	-
1989	5,340	296	419	697	894	873	759	544	343	244	3
1988	5,497	286	491	699	884	896	764	566	367	272	1
1987	5,310	279	415	706	884	918	751	531	322	251	-
1986	5,514	304	421	693	924	913	781	530	363	291	-
1985	5,616	294	447	752	911	930	844	582	327	249	-
1984	5,765	265	522	803	978	914	830	531	370	271	1
1983	5,790	295	475	819	949	977	799	586	307	283	3
1982	5,856	261	537	824	969	1,000	765	554	376	290	2
1981	5,831	286	559	854	943	986	864	527	303	265	-
1980	5,899	293	608	895	927	952	764	549	352	274	-
1979	5,877	304	634	826	969	960	809	521	320	262	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All races, female											
1998	2,684	84	142	206	232	298	358	359	371	523	-
1997	2,693	104	132	169	249	346	368	371	367	460	-
1996	2,639	77	118	199	258	316	365	380	350	440	-
1995	2,733	83	129	206	267	369	410	363	336	463	-
1994	2,688	90	114	186	254	412	412	395	316	407	-
1993	2,726	96	150	183	294	397	387	375	316	415	-
1992	2,741	97	131	180	302	393	412	400	317	397	-
1991	2,814	77	136	212	374	380	415	400	327	389	-
1990	2,769	68	136	218	309	393	447	414	312	382	-
1989	2,764	82	147	271	339	397	409	365	278	354	-
1988	2,702	79	155	223	348	393	434	377	275	305	-
1987	2,658	88	156	251	334	378	402	335	269	336	-
1986	2,778	83	161	263	402	420	414	353	238	343	-
1985	2,674	92	141	256	379	441	376	313	238	305	-
1984	2,697	92	166	273	419	433	358	317	232	289	-
1983	2,696	91	167	305	420	392	363	311	226	295	1
1982	2,672	105	190	289	402	401	341	319	244	273	2
1981	2,675	78	205	315	430	406	365	276	211	291	-
1980	2,654	97	209	303	393	431	348	261	267	249	1
1979	2,712	100	235	329	417	403	323	302	230	258	-
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
White, both sexes											
1998	6,612	295	438	674	716	863	960	866	733	814	-
1997	6,510	296	435	538	724	962	1,002	832	706	723	-
1996	6,463	276	385	578	779	959	964	846	673	707	1
1995	6,581	249	439	582	723	1,022	1,028	842	681	761	-
1994	6,489	262	427	576	804	1,018	951	898	652	649	-
1993	6,746	309	425	584	885	1,077	1,024	872	662	655	-
1992	6,676	259	396	609	887	1,040	1,077	889	620	663	-
1991	6,830	240	417	628	978	1,074	1,044	930	640	633	1
1990	6,918	220	398	653	1,014	1,090	1,089	918	640	649	-
1989	6,693	235	388	739	1,014	1,065	1,048	822	578	561	-
1988	6,816	219	450	723	1,026	1,104	1,069	864	586	542	1
1987	6,679	235	405	736	1,014	1,146	1,031	787	550	554	-
1986	6,879	246	385	765	1,090	1,161	1,062	794	553	578	-
1985	6,938	251	417	793	1,070	1,197	1,107	806	516	528	-
1984	7,078	220	502	846	1,161	1,197	1,044	776	555	531	-
1983	7,070	235	440	881	1,157	1,216	1,034	814	495	545	1
1982	7,158	228	521	874	1,161	1,250	994	795	572	534	4
1981	7,173	231	565	928	1,170	1,248	1,114	731	468	521	-
1980	7,181	247	585	966	1,121	1,217	1,000	749	574	490	-
1979	7,307	268	635	917	1,220	1,225	1,039	760	513	494	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
White, male											
1998	4,289	236	333	504	516	625	639	544	388	321	-
1997	4,193	219	337	410	525	675	676	494	362	293	-
1996	4,178	227	298	431	567	678	643	503	347	291	1
1995	4,257	196	343	431	523	706	671	515	371	320	-
1994	4,181	199	339	434	595	670	586	537	359	275	-
1993	4,396	242	322	446	640	728	672	529	367	271	-
1992	4,283	191	300	465	639	698	702	518	320	283	-
1991	4,405	187	310	466	677	746	677	563	335	267	1
1990	4,515	183	296	486	761	750	689	524	348	291	-
1989	4,293	182	285	515	731	723	665	487	314	226	-
1988	4,462	173	331	550	724	756	676	510	327	254	1
1987	4,344	171	289	537	734	801	662	472	293	238	-
1986	4,476	194	269	549	743	785	695	477	330	259	-
1985	4,597	186	309	587	730	803	763	516	297	234	-
1984	4,688	157	375	621	792	794	715	479	332	257	-
1983	4,725	180	325	627	785	856	705	526	281	263	1
1982	4,833	159	370	643	816	884	674	509	348	273	2
1981	4,824	177	409	662	797	871	777	475	275	244	-
1980	4,827	179	437	692	772	823	679	503	321	253	-
1979	4,914	199	457	649	845	860	734	476	297	241	2
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
White, female											
1998	2,323	59	105	170	200	238	321	322	345	493	-
1997	2,317	77	98	128	199	287	326	338	344	430	-
1996	2,285	49	87	147	212	281	321	343	326	416	-
1995	2,324	53	96	151	200	316	357	327	310	441	-
1994	2,308	63	88	142	209	348	365	361	293	374	-
1993	2,350	67	103	138	245	349	352	343	295	384	-
1992	2,393	68	96	144	248	342	375	371	300	380	-
1991	2,425	53	107	162	301	328	367	367	305	366	-
1990	2,403	37	102	167	253	340	400	394	292	358	-
1989	2,400	53	103	224	283	342	383	335	264	335	-
1988	2,354	46	119	173	302	348	393	354	259	288	-
1987	2,335	64	116	199	280	345	369	315	257	316	-
1986	2,403	52	116	216	347	376	367	317	223	319	-
1985	2,341	65	108	206	340	394	344	290	219	294	-
1984	2,390	63	127	225	369	403	329	297	223	274	-
1983	2,345	55	115	254	372	360	329	288	214	282	-
1982	2,325	69	151	231	345	366	320	286	224	261	2
1981	2,349	54	156	266	373	377	337	256	193	277	-
1980	2,354	68	148	274	349	394	321	246	253	237	-
1979	2,393	69	178	268	375	365	305	284	216	253	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All other, both sexes											
1998	1,353	152	153	191	168	171	151	98	69	64	-
1997	1,384	147	184	176	197	182	156	99	52	61	1
1996	1,391	144	161	217	186	182	162	104	55	50	-
1995	1,479	130	199	206	234	186	151	110	69	49	-
1994	1,426	156	177	205	201	213	138	95	55	52	1
1993	1,495	129	224	204	215	206	131	94	70	54	-
1992	1,431	129	176	210	243	195	141	89	53	32	-
1991	1,448	147	175	219	222	186	142	105	50	37	-
1990	1,487	143	167	225	246	218	140	78	67	48	-
1989	1,411	143	178	229	219	205	120	87	43	37	3
1988	1,383	146	196	199	206	185	129	79	56	35	-
1987	1,289	132	166	221	204	150	122	79	41	33	-
1986	1,413	141	197	191	236	172	133	89	48	56	-
1985	1,352	135	171	215	220	174	113	89	49	26	-
1984	1,384	137	186	230	236	150	144	72	47	29	1
1983	1,416	151	202	243	212	153	128	83	38	33	3
1982	1,370	138	206	239	210	151	112	78	48	29	-
1981	1,333	133	199	241	203	144	115	72	46	35	-
1980	1,372	143	232	232	199	166	112	61	45	33	1
1979	1,282	136	234	238	166	138	93	63	37	26	1
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All other, male											
1998	992	127	116	155	136	111	114	61	43	34	-
1997	1,008	120	150	135	147	123	114	66	29	31	1
1996	1,037	116	130	165	140	147	118	67	31	26	-
1995	1,070	100	166	151	167	133	98	74	43	27	-
1994	1,046	129	151	161	156	149	91	61	32	19	1
1993	1,119	100	177	159	166	158	96	62	49	23	-
1992	1,083	100	141	174	189	144	104	60	36	15	-
1991	1,059	123	146	169	149	134	94	72	28	14	-
1990	1,121	112	133	174	190	165	93	58	47	24	-
1989	1,047	114	134	182	163	150	94	57	29	18	3
1988	1,035	113	160	149	160	140	88	56	40	18	-
1987	966	108	126	169	150	117	89	59	29	13	-
1986	1,038	110	152	144	181	128	86	53	33	32	-
1985	1,019	108	138	165	181	127	81	66	30	15	-
1984	1,077	108	147	182	186	120	115	52	38	14	1
1983	1,065	115	150	192	164	121	94	60	26	20	2
1982	1,023	102	167	181	153	116	91	45	28	17	-
1981	1,007	109	150	192	146	115	87	52	28	21	-
1980	1,072	114	171	203	155	129	85	46	31	21	-
1979	963	105	177	177	124	100	75	45	23	21	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
All other, female											
1998	361	25	37	36	32	60	37	37	26	30	-
1997	376	27	34	41	50	59	42	33	23	30	-
1996	354	28	31	52	46	35	44	37	24	24	-
1995	409	30	33	55	67	53	53	36	26	22	-
1994	380	27	26	44	45	64	47	34	23	33	-
1993	376	29	47	45	49	48	35	32	21	31	-
1992	348	29	35	36	54	51	37	29	17	17	-
1991	389	24	29	50	73	52	48	33	22	23	-
1990	366	31	34	51	56	53	47	20	20	24	-
1989	364	29	44	47	56	55	26	30	14	19	-
1988	348	33	36	50	46	45	41	23	16	17	-
1987	323	24	40	52	54	33	33	20	12	20	-
1986	375	31	45	47	55	44	47	36	15	24	-
1985	333	27	33	50	39	47	32	23	19	11	-
1984	307	29	39	48	50	30	29	20	9	15	-
1983	351	36	52	51	48	32	34	23	12	13	1
1982	347	36	39	58	57	35	21	33	20	12	-
1981	326	24	49	49	57	29	28	20	18	14	-
1980	300	29	61	29	44	37	27	15	14	12	1
1979	319	31	57	61	42	38	18	18	14	5	-
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
Black, both sexes											
1998	1,155	132	132	164	149	152	131	88	52	53	-
1997	1,165	115	158	159	168	154	136	84	40	52	1
1996	1,181	117	147	192	157	165	136	85	40	41	-
1995	1,274	107	172	184	207	168	128	92	60	40	-
1994	1,242	137	159	187	178	186	119	78	43	44	1
1993	1,286	108	203	181	188	177	113	84	60	40	-
1992	1,248	116	154	193	218	169	131	79	35	28	-
1991	1,274	130	152	198	200	165	128	92	42	31	-
1990	1,311	126	152	206	223	193	121	66	55	40	-
1989	1,251	133	164	206	199	183	106	76	33	32	3
1988	1,242	132	177	184	197	166	112	72	45	30	-
1987	1,161	113	158	202	193	131	111	69	36	29	-
1986	1,289	134	184	181	216	155	116	79	40	52	-
1985	1,243	129	165	204	204	163	100	78	37	21	-
1984	1,248	127	169	215	211	140	127	64	42	25	1
1983	1,288	140	185	224	202	140	117	73	35	30	3
1982	1,257	123	197	226	201	139	101	67	42	23	-
1981	1,216	123	188	227	186	132	100	64	40	30	-
1980	1,287	136	219	218	190	155	109	56	40	28	1
1979	1,204	125	221	228	158	135	85	57	30	25	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
Black, male											
1998	849	112	104	133	125	97	100	55	29	27	-
1997	851	95	128	123	125	104	100	56	21	27	1
1996	892	94	122	148	119	135	96	61	24	20	-
1995	926	83	144	134	150	119	87	64	38	20	-
1994	920	115	136	151	137	131	80	50	27	16	1
1993	968	87	161	144	145	138	80	55	41	18	-
1992	949	93	126	159	168	123	95	54	23	14	-
1991	933	108	126	153	139	118	88	60	23	13	-
1990	1,000	102	121	164	171	145	80	51	40	20	-
1989	939	105	121	163	149	139	84	50	24	16	3
1988	939	101	145	138	152	128	77	54	34	15	-
1987	864	91	120	152	141	99	81	53	26	11	-
1986	956	107	143	140	164	112	75	47	29	30	-
1985	946	103	133	158	171	120	71	57	21	13	-
1984	985	103	133	170	169	112	102	47	34	12	1
1983	986	109	141	179	157	114	85	54	25	18	2
1982	945	91	159	172	147	110	82	36	26	15	-
1981	920	100	140	182	135	104	78	48	24	16	-
1980	1,005	109	163	192	151	119	82	41	27	16	-
1979	900	96	166	169	118	99	68	39	17	20	1
Malignant neoplasms of lip, oral cavity and pharynx (140-149)											
Black, female											
1998	306	20	28	31	24	55	31	33	23	26	-
1997	314	20	30	36	43	50	36	28	19	25	-
1996	289	23	25	44	38	30	40	24	16	21	-
1995	348	24	28	50	57	49	41	28	22	20	-
1994	322	22	23	36	41	55	39	28	16	28	-
1993	318	21	42	37	43	39	33	29	19	22	-
1992	299	23	28	34	50	46	36	25	12	14	-
1991	341	22	26	45	61	47	40	32	19	18	-
1990	311	24	31	42	52	48	41	15	15	20	-
1989	312	28	43	43	50	44	22	26	9	16	-
1988	303	31	32	46	45	38	35	18	11	15	-
1987	297	22	38	50	52	32	30	16	10	18	-
1986	333	27	41	41	52	43	41	32	11	22	-
1985	297	26	32	46	33	43	29	21	16	8	-
1984	263	24	36	45	42	28	25	17	8	13	-
1983	302	31	44	45	45	26	32	19	10	12	1
1982	312	32	38	54	54	29	19	31	16	8	-
1981	296	23	48	45	51	28	22	16	16	14	-
1980	282	27	56	26	39	36	27	15	13	12	1
1979	304	29	55	59	40	36	17	18	13	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
All races, both sexes											
1998	11,765	498	799	1,070	1,371	1,722	2,007	1,797	1,188	1,037	1
1997	11,277	426	741	1,094	1,363	1,744	1,925	1,652	1,103	947	-
1996	11,231	454	697	989	1,420	1,777	1,920	1,621	1,143	911	-
1995	10,969	402	684	985	1,358	1,800	1,907	1,546	1,145	818	1
1994	10,817	452	683	931	1,425	1,790	1,904	1,434	1,079	847	-
1993	10,450	382	655	972	1,478	1,737	1,749	1,443	954	783	2
1992	10,239	363	631	933	1,451	1,791	1,750	1,361	935	734	-
1991	9,968	298	608	1,027	1,475	1,802	1,603	1,320	851	729	1
1990	9,719	341	623	939	1,456	1,752	1,579	1,268	823	689	-
1989	9,637	350	596	994	1,402	1,770	1,534	1,264	811	623	2
1988	9,336	313	632	1,001	1,451	1,655	1,452	1,203	760	652	1
1987	9,089	306	591	1,037	1,472	1,625	1,465	1,130	681	568	3
1986	8,918	321	569	1,076	1,505	1,542	1,466	1,032	663	530	-
1985	8,613	301	569	1,078	1,476	1,454	1,358	981	666	526	1
1984	8,658	284	604	1,097	1,456	1,487	1,326	979	665	557	1
1983	8,366	304	623	1,109	1,362	1,455	1,269	939	606	503	2
1982	8,225	308	615	1,099	1,337	1,479	1,215	922	589	478	2
1981	8,084	287	660	1,101	1,340	1,395	1,227	825	620	452	2
1980	7,985	325	670	1,102	1,323	1,350	1,140	879	547	468	-
1979	7,669	296	722	1,060	1,260	1,299	1,074	808	511	450	1
Malignant neoplasm of esophagus (150)											
All races, male											
1998	8,876	419	670	880	1,128	1,359	1,558	1,312	773	539	1
1997	8,546	380	647	896	1,085	1,401	1,491	1,179	734	498	-
1996	8,400	393	582	823	1,169	1,403	1,417	1,147	735	475	-
1995	8,333	341	569	822	1,102	1,432	1,491	1,142	752	410	-
1994	8,191	387	577	781	1,141	1,406	1,470	1,058	686	449	-
1993	7,813	327	549	793	1,173	1,370	1,325	1,027	611	389	2
1992	7,653	309	520	755	1,138	1,406	1,348	981	609	347	-
1991	7,427	245	522	833	1,128	1,432	1,207	926	540	385	1
1990	7,213	298	518	749	1,122	1,352	1,186	887	532	354	-
1989	7,154	294	495	823	1,127	1,361	1,127	859	513	309	2
1988	6,916	262	531	808	1,149	1,258	1,084	858	476	307	1
1987	6,723	245	479	820	1,154	1,246	1,061	809	446	286	2
1986	6,535	255	454	832	1,182	1,200	1,046	692	416	281	-
1985	6,409	243	452	873	1,148	1,103	1,024	697	431	260	1
1984	6,313	230	470	862	1,099	1,134	962	674	411	289	1
1983	6,151	249	493	859	1,041	1,109	918	662	406	256	-
1982	5,967	258	478	838	1,001	1,104	875	648	375	241	-
1981	5,953	232	508	854	1,031	1,077	903	569	394	245	2
1980	5,848	255	521	845	995	1,032	821	623	337	272	-
1979	5,623	232	580	812	961	971	810	556	332	225	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
All races, female											
1998	2,889	79	129	190	243	363	449	485	415	498	-
1997	2,731	46	94	198	278	343	434	473	369	449	-
1996	2,831	61	115	166	251	374	503	474	408	436	-
1995	2,636	61	115	163	256	368	416	404	393	408	1
1994	2,626	65	106	150	284	384	434	376	393	398	-
1993	2,637	55	106	179	305	367	424	416	343	394	-
1992	2,586	54	111	178	313	385	402	380	326	387	-
1991	2,541	53	86	194	347	370	396	394	311	344	-
1990	2,506	43	105	190	334	400	393	381	291	335	-
1989	2,483	56	101	171	275	409	407	405	298	314	-
1988	2,420	51	101	193	302	397	368	345	284	345	-
1987	2,366	61	112	217	318	379	404	321	235	282	1
1986	2,383	66	115	244	323	342	420	340	247	249	-
1985	2,204	58	117	205	328	351	334	284	235	266	-
1984	2,345	54	134	235	357	353	364	305	254	268	-
1983	2,215	55	130	250	321	346	351	277	200	247	2
1982	2,258	50	137	261	336	375	340	274	214	237	2
1981	2,131	55	152	247	309	318	324	256	226	207	-
1980	2,137	70	149	257	328	318	319	256	210	196	-
1979	2,046	64	142	248	299	328	264	252	179	225	1
Malignant neoplasm of esophagus (150)											
White, both sexes											
1998	9,810	366	602	809	1,059	1,435	1,730	1,579	1,079	936	-
1997	9,348	325	546	819	1,060	1,419	1,641	1,463	985	857	-
1996	9,253	349	485	721	1,117	1,418	1,632	1,434	1,016	838	-
1995	8,956	279	464	705	1,048	1,470	1,612	1,362	1,027	752	1
1994	8,711	319	460	653	1,083	1,432	1,587	1,254	954	776	-
1993	8,312	244	434	667	1,100	1,364	1,474	1,262	845	710	-
1992	8,196	233	400	650	1,099	1,442	1,472	1,185	834	665	-
1991	7,865	189	380	697	1,106	1,429	1,330	1,157	756	650	-
1990	7,599	205	393	633	1,090	1,385	1,304	1,088	714	624	-
1989	7,469	211	366	674	1,043	1,346	1,267	1,068	728	565	1
1988	7,182	174	370	665	1,076	1,282	1,191	1,024	676	591	-
1987	7,022	173	357	730	1,061	1,283	1,240	964	588	498	1
1986	6,786	177	341	731	1,085	1,186	1,197	884	582	476	-
1985	6,539	164	331	709	1,065	1,140	1,101	838	591	472	-
1984	6,486	143	346	705	1,044	1,154	1,068	850	557	511	-
1983	6,292	159	366	720	969	1,154	1,034	799	535	458	-
1982	6,117	148	366	699	951	1,160	974	791	517	425	-
1981	6,017	142	364	713	944	1,103	1,002	700	559	402	1
1980	5,936	160	373	704	966	1,071	931	733	475	421	-
1979	5,715	163	400	695	907	1,012	894	689	451	403	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
White, male											
1998	7,481	315	521	678	890	1,160	1,372	1,167	709	479	-
1997	7,145	301	487	686	857	1,166	1,290	1,049	660	451	-
1996	6,964	312	419	617	936	1,142	1,225	1,017	645	437	-
1995	6,824	249	396	603	862	1,176	1,272	1,012	678	374	-
1994	6,625	285	403	560	894	1,134	1,248	918	604	409	-
1993	6,234	213	377	553	903	1,099	1,122	898	535	349	-
1992	6,163	208	336	553	882	1,148	1,137	865	542	306	-
1991	5,873	160	336	579	869	1,143	1,012	815	479	329	-
1990	5,661	183	331	522	855	1,095	986	756	465	319	-
1989	5,531	186	313	560	855	1,058	922	729	458	277	1
1988	5,260	147	307	550	859	998	874	723	418	270	-
1987	5,190	147	299	593	859	989	881	682	379	247	1
1986	4,946	146	278	576	860	930	843	589	360	252	-
1985	4,841	138	272	585	845	867	827	589	375	228	-
1984	4,689	121	279	577	794	878	776	576	330	257	-
1983	4,595	136	307	569	736	889	742	549	354	228	-
1982	4,373	129	292	545	711	854	699	550	316	205	-
1981	4,372	116	285	577	724	842	725	473	348	210	1
1980	4,328	137	299	559	733	813	667	508	286	239	-
1979	4,114	130	333	536	681	748	654	471	284	192	-
Malignant neoplasm of esophagus (150)											
White, female											
1998	2,329	51	81	131	169	275	358	412	370	457	-
1997	2,203	24	59	133	203	253	351	414	325	406	-
1996	2,289	37	66	104	181	276	407	417	371	401	-
1995	2,132	30	68	102	186	294	340	350	349	378	1
1994	2,086	34	57	93	189	298	339	336	350	367	-
1993	2,078	31	57	114	197	265	352	364	310	361	-
1992	2,033	25	64	97	217	294	335	320	292	359	-
1991	1,992	29	44	118	237	286	318	342	277	321	-
1990	1,938	22	62	111	235	290	318	332	249	305	-
1989	1,938	25	53	114	188	288	345	339	270	288	-
1988	1,922	27	63	115	217	284	317	301	258	321	-
1987	1,832	26	58	137	202	294	359	282	209	251	-
1986	1,840	31	63	155	225	256	354	295	222	224	-
1985	1,698	26	59	124	220	273	274	249	216	244	-
1984	1,797	22	67	128	250	276	292	274	227	254	-
1983	1,697	23	59	151	233	265	292	250	181	230	-
1982	1,744	19	74	154	240	306	275	241	201	220	-
1981	1,645	26	79	136	220	261	277	227	211	192	-
1980	1,608	23	74	145	233	258	264	225	189	182	-
1979	1,601	33	67	159	226	264	240	218	167	211	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
All other, both sexes											
1998	1,955	132	197	261	312	287	277	218	109	101	1
1997	1,929	101	195	275	303	325	284	189	118	90	-
1996	1,978	105	212	268	303	359	288	187	127	73	-
1995	2,013	123	220	280	310	330	295	184	118	66	-
1994	2,106	133	223	278	342	358	317	180	125	71	-
1993	2,138	138	221	305	378	373	275	181	109	73	2
1992	2,043	130	231	283	352	349	278	176	101	69	-
1991	2,103	109	228	330	369	373	273	163	95	79	1
1990	2,120	136	230	306	366	367	275	180	109	65	-
1989	2,168	139	230	320	359	424	267	196	83	58	1
1988	2,154	139	262	336	375	373	261	179	84	61	1
1987	2,067	133	234	307	411	342	225	166	93	70	2
1986	2,132	144	228	345	420	356	269	148	81	54	-
1985	2,074	137	238	369	411	314	257	143	75	54	1
1984	2,172	141	258	392	412	333	258	129	108	46	1
1983	2,074	145	257	389	393	301	235	140	71	45	2
1982	2,108	160	249	400	386	319	241	131	72	53	2
1981	2,067	145	296	388	396	292	225	125	61	50	1
1980	2,049	165	297	398	357	279	209	146	72	47	-
1979	1,954	133	322	365	353	287	180	119	60	47	1
Malignant neoplasm of esophagus (150)											
All other, male											
1998	1,395	104	149	202	238	199	186	145	64	60	1
1997	1,401	79	160	210	228	235	201	130	74	47	-
1996	1,436	81	163	206	233	261	192	130	90	38	-
1995	1,509	92	173	219	240	256	219	130	74	36	-
1994	1,566	102	174	221	247	272	222	140	82	40	-
1993	1,579	114	172	240	270	271	203	129	76	40	2
1992	1,490	101	184	202	256	258	211	116	67	41	-
1991	1,554	85	186	254	259	289	195	111	61	56	1
1990	1,552	115	187	227	267	257	200	131	67	35	-
1989	1,623	108	182	263	272	303	205	130	55	32	1
1988	1,656	115	224	258	290	260	210	135	58	37	1
1987	1,533	98	180	227	295	257	180	127	67	39	1
1986	1,589	109	176	256	322	270	203	103	56	29	-
1985	1,568	105	180	288	303	236	197	108	56	32	1
1984	1,624	109	191	285	305	256	186	98	81	32	1
1983	1,556	113	186	290	305	220	176	113	52	28	-
1982	1,594	129	186	293	290	250	176	98	59	36	-
1981	1,581	116	223	277	307	235	178	96	46	35	1
1980	1,520	118	222	286	262	219	154	115	51	33	-
1979	1,509	102	247	276	280	223	156	85	48	33	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
All other, female											
1998	560	28	48	59	74	88	91	73	45	41	-
1997	528	22	35	65	75	90	83	59	44	43	-
1996	542	24	49	62	70	98	96	57	37	35	-
1995	504	31	47	61	70	74	76	54	44	30	-
1994	540	31	49	57	95	86	95	40	43	31	-
1993	559	24	49	65	108	102	72	52	33	33	-
1992	553	29	47	81	96	91	67	60	34	28	-
1991	549	24	42	76	110	84	78	52	34	23	-
1990	568	21	43	79	99	110	75	49	42	30	-
1989	545	31	48	57	87	121	62	66	28	26	-
1988	498	24	38	78	85	113	51	44	26	24	-
1987	534	35	54	80	116	85	45	39	26	31	1
1986	543	35	52	89	98	86	66	45	25	25	-
1985	506	32	58	81	108	78	60	35	19	22	-
1984	548	32	67	107	107	77	72	31	27	14	-
1983	518	32	71	99	88	81	59	27	19	17	2
1982	514	31	63	107	96	69	65	33	13	17	2
1981	486	29	73	111	89	57	47	29	15	15	-
1980	529	47	75	112	95	60	55	31	21	14	-
1979	445	31	75	89	73	64	24	34	12	14	1
Malignant neoplasm of esophagus (150)											
Black, both sexes											
1998	1,753	119	189	240	281	255	248	189	92	83	1
1997	1,746	95	181	257	278	297	252	168	99	74	-
1996	1,827	101	204	255	283	328	265	166	110	64	-
1995	1,866	121	210	267	290	300	269	166	108	54	-
1994	1,948	128	211	263	325	329	285	167	110	59	-
1993	2,011	133	214	288	351	357	254	171	99	65	1
1992	1,897	126	217	272	327	329	257	158	86	54	-
1991	1,987	104	219	316	355	346	258	149	88	70	1
1990	1,974	132	222	294	344	346	249	159	93	55	-
1989	2,047	135	226	305	342	407	244	182	73	48	1
1988	2,050	138	253	325	358	349	247	169	75	54	1
1987	1,965	129	231	297	391	325	212	153	83	61	1
1986	2,060	143	224	331	410	343	258	142	74	50	-
1985	1,985	135	230	357	396	305	242	128	70	48	1
1984	2,087	140	253	378	400	321	243	124	95	40	1
1983	1,995	140	254	379	379	291	219	131	66	40	2
1982	2,016	154	243	387	370	307	229	123	63	47	2
1981	1,986	143	289	380	380	283	209	113	58	46	1
1980	1,972	159	292	391	346	268	199	136	64	39	-
1979	1,895	131	319	358	338	281	173	109	55	44	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of esophagus (150)											
Black, male											
1998	1,245	92	143	186	213	178	167	125	52	45	1
1997	1,271	73	147	195	209	217	176	117	64	39	-
1996	1,317	79	156	195	218	236	171	114	78	31	-
1995	1,395	90	164	211	225	233	197	118	65	27	-
1994	1,434	97	163	207	232	249	196	128	70	33	-
1993	1,480	109	166	225	250	260	186	121	68	36	1
1992	1,373	98	171	194	236	242	193	101	55	32	-
1991	1,465	81	180	242	248	266	187	99	56	50	1
1990	1,433	113	179	219	246	241	178	111	58	28	-
1989	1,524	104	180	251	257	287	188	119	48	24	1
1988	1,580	114	217	249	283	241	199	126	52	32	1
1987	1,452	94	177	217	281	243	170	115	59	34	1
1986	1,536	108	172	247	316	260	196	97	51	25	-
1985	1,498	103	174	277	295	230	186	93	51	28	1
1984	1,560	108	187	274	297	245	176	94	72	27	1
1983	1,490	110	185	281	294	210	163	105	48	23	-
1982	1,517	125	180	283	276	240	167	91	50	31	-
1981	1,515	115	216	269	294	227	165	87	43	34	1
1980	1,457	113	217	284	254	208	145	106	44	27	-
1979	1,464	100	245	271	268	219	150	77	43	32	-
Malignant neoplasm of esophagus (150)											
Black, female											
1998	508	27	46	54	68	77	81	64	40	38	-
1997	475	22	34	62	69	80	76	51	35	35	-
1996	510	22	48	60	65	92	94	52	32	33	-
1995	471	31	46	56	65	67	72	48	43	27	-
1994	514	31	48	56	93	80	89	39	40	26	-
1993	531	24	48	63	101	97	68	50	31	29	-
1992	524	28	46	78	91	87	64	57	31	22	-
1991	522	23	39	74	107	80	71	50	32	20	-
1990	541	19	43	75	98	105	71	48	35	27	-
1989	523	31	46	54	85	120	56	63	25	24	-
1988	470	24	36	76	75	108	48	43	23	22	-
1987	513	35	54	80	110	82	42	38	24	27	-
1986	524	35	52	84	94	83	62	45	23	25	-
1985	487	32	56	80	101	75	56	35	19	20	-
1984	527	32	66	104	103	76	67	30	23	13	-
1983	505	30	69	98	85	81	56	26	18	17	2
1982	499	29	63	104	94	67	62	32	13	16	2
1981	471	28	73	111	86	56	44	26	15	12	-
1980	515	46	75	107	92	60	54	30	20	12	-
1979	431	31	74	87	70	62	23	32	12	12	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
All races, both sexes											
1998	12,959	481	643	788	1,008	1,463	1,890	2,082	1,849	2,123	2
1997	13,062	412	597	806	1,043	1,508	1,879	2,159	1,883	2,114	1
1996	13,336	436	571	803	1,088	1,551	2,069	2,141	1,894	2,170	1
1995	13,645	481	633	843	1,180	1,622	2,107	2,106	1,937	2,140	-
1994	13,570	416	607	785	1,222	1,642	2,140	2,221	1,847	2,068	-
1993	13,850	402	628	842	1,288	1,764	2,160	2,177	1,896	2,079	1
1992	13,630	391	577	839	1,240	1,829	2,141	2,114	1,817	2,070	-
1991	14,225	406	615	905	1,423	1,899	2,132	2,288	1,896	2,020	-
1990	14,073	420	639	892	1,371	1,877	2,194	2,292	1,790	2,003	1
1989	14,186	396	618	929	1,430	1,935	2,241	2,192	1,902	1,988	1
1988	13,676	355	577	965	1,446	1,883	2,086	2,143	1,811	1,867	2
1987	13,742	343	548	952	1,468	1,963	2,159	2,210	1,752	1,791	2
1986	13,855	390	583	1,045	1,476	1,909	2,176	2,191	1,710	1,785	1
1985	13,951	357	665	1,067	1,534	1,917	2,240	2,080	1,755	1,803	2
1984	14,246	402	673	1,065	1,549	1,959	2,284	2,096	1,879	1,834	-
1983	14,152	368	647	1,112	1,524	1,990	2,276	2,209	1,815	1,721	2
1982	14,293	379	664	1,110	1,638	2,012	2,330	2,125	1,743	1,795	3
1981	14,428	397	701	1,183	1,550	2,028	2,327	2,177	1,799	1,781	1
1980	14,372	388	690	1,118	1,626	2,026	2,283	2,195	1,798	1,796	1
1979	14,509	363	730	1,205	1,616	2,125	2,345	2,167	1,792	1,724	-
Malignant neoplasm of stomach (151)											
All races, male											
1998	7,565	296	424	533	680	978	1,204	1,231	1,001	862	1
1997	7,766	287	426	559	709	1,020	1,192	1,300	1,036	862	-
1996	7,859	265	371	567	761	1,011	1,316	1,291	1,005	913	1
1995	8,010	313	434	571	816	1,074	1,338	1,251	998	883	-
1994	8,039	281	406	562	832	1,121	1,343	1,257	997	859	-
1993	8,229	277	416	563	884	1,172	1,427	1,262	1,003	836	-
1992	8,004	277	389	572	835	1,214	1,381	1,220	937	821	-
1991	8,569	273	433	623	1,020	1,310	1,330	1,370	966	826	-
1990	8,336	275	442	628	895	1,252	1,401	1,329	938	816	1
1989	8,520	278	442	668	992	1,294	1,437	1,292	996	801	-
1988	8,105	233	406	675	1,015	1,264	1,307	1,250	908	715	-
1987	8,241	225	375	672	1,007	1,303	1,376	1,315	863	752	-
1986	8,400	267	414	732	996	1,313	1,404	1,323	837	756	1
1985	8,345	226	457	727	1,032	1,283	1,443	1,199	899	744	1
1984	8,469	258	439	729	1,064	1,317	1,421	1,208	982	770	-
1983	8,362	233	430	772	1,038	1,279	1,388	1,292	901	748	2
1982	8,506	252	442	741	1,118	1,294	1,472	1,193	895	827	2
1981	8,631	274	459	812	1,044	1,358	1,421	1,291	914	765	1
1980	8,514	249	468	729	1,123	1,367	1,394	1,252	917	743	1
1979	8,650	250	497	819	1,089	1,349	1,439	1,249	914	791	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
All races, female											
1998	5,394	185	219	255	328	485	686	851	848	1,261	1
1997	5,296	125	171	247	334	488	687	859	847	1,252	1
1996	5,477	171	200	236	327	540	753	850	889	1,257	-
1995	5,635	168	199	272	364	548	769	855	939	1,257	-
1994	5,531	135	201	223	390	521	797	964	850	1,209	-
1993	5,621	125	212	279	404	592	733	915	893	1,243	1
1992	5,626	114	188	267	405	615	760	894	880	1,249	-
1991	5,656	133	182	282	403	589	802	918	930	1,194	-
1990	5,737	145	197	264	476	625	793	963	852	1,187	-
1989	5,666	118	176	261	438	641	804	900	906	1,187	1
1988	5,571	122	171	290	431	619	779	893	903	1,152	2
1987	5,501	118	173	280	461	660	783	895	889	1,039	2
1986	5,455	123	169	313	480	596	772	868	873	1,029	-
1985	5,606	131	208	340	502	634	797	881	856	1,059	1
1984	5,777	144	234	336	485	642	863	888	897	1,064	-
1983	5,790	135	217	340	486	711	888	917	914	973	-
1982	5,787	127	222	369	520	718	858	932	848	968	1
1981	5,797	123	242	371	506	670	906	886	885	1,016	-
1980	5,858	139	222	389	503	659	889	943	881	1,053	-
1979	5,859	113	233	386	527	776	906	918	878	933	-
Malignant neoplasm of stomach (151)											
White, both sexes											
1998	10,125	356	460	588	739	1,110	1,492	1,681	1,543	1,759	2
1997	10,157	293	438	593	769	1,162	1,477	1,714	1,553	1,731	-
1996	10,456	283	438	599	811	1,232	1,648	1,728	1,549	1,773	1
1995	10,700	340	469	626	859	1,270	1,651	1,716	1,598	1,775	-
1994	10,732	293	458	582	922	1,274	1,696	1,849	1,501	1,743	-
1993	11,041	302	479	625	997	1,415	1,730	1,776	1,569	1,725	-
1992	10,827	299	420	607	937	1,464	1,721	1,735	1,515	1,738	-
1991	11,407	294	462	678	1,101	1,486	1,752	1,901	1,597	1,689	-
1990	11,309	297	470	675	1,083	1,521	1,777	1,874	1,525	1,678	1
1989	11,449	292	458	723	1,128	1,535	1,788	1,832	1,626	1,698	1
1988	11,085	263	447	733	1,140	1,504	1,717	1,762	1,537	1,589	1
1987	11,194	257	404	755	1,173	1,578	1,762	1,865	1,450	1,557	2
1986	11,274	286	435	841	1,191	1,526	1,761	1,839	1,446	1,535	1
1985	11,463	276	517	854	1,235	1,552	1,852	1,750	1,489	1,557	1
1984	11,725	308	536	841	1,262	1,594	1,875	1,758	1,599	1,595	-
1983	11,619	279	503	871	1,217	1,609	1,853	1,883	1,551	1,498	-
1982	11,828	288	524	880	1,280	1,665	1,927	1,812	1,512	1,583	2
1981	11,958	298	545	932	1,271	1,667	1,900	1,865	1,554	1,577	-
1980	12,006	310	539	888	1,354	1,668	1,887	1,861	1,572	1,619	-
1979	12,076	281	567	946	1,322	1,728	1,954	1,827	1,583	1,557	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
White, male											
1998	5,931	211	312	408	499	745	968	990	843	727	1
1997	6,074	208	315	412	520	801	953	1,053	846	712	-
1996	6,188	175	292	424	573	816	1,045	1,039	827	755	1
1995	6,351	229	323	437	601	858	1,062	1,029	830	747	-
1994	6,379	202	311	420	626	876	1,070	1,058	819	723	-
1993	6,565	211	328	422	689	945	1,147	1,026	819	703	-
1992	6,385	216	280	427	646	975	1,107	1,009	778	702	-
1991	6,868	197	326	476	810	1,024	1,091	1,129	821	690	-
1990	6,690	190	324	487	711	1,032	1,134	1,082	790	679	1
1989	6,836	201	329	515	790	1,029	1,139	1,071	855	684	-
1988	6,499	181	313	512	802	998	1,066	1,016	761	598	-
1987	6,721	178	276	538	815	1,058	1,129	1,108	704	652	-
1986	6,831	201	314	603	809	1,056	1,139	1,107	700	640	1
1985	6,842	170	357	586	832	1,035	1,201	1,007	765	640	-
1984	6,949	199	356	581	866	1,066	1,169	1,012	825	660	-
1983	6,816	169	339	612	833	1,045	1,113	1,103	745	650	-
1982	6,951	194	343	589	878	1,065	1,208	998	762	710	2
1981	7,151	210	364	640	873	1,117	1,154	1,102	784	679	-
1980	7,049	199	373	579	933	1,128	1,157	1,042	787	654	-
1979	7,150	194	391	636	894	1,089	1,193	1,044	806	709	-
Malignant neoplasm of stomach (151)											
White, female											
1998	4,194	145	148	180	240	365	524	691	700	1,032	1
1997	4,083	85	123	181	249	361	524	661	707	1,019	-
1996	4,268	108	146	175	238	416	603	689	722	1,018	-
1995	4,349	111	146	189	258	412	589	687	768	1,028	-
1994	4,353	91	147	162	296	398	626	791	682	1,020	-
1993	4,476	91	151	203	308	470	583	750	750	1,022	-
1992	4,442	83	140	180	291	489	614	726	737	1,036	-
1991	4,539	97	136	202	291	462	661	772	776	999	-
1990	4,619	107	146	188	372	489	643	792	735	999	-
1989	4,613	91	129	208	338	506	649	761	771	1,014	1
1988	4,586	82	134	221	338	506	651	746	776	991	1
1987	4,473	79	128	217	358	520	633	757	746	905	2
1986	4,443	85	121	238	382	470	622	732	746	895	-
1985	4,621	106	160	268	403	517	651	743	724	917	1
1984	4,776	109	180	260	396	528	706	746	774	935	-
1983	4,803	110	164	259	384	564	740	780	806	848	-
1982	4,877	94	181	291	402	600	719	814	750	873	-
1981	4,807	88	181	292	398	550	746	763	770	898	-
1980	4,957	111	166	309	421	540	730	819	785	965	-
1979	4,926	87	176	310	428	639	761	783	777	848	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
All other, both sexes											
1998	2,834	125	183	200	269	353	398	401	306	364	-
1997	2,905	119	159	213	274	346	402	445	330	383	1
1996	2,880	153	133	204	277	319	421	413	345	397	-
1995	2,945	141	164	217	321	352	456	390	339	365	-
1994	2,838	123	149	203	300	368	444	372	346	325	-
1993	2,809	100	149	217	291	349	430	401	327	354	1
1992	2,803	92	157	232	303	365	420	379	302	332	-
1991	2,818	112	153	227	322	413	380	387	299	331	-
1990	2,764	123	169	217	288	356	417	418	265	325	-
1989	2,737	104	160	206	302	400	453	360	276	290	-
1988	2,591	92	130	232	306	379	369	381	274	278	1
1987	2,548	86	144	197	295	385	397	345	302	234	-
1986	2,581	104	148	204	285	383	415	352	264	250	-
1985	2,488	81	148	213	299	365	388	330	266	246	1
1984	2,521	94	137	224	287	365	409	338	280	239	-
1983	2,533	89	144	241	307	381	423	326	264	223	2
1982	2,465	91	140	230	358	347	403	313	231	212	1
1981	2,470	99	156	251	279	361	427	312	245	204	1
1980	2,366	78	151	230	272	358	396	334	226	177	1
1979	2,433	82	163	259	294	397	391	340	209	167	-
Malignant neoplasm of stomach (151)											
All other, male											
1998	1,634	85	112	125	181	233	236	241	158	135	-
1997	1,692	79	111	147	189	219	239	247	190	150	-
1996	1,671	90	79	143	188	195	271	252	178	158	-
1995	1,659	84	111	134	215	216	276	222	168	136	-
1994	1,660	79	95	142	206	245	273	199	178	136	-
1993	1,664	66	88	141	195	227	280	236	184	133	-
1992	1,619	61	109	145	189	239	274	211	159	119	-
1991	1,701	76	107	147	210	286	239	241	145	136	-
1990	1,646	85	118	141	184	220	267	247	148	137	-
1989	1,684	77	113	153	202	265	298	221	141	117	-
1988	1,606	52	93	163	213	266	241	234	147	117	-
1987	1,520	47	99	134	192	245	247	207	159	100	-
1986	1,569	66	100	129	187	257	265	216	137	116	-
1985	1,503	56	100	141	200	248	242	192	134	104	1
1984	1,520	59	83	148	198	251	252	196	157	110	-
1983	1,546	64	91	160	205	234	275	189	156	98	2
1982	1,555	58	99	152	240	229	264	195	133	117	-
1981	1,480	64	95	172	171	241	267	189	130	86	1
1980	1,465	50	95	150	190	239	237	210	130	89	1
1979	1,500	56	106	183	195	260	246	205	108	82	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
All other, female											
1998	1,200	40	71	75	88	120	162	160	148	229	-
1997	1,213	40	48	66	85	127	163	198	140	233	1
1996	1,209	63	54	61	89	124	150	161	167	239	-
1995	1,286	57	53	83	106	136	180	168	171	229	-
1994	1,178	44	54	61	94	123	171	173	168	189	-
1993	1,145	34	61	76	96	122	150	165	143	221	1
1992	1,184	31	48	87	114	126	146	168	143	213	-
1991	1,117	36	46	80	112	127	141	146	154	195	-
1990	1,118	38	51	76	104	136	150	171	117	188	-
1989	1,053	27	47	53	100	135	155	139	135	173	-
1988	985	40	37	69	93	113	128	147	127	161	1
1987	1,028	39	45	63	103	140	150	138	143	134	-
1986	1,012	38	48	75	98	126	150	136	127	134	-
1985	985	25	48	72	99	117	146	138	132	142	-
1984	1,001	35	54	76	89	114	157	142	123	129	-
1983	987	25	53	81	102	147	148	137	108	125	-
1982	910	33	41	78	118	118	139	118	98	95	1
1981	990	35	61	79	108	120	160	123	115	118	-
1980	901	28	56	80	82	119	159	124	96	88	-
1979	933	26	57	76	99	137	145	135	101	85	-
Malignant neoplasm of stomach (151)											
Black, both sexes											
1998	2,175	97	133	150	195	275	303	317	241	302	-
1997	2,201	84	124	156	197	268	299	345	252	312	1
1996	2,227	120	103	152	215	245	330	320	255	334	-
1995	2,274	99	124	160	252	273	359	287	262	308	-
1994	2,206	90	106	157	233	290	353	286	282	265	-
1993	2,241	79	115	166	233	278	343	318	265	299	1
1992	2,213	69	115	175	233	286	338	302	249	275	-
1991	2,338	92	119	187	274	336	318	326	267	268	-
1990	2,258	99	132	177	238	296	337	334	229	274	-
1989	2,249	76	124	182	255	335	368	302	236	238	-
1988	2,118	71	106	175	247	308	301	331	231	230	1
1987	2,132	66	112	160	243	330	338	290	261	209	-
1986	2,118	74	115	162	238	318	348	314	219	204	-
1985	2,095	62	125	179	250	301	331	283	238	216	1
1984	2,138	80	111	194	244	316	347	302	232	200	-
1983	2,182	67	120	206	272	338	384	280	223	190	2
1982	2,120	72	115	189	310	289	363	281	208	185	1
1981	2,124	76	134	217	243	319	380	267	209	178	-
1980	2,048	63	129	200	244	316	361	293	188	147	1
1979	2,135	75	134	231	265	353	355	294	181	147	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of stomach (151)											
Black, male											
1998	1,261	71	83	93	131	184	185	191	123	104	-
1997	1,297	59	89	108	143	179	186	182	143	118	-
1996	1,301	76	62	108	152	154	207	190	128	132	-
1995	1,289	68	90	108	176	170	207	163	121	107	-
1994	1,299	63	72	109	167	193	215	153	136	108	-
1993	1,339	50	72	119	166	179	222	186	142	108	-
1992	1,273	49	86	102	154	186	219	160	128	94	-
1991	1,422	63	87	129	181	236	201	195	132	104	-
1990	1,341	70	97	121	149	187	214	188	123	109	-
1989	1,390	59	93	139	174	215	234	187	117	96	-
1988	1,316	45	77	131	174	224	191	201	117	90	-
1987	1,285	34	86	107	164	218	208	174	133	89	-
1986	1,305	51	81	106	160	216	224	186	112	93	-
1985	1,263	44	89	119	167	204	205	160	121	88	1
1984	1,306	50	71	132	169	220	209	174	133	93	-
1983	1,342	53	81	134	183	213	246	162	125	83	2
1982	1,360	48	89	128	207	192	240	174	118	106	-
1981	1,283	54	84	150	150	215	234	160	107	75	-
1980	1,273	42	84	129	169	209	213	188	105	74	1
1979	1,313	51	89	164	179	227	223	170	92	71	-
Malignant neoplasm of stomach (151)											
Black, female											
1998	914	26	50	57	64	91	118	126	118	198	-
1997	904	25	35	48	54	89	113	163	109	194	1
1996	926	44	41	44	63	91	123	130	127	202	-
1995	985	31	34	52	76	103	152	124	141	201	-
1994	907	27	34	48	66	97	138	133	146	157	-
1993	902	29	43	47	67	99	121	132	123	191	1
1992	940	20	29	73	79	100	119	142	121	181	-
1991	916	29	32	58	93	100	117	131	135	164	-
1990	917	29	35	56	89	109	123	146	106	165	-
1989	859	17	31	43	81	120	134	115	119	142	-
1988	802	26	29	44	73	84	110	130	114	140	1
1987	847	32	26	53	79	112	130	116	128	120	-
1986	813	23	34	56	78	102	124	128	107	111	-
1985	832	18	36	60	83	97	126	123	117	128	-
1984	832	30	40	62	75	96	138	128	99	107	-
1983	840	14	39	72	89	125	138	118	98	107	-
1982	760	24	26	61	103	97	123	107	90	79	1
1981	841	22	50	67	93	104	146	107	102	103	-
1980	775	21	45	71	75	107	148	105	83	73	-
1979	822	24	45	67	86	126	132	124	89	76	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
All races, both sexes											
1998	56,785	1,455	2,427	3,307	4,503	6,440	8,359	9,213	8,561	10,885	1
1997	56,503	1,577	2,288	3,160	4,501	6,480	8,430	9,133	8,592	10,737	2
1996	56,497	1,511	2,038	3,099	4,680	6,788	8,709	9,070	8,559	10,478	1
1995	57,333	1,401	2,129	3,234	4,746	7,002	9,032	9,260	8,680	10,216	2
1994	57,062	1,415	2,234	3,185	4,782	7,204	8,926	9,102	8,637	9,961	1
1993	56,955	1,287	2,121	3,137	4,961	7,224	9,102	9,331	8,295	9,939	7
1992	56,494	1,258	2,005	3,157	5,042	7,344	8,738	9,268	8,509	9,597	5
1991	56,243	1,179	1,976	3,327	5,281	7,394	8,650	9,334	8,214	9,324	3
1990	56,525	1,208	2,018	3,179	5,343	7,759	8,755	9,439	8,248	9,092	4
1989	56,261	1,107	1,959	3,274	5,495	7,615	9,036	9,419	8,097	8,876	3
1988	55,920	1,160	1,948	3,400	5,681	7,749	9,021	9,162	7,679	8,601	-
1987	56,334	1,181	2,052	3,679	5,866	7,713	9,075	9,281	7,720	8,284	3
1986	55,816	1,124	2,038	3,664	5,824	7,812	9,225	8,968	7,638	8,155	5
1985	56,451	1,137	1,983	3,861	6,116	7,768	9,204	9,240	7,594	8,130	4
1984	56,332	1,046	2,142	4,026	6,081	7,948	9,167	9,080	7,433	8,034	2
1983	54,723	1,095	2,119	3,973	5,932	7,592	9,031	8,787	7,197	7,648	6
1982	53,943	1,056	2,166	3,989	5,843	7,643	8,726	8,681	7,115	7,365	6
1981	53,249	1,138	2,224	4,058	5,662	7,529	8,650	8,528	7,036	7,141	5
1980	53,272	1,116	2,292	4,140	5,650	7,754	8,669	8,415	6,979	6,922	9
1979	52,273	1,159	2,363	4,115	5,690	7,557	8,564	8,171	6,839	6,504	3
Malignant neoplasms of colon, rectum and anus (153-154)											
All races, male											
1998	27,940	792	1,397	1,920	2,615	3,694	4,530	4,732	3,795	3,594	1
1997	27,942	843	1,317	1,780	2,646	3,799	4,612	4,699	3,870	3,516	-
1996	27,851	844	1,144	1,757	2,753	3,902	4,812	4,449	3,859	3,503	-
1995	28,261	789	1,186	1,853	2,788	4,005	4,847	4,588	3,865	3,471	-
1994	28,260	759	1,303	1,808	2,729	4,107	4,860	4,598	3,870	3,339	1
1993	27,972	715	1,185	1,802	2,848	4,016	4,941	4,728	3,603	3,305	3
1992	28,042	703	1,098	1,780	2,950	4,198	4,716	4,675	3,747	3,323	5
1991	27,773	632	1,108	1,837	3,034	4,213	4,629	4,669	3,636	3,182	1
1990	28,173	645	1,126	1,825	3,071	4,398	4,705	4,787	3,670	3,151	3
1989	27,770	611	1,080	1,891	3,155	4,245	4,894	4,608	3,533	3,007	-
1988	27,584	637	1,049	1,864	3,248	4,245	4,856	4,539	3,406	2,899	-
1987	27,889	625	1,115	2,047	3,311	4,319	4,810	4,566	3,391	2,885	1
1986	27,469	597	1,081	2,042	3,251	4,282	4,957	4,394	3,351	2,785	1
1985	27,612	613	1,077	2,145	3,413	4,300	4,891	4,468	3,228	2,726	2
1984	27,456	538	1,152	2,217	3,357	4,297	4,805	4,401	3,187	2,741	1
1983	26,912	600	1,143	2,151	3,303	4,118	4,745	4,283	3,162	2,691	4
1982	26,284	534	1,137	2,184	3,203	4,122	4,539	4,125	3,025	2,674	4
1981	25,899	588	1,167	2,130	3,130	4,084	4,574	3,998	2,959	2,570	2
1980	26,005	596	1,189	2,213	3,033	4,250	4,421	4,086	2,939	2,580	4
1979	25,405	562	1,246	2,168	3,031	4,022	4,467	3,836	2,959	2,405	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
All races, female											
1998	28,845	663	1,030	1,387	1,888	2,746	3,829	4,481	4,766	7,291	-
1997	28,561	734	971	1,380	1,855	2,681	3,818	4,434	4,722	7,221	2
1996	28,646	667	894	1,342	1,927	2,886	3,897	4,621	4,700	6,975	1
1995	29,072	612	943	1,381	1,958	2,997	4,185	4,672	4,815	6,745	2
1994	28,802	656	931	1,377	2,053	3,097	4,066	4,504	4,767	6,622	-
1993	28,983	572	936	1,335	2,113	3,208	4,161	4,603	4,692	6,634	4
1992	28,452	555	907	1,377	2,092	3,146	4,022	4,593	4,762	6,274	-
1991	28,470	547	868	1,490	2,247	3,181	4,021	4,665	4,578	6,142	2
1990	28,352	563	892	1,354	2,272	3,361	4,050	4,652	4,578	5,941	1
1989	28,491	496	879	1,383	2,340	3,370	4,142	4,811	4,564	5,869	3
1988	28,336	523	899	1,536	2,433	3,504	4,165	4,623	4,273	5,702	-
1987	28,445	556	937	1,632	2,555	3,394	4,265	4,715	4,329	5,399	2
1986	28,347	527	957	1,622	2,573	3,530	4,268	4,574	4,287	5,370	4
1985	28,839	524	906	1,716	2,703	3,468	4,313	4,772	4,366	5,404	2
1984	28,876	508	990	1,809	2,724	3,651	4,362	4,679	4,246	5,293	1
1983	27,811	495	976	1,822	2,629	3,474	4,286	4,504	4,035	4,957	2
1982	27,659	522	1,029	1,805	2,640	3,521	4,187	4,556	4,090	4,691	2
1981	27,350	550	1,057	1,928	2,532	3,445	4,076	4,530	4,077	4,571	3
1980	27,267	520	1,103	1,927	2,617	3,504	4,248	4,329	4,040	4,342	5
1979	26,868	597	1,117	1,947	2,659	3,535	4,097	4,335	3,880	4,099	2
Malignant neoplasms of colon, rectum and anus (153-154)											
White, both sexes											
1998	49,130	1,138	1,909	2,745	3,710	5,488	7,234	8,159	7,692	9,840	-
1997	48,950	1,217	1,888	2,579	3,733	5,551	7,279	8,143	7,699	9,674	2
1996	49,179	1,181	1,632	2,515	3,922	5,844	7,614	8,064	7,696	9,531	1
1995	49,988	1,099	1,725	2,702	4,006	5,974	7,904	8,237	7,841	9,272	1
1994	49,961	1,117	1,802	2,634	4,060	6,226	7,886	8,165	7,780	9,097	-
1993	49,916	1,017	1,718	2,592	4,222	6,260	7,998	8,330	7,510	9,048	2
1992	49,715	1,022	1,647	2,638	4,337	6,352	7,706	8,326	7,725	8,770	2
1991	49,556	954	1,619	2,774	4,557	6,478	7,654	8,390	7,408	8,490	2
1990	49,858	971	1,629	2,659	4,624	6,755	7,817	8,443	7,484	8,328	1
1989	49,807	917	1,611	2,766	4,782	6,666	8,068	8,409	7,373	8,129	2
1988	49,726	981	1,623	2,894	4,937	6,793	8,128	8,284	7,043	7,885	-
1987	50,113	972	1,713	3,140	5,070	6,799	8,166	8,363	7,049	7,682	1
1986	49,842	924	1,718	3,200	5,096	6,954	8,285	8,076	6,971	7,527	4
1985	50,402	928	1,672	3,343	5,359	6,919	8,195	8,388	6,957	7,514	3
1984	50,573	855	1,826	3,465	5,358	7,134	8,252	8,304	6,795	7,467	1
1983	49,254	925	1,828	3,507	5,247	6,764	8,137	8,011	6,630	7,137	3
1982	48,547	879	1,851	3,456	5,178	6,898	7,847	7,928	6,560	6,870	4
1981	48,041	953	1,917	3,545	5,069	6,767	7,836	7,748	6,495	6,694	2
1980	48,196	960	1,966	3,665	5,039	6,964	7,844	7,717	6,473	6,470	5
1979	47,362	983	2,026	3,645	5,087	6,817	7,769	7,433	6,403	6,145	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
White, male											
1998	24,299	635	1,111	1,620	2,188	3,206	3,978	4,235	3,422	3,259	-
1997	24,290	652	1,099	1,440	2,222	3,305	4,042	4,214	3,517	3,163	-
1996	24,367	679	943	1,439	2,353	3,417	4,244	3,992	3,500	3,182	-
1995	24,720	633	974	1,584	2,375	3,476	4,271	4,105	3,504	3,136	-
1994	24,818	605	1,075	1,509	2,348	3,602	4,306	4,148	3,518	3,043	-
1993	24,640	571	989	1,507	2,453	3,539	4,413	4,274	3,272	2,978	-
1992	24,767	589	914	1,519	2,578	3,672	4,221	4,196	3,409	3,016	2
1991	24,613	511	908	1,550	2,648	3,748	4,166	4,248	3,294	2,866	-
1990	24,943	531	929	1,563	2,690	3,873	4,248	4,300	3,338	2,854	1
1989	24,648	510	881	1,628	2,809	3,773	4,402	4,117	3,218	2,724	-
1988	24,634	550	882	1,603	2,847	3,781	4,395	4,137	3,136	2,643	-
1987	24,901	528	943	1,773	2,896	3,858	4,372	4,131	3,103	2,651	1
1986	24,593	505	925	1,806	2,882	3,844	4,488	3,950	3,073	2,536	1
1985	24,782	513	925	1,896	3,024	3,865	4,386	4,087	2,957	2,523	2
1984	24,681	437	979	1,941	2,987	3,891	4,342	4,041	2,916	2,527	1
1983	24,299	513	999	1,912	2,944	3,702	4,324	3,917	2,915	2,503	1
1982	23,695	445	992	1,931	2,858	3,738	4,117	3,767	2,770	2,479	2
1981	23,426	483	1,030	1,896	2,828	3,693	4,187	3,625	2,720	2,402	2
1980	23,558	520	1,038	1,965	2,720	3,849	4,025	3,746	2,714	2,397	3
1979	23,103	489	1,087	1,941	2,744	3,667	4,086	3,515	2,748	2,255	1
Malignant neoplasms of colon, rectum and anus (153-154)											
White, female											
1998	24,831	503	798	1,125	1,522	2,282	3,256	3,924	4,270	6,581	-
1997	24,660	565	789	1,139	1,511	2,246	3,237	3,929	4,182	6,511	2
1996	24,812	502	689	1,076	1,569	2,427	3,370	4,072	4,196	6,349	1
1995	25,268	466	751	1,118	1,631	2,498	3,633	4,132	4,337	6,136	1
1994	25,143	512	727	1,125	1,712	2,624	3,580	4,017	4,262	6,054	-
1993	25,276	446	729	1,085	1,769	2,721	3,585	4,056	4,238	6,070	2
1992	24,948	433	733	1,119	1,759	2,680	3,485	4,130	4,316	5,754	-
1991	24,943	443	711	1,224	1,909	2,730	3,488	4,142	4,114	5,624	2
1990	24,915	440	700	1,096	1,934	2,882	3,569	4,143	4,146	5,474	-
1989	25,159	407	730	1,138	1,973	2,893	3,666	4,292	4,155	5,405	2
1988	25,092	431	741	1,291	2,090	3,012	3,733	4,147	3,907	5,242	-
1987	25,212	444	770	1,367	2,174	2,941	3,794	4,232	3,946	5,031	-
1986	25,249	419	793	1,394	2,214	3,110	3,797	4,126	3,898	4,991	3
1985	25,620	415	747	1,447	2,335	3,054	3,809	4,301	4,000	4,991	1
1984	25,892	418	847	1,524	2,371	3,243	3,910	4,263	3,879	4,940	-
1983	24,955	412	829	1,595	2,303	3,062	3,813	4,094	3,715	4,634	2
1982	24,852	434	859	1,525	2,320	3,160	3,730	4,161	3,790	4,391	2
1981	24,615	470	887	1,649	2,241	3,074	3,649	4,123	3,775	4,292	-
1980	24,638	440	928	1,700	2,319	3,115	3,819	3,971	3,759	4,073	2
1979	24,259	494	939	1,704	2,343	3,150	3,683	3,918	3,655	3,890	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
All other, both sexes											
1998	7,655	317	518	562	793	952	1,125	1,054	869	1,045	1
1997	7,553	360	400	581	768	929	1,151	990	893	1,063	-
1996	7,318	330	406	584	758	944	1,095	1,006	863	947	-
1995	7,345	302	404	532	740	1,028	1,128	1,023	839	944	1
1994	7,101	298	432	551	722	978	1,040	937	857	864	1
1993	7,039	270	403	545	739	964	1,104	1,001	785	891	5
1992	6,779	236	358	519	705	992	1,032	942	784	827	3
1991	6,687	225	357	553	724	916	996	944	806	834	1
1990	6,667	237	389	520	719	1,004	938	996	764	764	3
1989	6,454	190	348	508	713	949	968	1,010	724	747	1
1988	6,194	179	325	506	744	956	893	878	636	716	-
1987	6,221	209	339	539	796	914	909	918	671	602	2
1986	5,974	200	320	464	728	858	940	892	667	628	1
1985	6,049	209	311	518	757	849	1,009	852	637	616	1
1984	5,759	191	316	561	723	814	915	776	638	567	1
1983	5,469	170	291	466	685	828	894	776	567	511	3
1982	5,396	177	315	533	665	745	879	753	555	495	2
1981	5,208	185	307	513	593	762	814	780	541	447	3
1980	5,076	156	326	475	611	790	825	698	506	452	4
1979	4,911	176	337	470	603	740	795	738	436	359	1
Malignant neoplasms of colon, rectum and anus (153-154)											
All other, male											
1998	3,641	157	286	300	427	488	552	497	373	335	1
1997	3,652	191	218	340	424	494	570	485	353	353	-
1996	3,484	165	201	318	400	485	568	457	359	321	-
1995	3,541	156	212	269	413	529	576	483	361	335	-
1994	3,442	154	228	299	381	505	554	450	352	296	1
1993	3,332	144	196	295	395	477	528	454	331	327	3
1992	3,275	114	184	261	372	526	495	479	338	307	3
1991	3,160	121	200	287	386	465	463	421	342	316	1
1990	3,230	114	197	262	381	525	457	487	332	297	2
1989	3,122	101	199	263	346	472	492	491	315	283	-
1988	2,950	87	167	261	401	464	461	402	270	256	-
1987	2,988	97	172	274	415	461	438	435	288	234	-
1986	2,876	92	156	236	369	438	469	444	278	249	-
1985	2,830	100	152	249	389	435	505	381	271	203	-
1984	2,775	101	173	276	370	406	463	360	271	214	-
1983	2,613	87	144	239	359	416	421	366	247	188	3
1982	2,589	89	145	253	345	384	422	358	255	195	2
1981	2,473	105	137	234	302	391	387	373	239	168	-
1980	2,447	76	151	248	313	401	396	340	225	183	1
1979	2,302	73	159	227	287	355	381	321	211	150	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
All other, female											
1998	4,014	160	232	262	366	464	573	557	496	710	-
1997	3,901	169	182	241	344	435	581	505	540	710	-
1996	3,834	165	205	266	358	459	527	549	504	626	-
1995	3,804	146	192	263	327	499	552	540	478	609	1
1994	3,659	144	204	252	341	473	486	487	505	568	-
1993	3,707	126	207	250	344	487	576	547	454	564	2
1992	3,504	122	174	258	333	466	537	463	446	520	-
1991	3,527	104	157	266	338	451	533	523	464	518	-
1990	3,437	123	192	258	338	479	481	509	432	467	1
1989	3,332	89	149	245	367	477	476	519	409	464	1
1988	3,244	92	158	245	343	492	432	476	366	460	-
1987	3,233	112	167	265	381	453	471	483	383	368	2
1986	3,098	108	164	228	359	420	471	448	389	379	1
1985	3,219	109	159	269	368	414	504	471	366	413	1
1984	2,984	90	143	285	353	408	452	416	367	353	1
1983	2,856	83	147	227	326	412	473	410	320	323	-
1982	2,807	88	170	280	320	361	457	395	300	300	-
1981	2,735	80	170	279	291	371	427	407	302	279	3
1980	2,629	80	175	227	298	389	429	358	281	269	3
1979	2,609	103	178	243	316	385	414	417	225	209	1
Malignant neoplasms of colon, rectum and anus (153-154)											
Black, both sexes											
1998	6,578	263	441	480	687	805	976	922	747	917	1
1997	6,548	302	336	506	675	800	1,006	862	777	951	-
1996	6,368	278	344	515	643	818	969	872	752	845	-
1995	6,431	261	345	471	647	898	993	903	753	833	1
1994	6,228	253	376	493	627	861	897	829	753	779	-
1993	6,211	228	365	466	642	854	977	897	706	796	3
1992	5,996	196	309	452	602	873	910	850	726	752	1
1991	5,974	197	314	495	634	815	880	853	738	764	1
1990	5,998	213	340	464	641	895	847	915	692	702	3
1989	5,768	160	317	457	628	855	858	906	652	672	1
1988	5,578	149	279	448	662	886	804	811	573	645	-
1987	5,610	172	307	486	721	839	825	831	617	546	2
1986	5,441	178	289	415	650	780	865	814	618	585	1
1985	5,492	188	280	467	687	764	914	789	578	572	1
1984	5,259	174	284	504	654	742	853	724	589	507	1
1983	5,032	155	260	433	630	782	828	727	523	455	3
1982	4,979	163	286	494	616	684	822	697	509	458	1
1981	4,802	174	274	472	531	712	767	727	491	415	2
1980	4,710	145	297	438	571	730	781	645	470	418	4
1979	4,537	156	298	433	562	689	743	687	409	322	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of colon, rectum and anus (153-154)											
Black, male											
1998	3,071	130	242	256	366	411	469	416	308	291	1
1997	3,116	161	187	294	366	425	500	409	294	304	-
1996	3,000	143	176	284	336	415	495	390	305	273	-
1995	3,054	135	178	235	351	458	496	420	317	295	-
1994	2,985	132	197	262	326	448	472	388	307	258	-
1993	2,906	125	178	252	343	423	458	402	289	280	1
1992	2,854	93	158	225	321	458	428	418	309	274	1
1991	2,769	107	172	258	333	412	397	367	301	285	1
1990	2,863	101	174	237	336	460	407	436	297	262	2
1989	2,726	87	185	240	290	421	420	428	270	247	-
1988	2,605	70	145	232	352	417	407	357	242	221	-
1987	2,642	82	155	247	374	419	392	371	255	203	-
1986	2,564	87	140	208	325	390	419	392	245	228	-
1985	2,504	88	132	222	353	381	446	337	231	186	-
1984	2,502	93	155	247	325	369	430	326	245	189	-
1983	2,359	80	125	220	322	388	382	333	220	164	3
1982	2,342	81	129	230	314	344	384	322	230	178	1
1981	2,240	101	118	214	262	354	359	343	213	152	-
1980	2,233	72	139	227	293	363	367	303	199	167	1
1979	2,082	66	140	208	264	320	339	293	195	130	-
Malignant neoplasms of colon, rectum and anus (153-154)											
Black, female											
1998	3,507	133	199	224	321	394	507	506	439	626	-
1997	3,432	141	149	212	309	375	506	453	483	647	-
1996	3,368	135	168	231	307	403	474	482	447	572	-
1995	3,377	126	167	236	296	440	497	483	436	538	1
1994	3,243	121	179	231	301	413	425	441	446	521	-
1993	3,305	103	187	214	299	431	519	495	417	516	2
1992	3,142	103	151	227	281	415	482	432	417	478	-
1991	3,205	90	142	237	301	403	483	486	437	479	-
1990	3,135	112	166	227	305	435	440	479	395	440	1
1989	3,042	73	132	217	338	434	438	478	382	425	1
1988	2,973	79	134	216	310	469	397	454	331	424	-
1987	2,968	90	152	239	347	420	433	460	362	343	2
1986	2,877	91	149	207	325	390	446	422	373	357	1
1985	2,988	100	148	245	334	383	468	452	347	386	1
1984	2,757	81	129	257	329	373	423	398	344	318	1
1983	2,673	75	135	213	308	394	446	394	303	291	-
1982	2,637	82	157	264	302	340	438	375	279	280	-
1981	2,562	73	156	258	269	358	408	384	278	263	2
1980	2,477	73	158	211	278	367	414	342	271	251	3
1979	2,455	90	158	225	298	369	404	394	214	192	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All races, both sexes											
1998	12,381	648	797	876	1,182	1,597	1,916	1,961	1,520	1,303	-
1997	11,938	551	714	854	1,130	1,621	1,883	1,906	1,441	1,261	1
1996	11,584	487	612	786	1,122	1,651	1,941	1,735	1,367	1,288	-
1995	11,191	436	538	766	1,226	1,534	1,914	1,722	1,342	1,167	-
1994	10,427	387	508	723	1,114	1,462	1,723	1,609	1,258	1,069	2
1993	10,063	337	478	695	1,063	1,495	1,693	1,537	1,198	1,055	1
1992	9,555	326	435	655	1,102	1,393	1,540	1,496	1,140	952	1
1991	8,847	275	376	599	1,006	1,313	1,436	1,401	1,055	911	-
1990	8,511	262	403	602	997	1,287	1,401	1,295	982	836	-
1989	7,988	224	358	593	943	1,245	1,301	1,203	931	775	-
1988	7,575	188	323	642	922	1,186	1,205	1,114	868	722	-
1987	7,227	203	322	568	805	1,129	1,213	1,101	804	653	1
1986	6,983	177	352	551	834	1,057	1,144	1,100	765	606	3
1985	6,809	219	334	542	819	1,038	1,172	998	709	583	2
1984	6,521	160	346	536	825	1,018	1,021	939	702	553	1
1983	6,215	211	319	537	785	940	1,021	891	597	526	4
1982	6,143	157	309	528	750	968	992	893	643	536	3
1981	5,877	182	322	530	698	888	1,002	849	588	522	-
1980	5,618	173	289	522	699	876	919	755	569	480	1
1979	5,623	167	313	530	734	937	910	743	523	459	2
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All races, male											
1998	7,780	512	621	629	837	1,072	1,224	1,172	810	503	-
1997	7,348	430	509	624	768	1,061	1,205	1,089	721	538	1
1996	7,128	372	431	561	801	1,073	1,257	976	699	525	-
1995	6,826	327	367	525	859	978	1,210	978	733	477	-
1994	6,388	284	355	511	771	986	1,016	967	665	431	-
1993	6,068	235	338	465	721	990	1,055	876	601	429	1
1992	5,854	227	305	463	775	933	986	868	563	389	-
1991	5,320	172	254	414	681	878	894	802	521	375	-
1990	5,161	175	283	422	672	825	871	755	513	350	-
1989	4,887	158	226	415	658	828	861	700	483	296	-
1988	4,621	125	213	478	623	826	735	638	421	312	-
1987	4,342	145	214	379	556	755	751	621	396	237	-
1986	4,213	109	247	399	576	704	674	607	407	233	3
1985	4,025	155	207	371	525	661	712	568	371	220	1
1984	3,940	109	221	384	552	650	636	536	372	219	1
1983	3,775	139	209	366	544	632	628	526	293	204	2
1982	3,663	104	191	357	487	627	604	519	333	219	1
1981	3,539	122	204	358	465	584	637	508	267	219	-
1980	3,292	102	203	341	442	549	547	430	278	198	1
1979	3,321	96	211	342	485	623	555	421	247	165	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All races, female											
1998	4,601	136	176	247	345	525	692	789	710	800	-
1997	4,590	121	205	230	362	560	678	817	720	723	-
1996	4,456	115	181	225	321	578	684	759	668	763	-
1995	4,365	109	171	241	367	556	704	744	609	690	-
1994	4,039	103	153	212	343	476	707	642	593	638	2
1993	3,995	102	140	230	342	505	638	661	597	626	-
1992	3,701	99	130	192	327	460	554	628	577	563	1
1991	3,527	103	122	185	325	435	542	599	534	536	-
1990	3,350	87	120	180	325	462	530	540	469	486	-
1989	3,101	66	132	178	285	417	440	503	448	479	-
1988	2,954	63	110	164	299	360	470	476	447	410	-
1987	2,885	58	108	189	249	374	462	480	408	416	1
1986	2,770	68	105	152	258	353	470	493	358	373	-
1985	2,784	64	127	171	294	377	460	430	338	363	1
1984	2,581	51	125	152	273	368	385	403	330	334	-
1983	2,440	72	110	171	241	308	393	365	304	322	2
1982	2,480	53	118	171	263	341	388	374	310	317	2
1981	2,338	60	118	172	233	304	365	341	321	303	-
1980	2,326	71	86	181	257	327	372	325	291	282	-
1979	2,302	71	102	188	249	314	355	322	276	294	-
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
White, both sexes											
1998	9,964	428	528	653	909	1,286	1,616	1,691	1,342	1,136	-
1997	9,622	363	511	630	854	1,311	1,580	1,635	1,271	1,106	-
1996	9,433	322	425	586	855	1,340	1,684	1,494	1,210	1,128	-
1995	9,050	306	368	570	966	1,234	1,607	1,472	1,173	1,017	-
1994	8,459	248	351	535	897	1,178	1,462	1,385	1,101	936	1
1993	8,236	227	335	504	833	1,231	1,451	1,336	1,043	943	-
1992	7,763	215	309	484	854	1,159	1,291	1,283	995	861	1
1991	7,195	180	253	447	794	1,093	1,203	1,214	920	806	-
1990	6,899	179	276	444	753	1,068	1,188	1,133	844	734	-
1989	6,511	154	256	433	741	1,030	1,092	1,027	832	687	-
1988	6,205	126	222	487	748	958	1,030	964	774	644	-
1987	5,923	141	216	420	666	956	1,032	942	698	583	-
1986	5,735	117	256	439	657	882	979	956	665	535	2
1985	5,587	137	241	399	647	863	992	873	646	523	1
1984	5,320	97	260	408	664	848	832	809	624	497	1
1983	5,032	144	213	413	639	779	826	780	520	465	2
1982	5,046	100	225	418	605	792	844	756	564	488	1
1981	4,882	135	231	414	553	729	834	743	536	488	-
1980	4,654	121	213	407	565	715	786	662	503	441	1
1979	4,688	120	242	426	603	762	789	645	462	419	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
White, male											
1998	6,163	340	399	466	642	862	1,040	1,016	715	438	-
1997	5,850	274	362	441	580	865	1,013	951	651	471	-
1996	5,713	238	293	413	622	860	1,088	847	624	452	-
1995	5,468	216	242	398	684	798	1,031	832	643	409	-
1994	5,092	179	242	368	614	793	871	832	584	370	-
1993	4,907	152	224	324	550	824	916	776	533	385	-
1992	4,687	141	216	340	604	772	822	751	489	350	-
1991	4,278	105	161	314	543	741	744	707	450	323	-
1990	4,124	114	186	305	506	683	748	656	441	308	-
1989	3,902	102	150	297	502	692	728	606	424	257	-
1988	3,715	78	138	370	497	667	616	558	374	278	-
1987	3,509	95	138	278	460	634	650	533	340	210	-
1986	3,377	65	175	312	441	575	575	526	358	202	2
1985	3,212	88	136	269	415	551	597	489	329	190	-
1984	3,100	60	154	289	438	528	510	447	325	191	1
1983	2,977	90	133	275	434	516	501	455	254	179	1
1982	2,952	62	135	277	391	516	519	430	285	196	-
1981	2,865	87	137	274	367	478	516	443	242	205	-
1980	2,646	64	139	256	349	440	453	370	252	181	1
1979	2,703	65	157	266	403	503	476	361	208	148	1
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
White, female											
1998	3,801	88	129	187	267	424	576	675	627	698	-
1997	3,772	89	149	189	274	446	567	684	620	635	-
1996	3,720	84	132	173	233	480	596	647	586	676	-
1995	3,582	90	126	172	282	436	576	640	530	608	-
1994	3,367	69	109	167	283	385	591	553	517	566	1
1993	3,329	75	111	180	283	407	535	560	510	558	-
1992	3,076	74	93	144	250	387	469	532	506	511	1
1991	2,917	75	92	133	251	352	459	507	470	483	-
1990	2,775	65	90	139	247	385	440	477	403	426	-
1989	2,609	52	106	136	239	338	364	421	408	430	-
1988	2,490	48	84	117	251	291	414	406	400	366	-
1987	2,414	46	78	142	206	322	382	409	358	373	-
1986	2,358	52	81	127	216	307	404	430	307	333	-
1985	2,375	49	105	130	232	312	395	384	317	333	1
1984	2,220	37	106	119	226	320	322	362	299	306	-
1983	2,055	54	80	138	205	263	325	325	266	286	1
1982	2,094	38	90	141	214	276	325	326	279	292	1
1981	2,017	48	94	140	186	251	318	300	294	283	-
1980	2,008	57	74	151	216	275	333	292	251	260	-
1979	1,985	55	85	160	200	259	313	284	254	271	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All other, both sexes											
1998	2,417	220	269	223	273	311	300	270	178	167	-
1997	2,316	188	203	224	276	310	303	271	170	155	1
1996	2,151	165	187	200	267	311	257	241	157	160	-
1995	2,141	130	170	196	260	300	307	250	169	150	-
1994	1,968	139	157	188	217	284	261	224	157	133	1
1993	1,827	110	143	191	230	264	242	201	155	112	1
1992	1,792	111	126	171	248	234	249	213	145	91	-
1991	1,652	95	123	152	212	220	233	187	135	105	-
1990	1,612	83	127	158	244	219	213	162	138	102	-
1989	1,477	70	102	160	202	215	209	176	99	88	-
1988	1,370	62	101	155	174	228	175	150	94	78	-
1987	1,304	62	106	148	139	173	181	159	106	70	1
1986	1,248	60	96	112	177	175	165	144	100	71	1
1985	1,222	82	93	143	172	175	180	125	63	60	1
1984	1,201	63	86	128	161	170	189	130	78	56	-
1983	1,183	67	106	124	146	161	195	111	77	61	2
1982	1,097	57	84	110	145	176	148	137	79	48	2
1981	995	47	91	116	145	159	168	106	52	34	-
1980	964	52	76	115	134	161	133	93	66	39	-
1979	935	47	71	104	131	175	121	98	61	40	1
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All other, male											
1998	1,617	172	222	163	195	210	184	156	95	65	-
1997	1,498	156	147	183	188	196	192	138	70	67	1
1996	1,415	134	138	148	179	213	169	129	75	73	-
1995	1,358	111	125	127	175	180	179	146	90	68	-
1994	1,296	105	113	143	157	193	145	135	81	61	-
1993	1,161	83	114	141	171	166	139	100	68	44	1
1992	1,167	86	89	123	171	161	164	117	74	39	-
1991	1,042	67	93	100	138	137	150	95	71	52	-
1990	1,037	61	97	117	166	142	123	99	72	42	-
1989	985	56	76	118	156	136	133	94	59	39	-
1988	906	47	75	108	126	159	119	80	47	34	-
1987	833	50	76	101	96	121	101	88	56	27	-
1986	836	44	72	87	135	129	99	81	49	31	1
1985	813	67	71	102	110	110	115	79	42	30	1
1984	840	49	67	95	114	122	126	89	47	28	-
1983	798	49	76	91	110	116	127	71	39	25	1
1982	711	42	56	80	96	111	85	89	48	23	1
1981	674	35	67	84	98	106	121	65	25	14	-
1980	646	38	64	85	93	109	94	60	26	17	-
1979	618	31	54	76	82	120	79	60	39	17	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
All other, female											
1998	800	48	47	60	78	101	116	114	83	102	-
1997	818	32	56	41	88	114	111	133	100	88	-
1996	736	31	49	52	88	98	88	112	82	87	-
1995	783	19	45	69	85	120	128	104	79	82	-
1994	672	34	44	45	60	91	116	89	76	72	1
1993	666	27	29	50	59	98	103	101	87	68	-
1992	625	25	37	48	77	73	85	96	71	52	-
1991	610	28	30	52	74	83	83	92	64	53	-
1990	575	22	30	41	78	77	90	63	66	60	-
1989	492	14	26	42	46	79	76	82	40	49	-
1988	464	15	26	47	48	69	56	70	47	44	-
1987	471	12	30	47	43	52	80	71	50	43	1
1986	412	16	24	25	42	46	66	63	51	40	-
1985	409	15	22	41	62	65	65	46	21	30	-
1984	361	14	19	33	47	48	63	41	31	28	-
1983	385	18	30	33	36	45	68	40	38	36	1
1982	386	15	28	30	49	65	63	48	31	25	1
1981	321	12	24	32	47	53	47	41	27	20	-
1980	318	14	12	30	41	52	39	33	40	22	-
1979	317	16	17	28	49	55	42	38	22	23	-
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
Black, both sexes											
1998	1,568	140	180	141	165	195	192	182	124	119	-
1997	1,446	119	122	135	169	199	175	169	119	107	1
1996	1,377	102	111	114	158	200	176	165	107	117	-
1995	1,421	84	105	132	163	211	200	163	110	116	-
1994	1,274	87	95	112	141	167	179	155	114	98	-
1993	1,203	68	84	125	143	173	169	142	112	77	-
1992	1,180	71	70	91	167	161	183	144	102	63	-
1991	1,108	61	72	85	139	152	164	133	97	86	-
1990	1,091	48	59	89	162	156	164	128	98	82	-
1989	1,042	48	63	108	138	154	160	137	77	67	-
1988	962	41	51	97	129	171	134	117	75	62	-
1987	938	29	73	101	108	132	139	112	81	59	-
1986	919	34	58	82	145	132	131	123	72	56	1
1985	894	44	69	92	131	134	145	100	46	45	1
1984	923	37	64	89	125	147	158	108	61	45	-
1983	903	37	74	95	115	129	163	89	64	44	-
1982	863	40	62	89	107	145	128	114	62	39	1
1981	819	32	67	98	120	140	145	92	40	30	-
1980	806	40	64	102	112	135	111	81	59	32	-
1979	775	37	60	85	112	154	102	75	50	34	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
Black, male											
1998	1,039	115	147	102	120	132	117	101	73	39	-
1997	900	99	88	107	114	130	109	73	45	42	1
1996	874	86	81	82	103	133	114	89	47	47	-
1995	867	71	74	78	113	126	112	88	55	50	-
1994	812	68	67	83	104	112	90	95	56	41	-
1993	744	48	69	89	109	110	97	69	45	27	-
1992	757	56	49	65	114	108	121	80	49	25	-
1991	678	44	51	60	89	92	105	63	46	42	-
1990	653	34	43	64	107	91	89	78	47	33	-
1989	674	40	44	79	103	103	95	70	48	27	-
1988	606	30	35	65	93	115	89	60	35	27	-
1987	572	19	51	67	73	90	77	58	41	21	-
1986	596	25	43	66	109	95	76	66	32	24	1
1985	578	34	51	62	82	85	94	62	30	21	1
1984	628	24	51	64	85	104	103	72	36	25	-
1983	581	24	52	64	84	89	102	55	30	17	-
1982	550	31	44	66	69	90	70	73	35	18	-
1981	534	22	47	67	77	95	103	52	17	12	-
1980	526	28	54	74	75	87	77	51	23	14	-
1979	506	25	45	59	70	103	67	46	30	15	1
Malignant neoplasms of liver and intrahepatic bile ducts (155)											
Black, female											
1998	529	25	33	39	45	63	75	81	51	80	-
1997	546	20	34	28	55	69	66	96	74	65	-
1996	503	16	30	32	55	67	62	76	60	70	-
1995	554	13	31	54	50	85	88	75	55	66	-
1994	462	19	28	29	37	55	89	60	58	57	-
1993	459	20	15	36	34	63	72	73	67	50	-
1992	423	15	21	26	53	53	62	64	53	38	-
1991	430	17	21	25	50	60	59	70	51	44	-
1990	438	14	16	25	55	65	75	50	51	49	-
1989	368	8	19	29	35	51	65	67	29	40	-
1988	356	11	16	32	36	56	45	57	40	35	-
1987	366	10	22	34	35	42	62	54	40	38	-
1986	323	9	15	16	36	37	55	57	40	32	-
1985	316	10	18	30	49	49	51	38	16	24	-
1984	295	13	13	25	40	43	55	36	25	20	-
1983	322	13	22	31	31	40	61	34	34	27	-
1982	313	9	18	23	38	55	58	41	27	21	1
1981	285	10	20	31	43	45	42	40	23	18	-
1980	280	12	10	28	37	48	34	30	36	18	-
1979	269	12	15	26	42	51	35	29	20	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
All races, both sexes											
1998	28,335	799	1,346	1,900	2,517	3,591	4,559	4,857	4,062	4,076	-
1997	27,675	750	1,273	1,808	2,536	3,541	4,573	4,685	3,988	3,900	-
1996	27,260	754	1,148	1,721	2,470	3,567	4,635	4,722	3,858	3,797	4
1995	26,766	726	1,157	1,648	2,588	3,595	4,674	4,453	3,662	3,705	1
1994	26,834	664	1,127	1,723	2,531	3,801	4,704	4,503	3,721	3,483	-
1993	26,445	636	1,111	1,643	2,606	3,804	4,558	4,616	3,560	3,313	2
1992	26,071	645	1,025	1,692	2,708	3,856	4,411	4,364	3,508	3,285	1
1991	25,536	625	985	1,677	2,570	3,797	4,355	4,349	3,468	3,119	2
1990	25,082	647	967	1,663	2,693	3,871	4,218	4,147	3,401	2,956	1
1989	24,543	548	959	1,659	2,739	3,778	4,286	4,024	3,184	2,862	-
1988	23,848	555	996	1,701	2,736	3,638	4,062	4,007	2,985	2,686	-
1987	23,737	507	1,029	1,767	2,817	3,630	4,012	3,858	2,926	2,704	-
1986	23,458	499	943	1,759	2,862	3,713	4,123	3,755	2,740	2,582	2
1985	23,102	562	980	1,936	2,777	3,474	4,007	3,666	2,855	2,362	3
1984	23,155	542	1,031	1,909	2,945	3,520	3,972	3,727	2,725	2,305	1
1983	22,532	491	1,064	1,896	2,797	3,435	3,911	3,556	2,604	2,302	2
1982	22,010	527	1,106	1,906	2,794	3,475	3,785	3,405	2,496	2,082	3
1981	21,792	547	1,092	1,916	2,744	3,436	3,786	3,287	2,432	2,071	2
1980	21,319	518	1,149	2,014	2,694	3,431	3,624	3,193	2,342	1,944	-
1979	21,058	546	1,154	1,873	2,740	3,370	3,494	3,204	2,329	1,895	4
Malignant neoplasm of pancreas (157)											
All races, male											
1998	13,806	480	824	1,126	1,429	1,984	2,353	2,304	1,678	1,267	-
1997	13,470	480	787	1,085	1,474	1,942	2,359	2,123	1,632	1,211	-
1996	13,027	490	710	990	1,365	1,964	2,256	2,185	1,576	1,146	3
1995	12,826	446	705	961	1,443	1,912	2,338	2,022	1,470	1,167	1
1994	12,920	421	679	1,009	1,437	2,105	2,345	2,060	1,446	1,051	-
1993	12,669	397	648	945	1,488	1,966	2,279	2,090	1,453	1,037	-
1992	12,672	401	612	1,000	1,569	2,096	2,239	2,011	1,358	1,021	-
1991	12,375	374	609	972	1,475	2,069	2,214	1,969	1,390	940	2
1990	12,199	395	570	986	1,537	2,096	2,152	1,929	1,311	888	1
1989	11,965	347	583	949	1,550	2,046	2,162	1,808	1,295	923	-
1988	11,722	348	646	999	1,582	1,967	2,010	1,827	1,147	871	-
1987	11,550	328	622	1,024	1,562	1,945	1,961	1,774	1,227	822	-
1986	11,403	296	541	1,021	1,642	1,978	2,032	1,680	1,069	822	-
1985	11,542	342	582	1,134	1,617	1,895	2,006	1,704	1,165	803	1
1984	11,517	338	639	1,131	1,640	1,879	2,027	1,740	1,066	766	1
1983	11,240	297	630	1,157	1,622	1,867	1,954	1,623	1,056	755	1
1982	11,228	338	685	1,149	1,571	1,863	1,953	1,600	1,051	766	1
1981	11,191	332	655	1,134	1,593	1,891	1,970	1,590	999	719	2
1980	10,994	322	729	1,190	1,552	1,852	1,918	1,480	1,015	695	-
1979	11,023	327	710	1,095	1,617	1,892	1,878	1,549	985	688	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
All races, female											
1998	14,529	319	522	774	1,088	1,607	2,206	2,553	2,384	2,809	-
1997	14,205	270	486	723	1,062	1,599	2,214	2,562	2,356	2,689	-
1996	14,233	264	438	731	1,105	1,603	2,379	2,537	2,282	2,651	1
1995	13,940	280	452	687	1,145	1,683	2,336	2,431	2,192	2,538	-
1994	13,914	243	448	714	1,094	1,696	2,359	2,443	2,275	2,432	-
1993	13,776	239	463	698	1,118	1,838	2,279	2,526	2,107	2,276	2
1992	13,399	244	413	692	1,139	1,760	2,172	2,353	2,150	2,264	1
1991	13,161	251	376	705	1,095	1,728	2,141	2,380	2,078	2,179	-
1990	12,883	252	397	677	1,156	1,775	2,066	2,218	2,090	2,068	-
1989	12,578	201	376	710	1,189	1,732	2,124	2,216	1,889	1,939	-
1988	12,126	207	350	702	1,154	1,671	2,052	2,180	1,838	1,815	-
1987	12,187	179	407	743	1,255	1,685	2,051	2,084	1,699	1,882	-
1986	12,055	203	402	738	1,220	1,735	2,091	2,075	1,671	1,760	2
1985	11,560	220	398	802	1,160	1,579	2,001	1,962	1,690	1,559	2
1984	11,638	204	392	778	1,305	1,641	1,945	1,987	1,659	1,539	-
1983	11,292	194	434	739	1,175	1,568	1,957	1,933	1,548	1,547	1
1982	10,782	189	421	757	1,223	1,612	1,832	1,805	1,445	1,316	2
1981	10,601	215	437	782	1,151	1,545	1,816	1,697	1,433	1,352	-
1980	10,325	196	420	824	1,142	1,579	1,706	1,713	1,327	1,249	-
1979	10,035	219	444	778	1,123	1,478	1,616	1,655	1,344	1,207	2
Malignant neoplasm of pancreas (157)											
White, both sexes											
1998	24,397	635	1,108	1,563	2,065	3,058	3,958	4,274	3,630	3,620	-
1997	23,864	593	1,050	1,485	2,114	3,004	3,983	4,132	3,562	3,454	-
1996	23,513	607	944	1,396	2,060	3,016	4,033	4,172	3,425	3,405	2
1995	23,269	617	960	1,372	2,177	3,086	4,105	3,956	3,236	3,327	-
1994	23,104	522	921	1,389	2,124	3,246	4,094	3,948	3,305	3,120	-
1993	22,859	523	896	1,355	2,182	3,241	3,981	4,077	3,159	2,978	-
1992	22,519	510	843	1,368	2,298	3,288	3,851	3,846	3,118	2,942	1
1991	22,216	522	809	1,401	2,201	3,287	3,817	3,842	3,053	2,813	1
1990	21,728	523	787	1,365	2,295	3,341	3,637	3,649	3,037	2,682	1
1989	21,236	439	776	1,347	2,297	3,248	3,767	3,569	2,829	2,565	-
1988	20,679	453	801	1,420	2,320	3,117	3,536	3,519	2,696	2,432	-
1987	20,664	416	864	1,470	2,381	3,151	3,526	3,426	2,630	2,422	-
1986	20,470	401	783	1,487	2,467	3,213	3,617	3,324	2,438	2,340	1
1985	20,173	458	800	1,631	2,353	3,048	3,512	3,273	2,546	2,154	2
1984	20,153	419	841	1,591	2,508	3,053	3,472	3,317	2,456	2,115	1
1983	19,630	398	888	1,585	2,362	2,995	3,410	3,140	2,354	2,104	1
1982	19,299	439	919	1,610	2,411	3,045	3,318	3,047	2,267	1,895	2
1981	19,262	449	919	1,654	2,406	3,008	3,324	2,939	2,246	1,926	1
1980	18,738	431	974	1,712	2,306	2,990	3,185	2,862	2,141	1,795	-
1979	18,623	452	970	1,600	2,393	2,941	3,086	2,874	2,164	1,756	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
White, male											
1998	11,978	375	687	939	1,183	1,717	2,077	2,051	1,516	1,137	-
1997	11,702	383	658	897	1,251	1,681	2,082	1,890	1,474	1,094	-
1996	11,300	388	593	806	1,172	1,689	1,986	1,942	1,431	1,023	2
1995	11,212	385	594	810	1,238	1,648	2,070	1,804	1,321	1,053	-
1994	11,149	329	562	827	1,227	1,822	2,062	1,823	1,283	936	-
1993	10,983	325	522	791	1,277	1,705	2,009	1,848	1,276	937	-
1992	10,980	320	507	809	1,355	1,812	1,990	1,776	1,203	916	-
1991	10,802	314	499	816	1,284	1,800	1,966	1,769	1,216	848	1
1990	10,582	327	462	825	1,332	1,824	1,886	1,700	1,171	787	1
1989	10,386	276	469	791	1,315	1,788	1,914	1,626	1,153	818	-
1988	10,188	288	525	846	1,354	1,706	1,767	1,611	1,038	791	-
1987	10,040	274	527	854	1,343	1,689	1,722	1,572	1,103	732	-
1986	9,974	237	464	866	1,415	1,730	1,798	1,488	958	742	-
1985	10,070	280	480	973	1,374	1,663	1,763	1,522	1,040	732	-
1984	10,033	262	532	942	1,414	1,649	1,795	1,553	951	704	1
1983	9,749	243	527	972	1,382	1,636	1,719	1,430	928	683	1
1982	9,835	284	569	978	1,344	1,645	1,712	1,444	964	693	1
1981	9,840	275	556	987	1,399	1,651	1,723	1,420	912	669	1
1980	9,676	271	629	1,011	1,347	1,623	1,711	1,325	918	640	-
1979	9,794	271	616	949	1,436	1,665	1,660	1,394	918	641	1
Malignant neoplasm of pancreas (157)											
White, female											
1998	12,419	260	421	624	882	1,341	1,881	2,223	2,114	2,483	-
1997	12,162	210	392	588	863	1,323	1,901	2,242	2,088	2,360	-
1996	12,213	219	351	590	888	1,327	2,047	2,230	1,994	2,382	-
1995	12,057	232	366	562	939	1,438	2,035	2,152	1,915	2,274	-
1994	11,955	193	359	562	897	1,424	2,032	2,125	2,022	2,184	-
1993	11,876	198	374	564	905	1,536	1,972	2,229	1,883	2,041	-
1992	11,539	190	336	559	943	1,476	1,861	2,070	1,915	2,026	1
1991	11,414	208	310	585	917	1,487	1,851	2,073	1,837	1,965	-
1990	11,146	196	325	540	963	1,517	1,751	1,949	1,866	1,895	-
1989	10,850	163	307	556	982	1,460	1,853	1,943	1,676	1,747	-
1988	10,491	165	276	574	966	1,411	1,769	1,908	1,658	1,641	-
1987	10,624	142	337	616	1,038	1,462	1,804	1,854	1,527	1,690	-
1986	10,496	164	319	621	1,052	1,483	1,819	1,836	1,480	1,598	1
1985	10,103	178	320	658	979	1,385	1,749	1,751	1,506	1,422	2
1984	10,120	157	309	649	1,094	1,404	1,677	1,764	1,505	1,411	-
1983	9,881	155	361	613	980	1,359	1,691	1,710	1,426	1,421	-
1982	9,464	155	350	632	1,067	1,400	1,606	1,603	1,303	1,202	1
1981	9,422	174	363	667	1,007	1,357	1,601	1,519	1,334	1,257	-
1980	9,062	160	345	701	959	1,367	1,474	1,537	1,223	1,155	-
1979	8,829	181	354	651	957	1,276	1,426	1,480	1,246	1,115	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
All other, both sexes											
1998	3,938	164	238	337	452	533	601	583	432	456	-
1997	3,811	157	223	323	422	537	590	553	426	446	-
1996	3,747	147	204	325	410	551	602	550	433	392	2
1995	3,497	109	197	276	411	509	569	497	426	378	1
1994	3,730	142	206	334	407	555	610	555	416	363	-
1993	3,586	113	215	288	424	563	577	539	401	335	2
1992	3,552	135	182	324	410	568	560	518	390	343	-
1991	3,320	103	176	276	369	510	538	507	415	306	1
1990	3,354	124	180	298	398	530	581	498	364	274	-
1989	3,307	109	183	312	442	530	519	455	355	297	-
1988	3,169	102	195	281	416	521	526	488	289	254	-
1987	3,073	91	165	297	436	479	486	432	296	282	-
1986	2,988	98	160	272	395	500	506	431	302	242	1
1985	2,929	104	180	305	424	426	495	393	309	208	1
1984	3,002	123	190	318	437	467	500	410	269	190	-
1983	2,902	93	176	311	435	440	501	416	250	198	1
1982	2,711	88	187	296	383	430	467	358	229	187	1
1981	2,530	98	173	262	338	428	462	348	186	145	1
1980	2,581	87	175	302	388	441	439	331	201	149	-
1979	2,435	94	184	273	347	429	408	330	165	139	1
Malignant neoplasm of pancreas (157)											
All other, male											
1998	1,828	105	137	187	246	267	276	253	162	130	-
1997	1,768	97	129	188	223	261	277	233	158	117	-
1996	1,727	102	117	184	193	275	270	243	145	123	1
1995	1,614	61	111	151	205	264	268	218	149	114	1
1994	1,771	92	117	182	210	283	283	237	163	115	-
1993	1,686	72	126	154	211	261	270	242	177	100	-
1992	1,692	81	105	191	214	284	249	235	155	105	-
1991	1,573	60	110	156	191	269	248	200	174	92	1
1990	1,617	68	108	161	205	272	266	229	140	101	-
1989	1,579	71	114	158	235	258	248	182	142	105	-
1988	1,534	60	121	153	228	261	243	216	109	80	-
1987	1,510	54	95	170	219	256	239	202	124	90	-
1986	1,429	59	77	155	227	248	234	192	111	80	-
1985	1,472	62	102	161	243	232	243	182	125	71	1
1984	1,484	76	107	189	226	230	232	187	115	62	-
1983	1,491	54	103	185	240	231	235	193	128	72	-
1982	1,393	54	116	171	227	218	241	156	87	73	-
1981	1,351	57	99	147	194	240	247	170	87	50	1
1980	1,318	51	100	179	205	229	207	155	97	55	-
1979	1,229	56	94	146	181	227	218	155	67	47	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
All other, female											
1998	2,110	59	101	150	206	266	325	330	270	326	-
1997	2,043	60	94	135	199	276	313	320	268	329	-
1996	2,020	45	87	141	217	276	332	307	288	269	1
1995	1,883	48	86	125	206	245	301	279	277	264	-
1994	1,959	50	89	152	197	272	327	318	253	248	-
1993	1,900	41	89	134	213	302	307	297	224	235	2
1992	1,860	54	77	133	196	284	311	283	235	238	-
1991	1,747	43	66	120	178	241	290	307	241	214	-
1990	1,737	56	72	137	193	258	315	269	224	173	-
1989	1,728	38	69	154	207	272	271	273	213	192	-
1988	1,635	42	74	128	188	260	283	272	180	174	-
1987	1,563	37	70	127	217	223	247	230	172	192	-
1986	1,559	39	83	117	168	252	272	239	191	162	1
1985	1,457	42	78	144	181	194	252	211	184	137	-
1984	1,518	47	83	129	211	237	268	223	154	128	-
1983	1,411	39	73	126	195	209	266	223	122	126	1
1982	1,318	34	71	125	156	212	226	202	142	114	1
1981	1,179	41	74	115	144	188	215	178	99	95	-
1980	1,263	36	75	123	183	212	232	176	104	94	-
1979	1,206	38	90	127	166	202	190	175	98	92	-
Malignant neoplasm of pancreas (157)											
Black, both sexes											
1998	3,385	141	205	288	395	460	508	500	375	391	-
1997	3,319	133	192	285	380	464	503	482	365	399	-
1996	3,222	121	178	274	362	482	513	477	353	351	2
1995	3,032	96	169	238	358	439	498	426	369	335	1
1994	3,255	126	175	292	351	483	545	479	365	317	-
1993	3,176	104	193	255	375	494	514	476	355	298	1
1992	3,180	117	157	295	359	515	503	469	354	307	-
1991	2,933	92	154	249	331	451	476	439	362	274	1
1990	3,023	109	162	271	360	479	518	439	335	253	-
1989	2,954	92	163	274	402	479	460	410	329	256	-
1988	2,824	93	171	252	368	469	477	434	252	226	-
1987	2,778	76	149	257	397	435	443	407	270	250	-
1986	2,740	85	143	257	364	454	462	407	271	222	1
1985	2,677	96	165	282	384	395	454	364	283	183	1
1984	2,764	109	176	296	401	437	463	385	245	165	-
1983	2,674	86	155	291	404	408	454	384	231	187	1
1982	2,508	73	168	274	362	393	448	337	208	172	1
1981	2,356	88	160	249	320	402	434	319	180	132	-
1980	2,401	81	166	279	370	411	418	306	180	130	-
1979	2,277	88	169	260	331	403	381	303	154	129	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of pancreas (157)											
Black, male											
1998	1,543	89	119	155	212	230	229	215	136	101	-
1997	1,511	80	112	165	196	224	237	201	126	95	-
1996	1,472	85	104	154	169	242	235	203	112	105	1
1995	1,375	54	100	127	177	225	235	176	124	95	1
1994	1,522	84	98	155	182	244	249	205	136	92	-
1993	1,475	67	113	133	193	226	236	208	153	81	-
1992	1,503	71	88	173	192	256	219	208	141	91	-
1991	1,379	56	97	142	176	237	222	163	145	74	1
1990	1,442	61	98	145	189	247	235	198	125	85	-
1989	1,403	60	101	139	214	233	219	164	126	90	-
1988	1,355	56	111	140	201	228	221	185	92	70	-
1987	1,348	46	86	153	200	231	211	185	109	77	-
1986	1,294	52	67	146	210	223	213	178	91	72	-
1985	1,327	56	93	147	217	216	216	165	113	61	1
1984	1,347	68	99	174	202	217	211	170	101	52	-
1983	1,351	50	93	172	222	207	203	177	115	66	-
1982	1,281	45	103	159	215	202	227	145	77	65	-
1981	1,258	52	90	141	180	227	233	159	84	45	-
1980	1,216	49	92	166	193	215	200	137	83	45	-
1979	1,151	54	89	138	172	213	202	141	61	44	1
Malignant neoplasm of pancreas (157)											
Black, female											
1998	1,842	52	86	133	183	230	279	285	239	290	-
1997	1,808	53	80	120	184	240	266	281	239	304	-
1996	1,750	36	74	120	193	240	278	274	241	246	1
1995	1,657	42	69	111	181	214	263	250	245	240	-
1994	1,733	42	77	137	169	239	296	274	229	225	-
1993	1,701	37	80	122	182	268	278	268	202	217	1
1992	1,677	46	69	122	167	259	284	261	213	216	-
1991	1,554	36	57	107	155	214	254	276	217	200	-
1990	1,581	48	64	126	171	232	283	241	210	168	-
1989	1,551	32	62	135	188	246	241	246	203	166	-
1988	1,469	37	60	112	167	241	256	249	160	156	-
1987	1,430	30	63	104	197	204	232	222	161	173	-
1986	1,446	33	76	111	154	231	249	229	180	150	1
1985	1,350	40	72	135	167	179	238	199	170	122	-
1984	1,417	41	77	122	199	220	252	215	144	113	-
1983	1,323	36	62	119	182	201	251	207	116	121	1
1982	1,227	28	65	115	147	191	221	192	131	107	1
1981	1,098	36	70	108	140	175	201	160	96	87	-
1980	1,185	32	74	113	177	196	218	169	97	85	-
1979	1,126	34	80	122	159	190	179	162	93	85	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
All races, both sexes											
1998	3,866	157	279	402	500	616	655	545	381	243	-
1997	3,811	148	247	395	560	609	624	546	354	258	1
1996	3,918	159	272	412	578	649	654	524	337	241	-
1995	3,871	165	269	363	529	681	641	555	358	229	-
1994	3,947	139	257	390	577	697	701	479	363	249	-
1993	3,981	137	266	409	533	722	699	530	340	260	-
1992	3,966	133	267	386	597	738	667	539	325	219	-
1991	3,989	153	281	427	581	737	653	485	353	213	3
1990	3,710	114	269	372	611	701	624	473	280	186	1
1989	3,729	138	238	423	603	648	594	487	311	189	2
1988	3,667	133	240	435	602	680	578	457	262	201	1
1987	3,671	104	260	446	611	662	591	453	245	215	1
1986	3,611	115	231	481	628	646	538	444	274	172	1
1985	3,502	126	245	455	599	665	523	422	229	174	1
1984	3,626	116	280	503	652	670	520	395	269	155	1
1983	3,605	141	269	454	635	681	568	404	232	153	1
1982	3,636	100	310	492	626	698	599	381	213	144	-
1981	3,531	115	295	504	599	620	529	384	243	168	2
1980	3,412	134	272	521	552	669	507	347	233	114	1
1979	3,446	143	321	433	604	655	521	338	219	141	-
Malignant neoplasm of larynx (161)											
All races, male											
1998	3,002	131	229	328	394	486	505	404	274	180	-
1997	3,017	122	204	328	445	467	497	426	275	207	1
1996	3,095	130	222	345	461	505	508	406	258	186	-
1995	3,081	135	228	300	422	548	485	429	287	181	-
1994	3,127	117	211	315	463	543	542	360	300	204	-
1993	3,163	113	226	338	421	582	556	403	262	197	-
1992	3,114	115	208	302	469	570	528	404	266	174	-
1991	3,127	112	229	323	469	569	514	375	287	164	2
1990	2,977	97	224	303	481	552	498	369	236	155	1
1989	2,961	114	186	336	475	501	477	380	260	152	1
1988	2,965	115	194	350	476	545	480	373	207	160	-
1987	2,975	83	208	366	490	535	485	365	206	173	1
1986	2,906	87	192	382	496	516	430	370	220	145	1
1985	2,844	98	198	370	489	549	414	343	192	140	-
1984	2,960	93	228	415	530	541	417	328	230	128	1
1983	2,943	112	212	365	517	554	473	332	199	125	1
1982	2,971	85	246	417	485	563	501	321	173	126	-
1981	2,905	91	241	399	470	514	445	330	215	144	2
1980	2,845	101	225	417	455	570	426	296	215	91	1
1979	2,910	116	267	354	508	547	448	290	199	122	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
All races, female											
1998	864	26	50	74	106	130	150	141	107	63	-
1997	794	26	43	67	115	142	127	120	79	51	-
1996	823	29	50	67	117	144	146	118	79	55	-
1995	790	30	41	63	107	133	156	126	71	48	-
1994	820	22	46	75	114	154	159	119	63	45	-
1993	818	24	40	71	112	140	143	127	78	63	-
1992	852	18	59	84	128	168	139	135	59	45	-
1991	862	41	52	104	112	168	139	110	66	49	1
1990	733	17	45	69	130	149	126	104	44	31	-
1989	768	24	52	87	128	147	117	107	51	37	1
1988	702	18	46	85	126	135	98	84	55	41	1
1987	696	21	52	80	121	127	106	88	39	42	-
1986	705	28	39	99	132	130	108	74	54	27	-
1985	658	28	47	85	110	116	109	79	37	34	1
1984	666	23	52	88	122	129	103	67	39	27	-
1983	662	29	57	89	118	127	95	72	33	28	-
1982	665	15	64	75	141	135	98	60	40	18	-
1981	626	24	54	105	129	106	84	54	28	24	-
1980	567	33	47	104	97	99	81	51	18	23	-
1979	536	27	54	79	96	108	73	48	20	19	-
Malignant neoplasm of larynx (161)											
White, both sexes											
1998	3,106	88	203	287	383	491	548	476	348	223	-
1997	3,041	91	180	288	428	493	516	469	308	223	-
1996	3,133	100	195	302	437	519	543	454	302	220	-
1995	3,127	107	185	268	412	562	545	467	323	197	-
1994	3,174	86	181	281	443	577	598	416	320	214	-
1993	3,233	82	171	297	422	611	584	458	309	237	-
1992	3,220	84	176	273	485	604	579	474	292	200	-
1991	3,245	88	180	325	471	600	570	441	320	185	2
1990	3,046	71	180	281	495	594	541	413	258	164	1
1989	3,037	79	161	317	486	539	523	433	274	170	2
1988	2,975	90	159	339	474	554	502	387	240	180	-
1987	3,047	65	185	353	497	563	521	402	216	196	-
1986	2,975	76	160	384	503	552	474	387	239	158	-
1985	2,883	76	166	350	490	576	455	381	206	152	-
1984	2,995	82	197	391	519	566	454	357	247	142	1
1983	3,052	92	200	367	532	601	508	370	210	138	-
1982	3,088	69	234	401	524	617	536	344	195	128	-
1981	2,960	65	215	401	490	548	473	346	225	155	1
1980	2,849	93	199	414	471	565	454	304	218	98	-
1979	2,957	102	240	354	517	570	472	316	200	135	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
White, male											
1998	2,381	74	165	227	299	388	419	344	252	165	-
1997	2,388	76	145	239	333	386	404	358	240	179	-
1996	2,447	84	161	251	340	405	417	346	227	168	-
1995	2,483	88	156	223	321	458	417	357	258	155	-
1994	2,501	75	145	226	355	446	467	308	265	173	-
1993	2,560	67	143	252	334	487	468	342	240	178	-
1992	2,512	74	137	214	381	460	454	349	240	159	-
1991	2,540	63	145	248	384	467	445	338	260	139	2
1990	2,448	59	154	235	388	467	431	326	216	135	1
1989	2,377	62	125	245	385	408	415	333	226	136	1
1988	2,370	77	126	268	370	437	410	308	189	142	-
1987	2,460	53	146	295	393	452	427	323	181	157	-
1986	2,390	58	129	307	405	440	377	318	188	132	-
1985	2,341	61	132	281	401	471	360	314	172	124	-
1984	2,421	61	156	319	425	449	360	296	211	115	1
1983	2,492	79	159	292	432	484	421	304	180	113	-
1982	2,514	61	185	337	403	495	445	290	157	111	-
1981	2,453	54	181	322	388	453	390	297	199	134	1
1980	2,377	75	168	334	386	481	375	254	200	77	-
1979	2,488	86	198	289	433	469	404	270	182	116	-
Malignant neoplasm of larynx (161)											
White, female											
1998	725	14	38	60	84	103	129	132	96	58	-
1997	653	15	35	49	95	107	112	111	68	44	-
1996	686	16	34	51	97	114	126	108	75	52	-
1995	644	19	29	45	91	104	128	110	65	42	-
1994	673	11	36	55	88	131	131	108	55	41	-
1993	673	15	28	45	88	124	116	116	69	59	-
1992	708	10	39	59	104	144	125	125	52	41	-
1991	705	25	35	77	87	133	125	103	60	46	-
1990	598	12	26	46	107	127	110	87	42	29	-
1989	660	17	36	72	101	131	108	100	48	34	1
1988	605	13	33	71	104	117	92	79	51	38	-
1987	587	12	39	58	104	111	94	79	35	39	-
1986	585	18	31	77	98	112	97	69	51	26	-
1985	542	15	34	69	89	105	95	67	34	28	-
1984	574	21	41	72	94	117	94	61	36	27	-
1983	560	13	41	75	100	117	87	66	30	25	-
1982	574	8	49	64	121	122	91	54	38	17	-
1981	507	11	34	79	102	95	83	49	26	21	-
1980	472	18	31	80	85	84	79	50	18	21	-
1979	469	16	42	65	84	101	68	46	18	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
All other, both sexes											
1998	760	69	76	115	117	125	107	69	33	20	-
1997	770	57	67	107	132	116	108	77	46	35	1
1996	785	59	77	110	141	130	111	70	35	21	-
1995	744	58	84	95	117	119	96	88	35	32	-
1994	773	53	76	109	134	120	103	63	43	35	-
1993	748	55	95	112	111	111	115	72	31	23	-
1992	746	49	91	113	112	134	88	65	33	19	-
1991	744	65	101	102	110	137	83	44	33	28	1
1990	664	43	89	91	116	107	83	60	22	22	-
1989	692	59	77	106	117	109	71	54	37	19	-
1988	692	43	81	96	128	126	76	70	22	21	1
1987	624	39	75	93	114	99	70	51	29	19	1
1986	636	39	71	97	125	94	64	57	35	14	1
1985	619	50	79	105	109	89	68	41	23	22	1
1984	631	34	83	112	133	104	66	38	22	13	-
1983	553	49	69	87	103	80	60	34	22	15	1
1982	548	31	76	91	102	81	63	37	18	16	-
1981	571	50	80	103	109	72	56	38	18	13	1
1980	563	41	73	107	81	104	53	43	15	16	1
1979	489	41	81	79	87	85	49	22	19	6	-
Malignant neoplasm of larynx (161)											
All other, male											
1998	621	57	64	101	95	98	86	60	22	15	-
1997	629	46	59	89	112	81	93	68	35	28	1
1996	648	46	61	94	121	100	91	60	31	18	-
1995	598	47	72	77	101	90	68	72	29	26	-
1994	626	42	66	89	108	97	75	52	35	31	-
1993	603	46	83	86	87	95	88	61	22	19	-
1992	602	41	71	88	88	110	74	55	26	15	-
1991	587	49	84	75	85	102	69	37	27	25	-
1990	529	38	70	68	93	85	67	43	20	20	-
1989	584	52	61	91	90	93	62	47	34	16	-
1988	595	38	68	82	106	108	70	65	18	18	-
1987	515	30	62	71	97	83	58	42	25	16	1
1986	516	29	63	75	91	76	53	52	32	13	1
1985	503	37	66	89	88	78	54	29	20	16	-
1984	539	32	72	96	105	92	57	32	19	13	-
1983	451	33	53	73	85	70	52	28	19	12	1
1982	457	24	61	80	82	68	56	31	16	15	-
1981	452	37	60	77	82	61	55	33	16	10	1
1980	468	26	57	83	69	89	51	42	15	14	1
1979	422	30	69	65	75	78	44	20	17	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
All other, female											
1998	139	12	12	14	22	27	21	9	11	5	-
1997	141	11	8	18	20	35	15	9	11	7	-
1996	137	13	16	16	20	30	20	10	4	3	-
1995	146	11	12	18	16	29	28	16	6	6	-
1994	147	11	10	20	26	23	28	11	8	4	-
1993	145	9	12	26	24	16	27	11	9	4	-
1992	144	8	20	25	24	24	14	10	7	4	-
1991	157	16	17	27	25	35	14	7	6	3	1
1990	135	5	19	23	23	22	16	17	2	2	-
1989	108	7	16	15	27	16	9	7	3	3	-
1988	97	5	13	14	22	18	6	5	4	3	1
1987	109	9	13	22	17	16	12	9	4	3	-
1986	120	10	8	22	34	18	11	5	3	1	-
1985	116	13	13	16	21	11	14	12	3	6	1
1984	92	2	11	16	28	12	9	6	3	-	-
1983	102	16	16	14	18	10	8	6	3	3	-
1982	91	7	15	11	20	13	7	6	2	1	-
1981	119	13	20	26	27	11	1	5	2	3	-
1980	95	15	16	24	12	15	2	1	-	2	-
1979	67	11	12	14	12	7	5	2	2	-	-
Malignant neoplasm of larynx (161)											
Black, both sexes											
1998	713	66	70	111	111	119	101	59	29	18	-
1997	720	54	64	102	125	109	99	70	41	31	1
1996	756	56	74	109	137	125	109	63	34	19	-
1995	700	56	82	89	112	116	90	77	31	28	-
1994	727	52	75	103	126	112	95	59	40	31	-
1993	718	53	93	110	109	105	107	68	27	23	-
1992	720	49	90	113	110	128	85	61	28	14	-
1991	717	64	100	100	108	129	80	38	31	26	1
1990	631	43	88	87	108	103	76	54	21	21	-
1989	666	59	76	101	113	105	69	50	31	19	-
1988	664	42	80	94	123	120	75	61	21	20	1
1987	594	37	74	88	109	94	63	48	28	18	1
1986	615	39	69	95	119	92	59	56	33	14	1
1985	586	49	75	101	104	80	67	38	23	18	1
1984	608	34	81	112	129	100	60	34	20	12	-
1983	528	47	69	83	102	76	55	28	20	14	1
1982	533	31	72	91	97	81	61	36	17	14	-
1981	557	49	80	103	108	69	55	33	16	13	1
1980	542	40	71	101	80	99	51	42	14	14	1
1979	478	40	81	78	85	82	47	21	18	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of larynx (161)											
Black, male											
1998	582	54	60	97	90	93	81	51	20	13	-
1997	592	44	56	85	108	77	85	64	30	25	1
1996	621	44	58	93	118	95	89	53	30	16	-
1995	560	45	70	72	96	88	64	62	25	23	-
1994	588	41	65	84	100	91	68	49	34	28	-
1993	580	45	81	85	85	91	81	58	19	19	-
1992	584	41	70	88	87	107	72	52	22	11	-
1991	565	48	83	73	85	95	66	33	25	23	-
1990	499	38	69	64	87	81	61	37	19	19	-
1989	565	52	60	86	89	91	61	43	29	16	-
1988	571	37	67	80	103	102	69	56	18	17	-
1987	493	28	62	68	93	78	53	41	24	15	1
1986	499	29	61	74	87	74	49	51	30	13	1
1985	479	36	62	85	83	71	53	28	20	15	-
1984	520	32	70	96	103	88	51	28	19	12	-
1983	433	31	53	70	84	69	48	23	18	11	1
1982	443	24	57	80	78	68	54	30	15	13	-
1981	443	36	60	77	81	60	54	30	15	10	1
1980	449	26	55	78	68	84	49	41	14	12	1
1979	412	29	69	64	73	76	42	19	16	6	-
Malignant neoplasm of larynx (161)											
Black, female											
1998	131	12	10	14	21	26	20	8	9	5	-
1997	128	10	8	17	17	32	14	6	11	6	-
1996	135	12	16	16	19	30	20	10	4	3	-
1995	140	11	12	17	16	28	26	15	6	5	-
1994	139	11	10	19	26	21	27	10	6	3	-
1993	138	8	12	25	24	14	26	10	8	4	-
1992	136	8	20	25	23	21	13	9	6	3	-
1991	152	16	17	27	23	34	14	5	6	3	1
1990	132	5	19	23	21	22	15	17	2	2	-
1989	101	7	16	15	24	14	8	7	2	3	-
1988	93	5	13	14	20	18	6	5	3	3	1
1987	101	9	12	20	16	16	10	7	4	3	-
1986	116	10	8	21	32	18	10	5	3	1	-
1985	107	13	13	16	21	9	14	10	3	3	1
1984	88	2	11	16	26	12	9	6	1	-	-
1983	95	16	16	13	18	7	7	5	2	3	-
1982	90	7	15	11	19	13	7	6	2	1	-
1981	114	13	20	26	27	9	1	3	1	3	-
1980	93	14	16	23	12	15	2	1	-	2	-
1979	66	11	12	14	12	6	5	2	2	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
All races, both sexes											
1998	154,561	3,965	7,535	12,152	17,635	24,424	30,153	26,289	17,664	11,753	2
1997	153,310	3,934	7,712	11,837	17,464	25,198	29,510	25,926	17,207	11,521	1
1996	152,015	4,146	7,766	11,792	17,838	25,878	29,398	24,752	16,903	10,567	8
1995	151,200	4,129	7,681	11,937	18,260	26,123	29,606	24,212	16,070	10,303	5
1994	149,482	4,019	7,669	12,087	18,977	26,810	28,666	23,481	15,492	9,459	6
1993	148,855	4,206	7,534	12,406	19,672	26,981	28,242	23,014	14,745	9,202	7
1992	145,943	4,199	7,538	12,308	19,926	26,829	27,309	22,456	14,068	8,380	6
1991	143,758	4,143	7,383	12,642	20,391	26,693	26,318	21,781	13,475	8,040	5
1990	141,285	4,179	7,556	12,972	20,943	26,330	25,709	21,020	12,346	7,328	4
1989	137,150	4,073	7,560	13,034	20,717	25,534	24,979	20,004	11,718	6,771	4
1988	133,284	4,054	7,528	13,267	20,831	24,616	23,851	18,730	11,113	6,443	9
1987	130,009	3,952	7,561	13,516	20,467	24,216	23,281	18,018	10,148	6,047	6
1986	125,522	3,781	7,551	13,610	20,073	23,167	22,353	17,027	9,495	5,707	11
1985	122,556	3,835	7,591	13,880	20,158	22,471	21,569	16,099	8,970	5,206	11
1984	118,730	3,914	7,696	14,070	19,284	21,742	20,666	15,210	8,484	4,998	11
1983	115,023	3,793	7,878	13,882	18,477	21,129	19,957	14,332	8,123	4,687	13
1982	111,393	3,924	8,057	13,527	18,410	20,611	18,813	13,504	7,361	4,541	16
1981	106,561	3,963	8,246	13,440	17,193	19,835	17,697	12,507	6,826	4,179	16
1980	103,844	4,019	8,318	13,123	16,875	19,151	17,198	12,092	6,497	3,942	6
1979	98,451	4,062	8,199	12,603	16,280	18,134	15,751	11,010	6,280	3,515	10
Malignant neoplasms of trachea, bronchus and lung (162)											
All races, male											
1998	91,447	2,362	4,540	7,453	10,743	14,945	18,191	15,524	9,934	6,084	2
1997	91,352	2,373	4,659	7,202	10,845	15,580	17,829	15,342	9,793	6,047	1
1996	91,620	2,525	4,726	7,331	11,112	16,104	17,973	14,768	9,780	5,570	5
1995	91,856	2,447	4,694	7,388	11,434	16,435	18,349	14,531	9,373	5,530	2
1994	91,893	2,400	4,760	7,523	12,194	16,872	17,827	14,208	9,204	5,205	6
1993	92,564	2,524	4,693	7,893	12,650	17,246	17,674	14,163	8,943	5,035	4
1992	91,405	2,577	4,740	7,811	12,779	17,285	17,098	14,016	8,659	4,719	4
1991	91,690	2,573	4,625	8,128	13,363	17,380	17,019	13,718	8,536	4,642	3
1990	91,091	2,573	4,733	8,458	13,736	17,295	16,736	13,561	7,892	4,317	2
1989	89,052	2,555	4,820	8,531	13,634	16,813	16,361	13,007	7,606	4,060	2
1988	88,059	2,466	4,871	8,747	13,856	16,526	15,963	12,649	7,359	3,869	7
1987	87,261	2,508	4,887	9,064	13,828	16,406	15,954	12,240	6,958	3,725	6
1986	85,057	2,397	4,928	9,131	13,762	15,863	15,294	11,903	6,570	3,526	7
1985	83,854	2,433	4,933	9,424	13,789	15,536	15,007	11,464	6,235	3,297	7
1984	82,491	2,471	5,099	9,622	13,433	15,219	14,825	10,995	5,981	3,184	6
1983	80,338	2,437	5,227	9,433	12,869	14,993	14,386	10,547	5,770	2,998	7
1982	79,228	2,559	5,474	9,329	13,054	14,908	13,893	10,119	5,310	2,957	11
1981	76,764	2,617	5,703	9,408	12,381	14,565	13,240	9,431	4,979	2,733	14
1980	75,535	2,670	5,696	9,361	12,323	14,227	13,062	9,184	4,829	2,509	5
1979	72,803	2,702	5,733	9,079	12,104	13,782	12,200	8,520	4,656	2,368	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
All races, female											
1998	63,114	1,603	2,995	4,699	6,892	9,479	11,962	10,765	7,730	5,669	-
1997	61,958	1,561	3,053	4,635	6,619	9,618	11,681	10,584	7,414	5,474	-
1996	60,395	1,621	3,040	4,461	6,726	9,774	11,425	9,984	7,123	4,997	3
1995	59,344	1,682	2,987	4,549	6,826	9,688	11,257	9,681	6,697	4,773	3
1994	57,589	1,619	2,909	4,564	6,783	9,938	10,839	9,273	6,288	4,254	-
1993	56,291	1,682	2,841	4,513	7,022	9,735	10,568	8,851	5,802	4,167	3
1992	54,538	1,622	2,798	4,497	7,147	9,544	10,211	8,440	5,409	3,661	2
1991	52,068	1,570	2,758	4,514	7,028	9,313	9,299	8,063	4,939	3,398	2
1990	50,194	1,606	2,823	4,514	7,207	9,035	8,973	7,459	4,454	3,011	2
1989	48,098	1,518	2,740	4,503	7,083	8,721	8,618	6,997	4,112	2,711	2
1988	45,225	1,588	2,657	4,520	6,975	8,090	7,888	6,081	3,754	2,574	2
1987	42,748	1,444	2,674	4,452	6,639	7,810	7,327	5,778	3,190	2,322	-
1986	40,465	1,384	2,623	4,479	6,311	7,304	7,059	5,124	2,925	2,181	4
1985	38,702	1,402	2,658	4,456	6,369	6,935	6,562	4,635	2,735	1,909	4
1984	36,239	1,443	2,597	4,448	5,851	6,523	5,841	4,215	2,503	1,814	5
1983	34,685	1,356	2,651	4,449	5,608	6,136	5,571	3,785	2,353	1,689	6
1982	32,165	1,365	2,583	4,198	5,356	5,703	4,920	3,385	2,051	1,584	5
1981	29,797	1,346	2,543	4,032	4,812	5,270	4,457	3,076	1,847	1,446	2
1980	28,309	1,349	2,622	3,762	4,552	4,924	4,136	2,908	1,668	1,433	1
1979	25,648	1,360	2,466	3,524	4,176	4,352	3,551	2,490	1,624	1,147	3
Malignant neoplasms of trachea, bronchus and lung (162)											
White, both sexes											
1998	136,026	3,087	6,162	10,228	15,150	21,488	26,869	23,791	16,204	10,715	1
1997	135,060	3,094	6,393	9,948	15,005	22,260	26,276	23,508	15,783	10,503	1
1996	134,055	3,288	6,496	9,962	15,318	22,801	26,329	22,416	15,515	9,637	4
1995	133,366	3,331	6,374	10,089	15,682	22,982	26,513	22,054	14,750	9,370	4
1994	131,871	3,230	6,337	10,194	16,460	23,682	25,657	21,362	14,157	8,603	3
1993	131,337	3,330	6,246	10,523	17,089	23,887	25,256	20,986	13,469	8,359	3
1992	128,719	3,428	6,200	10,315	17,289	23,701	24,511	20,491	12,901	7,646	6
1991	126,872	3,338	6,060	10,717	17,789	23,582	23,675	19,839	12,289	7,342	4
1990	124,650	3,393	6,256	10,908	18,173	23,305	23,182	19,154	11,352	6,691	2
1989	121,023	3,312	6,228	11,080	18,078	22,573	22,495	18,138	10,770	6,197	2
1988	117,826	3,327	6,242	11,352	18,238	21,771	21,561	17,033	10,180	5,850	9
1987	114,888	3,256	6,225	11,589	17,874	21,454	20,994	16,427	9,308	5,520	4
1986	111,049	3,066	6,252	11,722	17,514	20,591	20,227	15,546	8,734	5,209	10
1985	108,529	3,186	6,310	11,856	17,634	20,141	19,421	14,738	8,251	4,770	10
1984	105,027	3,213	6,449	12,048	16,756	19,440	18,756	13,870	7,781	4,589	8
1983	101,857	3,062	6,607	11,927	16,180	18,926	17,988	13,141	7,447	4,350	3
1982	98,780	3,238	6,795	11,608	16,120	18,482	17,033	12,408	6,745	4,195	12
1981	94,441	3,240	6,947	11,568	15,092	17,742	16,059	11,500	6,265	3,876	9
1980	92,182	3,280	7,045	11,338	14,895	17,134	15,628	11,085	5,992	3,653	4
1979	87,532	3,315	6,982	10,860	14,425	16,255	14,404	10,123	5,797	3,269	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
White, male											
1998	79,649	1,799	3,626	6,193	9,106	13,052	16,125	13,910	9,035	5,506	1
1997	79,624	1,825	3,785	5,918	9,224	13,663	15,766	13,796	8,907	5,477	1
1996	79,809	1,953	3,871	6,063	9,392	14,070	15,967	13,255	8,903	5,032	2
1995	80,088	1,931	3,803	6,139	9,699	14,357	16,301	13,089	8,532	4,983	1
1994	80,096	1,876	3,825	6,256	10,479	14,758	15,796	12,812	8,325	4,679	3
1993	80,620	1,941	3,793	6,608	10,854	15,119	15,597	12,775	8,079	4,546	2
1992	79,728	2,056	3,818	6,443	10,980	15,141	15,189	12,691	7,835	4,290	4
1991	80,028	2,007	3,723	6,752	11,579	15,162	15,177	12,397	7,707	4,231	3
1990	79,488	2,023	3,844	7,015	11,819	15,134	14,951	12,232	7,206	3,923	1
1989	77,659	2,032	3,878	7,117	11,773	14,690	14,576	11,680	6,938	3,710	1
1988	77,063	1,974	3,970	7,415	12,040	14,431	14,311	11,384	6,717	3,474	7
1987	76,398	2,018	3,972	7,718	11,964	14,369	14,229	11,055	6,353	3,385	4
1986	74,517	1,905	4,001	7,810	11,862	13,917	13,711	10,792	6,023	3,197	7
1985	73,519	1,976	4,008	7,947	11,950	13,747	13,396	10,414	5,697	3,027	7
1984	72,239	1,986	4,223	8,126	11,527	13,440	13,323	9,975	5,459	2,898	5
1983	70,598	1,913	4,333	8,027	11,134	13,294	12,878	9,620	5,288	2,799	-
1982	69,686	2,063	4,539	7,908	11,309	13,244	12,444	9,260	4,876	2,733	7
1981	67,525	2,103	4,768	8,019	10,746	12,894	11,924	8,643	4,552	2,531	7
1980	66,651	2,160	4,771	8,020	10,791	12,618	11,823	8,382	4,452	2,322	3
1979	64,338	2,173	4,839	7,720	10,626	12,241	11,116	7,821	4,300	2,195	5
Malignant neoplasms of trachea, bronchus and lung (162)											
White, female											
1998	56,377	1,288	2,536	4,035	6,044	8,436	10,744	9,881	7,169	5,209	-
1997	55,436	1,269	2,608	4,030	5,781	8,597	10,510	9,712	6,876	5,026	-
1996	54,246	1,335	2,625	3,899	5,926	8,731	10,362	9,161	6,612	4,605	2
1995	53,278	1,400	2,571	3,950	5,983	8,625	10,212	8,965	6,218	4,387	3
1994	51,775	1,354	2,512	3,938	5,981	8,924	9,861	8,550	5,832	3,924	-
1993	50,717	1,389	2,453	3,915	6,235	8,768	9,659	8,211	5,390	3,813	1
1992	48,991	1,372	2,382	3,872	6,309	8,560	9,322	7,800	5,066	3,356	2
1991	46,844	1,331	2,337	3,965	6,210	8,420	8,498	7,442	4,582	3,111	1
1990	45,162	1,370	2,412	3,893	6,354	8,171	8,231	6,922	4,146	2,768	1
1989	43,364	1,280	2,350	3,963	6,305	7,883	7,919	6,458	3,832	2,487	1
1988	40,763	1,353	2,272	3,937	6,198	7,340	7,250	5,649	3,463	2,376	2
1987	38,490	1,238	2,253	3,871	5,910	7,085	6,765	5,372	2,955	2,135	-
1986	36,532	1,161	2,251	3,912	5,652	6,674	6,516	4,754	2,711	2,012	3
1985	35,010	1,210	2,302	3,909	5,684	6,394	6,025	4,324	2,554	1,743	3
1984	32,788	1,227	2,226	3,922	5,229	6,000	5,433	3,895	2,322	1,691	3
1983	31,259	1,149	2,274	3,900	5,046	5,632	5,110	3,521	2,159	1,551	3
1982	29,094	1,175	2,256	3,700	4,811	5,238	4,589	3,148	1,869	1,462	5
1981	26,916	1,137	2,179	3,549	4,346	4,848	4,135	2,857	1,713	1,345	2
1980	25,531	1,120	2,274	3,318	4,104	4,516	3,805	2,703	1,540	1,331	1
1979	23,194	1,142	2,143	3,140	3,799	4,014	3,288	2,302	1,497	1,074	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
All other, both sexes											
1998	18,535	878	1,373	1,924	2,485	2,936	3,284	2,498	1,460	1,038	1
1997	18,250	840	1,319	1,889	2,459	2,938	3,234	2,418	1,424	1,018	-
1996	17,960	858	1,270	1,830	2,520	3,077	3,069	2,336	1,388	930	4
1995	17,834	798	1,307	1,848	2,578	3,141	3,093	2,158	1,320	933	1
1994	17,611	789	1,332	1,893	2,517	3,128	3,009	2,119	1,335	856	3
1993	17,518	876	1,288	1,883	2,583	3,094	2,986	2,028	1,276	843	4
1992	17,224	771	1,338	1,993	2,637	3,128	2,798	1,965	1,167	734	-
1991	16,886	805	1,323	1,925	2,602	3,111	2,643	1,942	1,186	698	1
1990	16,635	786	1,300	2,064	2,770	3,025	2,527	1,866	994	637	2
1989	16,127	761	1,332	1,954	2,639	2,961	2,484	1,866	948	574	2
1988	15,458	727	1,286	1,915	2,593	2,845	2,290	1,697	933	593	-
1987	15,121	696	1,336	1,927	2,593	2,762	2,287	1,591	840	527	2
1986	14,473	715	1,299	1,888	2,559	2,576	2,126	1,481	761	498	1
1985	14,027	649	1,281	2,024	2,524	2,330	2,148	1,361	719	436	1
1984	13,703	701	1,247	2,022	2,528	2,302	1,910	1,340	703	409	3
1983	13,166	731	1,271	1,955	2,297	2,203	1,969	1,191	676	337	10
1982	12,613	686	1,262	1,919	2,290	2,129	1,780	1,096	616	346	4
1981	12,120	723	1,299	1,872	2,101	2,093	1,638	1,007	561	303	7
1980	11,662	739	1,273	1,785	1,980	2,017	1,570	1,007	505	289	2
1979	10,919	747	1,217	1,743	1,855	1,879	1,347	887	483	246	3
Malignant neoplasms of trachea, bronchus and lung (162)											
All other, male											
1998	11,798	563	914	1,260	1,637	1,893	2,066	1,614	899	578	1
1997	11,728	548	874	1,284	1,621	1,917	2,063	1,546	886	570	-
1996	11,811	572	855	1,268	1,720	2,034	2,006	1,513	877	538	3
1995	11,768	516	891	1,249	1,735	2,078	2,048	1,442	841	547	1
1994	11,797	524	935	1,267	1,715	2,114	2,031	1,396	879	526	3
1993	11,944	583	900	1,285	1,796	2,127	2,077	1,388	864	489	2
1992	11,677	521	922	1,368	1,799	2,144	1,909	1,325	824	429	-
1991	11,662	566	902	1,376	1,784	2,218	1,842	1,321	829	411	-
1990	11,603	550	889	1,443	1,917	2,161	1,785	1,329	686	394	1
1989	11,393	523	942	1,414	1,861	2,123	1,785	1,327	668	350	1
1988	10,996	492	901	1,332	1,816	2,095	1,652	1,265	642	395	-
1987	10,863	490	915	1,346	1,864	2,037	1,725	1,185	605	340	2
1986	10,540	492	927	1,321	1,900	1,946	1,583	1,111	547	329	-
1985	10,335	457	925	1,477	1,839	1,789	1,611	1,050	538	270	-
1984	10,252	485	876	1,496	1,906	1,779	1,502	1,020	522	286	1
1983	9,740	524	894	1,406	1,735	1,699	1,508	927	482	199	7
1982	9,542	496	935	1,421	1,745	1,664	1,449	859	434	224	4
1981	9,239	514	935	1,389	1,635	1,671	1,316	788	427	202	7
1980	8,884	510	925	1,341	1,532	1,609	1,239	802	377	187	2
1979	8,465	529	894	1,359	1,478	1,541	1,084	699	356	173	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
All other, female											
1998	6,737	315	459	664	848	1,043	1,218	884	561	460	-
1997	6,522	292	445	605	838	1,021	1,171	872	538	448	-
1996	6,149	286	415	562	800	1,043	1,063	823	511	392	1
1995	6,066	282	416	599	843	1,063	1,045	716	479	386	-
1994	5,814	265	397	626	802	1,014	978	723	456	330	-
1993	5,574	293	388	598	787	967	909	640	412	354	2
1992	5,547	250	416	625	838	984	889	640	343	305	-
1991	5,224	239	421	549	818	893	801	621	357	287	1
1990	5,032	236	411	621	853	864	742	537	308	243	1
1989	4,734	238	390	540	778	838	699	539	280	224	1
1988	4,462	235	385	583	777	750	638	432	291	198	-
1987	4,258	206	421	581	729	725	562	406	235	187	-
1986	3,933	223	372	567	659	630	543	370	214	169	1
1985	3,692	192	356	547	685	541	537	311	181	166	1
1984	3,451	216	371	526	622	523	408	320	181	123	2
1983	3,426	207	377	549	562	504	461	264	194	138	3
1982	3,071	190	327	498	545	465	331	237	182	122	-
1981	2,881	209	364	483	466	422	322	219	134	101	-
1980	2,778	229	348	444	448	408	331	205	128	102	-
1979	2,454	218	323	384	377	338	263	188	127	73	1
Malignant neoplasms of trachea, bronchus and lung (162)											
Black, both sexes											
1998	16,105	785	1,235	1,723	2,199	2,536	2,830	2,139	1,217	861	1
1997	16,014	750	1,195	1,695	2,173	2,587	2,838	2,107	1,197	855	-
1996	15,835	770	1,164	1,649	2,253	2,744	2,675	2,013	1,162	783	4
1995	15,790	718	1,183	1,685	2,326	2,812	2,722	1,862	1,125	773	1
1994	15,670	707	1,205	1,721	2,271	2,781	2,673	1,827	1,172	732	3
1993	15,641	793	1,184	1,693	2,329	2,778	2,663	1,774	1,106	719	4
1992	15,472	702	1,237	1,819	2,391	2,814	2,514	1,724	1,010	628	-
1991	15,201	739	1,217	1,761	2,380	2,808	2,365	1,716	1,036	600	1
1990	15,144	728	1,206	1,914	2,573	2,755	2,290	1,644	877	542	2
1989	14,703	693	1,228	1,804	2,425	2,739	2,252	1,670	839	510	2
1988	14,131	660	1,185	1,769	2,401	2,642	2,084	1,537	820	511	-
1987	13,886	646	1,230	1,798	2,419	2,548	2,096	1,433	737	458	2
1986	13,354	674	1,213	1,772	2,375	2,420	1,941	1,336	674	434	1
1985	13,009	601	1,217	1,915	2,363	2,176	1,962	1,228	638	389	1
1984	12,708	662	1,160	1,911	2,383	2,146	1,756	1,210	621	367	3
1983	12,282	685	1,202	1,849	2,172	2,067	1,831	1,092	592	304	9
1982	11,857	654	1,197	1,840	2,165	2,008	1,668	1,011	555	297	4
1981	11,372	677	1,246	1,789	1,998	1,975	1,521	920	494	268	4
1980	10,957	704	1,212	1,712	1,876	1,909	1,452	925	439	256	2
1979	10,247	718	1,155	1,663	1,753	1,780	1,251	811	418	220	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of trachea, bronchus and lung (162)											
Black, male											
1998	10,288	507	830	1,141	1,456	1,628	1,781	1,385	743	479	1
1997	10,336	494	793	1,148	1,444	1,697	1,808	1,350	756	474	-
1996	10,463	523	795	1,143	1,546	1,811	1,743	1,315	740	454	3
1995	10,537	464	820	1,152	1,579	1,876	1,816	1,259	735	454	1
1994	10,555	479	852	1,165	1,558	1,885	1,797	1,216	776	446	3
1993	10,733	533	829	1,167	1,630	1,930	1,863	1,215	744	421	2
1992	10,530	478	853	1,262	1,632	1,941	1,719	1,160	712	369	-
1991	10,545	528	837	1,263	1,641	2,000	1,653	1,165	729	352	-
1990	10,632	514	836	1,352	1,797	1,962	1,630	1,178	610	331	1
1989	10,457	480	876	1,315	1,702	1,971	1,642	1,205	594	314	1
1988	10,112	452	838	1,234	1,686	1,959	1,512	1,143	568	350	-
1987	10,065	458	848	1,270	1,749	1,899	1,595	1,079	534	297	2
1986	9,773	465	867	1,252	1,774	1,836	1,446	1,015	483	290	-
1985	9,615	429	881	1,400	1,718	1,681	1,481	953	477	237	-
1984	9,569	462	824	1,423	1,811	1,674	1,383	919	467	255	1
1983	9,117	495	849	1,338	1,634	1,609	1,403	845	425	182	7
1982	9,003	476	893	1,366	1,654	1,567	1,363	792	395	200	4
1981	8,697	484	894	1,327	1,559	1,581	1,233	729	379	176	4
1980	8,382	486	884	1,286	1,456	1,530	1,148	743	335	165	2
1979	7,991	511	846	1,298	1,404	1,478	1,011	639	315	152	2
Malignant neoplasms of trachea, bronchus and lung (162)											
Black, female											
1998	5,817	278	405	582	743	908	1,049	754	474	382	-
1997	5,678	256	402	547	729	890	1,030	757	441	381	-
1996	5,372	247	369	506	707	933	932	698	422	329	1
1995	5,253	254	363	533	747	936	906	603	390	319	-
1994	5,115	228	353	556	713	896	876	611	396	286	-
1993	4,908	260	355	526	699	848	800	559	362	298	2
1992	4,942	224	384	557	759	873	795	564	298	259	-
1991	4,656	211	380	498	739	808	712	551	307	248	1
1990	4,512	214	370	562	776	793	660	466	267	211	1
1989	4,246	213	352	489	723	768	610	465	245	196	1
1988	4,019	208	347	535	715	683	572	394	252	161	-
1987	3,821	188	382	528	670	649	501	354	203	161	-
1986	3,581	209	346	520	601	584	495	321	191	144	1
1985	3,394	172	336	515	645	495	481	275	161	152	1
1984	3,139	200	336	488	572	472	373	291	154	112	2
1983	3,165	190	353	511	538	458	428	247	167	122	2
1982	2,854	178	304	474	511	441	305	219	160	97	-
1981	2,675	193	352	462	439	394	288	191	115	92	-
1980	2,575	218	328	426	420	379	304	182	104	91	-
1979	2,256	207	309	365	349	302	240	172	103	68	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
All races, both sexes											
1998	7,431	505	586	586	668	824	936	906	739	733	-
1997	7,238	517	570	580	643	761	919	904	703	691	-
1996	7,279	533	553	561	596	802	960	820	721	691	-
1995	6,907	509	495	556	659	778	843	796	645	603	2
1994	6,680	477	529	486	623	747	887	774	610	553	-
1993	6,712	471	469	553	656	812	831	742	567	529	2
1992	6,568	450	469	512	701	792	757	751	533	502	-
1991	6,451	450	414	527	685	768	794	692	514	496	-
1990	6,420	436	438	545	695	751	784	666	488	494	1
1989	6,161	427	491	537	660	760	640	623	455	468	-
1988	5,979	422	453	499	667	693	695	634	443	408	2
1987	5,913	395	413	576	668	717	667	606	457	337	1
1986	5,673	368	444	537	621	612	654	530	426	365	-
1985	5,529	374	434	560	628	583	611	504	375	341	-
1984	5,378	335	453	551	548	585	614	477	363	328	1
1983	5,216	345	404	573	571	590	547	430	333	307	1
1982	5,106	338	437	553	592	558	563	438	299	277	-
1981	4,932	343	470	496	545	563	503	402	280	287	1
1980	4,679	297	449	517	510	520	456	375	310	264	-
1979	4,749	380	469	512	538	498	484	382	269	246	1
Malignant melanoma of skin (172)											
All races, male											
1998	4,747	327	389	394	449	575	622	574	458	362	-
1997	4,488	324	347	401	427	504	594	569	412	328	-
1996	4,490	318	368	374	410	536	598	511	422	344	-
1995	4,297	344	326	366	459	512	551	491	358	274	2
1994	4,117	303	351	329	413	486	566	482	341	258	-
1993	4,128	302	304	359	434	536	534	450	307	241	1
1992	4,045	295	293	348	481	517	474	471	269	223	-
1991	4,017	287	276	367	464	504	523	420	275	215	-
1990	3,844	267	270	349	464	468	472	401	245	214	1
1989	3,736	264	321	327	441	503	422	371	229	189	-
1988	3,585	250	273	329	442	446	425	373	229	188	2
1987	3,635	243	270	380	438	465	415	361	237	154	1
1986	3,378	232	283	353	395	383	383	289	220	165	-
1985	3,296	220	280	370	399	349	391	281	180	135	-
1984	3,207	198	289	362	353	383	365	281	174	126	-
1983	3,073	199	238	358	353	388	319	251	142	125	-
1982	3,036	196	278	363	356	343	342	238	153	111	-
1981	2,811	210	279	317	314	353	275	197	120	120	1
1980	2,750	175	290	327	317	317	273	198	137	118	-
1979	2,752	231	291	324	326	296	275	193	131	105	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
All races, female											
1998	2,684	178	197	192	219	249	314	332	281	371	-
1997	2,750	193	223	179	216	257	325	335	291	363	-
1996	2,789	215	185	187	186	266	362	309	299	347	-
1995	2,610	165	169	190	200	266	292	305	287	329	-
1994	2,563	174	178	157	210	261	321	292	269	295	-
1993	2,584	169	165	194	222	276	297	292	260	288	1
1992	2,523	155	176	164	220	275	283	280	264	279	-
1991	2,434	163	138	160	221	264	271	272	239	281	-
1990	2,576	169	168	196	231	283	312	265	243	280	-
1989	2,425	163	170	210	219	257	218	252	226	279	-
1988	2,394	172	180	170	225	247	270	261	214	220	-
1987	2,278	152	143	196	230	252	252	245	220	183	-
1986	2,295	136	161	184	226	229	271	241	206	200	-
1985	2,233	154	154	190	229	234	220	223	195	206	-
1984	2,171	137	164	189	195	202	249	196	189	202	1
1983	2,143	146	166	215	218	202	228	179	191	182	1
1982	2,070	142	159	190	236	215	221	200	146	166	-
1981	2,121	133	191	179	231	210	228	205	160	167	-
1980	1,929	122	159	190	193	203	183	177	173	146	-
1979	1,997	149	178	188	212	202	209	189	138	141	-
Malignant melanoma of skin (172)											
White, both sexes											
1998	7,275	496	576	576	652	809	916	889	719	714	-
1997	7,064	508	561	563	626	742	901	883	684	670	-
1996	7,141	528	541	550	585	788	942	804	704	677	-
1995	6,774	501	490	544	648	760	831	776	632	588	1
1994	6,556	472	525	475	612	735	871	756	597	537	-
1993	6,572	463	461	548	648	795	813	713	556	516	2
1992	6,423	445	460	504	687	770	732	730	523	490	-
1991	6,308	446	410	516	668	748	778	670	493	480	-
1990	6,290	432	430	536	680	732	769	650	476	481	-
1989	6,027	422	485	522	651	746	620	609	439	453	-
1988	5,877	420	447	493	654	677	678	619	437	397	2
1987	5,786	389	406	571	654	703	642	594	438	327	1
1986	5,565	366	435	529	608	592	646	511	415	360	-
1985	5,410	366	421	555	614	567	599	483	366	337	-
1984	5,264	334	446	542	535	570	598	463	354	317	-
1983	5,115	340	395	563	563	572	536	416	327	300	1
1982	5,000	333	425	543	583	546	542	428	293	271	-
1981	4,826	340	458	489	530	547	494	397	267	273	-
1980	4,572	293	443	505	493	499	443	369	302	258	-
1979	4,657	375	465	502	523	485	467	377	260	237	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
White, male											
1998	4,685	324	387	389	445	568	611	569	451	358	-
1997	4,421	320	343	396	418	499	587	561	406	323	-
1996	4,423	318	359	368	404	528	592	506	411	339	-
1995	4,237	339	325	363	452	504	544	480	352	271	1
1994	4,063	299	348	326	404	477	558	475	340	254	-
1993	4,078	300	300	357	430	529	527	438	306	239	1
1992	3,972	291	289	343	470	507	465	459	263	220	-
1991	3,951	284	274	361	452	500	514	410	267	210	-
1990	3,778	264	265	345	458	460	464	393	239	207	-
1989	3,676	260	317	323	437	496	413	364	223	185	-
1988	3,535	249	270	325	437	439	418	362	228	184	2
1987	3,578	238	268	379	431	456	403	358	229	150	1
1986	3,315	230	277	347	386	370	379	279	215	162	-
1985	3,238	216	273	369	390	343	388	269	175	134	-
1984	3,161	198	287	358	345	378	357	276	173	123	-
1983	3,019	197	232	353	350	379	312	245	138	123	-
1982	2,988	193	271	358	353	337	334	233	148	109	-
1981	2,762	208	274	312	307	343	271	195	116	117	-
1980	2,706	173	287	322	310	309	268	194	134	113	-
1979	2,706	229	289	319	321	287	267	191	126	100	1
Malignant melanoma of skin (172)											
White, female											
1998	2,590	172	189	187	207	241	305	320	268	356	-
1997	2,643	188	218	167	208	243	314	322	278	347	-
1996	2,718	210	182	182	181	260	350	298	293	338	-
1995	2,537	162	165	181	196	256	287	296	280	317	-
1994	2,493	173	177	149	208	258	313	281	257	283	-
1993	2,494	163	161	191	218	266	286	275	250	277	1
1992	2,451	154	171	161	217	263	267	271	260	270	-
1991	2,357	162	136	155	216	248	264	260	226	270	-
1990	2,512	168	165	191	222	272	305	257	237	274	-
1989	2,351	162	168	199	214	250	207	245	216	268	-
1988	2,342	171	177	168	217	238	260	257	209	213	-
1987	2,208	151	138	192	223	247	239	236	209	177	-
1986	2,250	136	158	182	222	222	267	232	200	198	-
1985	2,172	150	148	186	224	224	211	214	191	203	-
1984	2,103	136	159	184	190	192	241	187	181	194	-
1983	2,096	143	163	210	213	193	224	171	189	177	1
1982	2,012	140	154	185	230	209	208	195	145	162	-
1981	2,064	132	184	177	223	204	223	202	151	156	-
1980	1,866	120	156	183	183	190	175	175	168	145	-
1979	1,951	146	176	183	202	198	200	186	134	137	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
All other, both sexes											
1998	156	9	10	10	16	15	20	17	20	19	-
1997	174	9	9	17	17	19	18	21	19	21	-
1996	138	5	12	11	11	14	18	16	17	14	-
1995	133	8	5	12	11	18	12	20	13	15	1
1994	124	5	4	11	11	12	16	18	13	16	-
1993	140	8	8	5	8	17	18	29	11	13	-
1992	145	5	9	8	14	22	25	21	10	12	-
1991	143	4	4	11	17	20	16	22	21	16	-
1990	130	4	8	9	15	19	15	16	12	13	1
1989	134	5	6	15	9	14	20	14	16	15	-
1988	102	2	6	6	13	16	17	15	6	11	-
1987	127	6	7	5	14	14	25	12	19	10	-
1986	108	2	9	8	13	20	8	19	11	5	-
1985	119	8	13	5	14	16	12	21	9	4	-
1984	114	1	7	9	13	15	16	14	9	11	1
1983	101	5	9	10	8	18	11	14	6	7	-
1982	106	5	12	10	9	12	21	10	6	6	-
1981	106	3	12	7	15	16	9	5	13	14	1
1980	107	4	6	12	17	21	13	6	8	6	-
1979	92	5	4	10	15	13	17	5	9	9	-
Malignant melanoma of skin (172)											
All other, male											
1998	62	3	2	5	4	7	11	5	7	4	-
1997	67	4	4	5	9	5	7	8	6	5	-
1996	67	-	9	6	6	8	6	5	11	5	-
1995	60	5	1	3	7	8	7	11	6	3	1
1994	54	4	3	3	9	9	8	7	1	4	-
1993	50	2	4	2	4	7	7	12	1	2	-
1992	73	4	4	5	11	10	9	12	6	3	-
1991	66	3	2	6	12	4	9	10	8	5	-
1990	66	3	5	4	6	8	8	8	6	7	1
1989	60	4	4	4	4	7	9	7	6	4	-
1988	50	1	3	4	5	7	7	11	1	4	-
1987	57	5	2	1	7	9	12	3	8	4	-
1986	63	2	6	6	9	13	4	10	5	3	-
1985	58	4	7	1	9	6	3	12	5	1	-
1984	46	-	2	4	8	5	8	5	1	3	-
1983	54	2	6	5	3	9	7	6	4	2	-
1982	48	3	7	5	3	6	8	5	5	2	-
1981	49	2	5	5	7	10	4	2	4	3	1
1980	44	2	3	5	7	8	5	4	3	5	-
1979	46	2	2	5	5	9	8	2	5	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
All other, female											
1998	94	6	8	5	12	8	9	12	13	15	-
1997	107	5	5	12	8	14	11	13	13	16	-
1996	71	5	3	5	5	6	12	11	6	9	-
1995	73	3	4	9	4	10	5	9	7	12	-
1994	70	1	1	8	2	3	8	11	12	12	-
1993	90	6	4	3	4	10	11	17	10	11	-
1992	72	1	5	3	3	12	16	9	4	9	-
1991	77	1	2	5	5	16	7	12	13	11	-
1990	64	1	3	5	9	11	7	8	6	6	-
1989	74	1	2	11	5	7	11	7	10	11	-
1988	52	1	3	2	8	9	10	4	5	7	-
1987	70	1	5	4	7	5	13	9	11	6	-
1986	45	-	3	2	4	7	4	9	6	2	-
1985	61	4	6	4	5	10	9	9	4	3	-
1984	68	1	5	5	5	10	8	9	8	8	1
1983	47	3	3	5	5	9	4	8	2	5	-
1982	58	2	5	5	6	6	13	5	1	4	-
1981	57	1	7	2	8	6	5	3	9	11	-
1980	63	2	3	7	10	13	8	2	5	1	-
1979	46	3	2	5	10	4	9	3	4	4	-
Malignant melanoma of skin (172)											
Black, both sexes											
1998	121	8	6	6	13	10	14	12	18	18	-
1997	131	7	6	11	13	12	14	15	16	19	-
1996	100	3	7	7	8	11	12	12	13	13	-
1995	103	6	5	6	5	15	8	19	12	14	1
1994	98	5	4	9	9	9	12	11	11	15	-
1993	115	5	4	4	8	15	14	22	11	12	-
1992	110	3	4	6	11	17	24	16	6	9	-
1991	119	4	2	8	16	17	14	21	18	12	-
1990	106	3	6	9	13	17	11	15	9	13	1
1989	109	3	6	13	6	11	16	11	13	14	-
1988	88	2	5	4	11	15	14	12	5	11	-
1987	108	5	7	4	12	11	22	10	17	7	-
1986	91	2	8	7	11	18	8	14	8	4	-
1985	100	7	10	5	13	12	9	19	7	4	-
1984	94	1	6	5	10	12	14	14	7	11	1
1983	91	5	7	10	8	14	10	14	6	6	-
1982	94	5	11	9	5	11	21	8	6	5	-
1981	97	3	10	6	13	16	8	5	13	13	-
1980	98	4	5	10	17	20	12	6	8	4	-
1979	88	5	4	10	15	10	16	5	9	9	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant melanoma of skin (172)											
Black, male											
1998	44	2	1	3	3	4	7	2	6	4	-
1997	48	2	3	3	7	4	6	6	4	3	-
1996	47	-	5	4	6	6	3	3	7	4	-
1995	46	4	1	2	4	6	4	11	5	2	1
1994	41	4	3	2	8	7	4	3	1	4	-
1993	38	1	1	2	4	5	5	8	1	2	-
1992	53	2	2	4	8	8	8	10	3	-	-
1991	52	3	1	4	12	2	8	9	6	3	-
1990	51	2	3	4	6	8	5	7	3	7	1
1989	46	3	4	3	3	6	7	4	4	4	-
1988	41	1	2	3	4	7	5	8	1	4	-
1987	46	5	2	1	5	7	9	2	7	2	-
1986	56	2	5	6	8	12	4	7	4	3	-
1985	46	3	6	1	8	4	2	10	3	1	-
1984	36	-	2	3	5	4	6	5	1	3	-
1983	49	2	4	5	3	8	6	6	4	2	-
1982	40	3	6	4	2	5	8	4	5	1	-
1981	42	2	4	4	6	10	3	2	4	2	-
1980	39	2	3	4	7	8	4	4	3	3	-
1979	42	2	2	5	5	6	7	2	5	5	-
Malignant melanoma of skin (172)											
Black, female											
1998	77	6	5	3	10	6	7	10	12	14	-
1997	83	5	3	8	6	8	8	9	12	16	-
1996	53	3	2	3	2	5	9	9	6	9	-
1995	57	2	4	4	1	9	4	8	7	12	-
1994	57	1	1	7	1	2	8	8	10	11	-
1993	77	4	3	2	4	10	9	14	10	10	-
1992	57	1	2	2	3	9	16	6	3	9	-
1991	67	1	1	4	4	15	6	12	12	9	-
1990	55	1	3	5	7	9	6	8	6	6	-
1989	63	-	2	10	3	5	9	7	9	10	-
1988	47	1	3	1	7	8	9	4	4	7	-
1987	62	-	5	3	7	4	13	8	10	5	-
1986	35	-	3	1	3	6	4	7	4	1	-
1985	54	4	4	4	5	8	7	9	4	3	-
1984	58	1	4	2	5	8	8	9	6	8	1
1983	42	3	3	5	5	6	4	8	2	4	-
1982	54	2	5	5	3	6	13	4	1	4	-
1981	55	1	6	2	7	6	5	3	9	11	-
1980	59	2	2	6	10	12	8	2	5	1	-
1979	46	3	2	5	10	4	9	3	4	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
All races, both sexes											
1998	42,086	2,779	3,596	3,661	3,773	4,300	5,239	5,146	4,425	5,628	1
1997	42,297	2,931	3,600	3,641	3,797	4,534	5,197	5,057	4,396	5,515	-
1996	43,448	3,070	3,377	3,651	3,953	4,814	5,540	5,281	4,530	5,561	1
1995	44,209	3,063	3,558	3,651	4,161	5,114	5,745	5,267	4,512	5,362	1
1994	44,008	2,932	3,455	3,538	4,233	5,335	5,787	5,337	4,553	5,059	-
1993	43,910	2,831	3,358	3,617	4,390	5,445	5,650	5,444	4,346	5,129	1
1992	43,365	2,804	3,221	3,562	4,588	5,623	5,763	5,117	4,186	4,628	5
1991	43,849	2,697	3,154	3,841	4,959	5,722	5,514	5,188	4,201	4,546	1
1990	43,663	2,630	3,225	3,877	4,957	5,886	5,524	5,212	4,016	4,310	2
1989	43,138	2,573	3,161	3,973	5,047	5,760	5,536	5,152	3,883	4,107	1
1988	42,461	2,486	3,166	4,185	5,299	5,552	5,426	4,715	3,863	3,939	3
1987	41,211	2,374	3,079	4,118	5,366	5,463	5,290	4,581	3,516	3,666	3
1986	40,789	2,285	3,065	4,213	5,364	5,485	5,237	4,412	3,375	3,606	5
1985	40,383	2,322	3,140	4,497	5,461	5,269	5,074	4,394	3,305	3,432	3
1984	39,722	2,136	3,186	4,521	5,243	5,277	4,960	4,266	3,159	3,388	2
1983	38,247	2,082	3,223	4,545	5,190	5,053	4,807	3,980	2,984	3,205	2
1982	37,685	2,096	3,378	4,626	4,879	5,006	4,561	3,979	2,905	3,074	3
1981	36,737	2,130	3,493	4,555	4,749	4,992	4,485	3,684	2,758	2,867	4
1980	35,897	2,169	3,527	4,652	4,716	4,773	4,219	3,519	2,686	2,656	1
1979	34,622	2,214	3,580	4,392	4,469	4,623	4,062	3,266	2,615	2,509	5
Malignant neoplasm of breast (174-175)											
All races, male											
1998	349	14	22	26	27	37	38	64	57	47	-
1997	354	8	26	36	41	52	48	59	39	37	-
1996	357	12	19	28	34	40	61	63	40	45	-
1995	365	12	29	27	44	49	46	65	38	43	-
1994	364	16	19	24	44	54	66	44	40	38	-
1993	355	19	17	29	37	48	51	51	42	48	-
1992	297	10	16	21	28	47	45	48	32	40	-
1991	266	5	12	17	28	52	39	29	36	36	-
1990	272	16	17	23	39	43	43	36	24	20	-
1989	301	7	17	23	37	37	55	51	29	35	-
1988	289	13	20	13	29	40	47	47	34	33	-
1987	312	3	13	22	55	44	45	45	42	28	-
1986	250	6	16	21	30	47	35	37	24	24	-
1985	290	13	16	27	36	39	39	41	34	30	-
1984	250	12	12	25	26	40	38	35	25	23	-
1983	268	5	15	26	39	40	58	34	22	22	-
1982	271	5	10	26	30	40	45	42	41	22	-
1981	254	3	17	21	40	33	42	35	27	28	-
1980	256	9	17	25	49	39	33	41	18	18	-
1979	222	4	9	22	32	33	38	24	37	17	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
All races, female											
1998	41,737	2,765	3,574	3,635	3,746	4,263	5,201	5,082	4,368	5,581	1
1997	41,943	2,923	3,574	3,605	3,756	4,482	5,149	4,998	4,357	5,478	-
1996	43,091	3,058	3,358	3,623	3,919	4,774	5,479	5,218	4,490	5,516	1
1995	43,844	3,051	3,529	3,624	4,117	5,065	5,699	5,202	4,474	5,319	1
1994	43,644	2,916	3,436	3,514	4,189	5,281	5,721	5,293	4,513	5,021	-
1993	43,555	2,812	3,341	3,588	4,353	5,397	5,599	5,393	4,304	5,081	1
1992	43,068	2,794	3,205	3,541	4,560	5,576	5,718	5,069	4,154	4,588	5
1991	43,583	2,692	3,142	3,824	4,931	5,670	5,475	5,159	4,165	4,510	1
1990	43,391	2,614	3,208	3,854	4,918	5,843	5,481	5,176	3,992	4,290	2
1989	42,837	2,566	3,144	3,950	5,010	5,723	5,481	5,101	3,854	4,072	1
1988	42,172	2,473	3,146	4,172	5,270	5,512	5,379	4,668	3,829	3,906	3
1987	40,899	2,371	3,066	4,096	5,311	5,419	5,245	4,536	3,474	3,638	3
1986	40,539	2,279	3,049	4,192	5,334	5,438	5,202	4,375	3,351	3,582	5
1985	40,093	2,309	3,124	4,470	5,425	5,230	5,035	4,353	3,271	3,402	3
1984	39,472	2,124	3,174	4,496	5,217	5,237	4,922	4,231	3,134	3,365	2
1983	37,979	2,077	3,208	4,519	5,151	5,013	4,749	3,946	2,962	3,183	2
1982	37,414	2,091	3,368	4,600	4,849	4,966	4,516	3,937	2,864	3,052	3
1981	36,483	2,127	3,476	4,534	4,709	4,959	4,443	3,649	2,731	2,839	4
1980	35,641	2,160	3,510	4,627	4,667	4,734	4,186	3,478	2,668	2,638	1
1979	34,400	2,210	3,571	4,370	4,437	4,590	4,024	3,242	2,578	2,492	5
Malignant neoplasm of breast (174-175)											
White, both sexes											
1998	36,060	2,134	2,910	3,000	3,179	3,738	4,658	4,656	4,022	5,146	1
1997	36,194	2,280	2,966	3,006	3,166	3,903	4,570	4,546	3,993	5,046	-
1996	37,510	2,433	2,740	3,061	3,406	4,216	4,943	4,767	4,138	5,106	1
1995	38,236	2,413	2,894	3,051	3,526	4,511	5,115	4,793	4,113	4,946	1
1994	38,272	2,367	2,874	2,976	3,623	4,718	5,189	4,850	4,172	4,630	-
1993	38,341	2,326	2,768	3,065	3,804	4,836	5,086	4,939	3,985	4,729	-
1992	38,053	2,312	2,641	3,006	4,006	5,010	5,243	4,691	3,871	4,290	3
1991	38,478	2,212	2,628	3,285	4,317	5,141	4,955	4,741	3,869	4,202	-
1990	38,513	2,172	2,698	3,287	4,371	5,283	5,037	4,779	3,706	3,980	2
1989	38,253	2,101	2,673	3,429	4,501	5,170	5,059	4,736	3,618	3,810	1
1988	37,556	2,073	2,654	3,602	4,706	5,006	4,981	4,321	3,568	3,641	3
1987	36,569	1,943	2,598	3,582	4,771	4,986	4,812	4,240	3,261	3,409	2
1986	36,398	1,893	2,587	3,716	4,801	4,993	4,803	4,096	3,172	3,356	3
1985	36,146	1,948	2,719	3,939	4,903	4,808	4,642	4,067	3,083	3,202	3
1984	35,469	1,782	2,752	3,992	4,660	4,790	4,556	3,964	2,911	3,178	1
1983	34,324	1,746	2,778	4,044	4,683	4,600	4,412	3,681	2,783	3,005	2
1982	33,837	1,738	2,935	4,118	4,395	4,560	4,175	3,681	2,694	2,896	3
1981	33,124	1,794	3,102	4,074	4,304	4,569	4,132	3,439	2,550	2,681	2
1980	32,450	1,876	3,093	4,168	4,290	4,402	3,908	3,254	2,521	2,488	1
1979	31,364	1,870	3,160	3,989	4,062	4,245	3,775	3,028	2,472	2,361	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
White, male											
1998	301	11	19	20	23	33	31	57	52	40	-
1997	293	4	21	31	30	41	41	52	33	34	-
1996	292	10	14	24	25	31	53	53	34	38	-
1995	299	9	21	21	39	40	39	55	28	37	-
1994	312	13	18	23	36	43	55	36	38	33	-
1993	297	16	15	24	29	42	37	46	38	39	-
1992	256	8	11	17	25	42	37	43	27	38	-
1991	228	5	7	14	25	43	33	27	31	33	-
1990	227	11	14	17	32	36	36	35	20	18	-
1989	265	4	17	18	36	30	46	50	25	32	-
1988	229	9	12	11	27	33	44	37	27	21	-
1987	272	3	9	16	46	39	42	40	38	27	-
1986	215	6	13	20	25	39	30	30	22	21	-
1985	260	13	15	23	32	36	34	37	31	25	-
1984	213	8	12	23	20	32	31	30	23	22	-
1983	226	4	11	22	32	34	49	29	22	18	-
1982	236	5	10	21	25	35	42	38	34	17	-
1981	218	2	14	16	34	27	34	33	23	28	-
1980	227	7	16	20	45	37	28	37	16	16	-
1979	187	2	7	20	30	27	32	17	34	14	-
Malignant neoplasm of breast (174-175)											
White, female											
1998	35,759	2,123	2,891	2,980	3,156	3,705	4,627	4,599	3,970	5,106	1
1997	35,901	2,276	2,945	2,975	3,136	3,862	4,529	4,494	3,960	5,012	-
1996	37,218	2,423	2,726	3,037	3,381	4,185	4,890	4,714	4,104	5,068	1
1995	37,937	2,404	2,873	3,030	3,487	4,471	5,076	4,738	4,085	4,909	1
1994	37,960	2,354	2,856	2,953	3,587	4,675	5,134	4,814	4,134	4,597	-
1993	38,044	2,310	2,753	3,041	3,775	4,794	5,049	4,893	3,947	4,690	-
1992	37,797	2,304	2,630	2,989	3,981	4,968	5,206	4,648	3,844	4,252	3
1991	38,250	2,207	2,621	3,271	4,292	5,098	4,922	4,714	3,838	4,169	-
1990	38,286	2,161	2,684	3,270	4,339	5,247	5,001	4,744	3,686	3,962	2
1989	37,988	2,097	2,656	3,411	4,465	5,140	5,013	4,686	3,593	3,778	1
1988	37,327	2,064	2,642	3,591	4,679	4,973	4,937	4,284	3,541	3,620	3
1987	36,297	1,940	2,589	3,566	4,725	4,947	4,770	4,200	3,223	3,382	2
1986	36,183	1,887	2,574	3,696	4,776	4,954	4,773	4,066	3,150	3,335	3
1985	35,886	1,935	2,704	3,916	4,871	4,772	4,608	4,030	3,052	3,177	3
1984	35,256	1,774	2,740	3,969	4,640	4,758	4,525	3,934	2,888	3,156	1
1983	34,098	1,742	2,767	4,022	4,651	4,566	4,363	3,652	2,761	2,987	2
1982	33,601	1,733	2,925	4,097	4,370	4,525	4,133	3,643	2,660	2,879	3
1981	32,906	1,792	3,088	4,058	4,270	4,542	4,098	3,406	2,527	2,653	2
1980	32,223	1,869	3,077	4,148	4,245	4,365	3,880	3,217	2,505	2,472	1
1979	31,177	1,868	3,153	3,969	4,032	4,218	3,743	3,011	2,438	2,347	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
All other, both sexes											
1998	6,026	645	686	661	594	562	581	490	403	482	-
1997	6,103	651	634	635	631	631	627	511	403	469	-
1996	5,938	637	637	590	547	598	597	514	392	455	-
1995	5,973	650	664	600	635	603	630	474	399	416	-
1994	5,736	565	581	562	610	617	598	487	381	429	-
1993	5,569	505	590	552	586	609	564	505	361	400	1
1992	5,312	492	580	556	582	613	520	426	315	338	2
1991	5,371	485	526	556	642	581	559	447	332	344	1
1990	5,150	458	527	590	586	603	487	433	310	330	-
1989	4,885	472	488	544	546	590	477	416	265	297	-
1988	4,905	413	512	583	593	546	445	394	295	298	-
1987	4,642	431	481	536	595	477	478	341	255	257	1
1986	4,391	392	478	497	563	492	434	316	203	250	2
1985	4,237	374	421	558	558	461	432	327	222	230	-
1984	4,253	354	434	529	583	487	404	302	248	210	1
1983	3,923	336	445	501	507	453	395	299	201	200	-
1982	3,848	358	443	508	484	446	386	298	211	178	-
1981	3,613	336	391	481	445	423	353	245	208	186	2
1980	3,447	293	434	484	426	371	311	265	165	168	-
1979	3,258	344	420	403	407	378	287	238	143	148	-
Malignant neoplasm of breast (174-175)											
All other, male											
1998	48	3	3	6	4	4	7	7	5	7	-
1997	61	4	5	5	11	11	7	7	6	3	-
1996	65	2	5	4	9	9	8	10	6	7	-
1995	66	3	8	6	5	9	7	10	10	6	-
1994	52	3	1	1	8	11	11	8	2	5	-
1993	58	3	2	5	8	6	14	5	4	9	-
1992	41	2	5	4	3	5	8	5	5	2	-
1991	38	-	5	3	3	9	6	2	5	3	-
1990	45	5	3	6	7	7	7	1	4	2	-
1989	36	3	-	5	1	7	9	1	4	3	-
1988	60	4	8	2	2	7	3	10	7	12	-
1987	40	-	4	6	9	5	3	5	4	1	-
1986	35	-	3	1	5	8	5	7	2	3	-
1985	30	-	1	4	4	3	5	4	3	5	-
1984	37	4	-	2	6	8	7	5	2	1	-
1983	42	1	4	4	7	6	9	5	-	4	-
1982	35	-	-	5	5	5	3	4	7	5	-
1981	36	1	3	5	6	6	8	2	4	-	-
1980	29	2	1	5	4	2	5	4	2	2	-
1979	35	2	2	2	2	6	6	7	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
All other, female											
1998	5,978	642	683	655	590	558	574	483	398	475	-
1997	6,042	647	629	630	620	620	620	504	397	466	-
1996	5,873	635	632	586	538	589	589	504	386	448	-
1995	5,907	647	656	594	630	594	623	464	389	410	-
1994	5,684	562	580	561	602	606	587	479	379	424	-
1993	5,511	502	588	547	578	603	550	500	357	391	1
1992	5,271	490	575	552	579	608	512	421	310	336	2
1991	5,333	485	521	553	639	572	553	445	327	341	1
1990	5,105	453	524	584	579	596	480	432	306	328	-
1989	4,849	469	488	539	545	583	468	415	261	294	-
1988	4,845	409	504	581	591	539	442	384	288	286	-
1987	4,602	431	477	530	586	472	475	336	251	256	1
1986	4,356	392	475	496	558	484	429	309	201	247	2
1985	4,207	374	420	554	554	458	427	323	219	225	-
1984	4,216	350	434	527	577	479	397	297	246	209	1
1983	3,881	335	441	497	500	447	386	294	201	196	-
1982	3,813	358	443	503	479	441	383	294	204	173	-
1981	3,577	335	388	476	439	417	345	243	204	186	2
1980	3,418	291	433	479	422	369	306	261	163	166	-
1979	3,223	342	418	401	405	372	281	231	140	145	-
Malignant neoplasm of breast (174-175)											
Black, both sexes											
1998	5,329	550	596	546	524	494	531	445	376	454	-
1997	5,483	560	552	555	563	561	584	475	376	440	-
1996	5,316	561	546	529	495	526	536	470	363	427	-
1995	5,328	553	566	522	556	538	584	443	374	393	-
1994	5,134	495	490	497	547	549	557	443	352	400	-
1993	5,052	452	520	487	529	555	524	468	335	386	1
1992	4,816	427	514	491	523	561	479	393	299	317	2
1991	4,843	410	462	494	575	534	519	408	321	318	1
1990	4,703	396	468	528	540	551	455	404	296	317	-
1989	4,438	424	436	471	500	535	444	381	252	285	-
1988	4,525	370	467	522	540	508	410	366	281	291	-
1987	4,291	382	449	486	556	441	446	316	247	248	1
1986	4,076	361	437	454	518	464	410	298	192	238	2
1985	3,925	347	375	513	508	440	406	306	208	217	-
1984	3,981	328	401	489	542	461	388	287	243	200	1
1983	3,648	308	407	460	478	430	367	291	194	188	-
1982	3,602	322	406	474	452	417	366	293	206	172	-
1981	3,393	315	356	455	417	399	339	236	200	177	2
1980	3,244	269	397	453	402	358	303	256	161	161	-
1979	3,086	323	391	384	383	365	278	229	142	146	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of breast (174-175)											
Black, male											
1998	48	3	3	6	4	4	7	7	5	7	-
1997	59	4	5	5	9	11	7	7	6	3	-
1996	62	2	4	4	8	9	8	9	6	7	-
1995	63	3	8	6	5	8	7	10	9	6	-
1994	51	3	1	1	7	11	11	8	2	5	-
1993	54	3	2	5	8	6	14	4	3	7	-
1992	37	1	4	4	3	5	7	5	5	1	-
1991	34	-	5	3	1	7	6	2	5	3	-
1990	44	5	3	6	6	7	7	1	4	2	-
1989	35	3	-	4	1	7	9	1	4	3	-
1988	58	3	8	2	2	7	3	9	7	12	-
1987	39	-	4	6	9	5	2	5	4	1	-
1986	31	-	2	1	5	7	5	6	2	3	-
1985	29	-	1	4	4	2	5	4	3	5	-
1984	36	4	-	2	5	8	7	5	2	1	-
1983	38	1	4	4	5	5	9	5	-	3	-
1982	35	-	-	5	5	5	3	4	7	5	-
1981	35	1	3	5	6	5	8	2	4	-	-
1980	29	2	1	5	4	2	5	4	2	2	-
1979	34	2	2	2	2	5	6	7	3	3	-
Malignant neoplasm of breast (174-175)											
Black, female											
1998	5,281	547	593	540	520	490	524	438	371	447	-
1997	5,424	556	547	550	554	550	577	468	370	437	-
1996	5,254	559	542	525	487	517	528	461	357	420	-
1995	5,265	550	558	516	551	530	577	433	365	387	-
1994	5,083	492	489	496	540	538	546	435	350	395	-
1993	4,998	449	518	482	521	549	510	464	332	379	1
1992	4,779	426	510	487	520	556	472	388	294	316	2
1991	4,809	410	457	491	574	527	513	406	316	315	1
1990	4,659	391	465	522	534	544	448	403	292	315	-
1989	4,403	421	436	467	499	528	435	380	248	282	-
1988	4,467	367	459	520	538	501	407	357	274	279	-
1987	4,252	382	445	480	547	436	444	311	243	247	1
1986	4,045	361	435	453	513	457	405	292	190	235	2
1985	3,896	347	374	509	504	438	401	302	205	212	-
1984	3,945	324	401	487	537	453	381	282	241	199	1
1983	3,610	307	403	456	473	425	358	286	194	185	-
1982	3,567	322	406	469	447	412	363	289	199	167	-
1981	3,358	314	353	450	411	394	331	234	196	177	2
1980	3,215	267	396	448	398	356	298	252	159	159	-
1979	3,052	321	389	382	381	360	272	222	139	143	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of cervix uteri (180)											
All races, both sexes											
1998	4,340	504	471	355	329	351	334	338	268	319	-
1997	4,499	526	450	390	400	399	353	325	261	328	-
1996	4,542	427	434	370	404	437	394	332	270	338	2
1995	4,503	469	432	364	349	379	395	356	324	341	-
1994	4,602	484	437	390	390	460	440	348	311	264	-
1993	4,583	458	418	367	396	417	450	387	303	316	1
1992	4,641	423	407	395	383	443	433	366	343	302	-
1991	4,514	404	366	400	445	449	405	376	286	313	-
1990	4,627	411	421	395	446	449	421	339	341	300	-
1989	4,487	379	376	404	435	489	405	380	274	330	-
1988	4,443	384	381	376	476	475	423	365	300	301	-
1987	4,425	392	366	387	455	504	421	352	309	275	2
1986	4,558	385	403	447	518	489	404	371	294	309	1
1985	4,508	336	405	428	468	482	456	395	295	327	-
1984	4,564	354	391	495	511	473	445	375	302	296	-
1983	4,651	345	384	506	559	528	443	385	292	318	-
1982	4,631	349	446	525	520	505	458	400	298	294	1
1981	4,815	360	478	564	604	497	471	397	301	316	-
1980	4,874	412	458	550	575	537	519	420	297	273	1
1979	4,936	403	498	612	553	574	501	403	315	293	1
Malignant neoplasm of cervix uteri (180)											
All races, female											
1998	4,340	504	471	355	329	351	334	338	268	319	-
1997	4,499	526	450	390	400	399	353	325	261	328	-
1996	4,542	427	434	370	404	437	394	332	270	338	2
1995	4,503	469	432	364	349	379	395	356	324	341	-
1994	4,602	484	437	390	390	460	440	348	311	264	-
1993	4,583	458	418	367	396	417	450	387	303	316	1
1992	4,641	423	407	395	383	443	433	366	343	302	-
1991	4,514	404	366	400	445	449	405	376	286	313	-
1990	4,627	411	421	395	446	449	421	339	341	300	-
1989	4,487	379	376	404	435	489	405	380	274	330	-
1988	4,443	384	381	376	476	475	423	365	300	301	-
1987	4,425	392	366	387	455	504	421	352	309	275	2
1986	4,558	385	403	447	518	489	404	371	294	309	1
1985	4,508	336	405	428	468	482	456	395	295	327	-
1984	4,564	354	391	495	511	473	445	375	302	296	-
1983	4,651	345	384	506	559	528	443	385	292	318	-
1982	4,631	349	446	525	520	505	458	400	298	294	1
1981	4,815	360	478	564	604	497	471	397	301	316	-
1980	4,874	412	458	550	575	537	519	420	297	273	1
1979	4,936	403	498	612	553	574	501	403	315	293	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of cervix uteri (180)											
White, both sexes											
1998	3,272	374	365	271	245	245	258	262	200	237	-
1997	3,371	393	342	276	309	306	271	242	201	249	-
1996	3,479	320	329	274	287	326	302	276	218	270	1
1995	3,359	346	319	272	250	284	301	267	256	258	-
1994	3,524	362	341	290	298	353	345	270	241	213	-
1993	3,419	333	304	275	282	304	351	292	231	248	1
1992	3,433	309	286	290	267	316	336	279	269	242	-
1991	3,386	301	261	292	322	327	311	299	221	257	-
1990	3,511	292	307	296	323	341	327	266	276	250	-
1989	3,321	277	270	291	305	354	312	283	208	259	-
1988	3,343	296	258	273	359	365	330	262	235	242	-
1987	3,349	281	259	290	341	376	336	278	250	225	-
1986	3,391	256	286	329	381	371	309	284	242	250	-
1985	3,405	247	269	321	360	384	348	303	233	263	-
1984	3,463	249	259	378	396	366	349	299	244	236	-
1983	3,497	240	279	366	420	410	334	282	238	265	-
1982	3,522	255	313	389	411	397	340	304	241	241	1
1981	3,712	280	343	424	474	399	369	316	244	259	-
1980	3,691	275	336	429	441	416	401	328	249	216	1
1979	3,805	281	385	478	423	435	410	315	264	247	1
Malignant neoplasm of cervix uteri (180)											
White, female											
1998	3,272	374	365	271	245	245	258	262	200	237	-
1997	3,371	393	342	276	309	306	271	242	201	249	-
1996	3,479	320	329	274	287	326	302	276	218	270	1
1995	3,359	346	319	272	250	284	301	267	256	258	-
1994	3,524	362	341	290	298	353	345	270	241	213	-
1993	3,419	333	304	275	282	304	351	292	231	248	1
1992	3,433	309	286	290	267	316	336	279	269	242	-
1991	3,386	301	261	292	322	327	311	299	221	257	-
1990	3,511	292	307	296	323	341	327	266	276	250	-
1989	3,321	277	270	291	305	354	312	283	208	259	-
1988	3,343	296	258	273	359	365	330	262	235	242	-
1987	3,349	281	259	290	341	376	336	278	250	225	-
1986	3,391	256	286	329	381	371	309	284	242	250	-
1985	3,405	247	269	321	360	384	348	303	233	263	-
1984	3,463	249	259	378	396	366	349	299	244	236	-
1983	3,497	240	279	366	420	410	334	282	238	265	-
1982	3,522	255	313	389	411	397	340	304	241	241	1
1981	3,712	280	343	424	474	399	369	316	244	259	-
1980	3,691	275	336	429	441	416	401	328	249	216	1
1979	3,805	281	385	478	423	435	410	315	264	247	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of cervix uteri (180)											
All other, both sexes											
1998	1,068	130	106	84	84	106	76	76	68	82	-
1997	1,128	133	108	114	91	93	82	83	60	79	-
1996	1,063	107	105	96	117	111	92	56	52	68	1
1995	1,144	123	113	92	99	95	94	89	68	83	-
1994	1,078	122	96	100	92	107	95	78	70	51	-
1993	1,164	125	114	92	114	113	99	95	72	68	-
1992	1,208	114	121	105	116	127	97	87	74	60	-
1991	1,128	103	105	108	123	122	94	77	65	56	-
1990	1,116	119	114	99	123	108	94	73	65	50	-
1989	1,166	102	106	113	130	135	93	97	66	71	-
1988	1,100	88	123	103	117	110	93	103	65	59	-
1987	1,076	111	107	97	114	128	85	74	59	50	2
1986	1,167	129	117	118	137	118	95	87	52	59	1
1985	1,103	89	136	107	108	98	108	92	62	64	-
1984	1,101	105	132	117	115	107	96	76	58	60	-
1983	1,154	105	105	140	139	118	109	103	54	53	-
1982	1,109	94	133	136	109	108	118	96	57	53	-
1981	1,103	80	135	140	130	98	102	81	57	57	-
1980	1,183	137	122	121	134	121	118	92	48	57	-
1979	1,131	122	113	134	130	139	91	88	51	46	-
Malignant neoplasm of cervix uteri (180)											
All other, female											
1998	1,068	130	106	84	84	106	76	76	68	82	-
1997	1,128	133	108	114	91	93	82	83	60	79	-
1996	1,063	107	105	96	117	111	92	56	52	68	1
1995	1,144	123	113	92	99	95	94	89	68	83	-
1994	1,078	122	96	100	92	107	95	78	70	51	-
1993	1,164	125	114	92	114	113	99	95	72	68	-
1992	1,208	114	121	105	116	127	97	87	74	60	-
1991	1,128	103	105	108	123	122	94	77	65	56	-
1990	1,116	119	114	99	123	108	94	73	65	50	-
1989	1,166	102	106	113	130	135	93	97	66	71	-
1988	1,100	88	123	103	117	110	93	103	65	59	-
1987	1,076	111	107	97	114	128	85	74	59	50	2
1986	1,167	129	117	118	137	118	95	87	52	59	1
1985	1,103	89	136	107	108	98	108	92	62	64	-
1984	1,101	105	132	117	115	107	96	76	58	60	-
1983	1,154	105	105	140	139	118	109	103	54	53	-
1982	1,109	94	133	136	109	108	118	96	57	53	-
1981	1,103	80	135	140	130	98	102	81	57	57	-
1980	1,183	137	122	121	134	121	118	92	48	57	-
1979	1,131	122	113	134	130	139	91	88	51	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of cervix uteri (180)											
Black, both sexes											
1998	906	111	87	70	75	89	66	59	60	69	-
1997	969	110	87	97	75	82	73	68	53	73	-
1996	907	92	87	71	98	91	82	54	46	63	1
1995	995	98	95	82	79	82	85	77	60	75	-
1994	931	99	84	82	77	95	87	65	64	48	-
1993	1,035	107	99	80	98	99	90	89	69	63	-
1992	1,068	100	105	92	105	112	87	74	72	55	-
1991	983	91	89	90	107	108	81	73	60	51	-
1990	972	99	97	86	104	95	85	65	60	45	-
1989	1,049	89	93	100	117	120	88	90	62	69	-
1988	974	67	104	85	106	100	86	99	61	54	-
1987	969	104	95	86	102	112	82	72	52	46	2
1986	1,067	113	105	107	127	115	87	83	45	54	1
1985	1,008	78	123	99	99	89	103	87	57	61	-
1984	1,008	98	119	107	103	97	85	72	58	58	-
1983	1,062	94	99	122	128	110	103	96	51	50	-
1982	1,031	89	122	130	100	102	110	89	54	50	-
1981	1,023	76	120	130	125	96	98	78	51	53	-
1980	1,097	120	112	116	129	115	110	86	48	50	-
1979	1,061	109	105	121	125	136	86	85	49	42	-
Malignant neoplasm of cervix uteri (180)											
Black, female											
1998	906	111	87	70	75	89	66	59	60	69	-
1997	969	110	87	97	75	82	73	68	53	73	-
1996	907	92	87	71	98	91	82	54	46	63	1
1995	995	98	95	82	79	82	85	77	60	75	-
1994	931	99	84	82	77	95	87	65	64	48	-
1993	1,035	107	99	80	98	99	90	89	69	63	-
1992	1,068	100	105	92	105	112	87	74	72	55	-
1991	983	91	89	90	107	108	81	73	60	51	-
1990	972	99	97	86	104	95	85	65	60	45	-
1989	1,049	89	93	100	117	120	88	90	62	69	-
1988	974	67	104	85	106	100	86	99	61	54	-
1987	969	104	95	86	102	112	82	72	52	46	2
1986	1,067	113	105	107	127	115	87	83	45	54	1
1985	1,008	78	123	99	99	89	103	87	57	61	-
1984	1,008	98	119	107	103	97	85	72	58	58	-
1983	1,062	94	99	122	128	110	103	96	51	50	-
1982	1,031	89	122	130	100	102	110	89	54	50	-
1981	1,023	76	120	130	125	96	98	78	51	53	-
1980	1,097	120	112	116	129	115	110	86	48	50	-
1979	1,061	109	105	121	125	136	86	85	49	42	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
All races, both sexes											
1998	6,421	166	263	406	566	814	1,052	1,081	901	1,025	-
1997	6,189	151	231	392	569	751	1,053	1,017	850	1,023	1
1996	6,311	149	241	332	595	853	1,046	990	941	1,011	1
1995	6,237	174	241	326	576	847	1,051	1,025	867	991	-
1994	6,163	126	231	332	591	857	1,138	1,005	830	926	-
1993	6,098	138	210	336	590	853	1,076	1,047	846	861	1
1992	6,064	105	215	314	575	945	1,061	1,017	842	824	-
1991	5,925	109	190	299	611	949	1,052	958	775	819	-
1990	6,028	118	214	351	599	969	1,066	1,031	763	777	1
1989	5,850	129	190	316	625	994	1,002	967	730	763	-
1988	5,892	104	179	347	701	949	1,075	953	765	691	1
1987	5,894	113	189	351	652	982	1,067	994	726	690	-
1986	5,853	103	183	356	716	1,002	1,057	871	727	697	2
1985	5,959	112	211	422	770	982	1,060	928	671	679	-
1984	5,965	114	210	407	773	1,013	1,085	876	693	674	-
1983	6,017	114	196	453	802	984	1,066	911	692	689	-
1982	5,956	127	250	421	776	1,053	1,012	887	667	639	2
1981	5,883	111	248	467	784	1,028	974	852	649	634	-
1980	6,035	117	276	513	800	1,019	1,021	844	649	657	-
1979	5,775	112	298	490	801	990	964	810	634	549	1
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
All races, female											
1998	6,421	166	263	406	566	814	1,052	1,081	901	1,025	-
1997	6,189	151	231	392	569	751	1,053	1,017	850	1,023	1
1996	6,311	149	241	332	595	853	1,046	990	941	1,011	1
1995	6,237	174	241	326	576	847	1,051	1,025	867	991	-
1994	6,163	126	231	332	591	857	1,138	1,005	830	926	-
1993	6,098	138	210	336	590	853	1,076	1,047	846	861	1
1992	6,064	105	215	314	575	945	1,061	1,017	842	824	-
1991	5,925	109	190	299	611	949	1,052	958	775	819	-
1990	6,028	118	214	351	599	969	1,066	1,031	763	777	1
1989	5,850	129	190	316	625	994	1,002	967	730	763	-
1988	5,892	104	179	347	701	949	1,075	953	765	691	1
1987	5,894	113	189	351	652	982	1,067	994	726	690	-
1986	5,853	103	183	356	716	1,002	1,057	871	727	697	2
1985	5,959	112	211	422	770	982	1,060	928	671	679	-
1984	5,965	114	210	407	773	1,013	1,085	876	693	674	-
1983	6,017	114	196	453	802	984	1,066	911	692	689	-
1982	5,956	127	250	421	776	1,053	1,012	887	667	639	2
1981	5,883	111	248	467	784	1,028	974	852	649	634	-
1980	6,035	117	276	513	800	1,019	1,021	844	649	657	-
1979	5,775	112	298	490	801	990	964	810	634	549	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
White, both sexes											
1998	5,340	130	211	325	434	643	865	913	797	914	-
1997	5,160	129	186	311	444	586	868	877	737	903	-
1996	5,223	120	193	266	461	673	852	826	819	892	1
1995	5,293	145	209	261	442	679	888	897	771	884	-
1994	5,157	109	185	262	448	688	945	866	722	834	-
1993	5,081	114	167	264	467	681	869	902	741	772	-
1992	5,096	87	178	242	457	764	900	863	739	744	-
1991	4,943	91	157	222	479	762	874	835	673	726	-
1990	5,066	97	174	273	467	807	901	883	670	681	1
1989	4,911	111	148	261	498	822	834	813	637	687	-
1988	4,956	86	145	280	553	776	918	815	674	611	-
1987	5,002	93	151	283	532	827	922	849	633	610	-
1986	4,968	83	144	289	588	834	884	755	649	632	2
1985	5,091	94	174	356	634	823	904	805	592	611	-
1984	5,076	80	167	346	656	840	935	756	599	602	-
1983	5,187	88	168	370	686	850	902	786	621	633	-
1982	5,097	107	200	338	653	875	870	782	595	573	2
1981	5,075	95	198	388	657	871	835	758	587	583	-
1980	5,155	93	226	422	663	870	864	738	585	590	-
1979	4,942	87	253	407	654	831	821	713	568	509	1
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
White, female											
1998	5,340	130	211	325	434	643	865	913	797	914	-
1997	5,160	129	186	311	444	586	868	877	737	903	-
1996	5,223	120	193	266	461	673	852	826	819	892	1
1995	5,293	145	209	261	442	679	888	897	771	884	-
1994	5,157	109	185	262	448	688	945	866	722	834	-
1993	5,081	114	167	264	467	681	869	902	741	772	-
1992	5,096	87	178	242	457	764	900	863	739	744	-
1991	4,943	91	157	222	479	762	874	835	673	726	-
1990	5,066	97	174	273	467	807	901	883	670	681	1
1989	4,911	111	148	261	498	822	834	813	637	687	-
1988	4,956	86	145	280	553	776	918	815	674	611	-
1987	5,002	93	151	283	532	827	922	849	633	610	-
1986	4,968	83	144	289	588	834	884	755	649	632	2
1985	5,091	94	174	356	634	823	904	805	592	611	-
1984	5,076	80	167	346	656	840	935	756	599	602	-
1983	5,187	88	168	370	686	850	902	786	621	633	-
1982	5,097	107	200	338	653	875	870	782	595	573	2
1981	5,075	95	198	388	657	871	835	758	587	583	-
1980	5,155	93	226	422	663	870	864	738	585	590	-
1979	4,942	87	253	407	654	831	821	713	568	509	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
All other, both sexes											
1998	1,081	36	52	81	132	171	187	168	104	111	-
1997	1,029	22	45	81	125	165	185	140	113	120	1
1996	1,088	29	48	66	134	180	194	164	122	119	-
1995	944	29	32	65	134	168	163	128	96	107	-
1994	1,006	17	46	70	143	169	193	139	108	92	-
1993	1,017	24	43	72	123	172	207	145	105	89	1
1992	968	18	37	72	118	181	161	154	103	80	-
1991	982	18	33	77	132	187	178	123	102	93	-
1990	962	21	40	78	132	162	165	148	93	96	-
1989	939	18	42	55	127	172	168	154	93	76	-
1988	936	18	34	67	148	173	157	138	91	80	1
1987	892	20	38	68	120	155	145	145	93	80	-
1986	885	20	39	67	128	168	173	116	78	65	-
1985	868	18	37	66	136	159	156	123	79	68	-
1984	889	34	43	61	117	173	150	120	94	72	-
1983	830	26	28	83	116	134	164	125	71	56	-
1982	859	20	50	83	123	178	142	105	72	66	-
1981	808	16	50	79	127	157	139	94	62	51	-
1980	880	24	50	91	137	149	157	106	64	67	-
1979	833	25	45	83	147	159	143	97	66	40	-
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
All other, female											
1998	1,081	36	52	81	132	171	187	168	104	111	-
1997	1,029	22	45	81	125	165	185	140	113	120	1
1996	1,088	29	48	66	134	180	194	164	122	119	-
1995	944	29	32	65	134	168	163	128	96	107	-
1994	1,006	17	46	70	143	169	193	139	108	92	-
1993	1,017	24	43	72	123	172	207	145	105	89	1
1992	968	18	37	72	118	181	161	154	103	80	-
1991	982	18	33	77	132	187	178	123	102	93	-
1990	962	21	40	78	132	162	165	148	93	96	-
1989	939	18	42	55	127	172	168	154	93	76	-
1988	936	18	34	67	148	173	157	138	91	80	1
1987	892	20	38	68	120	155	145	145	93	80	-
1986	885	20	39	67	128	168	173	116	78	65	-
1985	868	18	37	66	136	159	156	123	79	68	-
1984	889	34	43	61	117	173	150	120	94	72	-
1983	830	26	28	83	116	134	164	125	71	56	-
1982	859	20	50	83	123	178	142	105	72	66	-
1981	808	16	50	79	127	157	139	94	62	51	-
1980	880	24	50	91	137	149	157	106	64	67	-
1979	833	25	45	83	147	159	143	97	66	40	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
Black, both sexes											
1998	970	28	36	68	121	161	173	154	95	105	-
1997	935	13	38	69	114	159	170	134	101	112	1
1996	977	19	40	54	119	169	179	151	109	111	-
1995	864	22	28	56	117	154	156	124	91	100	-
1994	929	13	39	63	133	161	180	129	101	86	-
1993	942	20	37	61	115	161	193	138	102	86	1
1992	889	11	34	62	111	162	148	148	100	75	-
1991	913	17	25	66	119	181	169	114	100	90	-
1990	893	17	35	70	121	150	152	143	90	94	-
1989	869	16	35	52	116	156	158	146	89	74	-
1988	880	13	30	58	138	166	153	128	86	79	1
1987	841	18	33	57	114	146	136	142	91	79	-
1986	835	18	35	55	118	164	167	112	75	63	-
1985	826	17	31	62	129	154	147	121	77	66	-
1984	837	31	38	56	105	166	146	114	90	71	-
1983	786	22	26	74	110	129	161	120	67	55	-
1982	807	14	41	76	116	171	136	100	72	64	-
1981	770	12	49	74	119	152	138	91	60	47	-
1980	848	22	47	87	133	144	154	101	62	64	-
1979	794	23	40	75	141	153	138	93	65	39	-
Malignant neoplasms of corpus uteri and uterus, part unspecified (179,182)											
Black, female											
1998	970	28	36	68	121	161	173	154	95	105	-
1997	935	13	38	69	114	159	170	134	101	112	1
1996	977	19	40	54	119	169	179	151	109	111	-
1995	864	22	28	56	117	154	156	124	91	100	-
1994	929	13	39	63	133	161	180	129	101	86	-
1993	942	20	37	61	115	161	193	138	102	86	1
1992	889	11	34	62	111	162	148	148	100	75	-
1991	913	17	25	66	119	181	169	114	100	90	-
1990	893	17	35	70	121	150	152	143	90	94	-
1989	869	16	35	52	116	156	158	146	89	74	-
1988	880	13	30	58	138	166	153	128	86	79	1
1987	841	18	33	57	114	146	136	142	91	79	-
1986	835	18	35	55	118	164	167	112	75	63	-
1985	826	17	31	62	129	154	147	121	77	66	-
1984	837	31	38	56	105	166	146	114	90	71	-
1983	786	22	26	74	110	129	161	120	67	55	-
1982	807	14	41	76	116	171	136	100	72	64	-
1981	770	12	49	74	119	152	138	91	60	47	-
1980	848	22	47	87	133	144	154	101	62	64	-
1979	794	23	40	75	141	153	138	93	65	39	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of ovary (183.0)											
All races, both sexes											
1998	13,391	549	876	1,080	1,269	1,639	2,048	2,082	1,653	1,590	1
1997	13,507	619	860	992	1,260	1,659	2,176	2,112	1,643	1,567	-
1996	13,161	623	760	993	1,314	1,704	2,053	1,991	1,663	1,463	-
1995	13,342	577	807	1,063	1,292	1,782	2,156	2,007	1,578	1,458	1
1994	13,500	573	839	1,061	1,386	1,892	2,118	2,000	1,598	1,428	-
1993	12,870	571	766	991	1,338	1,796	2,034	1,867	1,581	1,330	-
1992	13,182	534	785	1,030	1,462	1,916	2,153	1,928	1,448	1,300	1
1991	13,028	502	762	1,081	1,478	2,018	2,028	1,835	1,442	1,250	-
1990	12,566	542	715	1,011	1,476	1,971	1,985	1,838	1,278	1,148	-
1989	12,256	481	737	1,001	1,574	1,977	1,884	1,694	1,280	1,034	-
1988	12,214	474	715	1,117	1,589	1,939	1,914	1,636	1,224	1,008	1
1987	11,838	481	741	1,148	1,549	1,854	1,785	1,636	1,155	925	-
1986	11,729	509	703	1,135	1,626	1,807	1,877	1,458	1,085	930	1
1985	11,357	488	758	1,156	1,626	1,717	1,760	1,481	1,066	764	-
1984	11,208	494	767	1,210	1,536	1,692	1,670	1,413	1,050	797	-
1983	11,216	450	865	1,218	1,585	1,749	1,683	1,382	950	782	3
1982	11,059	505	799	1,301	1,593	1,682	1,580	1,385	933	727	2
1981	10,858	502	887	1,328	1,572	1,632	1,530	1,318	905	639	3
1980	10,731	499	974	1,256	1,529	1,588	1,514	1,271	864	682	-
1979	10,565	539	1,035	1,265	1,420	1,657	1,460	1,155	868	578	1
Malignant neoplasm of ovary (183.0)											
All races, female											
1998	13,391	549	876	1,080	1,269	1,639	2,048	2,082	1,653	1,590	1
1997	13,507	619	860	992	1,260	1,659	2,176	2,112	1,643	1,567	-
1996	13,161	623	760	993	1,314	1,704	2,053	1,991	1,663	1,463	-
1995	13,342	577	807	1,063	1,292	1,782	2,156	2,007	1,578	1,458	1
1994	13,500	573	839	1,061	1,386	1,892	2,118	2,000	1,598	1,428	-
1993	12,870	571	766	991	1,338	1,796	2,034	1,867	1,581	1,330	-
1992	13,182	534	785	1,030	1,462	1,916	2,153	1,928	1,448	1,300	1
1991	13,028	502	762	1,081	1,478	2,018	2,028	1,835	1,442	1,250	-
1990	12,566	542	715	1,011	1,476	1,971	1,985	1,838	1,278	1,148	-
1989	12,256	481	737	1,001	1,574	1,977	1,884	1,694	1,280	1,034	-
1988	12,214	474	715	1,117	1,589	1,939	1,914	1,636	1,224	1,008	1
1987	11,838	481	741	1,148	1,549	1,854	1,785	1,636	1,155	925	-
1986	11,729	509	703	1,135	1,626	1,807	1,877	1,458	1,085	930	1
1985	11,357	488	758	1,156	1,626	1,717	1,760	1,481	1,066	764	-
1984	11,208	494	767	1,210	1,536	1,692	1,670	1,413	1,050	797	-
1983	11,216	450	865	1,218	1,585	1,749	1,683	1,382	950	782	3
1982	11,059	505	799	1,301	1,593	1,682	1,580	1,385	933	727	2
1981	10,858	502	887	1,328	1,572	1,632	1,530	1,318	905	639	3
1980	10,731	499	974	1,256	1,529	1,588	1,514	1,271	864	682	-
1979	10,565	539	1,035	1,265	1,420	1,657	1,460	1,155	868	578	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of ovary (183.0)											
White, both sexes											
1998	12,115	474	784	958	1,122	1,462	1,864	1,923	1,551	1,491	1
1997	12,181	538	762	895	1,135	1,463	1,964	1,944	1,527	1,445	-
1996	11,913	539	674	876	1,196	1,518	1,858	1,856	1,543	1,365	-
1995	12,169	513	720	958	1,163	1,623	2,003	1,851	1,439	1,366	1
1994	12,256	513	762	940	1,234	1,715	1,930	1,838	1,487	1,330	-
1993	11,680	514	675	883	1,207	1,630	1,850	1,716	1,469	1,227	-
1992	11,947	467	698	917	1,293	1,725	1,977	1,797	1,339	1,199	-
1991	11,810	436	677	981	1,344	1,819	1,853	1,687	1,345	1,149	-
1990	11,469	491	647	906	1,349	1,790	1,824	1,709	1,171	1,065	-
1989	11,228	441	655	909	1,445	1,802	1,742	1,568	1,204	960	-
1988	11,129	412	646	997	1,446	1,764	1,752	1,518	1,130	955	-
1987	10,843	427	669	1,050	1,414	1,708	1,636	1,517	1,081	860	-
1986	10,751	458	623	1,031	1,472	1,651	1,727	1,378	1,016	869	-
1985	10,410	443	678	1,055	1,465	1,565	1,627	1,381	1,002	730	-
1984	10,275	445	695	1,088	1,408	1,557	1,542	1,311	975	746	-
1983	10,279	393	777	1,106	1,450	1,633	1,556	1,280	894	730	3
1982	10,174	458	721	1,199	1,462	1,550	1,446	1,288	879	691	2
1981	9,988	454	816	1,215	1,447	1,509	1,416	1,233	829	608	3
1980	9,852	447	891	1,145	1,394	1,454	1,386	1,194	817	641	-
1979	9,716	474	939	1,152	1,302	1,521	1,337	1,096	839	552	1
Malignant neoplasm of ovary (183.0)											
White, female											
1998	12,115	474	784	958	1,122	1,462	1,864	1,923	1,551	1,491	1
1997	12,181	538	762	895	1,135	1,463	1,964	1,944	1,527	1,445	-
1996	11,913	539	674	876	1,196	1,518	1,858	1,856	1,543	1,365	-
1995	12,169	513	720	958	1,163	1,623	2,003	1,851	1,439	1,366	1
1994	12,256	513	762	940	1,234	1,715	1,930	1,838	1,487	1,330	-
1993	11,680	514	675	883	1,207	1,630	1,850	1,716	1,469	1,227	-
1992	11,947	467	698	917	1,293	1,725	1,977	1,797	1,339	1,199	-
1991	11,810	436	677	981	1,344	1,819	1,853	1,687	1,345	1,149	-
1990	11,469	491	647	906	1,349	1,790	1,824	1,709	1,171	1,065	-
1989	11,228	441	655	909	1,445	1,802	1,742	1,568	1,204	960	-
1988	11,129	412	646	997	1,446	1,764	1,752	1,518	1,130	955	-
1987	10,843	427	669	1,050	1,414	1,708	1,636	1,517	1,081	860	-
1986	10,751	458	623	1,031	1,472	1,651	1,727	1,378	1,016	869	-
1985	10,410	443	678	1,055	1,465	1,565	1,627	1,381	1,002	730	-
1984	10,275	445	695	1,088	1,408	1,557	1,542	1,311	975	746	-
1983	10,279	393	777	1,106	1,450	1,633	1,556	1,280	894	730	3
1982	10,174	458	721	1,199	1,462	1,550	1,446	1,288	879	691	2
1981	9,988	454	816	1,215	1,447	1,509	1,416	1,233	829	608	3
1980	9,852	447	891	1,145	1,394	1,454	1,386	1,194	817	641	-
1979	9,716	474	939	1,152	1,302	1,521	1,337	1,096	839	552	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of ovary (183.0)											
All other, both sexes											
1998	1,276	75	92	122	147	177	184	159	102	99	-
1997	1,326	81	98	97	125	196	212	168	116	122	-
1996	1,248	84	86	117	118	186	195	135	120	98	-
1995	1,173	64	87	105	129	159	153	156	139	92	-
1994	1,244	60	77	121	152	177	188	162	111	98	-
1993	1,190	57	91	108	131	166	184	151	112	103	-
1992	1,235	67	87	113	169	191	176	131	109	101	1
1991	1,218	66	85	100	134	199	175	148	97	101	-
1990	1,097	51	68	105	127	181	161	129	107	83	-
1989	1,028	40	82	92	129	175	142	126	76	74	-
1988	1,085	62	69	120	143	175	162	118	94	53	1
1987	995	54	72	98	135	146	149	119	74	65	-
1986	978	51	80	104	154	156	150	80	69	61	1
1985	947	45	80	101	161	152	133	100	64	34	-
1984	933	49	72	122	128	135	128	102	75	51	-
1983	937	57	88	112	135	116	127	102	56	52	-
1982	885	47	78	102	131	132	134	97	54	36	-
1981	870	48	71	113	125	123	114	85	76	31	-
1980	879	52	83	111	135	134	128	77	47	41	-
1979	849	65	96	113	118	136	123	59	29	26	-
Malignant neoplasm of ovary (183.0)											
All other, female											
1998	1,276	75	92	122	147	177	184	159	102	99	-
1997	1,326	81	98	97	125	196	212	168	116	122	-
1996	1,248	84	86	117	118	186	195	135	120	98	-
1995	1,173	64	87	105	129	159	153	156	139	92	-
1994	1,244	60	77	121	152	177	188	162	111	98	-
1993	1,190	57	91	108	131	166	184	151	112	103	-
1992	1,235	67	87	113	169	191	176	131	109	101	1
1991	1,218	66	85	100	134	199	175	148	97	101	-
1990	1,097	51	68	105	127	181	161	129	107	83	-
1989	1,028	40	82	92	129	175	142	126	76	74	-
1988	1,085	62	69	120	143	175	162	118	94	53	1
1987	995	54	72	98	135	146	149	119	74	65	-
1986	978	51	80	104	154	156	150	80	69	61	1
1985	947	45	80	101	161	152	133	100	64	34	-
1984	933	49	72	122	128	135	128	102	75	51	-
1983	937	57	88	112	135	116	127	102	56	52	-
1982	885	47	78	102	131	132	134	97	54	36	-
1981	870	48	71	113	125	123	114	85	76	31	-
1980	879	52	83	111	135	134	128	77	47	41	-
1979	849	65	96	113	118	136	123	59	29	26	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of ovary (183.0)											
Black, both sexes											
1998	1,068	55	71	92	122	147	157	143	89	91	-
1997	1,117	62	82	80	102	165	187	146	98	111	-
1996	1,011	51	57	85	98	160	171	117	107	87	-
1995	992	44	61	81	107	143	140	141	122	82	-
1994	1,034	42	59	92	122	149	167	140	96	95	-
1993	1,000	39	64	79	107	143	167	138	100	98	-
1992	1,043	47	68	92	135	163	149	121	99	98	1
1991	1,045	49	63	75	115	176	159	135	95	92	-
1990	959	41	57	93	111	157	140	120	102	74	-
1989	890	37	68	74	103	154	131	119	68	68	-
1988	938	49	57	103	120	150	149	106	88	48	1
1987	880	47	61	79	123	134	138	110	67	58	-
1986	855	43	66	86	135	142	136	73	65	56	1
1985	834	40	67	87	140	140	124	91	57	33	-
1984	834	44	61	108	113	124	118	94	69	46	-
1983	837	43	75	101	123	106	119	98	49	48	-
1982	804	35	64	91	118	126	131	92	51	34	-
1981	775	33	55	104	116	112	109	81	72	26	-
1980	817	48	67	105	125	126	125	73	46	39	-
1979	782	55	79	103	111	132	119	56	29	25	-
Malignant neoplasm of ovary (183.0)											
Black, female											
1998	1,068	55	71	92	122	147	157	143	89	91	-
1997	1,117	62	82	80	102	165	187	146	98	111	-
1996	1,011	51	57	85	98	160	171	117	107	87	-
1995	992	44	61	81	107	143	140	141	122	82	-
1994	1,034	42	59	92	122	149	167	140	96	95	-
1993	1,000	39	64	79	107	143	167	138	100	98	-
1992	1,043	47	68	92	135	163	149	121	99	98	1
1991	1,045	49	63	75	115	176	159	135	95	92	-
1990	959	41	57	93	111	157	140	120	102	74	-
1989	890	37	68	74	103	154	131	119	68	68	-
1988	938	49	57	103	120	150	149	106	88	48	1
1987	880	47	61	79	123	134	138	110	67	58	-
1986	855	43	66	86	135	142	136	73	65	56	1
1985	834	40	67	87	140	140	124	91	57	33	-
1984	834	44	61	108	113	124	118	94	69	46	-
1983	837	43	75	101	123	106	119	98	49	48	-
1982	804	35	64	91	118	126	131	92	51	34	-
1981	775	33	55	104	116	112	109	81	72	26	-
1980	817	48	67	105	125	126	125	73	46	39	-
1979	782	55	79	103	111	132	119	56	29	25	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of prostate (185)											
All races, both sexes											
1998	32,203	82	291	673	1,429	2,768	4,795	6,750	7,050	8,323	-
1997	32,891	90	274	693	1,368	2,987	5,084	6,838	7,026	8,485	2
1996	34,123	89	294	682	1,500	3,227	5,677	7,139	7,222	8,257	1
1995	34,475	78	252	671	1,626	3,359	5,723	7,065	7,329	8,328	-
1994	34,902	96	274	662	1,666	3,495	5,966	7,249	7,429	8,034	1
1993	34,865	75	241	691	1,732	3,695	5,933	7,295	7,244	7,926	-
1992	34,240	75	234	685	1,889	3,883	5,950	7,256	6,928	7,302	2
1991	33,564	60	257	676	1,913	3,768	5,949	7,122	6,940	6,847	1
1990	32,378	47	257	770	1,955	3,930	5,768	6,894	6,452	6,276	2
1989	30,520	67	232	756	1,922	3,727	5,393	6,477	6,019	5,895	1
1988	28,982	64	247	705	1,822	3,558	5,254	6,188	5,682	5,433	2
1987	27,864	69	215	679	1,762	3,538	5,071	5,905	5,351	5,242	1
1986	27,262	58	223	717	1,851	3,439	5,059	5,752	5,162	4,974	1
1985	25,943	59	235	726	1,747	3,215	4,800	5,576	4,917	4,639	3
1984	25,407	57	210	734	1,746	3,130	4,635	5,384	4,654	4,817	8
1983	24,959	71	233	722	1,738	3,123	4,636	5,337	4,536	4,525	5
1982	24,021	52	234	728	1,667	3,097	4,324	5,093	4,433	4,358	8
1981	23,370	68	247	684	1,644	3,013	4,310	4,828	4,280	4,266	-
1980	22,881	67	241	710	1,596	3,035	4,129	4,767	4,178	4,135	1
1979	22,240	54	229	699	1,515	2,871	4,157	4,674	4,125	3,900	1
Malignant neoplasm of prostate (185)											
All races, male											
1998	32,203	82	291	673	1,429	2,768	4,795	6,750	7,050	8,323	-
1997	32,891	90	274	693	1,368	2,987	5,084	6,838	7,026	8,485	2
1996	34,123	89	294	682	1,500	3,227	5,677	7,139	7,222	8,257	1
1995	34,475	78	252	671	1,626	3,359	5,723	7,065	7,329	8,328	-
1994	34,902	96	274	662	1,666	3,495	5,966	7,249	7,429	8,034	1
1993	34,865	75	241	691	1,732	3,695	5,933	7,295	7,244	7,926	-
1992	34,240	75	234	685	1,889	3,883	5,950	7,256	6,928	7,302	2
1991	33,564	60	257	676	1,913	3,768	5,949	7,122	6,940	6,847	1
1990	32,378	47	257	770	1,955	3,930	5,768	6,894	6,452	6,276	2
1989	30,520	67	232	756	1,922	3,727	5,393	6,477	6,019	5,895	1
1988	28,982	64	247	705	1,822	3,558	5,254	6,188	5,682	5,433	2
1987	27,864	69	215	679	1,762	3,538	5,071	5,905	5,351	5,242	1
1986	27,262	58	223	717	1,851	3,439	5,059	5,752	5,162	4,974	1
1985	25,943	59	235	726	1,747	3,215	4,800	5,576	4,917	4,639	3
1984	25,407	57	210	734	1,746	3,130	4,635	5,384	4,654	4,817	8
1983	24,959	71	233	722	1,738	3,123	4,636	5,337	4,536	4,525	5
1982	24,021	52	234	728	1,667	3,097	4,324	5,093	4,433	4,358	8
1981	23,370	68	247	684	1,644	3,013	4,310	4,828	4,280	4,266	-
1980	22,881	67	241	710	1,596	3,035	4,129	4,767	4,178	4,135	1
1979	22,240	54	229	699	1,515	2,871	4,157	4,674	4,125	3,900	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,
sex and year

	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of prostate (185)											
White, both sexes											
1998	26,416	57	200	507	1,030	2,162	3,822	5,544	5,980	7,083	-
1997	27,052	66	199	504	1,006	2,317	4,045	5,652	6,036	7,196	1
1996	28,021	68	213	506	1,115	2,499	4,580	5,867	6,097	7,048	-
1995	28,495	56	195	510	1,203	2,630	4,585	5,950	6,190	7,145	-
1994	28,912	69	207	489	1,281	2,720	4,798	6,077	6,318	6,932	-
1993	28,847	46	178	517	1,318	2,904	4,850	6,165	6,095	6,750	-
1992	28,430	51	165	495	1,461	3,102	4,853	6,121	5,912	6,243	2
1991	27,971	49	184	502	1,513	3,004	4,897	5,987	5,896	5,914	-
1990	26,917	37	202	571	1,538	3,167	4,782	5,766	5,472	5,360	2
1989	25,469	48	168	570	1,550	2,997	4,452	5,398	5,149	5,110	-
1988	24,176	47	182	543	1,434	2,864	4,386	5,154	4,862	4,683	1
1987	23,169	55	161	508	1,404	2,832	4,191	4,917	4,545	4,529	-
1986	22,708	36	165	567	1,482	2,785	4,117	4,810	4,412	4,311	1
1985	21,472	40	157	552	1,387	2,598	3,924	4,647	4,162	3,983	2
1984	21,104	38	162	576	1,380	2,551	3,747	4,493	3,936	4,193	4
1983	20,787	50	177	549	1,380	2,523	3,790	4,450	3,842	3,994	3
1982	20,065	39	180	567	1,323	2,506	3,535	4,295	3,786	3,812	3
1981	19,405	49	184	528	1,283	2,383	3,504	4,052	3,673	3,725	-
1980	19,096	53	178	554	1,274	2,397	3,377	4,038	3,587	3,621	1
1979	18,654	31	176	551	1,192	2,276	3,425	3,934	3,584	3,474	-
Malignant neoplasm of prostate (185)											
White, male											
1998	26,416	57	200	507	1,030	2,162	3,822	5,544	5,980	7,083	-
1997	27,052	66	199	504	1,006	2,317	4,045	5,652	6,036	7,196	1
1996	28,021	68	213	506	1,115	2,499	4,580	5,867	6,097	7,048	-
1995	28,495	56	195	510	1,203	2,630	4,585	5,950	6,190	7,145	-
1994	28,912	69	207	489	1,281	2,720	4,798	6,077	6,318	6,932	-
1993	28,847	46	178	517	1,318	2,904	4,850	6,165	6,095	6,750	-
1992	28,430	51	165	495	1,461	3,102	4,853	6,121	5,912	6,243	2
1991	27,971	49	184	502	1,513	3,004	4,897	5,987	5,896	5,914	-
1990	26,917	37	202	571	1,538	3,167	4,782	5,766	5,472	5,360	2
1989	25,469	48	168	570	1,550	2,997	4,452	5,398	5,149	5,110	-
1988	24,176	47	182	543	1,434	2,864	4,386	5,154	4,862	4,683	1
1987	23,169	55	161	508	1,404	2,832	4,191	4,917	4,545	4,529	-
1986	22,708	36	165	567	1,482	2,785	4,117	4,810	4,412	4,311	1
1985	21,472	40	157	552	1,387	2,598	3,924	4,647	4,162	3,983	2
1984	21,104	38	162	576	1,380	2,551	3,747	4,493	3,936	4,193	4
1983	20,787	50	177	549	1,380	2,523	3,790	4,450	3,842	3,994	3
1982	20,065	39	180	567	1,323	2,506	3,535	4,295	3,786	3,812	3
1981	19,405	49	184	528	1,283	2,383	3,504	4,052	3,673	3,725	-
1980	19,096	53	178	554	1,274	2,397	3,377	4,038	3,587	3,621	1
1979	18,654	31	176	551	1,192	2,276	3,425	3,934	3,584	3,474	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of prostate (185)											
All other, both sexes											
1998	5,787	25	91	166	399	606	973	1,206	1,070	1,240	-
1997	5,839	24	75	189	362	670	1,039	1,186	990	1,289	1
1996	6,102	21	81	176	385	728	1,097	1,272	1,125	1,209	1
1995	5,980	22	57	161	423	729	1,138	1,115	1,139	1,183	-
1994	5,990	27	67	173	385	775	1,168	1,172	1,111	1,102	1
1993	6,018	29	63	174	414	791	1,083	1,130	1,149	1,176	-
1992	5,810	24	69	190	428	781	1,097	1,135	1,016	1,059	-
1991	5,593	11	73	174	400	764	1,052	1,135	1,044	933	1
1990	5,461	10	55	199	417	763	986	1,128	980	916	-
1989	5,051	19	64	186	372	730	941	1,079	870	785	1
1988	4,806	17	65	162	388	694	868	1,034	820	750	1
1987	4,695	14	54	171	358	706	880	988	806	713	1
1986	4,554	22	58	150	369	654	942	942	750	663	-
1985	4,471	19	78	174	360	617	876	929	755	656	1
1984	4,303	19	48	158	366	579	888	891	718	624	4
1983	4,172	21	56	173	358	600	846	887	694	531	2
1982	3,956	13	54	161	344	591	789	798	647	546	5
1981	3,965	19	63	156	361	630	806	776	607	541	-
1980	3,785	14	63	156	322	638	752	729	591	514	-
1979	3,586	23	53	148	323	595	732	740	541	426	1
Malignant neoplasm of prostate (185)											
All other, male											
1998	5,787	25	91	166	399	606	973	1,206	1,070	1,240	-
1997	5,839	24	75	189	362	670	1,039	1,186	990	1,289	1
1996	6,102	21	81	176	385	728	1,097	1,272	1,125	1,209	1
1995	5,980	22	57	161	423	729	1,138	1,115	1,139	1,183	-
1994	5,990	27	67	173	385	775	1,168	1,172	1,111	1,102	1
1993	6,018	29	63	174	414	791	1,083	1,130	1,149	1,176	-
1992	5,810	24	69	190	428	781	1,097	1,135	1,016	1,059	-
1991	5,593	11	73	174	400	764	1,052	1,135	1,044	933	1
1990	5,461	10	55	199	417	763	986	1,128	980	916	-
1989	5,051	19	64	186	372	730	941	1,079	870	785	1
1988	4,806	17	65	162	388	694	868	1,034	820	750	1
1987	4,695	14	54	171	358	706	880	988	806	713	1
1986	4,554	22	58	150	369	654	942	942	750	663	-
1985	4,471	19	78	174	360	617	876	929	755	656	1
1984	4,303	19	48	158	366	579	888	891	718	624	4
1983	4,172	21	56	173	358	600	846	887	694	531	2
1982	3,956	13	54	161	344	591	789	798	647	546	5
1981	3,965	19	63	156	361	630	806	776	607	541	-
1980	3,785	14	63	156	322	638	752	729	591	514	-
1979	3,586	23	53	148	323	595	732	740	541	426	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of prostate (185)											
Black, both sexes											
1998	5,436	24	87	158	375	581	925	1,124	1,004	1,148	-
1997	5,466	24	72	180	348	628	992	1,128	900	1,180	1
1996	5,717	20	76	164	364	696	1,024	1,210	1,054	1,103	1
1995	5,592	21	55	151	406	685	1,083	1,052	1,044	1,082	-
1994	5,650	27	66	167	367	734	1,115	1,104	1,042	1,018	1
1993	5,656	27	57	166	395	765	1,034	1,060	1,057	1,087	-
1992	5,485	24	65	183	417	750	1,035	1,066	955	980	-
1991	5,299	10	70	167	381	744	997	1,092	980	851	1
1990	5,181	10	50	195	400	725	944	1,063	922	865	-
1989	4,785	18	62	176	361	707	897	1,019	812	728	1
1988	4,582	17	63	153	376	670	834	988	777	697	1
1987	4,488	12	52	166	353	677	844	946	756	677	1
1986	4,358	22	56	143	355	639	912	899	709	619	-
1985	4,273	18	74	170	351	598	845	882	714	614	1
1984	4,130	19	48	152	359	565	853	843	682	597	4
1983	4,040	21	54	171	350	586	821	855	672	505	1
1982	3,811	13	53	157	332	577	761	771	614	520	5
1981	3,839	19	62	154	353	620	776	750	582	517	-
1980	3,670	13	61	151	318	625	728	709	566	493	-
1979	3,478	23	53	147	319	574	716	717	516	408	1
Malignant neoplasm of prostate (185)											
Black, male											
1998	5,436	24	87	158	375	581	925	1,124	1,004	1,148	-
1997	5,466	24	72	180	348	628	992	1,128	900	1,180	1
1996	5,717	20	76	164	364	696	1,024	1,210	1,054	1,103	1
1995	5,592	21	55	151	406	685	1,083	1,052	1,044	1,082	-
1994	5,650	27	66	167	367	734	1,115	1,104	1,042	1,018	1
1993	5,656	27	57	166	395	765	1,034	1,060	1,057	1,087	-
1992	5,485	24	65	183	417	750	1,035	1,066	955	980	-
1991	5,299	10	70	167	381	744	997	1,092	980	851	1
1990	5,181	10	50	195	400	725	944	1,063	922	865	-
1989	4,785	18	62	176	361	707	897	1,019	812	728	1
1988	4,582	17	63	153	376	670	834	988	777	697	1
1987	4,488	12	52	166	353	677	844	946	756	677	1
1986	4,358	22	56	143	355	639	912	899	709	619	-
1985	4,273	18	74	170	351	598	845	882	714	614	1
1984	4,130	19	48	152	359	565	853	843	682	597	4
1983	4,040	21	54	171	350	586	821	855	672	505	1
1982	3,811	13	53	157	332	577	761	771	614	520	5
1981	3,839	19	62	154	353	620	776	750	582	517	-
1980	3,670	13	61	151	318	625	728	709	566	493	-
1979	3,478	23	53	147	319	574	716	717	516	408	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All races, both sexes											
1998	11,484	453	754	986	1,151	1,495	1,791	1,699	1,381	1,262	-
1997	11,294	476	723	924	1,174	1,553	1,703	1,705	1,315	1,257	-
1996	11,095	458	685	907	1,188	1,528	1,770	1,611	1,312	1,153	-
1995	11,083	449	686	899	1,210	1,500	1,712	1,675	1,296	1,183	-
1994	10,750	432	650	865	1,192	1,574	1,706	1,601	1,197	1,082	1
1993	10,322	400	670	786	1,189	1,511	1,727	1,480	1,159	935	-
1992	10,429	441	588	867	1,198	1,490	1,684	1,548	1,171	1,001	2
1991	10,270	356	588	862	1,263	1,533	1,641	1,472	1,146	936	-
1990	9,843	365	571	854	1,170	1,544	1,567	1,432	1,052	838	-
1989	9,638	390	533	826	1,203	1,489	1,536	1,374	1,041	800	-
1988	9,203	353	532	783	1,203	1,400	1,494	1,265	966	742	-
1987	9,332	346	539	893	1,241	1,436	1,482	1,351	908	722	-
1986	8,987	323	547	850	1,253	1,308	1,422	1,234	926	674	1
1985	8,660	317	526	835	1,204	1,300	1,369	1,177	854	631	-
1984	8,411	293	548	849	1,109	1,265	1,374	1,156	782	625	-
1983	8,228	288	527	841	1,169	1,288	1,283	1,099	712	618	1
1982	8,084	328	525	854	1,071	1,271	1,263	1,087	714	565	-
1981	7,702	286	531	842	1,052	1,164	1,179	1,020	679	525	-
1980	7,537	292	569	841	1,075	1,201	1,158	940	625	454	-
1979	7,284	315	553	873	1,002	1,122	1,097	861	625	447	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All races, male											
1998	7,095	337	528	690	774	1,017	1,115	1,014	739	549	-
1997	6,914	329	489	632	787	1,013	1,085	1,027	701	565	-
1996	6,798	312	490	632	801	995	1,100	941	697	513	-
1995	6,639	316	479	601	796	960	1,025	1,000	680	503	-
1994	6,522	297	463	598	798	994	1,035	939	607	513	1
1993	6,358	280	472	538	808	976	1,103	857	595	420	-
1992	6,351	315	407	615	788	974	1,047	886	601	433	2
1991	6,179	247	402	539	840	973	1,031	838	593	410	-
1990	6,054	255	396	601	797	982	961	836	588	364	-
1989	5,879	276	359	564	800	930	965	801	549	356	-
1988	5,615	242	375	518	790	903	915	764	484	323	-
1987	5,704	227	387	624	828	910	884	780	480	319	-
1986	5,410	215	382	570	806	847	854	676	490	310	-
1985	5,288	228	363	573	784	842	806	695	454	278	-
1984	5,201	215	376	586	735	809	872	636	428	294	-
1983	5,033	196	358	567	773	852	767	639	366	270	1
1982	5,013	229	370	596	697	811	772	639	382	265	-
1981	4,728	191	372	561	743	714	702	598	345	238	-
1980	4,751	203	398	585	716	769	722	565	335	223	-
1979	4,550	225	387	599	661	746	670	491	332	215	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All races, female											
1998	4,389	116	226	296	377	478	676	685	642	713	-
1997	4,380	147	234	292	387	540	618	678	614	692	-
1996	4,297	146	195	275	387	533	670	670	615	640	-
1995	4,444	133	207	298	414	540	687	675	616	680	-
1994	4,228	135	187	267	394	580	671	662	590	569	-
1993	3,964	120	198	248	381	535	624	623	564	515	-
1992	4,078	126	181	252	410	516	637	662	570	568	-
1991	4,091	109	186	323	423	560	610	634	553	526	-
1990	3,789	110	175	253	373	562	606	596	464	474	-
1989	3,759	114	174	262	403	559	571	573	492	444	-
1988	3,588	111	157	265	413	497	579	501	482	419	-
1987	3,628	119	152	269	413	526	598	571	428	403	-
1986	3,577	108	165	280	447	461	568	558	436	364	1
1985	3,372	89	163	262	420	458	563	482	400	353	-
1984	3,210	78	172	263	374	456	502	520	354	331	-
1983	3,195	92	169	274	396	436	516	460	346	348	-
1982	3,071	99	155	258	374	460	491	448	332	300	-
1981	2,974	95	159	281	309	450	477	422	334	287	-
1980	2,786	89	171	256	359	432	436	375	290	231	-
1979	2,734	90	166	274	341	376	427	370	293	232	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
White, both sexes											
1998	10,286	389	642	871	1,034	1,340	1,619	1,543	1,294	1,168	-
1997	10,069	413	632	793	1,037	1,376	1,538	1,549	1,218	1,168	-
1996	9,946	391	591	818	1,053	1,386	1,601	1,468	1,210	1,062	-
1995	9,927	387	598	783	1,058	1,355	1,559	1,539	1,197	1,083	-
1994	9,637	366	566	748	1,078	1,412	1,538	1,469	1,117	995	1
1993	9,280	347	579	703	1,069	1,357	1,560	1,360	1,063	886	-
1992	9,441	383	518	773	1,085	1,360	1,530	1,440	1,084	930	2
1991	9,220	309	517	765	1,151	1,379	1,464	1,350	1,055	871	-
1990	8,854	310	496	758	1,037	1,398	1,425	1,328	982	779	-
1989	8,712	344	459	733	1,087	1,338	1,418	1,273	970	738	-
1988	8,334	314	461	687	1,108	1,277	1,362	1,171	896	701	-
1987	8,515	296	476	800	1,130	1,322	1,369	1,253	849	678	-
1986	8,165	281	487	761	1,142	1,202	1,316	1,145	863	636	1
1985	7,833	274	470	757	1,087	1,170	1,269	1,081	781	598	-
1984	7,663	247	484	769	993	1,167	1,283	1,077	735	586	-
1983	7,533	256	476	762	1,057	1,197	1,177	1,042	669	583	-
1982	7,364	278	473	774	977	1,151	1,163	1,022	665	540	-
1981	7,034	254	473	757	954	1,067	1,091	962	638	502	-
1980	6,934	262	510	752	989	1,115	1,080	879	590	437	-
1979	6,692	270	500	801	917	1,040	1,033	807	597	425	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
White, male											
1998	6,337	295	442	611	696	912	1,004	922	692	512	-
1997	6,150	284	426	542	697	896	986	936	659	517	-
1996	6,110	267	422	569	717	903	1,003	864	644	474	-
1995	5,974	276	420	517	701	874	940	926	636	462	-
1994	5,844	248	404	531	716	896	924	864	570	470	1
1993	5,738	248	410	484	726	891	1,001	789	552	394	-
1992	5,735	273	352	546	715	886	939	826	560	407	2
1991	5,552	219	356	480	762	884	926	765	553	384	-
1990	5,449	217	347	538	709	886	871	782	545	336	-
1989	5,314	249	313	502	733	831	887	736	510	326	-
1988	5,069	216	330	460	732	825	829	705	441	302	-
1987	5,192	197	347	565	749	825	817	730	444	293	-
1986	4,927	186	349	513	737	787	790	624	455	292	-
1985	4,760	197	320	522	705	756	745	642	406	259	-
1984	4,748	187	337	529	658	742	820	587	404	280	-
1983	4,600	179	324	517	691	794	708	607	340	253	-
1982	4,580	196	337	546	638	739	705	603	356	254	-
1981	4,307	169	337	502	667	650	652	570	327	224	-
1980	4,370	183	352	516	659	718	673	535	318	216	-
1979	4,191	193	354	552	605	684	634	465	320	203	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
White, female											
1998	3,949	94	200	260	338	428	615	621	602	656	-
1997	3,919	129	206	251	340	480	552	613	559	651	-
1996	3,836	124	169	249	336	483	598	604	566	588	-
1995	3,953	111	178	266	357	481	619	613	561	621	-
1994	3,793	118	162	217	362	516	614	605	547	525	-
1993	3,542	99	169	219	343	466	559	571	511	492	-
1992	3,706	110	166	227	370	474	591	614	524	523	-
1991	3,668	90	161	285	389	495	538	585	502	487	-
1990	3,405	93	149	220	328	512	554	546	437	443	-
1989	3,398	95	146	231	354	507	531	537	460	412	-
1988	3,265	98	131	227	376	452	533	466	455	399	-
1987	3,323	99	129	235	381	497	552	523	405	385	-
1986	3,238	95	138	248	405	415	526	521	408	344	1
1985	3,073	77	150	235	382	414	524	439	375	339	-
1984	2,915	60	147	240	335	425	463	490	331	306	-
1983	2,933	77	152	245	366	403	469	435	329	330	-
1982	2,784	82	136	228	339	412	458	419	309	286	-
1981	2,727	85	136	255	287	417	439	392	311	278	-
1980	2,564	79	158	236	330	397	407	344	272	221	-
1979	2,501	77	146	249	312	356	399	342	277	222	-

Source: CDC/NCHS, National Vital Statistics System
See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All other, both sexes											
1998	1,198	64	112	115	117	155	172	156	87	94	-
1997	1,225	63	91	131	137	177	165	156	97	89	-
1996	1,149	67	94	89	135	142	169	143	102	91	-
1995	1,156	62	88	116	152	145	153	136	99	100	-
1994	1,113	66	84	117	114	162	168	132	80	87	-
1993	1,042	53	91	83	120	154	167	120	96	49	-
1992	988	58	70	94	113	130	154	108	87	71	-
1991	1,050	47	71	97	112	154	177	122	91	65	-
1990	989	55	75	96	133	146	142	104	70	59	-
1989	926	46	74	93	116	151	118	101	71	62	-
1988	869	39	71	96	95	123	132	94	70	41	-
1987	817	50	63	93	111	114	113	98	59	44	-
1986	822	42	60	89	111	106	106	89	63	38	-
1985	827	43	56	78	117	130	100	96	73	33	-
1984	748	46	64	80	116	98	91	79	47	39	-
1983	695	32	51	79	112	91	106	57	43	35	1
1982	720	50	52	80	94	120	100	65	49	25	-
1981	668	32	58	85	98	97	88	58	41	23	-
1980	603	30	59	89	86	86	78	61	35	17	-
1979	592	45	53	72	85	82	64	54	28	22	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All other, male											
1998	758	42	86	79	78	105	111	92	47	37	-
1997	764	45	63	90	90	117	99	91	42	48	-
1996	688	45	68	63	84	92	97	77	53	39	-
1995	665	40	59	84	95	86	85	74	44	41	-
1994	678	49	59	67	82	98	111	75	37	43	-
1993	620	32	62	54	82	85	102	68	43	26	-
1992	616	42	55	69	73	88	108	60	41	26	-
1991	627	28	46	59	78	89	105	73	40	26	-
1990	605	38	49	63	88	96	90	54	43	28	-
1989	565	27	46	62	67	99	78	65	39	30	-
1988	546	26	45	58	58	78	86	59	43	21	-
1987	512	30	40	59	79	85	67	50	36	26	-
1986	483	29	33	57	69	60	64	52	35	18	-
1985	528	31	43	51	79	86	61	53	48	19	-
1984	453	28	39	57	77	67	52	49	24	14	-
1983	433	17	34	50	82	58	59	32	26	17	1
1982	433	33	33	50	59	72	67	36	26	11	-
1981	421	22	35	59	76	64	50	28	18	14	-
1980	381	20	46	69	57	51	49	30	17	7	-
1979	359	32	33	47	56	62	36	26	12	12	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
All other, female											
1998	440	22	26	36	39	50	61	64	40	57	-
1997	461	18	28	41	47	60	66	65	55	41	-
1996	461	22	26	26	51	50	72	66	49	52	-
1995	491	22	29	32	57	59	68	62	55	59	-
1994	435	17	25	50	32	64	57	57	43	44	-
1993	422	21	29	29	38	69	65	52	53	23	-
1992	372	16	15	25	40	42	46	48	46	45	-
1991	423	19	25	38	34	65	72	49	51	39	-
1990	384	17	26	33	45	50	52	50	27	31	-
1989	361	19	28	31	49	52	40	36	32	32	-
1988	323	13	26	38	37	45	46	35	27	20	-
1987	305	20	23	34	32	29	46	48	23	18	-
1986	339	13	27	32	42	46	42	37	28	20	-
1985	299	12	13	27	38	44	39	43	25	14	-
1984	295	18	25	23	39	31	39	30	23	25	-
1983	262	15	17	29	30	33	47	25	17	18	-
1982	287	17	19	30	35	48	33	29	23	14	-
1981	247	10	23	26	22	33	38	30	23	9	-
1980	222	10	13	20	29	35	29	31	18	10	-
1979	233	13	20	25	29	20	28	28	16	10	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
Black, both sexes											
1998	990	57	90	97	99	129	137	119	72	78	-
1997	1,037	48	76	113	112	146	145	127	82	78	-
1996	974	60	78	77	120	123	143	120	73	77	-
1995	999	53	75	98	127	123	133	116	86	91	-
1994	947	58	73	99	89	135	140	116	69	77	-
1993	908	47	79	72	107	132	145	100	83	42	-
1992	854	53	58	79	101	104	136	95	74	65	-
1991	918	42	67	80	95	131	156	108	78	57	-
1990	888	51	63	87	122	131	124	99	63	46	-
1989	822	43	67	84	100	135	101	95	58	53	-
1988	775	36	65	78	85	113	118	85	58	36	-
1987	727	48	56	84	102	102	96	86	48	39	-
1986	736	38	47	81	95	96	102	78	53	34	-
1985	748	38	46	66	108	121	90	86	67	32	-
1984	664	41	54	73	105	86	82	68	41	35	-
1983	633	28	46	73	98	84	99	52	37	33	1
1982	644	44	48	76	79	107	90	59	43	22	-
1981	601	27	50	77	91	88	84	53	38	18	-
1980	551	29	54	77	84	82	69	50	31	16	-
1979	532	43	51	63	73	73	60	47	24	18	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
Black, male											
1998	632	38	68	66	67	87	89	72	43	28	-
1997	636	33	54	76	73	96	87	69	35	41	-
1996	568	39	57	55	77	77	77	61	35	30	-
1995	570	33	49	71	79	71	73	66	41	36	-
1994	573	41	52	57	66	81	93	64	33	36	-
1993	545	28	54	48	74	71	88	57	41	24	-
1992	520	38	44	57	65	70	93	49	34	23	-
1991	551	25	43	48	67	75	92	65	36	22	-
1990	542	37	41	57	82	87	77	50	39	21	-
1989	501	26	43	55	58	89	66	61	32	25	-
1988	485	24	42	46	52	71	77	51	35	19	-
1987	448	28	35	53	72	77	55	40	29	22	-
1986	437	28	29	51	60	51	62	47	29	15	-
1985	477	28	34	45	72	83	56	46	44	18	-
1984	395	26	31	53	67	59	46	42	22	11	-
1983	394	15	30	47	70	53	56	29	23	17	1
1982	386	28	30	48	50	64	61	32	23	9	-
1981	373	18	30	52	70	57	48	26	15	12	-
1980	341	19	42	58	57	49	42	21	14	6	-
1979	325	30	33	43	47	55	33	24	10	10	-
Malignant neoplasms of kidney and renal pelvis (189.0,189.1)											
Black, female											
1998	358	19	22	31	32	42	48	47	29	50	-
1997	401	15	22	37	39	50	58	58	47	37	-
1996	406	21	21	22	43	46	66	59	38	47	-
1995	429	20	26	27	48	52	60	50	45	55	-
1994	374	17	21	42	23	54	47	52	36	41	-
1993	363	19	25	24	33	61	57	43	42	18	-
1992	334	15	14	22	36	34	43	46	40	42	-
1991	367	17	24	32	28	56	64	43	42	35	-
1990	346	14	22	30	40	44	47	49	24	25	-
1989	321	17	24	29	42	46	35	34	26	28	-
1988	290	12	23	32	33	42	41	34	23	17	-
1987	279	20	21	31	30	25	41	46	19	17	-
1986	299	10	18	30	35	45	40	31	24	19	-
1985	271	10	12	21	36	38	34	40	23	14	-
1984	269	15	23	20	38	27	36	26	19	24	-
1983	239	13	16	26	28	31	43	23	14	16	-
1982	258	16	18	28	29	43	29	27	20	13	-
1981	228	9	20	25	21	31	36	27	23	6	-
1980	210	10	12	19	27	33	27	29	17	10	-
1979	207	13	18	20	26	18	27	23	14	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
All races, both sexes											
1998	11,757	155	272	445	718	1,096	1,790	2,223	2,138	2,802	-
1997	11,582	151	293	400	685	1,177	1,730	2,135	2,206	2,686	-
1996	11,452	139	253	379	692	1,219	1,815	2,087	2,125	2,630	1
1995	11,084	144	217	397	650	1,200	1,752	2,066	2,036	2,489	1
1994	11,170	132	232	432	655	1,295	1,765	2,027	2,108	2,413	-
1993	10,962	151	233	366	739	1,301	1,729	1,976	1,965	2,379	-
1992	10,707	131	208	383	784	1,205	1,660	1,988	1,928	2,302	2
1991	10,406	98	223	386	802	1,226	1,681	1,886	1,841	2,166	-
1990	10,341	84	207	371	861	1,219	1,566	1,989	1,797	2,131	1
1989	10,121	106	184	421	810	1,208	1,645	1,896	1,772	1,989	-
1988	9,805	91	187	374	768	1,218	1,672	1,786	1,700	1,924	-
1987	9,589	83	205	401	754	1,137	1,566	1,753	1,691	1,912	-
1986	9,655	101	197	445	734	1,177	1,595	1,778	1,696	1,869	1
1985	9,787	89	221	429	810	1,205	1,670	1,759	1,598	1,923	2
1984	9,716	99	225	435	798	1,171	1,626	1,801	1,622	1,870	2
1983	9,844	102	193	498	803	1,229	1,712	1,765	1,612	1,846	1
1982	9,874	100	279	500	827	1,330	1,635	1,729	1,569	1,813	-
1981	9,815	78	244	485	830	1,295	1,663	1,805	1,563	1,768	2
1980	9,816	117	195	471	870	1,321	1,744	1,709	1,622	1,704	1
1979	9,694	128	277	454	830	1,309	1,623	1,801	1,566	1,631	-
Malignant neoplasm of bladder (188)											
All races, male											
1998	7,939	118	211	332	526	812	1,289	1,542	1,408	1,626	-
1997	7,662	100	220	297	505	848	1,223	1,495	1,420	1,480	-
1996	7,683	98	180	289	496	919	1,302	1,455	1,357	1,514	1
1995	7,521	101	165	283	487	890	1,290	1,455	1,330	1,422	-
1994	7,457	90	167	321	490	949	1,256	1,415	1,375	1,326	-
1993	7,474	100	174	280	556	970	1,266	1,376	1,318	1,351	-
1992	7,123	102	158	277	575	870	1,172	1,348	1,263	1,284	1
1991	7,027	68	162	284	612	904	1,238	1,289	1,218	1,191	-
1990	6,910	58	161	269	633	867	1,147	1,385	1,135	1,174	-
1989	6,843	73	129	321	598	878	1,199	1,357	1,144	1,087	-
1988	6,534	61	139	283	588	871	1,197	1,224	1,113	1,013	-
1987	6,490	56	129	298	577	863	1,124	1,248	1,125	1,015	-
1986	6,523	63	147	332	528	887	1,143	1,199	1,140	1,037	1
1985	6,679	64	155	327	609	900	1,189	1,232	1,041	1,107	2
1984	6,601	71	164	310	596	878	1,179	1,251	1,043	1,062	1
1983	6,735	66	142	361	622	893	1,233	1,213	1,085	1,068	1
1982	6,796	67	208	364	615	1,013	1,174	1,254	1,013	1,025	-
1981	6,711	58	188	366	617	947	1,235	1,231	1,001	1,013	2
1980	6,708	67	134	356	619	963	1,300	1,206	1,061	965	1
1979	6,709	92	206	333	609	979	1,227	1,265	1,014	939	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
All races, female											
1998	3,818	37	61	113	192	284	501	681	730	1,176	-
1997	3,920	51	73	103	180	329	507	640	786	1,206	-
1996	3,769	41	73	90	196	300	513	632	768	1,116	-
1995	3,563	43	52	114	163	310	462	611	706	1,067	1
1994	3,713	42	65	111	165	346	509	612	733	1,087	-
1993	3,488	51	59	86	183	331	463	600	647	1,028	-
1992	3,584	29	50	106	209	335	488	640	665	1,018	1
1991	3,379	30	61	102	190	322	443	597	623	975	-
1990	3,431	26	46	102	228	352	419	604	662	957	1
1989	3,278	33	55	100	212	330	446	539	628	902	-
1988	3,271	30	48	91	180	347	475	562	587	911	-
1987	3,099	27	76	103	177	274	442	505	566	897	-
1986	3,132	38	50	113	206	290	452	579	556	832	-
1985	3,108	25	66	102	201	305	481	527	557	816	-
1984	3,115	28	61	125	202	293	447	550	579	808	1
1983	3,109	36	51	137	181	336	479	552	527	778	-
1982	3,078	33	71	136	212	317	461	475	556	788	-
1981	3,104	20	56	119	213	348	428	574	562	755	-
1980	3,108	50	61	115	251	358	444	503	561	739	-
1979	2,985	36	71	121	221	330	396	536	552	692	-
Malignant neoplasm of bladder (188)											
White, both sexes											
1998	10,762	130	230	390	629	1,004	1,640	2,045	1,998	2,602	-
1997	10,608	134	262	354	606	1,069	1,579	1,970	2,047	2,483	-
1996	10,463	121	223	318	621	1,095	1,650	1,935	1,978	2,431	-
1995	10,151	126	193	361	562	1,089	1,610	1,925	1,886	2,296	-
1994	10,194	118	197	383	584	1,178	1,623	1,855	1,945	2,225	-
1993	10,025	131	191	316	666	1,187	1,587	1,811	1,814	2,225	-
1992	9,766	116	182	329	709	1,087	1,514	1,837	1,768	2,125	1
1991	9,492	90	185	325	725	1,124	1,516	1,725	1,706	2,016	-
1990	9,430	68	179	326	763	1,096	1,423	1,799	1,686	1,993	1
1989	9,193	92	157	371	729	1,084	1,475	1,730	1,647	1,830	-
1988	8,951	81	155	322	691	1,099	1,551	1,628	1,572	1,785	-
1987	8,742	79	170	348	675	1,029	1,422	1,602	1,554	1,794	-
1986	8,849	84	170	388	662	1,074	1,444	1,628	1,597	1,749	1
1985	8,949	74	189	381	726	1,087	1,513	1,619	1,488	1,803	-
1984	8,849	86	201	390	689	1,051	1,467	1,642	1,507	1,758	2
1983	8,999	90	163	434	720	1,104	1,556	1,626	1,502	1,740	1
1982	9,004	86	241	429	737	1,177	1,476	1,598	1,475	1,710	-
1981	8,973	68	205	428	734	1,157	1,500	1,673	1,458	1,682	1
1980	9,018	92	171	420	782	1,189	1,584	1,591	1,523	1,608	-
1979	8,862	107	238	385	743	1,177	1,495	1,633	1,476	1,545	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
White, male											
1998	7,397	99	186	294	478	764	1,202	1,444	1,332	1,539	-
1997	7,139	92	201	267	461	785	1,134	1,404	1,343	1,387	-
1996	7,147	87	159	249	460	842	1,207	1,372	1,284	1,430	-
1995	7,022	89	147	259	432	825	1,216	1,384	1,263	1,328	-
1994	6,925	80	147	294	448	873	1,179	1,318	1,284	1,248	-
1993	6,936	90	153	246	511	910	1,176	1,272	1,236	1,276	-
1992	6,593	90	142	246	531	807	1,089	1,266	1,166	1,196	-
1991	6,517	64	137	252	564	841	1,142	1,192	1,146	1,126	-
1990	6,403	49	145	239	580	795	1,060	1,264	1,084	1,116	-
1989	6,308	66	108	289	557	809	1,094	1,253	1,072	1,012	-
1988	6,077	55	123	253	543	804	1,129	1,131	1,049	956	-
1987	6,006	53	111	268	524	798	1,045	1,160	1,045	957	-
1986	6,093	51	132	306	485	830	1,061	1,107	1,094	985	1
1985	6,175	58	133	298	557	827	1,091	1,145	974	1,045	-
1984	6,093	65	150	285	531	799	1,082	1,152	982	1,008	1
1983	6,232	59	123	326	572	818	1,131	1,134	1,021	1,009	1
1982	6,295	60	186	325	570	905	1,082	1,172	973	972	-
1981	6,211	55	162	337	554	865	1,139	1,144	943	968	1
1980	6,270	58	119	321	570	881	1,210	1,141	1,011	925	-
1979	6,203	77	181	292	560	896	1,134	1,166	965	892	-
Malignant neoplasm of bladder (188)											
White, female											
1998	3,365	31	44	96	151	240	438	601	666	1,063	-
1997	3,469	42	61	87	145	284	445	566	704	1,096	-
1996	3,316	34	64	69	161	253	443	563	694	1,001	-
1995	3,129	37	46	102	130	264	394	541	623	968	-
1994	3,269	38	50	89	136	305	444	537	661	977	-
1993	3,089	41	38	70	155	277	411	539	578	949	-
1992	3,173	26	40	83	178	280	425	571	602	929	1
1991	2,975	26	48	73	161	283	374	533	560	890	-
1990	3,027	19	34	87	183	301	363	535	602	877	1
1989	2,885	26	49	82	172	275	381	477	575	818	-
1988	2,874	26	32	69	148	295	422	497	523	829	-
1987	2,736	26	59	80	151	231	377	442	509	837	-
1986	2,756	33	38	82	177	244	383	521	503	764	-
1985	2,774	16	56	83	169	260	422	474	514	758	-
1984	2,756	21	51	105	158	252	385	490	525	750	1
1983	2,767	31	40	108	148	286	425	492	481	731	-
1982	2,709	26	55	104	167	272	394	426	502	738	-
1981	2,762	13	43	91	180	292	361	529	515	714	-
1980	2,748	34	52	99	212	308	374	450	512	683	-
1979	2,659	30	57	93	183	281	361	467	511	653	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
All other, both sexes											
1998	995	25	42	55	89	92	150	178	140	200	-
1997	974	17	31	46	79	108	151	165	159	203	-
1996	989	18	30	61	71	124	165	152	147	199	1
1995	933	18	24	36	88	111	142	141	150	193	1
1994	976	14	35	49	71	117	142	172	163	188	-
1993	937	20	42	50	73	114	142	165	151	154	-
1992	941	15	26	54	75	118	146	151	160	177	1
1991	914	8	38	61	77	102	165	161	135	150	-
1990	911	16	28	45	98	123	143	190	111	138	-
1989	928	14	27	50	81	124	170	166	125	159	-
1988	854	10	32	52	77	119	121	158	128	139	-
1987	847	4	35	53	79	108	144	151	137	118	-
1986	806	17	27	57	72	103	151	150	99	120	-
1985	838	15	32	48	84	118	157	140	110	120	2
1984	867	13	24	45	109	120	159	159	115	112	-
1983	845	12	30	64	83	125	156	139	110	106	-
1982	870	14	38	71	90	153	159	131	94	103	-
1981	842	10	39	57	96	138	163	132	105	86	1
1980	798	25	24	51	88	132	160	118	99	96	1
1979	832	21	39	69	87	132	128	168	90	86	-
Malignant neoplasm of bladder (188)											
All other, male											
1998	542	19	25	38	48	48	87	98	76	87	-
1997	523	8	19	30	44	63	89	91	77	93	-
1996	536	11	21	40	36	77	95	83	73	84	1
1995	499	12	18	24	55	65	74	71	67	94	-
1994	532	10	20	27	42	76	77	97	91	78	-
1993	538	10	21	34	45	60	90	104	82	75	-
1992	530	12	16	31	44	63	83	82	97	88	1
1991	510	4	25	32	48	63	96	97	72	65	-
1990	507	9	16	30	53	72	87	121	51	58	-
1989	535	7	21	32	41	69	105	104	72	75	-
1988	457	6	16	30	45	67	68	93	64	57	-
1987	484	3	18	30	53	65	79	88	80	58	-
1986	430	12	15	26	43	57	82	92	46	52	-
1985	504	6	22	29	52	73	98	87	67	62	2
1984	508	6	14	25	65	79	97	99	61	54	-
1983	503	7	19	35	50	75	102	79	64	59	-
1982	501	7	22	39	45	108	92	82	40	53	-
1981	500	3	26	29	63	82	96	87	58	45	1
1980	438	9	15	35	49	82	90	65	50	40	1
1979	506	15	25	41	49	83	93	99	49	47	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
All other, female											
1998	453	6	17	17	41	44	63	80	64	113	-
1997	451	9	12	16	35	45	62	74	82	110	-
1996	453	7	9	21	35	47	70	69	74	115	-
1995	434	6	6	12	33	46	68	70	83	99	1
1994	444	4	15	22	29	41	65	75	72	110	-
1993	399	10	21	16	28	54	52	61	69	79	-
1992	411	3	10	23	31	55	63	69	63	89	-
1991	404	4	13	29	29	39	69	64	63	85	-
1990	404	7	12	15	45	51	56	69	60	80	-
1989	393	7	6	18	40	55	65	62	53	84	-
1988	397	4	16	22	32	52	53	65	64	82	-
1987	363	1	17	23	26	43	65	63	57	60	-
1986	376	5	12	31	29	46	69	58	53	68	-
1985	334	9	10	19	32	45	59	53	43	58	-
1984	359	7	10	20	44	41	62	60	54	58	-
1983	342	5	11	29	33	50	54	60	46	47	-
1982	369	7	16	32	45	45	67	49	54	50	-
1981	342	7	13	28	33	56	67	45	47	41	-
1980	360	16	9	16	39	50	70	53	49	56	-
1979	326	6	14	28	38	49	35	69	41	39	-
Malignant neoplasm of bladder (188)											
Black, both sexes											
1998	875	23	37	50	74	76	130	160	122	182	-
1997	862	17	29	38	74	99	128	147	138	181	-
1996	885	14	29	54	60	110	147	138	134	178	1
1995	845	14	21	31	76	102	132	131	137	174	1
1994	879	13	34	40	60	105	133	156	147	171	-
1993	838	17	39	47	64	106	125	148	130	138	-
1992	848	15	25	47	71	104	134	134	141	161	1
1991	841	8	37	58	68	91	152	146	123	144	-
1990	847	15	26	44	93	114	134	177	103	124	-
1989	853	14	21	48	75	113	157	159	115	143	-
1988	782	8	28	48	71	112	108	146	117	127	-
1987	785	4	34	46	75	97	138	139	129	106	-
1986	752	16	25	56	67	95	143	142	92	108	-
1985	785	14	31	46	81	112	148	131	104	104	2
1984	814	11	21	43	103	113	152	146	111	105	-
1983	802	12	27	60	82	121	152	131	99	101	-
1982	824	12	36	66	87	149	150	124	88	96	-
1981	807	10	38	56	94	134	156	124	100	82	-
1980	756	21	21	49	86	128	154	114	92	86	1
1979	785	21	39	65	83	129	119	160	83	76	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasm of bladder (188)											
Black, male											
1998	453	17	21	34	37	37	68	87	64	73	-
1997	456	8	17	24	42	57	72	80	67	84	-
1996	466	8	21	36	28	66	86	73	65	68	1
1995	441	11	16	19	47	59	66	66	58	82	-
1994	474	9	19	25	37	67	71	88	79	68	-
1993	470	7	19	31	40	54	74	95	70	64	-
1992	469	12	15	27	41	51	74	74	86	78	1
1991	471	4	25	32	41	57	90	90	65	60	-
1990	466	9	15	30	49	66	79	110	47	52	-
1989	484	7	17	31	36	59	98	99	66	65	-
1988	411	5	13	29	41	61	60	86	56	50	-
1987	444	3	17	25	50	57	76	82	74	50	-
1986	397	11	13	26	38	53	78	86	42	47	-
1985	465	6	21	28	49	69	90	79	63	52	2
1984	466	5	11	23	62	74	91	89	57	48	-
1983	475	7	16	34	50	71	100	72	58	56	-
1982	470	6	20	35	44	106	86	79	35	47	-
1981	474	3	25	29	62	78	91	80	54	43	-
1980	411	7	13	34	47	80	86	61	47	33	1
1979	482	15	25	37	47	81	88	95	46	44	-
Malignant neoplasm of bladder (188)											
Black, female											
1998	422	6	16	16	37	39	62	73	58	109	-
1997	406	9	12	14	32	42	56	67	71	97	-
1996	419	6	8	18	32	44	61	65	69	110	-
1995	404	3	5	12	29	43	66	65	79	92	1
1994	405	4	15	15	23	38	62	68	68	103	-
1993	368	10	20	16	24	52	51	53	60	74	-
1992	379	3	10	20	30	53	60	60	55	83	-
1991	370	4	12	26	27	34	62	56	58	84	-
1990	381	6	11	14	44	48	55	67	56	72	-
1989	369	7	4	17	39	54	59	60	49	78	-
1988	371	3	15	19	30	51	48	60	61	77	-
1987	341	1	17	21	25	40	62	57	55	56	-
1986	355	5	12	30	29	42	65	56	50	61	-
1985	320	8	10	18	32	43	58	52	41	52	-
1984	348	6	10	20	41	39	61	57	54	57	-
1983	327	5	11	26	32	50	52	59	41	45	-
1982	354	6	16	31	43	43	64	45	53	49	-
1981	333	7	13	27	32	56	65	44	46	39	-
1980	345	14	8	15	39	48	68	53	45	53	-
1979	303	6	14	28	36	48	31	65	37	32	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All races, both sexes											
1998	12,667	823	997	1,135	1,239	1,542	1,687	1,467	966	639	1
1997	12,474	782	959	1,075	1,259	1,522	1,672	1,479	944	619	-
1996	12,376	806	967	1,083	1,295	1,565	1,661	1,388	864	566	1
1995	12,063	851	856	1,022	1,238	1,567	1,651	1,340	877	543	1
1994	12,313	764	886	1,051	1,299	1,633	1,743	1,306	863	510	-
1993	11,993	723	856	1,054	1,212	1,600	1,696	1,325	816	496	-
1992	11,941	723	867	1,002	1,331	1,639	1,621	1,219	847	474	-
1991	11,952	677	836	1,059	1,428	1,597	1,512	1,272	823	469	-
1990	11,630	635	784	1,005	1,403	1,679	1,524	1,285	733	430	-
1989	11,130	548	790	1,009	1,397	1,560	1,484	1,198	698	365	1
1988	10,943	606	754	1,066	1,357	1,520	1,444	1,122	627	366	-
1987	10,833	561	773	1,057	1,439	1,552	1,429	1,072	582	340	-
1986	10,322	511	692	997	1,351	1,502	1,356	997	582	282	-
1985	10,265	512	720	1,042	1,424	1,385	1,370	983	523	281	-
1984	10,116	510	694	1,137	1,406	1,473	1,262	973	480	259	-
1983	9,674	464	733	1,094	1,358	1,377	1,174	877	448	228	1
1982	9,624	462	746	1,101	1,328	1,373	1,162	862	459	204	-
1981	9,382	456	764	1,158	1,339	1,324	1,149	777	404	180	-
1980	9,300	500	822	1,115	1,259	1,338	1,160	730	365	143	-
1979	8,936	528	802	1,098	1,207	1,308	1,041	671	307	116	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All races, male											
1998	6,936	503	594	676	703	869	886	736	448	229	1
1997	6,740	486	552	599	705	838	870	752	394	242	-
1996	6,733	498	571	642	726	851	901	675	385	208	1
1995	6,537	522	503	601	698	854	838	644	408	202	1
1994	6,702	455	549	635	747	880	925	618	365	171	-
1993	6,551	443	500	652	680	877	873	674	357	183	-
1992	6,432	430	522	585	760	888	843	591	360	183	-
1991	6,431	404	511	610	802	857	767	628	347	177	-
1990	6,339	389	487	570	817	938	788	615	324	144	-
1989	6,046	340	482	575	812	818	764	585	321	153	-
1988	6,076	377	446	619	795	889	761	554	279	147	-
1987	5,824	325	449	607	769	855	761	479	255	120	-
1986	5,669	334	418	581	771	844	688	479	242	110	-
1985	5,580	321	427	609	781	721	701	477	236	109	-
1984	5,515	308	406	652	761	826	661	469	210	97	-
1983	5,242	283	445	634	731	752	582	406	191	86	1
1982	5,224	268	433	622	736	731	575	437	209	76	-
1981	5,134	273	445	674	760	708	596	361	182	74	-
1980	5,242	307	490	668	726	752	588	353	164	54	-
1979	4,972	318	469	624	650	735	565	337	137	44	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All races, female											
1998	5,731	320	403	459	536	673	801	731	518	410	-
1997	5,734	296	407	476	554	684	802	727	550	377	-
1996	5,643	308	396	441	569	714	760	713	479	358	-
1995	5,526	329	353	421	540	713	813	696	469	341	-
1994	5,611	309	337	416	552	753	818	688	498	339	-
1993	5,442	280	356	402	532	723	823	651	459	313	-
1992	5,509	293	345	417	571	751	778	628	487	291	-
1991	5,521	273	325	449	626	740	745	644	476	292	-
1990	5,291	246	297	435	586	741	736	670	409	286	-
1989	5,084	208	308	434	585	742	720	613	377	212	1
1988	4,867	229	308	447	562	631	683	568	348	219	-
1987	5,009	236	324	450	670	697	668	593	327	220	-
1986	4,653	177	274	416	580	658	668	518	340	172	-
1985	4,685	191	293	433	643	664	669	506	287	172	-
1984	4,601	202	288	485	645	647	601	504	270	162	-
1983	4,432	181	288	460	627	625	592	471	257	142	-
1982	4,400	194	313	479	592	642	587	425	250	128	-
1981	4,248	183	319	484	579	616	553	416	222	106	-
1980	4,058	193	332	447	533	586	572	377	201	89	-
1979	3,964	210	333	474	557	573	476	334	170	72	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
White, both sexes											
1998	11,718	746	926	1,061	1,155	1,452	1,582	1,389	913	611	1
1997	11,518	718	884	987	1,157	1,433	1,568	1,405	909	577	-
1996	11,523	744	897	1,019	1,216	1,473	1,561	1,321	828	541	1
1995	11,165	780	805	920	1,159	1,476	1,567	1,273	828	514	-
1994	11,465	707	835	975	1,227	1,548	1,647	1,244	823	484	-
1993	11,132	657	802	989	1,142	1,490	1,598	1,252	778	468	-
1992	11,132	657	793	935	1,258	1,565	1,543	1,162	811	448	-
1991	11,133	634	770	989	1,340	1,508	1,428	1,215	787	448	-
1990	10,814	580	724	939	1,311	1,597	1,452	1,224	698	404	-
1989	10,372	512	727	947	1,307	1,485	1,398	1,140	674	349	1
1988	10,161	554	698	996	1,271	1,444	1,357	1,065	602	347	-
1987	10,109	518	719	970	1,354	1,475	1,363	1,023	561	324	-
1986	9,605	470	638	928	1,278	1,420	1,288	949	551	271	-
1985	9,612	466	660	975	1,352	1,305	1,315	942	502	275	-
1984	9,476	471	646	1,052	1,330	1,403	1,202	933	460	249	-
1983	9,071	445	683	1,030	1,264	1,309	1,123	841	427	218	-
1982	9,001	421	696	1,028	1,247	1,307	1,122	825	435	196	-
1981	8,772	425	716	1,094	1,266	1,257	1,094	749	389	169	-
1980	8,727	475	777	1,048	1,184	1,272	1,107	703	356	138	-
1979	8,353	484	741	1,043	1,153	1,240	991	635	296	111	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
White, male											
1998	6,415	452	552	635	654	822	831	695	428	220	1
1997	6,263	451	513	552	660	791	818	723	381	228	-
1996	6,287	464	532	610	682	803	852	641	368	200	1
1995	6,067	481	475	545	656	805	793	610	388	193	-
1994	6,246	429	511	581	704	845	874	592	349	165	-
1993	6,089	406	467	610	637	814	822	649	339	175	-
1992	6,014	397	478	550	725	852	806	562	346	176	-
1991	6,002	382	472	569	757	801	733	602	333	168	-
1990	5,890	355	450	535	766	892	742	589	308	139	-
1989	5,641	324	446	544	762	779	713	557	307	145	-
1988	5,659	341	419	581	743	839	728	528	270	139	-
1987	5,426	307	418	550	723	805	732	456	247	117	-
1986	5,288	306	389	542	726	798	647	454	229	106	-
1985	5,212	297	387	571	739	676	673	456	230	106	-
1984	5,145	285	372	597	718	786	632	451	202	95	-
1983	4,925	275	426	595	686	714	553	388	178	83	-
1982	4,883	246	398	576	691	692	558	416	199	74	-
1981	4,780	256	417	644	714	667	568	344	175	69	-
1980	4,933	295	461	632	685	713	562	338	158	52	-
1979	4,655	287	433	595	625	699	543	315	131	42	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
White, female											
1998	5,303	294	374	426	501	630	751	694	485	391	-
1997	5,255	267	371	435	497	642	750	682	528	349	-
1996	5,236	280	365	409	534	670	709	680	460	341	-
1995	5,098	299	330	375	503	671	774	663	440	321	-
1994	5,219	278	324	394	523	703	773	652	474	319	-
1993	5,043	251	335	379	505	676	776	603	439	293	-
1992	5,118	260	315	385	533	713	737	600	465	272	-
1991	5,131	252	298	420	583	707	695	613	454	280	-
1990	4,924	225	274	404	545	705	710	635	390	265	-
1989	4,731	188	281	403	545	706	685	583	367	204	1
1988	4,502	213	279	415	528	605	629	537	332	208	-
1987	4,683	211	301	420	631	670	631	567	314	207	-
1986	4,317	164	249	386	552	622	641	495	322	165	-
1985	4,400	169	273	404	613	629	642	486	272	169	-
1984	4,331	186	274	455	612	617	570	482	258	154	-
1983	4,146	170	257	435	578	595	570	453	249	135	-
1982	4,118	175	298	452	556	615	564	409	236	122	-
1981	3,992	169	299	450	552	590	526	405	214	100	-
1980	3,794	180	316	416	499	559	545	365	198	86	-
1979	3,698	197	308	448	528	541	448	320	165	69	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All other, both sexes											
1998	949	77	71	74	84	90	105	78	53	28	-
1997	956	64	75	88	102	89	104	74	35	42	-
1996	853	62	70	64	79	92	100	67	36	25	-
1995	898	71	51	102	79	91	84	67	49	29	1
1994	848	57	51	76	72	85	96	62	40	26	-
1993	861	66	54	65	70	110	98	73	38	28	-
1992	809	66	74	67	73	74	78	57	36	26	-
1991	819	43	66	70	88	89	84	57	36	21	-
1990	816	55	60	66	92	82	72	61	35	26	-
1989	758	36	63	62	90	75	86	58	24	16	-
1988	782	52	56	70	86	76	87	57	25	19	-
1987	724	43	54	87	85	77	66	49	21	16	-
1986	717	41	54	69	73	82	68	48	31	11	-
1985	653	46	60	67	72	80	55	41	21	6	-
1984	640	39	48	85	76	70	60	40	20	10	-
1983	603	19	50	64	94	68	51	36	21	10	1
1982	623	41	50	73	81	66	40	37	24	8	-
1981	610	31	48	64	73	67	55	28	15	11	-
1980	573	25	45	67	75	66	53	27	9	5	-
1979	583	44	61	55	54	68	50	36	11	5	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All other, male											
1998	521	51	42	41	49	47	55	41	20	9	-
1997	477	35	39	47	45	47	52	29	13	14	-
1996	446	34	39	32	44	48	49	34	17	8	-
1995	470	41	28	56	42	49	45	34	20	9	1
1994	456	26	38	54	43	35	51	26	16	6	-
1993	462	37	33	42	43	63	51	25	18	8	-
1992	418	33	44	35	35	36	37	29	14	7	-
1991	429	22	39	41	45	56	34	26	14	9	-
1990	449	34	37	35	51	46	46	26	16	5	-
1989	405	16	36	31	50	39	51	28	14	8	-
1988	417	36	27	38	52	50	33	26	9	8	-
1987	398	18	31	57	46	50	29	23	8	3	-
1986	381	28	29	39	45	46	41	25	13	4	-
1985	368	24	40	38	42	45	28	21	6	3	-
1984	370	23	34	55	43	40	29	18	8	2	-
1983	317	8	19	39	45	38	29	18	13	3	1
1982	341	22	35	46	45	39	17	21	10	2	-
1981	354	17	28	30	46	41	28	17	7	5	-
1980	309	12	29	36	41	39	26	15	6	2	-
1979	317	31	36	29	25	36	22	22	6	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
All other, female											
1998	428	26	29	33	35	43	50	37	33	19	-
1997	479	29	36	41	57	42	52	45	22	28	-
1996	407	28	31	32	35	44	51	33	19	17	-
1995	428	30	23	46	37	42	39	33	29	20	-
1994	392	31	13	22	29	50	45	36	24	20	-
1993	399	29	21	23	27	47	47	48	20	20	-
1992	391	33	30	32	38	38	41	28	22	19	-
1991	390	21	27	29	43	33	50	31	22	12	-
1990	367	21	23	31	41	36	26	35	19	21	-
1989	353	20	27	31	40	36	35	30	10	8	-
1988	365	16	29	32	34	26	54	31	16	11	-
1987	326	25	23	30	39	27	37	26	13	13	-
1986	336	13	25	30	28	36	27	23	18	7	-
1985	285	22	20	29	30	35	27	20	15	3	-
1984	270	16	14	30	33	30	31	22	12	8	-
1983	286	11	31	25	49	30	22	18	8	7	-
1982	282	19	15	27	36	27	23	16	14	6	-
1981	256	14	20	34	27	26	27	11	8	6	-
1980	264	13	16	31	34	27	27	12	3	3	-
1979	266	13	25	26	29	32	28	14	5	3	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
Black, both sexes											
1998	747	60	54	68	67	75	87	60	38	25	-
1997	765	45	59	73	84	72	90	61	26	35	-
1996	695	44	56	53	68	79	80	57	28	22	-
1995	720	55	37	78	57	79	70	54	42	23	1
1994	685	45	42	63	61	69	76	50	36	21	-
1993	697	54	44	52	64	95	79	60	31	21	-
1992	680	51	62	58	62	67	67	48	34	25	-
1991	681	32	51	54	75	77	74	48	32	18	-
1990	691	47	50	55	78	71	64	54	31	22	-
1989	634	31	56	49	78	64	73	49	21	11	-
1988	669	43	46	59	80	63	74	50	22	16	-
1987	641	40	49	74	77	69	61	45	19	15	-
1986	616	35	51	57	66	70	60	40	25	9	-
1985	559	41	55	59	64	73	48	32	17	5	-
1984	557	35	45	77	70	64	56	31	14	8	-
1983	541	17	48	61	82	60	47	33	18	10	1
1982	563	37	46	67	74	62	39	32	21	7	-
1981	542	29	45	57	63	60	51	22	14	10	-
1980	527	25	41	64	70	62	49	24	9	5	-
1979	539	41	58	54	47	61	49	33	9	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
Black, male											
1998	399	40	33	37	41	38	44	29	14	7	-
1997	381	25	29	38	39	37	47	24	8	11	-
1996	365	23	32	29	37	40	41	27	13	7	-
1995	377	32	19	41	29	42	38	27	18	7	1
1994	357	17	31	46	35	27	38	19	16	4	-
1993	379	31	27	34	41	52	44	21	14	7	-
1992	351	29	39	30	29	33	32	24	14	7	-
1991	350	18	30	33	37	46	31	21	12	7	-
1990	372	31	30	27	43	38	42	21	12	4	-
1989	335	14	31	23	44	33	42	25	12	5	-
1988	358	32	24	33	50	40	28	22	8	8	-
1987	351	17	28	50	43	46	27	19	6	3	-
1986	328	24	27	36	41	37	38	21	9	3	-
1985	309	22	37	33	38	40	24	14	5	2	-
1984	320	21	31	51	41	35	28	11	6	2	-
1983	277	8	18	36	40	32	25	16	10	3	1
1982	311	20	33	45	40	37	17	19	8	2	-
1981	308	15	25	24	38	38	25	14	6	5	-
1980	285	12	25	33	37	39	24	13	6	2	-
1979	292	30	35	29	19	31	21	20	4	2	-
Malignant neoplasms of meninges, brain and other parts of central nervous system (191-192)											
Black, female											
1998	348	20	21	31	26	37	43	31	24	18	-
1997	384	20	30	35	45	35	43	37	18	24	-
1996	330	21	24	24	31	39	39	30	15	15	-
1995	343	23	18	37	28	37	32	27	24	16	-
1994	328	28	11	17	26	42	38	31	20	17	-
1993	318	23	17	18	23	43	35	39	17	14	-
1992	329	22	23	28	33	34	35	24	20	18	-
1991	331	14	21	21	38	31	43	27	20	11	-
1990	319	16	20	28	35	33	22	33	19	18	-
1989	299	17	25	26	34	31	31	24	9	6	-
1988	311	11	22	26	30	23	46	28	14	8	-
1987	290	23	21	24	34	23	34	26	13	12	-
1986	288	11	24	21	25	33	22	19	16	6	-
1985	250	19	18	26	26	33	24	18	12	3	-
1984	237	14	14	26	29	29	28	20	8	6	-
1983	264	9	30	25	42	28	22	17	8	7	-
1982	252	17	13	22	34	25	22	13	13	5	-
1981	234	14	20	33	25	22	26	8	8	5	-
1980	242	13	16	31	33	23	25	11	3	3	-
1979	247	11	23	25	28	30	28	13	5	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All races, both sexes											
1998	55,538	1,740	2,332	3,048	4,184	5,949	8,449	9,493	8,045	7,735	2
1997	55,835	1,776	2,420	3,155	4,294	6,277	8,481	9,110	7,882	7,608	-
1996	54,930	1,778	2,255	3,001	4,279	6,240	8,501	8,949	7,740	7,313	1
1995	54,367	1,722	2,216	2,957	4,304	6,332	8,443	8,763	7,420	7,150	2
1994	53,080	1,735	2,240	2,877	4,414	6,254	8,227	8,296	7,199	6,825	1
1993	51,627	1,582	2,044	2,732	4,304	6,407	7,884	8,180	6,860	6,512	2
1992	50,384	1,579	1,991	2,763	4,230	6,274	7,623	7,932	6,610	6,082	1
1991	49,590	1,563	1,879	2,783	4,336	6,166	7,570	7,771	6,393	5,934	2
1990	47,741	1,409	1,909	2,739	4,281	6,108	7,143	7,325	6,178	5,538	3
1989	46,751	1,429	1,755	2,766	4,421	5,966	7,078	7,132	5,913	5,152	2
1988	44,762	1,343	1,783	2,733	4,214	5,746	6,580	6,796	5,602	4,951	1
1987	43,599	1,294	1,740	2,770	4,204	5,639	6,561	6,500	5,264	4,750	6
1986	43,531	1,235	1,757	2,768	4,208	5,804	6,518	6,460	5,094	4,485	1
1985	42,478	1,212	1,747	2,864	4,442	5,507	6,253	6,194	4,847	4,370	3
1984	41,202	1,202	1,837	2,976	4,300	5,304	6,046	5,768	4,716	4,061	1
1983	40,311	1,192	1,845	2,891	4,177	5,242	5,945	5,734	4,370	3,985	6
1982	39,461	1,153	1,940	2,810	4,029	5,096	5,658	5,615	4,313	3,803	2
1981	37,826	1,132	1,824	2,847	4,022	4,875	5,401	5,326	4,122	3,387	8
1980	37,971	1,190	1,927	3,037	3,950	4,772	5,547	4,993	4,029	3,368	1
1979	36,445	1,220	2,008	2,842	3,782	4,740	5,205	4,811	3,805	3,121	1
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All races, male											
1998	29,537	1,062	1,435	1,775	2,470	3,458	4,680	4,931	3,853	3,031	1
1997	29,754	1,120	1,476	1,835	2,468	3,656	4,688	4,755	3,841	2,941	-
1996	29,158	1,108	1,326	1,771	2,483	3,524	4,694	4,617	3,674	2,935	1
1995	28,887	1,048	1,349	1,708	2,527	3,629	4,592	4,480	3,507	2,819	-
1994	28,151	1,043	1,353	1,668	2,632	3,531	4,528	4,266	3,326	2,681	-
1993	27,146	957	1,223	1,626	2,426	3,651	4,230	4,176	3,165	2,472	1
1992	26,779	951	1,194	1,635	2,468	3,556	4,141	4,091	3,076	2,324	1
1991	25,977	968	1,107	1,636	2,467	3,450	4,091	3,869	2,889	2,224	1
1990	25,504	841	1,182	1,635	2,540	3,428	3,906	3,748	2,861	2,158	1
1989	24,801	872	1,036	1,631	2,533	3,360	3,802	3,690	2,693	1,945	1
1988	23,792	817	1,060	1,628	2,442	3,180	3,547	3,475	2,564	1,954	1
1987	22,939	760	1,027	1,614	2,400	3,212	3,509	3,288	2,353	1,752	5
1986	23,151	746	998	1,602	2,374	3,261	3,576	3,251	2,352	1,730	1
1985	22,542	704	1,014	1,619	2,535	3,084	3,361	3,116	2,244	1,748	1
1984	21,667	708	1,042	1,705	2,434	2,903	3,252	2,838	2,133	1,598	1
1983	21,509	687	1,063	1,696	2,408	2,896	3,214	2,924	1,963	1,663	3
1982	21,029	674	1,128	1,583	2,348	2,867	3,024	2,817	1,987	1,533	1
1981	20,100	661	1,052	1,638	2,289	2,687	2,918	2,614	1,915	1,376	3
1980	20,341	667	1,147	1,742	2,236	2,652	2,978	2,502	1,842	1,433	1
1979	19,826	735	1,134	1,640	2,170	2,710	2,817	2,441	1,764	1,402	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All races, female											
1998	26,001	678	897	1,273	1,714	2,491	3,769	4,562	4,192	4,704	1
1997	26,081	656	944	1,320	1,826	2,621	3,793	4,355	4,041	4,667	-
1996	25,772	670	929	1,230	1,796	2,716	3,807	4,332	4,066	4,378	-
1995	25,480	674	867	1,249	1,777	2,703	3,851	4,283	3,913	4,331	2
1994	24,929	692	887	1,209	1,782	2,723	3,699	4,030	3,873	4,144	1
1993	24,481	625	821	1,106	1,878	2,756	3,654	4,004	3,695	4,040	1
1992	23,605	628	797	1,128	1,762	2,718	3,482	3,841	3,534	3,758	-
1991	23,613	595	772	1,147	1,869	2,716	3,479	3,902	3,504	3,710	1
1990	22,237	568	727	1,104	1,741	2,680	3,237	3,577	3,317	3,380	2
1989	21,950	557	719	1,135	1,888	2,606	3,276	3,442	3,220	3,207	1
1988	20,970	526	723	1,105	1,772	2,566	3,033	3,321	3,038	2,997	-
1987	20,660	534	713	1,156	1,804	2,427	3,052	3,212	2,911	2,998	1
1986	20,380	489	759	1,166	1,834	2,543	2,942	3,209	2,742	2,755	-
1985	19,936	508	733	1,245	1,907	2,423	2,892	3,078	2,603	2,622	2
1984	19,535	494	795	1,271	1,866	2,401	2,794	2,930	2,583	2,463	-
1983	18,802	505	782	1,195	1,769	2,346	2,731	2,810	2,407	2,322	3
1982	18,432	479	812	1,227	1,681	2,229	2,634	2,798	2,326	2,270	1
1981	17,726	471	772	1,209	1,733	2,188	2,483	2,712	2,207	2,011	5
1980	17,630	523	780	1,295	1,714	2,120	2,569	2,491	2,187	1,935	-
1979	16,619	485	874	1,202	1,612	2,030	2,388	2,370	2,041	1,719	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
White, both sexes											
1998	49,365	1,405	1,941	2,545	3,600	5,270	7,642	8,723	7,443	7,147	1
1997	49,769	1,464	2,045	2,701	3,707	5,596	7,668	8,374	7,323	7,071	-
1996	48,912	1,484	1,919	2,566	3,695	5,531	7,658	8,218	7,155	6,773	1
1995	48,557	1,456	1,847	2,543	3,706	5,672	7,697	8,040	6,889	6,645	1
1994	47,532	1,443	1,919	2,461	3,832	5,617	7,511	7,672	6,651	6,330	-
1993	46,071	1,321	1,725	2,341	3,764	5,720	7,156	7,505	6,355	6,020	1
1992	45,120	1,358	1,676	2,346	3,719	5,648	6,951	7,294	6,130	5,663	1
1991	44,288	1,343	1,586	2,372	3,835	5,486	6,861	7,107	5,897	5,539	2
1990	42,755	1,191	1,624	2,346	3,772	5,467	6,500	6,736	5,737	5,173	2
1989	41,906	1,212	1,484	2,415	3,910	5,344	6,453	6,564	5,472	4,792	2
1988	40,197	1,133	1,523	2,359	3,793	5,138	6,004	6,243	5,184	4,615	1
1987	39,147	1,101	1,485	2,445	3,744	5,075	5,949	5,953	4,893	4,413	2
1986	39,102	1,071	1,500	2,412	3,723	5,242	5,909	5,944	4,704	4,185	1
1985	38,227	1,030	1,538	2,510	3,955	4,975	5,664	5,731	4,445	4,102	2
1984	37,154	1,022	1,588	2,610	3,843	4,832	5,503	5,307	4,397	3,837	1
1983	36,303	1,028	1,594	2,571	3,709	4,730	5,423	5,260	4,054	3,741	6
1982	35,613	1,007	1,676	2,486	3,603	4,612	5,153	5,172	3,992	3,592	1
1981	34,109	981	1,607	2,523	3,630	4,370	4,911	4,900	3,856	3,174	4
1980	34,386	1,029	1,692	2,671	3,556	4,308	5,106	4,607	3,776	3,182	-
1979	33,069	1,070	1,761	2,520	3,399	4,270	4,800	4,477	3,589	2,930	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
White, male											
1998	26,348	865	1,207	1,513	2,153	3,092	4,286	4,562	3,581	2,793	-
1997	26,532	931	1,260	1,589	2,141	3,292	4,267	4,380	3,579	2,731	-
1996	25,978	928	1,132	1,528	2,162	3,146	4,248	4,266	3,398	2,716	1
1995	25,853	898	1,130	1,495	2,203	3,276	4,201	4,129	3,270	2,608	-
1994	25,227	893	1,164	1,431	2,299	3,185	4,152	3,976	3,086	2,479	-
1993	24,216	814	1,046	1,406	2,155	3,267	3,848	3,829	2,958	2,271	-
1992	23,980	831	1,008	1,404	2,181	3,226	3,789	3,770	2,857	2,160	1
1991	23,178	832	942	1,407	2,205	3,093	3,732	3,527	2,671	2,065	1
1990	22,812	711	1,022	1,427	2,252	3,069	3,561	3,457	2,651	1,996	1
1989	22,146	744	877	1,426	2,262	3,025	3,475	3,397	2,474	1,784	1
1988	21,375	688	913	1,423	2,213	2,869	3,243	3,190	2,380	1,811	1
1987	20,562	651	891	1,433	2,158	2,891	3,199	2,995	2,170	1,608	2
1986	20,751	653	869	1,401	2,113	2,942	3,244	2,992	2,144	1,611	1
1985	20,268	601	892	1,431	2,263	2,802	3,055	2,890	2,046	1,632	1
1984	19,486	598	911	1,506	2,178	2,656	2,971	2,593	1,981	1,506	1
1983	19,327	585	929	1,514	2,124	2,623	2,946	2,677	1,810	1,546	3
1982	18,953	595	976	1,406	2,119	2,607	2,756	2,589	1,818	1,433	-
1981	18,139	575	938	1,465	2,076	2,429	2,665	2,402	1,774	1,285	1
1980	18,386	594	1,016	1,529	2,026	2,367	2,735	2,310	1,713	1,349	-
1979	17,992	648	1,010	1,463	1,944	2,460	2,603	2,267	1,653	1,316	1
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
White, female											
1998	23,017	540	734	1,032	1,447	2,178	3,356	4,161	3,862	4,354	1
1997	23,237	533	785	1,112	1,566	2,304	3,401	3,994	3,744	4,340	-
1996	22,934	556	787	1,038	1,533	2,385	3,410	3,952	3,757	4,057	-
1995	22,704	558	717	1,048	1,503	2,396	3,496	3,911	3,619	4,037	1
1994	22,305	550	755	1,030	1,533	2,432	3,359	3,696	3,565	3,851	-
1993	21,855	507	679	935	1,609	2,453	3,308	3,676	3,397	3,749	1
1992	21,140	527	668	942	1,538	2,422	3,162	3,524	3,273	3,503	-
1991	21,110	511	644	965	1,630	2,393	3,129	3,580	3,226	3,474	1
1990	19,943	480	602	919	1,520	2,398	2,939	3,279	3,086	3,177	1
1989	19,760	468	607	989	1,648	2,319	2,978	3,167	2,998	3,008	1
1988	18,822	445	610	936	1,580	2,269	2,761	3,053	2,804	2,804	-
1987	18,585	450	594	1,012	1,586	2,184	2,750	2,958	2,723	2,805	-
1986	18,351	418	631	1,011	1,610	2,300	2,665	2,952	2,560	2,574	-
1985	17,959	429	646	1,079	1,692	2,173	2,609	2,841	2,399	2,470	1
1984	17,668	424	677	1,104	1,665	2,176	2,532	2,714	2,416	2,331	-
1983	16,976	443	665	1,057	1,585	2,107	2,477	2,583	2,244	2,195	3
1982	16,660	412	700	1,080	1,484	2,005	2,397	2,583	2,174	2,159	1
1981	15,970	406	669	1,058	1,554	1,941	2,246	2,498	2,082	1,889	3
1980	16,000	435	676	1,142	1,530	1,941	2,371	2,297	2,063	1,833	-
1979	15,077	422	751	1,057	1,455	1,810	2,197	2,210	1,936	1,614	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All other, both sexes											
1998	6,173	335	391	503	584	679	807	770	602	588	1
1997	6,066	312	375	454	587	681	813	736	559	537	-
1996	6,018	294	336	435	584	709	843	731	585	540	-
1995	5,810	266	369	414	598	660	746	723	531	505	1
1994	5,548	292	321	416	582	637	716	624	548	495	1
1993	5,556	261	319	391	540	687	728	675	505	492	1
1992	5,264	221	315	417	511	626	672	638	480	419	-
1991	5,302	220	293	411	501	680	709	664	496	395	-
1990	4,986	218	285	393	509	641	643	589	441	365	1
1989	4,845	217	271	351	511	622	625	568	441	360	-
1988	4,565	210	260	374	421	608	576	553	418	336	-
1987	4,452	193	255	325	460	564	612	547	371	337	4
1986	4,429	164	257	356	485	562	609	516	390	300	-
1985	4,251	182	209	354	487	532	589	463	402	268	1
1984	4,048	180	249	366	457	472	543	461	319	224	-
1983	4,008	164	251	320	468	512	522	474	316	244	-
1982	3,848	146	264	324	426	484	505	443	321	211	1
1981	3,717	151	217	324	392	505	490	426	266	213	4
1980	3,585	161	235	366	394	464	441	386	253	186	1
1979	3,376	150	247	322	383	470	405	334	216	191	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All other, male											
1998	3,189	197	228	262	317	366	394	369	272	238	1
1997	3,222	189	216	246	327	364	421	375	262	210	-
1996	3,180	180	194	243	321	378	446	351	276	219	-
1995	3,034	150	219	213	324	353	391	351	237	211	-
1994	2,924	150	189	237	333	346	376	290	240	202	-
1993	2,930	143	177	220	271	384	382	347	207	201	1
1992	2,799	120	186	231	287	330	352	321	219	164	-
1991	2,799	136	165	229	262	357	359	342	218	159	-
1990	2,692	130	160	208	288	359	345	291	210	162	-
1989	2,655	128	159	205	271	335	327	293	219	161	-
1988	2,417	129	147	205	229	311	304	285	184	143	-
1987	2,377	109	136	181	242	321	310	293	183	144	3
1986	2,400	93	129	201	261	319	332	259	208	119	-
1985	2,274	103	122	188	272	282	306	226	198	116	-
1984	2,181	110	131	199	256	247	281	245	152	92	-
1983	2,182	102	134	182	284	273	268	247	153	117	-
1982	2,076	79	152	177	229	260	268	228	169	100	1
1981	1,961	86	114	173	213	258	253	212	141	91	2
1980	1,955	73	131	213	210	285	243	192	129	84	1
1979	1,834	87	124	177	226	250	214	174	111	86	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
All other, female											
1998	2,984	138	163	241	267	313	413	401	330	350	-
1997	2,844	123	159	208	260	317	392	361	297	327	-
1996	2,838	114	142	192	263	331	397	380	309	321	-
1995	2,776	116	150	201	274	307	355	372	294	294	1
1994	2,624	142	132	179	249	291	340	334	308	293	1
1993	2,626	118	142	171	269	303	346	328	298	291	-
1992	2,465	101	129	186	224	296	320	317	261	255	-
1991	2,503	84	128	182	239	323	350	322	278	236	-
1990	2,294	88	125	185	221	282	298	298	231	203	1
1989	2,190	89	112	146	240	287	298	275	222	199	-
1988	2,148	81	113	169	192	297	272	268	234	193	-
1987	2,075	84	119	144	218	243	302	254	188	193	1
1986	2,029	71	128	155	224	243	277	257	182	181	-
1985	1,977	79	87	166	215	250	283	237	204	152	1
1984	1,867	70	118	167	201	225	262	216	167	132	-
1983	1,826	62	117	138	184	239	254	227	163	127	-
1982	1,772	67	112	147	197	224	237	215	152	111	-
1981	1,756	65	103	151	179	247	237	214	125	122	2
1980	1,630	88	104	153	184	179	198	194	124	102	-
1979	1,542	63	123	145	157	220	191	160	105	105	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
Black, both sexes											
1998	5,222	295	332	441	509	569	674	627	524	495	1
1997	5,128	271	311	373	505	576	695	608	487	467	-
1996	5,148	247	293	383	518	612	691	630	515	467	-
1995	4,958	238	316	357	524	573	641	620	447	432	-
1994	4,746	254	274	366	494	565	621	543	466	432	1
1993	4,791	224	284	338	479	604	628	589	441	419	1
1992	4,575	188	275	364	449	554	578	554	428	375	-
1991	4,633	185	250	363	434	600	633	579	445	360	-
1990	4,364	190	256	351	454	563	576	520	395	327	1
1989	4,235	193	235	307	440	557	545	501	398	329	-
1988	4,022	179	224	337	374	543	519	493	370	307	-
1987	3,958	177	229	286	423	509	554	493	330	305	4
1986	3,973	142	230	318	446	514	553	459	353	276	-
1985	3,774	154	179	318	435	489	539	419	363	242	1
1984	3,643	159	212	333	415	431	489	418	298	200	-
1983	3,617	147	226	291	434	473	483	429	293	218	-
1982	3,494	130	237	296	398	444	472	404	293	190	1
1981	3,393	136	191	299	366	464	462	392	251	195	3
1980	3,271	145	214	335	369	419	410	352	243	174	1
1979	3,111	134	235	305	362	441	375	310	202	172	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
Black, male											
1998	2,677	173	199	227	275	309	324	292	234	188	1
1997	2,693	163	180	195	278	308	361	301	217	177	-
1996	2,700	155	167	208	286	331	348	300	236	186	-
1995	2,586	137	193	186	286	308	330	296	199	168	-
1994	2,473	131	161	211	279	320	320	249	188	167	-
1993	2,494	121	162	189	249	337	326	301	170	155	1
1992	2,391	100	157	201	251	293	297	272	187	138	-
1991	2,416	112	141	206	226	314	310	292	188	142	-
1990	2,346	114	146	191	258	314	306	251	187	142	-
1989	2,302	120	138	178	228	297	283	247	197	143	-
1988	2,121	112	127	185	200	274	270	253	156	134	-
1987	2,116	101	122	162	223	293	286	261	156	127	3
1986	2,124	81	112	182	238	292	296	225	177	106	-
1985	1,999	90	105	166	237	263	278	200	170	104	-
1984	1,953	98	112	180	234	225	244	221	146	80	-
1983	1,947	91	119	164	262	251	241	221	141	105	-
1982	1,865	73	135	163	208	235	250	201	148	91	1
1981	1,772	76	102	156	199	230	234	190	132	83	2
1980	1,772	64	119	195	197	258	221	166	123	76	1
1979	1,673	78	115	166	215	232	196	156	105	78	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (200-208)											
Black, female											
1998	2,545	122	133	214	234	260	350	335	290	307	-
1997	2,435	108	131	178	227	268	334	307	270	290	-
1996	2,448	92	126	175	232	281	343	330	279	281	-
1995	2,372	101	123	171	238	265	311	324	248	264	-
1994	2,273	123	113	155	215	245	301	294	278	265	1
1993	2,297	103	122	149	230	267	302	288	271	264	-
1992	2,184	88	118	163	198	261	281	282	241	237	-
1991	2,217	73	109	157	208	286	323	287	257	218	-
1990	2,018	76	110	160	196	249	270	269	208	185	1
1989	1,933	73	97	129	212	260	262	254	201	186	-
1988	1,901	67	97	152	174	269	249	240	214	173	-
1987	1,842	76	107	124	200	216	268	232	174	178	1
1986	1,849	61	118	136	208	222	257	234	176	170	-
1985	1,775	64	74	152	198	226	261	219	193	138	1
1984	1,690	61	100	153	181	206	245	197	152	120	-
1983	1,670	56	107	127	172	222	242	208	152	113	-
1982	1,629	57	102	133	190	209	222	203	145	99	-
1981	1,621	60	89	143	167	234	228	202	119	112	1
1980	1,499	81	95	140	172	161	189	186	120	98	-
1979	1,438	56	120	139	147	209	179	154	97	94	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
All races, both sexes											
1998	1,311	73	85	74	71	102	146	128	113	68	-
1997	1,443	84	75	83	80	119	132	140	112	106	-
1996	1,408	99	77	83	79	118	129	124	108	81	-
1995	1,431	92	81	83	71	105	147	141	111	82	-
1994	1,440	99	90	68	90	131	152	126	100	83	-
1993	1,574	78	80	88	96	128	157	153	125	96	-
1992	1,639	109	88	80	100	130	151	137	114	76	-
1991	1,625	82	80	90	103	145	155	137	118	77	-
1990	1,632	76	70	100	114	135	141	136	114	70	-
1989	1,721	99	77	95	113	155	148	155	94	80	-
1988	1,608	75	92	74	118	149	146	148	87	54	-
1987	1,755	105	89	110	134	162	148	166	82	78	-
1986	1,828	75	80	106	132	189	191	152	87	71	-
1985	1,778	84	74	107	146	154	162	133	109	72	-
1984	1,960	106	101	122	168	174	192	136	105	73	-
1983	1,892	90	111	117	161	160	170	158	113	67	1
1982	1,930	89	95	136	168	174	179	155	121	77	-
1981	1,997	88	112	145	159	196	175	195	102	72	-
1980	2,064	108	150	159	174	190	192	158	105	73	-
1979	2,021	113	139	147	167	192	195	153	113	56	-
Hodgkin's disease (201)											
All races, male											
1998	767	50	52	47	40	60	79	67	58	23	-
1997	760	58	45	45	40	70	67	65	50	31	-
1996	764	51	44	54	40	55	73	59	46	33	-
1995	799	57	55	54	35	57	78	51	51	35	-
1994	773	60	66	46	54	74	87	54	44	25	-
1993	900	49	61	59	56	81	84	77	50	43	-
1992	949	67	65	45	67	78	90	66	54	32	-
1991	928	59	61	60	56	90	79	66	44	29	-
1990	956	56	46	69	75	66	81	56	44	30	-
1989	1,007	65	48	61	69	93	80	78	37	32	-
1988	945	54	73	42	80	81	73	66	43	27	-
1987	1,023	72	55	76	77	94	81	78	41	23	-
1986	1,069	47	57	68	81	99	103	80	36	27	-
1985	1,033	58	51	63	97	82	87	55	46	23	-
1984	1,204	76	77	78	111	106	101	79	48	32	-
1983	1,093	60	68	78	105	90	93	75	51	25	1
1982	1,090	54	60	82	105	87	109	60	48	27	-
1981	1,122	53	75	87	102	101	90	82	45	27	-
1980	1,174	75	94	88	96	105	113	78	46	26	-
1979	1,202	90	96	89	100	104	100	74	51	25	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
All races, female											
1998	544	23	33	27	31	42	67	61	55	45	-
1997	683	26	30	38	40	49	65	75	62	75	-
1996	644	48	33	29	39	63	56	65	62	48	-
1995	632	35	26	29	36	48	69	90	60	47	-
1994	667	39	24	22	36	57	65	72	56	58	-
1993	674	29	19	29	40	47	73	76	75	53	-
1992	690	42	23	35	33	52	61	71	60	44	-
1991	697	23	19	30	47	55	76	71	74	48	-
1990	676	20	24	31	39	69	60	80	70	40	-
1989	714	34	29	34	44	62	68	77	57	48	-
1988	663	21	19	32	38	68	73	82	44	27	-
1987	732	33	34	34	57	68	67	88	41	55	-
1986	759	28	23	38	51	90	88	72	51	44	-
1985	745	26	23	44	49	72	75	78	63	49	-
1984	756	30	24	44	57	68	91	57	57	41	-
1983	799	30	43	39	56	70	77	83	62	42	-
1982	840	35	35	54	63	87	70	95	73	50	-
1981	875	35	37	58	57	95	85	113	57	45	-
1980	890	33	56	71	78	85	79	80	59	47	-
1979	819	23	43	58	67	88	95	79	62	31	-
Hodgkin's disease (201)											
White, both sexes											
1998	1,148	61	76	66	57	97	135	119	106	66	-
1997	1,262	74	61	72	69	115	124	131	108	103	-
1996	1,249	90	69	72	72	109	120	118	104	76	-
1995	1,262	84	70	72	62	101	142	132	105	80	-
1994	1,263	82	72	58	85	120	141	121	93	80	-
1993	1,405	71	71	77	85	118	145	142	119	88	-
1992	1,466	98	81	68	93	121	144	126	108	73	-
1991	1,422	70	71	77	89	132	145	128	113	75	-
1990	1,471	67	59	90	108	117	134	130	109	69	-
1989	1,533	86	69	80	96	140	141	150	89	74	-
1988	1,445	67	78	71	105	138	136	138	84	50	-
1987	1,574	96	82	93	116	148	139	158	76	71	-
1986	1,659	68	66	91	116	177	179	145	82	68	-
1985	1,588	75	67	89	129	144	149	123	103	69	-
1984	1,757	86	90	112	155	159	178	122	102	71	-
1983	1,730	83	97	111	152	150	159	151	108	66	1
1982	1,769	78	85	127	157	158	169	147	111	75	-
1981	1,820	74	104	126	147	180	160	183	102	70	-
1980	1,891	99	135	139	156	179	184	146	103	71	-
1979	1,854	105	130	131	155	179	186	144	109	52	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
White, male											
1998	668	40	46	43	32	58	70	63	55	23	-
1997	661	55	36	39	33	70	61	61	50	30	-
1996	672	46	41	48	34	50	67	54	42	33	-
1995	692	53	45	45	31	54	75	46	48	34	-
1994	668	52	51	37	52	69	81	52	39	25	-
1993	794	44	55	51	50	73	76	71	47	38	-
1992	847	64	59	40	62	72	85	61	48	29	-
1991	806	51	52	49	48	84	74	63	44	28	-
1990	853	47	40	63	70	61	75	50	43	30	-
1989	892	56	42	53	60	85	75	76	35	29	-
1988	851	47	65	40	72	76	68	62	41	24	-
1987	915	65	50	63	66	84	77	75	38	20	-
1986	972	42	50	59	71	91	97	78	35	26	-
1985	922	52	46	53	87	80	81	49	43	21	-
1984	1,066	62	70	70	100	97	93	69	48	31	-
1983	994	54	61	72	98	83	83	73	49	24	1
1982	1,005	48	55	75	98	79	106	56	44	27	-
1981	1,025	45	71	78	97	92	80	76	45	27	-
1980	1,066	69	81	75	84	97	109	73	45	24	-
1979	1,093	83	88	77	91	96	96	69	50	23	-
Hodgkin's disease (201)											
White, female											
1998	480	21	30	23	25	39	65	56	51	43	-
1997	601	19	25	33	36	45	63	70	58	73	-
1996	577	44	28	24	38	59	53	64	62	43	-
1995	570	31	25	27	31	47	67	86	57	46	-
1994	595	30	21	21	33	51	60	69	54	55	-
1993	611	27	16	26	35	45	69	71	72	50	-
1992	619	34	22	28	31	49	59	65	60	44	-
1991	616	19	19	28	41	48	71	65	69	47	-
1990	618	20	19	27	38	56	59	80	66	39	-
1989	641	30	27	27	36	55	66	74	54	45	-
1988	594	20	13	31	33	62	68	76	43	26	-
1987	659	31	32	30	50	64	62	83	38	51	-
1986	687	26	16	32	45	86	82	67	47	42	-
1985	666	23	21	36	42	64	68	74	60	48	-
1984	691	24	20	42	55	62	85	53	54	40	-
1983	736	29	36	39	54	67	76	78	59	42	-
1982	764	30	30	52	59	79	63	91	67	48	-
1981	795	29	33	48	50	88	80	107	57	43	-
1980	825	30	54	64	72	82	75	73	58	47	-
1979	761	22	42	54	64	83	90	75	59	29	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
All other, both sexes											
1998	163	12	9	8	14	5	11	9	7	2	-
1997	181	10	14	11	11	4	8	9	4	3	-
1996	159	9	8	11	7	9	9	6	4	5	-
1995	169	8	11	11	9	4	5	9	6	2	-
1994	177	17	18	10	5	11	11	5	7	3	-
1993	169	7	9	11	11	10	12	11	6	8	-
1992	173	11	7	12	7	9	7	11	6	3	-
1991	203	12	9	13	14	13	10	9	5	2	-
1990	161	9	11	10	6	18	7	6	5	1	-
1989	188	13	8	15	17	15	7	5	5	6	-
1988	163	8	14	3	13	11	10	10	3	4	-
1987	181	9	7	17	18	14	9	8	6	7	-
1986	169	7	14	15	16	12	12	7	5	3	-
1985	190	9	7	18	17	10	13	10	6	3	-
1984	203	20	11	10	13	15	14	14	3	2	-
1983	162	7	14	6	9	10	11	7	5	1	-
1982	161	11	10	9	11	16	10	8	10	2	-
1981	177	14	8	19	12	16	15	12	-	2	-
1980	173	9	15	20	18	11	8	12	2	2	-
1979	167	8	9	16	12	13	9	9	4	4	-
Hodgkin's disease (201)											
All other, male											
1998	99	10	6	4	8	2	9	4	3	-	-
1997	99	3	9	6	7	-	6	4	-	1	-
1996	92	5	3	6	6	5	6	5	4	-	-
1995	107	4	10	9	4	3	3	5	3	1	-
1994	105	8	15	9	2	5	6	2	5	-	-
1993	106	5	6	8	6	8	8	6	3	5	-
1992	102	3	6	5	5	6	5	5	6	3	-
1991	122	8	9	11	8	6	5	3	-	1	-
1990	103	9	6	6	5	5	6	6	1	-	-
1989	115	9	6	8	9	8	5	2	2	3	-
1988	94	7	8	2	8	5	5	4	2	3	-
1987	108	7	5	13	11	10	4	3	3	3	-
1986	97	5	7	9	10	8	6	2	1	1	-
1985	111	6	5	10	10	2	6	6	3	2	-
1984	138	14	7	8	11	9	8	10	-	1	-
1983	99	6	7	6	7	7	10	2	2	1	-
1982	85	6	5	7	7	8	3	4	4	-	-
1981	97	8	4	9	5	9	10	6	-	-	-
1980	108	6	13	13	12	8	4	5	1	2	-
1979	109	7	8	12	9	8	4	5	1	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
All other, female											
1998	64	2	3	4	6	3	2	5	4	2	-
1997	82	7	5	5	4	4	2	5	4	2	-
1996	67	4	5	5	1	4	3	1	-	5	-
1995	62	4	1	2	5	1	2	4	3	1	-
1994	72	9	3	1	3	6	5	3	2	3	-
1993	63	2	3	3	5	2	4	5	3	3	-
1992	71	8	1	7	2	3	2	6	-	-	-
1991	81	4	-	2	6	7	5	6	5	1	-
1990	58	-	5	4	1	13	1	-	4	1	-
1989	73	4	2	7	8	7	2	3	3	3	-
1988	69	1	6	1	5	6	5	6	1	1	-
1987	73	2	2	4	7	4	5	5	3	4	-
1986	72	2	7	6	6	4	6	5	4	2	-
1985	79	3	2	8	7	8	7	4	3	1	-
1984	65	6	4	2	2	6	6	4	3	1	-
1983	63	1	7	-	2	3	1	5	3	-	-
1982	76	5	5	2	4	8	7	4	6	2	-
1981	80	6	4	10	7	7	5	6	-	2	-
1980	65	3	2	7	6	3	4	7	1	-	-
1979	58	1	1	4	3	5	5	4	3	2	-
Hodgkin's disease (201)											
Black, both sexes											
1998	147	12	8	7	13	4	7	6	7	2	-
1997	164	8	14	11	9	4	8	7	3	3	-
1996	142	9	8	11	7	8	7	3	3	4	-
1995	157	8	11	11	8	4	5	7	5	2	-
1994	152	15	17	9	5	8	10	5	5	2	-
1993	154	7	8	10	9	8	11	10	5	7	-
1992	163	11	6	12	5	9	6	10	3	2	-
1991	184	11	9	12	12	10	9	7	4	2	-
1990	148	8	11	9	6	15	6	5	5	1	-
1989	173	12	7	15	12	13	7	4	4	5	-
1988	150	7	14	3	13	11	7	8	2	2	-
1987	161	9	7	14	14	13	5	7	5	7	-
1986	155	6	12	14	16	12	11	5	4	3	-
1985	174	8	6	18	16	8	10	9	4	2	-
1984	187	19	10	9	12	14	12	10	3	2	-
1983	152	6	12	5	9	10	11	5	5	1	-
1982	151	11	10	6	11	15	9	7	10	2	-
1981	161	14	8	17	11	14	14	10	-	2	-
1980	160	8	15	17	17	11	6	11	2	2	-
1979	159	8	9	14	12	12	8	9	4	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hodgkin's disease (201)											
Black, male											
1998	91	10	5	4	7	2	6	4	3	-	-
1997	92	3	9	6	6	-	6	3	-	1	-
1996	83	5	3	6	6	5	5	2	3	-	-
1995	98	4	10	9	3	3	3	3	2	1	-
1994	87	7	14	8	2	3	6	2	3	-	-
1993	94	5	5	7	4	7	7	5	2	4	-
1992	92	3	5	5	3	6	4	4	3	2	-
1991	110	7	9	10	7	3	5	3	-	1	-
1990	96	8	6	5	5	5	5	5	1	-	-
1989	107	9	5	8	5	7	5	1	2	3	-
1988	84	6	8	2	8	5	2	3	1	2	-
1987	98	7	5	11	9	9	3	2	2	3	-
1986	91	4	5	9	10	8	6	2	-	1	-
1985	104	5	5	10	10	2	6	6	1	1	-
1984	127	13	6	7	10	8	7	7	-	1	-
1983	90	5	5	5	7	7	10	1	2	1	-
1982	77	6	5	4	7	7	3	3	4	-	-
1981	85	8	4	7	4	7	10	4	-	-	-
1980	101	5	13	11	12	8	2	4	1	2	-
1979	102	7	8	10	9	7	3	5	1	2	-
Hodgkin's disease (201)											
Black, female											
1998	56	2	3	3	6	2	1	2	4	2	-
1997	72	5	5	5	3	4	2	4	3	2	-
1996	59	4	5	5	1	3	2	1	-	4	-
1995	59	4	1	2	5	1	2	4	3	1	-
1994	65	8	3	1	3	5	4	3	2	2	-
1993	60	2	3	3	5	1	4	5	3	3	-
1992	71	8	1	7	2	3	2	6	-	-	-
1991	74	4	-	2	5	7	4	4	4	1	-
1990	52	-	5	4	1	10	1	-	4	1	-
1989	66	3	2	7	7	6	2	3	2	2	-
1988	66	1	6	1	5	6	5	5	1	-	-
1987	63	2	2	3	5	4	2	5	3	4	-
1986	64	2	7	5	6	4	5	3	4	2	-
1985	70	3	1	8	6	6	4	3	3	1	-
1984	60	6	4	2	2	6	5	3	3	1	-
1983	62	1	7	-	2	3	1	4	3	-	-
1982	74	5	5	2	4	8	6	4	6	2	-
1981	76	6	4	10	7	7	4	6	-	2	-
1980	59	3	2	6	5	3	4	7	1	-	-
1979	57	1	1	4	3	5	5	4	3	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
All races, both sexes											
1998	23,544	796	1,086	1,368	1,852	2,557	3,542	4,052	3,503	3,189	1
1997	23,693	817	1,142	1,387	1,886	2,671	3,555	3,980	3,387	3,054	-
1996	22,934	827	1,014	1,328	1,825	2,590	3,555	3,775	3,282	2,897	1
1995	22,522	786	1,048	1,322	1,749	2,655	3,530	3,653	3,012	2,845	-
1994	21,930	791	970	1,267	1,885	2,662	3,459	3,452	3,005	2,599	-
1993	20,631	716	868	1,190	1,744	2,558	3,234	3,291	2,812	2,443	-
1992	20,179	709	875	1,132	1,788	2,545	3,147	3,229	2,608	2,290	1
1991	19,700	744	796	1,178	1,786	2,491	3,041	3,051	2,576	2,212	1
1990	18,601	629	821	1,143	1,734	2,489	2,809	2,901	2,383	2,053	-
1989	18,197	638	728	1,137	1,786	2,390	2,801	2,825	2,340	1,927	1
1988	17,259	570	748	1,120	1,737	2,314	2,608	2,647	2,144	1,784	1
1987	16,419	520	691	1,130	1,653	2,228	2,607	2,523	2,042	1,644	2
1986	16,351	533	732	1,113	1,687	2,343	2,550	2,468	1,849	1,489	-
1985	15,562	500	704	1,148	1,775	2,132	2,323	2,344	1,749	1,427	1
1984	14,692	449	708	1,148	1,652	2,032	2,280	2,082	1,693	1,309	1
1983	14,298	470	720	1,173	1,604	1,987	2,271	2,040	1,545	1,235	-
1982	13,803	454	795	1,141	1,564	1,885	2,054	1,992	1,492	1,141	1
1981	12,827	408	721	1,110	1,556	1,787	1,835	1,814	1,352	1,037	1
1980	12,761	445	753	1,172	1,449	1,724	1,993	1,703	1,319	973	1
1979	12,022	443	753	1,070	1,435	1,750	1,806	1,599	1,168	869	-
Non-Hodgkin's lymphoma (200,202)											
All races, male											
1998	12,279	523	696	815	1,101	1,436	1,862	1,978	1,570	1,225	-
1997	12,369	556	734	838	1,069	1,502	1,874	1,982	1,502	1,066	-
1996	11,962	548	612	808	1,079	1,408	1,877	1,839	1,478	1,060	1
1995	11,694	514	652	765	1,037	1,483	1,820	1,721	1,330	1,012	-
1994	11,373	511	593	748	1,097	1,422	1,779	1,686	1,282	986	-
1993	10,560	469	531	693	969	1,365	1,612	1,577	1,219	872	-
1992	10,542	463	560	673	1,028	1,358	1,625	1,567	1,145	819	1
1991	10,157	486	497	724	987	1,341	1,511	1,421	1,086	796	-
1990	9,795	409	527	700	1,038	1,328	1,450	1,401	1,039	746	-
1989	9,420	408	459	683	1,013	1,260	1,440	1,334	968	710	1
1988	8,892	372	452	679	1,001	1,215	1,324	1,241	887	655	1
1987	8,348	320	421	671	898	1,221	1,314	1,185	813	554	2
1986	8,485	351	438	654	936	1,269	1,313	1,131	778	519	-
1985	8,132	321	415	643	982	1,166	1,185	1,162	750	526	-
1984	7,450	284	397	669	909	1,038	1,174	925	696	484	1
1983	7,382	275	420	684	906	1,025	1,191	971	609	454	-
1982	7,141	280	466	620	911	991	1,034	946	599	419	-
1981	6,523	249	433	621	839	925	926	814	548	387	-
1980	6,573	250	467	675	812	905	981	742	550	361	1
1979	6,398	277	433	592	797	981	929	766	508	353	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
All races, female											
1998	11,265	273	390	553	751	1,121	1,680	2,074	1,933	1,964	1
1997	11,324	261	408	549	817	1,169	1,681	1,998	1,885	1,988	-
1996	10,972	279	402	520	746	1,182	1,678	1,936	1,804	1,837	-
1995	10,828	272	396	557	712	1,172	1,710	1,932	1,682	1,833	-
1994	10,557	280	377	519	788	1,240	1,680	1,766	1,723	1,613	-
1993	10,071	247	337	497	775	1,193	1,622	1,714	1,593	1,571	-
1992	9,637	246	315	459	760	1,187	1,522	1,662	1,463	1,471	-
1991	9,543	258	299	454	799	1,150	1,530	1,630	1,490	1,416	1
1990	8,806	220	294	443	696	1,161	1,359	1,500	1,344	1,307	-
1989	8,777	230	269	454	773	1,130	1,361	1,491	1,372	1,217	-
1988	8,367	198	296	441	736	1,099	1,284	1,406	1,257	1,129	-
1987	8,071	200	270	459	755	1,007	1,293	1,338	1,229	1,090	-
1986	7,866	182	294	459	751	1,074	1,237	1,337	1,071	970	-
1985	7,430	179	289	505	793	966	1,138	1,182	999	901	1
1984	7,242	165	311	479	743	994	1,106	1,157	997	825	-
1983	6,916	195	300	489	698	962	1,080	1,069	936	781	-
1982	6,662	174	329	521	653	894	1,020	1,046	893	722	1
1981	6,304	159	288	489	717	862	909	1,000	804	650	1
1980	6,188	195	286	497	637	819	1,012	961	769	612	-
1979	5,624	166	320	478	638	769	877	833	660	516	-
Non-Hodgkin's lymphoma (200,202)											
White, both sexes											
1998	21,603	662	937	1,207	1,682	2,343	3,308	3,822	3,330	3,041	1
1997	21,696	685	987	1,241	1,695	2,463	3,317	3,751	3,220	2,900	-
1996	21,062	704	890	1,190	1,648	2,399	3,316	3,577	3,098	2,762	1
1995	20,780	691	914	1,202	1,580	2,457	3,335	3,459	2,877	2,705	-
1994	20,269	670	878	1,136	1,692	2,493	3,264	3,298	2,845	2,477	-
1993	19,030	614	752	1,065	1,609	2,360	3,057	3,122	2,671	2,329	-
1992	18,608	626	782	1,016	1,621	2,373	2,973	3,047	2,486	2,178	1
1991	18,173	670	705	1,057	1,630	2,299	2,852	2,877	2,441	2,131	1
1990	17,185	547	718	1,016	1,599	2,312	2,643	2,751	2,285	1,962	-
1989	16,833	556	643	1,035	1,645	2,226	2,641	2,651	2,235	1,856	1
1988	16,006	489	655	1,002	1,626	2,152	2,470	2,521	2,043	1,699	1
1987	15,275	451	620	1,048	1,540	2,085	2,456	2,386	1,949	1,572	-
1986	15,178	477	651	1,013	1,590	2,191	2,386	2,335	1,761	1,425	-
1985	14,468	439	644	1,054	1,648	2,001	2,195	2,216	1,665	1,367	1
1984	13,718	394	648	1,048	1,535	1,921	2,149	1,985	1,627	1,270	1
1983	13,333	415	657	1,078	1,487	1,868	2,161	1,940	1,476	1,185	-
1982	12,890	418	720	1,067	1,453	1,773	1,933	1,890	1,418	1,102	1
1981	11,962	372	655	1,045	1,436	1,673	1,723	1,744	1,306	989	1
1980	11,909	404	687	1,076	1,357	1,617	1,889	1,612	1,263	931	-
1979	11,240	393	687	993	1,335	1,642	1,711	1,529	1,127	834	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
White, male											
1998	11,253	437	602	728	1,000	1,320	1,761	1,867	1,497	1,170	-
1997	11,241	463	634	758	968	1,393	1,748	1,858	1,421	1,004	-
1996	10,909	459	535	723	975	1,311	1,747	1,743	1,393	1,009	1
1995	10,732	451	561	701	938	1,376	1,722	1,635	1,265	959	-
1994	10,490	444	538	670	986	1,340	1,684	1,619	1,217	940	-
1993	9,658	402	460	632	899	1,254	1,516	1,487	1,160	815	-
1992	9,673	406	502	608	932	1,266	1,537	1,486	1,090	779	1
1991	9,336	436	441	657	896	1,247	1,418	1,339	1,027	769	-
1990	9,005	354	468	632	956	1,227	1,370	1,333	993	706	-
1989	8,643	357	408	623	937	1,175	1,360	1,251	922	671	1
1988	8,177	316	392	610	941	1,128	1,243	1,177	844	620	1
1987	7,709	279	380	624	843	1,135	1,233	1,111	779	517	-
1986	7,795	317	388	589	883	1,182	1,222	1,054	728	499	-
1985	7,547	281	384	593	912	1,093	1,122	1,106	711	508	-
1984	6,926	248	363	609	845	989	1,111	887	661	468	1
1983	6,823	235	388	622	833	965	1,132	916	581	427	-
1982	6,624	260	418	578	844	935	966	902	560	404	-
1981	6,066	228	398	585	777	875	862	783	522	367	-
1980	6,090	229	426	617	761	833	922	697	522	347	-
1979	5,966	249	395	552	734	927	875	737	487	338	-
Non-Hodgkin's lymphoma (200,202)											
White, female											
1998	10,350	225	335	479	682	1,023	1,547	1,955	1,833	1,871	1
1997	10,455	222	353	483	727	1,070	1,569	1,893	1,799	1,896	-
1996	10,153	245	355	467	673	1,088	1,569	1,834	1,705	1,753	-
1995	10,048	240	353	501	642	1,081	1,613	1,824	1,612	1,746	-
1994	9,779	226	340	466	706	1,153	1,580	1,679	1,628	1,537	-
1993	9,372	212	292	433	710	1,106	1,541	1,635	1,511	1,514	-
1992	8,935	220	280	408	689	1,107	1,436	1,561	1,396	1,399	-
1991	8,837	234	264	400	734	1,052	1,434	1,538	1,414	1,362	1
1990	8,180	193	250	384	643	1,085	1,273	1,418	1,292	1,256	-
1989	8,190	199	235	412	708	1,051	1,281	1,400	1,313	1,185	-
1988	7,829	173	263	392	685	1,024	1,227	1,344	1,199	1,079	-
1987	7,566	172	240	424	697	950	1,223	1,275	1,170	1,055	-
1986	7,383	160	263	424	707	1,009	1,164	1,281	1,033	926	-
1985	6,921	158	260	461	736	908	1,073	1,110	954	859	1
1984	6,792	146	285	439	690	932	1,038	1,098	966	802	-
1983	6,510	180	269	456	654	903	1,029	1,024	895	758	-
1982	6,266	158	302	489	609	838	967	988	858	698	1
1981	5,896	144	257	460	659	798	861	961	784	622	1
1980	5,819	175	261	459	596	784	967	915	741	584	-
1979	5,274	144	292	441	601	715	836	792	640	496	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
All other, both sexes											
1998	1,941	134	149	161	170	214	234	230	173	148	-
1997	1,997	132	155	146	191	208	238	229	167	154	-
1996	1,872	123	124	138	177	191	239	198	184	135	-
1995	1,742	95	134	120	169	198	195	194	135	140	-
1994	1,661	121	92	131	193	169	195	154	160	122	-
1993	1,601	102	116	125	135	198	177	169	141	114	-
1992	1,571	83	93	116	167	172	174	182	122	112	-
1991	1,527	74	91	121	156	192	189	174	135	81	-
1990	1,416	82	103	127	135	177	166	150	98	91	-
1989	1,364	82	85	102	141	164	160	174	105	71	-
1988	1,253	81	93	118	111	162	138	126	101	85	-
1987	1,144	69	71	82	113	143	151	137	93	72	2
1986	1,173	56	81	100	97	152	164	133	88	64	-
1985	1,094	61	60	94	127	131	128	128	84	60	-
1984	974	55	60	100	117	111	131	97	66	39	-
1983	965	55	63	95	117	119	110	100	69	50	-
1982	913	36	75	74	111	112	121	102	74	39	-
1981	865	36	66	65	120	114	112	70	46	48	-
1980	852	41	66	96	92	107	104	91	56	42	1
1979	782	50	66	77	100	108	95	70	41	35	-
Non-Hodgkin's lymphoma (200,202)											
All other, male											
1998	1,026	86	94	87	101	116	101	111	73	55	-
1997	1,128	93	100	80	101	109	126	124	81	62	-
1996	1,053	89	77	85	104	97	130	96	85	51	-
1995	962	63	91	64	99	107	98	86	65	53	-
1994	883	67	55	78	111	82	95	67	65	46	-
1993	902	67	71	61	70	111	96	90	59	57	-
1992	869	57	58	65	96	92	88	81	55	40	-
1991	821	50	56	67	91	94	93	82	59	27	-
1990	790	55	59	68	82	101	80	68	46	40	-
1989	777	51	51	60	76	85	80	83	46	39	-
1988	715	56	60	69	60	87	81	64	43	35	-
1987	639	41	41	47	55	86	81	74	34	37	2
1986	690	34	50	65	53	87	91	77	50	20	-
1985	585	40	31	50	70	73	63	56	39	18	-
1984	524	36	34	60	64	49	63	38	35	16	-
1983	559	40	32	62	73	60	59	55	28	27	-
1982	517	20	48	42	67	56	68	44	39	15	-
1981	457	21	35	36	62	50	64	31	26	20	-
1980	483	21	41	58	51	72	59	45	28	14	1
1979	432	28	38	40	63	54	54	29	21	15	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
All other, female											
1998	915	48	55	74	69	98	133	119	100	93	-
1997	869	39	55	66	90	99	112	105	86	92	-
1996	819	34	47	53	73	94	109	102	99	84	-
1995	780	32	43	56	70	91	97	108	70	87	-
1994	778	54	37	53	82	87	100	87	95	76	-
1993	699	35	45	64	65	87	81	79	82	57	-
1992	702	26	35	51	71	80	86	101	67	72	-
1991	706	24	35	54	65	98	96	92	76	54	-
1990	626	27	44	59	53	76	86	82	52	51	-
1989	587	31	34	42	65	79	80	91	59	32	-
1988	538	25	33	49	51	75	57	62	58	50	-
1987	505	28	30	35	58	57	70	63	59	35	-
1986	483	22	31	35	44	65	73	56	38	44	-
1985	509	21	29	44	57	58	65	72	45	42	-
1984	450	19	26	40	53	62	68	59	31	23	-
1983	406	15	31	33	44	59	51	45	41	23	-
1982	396	16	27	32	44	56	53	58	35	24	-
1981	408	15	31	29	58	64	48	39	20	28	-
1980	369	20	25	38	41	35	45	46	28	28	-
1979	350	22	28	37	37	54	41	41	20	20	-
Non-Hodgkin's lymphoma (200,202)											
Black, both sexes											
1998	1,518	118	130	139	141	162	171	153	129	110	-
1997	1,583	112	125	110	155	162	188	162	131	114	-
1996	1,517	109	106	116	152	156	169	148	141	105	-
1995	1,399	84	114	92	133	162	150	147	100	106	-
1994	1,310	102	76	109	149	136	155	113	121	87	-
1993	1,312	92	97	109	114	163	134	136	110	83	-
1992	1,282	70	80	93	136	141	137	136	95	89	-
1991	1,260	58	73	106	121	153	157	141	111	69	-
1990	1,172	70	91	108	110	146	126	123	78	72	-
1989	1,135	71	71	86	115	137	129	141	83	56	-
1988	1,047	72	78	105	91	131	112	102	79	74	-
1987	971	61	63	71	96	125	123	114	73	56	2
1986	1,006	52	73	82	85	130	139	107	72	54	-
1985	926	51	47	78	110	110	112	111	68	51	-
1984	840	47	47	88	101	92	112	84	60	30	-
1983	841	48	55	87	105	110	96	84	61	38	-
1982	792	33	64	65	98	99	105	87	61	33	-
1981	761	29	56	59	109	101	101	58	41	42	-
1980	742	35	62	84	81	86	89	79	52	37	1
1979	682	43	59	73	87	98	78	56	40	23	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Non-Hodgkin's lymphoma (200,202)											
Black, male											
1998	801	78	83	73	84	86	78	70	51	36	-
1997	881	78	82	56	82	85	96	84	58	43	-
1996	851	79	65	71	90	85	86	65	62	37	-
1995	780	58	78	51	83	89	73	63	47	35	-
1994	692	58	45	65	84	72	74	46	41	29	-
1993	726	59	64	55	60	91	67	70	41	35	-
1992	704	47	48	50	76	79	67	57	41	28	-
1991	670	41	45	57	71	77	70	61	47	22	-
1990	651	48	56	59	66	83	57	53	38	30	-
1989	643	45	44	53	62	71	60	62	35	29	-
1988	600	52	49	62	51	68	66	50	30	33	-
1987	544	37	36	43	47	74	68	58	23	28	2
1986	576	31	44	55	47	72	74	59	35	16	-
1985	489	34	25	40	60	64	51	49	27	14	-
1984	453	30	29	53	55	42	53	30	34	12	-
1983	481	36	28	55	63	57	50	46	25	21	-
1982	437	19	41	37	55	47	61	32	28	14	-
1981	403	17	30	31	57	43	58	25	25	17	-
1980	421	20	39	51	43	59	50	36	27	12	1
1979	373	26	33	39	55	49	41	20	20	9	-
Non-Hodgkin's lymphoma (200,202)											
Black, female											
1998	717	40	47	66	57	76	93	83	78	74	-
1997	702	34	43	54	73	77	92	78	73	71	-
1996	666	30	41	45	62	71	83	83	79	68	-
1995	619	26	36	41	50	73	77	84	53	71	-
1994	618	44	31	44	65	64	81	67	80	58	-
1993	586	33	33	54	54	72	67	66	69	48	-
1992	578	23	32	43	60	62	70	79	54	61	-
1991	590	17	28	49	50	76	87	80	64	47	-
1990	521	22	35	49	44	63	69	70	40	42	-
1989	492	26	27	33	53	66	69	79	48	27	-
1988	447	20	29	43	40	63	46	52	49	41	-
1987	427	24	27	28	49	51	55	56	50	28	-
1986	430	21	29	27	38	58	65	48	37	38	-
1985	437	17	22	38	50	46	61	62	41	37	-
1984	387	17	18	35	46	50	59	54	26	18	-
1983	360	12	27	32	42	53	46	38	36	17	-
1982	355	14	23	28	43	52	44	55	33	19	-
1981	358	12	26	28	52	58	43	33	16	25	-
1980	321	15	23	33	38	27	39	43	25	25	-
1979	309	17	26	34	32	49	37	36	20	14	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
All races, both sexes											
1998	20,324	611	744	956	1,370	2,001	2,952	3,343	2,912	3,100	1
1997	20,313	610	794	995	1,429	2,130	2,929	3,189	2,810	3,087	-
1996	20,340	635	752	977	1,400	2,166	2,998	3,192	2,842	3,031	-
1995	20,148	594	686	944	1,474	2,102	2,973	3,168	2,805	2,961	2
1994	19,669	586	748	941	1,481	2,106	2,840	2,936	2,642	2,893	-
1993	19,530	580	720	919	1,502	2,232	2,774	2,916	2,514	2,777	1
1992	19,272	575	688	976	1,491	2,218	2,671	2,878	2,509	2,650	-
1991	18,945	553	664	957	1,501	2,142	2,730	2,874	2,395	2,579	-
1990	18,574	525	682	960	1,483	2,145	2,609	2,756	2,367	2,393	3
1989	18,246	514	620	947	1,580	2,116	2,650	2,622	2,291	2,242	-
1988	17,577	507	627	952	1,457	1,970	2,396	2,580	2,234	2,233	-
1987	17,284	485	639	969	1,489	1,993	2,333	2,467	2,078	2,162	2
1986	17,365	474	627	1,021	1,511	2,003	2,370	2,439	2,120	2,079	1
1985	17,319	484	638	1,037	1,576	2,035	2,337	2,431	2,004	2,092	2
1984	17,080	481	707	1,084	1,537	1,988	2,287	2,298	1,973	1,988	-
1983	16,820	483	691	996	1,511	1,948	2,255	2,314	1,822	1,998	4
1982	16,704	455	677	983	1,457	1,924	2,178	2,332	1,889	1,918	-
1981	16,273	475	671	1,057	1,452	1,841	2,189	2,196	1,864	1,717	4
1980	16,533	494	716	1,098	1,471	1,802	2,231	2,097	1,824	1,751	-
1979	15,988	498	756	1,050	1,383	1,789	2,103	2,030	1,794	1,671	1
Leukemia (204-208)											
All races, male											
1998	11,201	340	434	567	819	1,250	1,783	1,918	1,507	1,213	1
1997	11,265	349	450	577	856	1,343	1,748	1,811	1,505	1,286	-
1996	11,179	384	426	573	855	1,311	1,781	1,785	1,418	1,291	-
1995	11,229	328	402	555	898	1,307	1,797	1,811	1,442	1,251	-
1994	10,840	328	431	551	928	1,269	1,684	1,666	1,340	1,161	-
1993	10,750	310	423	564	900	1,388	1,631	1,647	1,278	1,094	1
1992	10,609	316	381	589	890	1,383	1,546	1,637	1,260	1,062	-
1991	10,194	319	372	542	909	1,250	1,644	1,543	1,167	984	-
1990	10,192	281	405	567	898	1,311	1,535	1,514	1,180	976	1
1989	10,039	295	346	544	949	1,311	1,528	1,516	1,138	850	-
1988	9,724	288	357	567	865	1,187	1,407	1,476	1,099	908	-
1987	9,387	255	348	552	895	1,188	1,348	1,360	1,047	836	1
1986	9,565	264	342	590	883	1,193	1,408	1,363	1,060	859	1
1985	9,442	256	371	591	938	1,196	1,347	1,306	1,000	874	1
1984	9,276	257	392	623	904	1,167	1,304	1,225	979	824	-
1983	9,312	269	392	581	911	1,179	1,302	1,273	896	883	2
1982	9,244	256	381	579	879	1,170	1,256	1,276	971	826	-
1981	9,078	265	360	638	877	1,130	1,291	1,186	956	736	2
1980	9,193	270	416	629	867	1,075	1,307	1,157	894	798	-
1979	8,893	265	405	623	812	1,087	1,234	1,098	886	784	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
All races, female											
1998	9,123	271	310	389	551	751	1,169	1,425	1,405	1,887	-
1997	9,048	261	344	418	573	787	1,181	1,378	1,305	1,801	-
1996	9,161	251	326	404	545	855	1,217	1,407	1,424	1,740	-
1995	8,919	266	284	389	576	795	1,176	1,357	1,363	1,710	2
1994	8,829	258	317	390	553	837	1,156	1,270	1,302	1,732	-
1993	8,780	270	297	355	602	844	1,143	1,269	1,236	1,683	-
1992	8,663	259	307	387	601	835	1,125	1,241	1,249	1,588	-
1991	8,751	234	292	415	592	892	1,086	1,331	1,228	1,595	-
1990	8,382	244	277	393	585	834	1,074	1,242	1,187	1,417	2
1989	8,207	219	274	403	631	805	1,122	1,106	1,153	1,392	-
1988	7,853	219	270	385	592	783	989	1,104	1,135	1,325	-
1987	7,897	230	291	417	594	805	985	1,107	1,031	1,326	1
1986	7,800	210	285	431	628	810	962	1,076	1,060	1,220	-
1985	7,877	228	267	446	638	839	990	1,125	1,004	1,218	1
1984	7,804	224	315	461	633	821	983	1,073	994	1,164	-
1983	7,508	214	299	415	600	769	953	1,041	926	1,115	2
1982	7,460	199	296	404	578	754	922	1,056	918	1,092	-
1981	7,195	210	311	419	575	711	898	1,010	908	981	2
1980	7,340	224	300	469	604	727	924	940	930	953	-
1979	7,095	233	351	427	571	702	869	932	908	887	-
Leukemia (204-208)											
White, both sexes											
1998	18,214	493	627	801	1,193	1,800	2,711	3,105	2,710	2,887	-
1997	18,283	523	681	871	1,245	1,930	2,656	2,965	2,635	2,905	-
1996	18,234	537	647	843	1,233	1,928	2,733	2,959	2,639	2,815	-
1995	18,109	500	579	819	1,300	1,899	2,721	2,933	2,615	2,782	1
1994	17,708	498	652	825	1,290	1,914	2,599	2,718	2,471	2,702	-
1993	17,504	485	623	798	1,317	2,023	2,545	2,704	2,329	2,594	-
1992	17,405	500	585	842	1,340	2,017	2,454	2,671	2,348	2,494	-
1991	17,064	472	559	819	1,350	1,932	2,522	2,672	2,227	2,408	-
1990	16,722	451	590	834	1,310	1,936	2,410	2,551	2,212	2,250	2
1989	16,434	448	534	828	1,410	1,916	2,420	2,446	2,128	2,097	-
1988	15,854	438	553	841	1,326	1,772	2,201	2,377	2,076	2,090	-
1987	15,569	418	547	848	1,338	1,813	2,135	2,278	1,942	2,030	1
1986	15,709	420	548	903	1,332	1,824	2,193	2,276	1,969	1,951	1
1985	15,703	418	563	912	1,414	1,857	2,130	2,290	1,856	1,985	1
1984	15,479	422	604	966	1,394	1,820	2,086	2,142	1,854	1,891	-
1983	15,231	421	599	899	1,356	1,765	2,061	2,145	1,716	1,890	4
1982	15,152	402	585	869	1,295	1,772	2,007	2,156	1,775	1,838	-
1981	14,786	415	594	937	1,344	1,669	2,030	2,038	1,751	1,632	-
1980	15,081	423	634	980	1,344	1,652	2,075	1,957	1,726	1,669	-
1979	14,607	435	673	943	1,260	1,632	1,956	1,910	1,699	1,585	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
White, male											
1998	10,070	273	372	485	721	1,135	1,651	1,794	1,413	1,124	-
1997	10,171	301	389	514	743	1,232	1,593	1,685	1,424	1,215	-
1996	10,055	329	371	500	754	1,175	1,632	1,666	1,320	1,196	-
1995	10,143	284	340	482	794	1,183	1,653	1,688	1,356	1,176	-
1994	9,772	285	380	485	812	1,157	1,545	1,548	1,261	1,082	-
1993	9,658	267	368	487	800	1,265	1,515	1,527	1,189	1,028	-
1992	9,595	274	319	513	811	1,272	1,427	1,511	1,186	1,002	-
1991	9,165	267	315	465	835	1,137	1,530	1,423	1,084	911	-
1990	9,189	241	352	500	805	1,182	1,414	1,405	1,105	917	1
1989	9,011	261	299	473	852	1,188	1,398	1,417	1,039	787	-
1988	8,785	247	319	503	786	1,075	1,297	1,364	1,034	841	-
1987	8,480	219	307	485	810	1,082	1,250	1,261	974	778	1
1986	8,685	233	305	529	787	1,088	1,307	1,277	988	809	1
1985	8,548	223	321	522	850	1,102	1,232	1,222	920	823	1
1984	8,409	221	341	565	821	1,072	1,197	1,130	924	782	-
1983	8,461	238	340	534	815	1,072	1,205	1,187	844	833	2
1982	8,402	226	335	518	789	1,080	1,165	1,176	909	785	-
1981	8,276	230	322	573	814	1,037	1,209	1,107	896	697	-
1980	8,396	238	374	560	800	978	1,212	1,092	840	755	-
1979	8,138	232	368	560	742	1,001	1,147	1,034	833	747	1
Leukemia (204-208)											
White, female											
1998	8,144	220	255	316	472	665	1,060	1,311	1,297	1,763	-
1997	8,112	222	292	357	502	698	1,063	1,280	1,211	1,690	-
1996	8,179	208	276	343	479	753	1,101	1,293	1,319	1,619	-
1995	7,966	216	239	337	506	716	1,068	1,245	1,259	1,606	1
1994	7,936	213	272	340	478	757	1,054	1,170	1,210	1,620	-
1993	7,846	218	255	311	517	758	1,030	1,177	1,140	1,566	-
1992	7,810	226	266	329	529	745	1,027	1,160	1,162	1,492	-
1991	7,899	205	244	354	515	795	992	1,249	1,143	1,497	-
1990	7,533	210	238	334	505	754	996	1,146	1,107	1,333	1
1989	7,423	187	235	355	558	728	1,022	1,029	1,089	1,310	-
1988	7,069	191	234	338	540	697	904	1,013	1,042	1,249	-
1987	7,089	199	240	363	528	731	885	1,017	968	1,252	-
1986	7,024	187	243	374	545	736	886	999	981	1,142	-
1985	7,155	195	242	390	564	755	898	1,068	936	1,162	-
1984	7,070	201	263	401	573	748	889	1,012	930	1,109	-
1983	6,770	183	259	365	541	693	856	958	872	1,057	2
1982	6,750	176	250	351	506	692	842	980	866	1,053	-
1981	6,510	185	272	364	530	632	821	931	855	935	-
1980	6,685	185	260	420	544	674	863	865	886	914	-
1979	6,469	203	305	383	518	631	809	876	866	838	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
All other, both sexes											
1998	2,110	118	117	155	177	201	241	238	202	213	1
1997	2,030	87	113	124	184	200	273	224	175	182	-
1996	2,106	98	105	134	167	238	265	233	203	216	-
1995	2,039	94	107	125	174	203	252	235	190	179	1
1994	1,961	88	96	116	191	192	241	218	171	191	-
1993	2,026	95	97	121	185	209	229	212	185	183	1
1992	1,867	75	103	134	151	201	217	207	161	156	-
1991	1,881	81	105	138	151	210	208	202	168	171	-
1990	1,852	74	92	126	173	209	199	205	155	143	1
1989	1,812	66	86	119	170	200	230	176	163	145	-
1988	1,723	69	74	111	131	198	195	203	158	143	-
1987	1,715	67	92	121	151	180	198	189	136	132	1
1986	1,656	54	79	118	179	179	177	163	151	128	-
1985	1,616	66	75	125	162	178	207	141	148	107	1
1984	1,601	59	103	118	143	168	201	156	119	97	-
1983	1,589	62	92	97	155	183	194	169	106	108	-
1982	1,552	53	92	114	162	152	171	176	114	80	-
1981	1,487	60	77	120	108	172	159	158	113	85	4
1980	1,452	71	82	118	127	150	156	140	98	82	-
1979	1,381	63	83	107	123	157	147	120	95	86	-
Leukemia (204-208)											
All other, male											
1998	1,131	67	62	82	98	115	132	124	94	89	1
1997	1,094	48	61	63	113	111	155	126	81	71	-
1996	1,124	55	55	73	101	136	149	119	98	95	-
1995	1,086	44	62	73	104	124	144	123	86	75	-
1994	1,068	43	51	66	116	112	139	118	79	79	-
1993	1,092	43	55	77	100	123	116	120	89	66	1
1992	1,014	42	62	76	79	111	119	126	74	60	-
1991	1,029	52	57	77	74	113	114	120	83	73	-
1990	1,003	40	53	67	93	129	121	109	75	59	-
1989	1,028	34	47	71	97	123	130	99	99	63	-
1988	939	41	38	64	79	112	110	112	65	67	-
1987	907	36	41	67	85	106	98	99	73	58	-
1986	880	31	37	61	96	105	101	86	72	50	-
1985	894	33	50	69	88	94	115	84	80	51	-
1984	867	36	51	58	83	95	107	95	55	42	-
1983	851	31	52	47	96	107	97	86	52	50	-
1982	842	30	46	61	90	90	91	100	62	41	-
1981	802	35	38	65	63	93	82	79	60	39	2
1980	797	32	42	69	67	97	95	65	54	43	-
1979	755	33	37	63	70	86	87	64	53	37	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
All other, female											
1998	979	51	55	73	79	86	109	114	108	124	-
1997	936	39	52	61	71	89	118	98	94	111	-
1996	982	43	50	61	66	102	116	114	105	121	-
1995	953	50	45	52	70	79	108	112	104	104	1
1994	893	45	45	50	75	80	102	100	92	112	-
1993	934	52	42	44	85	86	113	92	96	117	-
1992	853	33	41	58	72	90	98	81	87	96	-
1991	852	29	48	61	77	97	94	82	85	98	-
1990	849	34	39	59	80	80	78	96	80	84	1
1989	784	32	39	48	73	77	100	77	64	82	-
1988	784	28	36	47	52	86	85	91	93	76	-
1987	808	31	51	54	66	74	100	90	63	74	1
1986	776	23	42	57	83	74	76	77	79	78	-
1985	722	33	25	56	74	84	92	57	68	56	1
1984	734	23	52	60	60	73	94	61	64	55	-
1983	738	31	40	50	59	76	97	83	54	58	-
1982	710	23	46	53	72	62	80	76	52	39	-
1981	685	25	39	55	45	79	77	79	53	46	2
1980	655	39	40	49	60	53	61	75	44	39	-
1979	626	30	46	44	53	71	60	56	42	49	-
Leukemia (204-208)											
Black, both sexes											
1998	1,752	99	89	130	150	165	201	198	183	178	1
1997	1,673	73	86	98	157	164	231	191	154	160	-
1996	1,763	75	90	114	143	198	217	206	188	190	-
1995	1,686	82	85	108	151	179	213	198	158	155	-
1994	1,634	74	74	99	161	169	204	191	141	172	-
1993	1,681	70	85	96	162	183	194	181	162	154	1
1992	1,587	58	84	119	132	174	182	183	149	146	-
1991	1,609	67	84	117	135	185	187	173	154	157	-
1990	1,591	67	81	111	151	184	183	183	143	133	1
1989	1,540	57	74	99	144	173	196	161	153	132	-
1988	1,478	53	59	95	113	175	175	186	140	134	-
1987	1,479	59	81	103	139	151	183	171	124	123	1
1986	1,460	40	69	105	161	163	157	147	144	119	-
1985	1,401	53	63	111	139	165	189	123	138	100	1
1984	1,407	49	81	103	126	151	184	141	110	88	-
1983	1,394	53	81	83	143	164	180	147	98	99	-
1982	1,378	42	79	102	151	139	160	160	102	71	-
1981	1,337	53	66	107	101	153	153	147	107	78	3
1980	1,310	64	70	105	120	133	147	130	95	78	-
1979	1,262	58	78	98	119	149	137	114	89	82	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Leukemia (204-208)											
Black, male											
1998	936	55	50	71	84	96	104	101	82	71	1
1997	905	41	47	51	96	90	136	105	68	62	-
1996	931	43	46	60	85	112	115	109	88	83	-
1995	908	39	52	68	89	113	120	103	71	62	-
1994	885	36	39	59	95	104	119	104	60	66	-
1993	906	31	49	62	91	112	101	104	75	51	1
1992	840	33	50	68	71	95	99	106	65	51	-
1991	869	41	45	68	66	99	100	102	73	66	-
1990	854	35	46	64	82	112	109	93	68	55	-
1989	868	33	40	57	77	105	112	87	92	57	-
1988	806	32	33	54	64	97	98	102	54	66	-
1987	781	32	34	58	77	93	93	90	65	52	-
1986	771	24	31	54	85	98	88	76	67	46	-
1985	769	29	41	61	72	87	104	72	74	48	-
1984	754	32	38	50	74	84	93	87	53	37	-
1983	741	25	45	41	90	92	87	75	49	47	-
1982	745	26	38	55	84	82	83	89	54	37	-
1981	709	29	31	58	59	78	76	70	56	36	2
1980	714	26	34	61	65	87	87	57	52	40	-
1979	677	28	33	56	68	80	83	58	50	36	-
Leukemia (204-208)											
Black, female											
1998	816	44	39	59	66	69	97	97	101	107	-
1997	768	32	39	47	61	74	95	86	86	98	-
1996	832	32	44	54	58	86	102	97	100	107	-
1995	778	43	33	40	62	66	93	95	87	93	-
1994	749	38	35	40	66	65	85	87	81	106	-
1993	775	39	36	34	71	71	93	77	87	103	-
1992	747	25	34	51	61	79	83	77	84	95	-
1991	740	26	39	49	69	86	87	71	81	91	-
1990	737	32	35	47	69	72	74	90	75	78	1
1989	672	24	34	42	67	68	84	74	61	75	-
1988	672	21	26	41	49	78	77	84	86	68	-
1987	698	27	47	45	62	58	90	81	59	71	1
1986	689	16	38	51	76	65	69	71	77	73	-
1985	632	24	22	50	67	78	85	51	64	52	1
1984	653	17	43	53	52	67	91	54	57	51	-
1983	653	28	36	42	53	72	93	72	49	52	-
1982	633	16	41	47	67	57	77	71	48	34	-
1981	628	24	35	49	42	75	77	77	51	42	1
1980	596	38	36	44	55	46	60	73	43	38	-
1979	585	30	45	42	51	69	54	56	39	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
All races, both sexes											
1998	10,359	260	417	650	891	1,289	1,809	1,970	1,517	1,378	-
1997	10,386	265	409	690	899	1,357	1,865	1,801	1,573	1,361	-
1996	10,248	217	412	613	975	1,366	1,819	1,858	1,508	1,304	-
1995	10,266	250	401	608	1,010	1,470	1,793	1,801	1,492	1,262	-
1994	10,041	259	432	601	958	1,355	1,776	1,782	1,452	1,250	1
1993	9,892	208	376	535	962	1,489	1,719	1,820	1,409	1,196	1
1992	9,294	186	340	575	851	1,381	1,654	1,688	1,379	1,066	-
1991	9,320	184	339	558	946	1,388	1,644	1,709	1,304	1,066	1
1990	8,934	179	336	536	950	1,339	1,584	1,532	1,314	1,022	-
1989	8,587	178	330	587	942	1,305	1,479	1,530	1,188	903	1
1988	8,318	191	316	587	902	1,313	1,430	1,421	1,137	880	-
1987	8,141	184	321	561	928	1,256	1,473	1,344	1,062	866	2
1986	7,987	153	318	528	878	1,269	1,407	1,401	1,038	846	-
1985	7,819	144	331	572	945	1,186	1,431	1,286	985	779	-
1984	7,470	166	321	622	943	1,110	1,287	1,252	945	691	-
1983	7,301	149	323	605	901	1,147	1,249	1,222	890	685	1
1982	7,024	155	373	550	840	1,113	1,247	1,136	811	667	1
1981	6,729	161	320	535	855	1,051	1,202	1,121	804	561	3
1980	6,613	143	308	608	856	1,056	1,131	1,035	781	571	-
1979	6,414	166	360	575	797	1,009	1,101	1,029	730	525	-
Multiple myeloma and immunoproliferative neoplasms (203)											
All races, male											
1998	5,290	149	253	346	510	712	956	968	718	570	-
1997	5,360	157	247	375	503	741	999	897	784	558	-
1996	5,253	125	244	336	509	750	963	934	732	551	-
1995	5,165	149	240	334	557	782	897	897	684	521	-
1994	5,165	144	263	323	553	766	978	860	660	509	-
1993	4,936	129	208	310	501	817	903	875	618	463	-
1992	4,679	105	188	328	483	737	880	821	617	411	-
1991	4,698	104	177	310	515	769	857	839	592	415	1
1990	4,561	95	204	299	529	723	840	777	598	406	-
1989	4,335	104	183	343	502	696	754	762	550	353	-
1988	4,231	103	178	340	496	697	743	692	535	364	-
1987	4,181	113	203	315	530	709	766	665	452	339	2
1986	4,032	84	161	290	474	700	752	677	478	325	-
1985	3,935	69	177	322	518	640	742	593	448	325	-
1984	3,737	91	176	335	510	592	673	609	410	258	-
1983	3,722	83	183	353	486	602	628	605	407	301	-
1982	3,554	84	221	302	453	619	625	535	369	261	1
1981	3,377	94	184	292	471	531	611	532	366	226	1
1980	3,401	72	170	350	461	567	577	525	352	248	-
1979	3,333	103	200	336	461	538	554	503	319	240	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
All races, female											
1998	5,069	111	164	304	381	577	853	1,002	799	808	-
1997	5,026	108	162	315	396	616	866	904	789	803	-
1996	4,995	92	168	277	466	616	856	924	776	753	-
1995	5,101	101	161	274	453	688	896	904	808	741	-
1994	4,876	115	169	278	405	589	798	922	792	741	1
1993	4,956	79	168	225	461	672	816	945	791	733	1
1992	4,615	81	152	247	368	644	774	867	762	655	-
1991	4,622	80	162	248	431	619	787	870	712	651	-
1990	4,373	84	132	237	421	616	744	755	716	616	-
1989	4,252	74	147	244	440	609	725	768	638	550	1
1988	4,087	88	138	247	406	616	687	729	602	516	-
1987	3,960	71	118	246	398	547	707	679	610	527	-
1986	3,955	69	157	238	404	569	655	724	560	521	-
1985	3,884	75	154	250	427	546	689	693	537	454	-
1984	3,733	75	145	287	433	518	614	643	535	433	-
1983	3,579	66	140	252	415	545	621	617	483	384	1
1982	3,470	71	152	248	387	494	622	601	442	406	-
1981	3,352	67	136	243	384	520	591	589	438	335	2
1980	3,212	71	138	258	395	489	554	510	429	323	-
1979	3,081	63	160	239	336	471	547	526	411	285	-
Multiple myeloma and immunoproliferative neoplasms (203)											
White, both sexes											
1998	8,400	189	301	471	668	1,030	1,488	1,677	1,297	1,153	-
1997	8,528	182	316	517	698	1,088	1,571	1,527	1,360	1,163	-
1996	8,367	153	313	461	742	1,095	1,489	1,564	1,314	1,120	-
1995	8,406	181	284	450	764	1,215	1,499	1,516	1,292	1,078	-
1994	8,292	193	317	442	765	1,090	1,507	1,535	1,242	1,071	-
1993	8,132	151	279	401	753	1,219	1,409	1,537	1,236	1,009	1
1992	7,641	134	228	420	665	1,137	1,380	1,450	1,188	918	-
1991	7,629	131	251	419	766	1,123	1,342	1,430	1,116	925	1
1990	7,377	126	257	406	755	1,102	1,313	1,304	1,131	892	-
1989	7,106	122	238	472	759	1,062	1,251	1,317	1,020	765	1
1988	6,892	139	237	445	736	1,076	1,197	1,207	981	776	-
1987	6,729	136	236	456	750	1,029	1,219	1,131	926	740	1
1986	6,556	106	235	405	685	1,050	1,151	1,188	892	741	-
1985	6,468	98	264	455	764	973	1,190	1,102	821	681	-
1984	6,200	120	246	484	759	932	1,090	1,058	814	605	-
1983	6,009	109	241	483	714	947	1,042	1,024	754	600	1
1982	5,802	109	286	423	698	909	1,044	979	688	577	-
1981	5,541	120	254	415	703	848	998	935	697	483	3
1980	5,505	103	236	476	699	860	958	892	684	511	-
1979	5,368	137	271	453	649	817	947	894	654	459	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
White, male											
1998	4,357	115	187	257	400	579	804	838	616	476	-
1997	4,459	112	201	278	397	597	865	776	684	482	-
1996	4,342	94	185	257	399	610	802	803	643	478	-
1995	4,286	110	184	267	440	663	751	760	601	439	-
1994	4,297	112	195	239	449	619	842	757	569	432	-
1993	4,106	101	163	236	406	675	741	744	562	390	-
1992	3,865	87	128	243	376	616	740	712	533	350	-
1991	3,871	78	134	236	426	625	710	702	516	357	1
1990	3,765	69	162	232	421	599	702	669	510	343	-
1989	3,600	70	128	277	413	577	642	653	478	297	-
1988	3,562	78	137	270	414	590	635	587	461	326	-
1987	3,458	88	154	261	439	590	639	548	379	293	1
1986	3,299	61	126	224	372	581	618	583	393	277	-
1985	3,251	45	141	263	414	527	620	513	372	280	-
1984	3,085	67	137	262	412	498	570	507	348	225	-
1983	3,049	58	140	286	378	503	526	501	336	262	-
1982	2,922	61	168	235	388	513	519	455	305	217	-
1981	2,772	72	147	229	388	425	514	436	311	194	1
1980	2,834	58	135	277	381	459	492	448	306	223	-
1979	2,795	84	159	274	377	436	485	427	283	208	-
Multiple myeloma and immunoproliferative neoplasms (203)											
White, female											
1998	4,043	74	114	214	268	451	684	839	681	677	-
1997	4,069	70	115	239	301	491	706	751	676	681	-
1996	4,025	59	128	204	343	485	687	761	671	642	-
1995	4,120	71	100	183	324	552	748	756	691	639	-
1994	3,995	81	122	203	316	471	665	778	673	639	-
1993	4,026	50	116	165	347	544	668	793	674	619	1
1992	3,776	47	100	177	289	521	640	738	655	568	-
1991	3,758	53	117	183	340	498	632	728	600	568	-
1990	3,612	57	95	174	334	503	611	635	621	549	-
1989	3,506	52	110	195	346	485	609	664	542	468	1
1988	3,330	61	100	175	322	486	562	620	520	450	-
1987	3,271	48	82	195	311	439	580	583	547	447	-
1986	3,257	45	109	181	313	469	533	605	499	464	-
1985	3,217	53	123	192	350	446	570	589	449	401	-
1984	3,115	53	109	222	347	434	520	551	466	380	-
1983	2,960	51	101	197	336	444	516	523	418	338	1
1982	2,880	48	118	188	310	396	525	524	383	360	-
1981	2,769	48	107	186	315	423	484	499	386	289	2
1980	2,671	45	101	199	318	401	466	444	378	288	-
1979	2,573	53	112	179	272	381	462	467	371	251	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
All other, both sexes											
1998	1,959	71	116	179	223	259	321	293	220	225	-
1997	1,858	83	93	173	201	269	294	274	213	198	-
1996	1,881	64	99	152	233	271	330	294	194	184	-
1995	1,860	69	117	158	246	255	294	285	200	184	-
1994	1,749	66	115	159	193	265	269	247	210	179	1
1993	1,760	57	97	134	209	270	310	283	173	187	-
1992	1,653	52	112	155	186	244	274	238	191	148	-
1991	1,691	53	88	139	180	265	302	279	188	141	-
1990	1,557	53	79	130	195	237	271	228	183	130	-
1989	1,481	56	92	115	183	243	228	213	168	138	-
1988	1,426	52	79	142	166	237	233	214	156	104	-
1987	1,412	48	85	105	178	227	254	213	136	126	1
1986	1,431	47	83	123	193	219	256	213	146	105	-
1985	1,351	46	67	117	181	213	241	184	164	98	-
1984	1,270	46	75	138	184	178	197	194	131	86	-
1983	1,292	40	82	122	187	200	207	198	136	85	-
1982	1,222	46	87	127	142	204	203	157	123	90	1
1981	1,188	41	66	120	152	203	204	186	107	78	-
1980	1,108	40	72	132	157	196	173	143	97	60	-
1979	1,046	29	89	122	148	192	154	135	76	66	-
Multiple myeloma and immunoproliferative neoplasms (203)											
All other, male											
1998	933	34	66	89	110	133	152	130	102	94	-
1997	901	45	46	97	106	144	134	121	100	76	-
1996	911	31	59	79	110	140	161	131	89	73	-
1995	879	39	56	67	117	119	146	137	83	82	-
1994	868	32	68	84	104	147	136	103	91	77	-
1993	830	28	45	74	95	142	162	131	56	73	-
1992	814	18	60	85	107	121	140	109	84	61	-
1991	827	26	43	74	89	144	147	137	76	58	-
1990	796	26	42	67	108	124	138	108	88	63	-
1989	735	34	55	66	89	119	112	109	72	56	-
1988	669	25	41	70	82	107	108	105	74	38	-
1987	723	25	49	54	91	119	127	117	73	46	1
1986	733	23	35	66	102	119	134	94	85	48	-
1985	684	24	36	59	104	113	122	80	76	45	-
1984	652	24	39	73	98	94	103	102	62	33	-
1983	673	25	43	67	108	99	102	104	71	39	-
1982	632	23	53	67	65	106	106	80	64	44	1
1981	605	22	37	63	83	106	97	96	55	32	-
1980	567	14	35	73	80	108	85	77	46	25	-
1979	538	19	41	62	84	102	69	76	36	32	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
All other, female											
1998	1,026	37	50	90	113	126	169	163	118	131	-
1997	957	38	47	76	95	125	160	153	113	122	-
1996	970	33	40	73	123	131	169	163	105	111	-
1995	981	30	61	91	129	136	148	148	117	102	-
1994	881	34	47	75	89	118	133	144	119	102	1
1993	930	29	52	60	114	128	148	152	117	114	-
1992	839	34	52	70	79	123	134	129	107	87	-
1991	864	27	45	65	91	121	155	142	112	83	-
1990	761	27	37	63	87	113	133	120	95	67	-
1989	746	22	37	49	94	124	116	104	96	82	-
1988	757	27	38	72	84	130	125	109	82	66	-
1987	689	23	36	51	87	108	127	96	63	80	-
1986	698	24	48	57	91	100	122	119	61	57	-
1985	667	22	31	58	77	100	119	104	88	53	-
1984	618	22	36	65	86	84	94	92	69	53	-
1983	619	15	39	55	79	101	105	94	65	46	-
1982	590	23	34	60	77	98	97	77	59	46	-
1981	583	19	29	57	69	97	107	90	52	46	-
1980	541	26	37	59	77	88	88	66	51	35	-
1979	508	10	48	60	64	90	85	59	40	34	-
Multiple myeloma and immunoproliferative neoplasms (203)											
Black, both sexes											
1998	1,805	66	105	165	205	238	295	270	205	205	-
1997	1,708	78	86	154	184	246	268	248	199	190	-
1996	1,726	54	89	142	216	250	298	273	183	168	-
1995	1,716	64	106	146	232	228	273	268	184	169	-
1994	1,650	63	107	149	179	252	252	234	199	171	1
1993	1,644	55	94	123	194	250	289	262	164	175	-
1992	1,543	49	105	140	176	230	253	225	181	138	-
1991	1,580	49	84	128	166	252	280	258	176	132	-
1990	1,453	45	73	123	187	218	261	209	169	121	-
1989	1,387	53	83	107	169	234	213	195	158	136	-
1988	1,347	47	73	134	157	226	225	197	149	97	-
1987	1,347	48	78	98	174	220	243	201	128	119	1
1986	1,352	44	76	117	184	209	246	200	133	100	-
1985	1,273	42	63	111	170	206	228	176	153	89	-
1984	1,209	44	74	133	176	174	181	183	125	80	-
1983	1,230	40	78	116	177	189	196	193	129	80	-
1982	1,173	44	84	123	138	191	198	150	120	84	1
1981	1,134	40	61	116	145	196	194	177	103	73	-
1980	1,059	38	67	129	151	189	168	132	94	57	-
1979	1,008	25	89	120	144	182	152	131	69	63	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Multiple myeloma and immunoproliferative neoplasms (203)											
Black, male											
1998	849	30	61	79	100	125	136	117	98	81	-
1997	815	41	42	82	94	133	123	109	91	71	-
1996	835	28	53	71	105	129	142	124	83	66	-
1995	800	36	53	58	111	103	134	127	79	70	-
1994	809	30	63	79	98	141	121	97	84	72	-
1993	768	26	44	65	94	127	151	122	52	65	-
1992	755	17	54	78	101	113	127	105	78	57	-
1991	767	23	42	71	82	135	135	126	68	53	-
1990	745	23	38	63	105	114	135	100	80	57	-
1989	684	33	49	60	84	114	106	97	68	54	-
1988	631	22	37	67	77	104	104	98	71	33	-
1987	693	25	47	50	90	117	122	111	66	44	1
1986	686	22	32	64	96	114	128	88	75	43	-
1985	637	22	34	55	95	110	117	73	68	41	-
1984	619	23	39	70	95	91	91	97	59	30	-
1983	635	25	41	63	102	95	94	99	65	36	-
1982	606	22	51	67	62	99	103	77	62	40	1
1981	575	22	37	60	79	102	90	91	51	30	-
1980	536	13	33	72	77	104	82	69	43	22	-
1979	521	17	41	61	83	96	69	73	34	31	-
Multiple myeloma and immunoproliferative neoplasms (203)											
Black, female											
1998	956	36	44	86	105	113	159	153	107	124	-
1997	893	37	44	72	90	113	145	139	108	119	-
1996	891	26	36	71	111	121	156	149	100	102	-
1995	916	28	53	88	121	125	139	141	105	99	-
1994	841	33	44	70	81	111	131	137	115	99	1
1993	876	29	50	58	100	123	138	140	112	110	-
1992	788	32	51	62	75	117	126	120	103	81	-
1991	813	26	42	57	84	117	145	132	108	79	-
1990	708	22	35	60	82	104	126	109	89	64	-
1989	703	20	34	47	85	120	107	98	90	82	-
1988	716	25	36	67	80	122	121	99	78	64	-
1987	654	23	31	48	84	103	121	90	62	75	-
1986	666	22	44	53	88	95	118	112	58	57	-
1985	636	20	29	56	75	96	111	103	85	48	-
1984	590	21	35	63	81	83	90	86	66	50	-
1983	595	15	37	53	75	94	102	94	64	44	-
1982	567	22	33	56	76	92	95	73	58	44	-
1981	559	18	24	56	66	94	104	86	52	43	-
1980	523	25	34	57	74	85	86	63	51	35	-
1979	487	8	48	59	61	86	83	58	35	32	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All races, both sexes											
1998	55,597	1,961	2,725	3,630	4,733	6,252	8,065	8,593	7,438	8,562	2
1997	56,301	1,839	2,759	3,537	4,900	6,579	8,352	8,622	7,497	8,575	1
1996	57,121	1,966	2,753	3,502	4,993	6,956	8,648	8,695	7,505	8,452	2
1995	57,150	1,848	2,694	3,587	4,932	7,211	8,699	8,511	7,603	8,334	1
1994	57,088	1,872	2,593	3,595	5,275	7,313	8,673	8,422	7,399	8,072	3
1993	57,122	1,872	2,568	3,613	5,440	7,354	8,740	8,408	7,365	7,950	-
1992	55,052	1,784	2,419	3,505	5,465	7,209	8,312	8,115	7,024	7,538	4
1991	54,264	1,631	2,424	3,536	5,447	7,416	8,327	7,874	6,599	7,252	2
1990	52,775	1,587	2,244	3,545	5,442	7,208	7,757	7,834	6,561	6,934	6
1989	54,502	1,658	2,434	3,750	5,822	7,497	8,064	7,992	6,585	6,981	2
1988	54,859	1,581	2,448	4,059	6,082	7,514	8,134	7,982	6,556	6,802	2
1987	54,652	1,484	2,491	4,084	6,224	7,558	8,144	7,835	6,437	6,680	4
1986	54,562	1,569	2,496	4,148	6,280	7,330	8,270	7,741	6,375	6,387	10
1985	53,420	1,471	2,636	4,198	6,293	7,138	7,885	7,407	6,086	6,357	8
1984	52,073	1,444	2,518	4,273	6,026	7,031	7,775	7,301	5,888	5,978	10
1983	51,521	1,525	2,604	4,383	5,851	6,890	7,728	7,179	5,688	6,024	9
1982	50,123	1,504	2,596	4,174	5,829	6,799	7,344	6,982	5,530	5,700	9
1981	48,746	1,437	2,754	4,291	5,452	6,794	7,134	6,595	5,426	5,390	6
1980	48,413	1,507	2,865	4,200	5,426	6,725	7,042	6,519	5,257	5,195	4
1979	46,531	1,441	2,707	4,183	5,267	6,423	6,762	6,349	5,143	4,860	5
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All races, male											
1998	27,562	1,122	1,527	2,063	2,652	3,500	4,214	4,231	3,282	2,879	1
1997	27,665	1,037	1,541	1,949	2,704	3,641	4,341	4,204	3,239	2,880	1
1996	28,361	1,092	1,527	1,954	2,766	3,870	4,597	4,296	3,303	2,787	1
1995	28,370	1,003	1,499	2,043	2,772	3,968	4,582	4,183	3,311	2,795	1
1994	28,205	1,011	1,471	2,018	2,982	4,021	4,603	4,010	3,162	2,640	1
1993	28,505	1,056	1,451	2,053	3,068	4,017	4,680	4,136	3,173	2,626	-
1992	27,461	1,027	1,361	1,979	3,099	3,955	4,265	3,983	3,007	2,592	2
1991	27,174	925	1,363	2,069	3,127	4,070	4,352	3,760	2,854	2,362	2
1990	26,196	885	1,274	2,042	3,053	3,926	4,051	3,741	2,719	2,353	3
1989	27,534	952	1,384	2,129	3,319	4,174	4,284	3,896	2,853	2,336	1
1988	27,746	923	1,365	2,332	3,413	4,173	4,284	3,858	2,841	2,359	2
1987	27,594	819	1,382	2,348	3,504	4,167	4,376	3,809	2,764	2,186	2
1986	27,419	861	1,365	2,391	3,425	4,009	4,277	3,754	2,694	2,143	6
1985	27,049	789	1,482	2,364	3,530	3,973	4,105	3,578	2,587	2,173	2
1984	26,031	776	1,402	2,430	3,317	3,859	4,017	3,491	2,425	1,940	5
1983	26,053	867	1,466	2,485	3,306	3,773	4,006	3,530	2,376	2,051	4
1982	24,771	812	1,460	2,295	3,209	3,661	3,710	3,255	2,218	1,932	7
1981	24,064	804	1,490	2,414	3,009	3,640	3,653	2,954	2,205	1,799	5
1980	24,087	821	1,513	2,335	3,019	3,654	3,505	3,069	2,113	1,841	1
1979	23,132	808	1,457	2,264	2,905	3,437	3,503	2,898	2,101	1,686	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All races, female											
1998	28,035	839	1,198	1,567	2,081	2,752	3,851	4,362	4,156	5,683	1
1997	28,636	802	1,218	1,588	2,196	2,938	4,011	4,418	4,258	5,695	-
1996	28,760	874	1,226	1,548	2,227	3,086	4,051	4,399	4,202	5,665	1
1995	28,780	845	1,195	1,544	2,160	3,243	4,117	4,328	4,292	5,539	-
1994	28,883	861	1,122	1,577	2,293	3,292	4,070	4,412	4,237	5,432	2
1993	28,617	816	1,117	1,560	2,372	3,337	4,060	4,272	4,192	5,324	-
1992	27,591	757	1,058	1,526	2,366	3,254	4,047	4,132	4,017	4,946	2
1991	27,090	706	1,061	1,467	2,320	3,346	3,975	4,114	3,745	4,890	-
1990	26,579	702	970	1,503	2,389	3,282	3,706	4,093	3,842	4,581	3
1989	26,968	706	1,050	1,621	2,503	3,323	3,780	4,096	3,732	4,645	1
1988	27,113	658	1,083	1,727	2,669	3,341	3,850	4,124	3,715	4,443	-
1987	27,058	665	1,109	1,736	2,720	3,391	3,768	4,026	3,673	4,494	2
1986	27,143	708	1,131	1,757	2,855	3,321	3,993	3,987	3,681	4,244	4
1985	26,371	682	1,154	1,834	2,763	3,165	3,780	3,829	3,499	4,184	6
1984	26,042	668	1,116	1,843	2,709	3,172	3,758	3,810	3,463	4,038	5
1983	25,468	658	1,138	1,898	2,545	3,117	3,722	3,649	3,312	3,973	5
1982	25,352	692	1,136	1,879	2,620	3,138	3,634	3,727	3,312	3,768	2
1981	24,682	633	1,264	1,877	2,443	3,154	3,481	3,641	3,221	3,591	1
1980	24,326	686	1,352	1,865	2,407	3,071	3,537	3,450	3,144	3,354	3
1979	23,399	633	1,250	1,919	2,362	2,986	3,259	3,451	3,042	3,174	4
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
White, both sexes											
1998	48,360	1,491	2,191	2,975	3,933	5,398	7,099	7,751	6,801	7,860	2
1997	48,983	1,405	2,225	2,927	4,070	5,699	7,411	7,751	6,823	7,801	-
1996	49,790	1,517	2,269	2,876	4,242	6,038	7,625	7,824	6,815	7,722	2
1995	49,734	1,457	2,184	2,972	4,152	6,283	7,653	7,646	6,844	7,632	-
1994	49,791	1,490	2,072	2,950	4,452	6,354	7,687	7,643	6,691	7,397	2
1993	49,776	1,480	2,057	2,947	4,625	6,367	7,738	7,571	6,705	7,288	-
1992	48,055	1,413	1,946	2,850	4,651	6,255	7,337	7,329	6,388	6,920	3
1991	47,291	1,307	1,946	2,909	4,657	6,450	7,447	7,031	5,941	6,604	-
1990	45,959	1,264	1,784	2,930	4,607	6,214	6,892	7,029	5,953	6,376	1
1989	47,539	1,296	1,980	3,083	4,965	6,505	7,165	7,175	5,988	6,409	2
1988	47,966	1,239	1,976	3,398	5,177	6,582	7,224	7,157	5,976	6,250	-
1987	47,712	1,156	1,989	3,429	5,339	6,621	7,252	7,019	5,818	6,078	3
1986	47,840	1,242	2,042	3,479	5,346	6,461	7,399	6,973	5,822	5,833	6
1985	46,658	1,179	2,096	3,495	5,451	6,242	7,000	6,619	5,489	5,848	6
1984	45,481	1,142	2,037	3,548	5,199	6,154	6,852	6,557	5,346	5,523	4
1983	45,108	1,196	2,136	3,667	5,037	6,065	6,791	6,428	5,188	5,592	5
1982	44,112	1,190	2,129	3,545	5,046	5,983	6,544	6,310	5,036	5,330	6
1981	42,922	1,173	2,292	3,675	4,706	5,946	6,387	5,955	4,925	5,021	3
1980	42,539	1,216	2,333	3,572	4,659	5,904	6,266	5,893	4,814	4,821	4
1979	40,962	1,184	2,264	3,549	4,549	5,606	5,959	5,729	4,743	4,520	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
White, male											
1998	24,039	869	1,221	1,710	2,239	3,035	3,744	3,846	3,038	2,646	1
1997	24,076	799	1,231	1,613	2,254	3,178	3,892	3,808	2,985	2,632	-
1996	24,728	841	1,247	1,607	2,365	3,375	4,083	3,899	3,028	2,548	1
1995	24,650	783	1,222	1,692	2,364	3,459	4,064	3,784	3,017	2,551	-
1994	24,625	787	1,169	1,668	2,547	3,528	4,105	3,685	2,871	2,427	1
1993	24,802	823	1,160	1,677	2,622	3,486	4,152	3,770	2,900	2,394	-
1992	24,014	816	1,103	1,620	2,666	3,474	3,782	3,618	2,745	2,391	1
1991	23,603	729	1,088	1,694	2,702	3,561	3,907	3,360	2,570	2,134	-
1990	22,681	704	996	1,709	2,576	3,394	3,590	3,361	2,460	2,162	1
1989	23,850	744	1,118	1,733	2,831	3,651	3,815	3,492	2,574	2,102	1
1988	24,203	712	1,088	1,967	2,922	3,665	3,829	3,450	2,586	2,173	-
1987	23,989	619	1,094	1,983	3,017	3,671	3,909	3,412	2,492	1,945	1
1986	23,980	669	1,118	2,005	2,946	3,560	3,830	3,390	2,449	1,936	4
1985	23,490	631	1,190	1,963	3,072	3,488	3,621	3,189	2,316	1,968	2
1984	22,619	613	1,131	2,008	2,849	3,409	3,540	3,131	2,204	1,771	2
1983	22,686	686	1,219	2,087	2,833	3,335	3,508	3,136	2,151	1,890	1
1982	21,638	641	1,189	1,943	2,787	3,225	3,285	2,936	2,010	1,785	5
1981	21,092	647	1,246	2,078	2,595	3,199	3,264	2,678	1,980	1,653	2
1980	21,088	664	1,229	2,008	2,601	3,217	3,112	2,782	1,908	1,692	1
1979	20,252	673	1,208	1,927	2,519	3,025	3,081	2,599	1,904	1,561	1
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
White, female											
1998	24,321	622	970	1,265	1,694	2,363	3,355	3,905	3,763	5,214	1
1997	24,907	606	994	1,314	1,816	2,521	3,519	3,943	3,838	5,169	-
1996	25,062	676	1,022	1,269	1,877	2,663	3,542	3,925	3,787	5,174	1
1995	25,084	674	962	1,280	1,788	2,824	3,589	3,862	3,827	5,081	-
1994	25,166	703	903	1,282	1,905	2,826	3,582	3,958	3,820	4,970	1
1993	24,974	657	897	1,270	2,003	2,881	3,586	3,801	3,805	4,894	-
1992	24,041	597	843	1,230	1,985	2,781	3,555	3,711	3,643	4,529	2
1991	23,688	578	858	1,215	1,955	2,889	3,540	3,671	3,371	4,470	-
1990	23,278	560	788	1,221	2,031	2,820	3,302	3,668	3,493	4,214	-
1989	23,689	552	862	1,350	2,134	2,854	3,350	3,683	3,414	4,307	1
1988	23,763	527	888	1,431	2,255	2,917	3,395	3,707	3,390	4,077	-
1987	23,723	537	895	1,446	2,322	2,950	3,343	3,607	3,326	4,133	2
1986	23,860	573	924	1,474	2,400	2,901	3,569	3,583	3,373	3,897	2
1985	23,168	548	906	1,532	2,379	2,754	3,379	3,430	3,173	3,880	4
1984	22,862	529	906	1,540	2,350	2,745	3,312	3,426	3,142	3,752	2
1983	22,422	510	917	1,580	2,204	2,730	3,283	3,292	3,037	3,702	4
1982	22,474	549	940	1,602	2,259	2,758	3,259	3,374	3,026	3,545	1
1981	21,830	526	1,046	1,597	2,111	2,747	3,123	3,277	2,945	3,368	1
1980	21,451	552	1,104	1,564	2,058	2,687	3,154	3,111	2,906	3,129	3
1979	20,710	511	1,056	1,622	2,030	2,581	2,878	3,130	2,839	2,959	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All other, both sexes											
1998	7,237	470	534	655	800	854	966	842	637	702	-
1997	7,318	434	534	610	830	880	941	871	674	774	1
1996	7,331	449	484	626	751	918	1,023	871	690	730	-
1995	7,416	391	510	615	780	928	1,046	865	759	702	1
1994	7,297	382	521	645	823	959	986	779	708	675	1
1993	7,346	392	511	666	815	987	1,002	837	660	662	-
1992	6,997	371	473	655	814	954	975	786	636	618	1
1991	6,973	324	478	627	790	966	880	843	658	648	2
1990	6,816	323	460	615	835	994	865	805	608	558	5
1989	6,963	362	454	667	857	992	899	817	597	572	-
1988	6,893	342	472	661	905	932	910	825	580	552	2
1987	6,940	328	502	655	885	937	892	816	619	602	1
1986	6,722	327	454	669	934	869	871	768	553	554	4
1985	6,762	292	540	703	842	896	885	788	597	509	2
1984	6,592	302	481	725	827	877	923	744	542	455	6
1983	6,413	329	468	716	814	825	937	751	500	432	4
1982	6,011	314	467	629	783	816	800	672	494	370	3
1981	5,824	264	462	616	746	848	747	640	501	369	3
1980	5,874	291	532	628	767	821	776	626	443	374	-
1979	5,569	257	443	634	718	817	803	620	400	340	1
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All other, male											
1998	3,523	253	306	353	413	465	470	385	244	233	-
1997	3,589	238	310	336	450	463	449	396	254	248	1
1996	3,633	251	280	347	401	495	514	397	275	239	-
1995	3,720	220	277	351	408	509	518	399	294	244	1
1994	3,580	224	302	350	435	493	498	325	291	213	-
1993	3,703	233	291	376	446	531	528	366	273	232	-
1992	3,447	211	258	359	433	481	483	365	262	201	1
1991	3,571	196	275	375	425	509	445	400	284	228	2
1990	3,515	181	278	333	477	532	461	380	259	191	2
1989	3,684	208	266	396	488	523	469	404	279	234	-
1988	3,543	211	277	365	491	508	455	408	255	186	2
1987	3,605	200	288	365	487	496	467	397	272	241	1
1986	3,439	192	247	386	479	449	447	364	245	207	2
1985	3,559	158	292	401	458	485	484	389	271	205	-
1984	3,412	163	271	422	468	450	477	360	221	169	3
1983	3,367	181	247	398	473	438	498	394	225	161	3
1982	3,133	171	271	352	422	436	425	319	208	147	2
1981	2,972	157	244	336	414	441	389	276	225	146	3
1980	2,999	157	284	327	418	437	393	287	205	149	-
1979	2,880	135	249	337	386	412	422	299	197	125	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
All other, female											
1998	3,714	217	228	302	387	389	496	457	393	469	-
1997	3,729	196	224	274	380	417	492	475	420	526	-
1996	3,698	198	204	279	350	423	509	474	415	491	-
1995	3,696	171	233	264	372	419	528	466	465	458	-
1994	3,717	158	219	295	388	466	488	454	417	462	1
1993	3,643	159	220	290	369	456	474	471	387	430	-
1992	3,550	160	215	296	381	473	492	421	374	417	-
1991	3,402	128	203	252	365	457	435	443	374	420	-
1990	3,301	142	182	282	358	462	404	425	349	367	3
1989	3,279	154	188	271	369	469	430	413	318	338	-
1988	3,350	131	195	296	414	424	455	417	325	366	-
1987	3,335	128	214	290	398	441	425	419	347	361	-
1986	3,283	135	207	283	455	420	424	404	308	347	2
1985	3,203	134	248	302	384	411	401	399	326	304	2
1984	3,180	139	210	303	359	427	446	384	321	286	3
1983	3,046	148	221	318	341	387	439	357	275	271	1
1982	2,878	143	196	277	361	380	375	353	286	223	1
1981	2,852	107	218	280	332	407	358	364	276	223	-
1980	2,875	134	248	301	349	384	383	339	238	225	-
1979	2,689	122	194	297	332	405	381	321	203	215	1
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
Black, both sexes											
1998	6,097	399	453	567	681	723	814	684	535	606	-
1997	6,281	385	471	531	714	743	803	739	580	668	1
1996	6,243	381	409	553	655	787	864	729	586	632	-
1995	6,419	345	446	539	680	809	904	772	640	595	1
1994	6,356	331	465	568	730	841	861	657	612	578	1
1993	6,352	330	459	579	718	853	871	710	577	571	-
1992	6,127	334	411	585	714	840	864	691	544	528	-
1991	6,114	278	423	549	706	882	765	723	583	564	2
1990	6,043	282	409	549	764	891	764	709	531	488	4
1989	6,200	317	407	618	775	871	807	731	514	507	-
1988	6,164	300	430	590	812	847	824	733	506	495	2
1987	6,275	299	461	600	822	855	801	741	552	535	1
1986	6,084	295	414	619	863	798	800	684	499	497	4
1985	6,093	272	487	637	769	829	799	692	523	468	2
1984	5,969	277	440	659	758	792	848	682	486	401	5
1983	5,813	302	421	655	750	751	869	685	448	381	2
1982	5,509	293	422	575	733	752	732	611	455	341	3
1981	5,355	247	430	585	700	779	692	576	453	331	2
1980	5,435	268	504	592	707	758	721	573	406	343	-
1979	5,177	243	415	589	675	775	747	571	364	317	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
Black, male											
1998	2,956	215	265	310	362	397	401	308	194	187	-
1997	3,106	214	278	297	398	397	382	336	214	207	1
1996	3,108	216	242	317	353	426	444	325	229	192	-
1995	3,202	197	246	305	365	443	440	352	239	190	1
1994	3,127	191	275	311	397	426	444	273	241	177	-
1993	3,210	201	259	334	398	464	458	304	236	194	-
1992	3,026	196	233	328	382	424	430	320	216	157	-
1991	3,165	172	253	333	390	469	392	347	240	200	2
1990	3,130	163	247	306	437	476	414	333	220	166	2
1989	3,286	184	247	371	443	455	416	356	240	204	-
1988	3,181	188	259	331	446	464	418	361	219	158	2
1987	3,275	185	266	343	457	456	424	359	240	206	1
1986	3,109	176	229	358	448	415	406	316	209	185	2
1985	3,220	148	276	362	427	450	433	335	239	185	-
1984	3,098	153	248	387	438	409	437	327	192	150	2
1983	3,067	172	223	372	434	403	459	352	202	143	1
1982	2,867	161	242	326	393	407	384	286	194	136	2
1981	2,744	148	236	320	389	406	354	251	203	129	2
1980	2,773	145	269	307	389	404	365	261	187	134	-
1979	2,686	130	234	316	364	394	390	283	177	117	-
All other and unspecified malignant neoplasms (152,156,158-160,163-171,173,181,183.2-184,186-187,189.2-190,193-199)											
Black, female											
1998	3,141	184	188	257	319	326	413	376	341	419	-
1997	3,175	171	193	234	316	346	421	403	366	461	-
1996	3,135	165	167	236	302	361	420	404	357	440	-
1995	3,217	148	200	234	315	366	464	420	401	405	-
1994	3,229	140	190	257	333	415	417	384	371	401	1
1993	3,142	129	200	245	320	389	413	406	341	377	-
1992	3,101	138	178	257	332	416	434	371	328	371	-
1991	2,949	106	170	216	316	413	373	376	343	364	-
1990	2,913	119	162	243	327	415	350	376	311	322	2
1989	2,914	133	160	247	332	416	391	375	274	303	-
1988	2,983	112	171	259	366	383	406	372	287	337	-
1987	3,000	114	195	257	365	399	377	382	312	329	-
1986	2,975	119	185	261	415	383	394	368	290	312	2
1985	2,873	124	211	275	342	379	366	357	284	283	2
1984	2,871	124	192	272	320	383	411	355	294	251	3
1983	2,746	130	198	283	316	348	410	333	246	238	1
1982	2,642	132	180	249	340	345	348	325	261	205	1
1981	2,611	99	194	265	311	373	338	325	250	202	-
1980	2,662	123	235	285	318	354	356	312	219	209	-
1979	2,491	113	181	273	311	381	357	288	187	200	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All races, both sexes											
1998	7,933	220	297	342	431	587	933	1,097	1,261	1,941	-
1997	7,659	232	271	332	394	647	870	1,152	1,201	1,738	-
1996	7,647	251	248	334	456	607	907	1,144	1,193	1,672	1
1995	7,759	221	281	316	488	694	918	1,100	1,197	1,668	-
1994	7,517	238	270	318	435	642	852	1,091	1,136	1,605	-
1993	7,441	213	264	308	463	708	872	1,069	1,142	1,534	2
1992	7,070	204	234	289	468	673	898	1,040	1,035	1,390	-
1991	6,994	179	207	307	480	694	895	1,003	1,094	1,337	-
1990	6,752	169	230	286	469	681	848	1,034	972	1,214	1
1989	6,701	178	232	309	504	682	865	980	1,019	1,187	1
1988	6,735	186	227	321	521	688	878	990	984	1,140	1
1987	6,570	164	204	321	520	771	875	983	946	1,056	-
1986	6,730	187	252	365	553	656	885	1,020	943	1,046	2
1985	6,699	173	248	386	567	710	955	938	898	1,038	2
1984	6,436	167	198	387	546	723	898	972	844	951	-
1983	6,483	174	274	380	580	737	831	966	834	941	2
1982	6,460	199	279	434	629	769	892	921	756	818	1
1981	6,235	180	306	378	609	739	881	911	742	746	2
1980	6,193	240	301	459	598	748	821	796	734	698	-
1979	5,904	219	289	492	560	743	807	779	609	644	1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All races, male											
1998	3,516	116	171	188	209	305	463	533	498	579	-
1997	3,366	122	141	183	211	324	418	548	495	506	-
1996	3,412	139	126	191	254	312	419	527	472	512	1
1995	3,474	116	162	171	248	369	431	499	462	507	-
1994	3,404	117	151	186	235	349	422	474	458	481	-
1993	3,424	119	127	168	274	367	452	492	443	489	1
1992	3,256	113	131	149	268	359	439	490	404	442	-
1991	3,179	103	117	167	262	358	438	452	445	395	-
1990	3,012	82	112	138	240	361	442	429	389	378	-
1989	3,048	91	134	159	252	342	421	429	433	392	-
1988	3,012	96	123	160	258	363	398	440	379	367	1
1987	3,065	89	110	168	282	404	464	450	397	328	-
1986	3,114	106	138	187	298	296	428	466	381	358	2
1985	3,008	84	144	203	280	349	436	420	341	336	1
1984	2,920	84	108	197	285	345	429	435	329	315	-
1983	2,985	88	133	208	306	372	391	424	344	307	2
1982	2,977	93	149	232	315	376	417	402	316	293	1
1981	2,934	88	160	198	312	388	435	418	301	249	-
1980	2,879	107	142	236	310	370	385	367	317	232	-
1979	2,704	108	135	249	278	365	381	343	254	220	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All races, female											
1998	4,417	104	126	154	222	282	470	564	763	1,362	-
1997	4,293	110	130	149	183	323	452	604	706	1,232	-
1996	4,235	112	122	143	202	295	488	617	721	1,160	-
1995	4,285	105	119	145	240	325	487	601	735	1,161	-
1994	4,113	121	119	132	200	293	430	617	678	1,124	-
1993	4,017	94	137	140	189	341	420	577	699	1,045	1
1992	3,814	91	103	140	200	314	459	550	631	948	-
1991	3,815	76	90	140	218	336	457	551	649	942	-
1990	3,740	87	118	148	229	320	406	605	583	836	1
1989	3,653	87	98	150	252	340	444	551	586	795	1
1988	3,723	90	104	161	263	325	480	550	605	773	-
1987	3,505	75	94	153	238	367	411	533	549	728	-
1986	3,616	81	114	178	255	360	457	554	562	688	-
1985	3,691	89	104	183	287	361	519	518	557	702	1
1984	3,516	83	90	190	261	378	469	537	515	636	-
1983	3,498	86	141	172	274	365	440	542	490	634	-
1982	3,483	106	130	202	314	393	475	519	440	525	-
1981	3,301	92	146	180	297	351	446	493	441	497	2
1980	3,314	133	159	223	288	378	436	429	417	466	-
1979	3,200	111	154	243	282	378	426	436	355	424	1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
White, both sexes											
1998	6,940	170	240	287	365	509	805	989	1,163	1,790	-
1997	6,682	187	221	282	326	541	764	1,035	1,087	1,606	-
1996	6,657	193	212	287	383	536	795	1,015	1,065	1,548	-
1995	6,748	180	234	269	414	585	803	1,007	1,089	1,498	-
1994	6,595	186	225	265	360	561	763	983	1,031	1,486	-
1993	6,539	165	221	257	390	613	776	964	1,038	1,427	1
1992	6,191	173	190	235	404	570	788	929	934	1,286	-
1991	6,136	143	172	251	405	591	794	906	997	1,243	-
1990	5,908	132	177	235	408	580	756	926	873	1,132	-
1989	5,858	149	191	251	444	585	760	880	943	1,077	1
1988	5,870	150	188	269	437	584	777	869	898	1,051	-
1987	5,802	142	173	267	446	676	785	881	868	970	-
1986	5,895	148	200	311	461	571	782	907	867	970	2
1985	5,839	134	203	320	484	609	826	849	822	968	1
1984	5,656	131	158	337	469	638	785	882	766	873	-
1983	5,728	144	230	322	514	646	720	878	770	883	1
1982	5,654	165	229	357	535	674	791	830	699	772	1
1981	5,508	145	257	327	522	664	770	840	687	688	1
1980	5,383	188	241	403	499	615	731	726	670	651	-
1979	5,201	173	234	424	484	652	730	717	570	610	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
White, male											
1998	3,079	89	136	158	184	264	415	486	460	535	-
1997	2,926	102	120	155	171	273	370	489	450	467	-
1996	2,973	111	104	162	214	279	377	477	413	474	-
1995	3,025	95	141	151	214	320	379	467	418	445	-
1994	2,997	97	125	155	199	309	381	439	415	447	-
1993	2,997	92	104	140	235	321	408	444	405	457	1
1992	2,843	93	106	124	238	305	388	442	359	411	-
1991	2,783	80	97	142	223	310	395	415	404	357	-
1990	2,658	69	86	116	213	310	399	392	358	360	-
1989	2,685	77	112	126	223	299	375	387	409	356	-
1988	2,639	80	102	137	220	315	361	382	354	333	-
1987	2,683	79	92	142	242	356	416	399	360	295	-
1986	2,740	86	108	169	255	258	378	424	352	331	2
1985	2,632	64	121	179	245	302	383	373	315	316	1
1984	2,568	70	83	178	239	304	381	394	306	287	-
1983	2,639	78	110	180	273	330	338	393	313	286	1
1982	2,617	78	125	190	272	331	381	366	288	283	1
1981	2,625	77	132	175	274	354	387	381	282	233	-
1980	2,514	85	118	205	261	310	344	334	285	218	-
1979	2,406	92	117	217	237	329	351	321	240	210	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
White, female											
1998	3,861	81	104	129	181	245	390	503	703	1,255	-
1997	3,756	85	101	127	155	268	394	546	637	1,139	-
1996	3,684	82	108	125	169	257	418	538	652	1,074	-
1995	3,723	85	93	118	200	265	424	540	671	1,053	-
1994	3,598	89	100	110	161	252	382	544	616	1,039	-
1993	3,542	73	117	117	155	292	368	520	633	970	-
1992	3,348	80	84	111	166	265	400	487	575	875	-
1991	3,353	63	75	109	182	281	399	491	593	886	-
1990	3,250	63	91	119	195	270	357	534	515	772	-
1989	3,173	72	79	125	221	286	385	493	534	721	1
1988	3,231	70	86	132	217	269	416	487	544	718	-
1987	3,119	63	81	125	204	320	369	482	508	675	-
1986	3,155	62	92	142	206	313	404	483	515	639	-
1985	3,207	70	82	141	239	307	443	476	507	652	-
1984	3,088	61	75	159	230	334	404	488	460	586	-
1983	3,089	66	120	142	241	316	382	485	457	597	-
1982	3,037	87	104	167	263	343	410	464	411	489	-
1981	2,883	68	125	152	248	310	383	459	405	455	1
1980	2,869	103	123	198	238	305	387	392	385	433	-
1979	2,795	81	117	207	247	323	379	396	330	400	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All other, both sexes											
1998	993	50	57	55	66	78	128	108	98	151	-
1997	977	45	50	50	68	106	106	117	114	132	-
1996	990	58	36	47	73	71	112	129	128	124	1
1995	1,011	41	47	47	74	109	115	93	108	170	-
1994	922	52	45	53	75	81	89	108	105	119	-
1993	902	48	43	51	73	95	96	105	104	107	1
1992	879	31	44	54	64	103	110	111	101	104	-
1991	858	36	35	56	75	103	101	97	97	94	-
1990	844	37	53	51	61	101	92	108	99	82	1
1989	843	29	41	58	60	97	105	100	76	110	-
1988	865	36	39	52	84	104	101	121	86	89	1
1987	768	22	31	54	74	95	90	102	78	86	-
1986	835	39	52	54	92	85	103	113	76	76	-
1985	860	39	45	66	83	101	129	89	76	70	1
1984	780	36	40	50	77	85	113	90	78	78	-
1983	755	30	44	58	66	91	111	88	64	58	1
1982	806	34	50	77	94	95	101	91	57	46	-
1981	727	35	49	51	87	75	111	71	55	58	1
1980	810	52	60	56	99	133	90	70	64	47	-
1979	703	46	55	68	76	91	77	62	39	34	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All other, male											
1998	437	27	35	30	25	41	48	47	38	44	-
1997	440	20	21	28	40	51	48	59	45	39	-
1996	439	28	22	29	40	33	42	50	59	38	1
1995	449	21	21	20	34	49	52	32	44	62	-
1994	407	20	26	31	36	40	41	35	43	34	-
1993	427	27	23	28	39	46	44	48	38	32	-
1992	413	20	25	25	30	54	51	48	45	31	-
1991	396	23	20	25	39	48	43	37	41	38	-
1990	354	13	26	22	27	51	43	37	31	18	-
1989	363	14	22	33	29	43	46	42	24	36	-
1988	373	16	21	23	38	48	37	58	25	34	1
1987	382	10	18	26	40	48	48	51	37	33	-
1986	374	20	30	18	43	38	50	42	29	27	-
1985	376	20	23	24	35	47	53	47	26	20	-
1984	352	14	25	19	46	41	48	41	23	28	-
1983	346	10	23	28	33	42	53	31	31	21	1
1982	360	15	24	42	43	45	36	36	28	10	-
1981	309	11	28	23	38	34	48	37	19	16	-
1980	365	22	24	31	49	60	41	33	32	14	-
1979	298	16	18	32	41	36	30	22	14	10	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
All other, female											
1998	556	23	22	25	41	37	80	61	60	107	-
1997	537	25	29	22	28	55	58	58	69	93	-
1996	551	30	14	18	33	38	70	79	69	86	-
1995	562	20	26	27	40	60	63	61	64	108	-
1994	515	32	19	22	39	41	48	73	62	85	-
1993	475	21	20	23	34	49	52	57	66	75	1
1992	466	11	19	29	34	49	59	63	56	73	-
1991	462	13	15	31	36	55	58	60	56	56	-
1990	490	24	27	29	34	50	49	71	68	64	1
1989	480	15	19	25	31	54	59	58	52	74	-
1988	492	20	18	29	46	56	64	63	61	55	-
1987	386	12	13	28	34	47	42	51	41	53	-
1986	461	19	22	36	49	47	53	71	47	49	-
1985	484	19	22	42	48	54	76	42	50	50	1
1984	428	22	15	31	31	44	65	49	55	50	-
1983	409	20	21	30	33	49	58	57	33	37	-
1982	446	19	26	35	51	50	65	55	29	36	-
1981	418	24	21	28	49	41	63	34	36	42	1
1980	445	30	36	25	50	73	49	37	32	33	-
1979	405	30	37	36	35	55	47	40	25	24	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
Black, both sexes											
1998	855	45	44	47	58	67	112	93	84	128	-
1997	861	41	41	46	64	92	93	106	98	117	-
1996	855	51	35	38	68	62	96	105	106	108	1
1995	878	35	39	42	65	99	100	85	95	144	-
1994	804	43	43	46	69	72	80	96	90	100	-
1993	798	42	35	40	67	87	88	91	94	93	1
1992	778	28	38	50	53	90	99	103	87	93	-
1991	777	32	32	50	71	98	93	85	86	84	-
1990	747	31	44	43	54	93	84	91	91	76	1
1989	742	27	36	55	53	86	92	90	68	97	-
1988	784	33	36	48	78	99	96	107	77	82	1
1987	689	20	29	49	68	84	81	93	73	73	-
1986	756	37	45	51	83	82	91	107	68	72	-
1985	776	37	43	63	77	95	114	81	66	59	1
1984	697	34	39	47	67	79	98	78	70	69	-
1983	684	28	40	52	60	86	100	82	58	51	1
1982	748	29	45	73	87	89	98	86	55	41	-
1981	665	32	46	47	79	68	100	69	48	51	1
1980	749	51	57	51	94	126	80	68	59	39	-
1979	654	42	51	65	70	84	72	57	34	33	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
Black, male											
1998	366	24	28	26	21	33	41	39	32	32	-
1997	376	17	16	24	37	46	41	55	39	27	-
1996	372	25	22	24	38	28	33	40	46	30	1
1995	394	19	18	20	29	45	48	31	37	52	-
1994	352	14	25	26	34	36	36	30	37	26	-
1993	374	22	17	23	37	43	41	42	32	24	-
1992	360	17	22	23	24	44	48	43	36	28	-
1991	354	22	19	23	38	44	40	30	37	32	-
1990	316	11	24	18	26	46	41	30	29	16	-
1989	321	12	19	32	27	40	38	37	21	31	-
1988	330	14	20	21	35	46	35	48	20	30	1
1987	339	10	17	23	37	40	43	47	33	28	-
1986	332	18	26	17	37	37	46	40	23	26	-
1985	332	19	22	23	31	44	46	43	21	14	-
1984	308	12	25	18	40	38	40	32	20	24	-
1983	307	9	22	24	30	39	47	28	28	17	1
1982	331	12	22	41	38	42	35	33	27	9	-
1981	273	10	27	20	33	30	41	36	15	12	-
1980	339	21	23	28	48	59	35	32	29	11	-
1979	276	14	18	30	37	33	29	21	12	10	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (210-239)											
Black, female											
1998	489	21	16	21	37	34	71	54	52	96	-
1997	485	24	25	22	27	46	52	51	59	90	-
1996	483	26	13	14	30	34	63	65	60	78	-
1995	484	16	21	22	36	54	52	54	58	92	-
1994	452	29	18	20	35	36	44	66	53	74	-
1993	424	20	18	17	30	44	47	49	62	69	1
1992	418	11	16	27	29	46	51	60	51	65	-
1991	423	10	13	27	33	54	53	55	49	52	-
1990	431	20	20	25	28	47	43	61	62	60	1
1989	421	15	17	23	26	46	54	53	47	66	-
1988	454	19	16	27	43	53	61	59	57	52	-
1987	350	10	12	26	31	44	38	46	40	45	-
1986	424	19	19	34	46	45	45	67	45	46	-
1985	444	18	21	40	46	51	68	38	45	45	1
1984	389	22	14	29	27	41	58	46	50	45	-
1983	377	19	18	28	30	47	53	54	30	34	-
1982	417	17	23	32	49	47	63	53	28	32	-
1981	392	22	19	27	46	38	59	33	33	39	1
1980	410	30	34	23	46	67	45	36	30	28	-
1979	378	28	33	35	33	51	43	36	22	23	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
All races, both sexes											
1998	4,544	101	101	91	144	214	358	555	706	1,693	-
1997	4,471	98	92	83	138	207	357	532	726	1,687	-
1996	4,332	108	92	119	140	210	378	501	665	1,532	1
1995	4,556	84	87	103	146	230	373	531	709	1,627	-
1994	4,380	99	99	83	148	212	377	537	661	1,455	-
1993	4,315	78	87	103	133	223	341	514	682	1,496	1
1992	4,246	85	85	90	134	235	381	520	614	1,379	-
1991	4,121	87	79	92	161	233	346	553	642	1,331	-
1990	4,092	61	95	98	174	250	354	496	628	1,299	-
1989	3,993	86	80	96	168	276	364	468	641	1,207	1
1988	3,816	75	60	120	145	225	354	533	595	1,186	-
1987	3,813	59	73	115	151	233	364	504	573	1,161	-
1986	3,708	56	60	98	146	222	353	495	628	1,100	-
1985	3,694	59	74	122	170	235	360	506	586	1,047	-
1984	3,527	55	70	96	164	266	342	475	535	989	1
1983	3,487	55	65	132	183	259	340	479	517	898	2
1982	3,292	44	62	90	172	256	329	438	526	832	-
1981	3,300	69	75	115	160	285	357	438	473	815	-
1980	3,238	54	83	105	177	249	340	446	476	754	-
1979	3,171	57	85	117	155	218	351	388	484	728	1
Anemias (280-285)											
All races, male											
1998	1,873	52	51	55	75	106	193	241	306	498	-
1997	1,805	47	44	43	71	92	164	266	294	493	-
1996	1,820	54	46	74	72	105	202	240	302	428	1
1995	1,855	47	42	50	61	127	182	224	297	476	-
1994	1,909	48	56	39	77	103	196	254	266	487	-
1993	1,789	32	42	51	72	109	161	248	256	463	-
1992	1,825	48	39	38	66	114	192	225	248	456	-
1991	1,758	42	35	48	76	113	166	236	263	452	-
1990	1,770	34	52	50	74	113	187	236	250	412	-
1989	1,688	51	36	44	86	123	167	206	271	376	1
1988	1,614	42	31	58	74	115	190	244	224	347	-
1987	1,670	31	39	62	90	100	203	236	238	348	-
1986	1,633	37	31	45	67	109	169	236	259	384	-
1985	1,633	31	36	54	98	107	178	231	258	345	-
1984	1,545	22	42	44	88	140	164	222	230	323	1
1983	1,571	27	36	64	95	128	170	218	206	323	-
1982	1,452	22	29	44	94	125	153	212	214	271	-
1981	1,519	34	34	53	95	132	178	217	214	283	-
1980	1,421	25	38	55	84	128	156	185	201	257	-
1979	1,475	27	42	59	79	125	176	177	213	258	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
All races, female											
1998	2,671	49	50	36	69	108	165	314	400	1,195	-
1997	2,666	51	48	40	67	115	193	266	432	1,194	-
1996	2,512	54	46	45	68	105	176	261	363	1,104	-
1995	2,701	37	45	53	85	103	191	307	412	1,151	-
1994	2,471	51	43	44	71	109	181	283	395	968	-
1993	2,526	46	45	52	61	114	180	266	426	1,033	1
1992	2,421	37	46	52	68	121	189	295	366	923	-
1991	2,363	45	44	44	85	120	180	317	379	879	-
1990	2,322	27	43	48	100	137	167	260	378	887	-
1989	2,305	35	44	52	82	153	197	262	370	831	-
1988	2,202	33	29	62	71	110	164	289	371	839	-
1987	2,143	28	34	53	61	133	161	268	335	813	-
1986	2,075	19	29	53	79	113	184	259	369	716	-
1985	2,061	28	38	68	72	128	182	275	328	702	-
1984	1,982	33	28	52	76	126	178	253	305	666	-
1983	1,916	28	29	68	88	131	170	261	311	575	2
1982	1,840	22	33	46	78	131	176	226	312	561	-
1981	1,781	35	41	62	65	153	179	221	259	532	-
1980	1,817	29	45	50	93	121	184	261	275	497	-
1979	1,696	30	43	58	76	93	175	211	271	470	1
Anemias (280-285)											
White, both sexes											
1998	3,503	50	49	55	101	163	296	479	635	1,519	-
1997	3,473	30	58	45	106	155	295	476	652	1,519	-
1996	3,340	43	49	76	100	173	330	435	591	1,377	-
1995	3,461	35	46	71	100	194	301	443	621	1,465	-
1994	3,332	37	55	61	108	163	316	473	586	1,315	-
1993	3,308	30	46	63	100	180	286	439	618	1,346	-
1992	3,241	44	50	61	111	192	314	435	552	1,261	-
1991	3,180	30	49	61	126	176	291	482	575	1,197	-
1990	3,136	26	53	68	137	196	299	418	571	1,182	-
1989	3,089	43	41	69	115	228	310	409	581	1,094	-
1988	3,034	36	38	83	119	182	316	465	533	1,083	-
1987	3,027	29	51	86	123	189	313	451	527	1,074	-
1986	2,949	30	40	75	118	187	319	428	568	1,006	-
1985	2,931	25	49	95	128	196	306	446	529	964	-
1984	2,844	24	52	75	132	221	292	424	494	913	1
1983	2,786	30	37	92	150	217	299	438	472	836	1
1982	2,620	21	43	69	145	210	290	389	470	766	-
1981	2,658	43	48	90	131	240	319	394	430	764	-
1980	2,550	30	57	76	140	208	286	403	431	702	-
1979	2,504	36	59	87	125	182	304	342	440	672	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
White, male											
1998	1,391	30	25	32	49	84	157	209	279	446	-
1997	1,353	19	24	26	55	73	143	238	267	439	-
1996	1,372	22	25	50	53	91	179	208	274	389	-
1995	1,362	20	25	35	48	111	143	188	267	433	-
1994	1,423	21	34	29	59	80	162	227	238	444	-
1993	1,336	12	19	34	59	87	135	217	237	424	-
1992	1,349	29	24	27	60	91	161	190	226	422	-
1991	1,305	15	20	32	60	92	142	201	228	412	-
1990	1,274	15	30	33	55	93	155	191	224	372	-
1989	1,267	29	20	30	61	105	144	185	243	340	-
1988	1,247	22	18	44	64	92	177	212	200	316	-
1987	1,269	15	30	48	78	80	178	213	213	321	-
1986	1,272	20	23	36	56	93	157	208	236	348	-
1985	1,264	11	25	39	77	89	157	207	234	318	-
1984	1,203	10	31	37	70	114	140	200	211	297	1
1983	1,217	11	20	47	77	108	150	196	186	304	-
1982	1,132	10	21	34	81	106	138	188	197	249	-
1981	1,193	21	23	43	78	113	161	193	189	262	-
1980	1,085	15	32	39	69	105	131	171	181	241	-
1979	1,163	18	29	48	66	109	155	162	199	239	-
Anemias (280-285)											
White, female											
1998	2,112	20	24	23	52	79	139	270	356	1,073	-
1997	2,120	11	34	19	51	82	152	238	385	1,080	-
1996	1,968	21	24	26	47	82	151	227	317	988	-
1995	2,099	15	21	36	52	83	158	255	354	1,032	-
1994	1,909	16	21	32	49	83	154	246	348	871	-
1993	1,972	18	27	29	41	93	151	222	381	922	-
1992	1,892	15	26	34	51	101	153	245	326	839	-
1991	1,875	15	29	29	66	84	149	281	347	785	-
1990	1,862	11	23	35	82	103	144	227	347	810	-
1989	1,822	14	21	39	54	123	166	224	338	754	-
1988	1,787	14	20	39	55	90	139	253	333	767	-
1987	1,758	14	21	38	45	109	135	238	314	753	-
1986	1,677	10	17	39	62	94	162	220	332	658	-
1985	1,667	14	24	56	51	107	149	239	295	646	-
1984	1,641	14	21	38	62	107	152	224	283	616	-
1983	1,569	19	17	45	73	109	149	242	286	532	1
1982	1,488	11	22	35	64	104	152	201	273	517	-
1981	1,465	22	25	47	53	127	158	201	241	502	-
1980	1,465	15	25	37	71	103	155	232	250	461	-
1979	1,341	18	30	39	59	73	149	180	241	433	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
All other, both sexes											
1998	1,041	51	52	36	43	51	62	76	71	174	-
1997	998	68	34	38	32	52	62	56	74	168	-
1996	992	65	43	43	40	37	48	66	74	155	1
1995	1,095	49	41	32	46	36	72	88	88	162	-
1994	1,048	62	44	22	40	49	61	64	75	140	-
1993	1,007	48	41	40	33	43	55	75	64	150	1
1992	1,005	41	35	29	23	43	67	85	62	118	-
1991	941	57	30	31	35	57	55	71	67	134	-
1990	956	35	42	30	37	54	55	78	57	117	-
1989	904	43	39	27	53	48	54	59	60	113	1
1988	782	39	22	37	26	43	38	68	62	103	-
1987	786	30	22	29	28	44	51	53	46	87	-
1986	759	26	20	23	28	35	34	67	60	94	-
1985	763	34	25	27	42	39	54	60	57	83	-
1984	683	31	18	21	32	45	50	51	41	76	-
1983	701	25	28	40	33	42	41	41	45	62	1
1982	672	23	19	21	27	46	39	49	56	66	-
1981	642	26	27	25	29	45	38	44	43	51	-
1980	688	24	26	29	37	41	54	43	45	52	-
1979	667	21	26	30	30	36	47	46	44	56	1
Anemias (280-285)											
All other, male											
1998	482	22	26	23	26	22	36	32	27	52	-
1997	452	28	20	17	16	19	21	28	27	54	-
1996	448	32	21	24	19	14	23	32	28	39	1
1995	493	27	17	15	13	16	39	36	30	43	-
1994	486	27	22	10	18	23	34	27	28	43	-
1993	453	20	23	17	13	22	26	31	19	39	-
1992	476	19	15	11	6	23	31	35	22	34	-
1991	453	27	15	16	16	21	24	35	35	40	-
1990	496	19	22	17	19	20	32	45	26	40	-
1989	421	22	16	14	25	18	23	21	28	36	1
1988	367	20	13	14	10	23	13	32	24	31	-
1987	401	16	9	14	12	20	25	23	25	27	-
1986	361	17	8	9	11	16	12	28	23	36	-
1985	369	20	11	15	21	18	21	24	24	27	-
1984	342	12	11	7	18	26	24	22	19	26	-
1983	354	16	16	17	18	20	20	22	20	19	-
1982	320	12	8	10	13	19	15	24	17	22	-
1981	326	13	11	10	17	19	17	24	25	21	-
1980	336	10	6	16	15	23	25	14	20	16	-
1979	312	9	13	11	13	16	21	15	14	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
All other, female											
1998	559	29	26	13	17	29	26	44	44	122	-
1997	546	40	14	21	16	33	41	28	47	114	-
1996	544	33	22	19	21	23	25	34	46	116	-
1995	602	22	24	17	33	20	33	52	58	119	-
1994	562	35	22	12	22	26	27	37	47	97	-
1993	554	28	18	23	20	21	29	44	45	111	1
1992	529	22	20	18	17	20	36	50	40	84	-
1991	488	30	15	15	19	36	31	36	32	94	-
1990	460	16	20	13	18	34	23	33	31	77	-
1989	483	21	23	13	28	30	31	38	32	77	-
1988	415	19	9	23	16	20	25	36	38	72	-
1987	385	14	13	15	16	24	26	30	21	60	-
1986	398	9	12	14	17	19	22	39	37	58	-
1985	394	14	14	12	21	21	33	36	33	56	-
1984	341	19	7	14	14	19	26	29	22	50	-
1983	347	9	12	23	15	22	21	19	25	43	1
1982	352	11	11	11	14	27	24	25	39	44	-
1981	316	13	16	15	12	26	21	20	18	30	-
1980	352	14	20	13	22	18	29	29	25	36	-
1979	355	12	13	19	17	20	26	31	30	37	1
Anemias (280-285)											
Black, both sexes											
1998	953	49	49	34	41	42	46	68	61	156	-
1997	927	66	34	35	30	48	54	49	65	152	-
1996	929	62	41	40	38	31	43	58	68	141	1
1995	1,014	47	41	31	41	33	61	76	73	148	-
1994	991	58	44	22	37	48	53	61	68	124	-
1993	947	45	41	37	30	39	48	70	58	137	-
1992	947	41	32	26	22	37	63	79	59	106	-
1991	883	57	28	26	28	55	50	66	63	126	-
1990	887	32	40	24	34	49	49	71	54	106	-
1989	852	41	38	25	49	45	46	54	56	106	1
1988	731	39	22	36	26	38	34	61	57	93	-
1987	739	29	21	26	24	43	47	49	40	77	-
1986	706	25	20	23	26	33	31	57	54	83	-
1985	717	34	24	24	40	36	50	56	51	73	-
1984	640	30	17	20	28	39	45	46	40	72	-
1983	665	23	27	39	32	39	38	38	43	56	1
1982	632	23	19	19	27	42	33	46	51	59	-
1981	615	26	27	21	27	43	37	43	40	48	-
1980	660	23	26	28	36	35	53	38	42	50	-
1979	637	17	26	30	29	35	45	45	43	51	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Anemias (280-285)											
Black, male											
1998	429	20	25	21	24	15	26	28	22	44	-
1997	423	28	20	17	14	18	19	26	24	48	-
1996	419	30	20	21	18	13	20	28	27	32	1
1995	450	25	17	14	12	15	32	32	24	33	-
1994	461	24	22	10	17	22	31	25	25	36	-
1993	424	19	23	16	12	18	23	31	16	32	-
1992	450	19	13	10	6	21	29	30	20	28	-
1991	420	27	14	15	14	19	21	32	32	36	-
1990	460	19	21	13	17	18	29	41	23	35	-
1989	401	21	15	13	24	16	18	19	26	35	1
1988	337	20	13	13	10	18	12	26	22	26	-
1987	378	16	9	12	9	20	22	21	22	22	-
1986	334	16	8	9	10	16	10	23	20	31	-
1985	345	20	10	13	20	18	19	22	20	21	-
1984	315	11	11	7	15	22	22	18	18	23	-
1983	334	14	15	17	17	18	19	20	20	16	-
1982	296	12	8	9	13	16	12	22	16	17	-
1981	308	13	11	7	16	18	16	23	23	20	-
1980	321	10	6	15	15	19	25	11	19	14	-
1979	298	8	13	11	12	16	20	15	13	17	-
Anemias (280-285)											
Black, female											
1998	524	29	24	13	17	27	20	40	39	112	-
1997	504	38	14	18	16	30	35	23	41	104	-
1996	510	32	21	19	20	18	23	30	41	109	-
1995	564	22	24	17	29	18	29	44	49	115	-
1994	530	34	22	12	20	26	22	36	43	88	-
1993	523	26	18	21	18	21	25	39	42	105	-
1992	497	22	19	16	16	16	34	49	39	78	-
1991	463	30	14	11	14	36	29	34	31	90	-
1990	427	13	19	11	17	31	20	30	31	71	-
1989	451	20	23	12	25	29	28	35	30	71	-
1988	394	19	9	23	16	20	22	35	35	67	-
1987	361	13	12	14	15	23	25	28	18	55	-
1986	372	9	12	14	16	17	21	34	34	52	-
1985	372	14	14	11	20	18	31	34	31	52	-
1984	325	19	6	13	13	17	23	28	22	49	-
1983	331	9	12	22	15	21	19	18	23	40	1
1982	336	11	11	10	14	26	21	24	35	42	-
1981	307	13	16	14	11	25	21	20	17	28	-
1980	339	13	20	13	21	16	28	27	23	36	-
1979	339	9	13	19	17	19	25	30	30	34	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
All races, both sexes											
1998	64,751	1,784	2,602	3,687	5,018	7,042	9,448	10,644	9,885	11,955	-
1997	62,636	1,767	2,568	3,512	4,858	6,955	9,356	10,223	9,369	11,386	2
1996	61,767	1,777	2,481	3,418	5,011	7,274	9,279	9,962	9,027	10,834	1
1995	59,254	1,633	2,363	3,355	4,833	7,097	9,134	9,426	8,709	10,086	1
1994	56,692	1,495	2,194	3,060	4,724	6,881	8,863	8,845	8,334	9,677	2
1993	53,894	1,407	2,042	3,030	4,448	6,631	8,353	8,509	7,813	9,196	9
1992	50,067	1,374	1,829	2,802	4,307	6,253	7,723	8,029	7,062	8,261	2
1991	48,951	1,187	1,847	2,761	4,250	6,440	7,394	7,736	6,939	8,019	2
1990	47,664	1,187	1,647	2,553	4,416	6,306	6,974	7,679	6,858	7,706	4
1989	46,833	1,178	1,606	2,624	4,318	6,045	7,123	7,535	6,625	7,470	1
1988	40,368	993	1,509	2,305	3,804	5,170	5,922	6,370	5,537	6,548	-
1987	38,532	901	1,357	2,263	3,651	4,952	5,837	6,102	5,368	6,118	2
1986	37,184	842	1,316	2,234	3,546	4,662	5,607	5,858	5,190	5,939	6
1985	36,969	773	1,218	2,182	3,637	4,597	5,562	6,058	5,250	5,817	8
1984	35,787	746	1,259	2,045	3,450	4,423	5,511	5,801	5,056	5,690	1
1983	36,246	784	1,294	2,378	3,522	4,574	5,611	5,838	5,085	5,328	4
1982	34,583	748	1,312	2,264	3,379	4,419	5,292	5,428	4,911	5,170	8
1981	34,642	763	1,404	2,273	3,347	4,532	5,309	5,508	4,713	5,120	8
1980	34,851	773	1,415	2,364	3,425	4,655	5,456	5,484	4,650	4,971	5
1979	33,192	754	1,303	2,297	3,240	4,349	5,049	5,236	4,666	4,648	6
Diabetes mellitus (250)											
All races, male											
1998	29,584	1,062	1,426	2,007	2,644	3,588	4,572	4,842	4,095	3,765	-
1997	28,187	990	1,452	1,795	2,418	3,486	4,544	4,634	3,786	3,516	2
1996	27,646	1,059	1,376	1,803	2,564	3,519	4,373	4,447	3,652	3,253	-
1995	26,124	954	1,338	1,776	2,394	3,437	4,127	4,222	3,416	2,942	-
1994	24,758	848	1,208	1,576	2,366	3,288	4,040	3,825	3,169	2,876	1
1993	23,430	792	1,125	1,597	2,187	3,159	3,854	3,627	2,990	2,671	3
1992	21,672	813	993	1,431	2,126	2,908	3,468	3,396	2,651	2,442	1
1991	21,096	687	976	1,400	2,066	3,011	3,373	3,306	2,646	2,257	2
1990	20,266	646	901	1,316	2,120	2,927	3,116	3,295	2,456	2,132	2
1989	19,730	681	875	1,349	2,033	2,746	3,175	3,077	2,360	2,041	-
1988	16,975	591	764	1,130	1,810	2,377	2,535	2,571	1,991	1,891	-
1987	16,237	525	706	1,187	1,735	2,344	2,568	2,429	1,898	1,668	1
1986	15,369	491	641	1,136	1,646	2,127	2,437	2,311	1,815	1,640	3
1985	15,263	436	640	1,079	1,688	2,069	2,430	2,393	1,846	1,590	2
1984	14,859	441	687	1,017	1,594	2,028	2,308	2,335	1,808	1,613	-
1983	14,668	430	652	1,172	1,613	2,041	2,337	2,261	1,681	1,474	2
1982	14,171	411	675	1,115	1,567	2,012	2,250	2,161	1,669	1,387	-
1981	14,211	436	734	1,120	1,588	1,952	2,249	2,133	1,584	1,487	3
1980	14,325	386	708	1,162	1,589	2,037	2,279	2,133	1,631	1,481	1
1979	13,795	400	667	1,147	1,552	1,997	2,136	2,013	1,614	1,359	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
All races, female											
1998	35,167	722	1,176	1,680	2,374	3,454	4,876	5,802	5,790	8,190	-
1997	34,449	777	1,116	1,717	2,440	3,469	4,812	5,589	5,583	7,870	-
1996	34,121	718	1,105	1,615	2,447	3,755	4,906	5,515	5,375	7,581	1
1995	33,130	679	1,025	1,579	2,439	3,660	5,007	5,204	5,293	7,144	1
1994	31,934	647	986	1,484	2,358	3,593	4,823	5,020	5,165	6,801	1
1993	30,464	615	917	1,433	2,261	3,472	4,499	4,882	4,823	6,525	6
1992	28,395	561	836	1,371	2,181	3,345	4,255	4,633	4,411	5,819	1
1991	27,855	500	871	1,361	2,184	3,429	4,021	4,430	4,293	5,762	-
1990	27,398	541	746	1,237	2,296	3,379	3,858	4,384	4,402	5,574	2
1989	27,103	497	731	1,275	2,285	3,299	3,948	4,458	4,265	5,429	1
1988	23,393	402	745	1,175	1,994	2,793	3,387	3,799	3,546	4,657	-
1987	22,295	376	651	1,076	1,916	2,608	3,269	3,673	3,470	4,450	1
1986	21,815	351	675	1,098	1,900	2,535	3,170	3,547	3,375	4,299	3
1985	21,706	337	578	1,103	1,949	2,528	3,132	3,665	3,404	4,227	6
1984	20,928	305	572	1,028	1,856	2,395	3,203	3,466	3,248	4,077	1
1983	21,578	354	642	1,206	1,909	2,533	3,274	3,577	3,404	3,854	2
1982	20,412	337	637	1,149	1,812	2,407	3,042	3,267	3,242	3,783	8
1981	20,431	327	670	1,153	1,759	2,580	3,060	3,375	3,129	3,633	5
1980	20,526	387	707	1,202	1,836	2,618	3,177	3,351	3,019	3,490	4
1979	19,397	354	636	1,150	1,688	2,352	2,913	3,223	3,052	3,289	4
Diabetes mellitus (250)											
White, both sexes											
1998	51,706	1,251	1,853	2,599	3,735	5,421	7,464	8,816	8,396	10,293	-
1997	49,850	1,247	1,820	2,512	3,512	5,331	7,457	8,443	7,960	9,719	2
1996	49,511	1,253	1,766	2,488	3,703	5,669	7,368	8,288	7,658	9,378	-
1995	47,475	1,161	1,685	2,406	3,536	5,528	7,315	7,869	7,374	8,722	-
1994	45,650	1,085	1,560	2,199	3,554	5,393	7,135	7,378	7,048	8,401	1
1993	43,461	1,004	1,467	2,164	3,343	5,161	6,747	7,188	6,608	8,008	3
1992	40,442	1,005	1,287	2,037	3,177	4,870	6,331	6,746	6,007	7,192	1
1991	39,600	865	1,272	2,022	3,184	5,030	6,076	6,460	5,954	6,992	1
1990	38,696	840	1,144	1,865	3,332	4,979	5,674	6,474	5,882	6,770	1
1989	38,053	845	1,149	1,904	3,245	4,737	5,871	6,268	5,708	6,582	-
1988	32,692	718	1,061	1,698	2,874	4,057	4,816	5,275	4,783	5,774	-
1987	31,395	626	976	1,684	2,753	3,899	4,792	5,128	4,566	5,439	-
1986	30,284	598	916	1,632	2,693	3,729	4,577	4,954	4,448	5,218	3
1985	30,305	521	868	1,623	2,806	3,633	4,584	5,063	4,551	5,223	4
1984	29,466	545	896	1,528	2,632	3,496	4,503	4,948	4,418	5,145	-
1983	29,987	558	939	1,746	2,723	3,727	4,659	4,990	4,422	4,827	1
1982	28,910	542	963	1,724	2,630	3,595	4,466	4,700	4,312	4,692	5
1981	28,750	553	1,032	1,723	2,624	3,615	4,411	4,721	4,175	4,635	4
1980	28,868	552	1,042	1,772	2,676	3,699	4,566	4,697	4,124	4,498	1
1979	27,624	527	917	1,738	2,571	3,484	4,202	4,505	4,246	4,199	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
White, male											
1998	24,249	765	1,042	1,480	2,059	2,892	3,778	4,175	3,627	3,309	-
1997	22,968	703	1,061	1,346	1,832	2,784	3,772	3,980	3,306	3,079	2
1996	22,829	774	1,029	1,393	2,001	2,909	3,632	3,827	3,218	2,877	-
1995	21,407	703	972	1,336	1,839	2,782	3,458	3,645	3,000	2,569	-
1994	20,391	649	886	1,181	1,856	2,690	3,365	3,298	2,747	2,562	1
1993	19,311	583	848	1,183	1,729	2,573	3,232	3,197	2,593	2,337	3
1992	17,951	604	732	1,100	1,644	2,392	2,942	2,947	2,320	2,172	1
1991	17,473	512	715	1,079	1,614	2,452	2,896	2,868	2,305	2,015	1
1990	16,817	463	662	1,000	1,674	2,418	2,640	2,872	2,158	1,903	1
1989	16,282	500	650	1,010	1,590	2,231	2,687	2,662	2,081	1,816	-
1988	14,008	445	557	871	1,422	1,956	2,135	2,195	1,746	1,692	-
1987	13,553	363	527	928	1,379	1,925	2,205	2,121	1,668	1,503	-
1986	12,788	372	471	889	1,312	1,774	2,061	2,000	1,580	1,466	1
1985	12,758	308	485	857	1,359	1,715	2,087	2,045	1,628	1,432	-
1984	12,450	332	505	808	1,259	1,655	1,983	2,032	1,618	1,470	-
1983	12,337	307	505	904	1,298	1,745	2,004	1,964	1,485	1,353	-
1982	12,092	307	534	900	1,303	1,717	1,969	1,897	1,483	1,261	-
1981	12,066	330	568	904	1,334	1,635	1,937	1,858	1,431	1,362	2
1980	12,125	290	563	902	1,315	1,709	1,973	1,848	1,466	1,361	-
1979	11,688	289	507	909	1,303	1,676	1,845	1,750	1,483	1,231	1
Diabetes mellitus (250)											
White, female											
1998	27,457	486	811	1,119	1,676	2,529	3,686	4,641	4,769	6,984	-
1997	26,882	544	759	1,166	1,680	2,547	3,685	4,463	4,654	6,640	-
1996	26,682	479	737	1,095	1,702	2,760	3,736	4,461	4,440	6,501	-
1995	26,068	458	713	1,070	1,697	2,746	3,857	4,224	4,374	6,153	-
1994	25,259	436	674	1,018	1,698	2,703	3,770	4,080	4,301	5,839	-
1993	24,150	421	619	981	1,614	2,588	3,515	3,991	4,015	5,671	-
1992	22,491	401	555	937	1,533	2,478	3,389	3,799	3,687	5,020	-
1991	22,127	353	557	943	1,570	2,578	3,180	3,592	3,649	4,977	-
1990	21,879	377	482	865	1,658	2,561	3,034	3,602	3,724	4,867	-
1989	21,771	345	499	894	1,655	2,506	3,184	3,606	3,627	4,766	-
1988	18,684	273	504	827	1,452	2,101	2,681	3,080	3,037	4,082	-
1987	17,842	263	449	756	1,374	1,974	2,587	3,007	2,898	3,936	-
1986	17,496	226	445	743	1,381	1,955	2,516	2,954	2,868	3,752	2
1985	17,547	213	383	766	1,447	1,918	2,497	3,018	2,923	3,791	4
1984	17,016	213	391	720	1,373	1,841	2,520	2,916	2,800	3,675	-
1983	17,650	251	434	842	1,425	1,982	2,655	3,026	2,937	3,474	1
1982	16,818	235	429	824	1,327	1,878	2,497	2,803	2,829	3,431	5
1981	16,684	223	464	819	1,290	1,980	2,474	2,863	2,744	3,273	2
1980	16,743	262	479	870	1,361	1,990	2,593	2,849	2,658	3,137	1
1979	15,936	238	410	829	1,268	1,808	2,357	2,755	2,763	2,968	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
All other, both sexes											
1998	13,045	533	749	1,088	1,283	1,621	1,984	1,828	1,489	1,662	-
1997	12,786	520	748	1,000	1,346	1,624	1,899	1,780	1,409	1,667	-
1996	12,256	524	715	930	1,308	1,605	1,911	1,674	1,369	1,456	1
1995	11,779	472	678	949	1,297	1,569	1,819	1,557	1,335	1,364	1
1994	11,042	410	634	861	1,170	1,488	1,728	1,467	1,286	1,276	1
1993	10,433	403	575	866	1,105	1,470	1,606	1,321	1,205	1,188	6
1992	9,625	369	542	765	1,130	1,383	1,392	1,283	1,055	1,069	1
1991	9,351	322	575	739	1,066	1,410	1,318	1,276	985	1,027	1
1990	8,968	347	503	688	1,084	1,327	1,300	1,205	976	936	3
1989	8,780	333	457	720	1,073	1,308	1,252	1,267	917	888	1
1988	7,676	275	448	607	930	1,113	1,106	1,095	754	774	-
1987	7,137	275	381	579	898	1,053	1,045	974	802	679	2
1986	6,900	244	400	602	853	933	1,030	904	742	721	3
1985	6,664	252	350	559	831	964	978	995	699	594	4
1984	6,321	201	363	517	818	927	1,008	853	638	545	1
1983	6,259	226	355	632	799	847	952	848	663	501	3
1982	5,673	206	349	540	749	824	826	728	599	478	3
1981	5,892	210	372	550	723	917	898	787	538	485	4
1980	5,983	221	373	592	749	956	890	787	526	473	4
1979	5,568	227	386	559	669	865	847	731	420	449	1
Diabetes mellitus (250)											
All other, male											
1998	5,335	297	384	527	585	696	794	667	468	456	-
1997	5,219	287	391	449	586	702	772	654	480	437	-
1996	4,817	285	347	410	563	610	741	620	434	376	-
1995	4,717	251	366	440	555	655	669	577	416	373	-
1994	4,367	199	322	395	510	598	675	527	422	314	-
1993	4,119	209	277	414	458	586	622	430	397	334	-
1992	3,721	209	261	331	482	516	526	449	331	270	-
1991	3,623	175	261	321	452	559	477	438	341	242	1
1990	3,449	183	239	316	446	509	476	423	298	229	1
1989	3,448	181	225	339	443	515	488	415	279	225	-
1988	2,967	146	207	259	388	421	400	376	245	199	-
1987	2,684	162	179	259	356	419	363	308	230	165	1
1986	2,581	119	170	247	334	353	376	311	235	174	2
1985	2,505	128	155	222	329	354	343	348	218	158	2
1984	2,409	109	182	209	335	373	325	303	190	143	-
1983	2,331	123	147	268	315	296	333	297	196	121	2
1982	2,079	104	141	215	264	295	281	264	186	126	-
1981	2,145	106	166	216	254	317	312	275	153	125	1
1980	2,200	96	145	260	274	328	306	285	165	120	1
1979	2,107	111	160	238	249	321	291	263	131	128	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
All other, female											
1998	7,710	236	365	561	698	925	1,190	1,161	1,021	1,206	-
1997	7,567	233	357	551	760	922	1,127	1,126	929	1,230	-
1996	7,439	239	368	520	745	995	1,170	1,054	935	1,080	1
1995	7,062	221	312	509	742	914	1,150	980	919	991	1
1994	6,675	211	312	466	660	890	1,053	940	864	962	1
1993	6,314	194	298	452	647	884	984	891	808	854	6
1992	5,904	160	281	434	648	867	866	834	724	799	1
1991	5,728	147	314	418	614	851	841	838	644	785	-
1990	5,519	164	264	372	638	818	824	782	678	707	2
1989	5,332	152	232	381	630	793	764	852	638	663	1
1988	4,709	129	241	348	542	692	706	719	509	575	-
1987	4,453	113	202	320	542	634	682	666	572	514	1
1986	4,319	125	230	355	519	580	654	593	507	547	1
1985	4,159	124	195	337	502	610	635	647	481	436	2
1984	3,912	92	181	308	483	554	683	550	448	402	1
1983	3,928	103	208	364	484	551	619	551	467	380	1
1982	3,594	102	208	325	485	529	545	464	413	352	3
1981	3,747	104	206	334	469	600	586	512	385	360	3
1980	3,783	125	228	332	475	628	584	502	361	353	3
1979	3,461	116	226	321	420	544	556	468	289	321	-
Diabetes mellitus (250)											
Black, both sexes											
1998	11,378	474	631	953	1,109	1,409	1,728	1,573	1,282	1,466	-
1997	11,130	471	661	875	1,171	1,398	1,627	1,513	1,224	1,454	-
1996	10,800	471	633	806	1,163	1,400	1,665	1,466	1,219	1,299	-
1995	10,402	435	599	829	1,137	1,381	1,601	1,348	1,167	1,224	1
1994	9,849	367	562	768	1,037	1,320	1,532	1,310	1,118	1,153	1
1993	9,380	366	523	772	976	1,319	1,458	1,175	1,077	1,054	5
1992	8,653	345	492	664	1,008	1,237	1,248	1,153	935	967	1
1991	8,517	294	525	681	960	1,269	1,206	1,150	874	959	1
1990	8,114	314	457	632	964	1,196	1,177	1,091	882	839	3
1989	7,955	307	406	651	979	1,196	1,127	1,146	797	807	1
1988	6,972	240	411	559	839	1,005	1,007	994	679	693	-
1987	6,497	258	350	511	826	960	943	883	737	608	2
1986	6,299	228	366	548	778	844	952	818	663	659	3
1985	6,104	233	320	499	754	878	900	922	647	533	4
1984	5,810	190	330	471	749	850	930	782	595	481	1
1983	5,790	216	334	582	743	786	871	765	624	458	2
1982	5,227	184	324	496	700	770	760	663	542	435	3
1981	5,461	193	343	506	671	856	829	735	490	448	4
1980	5,544	201	343	546	682	889	831	733	488	441	4
1979	5,174	212	358	523	633	808	794	671	379	405	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diabetes mellitus (250)											
Black, male											
1998	4,511	256	318	446	487	587	679	550	395	372	-
1997	4,440	263	337	379	501	599	646	539	401	347	-
1996	4,132	260	302	345	479	522	624	533	378	312	-
1995	4,110	229	316	379	487	576	579	493	353	314	-
1994	3,834	173	288	349	457	523	590	456	350	267	-
1993	3,648	188	249	367	402	515	560	376	332	284	-
1992	3,271	196	230	284	424	449	461	393	280	226	-
1991	3,267	164	239	291	404	503	435	389	286	221	1
1990	3,049	161	212	285	395	457	414	368	255	198	1
1989	3,072	165	200	301	406	472	433	360	225	190	-
1988	2,640	124	191	235	350	377	350	334	206	163	-
1987	2,388	150	160	228	322	375	318	266	203	141	1
1986	2,295	111	153	225	301	304	334	268	199	150	2
1985	2,230	118	141	193	292	314	301	310	194	132	2
1984	2,160	102	162	187	298	332	289	269	174	117	-
1983	2,123	118	139	247	293	271	298	252	180	107	2
1982	1,878	93	130	197	238	273	252	231	159	113	-
1981	1,925	96	148	193	231	296	264	251	129	110	1
1980	2,010	85	131	239	250	301	276	262	153	111	1
1979	1,915	104	143	220	232	295	268	231	116	107	1
Diabetes mellitus (250)											
Black, female											
1998	6,867	218	313	507	622	822	1,049	1,023	887	1,094	-
1997	6,690	208	324	496	670	799	981	974	823	1,107	-
1996	6,668	211	331	461	684	878	1,041	933	841	987	-
1995	6,292	206	283	450	650	805	1,022	855	814	910	1
1994	6,015	194	274	419	580	797	942	854	768	886	1
1993	5,732	178	274	405	574	804	898	799	745	770	5
1992	5,382	149	262	380	584	788	787	760	655	741	1
1991	5,250	130	286	390	556	766	771	761	588	738	-
1990	5,065	153	245	347	569	739	763	723	627	641	2
1989	4,883	142	206	350	573	724	694	786	572	617	1
1988	4,332	116	220	324	489	628	657	660	473	530	-
1987	4,109	108	190	283	504	585	625	617	534	467	1
1986	4,004	117	213	323	477	540	618	550	464	509	1
1985	3,874	115	179	306	462	564	599	612	453	401	2
1984	3,650	88	168	284	451	518	641	513	421	364	1
1983	3,667	98	195	335	450	515	573	513	444	351	-
1982	3,349	91	194	299	462	497	508	432	383	322	3
1981	3,536	97	195	313	440	560	565	484	361	338	3
1980	3,534	116	212	307	432	588	555	471	335	330	3
1979	3,259	108	215	303	401	513	526	440	263	298	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
All races, both sexes											
1998	4,077	23	33	49	70	136	208	452	714	2,311	-
1997	3,928	32	38	52	75	140	227	407	684	2,192	-
1996	3,706	28	39	51	60	118	242	424	600	2,078	1
1995	3,584	32	32	52	71	124	239	373	629	1,961	-
1994	3,451	26	27	44	71	125	223	382	592	1,878	1
1993	3,451	27	30	41	59	145	226	385	627	1,818	1
1992	3,135	28	32	39	64	119	214	347	553	1,644	-
1991	2,991	26	19	43	61	107	210	371	552	1,524	1
1990	3,017	22	23	49	76	137	194	377	531	1,532	1
1989	3,046	24	18	54	65	135	205	368	566	1,526	2
1988	2,985	17	19	38	82	135	220	355	552	1,468	-
1987	2,889	10	29	41	67	121	225	359	535	1,408	-
1986	2,862	18	23	50	86	137	260	345	481	1,377	1
1985	2,874	20	32	59	77	146	223	336	543	1,352	2
1984	2,600	12	27	47	78	132	234	323	455	1,222	-
1983	2,656	18	27	42	93	130	240	328	492	1,198	2
1982	2,430	18	26	56	77	131	231	347	442	1,015	3
1981	2,331	18	30	56	91	150	216	286	395	991	2
1980	2,364	28	29	58	95	148	224	311	417	960	-
1979	2,210	26	45	63	105	124	214	314	384	826	2
Nutritional deficiencies (260-269)											
All races, male											
1998	1,278	10	17	24	41	70	95	192	240	544	-
1997	1,213	19	16	27	38	65	109	172	219	519	-
1996	1,225	16	23	27	34	66	108	177	205	538	1
1995	1,193	18	14	30	39	54	104	140	236	510	-
1994	1,186	15	14	26	42	64	110	160	220	485	1
1993	1,257	10	17	23	28	98	116	168	248	501	-
1992	1,089	11	21	22	33	69	98	146	211	426	-
1991	1,085	16	12	22	31	50	104	160	222	426	1
1990	1,108	14	14	21	34	70	93	166	210	445	1
1989	1,098	14	9	31	34	67	94	151	214	440	2
1988	1,096	9	15	22	41	74	109	139	226	405	-
1987	1,107	4	15	28	38	73	98	171	215	413	-
1986	1,088	9	11	28	44	76	133	159	194	394	-
1985	1,086	13	12	33	46	72	114	146	212	395	-
1984	977	5	16	27	41	78	120	136	163	352	-
1983	1,065	11	11	21	56	72	136	154	181	374	1
1982	994	7	16	30	43	70	123	162	183	318	2
1981	941	12	20	27	46	83	99	135	161	308	2
1980	962	13	16	27	44	89	108	145	156	312	-
1979	927	11	28	23	54	73	87	145	163	275	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
All races, female											
1998	2,799	13	16	25	29	66	113	260	474	1,767	-
1997	2,715	13	22	25	37	75	118	235	465	1,673	-
1996	2,481	12	16	24	26	52	134	247	395	1,540	-
1995	2,391	14	18	22	32	70	135	233	393	1,451	-
1994	2,265	11	13	18	29	61	113	222	372	1,393	-
1993	2,194	17	13	18	31	47	110	217	379	1,317	1
1992	2,046	17	11	17	31	50	116	201	342	1,218	-
1991	1,906	10	7	21	30	57	106	211	330	1,098	-
1990	1,909	8	9	28	42	67	101	211	321	1,087	-
1989	1,948	10	9	23	31	68	111	217	352	1,086	-
1988	1,889	8	4	16	41	61	111	216	326	1,063	-
1987	1,782	6	14	13	29	48	127	188	320	995	-
1986	1,774	9	12	22	42	61	127	186	287	983	1
1985	1,788	7	20	26	31	74	109	190	331	957	2
1984	1,623	7	11	20	37	54	114	187	292	870	-
1983	1,591	7	16	21	37	58	104	174	311	824	1
1982	1,436	11	10	26	34	61	108	185	259	697	1
1981	1,390	6	10	29	45	67	117	151	234	683	-
1980	1,402	15	13	31	51	59	116	166	261	648	-
1979	1,283	15	17	40	51	51	127	169	221	551	1
Nutritional deficiencies (260-269)											
White, both sexes											
1998	3,539	18	30	41	57	106	171	396	611	2,049	-
1997	3,447	21	27	41	56	107	185	358	608	1,990	-
1996	3,255	21	28	35	50	86	204	373	528	1,886	1
1995	3,118	25	21	39	54	96	203	305	545	1,778	-
1994	3,007	19	18	33	57	102	196	328	516	1,695	1
1993	2,986	20	22	31	40	109	183	326	545	1,660	-
1992	2,642	18	22	25	47	91	168	278	467	1,466	-
1991	2,558	23	15	33	48	79	161	321	486	1,330	1
1990	2,546	9	15	30	55	106	146	318	444	1,369	1
1989	2,571	20	15	37	47	104	163	300	484	1,345	1
1988	2,526	12	12	21	59	105	172	306	481	1,299	-
1987	2,478	9	16	30	49	96	184	307	466	1,271	-
1986	2,438	10	17	34	73	106	216	274	411	1,242	1
1985	2,490	11	26	45	60	113	178	288	470	1,243	2
1984	2,231	11	22	35	52	107	184	254	398	1,117	-
1983	2,252	12	22	31	69	105	183	274	425	1,081	1
1982	2,049	12	19	42	60	103	182	287	382	905	1
1981	1,989	13	17	46	74	124	172	246	348	881	1
1980	2,008	20	21	41	74	118	190	258	366	869	-
1979	1,862	17	33	46	90	87	172	269	333	748	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
White, male											
1998	1,067	6	14	19	34	53	76	162	196	473	-
1997	1,003	12	13	19	25	46	87	146	188	451	-
1996	1,035	12	18	15	28	43	90	154	178	472	1
1995	999	15	7	19	28	40	91	117	196	452	-
1994	995	10	9	21	32	50	101	133	189	426	1
1993	1,046	5	11	16	17	77	92	135	218	450	-
1992	872	7	13	10	24	52	74	118	172	369	-
1991	868	14	10	16	21	33	80	137	185	341	1
1990	903	4	9	14	23	58	72	139	173	382	1
1989	895	10	9	20	23	46	77	121	181	378	1
1988	884	5	9	11	28	55	84	117	198	344	-
1987	901	4	6	18	28	56	76	140	187	359	-
1986	883	3	6	20	34	62	107	122	161	342	-
1985	916	7	8	24	35	59	86	123	181	363	-
1984	799	4	13	19	26	68	96	101	138	307	-
1983	853	7	7	14	40	57	101	119	152	331	-
1982	818	5	11	25	35	58	98	125	156	279	-
1981	786	9	11	22	36	68	78	110	145	274	1
1980	779	6	10	17	29	68	88	123	132	276	-
1979	750	7	18	17	46	50	69	123	135	246	1
Nutritional deficiencies (260-269)											
White, female											
1998	2,472	12	16	22	23	53	95	234	415	1,576	-
1997	2,444	9	14	22	31	61	98	212	420	1,539	-
1996	2,220	9	10	20	22	43	114	219	350	1,414	-
1995	2,119	10	14	20	26	56	112	188	349	1,326	-
1994	2,012	9	9	12	25	52	95	195	327	1,269	-
1993	1,940	15	11	15	23	32	91	191	327	1,210	-
1992	1,770	11	9	15	23	39	94	160	295	1,097	-
1991	1,690	9	5	17	27	46	81	184	301	989	-
1990	1,643	5	6	16	32	48	74	179	271	987	-
1989	1,676	10	6	17	24	58	86	179	303	967	-
1988	1,642	7	3	10	31	50	88	189	283	955	-
1987	1,577	5	10	12	21	40	108	167	279	912	-
1986	1,555	7	11	14	39	44	109	152	250	900	1
1985	1,574	4	18	21	25	54	92	165	289	880	2
1984	1,432	7	9	16	26	39	88	153	260	810	-
1983	1,399	5	15	17	29	48	82	155	273	750	1
1982	1,231	7	8	17	25	45	84	162	226	626	1
1981	1,203	4	6	24	38	56	94	136	203	607	-
1980	1,229	14	11	24	45	50	102	135	234	593	-
1979	1,112	10	15	29	44	37	103	146	198	502	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
All other, both sexes											
1998	538	5	3	8	13	30	37	56	103	262	-
1997	481	11	11	11	19	33	42	49	76	202	-
1996	451	7	11	16	10	32	38	51	72	192	-
1995	466	7	11	13	17	28	36	68	84	183	-
1994	444	7	9	11	14	23	27	54	76	183	-
1993	465	7	8	10	19	36	43	59	82	158	1
1992	493	10	10	14	17	28	46	69	86	178	-
1991	433	3	4	10	13	28	49	50	66	194	-
1990	471	13	8	19	21	31	48	59	87	163	-
1989	475	4	3	17	18	31	42	68	82	181	1
1988	459	5	7	17	23	30	48	49	71	169	-
1987	411	1	13	11	18	25	41	52	69	137	-
1986	424	8	6	16	13	31	44	71	70	135	-
1985	384	9	6	14	17	33	45	48	73	109	-
1984	369	1	5	12	26	25	50	69	57	105	-
1983	404	6	5	11	24	25	57	54	67	117	1
1982	381	6	7	14	17	28	49	60	60	110	2
1981	342	5	13	10	17	26	44	40	47	110	1
1980	356	8	8	17	21	30	34	53	51	91	-
1979	348	9	12	17	15	37	42	45	51	78	-
Nutritional deficiencies (260-269)											
All other, male											
1998	211	4	3	5	7	17	19	30	44	71	-
1997	210	7	3	8	13	19	22	26	31	68	-
1996	190	4	5	12	6	23	18	23	27	66	-
1995	194	3	7	11	11	14	13	23	40	58	-
1994	191	5	5	5	10	14	9	27	31	59	-
1993	211	5	6	7	11	21	24	33	30	51	-
1992	217	4	8	12	9	17	24	28	39	57	-
1991	217	2	2	6	10	17	24	23	37	85	-
1990	205	10	5	7	11	12	21	27	37	63	-
1989	203	4	-	11	11	21	17	30	33	62	1
1988	212	4	6	11	13	19	25	22	28	61	-
1987	206	-	9	10	10	17	22	31	28	54	-
1986	205	6	5	8	10	14	26	37	33	52	-
1985	170	6	4	9	11	13	28	23	31	32	-
1984	178	1	3	8	15	10	24	35	25	45	-
1983	212	4	4	7	16	15	35	35	29	43	1
1982	176	2	5	5	8	12	25	37	27	39	2
1981	155	3	9	5	10	15	21	25	16	34	1
1980	183	7	6	10	15	21	20	22	24	36	-
1979	177	4	10	6	8	23	18	22	28	29	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
All other, female											
1998	327	1	-	3	6	13	18	26	59	191	-
1997	271	4	8	3	6	14	20	23	45	134	-
1996	261	3	6	4	4	9	20	28	45	126	-
1995	272	4	4	2	6	14	23	45	44	125	-
1994	253	2	4	6	4	9	18	27	45	124	-
1993	254	2	2	3	8	15	19	26	52	107	1
1992	276	6	2	2	8	11	22	41	47	121	-
1991	216	1	2	4	3	11	25	27	29	109	-
1990	266	3	3	12	10	19	27	32	50	100	-
1989	272	-	3	6	7	10	25	38	49	119	-
1988	247	1	1	6	10	11	23	27	43	108	-
1987	205	1	4	1	8	8	19	21	41	83	-
1986	219	2	1	8	3	17	18	34	37	83	-
1985	214	3	2	5	6	20	17	25	42	77	-
1984	191	-	2	4	11	15	26	34	32	60	-
1983	192	2	1	4	8	10	22	19	38	74	-
1982	205	4	2	9	9	16	24	23	33	71	-
1981	187	2	4	5	7	11	23	15	31	76	-
1980	173	1	2	7	6	9	14	31	27	55	-
1979	171	5	2	11	7	14	24	23	23	49	-
Nutritional deficiencies (260-269)											
Black, both sexes											
1998	480	5	3	8	12	29	34	51	94	223	-
1997	429	10	7	10	19	31	37	47	65	178	-
1996	396	6	8	14	8	30	33	51	63	162	-
1995	418	7	11	13	16	27	34	61	75	155	-
1994	411	7	8	9	14	21	27	52	72	163	-
1993	424	6	7	9	18	34	42	53	76	138	1
1992	445	10	9	14	17	26	40	64	73	157	-
1991	393	3	4	10	12	27	42	46	61	172	-
1990	433	13	8	18	20	29	45	56	78	147	-
1989	443	4	3	15	17	29	41	64	75	167	1
1988	420	5	6	17	23	28	44	44	63	155	-
1987	383	1	12	9	15	23	38	47	68	126	-
1986	392	8	6	16	13	30	40	66	65	118	-
1985	349	8	4	13	17	32	40	43	63	99	-
1984	338	1	5	12	25	24	46	62	55	90	-
1983	372	6	5	11	24	25	51	51	65	98	1
1982	346	5	7	13	17	27	48	58	51	93	2
1981	310	5	13	10	17	23	42	36	43	92	1
1980	333	7	8	17	20	29	32	49	43	86	-
1979	325	9	12	15	14	34	41	42	46	74	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nutritional deficiencies (260-269)											
Black, male											
1998	188	4	3	5	7	17	18	27	38	58	-
1997	185	6	3	7	13	18	19	25	28	55	-
1996	167	3	4	12	4	22	16	23	24	53	-
1995	176	3	7	11	10	14	12	20	35	50	-
1994	177	5	5	5	10	12	9	25	30	51	-
1993	193	4	6	7	10	20	23	28	29	44	-
1992	195	4	7	12	9	15	20	27	32	50	-
1991	193	2	2	6	9	16	19	20	33	75	-
1990	189	10	5	7	11	12	21	25	33	54	-
1989	188	4	-	10	11	19	16	29	30	55	1
1988	195	4	5	11	13	17	24	21	25	56	-
1987	194	-	9	9	8	15	20	31	27	50	-
1986	190	6	5	8	10	13	23	35	32	44	-
1985	149	5	3	8	11	13	25	21	25	25	-
1984	161	1	3	8	14	10	22	30	23	38	-
1983	199	4	4	7	16	15	32	32	28	38	1
1982	159	2	5	5	8	12	24	36	20	32	2
1981	137	3	9	5	10	13	19	22	14	26	1
1980	172	6	6	10	15	20	19	21	20	33	-
1979	165	4	10	6	8	21	17	21	25	28	-
Nutritional deficiencies (260-269)											
Black, female											
1998	292	1	-	3	5	12	16	24	56	165	-
1997	244	4	4	3	6	13	18	22	37	123	-
1996	229	3	4	2	4	8	17	28	39	109	-
1995	242	4	4	2	6	13	22	41	40	105	-
1994	234	2	3	4	4	9	18	27	42	112	-
1993	231	2	1	2	8	14	19	25	47	94	1
1992	250	6	2	2	8	11	20	37	41	107	-
1991	200	1	2	4	3	11	23	26	28	97	-
1990	244	3	3	11	9	17	24	31	45	93	-
1989	255	-	3	5	6	10	25	35	45	112	-
1988	225	1	1	6	10	11	20	23	38	99	-
1987	189	1	3	-	7	8	18	16	41	76	-
1986	202	2	1	8	3	17	17	31	33	74	-
1985	200	3	1	5	6	19	15	22	38	74	-
1984	177	-	2	4	11	14	24	32	32	52	-
1983	173	2	1	4	8	10	19	19	37	60	-
1982	187	3	2	8	9	15	24	22	31	61	-
1981	173	2	4	5	7	10	23	14	29	66	-
1980	161	1	2	7	5	9	13	28	23	53	-
1979	160	5	2	9	6	13	24	21	21	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
All races, both sexes											
1998	3,881	21	30	47	64	129	199	418	682	2,214	-
1997	3,744	32	35	47	69	132	214	385	655	2,101	-
1996	3,531	26	39	44	56	115	228	401	569	2,000	1
1995	3,447	29	29	48	66	119	229	360	614	1,888	-
1994	3,274	25	24	41	65	119	213	363	559	1,792	1
1993	3,275	23	28	38	53	137	212	363	593	1,740	-
1992	2,998	27	27	35	61	114	209	330	538	1,573	-
1991	2,852	22	19	40	56	101	202	354	524	1,467	1
1990	2,868	22	21	48	72	127	176	361	509	1,464	1
1989	2,906	23	17	52	55	129	197	353	546	1,458	2
1988	2,842	16	15	35	80	125	211	329	532	1,406	-
1987	2,741	10	24	41	60	115	207	344	515	1,345	-
1986	2,692	17	18	43	79	127	239	330	455	1,308	1
1985	2,720	15	29	57	69	135	211	321	521	1,296	2
1984	2,465	10	24	46	72	122	224	302	443	1,163	-
1983	2,515	16	23	36	85	123	224	315	475	1,145	2
1982	2,314	17	24	51	71	126	218	335	421	975	3
1981	2,211	15	27	53	75	140	207	279	376	953	2
1980	2,275	24	28	55	86	143	210	301	407	934	-
1979	2,096	25	41	55	95	115	199	296	370	808	2
Malnutrition (260-263)											
All races, male											
1998	1,216	9	16	23	38	66	90	181	232	519	-
1997	1,159	19	15	24	33	63	105	167	206	498	-
1996	1,155	15	23	24	31	63	100	164	195	517	1
1995	1,131	16	13	27	36	51	97	136	227	485	-
1994	1,112	14	12	24	37	60	104	149	206	463	1
1993	1,186	8	16	21	26	90	109	165	228	478	-
1992	1,034	10	19	19	31	65	96	135	202	411	-
1991	1,027	14	12	21	27	45	100	152	209	412	1
1990	1,052	14	13	20	32	64	86	156	204	424	1
1989	1,047	14	8	29	30	63	92	142	208	422	2
1988	1,039	8	11	19	41	66	104	135	215	388	-
1987	1,049	4	13	28	33	67	92	163	207	399	-
1986	1,019	8	10	25	41	69	121	151	184	373	-
1985	1,013	10	10	31	39	64	106	137	207	377	-
1984	922	4	14	26	36	72	115	126	159	337	-
1983	1,003	11	7	17	53	66	128	150	175	355	1
1982	944	7	15	26	38	66	115	158	173	310	2
1981	890	9	19	25	39	76	96	132	151	297	2
1980	931	12	16	27	41	86	100	142	155	306	-
1979	866	11	25	21	49	66	79	134	156	270	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
All races, female											
1998	2,665	12	14	24	26	63	109	237	450	1,695	-
1997	2,585	13	20	23	36	69	109	218	449	1,603	-
1996	2,376	11	16	20	25	52	128	237	374	1,483	-
1995	2,316	13	16	21	30	68	132	224	387	1,403	-
1994	2,162	11	12	17	28	59	109	214	353	1,329	-
1993	2,089	15	12	17	27	47	103	198	365	1,262	-
1992	1,964	17	8	16	30	49	113	195	336	1,162	-
1991	1,825	8	7	19	29	56	102	202	315	1,055	-
1990	1,816	8	8	28	40	63	90	205	305	1,040	-
1989	1,859	9	9	23	25	66	105	211	338	1,036	-
1988	1,803	8	4	16	39	59	107	194	317	1,018	-
1987	1,692	6	11	13	27	48	115	181	308	946	-
1986	1,673	9	8	18	38	58	118	179	271	935	1
1985	1,707	5	19	26	30	71	105	184	314	919	2
1984	1,543	6	10	20	36	50	109	176	284	826	-
1983	1,512	5	16	19	32	57	96	165	300	790	1
1982	1,370	10	9	25	33	60	103	177	248	665	1
1981	1,321	6	8	28	36	64	111	147	225	656	-
1980	1,344	12	12	28	45	57	110	159	252	628	-
1979	1,230	14	16	34	46	49	120	162	214	538	1
Malnutrition (260-263)											
White, both sexes											
1998	3,361	16	27	39	51	103	162	365	580	1,960	-
1997	3,284	21	24	37	51	99	175	338	582	1,908	-
1996	3,102	20	28	30	46	83	191	350	504	1,815	1
1995	3,002	23	19	36	49	92	195	295	534	1,710	-
1994	2,857	19	18	32	53	97	187	311	485	1,617	1
1993	2,833	17	20	28	35	103	172	309	511	1,590	-
1992	2,530	17	17	23	45	89	164	263	456	1,405	-
1991	2,434	19	15	30	43	74	154	307	461	1,278	1
1990	2,412	9	14	30	52	98	129	303	424	1,307	1
1989	2,452	19	14	36	39	99	157	288	468	1,282	1
1988	2,401	11	10	19	58	98	164	282	466	1,239	-
1987	2,349	9	13	30	42	92	166	293	450	1,213	-
1986	2,294	9	14	30	66	96	195	263	392	1,181	1
1985	2,364	8	23	43	54	103	169	279	449	1,189	2
1984	2,106	9	19	34	47	98	175	235	388	1,059	-
1983	2,133	11	19	27	62	101	170	263	409	1,032	1
1982	1,944	12	17	37	54	99	169	275	366	869	1
1981	1,890	11	15	44	62	116	166	239	329	850	1
1980	1,934	16	20	38	66	114	177	249	357	849	-
1979	1,760	17	29	38	80	80	158	255	320	731	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
White, male											
1998	1,014	5	13	18	31	52	71	152	188	451	-
1997	959	12	12	17	21	44	84	142	177	434	-
1996	977	11	18	13	25	40	83	141	171	455	1
1995	951	13	7	17	25	37	86	114	190	430	-
1994	934	10	9	20	29	47	95	122	175	406	1
1993	985	4	10	14	16	71	87	132	198	430	-
1992	828	6	11	9	22	51	72	108	165	356	-
1991	817	12	10	15	17	29	77	132	172	328	1
1990	855	4	8	14	22	53	65	129	168	365	1
1989	853	10	8	19	19	43	77	114	176	361	1
1988	837	4	7	9	28	50	80	113	189	327	-
1987	853	4	5	18	23	52	70	133	180	346	-
1986	823	2	6	18	31	55	95	115	154	324	-
1985	860	5	6	22	30	52	81	118	176	346	-
1984	751	3	11	18	22	63	92	93	135	292	-
1983	805	7	4	12	37	54	96	115	146	314	-
1982	772	5	10	21	30	55	90	121	148	272	-
1981	743	7	10	21	31	63	75	107	135	264	1
1980	755	5	10	17	26	65	81	120	131	273	-
1979	697	7	15	15	41	45	62	115	128	241	1
Malnutrition (260-263)											
White, female											
1998	2,347	11	14	21	20	51	91	213	392	1,509	-
1997	2,325	9	12	20	30	55	91	196	405	1,474	-
1996	2,125	9	10	17	21	43	108	209	333	1,360	-
1995	2,051	10	12	19	24	55	109	181	344	1,280	-
1994	1,923	9	9	12	24	50	92	189	310	1,211	-
1993	1,848	13	10	14	19	32	85	177	313	1,160	-
1992	1,702	11	6	14	23	38	92	155	291	1,049	-
1991	1,617	7	5	15	26	45	77	175	289	950	-
1990	1,557	5	6	16	30	45	64	174	256	942	-
1989	1,599	9	6	17	20	56	80	174	292	921	-
1988	1,564	7	3	10	30	48	84	169	277	912	-
1987	1,496	5	8	12	19	40	96	160	270	867	-
1986	1,471	7	8	12	35	41	100	148	238	857	1
1985	1,504	3	17	21	24	51	88	161	273	843	2
1984	1,355	6	8	16	25	35	83	142	253	767	-
1983	1,328	4	15	15	25	47	74	148	263	718	1
1982	1,172	7	7	16	24	44	79	154	218	597	1
1981	1,147	4	5	23	31	53	91	132	194	586	-
1980	1,179	11	10	21	40	49	96	129	226	576	-
1979	1,063	10	14	23	39	35	96	140	192	490	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
All other, both sexes											
1998	520	5	3	8	13	26	37	53	102	254	-
1997	460	11	11	10	18	33	39	47	73	193	-
1996	429	6	11	14	10	32	37	51	65	185	-
1995	445	6	10	12	17	27	34	65	80	178	-
1994	417	6	6	9	12	22	26	52	74	175	-
1993	442	6	8	10	18	34	40	54	82	150	-
1992	468	10	10	12	16	25	45	67	82	168	-
1991	418	3	4	10	13	27	48	47	63	189	-
1990	456	13	7	18	20	29	47	58	85	157	-
1989	454	4	3	16	16	30	40	65	78	176	1
1988	441	5	5	16	22	27	47	47	66	167	-
1987	392	1	11	11	18	23	41	51	65	132	-
1986	398	8	4	13	13	31	44	67	63	127	-
1985	356	7	6	14	15	32	42	42	72	107	-
1984	359	1	5	12	25	24	49	67	55	104	-
1983	382	5	4	9	23	22	54	52	66	113	1
1982	370	5	7	14	17	27	49	60	55	106	2
1981	321	4	12	9	13	24	41	40	47	103	1
1980	341	8	8	17	20	29	33	52	50	85	-
1979	336	8	12	17	15	35	41	41	50	77	-
Malnutrition (260-263)											
All other, male											
1998	202	4	3	5	7	14	19	29	44	68	-
1997	200	7	3	7	12	19	21	25	29	64	-
1996	178	4	5	11	6	23	17	23	24	62	-
1995	180	3	6	10	11	14	11	22	37	55	-
1994	178	4	3	4	8	13	9	27	31	57	-
1993	201	4	6	7	10	19	22	33	30	48	-
1992	206	4	8	10	9	14	24	27	37	55	-
1991	210	2	2	6	10	16	23	20	37	84	-
1990	197	10	5	6	10	11	21	27	36	59	-
1989	194	4	-	10	11	20	15	28	32	61	1
1988	202	4	4	10	13	16	24	22	26	61	-
1987	196	-	8	10	10	15	22	30	27	53	-
1986	196	6	4	7	10	14	26	36	30	49	-
1985	153	5	4	9	9	12	25	19	31	31	-
1984	171	1	3	8	14	9	23	33	24	45	-
1983	198	4	3	5	16	12	32	35	29	41	1
1982	172	2	5	5	8	11	25	37	25	38	2
1981	147	2	9	4	8	13	21	25	16	33	1
1980	176	7	6	10	15	21	19	22	24	33	-
1979	169	4	10	6	8	21	17	19	28	29	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
All other, female											
1998	318	1	-	3	6	12	18	24	58	186	-
1997	260	4	8	3	6	14	18	22	44	129	-
1996	251	2	6	3	4	9	20	28	41	123	-
1995	265	3	4	2	6	13	23	43	43	123	-
1994	239	2	3	5	4	9	17	25	43	118	-
1993	241	2	2	3	8	15	18	21	52	102	-
1992	262	6	2	2	7	11	21	40	45	113	-
1991	208	1	2	4	3	11	25	27	26	105	-
1990	259	3	2	12	10	18	26	31	49	98	-
1989	260	-	3	6	5	10	25	37	46	115	-
1988	239	1	1	6	9	11	23	25	40	106	-
1987	196	1	3	1	8	8	19	21	38	79	-
1986	202	2	-	6	3	17	18	31	33	78	-
1985	203	2	2	5	6	20	17	23	41	76	-
1984	188	-	2	4	11	15	26	34	31	59	-
1983	184	1	1	4	7	10	22	17	37	72	-
1982	198	3	2	9	9	16	24	23	30	68	-
1981	174	2	3	5	5	11	20	15	31	70	-
1980	165	1	2	7	5	8	14	30	26	52	-
1979	167	4	2	11	7	14	24	22	22	48	-
Malnutrition (260-263)											
Black, both sexes											
1998	465	5	3	8	12	26	34	49	93	216	-
1997	411	10	7	9	18	31	34	45	62	172	-
1996	376	5	8	12	8	30	32	51	57	156	-
1995	398	6	10	12	16	26	32	58	71	151	-
1994	385	6	5	7	12	20	26	50	70	156	-
1993	403	5	7	9	17	32	39	48	76	131	-
1992	422	10	9	12	16	23	39	62	70	148	-
1991	378	3	4	10	12	26	41	43	58	167	-
1990	420	13	7	17	19	27	44	56	76	142	-
1989	425	4	3	14	16	29	40	61	71	162	1
1988	405	5	5	16	22	26	43	43	58	153	-
1987	365	1	10	9	15	21	38	46	64	122	-
1986	367	8	4	13	13	30	40	62	58	111	-
1985	323	6	4	13	15	31	37	39	62	97	-
1984	328	1	5	12	24	23	45	60	53	89	-
1983	351	5	4	9	23	22	48	49	64	95	1
1982	337	5	7	13	17	26	48	58	46	90	2
1981	292	4	12	9	13	22	39	36	43	87	1
1980	321	7	8	17	20	28	31	48	42	82	-
1979	315	8	12	15	14	33	40	39	45	73	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Malnutrition (260-263)											
Black, male											
1998	180	4	3	5	7	14	18	27	38	55	-
1997	177	6	3	6	12	18	18	24	26	53	-
1996	156	3	4	11	4	22	15	23	22	49	-
1995	163	3	6	10	10	14	10	19	32	48	-
1994	164	4	3	4	8	11	9	25	30	49	-
1993	184	3	6	7	9	18	21	28	29	41	-
1992	184	4	7	10	9	12	20	26	30	48	-
1991	186	2	2	6	9	15	18	17	33	74	-
1990	182	10	5	6	10	11	21	25	32	51	-
1989	181	4	-	9	11	19	15	27	29	54	1
1988	187	4	4	10	13	15	23	21	23	56	-
1987	184	-	8	9	8	13	20	30	26	49	-
1986	182	6	4	7	10	13	23	34	29	42	-
1985	133	4	3	8	9	12	22	18	25	24	-
1984	154	1	3	8	13	9	21	28	22	38	-
1983	186	4	3	5	16	12	29	32	28	37	1
1982	155	2	5	5	8	11	24	36	18	31	2
1981	130	2	9	4	8	12	19	22	14	25	1
1980	166	6	6	10	15	20	18	21	20	31	-
1979	159	4	10	6	8	20	16	19	25	28	-
Malnutrition (260-263)											
Black, female											
1998	285	1	-	3	5	12	16	22	55	161	-
1997	234	4	4	3	6	13	16	21	36	119	-
1996	220	2	4	1	4	8	17	28	35	107	-
1995	235	3	4	2	6	12	22	39	39	103	-
1994	221	2	2	3	4	9	17	25	40	107	-
1993	219	2	1	2	8	14	18	20	47	90	-
1992	238	6	2	2	7	11	19	36	40	100	-
1991	192	1	2	4	3	11	23	26	25	93	-
1990	238	3	2	11	9	16	23	31	44	91	-
1989	244	-	3	5	5	10	25	34	42	108	-
1988	218	1	1	6	9	11	20	22	35	97	-
1987	181	1	2	-	7	8	18	16	38	73	-
1986	185	2	-	6	3	17	17	28	29	69	-
1985	190	2	1	5	6	19	15	21	37	73	-
1984	174	-	2	4	11	14	24	32	31	51	-
1983	165	1	1	4	7	10	19	17	36	58	-
1982	182	3	2	8	9	15	24	22	28	59	-
1981	162	2	3	5	5	10	20	14	29	62	-
1980	155	1	2	7	5	8	13	27	22	51	-
1979	156	4	2	9	6	13	24	20	20	45	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
All races, both sexes											
1998	196	2	3	2	6	7	9	34	32	97	-
1997	184	-	3	5	6	8	13	22	29	91	-
1996	175	2	-	7	4	3	14	23	31	78	-
1995	137	3	3	4	5	5	10	13	15	73	-
1994	177	1	3	3	6	6	10	19	33	86	-
1993	176	4	2	3	6	8	14	22	34	78	1
1992	137	1	5	4	3	5	5	17	15	71	-
1991	139	4	-	3	5	6	8	17	28	57	-
1990	149	-	2	1	4	10	18	16	22	68	-
1989	140	1	1	2	10	6	8	15	20	68	-
1988	143	1	4	3	2	10	9	26	20	62	-
1987	148	-	5	-	7	6	18	15	20	63	-
1986	170	1	5	7	7	10	21	15	26	69	-
1985	154	5	3	2	8	11	12	15	22	56	-
1984	135	2	3	1	6	10	10	21	12	59	-
1983	141	2	4	6	8	7	16	13	17	53	-
1982	116	1	2	5	6	5	13	12	21	40	-
1981	120	3	3	3	16	10	9	7	19	38	-
1980	89	4	1	3	9	5	14	10	10	26	-
1979	114	1	4	8	10	9	15	18	14	18	-
Other nutritional deficiencies (264-269)											
All races, male											
1998	62	1	1	1	3	4	5	11	8	25	-
1997	54	-	1	3	5	2	4	5	13	21	-
1996	70	1	-	3	3	3	8	13	10	21	-
1995	62	2	1	3	3	3	7	4	9	25	-
1994	74	1	2	2	5	4	6	11	14	22	-
1993	71	2	1	2	2	8	7	3	20	23	-
1992	55	1	2	3	2	4	2	11	9	15	-
1991	58	2	-	1	4	5	4	8	13	14	-
1990	56	-	1	1	2	6	7	10	6	21	-
1989	51	-	1	2	4	4	2	9	6	18	-
1988	57	1	4	3	-	8	5	4	11	17	-
1987	58	-	2	-	5	6	6	8	8	14	-
1986	69	1	1	3	3	7	12	8	10	21	-
1985	73	3	2	2	7	8	8	9	5	18	-
1984	55	1	2	1	5	6	5	10	4	15	-
1983	62	-	4	4	3	6	8	4	6	19	-
1982	50	-	1	4	5	4	8	4	10	8	-
1981	51	3	1	2	7	7	3	3	10	11	-
1980	31	1	-	-	3	3	8	3	1	6	-
1979	61	-	3	2	5	7	8	11	7	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
All races, female											
1998	134	1	2	1	3	3	4	23	24	72	-
1997	130	-	2	2	1	6	9	17	16	70	-
1996	105	1	-	4	1	-	6	10	21	57	-
1995	75	1	2	1	2	2	3	9	6	48	-
1994	103	-	1	1	1	2	4	8	19	64	-
1993	105	2	1	1	4	-	7	19	14	55	1
1992	82	-	3	1	1	1	3	6	6	56	-
1991	81	2	-	2	1	1	4	9	15	43	-
1990	93	-	1	-	2	4	11	6	16	47	-
1989	89	1	-	-	6	2	6	6	14	50	-
1988	86	-	-	-	2	2	4	22	9	45	-
1987	90	-	3	-	2	-	12	7	12	49	-
1986	101	-	4	4	4	3	9	7	16	48	-
1985	81	2	1	-	1	3	4	6	17	38	-
1984	80	1	1	-	1	4	5	11	8	44	-
1983	79	2	-	2	5	1	8	9	11	34	-
1982	66	1	1	1	1	1	5	8	11	32	-
1981	69	-	2	1	9	3	6	4	9	27	-
1980	58	3	1	3	6	2	6	7	9	20	-
1979	53	1	1	6	5	2	7	7	7	13	-
Other nutritional deficiencies (264-269)											
White, both sexes											
1998	178	2	3	2	6	3	9	31	31	89	-
1997	163	-	3	4	5	8	10	20	26	82	-
1996	153	1	-	5	4	3	13	23	24	71	-
1995	116	2	2	3	5	4	8	10	11	68	-
1994	150	-	-	1	4	5	9	17	31	78	-
1993	153	3	2	3	5	6	11	17	34	70	-
1992	112	1	5	2	2	2	4	15	11	61	-
1991	124	4	-	3	5	5	7	14	25	52	-
1990	134	-	1	-	3	8	17	15	20	62	-
1989	119	1	1	1	8	5	6	12	16	63	-
1988	125	1	2	2	1	7	8	24	15	60	-
1987	129	-	3	-	7	4	18	14	16	58	-
1986	144	1	3	4	7	10	21	11	19	61	-
1985	126	3	3	2	6	10	9	9	21	54	-
1984	125	2	3	1	5	9	9	19	10	58	-
1983	119	1	3	4	7	4	13	11	16	49	-
1982	105	-	2	5	6	4	13	12	16	36	-
1981	99	2	2	2	12	8	6	7	19	31	-
1980	74	4	1	3	8	4	13	9	9	20	-
1979	102	-	4	8	10	7	14	14	13	17	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
White, male											
1998	53	1	1	1	3	1	5	10	8	22	-
1997	44	-	1	2	4	2	3	4	11	17	-
1996	58	1	-	2	3	3	7	13	7	17	-
1995	48	2	-	2	3	3	5	3	6	22	-
1994	61	-	-	1	3	3	6	11	14	20	-
1993	61	1	1	2	1	6	5	3	20	20	-
1992	44	1	2	1	2	1	2	10	7	13	-
1991	51	2	-	1	4	4	3	5	13	13	-
1990	48	-	1	-	1	5	7	10	5	17	-
1989	42	-	1	1	4	3	-	7	5	17	-
1988	47	1	2	2	-	5	4	4	9	17	-
1987	48	-	1	-	5	4	6	7	7	13	-
1986	60	1	-	2	3	7	12	7	7	18	-
1985	56	2	2	2	5	7	5	5	5	17	-
1984	48	1	2	1	4	5	4	8	3	15	-
1983	48	-	3	2	3	3	5	4	6	17	-
1982	46	-	1	4	5	3	8	4	8	7	-
1981	43	2	1	1	5	5	3	3	10	10	-
1980	24	1	-	-	3	3	7	3	1	3	-
1979	53	-	3	2	5	5	7	8	7	5	-
Other nutritional deficiencies (264-269)											
White, female											
1998	125	1	2	1	3	2	4	21	23	67	-
1997	119	-	2	2	1	6	7	16	15	65	-
1996	95	-	-	3	1	-	6	10	17	54	-
1995	68	-	2	1	2	1	3	7	5	46	-
1994	89	-	-	-	1	2	3	6	17	58	-
1993	92	2	1	1	4	-	6	14	14	50	-
1992	68	-	3	1	-	1	2	5	4	48	-
1991	73	2	-	2	1	1	4	9	12	39	-
1990	86	-	-	-	2	3	10	5	15	45	-
1989	77	1	-	-	4	2	6	5	11	46	-
1988	78	-	-	-	1	2	4	20	6	43	-
1987	81	-	2	-	2	-	12	7	9	45	-
1986	84	-	3	2	4	3	9	4	12	43	-
1985	70	1	1	-	1	3	4	4	16	37	-
1984	77	1	1	-	1	4	5	11	7	43	-
1983	71	1	-	2	4	1	8	7	10	32	-
1982	59	-	1	1	1	1	5	8	8	29	-
1981	56	-	1	1	7	3	3	4	9	21	-
1980	50	3	1	3	5	1	6	6	8	17	-
1979	49	-	1	6	5	2	7	6	6	12	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
All other, both sexes											
1998	18	-	-	-	-	4	-	3	1	8	-
1997	21	-	-	1	1	-	3	2	3	9	-
1996	22	1	-	2	-	-	1	-	7	7	-
1995	21	1	1	1	-	1	2	3	4	5	-
1994	27	1	3	2	2	1	1	2	2	8	-
1993	23	1	-	-	1	2	3	5	-	8	1
1992	25	-	-	2	1	3	1	2	4	10	-
1991	15	-	-	-	-	1	1	3	3	5	-
1990	15	-	1	1	1	2	1	1	2	6	-
1989	21	-	-	1	2	1	2	3	4	5	-
1988	18	-	2	1	1	3	1	2	5	2	-
1987	19	-	2	-	-	2	-	1	4	5	-
1986	26	-	2	3	-	-	-	4	7	8	-
1985	28	2	-	-	2	1	3	6	1	2	-
1984	10	-	-	-	1	1	1	2	2	1	-
1983	22	1	1	2	1	3	3	2	1	4	-
1982	11	1	-	-	-	1	-	-	5	4	-
1981	21	1	1	1	4	2	3	-	-	7	-
1980	15	-	-	-	1	1	1	1	1	6	-
1979	12	1	-	-	-	2	1	4	1	1	-
Other nutritional deficiencies (264-269)											
All other, male											
1998	9	-	-	-	-	3	-	1	-	3	-
1997	10	-	-	1	1	-	1	1	2	4	-
1996	12	-	-	1	-	-	1	-	3	4	-
1995	14	-	1	1	-	-	2	1	3	3	-
1994	13	1	2	1	2	1	-	-	-	2	-
1993	10	1	-	-	1	2	2	-	-	3	-
1992	11	-	-	2	-	3	-	1	2	2	-
1991	7	-	-	-	-	1	1	3	-	1	-
1990	8	-	-	1	1	1	-	-	1	4	-
1989	9	-	-	1	-	1	2	2	1	1	-
1988	10	-	2	1	-	3	1	-	2	-	-
1987	10	-	1	-	-	2	-	1	1	1	-
1986	9	-	1	1	-	-	-	1	3	3	-
1985	17	1	-	-	2	1	3	4	-	1	-
1984	7	-	-	-	1	1	1	2	1	-	-
1983	14	-	1	2	-	3	3	-	-	2	-
1982	4	-	-	-	-	1	-	-	2	1	-
1981	8	1	-	1	2	2	-	-	-	1	-
1980	7	-	-	-	-	-	1	-	-	3	-
1979	8	-	-	-	-	2	1	3	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
All other, female											
1998	9	-	-	-	-	1	-	2	1	5	-
1997	11	-	-	-	-	-	2	1	1	5	-
1996	10	1	-	1	-	-	-	-	4	3	-
1995	7	1	-	-	-	1	-	2	1	2	-
1994	14	-	1	1	-	-	1	2	2	6	-
1993	13	-	-	-	-	-	1	5	-	5	1
1992	14	-	-	-	1	-	1	1	2	8	-
1991	8	-	-	-	-	-	-	-	3	4	-
1990	7	-	1	-	-	1	1	1	1	2	-
1989	12	-	-	-	2	-	-	1	3	4	-
1988	8	-	-	-	1	-	-	2	3	2	-
1987	9	-	1	-	-	-	-	-	3	4	-
1986	17	-	1	2	-	-	-	3	4	5	-
1985	11	1	-	-	-	-	-	2	1	1	-
1984	3	-	-	-	-	-	-	-	1	1	-
1983	8	1	-	-	1	-	-	2	1	2	-
1982	7	1	-	-	-	-	-	-	3	3	-
1981	13	-	1	-	2	-	3	-	-	6	-
1980	8	-	-	-	1	1	-	1	1	3	-
1979	4	1	-	-	-	-	-	1	1	1	-
Other nutritional deficiencies (264-269)											
Black, both sexes											
1998	15	-	-	-	-	3	-	2	1	7	-
1997	18	-	-	1	1	-	3	2	3	6	-
1996	20	1	-	2	-	-	1	-	6	6	-
1995	20	1	1	1	-	1	2	3	4	4	-
1994	26	1	3	2	2	1	1	2	2	7	-
1993	21	1	-	-	1	2	3	5	-	7	1
1992	23	-	-	2	1	3	1	2	3	9	-
1991	15	-	-	-	-	1	1	3	3	5	-
1990	13	-	1	1	1	2	1	-	2	5	-
1989	18	-	-	1	1	-	1	3	4	5	-
1988	15	-	1	1	1	2	1	1	5	2	-
1987	18	-	2	-	-	2	-	1	4	4	-
1986	25	-	2	3	-	-	-	4	7	7	-
1985	26	2	-	-	2	1	3	4	1	2	-
1984	10	-	-	-	1	1	1	2	2	1	-
1983	21	1	1	2	1	3	3	2	1	3	-
1982	9	-	-	-	-	1	-	-	5	3	-
1981	18	1	1	1	4	1	3	-	-	5	-
1980	12	-	-	-	-	1	1	1	1	4	-
1979	10	1	-	-	-	1	1	3	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other nutritional deficiencies (264-269)											
Black, male											
1998	8	-	-	-	-	3	-	-	-	3	-
1997	8	-	-	1	1	-	1	1	2	2	-
1996	11	-	-	1	-	-	1	-	2	4	-
1995	13	-	1	1	-	-	2	1	3	2	-
1994	13	1	2	1	2	1	-	-	-	2	-
1993	9	1	-	-	1	2	2	-	-	3	-
1992	11	-	-	2	-	3	-	1	2	2	-
1991	7	-	-	-	-	1	1	3	-	1	-
1990	7	-	-	1	1	1	-	-	1	3	-
1989	7	-	-	1	-	-	1	2	1	1	-
1988	8	-	1	1	-	2	1	-	2	-	-
1987	10	-	1	-	-	2	-	1	1	1	-
1986	8	-	1	1	-	-	-	1	3	2	-
1985	16	1	-	-	2	1	3	3	-	1	-
1984	7	-	-	-	1	1	1	2	1	-	-
1983	13	-	1	2	-	3	3	-	-	1	-
1982	4	-	-	-	-	1	-	-	2	1	-
1981	7	1	-	1	2	1	-	-	-	1	-
1980	6	-	-	-	-	-	1	-	-	2	-
1979	6	-	-	-	-	1	1	2	-	-	-
Other nutritional deficiencies (264-269)											
Black, female											
1998	7	-	-	-	-	-	-	2	1	4	-
1997	10	-	-	-	-	-	2	1	1	4	-
1996	9	1	-	1	-	-	-	-	4	2	-
1995	7	1	-	-	-	1	-	2	1	2	-
1994	13	-	1	1	-	-	1	2	2	5	-
1993	12	-	-	-	-	-	1	5	-	4	1
1992	12	-	-	-	1	-	1	1	1	7	-
1991	8	-	-	-	-	-	-	-	3	4	-
1990	6	-	1	-	-	1	1	-	1	2	-
1989	11	-	-	-	1	-	-	1	3	4	-
1988	7	-	-	-	1	-	-	1	3	2	-
1987	8	-	1	-	-	-	-	-	3	3	-
1986	17	-	1	2	-	-	-	3	4	5	-
1985	10	1	-	-	-	-	-	1	1	1	-
1984	3	-	-	-	-	-	-	-	1	1	-
1983	8	1	-	-	1	-	-	2	1	2	-
1982	5	-	-	-	-	-	-	-	3	2	-
1981	11	-	1	-	2	-	3	-	-	4	-
1980	6	-	-	-	-	1	-	1	1	2	-
1979	4	1	-	-	-	-	-	1	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
All races, both sexes											
1998	768	49	43	39	39	49	47	56	50	41	-
1997	781	44	43	45	40	51	48	74	45	53	-
1996	758	53	36	34	42	47	60	63	53	49	3
1995	761	51	37	38	56	50	48	43	44	54	-
1994	770	41	39	42	50	40	71	55	51	49	-
1993	788	42	32	36	39	71	58	66	60	50	-
1992	798	27	39	47	44	56	63	66	52	58	-
1991	887	33	29	38	58	76	71	69	71	75	-
1990	1,017	37	44	49	67	70	76	67	61	61	-
1989	1,068	38	45	38	68	87	74	59	63	65	2
1988	1,156	39	36	51	71	86	103	92	70	80	-
1987	1,161	38	53	50	77	72	88	61	71	83	-
1986	1,160	33	36	61	93	81	94	85	67	70	-
1985	1,206	23	45	67	62	75	98	102	73	64	-
1984	1,194	27	43	72	73	62	73	82	74	63	1
1983	1,287	44	36	54	82	88	84	81	61	60	3
1982	1,282	33	49	61	77	75	74	75	53	82	-
1981	1,405	51	51	82	80	77	70	73	57	68	1
1980	1,415	28	50	62	84	79	94	57	61	45	-
1979	1,393	38	48	68	81	88	81	64	57	46	-
Meningitis (320-322)											
All races, male											
1998	424	34	27	22	18	29	29	29	22	8	-
1997	403	33	28	27	23	30	26	34	18	15	-
1996	423	35	28	19	27	27	30	31	21	14	3
1995	408	26	28	24	28	29	22	21	16	14	-
1994	418	26	24	23	23	20	30	27	16	21	-
1993	420	26	16	18	22	42	27	31	26	15	-
1992	402	15	24	35	21	24	32	26	22	17	-
1991	442	20	17	20	37	33	26	25	26	26	-
1990	545	23	27	23	35	32	35	26	22	22	-
1989	588	28	24	21	37	51	29	19	25	23	1
1988	601	27	24	36	41	40	41	34	20	28	-
1987	645	23	33	28	53	34	54	29	24	22	-
1986	624	16	16	31	53	44	36	42	28	20	-
1985	634	15	21	35	31	36	53	42	25	20	-
1984	650	13	27	41	39	36	35	48	27	15	-
1983	670	26	21	29	48	41	37	30	19	20	2
1982	700	16	30	36	47	40	36	22	17	31	-
1981	776	30	36	46	42	41	31	31	15	24	1
1980	796	18	32	36	58	38	46	30	28	13	-
1979	765	20	28	46	47	47	36	29	24	15	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
All races, female											
1998	344	15	16	17	21	20	18	27	28	33	-
1997	378	11	15	18	17	21	22	40	27	38	-
1996	335	18	8	15	15	20	30	32	32	35	-
1995	353	25	9	14	28	21	26	22	28	40	-
1994	352	15	15	19	27	20	41	28	35	28	-
1993	368	16	16	18	17	29	31	35	34	35	-
1992	396	12	15	12	23	32	31	40	30	41	-
1991	445	13	12	18	21	43	45	44	45	49	-
1990	472	14	17	26	32	38	41	41	39	39	-
1989	480	10	21	17	31	36	45	40	38	42	1
1988	555	12	12	15	30	46	62	58	50	52	-
1987	516	15	20	22	24	38	34	32	47	61	-
1986	536	17	20	30	40	37	58	43	39	50	-
1985	572	8	24	32	31	39	45	60	48	44	-
1984	544	14	16	31	34	26	38	34	47	48	1
1983	617	18	15	25	34	47	47	51	42	40	1
1982	582	17	19	25	30	35	38	53	36	51	-
1981	629	21	15	36	38	36	39	42	42	44	-
1980	619	10	18	26	26	41	48	27	33	32	-
1979	628	18	20	22	34	41	45	35	33	31	-
Meningitis (320-322)											
White, both sexes											
1998	545	31	32	26	27	37	36	50	43	35	-
1997	546	25	31	35	32	35	38	62	38	46	-
1996	529	32	27	23	31	38	49	55	46	38	2
1995	541	32	28	21	45	44	37	32	36	46	-
1994	542	25	21	31	43	33	62	39	46	39	-
1993	547	29	22	24	25	52	45	60	48	40	-
1992	569	18	24	30	37	41	52	59	40	53	-
1991	618	25	22	32	39	60	55	58	55	55	-
1990	705	21	25	36	54	60	60	50	47	48	-
1989	703	18	25	21	45	66	53	48	55	52	1
1988	808	18	19	36	52	62	85	69	53	63	-
1987	812	24	31	31	57	56	72	47	60	64	-
1986	827	19	26	45	65	64	74	67	56	56	-
1985	871	12	30	55	42	60	78	83	63	57	-
1984	846	18	24	48	51	43	63	65	61	54	-
1983	935	26	28	42	63	61	61	70	49	53	3
1982	912	19	31	45	54	61	60	56	42	71	-
1981	987	28	35	56	60	56	49	55	47	61	-
1980	1,020	16	33	48	66	50	74	45	51	36	-
1979	951	19	33	44	58	61	67	51	49	33	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
White, male											
1998	297	23	18	16	11	20	22	27	19	5	-
1997	275	20	19	23	17	19	24	28	15	14	-
1996	294	21	22	13	18	23	25	29	17	10	2
1995	290	13	22	14	21	27	20	15	14	12	-
1994	285	16	10	15	23	18	26	16	16	15	-
1993	286	17	10	9	16	32	21	29	18	13	-
1992	280	11	12	21	16	21	28	24	16	13	-
1991	306	16	13	17	24	28	21	21	19	20	-
1990	371	12	17	15	29	27	27	20	15	16	-
1989	383	13	12	13	24	39	23	15	23	21	1
1988	402	10	12	26	31	30	34	25	14	21	-
1987	460	15	20	18	40	27	47	23	17	21	-
1986	445	10	10	24	35	36	28	32	25	18	-
1985	451	10	15	30	22	28	43	29	22	19	-
1984	472	10	14	26	29	24	29	39	21	15	-
1983	486	15	17	22	37	32	29	24	12	20	2
1982	492	7	18	26	33	34	29	16	13	25	-
1981	547	23	27	33	31	26	24	23	11	21	-
1980	577	9	24	26	48	23	36	24	25	10	-
1979	521	13	18	30	31	35	31	22	20	8	-
Meningitis (320-322)											
White, female											
1998	248	8	14	10	16	17	14	23	24	30	-
1997	271	5	12	12	15	16	14	34	23	32	-
1996	235	11	5	10	13	15	24	26	29	28	-
1995	251	19	6	7	24	17	17	17	22	34	-
1994	257	9	11	16	20	15	36	23	30	24	-
1993	261	12	12	15	9	20	24	31	30	27	-
1992	289	7	12	9	21	20	24	35	24	40	-
1991	312	9	9	15	15	32	34	37	36	35	-
1990	334	9	8	21	25	33	33	30	32	32	-
1989	320	5	13	8	21	27	30	33	32	31	-
1988	406	8	7	10	21	32	51	44	39	42	-
1987	352	9	11	13	17	29	25	24	43	43	-
1986	382	9	16	21	30	28	46	35	31	38	-
1985	420	2	15	25	20	32	35	54	41	38	-
1984	374	8	10	22	22	19	34	26	40	39	-
1983	449	11	11	20	26	29	32	46	37	33	1
1982	420	12	13	19	21	27	31	40	29	46	-
1981	440	5	8	23	29	30	25	32	36	40	-
1980	443	7	9	22	18	27	38	21	26	26	-
1979	430	6	15	14	27	26	36	29	29	25	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
All other, both sexes											
1998	223	18	11	13	12	12	11	6	7	6	-
1997	235	19	12	10	8	16	10	12	7	7	-
1996	229	21	9	11	11	9	11	8	7	11	1
1995	220	19	9	17	11	6	11	11	8	8	-
1994	228	16	18	11	7	7	9	16	5	10	-
1993	241	13	10	12	14	19	13	6	12	10	-
1992	229	9	15	17	7	15	11	7	12	5	-
1991	269	8	7	6	19	16	16	11	16	20	-
1990	312	16	19	13	13	10	16	17	14	13	-
1989	365	20	20	17	23	21	21	11	8	13	1
1988	348	21	17	15	19	24	18	23	17	17	-
1987	349	14	22	19	20	16	16	14	11	19	-
1986	333	14	10	16	28	17	20	18	11	14	-
1985	335	11	15	12	20	15	20	19	10	7	-
1984	348	9	19	24	22	19	10	17	13	9	1
1983	352	18	8	12	19	27	23	11	12	7	-
1982	370	14	18	16	23	14	14	19	11	11	-
1981	418	23	16	26	20	21	21	18	10	7	1
1980	395	12	17	14	18	29	20	12	10	9	-
1979	442	19	15	24	23	27	14	13	8	13	-
Meningitis (320-322)											
All other, male											
1998	127	11	9	6	7	9	7	2	3	3	-
1997	128	13	9	4	6	11	2	6	3	1	-
1996	129	14	6	6	9	4	5	2	4	4	1
1995	118	13	6	10	7	2	2	6	2	2	-
1994	133	10	14	8	-	2	4	11	-	6	-
1993	134	9	6	9	6	10	6	2	8	2	-
1992	122	4	12	14	5	3	4	2	6	4	-
1991	136	4	4	3	13	5	5	4	7	6	-
1990	174	11	10	8	6	5	8	6	7	6	-
1989	205	15	12	8	13	12	6	4	2	2	-
1988	199	17	12	10	10	10	7	9	6	7	-
1987	185	8	13	10	13	7	7	6	7	1	-
1986	179	6	6	7	18	8	8	10	3	2	-
1985	183	5	6	5	9	8	10	13	3	1	-
1984	178	3	13	15	10	12	6	9	6	-	-
1983	184	11	4	7	11	9	8	6	7	-	-
1982	208	9	12	10	14	6	7	6	4	6	-
1981	229	7	9	13	11	15	7	8	4	3	1
1980	219	9	8	10	10	15	10	6	3	3	-
1979	244	7	10	16	16	12	5	7	4	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
All other, female											
1998	96	7	2	7	5	3	4	4	4	3	-
1997	107	6	3	6	2	5	8	6	4	6	-
1996	100	7	3	5	2	5	6	6	3	7	-
1995	102	6	3	7	4	4	9	5	6	6	-
1994	95	6	4	3	7	5	5	5	5	4	-
1993	107	4	4	3	8	9	7	4	4	8	-
1992	107	5	3	3	2	12	7	5	6	1	-
1991	133	4	3	3	6	11	11	7	9	14	-
1990	138	5	9	5	7	5	8	11	7	7	-
1989	160	5	8	9	10	9	15	7	6	11	1
1988	149	4	5	5	9	14	11	14	11	10	-
1987	164	6	9	9	7	9	9	8	4	18	-
1986	154	8	4	9	10	9	12	8	8	12	-
1985	152	6	9	7	11	7	10	6	7	6	-
1984	170	6	6	9	12	7	4	8	7	9	1
1983	168	7	4	5	8	18	15	5	5	7	-
1982	162	5	6	6	9	8	7	13	7	5	-
1981	189	16	7	13	9	6	14	10	6	4	-
1980	176	3	9	4	8	14	10	6	7	6	-
1979	198	12	5	8	7	15	9	6	4	6	-
Meningitis (320-322)											
Black, both sexes											
1998	195	16	11	9	12	12	11	5	4	6	-
1997	204	18	11	7	8	13	8	11	5	6	-
1996	211	19	8	9	11	7	10	7	5	11	1
1995	197	17	9	15	11	6	7	10	7	7	-
1994	210	15	18	10	6	7	8	13	3	9	-
1993	216	12	9	11	12	15	10	6	11	9	-
1992	204	9	13	16	7	12	10	6	12	5	-
1991	245	8	6	6	16	15	14	11	15	19	-
1990	284	16	17	13	13	9	13	16	13	10	-
1989	331	20	19	16	19	19	17	11	8	12	1
1988	326	20	17	14	18	21	17	22	16	16	-
1987	319	11	22	15	20	15	15	14	10	19	-
1986	305	12	10	12	24	17	19	18	10	13	-
1985	301	11	15	10	20	14	20	15	8	7	-
1984	319	9	19	22	21	18	10	16	13	9	-
1983	332	17	7	11	17	27	23	11	11	6	-
1982	339	14	17	14	23	12	14	19	10	10	-
1981	380	21	14	22	18	18	20	15	10	7	1
1980	369	11	16	14	17	27	20	12	9	7	-
1979	404	16	13	21	23	22	14	12	8	13	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Meningitis (320-322)											
Black, male											
1998	114	10	9	6	7	9	7	1	2	3	-
1997	112	13	8	2	6	8	1	5	3	1	-
1996	119	12	5	6	9	3	5	2	2	4	1
1995	106	11	6	9	7	2	1	5	2	1	-
1994	122	9	14	8	-	2	4	8	-	5	-
1993	119	9	6	9	6	6	5	2	7	1	-
1992	107	4	10	14	5	1	3	1	6	4	-
1991	123	4	3	3	10	4	3	4	6	6	-
1990	159	11	8	8	6	4	6	6	7	5	-
1989	186	15	11	8	11	10	5	4	2	2	-
1988	185	16	12	9	9	8	6	9	5	6	-
1987	169	6	13	9	13	7	6	6	6	1	-
1986	165	6	6	6	16	8	7	10	2	2	-
1985	163	5	6	4	9	8	10	10	2	1	-
1984	164	3	13	13	10	11	6	8	6	-	-
1983	175	10	4	7	10	9	8	6	7	-	-
1982	196	9	11	9	14	6	7	6	4	6	-
1981	204	6	8	11	9	13	6	6	4	3	1
1980	203	8	7	10	9	13	10	6	3	3	-
1979	222	6	9	15	16	10	5	6	4	7	-
Meningitis (320-322)											
Black, female											
1998	81	6	2	3	5	3	4	4	2	3	-
1997	92	5	3	5	2	5	7	6	2	5	-
1996	92	7	3	3	2	4	5	5	3	7	-
1995	91	6	3	6	4	4	6	5	5	6	-
1994	88	6	4	2	6	5	4	5	3	4	-
1993	97	3	3	2	6	9	5	4	4	8	-
1992	97	5	3	2	2	11	7	5	6	1	-
1991	122	4	3	3	6	11	11	7	9	13	-
1990	125	5	9	5	7	5	7	10	6	5	-
1989	145	5	8	8	8	9	12	7	6	10	1
1988	141	4	5	5	9	13	11	13	11	10	-
1987	150	5	9	6	7	8	9	8	4	18	-
1986	140	6	4	6	8	9	12	8	8	11	-
1985	138	6	9	6	11	6	10	5	6	6	-
1984	155	6	6	9	11	7	4	8	7	9	-
1983	157	7	3	4	7	18	15	5	4	6	-
1982	143	5	6	5	9	6	7	13	6	4	-
1981	176	15	6	11	9	5	14	9	6	4	-
1980	166	3	9	4	8	14	10	6	6	4	-
1979	182	10	4	6	7	12	9	6	4	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
All races, both sexes											
1998	13,167	13	22	56	167	483	1,465	2,810	3,594	4,553	-
1997	12,361	12	24	63	173	502	1,287	2,672	3,451	4,169	-
1996	11,845	15	18	54	162	439	1,337	2,580	3,275	3,964	-
1995	10,820	9	22	62	146	440	1,253	2,311	3,005	3,559	-
1994	9,957	10	28	53	153	454	1,208	2,138	2,770	3,140	-
1993	9,107	14	17	35	138	394	1,103	2,028	2,506	2,868	-
1992	7,769	8	17	42	155	421	973	1,795	2,086	2,264	1
1991	7,455	2	7	42	131	405	980	1,708	2,003	2,170	-
1990	7,254	4	19	31	126	422	969	1,739	1,954	1,985	1
1989	6,984	5	14	46	137	449	954	1,641	1,879	1,855	-
1988	6,397	6	12	44	155	400	887	1,484	1,729	1,670	-
1987	6,288	1	18	37	151	411	927	1,569	1,612	1,559	-
1986	5,579	5	13	43	155	403	814	1,376	1,449	1,317	-
1985	5,173	1	15	43	117	365	792	1,266	1,355	1,214	-
1984	4,937	4	11	64	143	371	818	1,281	1,217	1,022	-
1983	4,567	4	9	75	165	362	767	1,193	1,105	882	-
1982	4,106	5	19	42	151	345	711	1,057	968	805	1
1981	3,948	4	16	47	164	360	768	1,005	906	671	1
1980	3,770	3	22	65	137	371	689	997	845	638	-
1979	3,510	5	13	59	155	374	699	904	748	550	-
Parkinson's disease (332)											
All races, male											
1998	7,494	8	16	39	95	327	949	1,759	2,086	2,212	-
1997	7,024	5	12	35	98	338	832	1,686	2,029	1,983	-
1996	6,908	8	12	32	96	285	920	1,663	1,916	1,975	-
1995	6,177	5	15	41	94	278	814	1,431	1,740	1,750	-
1994	5,807	7	17	32	97	300	800	1,339	1,630	1,583	-
1993	5,328	6	10	20	76	261	736	1,323	1,463	1,431	-
1992	4,541	7	8	24	95	277	648	1,131	1,226	1,120	-
1991	4,360	1	3	34	95	251	621	1,128	1,165	1,059	-
1990	4,244	2	9	19	76	272	630	1,097	1,159	979	1
1989	3,996	2	6	26	91	296	628	1,024	1,060	861	-
1988	3,661	5	4	25	87	253	575	924	980	801	-
1987	3,622	1	11	21	92	255	604	989	898	750	-
1986	3,185	4	6	22	88	271	528	821	806	637	-
1985	2,885	-	10	29	72	224	512	744	722	570	-
1984	2,782	1	6	42	93	238	499	743	662	493	-
1983	2,602	2	4	46	109	227	458	721	605	426	-
1982	2,320	4	10	28	87	207	442	633	525	381	1
1981	2,204	4	8	27	99	232	464	575	490	303	-
1980	2,094	2	15	41	82	236	440	538	443	294	-
1979	1,923	4	8	33	96	233	409	511	379	249	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
All races, female											
1998	5,673	5	6	17	72	156	516	1,051	1,508	2,341	-
1997	5,337	7	12	28	75	164	455	986	1,422	2,186	-
1996	4,937	7	6	22	66	154	417	917	1,359	1,989	-
1995	4,643	4	7	21	52	162	439	880	1,265	1,809	-
1994	4,150	3	11	21	56	154	408	799	1,140	1,557	-
1993	3,779	8	7	15	62	133	367	705	1,043	1,437	-
1992	3,228	1	9	18	60	144	325	664	860	1,144	1
1991	3,095	1	4	8	36	154	359	580	838	1,111	-
1990	3,010	2	10	12	50	150	339	642	795	1,006	-
1989	2,988	3	8	20	46	153	326	617	819	994	-
1988	2,736	1	8	19	68	147	312	560	749	869	-
1987	2,666	-	7	16	59	156	323	580	714	809	-
1986	2,394	1	7	21	67	132	286	555	643	680	-
1985	2,288	1	5	14	45	141	280	522	633	644	-
1984	2,155	3	5	22	50	133	319	538	555	529	-
1983	1,965	2	5	29	56	135	309	472	500	456	-
1982	1,786	1	9	14	64	138	269	424	443	424	-
1981	1,744	-	8	20	65	128	304	430	416	368	1
1980	1,676	1	7	24	55	135	249	459	402	344	-
1979	1,587	1	5	26	59	141	290	393	369	301	-
Parkinson's disease (332)											
White, both sexes											
1998	12,556	12	20	47	150	452	1,392	2,669	3,464	4,348	-
1997	11,838	12	23	59	164	473	1,220	2,551	3,329	4,000	-
1996	11,351	15	16	49	145	411	1,256	2,464	3,170	3,824	-
1995	10,349	8	20	57	134	404	1,177	2,217	2,884	3,442	-
1994	9,514	8	27	47	145	419	1,137	2,039	2,661	3,029	-
1993	8,723	9	12	29	128	374	1,044	1,955	2,407	2,761	-
1992	7,434	8	15	40	148	391	927	1,714	1,999	2,186	1
1991	7,125	2	7	36	117	373	926	1,621	1,939	2,098	-
1990	6,918	2	18	28	119	384	919	1,650	1,875	1,919	-
1989	6,652	4	11	45	124	422	893	1,556	1,807	1,787	-
1988	6,118	6	11	39	144	372	836	1,414	1,669	1,617	-
1987	6,024	1	14	35	134	389	884	1,497	1,551	1,516	-
1986	5,350	5	12	40	149	378	767	1,322	1,401	1,272	-
1985	4,974	1	14	41	110	336	756	1,217	1,315	1,180	-
1984	4,725	3	10	58	129	352	767	1,230	1,174	997	-
1983	4,347	3	7	73	150	332	721	1,142	1,058	856	-
1982	3,943	4	19	37	142	324	671	1,018	944	781	1
1981	3,805	4	13	42	162	344	729	971	882	652	1
1980	3,626	3	20	60	130	354	667	955	815	619	-
1979	3,361	5	13	56	144	350	658	871	729	532	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
White, male											
1998	7,155	7	14	33	89	310	905	1,677	2,016	2,103	-
1997	6,707	5	11	32	95	316	788	1,612	1,953	1,890	-
1996	6,620	8	10	28	85	268	864	1,594	1,856	1,906	-
1995	5,906	4	14	37	86	256	767	1,374	1,676	1,688	-
1994	5,528	6	16	27	91	274	750	1,277	1,564	1,521	-
1993	5,080	3	8	17	68	248	696	1,276	1,398	1,364	-
1992	4,338	7	8	23	91	256	620	1,079	1,174	1,077	-
1991	4,167	1	3	30	85	233	587	1,074	1,127	1,025	-
1990	4,042	1	9	18	71	245	597	1,045	1,111	945	-
1989	3,795	2	6	25	82	275	586	968	1,021	828	-
1988	3,494	5	4	22	79	239	542	881	940	775	-
1987	3,456	1	9	21	82	240	576	942	855	729	-
1986	3,030	4	5	21	83	250	492	788	779	606	-
1985	2,766	-	10	27	66	207	491	710	701	552	-
1984	2,661	1	5	39	85	226	468	715	636	482	-
1983	2,471	1	4	45	100	205	432	691	578	411	-
1982	2,221	4	10	23	81	196	414	607	513	370	1
1981	2,122	4	8	26	98	226	436	550	477	296	-
1980	2,005	2	13	38	79	223	425	514	425	283	-
1979	1,837	4	8	33	89	220	386	492	366	238	-
Parkinson's disease (332)											
White, female											
1998	5,401	5	6	14	61	142	487	992	1,448	2,245	-
1997	5,131	7	12	27	69	157	432	939	1,376	2,110	-
1996	4,731	7	6	21	60	143	392	870	1,314	1,918	-
1995	4,443	4	6	20	48	148	410	843	1,208	1,754	-
1994	3,986	2	11	20	54	145	387	762	1,097	1,508	-
1993	3,643	6	4	12	60	126	348	679	1,009	1,397	-
1992	3,096	1	7	17	57	135	307	635	825	1,109	1
1991	2,958	1	4	6	32	140	339	547	812	1,073	-
1990	2,876	1	9	10	48	139	322	605	764	974	-
1989	2,857	2	5	20	42	147	307	588	786	959	-
1988	2,624	1	7	17	65	133	294	533	729	842	-
1987	2,568	-	5	14	52	149	308	555	696	787	-
1986	2,320	1	7	19	66	128	275	534	622	666	-
1985	2,208	1	4	14	44	129	265	507	614	628	-
1984	2,064	2	5	19	44	126	299	515	538	515	-
1983	1,876	2	3	28	50	127	289	451	480	445	-
1982	1,722	-	9	14	61	128	257	411	431	411	-
1981	1,683	-	5	16	64	118	293	421	405	356	1
1980	1,621	1	7	22	51	131	242	441	390	336	-
1979	1,524	1	5	23	55	130	272	379	363	294	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
All other, both sexes											
1998	611	1	2	9	17	31	73	141	130	205	-
1997	523	-	1	4	9	29	67	121	122	169	-
1996	494	-	2	5	17	28	81	116	105	140	-
1995	471	1	2	5	12	36	76	94	121	117	-
1994	443	2	1	6	8	35	71	99	109	111	-
1993	384	5	5	6	10	20	59	73	99	107	-
1992	335	-	2	2	7	30	46	81	87	78	-
1991	330	-	-	6	14	32	54	87	64	72	-
1990	336	2	1	3	7	38	50	89	79	66	1
1989	332	1	3	1	13	27	61	85	72	68	-
1988	279	-	1	5	11	28	51	70	60	53	-
1987	264	-	4	2	17	22	43	72	61	43	-
1986	229	-	1	3	6	25	47	54	48	45	-
1985	199	-	1	2	7	29	36	49	40	34	-
1984	212	1	1	6	14	19	51	51	43	25	-
1983	220	1	2	2	15	30	46	51	47	26	-
1982	163	1	-	5	9	21	40	39	24	24	-
1981	143	-	3	5	2	16	39	34	24	19	-
1980	144	-	2	5	7	17	22	42	30	19	-
1979	149	-	-	3	11	24	41	33	19	18	-
Parkinson's disease (332)											
All other, male											
1998	339	1	2	6	6	17	44	82	70	109	-
1997	317	-	1	3	3	22	44	74	76	93	-
1996	288	-	2	4	11	17	56	69	60	69	-
1995	271	1	1	4	8	22	47	57	64	62	-
1994	279	1	1	5	6	26	50	62	66	62	-
1993	248	3	2	3	8	13	40	47	65	67	-
1992	203	-	-	1	4	21	28	52	52	43	-
1991	193	-	-	4	10	18	34	54	38	34	-
1990	202	1	-	1	5	27	33	52	48	34	1
1989	201	-	-	1	9	21	42	56	39	33	-
1988	167	-	-	3	8	14	33	43	40	26	-
1987	166	-	2	-	10	15	28	47	43	21	-
1986	155	-	1	1	5	21	36	33	27	31	-
1985	119	-	-	2	6	17	21	34	21	18	-
1984	121	-	1	3	8	12	31	28	26	11	-
1983	131	1	-	1	9	22	26	30	27	15	-
1982	99	-	-	5	6	11	28	26	12	11	-
1981	82	-	-	1	1	6	28	25	13	7	-
1980	89	-	2	3	3	13	15	24	18	11	-
1979	86	-	-	-	7	13	23	19	13	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
All other, female											
1998	272	-	-	3	11	14	29	59	60	96	-
1997	206	-	-	1	6	7	23	47	46	76	-
1996	206	-	-	1	6	11	25	47	45	71	-
1995	200	-	1	1	4	14	29	37	57	55	-
1994	164	1	-	1	2	9	21	37	43	49	-
1993	136	2	3	3	2	7	19	26	34	40	-
1992	132	-	2	1	3	9	18	29	35	35	-
1991	137	-	-	2	4	14	20	33	26	38	-
1990	134	1	1	2	2	11	17	37	31	32	-
1989	131	1	3	-	4	6	19	29	33	35	-
1988	112	-	1	2	3	14	18	27	20	27	-
1987	98	-	2	2	7	7	15	25	18	22	-
1986	74	-	-	2	1	4	11	21	21	14	-
1985	80	-	1	-	1	12	15	15	19	16	-
1984	91	1	-	3	6	7	20	23	17	14	-
1983	89	-	2	1	6	8	20	21	20	11	-
1982	64	1	-	-	3	10	12	13	12	13	-
1981	61	-	3	4	1	10	11	9	11	12	-
1980	55	-	-	2	4	4	7	18	12	8	-
1979	63	-	-	3	4	11	18	14	6	7	-
Parkinson's disease (332)											
Black, both sexes											
1998	454	1	2	7	12	21	60	109	98	143	-
1997	369	-	1	1	7	23	51	76	87	122	-
1996	370	-	2	5	12	19	63	84	82	103	-
1995	349	1	2	3	10	30	54	67	90	85	-
1994	342	2	1	5	7	25	55	66	86	94	-
1993	291	5	4	5	9	15	48	56	78	71	-
1992	252	-	1	1	6	23	30	63	70	56	-
1991	258	-	-	6	13	22	45	61	55	55	-
1990	245	2	1	1	6	28	37	66	57	46	1
1989	265	1	2	1	11	23	50	68	57	51	-
1988	216	-	1	3	9	18	40	57	47	41	-
1987	200	-	3	1	12	16	34	55	48	31	-
1986	184	-	1	1	4	20	43	45	35	35	-
1985	158	-	-	1	6	25	30	39	29	27	-
1984	166	-	-	5	13	17	41	41	31	17	-
1983	176	1	2	2	12	26	33	43	40	17	-
1982	141	1	-	4	9	19	34	32	21	21	-
1981	125	-	2	5	2	13	35	31	22	15	-
1980	123	-	2	5	6	15	22	34	24	15	-
1979	119	-	-	2	9	21	35	27	15	10	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Parkinson's disease (332)											
Black, male											
1998	251	1	2	4	4	12	38	63	53	73	-
1997	215	-	1	1	2	18	34	45	53	60	-
1996	209	-	2	4	7	10	45	49	48	44	-
1995	202	1	1	2	7	19	30	41	51	45	-
1994	210	1	1	4	6	20	35	41	51	51	-
1993	185	3	2	3	7	10	31	37	52	40	-
1992	149	-	-	-	4	16	19	41	39	28	-
1991	145	-	-	4	10	12	32	33	30	23	-
1990	146	1	-	-	4	21	25	37	34	23	1
1989	158	-	-	1	8	18	32	45	29	25	-
1988	129	-	-	2	6	10	25	37	30	19	-
1987	126	-	2	-	6	11	24	36	32	15	-
1986	121	-	1	1	4	16	34	24	19	22	-
1985	91	-	-	1	5	15	15	27	13	15	-
1984	96	-	-	2	8	10	25	23	19	8	-
1983	102	1	-	1	9	18	17	26	20	10	-
1982	83	-	-	4	6	10	22	21	11	9	-
1981	74	-	-	1	1	6	25	23	12	6	-
1980	78	-	2	3	2	12	15	19	15	10	-
1979	65	-	-	-	5	12	18	16	10	4	-
Parkinson's disease (332)											
Black, female											
1998	203	-	-	3	8	9	22	46	45	70	-
1997	154	-	-	-	5	5	17	31	34	62	-
1996	161	-	-	1	5	9	18	35	34	59	-
1995	147	-	1	1	3	11	24	26	39	40	-
1994	132	1	-	1	1	5	20	25	35	43	-
1993	106	2	2	2	2	5	17	19	26	31	-
1992	103	-	1	1	2	7	11	22	31	28	-
1991	113	-	-	2	3	10	13	28	25	32	-
1990	99	1	1	1	2	7	12	29	23	23	-
1989	107	1	2	-	3	5	18	23	28	26	-
1988	87	-	1	1	3	8	15	20	17	22	-
1987	74	-	1	1	6	5	10	19	16	16	-
1986	63	-	-	-	-	4	9	21	16	13	-
1985	67	-	-	-	1	10	15	12	16	12	-
1984	70	-	-	3	5	7	16	18	12	9	-
1983	74	-	2	1	3	8	16	17	20	7	-
1982	58	1	-	-	3	9	12	11	10	12	-
1981	51	-	2	4	1	7	10	8	10	9	-
1980	45	-	-	2	4	3	7	15	9	5	-
1979	54	-	-	2	4	9	17	11	5	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
All races, both sexes											
1998	22,725	16	23	65	194	553	1,354	3,204	5,163	12,142	1
1997	22,475	13	32	79	187	549	1,445	3,282	5,294	11,584	1
1996	21,397	10	19	74	212	600	1,449	3,231	5,088	10,709	-
1995	20,606	10	27	79	251	581	1,502	3,135	5,047	9,965	1
1994	18,584	10	26	75	248	622	1,441	2,897	4,388	8,869	1
1993	16,754	10	20	70	227	516	1,291	2,714	4,133	7,768	-
1992	14,435	9	17	47	224	550	1,229	2,368	3,509	6,480	-
1991	14,112	10	18	73	234	564	1,253	2,378	3,546	6,026	1
1990	13,744	9	28	74	236	577	1,270	2,411	3,457	5,675	1
1989	13,110	3	17	90	264	592	1,307	2,538	3,239	5,058	-
1988	12,281	10	22	112	290	650	1,333	2,404	3,091	4,363	1
1987	11,311	8	18	100	305	644	1,287	2,367	2,757	3,821	1
1986	9,426	10	17	106	304	610	1,249	2,002	2,278	2,843	2
1985	8,258	10	25	113	286	623	1,146	1,861	1,892	2,297	-
1984	6,250	9	21	100	264	575	965	1,437	1,318	1,551	-
1983	4,434	5	25	100	260	528	755	983	821	951	1
1982	2,627	11	22	90	239	347	476	564	436	437	1
1981	1,757	11	24	87	159	269	339	323	255	287	-
1980	1,314	11	17	70	173	210	287	216	154	170	-
1979	857	8	18	58	117	184	145	141	89	94	-
Alzheimer's disease (331.0)											
All races, male											
1998	7,054	6	11	29	91	257	576	1,326	1,843	2,911	-
1997	7,038	5	16	41	86	252	633	1,337	1,901	2,762	1
1996	6,971	5	6	30	92	281	644	1,338	1,911	2,661	-
1995	6,999	3	13	37	106	297	677	1,375	1,919	2,569	1
1994	6,377	5	11	41	113	301	665	1,266	1,650	2,320	-
1993	5,821	2	6	35	100	250	639	1,194	1,600	1,991	-
1992	5,097	4	6	24	106	264	568	1,064	1,378	1,682	-
1991	5,070	3	6	35	110	272	598	1,046	1,370	1,626	1
1990	4,992	5	16	34	111	278	637	1,023	1,348	1,536	-
1989	4,970	3	7	42	133	301	620	1,188	1,295	1,379	-
1988	4,732	7	12	53	139	308	630	1,104	1,241	1,235	-
1987	4,545	6	13	46	151	313	668	1,077	1,164	1,106	-
1986	3,862	4	8	47	151	286	654	928	937	845	-
1985	3,523	3	13	45	137	325	612	883	790	713	-
1984	2,803	5	11	44	124	298	511	701	586	519	-
1983	2,021	2	12	48	114	275	378	495	380	313	1
1982	1,208	6	8	39	125	174	242	292	185	135	1
1981	831	4	15	45	80	151	171	167	107	89	-
1980	653	6	4	34	82	120	172	107	63	64	-
1979	429	2	11	27	58	96	78	68	39	48	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
All races, female											
1998	15,671	10	12	36	103	296	778	1,878	3,320	9,231	1
1997	15,437	8	16	38	101	297	812	1,945	3,393	8,822	-
1996	14,426	5	13	44	120	319	805	1,893	3,177	8,048	-
1995	13,607	7	14	42	145	284	825	1,760	3,128	7,396	-
1994	12,207	5	15	34	135	321	776	1,631	2,738	6,549	1
1993	10,933	8	14	35	127	266	652	1,520	2,533	5,777	-
1992	9,338	5	11	23	118	286	661	1,304	2,131	4,798	-
1991	9,042	7	12	38	124	292	655	1,332	2,176	4,400	-
1990	8,752	4	12	40	125	299	633	1,388	2,109	4,139	1
1989	8,140	-	10	48	131	291	687	1,350	1,944	3,679	-
1988	7,549	3	10	59	151	342	703	1,300	1,850	3,128	1
1987	6,766	2	5	54	154	331	619	1,290	1,593	2,715	1
1986	5,564	6	9	59	153	324	595	1,074	1,341	1,998	2
1985	4,735	7	12	68	149	298	534	978	1,102	1,584	-
1984	3,447	4	10	56	140	277	454	736	732	1,032	-
1983	2,413	3	13	52	146	253	377	488	441	638	-
1982	1,419	5	14	51	114	173	234	272	251	302	-
1981	926	7	9	42	79	118	168	156	148	198	-
1980	661	5	13	36	91	90	115	109	91	106	-
1979	428	6	7	31	59	88	67	73	50	46	-
Alzheimer's disease (331.0)											
White, both sexes											
1998	21,366	13	23	62	176	513	1,240	2,982	4,855	11,491	1
1997	21,192	13	30	72	172	506	1,326	3,064	5,013	10,988	-
1996	20,198	10	19	67	194	554	1,326	3,047	4,815	10,163	-
1995	19,405	9	23	73	228	540	1,392	2,945	4,731	9,455	1
1994	17,550	9	20	67	222	585	1,339	2,691	4,142	8,470	1
1993	15,826	9	17	64	202	470	1,215	2,530	3,918	7,397	-
1992	13,634	9	16	43	208	515	1,132	2,216	3,317	6,176	-
1991	13,401	9	18	67	210	519	1,184	2,236	3,367	5,782	1
1990	13,020	9	26	67	219	537	1,194	2,268	3,270	5,423	1
1989	12,443	3	16	82	235	552	1,229	2,385	3,078	4,861	-
1988	11,730	10	21	98	275	614	1,246	2,283	2,982	4,196	1
1987	10,801	8	15	91	283	594	1,230	2,233	2,661	3,682	1
1986	9,037	9	16	98	278	579	1,179	1,924	2,190	2,757	2
1985	7,917	9	23	104	263	587	1,077	1,805	1,826	2,218	-
1984	6,032	8	20	97	248	550	928	1,388	1,273	1,510	-
1983	4,256	5	24	94	246	507	725	945	792	912	1
1982	2,533	11	20	87	236	328	465	541	417	424	-
1981	1,691	10	23	79	156	261	326	313	244	276	-
1980	1,258	10	14	66	166	202	274	209	149	162	-
1979	818	7	17	53	109	174	140	137	88	90	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
White, male											
1998	6,643	5	11	28	85	240	532	1,235	1,754	2,749	-
1997	6,579	5	16	37	81	233	583	1,242	1,794	2,584	-
1996	6,564	5	6	29	82	258	596	1,271	1,814	2,501	-
1995	6,579	2	13	34	99	279	622	1,297	1,797	2,433	1
1994	6,000	5	8	36	100	284	621	1,182	1,552	2,210	-
1993	5,466	2	5	32	89	229	599	1,120	1,511	1,875	-
1992	4,795	4	5	23	100	249	524	994	1,304	1,591	-
1991	4,787	2	6	33	100	255	561	983	1,294	1,549	1
1990	4,675	5	15	30	103	256	594	952	1,253	1,463	-
1989	4,692	3	7	40	119	279	580	1,108	1,231	1,323	-
1988	4,515	7	12	49	135	288	595	1,049	1,191	1,186	-
1987	4,329	6	11	42	137	292	637	1,026	1,119	1,058	-
1986	3,708	4	8	44	137	270	624	894	899	826	-
1985	3,383	3	11	43	122	302	581	860	765	694	-
1984	2,712	4	10	43	118	289	491	681	568	504	-
1983	1,935	2	12	45	109	262	364	471	365	301	1
1982	1,166	6	6	37	123	164	237	283	178	131	-
1981	797	4	14	40	78	146	163	162	102	86	-
1980	619	5	3	32	78	115	164	101	61	59	-
1979	408	2	10	25	54	92	74	67	38	44	-
Alzheimer's disease (331.0)											
White, female											
1998	14,723	8	12	34	91	273	708	1,747	3,101	8,742	1
1997	14,613	8	14	35	91	273	743	1,822	3,219	8,404	-
1996	13,634	5	13	38	112	296	730	1,776	3,001	7,662	-
1995	12,826	7	10	39	129	261	770	1,648	2,934	7,022	-
1994	11,550	4	12	31	122	301	718	1,509	2,590	6,260	1
1993	10,360	7	12	32	113	241	616	1,410	2,407	5,522	-
1992	8,839	5	11	20	108	266	608	1,222	2,013	4,585	-
1991	8,614	7	12	34	110	264	623	1,253	2,073	4,233	-
1990	8,345	4	11	37	116	281	600	1,316	2,017	3,960	1
1989	7,751	-	9	42	116	273	649	1,277	1,847	3,538	-
1988	7,215	3	9	49	140	326	651	1,234	1,791	3,010	1
1987	6,472	2	4	49	146	302	593	1,207	1,542	2,624	1
1986	5,329	5	8	54	141	309	555	1,030	1,291	1,931	2
1985	4,534	6	12	61	141	285	496	945	1,061	1,524	-
1984	3,320	4	10	54	130	261	437	707	705	1,006	-
1983	2,321	3	12	49	137	245	361	474	427	611	-
1982	1,367	5	14	50	113	164	228	258	239	293	-
1981	894	6	9	39	78	115	163	151	142	190	-
1980	639	5	11	34	88	87	110	108	88	103	-
1979	410	5	7	28	55	82	66	70	50	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
All other, both sexes											
1998	1,359	3	-	3	18	40	114	222	308	651	-
1997	1,283	-	2	7	15	43	119	218	281	596	1
1996	1,199	-	-	7	18	46	123	184	273	546	-
1995	1,201	1	4	6	23	41	110	190	316	510	-
1994	1,034	1	6	8	26	37	102	206	246	399	-
1993	928	1	3	6	25	46	76	184	215	371	-
1992	801	-	1	4	16	35	97	152	192	304	-
1991	711	1	-	6	24	45	69	142	179	244	-
1990	724	-	2	7	17	40	76	143	187	252	-
1989	667	-	1	8	29	40	78	153	161	197	-
1988	551	-	1	14	15	36	87	121	109	167	-
1987	510	-	3	9	22	50	57	134	96	139	-
1986	389	1	1	8	26	31	70	78	88	86	-
1985	341	1	2	9	23	36	69	56	66	79	-
1984	218	1	1	3	16	25	37	49	45	41	-
1983	178	-	1	6	14	21	30	38	29	39	-
1982	94	-	2	3	3	19	11	23	19	13	1
1981	66	1	1	8	3	8	13	10	11	11	-
1980	56	1	3	4	7	8	13	7	5	8	-
1979	39	1	1	5	8	10	5	4	1	4	-
Alzheimer's disease (331.0)											
All other, male											
1998	411	1	-	1	6	17	44	91	89	162	-
1997	459	-	-	4	5	19	50	95	107	178	1
1996	407	-	-	1	10	23	48	67	97	160	-
1995	420	1	-	3	7	18	55	78	122	136	-
1994	377	-	3	5	13	17	44	84	98	110	-
1993	355	-	1	3	11	21	40	74	89	116	-
1992	302	-	1	1	6	15	44	70	74	91	-
1991	283	1	-	2	10	17	37	63	76	77	-
1990	317	-	1	4	8	22	43	71	95	73	-
1989	278	-	-	2	14	22	40	80	64	56	-
1988	217	-	-	4	4	20	35	55	50	49	-
1987	216	-	2	4	14	21	31	51	45	48	-
1986	154	-	-	3	14	16	30	34	38	19	-
1985	140	-	2	2	15	23	31	23	25	19	-
1984	91	1	1	1	6	9	20	20	18	15	-
1983	86	-	-	3	5	13	14	24	15	12	-
1982	42	-	2	2	2	10	5	9	7	4	1
1981	34	-	1	5	2	5	8	5	5	3	-
1980	34	1	1	2	4	5	8	6	2	5	-
1979	21	-	1	2	4	4	4	1	1	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
All other, female											
1998	948	2	-	2	12	23	70	131	219	489	-
1997	824	-	2	3	10	24	69	123	174	418	-
1996	792	-	-	6	8	23	75	117	176	386	-
1995	781	-	4	3	16	23	55	112	194	374	-
1994	657	1	3	3	13	20	58	122	148	289	-
1993	573	1	2	3	14	25	36	110	126	255	-
1992	499	-	-	3	10	20	53	82	118	213	-
1991	428	-	-	4	14	28	32	79	103	167	-
1990	407	-	1	3	9	18	33	72	92	179	-
1989	389	-	1	6	15	18	38	73	97	141	-
1988	334	-	1	10	11	16	52	66	59	118	-
1987	294	-	1	5	8	29	26	83	51	91	-
1986	235	1	1	5	12	15	40	44	50	67	-
1985	201	1	-	7	8	13	38	33	41	60	-
1984	127	-	-	2	10	16	17	29	27	26	-
1983	92	-	1	3	9	8	16	14	14	27	-
1982	52	-	-	1	1	9	6	14	12	9	-
1981	32	1	-	3	1	3	5	5	6	8	-
1980	22	-	2	2	3	3	5	1	3	3	-
1979	18	1	-	3	4	6	1	3	-	-	-
Alzheimer's disease (331.0)											
Black, both sexes											
1998	1,209	3	-	3	15	31	99	209	283	566	-
1997	1,148	-	2	5	14	39	105	201	255	525	1
1996	1,083	-	-	7	18	45	110	166	246	490	-
1995	1,086	1	3	5	19	36	102	170	286	464	-
1994	926	1	4	6	25	33	87	188	224	355	-
1993	827	1	2	5	24	38	70	164	201	321	-
1992	730	-	1	4	14	32	90	142	172	275	-
1991	660	1	-	6	22	41	67	132	166	224	-
1990	662	-	1	7	15	35	72	129	174	229	-
1989	618	-	1	5	27	36	74	147	149	179	-
1988	499	-	1	14	14	31	81	112	100	146	-
1987	463	-	1	9	15	46	55	127	88	122	-
1986	347	1	-	7	21	28	66	73	77	74	-
1985	310	1	2	8	22	32	63	50	59	73	-
1984	197	1	1	3	15	23	32	45	40	37	-
1983	156	-	1	6	13	19	28	31	26	32	-
1982	87	-	2	3	3	17	10	22	17	13	-
1981	61	1	1	7	3	8	12	8	10	11	-
1980	53	1	3	3	7	8	11	7	5	8	-
1979	33	1	1	4	5	8	5	4	1	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alzheimer's disease (331.0)											
Black, male											
1998	352	1	-	1	5	15	37	86	79	128	-
1997	404	-	-	3	5	19	43	89	98	146	1
1996	350	-	-	1	10	22	43	56	90	128	-
1995	374	1	-	3	7	15	49	71	108	120	-
1994	329	-	2	4	12	16	37	75	86	94	-
1993	308	-	1	3	11	16	38	64	83	92	-
1992	267	-	1	1	6	14	40	66	62	77	-
1991	265	1	-	2	9	17	37	61	69	69	-
1990	281	-	-	4	7	19	39	62	87	63	-
1989	259	-	-	1	14	19	38	77	60	50	-
1988	193	-	-	4	4	17	31	52	43	42	-
1987	193	-	1	4	11	19	30	47	42	39	-
1986	130	-	-	3	10	16	27	31	30	13	-
1985	121	-	2	1	14	19	26	21	21	17	-
1984	81	1	1	1	6	9	16	19	14	14	-
1983	74	-	-	3	5	11	12	19	15	9	-
1982	37	-	2	2	2	8	5	8	6	4	-
1981	30	-	1	5	2	5	7	3	4	3	-
1980	31	1	1	1	4	5	6	6	2	5	-
1979	18	-	1	1	4	2	4	1	1	4	-
Alzheimer's disease (331.0)											
Black, female											
1998	857	2	-	2	10	16	62	123	204	438	-
1997	744	-	2	2	9	20	62	112	157	379	-
1996	733	-	-	6	8	23	67	110	156	362	-
1995	712	-	3	2	12	21	53	99	178	344	-
1994	597	1	2	2	13	17	50	113	138	261	-
1993	519	1	1	2	13	22	32	100	118	229	-
1992	463	-	-	3	8	18	50	76	110	198	-
1991	395	-	-	4	13	24	30	71	97	155	-
1990	381	-	1	3	8	16	33	67	87	166	-
1989	359	-	1	4	13	17	36	70	89	129	-
1988	306	-	1	10	10	14	50	60	57	104	-
1987	270	-	-	5	4	27	25	80	46	83	-
1986	217	1	-	4	11	12	39	42	47	61	-
1985	189	1	-	7	8	13	37	29	38	56	-
1984	116	-	-	2	9	14	16	26	26	23	-
1983	82	-	1	3	8	8	16	12	11	23	-
1982	50	-	-	1	1	9	5	14	11	9	-
1981	31	1	-	2	1	3	5	5	6	8	-
1980	22	-	2	2	3	3	5	1	3	3	-
1979	15	1	-	3	1	6	1	3	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Major cardiovascular diseases (390-434,436-448)											
All races, both sexes											
1998	940,159	17,524	25,058	32,876	45,917	67,771	102,198	137,688	162,248	324,635	65
1997	943,749	17,672	25,098	32,706	46,903	70,736	104,763	138,908	161,710	321,143	75
1996	949,769	18,189	24,402	32,219	48,770	73,418	107,911	139,268	163,776	317,870	94
1995	951,015	17,370	24,278	32,461	49,560	76,402	110,844	138,313	163,547	313,533	104
1994	940,326	16,764	23,716	32,556	50,356	77,967	111,757	137,281	161,723	303,890	82
1993	947,760	16,146	23,126	33,133	52,611	81,673	113,192	140,963	162,043	300,832	97
1992	913,588	15,711	21,919	32,761	53,466	80,836	111,977	137,519	155,677	280,266	106
1991	916,001	15,104	21,364	33,416	55,399	84,030	112,272	139,633	155,503	276,116	124
1990	915,671	14,787	21,454	34,384	57,423	86,352	112,354	141,378	155,212	270,045	123
1989	931,516	14,630	22,200	36,032	60,123	89,052	114,375	144,617	157,210	270,837	116
1988	969,028	14,812	22,822	38,790	64,285	93,590	121,695	150,063	161,876	277,831	122
1987	963,245	14,656	23,942	39,928	65,987	94,266	123,200	148,872	158,448	270,492	130
1986	967,866	14,471	24,372	41,852	68,753	95,178	125,388	150,175	157,591	266,161	419
1985	977,478	14,895	25,648	44,434	70,380	96,432	127,790	151,344	158,342	265,060	359
1984	973,011	15,144	26,364	45,573	72,080	96,995	128,446	150,436	157,132	258,176	147
1983	980,736	15,353	27,881	47,569	72,901	99,116	130,184	151,414	157,538	256,596	163
1982	967,546	15,866	28,651	48,282	73,000	100,173	129,275	149,367	154,549	246,519	221
1981	972,675	16,664	30,295	50,426	73,658	101,463	129,713	147,084	155,832	245,465	174
1980	988,196	16,873	31,447	52,135	74,100	104,196	131,539	151,674	158,899	245,214	174
1979	958,001	17,570	32,251	52,035	74,360	101,794	128,465	147,101	153,320	229,412	173
Major cardiovascular diseases (390-434,436-448)											
All races, male											
1998	441,133	12,445	17,595	22,280	30,074	41,940	58,608	72,016	72,907	97,319	41
1997	445,592	12,615	17,675	22,277	30,786	44,081	60,399	72,481	72,525	96,810	54
1996	448,717	12,880	17,079	21,956	32,126	45,838	62,230	72,905	72,606	95,255	60
1995	450,389	12,513	17,188	22,143	32,557	47,534	64,337	72,057	72,410	93,052	58
1994	447,097	12,000	16,822	22,277	33,264	48,884	64,800	71,364	71,136	90,453	49
1993	452,290	11,559	16,344	22,686	34,758	51,377	65,344	73,333	71,258	89,552	54
1992	439,184	11,363	15,407	22,418	35,362	50,603	64,804	71,305	68,033	84,041	65
1991	441,560	10,877	15,076	22,986	36,713	52,703	65,254	72,329	67,584	82,422	84
1990	442,609	10,597	15,168	23,717	37,918	53,958	65,079	72,786	67,566	80,703	71
1989	450,880	10,649	15,695	24,934	39,850	55,949	66,043	74,009	67,706	80,801	66
1988	470,728	10,627	16,090	26,721	42,452	58,451	70,627	76,786	69,538	83,568	68
1987	469,929	10,511	17,036	27,641	43,565	58,774	71,103	76,034	67,812	81,452	78
1986	475,446	10,338	17,357	29,283	45,601	59,656	72,343	76,474	67,546	80,611	145
1985	484,897	10,850	18,438	31,087	47,120	60,679	74,144	77,144	68,133	81,662	130
1984	484,863	11,056	18,938	31,788	48,274	61,069	73,815	76,348	67,197	80,973	81
1983	492,173	11,145	20,029	33,334	48,682	62,710	75,445	77,294	67,082	81,355	81
1982	489,049	11,548	20,748	33,953	49,013	63,596	74,651	75,938	65,399	79,372	121
1981	494,051	12,094	21,884	35,565	49,338	64,107	75,009	75,082	66,408	79,727	97
1980	503,599	12,184	22,888	36,880	49,908	65,806	75,935	76,421	67,417	81,260	102
1979	494,112	12,866	23,492	37,037	50,468	64,833	74,345	74,198	65,318	76,911	110

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Major cardiovascular diseases (390-434,436-448)											
All races, female											
1998	499,026	5,079	7,463	10,596	15,843	25,831	43,590	65,672	89,341	227,316	24
1997	498,157	5,057	7,423	10,429	16,117	26,655	44,364	66,427	89,185	224,333	21
1996	501,052	5,309	7,323	10,263	16,644	27,580	45,681	66,363	91,170	222,615	34
1995	500,626	4,857	7,090	10,318	17,003	28,868	46,507	66,256	91,137	220,481	46
1994	493,229	4,764	6,894	10,279	17,092	29,083	46,957	65,917	90,587	213,437	33
1993	495,470	4,587	6,782	10,447	17,853	30,296	47,848	67,630	90,785	211,280	43
1992	474,404	4,348	6,512	10,343	18,104	30,233	47,173	66,214	87,644	196,225	41
1991	474,441	4,227	6,288	10,430	18,686	31,327	47,018	67,304	87,919	193,694	40
1990	473,062	4,190	6,286	10,667	19,505	32,394	47,275	68,592	87,646	189,342	52
1989	480,636	3,981	6,505	11,098	20,273	33,103	48,332	70,608	89,504	190,036	50
1988	498,300	4,185	6,732	12,069	21,833	35,139	51,068	73,277	92,338	194,263	54
1987	493,316	4,145	6,906	12,287	22,422	35,492	52,097	72,838	90,636	189,040	52
1986	492,420	4,133	7,015	12,569	23,152	35,522	53,045	73,701	90,045	185,550	274
1985	492,581	4,045	7,210	13,347	23,260	35,753	53,646	74,200	90,209	183,398	229
1984	488,148	4,088	7,426	13,785	23,806	35,926	54,631	74,088	89,935	177,203	66
1983	488,563	4,208	7,852	14,235	24,219	36,406	54,739	74,120	90,456	175,241	82
1982	478,497	4,318	7,903	14,329	23,987	36,577	54,624	73,429	89,150	167,147	100
1981	478,624	4,570	8,411	14,861	24,320	37,356	54,704	72,002	89,424	165,738	77
1980	484,597	4,689	8,559	15,255	24,192	38,390	55,604	75,253	91,482	163,954	72
1979	463,889	4,704	8,759	14,998	23,892	36,961	54,120	72,903	88,002	152,501	63
Major cardiovascular diseases (390-434,436-448)											
White, both sexes											
1998	821,466	12,572	19,095	25,625	36,573	56,258	87,779	121,559	146,848	298,456	35
1997	827,096	12,824	19,159	25,416	37,597	59,241	90,088	123,478	146,755	295,822	51
1996	833,005	13,379	18,658	25,217	39,200	61,649	93,112	123,656	148,508	293,065	68
1995	834,009	12,762	18,659	25,375	40,168	64,043	95,710	122,962	148,046	289,193	74
1994	826,140	12,392	18,158	25,563	41,268	65,639	96,935	122,293	146,399	280,881	53
1993	832,732	12,075	17,700	25,882	43,010	68,939	98,323	125,705	146,755	277,821	51
1992	803,705	11,774	16,846	25,730	43,937	68,459	97,568	122,695	141,391	259,179	68
1991	805,982	11,317	16,395	26,517	45,690	71,171	97,876	124,440	141,019	255,553	90
1990	807,379	11,127	16,513	27,437	47,617	73,304	98,475	126,146	141,277	250,113	68
1989	820,516	10,899	16,954	28,799	49,571	75,680	100,229	128,623	143,164	251,003	76
1988	856,018	11,161	17,660	31,367	53,244	80,032	107,086	133,699	147,280	258,196	78
1987	853,140	11,144	18,718	32,639	55,218	81,035	108,328	133,145	144,590	251,686	83
1986	858,462	11,091	19,074	34,387	57,831	82,219	110,515	134,400	144,002	247,905	351
1985	868,777	11,422	20,390	36,589	59,465	83,471	112,404	135,894	144,750	247,775	304
1984	867,015	11,684	21,104	37,704	61,243	84,376	113,090	135,817	143,616	242,014	93
1983	874,744	11,927	22,424	39,431	61,944	86,474	114,636	136,723	144,352	240,738	100
1982	865,464	12,325	23,264	40,319	62,437	87,561	114,626	135,232	142,177	231,548	144
1981	870,013	13,019	24,629	42,139	63,155	88,726	115,025	133,338	143,477	230,445	105
1980	883,219	13,206	25,606	43,818	63,576	90,669	116,738	136,734	146,867	230,139	105
1979	858,578	13,897	26,370	43,779	64,281	88,636	114,244	132,697	142,821	216,028	113

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Major cardiovascular diseases (390-434,436-448)											
White, male											
1998	384,394	9,346	13,840	17,858	24,581	35,636	51,396	64,376	66,681	89,208	21
1997	389,572	9,509	13,931	17,808	25,291	37,788	53,011	65,251	66,474	88,968	37
1996	392,777	9,839	13,511	17,712	26,508	39,411	54,720	65,561	66,510	87,476	43
1995	393,746	9,559	13,602	17,796	27,129	40,754	56,592	64,824	66,112	85,423	42
1994	392,014	9,213	13,323	18,095	27,967	42,208	57,185	64,463	64,832	83,216	31
1993	396,730	9,007	12,935	18,279	29,193	44,381	57,773	66,370	64,990	82,275	28
1992	385,692	8,869	12,311	18,246	29,914	43,817	57,367	64,357	62,122	77,291	41
1991	387,689	8,459	11,995	18,807	31,108	45,682	57,919	65,306	61,474	75,730	59
1990	389,712	8,284	12,104	19,472	32,431	46,905	57,977	65,725	61,755	74,180	36
1989	396,606	8,240	12,386	20,604	33,790	48,593	58,934	66,698	61,933	74,360	42
1988	415,103	8,314	12,945	22,260	36,125	51,058	63,161	69,190	63,375	77,128	46
1987	415,268	8,311	13,776	23,285	37,378	51,481	63,556	68,547	61,957	75,247	50
1986	421,089	8,244	14,084	24,736	39,305	52,562	64,867	69,060	61,784	74,477	113
1985	430,629	8,640	15,179	26,489	40,779	53,564	66,294	69,884	62,383	75,811	100
1984	431,771	8,854	15,696	27,114	41,984	54,178	65,998	69,549	61,491	75,367	51
1983	438,960	8,988	16,674	28,488	42,398	55,851	67,480	70,365	61,536	75,790	51
1982	437,161	9,320	17,496	29,154	42,930	56,602	67,022	69,161	60,189	74,023	74
1981	441,923	9,823	18,416	30,538	43,281	57,216	67,465	68,578	61,084	74,362	58
1980	449,980	9,923	19,292	31,847	43,801	58,493	68,334	69,349	62,177	75,651	61
1979	442,730	10,595	19,848	31,909	44,647	57,591	67,047	67,358	60,667	71,947	73
Major cardiovascular diseases (390-434,436-448)											
White, female											
1998	437,072	3,226	5,255	7,767	11,992	20,622	36,383	57,183	80,167	209,248	14
1997	437,524	3,315	5,228	7,608	12,306	21,453	37,077	58,227	80,281	206,854	14
1996	440,228	3,540	5,147	7,505	12,692	22,238	38,392	58,095	81,998	205,589	25
1995	440,263	3,203	5,057	7,579	13,039	23,289	39,118	58,138	81,934	203,770	32
1994	434,126	3,179	4,835	7,468	13,301	23,431	39,750	57,830	81,567	197,665	22
1993	436,002	3,068	4,765	7,603	13,817	24,558	40,550	59,335	81,765	195,546	23
1992	418,013	2,905	4,535	7,484	14,023	24,642	40,201	58,338	79,269	181,888	27
1991	418,293	2,858	4,400	7,710	14,582	25,489	39,957	59,134	79,545	179,823	31
1990	417,667	2,843	4,409	7,965	15,186	26,399	40,498	60,421	79,522	175,933	32
1989	423,910	2,659	4,568	8,195	15,781	27,087	41,295	61,925	81,231	176,643	34
1988	440,915	2,847	4,715	9,107	17,119	28,974	43,925	64,509	83,905	181,068	32
1987	437,872	2,833	4,942	9,354	17,840	29,554	44,772	64,598	82,633	176,439	33
1986	437,373	2,847	4,990	9,651	18,526	29,657	45,648	65,340	82,218	173,428	238
1985	438,148	2,782	5,211	10,100	18,686	29,907	46,110	66,010	82,367	171,964	204
1984	435,244	2,830	5,408	10,590	19,259	30,198	47,092	66,268	82,125	166,647	42
1983	435,784	2,939	5,750	10,943	19,546	30,623	47,156	66,358	82,816	164,948	49
1982	428,303	3,005	5,768	11,165	19,507	30,959	47,604	66,071	81,988	157,525	70
1981	428,090	3,196	6,213	11,601	19,874	31,510	47,560	64,760	82,393	156,083	47
1980	433,239	3,283	6,314	11,971	19,775	32,176	48,404	67,385	84,690	154,488	44
1979	415,848	3,302	6,522	11,870	19,634	31,045	47,197	65,339	82,154	144,081	40

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Major cardiovascular diseases (390-434,436-448)											
All other, both sexes											
1998	118,693	4,952	5,963	7,251	9,344	11,513	14,419	16,129	15,400	26,179	30
1997	116,653	4,848	5,939	7,290	9,306	11,495	14,675	15,430	14,955	25,321	24
1996	116,764	4,810	5,744	7,002	9,570	11,769	14,799	15,612	15,268	24,805	26
1995	117,006	4,608	5,619	7,086	9,392	12,359	15,134	15,351	15,501	24,340	30
1994	114,186	4,372	5,558	6,993	9,088	12,328	14,822	14,988	15,324	23,009	29
1993	115,028	4,071	5,426	7,251	9,601	12,734	14,869	15,258	15,288	23,011	46
1992	109,883	3,937	5,073	7,031	9,529	12,377	14,409	14,824	14,286	21,087	38
1991	110,019	3,787	4,969	6,899	9,709	12,859	14,396	15,193	14,484	20,563	34
1990	108,292	3,660	4,941	6,947	9,806	13,048	13,879	15,232	13,935	19,932	55
1989	111,000	3,731	5,246	7,233	10,552	13,372	14,146	15,994	14,046	19,834	40
1988	113,010	3,651	5,162	7,423	11,041	13,558	14,609	16,364	14,596	19,635	44
1987	110,105	3,512	5,224	7,289	10,769	13,231	14,872	15,727	13,858	18,806	47
1986	109,404	3,380	5,298	7,465	10,922	12,959	14,873	15,775	13,589	18,256	68
1985	108,701	3,473	5,258	7,845	10,915	12,961	15,386	15,450	13,592	17,285	55
1984	105,996	3,460	5,260	7,869	10,837	12,619	15,356	14,619	13,516	16,162	54
1983	105,992	3,426	5,457	8,138	10,957	12,642	15,548	14,691	13,186	15,858	63
1982	102,082	3,541	5,387	7,963	10,563	12,612	14,649	14,135	12,372	14,971	77
1981	102,662	3,645	5,666	8,287	10,503	12,737	14,688	13,746	12,355	15,020	69
1980	104,977	3,667	5,841	8,317	10,524	13,527	14,801	14,940	12,032	15,075	69
1979	99,423	3,673	5,881	8,256	10,079	13,158	14,221	14,404	10,499	13,384	60
Major cardiovascular diseases (390-434,436-448)											
All other, male											
1998	56,739	3,099	3,755	4,422	5,493	6,304	7,212	7,640	6,226	8,111	20
1997	56,020	3,106	3,744	4,469	5,495	6,293	7,388	7,230	6,051	7,842	17
1996	55,940	3,041	3,568	4,244	5,618	6,427	7,510	7,344	6,096	7,779	17
1995	56,643	2,954	3,586	4,347	5,428	6,780	7,745	7,233	6,298	7,629	16
1994	55,083	2,787	3,499	4,182	5,297	6,676	7,615	6,901	6,304	7,237	18
1993	55,560	2,552	3,409	4,407	5,565	6,996	7,571	6,963	6,268	7,277	26
1992	53,492	2,494	3,096	4,172	5,448	6,786	7,437	6,948	5,911	6,750	24
1991	53,871	2,418	3,081	4,179	5,605	7,021	7,335	7,023	6,110	6,692	25
1990	52,897	2,313	3,064	4,245	5,487	7,053	7,102	7,061	5,811	6,523	35
1989	54,274	2,409	3,309	4,330	6,060	7,356	7,109	7,311	5,773	6,441	24
1988	55,625	2,313	3,145	4,461	6,327	7,393	7,466	7,596	6,163	6,440	22
1987	54,661	2,200	3,260	4,356	6,187	7,293	7,547	7,487	5,855	6,205	28
1986	54,357	2,094	3,273	4,547	6,296	7,094	7,476	7,414	5,762	6,134	32
1985	54,268	2,210	3,259	4,598	6,341	7,115	7,850	7,260	5,750	5,851	30
1984	53,092	2,202	3,242	4,674	6,290	6,891	7,817	6,799	5,706	5,606	30
1983	53,213	2,157	3,355	4,846	6,284	6,859	7,965	6,929	5,546	5,565	30
1982	51,888	2,228	3,252	4,799	6,083	6,994	7,629	6,777	5,210	5,349	47
1981	52,128	2,271	3,468	5,027	6,057	6,891	7,544	6,504	5,324	5,365	39
1980	53,619	2,261	3,596	5,033	6,107	7,313	7,601	7,072	5,240	5,609	41
1979	51,382	2,271	3,644	5,128	5,821	7,242	7,298	6,840	4,651	4,964	37

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Major cardiovascular diseases (390-434,436-448)											
All other, female											
1998	61,954	1,853	2,208	2,829	3,851	5,209	7,207	8,489	9,174	18,068	10
1997	60,633	1,742	2,195	2,821	3,811	5,202	7,287	8,200	8,904	17,479	7
1996	60,824	1,769	2,176	2,758	3,952	5,342	7,289	8,268	9,172	17,026	9
1995	60,363	1,654	2,033	2,739	3,964	5,579	7,389	8,118	9,203	16,711	14
1994	59,103	1,585	2,059	2,811	3,791	5,652	7,207	8,087	9,020	15,772	11
1993	59,468	1,519	2,017	2,844	4,036	5,738	7,298	8,295	9,020	15,734	20
1992	56,391	1,443	1,977	2,859	4,081	5,591	6,972	7,876	8,375	14,337	14
1991	56,148	1,369	1,888	2,720	4,104	5,838	7,061	8,170	8,374	13,871	9
1990	55,395	1,347	1,877	2,702	4,319	5,995	6,777	8,171	8,124	13,409	20
1989	56,726	1,322	1,937	2,903	4,492	6,016	7,037	8,683	8,273	13,393	16
1988	57,385	1,338	2,017	2,962	4,714	6,165	7,143	8,768	8,433	13,195	22
1987	55,444	1,312	1,964	2,933	4,582	5,938	7,325	8,240	8,003	12,601	19
1986	55,047	1,286	2,025	2,918	4,626	5,865	7,397	8,361	7,827	12,122	36
1985	54,433	1,263	1,999	3,247	4,574	5,846	7,536	8,190	7,842	11,434	25
1984	52,904	1,258	2,018	3,195	4,547	5,728	7,539	7,820	7,810	10,556	24
1983	52,779	1,269	2,102	3,292	4,673	5,783	7,583	7,762	7,640	10,293	33
1982	50,194	1,313	2,135	3,164	4,480	5,618	7,020	7,358	7,162	9,622	30
1981	50,534	1,374	2,198	3,260	4,446	5,846	7,144	7,242	7,031	9,655	30
1980	51,358	1,406	2,245	3,284	4,417	6,214	7,200	7,868	6,792	9,466	28
1979	48,041	1,402	2,237	3,128	4,258	5,916	6,923	7,564	5,848	8,420	23
Major cardiovascular diseases (390-434,436-448)											
Black, both sexes											
1998	103,422	4,513	5,315	6,474	8,254	10,143	12,495	13,852	13,135	22,390	28
1997	101,956	4,388	5,332	6,446	8,258	10,064	12,847	13,313	12,803	21,837	22
1996	102,588	4,360	5,154	6,269	8,529	10,379	12,987	13,558	13,206	21,447	26
1995	103,492	4,210	5,047	6,350	8,386	11,016	13,317	13,498	13,479	21,234	28
1994	101,164	3,956	5,014	6,254	8,145	10,977	13,100	13,197	13,407	20,096	27
1993	102,707	3,723	4,936	6,604	8,653	11,424	13,333	13,516	13,444	20,228	40
1992	98,267	3,592	4,620	6,317	8,634	11,123	12,888	13,140	12,612	18,609	37
1991	99,026	3,489	4,549	6,285	8,795	11,643	13,006	13,564	12,919	18,166	31
1990	98,043	3,372	4,531	6,327	9,002	11,875	12,558	13,745	12,490	17,800	51
1989	100,876	3,447	4,790	6,618	9,712	12,236	12,928	14,517	12,607	17,681	38
1988	103,407	3,390	4,758	6,887	10,167	12,457	13,380	14,937	13,230	17,708	41
1987	100,896	3,265	4,804	6,697	9,993	12,270	13,617	14,366	12,631	16,967	44
1986	100,875	3,141	4,892	6,988	10,180	12,077	13,725	14,484	12,378	16,609	59
1985	100,268	3,243	4,933	7,319	10,174	12,060	14,224	14,147	12,427	15,677	45
1984	98,136	3,216	4,913	7,319	10,171	11,771	14,336	13,476	12,406	14,684	49
1983	98,496	3,217	5,124	7,632	10,297	11,836	14,515	13,622	12,145	14,494	42
1982	95,057	3,318	5,079	7,513	9,951	11,826	13,685	13,087	11,435	13,706	66
1981	96,269	3,426	5,367	7,827	9,994	12,044	13,843	12,808	11,503	13,832	60
1980	98,478	3,472	5,532	7,874	10,030	12,831	13,887	13,955	11,176	13,868	65
1979	93,349	3,498	5,601	7,845	9,572	12,450	13,406	13,472	9,722	12,293	55

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Major cardiovascular diseases (390-434,436-448)											
Black, male											
1998	48,372	2,775	3,303	3,882	4,776	5,528	6,159	6,364	5,130	6,455	18
1997	47,934	2,771	3,327	3,890	4,829	5,446	6,359	6,098	5,003	6,268	15
1996	48,175	2,709	3,172	3,751	4,950	5,623	6,493	6,242	5,115	6,277	17
1995	49,166	2,676	3,207	3,863	4,817	5,982	6,684	6,233	5,266	6,244	14
1994	47,813	2,482	3,118	3,700	4,712	5,883	6,585	5,966	5,334	5,907	17
1993	48,649	2,309	3,082	3,996	4,989	6,212	6,676	5,984	5,262	6,054	23
1992	46,954	2,258	2,777	3,700	4,892	6,033	6,566	6,006	5,001	5,670	23
1991	47,645	2,212	2,801	3,773	5,034	6,279	6,537	6,129	5,208	5,633	23
1990	47,194	2,115	2,785	3,854	5,003	6,362	6,340	6,237	5,028	5,618	32
1989	48,499	2,209	3,004	3,952	5,531	6,673	6,394	6,460	4,938	5,484	23
1988	50,080	2,134	2,880	4,107	5,776	6,702	6,742	6,799	5,364	5,576	22
1987	49,193	2,033	2,973	3,959	5,676	6,684	6,783	6,628	5,137	5,416	25
1986	49,371	1,924	3,002	4,225	5,818	6,539	6,792	6,625	5,061	5,441	29
1985	49,367	2,038	3,038	4,270	5,870	6,553	7,135	6,444	5,085	5,199	26
1984	48,482	2,037	3,011	4,335	5,869	6,375	7,179	6,076	5,047	4,984	28
1983	48,799	2,025	3,127	4,519	5,847	6,344	7,302	6,269	4,982	4,999	23
1982	47,623	2,081	3,040	4,501	5,683	6,496	6,995	6,057	4,694	4,797	39
1981	48,305	2,124	3,261	4,719	5,735	6,452	6,963	5,918	4,842	4,854	34
1980	49,603	2,126	3,380	4,713	5,777	6,860	6,976	6,452	4,768	5,024	39
1979	47,655	2,155	3,440	4,831	5,474	6,758	6,755	6,241	4,247	4,468	34
Major cardiovascular diseases (390-434,436-448)											
Black, female											
1998	55,050	1,738	2,012	2,592	3,478	4,615	6,336	7,488	8,005	15,935	10
1997	54,022	1,617	2,005	2,556	3,429	4,618	6,488	7,215	7,800	15,569	7
1996	54,413	1,651	1,982	2,518	3,579	4,756	6,494	7,316	8,091	15,170	9
1995	54,326	1,534	1,840	2,487	3,569	5,034	6,633	7,265	8,213	14,990	14
1994	53,351	1,474	1,896	2,554	3,433	5,094	6,515	7,231	8,073	14,189	10
1993	54,058	1,414	1,854	2,608	3,664	5,212	6,657	7,532	8,182	14,174	17
1992	51,313	1,334	1,843	2,617	3,742	5,090	6,322	7,134	7,611	12,939	14
1991	51,381	1,277	1,748	2,512	3,761	5,364	6,469	7,435	7,711	12,533	8
1990	50,849	1,257	1,746	2,473	3,999	5,513	6,218	7,508	7,462	12,182	19
1989	52,377	1,238	1,786	2,666	4,181	5,563	6,534	8,057	7,669	12,197	15
1988	53,327	1,256	1,878	2,780	4,391	5,755	6,638	8,138	7,866	12,132	19
1987	51,703	1,232	1,831	2,738	4,317	5,586	6,834	7,738	7,494	11,551	19
1986	51,504	1,217	1,890	2,763	4,362	5,538	6,933	7,859	7,317	11,168	30
1985	50,901	1,205	1,895	3,049	4,304	5,507	7,089	7,703	7,342	10,478	19
1984	49,654	1,179	1,902	2,984	4,302	5,396	7,157	7,400	7,359	9,700	21
1983	49,697	1,192	1,997	3,113	4,450	5,492	7,213	7,353	7,163	9,495	19
1982	47,434	1,237	2,039	3,012	4,268	5,330	6,690	7,030	6,741	8,909	27
1981	47,964	1,302	2,106	3,108	4,259	5,592	6,880	6,890	6,661	8,978	26
1980	48,875	1,346	2,152	3,161	4,253	5,971	6,911	7,503	6,408	8,844	26
1979	45,694	1,343	2,161	3,014	4,098	5,692	6,651	7,231	5,475	7,825	21

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
All races, both sexes											
1998	724,859	14,298	20,758	27,272	37,796	54,827	80,468	105,180	121,589	243,609	56
1997	726,974	14,461	20,816	27,256	38,702	57,176	82,240	106,290	121,226	239,981	68
1996	733,361	14,768	20,266	26,872	40,463	59,580	85,335	106,781	122,981	237,522	81
1995	737,563	14,218	20,280	27,041	41,199	62,231	87,827	106,766	123,348	235,254	93
1994	732,409	13,824	19,797	27,393	41,942	63,789	89,159	106,184	122,474	228,724	71
1993	743,460	13,292	19,387	27,903	44,140	67,386	90,717	110,027	123,993	227,632	80
1992	717,706	13,009	18,404	27,678	44,838	66,644	89,885	107,365	119,373	212,047	88
1991	720,862	12,468	17,906	28,310	46,675	69,101	90,300	109,493	119,389	208,984	107
1990	720,058	12,175	18,029	29,097	48,449	71,119	90,254	110,825	119,019	203,641	104
1989	733,867	12,222	18,700	30,634	50,717	73,686	92,101	113,653	120,665	203,863	92
1988	765,156	12,378	19,380	33,171	54,343	77,645	98,481	118,069	124,047	209,252	108
1987	760,353	12,344	20,364	34,199	55,818	78,513	99,559	117,003	121,098	202,816	107
1986	765,490	12,186	20,796	35,945	58,367	79,264	101,508	118,292	120,664	199,281	327
1985	771,169	12,576	21,969	38,146	59,919	80,350	103,383	118,683	120,733	196,933	278
1984	765,114	12,754	22,482	39,200	61,291	80,756	103,766	117,610	119,018	190,266	112
1983	770,432	12,868	23,991	40,981	61,964	82,607	105,412	117,875	118,748	188,303	127
1982	755,592	13,400	24,534	41,511	62,122	82,996	103,708	115,577	115,423	178,920	176
1981	753,884	14,130	25,909	43,279	62,336	83,786	103,165	112,893	115,248	175,804	146
1980	761,085	14,182	26,896	44,742	62,502	85,701	104,166	115,172	116,154	174,213	147
1979	733,235	14,792	27,556	44,413	62,623	83,097	100,922	111,107	111,177	160,621	143
Diseases of heart (390-398,402,404,410-429)											
All races, male											
1998	353,897	10,555	15,160	19,050	25,373	34,860	47,415	56,642	56,547	75,173	37
1997	356,598	10,752	15,202	19,105	26,053	36,558	48,699	57,116	55,992	74,036	49
1996	360,075	10,918	14,708	18,825	27,313	38,189	50,471	57,409	56,072	73,108	54
1995	362,714	10,676	14,920	19,034	27,707	39,689	52,263	57,071	56,149	71,584	51
1994	361,276	10,295	14,597	19,297	28,434	40,903	53,025	56,457	55,203	69,795	47
1993	367,479	9,965	14,227	19,702	29,856	43,255	53,546	58,409	55,847	69,386	46
1992	357,545	9,833	13,440	19,548	30,408	42,607	53,174	56,950	53,430	65,053	53
1991	359,814	9,410	13,121	20,035	31,692	44,230	53,564	57,850	52,993	63,982	74
1990	360,788	9,169	13,218	20,665	32,819	45,447	53,392	58,130	53,022	62,406	59
1989	368,202	9,304	13,747	21,872	34,363	47,288	54,331	59,224	53,110	62,375	57
1988	385,402	9,300	14,183	23,496	36,674	49,584	58,304	61,575	54,513	64,605	65
1987	385,217	9,263	15,032	24,344	37,706	49,991	58,656	60,965	53,033	62,828	66
1986	390,912	9,120	15,404	25,925	39,640	50,797	59,833	61,522	52,783	62,184	119
1985	398,208	9,590	16,371	27,410	41,028	51,618	61,091	61,742	53,377	62,796	107
1984	397,097	9,726	16,814	28,163	42,089	51,822	60,809	61,139	52,058	61,569	68
1983	402,727	9,805	17,884	29,556	42,409	53,458	62,255	61,302	51,725	61,628	71
1982	398,570	10,218	18,507	30,083	42,803	53,824	61,049	59,991	50,186	59,440	101
1981	400,535	10,769	19,475	31,540	42,827	54,114	61,004	58,822	50,527	59,076	79
1980	405,661	10,819	20,293	32,680	43,133	55,378	61,375	59,380	50,545	59,652	87
1979	396,372	11,419	20,876	32,708	43,621	54,258	59,741	57,398	48,531	55,706	91

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,
sex and year

	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
All races, female											
1998	370,962	3,743	5,598	8,222	12,423	19,967	33,053	48,538	65,042	168,436	19
1997	370,376	3,709	5,614	8,151	12,649	20,618	33,541	49,174	65,234	165,945	19
1996	373,286	3,850	5,558	8,047	13,150	21,391	34,864	49,372	66,909	164,414	27
1995	374,849	3,542	5,360	8,007	13,492	22,542	35,564	49,695	67,199	163,670	42
1994	371,133	3,529	5,200	8,096	13,508	22,886	36,134	49,727	67,271	158,929	24
1993	375,981	3,327	5,160	8,201	14,284	24,131	37,171	51,618	68,146	158,246	34
1992	360,161	3,176	4,964	8,130	14,430	24,037	36,711	50,415	65,943	146,994	35
1991	361,048	3,058	4,785	8,275	14,983	24,871	36,736	51,643	66,396	145,002	33
1990	359,270	3,006	4,811	8,432	15,630	25,672	36,862	52,695	65,997	141,235	45
1989	365,665	2,918	4,953	8,762	16,354	26,398	37,770	54,429	67,555	141,488	35
1988	379,754	3,078	5,197	9,675	17,669	28,061	40,177	56,494	69,534	144,647	43
1987	375,136	3,081	5,332	9,855	18,112	28,522	40,903	56,038	68,065	139,988	41
1986	374,578	3,066	5,392	10,020	18,727	28,467	41,675	56,770	67,881	137,097	208
1985	372,961	2,986	5,598	10,736	18,891	28,732	42,292	56,941	67,356	134,137	171
1984	368,017	3,028	5,668	11,037	19,202	28,934	42,957	56,471	66,960	128,697	44
1983	367,705	3,063	6,107	11,425	19,555	29,149	43,157	56,573	67,023	126,675	56
1982	357,022	3,182	6,027	11,428	19,319	29,172	42,659	55,586	65,237	119,480	75
1981	353,349	3,361	6,434	11,739	19,509	29,672	42,161	54,071	64,721	116,728	67
1980	355,424	3,363	6,603	12,062	19,369	30,323	42,791	55,792	65,609	114,561	60
1979	336,863	3,373	6,680	11,705	19,002	28,839	41,181	53,709	62,646	104,915	52
Diseases of heart (390-398,402,404,410-429)											
White, both sexes											
1998	635,549	10,568	16,165	21,608	30,487	45,974	69,475	93,224	110,409	224,310	31
1997	639,225	10,839	16,242	21,565	31,441	48,297	71,174	94,829	110,259	221,244	46
1996	645,514	11,188	15,798	21,388	32,914	50,587	74,106	95,198	111,839	219,280	60
1995	649,089	10,699	15,903	21,541	33,767	52,653	76,258	95,321	111,975	217,313	69
1994	646,056	10,495	15,525	21,856	34,784	54,177	77,859	95,049	111,307	211,682	46
1993	655,365	10,232	15,111	22,138	36,489	57,357	79,283	98,580	112,521	210,410	43
1992	633,487	10,013	14,412	22,089	37,262	56,941	78,837	96,154	108,671	196,198	54
1991	636,805	9,594	14,061	22,836	38,963	59,032	79,188	98,005	108,605	193,667	79
1990	637,364	9,446	14,172	23,571	40,630	60,874	79,629	99,299	108,657	188,775	60
1989	648,866	9,336	14,607	24,819	42,314	63,155	81,251	101,628	110,186	189,063	60
1988	678,526	9,535	15,324	27,239	45,579	67,020	87,264	105,644	113,155	194,628	71
1987	675,732	9,634	16,236	28,413	47,232	68,024	88,150	105,182	110,676	188,729	70
1986	681,363	9,544	16,632	29,986	49,625	69,052	90,097	106,386	110,509	185,596	281
1985	688,152	9,904	17,794	31,829	51,210	70,147	91,667	107,100	110,747	184,260	238
1984	684,511	10,083	18,328	32,901	52,675	70,958	92,186	106,734	109,088	178,344	73
1983	689,749	10,276	19,633	34,451	53,300	72,772	93,574	106,903	109,023	176,759	82
1982	678,837	10,650	20,297	35,160	53,807	73,324	92,683	105,291	106,504	168,152	111
1981	677,449	11,299	21,526	36,733	54,044	74,090	92,321	102,919	106,438	165,216	88
1980	683,347	11,376	22,321	38,156	54,319	75,490	93,293	104,480	107,642	163,624	90
1979	660,154	11,990	22,979	37,963	54,787	73,213	90,582	100,787	103,806	151,435	93

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
White, male											
1998	309,952	8,137	12,164	15,531	21,000	29,909	41,819	50,851	51,925	69,049	19
1997	313,316	8,345	12,243	15,548	21,652	31,595	43,010	51,683	51,463	68,126	34
1996	316,889	8,577	11,874	15,459	22,816	33,162	44,697	51,858	51,497	67,297	40
1995	318,751	8,371	12,060	15,579	23,306	34,305	46,243	51,571	51,495	65,830	38
1994	318,543	8,121	11,803	15,943	24,166	35,584	47,108	51,278	50,575	64,297	30
1993	323,802	7,968	11,456	16,107	25,337	37,646	47,598	53,133	51,065	63,824	24
1992	315,483	7,844	10,933	16,163	26,003	37,227	47,357	51,620	48,903	59,851	32
1991	317,642	7,520	10,657	16,638	27,123	38,633	47,810	52,464	48,401	58,919	52
1990	319,362	7,368	10,758	17,187	28,354	39,841	47,864	52,731	48,624	57,433	32
1989	325,397	7,374	11,069	18,284	29,432	41,406	48,771	53,603	48,682	57,455	35
1988	341,519	7,446	11,636	19,816	31,538	43,640	52,459	55,707	49,840	59,656	44
1987	342,063	7,504	12,358	20,830	32,662	44,092	52,756	55,249	48,550	58,061	42
1986	347,967	7,426	12,749	22,214	34,482	45,110	53,966	55,857	48,410	57,457	94
1985	355,374	7,815	13,692	23,617	35,851	45,931	54,996	56,211	48,964	58,354	84
1984	355,454	7,965	14,158	24,359	36,961	46,361	54,800	55,926	47,765	57,316	44
1983	360,951	8,095	15,105	25,559	37,318	48,002	56,085	56,050	47,503	57,483	47
1982	358,308	8,418	15,857	26,144	37,905	48,356	55,231	54,961	46,313	55,465	62
1981	360,386	8,940	16,681	27,428	37,953	48,752	55,285	54,024	46,630	55,186	47
1980	364,679	8,998	17,371	28,561	38,296	49,725	55,713	54,219	46,764	55,608	51
1979	357,378	9,591	17,951	28,593	39,013	48,710	54,317	52,395	45,188	52,189	62
Diseases of heart (390-398,402,404,410-429)											
White, female											
1998	325,597	2,431	4,001	6,077	9,487	16,065	27,656	42,373	58,484	155,261	12
1997	325,909	2,494	3,999	6,017	9,789	16,702	28,164	43,146	58,796	153,118	12
1996	328,625	2,611	3,924	5,929	10,098	17,425	29,409	43,340	60,342	151,983	20
1995	330,338	2,328	3,843	5,962	10,461	18,348	30,015	43,750	60,480	151,483	31
1994	327,513	2,374	3,722	5,913	10,618	18,593	30,751	43,771	60,732	147,385	16
1993	331,563	2,264	3,655	6,031	11,152	19,711	31,685	45,447	61,456	146,586	19
1992	318,004	2,169	3,479	5,926	11,259	19,714	31,480	44,534	59,768	136,347	22
1991	319,163	2,074	3,404	6,198	11,840	20,399	31,378	45,541	60,204	134,748	27
1990	318,002	2,078	3,414	6,384	12,276	21,033	31,765	46,568	60,033	131,342	28
1989	323,469	1,962	3,538	6,535	12,882	21,749	32,480	48,025	61,504	131,608	25
1988	337,007	2,089	3,688	7,423	14,041	23,380	34,805	49,937	63,315	134,972	27
1987	333,669	2,130	3,878	7,583	14,570	23,932	35,394	49,933	62,126	130,668	28
1986	333,396	2,118	3,883	7,772	15,143	23,942	36,131	50,529	62,099	128,139	187
1985	332,778	2,089	4,102	8,212	15,359	24,216	36,671	50,889	61,783	125,906	154
1984	329,057	2,118	4,170	8,542	15,714	24,597	37,386	50,808	61,323	121,028	29
1983	328,798	2,181	4,528	8,892	15,982	24,770	37,489	50,853	61,520	119,276	35
1982	320,529	2,232	4,440	9,016	15,902	24,968	37,452	50,330	60,191	112,687	49
1981	317,063	2,359	4,845	9,305	16,091	25,338	37,036	48,895	59,808	110,030	41
1980	318,668	2,378	4,950	9,595	16,023	25,765	37,580	50,261	60,878	108,016	39
1979	302,776	2,399	5,028	9,370	15,774	24,503	36,265	48,392	58,618	99,246	31

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
All other, both sexes											
1998	89,310	3,730	4,593	5,664	7,309	8,853	10,993	11,956	11,180	19,299	25
1997	87,749	3,622	4,574	5,691	7,261	8,879	11,066	11,461	10,967	18,737	22
1996	87,847	3,580	4,468	5,484	7,549	8,993	11,229	11,583	11,142	18,242	21
1995	88,474	3,519	4,377	5,500	7,432	9,578	11,569	11,445	11,373	17,941	24
1994	86,353	3,329	4,272	5,537	7,158	9,612	11,300	11,135	11,167	17,042	25
1993	88,095	3,060	4,276	5,765	7,651	10,029	11,434	11,447	11,472	17,222	37
1992	84,219	2,996	3,992	5,589	7,576	9,703	11,048	11,211	10,702	15,849	34
1991	84,057	2,874	3,845	5,474	7,712	10,069	11,112	11,488	10,784	15,317	28
1990	82,694	2,729	3,857	5,526	7,819	10,245	10,625	11,526	10,362	14,866	44
1989	85,001	2,886	4,093	5,815	8,403	10,531	10,850	12,025	10,479	14,800	32
1988	86,630	2,843	4,056	5,932	8,764	10,625	11,217	12,425	10,892	14,624	37
1987	84,621	2,710	4,128	5,786	8,586	10,489	11,409	11,821	10,422	14,087	37
1986	84,127	2,642	4,164	5,959	8,742	10,212	11,411	11,906	10,155	13,685	46
1985	83,017	2,672	4,175	6,317	8,709	10,203	11,716	11,583	9,986	12,673	40
1984	80,603	2,671	4,154	6,299	8,616	9,798	11,580	10,876	9,930	11,922	39
1983	80,683	2,592	4,358	6,530	8,664	9,835	11,838	10,972	9,725	11,544	45
1982	76,755	2,750	4,237	6,351	8,315	9,672	11,025	10,286	8,919	10,768	65
1981	76,435	2,831	4,383	6,546	8,292	9,696	10,844	9,974	8,810	10,588	58
1980	77,738	2,806	4,575	6,586	8,183	10,211	10,873	10,692	8,512	10,589	57
1979	73,081	2,802	4,577	6,450	7,836	9,884	10,340	10,320	7,371	9,186	50
Diseases of heart (390-398,402,404,410-429)											
All other, male											
1998	43,945	2,418	2,996	3,519	4,373	4,951	5,596	5,791	4,622	6,124	18
1997	43,282	2,407	2,959	3,557	4,401	4,963	5,689	5,433	4,529	5,910	15
1996	43,186	2,341	2,834	3,366	4,497	5,027	5,774	5,551	4,575	5,811	14
1995	43,963	2,305	2,860	3,455	4,401	5,384	6,020	5,500	4,654	5,754	13
1994	42,733	2,174	2,794	3,354	4,268	5,319	5,917	5,179	4,628	5,498	17
1993	43,677	1,997	2,771	3,595	4,519	5,609	5,948	5,276	4,782	5,562	22
1992	42,062	1,989	2,507	3,385	4,405	5,380	5,817	5,330	4,527	5,202	21
1991	42,172	1,890	2,464	3,397	4,569	5,597	5,754	5,386	4,592	5,063	22
1990	41,426	1,801	2,460	3,478	4,465	5,606	5,528	5,399	4,398	4,973	27
1989	42,805	1,930	2,678	3,588	4,931	5,882	5,560	5,621	4,428	4,920	22
1988	43,883	1,854	2,547	3,680	5,136	5,944	5,845	5,868	4,673	4,949	21
1987	43,154	1,759	2,674	3,514	5,044	5,899	5,900	5,716	4,483	4,767	24
1986	42,945	1,694	2,655	3,711	5,158	5,687	5,867	5,665	4,373	4,727	25
1985	42,834	1,775	2,679	3,793	5,177	5,687	6,095	5,531	4,413	4,442	23
1984	41,643	1,761	2,656	3,804	5,128	5,461	6,009	5,213	4,293	4,253	24
1983	41,776	1,710	2,779	3,997	5,091	5,456	6,170	5,252	4,222	4,145	24
1982	40,262	1,800	2,650	3,939	4,898	5,468	5,818	5,030	3,873	3,975	39
1981	40,149	1,829	2,794	4,112	4,874	5,362	5,719	4,798	3,897	3,890	32
1980	40,982	1,821	2,922	4,119	4,837	5,653	5,662	5,161	3,781	4,044	36
1979	38,994	1,828	2,925	4,115	4,608	5,548	5,424	5,003	3,343	3,517	29

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
All other, female											
1998	45,365	1,312	1,597	2,145	2,936	3,902	5,397	6,165	6,558	13,175	7
1997	44,467	1,215	1,615	2,134	2,860	3,916	5,377	6,028	6,438	12,827	7
1996	44,661	1,239	1,634	2,118	3,052	3,966	5,455	6,032	6,567	12,431	7
1995	44,511	1,214	1,517	2,045	3,031	4,194	5,549	5,945	6,719	12,187	11
1994	43,620	1,155	1,478	2,183	2,890	4,293	5,383	5,956	6,539	11,544	8
1993	44,418	1,063	1,505	2,170	3,132	4,420	5,486	6,171	6,690	11,660	15
1992	42,157	1,007	1,485	2,204	3,171	4,323	5,231	5,881	6,175	10,647	13
1991	41,885	984	1,381	2,077	3,143	4,472	5,358	6,102	6,192	10,254	6
1990	41,268	928	1,397	2,048	3,354	4,639	5,097	6,127	5,964	9,893	17
1989	42,196	956	1,415	2,227	3,472	4,649	5,290	6,404	6,051	9,880	10
1988	42,747	989	1,509	2,252	3,628	4,681	5,372	6,557	6,219	9,675	16
1987	41,467	951	1,454	2,272	3,542	4,590	5,509	6,105	5,939	9,320	13
1986	41,182	948	1,509	2,248	3,584	4,525	5,544	6,241	5,782	8,958	21
1985	40,183	897	1,496	2,524	3,532	4,516	5,621	6,052	5,573	8,231	17
1984	38,960	910	1,498	2,495	3,488	4,337	5,571	5,663	5,637	7,669	15
1983	38,907	882	1,579	2,533	3,573	4,379	5,668	5,720	5,503	7,399	21
1982	36,493	950	1,587	2,412	3,417	4,204	5,207	5,256	5,046	6,793	26
1981	36,286	1,002	1,589	2,434	3,418	4,334	5,125	5,176	4,913	6,698	26
1980	36,756	985	1,653	2,467	3,346	4,558	5,211	5,531	4,731	6,545	21
1979	34,087	974	1,652	2,335	3,228	4,336	4,916	5,317	4,028	5,669	21
Diseases of heart (390-398,402,404,410-429)											
Black, both sexes											
1998	78,294	3,416	4,097	5,081	6,492	7,886	9,594	10,355	9,642	16,517	23
1997	77,174	3,267	4,132	5,065	6,491	7,809	9,754	9,991	9,488	16,208	20
1996	77,641	3,263	4,034	4,918	6,767	7,967	9,924	10,137	9,677	15,880	21
1995	78,643	3,232	3,950	4,945	6,688	8,578	10,222	10,144	9,937	15,682	22
1994	76,852	3,009	3,872	4,972	6,431	8,598	10,039	9,866	9,841	14,932	24
1993	79,011	2,806	3,910	5,268	6,941	9,077	10,295	10,186	10,107	15,190	31
1992	75,600	2,743	3,652	5,060	6,903	8,771	9,917	9,948	9,475	14,038	33
1991	76,014	2,671	3,526	4,998	7,019	9,175	10,092	10,307	9,651	13,597	26
1990	75,111	2,517	3,547	5,044	7,207	9,356	9,660	10,427	9,318	13,310	41
1989	77,431	2,674	3,757	5,346	7,737	9,667	9,920	10,917	9,427	13,228	31
1988	79,466	2,655	3,755	5,500	8,087	9,773	10,289	11,359	9,906	13,240	37
1987	77,747	2,517	3,813	5,324	7,975	9,745	10,489	10,846	9,533	12,729	35
1986	77,726	2,447	3,864	5,594	8,164	9,544	10,556	10,953	9,255	12,463	41
1985	76,684	2,486	3,919	5,891	8,130	9,494	10,863	10,592	9,159	11,527	33
1984	74,698	2,473	3,891	5,866	8,090	9,133	10,803	10,046	9,112	10,859	35
1983	75,008	2,436	4,094	6,119	8,133	9,226	11,050	10,166	8,961	10,557	32
1982	71,468	2,577	3,992	5,995	7,824	9,057	10,285	9,522	8,243	9,887	55
1981	71,697	2,664	4,158	6,203	7,879	9,178	10,198	9,281	8,198	9,771	50
1980	72,956	2,656	4,327	6,243	7,802	9,684	10,183	9,984	7,924	9,762	54
1979	68,579	2,682	4,360	6,124	7,428	9,337	9,724	9,637	6,825	8,432	47

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of heart (390-398,402,404,410-429)											
Black, male											
1998	37,662	2,158	2,636	3,097	3,809	4,380	4,800	4,860	3,847	4,894	16
1997	37,212	2,129	2,634	3,101	3,884	4,309	4,906	4,627	3,801	4,741	13
1996	37,335	2,086	2,514	2,974	3,967	4,400	5,026	4,752	3,838	4,730	14
1995	38,389	2,092	2,557	3,080	3,929	4,761	5,231	4,780	3,915	4,741	11
1994	37,241	1,922	2,494	2,967	3,794	4,702	5,132	4,504	3,964	4,521	16
1993	38,357	1,801	2,500	3,248	4,079	5,002	5,264	4,568	4,013	4,642	19
1992	37,040	1,792	2,250	3,013	3,967	4,805	5,150	4,624	3,853	4,388	20
1991	37,454	1,738	2,234	3,067	4,112	5,030	5,156	4,708	3,930	4,306	21
1990	37,038	1,638	2,227	3,149	4,087	5,064	4,956	4,785	3,823	4,294	25
1989	38,321	1,771	2,434	3,273	4,486	5,352	5,001	4,966	3,811	4,203	22
1988	39,584	1,706	2,335	3,384	4,688	5,388	5,282	5,256	4,089	4,316	21
1987	38,934	1,616	2,436	3,187	4,642	5,423	5,315	5,079	3,957	4,175	22
1986	39,076	1,543	2,444	3,455	4,770	5,255	5,336	5,078	3,849	4,200	23
1985	38,982	1,628	2,490	3,514	4,782	5,241	5,537	4,894	3,938	3,964	19
1984	38,034	1,620	2,468	3,528	4,777	5,036	5,511	4,680	3,793	3,790	22
1983	38,311	1,598	2,587	3,717	4,737	5,064	5,655	4,744	3,796	3,722	19
1982	36,925	1,676	2,469	3,699	4,579	5,061	5,309	4,492	3,493	3,572	32
1981	37,188	1,708	2,630	3,866	4,604	5,027	5,269	4,354	3,539	3,517	28
1980	37,877	1,702	2,736	3,858	4,572	5,297	5,177	4,718	3,441	3,615	35
1979	36,122	1,739	2,750	3,870	4,321	5,169	5,015	4,545	3,056	3,169	27
Diseases of heart (390-398,402,404,410-429)											
Black, female											
1998	40,632	1,258	1,461	1,984	2,683	3,506	4,794	5,495	5,795	11,623	7
1997	39,962	1,138	1,498	1,964	2,607	3,500	4,848	5,364	5,687	11,467	7
1996	40,306	1,177	1,520	1,944	2,800	3,567	4,898	5,385	5,839	11,150	7
1995	40,254	1,140	1,393	1,865	2,759	3,817	4,991	5,364	6,022	10,941	11
1994	39,611	1,087	1,378	2,005	2,637	3,896	4,907	5,362	5,877	10,411	8
1993	40,654	1,005	1,410	2,020	2,862	4,075	5,031	5,618	6,094	10,548	12
1992	38,560	951	1,402	2,047	2,936	3,966	4,767	5,324	5,622	9,650	13
1991	38,560	933	1,292	1,931	2,907	4,145	4,936	5,599	5,721	9,291	5
1990	38,073	879	1,320	1,895	3,120	4,292	4,704	5,642	5,495	9,016	16
1989	39,110	903	1,323	2,073	3,251	4,315	4,919	5,951	5,616	9,025	9
1988	39,882	949	1,420	2,116	3,399	4,385	5,007	6,103	5,817	8,924	16
1987	38,813	901	1,377	2,137	3,333	4,322	5,174	5,767	5,576	8,554	13
1986	38,650	904	1,420	2,139	3,394	4,289	5,220	5,875	5,406	8,263	18
1985	37,702	858	1,429	2,377	3,348	4,253	5,326	5,698	5,221	7,563	14
1984	36,664	853	1,423	2,338	3,313	4,097	5,292	5,366	5,319	7,069	13
1983	36,697	838	1,507	2,402	3,396	4,162	5,395	5,422	5,165	6,835	13
1982	34,543	901	1,523	2,296	3,245	3,996	4,976	5,030	4,750	6,315	23
1981	34,509	956	1,528	2,337	3,275	4,151	4,929	4,927	4,659	6,254	22
1980	35,079	954	1,591	2,385	3,230	4,387	5,006	5,266	4,483	6,147	19
1979	32,457	943	1,610	2,254	3,107	4,168	4,709	5,092	3,769	5,263	20

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All races, both sexes											
1998	4,792	141	187	232	317	445	623	759	759	1,012	-
1997	5,014	161	192	234	304	466	658	818	784	1,039	-
1996	5,006	155	181	235	317	493	702	793	837	951	-
1995	5,147	140	178	252	338	493	745	857	825	972	1
1994	5,415	168	184	259	386	597	835	850	859	903	1
1993	5,743	167	203	296	396	649	855	922	881	1,005	-
1992	5,702	156	229	296	435	671	811	975	870	884	1
1991	5,951	170	204	296	507	688	876	1,024	899	925	-
1990	6,018	148	226	340	518	765	859	960	911	884	-
1989	6,087	181	216	335	523	755	911	1,026	895	827	-
1988	6,421	192	244	423	571	840	983	1,053	899	809	1
1987	6,389	191	246	399	625	898	976	1,051	812	751	-
1986	6,441	198	279	437	683	826	1,022	1,040	789	714	2
1985	6,593	193	333	508	675	849	1,057	1,022	796	663	-
1984	6,912	221	311	527	796	941	1,133	1,061	754	652	-
1983	6,895	264	392	582	794	948	1,057	969	730	649	1
1982	7,053	228	361	597	885	1,052	1,192	1,021	636	587	2
1981	7,378	281	423	691	840	1,122	1,223	1,011	665	570	-
1980	7,836	328	510	768	928	1,234	1,207	1,018	694	535	1
1979	7,795	344	541	766	956	1,207	1,251	975	593	554	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All races, male											
1998	1,422	59	76	76	113	155	177	231	198	210	-
1997	1,553	59	76	103	101	139	218	248	227	234	-
1996	1,482	48	76	75	109	167	215	246	219	186	-
1995	1,534	45	64	99	118	148	229	233	231	219	-
1994	1,679	67	77	90	149	212	274	242	223	188	-
1993	1,790	57	71	115	149	234	258	282	226	233	-
1992	1,714	71	92	103	156	224	239	272	214	181	-
1991	1,786	64	67	115	182	225	277	273	224	200	-
1990	1,830	67	64	131	203	281	280	245	215	157	-
1989	1,877	71	78	135	199	253	285	292	216	168	-
1988	1,945	85	93	154	194	281	296	282	207	174	1
1987	1,934	68	80	162	231	296	291	292	185	134	-
1986	2,054	72	103	180	258	287	314	288	189	147	-
1985	2,142	74	134	192	251	304	336	294	191	137	-
1984	2,279	79	140	197	319	303	379	306	190	135	-
1983	2,199	91	143	214	304	336	297	252	188	133	1
1982	2,336	103	151	235	348	379	371	267	140	114	1
1981	2,469	113	171	267	302	393	383	301	175	129	-
1980	2,660	123	213	292	338	437	362	288	190	133	1
1979	2,651	133	222	323	346	393	389	286	165	106	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All races, female											
1998	3,370	82	111	156	204	290	446	528	561	802	-
1997	3,461	102	116	131	203	327	440	570	557	805	-
1996	3,524	107	105	160	208	326	487	547	618	765	-
1995	3,613	95	114	153	220	345	516	624	594	753	1
1994	3,736	101	107	169	237	385	561	608	636	715	1
1993	3,953	110	132	181	247	415	597	640	655	772	-
1992	3,988	85	137	193	279	447	572	703	656	703	1
1991	4,165	106	137	181	325	463	599	751	675	725	-
1990	4,188	81	162	209	315	484	579	715	696	727	-
1989	4,210	110	138	200	324	502	626	734	679	659	-
1988	4,476	107	151	269	377	559	687	771	692	635	-
1987	4,455	123	166	237	394	602	685	759	627	617	-
1986	4,387	126	176	257	425	539	708	752	600	567	2
1985	4,451	119	199	316	424	545	721	728	605	526	-
1984	4,633	142	171	330	477	638	754	755	564	517	-
1983	4,696	173	249	368	490	612	760	717	542	516	-
1982	4,717	125	210	362	537	673	821	754	496	473	1
1981	4,909	168	252	424	538	729	840	710	490	441	-
1980	5,176	205	297	476	590	797	845	730	504	402	-
1979	5,144	211	319	443	610	814	862	689	428	448	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
White, both sexes											
1998	4,334	100	150	190	269	402	576	718	732	964	-
1997	4,547	118	151	190	266	434	611	778	753	1,005	-
1996	4,534	124	144	196	270	435	650	759	805	909	-
1995	4,661	102	132	207	289	446	694	814	795	943	1
1994	4,923	118	148	211	345	536	785	824	832	868	1
1993	5,231	128	170	248	358	592	809	877	831	968	-
1992	5,161	112	180	250	387	606	759	924	830	848	-
1991	5,426	122	162	263	456	633	815	963	862	887	-
1990	5,483	117	185	294	456	704	814	916	879	842	-
1989	5,523	142	176	285	463	688	851	973	860	796	-
1988	5,874	150	197	380	518	777	931	1,012	866	772	-
1987	5,875	159	211	350	574	837	935	1,004	783	727	-
1986	5,872	168	236	382	618	774	960	1,003	758	691	2
1985	6,026	158	278	448	619	792	1,010	981	769	629	-
1984	6,355	179	281	469	743	871	1,064	1,032	734	627	-
1983	6,341	220	348	518	744	902	1,001	925	710	622	1
1982	6,504	191	321	532	811	995	1,138	982	613	564	2
1981	6,798	237	375	624	772	1,063	1,170	967	640	549	-
1980	7,203	275	453	698	862	1,168	1,160	984	670	514	1
1979	7,195	289	481	703	896	1,151	1,200	927	574	535	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
White, male											
1998	1,288	44	63	65	97	142	169	219	190	201	-
1997	1,389	44	58	81	91	128	203	236	219	228	-
1996	1,331	39	63	65	96	145	193	236	211	176	-
1995	1,375	34	49	81	105	138	213	221	228	210	-
1994	1,513	47	67	80	136	186	259	234	215	182	-
1993	1,621	42	65	101	137	213	243	271	213	225	-
1992	1,537	51	76	92	140	203	223	255	201	174	-
1991	1,607	45	55	102	163	203	258	253	215	188	-
1990	1,655	53	57	119	175	264	269	231	207	146	-
1989	1,674	61	63	118	173	235	262	276	208	163	-
1988	1,746	69	75	142	179	259	278	268	198	162	-
1987	1,767	64	68	148	214	283	280	271	177	125	-
1986	1,842	60	90	162	232	272	294	274	181	140	-
1985	1,934	61	116	174	230	280	319	283	181	126	-
1984	2,087	64	127	183	300	283	359	297	182	129	-
1983	1,996	79	129	198	283	318	276	239	182	127	1
1982	2,133	90	136	214	320	355	350	253	136	110	1
1981	2,254	97	155	241	277	371	362	288	167	123	-
1980	2,424	109	189	264	314	413	346	280	182	128	1
1979	2,414	110	199	308	321	370	372	272	160	100	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
White, female											
1998	3,046	56	87	125	172	260	407	499	542	763	-
1997	3,158	74	93	109	175	306	408	542	534	777	-
1996	3,203	85	81	131	174	290	457	523	594	733	-
1995	3,286	68	83	126	184	308	481	593	567	733	1
1994	3,410	71	81	131	209	350	526	590	617	686	1
1993	3,610	86	105	147	221	379	566	606	618	743	-
1992	3,624	61	104	158	247	403	536	669	629	674	-
1991	3,819	77	107	161	293	430	557	710	647	699	-
1990	3,828	64	128	175	281	440	545	685	672	696	-
1989	3,849	81	113	167	290	453	589	697	652	633	-
1988	4,128	81	122	238	339	518	653	744	668	610	-
1987	4,108	95	143	202	360	554	655	733	606	602	-
1986	4,030	108	146	220	386	502	666	729	577	551	2
1985	4,092	97	162	274	389	512	691	698	588	503	-
1984	4,268	115	154	286	443	588	705	735	552	498	-
1983	4,345	141	219	320	461	584	725	686	528	495	-
1982	4,371	101	185	318	491	640	788	729	477	454	1
1981	4,544	140	220	383	495	692	808	679	473	426	-
1980	4,779	166	264	434	548	755	814	704	488	386	-
1979	4,781	179	282	395	575	781	828	655	414	435	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All other, both sexes											
1998	458	41	37	42	48	43	47	41	27	48	-
1997	467	43	41	44	38	32	47	40	31	34	-
1996	472	31	37	39	47	58	52	34	32	42	-
1995	486	38	46	45	49	47	51	43	30	29	-
1994	492	50	36	48	41	61	50	26	27	35	-
1993	512	39	33	48	38	57	46	45	50	37	-
1992	541	44	49	46	48	65	52	51	40	36	1
1991	525	48	42	33	51	55	61	61	37	38	-
1990	535	31	41	46	62	61	45	44	32	42	-
1989	564	39	40	50	60	67	60	53	35	31	-
1988	547	42	47	43	53	63	52	41	33	37	1
1987	514	32	35	49	51	61	41	47	29	24	-
1986	569	30	43	55	65	52	62	37	31	23	-
1985	567	35	55	60	56	57	47	41	27	34	-
1984	557	42	30	58	53	70	69	29	20	25	-
1983	554	44	44	64	50	46	56	44	20	27	-
1982	549	37	40	65	74	57	54	39	23	23	-
1981	580	44	48	67	68	59	53	44	25	21	-
1980	633	53	57	70	66	66	47	34	24	21	-
1979	600	55	60	63	60	56	51	48	19	19	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All other, male											
1998	134	15	13	11	16	13	8	12	8	9	-
1997	164	15	18	22	10	11	15	12	8	6	-
1996	151	9	13	10	13	22	22	10	8	10	-
1995	159	11	15	18	13	10	16	12	3	9	-
1994	166	20	10	10	13	26	15	8	8	6	-
1993	169	15	6	14	12	21	15	11	13	8	-
1992	177	20	16	11	16	21	16	17	13	7	-
1991	179	19	12	13	19	22	19	20	9	12	-
1990	175	14	7	12	28	17	11	14	8	11	-
1989	203	10	15	17	26	18	23	16	8	5	-
1988	199	16	18	12	15	22	18	14	9	12	1
1987	167	4	12	14	17	13	11	21	8	9	-
1986	212	12	13	18	26	15	20	14	8	7	-
1985	208	13	18	18	21	24	17	11	10	11	-
1984	192	15	13	14	19	20	20	9	8	6	-
1983	203	12	14	16	21	18	21	13	6	6	-
1982	203	13	15	21	28	24	21	14	4	4	-
1981	215	16	16	26	25	22	21	13	8	6	-
1980	236	14	24	28	24	24	16	8	8	5	-
1979	237	23	23	15	25	23	17	14	5	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
All other, female											
1998	324	26	24	31	32	30	39	29	19	39	-
1997	303	28	23	22	28	21	32	28	23	28	-
1996	321	22	24	29	34	36	30	24	24	32	-
1995	327	27	31	27	36	37	35	31	27	20	-
1994	326	30	26	38	28	35	35	18	19	29	-
1993	343	24	27	34	26	36	31	34	37	29	-
1992	364	24	33	35	32	44	36	34	27	29	1
1991	346	29	30	20	32	33	42	41	28	26	-
1990	360	17	34	34	34	44	34	30	24	31	-
1989	361	29	25	33	34	49	37	37	27	26	-
1988	348	26	29	31	38	41	34	27	24	25	-
1987	347	28	23	35	34	48	30	26	21	15	-
1986	357	18	30	37	39	37	42	23	23	16	-
1985	359	22	37	42	35	33	30	30	17	23	-
1984	365	27	17	44	34	50	49	20	12	19	-
1983	351	32	30	48	29	28	35	31	14	21	-
1982	346	24	25	44	46	33	33	25	19	19	-
1981	365	28	32	41	43	37	32	31	17	15	-
1980	397	39	33	42	42	42	31	26	16	16	-
1979	363	32	37	48	35	33	34	34	14	13	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
Black, both sexes											
1998	356	33	30	32	39	34	32	30	20	33	-
1997	353	38	36	36	30	23	30	30	17	24	-
1996	355	26	28	29	35	38	35	24	23	30	-
1995	368	31	37	31	36	35	35	33	23	15	-
1994	370	37	27	36	35	49	38	19	14	26	-
1993	410	36	30	43	26	41	33	33	32	31	-
1992	431	35	44	32	39	48	43	35	29	32	1
1991	413	39	37	26	44	41	37	46	24	34	-
1990	436	26	35	36	48	50	39	29	27	35	-
1989	448	31	30	43	46	50	44	44	30	25	-
1988	426	38	39	33	41	47	37	33	22	28	1
1987	412	26	30	39	42	51	29	32	24	14	-
1986	470	26	37	46	58	36	52	28	26	18	-
1985	472	28	49	52	49	46	37	34	22	23	-
1984	458	32	28	49	41	61	60	21	12	21	-
1983	465	40	36	53	43	39	46	37	16	21	-
1982	458	30	35	51	62	46	50	36	22	21	-
1981	479	38	39	58	58	46	44	37	19	18	-
1980	556	51	51	63	58	57	42	31	21	19	-
1979	531	50	58	56	51	51	39	42	19	18	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
Black, male											
1998	101	10	9	8	13	10	4	11	6	4	-
1997	124	12	17	19	7	7	9	6	7	2	-
1996	106	7	9	5	10	15	16	7	4	5	-
1995	123	9	13	14	11	9	11	5	3	4	-
1994	131	17	8	7	12	22	12	7	1	5	-
1993	133	13	6	14	9	12	9	8	8	7	-
1992	131	17	13	6	12	14	12	8	8	7	-
1991	138	14	12	10	17	15	13	13	6	11	-
1990	146	14	5	10	23	14	10	11	7	9	-
1989	168	9	11	15	22	15	18	14	6	4	-
1988	154	15	16	9	12	18	12	12	7	6	1
1987	132	1	10	10	13	12	8	12	6	7	-
1986	177	11	12	15	24	11	15	11	5	5	-
1985	174	9	16	17	18	21	12	8	8	8	-
1984	157	11	13	13	13	17	17	5	4	5	-
1983	166	10	10	15	19	13	16	11	3	5	-
1982	158	10	14	15	22	19	17	12	3	3	-
1981	173	15	11	24	23	15	17	9	6	4	-
1980	206	14	21	23	19	21	15	8	7	5	-
1979	211	22	21	14	21	22	13	13	5	6	-
Acute rheumatic fever and chronic rheumatic heart diseases (390-398)											
Black, female											
1998	255	23	21	24	26	24	28	19	14	29	-
1997	229	26	19	17	23	16	21	24	10	22	-
1996	249	19	19	24	25	23	19	17	19	25	-
1995	245	22	24	17	25	26	24	28	20	11	-
1994	239	20	19	29	23	27	26	12	13	21	-
1993	277	23	24	29	17	29	24	25	24	24	-
1992	300	18	31	26	27	34	31	27	21	25	1
1991	275	25	25	16	27	26	24	33	18	23	-
1990	290	12	30	26	25	36	29	18	20	26	-
1989	280	22	19	28	24	35	26	30	24	21	-
1988	272	23	23	24	29	29	25	21	15	22	-
1987	280	25	20	29	29	39	21	20	18	7	-
1986	293	15	25	31	34	25	37	17	21	13	-
1985	298	19	33	35	31	25	25	26	14	15	-
1984	301	21	15	36	28	44	43	16	8	16	-
1983	299	30	26	38	24	26	30	26	13	16	-
1982	300	20	21	36	40	27	33	24	19	18	-
1981	306	23	28	34	35	31	27	28	13	14	-
1980	350	37	30	40	39	36	27	23	14	14	-
1979	320	28	37	42	30	29	26	29	14	12	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Hypertensive heart disease (402)											
All races, both sexes											
1998	27,715	1,059	1,303	1,498	1,836	2,145	2,752	3,404	3,869	8,610	5
1997	26,607	994	1,276	1,352	1,633	2,081	2,751	3,358	3,696	8,298	6
1996	26,188	909	1,193	1,337	1,699	2,020	2,759	3,297	3,857	8,006	5
1995	25,014	894	1,062	1,219	1,648	2,209	2,677	3,142	3,693	7,404	4
1994	23,943	877	1,012	1,216	1,564	1,993	2,673	2,999	3,637	6,900	4
1993	23,037	703	981	1,179	1,573	2,097	2,647	3,054	3,491	6,369	2
1992	21,853	629	877	1,247	1,655	2,084	2,612	2,879	3,261	5,704	5
1991	21,365	641	800	1,145	1,648	2,065	2,493	2,993	3,243	5,495	7
1990	21,072	602	797	1,151	1,659	2,256	2,489	2,988	3,111	5,239	5
1989	20,963	626	793	1,229	1,717	2,231	2,466	3,077	3,097	4,974	6
1988	20,788	555	819	1,225	1,806	2,329	2,594	3,004	3,028	4,716	5
1987	20,678	536	838	1,257	1,876	2,340	2,681	2,922	3,030	4,526	2
1986	20,509	513	836	1,255	1,822	2,256	2,706	3,060	2,938	4,445	9
1985	20,882	525	851	1,337	1,919	2,334	2,748	3,043	3,024	4,426	7
1984	20,712	513	852	1,332	1,898	2,376	2,842	2,979	3,017	4,261	5
1983	20,984	536	848	1,433	1,995	2,379	2,991	3,041	3,005	4,124	2
1982	20,779	570	874	1,416	1,890	2,408	2,850	3,120	3,022	4,009	14
1981	20,793	576	961	1,501	1,935	2,395	2,884	3,012	2,942	3,963	4
1980	21,095	506	936	1,499	1,922	2,554	2,863	3,150	3,088	3,949	5
1979	20,992	593	1,013	1,525	1,942	2,596	2,872	3,068	2,982	3,742	14
Hypertensive heart disease (402)											
All races, male											
1998	11,937	728	885	990	1,155	1,220	1,427	1,473	1,339	1,854	1
1997	11,386	680	870	878	1,014	1,239	1,377	1,412	1,296	1,795	4
1996	11,129	608	800	893	1,036	1,187	1,396	1,412	1,292	1,742	3
1995	10,638	631	699	810	1,023	1,254	1,326	1,327	1,257	1,575	2
1994	10,055	596	669	757	970	1,110	1,325	1,262	1,230	1,401	3
1993	9,771	475	647	753	945	1,191	1,269	1,306	1,140	1,401	1
1992	9,211	408	558	774	990	1,153	1,278	1,178	1,079	1,160	2
1991	8,995	436	513	740	988	1,109	1,200	1,246	1,071	1,131	7
1990	8,822	369	506	732	968	1,215	1,179	1,186	1,044	1,086	3
1989	8,923	404	508	795	1,042	1,239	1,204	1,249	968	1,005	3
1988	8,850	349	533	766	1,043	1,285	1,268	1,187	954	992	4
1987	8,855	343	534	778	1,116	1,292	1,265	1,161	992	907	1
1986	8,614	330	508	801	1,039	1,235	1,292	1,196	910	855	6
1985	8,970	335	536	810	1,122	1,266	1,316	1,254	935	917	2
1984	8,892	332	541	836	1,120	1,239	1,354	1,190	917	917	2
1983	9,106	344	545	895	1,201	1,292	1,341	1,218	934	888	1
1982	8,896	358	542	852	1,098	1,276	1,322	1,236	924	889	6
1981	8,994	338	617	925	1,086	1,312	1,342	1,190	903	874	3
1980	8,870	303	555	913	1,130	1,319	1,312	1,200	906	839	2
1979	8,902	373	609	919	1,086	1,344	1,286	1,163	874	830	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart disease (402)											
All races, female											
1998	15,778	331	418	508	681	925	1,325	1,931	2,530	6,756	4
1997	15,221	314	406	474	619	842	1,374	1,946	2,400	6,503	2
1996	15,059	301	393	444	663	833	1,363	1,885	2,565	6,264	2
1995	14,376	263	363	409	625	955	1,351	1,815	2,436	5,829	2
1994	13,888	281	343	459	594	883	1,348	1,737	2,407	5,499	1
1993	13,266	228	334	426	628	906	1,378	1,748	2,351	4,968	1
1992	12,642	221	319	473	665	931	1,334	1,701	2,182	4,544	3
1991	12,370	205	287	405	660	956	1,293	1,747	2,172	4,364	-
1990	12,250	233	291	419	691	1,041	1,310	1,802	2,067	4,153	2
1989	12,040	222	285	434	675	992	1,262	1,828	2,129	3,969	3
1988	11,938	206	286	459	763	1,044	1,326	1,817	2,074	3,724	1
1987	11,823	193	304	479	760	1,048	1,416	1,761	2,038	3,619	1
1986	11,895	183	328	454	783	1,021	1,414	1,864	2,028	3,590	3
1985	11,912	190	315	527	797	1,068	1,432	1,789	2,089	3,509	5
1984	11,820	181	311	496	778	1,137	1,488	1,789	2,100	3,344	3
1983	11,878	192	303	538	794	1,087	1,650	1,823	2,071	3,236	1
1982	11,883	212	332	564	792	1,132	1,528	1,884	2,098	3,120	8
1981	11,799	238	344	576	849	1,083	1,542	1,822	2,039	3,089	1
1980	12,225	203	381	586	792	1,235	1,551	1,950	2,182	3,110	3
1979	12,090	220	404	606	856	1,252	1,586	1,905	2,108	2,912	7
Hypertensive heart disease (402)											
White, both sexes											
1998	20,047	554	733	853	1,075	1,377	1,898	2,540	3,111	7,346	2
1997	19,377	520	718	760	953	1,300	1,900	2,546	2,985	7,123	5
1996	19,197	465	661	790	1,049	1,349	1,915	2,482	3,116	6,849	3
1995	18,166	461	553	701	950	1,457	1,914	2,382	2,944	6,314	2
1994	17,395	408	553	676	930	1,296	1,918	2,260	2,931	5,933	1
1993	16,620	341	494	651	967	1,349	1,842	2,324	2,770	5,448	1
1992	15,685	300	452	670	999	1,355	1,877	2,180	2,610	4,851	1
1991	15,409	297	423	630	1,025	1,346	1,763	2,245	2,622	4,695	5
1990	15,165	273	401	655	982	1,494	1,687	2,284	2,550	4,498	1
1989	15,012	278	397	675	1,030	1,481	1,737	2,314	2,523	4,250	2
1988	14,901	256	417	698	1,102	1,571	1,847	2,256	2,426	4,034	3
1987	15,119	283	408	753	1,174	1,603	1,936	2,263	2,493	3,909	2
1986	14,982	222	420	740	1,148	1,573	1,950	2,348	2,408	3,875	4
1985	15,303	248	443	797	1,205	1,630	2,008	2,333	2,460	3,878	5
1984	15,237	233	443	753	1,222	1,659	2,085	2,350	2,491	3,734	2
1983	15,463	254	441	874	1,304	1,699	2,128	2,409	2,462	3,629	1
1982	15,392	270	482	836	1,255	1,740	2,048	2,493	2,491	3,521	6
1981	15,474	271	525	921	1,300	1,704	2,160	2,404	2,456	3,477	-
1980	15,682	233	504	922	1,278	1,783	2,144	2,488	2,627	3,466	3
1979	15,518	297	550	915	1,299	1,831	2,116	2,425	2,526	3,300	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Hypertensive heart disease (402)											
White, male											
1998	8,223	427	527	607	726	795	1,023	1,098	1,049	1,524	-
1997	7,916	391	518	528	630	817	965	1,076	1,069	1,475	3
1996	7,735	342	476	550	676	822	985	1,041	1,039	1,407	1
1995	7,371	356	389	494	626	856	967	998	1,005	1,290	1
1994	6,996	316	413	445	608	750	966	947	989	1,172	-
1993	6,742	249	356	464	611	792	901	991	907	1,131	1
1992	6,312	220	312	461	635	785	940	903	838	913	-
1991	6,218	228	287	435	634	762	875	950	852	915	5
1990	6,077	181	278	447	616	828	831	925	815	886	-
1989	6,082	194	272	463	651	863	858	948	778	805	1
1988	6,099	183	306	460	663	898	922	897	750	794	2
1987	6,245	202	281	477	741	926	948	906	791	746	1
1986	6,043	163	264	526	689	871	978	903	727	697	1
1985	6,322	174	307	524	736	902	978	966	740	759	1
1984	6,302	171	307	508	749	904	997	944	740	776	-
1983	6,458	185	310	568	829	964	955	962	740	738	-
1982	6,341	182	325	541	749	940	968	984	732	751	3
1981	6,462	173	367	599	757	957	1,005	949	742	727	-
1980	6,386	159	325	587	773	932	1,016	947	750	732	1
1979	6,325	204	356	575	775	979	945	903	697	705	4
Hypertensive heart disease (402)											
White, female											
1998	11,824	127	206	246	349	582	875	1,442	2,062	5,822	2
1997	11,461	129	200	232	323	483	935	1,470	1,916	5,648	2
1996	11,462	123	185	240	373	527	930	1,441	2,077	5,442	2
1995	10,795	105	164	207	324	601	947	1,384	1,939	5,024	1
1994	10,399	92	140	231	322	546	952	1,313	1,942	4,761	1
1993	9,878	92	138	187	356	557	941	1,333	1,863	4,317	-
1992	9,373	80	140	209	364	570	937	1,277	1,772	3,938	1
1991	9,191	69	136	195	391	584	888	1,295	1,770	3,780	-
1990	9,088	92	123	208	366	666	856	1,359	1,735	3,612	1
1989	8,930	84	125	212	379	618	879	1,366	1,745	3,445	1
1988	8,802	73	111	238	439	673	925	1,359	1,676	3,240	1
1987	8,874	81	127	276	433	677	988	1,357	1,702	3,163	1
1986	8,939	59	156	214	459	702	972	1,445	1,681	3,178	3
1985	8,981	74	136	273	469	728	1,030	1,367	1,720	3,119	4
1984	8,935	62	136	245	473	755	1,088	1,406	1,751	2,958	2
1983	9,005	69	131	306	475	735	1,173	1,447	1,722	2,891	1
1982	9,051	88	157	295	506	800	1,080	1,509	1,759	2,770	3
1981	9,012	98	158	322	543	747	1,155	1,455	1,714	2,750	-
1980	9,296	74	179	335	505	851	1,128	1,541	1,877	2,734	2
1979	9,193	93	194	340	524	852	1,171	1,522	1,829	2,595	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart disease (402)											
All other, both sexes											
1998	7,668	505	570	645	761	768	854	864	758	1,264	3
1997	7,230	474	558	592	680	781	851	812	711	1,175	1
1996	6,991	444	532	547	650	671	844	815	741	1,157	2
1995	6,848	433	509	518	698	752	763	760	749	1,090	2
1994	6,548	469	459	540	634	697	755	739	706	967	3
1993	6,417	362	487	528	606	748	805	730	721	921	1
1992	6,168	329	425	577	656	729	735	699	651	853	4
1991	5,956	344	377	515	623	719	730	748	621	800	2
1990	5,907	329	396	496	677	762	802	704	561	741	4
1989	5,951	348	396	554	687	750	729	763	574	724	4
1988	5,887	299	402	527	704	758	747	748	602	682	2
1987	5,559	253	430	504	702	737	745	659	537	617	-
1986	5,527	291	416	515	674	683	756	712	530	570	5
1985	5,579	277	408	540	714	704	740	710	564	548	2
1984	5,475	280	409	579	676	717	757	629	526	527	3
1983	5,521	282	407	559	691	680	863	632	543	495	1
1982	5,387	300	392	580	635	668	802	627	531	488	8
1981	5,319	305	436	580	635	691	724	608	486	486	4
1980	5,413	273	432	577	644	771	719	662	461	483	2
1979	5,474	296	463	610	643	765	756	643	456	442	7
Hypertensive heart disease (402)											
All other, male											
1998	3,714	301	358	383	429	425	404	375	290	330	1
1997	3,470	289	352	350	384	422	412	336	227	320	1
1996	3,394	266	324	343	360	365	411	371	253	335	2
1995	3,267	275	310	316	397	398	359	329	252	285	1
1994	3,059	280	256	312	362	360	359	315	241	229	3
1993	3,029	226	291	289	334	399	368	315	233	270	-
1992	2,899	188	246	313	355	368	338	275	241	247	2
1991	2,777	208	226	305	354	347	325	296	219	216	2
1990	2,745	188	228	285	352	387	348	261	229	200	3
1989	2,841	210	236	332	391	376	346	301	190	200	2
1988	2,751	166	227	306	380	387	346	290	204	198	2
1987	2,610	141	253	301	375	366	317	255	201	161	-
1986	2,571	167	244	275	350	364	314	293	183	158	5
1985	2,648	161	229	286	386	364	338	288	195	158	1
1984	2,590	161	234	328	371	335	357	246	177	141	2
1983	2,648	159	235	327	372	328	386	256	194	150	1
1982	2,555	176	217	311	349	336	354	252	192	138	3
1981	2,532	165	250	326	329	355	337	241	161	147	3
1980	2,484	144	230	326	357	387	296	253	156	107	1
1979	2,577	169	253	344	311	365	341	260	177	125	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart disease (402)											
All other, female											
1998	3,954	204	212	262	332	343	450	489	468	934	2
1997	3,760	185	206	242	296	359	439	476	484	855	-
1996	3,597	178	208	204	290	306	433	444	488	822	-
1995	3,581	158	199	202	301	354	404	431	497	805	1
1994	3,489	189	203	228	272	337	396	424	465	738	-
1993	3,388	136	196	239	272	349	437	415	488	651	1
1992	3,269	141	179	264	301	361	397	424	410	606	2
1991	3,179	136	151	210	269	372	405	452	402	584	-
1990	3,162	141	168	211	325	375	454	443	332	541	1
1989	3,110	138	160	222	296	374	383	462	384	524	2
1988	3,136	133	175	221	324	371	401	458	398	484	-
1987	2,949	112	177	203	327	371	428	404	336	456	-
1986	2,956	124	172	240	324	319	442	419	347	412	-
1985	2,931	116	179	254	328	340	402	422	369	390	1
1984	2,885	119	175	251	305	382	400	383	349	386	1
1983	2,873	123	172	232	319	352	477	376	349	345	-
1982	2,832	124	175	269	286	332	448	375	339	350	5
1981	2,787	140	186	254	306	336	387	367	325	339	1
1980	2,929	129	202	251	287	384	423	409	305	376	1
1979	2,897	127	210	266	332	400	415	383	279	317	4
Hypertensive heart disease (402)											
Black, both sexes											
1998	7,158	490	540	618	721	727	776	817	697	1,128	3
1997	6,708	450	535	561	633	743	775	744	645	1,048	1
1996	6,564	427	509	520	617	635	795	758	687	1,050	2
1995	6,408	415	491	492	658	713	713	711	683	973	2
1994	6,160	455	437	515	601	668	704	686	648	887	2
1993	6,057	356	468	502	576	710	757	683	672	839	1
1992	5,854	319	412	551	636	694	696	656	607	781	4
1991	5,639	336	367	501	585	688	700	707	569	715	2
1990	5,636	322	384	482	652	719	774	663	523	688	4
1989	5,680	340	383	537	665	721	700	716	538	662	4
1988	5,626	294	392	516	673	724	716	697	563	642	2
1987	5,276	242	411	485	676	704	706	623	505	558	-
1986	5,312	281	404	505	658	662	723	678	499	531	3
1985	5,342	266	401	523	696	684	702	670	525	507	1
1984	5,276	272	396	561	660	692	726	599	502	504	3
1983	5,339	272	394	544	675	665	835	609	517	470	1
1982	5,197	294	385	569	610	648	771	602	502	460	7
1981	5,160	296	429	569	621	674	700	582	461	467	4
1980	5,228	266	420	564	633	750	693	629	432	463	2
1979	5,311	288	457	597	624	744	729	627	439	415	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart disease (402)											
Black, male											
1998	3,463	289	335	368	407	403	367	354	262	287	1
1997	3,190	271	334	329	353	400	371	302	198	271	1
1996	3,160	254	309	320	343	340	383	344	232	286	2
1995	3,039	264	295	297	372	376	332	307	226	235	1
1994	2,855	269	244	296	339	339	330	291	219	199	2
1993	2,843	222	274	276	318	373	345	288	209	243	-
1992	2,724	184	237	297	342	347	315	254	214	216	2
1991	2,635	204	219	292	335	333	312	279	197	188	2
1990	2,608	183	219	278	337	364	335	242	206	184	3
1989	2,688	203	224	320	375	361	328	271	176	175	2
1988	2,607	161	220	297	359	366	326	270	187	181	2
1987	2,465	134	240	289	363	347	293	236	184	145	-
1986	2,465	160	239	271	343	353	299	270	174	142	3
1985	2,515	154	225	277	378	350	313	268	172	140	1
1984	2,474	155	226	317	358	322	335	226	163	138	2
1983	2,546	152	226	319	360	321	370	244	182	140	1
1982	2,449	174	212	304	336	320	336	235	175	132	3
1981	2,444	160	246	322	321	341	322	225	149	141	3
1980	2,388	141	223	317	351	371	281	239	142	104	1
1979	2,482	163	248	332	300	352	321	251	173	117	3
Hypertensive heart disease (402)											
Black, female											
1998	3,695	201	205	250	314	324	409	463	435	841	2
1997	3,518	179	201	232	280	343	404	442	447	777	-
1996	3,404	173	200	200	274	295	412	414	455	764	-
1995	3,369	151	196	195	286	337	381	404	457	738	1
1994	3,305	186	193	219	262	329	374	395	429	688	-
1993	3,214	134	194	226	258	337	412	395	463	596	1
1992	3,130	135	175	254	294	347	381	402	393	565	2
1991	3,004	132	148	209	250	355	388	428	372	527	-
1990	3,028	139	165	204	315	355	439	421	317	504	1
1989	2,992	137	159	217	290	360	372	445	362	487	2
1988	3,019	133	172	219	314	358	390	427	376	461	-
1987	2,811	108	171	196	313	357	413	387	321	413	-
1986	2,847	121	165	234	315	309	424	408	325	389	-
1985	2,827	112	176	246	318	334	389	402	353	367	-
1984	2,802	117	170	244	302	370	391	373	339	366	1
1983	2,793	120	168	225	315	344	465	365	335	330	-
1982	2,748	120	173	265	274	328	435	367	327	328	4
1981	2,716	136	183	247	300	333	378	357	312	326	1
1980	2,840	125	197	247	282	379	412	390	290	359	1
1979	2,829	125	209	265	324	392	408	376	266	298	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
All races, both sexes											
1998	2,412	48	59	66	99	164	248	362	471	814	-
1997	2,424	43	40	78	93	148	254	382	450	860	-
1996	2,501	56	46	84	124	155	246	392	428	895	-
1995	2,484	46	54	65	118	155	269	402	466	839	-
1994	2,494	43	60	73	113	201	269	376	474	810	-
1993	2,464	35	50	74	118	175	292	352	498	791	-
1992	2,492	36	42	92	137	192	283	365	479	808	-
1991	2,445	44	50	67	121	201	291	397	438	766	-
1990	2,377	31	54	83	128	228	273	347	467	682	-
1989	2,385	29	42	64	134	187	288	426	428	735	-
1988	2,450	30	35	75	150	208	260	411	466	748	-
1987	2,521	29	46	80	140	190	308	422	448	784	-
1986	2,496	19	45	66	146	219	311	385	432	804	-
1985	2,800	36	52	90	160	209	334	464	534	876	4
1984	2,855	25	45	83	123	266	347	463	554	907	-
1983	3,073	26	43	88	165	257	357	499	578	1,014	-
1982	3,028	26	42	97	144	232	362	520	556	993	1
1981	3,347	29	60	125	141	278	416	555	627	1,051	1
1980	3,711	42	68	128	193	315	450	608	669	1,190	-
1979	3,649	38	65	124	172	330	457	619	657	1,118	1
Hypertensive heart and renal disease (404)											
All races, male											
1998	996	35	39	42	56	93	110	163	168	244	-
1997	1,051	31	28	46	49	78	126	192	179	267	-
1996	1,002	39	30	40	67	69	96	168	180	261	-
1995	997	24	34	26	56	74	121	182	171	269	-
1994	1,066	36	38	40	54	97	129	190	190	243	-
1993	1,023	19	33	41	54	82	137	164	196	243	-
1992	1,024	27	22	50	64	82	127	160	182	274	-
1991	996	30	24	40	56	94	126	183	159	236	-
1990	959	22	32	45	58	115	116	140	170	212	-
1989	954	19	27	37	67	82	132	180	164	210	-
1988	994	19	17	39	67	99	120	178	176	236	-
1987	997	21	33	50	65	95	130	182	165	209	-
1986	1,017	7	28	33	70	117	158	173	179	204	-
1985	1,152	23	29	46	82	112	161	217	209	248	-
1984	1,151	12	30	51	57	143	160	200	222	249	-
1983	1,221	18	20	48	81	133	172	218	204	298	-
1982	1,222	17	24	56	77	95	171	249	189	306	1
1981	1,351	16	32	62	79	140	199	234	236	314	-
1980	1,492	19	34	72	95	145	218	253	241	390	-
1979	1,507	19	33	60	86	164	212	288	244	360	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
All races, female											
1998	1,416	13	20	24	43	71	138	199	303	570	-
1997	1,373	12	12	32	44	70	128	190	271	593	-
1996	1,499	17	16	44	57	86	150	224	248	634	-
1995	1,487	22	20	39	62	81	148	220	295	570	-
1994	1,428	7	22	33	59	104	140	186	284	567	-
1993	1,441	16	17	33	64	93	155	188	302	548	-
1992	1,468	9	20	42	73	110	156	205	297	534	-
1991	1,449	14	26	27	65	107	165	214	279	530	-
1990	1,418	9	22	38	70	113	157	207	297	470	-
1989	1,431	10	15	27	67	105	156	246	264	525	-
1988	1,456	11	18	36	83	109	140	233	290	512	-
1987	1,524	8	13	30	75	95	178	240	283	575	-
1986	1,479	12	17	33	76	102	153	212	253	600	-
1985	1,648	13	23	44	78	97	173	247	325	628	4
1984	1,704	13	15	32	66	123	187	263	332	658	-
1983	1,852	8	23	40	84	124	185	281	374	716	-
1982	1,806	9	18	41	67	137	191	271	367	687	-
1981	1,996	13	28	63	62	138	217	321	391	737	1
1980	2,219	23	34	56	98	170	232	355	428	800	-
1979	2,142	19	32	64	86	166	245	331	413	758	1
Hypertensive heart and renal disease (404)											
White, both sexes											
1998	1,648	15	25	19	40	94	162	246	365	661	-
1997	1,722	15	19	34	48	85	173	283	349	694	-
1996	1,748	23	19	35	64	91	159	255	342	742	-
1995	1,765	18	16	30	60	95	173	304	368	676	-
1994	1,757	15	16	23	58	125	175	281	366	669	-
1993	1,745	13	17	34	68	113	196	250	381	656	-
1992	1,743	13	16	32	65	109	190	264	387	660	-
1991	1,714	10	14	28	64	114	181	286	335	656	-
1990	1,648	12	14	33	70	142	184	250	363	561	-
1989	1,693	5	14	26	76	110	191	322	316	616	-
1988	1,771	11	15	39	82	134	169	298	358	649	-
1987	1,828	9	16	36	67	113	214	321	353	673	-
1986	1,831	4	19	35	84	122	212	294	343	695	-
1985	2,089	9	16	46	87	129	235	365	442	745	1
1984	2,165	12	18	45	65	174	223	346	456	807	-
1983	2,365	11	15	44	91	158	258	391	482	899	-
1982	2,354	7	15	51	88	154	252	406	472	892	1
1981	2,642	12	31	61	81	188	298	450	542	954	-
1980	2,909	15	28	75	116	206	314	464	579	1,094	-
1979	2,880	8	26	63	103	213	346	504	576	1,016	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
White, male											
1998	669	13	15	16	24	59	74	112	136	207	-
1997	741	11	16	21	26	48	88	152	144	218	-
1996	680	18	12	17	32	41	65	117	148	215	-
1995	703	11	11	12	28	48	80	142	139	218	-
1994	744	13	11	13	30	64	88	152	154	199	-
1993	727	8	11	19	34	61	95	125	160	204	-
1992	701	10	10	16	34	48	92	116	150	221	-
1991	682	5	7	20	34	57	76	136	131	200	-
1990	648	9	10	17	31	75	85	95	136	179	-
1989	658	2	6	15	45	55	93	130	115	183	-
1988	702	7	8	17	42	67	76	140	139	196	-
1987	685	5	14	25	31	61	95	135	126	175	-
1986	721	-	12	17	46	70	112	139	139	171	-
1985	863	8	8	26	48	75	130	176	168	217	-
1984	853	6	12	29	34	103	108	152	181	215	-
1983	923	10	5	23	48	84	134	177	174	259	-
1982	948	4	9	35	50	65	136	195	165	277	1
1981	1,059	7	20	32	47	111	142	194	210	279	-
1980	1,175	6	15	50	69	108	162	199	211	347	-
1979	1,179	1	16	35	51	112	169	239	216	323	-
Hypertensive heart and renal disease (404)											
White, female											
1998	979	2	10	3	16	35	88	134	229	454	-
1997	981	4	3	13	22	37	85	131	205	476	-
1996	1,068	5	7	18	32	50	94	138	194	527	-
1995	1,062	7	5	18	32	47	93	162	229	458	-
1994	1,013	2	5	10	28	61	87	129	212	470	-
1993	1,018	5	6	15	34	52	101	125	221	452	-
1992	1,042	3	6	16	31	61	98	148	237	439	-
1991	1,032	5	7	8	30	57	105	150	204	456	-
1990	1,000	3	4	16	39	67	99	155	227	382	-
1989	1,035	3	8	11	31	55	98	192	201	433	-
1988	1,069	4	7	22	40	67	93	158	219	453	-
1987	1,143	4	2	11	36	52	119	186	227	498	-
1986	1,110	4	7	18	38	52	100	155	204	524	-
1985	1,226	1	8	20	39	54	105	189	274	528	1
1984	1,312	6	6	16	31	71	115	194	275	592	-
1983	1,442	1	10	21	43	74	124	214	308	640	-
1982	1,406	3	6	16	38	89	116	211	307	615	-
1981	1,583	5	11	29	34	77	156	256	332	675	-
1980	1,734	9	13	25	47	98	152	265	368	747	-
1979	1,701	7	10	28	52	101	177	265	360	693	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
All other, both sexes											
1998	764	33	34	47	59	70	86	116	106	153	-
1997	702	28	21	44	45	63	81	99	101	166	-
1996	753	33	27	49	60	64	87	137	86	153	-
1995	719	28	38	35	58	60	96	98	98	163	-
1994	737	28	44	50	55	76	94	95	108	141	-
1993	719	22	33	40	50	62	96	102	117	135	-
1992	749	23	26	60	72	83	93	101	92	148	-
1991	731	34	36	39	57	87	110	111	103	110	-
1990	729	19	40	50	58	86	89	97	104	121	-
1989	692	24	28	38	58	77	97	104	112	119	-
1988	679	19	20	36	68	74	91	113	108	99	-
1987	693	20	30	44	73	77	94	101	95	111	-
1986	665	15	26	31	62	97	99	91	89	109	-
1985	711	27	36	44	73	80	99	99	92	131	3
1984	690	13	27	38	58	92	124	117	98	100	-
1983	708	15	28	44	74	99	99	108	96	115	-
1982	674	19	27	46	56	78	110	114	84	101	-
1981	705	17	29	64	60	90	118	105	85	97	1
1980	802	27	40	53	77	109	136	144	90	96	-
1979	769	30	39	61	69	117	111	115	81	102	1
Hypertensive heart and renal disease (404)											
All other, male											
1998	327	22	24	26	32	34	36	51	32	37	-
1997	310	20	12	25	23	30	38	40	35	49	-
1996	322	21	18	23	35	28	31	51	32	46	-
1995	294	13	23	14	28	26	41	40	32	51	-
1994	322	23	27	27	24	33	41	38	36	44	-
1993	296	11	22	22	20	21	42	39	36	39	-
1992	323	17	12	34	30	34	35	44	32	53	-
1991	314	25	17	20	22	37	50	47	28	36	-
1990	311	13	22	28	27	40	31	45	34	33	-
1989	296	17	21	22	22	27	39	50	49	27	-
1988	292	12	9	22	25	32	44	38	37	40	-
1987	312	16	19	25	34	34	35	47	39	34	-
1986	296	7	16	16	24	47	46	34	40	33	-
1985	289	15	21	20	34	37	31	41	41	31	-
1984	298	6	18	22	23	40	52	48	41	34	-
1983	298	8	15	25	33	49	38	41	30	39	-
1982	274	13	15	21	27	30	35	54	24	29	-
1981	292	9	12	30	32	29	57	40	26	35	-
1980	317	13	19	22	26	37	56	54	30	43	-
1979	328	18	17	25	35	52	43	49	28	37	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
All other, female											
1998	437	11	10	21	27	36	50	65	74	116	-
1997	392	8	9	19	22	33	43	59	66	117	-
1996	431	12	9	26	25	36	56	86	54	107	-
1995	425	15	15	21	30	34	55	58	66	112	-
1994	415	5	17	23	31	43	53	57	72	97	-
1993	423	11	11	18	30	41	54	63	81	96	-
1992	426	6	14	26	42	49	58	57	60	95	-
1991	417	9	19	19	35	50	60	64	75	74	-
1990	418	6	18	22	31	46	58	52	70	88	-
1989	396	7	7	16	36	50	58	54	63	92	-
1988	387	7	11	14	43	42	47	75	71	59	-
1987	381	4	11	19	39	43	59	54	56	77	-
1986	369	8	10	15	38	50	53	57	49	76	-
1985	422	12	15	24	39	43	68	58	51	100	3
1984	392	7	9	16	35	52	72	69	57	66	-
1983	410	7	13	19	41	50	61	67	66	76	-
1982	400	6	12	25	29	48	75	60	60	72	-
1981	413	8	17	34	28	61	61	65	59	62	1
1980	485	14	21	31	51	72	80	90	60	53	-
1979	441	12	22	36	34	65	68	66	53	65	1
Hypertensive heart and renal disease (404)											
Black, both sexes											
1998	700	32	32	44	55	66	76	104	94	139	-
1997	653	25	21	42	41	57	75	90	94	154	-
1996	689	33	27	48	56	59	79	125	75	133	-
1995	672	26	38	33	53	58	89	89	90	152	-
1994	679	26	42	46	47	75	88	88	96	126	-
1993	683	21	33	37	49	60	92	98	106	128	-
1992	707	23	25	54	68	79	88	91	88	140	-
1991	695	34	35	38	56	84	102	105	97	102	-
1990	707	19	40	50	56	82	87	92	97	119	-
1989	656	24	28	36	53	73	95	99	103	112	-
1988	658	18	19	35	67	71	90	111	104	92	-
1987	662	19	29	44	72	74	89	96	86	106	-
1986	634	15	26	31	61	94	93	84	84	103	-
1985	675	26	35	44	70	76	95	94	84	123	2
1984	665	11	26	38	58	87	120	116	94	92	-
1983	686	15	28	43	71	97	96	103	93	110	-
1982	653	19	25	45	54	74	110	108	80	99	-
1981	683	16	28	62	60	90	114	102	83	90	1
1980	777	27	40	51	76	106	129	142	86	90	-
1979	748	29	39	61	67	111	109	111	79	99	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hypertensive heart and renal disease (404)											
Black, male											
1998	293	21	22	24	31	33	30	45	26	30	-
1997	287	19	12	25	21	27	35	37	32	41	-
1996	298	21	18	23	33	28	29	46	25	41	-
1995	274	11	23	14	25	25	38	37	29	46	-
1994	294	22	26	25	21	32	37	34	30	38	-
1993	276	11	22	19	20	19	41	37	31	35	-
1992	294	17	11	29	28	30	32	36	29	50	-
1991	298	25	16	19	22	35	47	44	27	32	-
1990	300	13	22	28	25	37	31	43	32	31	-
1989	280	17	21	21	20	25	38	48	43	25	-
1988	277	11	9	21	24	30	44	36	34	35	-
1987	299	15	19	25	34	32	34	44	35	32	-
1986	279	7	16	16	23	46	43	29	36	32	-
1985	274	14	21	20	33	35	28	40	37	28	-
1984	287	5	17	22	23	38	50	48	39	31	-
1983	289	8	15	25	30	48	37	40	29	37	-
1982	266	13	14	20	26	29	35	51	23	29	-
1981	280	9	12	29	32	29	54	38	25	30	-
1980	305	13	19	20	25	36	54	53	27	41	-
1979	318	17	17	25	34	50	42	47	26	37	-
Hypertensive heart and renal disease (404)											
Black, female											
1998	407	11	10	20	24	33	46	59	68	109	-
1997	366	6	9	17	20	30	40	53	62	113	-
1996	391	12	9	25	23	31	50	79	50	92	-
1995	398	15	15	19	28	33	51	52	61	106	-
1994	385	4	16	21	26	43	51	54	66	88	-
1993	407	10	11	18	29	41	51	61	75	93	-
1992	413	6	14	25	40	49	56	55	59	90	-
1991	397	9	19	19	34	49	55	61	70	70	-
1990	407	6	18	22	31	45	56	49	65	88	-
1989	376	7	7	15	33	48	57	51	60	87	-
1988	381	7	10	14	43	41	46	75	70	57	-
1987	363	4	10	19	38	42	55	52	51	74	-
1986	355	8	10	15	38	48	50	55	48	71	-
1985	401	12	14	24	37	41	67	54	47	95	2
1984	378	6	9	16	35	49	70	68	55	61	-
1983	397	7	13	18	41	49	59	63	64	73	-
1982	387	6	11	25	28	45	75	57	57	70	-
1981	403	7	16	33	28	61	60	64	58	60	1
1980	472	14	21	31	51	70	75	89	59	49	-
1979	430	12	22	36	33	61	67	64	53	62	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Ischemic heart diseases (410-414,429.2)											
All races, both sexes											
1998	526,617	9,388	14,823	20,001	28,177	41,355	60,885	78,875	90,037	174,097	41
1997	533,061	9,761	15,060	20,191	29,221	43,524	62,509	80,361	90,140	173,376	54
1996	544,638	10,168	14,802	20,041	30,976	45,762	65,742	81,744	92,511	173,601	58
1995	552,464	9,750	14,998	20,400	31,349	48,222	68,069	82,186	93,498	174,538	80
1994	552,678	9,621	14,747	20,715	32,415	49,551	69,017	82,366	93,616	171,387	60
1993	562,703	9,408	14,296	21,023	34,112	52,312	70,430	84,947	95,211	171,627	64
1992	549,007	9,300	13,726	21,093	34,562	52,060	70,302	83,823	92,646	162,235	74
1991	554,817	8,972	13,506	21,708	36,229	54,234	71,003	85,756	93,314	160,924	88
1990	558,291	8,817	13,638	22,382	37,752	55,897	71,579	87,643	93,448	158,357	81
1989	568,901	8,858	14,052	23,426	39,576	57,932	72,679	89,500	94,950	159,032	76
1988	582,419	8,888	14,240	24,881	41,378	59,662	76,407	91,481	96,063	160,332	82
1987	584,580	8,888	15,193	26,151	43,029	60,781	77,829	91,570	94,375	157,199	85
1986	593,396	8,857	15,670	27,678	45,348	61,947	79,790	93,004	94,841	156,245	285
1985	608,372	9,327	16,771	29,693	47,217	63,893	82,583	95,089	96,536	157,418	231
1984	613,178	9,634	17,428	31,004	48,992	64,887	84,182	95,883	96,835	154,724	83
1983	624,910	9,819	18,878	32,695	49,943	67,317	86,305	97,077	97,940	155,325	103
1982	620,841	10,278	19,639	33,564	50,727	68,401	86,076	96,217	96,531	149,600	135
1981	624,254	10,957	20,673	34,944	51,382	69,661	86,332	94,854	97,127	148,587	124
1980	635,677	11,162	21,680	36,631	51,849	71,497	87,751	97,554	99,094	148,795	120
1979	617,454	11,702	22,381	36,773	52,431	69,923	85,846	94,973	95,391	138,213	111
Ischemic heart diseases (410-414,429.2)											
All races, male											
1998	267,182	7,365	11,426	14,644	19,501	27,242	37,159	43,977	43,341	55,650	31
1997	271,059	7,718	11,529	14,721	20,352	28,741	38,274	44,543	43,117	55,169	41
1996	276,545	7,952	11,254	14,626	21,631	30,154	40,019	45,231	43,542	54,980	40
1995	280,110	7,708	11,564	14,949	21,831	31,624	41,716	45,129	43,846	54,405	43
1994	280,975	7,572	11,332	15,122	22,683	32,771	42,234	45,020	43,399	53,687	40
1993	286,382	7,471	10,964	15,424	23,834	34,602	42,751	46,437	44,145	53,454	36
1992	280,945	7,450	10,503	15,445	24,240	34,328	42,701	45,728	42,506	50,781	47
1991	284,012	7,107	10,354	15,942	25,323	35,745	43,216	46,356	42,415	50,286	61
1990	286,592	7,002	10,481	16,506	26,396	36,722	43,372	47,010	42,515	49,536	46
1989	292,072	7,113	10,816	17,343	27,630	38,127	43,929	47,714	42,755	49,500	49
1988	299,808	7,080	10,888	18,306	28,815	39,206	46,144	48,811	43,149	50,123	49
1987	302,713	7,072	11,721	19,322	30,045	39,697	46,971	48,685	42,199	49,300	52
1986	309,249	7,035	12,210	20,710	31,722	40,651	48,064	49,309	42,242	49,329	103
1985	320,412	7,510	13,068	22,144	33,260	41,970	49,885	50,368	43,512	50,832	88
1984	323,878	7,714	13,625	22,992	34,550	42,585	50,300	50,764	43,129	50,442	53
1983	332,378	7,918	14,687	24,396	35,093	44,474	52,001	51,438	43,363	51,237	58
1982	333,524	8,268	15,399	25,190	35,879	45,415	51,799	50,919	42,702	50,037	79
1981	337,322	8,823	16,166	26,359	36,296	46,069	52,107	50,272	43,127	50,218	68
1980	344,600	8,973	17,050	27,647	36,731	47,405	52,727	51,144	43,675	51,417	73
1979	339,843	9,500	17,670	27,947	37,605	46,762	51,959	49,946	42,175	48,331	76

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Ischemic heart diseases (410-414,429.2)											
Black, male											
1998	24,398	1,167	1,637	2,042	2,526	3,001	3,379	3,402	2,660	3,372	12
1997	24,489	1,233	1,633	2,038	2,660	2,946	3,458	3,329	2,693	3,307	10
1996	24,530	1,173	1,585	1,995	2,717	3,053	3,504	3,350	2,722	3,262	8
1995	25,542	1,221	1,623	2,100	2,665	3,351	3,775	3,428	2,774	3,368	9
1994	24,923	1,096	1,658	1,975	2,624	3,337	3,660	3,205	2,860	3,262	14
1993	25,740	1,017	1,568	2,140	2,883	3,530	3,736	3,244	2,927	3,367	15
1992	24,794	1,042	1,447	2,024	2,741	3,435	3,677	3,347	2,718	3,141	16
1991	25,310	1,003	1,481	2,080	2,881	3,566	3,712	3,276	2,882	3,154	19
1990	25,060	960	1,474	2,154	2,835	3,578	3,572	3,462	2,725	3,139	17
1989	25,776	1,075	1,564	2,187	3,150	3,770	3,517	3,534	2,725	3,035	17
1988	25,676	1,021	1,440	2,230	3,166	3,659	3,574	3,597	2,832	2,937	15
1987	25,738	1,008	1,536	2,118	3,138	3,697	3,753	3,524	2,765	2,918	16
1986	25,853	909	1,583	2,298	3,264	3,620	3,693	3,520	2,657	3,011	18
1985	26,331	1,022	1,588	2,391	3,304	3,679	3,934	3,483	2,798	2,858	15
1984	26,240	1,040	1,663	2,422	3,407	3,560	3,950	3,428	2,771	2,766	15
1983	26,822	1,063	1,809	2,588	3,375	3,646	4,107	3,486	2,819	2,809	15
1982	26,457	1,084	1,751	2,678	3,272	3,721	3,939	3,408	2,653	2,738	23
1981	26,947	1,159	1,854	2,767	3,390	3,774	3,995	3,339	2,713	2,702	24
1980	27,625	1,179	1,944	2,796	3,356	3,962	3,931	3,654	2,693	2,857	30
1979	26,773	1,180	2,045	2,888	3,254	3,885	3,877	3,525	2,388	2,496	21
Ischemic heart diseases (410-414,429.2)											
Black, female											
1998	25,718	583	783	1,122	1,675	2,279	3,205	3,679	3,908	7,890	3
1997	25,744	535	812	1,199	1,657	2,290	3,302	3,673	3,867	7,835	7
1996	26,218	573	858	1,183	1,808	2,408	3,346	3,739	4,026	7,677	4
1995	26,452	573	784	1,138	1,732	2,589	3,431	3,728	4,239	7,658	10
1994	26,093	536	780	1,208	1,719	2,636	3,366	3,749	4,110	7,353	8
1993	26,920	513	801	1,244	1,873	2,730	3,455	3,943	4,209	7,550	10
1992	25,490	477	798	1,241	1,884	2,644	3,270	3,684	3,971	6,936	9
1991	25,788	476	741	1,216	1,934	2,805	3,410	3,934	4,069	6,661	5
1990	25,321	473	758	1,155	2,052	2,944	3,214	3,895	3,936	6,404	12
1989	25,936	459	750	1,276	2,122	2,883	3,401	4,145	3,928	6,448	6
1988	25,842	473	820	1,277	2,157	2,864	3,355	4,158	3,999	6,224	10
1987	25,299	463	774	1,327	2,122	2,886	3,397	3,931	3,838	6,081	8
1986	25,362	475	807	1,294	2,190	2,899	3,478	4,040	3,779	5,898	13
1985	25,142	459	816	1,489	2,202	2,910	3,656	3,983	3,705	5,434	8
1984	24,929	456	864	1,527	2,220	2,789	3,703	3,853	3,841	5,177	8
1983	25,415	459	924	1,593	2,310	2,907	3,875	3,957	3,768	5,103	8
1982	24,235	535	976	1,504	2,302	2,849	3,545	3,646	3,527	4,851	13
1981	24,406	513	945	1,529	2,290	2,995	3,655	3,624	3,514	4,846	18
1980	25,028	561	1,014	1,638	2,308	3,133	3,702	3,939	3,439	4,756	15
1979	23,262	564	1,029	1,531	2,174	3,039	3,470	3,863	2,877	4,175	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute myocardial infarction (410)											
All races, female											
1998	96,556	958	1,592	2,543	3,976	6,259	10,255	14,651	17,985	37,383	3
1997	97,041	968	1,667	2,652	4,140	6,694	10,542	14,952	18,238	36,217	4
1996	100,793	1,065	1,708	2,604	4,482	7,124	11,464	15,390	19,195	36,748	3
1995	102,242	961	1,673	2,615	4,579	7,764	11,830	15,728	19,248	36,839	7
1994	104,105	1,034	1,705	2,793	4,772	7,982	12,343	16,106	19,866	36,470	2
1993	105,724	980	1,629	2,784	4,994	8,368	12,678	16,731	20,241	36,329	4
1992	106,262	976	1,656	2,949	5,167	8,624	13,195	17,078	20,742	34,851	4
1991	109,111	977	1,687	3,104	5,554	9,164	13,441	18,013	21,079	35,111	10
1990	109,525	942	1,607	3,063	5,847	9,698	13,811	18,652	21,125	33,858	7
1989	112,652	918	1,732	3,179	6,244	10,237	14,215	19,386	21,726	34,101	9
1988	112,066	973	1,758	3,460	6,627	10,529	14,965	19,356	21,415	31,994	10
1987	113,290	1,024	1,934	3,651	6,969	11,029	15,424	19,848	21,477	30,936	10
1986	115,207	1,046	1,915	3,881	7,375	11,421	16,140	20,467	21,433	30,471	39
1985	118,339	1,023	2,123	4,350	7,788	11,923	16,903	21,334	21,434	30,370	24
1984	119,188	1,128	2,196	4,641	8,106	12,151	17,620	21,338	21,693	29,275	9
1983	120,621	1,143	2,427	4,801	8,531	12,564	17,943	21,654	21,478	28,985	13
1982	120,659	1,178	2,541	5,023	8,564	12,898	18,210	21,666	21,313	28,119	18
1981	119,606	1,299	2,691	5,090	8,715	13,322	17,877	20,746	21,266	27,527	20
1980	121,099	1,343	2,703	5,295	8,736	13,561	18,270	21,616	21,445	27,053	15
1979	119,408	1,346	2,938	5,525	8,907	13,283	17,999	21,396	21,084	25,759	12
Acute myocardial infarction (410)											
White, both sexes											
1998	181,298	3,521	5,687	7,964	10,908	15,744	22,670	28,806	31,265	51,520	5
1997	184,132	3,720	5,907	7,983	11,376	16,883	23,553	29,616	31,672	50,235	14
1996	191,235	3,986	5,982	7,973	12,272	17,986	25,340	30,487	32,935	50,840	9
1995	195,726	3,909	6,133	8,280	12,760	19,176	26,497	31,103	33,427	50,881	11
1994	199,856	3,945	6,037	8,668	13,403	20,177	27,604	31,788	34,147	50,570	8
1993	204,386	3,917	5,915	8,666	14,246	21,406	28,275	33,314	34,909	50,229	8
1992	206,629	3,949	5,866	8,995	14,621	21,762	29,466	33,874	35,442	48,979	8
1991	212,058	3,887	5,922	9,533	15,498	23,266	30,098	35,306	35,920	48,997	18
1990	216,252	3,910	5,954	9,952	16,570	24,423	30,963	36,660	36,484	47,720	9
1989	223,228	3,940	6,194	10,452	17,669	25,854	32,239	37,908	37,287	47,979	16
1988	225,262	3,980	6,443	11,088	18,571	26,873	33,823	38,247	36,758	45,567	14
1987	230,809	4,147	7,127	12,207	19,958	27,988	35,151	39,158	36,696	44,232	16
1986	238,218	4,247	7,417	13,219	21,488	29,379	36,840	40,516	36,917	43,749	54
1985	250,527	4,504	8,312	14,839	23,261	31,351	39,055	42,580	37,535	44,515	34
1984	255,496	4,843	8,850	15,845	24,730	32,261	40,414	43,141	37,745	43,135	20
1983	262,549	5,056	9,765	16,884	25,846	34,052	41,988	43,607	37,550	43,134	19
1982	267,971	5,414	10,558	17,926	26,936	35,655	42,942	43,933	37,410	42,275	33
1981	269,389	5,934	11,124	18,745	27,389	36,459	42,851	42,868	37,432	41,567	32
1980	275,957	6,131	11,815	20,003	28,289	37,938	43,849	43,629	37,860	41,514	14
1979	277,070	6,554	12,565	20,869	29,443	37,792	43,807	43,528	37,389	39,846	30

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute myocardial infarction (410)											
All other, both sexes											
1998	22,253	841	1,147	1,450	1,905	2,367	2,949	3,212	3,005	4,518	2
1997	22,080	854	1,144	1,528	1,926	2,478	3,051	3,081	2,892	4,266	2
1996	22,297	839	1,122	1,498	2,048	2,519	3,093	3,176	2,891	4,269	4
1995	22,503	817	1,103	1,441	1,991	2,726	3,176	3,200	2,981	4,181	7
1994	22,543	835	1,166	1,492	1,998	2,776	3,292	3,158	2,926	3,982	3
1993	23,070	752	1,105	1,570	2,165	2,821	3,254	3,209	3,068	4,162	6
1992	22,754	778	1,110	1,633	2,186	2,841	3,254	3,225	2,963	3,840	7
1991	23,197	739	1,036	1,644	2,363	2,974	3,310	3,407	3,024	3,757	6
1990	22,756	691	1,106	1,628	2,307	3,124	3,229	3,452	2,858	3,508	6
1989	23,722	749	1,157	1,723	2,539	3,317	3,279	3,578	2,913	3,598	6
1988	22,688	684	1,077	1,679	2,537	3,152	3,191	3,462	2,876	3,139	5
1987	22,733	757	1,141	1,643	2,559	3,207	3,340	3,319	2,741	3,131	9
1986	22,784	723	1,218	1,749	2,683	3,173	3,365	3,420	2,658	2,858	4
1985	23,672	773	1,180	1,980	2,803	3,324	3,652	3,436	2,698	2,864	7
1984	23,626	815	1,351	2,042	2,844	3,222	3,628	3,294	2,753	2,725	7
1983	23,751	811	1,401	2,122	2,849	3,332	3,798	3,304	2,627	2,547	14
1982	23,060	830	1,393	2,210	2,848	3,299	3,646	3,072	2,378	2,391	12
1981	23,115	917	1,455	2,264	2,825	3,437	3,564	2,961	2,381	2,335	16
1980	23,627	941	1,545	2,258	2,802	3,462	3,571	3,293	2,290	2,426	16
1979	23,392	951	1,624	2,350	2,862	3,510	3,417	3,297	2,048	2,256	12
Acute myocardial infarction (410)											
All other, male											
1998	11,074	567	775	925	1,152	1,297	1,497	1,574	1,234	1,474	1
1997	10,933	590	784	955	1,142	1,377	1,532	1,398	1,191	1,393	1
1996	11,041	571	740	937	1,202	1,371	1,546	1,482	1,215	1,401	2
1995	11,271	551	719	923	1,191	1,463	1,645	1,528	1,232	1,410	4
1994	11,351	575	795	884	1,152	1,497	1,747	1,444	1,268	1,383	2
1993	11,601	498	744	980	1,305	1,537	1,698	1,449	1,316	1,403	4
1992	11,593	538	718	992	1,279	1,582	1,693	1,530	1,270	1,358	4
1991	11,713	517	676	1,014	1,374	1,613	1,697	1,583	1,276	1,312	4
1990	11,633	481	741	1,049	1,269	1,686	1,674	1,651	1,226	1,260	2
1989	12,070	524	753	1,064	1,480	1,824	1,657	1,651	1,244	1,262	2
1988	11,606	458	673	1,030	1,425	1,754	1,653	1,637	1,251	1,107	2
1987	11,819	507	737	1,022	1,474	1,759	1,774	1,595	1,180	1,124	7
1986	11,835	469	789	1,096	1,541	1,735	1,716	1,619	1,167	1,010	2
1985	12,549	521	750	1,185	1,643	1,875	1,930	1,680	1,227	1,056	3
1984	12,679	549	888	1,214	1,710	1,812	1,865	1,658	1,233	1,056	5
1983	12,781	556	935	1,302	1,677	1,881	2,008	1,600	1,173	1,003	10
1982	12,624	565	879	1,430	1,648	1,858	1,956	1,542	1,089	971	8
1981	12,796	649	958	1,450	1,665	1,938	1,962	1,494	1,079	888	7
1980	13,015	642	1,014	1,427	1,647	1,954	1,941	1,648	1,022	985	9
1979	13,164	662	1,059	1,537	1,707	2,029	1,891	1,656	951	917	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute myocardial infarction (410)											
All other, female											
1998	11,179	274	372	525	753	1,070	1,452	1,638	1,771	3,044	1
1997	11,147	264	360	573	784	1,101	1,519	1,683	1,701	2,873	1
1996	11,256	268	382	561	846	1,148	1,547	1,694	1,676	2,868	2
1995	11,232	266	384	518	800	1,263	1,531	1,672	1,749	2,771	3
1994	11,192	260	371	608	846	1,279	1,545	1,714	1,658	2,599	1
1993	11,469	254	361	590	860	1,284	1,556	1,760	1,752	2,759	2
1992	11,161	240	392	641	907	1,259	1,561	1,695	1,693	2,482	3
1991	11,484	222	360	630	989	1,361	1,613	1,824	1,748	2,445	2
1990	11,123	210	365	579	1,038	1,438	1,555	1,801	1,632	2,248	4
1989	11,652	225	404	659	1,059	1,493	1,622	1,927	1,669	2,336	4
1988	11,082	226	404	649	1,112	1,398	1,538	1,825	1,625	2,032	3
1987	10,914	250	404	621	1,085	1,448	1,566	1,724	1,561	2,007	2
1986	10,949	254	429	653	1,142	1,438	1,649	1,801	1,491	1,848	2
1985	11,123	252	430	795	1,160	1,449	1,722	1,756	1,471	1,808	4
1984	10,947	266	463	828	1,134	1,410	1,763	1,636	1,520	1,669	2
1983	10,970	255	466	820	1,172	1,451	1,790	1,704	1,454	1,544	4
1982	10,436	265	514	780	1,200	1,441	1,690	1,530	1,289	1,420	4
1981	10,319	268	497	814	1,160	1,499	1,602	1,467	1,302	1,447	9
1980	10,612	299	531	831	1,155	1,508	1,630	1,645	1,268	1,441	7
1979	10,228	289	565	813	1,155	1,481	1,526	1,641	1,097	1,339	4
Acute myocardial infarction (410)											
Black, both sexes											
1998	19,198	753	993	1,265	1,650	2,042	2,532	2,736	2,580	3,874	1
1997	19,244	739	1,012	1,326	1,678	2,123	2,671	2,686	2,501	3,736	2
1996	19,492	750	975	1,334	1,801	2,216	2,690	2,747	2,505	3,733	4
1995	19,766	732	956	1,259	1,736	2,415	2,782	2,786	2,608	3,700	5
1994	19,811	732	1,043	1,324	1,763	2,437	2,860	2,765	2,559	3,506	3
1993	20,399	654	966	1,410	1,925	2,506	2,887	2,806	2,710	3,666	6
1992	20,104	696	979	1,429	1,946	2,516	2,866	2,825	2,607	3,406	6
1991	20,654	668	917	1,470	2,103	2,648	2,952	3,013	2,687	3,351	5
1990	20,436	620	996	1,447	2,102	2,828	2,887	3,082	2,566	3,139	6
1989	21,198	668	1,022	1,546	2,288	2,984	2,936	3,175	2,576	3,209	5
1988	20,393	615	983	1,514	2,268	2,850	2,861	3,108	2,563	2,807	5
1987	20,453	682	1,035	1,481	2,313	2,918	3,006	2,984	2,456	2,769	8
1986	20,684	641	1,100	1,595	2,431	2,926	3,069	3,098	2,359	2,599	4
1985	21,447	691	1,062	1,805	2,560	3,029	3,320	3,070	2,423	2,609	6
1984	21,498	737	1,240	1,876	2,623	2,953	3,323	2,981	2,465	2,435	4
1983	21,775	749	1,286	1,959	2,639	3,073	3,489	3,008	2,379	2,318	12
1982	21,099	758	1,284	2,060	2,656	3,038	3,322	2,773	2,152	2,151	9
1981	21,363	839	1,363	2,112	2,644	3,198	3,299	2,712	2,175	2,124	14
1980	21,892	867	1,443	2,117	2,642	3,222	3,291	3,027	2,100	2,221	15
1979	21,714	898	1,530	2,222	2,677	3,282	3,152	3,024	1,862	2,060	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute myocardial infarction (410)											
Black, male											
1998	9,273	491	663	787	975	1,113	1,258	1,290	1,008	1,184	-
1997	9,215	496	678	802	970	1,161	1,303	1,189	984	1,119	1
1996	9,357	492	624	821	1,027	1,189	1,323	1,243	1,009	1,140	2
1995	9,594	487	615	798	1,009	1,269	1,407	1,297	1,017	1,168	2
1994	9,736	489	693	775	993	1,297	1,468	1,235	1,080	1,176	2
1993	9,933	419	637	864	1,147	1,334	1,476	1,212	1,107	1,148	4
1992	9,970	468	621	844	1,110	1,371	1,464	1,310	1,088	1,134	3
1991	10,158	464	582	892	1,196	1,419	1,487	1,361	1,071	1,113	3
1990	10,260	420	651	912	1,146	1,515	1,474	1,440	1,069	1,101	2
1989	10,545	458	655	937	1,318	1,613	1,460	1,425	1,055	1,077	2
1988	10,197	404	600	916	1,260	1,544	1,462	1,443	1,057	950	2
1987	10,344	447	655	907	1,308	1,567	1,564	1,361	1,018	941	6
1986	10,525	404	700	979	1,378	1,574	1,541	1,418	997	900	2
1985	11,103	458	656	1,071	1,473	1,691	1,706	1,446	1,051	934	3
1984	11,321	495	808	1,099	1,556	1,632	1,670	1,458	1,059	923	3
1983	11,529	513	848	1,181	1,527	1,715	1,809	1,420	1,040	898	9
1982	11,345	507	797	1,325	1,521	1,692	1,729	1,328	966	865	5
1981	11,638	584	885	1,332	1,543	1,784	1,770	1,330	962	797	7
1980	11,834	581	933	1,316	1,535	1,788	1,736	1,478	922	872	9
1979	12,016	619	976	1,435	1,579	1,854	1,717	1,468	851	816	7
Acute myocardial infarction (410)											
Black, female											
1998	9,925	262	330	478	675	929	1,274	1,446	1,572	2,690	1
1997	10,029	243	334	524	708	962	1,368	1,497	1,517	2,617	1
1996	10,135	258	351	513	774	1,027	1,367	1,504	1,496	2,593	2
1995	10,172	245	341	461	727	1,146	1,375	1,489	1,591	2,532	3
1994	10,075	243	350	549	770	1,140	1,392	1,530	1,479	2,330	1
1993	10,466	235	329	546	778	1,172	1,411	1,594	1,603	2,518	2
1992	10,134	228	358	585	836	1,145	1,402	1,515	1,519	2,272	3
1991	10,496	204	335	578	907	1,229	1,465	1,652	1,616	2,238	2
1990	10,176	200	345	535	956	1,313	1,413	1,642	1,497	2,038	4
1989	10,653	210	367	609	970	1,371	1,476	1,750	1,521	2,132	3
1988	10,196	211	383	598	1,008	1,306	1,399	1,665	1,506	1,857	3
1987	10,109	235	380	574	1,005	1,351	1,442	1,623	1,438	1,828	2
1986	10,159	237	400	616	1,053	1,352	1,528	1,680	1,362	1,699	2
1985	10,344	233	406	734	1,087	1,338	1,614	1,624	1,372	1,675	3
1984	10,177	242	432	777	1,067	1,321	1,653	1,523	1,406	1,512	1
1983	10,246	236	438	778	1,112	1,358	1,680	1,588	1,339	1,420	3
1982	9,754	251	487	735	1,135	1,346	1,593	1,445	1,186	1,286	4
1981	9,725	255	478	780	1,101	1,414	1,529	1,382	1,213	1,327	7
1980	10,058	286	510	801	1,107	1,434	1,555	1,549	1,178	1,349	6
1979	9,698	279	554	787	1,098	1,428	1,435	1,556	1,011	1,244	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
All races, both sexes											
1998	2,907	111	152	179	225	262	307	378	393	781	-
1997	3,046	118	196	190	266	292	350	373	351	787	4
1996	2,889	104	156	185	231	289	318	391	327	745	2
1995	2,740	91	159	175	221	265	332	304	376	707	7
1994	2,862	106	157	150	246	257	338	327	399	750	4
1993	2,968	110	149	175	253	317	358	374	414	664	5
1992	3,179	128	147	199	286	355	389	405	404	710	5
1991	3,300	125	159	211	317	387	416	409	420	690	7
1990	3,333	137	183	228	332	402	409	466	398	642	1
1989	3,376	108	138	225	358	406	416	442	437	718	1
1988	3,408	112	157	222	319	413	443	459	428	708	2
1987	3,532	121	181	266	362	441	422	521	427	654	3
1986	3,856	126	186	319	440	485	531	500	445	647	2
1985	3,969	125	249	321	477	500	530	500	464	664	2
1984	4,034	108	228	334	461	528	537	530	482	671	2
1983	4,059	143	264	324	468	538	549	526	444	657	1
1982	4,120	159	253	401	474	521	600	500	447	593	3
1981	4,549	145	282	435	538	594	613	550	528	689	-
1980	4,773	166	314	453	534	641	630	605	559	695	4
1979	4,762	162	305	456	579	627	637	637	526	664	2
Other acute ischemic heart diseases (411)											
All races, male											
1998	1,563	89	120	129	164	177	187	204	166	234	-
1997	1,691	100	153	140	188	191	221	201	156	244	4
1996	1,591	79	119	134	174	189	200	211	153	220	-
1995	1,517	74	130	138	153	186	210	164	170	211	5
1994	1,562	86	117	105	188	177	209	174	175	236	1
1993	1,662	91	121	124	177	211	223	187	198	215	2
1992	1,770	106	118	149	204	215	248	205	180	218	3
1991	1,941	106	121	156	224	283	277	212	188	229	6
1990	1,968	114	142	175	231	289	265	250	189	206	1
1989	1,902	86	105	176	254	279	255	225	200	224	1
1988	1,955	88	115	167	238	275	276	263	183	228	1
1987	2,073	95	143	196	265	300	261	300	207	208	-
1986	2,352	103	150	243	327	336	337	300	216	198	2
1985	2,381	102	199	247	338	342	343	273	213	216	2
1984	2,459	89	176	253	336	390	336	279	230	241	2
1983	2,501	113	200	254	335	393	341	301	225	232	-
1982	2,611	128	205	313	352	371	386	285	213	213	3
1981	2,801	119	225	332	399	423	390	293	242	251	-
1980	2,975	144	231	350	399	450	414	326	260	257	3
1979	2,992	135	239	362	434	435	399	356	244	249	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
All races, female											
1998	1,344	22	32	50	61	85	120	174	227	547	-
1997	1,355	18	43	50	78	101	129	172	195	543	-
1996	1,298	25	37	51	57	100	118	180	174	525	2
1995	1,223	17	29	37	68	79	122	140	206	496	2
1994	1,300	20	40	45	58	80	129	153	224	514	3
1993	1,306	19	28	51	76	106	135	187	216	449	3
1992	1,409	22	29	50	82	140	141	200	224	492	2
1991	1,359	19	38	55	93	104	139	197	232	461	1
1990	1,365	23	41	53	101	113	144	216	209	436	-
1989	1,474	22	33	49	104	127	161	217	237	494	-
1988	1,453	24	42	55	81	138	167	196	245	480	1
1987	1,459	26	38	70	97	141	161	221	220	446	3
1986	1,504	23	36	76	113	149	194	200	229	449	-
1985	1,588	23	50	74	139	158	187	227	251	448	-
1984	1,575	19	52	81	125	138	201	251	252	430	-
1983	1,558	30	64	70	133	145	208	225	219	425	1
1982	1,509	31	48	88	122	150	214	215	234	380	-
1981	1,748	26	57	103	139	171	223	257	286	438	-
1980	1,798	22	83	103	135	191	216	279	299	438	1
1979	1,770	27	66	94	145	192	238	281	282	415	1
Other acute ischemic heart diseases (411)											
White, both sexes											
1998	2,448	82	122	138	181	207	253	320	351	706	-
1997	2,596	99	159	146	218	243	292	323	310	705	2
1996	2,454	76	129	144	183	249	269	332	296	664	1
1995	2,356	69	133	144	176	223	287	260	340	646	7
1994	2,442	80	121	118	200	226	285	290	338	682	4
1993	2,541	83	122	143	213	261	311	335	361	604	1
1992	2,699	108	113	159	235	313	334	345	349	623	2
1991	2,807	107	131	178	260	314	360	351	360	623	7
1990	2,807	108	147	190	264	330	349	397	353	571	1
1989	2,843	79	105	165	292	329	362	360	388	662	1
1988	2,915	81	119	194	270	344	383	393	387	640	-
1987	3,007	97	142	213	295	377	360	449	387	589	2
1986	3,296	100	149	266	378	389	453	435	398	591	2
1985	3,394	97	199	264	401	429	443	444	422	601	2
1984	3,387	81	177	272	378	441	442	465	412	608	-
1983	3,405	122	210	258	386	441	458	453	384	579	1
1982	3,488	120	193	321	398	440	532	429	399	525	1
1981	3,835	115	220	346	447	512	499	474	461	629	-
1980	4,048	113	243	363	445	525	551	539	492	642	4
1979	4,020	133	241	363	464	528	543	545	468	603	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
White, male											
1998	1,293	68	97	98	133	142	155	175	145	210	-
1997	1,447	86	127	111	162	161	193	176	135	216	2
1996	1,342	62	100	111	140	161	173	172	137	199	-
1995	1,308	57	111	113	128	160	186	137	157	199	5
1994	1,331	64	93	85	154	159	180	155	151	211	1
1993	1,417	69	100	109	152	176	195	166	170	193	-
1992	1,495	91	95	123	171	186	212	170	154	193	1
1991	1,654	94	104	136	187	232	248	183	167	195	6
1990	1,667	91	115	148	186	244	232	219	171	180	1
1989	1,608	69	84	133	216	230	232	187	173	202	1
1988	1,661	66	94	148	206	228	236	224	165	206	-
1987	1,750	77	113	163	218	257	226	255	188	182	-
1986	2,014	86	123	207	281	276	287	266	195	179	2
1985	2,060	84	168	208	291	306	293	244	198	193	2
1984	2,081	69	141	215	276	334	281	253	205	210	-
1983	2,132	100	164	206	284	335	298	267	192	201	-
1982	2,230	99	162	261	305	324	346	245	193	183	1
1981	2,383	97	182	270	338	376	328	250	211	231	-
1980	2,522	102	186	283	334	375	374	289	230	238	3
1979	2,532	114	194	299	354	369	342	310	212	227	1
Other acute ischemic heart diseases (411)											
White, female											
1998	1,155	14	25	40	48	65	98	145	206	496	-
1997	1,149	13	32	35	56	82	99	147	175	489	-
1996	1,112	14	29	33	43	88	96	160	159	465	1
1995	1,048	12	22	31	48	63	101	123	183	447	2
1994	1,111	16	28	33	46	67	105	135	187	471	3
1993	1,124	14	22	34	61	85	116	169	191	411	1
1992	1,204	17	18	36	64	127	122	175	195	430	1
1991	1,153	13	27	42	73	82	112	168	193	428	1
1990	1,140	17	32	42	78	86	117	178	182	391	-
1989	1,235	10	21	32	76	99	130	173	215	460	-
1988	1,254	15	25	46	64	116	147	169	222	434	-
1987	1,257	20	29	50	77	120	134	194	199	407	2
1986	1,282	14	26	59	97	113	166	169	203	412	-
1985	1,334	13	31	56	110	123	150	200	224	408	-
1984	1,306	12	36	57	102	107	161	212	207	398	-
1983	1,273	22	46	52	102	106	160	186	192	378	1
1982	1,258	21	31	60	93	116	186	184	206	342	-
1981	1,452	18	38	76	109	136	171	224	250	398	-
1980	1,526	11	57	80	111	150	177	250	262	404	1
1979	1,488	19	47	64	110	159	201	235	256	376	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
All other, both sexes											
1998	459	29	30	41	44	55	54	58	42	75	-
1997	450	19	37	44	48	49	58	50	41	82	2
1996	435	28	27	41	48	40	49	59	31	81	1
1995	384	22	26	31	45	42	45	44	36	61	-
1994	420	26	36	32	46	31	53	37	61	68	-
1993	427	27	27	32	40	56	47	39	53	60	4
1992	480	20	34	40	51	42	55	60	55	87	3
1991	493	18	28	33	57	73	56	58	60	67	-
1990	526	29	36	38	68	72	60	69	45	71	-
1989	533	29	33	60	66	77	54	82	49	56	-
1988	493	31	38	28	49	69	60	66	41	68	2
1987	525	24	39	53	67	64	62	72	40	65	1
1986	560	26	37	53	62	96	78	65	47	56	-
1985	575	28	50	57	76	71	87	56	42	63	-
1984	647	27	51	62	83	87	95	65	70	63	2
1983	654	21	54	66	82	97	91	73	60	78	-
1982	632	39	60	80	76	81	68	71	48	68	2
1981	714	30	62	89	91	82	114	76	67	60	-
1980	725	53	71	90	89	116	79	66	67	53	-
1979	742	29	64	93	115	99	94	92	58	61	-
Other acute ischemic heart diseases (411)											
All other, male											
1998	270	21	23	31	31	35	32	29	21	24	-
1997	244	14	26	29	26	30	28	25	21	28	2
1996	249	17	19	23	34	28	27	39	16	21	-
1995	209	17	19	25	25	26	24	27	13	12	-
1994	231	22	24	20	34	18	29	19	24	25	-
1993	245	22	21	15	25	35	28	21	28	22	2
1992	275	15	23	26	33	29	36	35	26	25	2
1991	287	12	17	20	37	51	29	29	21	34	-
1990	301	23	27	27	45	45	33	31	18	26	-
1989	294	17	21	43	38	49	23	38	27	22	-
1988	294	22	21	19	32	47	40	39	18	22	1
1987	323	18	30	33	47	43	35	45	19	26	-
1986	338	17	27	36	46	60	50	34	21	19	-
1985	321	18	31	39	47	36	50	29	15	23	-
1984	378	20	35	38	60	56	55	26	25	31	2
1983	369	13	36	48	51	58	43	34	33	31	-
1982	381	29	43	52	47	47	40	40	20	30	2
1981	418	22	43	62	61	47	62	43	31	20	-
1980	453	42	45	67	65	75	40	37	30	19	-
1979	460	21	45	63	80	66	57	46	32	22	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
All other, female											
1998	189	8	7	10	13	20	22	29	21	51	-
1997	206	5	11	15	22	19	30	25	20	54	-
1996	186	11	8	18	14	12	22	20	15	60	1
1995	175	5	7	6	20	16	21	17	23	49	-
1994	189	4	12	12	12	13	24	18	37	43	-
1993	182	5	6	17	15	21	19	18	25	38	2
1992	205	5	11	14	18	13	19	25	29	62	1
1991	206	6	11	13	20	22	27	29	39	33	-
1990	225	6	9	11	23	27	27	38	27	45	-
1989	239	12	12	17	28	28	31	44	22	34	-
1988	199	9	17	9	17	22	20	27	23	46	1
1987	202	6	9	20	20	21	27	27	21	39	1
1986	222	9	10	17	16	36	28	31	26	37	-
1985	254	10	19	18	29	35	37	27	27	40	-
1984	269	7	16	24	23	31	40	39	45	32	-
1983	285	8	18	18	31	39	48	39	27	47	-
1982	251	10	17	28	29	34	28	31	28	38	-
1981	296	8	19	27	30	35	52	33	36	40	-
1980	272	11	26	23	24	41	39	29	37	34	-
1979	282	8	19	30	35	33	37	46	26	39	-
Other acute ischemic heart diseases (411)											
Black, both sexes											
1998	407	25	27	32	39	49	49	53	38	65	-
1997	396	18	30	39	43	45	51	47	34	72	1
1996	398	25	26	36	47	35	45	56	26	73	1
1995	351	20	24	27	43	33	43	42	31	57	-
1994	361	22	32	26	39	28	45	31	51	59	-
1993	386	21	26	31	38	48	40	37	50	53	3
1992	428	13	25	37	46	38	53	58	50	79	3
1991	450	13	25	26	53	68	54	53	57	63	-
1990	479	26	35	33	66	62	58	64	37	65	-
1989	496	29	29	55	59	73	51	76	47	51	-
1988	465	28	36	27	47	67	58	63	38	62	2
1987	482	23	34	47	63	60	61	67	36	58	1
1986	517	25	36	48	61	86	70	58	44	53	-
1985	549	27	48	55	74	69	85	52	40	57	-
1984	615	27	50	56	82	82	87	63	68	60	2
1983	624	20	52	61	81	93	87	71	56	75	-
1982	598	38	57	75	72	76	63	68	46	65	1
1981	673	27	57	84	84	77	113	71	65	56	-
1980	685	50	67	85	87	110	73	61	62	51	-
1979	709	27	61	91	109	96	91	89	54	57	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute ischemic heart diseases (411)											
Black, male											
1998	234	17	20	22	29	31	27	26	19	21	-
1997	210	13	19	27	24	28	24	24	16	23	1
1996	230	15	18	19	33	25	26	37	13	20	-
1995	190	15	18	21	24	21	22	27	13	9	-
1994	196	18	21	15	29	16	24	15	21	22	-
1993	216	16	21	14	23	27	24	19	27	19	2
1992	237	8	15	24	28	27	34	33	24	23	2
1991	259	10	14	16	34	48	27	26	19	32	-
1990	272	21	26	24	43	40	32	28	13	24	-
1989	271	17	18	38	33	46	22	34	27	20	-
1988	274	19	19	18	31	46	38	36	16	21	1
1987	290	17	26	27	44	39	34	41	17	22	-
1986	307	17	26	31	45	53	45	28	20	17	-
1985	304	17	29	38	46	35	48	27	14	20	-
1984	354	20	34	32	59	51	50	25	24	30	2
1983	350	13	35	44	51	57	39	33	30	29	-
1982	356	29	40	48	44	42	36	38	20	29	1
1981	394	19	39	59	58	45	61	40	30	19	-
1980	420	39	41	63	63	70	35	33	27	18	-
1979	435	19	42	61	77	63	55	44	29	20	-
Other acute ischemic heart diseases (411)											
Black, female											
1998	173	8	7	10	10	18	22	27	19	44	-
1997	186	5	11	12	19	17	27	23	18	49	-
1996	168	10	8	17	14	10	19	19	13	53	1
1995	161	5	6	6	19	12	21	15	18	48	-
1994	165	4	11	11	10	12	21	16	30	37	-
1993	170	5	5	17	15	21	16	18	23	34	1
1992	191	5	10	13	18	11	19	25	26	56	1
1991	191	3	11	10	19	20	27	27	38	31	-
1990	207	5	9	9	23	22	26	36	24	41	-
1989	225	12	11	17	26	27	29	42	20	31	-
1988	191	9	17	9	16	21	20	27	22	41	1
1987	192	6	8	20	19	21	27	26	19	36	1
1986	210	8	10	17	16	33	25	30	24	36	-
1985	245	10	19	17	28	34	37	25	26	37	-
1984	261	7	16	24	23	31	37	38	44	30	-
1983	274	7	17	17	30	36	48	38	26	46	-
1982	242	9	17	27	28	34	27	30	26	36	-
1981	279	8	18	25	26	32	52	31	35	37	-
1980	265	11	26	22	24	40	38	28	35	33	-
1979	274	8	19	30	32	33	36	45	25	37	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,
sex and year

	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other forms of chronic ischemic heart disease (412-414,429.2)											
All other, both sexes											
1998	35,441	1,098	1,599	2,104	2,855	3,602	4,627	5,063	4,698	8,672	15
1997	35,368	1,137	1,594	2,141	2,920	3,545	4,662	4,941	4,724	8,602	14
1996	35,502	1,085	1,604	2,065	3,015	3,690	4,712	4,972	4,937	8,297	7
1995	36,281	1,146	1,581	2,172	2,926	3,956	5,010	4,916	5,055	8,412	14
1994	35,072	999	1,523	2,093	2,865	3,953	4,681	4,746	4,964	8,093	19
1993	35,843	953	1,505	2,156	3,093	4,118	4,760	4,889	5,030	8,197	20
1992	33,587	900	1,383	1,992	2,917	3,930	4,524	4,723	4,639	7,525	16
1991	33,452	874	1,413	1,994	2,926	4,032	4,538	4,668	4,723	7,261	20
1990	32,711	867	1,312	2,011	2,967	4,006	4,270	4,668	4,551	7,123	26
1989	33,133	911	1,366	2,052	3,183	3,945	4,315	4,873	4,493	7,001	18
1988	33,648	924	1,354	2,127	3,261	3,953	4,388	5,056	4,664	6,984	18
1987	32,888	829	1,359	2,086	3,104	3,880	4,461	4,833	4,511	6,820	16
1986	32,645	779	1,360	2,070	3,178	3,773	4,388	4,789	4,425	6,902	30
1985	32,003	813	1,375	2,174	3,075	3,752	4,525	4,743	4,408	6,232	20
1984	31,445	794	1,314	2,179	3,115	3,580	4,558	4,581	4,451	5,999	17
1983	32,163	808	1,486	2,302	3,168	3,620	4,710	4,699	4,512	6,048	20
1982	31,138	872	1,472	2,161	3,032	3,680	4,375	4,532	4,298	5,866	31
1981	31,227	843	1,459	2,220	3,089	3,676	4,505	4,487	4,267	5,809	30
1980	32,023	865	1,532	2,358	3,086	3,922	4,536	4,805	4,256	5,817	31
1979	29,452	858	1,563	2,232	2,777	3,746	4,340	4,572	3,603	4,976	24
Other forms of chronic ischemic heart disease (412-414,429.2)											
All other, male											
1998	17,789	763	1,107	1,416	1,781	2,117	2,459	2,557	2,008	2,788	13
1997	17,885	832	1,078	1,417	1,904	2,073	2,517	2,505	2,037	2,749	8
1996	17,654	763	1,070	1,340	1,886	2,149	2,521	2,462	2,059	2,636	6
1995	18,328	807	1,117	1,444	1,828	2,376	2,727	2,453	2,096	2,699	7
1994	17,534	695	1,072	1,379	1,824	2,303	2,522	2,281	2,073	2,575	12
1993	17,919	662	1,004	1,422	1,901	2,437	2,554	2,324	2,171	2,629	11
1992	16,816	639	923	1,297	1,786	2,289	2,482	2,336	1,951	2,377	11
1991	16,979	595	982	1,315	1,836	2,361	2,461	2,208	2,105	2,378	16
1990	16,460	581	884	1,340	1,811	2,273	2,332	2,263	1,915	2,374	17
1989	16,858	661	969	1,333	1,987	2,333	2,275	2,352	1,935	2,296	15
1988	17,073	662	901	1,417	2,063	2,302	2,324	2,406	2,035	2,278	12
1987	16,828	589	949	1,311	1,949	2,272	2,409	2,391	1,970	2,231	11
1986	16,647	539	937	1,378	2,003	2,177	2,345	2,311	1,877	2,368	16
1985	16,466	593	963	1,396	1,931	2,123	2,409	2,276	1,939	2,153	15
1984	16,045	578	885	1,397	1,923	2,053	2,438	2,171	1,924	2,052	10
1983	16,370	583	996	1,466	1,934	2,030	2,462	2,264	1,959	2,090	9
1982	16,146	591	977	1,388	1,839	2,151	2,360	2,269	1,854	2,083	20
1981	16,127	582	995	1,472	1,882	2,068	2,357	2,170	1,891	2,081	18
1980	16,642	596	1,037	1,510	1,869	2,215	2,349	2,342	1,915	2,210	22
1979	15,508	575	1,090	1,492	1,709	2,105	2,274	2,216	1,646	1,851	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Other forms of chronic ischemic heart disease (412-414,429.2)											
All other, female											
1998	17,652	335	492	688	1,074	1,485	2,168	2,506	2,690	5,884	2
1997	17,483	305	516	724	1,016	1,472	2,145	2,436	2,687	5,853	6
1996	17,848	322	534	725	1,129	1,541	2,191	2,510	2,878	5,661	1
1995	17,953	339	464	728	1,098	1,580	2,283	2,463	2,959	5,713	7
1994	17,538	304	451	714	1,041	1,650	2,159	2,465	2,891	5,518	7
1993	17,924	291	501	734	1,192	1,681	2,206	2,565	2,859	5,568	9
1992	16,771	261	460	695	1,131	1,641	2,042	2,387	2,688	5,148	5
1991	16,473	279	431	679	1,090	1,671	2,077	2,460	2,618	4,883	4
1990	16,251	286	428	671	1,156	1,733	1,938	2,405	2,636	4,749	9
1989	16,275	250	397	719	1,196	1,612	2,040	2,521	2,558	4,705	3
1988	16,575	262	453	710	1,198	1,651	2,064	2,650	2,629	4,706	6
1987	16,060	240	410	775	1,155	1,608	2,052	2,442	2,541	4,589	5
1986	15,998	240	423	692	1,175	1,596	2,043	2,478	2,548	4,534	14
1985	15,537	220	412	778	1,144	1,629	2,116	2,467	2,469	4,079	5
1984	15,400	216	429	782	1,192	1,527	2,120	2,410	2,527	3,947	7
1983	15,793	225	490	836	1,234	1,590	2,248	2,435	2,553	3,958	11
1982	14,992	281	495	773	1,193	1,529	2,015	2,263	2,444	3,783	11
1981	15,100	261	464	748	1,207	1,608	2,148	2,317	2,376	3,728	12
1980	15,381	269	495	848	1,217	1,707	2,187	2,463	2,341	3,607	9
1979	13,944	283	473	740	1,068	1,641	2,066	2,356	1,957	3,125	9
Other forms of chronic ischemic heart disease (412-414,429.2)											
Black, both sexes											
1998	30,511	972	1,400	1,867	2,512	3,189	4,003	4,292	3,950	7,323	14
1997	30,593	1,011	1,403	1,872	2,596	3,068	4,038	4,269	4,025	7,334	14
1996	30,858	971	1,442	1,808	2,677	3,210	4,115	4,286	4,217	7,133	7
1995	31,877	1,042	1,427	1,952	2,618	3,492	4,381	4,328	4,374	7,269	14
1994	30,844	878	1,363	1,833	2,541	3,508	4,121	4,158	4,360	7,050	19
1993	31,875	855	1,377	1,943	2,793	3,706	4,264	4,344	4,376	7,198	16
1992	29,752	810	1,241	1,799	2,633	3,525	4,028	4,148	4,032	6,592	16
1991	29,994	798	1,280	1,800	2,659	3,655	4,116	4,144	4,207	6,401	19
1990	29,466	787	1,201	1,829	2,719	3,632	3,841	4,211	4,058	6,339	23
1989	30,018	837	1,263	1,862	2,925	3,596	3,931	4,428	4,030	6,223	18
1988	30,660	851	1,241	1,966	3,008	3,606	4,010	4,584	4,230	6,292	18
1987	30,102	766	1,241	1,917	2,884	3,605	4,083	4,404	4,111	6,172	15
1986	30,014	718	1,254	1,949	2,962	3,507	4,032	4,404	4,033	6,257	27
1985	29,477	763	1,294	2,020	2,872	3,491	4,185	4,344	4,040	5,626	17
1984	29,056	732	1,237	2,017	2,922	3,314	4,243	4,237	4,079	5,448	17
1983	29,838	753	1,395	2,161	2,965	3,387	4,406	4,364	4,152	5,519	11
1982	28,995	823	1,386	2,047	2,846	3,456	4,099	4,213	3,982	5,373	26
1981	29,317	806	1,379	2,100	2,952	3,494	4,238	4,180	3,987	5,368	28
1980	30,076	823	1,448	2,232	2,935	3,763	4,269	4,505	3,970	5,341	30
1979	27,612	819	1,483	2,106	2,642	3,546	4,104	4,275	3,349	4,554	23

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other forms of chronic ischemic heart disease (412-414,429.2)											
Black, male											
1998	14,891	659	954	1,233	1,522	1,857	2,094	2,086	1,633	2,167	12
1997	15,064	724	936	1,209	1,666	1,757	2,131	2,116	1,693	2,165	8
1996	14,943	666	943	1,155	1,657	1,839	2,155	2,070	1,700	2,102	6
1995	15,758	719	990	1,281	1,632	2,061	2,346	2,104	1,744	2,191	7
1994	14,991	589	944	1,185	1,602	2,024	2,168	1,955	1,759	2,064	12
1993	15,591	582	910	1,262	1,713	2,169	2,236	2,013	1,793	2,200	9
1992	14,587	566	811	1,156	1,603	2,037	2,179	2,004	1,606	1,984	11
1991	14,893	529	885	1,172	1,651	2,099	2,198	1,889	1,792	2,009	16
1990	14,528	519	797	1,218	1,646	2,023	2,066	1,994	1,643	2,014	15
1989	14,960	600	891	1,212	1,799	2,111	2,035	2,075	1,643	1,938	15
1988	15,205	598	821	1,296	1,875	2,069	2,074	2,118	1,759	1,966	12
1987	15,104	544	855	1,184	1,786	2,091	2,155	2,122	1,730	1,955	10
1986	15,021	488	857	1,288	1,841	1,993	2,107	2,074	1,640	2,094	16
1985	14,924	547	903	1,282	1,785	1,953	2,180	2,010	1,733	1,904	12
1984	14,565	525	821	1,291	1,792	1,877	2,230	1,945	1,688	1,813	10
1983	14,943	537	926	1,363	1,797	1,874	2,259	2,033	1,749	1,882	6
1982	14,756	548	914	1,305	1,707	1,987	2,174	2,042	1,667	1,844	17
1981	14,915	556	930	1,376	1,789	1,945	2,164	1,969	1,721	1,886	17
1980	15,371	559	970	1,417	1,758	2,104	2,160	2,143	1,744	1,967	21
1979	14,322	542	1,027	1,392	1,598	1,968	2,105	2,013	1,508	1,660	14
Other forms of chronic ischemic heart disease (412-414,429.2)											
Black, female											
1998	15,620	313	446	634	990	1,332	1,909	2,206	2,317	5,156	2
1997	15,529	287	467	663	930	1,311	1,907	2,153	2,332	5,169	6
1996	15,915	305	499	653	1,020	1,371	1,960	2,216	2,517	5,031	1
1995	16,119	323	437	671	986	1,431	2,035	2,224	2,630	5,078	7
1994	15,853	289	419	648	939	1,484	1,953	2,203	2,601	4,986	7
1993	16,284	273	467	681	1,080	1,537	2,028	2,331	2,583	4,998	7
1992	15,165	244	430	643	1,030	1,488	1,849	2,144	2,426	4,608	5
1991	15,101	269	395	628	1,008	1,556	1,918	2,255	2,415	4,392	3
1990	14,938	268	404	611	1,073	1,609	1,775	2,217	2,415	4,325	8
1989	15,058	237	372	650	1,126	1,485	1,896	2,353	2,387	4,285	3
1988	15,455	253	420	670	1,133	1,537	1,936	2,466	2,471	4,326	6
1987	14,998	222	386	733	1,098	1,514	1,928	2,282	2,381	4,217	5
1986	14,993	230	397	661	1,121	1,514	1,925	2,330	2,393	4,163	11
1985	14,553	216	391	738	1,087	1,538	2,005	2,334	2,307	3,722	5
1984	14,491	207	416	726	1,130	1,437	2,013	2,292	2,391	3,635	7
1983	14,895	216	469	798	1,168	1,513	2,147	2,331	2,403	3,637	5
1982	14,239	275	472	742	1,139	1,469	1,925	2,171	2,315	3,529	9
1981	14,402	250	449	724	1,163	1,549	2,074	2,211	2,266	3,482	11
1980	14,705	264	478	815	1,177	1,659	2,109	2,362	2,226	3,374	9
1979	13,290	277	456	714	1,044	1,578	1,999	2,262	1,841	2,894	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerotic cardiovascular disease, so described (429.2)											
All races, female											
1998	32,968	293	504	784	1,197	1,745	2,795	4,067	5,556	15,756	4
1997	33,233	295	507	784	1,197	1,765	2,834	4,213	5,576	15,803	4
1996	34,005	304	461	747	1,255	1,879	2,976	4,206	5,852	16,069	7
1995	35,886	302	497	742	1,278	2,004	3,150	4,475	6,157	17,018	15
1994	35,595	301	484	734	1,307	1,993	3,204	4,488	6,279	16,518	11
1993	36,620	277	474	776	1,407	2,154	3,332	4,603	6,268	17,088	8
1992	34,924	211	433	667	1,348	2,081	3,157	4,464	6,063	16,265	11
1991	35,103	227	390	660	1,278	2,227	3,126	4,585	6,231	16,152	9
1990	35,125	225	433	735	1,391	2,164	3,123	4,619	6,253	15,992	16
1989	36,332	203	408	752	1,430	2,149	3,222	4,863	6,591	16,543	12
1988	37,524	178	409	715	1,452	2,253	3,370	4,986	6,687	17,302	11
1987	37,486	184	366	752	1,424	2,265	3,416	4,990	6,740	17,168	8
1986	37,602	179	379	723	1,465	2,092	3,378	5,015	6,818	17,356	31
1985	37,336	157	333	716	1,457	2,159	3,466	5,124	6,954	16,795	37
1984	37,473	155	363	805	1,475	2,185	3,522	5,024	7,066	16,738	5
1983	37,562	149	407	776	1,471	2,285	3,502	5,163	7,159	16,493	16
1982	35,351	170	352	749	1,359	2,136	3,210	4,927	6,891	15,435	7
1981	35,703	155	416	721	1,377	2,136	3,498	5,031	6,881	15,339	9
1980	36,096	211	452	819	1,448	2,230	3,472	5,133	7,106	15,077	9
1979	33,532	196	441	768	1,337	2,074	3,298	4,902	6,778	13,605	12
Atherosclerotic cardiovascular disease, so described (429.2)											
White, both sexes											
1998	56,113	1,133	1,921	2,323	3,073	4,226	5,745	7,531	9,057	20,132	12
1997	56,622	1,164	1,785	2,301	3,210	4,298	5,941	7,739	8,963	20,264	13
1996	57,946	1,172	1,674	2,251	3,288	4,492	6,136	7,718	9,320	20,892	23
1995	59,967	1,035	1,765	2,205	3,328	4,723	6,626	8,032	9,681	21,613	28
1994	60,244	1,015	1,730	2,272	3,481	4,911	6,668	8,113	9,786	21,324	13
1993	61,027	995	1,561	2,287	3,547	5,179	6,783	8,221	9,838	21,738	11
1992	58,108	953	1,448	2,069	3,521	4,942	6,408	7,847	9,480	20,582	20
1991	58,519	891	1,339	2,016	3,583	5,134	6,447	8,045	9,686	20,553	19
1990	58,439	803	1,382	2,050	3,696	5,110	6,437	8,170	9,658	20,381	24
1989	59,705	754	1,324	2,214	3,674	5,152	6,416	8,343	10,022	21,093	22
1988	61,916	726	1,252	2,254	3,764	5,283	6,885	8,684	10,294	22,136	18
1987	61,744	667	1,273	2,312	3,707	5,272	6,959	8,648	10,226	21,986	17
1986	62,009	676	1,230	2,331	3,789	5,075	6,822	8,694	10,388	22,317	48
1985	61,607	641	1,163	2,206	3,788	5,007	6,740	8,732	10,680	22,048	54
1984	62,003	595	1,214	2,279	3,800	5,083	6,982	8,741	10,608	22,115	13
1983	62,415	615	1,304	2,290	3,675	5,099	6,785	8,989	11,027	22,053	21
1982	58,834	578	1,175	2,174	3,466	4,779	6,464	8,660	10,487	20,544	11
1981	59,847	612	1,341	2,229	3,494	4,977	6,720	8,778	10,588	20,612	11
1980	60,243	613	1,387	2,391	3,465	4,965	6,702	8,915	10,831	20,507	15
1979	56,935	654	1,395	2,326	3,453	4,728	6,382	8,460	10,434	18,637	13

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerotic cardiovascular disease, so described (429.2)											
White, male											
1998	27,993	942	1,578	1,777	2,239	2,914	3,568	4,117	4,194	5,875	9
1997	28,063	959	1,449	1,754	2,330	2,940	3,700	4,178	4,084	5,873	12
1996	28,790	970	1,376	1,735	2,416	3,069	3,786	4,167	4,216	6,213	16
1995	29,230	848	1,405	1,700	2,395	3,191	4,117	4,287	4,347	6,160	15
1994	29,716	825	1,422	1,765	2,545	3,409	4,110	4,349	4,318	6,212	8
1993	29,810	829	1,259	1,768	2,601	3,531	4,148	4,384	4,396	6,165	7
1992	28,152	816	1,161	1,624	2,580	3,358	3,866	4,110	4,179	5,734	12
1991	28,349	755	1,096	1,564	2,650	3,426	3,954	4,211	4,208	5,798	13
1990	28,261	680	1,099	1,539	2,703	3,506	3,933	4,264	4,154	5,750	13
1989	28,485	638	1,064	1,710	2,661	3,500	3,850	4,285	4,231	5,935	11
1988	29,551	622	995	1,764	2,711	3,583	4,176	4,496	4,409	6,245	11
1987	29,282	554	1,029	1,801	2,646	3,531	4,177	4,448	4,310	6,200	13
1986	29,318	562	994	1,825	2,697	3,484	4,070	4,435	4,372	6,326	23
1985	28,937	532	946	1,709	2,702	3,332	3,945	4,386	4,465	6,443	18
1984	29,179	487	978	1,738	2,695	3,365	4,101	4,445	4,321	6,550	9
1983	29,612	522	1,053	1,772	2,622	3,328	3,965	4,554	4,614	6,706	10
1982	27,772	483	951	1,646	2,466	3,103	3,811	4,376	4,308	6,195	7
1981	28,379	527	1,064	1,721	2,465	3,278	3,862	4,378	4,380	6,288	7
1980	28,719	485	1,096	1,831	2,386	3,263	3,895	4,496	4,398	6,476	10
1979	27,543	552	1,091	1,793	2,447	3,143	3,724	4,219	4,217	5,963	5
Atherosclerotic cardiovascular disease, so described (429.2)											
White, female											
1998	28,120	191	343	546	834	1,312	2,177	3,414	4,863	14,257	3
1997	28,559	205	336	547	880	1,358	2,241	3,561	4,879	14,391	1
1996	29,156	202	298	516	872	1,423	2,350	3,551	5,104	14,679	7
1995	30,737	187	360	505	933	1,532	2,509	3,745	5,334	15,453	13
1994	30,528	190	308	507	936	1,502	2,558	3,764	5,468	15,112	5
1993	31,217	166	302	519	946	1,648	2,635	3,837	5,442	15,573	4
1992	29,956	137	287	445	941	1,584	2,542	3,737	5,301	14,848	8
1991	30,170	136	243	452	933	1,708	2,493	3,834	5,478	14,755	6
1990	30,178	123	283	511	993	1,604	2,504	3,906	5,504	14,631	11
1989	31,220	116	260	504	1,013	1,652	2,566	4,058	5,791	15,158	11
1988	32,365	104	257	490	1,053	1,700	2,709	4,188	5,885	15,891	7
1987	32,462	113	244	511	1,061	1,741	2,782	4,200	5,916	15,786	4
1986	32,691	114	236	506	1,092	1,591	2,752	4,259	6,016	15,991	25
1985	32,670	109	217	497	1,086	1,675	2,795	4,346	6,215	15,605	36
1984	32,824	108	236	541	1,105	1,718	2,881	4,296	6,287	15,565	4
1983	32,803	93	251	518	1,053	1,771	2,820	4,435	6,413	15,347	11
1982	31,062	95	224	528	1,000	1,676	2,653	4,284	6,179	14,349	4
1981	31,468	85	277	508	1,029	1,699	2,858	4,400	6,208	14,324	4
1980	31,524	128	291	560	1,079	1,702	2,807	4,419	6,433	14,031	5
1979	29,392	102	304	533	1,006	1,585	2,658	4,241	6,217	12,674	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerotic cardiovascular disease, so described (429.2)											
All other, both sexes											
1998	10,663	370	622	776	1,038	1,197	1,447	1,415	1,241	2,200	10
1997	10,338	383	588	759	1,062	1,115	1,377	1,341	1,273	2,110	9
1996	10,568	388	591	762	1,115	1,217	1,451	1,372	1,306	2,057	4
1995	11,210	387	563	813	1,058	1,314	1,559	1,497	1,424	2,277	7
1994	10,976	369	636	725	1,085	1,336	1,469	1,435	1,465	2,123	12
1993	11,613	359	592	845	1,222	1,453	1,583	1,498	1,481	2,237	11
1992	10,848	305	500	740	1,156	1,342	1,527	1,450	1,383	2,127	8
1991	10,860	308	517	732	1,085	1,405	1,521	1,460	1,380	2,148	14
1990	10,681	312	501	768	1,139	1,409	1,381	1,417	1,404	2,070	16
1989	11,175	315	516	758	1,233	1,426	1,502	1,608	1,444	2,089	10
1988	10,911	286	484	781	1,193	1,363	1,426	1,599	1,430	2,090	12
1987	10,698	265	470	714	1,098	1,341	1,427	1,567	1,441	2,079	11
1986	10,658	240	495	767	1,103	1,294	1,435	1,541	1,423	2,107	17
1985	9,960	214	428	673	1,080	1,242	1,492	1,464	1,328	1,836	9
1984	9,887	226	402	731	1,070	1,177	1,431	1,393	1,414	1,808	2
1983	10,003	249	495	782	1,079	1,202	1,491	1,430	1,345	1,732	8
1982	9,221	255	412	672	975	1,152	1,312	1,326	1,274	1,637	8
1981	9,249	241	488	692	961	1,095	1,356	1,335	1,221	1,645	8
1980	9,679	279	489	769	1,028	1,190	1,393	1,412	1,222	1,677	10
1979	9,154	299	524	773	930	1,174	1,366	1,341	1,067	1,464	7
Atherosclerotic cardiovascular disease, so described (429.2)											
All other, male											
1998	5,815	268	461	538	675	764	829	762	548	701	9
1997	5,664	293	417	522	745	708	784	689	576	698	6
1996	5,719	286	428	531	732	761	825	717	558	667	4
1995	6,061	272	426	576	713	842	918	767	601	712	5
1994	5,909	258	460	498	714	845	823	711	654	717	6
1993	6,210	248	420	588	761	947	886	732	655	722	7
1992	5,880	231	354	518	749	845	912	723	621	710	5
1991	5,927	217	370	524	740	886	888	709	627	751	11
1990	5,734	210	351	544	741	849	762	704	655	709	11
1989	6,063	228	368	510	816	929	846	803	644	704	9
1988	5,752	212	332	556	794	810	765	801	628	679	8
1987	5,674	194	348	473	735	817	793	777	617	697	7
1986	5,747	175	352	550	730	793	809	785	621	742	11
1985	5,294	166	312	454	709	758	821	686	589	646	8
1984	5,238	179	275	467	700	710	790	665	635	635	1
1983	5,244	193	339	524	661	688	809	702	599	586	3
1982	4,932	180	284	451	616	692	755	683	562	551	5
1981	5,014	171	349	479	613	658	716	704	548	630	3
1980	5,107	196	328	510	659	662	728	698	549	631	6
1979	5,014	205	387	538	599	685	726	680	506	533	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerotic cardiovascular disease, so described (429.2)											
All other, female											
1998	4,848	102	161	238	363	433	618	653	693	1,499	1
1997	4,674	90	171	237	317	407	593	652	697	1,412	3
1996	4,849	102	163	231	383	456	626	655	748	1,390	-
1995	5,149	115	137	237	345	472	641	730	823	1,565	2
1994	5,067	111	176	227	371	491	646	724	811	1,406	6
1993	5,403	111	172	257	461	506	697	766	826	1,515	4
1992	4,968	74	146	222	407	497	615	727	762	1,417	3
1991	4,933	91	147	208	345	519	633	751	753	1,397	3
1990	4,947	102	150	224	398	560	619	713	749	1,361	5
1989	5,112	87	148	248	417	497	656	805	800	1,385	1
1988	5,159	74	152	225	399	553	661	798	802	1,411	4
1987	5,024	71	122	241	363	524	634	790	824	1,382	4
1986	4,911	65	143	217	373	501	626	756	802	1,365	6
1985	4,666	48	116	219	371	484	671	778	739	1,190	1
1984	4,649	47	127	264	370	467	641	728	779	1,173	1
1983	4,759	56	156	258	418	514	682	728	746	1,146	5
1982	4,289	75	128	221	359	460	557	643	712	1,086	3
1981	4,235	70	139	213	348	437	640	631	673	1,015	5
1980	4,572	83	161	259	369	528	665	714	673	1,046	4
1979	4,140	94	137	235	331	489	640	661	561	931	4
Atherosclerotic cardiovascular disease, so described (429.2)											
Black, both sexes											
1998	9,519	334	549	717	926	1,095	1,312	1,259	1,077	1,934	9
1997	9,247	344	524	682	949	1,016	1,230	1,198	1,138	1,868	9
1996	9,456	349	537	686	1,021	1,102	1,301	1,221	1,159	1,809	4
1995	10,152	358	518	735	960	1,194	1,406	1,364	1,289	2,036	7
1994	9,927	333	580	640	977	1,219	1,346	1,295	1,344	1,890	12
1993	10,608	331	549	776	1,131	1,319	1,455	1,366	1,330	2,029	10
1992	9,887	283	449	682	1,081	1,247	1,392	1,314	1,233	1,924	8
1991	9,944	281	478	676	1,012	1,314	1,406	1,328	1,248	1,910	14
1990	9,810	290	457	717	1,054	1,303	1,266	1,314	1,273	1,886	14
1989	10,336	296	488	696	1,143	1,320	1,394	1,507	1,313	1,914	10
1988	10,099	270	451	733	1,117	1,267	1,320	1,461	1,324	1,913	12
1987	9,967	251	435	660	1,028	1,263	1,330	1,458	1,339	1,926	11
1986	10,014	222	465	724	1,062	1,222	1,349	1,460	1,327	1,946	15
1985	9,312	203	402	637	1,016	1,172	1,407	1,355	1,241	1,686	8
1984	9,307	212	380	686	1,020	1,106	1,353	1,323	1,322	1,681	2
1983	9,486	239	463	744	1,031	1,150	1,424	1,354	1,259	1,633	5
1982	8,780	248	393	646	935	1,115	1,260	1,272	1,200	1,526	6
1981	8,872	238	473	665	933	1,057	1,295	1,257	1,173	1,572	8
1980	9,288	275	473	742	985	1,164	1,341	1,354	1,168	1,573	10
1979	8,806	292	511	751	892	1,144	1,328	1,276	1,009	1,393	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerotic cardiovascular disease, so described (429.2)											
Black, male											
1998	5,091	237	402	492	587	697	750	656	470	568	8
1997	4,996	260	370	465	659	639	683	611	514	593	6
1996	5,048	251	384	478	667	686	738	629	489	547	4
1995	5,420	247	388	518	642	761	823	690	534	607	5
1994	5,245	224	420	433	630	769	743	631	596	597	6
1993	5,610	222	391	534	705	857	804	652	572	641	7
1992	5,287	212	312	477	702	783	819	641	535	620	5
1991	5,344	195	340	479	684	819	811	620	544	648	11
1990	5,202	196	315	509	681	773	690	645	581	630	9
1989	5,533	214	346	469	754	859	781	735	560	619	9
1988	5,246	198	310	522	743	746	698	709	564	596	8
1987	5,217	184	321	432	680	772	725	711	552	633	7
1986	5,355	161	327	520	696	742	751	739	571	669	11
1985	4,906	156	293	425	666	708	763	621	544	586	7
1984	4,866	167	256	436	661	662	741	621	579	571	1
1983	4,911	185	313	495	630	652	759	649	547	545	2
1982	4,655	175	268	432	589	667	719	643	521	500	4
1981	4,763	169	338	458	593	633	674	646	516	596	3
1980	4,865	192	314	495	633	640	692	661	514	584	6
1979	4,795	199	379	522	569	666	699	633	475	503	3
Atherosclerotic cardiovascular disease, so described (429.2)											
Black, female											
1998	4,428	97	147	225	339	398	562	603	607	1,366	1
1997	4,251	84	154	217	290	377	547	587	624	1,275	3
1996	4,408	98	153	208	354	416	563	592	670	1,262	-
1995	4,732	111	130	217	318	433	583	674	755	1,429	2
1994	4,682	109	160	207	347	450	603	664	748	1,293	6
1993	4,998	109	158	242	426	462	651	714	758	1,388	3
1992	4,600	71	137	205	379	464	573	673	698	1,304	3
1991	4,600	86	138	197	328	495	595	708	704	1,262	3
1990	4,608	94	142	208	373	530	576	669	692	1,256	5
1989	4,803	82	142	227	389	461	613	772	753	1,295	1
1988	4,853	72	141	211	374	521	622	752	760	1,317	4
1987	4,750	67	114	228	348	491	605	747	787	1,293	4
1986	4,659	61	138	204	366	480	598	721	756	1,277	4
1985	4,406	47	109	212	350	464	644	734	697	1,100	1
1984	4,441	45	124	250	359	444	612	702	743	1,110	1
1983	4,575	54	150	249	401	498	665	705	712	1,088	3
1982	4,125	73	125	214	346	448	541	629	679	1,026	2
1981	4,109	69	135	207	340	424	621	611	657	976	5
1980	4,423	83	159	247	352	524	649	693	654	989	4
1979	4,011	93	132	229	323	478	629	643	534	890	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
All races, both sexes											
1998	253,383	3,412	5,294	7,309	11,028	17,559	27,767	37,533	45,076	94,946	12
1997	256,843	3,522	5,440	7,430	11,381	18,458	28,237	38,211	44,989	95,714	12
1996	259,703	3,679	5,277	7,372	12,022	19,259	29,404	38,600	45,732	94,798	16
1995	260,318	3,511	5,275	7,486	11,991	20,018	29,879	38,050	45,609	94,879	20
1994	256,197	3,351	5,021	7,408	12,202	20,094	29,646	37,545	44,893	92,638	20
1993	259,639	3,275	4,974	7,480	12,679	21,136	30,177	38,331	45,501	92,597	23
1992	247,489	3,187	4,655	7,457	12,792	20,818	29,258	37,022	42,974	85,997	26
1991	246,883	3,022	4,533	7,572	13,383	21,068	29,211	37,129	42,884	84,779	24
1990	246,830	2,964	4,512	7,756	13,708	21,429	29,160	37,478	42,646	84,036	25
1989	247,695	2,992	4,723	8,054	14,103	21,777	28,827	37,621	42,847	83,555	21
1988	258,234	3,100	4,827	8,857	14,994	22,578	30,639	39,030	44,277	86,692	31
1987	255,064	2,931	5,001	9,009	15,345	22,532	30,530	38,357	42,844	85,117	29
1986	255,871	2,845	5,124	9,293	15,845	22,541	30,797	38,333	43,010	84,567	160
1985	258,637	3,070	5,439	9,674	15,808	22,469	31,114	38,377	43,831	85,491	125
1984	258,132	3,047	5,383	9,773	16,087	22,616	31,190	38,784	43,833	84,270	39
1983	262,133	2,945	5,649	10,293	16,026	23,094	31,694	39,221	44,947	85,202	40
1982	257,635	3,042	5,848	10,181	16,028	22,995	31,112	38,726	44,535	82,160	68
1981	258,105	3,108	5,983	10,579	16,175	23,099	31,228	38,362	44,977	81,739	57
1980	261,398	3,032	6,130	10,757	15,731	23,301	31,606	39,700	46,332	81,976	61
1979	246,141	3,082	5,968	9,999	15,164	22,092	30,237	37,710	43,927	75,346	47
All other forms of chronic ischemic heart disease (412-414)											
All races, male											
1998	124,816	2,662	4,025	5,329	7,586	11,535	17,211	21,527	22,148	30,185	9
1997	126,470	2,760	4,126	5,446	7,927	12,235	17,507	21,730	21,975	30,070	7
1996	127,706	2,857	3,935	5,359	8,471	12,754	18,239	21,863	21,984	29,519	10
1995	127,315	2,749	4,040	5,429	8,398	13,267	18,628	21,336	21,568	29,099	7
1994	125,494	2,657	3,835	5,387	8,607	13,369	18,539	20,946	21,045	28,440	16
1993	126,968	2,614	3,773	5,492	8,878	14,054	18,643	21,342	21,160	28,290	10
1992	122,022	2,546	3,550	5,475	9,067	13,931	18,150	20,669	19,863	26,151	16
1991	121,651	2,380	3,496	5,625	9,402	14,074	18,130	20,524	19,527	25,865	17
1990	121,146	2,339	3,436	5,731	9,691	14,229	18,031	20,332	19,300	25,501	13
1989	121,324	2,390	3,660	5,951	9,935	14,485	17,675	20,301	19,206	25,161	15
1988	126,666	2,467	3,684	6,512	10,591	15,042	18,878	20,898	19,710	26,259	20
1987	125,432	2,349	3,867	6,653	10,851	14,883	18,673	20,531	19,105	25,768	17
1986	126,037	2,271	3,994	7,005	11,172	14,907	18,783	20,320	18,891	25,927	48
1985	127,940	2,456	4,242	7,265	11,235	14,786	18,972	20,341	19,446	26,518	43
1984	127,068	2,429	4,191	7,288	11,351	14,788	18,651	20,278	19,138	26,431	23
1983	129,342	2,366	4,356	7,641	11,311	15,245	19,043	20,624	19,226	27,017	25
1982	127,837	2,411	4,549	7,667	11,225	15,193	18,469	20,236	19,144	26,531	37
1981	128,230	2,454	4,640	7,908	11,320	15,136	18,601	19,814	19,410	26,674	30
1980	129,314	2,419	4,738	7,990	10,932	15,191	18,540	20,318	19,763	27,166	39
1979	123,240	2,449	4,702	7,560	10,727	14,480	17,885	19,262	18,855	25,243	37

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
All races, female											
1998	128,567	750	1,269	1,980	3,442	6,024	10,556	16,006	22,928	64,761	3
1997	130,373	762	1,314	1,984	3,454	6,223	10,730	16,481	23,014	65,644	5
1996	131,997	822	1,342	2,013	3,551	6,505	11,165	16,737	23,748	65,279	6
1995	133,003	762	1,235	2,057	3,593	6,751	11,251	16,714	24,041	65,780	13
1994	130,703	694	1,186	2,021	3,595	6,725	11,107	16,599	23,848	64,198	4
1993	132,671	661	1,201	1,988	3,801	7,082	11,534	16,989	24,341	64,307	13
1992	125,467	641	1,105	1,982	3,725	6,887	11,108	16,353	23,111	59,846	10
1991	125,232	642	1,037	1,947	3,981	6,994	11,081	16,605	23,357	58,914	7
1990	125,684	625	1,076	2,025	4,017	7,200	11,129	17,146	23,346	58,535	12
1989	126,371	602	1,063	2,103	4,168	7,292	11,152	17,320	23,641	58,394	6
1988	131,568	633	1,143	2,345	4,403	7,536	11,761	18,132	24,567	60,433	11
1987	129,632	582	1,134	2,356	4,494	7,649	11,857	17,826	23,739	59,349	12
1986	129,834	574	1,130	2,288	4,673	7,634	12,014	18,013	24,119	58,640	112
1985	130,697	614	1,197	2,409	4,573	7,683	12,142	18,036	24,385	58,973	82
1984	131,064	618	1,192	2,485	4,736	7,828	12,539	18,506	24,695	57,839	16
1983	132,791	579	1,293	2,652	4,715	7,849	12,651	18,597	25,721	58,185	15
1982	129,798	631	1,299	2,514	4,803	7,802	12,643	18,490	25,391	55,629	31
1981	129,875	654	1,343	2,671	4,855	7,963	12,627	18,548	25,567	55,065	27
1980	132,084	613	1,392	2,767	4,799	8,110	13,066	19,382	26,569	54,810	22
1979	122,901	633	1,266	2,439	4,437	7,612	12,352	18,448	25,072	50,103	10
All other forms of chronic ischemic heart disease (412-414)											
White, both sexes											
1998	228,605	2,684	4,317	5,981	9,211	15,154	24,587	33,885	41,619	88,474	7
1997	231,813	2,768	4,434	6,048	9,523	16,028	24,952	34,611	41,538	89,222	7
1996	234,769	2,982	4,264	6,069	10,122	16,786	26,143	35,000	42,101	88,558	13
1995	235,247	2,752	4,257	6,127	10,123	17,376	26,428	34,631	41,978	88,744	13
1994	232,101	2,721	4,134	6,040	10,422	17,477	26,434	34,234	41,394	86,668	13
1993	235,409	2,681	4,061	6,169	10,808	18,471	27,000	34,940	41,952	86,637	14
1992	224,750	2,592	3,772	6,205	11,031	18,230	26,261	33,749	39,718	80,599	18
1991	224,291	2,456	3,637	6,310	11,542	18,441	26,194	33,921	39,541	79,666	18
1990	224,800	2,409	3,701	6,513	11,880	18,832	26,271	34,227	39,499	78,983	15
1989	225,737	2,396	3,873	6,760	12,153	19,258	26,014	34,356	39,798	78,643	13
1988	235,497	2,462	3,957	7,511	12,926	19,988	27,677	35,573	41,043	81,798	25
1987	232,874	2,367	4,112	7,637	13,339	19,993	27,496	35,091	39,774	80,376	24
1986	233,884	2,306	4,259	7,990	13,770	20,062	27,844	35,085	40,008	79,772	147
1985	236,594	2,471	4,492	8,173	13,813	19,959	28,081	35,098	40,751	81,095	114
1984	236,574	2,479	4,471	8,325	14,042	20,213	28,063	35,596	40,796	80,079	24
1983	239,973	2,386	4,658	8,773	13,937	20,676	28,475	35,952	41,780	80,886	28
1982	235,718	2,425	4,788	8,692	13,971	20,467	28,049	35,520	41,511	77,931	45
1981	236,127	2,506	5,012	9,051	14,047	20,518	28,079	35,210	41,931	77,575	35
1980	239,054	2,446	5,087	9,168	13,673	20,569	28,463	36,307	43,298	77,836	40
1979	225,843	2,523	4,929	8,540	13,317	19,520	27,263	34,479	41,391	71,834	30

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
White, male											
1998	112,842	2,167	3,379	4,451	6,480	10,182	15,581	19,732	20,688	28,098	5
1997	114,249	2,221	3,465	4,551	6,768	10,870	15,774	19,914	20,514	28,019	5
1996	115,771	2,380	3,293	4,550	7,317	11,366	16,543	20,118	20,483	27,550	8
1995	115,048	2,214	3,349	4,561	7,283	11,733	16,819	19,650	20,073	27,112	5
1994	113,869	2,220	3,223	4,506	7,497	11,911	16,840	19,376	19,626	26,582	10
1993	115,259	2,200	3,189	4,658	7,738	12,564	16,975	19,750	19,644	26,383	6
1992	111,086	2,138	2,981	4,696	8,030	12,487	16,580	19,056	18,533	24,484	10
1991	110,599	2,002	2,884	4,834	8,306	12,599	16,557	19,025	18,049	24,238	12
1990	110,420	1,968	2,903	4,935	8,621	12,805	16,461	18,773	18,040	23,836	7
1989	110,529	1,957	3,059	5,128	8,764	13,081	16,246	18,752	17,915	23,569	9
1988	115,345	2,017	3,115	5,651	9,322	13,550	17,319	19,293	18,303	24,660	16
1987	114,278	1,954	3,266	5,815	9,637	13,428	17,057	18,917	17,752	24,234	13
1986	115,137	1,907	3,409	6,177	9,899	13,523	17,247	18,794	17,635	24,301	43
1985	116,768	2,029	3,591	6,323	10,013	13,421	17,384	18,751	18,096	25,011	36
1984	116,261	2,030	3,581	6,358	10,128	13,445	17,003	18,772	17,849	25,014	14
1983	118,216	1,976	3,699	6,699	10,038	13,903	17,390	19,062	17,866	25,513	19
1982	116,623	2,000	3,856	6,730	10,002	13,734	16,864	18,650	17,852	24,999	22
1981	117,117	2,043	3,994	6,915	10,051	13,726	16,960	18,348	18,067	25,223	15
1980	117,779	2,019	4,029	6,990	9,722	13,638	16,919	18,674	18,397	25,587	23
1979	112,746	2,079	3,999	6,606	9,617	13,060	16,337	17,726	17,715	23,925	25
All other forms of chronic ischemic heart disease (412-414)											
White, female											
1998	115,763	517	938	1,530	2,731	4,972	9,006	14,153	20,931	60,376	2
1997	117,564	547	969	1,497	2,755	5,158	9,178	14,697	21,024	61,203	2
1996	118,998	602	971	1,519	2,805	5,420	9,600	14,882	21,618	61,008	5
1995	120,199	538	908	1,566	2,840	5,643	9,609	14,981	21,905	61,632	8
1994	118,232	501	911	1,534	2,925	5,566	9,594	14,858	21,768	60,086	3
1993	120,150	481	872	1,511	3,070	5,907	10,025	15,190	22,308	60,254	8
1992	113,664	454	791	1,509	3,001	5,743	9,681	14,693	21,185	56,115	8
1991	113,692	454	753	1,476	3,236	5,842	9,637	14,896	21,492	55,428	6
1990	114,380	441	798	1,578	3,259	6,027	9,810	15,454	21,459	55,147	8
1989	115,208	439	814	1,632	3,389	6,177	9,768	15,604	21,883	55,074	4
1988	120,152	445	842	1,860	3,604	6,438	10,358	16,280	22,740	57,138	9
1987	118,596	413	846	1,822	3,702	6,565	10,439	16,174	22,022	56,142	11
1986	118,747	399	850	1,813	3,871	6,539	10,597	16,291	22,373	55,471	104
1985	119,826	442	901	1,850	3,800	6,538	10,697	16,347	22,655	56,084	78
1984	120,313	449	890	1,967	3,914	6,768	11,060	16,824	22,947	55,065	10
1983	121,757	410	959	2,074	3,899	6,773	11,085	16,890	23,914	55,373	9
1982	119,095	425	932	1,962	3,969	6,733	11,185	16,870	23,659	52,932	23
1981	119,010	463	1,018	2,136	3,996	6,792	11,119	16,862	23,864	52,352	20
1980	121,275	427	1,058	2,178	3,951	6,931	11,544	17,633	24,901	52,249	17
1979	113,097	444	930	1,934	3,700	6,460	10,926	16,753	23,676	47,909	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
All other, both sexes											
1998	24,778	728	977	1,328	1,817	2,405	3,180	3,648	3,457	6,472	5
1997	25,030	754	1,006	1,382	1,858	2,430	3,285	3,600	3,451	6,492	5
1996	24,934	697	1,013	1,303	1,900	2,473	3,261	3,600	3,631	6,240	3
1995	25,071	759	1,018	1,359	1,868	2,642	3,451	3,419	3,631	6,135	7
1994	24,096	630	887	1,368	1,780	2,617	3,212	3,311	3,499	5,970	7
1993	24,230	594	913	1,311	1,871	2,665	3,177	3,391	3,549	5,960	9
1992	22,739	595	883	1,252	1,761	2,588	2,997	3,273	3,256	5,398	8
1991	22,592	566	896	1,262	1,841	2,627	3,017	3,208	3,343	5,113	6
1990	22,030	555	811	1,243	1,828	2,597	2,889	3,251	3,147	5,053	10
1989	21,958	596	850	1,294	1,950	2,519	2,813	3,265	3,049	4,912	8
1988	22,737	638	870	1,346	2,068	2,590	2,962	3,457	3,234	4,894	6
1987	22,190	564	889	1,372	2,006	2,539	3,034	3,266	3,070	4,741	5
1986	21,987	539	865	1,303	2,075	2,479	2,953	3,248	3,002	4,795	13
1985	22,043	599	947	1,501	1,995	2,510	3,033	3,279	3,080	4,396	11
1984	21,558	568	912	1,448	2,045	2,403	3,127	3,188	3,037	4,191	15
1983	22,160	559	991	1,520	2,089	2,418	3,219	3,269	3,167	4,316	12
1982	21,917	617	1,060	1,489	2,057	2,528	3,063	3,206	3,024	4,229	23
1981	21,978	602	971	1,528	2,128	2,581	3,149	3,152	3,046	4,164	22
1980	22,344	586	1,043	1,589	2,058	2,732	3,143	3,393	3,034	4,140	21
1979	20,298	559	1,039	1,459	1,847	2,572	2,974	3,231	2,536	3,512	17
All other forms of chronic ischemic heart disease (412-414)											
All other, male											
1998	11,974	495	646	878	1,106	1,353	1,630	1,795	1,460	2,087	4
1997	12,221	539	661	895	1,159	1,365	1,733	1,816	1,461	2,051	2
1996	11,935	477	642	809	1,154	1,388	1,696	1,745	1,501	1,969	2
1995	12,267	535	691	868	1,115	1,534	1,809	1,686	1,495	1,987	2
1994	11,625	437	612	881	1,110	1,458	1,699	1,570	1,419	1,858	6
1993	11,709	414	584	834	1,140	1,490	1,668	1,592	1,516	1,907	4
1992	10,936	408	569	779	1,037	1,444	1,570	1,613	1,330	1,667	6
1991	11,052	378	612	791	1,096	1,475	1,573	1,499	1,478	1,627	5
1990	10,726	371	533	796	1,070	1,424	1,570	1,559	1,260	1,665	6
1989	10,795	433	601	823	1,171	1,404	1,429	1,549	1,291	1,592	6
1988	11,321	450	569	861	1,269	1,492	1,559	1,605	1,407	1,599	4
1987	11,154	395	601	838	1,214	1,455	1,616	1,614	1,353	1,534	4
1986	10,900	364	585	828	1,273	1,384	1,536	1,526	1,256	1,626	5
1985	11,172	427	651	942	1,222	1,365	1,588	1,590	1,350	1,507	7
1984	10,807	399	610	930	1,223	1,343	1,648	1,506	1,289	1,417	9
1983	11,126	390	657	942	1,273	1,342	1,653	1,562	1,360	1,504	6
1982	11,214	411	693	937	1,223	1,459	1,605	1,586	1,292	1,532	15
1981	11,113	411	646	993	1,269	1,410	1,641	1,466	1,343	1,451	15
1980	11,535	400	709	1,000	1,210	1,553	1,621	1,644	1,366	1,579	16
1979	10,494	370	703	954	1,110	1,420	1,548	1,536	1,140	1,318	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
All other, female											
1998	12,804	233	331	450	711	1,052	1,550	1,853	1,997	4,385	1
1997	12,809	215	345	487	699	1,065	1,552	1,784	1,990	4,441	3
1996	12,999	220	371	494	746	1,085	1,565	1,855	2,130	4,271	1
1995	12,804	224	327	491	753	1,108	1,642	1,733	2,136	4,148	5
1994	12,471	193	275	487	670	1,159	1,513	1,741	2,080	4,112	1
1993	12,521	180	329	477	731	1,175	1,509	1,799	2,033	4,053	5
1992	11,803	187	314	473	724	1,144	1,427	1,660	1,926	3,731	2
1991	11,540	188	284	471	745	1,152	1,444	1,709	1,865	3,486	1
1990	11,304	184	278	447	758	1,173	1,319	1,692	1,887	3,388	4
1989	11,163	163	249	471	779	1,115	1,384	1,716	1,758	3,320	2
1988	11,416	188	301	485	799	1,098	1,403	1,852	1,827	3,295	2
1987	11,036	169	288	534	792	1,084	1,418	1,652	1,717	3,207	1
1986	11,087	175	280	475	802	1,095	1,417	1,722	1,746	3,169	8
1985	10,871	172	296	559	773	1,145	1,445	1,689	1,730	2,889	4
1984	10,751	169	302	518	822	1,060	1,479	1,682	1,748	2,774	6
1983	11,034	169	334	578	816	1,076	1,566	1,707	1,807	2,812	6
1982	10,703	206	367	552	834	1,069	1,458	1,620	1,732	2,697	8
1981	10,865	191	325	535	859	1,171	1,508	1,686	1,703	2,713	7
1980	10,809	186	334	589	848	1,179	1,522	1,749	1,668	2,561	5
1979	9,804	189	336	505	737	1,152	1,426	1,695	1,396	2,194	5
All other forms of chronic ischemic heart disease (412-414)											
Black, both sexes											
1998	20,992	638	851	1,150	1,586	2,094	2,691	3,033	2,873	5,389	5
1997	21,346	667	879	1,190	1,647	2,052	2,808	3,071	2,887	5,466	5
1996	21,402	622	905	1,122	1,656	2,108	2,814	3,065	3,058	5,324	3
1995	21,725	684	909	1,217	1,658	2,298	2,975	2,964	3,085	5,233	7
1994	20,917	545	783	1,193	1,564	2,289	2,775	2,863	3,016	5,160	7
1993	21,267	524	828	1,167	1,662	2,387	2,809	2,978	3,046	5,169	6
1992	19,865	527	792	1,117	1,552	2,278	2,636	2,834	2,799	4,668	8
1991	20,050	517	802	1,124	1,647	2,341	2,710	2,816	2,959	4,491	5
1990	19,656	497	744	1,112	1,665	2,329	2,575	2,897	2,785	4,453	9
1989	19,682	541	775	1,166	1,782	2,276	2,537	2,921	2,717	4,309	8
1988	20,561	581	790	1,233	1,891	2,339	2,690	3,123	2,906	4,379	6
1987	20,135	515	806	1,257	1,856	2,342	2,753	2,946	2,772	4,246	4
1986	20,000	496	789	1,225	1,900	2,285	2,683	2,944	2,706	4,311	12
1985	20,165	560	892	1,383	1,856	2,319	2,778	2,989	2,799	3,940	9
1984	19,749	520	857	1,331	1,902	2,208	2,890	2,914	2,757	3,767	15
1983	20,352	514	932	1,417	1,934	2,237	2,982	3,010	2,893	3,886	6
1982	20,215	575	993	1,401	1,911	2,341	2,839	2,941	2,782	3,847	20
1981	20,445	568	906	1,435	2,019	2,437	2,943	2,923	2,814	3,796	20
1980	20,788	548	975	1,490	1,950	2,599	2,928	3,151	2,802	3,768	20
1979	18,806	527	972	1,355	1,750	2,402	2,776	2,999	2,340	3,161	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of chronic ischemic heart disease (412-414)											
Black, male											
1998	9,800	422	552	741	935	1,160	1,344	1,430	1,163	1,599	4
1997	10,068	464	566	744	1,007	1,118	1,448	1,505	1,179	1,572	2
1996	9,895	415	559	677	990	1,153	1,417	1,441	1,211	1,555	2
1995	10,338	472	602	763	990	1,300	1,523	1,414	1,210	1,584	2
1994	9,746	365	524	752	972	1,255	1,425	1,324	1,163	1,467	6
1993	9,981	360	519	728	1,008	1,312	1,432	1,361	1,221	1,559	2
1992	9,300	354	499	679	901	1,254	1,360	1,363	1,071	1,364	6
1991	9,549	334	545	693	967	1,280	1,387	1,269	1,248	1,361	5
1990	9,326	323	482	709	965	1,250	1,376	1,349	1,062	1,384	6
1989	9,427	386	545	743	1,045	1,252	1,254	1,340	1,083	1,319	6
1988	9,959	400	511	774	1,132	1,323	1,376	1,409	1,195	1,370	4
1987	9,887	360	534	752	1,106	1,319	1,430	1,411	1,178	1,322	3
1986	9,666	327	530	768	1,145	1,251	1,356	1,335	1,069	1,425	5
1985	10,018	391	610	857	1,119	1,245	1,417	1,389	1,189	1,318	5
1984	9,699	358	565	855	1,131	1,215	1,489	1,324	1,109	1,242	9
1983	10,032	352	613	868	1,167	1,222	1,500	1,384	1,202	1,337	4
1982	10,101	373	646	873	1,118	1,320	1,455	1,399	1,146	1,344	13
1981	10,152	387	592	918	1,196	1,312	1,490	1,323	1,205	1,290	14
1980	10,506	367	656	922	1,125	1,464	1,468	1,482	1,230	1,383	15
1979	9,527	343	648	870	1,029	1,302	1,406	1,380	1,033	1,157	11
All other forms of chronic ischemic heart disease (412-414)											
Black, female											
1998	11,192	216	299	409	651	934	1,347	1,603	1,710	3,790	1
1997	11,278	203	313	446	640	934	1,360	1,566	1,708	3,894	3
1996	11,507	207	346	445	666	955	1,397	1,624	1,847	3,769	1
1995	11,387	212	307	454	668	998	1,452	1,550	1,875	3,649	5
1994	11,171	180	259	441	592	1,034	1,350	1,539	1,853	3,693	1
1993	11,286	164	309	439	654	1,075	1,377	1,617	1,825	3,610	4
1992	10,565	173	293	438	651	1,024	1,276	1,471	1,728	3,304	2
1991	10,501	183	257	431	680	1,061	1,323	1,547	1,711	3,130	-
1990	10,330	174	262	403	700	1,079	1,199	1,548	1,723	3,069	3
1989	10,255	155	230	423	737	1,024	1,283	1,581	1,634	2,990	2
1988	10,602	181	279	459	759	1,016	1,314	1,714	1,711	3,009	2
1987	10,248	155	272	505	750	1,023	1,323	1,535	1,594	2,924	1
1986	10,334	169	259	457	755	1,034	1,327	1,609	1,637	2,886	7
1985	10,147	169	282	526	737	1,074	1,361	1,600	1,610	2,622	4
1984	10,050	162	292	476	771	993	1,401	1,590	1,648	2,525	6
1983	10,320	162	319	549	767	1,015	1,482	1,626	1,691	2,549	2
1982	10,114	202	347	528	793	1,021	1,384	1,542	1,636	2,503	7
1981	10,293	181	314	517	823	1,125	1,453	1,600	1,609	2,506	6
1980	10,282	181	319	568	825	1,135	1,460	1,669	1,572	2,385	5
1979	9,279	184	324	485	721	1,100	1,370	1,619	1,307	2,004	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
All races, both sexes											
1998	163,323	3,662	4,386	5,475	7,367	10,718	15,960	21,780	26,453	59,076	10
1997	159,868	3,502	4,248	5,401	7,451	10,957	16,068	21,371	26,156	56,408	8
1996	155,028	3,480	4,044	5,175	7,347	11,150	15,886	20,555	25,348	54,069	18
1995	152,454	3,388	3,988	5,105	7,746	11,152	16,067	20,179	24,866	51,501	8
1994	147,879	3,115	3,794	5,130	7,464	11,447	16,365	19,593	23,888	48,724	6
1993	149,513	2,979	3,857	5,331	7,941	12,153	16,493	20,752	23,912	47,840	14
1992	138,652	2,888	3,530	4,950	8,049	11,637	15,877	19,323	22,117	42,416	8
1991	136,284	2,641	3,346	5,094	8,170	11,913	15,637	19,323	21,495	40,874	12
1990	132,300	2,577	3,314	5,141	8,392	11,973	15,054	18,887	21,082	38,479	18
1989	135,531	2,528	3,597	5,580	8,767	12,581	15,757	19,624	21,295	38,295	10
1988	153,078	2,713	4,042	6,567	10,438	14,606	18,237	22,120	23,591	42,647	20
1987	146,185	2,700	4,041	6,312	10,148	14,304	17,765	21,038	22,433	39,556	20
1986	142,648	2,599	3,966	6,509	10,368	14,016	17,679	20,803	21,664	37,073	31
1985	132,522	2,495	3,962	6,518	9,948	13,065	16,661	19,065	19,843	33,550	36
1984	121,457	2,361	3,846	6,254	9,482	12,286	15,262	17,224	17,858	29,722	24
1983	114,570	2,223	3,830	6,183	9,067	11,706	14,702	16,289	16,495	27,191	21
1982	103,891	2,298	3,618	5,837	8,476	10,903	13,228	14,699	14,678	23,731	24
1981	98,112	2,287	3,792	6,018	8,038	10,330	12,310	13,461	13,887	21,633	17
1980	92,766	2,144	3,702	5,716	7,610	10,101	11,895	12,842	12,609	19,744	21
1979	83,345	2,115	3,556	5,225	7,122	9,041	10,496	11,472	11,554	16,994	17
Other heart diseases (415-429.1,429.3-429.9)											
All races, male											
1998	72,360	2,368	2,734	3,298	4,548	6,150	8,542	10,798	11,501	17,215	5
1997	71,549	2,264	2,699	3,357	4,537	6,361	8,704	10,721	11,173	16,571	4
1996	69,917	2,271	2,548	3,191	4,470	6,612	8,745	10,352	10,839	15,939	11
1995	69,435	2,268	2,559	3,150	4,679	6,589	8,871	10,200	10,644	15,116	6
1994	67,501	2,024	2,481	3,288	4,578	6,713	9,063	9,743	10,161	14,276	4
1993	68,513	1,943	2,512	3,369	4,874	7,146	9,131	10,220	10,140	14,055	9
1992	64,651	1,877	2,265	3,176	4,958	6,820	8,829	9,612	9,449	12,657	4
1991	64,025	1,773	2,163	3,198	5,143	7,057	8,745	9,792	9,124	12,129	6
1990	62,585	1,709	2,135	3,251	5,194	7,114	8,445	9,549	9,078	11,415	10
1989	64,376	1,697	2,318	3,562	5,425	7,587	8,781	9,789	9,007	11,492	5
1988	73,805	1,767	2,652	4,231	6,555	8,713	10,476	11,117	10,027	13,080	11
1987	70,718	1,759	2,664	4,032	6,249	8,611	9,999	10,645	9,492	12,278	13
1986	69,978	1,676	2,555	4,201	6,551	8,507	10,005	10,556	9,263	11,649	10
1985	65,532	1,648	2,604	4,218	6,313	7,966	9,393	9,609	8,530	10,662	17
1984	60,897	1,589	2,478	4,087	6,043	7,552	8,616	8,679	7,600	9,826	13
1983	57,823	1,434	2,489	4,003	5,730	7,223	8,444	8,176	7,036	9,072	11
1982	52,592	1,472	2,391	3,750	5,401	6,659	7,386	7,320	6,231	8,094	14
1981	50,399	1,479	2,489	3,927	5,064	6,200	6,973	6,825	6,086	7,541	8
1980	48,039	1,401	2,441	3,756	4,839	6,072	6,756	6,495	5,533	6,873	11
1979	43,469	1,394	2,342	3,459	4,498	5,595	5,895	5,715	5,073	6,079	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
All races, female											
1998	90,963	1,294	1,652	2,177	2,819	4,568	7,418	10,982	14,952	41,861	5
1997	88,319	1,238	1,549	2,044	2,914	4,596	7,364	10,650	14,983	39,837	4
1996	85,111	1,209	1,496	1,984	2,877	4,538	7,141	10,203	14,509	38,130	7
1995	83,019	1,120	1,429	1,955	3,067	4,563	7,196	9,979	14,222	36,385	2
1994	80,378	1,091	1,313	1,842	2,886	4,734	7,302	9,850	13,727	34,448	2
1993	81,000	1,036	1,345	1,962	3,067	5,007	7,362	10,532	13,772	33,785	5
1992	74,001	1,011	1,265	1,774	3,091	4,817	7,048	9,711	12,668	29,759	4
1991	72,259	868	1,183	1,896	3,027	4,856	6,892	9,531	12,371	28,745	6
1990	69,715	868	1,179	1,890	3,198	4,859	6,609	9,338	12,004	27,064	8
1989	71,155	831	1,279	2,018	3,342	4,994	6,976	9,835	12,288	26,803	5
1988	79,273	946	1,390	2,336	3,883	5,893	7,761	11,003	13,564	29,567	9
1987	75,467	941	1,377	2,280	3,899	5,693	7,766	10,393	12,941	27,278	7
1986	72,670	923	1,411	2,308	3,817	5,509	7,674	10,247	12,401	25,424	21
1985	66,990	847	1,358	2,300	3,635	5,099	7,268	9,456	11,313	22,888	19
1984	60,560	772	1,368	2,167	3,439	4,734	6,646	8,545	10,258	19,896	11
1983	56,747	789	1,341	2,180	3,337	4,483	6,258	8,113	9,459	18,119	10
1982	51,299	826	1,227	2,087	3,075	4,244	5,842	7,379	8,447	15,637	10
1981	47,713	808	1,303	2,091	2,974	4,130	5,337	6,636	7,801	14,092	9
1980	44,727	743	1,261	1,960	2,771	4,029	5,139	6,347	7,076	12,871	10
1979	39,876	721	1,214	1,766	2,624	3,446	4,601	5,757	6,481	10,915	9
Other heart diseases (415-429.1,429.3-429.9)											
White, both sexes											
1998	141,056	2,479	3,210	4,140	5,730	8,770	13,584	19,178	23,909	54,507	5
1997	138,416	2,435	3,069	4,103	5,847	9,026	13,752	18,933	23,689	51,996	5
1996	133,631	2,360	2,925	3,930	5,666	9,199	13,494	18,165	22,924	49,826	11
1995	131,201	2,353	2,914	3,847	6,081	9,157	13,639	17,795	22,442	47,496	7
1994	127,338	2,193	2,786	3,848	5,945	9,429	13,990	17,259	21,513	44,968	6
1993	128,406	2,074	2,771	3,940	6,282	9,986	14,067	18,319	21,479	44,130	8
1992	118,712	1,986	2,565	3,709	6,403	9,624	13,542	16,971	19,855	39,056	5
1991	116,581	1,824	2,433	3,878	6,535	9,784	13,330	16,888	19,279	37,590	12
1990	112,770	1,814	2,388	3,884	6,712	9,839	12,924	16,395	18,871	35,219	10
1989	115,125	1,742	2,524	4,242	6,957	10,283	13,441	17,052	18,992	35,024	6
1988	130,390	1,869	2,924	5,075	8,346	12,050	15,549	19,181	21,023	39,032	11
1987	124,476	1,905	2,947	4,905	8,118	11,841	15,099	18,248	19,964	36,237	9
1986	121,271	1,821	2,902	5,023	8,350	11,678	15,016	18,011	19,289	33,906	24
1985	112,612	1,776	2,891	5,056	8,036	10,850	14,095	16,567	17,688	30,749	28
1984	103,294	1,661	2,874	4,913	7,695	10,256	12,913	15,063	15,846	27,239	14
1983	97,238	1,612	2,892	4,810	7,317	9,745	12,481	14,177	14,628	24,957	11
1982	88,576	1,645	2,765	4,628	6,882	9,094	11,258	12,868	13,121	21,900	12
1981	83,337	1,612	2,898	4,756	6,514	8,669	10,544	11,768	12,388	19,853	10
1980	78,251	1,550	2,804	4,536	6,191	8,336	10,110	11,154	11,285	18,051	13
1979	70,693	1,532	2,792	4,184	5,812	7,450	8,925	9,919	10,448	15,664	11

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
White, male											
1998	61,723	1,639	2,038	2,571	3,616	5,120	7,382	9,605	10,472	15,753	2
1997	61,273	1,617	2,010	2,598	3,625	5,341	7,557	9,604	10,163	15,206	1
1996	59,542	1,577	1,898	2,501	3,503	5,548	7,529	9,216	9,847	14,577	7
1995	59,000	1,637	1,902	2,435	3,760	5,504	7,663	9,089	9,618	13,828	5
1994	57,431	1,465	1,871	2,566	3,719	5,631	7,859	8,669	9,183	13,040	4
1993	58,095	1,380	1,829	2,516	3,952	5,987	7,888	9,103	9,155	12,864	4
1992	54,672	1,305	1,696	2,464	4,052	5,763	7,612	8,519	8,455	11,522	2
1991	54,102	1,259	1,629	2,488	4,216	5,891	7,572	8,589	8,190	11,054	6
1990	52,784	1,208	1,584	2,514	4,261	5,956	7,346	8,415	8,110	10,346	5
1989	54,133	1,206	1,655	2,785	4,438	6,332	7,584	8,576	8,032	10,384	2
1988	62,137	1,249	1,954	3,357	5,359	7,313	9,056	9,673	8,908	11,788	8
1987	59,623	1,275	1,990	3,224	5,101	7,199	8,680	9,283	8,426	11,096	7
1986	58,932	1,193	1,926	3,309	5,383	7,218	8,629	9,196	8,186	10,517	8
1985	55,179	1,194	1,937	3,369	5,198	6,738	8,073	8,403	7,544	9,652	13
1984	51,436	1,157	1,895	3,296	5,021	6,407	7,394	7,624	6,715	8,893	8
1983	48,716	1,055	1,941	3,190	4,727	6,131	7,232	7,132	6,209	8,246	7
1982	44,513	1,059	1,887	3,034	4,441	5,637	6,334	6,461	5,541	7,374	8
1981	42,630	1,093	1,969	3,181	4,184	5,297	6,050	6,028	5,385	6,828	4
1980	40,204	1,031	1,888	3,017	3,990	5,111	5,792	5,676	4,913	6,198	7
1979	36,749	1,034	1,904	2,820	3,757	4,687	5,094	4,953	4,569	5,520	5
Other heart diseases (415-429.1,429.3-429.9)											
White, female											
1998	79,333	840	1,172	1,569	2,114	3,650	6,202	9,573	13,437	38,754	3
1997	77,143	818	1,059	1,505	2,222	3,685	6,195	9,329	13,526	36,790	4
1996	74,089	783	1,027	1,429	2,163	3,651	5,965	8,949	13,077	35,249	4
1995	72,201	716	1,012	1,412	2,321	3,653	5,976	8,706	12,824	33,668	2
1994	69,907	728	915	1,282	2,226	3,798	6,131	8,590	12,330	31,928	2
1993	70,311	694	942	1,424	2,330	3,999	6,179	9,216	12,324	31,266	4
1992	64,040	681	869	1,245	2,351	3,861	5,930	8,452	11,400	27,534	3
1991	62,479	565	804	1,390	2,319	3,893	5,758	8,299	11,089	26,536	6
1990	59,986	606	804	1,370	2,451	3,883	5,578	7,980	10,761	24,873	5
1989	60,992	536	869	1,457	2,519	3,951	5,857	8,476	10,960	24,640	4
1988	68,253	620	970	1,718	2,987	4,737	6,493	9,508	12,115	27,244	3
1987	64,853	630	957	1,681	3,017	4,642	6,419	8,965	11,538	25,141	2
1986	62,339	628	976	1,714	2,967	4,460	6,387	8,815	11,103	23,389	16
1985	57,433	582	954	1,687	2,838	4,112	6,022	8,164	10,144	21,097	15
1984	51,858	504	979	1,617	2,674	3,849	5,519	7,439	9,131	18,346	6
1983	48,522	557	951	1,620	2,590	3,614	5,249	7,045	8,419	16,711	4
1982	44,063	586	878	1,594	2,441	3,457	4,924	6,407	7,580	14,526	4
1981	40,707	519	929	1,575	2,330	3,372	4,494	5,740	7,003	13,025	6
1980	38,047	519	916	1,519	2,201	3,225	4,318	5,478	6,372	11,853	6
1979	33,944	498	888	1,364	2,055	2,763	3,831	4,966	5,879	10,144	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
All other, both sexes											
1998	22,267	1,183	1,176	1,335	1,637	1,948	2,376	2,602	2,544	4,569	5
1997	21,452	1,067	1,179	1,298	1,604	1,931	2,316	2,438	2,467	4,412	3
1996	21,397	1,120	1,119	1,245	1,681	1,951	2,392	2,390	2,424	4,243	7
1995	21,253	1,035	1,074	1,258	1,665	1,995	2,428	2,384	2,424	4,005	1
1994	20,541	922	1,008	1,282	1,519	2,018	2,375	2,334	2,375	3,756	-
1993	21,107	905	1,086	1,391	1,659	2,167	2,426	2,433	2,433	3,710	6
1992	19,940	902	965	1,241	1,646	2,013	2,335	2,352	2,262	3,360	3
1991	19,703	817	913	1,216	1,635	2,129	2,307	2,435	2,216	3,284	-
1990	19,530	763	926	1,257	1,680	2,134	2,130	2,492	2,211	3,260	8
1989	20,406	786	1,073	1,338	1,810	2,298	2,316	2,572	2,303	3,271	4
1988	22,688	844	1,118	1,492	2,092	2,556	2,688	2,939	2,568	3,615	9
1987	21,709	795	1,094	1,407	2,030	2,463	2,666	2,790	2,469	3,319	11
1986	21,377	778	1,064	1,486	2,018	2,338	2,663	2,792	2,375	3,167	7
1985	19,910	719	1,071	1,462	1,912	2,215	2,566	2,498	2,155	2,801	8
1984	18,163	700	972	1,341	1,787	2,030	2,349	2,161	2,012	2,483	10
1983	17,332	611	938	1,373	1,750	1,961	2,221	2,112	1,867	2,234	10
1982	15,315	653	853	1,209	1,594	1,809	1,970	1,831	1,557	1,831	12
1981	14,775	675	894	1,262	1,524	1,661	1,766	1,693	1,499	1,780	7
1980	14,515	594	898	1,180	1,419	1,765	1,785	1,688	1,324	1,693	8
1979	12,652	583	764	1,041	1,310	1,591	1,571	1,553	1,106	1,330	6
Other heart diseases (415-429.1,429.3-429.9)											
All other, male											
1998	10,637	729	696	727	932	1,030	1,160	1,193	1,029	1,462	3
1997	10,276	647	689	759	912	1,020	1,147	1,117	1,010	1,365	3
1996	10,375	694	650	690	967	1,064	1,216	1,136	992	1,362	4
1995	10,435	631	657	715	919	1,085	1,208	1,111	1,026	1,288	1
1994	10,070	559	610	722	859	1,082	1,204	1,074	978	1,236	-
1993	10,418	563	683	853	922	1,159	1,243	1,117	985	1,191	5
1992	9,979	572	569	712	906	1,057	1,217	1,093	994	1,135	2
1991	9,923	514	534	710	927	1,166	1,173	1,203	934	1,075	-
1990	9,801	501	551	737	933	1,158	1,099	1,134	968	1,069	5
1989	10,243	491	663	777	987	1,255	1,197	1,213	975	1,108	3
1988	11,668	518	698	874	1,196	1,400	1,420	1,444	1,119	1,292	3
1987	11,095	484	674	808	1,148	1,412	1,319	1,362	1,066	1,182	6
1986	11,046	483	629	892	1,168	1,289	1,376	1,360	1,077	1,132	2
1985	10,353	454	667	849	1,115	1,228	1,320	1,206	986	1,010	4
1984	9,461	432	583	791	1,022	1,145	1,222	1,055	885	933	5
1983	9,107	379	548	813	1,003	1,092	1,212	1,044	827	826	4
1982	8,079	413	504	716	960	1,022	1,052	859	690	720	6
1981	7,769	386	520	746	880	903	923	797	701	713	4
1980	7,835	370	553	739	849	961	964	819	620	675	4
1979	6,720	360	438	639	741	908	801	762	504	559	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
All other, female											
1998	11,630	454	480	608	705	918	1,216	1,409	1,515	3,107	2
1997	11,176	420	490	539	692	911	1,169	1,321	1,457	3,047	-
1996	11,022	426	469	555	714	887	1,176	1,254	1,432	2,881	3
1995	10,818	404	417	543	746	910	1,220	1,273	1,398	2,717	-
1994	10,471	363	398	560	660	936	1,171	1,260	1,397	2,520	-
1993	10,689	342	403	538	737	1,008	1,183	1,316	1,448	2,519	1
1992	9,961	330	396	529	740	956	1,118	1,259	1,268	2,225	1
1991	9,780	303	379	506	708	963	1,134	1,232	1,282	2,209	-
1990	9,729	262	375	520	747	976	1,031	1,358	1,243	2,191	3
1989	10,163	295	410	561	823	1,043	1,119	1,359	1,328	2,163	1
1988	11,020	326	420	618	896	1,156	1,268	1,495	1,449	2,323	6
1987	10,614	311	420	599	882	1,051	1,347	1,428	1,403	2,137	5
1986	10,331	295	435	594	850	1,049	1,287	1,432	1,298	2,035	5
1985	9,557	265	404	613	797	987	1,246	1,292	1,169	1,791	4
1984	8,702	268	389	550	765	885	1,127	1,106	1,127	1,550	5
1983	8,225	232	390	560	747	869	1,009	1,068	1,040	1,408	6
1982	7,236	240	349	493	634	787	918	972	867	1,111	6
1981	7,006	289	374	516	644	758	843	896	798	1,067	3
1980	6,680	224	345	441	570	804	821	869	704	1,018	4
1979	5,932	223	326	402	569	683	770	791	602	771	3
Other heart diseases (415-429.1,429.3-429.9)											
Black, both sexes											
1998	19,964	1,111	1,075	1,223	1,476	1,779	2,126	2,323	2,263	3,955	5
1997	19,227	986	1,095	1,189	1,470	1,750	2,114	2,125	2,172	3,840	2
1996	19,285	1,031	1,027	1,143	1,534	1,774	2,165	2,141	2,144	3,728	7
1995	19,201	966	977	1,151	1,544	1,832	2,179	2,155	2,128	3,516	1
1994	18,627	859	928	1,192	1,405	1,833	2,183	2,119	2,113	3,278	-
1993	19,201	863	1,010	1,302	1,534	2,006	2,222	2,185	2,161	3,275	5
1992	18,324	847	926	1,158	1,535	1,871	2,143	2,135	2,062	3,008	3
1991	18,169	783	865	1,137	1,519	1,991	2,131	2,239	2,010	2,931	-
1990	17,951	717	856	1,167	1,564	1,983	1,974	2,286	2,010	2,925	8
1989	18,935	745	1,002	1,267	1,701	2,170	2,163	2,379	2,103	2,946	4
1988	21,238	811	1,045	1,409	1,983	2,408	2,517	2,763	2,386	3,317	9
1987	20,360	759	1,033	1,311	1,925	2,333	2,515	2,640	2,315	3,052	11
1986	20,095	741	1,007	1,420	1,933	2,233	2,517	2,603	2,210	2,902	7
1985	18,722	685	1,030	1,392	1,809	2,099	2,439	2,328	2,025	2,582	7
1984	17,130	662	914	1,269	1,704	1,944	2,244	2,029	1,892	2,299	9
1983	16,281	587	903	1,298	1,659	1,872	2,091	1,974	1,748	2,044	8
1982	14,468	615	820	1,148	1,524	1,719	1,870	1,722	1,459	1,718	12
1981	14,022	642	863	1,218	1,460	1,599	1,690	1,597	1,408	1,648	3
1980	13,742	572	858	1,131	1,371	1,676	1,686	1,589	1,253	1,577	7
1979	11,954	571	732	991	1,258	1,507	1,500	1,469	1,023	1,229	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other heart diseases (415-429.1,429.3-429.9)											
Black, male											
1998	9,407	671	633	655	832	933	1,020	1,048	893	1,201	3
1997	9,122	594	638	690	843	929	1,033	953	871	1,120	2
1996	9,241	631	593	631	864	964	1,094	1,005	855	1,136	4
1995	9,411	587	603	655	856	1,000	1,075	1,003	883	1,088	1
1994	9,038	518	558	664	798	972	1,093	967	854	1,017	-
1993	9,365	538	630	799	849	1,068	1,133	991	838	990	4
1992	9,097	532	542	657	844	979	1,114	979	884	974	2
1991	9,073	492	506	666	857	1,081	1,072	1,096	818	921	-
1990	8,924	468	507	679	867	1,071	1,008	1,027	853	931	5
1989	9,409	467	614	730	919	1,181	1,100	1,099	861	964	3
1988	10,870	498	650	827	1,127	1,315	1,326	1,341	1,029	1,157	3
1987	10,300	458	631	745	1,094	1,335	1,227	1,263	967	1,073	6
1986	10,302	456	594	855	1,116	1,225	1,286	1,248	977	1,010	2
1985	9,688	429	640	809	1,049	1,156	1,250	1,095	923	930	3
1984	8,876	409	549	754	976	1,099	1,159	973	816	850	5
1983	8,488	365	527	770	953	1,036	1,125	963	763	731	3
1982	7,595	395	478	682	923	972	982	786	639	670	6
1981	7,344	365	507	724	838	868	881	743	646	640	1
1980	7,353	355	529	702	821	907	896	764	572	608	4
1979	6,338	357	419	611	712	860	762	709	464	513	3
Other heart diseases (415-429.1,429.3-429.9)											
Black, female											
1998	10,557	440	442	568	644	846	1,106	1,275	1,370	2,754	2
1997	10,105	392	457	499	627	821	1,081	1,172	1,301	2,720	-
1996	10,044	400	434	512	670	810	1,071	1,136	1,289	2,592	3
1995	9,790	379	374	496	688	832	1,104	1,152	1,245	2,428	-
1994	9,589	341	370	528	607	861	1,090	1,152	1,259	2,261	-
1993	9,836	325	380	503	685	938	1,089	1,194	1,323	2,285	1
1992	9,227	315	384	501	691	892	1,029	1,156	1,178	2,034	1
1991	9,096	291	359	471	662	910	1,059	1,143	1,192	2,010	-
1990	9,027	249	349	488	697	912	966	1,259	1,157	1,994	3
1989	9,526	278	388	537	782	989	1,063	1,280	1,242	1,982	1
1988	10,368	313	395	582	856	1,093	1,191	1,422	1,357	2,160	6
1987	10,060	301	402	566	831	998	1,288	1,377	1,348	1,979	5
1986	9,793	285	413	565	817	1,008	1,231	1,355	1,233	1,892	5
1985	9,034	256	390	583	760	943	1,189	1,233	1,102	1,652	4
1984	8,254	253	365	515	728	845	1,085	1,056	1,076	1,449	4
1983	7,793	222	376	528	706	836	966	1,011	985	1,313	5
1982	6,873	220	342	466	601	747	888	936	820	1,048	6
1981	6,678	277	356	494	622	731	809	854	762	1,008	2
1980	6,389	217	329	429	550	769	790	825	681	969	3
1979	5,616	214	313	380	546	647	738	760	559	716	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
All races, both sexes											
1998	1,055	74	71	71	80	88	146	125	127	116	-
1997	981	51	66	57	80	102	115	140	103	95	-
1996	843	60	50	47	43	86	88	115	99	96	-
1995	979	59	52	66	82	90	115	123	110	105	-
1994	820	39	50	45	56	67	112	107	95	72	1
1993	892	44	45	50	56	95	104	113	107	109	-
1992	886	58	32	57	66	80	106	114	105	97	-
1991	871	46	39	61	74	95	117	119	78	88	-
1990	921	47	38	48	72	119	109	111	90	105	-
1989	955	39	46	48	73	84	102	114	106	100	-
1988	962	39	42	47	85	97	117	119	94	87	2
1987	876	35	52	57	84	82	118	113	79	72	-
1986	886	36	28	59	83	96	116	103	100	75	-
1985	927	48	46	64	82	100	122	127	83	77	-
1984	883	30	45	83	76	106	113	104	69	84	-
1983	815	30	57	74	93	91	100	88	58	70	-
1982	861	40	46	76	101	98	100	97	56	67	-
1981	765	38	51	69	72	101	83	95	44	58	-
1980	743	29	42	77	91	82	89	76	71	36	-
1979	769	43	47	60	82	100	96	72	64	46	-
Acute and subacute endocarditis (421)											
All races, male											
1998	604	48	55	41	41	52	83	66	66	45	-
1997	550	31	38	31	41	60	66	79	57	31	-
1996	480	46	35	30	27	49	48	56	47	35	-
1995	552	44	36	44	45	46	59	65	54	37	-
1994	458	30	38	32	31	34	58	58	52	24	1
1993	471	24	24	28	34	54	61	59	43	44	-
1992	512	45	21	43	49	46	56	54	53	38	-
1991	502	32	27	37	47	53	72	59	33	35	-
1990	541	31	27	31	44	74	66	62	40	45	-
1989	535	28	28	32	52	58	57	45	43	36	-
1988	565	28	27	38	49	57	60	60	45	34	2
1987	519	27	42	35	62	46	70	54	31	30	-
1986	527	28	24	43	56	62	66	48	45	34	-
1985	555	30	36	39	56	67	75	68	35	28	-
1984	510	22	31	50	40	71	72	47	30	28	-
1983	490	20	41	44	67	60	57	43	25	28	-
1982	528	28	33	46	63	62	67	48	27	27	-
1981	459	26	33	48	48	64	51	48	15	23	-
1980	439	23	25	54	68	50	46	38	32	15	-
1979	485	29	34	47	48	69	63	38	27	22	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
All races, female											
1998	451	26	16	30	39	36	63	59	61	71	-
1997	431	20	28	26	39	42	49	61	46	64	-
1996	363	14	15	17	16	37	40	59	52	61	-
1995	427	15	16	22	37	44	56	58	56	68	-
1994	362	9	12	13	25	33	54	49	43	48	-
1993	421	20	21	22	22	41	43	54	64	65	-
1992	374	13	11	14	17	34	50	60	52	59	-
1991	369	14	12	24	27	42	45	60	45	53	-
1990	380	16	11	17	28	45	43	49	50	60	-
1989	420	11	18	16	21	26	45	69	63	64	-
1988	397	11	15	9	36	40	57	59	49	53	-
1987	357	8	10	22	22	36	48	59	48	42	-
1986	359	8	4	16	27	34	50	55	55	41	-
1985	372	18	10	25	26	33	47	59	48	49	-
1984	373	8	14	33	36	35	41	57	39	56	-
1983	325	10	16	30	26	31	43	45	33	42	-
1982	333	12	13	30	38	36	33	49	29	40	-
1981	306	12	18	21	24	37	32	47	29	35	-
1980	304	6	17	23	23	32	43	38	39	21	-
1979	284	14	13	13	34	31	33	34	37	24	-
Acute and subacute endocarditis (421)											
White, both sexes											
1998	827	41	40	48	66	73	124	108	119	101	-
1997	795	34	45	46	62	83	91	128	94	94	-
1996	659	37	28	34	24	70	73	103	95	84	-
1995	798	38	35	51	65	73	102	112	103	99	-
1994	655	24	35	31	47	54	103	101	83	65	1
1993	702	28	28	34	44	81	90	102	97	103	-
1992	699	32	17	40	57	70	93	102	92	89	-
1991	695	28	26	44	60	78	101	111	73	83	-
1990	724	33	24	40	56	99	94	96	83	98	-
1989	717	20	27	37	59	74	88	109	94	88	-
1988	744	23	28	33	68	85	105	102	89	80	1
1987	692	23	33	44	66	71	109	103	73	63	-
1986	710	24	23	49	70	84	102	92	91	67	-
1985	746	31	35	50	76	90	106	118	76	68	-
1984	714	14	33	68	70	98	98	93	67	77	-
1983	648	20	41	59	79	79	89	77	53	65	-
1982	698	25	32	61	86	87	89	90	50	61	-
1981	627	26	39	53	65	88	77	88	40	52	-
1980	628	23	34	67	76	75	85	71	63	36	-
1979	626	29	38	51	73	85	86	60	60	41	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
White, male											
1998	467	28	31	29	32	41	73	58	62	42	-
1997	448	23	26	25	34	50	53	75	53	31	-
1996	382	32	22	23	15	41	42	53	45	31	-
1995	432	29	25	32	34	36	51	58	51	35	-
1994	370	20	25	22	27	27	52	55	49	22	1
1993	362	15	16	19	26	48	53	53	37	41	-
1992	399	25	12	30	44	40	51	48	47	34	-
1991	400	20	20	26	42	44	67	58	32	35	-
1990	431	22	15	26	38	60	61	58	37	44	-
1989	393	16	18	24	44	52	51	41	36	31	-
1988	418	17	17	28	38	49	51	48	44	31	1
1987	403	19	28	28	52	39	65	49	27	27	-
1986	419	18	20	36	48	56	57	47	41	29	-
1985	438	18	27	33	50	57	65	64	31	24	-
1984	397	10	23	40	35	65	65	42	29	26	-
1983	387	13	31	37	56	54	50	40	24	27	-
1982	421	17	22	39	56	57	58	44	23	23	-
1981	370	16	25	38	41	56	47	44	13	22	-
1980	363	18	21	48	55	43	42	35	28	15	-
1979	386	19	27	39	43	59	55	28	26	19	-
Acute and subacute endocarditis (421)											
White, female											
1998	360	13	9	19	34	32	51	50	57	59	-
1997	347	11	19	21	28	33	38	53	41	63	-
1996	277	5	6	11	9	29	31	50	50	53	-
1995	366	9	10	19	31	37	51	54	52	64	-
1994	285	4	10	9	20	27	51	46	34	43	-
1993	340	13	12	15	18	33	37	49	60	62	-
1992	300	7	5	10	13	30	42	54	45	55	-
1991	295	8	6	18	18	34	34	53	41	48	-
1990	293	11	9	14	18	39	33	38	46	54	-
1989	324	4	9	13	15	22	37	68	58	57	-
1988	326	6	11	5	30	36	54	54	45	49	-
1987	289	4	5	16	14	32	44	54	46	36	-
1986	291	6	3	13	22	28	45	45	50	38	-
1985	308	13	8	17	26	33	41	54	45	44	-
1984	317	4	10	28	35	33	33	51	38	51	-
1983	261	7	10	22	23	25	39	37	29	38	-
1982	277	8	10	22	30	30	31	46	27	38	-
1981	257	10	14	15	24	32	30	44	27	30	-
1980	265	5	13	19	21	32	43	36	35	21	-
1979	240	10	11	12	30	26	31	32	34	22	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
All other, both sexes											
1998	228	33	31	23	14	15	22	17	8	15	-
1997	186	17	21	11	18	19	24	12	9	1	-
1996	184	23	22	13	19	16	15	12	4	12	-
1995	181	21	17	15	17	17	13	11	7	6	-
1994	165	15	15	14	9	13	9	6	12	7	-
1993	190	16	17	16	12	14	14	11	10	6	-
1992	187	26	15	17	9	10	13	12	13	8	-
1991	176	18	13	17	14	17	16	8	5	5	-
1990	197	14	14	8	16	20	15	15	7	7	-
1989	238	19	19	11	14	10	14	5	12	12	-
1988	218	16	14	14	17	12	12	17	5	7	1
1987	184	12	19	13	18	11	9	10	6	9	-
1986	176	12	5	10	13	12	14	11	9	8	-
1985	181	17	11	14	6	10	16	9	7	9	-
1984	169	16	12	15	6	8	15	11	2	7	-
1983	167	10	16	15	14	12	11	11	5	5	-
1982	163	15	14	15	15	11	11	7	6	6	-
1981	138	12	12	16	7	13	6	7	4	6	-
1980	115	6	8	10	15	7	4	5	8	-	-
1979	143	14	9	9	9	15	10	12	4	5	-
Acute and subacute endocarditis (421)											
All other, male											
1998	137	20	24	12	9	11	10	8	4	3	-
1997	102	8	12	6	7	10	13	4	4	-	-
1996	98	14	13	7	12	8	6	3	2	4	-
1995	120	15	11	12	11	10	8	7	3	2	-
1994	88	10	13	10	4	7	6	3	3	2	-
1993	109	9	8	9	8	6	8	6	6	3	-
1992	113	20	9	13	5	6	5	6	6	4	-
1991	102	12	7	11	5	9	5	1	1	-	-
1990	110	9	12	5	6	14	5	4	3	1	-
1989	142	12	10	8	8	6	6	4	7	5	-
1988	147	11	10	10	11	8	9	12	1	3	1
1987	116	8	14	7	10	7	5	5	4	3	-
1986	108	10	4	7	8	6	9	1	4	5	-
1985	117	12	9	6	6	10	10	4	4	4	-
1984	113	12	8	10	5	6	7	5	1	2	-
1983	103	7	10	7	11	6	7	3	1	1	-
1982	107	11	11	7	7	5	9	4	4	4	-
1981	89	10	8	10	7	8	4	4	2	1	-
1980	76	5	4	6	13	7	4	3	4	-	-
1979	99	10	7	8	5	10	8	10	1	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
All other, female											
1998	91	13	7	11	5	4	12	9	4	12	-
1997	84	9	9	5	11	9	11	8	5	1	-
1996	86	9	9	6	7	8	9	9	2	8	-
1995	61	6	6	3	6	7	5	4	4	4	-
1994	77	5	2	4	5	6	3	3	9	5	-
1993	81	7	9	7	4	8	6	5	4	3	-
1992	74	6	6	4	4	4	8	6	7	4	-
1991	74	6	6	6	9	8	11	7	4	5	-
1990	87	5	2	3	10	6	10	11	4	6	-
1989	96	7	9	3	6	4	8	1	5	7	-
1988	71	5	4	4	6	4	3	5	4	4	-
1987	68	4	5	6	8	4	4	5	2	6	-
1986	68	2	1	3	5	6	5	10	5	3	-
1985	64	5	2	8	-	-	6	5	3	5	-
1984	56	4	4	5	1	2	8	6	1	5	-
1983	64	3	6	8	3	6	4	8	4	4	-
1982	56	4	3	8	8	6	2	3	2	2	-
1981	49	2	4	6	-	5	2	3	2	5	-
1980	39	1	4	4	2	-	-	2	4	-	-
1979	44	4	2	1	4	5	2	2	3	2	-
Acute and subacute endocarditis (421)											
Black, both sexes											
1998	214	32	30	21	13	14	18	17	7	15	-
1997	165	14	17	9	15	19	24	10	8	1	-
1996	165	21	21	11	17	14	15	8	4	10	-
1995	162	18	16	11	15	15	12	9	6	5	-
1994	138	13	13	14	8	9	6	3	8	5	-
1993	173	14	16	14	11	14	14	8	9	5	-
1992	181	26	15	16	9	9	12	11	13	8	-
1991	160	18	11	15	11	16	13	7	4	4	-
1990	183	14	14	8	15	19	12	14	6	7	-
1989	220	19	19	11	13	8	13	5	8	11	-
1988	205	16	14	13	17	11	12	13	4	7	1
1987	174	12	19	13	16	11	7	10	5	7	-
1986	170	12	5	9	12	12	14	11	7	8	-
1985	169	17	10	11	5	9	15	9	7	8	-
1984	156	16	12	13	6	8	14	8	1	6	-
1983	153	9	15	13	14	12	10	10	4	5	-
1982	153	13	14	14	14	9	10	7	6	5	-
1981	133	12	12	16	6	13	5	6	3	6	-
1980	101	6	7	10	14	4	2	4	6	-	-
1979	131	13	8	7	9	14	9	11	3	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and subacute endocarditis (421)											
Black, male											
1998	128	19	23	10	8	11	9	8	4	3	-
1997	92	6	11	5	6	10	13	4	3	-	-
1996	87	12	12	6	10	7	6	2	2	3	-
1995	107	13	11	8	10	10	8	5	2	1	-
1994	76	10	11	10	3	5	4	2	3	1	-
1993	96	7	8	7	7	6	8	4	5	2	-
1992	110	20	9	12	5	6	5	5	6	4	-
1991	94	12	6	9	3	8	5	1	1	-	-
1990	102	9	12	5	6	14	4	4	2	1	-
1989	132	12	10	8	8	5	6	4	3	4	-
1988	138	11	10	9	11	7	9	8	1	3	1
1987	110	8	14	7	9	7	4	5	3	2	-
1986	102	10	4	6	7	6	9	1	2	5	-
1985	108	12	9	5	5	9	9	4	4	3	-
1984	108	12	8	9	5	6	7	4	1	2	-
1983	95	7	9	7	11	6	6	3	-	1	-
1982	101	11	11	7	6	4	8	4	4	3	-
1981	86	10	8	10	6	8	3	4	1	1	-
1980	64	5	4	6	12	4	2	2	3	-	-
1979	89	10	6	6	5	9	7	9	1	3	-
Acute and subacute endocarditis (421)											
Black, female											
1998	86	13	7	11	5	3	9	9	3	12	-
1997	73	8	6	4	9	9	11	6	5	1	-
1996	78	9	9	5	7	7	9	6	2	7	-
1995	55	5	5	3	5	5	4	4	4	4	-
1994	62	3	2	4	5	4	2	1	5	4	-
1993	77	7	8	7	4	8	6	4	4	3	-
1992	71	6	6	4	4	3	7	6	7	4	-
1991	66	6	5	6	8	8	8	6	3	4	-
1990	81	5	2	3	9	5	8	10	4	6	-
1989	88	7	9	3	5	3	7	1	5	7	-
1988	67	5	4	4	6	4	3	5	3	4	-
1987	64	4	5	6	7	4	3	5	2	5	-
1986	68	2	1	3	5	6	5	10	5	3	-
1985	61	5	1	6	-	-	6	5	3	5	-
1984	48	4	4	4	1	2	7	4	-	4	-
1983	58	2	6	6	3	6	4	7	4	4	-
1982	52	2	3	7	8	5	2	3	2	2	-
1981	47	2	4	6	-	5	2	2	2	5	-
1980	37	1	3	4	2	-	-	2	3	-	-
1979	42	3	2	1	4	5	2	2	2	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
All races, both sexes											
1998	818	59	47	42	55	65	68	83	72	61	-
1997	854	47	40	52	71	72	74	79	69	59	-
1996	726	47	39	38	35	61	80	68	68	73	-
1995	770	50	53	40	56	65	86	65	49	54	-
1994	785	54	45	37	59	56	77	65	55	54	-
1993	847	44	41	62	56	81	89	80	59	55	1
1992	811	41	35	48	53	80	82	79	60	46	-
1991	775	37	33	51	54	73	86	58	53	58	-
1990	815	41	37	50	70	93	64	71	55	46	-
1989	828	45	34	59	73	92	78	75	56	53	-
1988	847	44	42	42	86	72	79	81	63	58	-
1987	899	35	51	60	86	92	93	71	69	57	-
1986	966	45	44	65	97	106	93	83	64	53	-
1985	878	35	54	65	82	89	85	74	56	63	-
1984	929	35	54	67	95	96	92	68	57	61	1
1983	783	42	57	67	74	94	96	58	39	31	-
1982	803	28	61	66	75	77	83	66	46	39	-
1981	858	48	56	77	91	85	78	77	51	43	-
1980	876	42	59	74	71	111	68	78	42	41	-
1979	744	23	40	72	81	83	58	53	44	29	-
Diseases of pericardium and acute myocarditis (420,422-423)											
All races, male											
1998	405	29	27	18	27	37	34	42	38	16	-
1997	452	28	23	33	46	42	41	44	26	24	-
1996	383	32	25	19	18	36	41	32	26	29	-
1995	389	23	32	28	32	36	44	29	19	18	-
1994	404	27	22	21	27	39	44	29	27	21	-
1993	416	30	25	32	27	36	48	36	22	16	-
1992	444	21	20	30	30	47	36	31	36	21	-
1991	411	24	21	27	30	42	47	27	24	21	-
1990	449	24	22	28	43	56	39	38	30	18	-
1989	447	28	21	32	45	61	44	32	23	18	-
1988	447	29	23	23	55	35	43	44	23	18	-
1987	488	15	29	35	47	54	58	34	26	23	-
1986	505	23	27	37	58	58	47	40	25	17	-
1985	471	24	38	35	42	45	47	41	22	19	-
1984	472	20	31	42	51	46	48	31	23	29	1
1983	433	19	31	41	52	44	55	26	18	15	-
1982	453	21	37	36	43	45	47	36	20	16	-
1981	456	26	34	53	50	40	36	41	25	14	-
1980	491	26	37	45	46	61	36	38	18	25	-
1979	398	9	21	45	50	41	29	24	20	17	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
All races, female											
1998	413	30	20	24	28	28	34	41	34	45	-
1997	402	19	17	19	25	30	33	35	43	35	-
1996	343	15	14	19	17	25	39	36	42	44	-
1995	381	27	21	12	24	29	42	36	30	36	-
1994	381	27	23	16	32	17	33	36	28	33	-
1993	431	14	16	30	29	45	41	44	37	39	1
1992	367	20	15	18	23	33	46	48	24	25	-
1991	364	13	12	24	24	31	39	31	29	37	-
1990	366	17	15	22	27	37	25	33	25	28	-
1989	381	17	13	27	28	31	34	43	33	35	-
1988	400	15	19	19	31	37	36	37	40	40	-
1987	411	20	22	25	39	38	35	37	43	34	-
1986	461	22	17	28	39	48	46	43	39	36	-
1985	407	11	16	30	40	44	38	33	34	44	-
1984	457	15	23	25	44	50	44	37	34	32	-
1983	350	23	26	26	22	50	41	32	21	16	-
1982	350	7	24	30	32	32	36	30	26	23	-
1981	402	22	22	24	41	45	42	36	26	29	-
1980	385	16	22	29	25	50	32	40	24	16	-
1979	346	14	19	27	31	42	29	29	24	12	-
Diseases of pericardium and acute myocarditis (420,422-423)											
White, both sexes											
1998	641	45	35	34	46	53	62	76	61	58	-
1997	694	34	33	41	65	66	67	70	66	54	-
1996	586	27	32	33	30	55	71	61	65	66	-
1995	594	44	41	30	47	53	68	59	42	50	-
1994	617	45	36	34	55	45	69	58	53	49	-
1993	660	29	29	42	47	72	82	70	52	47	1
1992	616	28	25	36	43	73	70	68	57	41	-
1991	587	25	24	42	38	59	78	51	45	55	-
1990	638	36	31	40	60	79	53	63	50	44	-
1989	651	35	31	52	57	73	66	67	52	47	-
1988	679	32	34	33	72	62	72	67	54	52	-
1987	719	21	37	52	74	81	85	60	64	53	-
1986	754	35	33	47	86	87	77	73	54	48	-
1985	678	27	38	49	65	79	75	60	50	58	-
1984	724	29	34	53	76	87	79	62	51	56	-
1983	621	30	48	50	59	82	84	53	34	29	-
1982	637	21	44	49	61	69	72	61	42	37	-
1981	694	34	37	64	78	73	72	70	46	39	-
1980	679	31	47	60	55	96	56	67	38	35	-
1979	591	20	29	55	71	71	48	46	39	25	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
White, male											
1998	314	19	21	14	24	29	31	39	33	15	-
1997	371	20	19	25	41	38	39	40	26	23	-
1996	314	21	20	16	16	34	37	31	25	28	-
1995	298	20	22	20	25	31	35	25	16	18	-
1994	325	21	20	18	25	33	39	27	26	17	-
1993	320	17	20	22	24	33	46	33	19	15	-
1992	339	10	17	22	25	43	33	26	35	19	-
1991	317	18	15	22	23	34	44	24	20	21	-
1990	357	23	19	22	37	46	31	35	28	18	-
1989	350	25	18	28	34	51	38	27	21	17	-
1988	360	22	18	19	48	32	38	38	18	17	-
1987	390	10	20	30	38	49	53	30	24	21	-
1986	408	19	22	30	53	47	43	34	19	14	-
1985	369	20	26	29	38	39	42	33	20	16	-
1984	367	18	20	33	41	42	43	28	21	26	-
1983	348	15	27	30	44	38	47	22	18	14	-
1982	367	17	26	29	34	41	40	31	17	15	-
1981	366	18	21	44	45	35	34	38	22	13	-
1980	370	17	27	36	35	51	29	31	16	20	-
1979	303	8	15	34	42	34	22	20	17	14	-
Diseases of pericardium and acute myocarditis (420,422-423)											
White, female											
1998	327	26	14	20	22	24	31	37	28	43	-
1997	323	14	14	16	24	28	28	30	40	31	-
1996	272	6	12	17	14	21	34	30	40	38	-
1995	296	24	19	10	22	22	33	34	26	32	-
1994	292	24	16	16	30	12	30	31	27	32	-
1993	340	12	9	20	23	39	36	37	33	32	1
1992	277	18	8	14	18	30	37	42	22	22	-
1991	270	7	9	20	15	25	34	27	25	34	-
1990	281	13	12	18	23	33	22	28	22	26	-
1989	301	10	13	24	23	22	28	40	31	30	-
1988	319	10	16	14	24	30	34	29	36	35	-
1987	329	11	17	22	36	32	32	30	40	32	-
1986	346	16	11	17	33	40	34	39	35	34	-
1985	309	7	12	20	27	40	33	27	30	42	-
1984	357	11	14	20	35	45	36	34	30	30	-
1983	273	15	21	20	15	44	37	31	16	15	-
1982	270	4	18	20	27	28	32	30	25	22	-
1981	328	16	16	20	33	38	38	32	24	26	-
1980	309	14	20	24	20	45	27	36	22	15	-
1979	288	12	14	21	29	37	26	26	22	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
All other, both sexes											
1998	177	14	12	8	9	12	6	7	11	3	-
1997	160	13	7	11	6	6	7	9	3	5	-
1996	140	20	7	5	5	6	9	7	3	7	-
1995	176	6	12	10	9	12	18	6	7	4	-
1994	168	9	9	3	4	11	8	7	2	5	-
1993	187	15	12	20	9	9	7	10	7	8	-
1992	195	13	10	12	10	7	12	11	3	5	-
1991	188	12	9	9	16	14	8	7	8	3	-
1990	177	5	6	10	10	14	11	8	5	2	-
1989	177	10	3	7	16	19	12	8	4	6	-
1988	168	12	8	9	14	10	7	14	9	6	-
1987	180	14	14	8	12	11	8	11	5	4	-
1986	212	10	11	18	11	19	16	10	10	5	-
1985	200	8	16	16	17	10	10	14	6	5	-
1984	205	6	20	14	19	9	13	6	6	5	1
1983	162	12	9	17	15	12	12	5	5	2	-
1982	166	7	17	17	14	8	11	5	4	2	-
1981	164	14	19	13	13	12	6	7	5	4	-
1980	197	11	12	14	16	15	12	11	4	6	-
1979	153	3	11	17	10	12	10	7	5	4	-
Diseases of pericardium and acute myocarditis (420,422-423)											
All other, male											
1998	91	10	6	4	3	8	3	3	5	1	-
1997	81	8	4	8	5	4	2	4	-	1	-
1996	69	11	5	3	2	2	4	1	1	1	-
1995	91	3	10	8	7	5	9	4	3	-	-
1994	79	6	2	3	2	6	5	2	1	4	-
1993	96	13	5	10	3	3	2	3	3	1	-
1992	105	11	3	8	5	4	3	5	1	2	-
1991	94	6	6	5	7	8	3	3	4	-	-
1990	92	1	3	6	6	10	8	3	2	-	-
1989	97	3	3	4	11	10	6	5	2	1	-
1988	87	7	5	4	7	3	5	6	5	1	-
1987	98	5	9	5	9	5	5	4	2	2	-
1986	97	4	5	7	5	11	4	6	6	3	-
1985	102	4	12	6	4	6	5	8	2	3	-
1984	105	2	11	9	10	4	5	3	2	3	1
1983	85	4	4	11	8	6	8	4	-	1	-
1982	86	4	11	7	9	4	7	5	3	1	-
1981	90	8	13	9	5	5	2	3	3	1	-
1980	121	9	10	9	11	10	7	7	2	5	-
1979	95	1	6	11	8	7	7	4	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
All other, female											
1998	86	4	6	4	6	4	3	4	6	2	-
1997	79	5	3	3	1	2	5	5	3	4	-
1996	71	9	2	2	3	4	5	6	2	6	-
1995	85	3	2	2	2	7	9	2	4	4	-
1994	89	3	7	-	2	5	3	5	1	1	-
1993	91	2	7	10	6	6	5	7	4	7	-
1992	90	2	7	4	5	3	9	6	2	3	-
1991	94	6	3	4	9	6	5	4	4	3	-
1990	85	4	3	4	4	4	3	5	3	2	-
1989	80	7	-	3	5	9	6	3	2	5	-
1988	81	5	3	5	7	7	2	8	4	5	-
1987	82	9	5	3	3	6	3	7	3	2	-
1986	115	6	6	11	6	8	12	4	4	2	-
1985	98	4	4	10	13	4	5	6	4	2	-
1984	100	4	9	5	9	5	8	3	4	2	-
1983	77	8	5	6	7	6	4	1	5	1	-
1982	80	3	6	10	5	4	4	-	1	1	-
1981	74	6	6	4	8	7	4	4	2	3	-
1980	76	2	2	5	5	5	5	4	2	1	-
1979	58	2	5	6	2	5	3	3	2	1	-
Diseases of pericardium and acute myocarditis (420,422-423)											
Black, both sexes											
1998	158	12	11	8	7	9	6	6	10	3	-
1997	144	12	6	10	6	6	7	7	1	5	-
1996	119	15	6	5	5	4	9	6	3	7	-
1995	151	5	9	7	9	11	16	6	4	4	-
1994	147	8	8	3	4	10	7	5	2	4	-
1993	155	12	10	17	8	8	7	6	7	7	-
1992	171	11	10	9	10	6	10	8	3	4	-
1991	169	12	8	9	14	12	7	5	7	3	-
1990	156	5	4	8	10	14	9	7	5	2	-
1989	166	10	3	7	15	18	11	7	4	5	-
1988	154	12	7	7	13	9	7	14	8	6	-
1987	153	13	13	5	12	7	7	10	4	4	-
1986	193	9	10	17	11	18	14	10	9	5	-
1985	183	7	16	15	16	8	9	14	4	5	-
1984	188	6	20	13	18	9	12	5	6	3	1
1983	147	9	8	17	14	11	11	5	5	2	-
1982	153	7	17	17	13	7	9	4	3	2	-
1981	151	13	19	13	10	11	5	6	5	4	-
1980	178	9	9	12	16	15	11	8	4	6	-
1979	141	3	11	17	7	12	8	7	5	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of pericardium and acute myocarditis (420,422-423)											
Black, male											
1998	77	9	5	4	2	5	3	2	4	1	-
1997	75	7	4	8	5	4	2	3	-	1	-
1996	58	8	5	3	2	-	4	1	1	1	-
1995	75	3	7	5	7	5	8	4	2	-	-
1994	67	5	2	3	2	6	5	1	1	3	-
1993	76	10	4	8	3	3	2	2	3	1	-
1992	93	9	3	5	5	3	3	4	1	1	-
1991	86	6	5	5	6	8	2	2	3	-	-
1990	79	1	1	5	6	10	6	3	2	-	-
1989	91	3	3	4	10	10	5	4	2	1	-
1988	80	7	4	3	7	3	5	6	4	1	-
1987	87	5	9	4	9	4	4	3	2	2	-
1986	87	3	5	7	5	10	3	6	5	3	-
1985	92	3	12	5	4	5	4	8	2	3	-
1984	98	2	11	9	9	4	5	3	2	2	1
1983	78	4	3	11	8	5	7	4	-	1	-
1982	77	4	11	7	8	3	6	4	2	1	-
1981	83	7	13	9	3	5	2	3	3	1	-
1980	110	7	9	8	11	10	6	5	2	5	-
1979	87	1	6	11	5	7	6	4	3	2	-
Diseases of pericardium and acute myocarditis (420,422-423)											
Black, female											
1998	81	3	6	4	5	4	3	4	6	2	-
1997	69	5	2	2	1	2	5	4	1	4	-
1996	61	7	1	2	3	4	5	5	2	6	-
1995	76	2	2	2	2	6	8	2	2	4	-
1994	80	3	6	-	2	4	2	4	1	1	-
1993	79	2	6	9	5	5	5	4	4	6	-
1992	78	2	7	4	5	3	7	4	2	3	-
1991	83	6	3	4	8	4	5	3	4	3	-
1990	77	4	3	3	4	4	3	4	3	2	-
1989	75	7	-	3	5	8	6	3	2	4	-
1988	74	5	3	4	6	6	2	8	4	5	-
1987	66	8	4	1	3	3	3	7	2	2	-
1986	106	6	5	10	6	8	11	4	4	2	-
1985	91	4	4	10	12	3	5	6	2	2	-
1984	90	4	9	4	9	5	7	2	4	1	-
1983	69	5	5	6	6	6	4	1	5	1	-
1982	76	3	6	10	5	4	3	-	1	1	-
1981	68	6	6	4	7	6	3	3	2	3	-
1980	68	2	-	4	5	5	5	3	2	1	-
1979	54	2	5	6	2	5	2	3	2	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
All races, both sexes											
1998	50,228	248	404	717	1,244	2,220	3,964	6,278	9,180	25,614	1
1997	48,930	204	461	700	1,182	2,129	3,920	6,244	8,992	24,696	3
1996	47,230	252	412	639	1,164	2,233	3,906	5,917	8,711	23,603	9
1995	46,484	263	383	662	1,215	2,221	3,814	5,870	8,708	22,983	1
1994	45,155	205	387	614	1,216	2,273	3,919	5,742	8,461	21,934	2
1993	45,613	242	417	718	1,300	2,403	4,106	6,098	8,433	21,483	6
1992	40,097	227	341	626	1,280	2,258	3,768	5,394	7,390	18,445	4
1991	39,206	223	307	581	1,323	2,270	3,635	5,349	7,248	17,866	4
1990	37,927	213	308	619	1,347	2,249	3,460	5,279	7,204	16,894	2
1989	38,817	210	391	723	1,429	2,421	3,745	5,661	7,333	16,548	2
1988	42,940	232	448	847	1,700	2,744	4,228	6,176	8,095	18,030	3
1987	41,210	233	419	811	1,630	2,808	4,278	6,046	7,721	16,828	1
1986	40,330	243	440	865	1,692	2,731	4,302	6,096	7,571	15,924	19
1985	37,747	232	472	913	1,656	2,629	4,105	5,573	7,096	14,609	18
1984	35,355	214	407	896	1,622	2,522	3,828	5,398	6,621	13,389	6
1983	33,227	221	473	865	1,558	2,510	3,689	5,026	6,069	12,374	3
1982	30,657	228	423	848	1,527	2,337	3,487	4,649	5,732	10,996	6
1981	28,972	239	491	913	1,489	2,346	3,359	4,385	5,338	9,946	7
1980	27,415	258	504	879	1,405	2,397	3,291	4,281	4,852	9,064	8
1979	25,074	246	512	917	1,444	2,272	2,979	3,848	4,539	7,879	4
Heart failure (428)											
All races, male											
1998	19,074	158	245	402	738	1,256	2,050	2,970	3,874	7,164	-
1997	18,621	133	293	407	694	1,171	1,987	2,994	3,736	6,940	-
1996	18,224	153	264	373	669	1,280	2,094	2,890	3,560	6,714	6
1995	17,752	172	239	407	709	1,242	1,998	2,756	3,592	6,429	-
1994	17,319	134	241	364	725	1,249	2,072	2,686	3,364	6,225	1
1993	17,586	159	258	423	773	1,317	2,084	2,838	3,490	6,017	2
1992	15,646	145	209	368	723	1,210	2,010	2,611	2,979	5,173	2
1991	15,348	141	176	344	802	1,299	1,883	2,570	2,874	5,025	1
1990	15,089	136	188	385	780	1,294	1,871	2,481	2,949	4,784	1
1989	15,556	153	243	427	793	1,399	1,960	2,683	2,947	4,744	1
1988	17,612	132	289	518	980	1,535	2,312	2,921	3,300	5,347	-
1987	16,941	144	256	482	955	1,582	2,279	2,920	3,098	4,948	-
1986	16,955	167	259	540	990	1,576	2,323	2,919	3,135	4,769	2
1985	15,860	146	299	564	962	1,489	2,201	2,656	2,908	4,356	7
1984	15,227	134	255	576	970	1,472	2,050	2,586	2,673	4,236	2
1983	14,515	139	301	535	937	1,443	2,054	2,474	2,465	3,905	1
1982	13,590	138	281	513	946	1,375	1,827	2,239	2,366	3,649	5
1981	13,049	158	319	552	893	1,333	1,823	2,167	2,202	3,311	3
1980	12,599	169	326	561	857	1,386	1,794	2,123	2,021	3,065	4
1979	11,664	170	331	587	877	1,350	1,603	1,849	1,909	2,721	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
All races, female											
1998	31,154	90	159	315	506	964	1,914	3,308	5,306	18,450	1
1997	30,309	71	168	293	488	958	1,933	3,250	5,256	17,756	3
1996	29,006	99	148	266	495	953	1,812	3,027	5,151	16,889	3
1995	28,732	91	144	255	506	979	1,816	3,114	5,116	16,554	1
1994	27,836	71	146	250	491	1,024	1,847	3,056	5,097	15,709	1
1993	28,027	83	159	295	527	1,086	2,022	3,260	4,943	15,466	4
1992	24,451	82	132	258	557	1,048	1,758	2,783	4,411	13,272	2
1991	23,858	82	131	237	521	971	1,752	2,779	4,374	12,841	3
1990	22,838	77	120	234	567	955	1,589	2,798	4,255	12,110	1
1989	23,261	57	148	296	636	1,022	1,785	2,978	4,386	11,804	1
1988	25,328	100	159	329	720	1,209	1,916	3,255	4,795	12,683	3
1987	24,269	89	163	329	675	1,226	1,999	3,126	4,623	11,880	1
1986	23,375	76	181	325	702	1,155	1,979	3,177	4,436	11,155	17
1985	21,887	86	173	349	694	1,140	1,904	2,917	4,188	10,253	11
1984	20,128	80	152	320	652	1,050	1,778	2,812	3,948	9,153	4
1983	18,712	82	172	330	621	1,067	1,635	2,552	3,604	8,469	2
1982	17,067	90	142	335	581	962	1,660	2,410	3,366	7,347	1
1981	15,923	81	172	361	596	1,013	1,536	2,218	3,136	6,635	4
1980	14,816	89	178	318	548	1,011	1,497	2,158	2,831	5,999	4
1979	13,410	76	181	330	567	922	1,376	1,999	2,630	5,158	2
Heart failure (428)											
White, both sexes											
1998	45,260	150	288	527	957	1,836	3,424	5,595	8,366	23,888	1
1997	44,148	141	319	510	900	1,770	3,401	5,582	8,245	23,006	3
1996	42,563	153	312	470	900	1,864	3,368	5,284	7,964	21,989	6
1995	41,907	172	267	492	965	1,874	3,285	5,211	8,008	21,414	1
1994	40,729	138	288	449	966	1,917	3,374	5,103	7,727	20,514	2
1993	40,967	161	282	532	1,029	1,976	3,505	5,481	7,679	20,053	4
1992	35,953	156	243	471	1,031	1,864	3,249	4,786	6,735	17,199	2
1991	35,055	149	216	447	1,069	1,854	3,135	4,733	6,548	16,631	4
1990	33,802	143	220	458	1,077	1,837	3,007	4,628	6,524	15,672	1
1989	34,408	152	272	532	1,126	2,001	3,189	4,969	6,585	15,363	2
1988	38,237	155	316	656	1,367	2,294	3,644	5,401	7,325	16,780	1
1987	36,746	165	308	661	1,301	2,314	3,679	5,326	7,007	15,690	-
1986	35,712	167	321	642	1,338	2,274	3,683	5,376	6,829	14,791	14
1985	33,400	166	328	696	1,325	2,185	3,480	4,916	6,412	13,594	13
1984	31,308	143	301	694	1,325	2,097	3,287	4,750	5,951	12,454	3
1983	29,428	168	359	670	1,239	2,091	3,148	4,457	5,467	11,540	-
1982	27,162	147	315	662	1,257	1,926	3,005	4,127	5,161	10,266	3
1981	25,592	167	378	712	1,176	1,971	2,876	3,850	4,854	9,275	3
1980	24,009	186	381	688	1,130	1,972	2,813	3,737	4,385	8,402	4
1979	21,948	178	400	700	1,148	1,862	2,536	3,351	4,137	7,325	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
White, male											
1998	17,035	97	179	310	584	1,058	1,802	2,680	3,561	6,623	-
1997	16,715	96	214	304	543	992	1,756	2,713	3,454	6,456	-
1996	16,317	95	204	292	526	1,092	1,840	2,601	3,296	6,213	4
1995	15,872	117	167	319	574	1,065	1,756	2,466	3,320	5,957	-
1994	15,481	94	187	279	584	1,079	1,811	2,417	3,083	5,787	1
1993	15,660	107	176	323	620	1,114	1,796	2,577	3,212	5,572	1
1992	13,921	102	151	292	589	1,039	1,747	2,350	2,733	4,785	1
1991	13,622	100	129	275	671	1,087	1,661	2,279	2,604	4,658	1
1990	13,314	95	145	303	643	1,083	1,656	2,196	2,651	4,396	-
1989	13,678	116	175	332	657	1,171	1,703	2,376	2,657	4,360	1
1988	15,570	86	210	419	822	1,304	2,043	2,583	2,984	4,920	-
1987	14,955	108	195	407	788	1,313	2,003	2,586	2,812	4,540	-
1986	14,932	118	205	421	816	1,341	2,042	2,601	2,829	4,389	1
1985	13,911	108	222	444	797	1,277	1,902	2,368	2,593	4,018	5
1984	13,396	97	194	472	829	1,258	1,790	2,292	2,397	3,873	-
1983	12,755	102	235	433	768	1,223	1,775	2,228	2,203	3,615	-
1982	11,967	96	218	415	790	1,140	1,610	2,006	2,136	3,383	3
1981	11,449	118	259	446	721	1,151	1,580	1,927	1,981	3,052	1
1980	10,965	128	260	442	700	1,164	1,551	1,881	1,822	2,818	2
1979	10,181	125	277	471	719	1,126	1,414	1,615	1,734	2,500	1
Heart failure (428)											
White, female											
1998	28,225	53	109	217	373	778	1,622	2,915	4,805	17,265	1
1997	27,433	45	105	206	357	778	1,645	2,869	4,791	16,550	3
1996	26,246	58	108	178	374	772	1,528	2,683	4,668	15,776	2
1995	26,035	55	100	173	391	809	1,529	2,745	4,688	15,457	1
1994	25,248	44	101	170	382	838	1,563	2,686	4,644	14,727	1
1993	25,307	54	106	209	409	862	1,709	2,904	4,467	14,481	3
1992	22,032	54	92	179	442	825	1,502	2,436	4,002	12,414	1
1991	21,433	49	87	172	398	767	1,474	2,454	3,944	11,973	3
1990	20,488	48	75	155	434	754	1,351	2,432	3,873	11,276	1
1989	20,730	36	97	200	469	830	1,486	2,593	3,928	11,003	1
1988	22,667	69	106	237	545	990	1,601	2,818	4,341	11,860	1
1987	21,791	57	113	254	513	1,001	1,676	2,740	4,195	11,150	-
1986	20,780	49	116	221	522	933	1,641	2,775	4,000	10,402	13
1985	19,489	58	106	252	528	908	1,578	2,548	3,819	9,576	8
1984	17,912	46	107	222	496	839	1,497	2,458	3,554	8,581	3
1983	16,673	66	124	237	471	868	1,373	2,229	3,264	7,925	-
1982	15,195	51	97	247	467	786	1,395	2,121	3,025	6,883	-
1981	14,143	49	119	266	455	820	1,296	1,923	2,873	6,223	2
1980	13,044	58	121	246	430	808	1,262	1,856	2,563	5,584	2
1979	11,767	53	123	229	429	736	1,122	1,736	2,403	4,825	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
All other, both sexes											
1998	4,968	98	116	190	287	384	540	683	814	1,726	-
1997	4,782	63	142	190	282	359	519	662	747	1,690	-
1996	4,667	99	100	169	264	369	538	633	747	1,614	3
1995	4,577	91	116	170	250	347	529	659	700	1,569	-
1994	4,426	67	99	165	250	356	545	639	734	1,420	-
1993	4,646	81	135	186	271	427	601	617	754	1,430	2
1992	4,144	71	98	155	249	394	519	608	655	1,246	2
1991	4,151	74	91	134	254	416	500	616	700	1,235	-
1990	4,125	70	88	161	270	412	453	651	680	1,222	1
1989	4,409	58	119	191	303	420	556	692	748	1,185	-
1988	4,703	77	132	191	333	450	584	775	770	1,250	2
1987	4,464	68	111	150	329	494	599	720	714	1,138	1
1986	4,618	76	119	223	354	457	619	720	742	1,133	5
1985	4,347	66	144	217	331	444	625	657	684	1,015	5
1984	4,047	71	106	202	297	425	541	648	670	935	3
1983	3,799	53	114	195	319	419	541	569	602	834	3
1982	3,495	81	108	186	270	411	482	522	571	730	3
1981	3,380	72	113	201	313	375	483	535	484	671	4
1980	3,406	72	123	191	275	425	478	544	467	662	4
1979	3,126	68	112	217	296	410	443	497	402	554	2
Heart failure (428)											
All other, male											
1998	2,039	61	66	92	154	198	248	290	313	541	-
1997	1,906	37	79	103	151	179	231	281	282	484	-
1996	1,907	58	60	81	143	188	254	289	264	501	2
1995	1,880	55	72	88	135	177	242	290	272	472	-
1994	1,838	40	54	85	141	170	261	269	281	438	-
1993	1,926	52	82	100	153	203	288	261	278	445	1
1992	1,725	43	58	76	134	171	263	261	246	388	1
1991	1,726	41	47	69	131	212	222	291	270	367	-
1990	1,775	41	43	82	137	211	215	285	298	388	1
1989	1,878	37	68	95	136	228	257	307	290	384	-
1988	2,042	46	79	99	158	231	269	338	316	427	-
1987	1,986	36	61	75	167	269	276	334	286	408	-
1986	2,023	49	54	119	174	235	281	318	306	380	1
1985	1,949	38	77	120	165	212	299	288	315	338	2
1984	1,831	37	61	104	141	214	260	294	276	363	2
1983	1,760	37	66	102	169	220	279	246	262	290	1
1982	1,623	42	63	98	156	235	217	233	230	266	2
1981	1,600	40	60	106	172	182	243	240	221	259	2
1980	1,634	41	66	119	157	222	243	242	199	247	2
1979	1,483	45	54	116	158	224	189	234	175	221	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
All other, female											
1998	2,929	37	50	98	133	186	292	393	501	1,185	-
1997	2,876	26	63	87	131	180	288	381	465	1,206	-
1996	2,760	41	40	88	121	181	284	344	483	1,113	1
1995	2,697	36	44	82	115	170	287	369	428	1,097	-
1994	2,588	27	45	80	109	186	284	370	453	982	-
1993	2,720	29	53	86	118	224	313	356	476	985	1
1992	2,419	28	40	79	115	223	256	347	409	858	1
1991	2,425	33	44	65	123	204	278	325	430	868	-
1990	2,350	29	45	79	133	201	238	366	382	834	-
1989	2,531	21	51	96	167	192	299	385	458	801	-
1988	2,661	31	53	92	175	219	315	437	454	823	2
1987	2,478	32	50	75	162	225	323	386	428	730	1
1986	2,595	27	65	104	180	222	338	402	436	753	4
1985	2,398	28	67	97	166	232	326	369	369	677	3
1984	2,216	34	45	98	156	211	281	354	394	572	1
1983	2,039	16	48	93	150	199	262	323	340	544	2
1982	1,872	39	45	88	114	176	265	289	341	464	1
1981	1,780	32	53	95	141	193	240	295	263	412	2
1980	1,772	31	57	72	118	203	235	302	268	415	2
1979	1,643	23	58	101	138	186	254	263	227	333	1
Heart failure (428)											
Black, both sexes											
1998	4,493	97	113	174	269	352	494	613	743	1,515	-
1997	4,292	61	132	174	266	334	489	592	647	1,484	-
1996	4,217	94	92	151	239	342	489	582	668	1,433	3
1995	4,156	85	104	158	230	322	483	598	629	1,408	-
1994	4,016	62	92	149	237	331	507	578	656	1,260	-
1993	4,247	77	128	176	258	398	558	561	684	1,274	2
1992	3,783	69	96	145	231	363	477	552	598	1,113	2
1991	3,800	70	85	125	237	389	462	560	636	1,113	-
1990	3,797	68	83	150	253	384	422	602	622	1,104	1
1989	4,082	52	108	187	287	402	520	638	682	1,070	-
1988	4,372	75	123	183	317	422	545	732	708	1,131	2
1987	4,164	63	103	139	310	464	556	679	670	1,045	1
1986	4,324	72	111	214	337	430	588	677	695	1,029	5
1985	4,077	64	137	207	313	432	597	612	638	927	4
1984	3,810	64	103	192	287	409	518	599	626	867	3
1983	3,561	53	110	186	299	400	508	535	561	765	2
1982	3,279	75	104	178	255	384	464	480	528	686	3
1981	3,183	69	110	197	298	361	458	502	454	613	2
1980	3,216	70	119	183	264	402	453	510	443	611	3
1979	2,949	68	107	203	286	384	424	470	378	510	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Heart failure (428)											
Black, male											
1998	1,817	61	65	83	145	178	221	260	275	457	-
1997	1,694	36	74	93	142	166	217	243	241	414	-
1996	1,707	56	55	72	127	174	233	267	239	420	2
1995	1,702	51	64	81	126	170	219	267	237	414	-
1994	1,654	35	49	78	134	158	239	243	251	374	-
1993	1,729	51	77	95	147	189	269	234	239	369	1
1992	1,555	43	56	70	123	160	240	234	215	335	1
1991	1,565	39	45	63	124	196	204	267	236	320	-
1990	1,631	40	41	79	127	198	200	260	266	351	1
1989	1,711	35	59	93	127	220	237	276	255	334	-
1988	1,889	45	73	94	147	220	250	319	285	379	-
1987	1,834	32	56	69	162	254	252	307	260	373	-
1986	1,869	47	52	116	163	217	261	292	280	335	1
1985	1,825	37	72	113	157	208	285	262	294	311	1
1984	1,706	33	60	101	137	208	247	261	250	331	2
1983	1,649	37	66	97	161	214	260	232	241	257	1
1982	1,509	40	60	93	148	218	205	208	212	247	2
1981	1,497	39	59	106	163	176	230	223	204	227	1
1980	1,522	39	65	112	149	207	228	223	185	220	2
1979	1,400	45	51	111	154	211	178	222	163	201	1
Heart failure (428)											
Black, female											
1998	2,676	36	48	91	124	174	273	353	468	1,058	-
1997	2,598	25	58	81	124	168	272	349	406	1,070	-
1996	2,510	38	37	79	112	168	256	315	429	1,013	1
1995	2,454	34	40	77	104	152	264	331	392	994	-
1994	2,362	27	43	71	103	173	268	335	405	886	-
1993	2,518	26	51	81	111	209	289	327	445	905	1
1992	2,228	26	40	75	108	203	237	318	383	778	1
1991	2,235	31	40	62	113	193	258	293	400	793	-
1990	2,166	28	42	71	126	186	222	342	356	753	-
1989	2,371	17	49	94	160	182	283	362	427	736	-
1988	2,483	30	50	89	170	202	295	413	423	752	2
1987	2,330	31	47	70	148	210	304	372	410	672	1
1986	2,455	25	59	98	174	213	327	385	415	694	4
1985	2,252	27	65	94	156	224	312	350	344	616	3
1984	2,104	31	43	91	150	201	271	338	376	536	1
1983	1,912	16	44	89	138	186	248	303	320	508	1
1982	1,770	35	44	85	107	166	259	272	316	439	1
1981	1,686	30	51	91	135	185	228	279	250	386	1
1980	1,694	31	54	71	115	195	225	287	258	391	1
1979	1,549	23	56	92	132	173	246	248	215	309	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All races, both sexes											
1998	111,222	3,281	3,864	4,645	5,988	8,345	11,782	15,294	17,074	33,285	9
1997	109,103	3,200	3,681	4,592	6,118	8,654	11,959	14,908	16,992	31,558	5
1996	106,229	3,121	3,543	4,451	6,105	8,770	11,812	14,455	16,470	30,297	9
1995	104,221	3,016	3,500	4,337	6,393	8,776	12,052	14,121	15,999	28,359	7
1994	101,119	2,817	3,312	4,434	6,133	9,051	12,257	13,679	15,277	26,664	3
1993	102,161	2,649	3,354	4,501	6,529	9,574	12,194	14,461	15,313	26,193	7
1992	96,858	2,562	3,122	4,219	6,650	9,219	11,921	13,736	14,562	23,828	4
1991	95,432	2,335	2,967	4,401	6,719	9,475	11,799	13,797	14,116	22,862	8
1990	92,637	2,276	2,931	4,424	6,903	9,512	11,421	13,426	13,733	21,434	16
1989	94,931	2,234	3,126	4,750	7,192	9,984	11,832	13,774	13,800	21,594	8
1988	108,329	2,398	3,510	5,631	8,567	11,693	13,813	15,744	15,339	24,472	15
1987	103,200	2,397	3,519	5,384	8,348	11,322	13,276	14,808	14,564	22,599	19
1986	100,466	2,275	3,454	5,520	8,496	11,083	13,168	14,521	13,929	21,021	12
1985	92,970	2,180	3,390	5,476	8,128	10,247	12,349	13,291	12,608	18,801	18
1984	84,290	2,082	3,340	5,208	7,689	9,562	11,229	11,654	11,111	16,188	17
1983	79,745	1,930	3,243	5,177	7,342	9,011	10,817	11,117	10,329	14,716	18
1982	71,570	2,002	3,088	4,847	6,773	8,391	9,558	9,887	8,844	12,629	18
1981	67,517	1,962	3,194	4,959	6,386	7,798	8,790	8,904	8,454	11,586	10
1980	63,732	1,815	3,097	4,686	6,043	7,511	8,447	8,407	7,644	10,603	13
1979	56,758	1,803	2,957	4,176	5,515	6,586	7,363	7,499	6,907	9,040	13
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All races, male											
1998	52,277	2,133	2,407	2,837	3,742	4,805	6,375	7,720	7,523	9,990	5
1997	51,926	2,072	2,345	2,886	3,756	5,088	6,610	7,604	7,354	9,576	4
1996	50,830	2,040	2,224	2,769	3,756	5,247	6,562	7,374	7,206	9,161	5
1995	50,742	2,029	2,252	2,671	3,893	5,265	6,770	7,350	6,979	8,632	6
1994	49,320	1,833	2,180	2,871	3,795	5,391	6,889	6,970	6,718	8,006	2
1993	50,040	1,730	2,205	2,886	4,040	5,739	6,938	7,287	6,585	7,978	7
1992	48,049	1,666	2,015	2,735	4,156	5,517	6,727	6,916	6,381	7,425	2
1991	47,764	1,576	1,939	2,790	4,264	5,663	6,743	7,136	6,193	7,048	5
1990	46,506	1,518	1,898	2,807	4,327	5,690	6,469	6,968	6,059	6,568	9
1989	47,838	1,488	2,026	3,071	4,535	6,069	6,720	7,029	5,994	6,694	4
1988	55,181	1,578	2,313	3,652	5,471	7,086	8,061	8,092	6,659	7,681	9
1987	52,770	1,573	2,337	3,480	5,185	6,929	7,592	7,637	6,337	7,277	13
1986	51,991	1,458	2,245	3,581	5,447	6,811	7,569	7,549	6,058	6,829	8
1985	48,646	1,448	2,231	3,580	5,253	6,365	7,070	6,844	5,565	6,259	10
1984	44,688	1,413	2,161	3,419	4,982	5,963	6,446	6,015	4,874	5,533	10
1983	42,385	1,256	2,116	3,383	4,674	5,676	6,278	5,633	4,528	5,124	10
1982	38,021	1,285	2,040	3,155	4,349	5,177	5,445	4,997	3,818	4,402	9
1981	36,435	1,269	2,103	3,274	4,073	4,763	5,063	4,569	3,844	4,193	5
1980	34,510	1,183	2,053	3,096	3,868	4,575	4,880	4,296	3,462	3,768	7
1979	30,922	1,186	1,956	2,780	3,523	4,135	4,200	3,804	3,117	3,319	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All races, female											
1998	58,945	1,148	1,457	1,808	2,246	3,540	5,407	7,574	9,551	23,295	4
1997	57,177	1,128	1,336	1,706	2,362	3,566	5,349	7,304	9,638	21,982	1
1996	55,399	1,081	1,319	1,682	2,349	3,523	5,250	7,081	9,264	21,136	4
1995	53,479	987	1,248	1,666	2,500	3,511	5,282	6,771	9,020	19,727	1
1994	51,799	984	1,132	1,563	2,338	3,660	5,368	6,709	8,559	18,658	1
1993	52,121	919	1,149	1,615	2,489	3,835	5,256	7,174	8,728	18,215	-
1992	48,809	896	1,107	1,484	2,494	3,702	5,194	6,820	8,181	16,403	2
1991	47,668	759	1,028	1,611	2,455	3,812	5,056	6,661	7,923	15,814	3
1990	46,131	758	1,033	1,617	2,576	3,822	4,952	6,458	7,674	14,866	7
1989	47,093	746	1,100	1,679	2,657	3,915	5,112	6,745	7,806	14,900	4
1988	53,148	820	1,197	1,979	3,096	4,607	5,752	7,652	8,680	16,791	6
1987	50,430	824	1,182	1,904	3,163	4,393	5,684	7,171	8,227	15,322	6
1986	48,475	817	1,209	1,939	3,049	4,272	5,599	6,972	7,871	14,192	4
1985	44,324	732	1,159	1,896	2,875	3,882	5,279	6,447	7,043	12,542	8
1984	39,602	669	1,179	1,789	2,707	3,599	4,783	5,639	6,237	10,655	7
1983	37,360	674	1,127	1,794	2,668	3,335	4,539	5,484	5,801	9,592	8
1982	33,549	717	1,048	1,692	2,424	3,214	4,113	4,890	5,026	8,227	9
1981	31,082	693	1,091	1,685	2,313	3,035	3,727	4,335	4,610	7,393	5
1980	29,222	632	1,044	1,590	2,175	2,936	3,567	4,111	4,182	6,835	6
1979	25,836	617	1,001	1,396	1,992	2,451	3,163	3,695	3,790	5,721	7
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
White, both sexes											
1998	94,328	2,243	2,847	3,531	4,661	6,808	9,974	13,399	15,363	30,460	4
1997	92,779	2,226	2,672	3,506	4,820	7,107	10,193	13,153	15,284	28,842	2
1996	89,823	2,143	2,553	3,393	4,712	7,210	9,982	12,717	14,800	27,687	5
1995	87,902	2,099	2,571	3,274	5,004	7,157	10,184	12,413	14,289	25,933	6
1994	85,337	1,986	2,427	3,334	4,877	7,413	10,444	11,997	13,650	24,340	3
1993	86,077	1,856	2,432	3,332	5,162	7,857	10,390	12,666	13,651	23,927	3
1992	81,444	1,770	2,280	3,162	5,272	7,617	10,130	12,015	12,971	21,727	3
1991	80,244	1,622	2,167	3,345	5,368	7,793	10,016	11,993	12,613	20,821	8
1990	77,606	1,602	2,113	3,346	5,519	7,824	9,770	11,608	12,214	19,405	9
1989	79,349	1,535	2,194	3,621	5,715	8,135	10,098	11,907	12,261	19,526	4
1988	90,730	1,659	2,546	4,353	6,839	9,609	11,728	13,611	13,555	22,120	9
1987	86,319	1,696	2,569	4,148	6,677	9,375	11,226	12,759	12,820	20,431	9
1986	84,095	1,595	2,525	4,285	6,856	9,233	11,154	12,470	12,315	19,000	10
1985	77,788	1,552	2,490	4,261	6,570	8,496	10,434	11,473	11,150	17,029	15
1984	70,548	1,475	2,506	4,098	6,224	7,974	9,449	10,158	9,777	14,652	11
1983	66,541	1,394	2,444	4,031	5,940	7,493	9,160	9,590	9,074	13,323	11
1982	60,079	1,452	2,374	3,856	5,478	7,012	8,092	8,590	7,868	11,536	9
1981	56,424	1,385	2,444	3,927	5,195	6,537	7,519	7,760	7,448	10,487	7
1980	52,935	1,310	2,342	3,721	4,930	6,193	7,156	7,279	6,799	9,578	9
1979	47,528	1,305	2,325	3,378	4,520	5,432	6,255	6,462	6,212	8,273	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
White, male											
1998	43,907	1,495	1,807	2,218	2,976	3,992	5,476	6,828	6,816	9,073	2
1997	43,739	1,478	1,751	2,244	3,007	4,261	5,709	6,776	6,630	8,696	1
1996	42,529	1,429	1,652	2,170	2,946	4,381	5,610	6,531	6,481	8,305	3
1995	42,398	1,471	1,688	2,064	3,127	4,372	5,821	6,540	6,231	7,818	5
1994	41,255	1,330	1,639	2,247	3,083	4,492	5,957	6,170	6,025	7,214	2
1993	41,753	1,241	1,617	2,152	3,282	4,792	5,993	6,440	5,887	7,236	3
1992	40,013	1,168	1,516	2,120	3,394	4,641	5,781	6,095	5,640	6,684	1
1991	39,763	1,121	1,465	2,165	3,480	4,726	5,800	6,228	5,534	6,340	5
1990	38,682	1,068	1,405	2,163	3,543	4,767	5,598	6,126	5,394	5,888	5
1989	39,712	1,049	1,444	2,401	3,703	5,058	5,792	6,132	5,318	5,976	1
1988	45,789	1,124	1,709	2,891	4,451	5,928	6,924	7,004	5,862	6,820	7
1987	43,875	1,138	1,747	2,759	4,223	5,798	6,559	6,618	5,563	6,508	7
1986	43,173	1,038	1,679	2,822	4,466	5,774	6,487	6,514	5,297	6,085	7
1985	40,461	1,048	1,662	2,863	4,313	5,365	6,064	5,938	4,900	5,594	8
1984	37,276	1,032	1,658	2,751	4,116	5,042	5,496	5,262	4,268	4,968	8
1983	35,226	925	1,648	2,690	3,859	4,816	5,360	4,842	3,964	4,590	7
1982	31,758	929	1,621	2,551	3,561	4,399	4,626	4,380	3,365	3,953	5
1981	30,445	941	1,664	2,653	3,377	4,055	4,389	4,019	3,369	3,741	3
1980	28,506	868	1,580	2,491	3,200	3,853	4,170	3,729	3,047	3,345	5
1979	25,879	882	1,585	2,276	2,953	3,468	3,603	3,290	2,792	2,987	4
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
White, female											
1998	50,421	748	1,040	1,313	1,685	2,816	4,498	6,571	8,547	21,387	2
1997	49,040	748	921	1,262	1,813	2,846	4,484	6,377	8,654	20,146	1
1996	47,294	714	901	1,223	1,766	2,829	4,372	6,186	8,319	19,382	2
1995	45,504	628	883	1,210	1,877	2,785	4,363	5,873	8,058	18,115	1
1994	44,082	656	788	1,087	1,794	2,921	4,487	5,827	7,625	17,126	1
1993	44,324	615	815	1,180	1,880	3,065	4,397	6,226	7,764	16,691	-
1992	41,431	602	764	1,042	1,878	2,976	4,349	5,920	7,331	15,043	2
1991	40,481	501	702	1,180	1,888	3,067	4,216	5,765	7,079	14,481	3
1990	38,924	534	708	1,183	1,976	3,057	4,172	5,482	6,820	13,517	4
1989	39,637	486	750	1,220	2,012	3,077	4,306	5,775	6,943	13,550	3
1988	44,941	535	837	1,462	2,388	3,681	4,804	6,607	7,693	15,300	2
1987	42,444	558	822	1,389	2,454	3,577	4,667	6,141	7,257	13,923	2
1986	40,922	557	846	1,463	2,390	3,459	4,667	5,956	7,018	12,915	3
1985	37,327	504	828	1,398	2,257	3,131	4,370	5,535	6,250	11,435	7
1984	33,272	443	848	1,347	2,108	2,932	3,953	4,896	5,509	9,684	3
1983	31,315	469	796	1,341	2,081	2,677	3,800	4,748	5,110	8,733	4
1982	28,321	523	753	1,305	1,917	2,613	3,466	4,210	4,503	7,583	4
1981	25,979	444	780	1,274	1,818	2,482	3,130	3,741	4,079	6,746	4
1980	24,429	442	762	1,230	1,730	2,340	2,986	3,550	3,752	6,233	4
1979	21,649	423	740	1,102	1,567	1,964	2,652	3,172	3,420	5,286	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All other, both sexes											
1998	16,894	1,038	1,017	1,114	1,327	1,537	1,808	1,895	1,711	2,825	5
1997	16,324	974	1,009	1,086	1,298	1,547	1,766	1,755	1,708	2,716	3
1996	16,406	978	990	1,058	1,393	1,560	1,830	1,738	1,670	2,610	4
1995	16,319	917	929	1,063	1,389	1,619	1,868	1,708	1,710	2,426	1
1994	15,782	831	885	1,100	1,256	1,638	1,813	1,682	1,627	2,324	-
1993	16,084	793	922	1,169	1,367	1,717	1,804	1,795	1,662	2,266	4
1992	15,414	792	842	1,057	1,378	1,602	1,791	1,721	1,591	2,101	1
1991	15,188	713	800	1,056	1,351	1,682	1,783	1,804	1,503	2,041	-
1990	15,031	674	818	1,078	1,384	1,688	1,651	1,818	1,519	2,029	7
1989	15,582	699	932	1,129	1,477	1,849	1,734	1,867	1,539	2,068	4
1988	17,599	739	964	1,278	1,728	2,084	2,085	2,133	1,784	2,352	6
1987	16,881	701	950	1,236	1,671	1,947	2,050	2,049	1,744	2,168	10
1986	16,371	680	929	1,235	1,640	1,850	2,014	2,051	1,614	2,021	2
1985	15,182	628	900	1,215	1,558	1,751	1,915	1,818	1,458	1,772	3
1984	13,742	607	834	1,110	1,465	1,588	1,780	1,496	1,334	1,536	6
1983	13,204	536	799	1,146	1,402	1,518	1,657	1,527	1,255	1,393	7
1982	11,491	550	714	991	1,295	1,379	1,466	1,297	976	1,093	9
1981	11,093	577	750	1,032	1,191	1,261	1,271	1,144	1,006	1,099	3
1980	10,797	505	755	965	1,113	1,318	1,291	1,128	845	1,025	4
1979	9,230	498	632	798	995	1,154	1,108	1,037	695	767	4
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All other, male											
1998	8,370	638	600	619	766	813	899	892	707	917	3
1997	8,187	594	594	642	749	827	901	828	724	880	3
1996	8,301	611	572	599	810	866	952	843	725	856	2
1995	8,344	558	564	607	766	893	949	810	748	814	1
1994	8,065	503	541	624	712	899	932	800	693	792	-
1993	8,287	489	588	734	758	947	945	847	698	742	4
1992	8,036	498	499	615	762	876	946	821	741	741	1
1991	8,001	455	474	625	784	937	943	908	659	708	-
1990	7,824	450	493	644	784	923	871	842	665	680	4
1989	8,126	439	582	670	832	1,011	928	897	676	718	3
1988	9,392	454	604	761	1,020	1,158	1,137	1,088	797	861	2
1987	8,895	435	590	721	962	1,131	1,033	1,019	774	769	6
1986	8,818	420	566	759	981	1,037	1,082	1,035	761	744	1
1985	8,185	400	569	717	940	1,000	1,006	906	665	665	2
1984	7,412	381	503	668	866	921	950	753	606	565	2
1983	7,159	331	468	693	815	860	918	791	564	534	3
1982	6,263	356	419	604	788	778	819	617	453	449	4
1981	5,990	328	439	621	696	708	674	550	475	452	2
1980	6,004	315	473	605	668	722	710	567	415	423	2
1979	5,043	304	371	504	570	667	597	514	325	332	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
All other, female											
1998	8,524	400	417	495	561	724	909	1,003	1,004	1,908	2
1997	8,137	380	415	444	549	720	865	927	984	1,836	-
1996	8,105	367	418	459	583	694	878	895	945	1,754	2
1995	7,975	359	365	456	623	726	919	898	962	1,612	-
1994	7,717	328	344	476	544	739	881	882	934	1,532	-
1993	7,797	304	334	435	609	770	859	948	964	1,524	-
1992	7,378	294	343	442	616	726	845	900	850	1,360	-
1991	7,187	258	326	431	567	745	840	896	844	1,333	-
1990	7,207	224	325	434	600	765	780	976	854	1,349	3
1989	7,456	260	350	459	645	838	806	970	863	1,350	1
1988	8,207	285	360	517	708	926	948	1,045	987	1,491	4
1987	7,986	266	360	515	709	816	1,017	1,030	970	1,399	4
1986	7,553	260	363	476	659	813	932	1,016	853	1,277	1
1985	6,997	228	331	498	618	751	909	912	793	1,107	1
1984	6,330	226	331	442	599	667	830	743	728	971	4
1983	6,045	205	331	453	587	658	739	736	691	859	4
1982	5,228	194	295	387	507	601	647	680	523	644	5
1981	5,103	249	311	411	495	553	597	594	531	647	1
1980	4,793	190	282	360	445	596	581	561	430	602	2
1979	4,187	194	261	294	425	487	511	523	370	435	2
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
Black, both sexes											
1998	15,099	970	921	1,020	1,187	1,404	1,608	1,687	1,503	2,422	5
1997	14,626	899	940	996	1,183	1,391	1,594	1,516	1,516	2,350	2
1996	14,784	901	908	976	1,273	1,414	1,652	1,545	1,469	2,278	4
1995	14,732	858	848	975	1,290	1,484	1,668	1,542	1,489	2,099	1
1994	14,326	776	815	1,026	1,156	1,483	1,663	1,533	1,447	2,009	-
1993	14,626	760	856	1,095	1,257	1,586	1,643	1,610	1,461	1,989	3
1992	14,189	741	805	988	1,285	1,493	1,644	1,564	1,448	1,883	1
1991	14,040	683	761	988	1,257	1,574	1,649	1,667	1,363	1,811	-
1990	13,815	630	755	1,001	1,286	1,566	1,531	1,663	1,377	1,812	7
1989	14,467	664	872	1,062	1,386	1,742	1,619	1,729	1,409	1,860	4
1988	16,507	708	901	1,206	1,636	1,966	1,953	2,004	1,666	2,173	6
1987	15,869	671	898	1,154	1,587	1,851	1,945	1,941	1,636	1,996	10
1986	15,408	648	881	1,180	1,573	1,773	1,901	1,905	1,499	1,860	2
1985	14,293	597	867	1,159	1,475	1,650	1,818	1,693	1,376	1,642	3
1984	12,976	576	779	1,051	1,393	1,518	1,700	1,417	1,259	1,423	5
1983	12,420	516	770	1,082	1,332	1,449	1,562	1,424	1,178	1,272	6
1982	10,883	520	685	939	1,242	1,319	1,387	1,231	922	1,025	9
1981	10,555	548	722	992	1,146	1,214	1,222	1,083	946	1,025	1
1980	10,247	487	723	926	1,077	1,255	1,220	1,067	800	960	4
1979	8,733	487	606	764	956	1,097	1,059	981	637	711	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
Black, male											
1998	7,385	582	540	558	677	739	787	778	610	740	3
1997	7,261	545	549	584	690	749	801	703	627	705	2
1996	7,389	555	521	550	725	783	851	735	613	712	2
1995	7,527	520	521	561	713	815	840	727	642	673	1
1994	7,241	468	496	573	659	803	845	721	599	639	-
1993	7,464	470	541	689	692	870	854	751	591	618	3
1992	7,339	460	474	570	711	810	866	736	662	634	1
1991	7,328	435	450	589	724	869	861	826	578	601	-
1990	7,112	418	453	590	728	849	798	760	583	579	4
1989	7,475	417	542	625	774	946	852	815	601	625	3
1988	8,763	435	563	721	962	1,085	1,062	1,008	739	774	2
1987	8,269	413	552	665	914	1,070	967	948	702	696	6
1986	8,244	396	533	726	941	992	1,013	949	690	667	1
1985	7,663	377	547	686	883	934	952	821	623	613	2
1984	6,964	362	470	635	825	881	900	705	563	515	2
1983	6,666	317	449	655	773	811	852	724	522	472	2
1982	5,908	340	396	575	761	747	763	570	421	419	4
1981	5,678	309	427	599	666	679	646	513	438	411	-
1980	5,657	304	451	576	649	686	660	534	382	383	2
1979	4,762	301	356	483	548	633	571	474	297	307	2
All other forms of heart disease (415-417,424-427,429.0-429.1,429.3-429.9)											
Black, female											
1998	7,714	388	381	462	510	665	821	909	893	1,682	2
1997	7,365	354	391	412	493	642	793	813	889	1,645	-
1996	7,395	346	387	426	548	631	801	810	856	1,566	2
1995	7,205	338	327	414	577	669	828	815	847	1,426	-
1994	7,085	308	319	453	497	680	818	812	848	1,370	-
1993	7,162	290	315	406	565	716	789	859	870	1,371	-
1992	6,850	281	331	418	574	683	778	828	786	1,249	-
1991	6,712	248	311	399	533	705	788	841	785	1,210	-
1990	6,703	212	302	411	558	717	733	903	794	1,233	3
1989	6,992	247	330	437	612	796	767	914	808	1,235	1
1988	7,744	273	338	485	674	881	891	996	927	1,399	4
1987	7,600	258	346	489	673	781	978	993	934	1,300	4
1986	7,164	252	348	454	632	781	888	956	809	1,193	1
1985	6,630	220	320	473	592	716	866	872	753	1,029	1
1984	6,012	214	309	416	568	637	800	712	696	908	3
1983	5,754	199	321	427	559	638	710	700	656	800	4
1982	4,975	180	289	364	481	572	624	661	501	606	5
1981	4,877	239	295	393	480	535	576	570	508	614	1
1980	4,590	183	272	350	428	569	560	533	418	577	2
1979	3,971	186	250	281	408	464	488	507	340	404	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All races, both sexes											
1998	14,308	255	338	450	676	949	1,546	2,010	2,565	5,203	-
1997	13,534	243	343	392	611	932	1,484	2,016	2,451	4,744	1
1996	12,945	251	330	409	594	961	1,438	1,965	2,301	4,396	-
1995	12,483	233	315	389	650	957	1,477	1,792	2,193	4,138	2
1994	11,765	197	292	386	625	915	1,417	1,794	2,113	3,706	-
1993	11,243	185	301	411	636	932	1,334	1,673	1,897	3,537	2
1992	10,265	168	243	363	574	904	1,295	1,570	1,832	3,039	1
1991	9,524	181	231	332	548	870	1,200	1,469	1,676	2,746	2
1990	9,169	160	237	368	527	821	1,167	1,462	1,541	2,639	2
1989	8,766	149	232	309	551	836	1,081	1,330	1,527	2,479	1
1988	8,456	121	250	330	578	823	1,073	1,297	1,438	2,312	1
1987	8,139	134	205	320	545	828	1,097	1,224	1,345	2,218	1
1986	7,984	119	240	336	542	766	1,051	1,267	1,360	2,108	4
1985	7,751	129	217	340	557	737	1,020	1,235	1,286	2,032	2
1984	7,774	133	191	368	569	738	1,138	1,240	1,209	2,004	1
1983	7,739	136	196	342	554	724	1,019	1,225	1,250	2,095	4
1982	7,571	118	216	341	553	718	1,021	1,155	1,240	2,027	3
1981	7,587	143	240	372	523	739	955	1,162	1,248	1,985	-
1980	7,827	151	257	380	551	724	1,018	1,203	1,246	2,050	2
1979	7,275	149	244	376	504	722	872	1,094	1,147	1,943	1
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All races, male											
1998	5,526	166	220	280	352	482	705	859	928	1,337	-
1997	5,419	163	212	228	379	451	706	877	946	1,259	1
1996	5,160	153	205	251	324	493	687	854	825	1,200	-
1995	5,025	149	200	217	371	454	723	759	820	1,114	1
1994	4,794	140	193	235	357	473	652	760	791	997	-
1993	4,512	109	180	229	350	462	609	722	706	935	2
1992	4,285	104	145	206	295	471	608	709	709	861	1
1991	3,956	101	132	203	299	424	560	616	632	821	1
1990	3,833	99	139	205	277	417	547	618	562	806	1
1989	3,771	97	154	185	298	440	516	606	608	689	-
1988	3,555	68	141	181	297	417	513	575	584	641	1
1987	3,530	80	122	200	326	425	519	545	515	659	-
1986	3,403	71	134	190	299	406	492	557	505	633	-
1985	3,385	79	128	197	313	379	499	558	498	618	-
1984	3,435	87	109	188	308	392	567	539	453	671	-
1983	3,476	79	118	197	302	377	488	557	512	713	4
1982	3,432	68	119	196	311	392	497	525	480	726	1
1981	3,477	79	142	220	280	379	468	534	539	715	-
1980	3,581	85	141	222	318	345	504	516	514	793	2
1979	3,377	77	135	218	264	410	427	495	488	726	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All races, female											
1998	8,782	89	118	170	324	467	841	1,151	1,637	3,866	-
1997	8,115	80	131	164	232	481	778	1,139	1,505	3,485	-
1996	7,785	98	125	158	270	468	751	1,111	1,476	3,196	-
1995	7,458	84	115	172	279	503	754	1,033	1,373	3,024	1
1994	6,971	57	99	151	268	442	765	1,034	1,322	2,709	-
1993	6,731	76	121	182	286	470	725	951	1,191	2,602	-
1992	5,980	64	98	157	279	433	687	861	1,123	2,178	-
1991	5,568	80	99	129	249	446	640	853	1,044	1,925	1
1990	5,336	61	98	163	250	404	620	844	979	1,833	1
1989	4,995	52	78	124	253	396	565	724	919	1,790	1
1988	4,901	53	109	149	281	406	560	722	854	1,671	-
1987	4,609	54	83	120	219	403	578	679	830	1,559	1
1986	4,581	48	106	146	243	360	559	710	855	1,475	4
1985	4,366	50	89	143	244	358	521	677	788	1,414	2
1984	4,339	46	82	180	261	346	571	701	756	1,333	1
1983	4,263	57	78	145	252	347	531	668	738	1,382	-
1982	4,139	50	97	145	242	326	524	630	760	1,301	2
1981	4,110	64	98	152	243	360	487	628	709	1,270	-
1980	4,246	66	116	158	233	379	514	687	732	1,257	-
1979	3,898	72	109	158	240	312	445	599	659	1,217	-
Essential (primary) hypertension and hypertensive renal disease (401,403)											
White, both sexes											
1998	10,818	116	181	258	394	598	1,130	1,537	2,077	4,383	-
1997	10,376	97	183	231	388	640	1,069	1,567	2,037	4,034	-
1996	9,864	114	166	240	356	653	1,078	1,538	1,885	3,704	-
1995	9,452	113	162	201	402	649	1,076	1,390	1,771	3,539	-
1994	8,902	84	147	219	418	613	1,058	1,387	1,680	3,165	-
1993	8,438	82	172	231	412	640	969	1,288	1,531	2,967	1
1992	7,750	65	116	211	360	593	941	1,246	1,497	2,607	-
1991	7,160	92	103	173	358	584	894	1,125	1,349	2,372	1
1990	6,960	73	111	213	335	560	874	1,140	1,277	2,286	1
1989	6,649	66	121	185	349	563	800	1,017	1,255	2,173	1
1988	6,433	60	107	195	340	589	801	1,027	1,198	2,001	-
1987	6,250	62	110	177	374	603	810	978	1,093	1,939	-
1986	6,171	64	133	199	362	546	809	1,005	1,123	1,851	3
1985	5,922	54	112	201	356	523	767	982	1,074	1,777	1
1984	5,998	61	112	230	370	542	836	969	1,005	1,796	-
1983	6,092	62	110	191	387	532	771	1,006	1,057	1,884	-
1982	6,006	57	118	220	380	534	781	942	1,035	1,851	3
1981	5,894	66	121	228	347	527	721	952	1,060	1,778	-
1980	6,130	71	124	244	365	499	780	969	1,096	1,865	1
1979	5,713	73	121	230	342	503	677	881	1,027	1,760	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
White, male											
1998	4,105	83	120	167	231	307	524	665	767	1,144	-
1997	4,038	69	114	144	251	312	542	675	792	1,056	-
1996	3,814	76	103	156	198	353	522	675	679	979	-
1995	3,697	74	113	125	243	322	536	595	665	927	-
1994	3,547	61	104	136	245	343	500	590	639	848	-
1993	3,370	55	115	142	228	333	451	583	579	780	1
1992	3,228	49	74	126	207	324	452	575	592	741	-
1991	2,938	55	67	115	209	301	424	489	513	701	1
1990	2,862	49	74	130	188	287	424	499	462	688	1
1989	2,819	46	84	125	206	297	395	477	508	600	-
1988	2,749	37	63	133	200	315	409	477	496	553	-
1987	2,692	42	67	110	241	321	403	437	424	581	-
1986	2,639	42	73	121	206	317	403	452	417	557	-
1985	2,588	40	70	129	219	280	388	459	424	535	-
1984	2,651	41	72	120	210	307	423	442	382	601	-
1983	2,724	41	71	116	224	287	381	472	432	637	-
1982	2,740	37	69	144	219	308	401	436	404	666	1
1981	2,718	38	81	151	191	285	368	450	459	643	-
1980	2,813	41	74	153	228	262	385	429	449	718	1
1979	2,645	44	74	147	191	300	337	406	435	651	1
Essential (primary) hypertension and hypertensive renal disease (401,403)											
White, female											
1998	6,713	33	61	91	163	291	606	872	1,310	3,239	-
1997	6,338	28	69	87	137	328	527	892	1,245	2,978	-
1996	6,050	38	63	84	158	300	556	863	1,206	2,725	-
1995	5,755	39	49	76	159	327	540	795	1,106	2,612	-
1994	5,355	23	43	83	173	270	558	797	1,041	2,317	-
1993	5,068	27	57	89	184	307	518	705	952	2,187	-
1992	4,522	16	42	85	153	269	489	671	905	1,866	-
1991	4,222	37	36	58	149	283	470	636	836	1,671	-
1990	4,098	24	37	83	147	273	450	641	815	1,598	-
1989	3,830	20	37	60	143	266	405	540	747	1,573	1
1988	3,684	23	44	62	140	274	392	550	702	1,448	-
1987	3,558	20	43	67	133	282	407	541	669	1,358	-
1986	3,532	22	60	78	156	229	406	553	706	1,294	3
1985	3,334	14	42	72	137	243	379	523	650	1,242	1
1984	3,347	20	40	110	160	235	413	527	623	1,195	-
1983	3,368	21	39	75	163	245	390	534	625	1,247	-
1982	3,266	20	49	76	161	226	380	506	631	1,185	2
1981	3,176	28	40	77	156	242	353	502	601	1,135	-
1980	3,317	30	50	91	137	237	395	540	647	1,147	-
1979	3,068	29	47	83	151	203	340	475	592	1,109	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All other, both sexes											
1998	3,490	139	157	192	282	351	416	473	488	820	-
1997	3,158	146	160	161	223	292	415	449	414	710	1
1996	3,081	137	164	169	238	308	360	427	416	692	-
1995	3,031	120	153	188	248	308	401	402	422	599	2
1994	2,863	113	145	167	207	302	359	407	433	541	-
1993	2,805	103	129	180	224	292	365	385	366	570	1
1992	2,515	103	127	152	214	311	354	324	335	432	1
1991	2,364	89	128	159	190	286	306	344	327	374	1
1990	2,209	87	126	155	192	261	293	322	264	353	1
1989	2,117	83	111	124	202	273	281	313	272	306	-
1988	2,023	61	143	135	238	234	272	270	240	311	1
1987	1,889	72	95	143	171	225	287	246	252	279	1
1986	1,813	55	107	137	180	220	242	262	237	257	1
1985	1,829	75	105	139	201	214	253	253	212	255	1
1984	1,776	72	79	138	199	196	302	271	204	208	1
1983	1,647	74	86	151	167	192	248	219	193	211	4
1982	1,565	61	98	121	173	184	240	213	205	176	-
1981	1,693	77	119	144	176	212	234	210	188	207	-
1980	1,697	80	133	136	186	225	238	234	150	185	1
1979	1,562	76	123	146	162	219	195	213	120	183	-
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All other, male											
1998	1,421	83	100	113	121	175	181	194	161	193	-
1997	1,381	94	98	84	128	139	164	202	154	203	1
1996	1,346	77	102	95	126	140	165	179	146	221	-
1995	1,328	75	87	92	128	132	187	164	155	187	1
1994	1,247	79	89	99	112	130	152	170	152	149	-
1993	1,142	54	65	87	122	129	158	139	127	155	1
1992	1,057	55	71	80	88	147	156	134	117	120	1
1991	1,018	46	65	88	90	123	136	127	119	120	-
1990	971	50	65	75	89	130	123	119	100	118	-
1989	952	51	70	60	92	143	121	129	100	89	-
1988	806	31	78	48	97	102	104	98	88	88	1
1987	838	38	55	90	85	104	116	108	91	78	-
1986	764	29	61	69	93	89	89	105	88	76	-
1985	797	39	58	68	94	99	111	99	74	83	-
1984	784	46	37	68	98	85	144	97	71	70	-
1983	752	38	47	81	78	90	107	85	80	76	4
1982	692	31	50	52	92	84	96	89	76	60	-
1981	759	41	61	69	89	94	100	84	80	72	-
1980	768	44	67	69	90	83	119	87	65	75	1
1979	732	33	61	71	73	110	90	89	53	75	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
All other, female											
1998	2,069	56	57	79	161	176	235	279	327	627	-
1997	1,777	52	62	77	95	153	251	247	260	507	-
1996	1,735	60	62	74	112	168	195	248	270	471	-
1995	1,703	45	66	96	120	176	214	238	267	412	1
1994	1,616	34	56	68	95	172	207	237	281	392	-
1993	1,663	49	64	93	102	163	207	246	239	415	-
1992	1,458	48	56	72	126	164	198	190	218	312	-
1991	1,346	43	63	71	100	163	170	217	208	254	1
1990	1,238	37	61	80	103	131	170	203	164	235	1
1989	1,165	32	41	64	110	130	160	184	172	217	-
1988	1,217	30	65	87	141	132	168	172	152	223	-
1987	1,051	34	40	53	86	121	171	138	161	201	1
1986	1,049	26	46	68	87	131	153	157	149	181	1
1985	1,032	36	47	71	107	115	142	154	138	172	1
1984	992	26	42	70	101	111	158	174	133	138	1
1983	895	36	39	70	89	102	141	134	113	135	-
1982	873	30	48	69	81	100	144	124	129	116	-
1981	934	36	58	75	87	118	134	126	108	135	-
1980	929	36	66	67	96	142	119	147	85	110	-
1979	830	43	62	75	89	109	105	124	67	108	-
Essential (primary) hypertension and hypertensive renal disease (401,403)											
Black, both sexes											
1998	3,179	131	147	189	263	323	389	421	420	734	-
1997	2,919	140	155	152	216	271	384	404	383	638	1
1996	2,845	133	158	163	222	285	327	396	377	623	-
1995	2,804	119	148	178	228	290	370	369	377	544	2
1994	2,654	109	137	160	193	283	335	375	394	486	-
1993	2,622	100	120	174	214	275	341	352	344	513	1
1992	2,327	98	124	142	205	288	327	297	297	389	1
1991	2,224	87	125	154	179	269	287	332	300	337	1
1990	2,109	86	122	149	183	247	283	305	256	331	1
1989	1,997	80	105	121	196	257	268	293	250	279	-
1988	1,926	57	137	131	232	226	252	258	223	291	1
1987	1,804	72	91	140	167	222	272	233	236	257	1
1986	1,732	55	102	132	175	210	235	248	223	242	1
1985	1,757	75	103	129	199	210	241	242	199	243	1
1984	1,711	67	78	138	194	194	293	256	191	195	1
1983	1,582	71	83	149	164	183	239	212	184	197	2
1982	1,489	59	97	117	166	181	231	204	191	157	-
1981	1,633	74	116	138	170	209	229	200	176	198	-
1980	1,627	77	127	128	182	222	229	220	143	171	1
1979	1,509	73	122	140	160	213	190	201	117	173	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Essential (primary) hypertension and hypertensive renal disease (401,403)											
Black, male											
1998	1,276	79	91	112	112	163	169	165	130	162	-
1997	1,257	90	96	77	125	124	151	180	138	169	1
1996	1,233	74	97	91	119	131	151	168	130	186	-
1995	1,216	74	83	86	115	124	172	148	136	162	1
1994	1,147	75	85	93	106	121	140	157	135	124	-
1993	1,067	51	63	86	116	125	150	122	115	134	1
1992	969	52	68	75	85	136	142	121	96	107	1
1991	945	44	63	86	83	113	128	122	107	100	-
1990	919	50	62	71	82	124	121	109	96	109	-
1989	892	49	65	59	90	135	112	119	92	76	-
1988	747	29	72	46	94	96	93	89	76	80	1
1987	795	38	53	90	82	103	107	99	84	69	-
1986	719	29	56	66	89	83	86	97	82	70	-
1985	764	39	56	63	94	97	107	95	67	78	-
1984	750	43	36	68	96	84	140	86	64	66	-
1983	719	36	46	80	77	84	104	82	76	69	2
1982	650	30	50	50	85	83	95	81	71	50	-
1981	727	39	59	66	83	94	97	78	73	70	-
1980	728	43	64	64	88	81	111	80	60	69	1
1979	705	31	61	65	73	107	87	83	53	71	-
Essential (primary) hypertension and hypertensive renal disease (401,403)											
Black, female											
1998	1,903	52	56	77	151	160	220	256	290	572	-
1997	1,662	50	59	75	91	147	233	224	245	469	-
1996	1,612	59	61	72	103	154	176	228	247	437	-
1995	1,588	45	65	92	113	166	198	221	241	382	1
1994	1,507	34	52	67	87	162	195	218	259	362	-
1993	1,555	49	57	88	98	150	191	230	229	379	-
1992	1,358	46	56	67	120	152	185	176	201	282	-
1991	1,279	43	62	68	96	156	159	210	193	237	1
1990	1,190	36	60	78	101	123	162	196	160	222	1
1989	1,105	31	40	62	106	122	156	174	158	203	-
1988	1,179	28	65	85	138	130	159	169	147	211	-
1987	1,009	34	38	50	85	119	165	134	152	188	1
1986	1,013	26	46	66	86	127	149	151	141	172	1
1985	993	36	47	66	105	113	134	147	132	165	1
1984	961	24	42	70	98	110	153	170	127	129	1
1983	863	35	37	69	87	99	135	130	108	128	-
1982	839	29	47	67	81	98	136	123	120	107	-
1981	906	35	57	72	87	115	132	122	103	128	-
1980	899	34	63	64	94	141	118	140	83	102	-
1979	804	42	61	75	87	106	103	118	64	102	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
All races, both sexes											
1998	158,042	2,437	3,265	4,059	5,583	8,776	15,095	23,648	30,652	60,589	9
1997	159,392	2,476	3,217	3,965	5,700	9,244	15,666	23,597	30,366	61,114	5
1996	159,547	2,647	3,139	3,936	5,731	9,544	15,719	23,582	30,824	60,445	11
1995	157,600	2,444	3,021	3,935	5,785	9,640	16,056	23,031	30,513	59,156	7
1994	152,939	2,338	3,005	3,725	5,841	9,630	15,710	22,678	29,678	56,306	7
1993	149,780	2,243	2,810	3,761	5,847	9,600	15,676	22,563	28,708	54,705	9
1992	143,449	2,121	2,663	3,707	5,988	9,602	15,339	22,019	27,314	50,825	13
1991	143,140	2,050	2,664	3,700	6,026	10,128	15,359	22,025	27,319	49,996	14
1990	143,752	2,064	2,625	3,868	6,243	10,469	15,547	22,433	27,427	49,215	13
1989	145,229	1,926	2,654	3,956	6,568	10,603	15,663	22,811	27,814	49,470	18
1988	150,145	2,021	2,607	4,178	7,003	11,117	16,536	23,823	28,778	50,140	9
1987	149,469	1,859	2,803	4,303	7,165	10,885	16,836	23,688	28,524	49,516	16
1986	149,269	1,864	2,774	4,435	7,331	11,219	17,186	23,739	28,115	48,753	65
1985	152,649	1,884	2,881	4,658	7,460	11,347	17,717	24,579	28,766	49,566	53
1984	153,920	1,964	3,114	4,746	7,682	11,514	18,053	24,628	29,140	49,250	21
1983	155,236	2,063	3,034	4,912	7,878	11,889	18,131	25,240	29,466	48,954	27
1982	157,388	2,019	3,281	5,126	7,878	12,351	18,863	25,674	29,874	48,625	24
1981	163,187	2,101	3,495	5,422	8,351	12,879	19,871	26,042	31,119	50,013	18
1980	169,876	2,170	3,572	5,604	8,533	13,626	20,503	28,048	32,770	51,157	18
1979	169,207	2,290	3,762	5,861	8,734	13,810	20,948	27,783	32,442	49,662	23
Cerebrovascular diseases (430-434,436-438)											
All races, male											
1998	61,000	1,366	1,752	2,231	3,106	4,561	7,433	10,693	11,862	15,933	4
1997	62,447	1,383	1,777	2,213	3,123	4,907	7,738	10,682	11,901	16,636	3
1996	62,344	1,452	1,709	2,187	3,139	5,027	7,835	10,783	11,989	16,133	4
1995	61,429	1,377	1,622	2,170	3,184	5,023	7,961	10,432	11,913	15,580	5
1994	60,089	1,319	1,618	2,064	3,167	5,131	7,778	10,455	11,561	14,861	-
1993	58,926	1,213	1,527	2,041	3,166	5,147	7,800	10,312	11,102	14,548	3
1992	56,522	1,161	1,426	1,997	3,266	5,035	7,605	9,933	10,447	13,591	8
1991	56,585	1,099	1,454	2,020	3,205	5,411	7,705	10,029	10,534	13,092	8
1990	56,569	1,085	1,419	2,113	3,290	5,486	7,685	10,173	10,466	12,858	7
1989	57,206	1,028	1,397	2,116	3,580	5,540	7,741	10,363	10,406	13,047	6
1988	59,630	1,080	1,388	2,286	3,824	5,778	8,230	10,725	10,840	13,362	-
1987	58,914	951	1,503	2,348	3,822	5,623	8,283	10,553	10,744	13,048	8
1986	59,019	959	1,439	2,360	3,892	5,770	8,356	10,500	10,746	12,972	16
1985	60,622	992	1,535	2,559	4,027	5,929	8,819	11,005	10,703	13,118	16
1984	61,541	1,045	1,630	2,571	4,085	6,106	8,916	10,759	10,947	13,510	8
1983	62,685	1,073	1,575	2,653	4,201	6,204	9,082	11,328	11,100	13,593	6
1982	63,725	1,047	1,706	2,779	4,158	6,537	9,439	11,391	11,031	13,750	13
1981	66,276	1,066	1,819	2,866	4,460	6,831	9,839	11,674	11,538	14,212	13
1980	69,823	1,048	1,951	2,960	4,613	7,192	10,283	12,468	12,367	14,950	11
1979	70,317	1,150	1,992	3,156	4,808	7,318	10,446	12,321	12,333	14,764	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
All races, female											
1998	97,042	1,071	1,513	1,828	2,477	4,215	7,662	12,955	18,790	44,656	5
1997	96,945	1,093	1,440	1,752	2,577	4,337	7,928	12,915	18,465	44,478	2
1996	97,203	1,195	1,430	1,749	2,592	4,517	7,884	12,799	18,835	44,312	7
1995	96,171	1,067	1,399	1,765	2,601	4,617	8,095	12,599	18,600	43,576	2
1994	92,850	1,019	1,387	1,661	2,674	4,499	7,932	12,223	18,117	41,445	7
1993	90,854	1,030	1,283	1,720	2,681	4,453	7,876	12,251	17,606	40,157	6
1992	86,927	960	1,237	1,710	2,722	4,567	7,734	12,086	16,867	37,234	5
1991	86,555	951	1,210	1,680	2,821	4,717	7,654	11,996	16,785	36,904	6
1990	87,183	979	1,206	1,755	2,953	4,983	7,862	12,260	16,961	36,357	6
1989	88,023	898	1,257	1,840	2,988	5,063	7,922	12,448	17,408	36,423	12
1988	90,515	941	1,219	1,892	3,179	5,339	8,306	13,098	17,938	36,778	9
1987	90,555	908	1,300	1,955	3,343	5,262	8,553	13,135	17,780	36,468	8
1986	90,250	905	1,335	2,075	3,439	5,449	8,830	13,239	17,369	35,781	49
1985	92,027	892	1,346	2,099	3,433	5,418	8,898	13,574	18,063	36,448	37
1984	92,379	919	1,484	2,175	3,597	5,408	9,137	13,869	18,193	35,740	13
1983	92,551	990	1,459	2,259	3,677	5,685	9,049	13,912	18,366	35,361	21
1982	93,663	972	1,575	2,347	3,720	5,814	9,424	14,283	18,843	34,875	11
1981	96,911	1,035	1,676	2,556	3,891	6,048	10,032	14,368	19,581	35,801	5
1980	100,053	1,122	1,621	2,644	3,920	6,434	10,220	15,580	20,403	36,207	7
1979	98,890	1,140	1,770	2,705	3,926	6,492	10,502	15,462	20,109	34,898	8
Cerebrovascular diseases (430-434,436-438)											
White, both sexes											
1998	136,485	1,508	2,209	2,881	4,117	6,840	12,598	20,563	27,522	55,685	4
1997	137,954	1,548	2,194	2,752	4,159	7,327	13,041	20,637	27,395	56,304	4
1996	137,940	1,692	2,161	2,793	4,232	7,462	13,061	20,590	27,705	55,645	7
1995	136,124	1,606	2,052	2,744	4,351	7,574	13,434	20,059	27,386	54,337	3
1994	132,159	1,518	2,000	2,649	4,392	7,633	13,079	19,822	26,574	51,953	4
1993	129,690	1,457	1,923	2,665	4,383	7,611	13,128	19,658	25,857	50,470	3
1992	124,080	1,398	1,848	2,605	4,511	7,637	12,821	19,237	24,621	46,942	10
1991	123,400	1,326	1,798	2,613	4,507	8,032	12,871	19,212	24,503	46,046	9
1990	124,217	1,310	1,797	2,779	4,721	8,331	13,072	19,598	24,682	45,480	4
1989	125,222	1,245	1,755	2,856	4,907	8,424	13,138	19,732	25,061	45,715	11
1988	129,726	1,342	1,770	2,989	5,298	8,829	13,896	20,759	25,904	46,413	4
1987	129,707	1,200	1,916	3,117	5,458	8,788	14,154	20,661	25,879	45,976	8
1986	129,648	1,265	1,878	3,265	5,645	9,068	14,478	20,724	25,470	45,341	46
1985	132,661	1,229	2,020	3,463	5,744	9,219	14,850	21,557	25,930	46,132	44
1984	134,022	1,326	2,211	3,500	5,967	9,277	15,114	21,662	26,299	46,113	11
1983	135,263	1,361	2,136	3,648	6,059	9,649	15,177	22,307	26,773	45,674	16
1982	137,343	1,360	2,343	3,823	6,090	10,023	16,005	22,574	27,172	45,491	15
1981	142,472	1,431	2,450	4,023	6,602	10,474	16,779	23,023	28,338	46,700	12
1980	148,413	1,466	2,572	4,212	6,670	10,981	17,353	24,668	29,987	47,869	11
1979	148,157	1,560	2,705	4,394	6,930	11,199	17,769	24,481	29,969	46,493	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
White, male											
1998	51,640	862	1,190	1,574	2,263	3,588	6,253	9,335	10,687	14,501	2
1997	53,052	874	1,196	1,517	2,303	3,911	6,492	9,366	10,789	15,221	2
1996	52,895	916	1,159	1,527	2,317	3,980	6,538	9,459	10,875	14,707	2
1995	51,929	893	1,053	1,487	2,427	3,984	6,697	9,110	10,667	14,191	3
1994	50,886	846	1,078	1,456	2,397	4,123	6,506	9,194	10,315	13,587	-
1993	50,106	784	1,029	1,435	2,396	4,136	6,608	9,016	10,004	13,308	1
1992	47,962	773	996	1,388	2,467	4,020	6,400	8,707	9,434	12,446	6
1991	47,720	676	976	1,435	2,424	4,333	6,497	8,795	9,388	11,877	5
1990	47,912	676	958	1,525	2,504	4,392	6,509	8,921	9,398	11,742	-
1989	48,459	647	916	1,548	2,700	4,453	6,558	9,047	9,414	11,902	4
1988	50,573	688	952	1,645	2,926	4,659	6,974	9,387	9,705	12,274	-
1987	50,099	598	1,040	1,689	2,947	4,573	7,028	9,216	9,692	11,981	5
1986	50,224	640	954	1,722	3,039	4,670	7,097	9,155	9,690	11,935	9
1985	51,816	636	1,082	1,935	3,131	4,847	7,473	9,654	9,672	12,100	11
1984	52,687	700	1,146	1,878	3,198	4,988	7,550	9,530	9,858	12,519	4
1983	53,822	700	1,121	1,997	3,262	5,096	7,673	10,029	10,102	12,558	4
1982	54,606	692	1,231	2,075	3,236	5,358	8,017	10,007	9,994	12,736	6
1981	56,857	698	1,276	2,128	3,541	5,612	8,392	10,313	10,441	13,129	9
1980	59,953	697	1,424	2,239	3,621	5,871	8,746	10,950	11,224	13,848	8
1979	60,501	776	1,410	2,330	3,825	5,988	8,949	10,847	11,326	13,696	9
Cerebrovascular diseases (430-434,436-438)											
White, female											
1998	84,845	646	1,019	1,307	1,854	3,252	6,345	11,228	16,835	41,184	2
1997	84,902	674	998	1,235	1,856	3,416	6,549	11,271	16,606	41,083	2
1996	85,045	776	1,002	1,266	1,915	3,482	6,523	11,131	16,830	40,938	5
1995	84,195	713	999	1,257	1,924	3,590	6,737	10,949	16,719	40,146	-
1994	81,273	672	922	1,193	1,995	3,510	6,573	10,628	16,259	38,366	4
1993	79,584	673	894	1,230	1,987	3,475	6,520	10,642	15,853	37,162	2
1992	76,118	625	852	1,217	2,044	3,617	6,421	10,530	15,187	34,496	4
1991	75,680	650	822	1,178	2,083	3,699	6,374	10,417	15,115	34,169	4
1990	76,305	634	839	1,254	2,217	3,939	6,563	10,677	15,284	33,738	4
1989	76,763	598	839	1,308	2,207	3,971	6,580	10,685	15,647	33,813	7
1988	79,153	654	818	1,344	2,372	4,170	6,922	11,372	16,199	34,139	4
1987	79,608	602	876	1,428	2,511	4,215	7,126	11,445	16,187	33,995	3
1986	79,424	625	924	1,543	2,606	4,398	7,381	11,569	15,780	33,406	37
1985	80,845	593	938	1,528	2,613	4,372	7,377	11,903	16,258	34,032	33
1984	81,335	626	1,065	1,622	2,769	4,289	7,564	12,132	16,441	33,594	7
1983	81,441	661	1,015	1,651	2,797	4,553	7,504	12,278	16,671	33,116	12
1982	82,737	668	1,112	1,748	2,854	4,665	7,988	12,567	17,178	32,755	9
1981	85,615	733	1,174	1,895	3,061	4,862	8,387	12,710	17,897	33,571	3
1980	88,460	769	1,148	1,973	3,049	5,110	8,607	13,718	18,763	34,021	3
1979	87,656	784	1,295	2,064	3,105	5,211	8,820	13,634	18,643	32,797	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
All other, both sexes											
1998	21,557	929	1,056	1,178	1,466	1,936	2,497	3,085	3,130	4,904	5
1997	21,438	928	1,023	1,213	1,541	1,917	2,625	2,960	2,971	4,810	1
1996	21,607	955	978	1,143	1,499	2,082	2,658	2,992	3,119	4,800	4
1995	21,476	838	969	1,191	1,434	2,066	2,622	2,972	3,127	4,819	4
1994	20,780	820	1,005	1,076	1,449	1,997	2,631	2,856	3,104	4,353	3
1993	20,090	786	887	1,096	1,464	1,989	2,548	2,905	2,851	4,235	6
1992	19,369	723	815	1,102	1,477	1,965	2,518	2,782	2,693	3,883	3
1991	19,740	724	866	1,087	1,519	2,096	2,488	2,813	2,816	3,950	5
1990	19,535	754	828	1,089	1,522	2,138	2,475	2,835	2,745	3,735	9
1989	20,007	681	899	1,100	1,661	2,179	2,525	3,079	2,753	3,755	7
1988	20,419	679	837	1,189	1,705	2,288	2,640	3,064	2,874	3,727	5
1987	19,762	659	887	1,186	1,707	2,097	2,682	3,027	2,645	3,540	8
1986	19,621	599	896	1,170	1,686	2,151	2,708	3,015	2,645	3,412	19
1985	19,988	655	861	1,195	1,716	2,128	2,867	3,022	2,836	3,434	9
1984	19,898	638	903	1,246	1,715	2,237	2,939	2,966	2,841	3,137	10
1983	19,973	702	898	1,264	1,819	2,240	2,954	2,933	2,693	3,280	11
1982	20,045	659	938	1,303	1,788	2,328	2,858	3,100	2,702	3,134	9
1981	20,715	670	1,045	1,399	1,749	2,405	3,092	3,019	2,781	3,313	6
1980	21,463	704	1,000	1,392	1,863	2,645	3,150	3,380	2,783	3,288	7
1979	21,050	730	1,057	1,467	1,804	2,611	3,179	3,302	2,473	3,169	7
Cerebrovascular diseases (430-434,436-438)											
All other, male											
1998	9,360	504	562	657	843	973	1,180	1,358	1,175	1,432	2
1997	9,395	509	581	696	820	996	1,246	1,316	1,112	1,415	1
1996	9,449	536	550	660	822	1,047	1,297	1,324	1,114	1,426	2
1995	9,500	484	569	683	757	1,039	1,264	1,322	1,246	1,389	2
1994	9,203	473	540	608	770	1,008	1,272	1,261	1,246	1,274	-
1993	8,820	429	498	606	770	1,011	1,192	1,296	1,098	1,240	2
1992	8,560	388	430	609	799	1,015	1,205	1,226	1,013	1,145	2
1991	8,865	423	478	585	781	1,078	1,208	1,234	1,146	1,215	3
1990	8,657	409	461	588	786	1,094	1,176	1,252	1,068	1,116	7
1989	8,747	381	481	568	880	1,087	1,183	1,316	992	1,145	2
1988	9,057	392	436	641	898	1,119	1,256	1,338	1,135	1,088	-
1987	8,815	353	463	659	875	1,050	1,255	1,337	1,052	1,067	3
1986	8,795	319	485	638	853	1,100	1,259	1,345	1,056	1,037	7
1985	8,806	356	453	624	896	1,082	1,346	1,351	1,031	1,018	5
1984	8,854	345	484	693	887	1,118	1,366	1,229	1,089	991	4
1983	8,863	373	454	656	939	1,108	1,409	1,299	998	1,035	2
1982	9,119	355	475	704	922	1,179	1,422	1,384	1,037	1,014	7
1981	9,419	368	543	738	919	1,219	1,447	1,361	1,097	1,083	4
1980	9,870	351	527	721	992	1,321	1,537	1,518	1,143	1,102	3
1979	9,816	374	582	826	983	1,330	1,497	1,474	1,007	1,068	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
All other, female											
1998	12,197	425	494	521	623	963	1,317	1,727	1,955	3,472	3
1997	12,043	419	442	517	721	921	1,379	1,644	1,859	3,395	-
1996	12,158	419	428	483	677	1,035	1,361	1,668	2,005	3,374	2
1995	11,976	354	400	508	677	1,027	1,358	1,650	1,881	3,430	2
1994	11,577	347	465	468	679	989	1,359	1,595	1,858	3,079	3
1993	11,270	357	389	490	694	978	1,356	1,609	1,753	2,995	4
1992	10,809	335	385	493	678	950	1,313	1,556	1,680	2,738	1
1991	10,875	301	388	502	738	1,018	1,280	1,579	1,670	2,735	2
1990	10,878	345	367	501	736	1,044	1,299	1,583	1,677	2,619	2
1989	11,260	300	418	532	781	1,092	1,342	1,763	1,761	2,610	5
1988	11,362	287	401	548	807	1,169	1,384	1,726	1,739	2,639	5
1987	10,947	306	424	527	832	1,047	1,427	1,690	1,593	2,473	5
1986	10,826	280	411	532	833	1,051	1,449	1,670	1,589	2,375	12
1985	11,182	299	408	571	820	1,046	1,521	1,671	1,805	2,416	4
1984	11,044	293	419	553	828	1,119	1,573	1,737	1,752	2,146	6
1983	11,110	329	444	608	880	1,132	1,545	1,634	1,695	2,245	9
1982	10,926	304	463	599	866	1,149	1,436	1,716	1,665	2,120	2
1981	11,296	302	502	661	830	1,186	1,645	1,658	1,684	2,230	2
1980	11,593	353	473	671	871	1,324	1,613	1,862	1,640	2,186	4
1979	11,234	356	475	641	821	1,281	1,682	1,828	1,466	2,101	1
Cerebrovascular diseases (430-434,436-438)											
Black, both sexes											
1998	18,201	822	937	1,010	1,251	1,608	2,082	2,554	2,569	4,129	5
1997	18,106	841	883	1,039	1,313	1,635	2,218	2,428	2,416	4,052	1
1996	18,447	837	843	1,004	1,295	1,785	2,261	2,504	2,658	4,020	4
1995	18,504	742	842	1,040	1,224	1,785	2,262	2,518	2,655	4,147	4
1994	17,996	741	886	944	1,279	1,725	2,257	2,442	2,651	3,725	2
1993	17,535	704	782	974	1,277	1,706	2,246	2,522	2,469	3,653	6
1992	17,016	648	717	962	1,293	1,717	2,217	2,435	2,372	3,358	3
1991	17,345	639	771	973	1,339	1,843	2,189	2,439	2,485	3,407	4
1990	17,382	690	739	976	1,367	1,916	2,183	2,528	2,415	3,280	8
1989	17,954	621	799	969	1,518	1,968	2,306	2,781	2,449	3,295	6
1988	18,460	616	754	1,104	1,550	2,083	2,407	2,766	2,576	3,303	2
1987	17,882	609	792	1,077	1,570	1,922	2,409	2,726	2,382	3,166	7
1986	17,936	560	809	1,078	1,555	1,976	2,466	2,752	2,421	3,078	15
1985	18,315	617	802	1,116	1,580	1,980	2,618	2,774	2,559	3,084	7
1984	18,346	601	830	1,148	1,598	2,083	2,757	2,712	2,615	2,830	9
1983	18,538	654	839	1,185	1,714	2,080	2,761	2,725	2,483	2,995	8
1982	18,676	613	883	1,232	1,692	2,186	2,686	2,869	2,492	2,853	9
1981	19,392	625	983	1,300	1,677	2,255	2,940	2,821	2,593	3,030	5
1980	20,111	664	954	1,316	1,770	2,512	2,973	3,169	2,566	2,999	7
1979	19,817	683	1,002	1,398	1,722	2,477	3,033	3,105	2,298	2,913	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cerebrovascular diseases (430-434,436-438)											
Black, male											
1998	7,746	449	494	555	719	809	977	1,096	942	1,108	2
1997	7,785	466	504	600	700	843	1,050	1,056	854	1,094	1
1996	7,963	470	490	578	710	907	1,085	1,089	939	1,105	2
1995	7,994	433	506	590	648	899	1,061	1,092	1,018	1,098	2
1994	7,815	429	474	540	679	869	1,076	1,053	1,014	1,006	-
1993	7,592	390	448	550	665	875	1,036	1,090	912	1,013	2
1992	7,409	357	380	539	700	883	1,057	1,036	857	932	2
1991	7,676	377	432	525	691	946	1,053	1,061	965	976	2
1990	7,638	382	422	542	702	979	1,027	1,097	902	942	6
1989	7,730	350	432	509	812	971	1,069	1,161	823	957	1
1988	8,091	363	400	593	824	1,011	1,138	1,192	972	910	-
1987	7,844	331	422	597	786	944	1,115	1,174	909	921	2
1986	7,934	303	441	589	783	1,001	1,137	1,190	930	904	6
1985	7,992	333	427	587	837	997	1,226	1,210	880	885	5
1984	8,084	327	447	640	833	1,044	1,262	1,080	974	878	4
1983	8,126	355	424	618	871	1,011	1,292	1,184	897	928	2
1982	8,405	333	448	656	862	1,109	1,327	1,238	927	914	7
1981	8,750	347	509	687	885	1,131	1,348	1,248	1,007	979	3
1980	9,187	337	504	679	941	1,248	1,430	1,388	1,043	993	3
1979	9,176	352	557	788	934	1,246	1,406	1,367	923	957	6
Cerebrovascular diseases (430-434,436-438)											
Black, female											
1998	10,455	373	443	455	532	799	1,105	1,458	1,627	3,021	3
1997	10,321	375	379	439	613	792	1,168	1,372	1,562	2,958	-
1996	10,484	367	353	426	585	878	1,176	1,415	1,719	2,915	2
1995	10,510	309	336	450	576	886	1,201	1,426	1,637	3,049	2
1994	10,181	312	412	404	600	856	1,181	1,389	1,637	2,719	2
1993	9,943	314	334	424	612	831	1,210	1,432	1,557	2,640	4
1992	9,607	291	337	423	593	834	1,160	1,399	1,515	2,426	1
1991	9,669	262	339	448	648	897	1,136	1,378	1,520	2,431	2
1990	9,744	308	317	434	665	937	1,156	1,431	1,513	2,338	2
1989	10,224	271	367	460	706	997	1,237	1,620	1,626	2,338	5
1988	10,369	253	354	511	726	1,072	1,269	1,574	1,604	2,393	2
1987	10,038	278	370	480	784	978	1,294	1,552	1,473	2,245	5
1986	10,002	257	368	489	772	975	1,329	1,562	1,491	2,174	9
1985	10,323	284	375	529	743	983	1,392	1,564	1,679	2,199	2
1984	10,262	274	383	508	765	1,039	1,495	1,632	1,641	1,952	5
1983	10,412	299	415	567	843	1,069	1,469	1,541	1,586	2,067	6
1982	10,271	280	435	576	830	1,077	1,359	1,631	1,565	1,939	2
1981	10,642	278	474	613	792	1,124	1,592	1,573	1,586	2,051	2
1980	10,924	327	450	637	829	1,264	1,543	1,781	1,523	2,006	4
1979	10,641	331	445	610	788	1,231	1,627	1,738	1,375	1,956	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
All races, both sexes											
1998	15,279	64	108	200	360	597	1,090	1,781	2,626	8,398	-
1997	16,057	75	133	192	328	664	1,141	1,930	2,808	8,730	1
1996	16,740	72	132	177	394	623	1,262	2,012	2,903	9,091	1
1995	16,723	72	124	207	388	678	1,229	2,022	3,013	8,941	1
1994	17,116	52	99	215	352	736	1,286	1,966	3,006	9,336	1
1993	17,272	49	122	179	396	727	1,277	2,041	3,016	9,399	3
1992	16,831	56	106	192	422	747	1,287	1,999	2,910	9,052	2
1991	17,420	69	88	204	438	776	1,350	2,183	3,036	9,223	1
1990	18,047	54	108	187	487	833	1,387	2,156	3,175	9,607	3
1989	19,357	56	112	233	488	882	1,466	2,354	3,407	10,304	1
1988	22,086	47	129	245	542	1,000	1,699	2,668	4,011	11,677	3
1987	22,474	51	123	291	579	1,078	1,797	2,791	4,053	11,653	2
1986	22,706	58	132	270	619	984	1,781	2,780	4,001	12,009	21
1985	23,926	49	133	319	573	1,076	1,822	2,935	4,328	12,609	25
1984	24,462	47	106	280	630	1,053	1,858	3,103	4,507	12,815	5
1983	26,371	68	174	325	667	1,109	2,082	3,286	4,988	13,626	4
1982	26,823	73	146	329	662	1,238	2,094	3,402	5,084	13,720	14
1981	28,004	52	165	328	661	1,251	2,147	3,596	5,380	14,346	10
1980	29,449	78	192	348	687	1,313	2,366	3,841	5,855	14,706	6
1979	28,801	66	142	346	677	1,368	2,368	3,756	5,769	14,260	3
Atherosclerosis (440)											
All races, male											
1998	5,719	40	76	133	230	358	625	877	1,093	2,251	-
1997	6,019	58	89	142	192	419	618	940	1,189	2,332	1
1996	6,281	53	94	112	266	373	661	985	1,256	2,437	1
1995	6,404	52	89	132	236	413	696	1,041	1,236	2,469	1
1994	6,420	37	68	134	230	440	666	968	1,282	2,549	-
1993	6,410	27	85	114	253	422	709	1,030	1,202	2,523	1
1992	6,328	45	73	120	245	428	716	999	1,199	2,466	1
1991	6,636	54	57	122	291	464	735	1,129	1,251	2,496	1
1990	6,859	31	79	126	311	486	774	1,086	1,276	2,654	3
1989	7,254	39	65	149	307	534	789	1,090	1,454	2,785	-
1988	8,327	31	88	156	345	578	973	1,291	1,639	3,178	1
1987	8,428	42	81	185	363	647	1,010	1,346	1,581	3,135	1
1986	8,587	47	88	182	403	599	998	1,339	1,620	3,267	9
1985	9,062	33	95	227	367	646	1,040	1,383	1,694	3,530	6
1984	9,240	39	67	171	371	626	1,005	1,476	1,838	3,604	2
1983	10,042	51	118	218	400	655	1,123	1,604	1,969	3,876	-
1982	10,583	53	99	212	413	759	1,163	1,686	2,043	4,112	4
1981	10,969	31	108	214	417	735	1,192	1,743	2,169	4,310	5
1980	11,601	53	128	227	439	810	1,305	1,746	2,345	4,508	1
1979	11,426	51	101	221	415	776	1,305	1,773	2,314	4,439	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
All races, female											
1998	9,560	24	32	67	130	239	465	904	1,533	6,147	-
1997	10,038	17	44	50	136	245	523	990	1,619	6,398	-
1996	10,459	19	38	65	128	250	601	1,027	1,647	6,654	-
1995	10,319	20	35	75	152	265	533	981	1,777	6,472	-
1994	10,696	15	31	81	122	296	620	998	1,724	6,787	1
1993	10,862	22	37	65	143	305	568	1,011	1,814	6,876	2
1992	10,503	11	33	72	177	319	571	1,000	1,711	6,586	1
1991	10,784	15	31	82	147	312	615	1,054	1,785	6,727	-
1990	11,188	23	29	61	176	347	613	1,070	1,899	6,953	-
1989	12,103	17	47	84	181	348	677	1,264	1,953	7,519	1
1988	13,759	16	41	89	197	422	726	1,377	2,372	8,499	2
1987	14,046	9	42	106	216	431	787	1,445	2,472	8,518	1
1986	14,119	11	44	88	216	385	783	1,441	2,381	8,742	12
1985	14,864	16	38	92	206	430	782	1,552	2,634	9,079	19
1984	15,222	8	39	109	259	427	853	1,627	2,669	9,211	3
1983	16,329	17	56	107	267	454	959	1,682	3,019	9,750	4
1982	16,240	20	47	117	249	479	931	1,716	3,041	9,608	10
1981	17,035	21	57	114	244	516	955	1,853	3,211	10,036	5
1980	17,848	25	64	121	248	503	1,061	2,095	3,510	10,198	5
1979	17,375	15	41	125	262	592	1,063	1,983	3,455	9,821	3
Atherosclerosis (440)											
White, both sexes											
1998	14,050	51	86	165	305	517	971	1,620	2,429	7,865	-
1997	14,750	56	98	155	278	569	994	1,761	2,603	8,195	1
1996	15,379	57	108	133	316	546	1,122	1,832	2,687	8,522	1
1995	15,378	63	104	166	321	574	1,078	1,833	2,772	8,426	1
1994	15,660	35	76	162	292	632	1,132	1,763	2,764	8,753	1
1993	15,905	38	100	140	327	626	1,117	1,870	2,767	8,878	2
1992	15,493	42	84	154	354	641	1,139	1,826	2,667	8,536	2
1991	15,981	55	72	159	356	676	1,191	1,972	2,777	8,681	1
1990	16,547	43	90	149	408	717	1,226	1,949	2,900	9,022	2
1989	17,791	47	94	194	405	770	1,295	2,130	3,149	9,663	-
1988	20,261	38	99	209	443	859	1,514	2,351	3,683	11,009	2
1987	20,726	36	103	238	485	939	1,593	2,492	3,755	11,034	1
1986	20,858	43	105	214	519	844	1,553	2,487	3,679	11,353	19
1985	22,021	40	107	256	489	930	1,587	2,634	3,996	11,917	20
1984	22,645	39	88	233	542	901	1,625	2,836	4,176	12,156	3
1983	24,451	58	139	267	558	961	1,840	2,967	4,604	13,022	1
1982	24,789	53	117	282	570	1,075	1,857	3,074	4,703	12,997	11
1981	25,824	38	132	273	554	1,062	1,870	3,256	4,971	13,613	5
1980	27,069	60	142	283	572	1,121	2,080	3,443	5,448	13,877	2
1979	26,602	55	115	277	561	1,164	2,074	3,397	5,372	13,545	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
White, male											
1998	5,213	32	61	110	203	310	552	804	1,005	2,108	-
1997	5,506	46	65	118	168	374	544	854	1,112	2,194	1
1996	5,746	46	81	88	220	338	594	909	1,161	2,273	1
1995	5,840	44	74	114	200	359	615	958	1,138	2,304	1
1994	5,838	26	57	105	194	385	596	877	1,178	2,385	-
1993	5,853	22	73	92	209	366	627	956	1,111	2,369	-
1992	5,756	35	56	98	205	374	633	917	1,098	2,309	1
1991	6,033	42	44	98	246	417	667	1,021	1,136	2,332	1
1990	6,232	25	65	105	274	427	692	984	1,175	2,455	2
1989	6,652	33	60	128	263	471	707	1,009	1,345	2,602	-
1988	7,529	27	63	137	294	500	880	1,143	1,493	2,953	1
1987	7,686	31	71	156	310	573	888	1,208	1,470	2,944	-
1986	7,767	33	74	143	342	523	883	1,201	1,479	3,051	9
1985	8,251	26	79	189	322	558	918	1,251	1,564	3,305	4
1984	8,447	32	60	144	325	537	884	1,344	1,689	3,397	1
1983	9,165	46	95	186	339	575	1,002	1,441	1,808	3,650	-
1982	9,689	42	80	191	365	664	1,031	1,518	1,889	3,872	3
1981	10,028	25	90	181	354	649	1,045	1,591	1,999	4,055	2
1980	10,543	40	96	189	366	705	1,162	1,573	2,180	4,202	-
1979	10,460	41	84	179	358	679	1,153	1,603	2,145	4,191	-
Atherosclerosis (440)											
White, female											
1998	8,837	19	25	55	102	207	419	816	1,424	5,757	-
1997	9,244	10	33	37	110	195	450	907	1,491	6,001	-
1996	9,633	11	27	45	96	208	528	923	1,526	6,249	-
1995	9,538	19	30	52	121	215	463	875	1,634	6,122	-
1994	9,822	9	19	57	98	247	536	886	1,586	6,368	1
1993	10,052	16	27	48	118	260	490	914	1,656	6,509	2
1992	9,737	7	28	56	149	267	506	909	1,569	6,227	1
1991	9,948	13	28	61	110	259	524	951	1,641	6,349	-
1990	10,315	18	25	44	134	290	534	965	1,725	6,567	-
1989	11,139	14	34	66	142	299	588	1,121	1,804	7,061	-
1988	12,732	11	36	72	149	359	634	1,208	2,190	8,056	1
1987	13,040	5	32	82	175	366	705	1,284	2,285	8,090	1
1986	13,091	10	31	71	177	321	670	1,286	2,200	8,302	10
1985	13,770	14	28	67	167	372	669	1,383	2,432	8,612	16
1984	14,198	7	28	89	217	364	741	1,492	2,487	8,759	2
1983	15,286	12	44	81	219	386	838	1,526	2,796	9,372	1
1982	15,100	11	37	91	205	411	826	1,556	2,814	9,125	8
1981	15,796	13	42	92	200	413	825	1,665	2,972	9,558	3
1980	16,526	20	46	94	206	416	918	1,870	3,268	9,675	2
1979	16,142	14	31	98	203	485	921	1,794	3,227	9,354	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
All other, both sexes											
1998	1,229	13	22	35	55	80	119	161	197	533	-
1997	1,307	19	35	37	50	95	147	169	205	535	-
1996	1,361	15	24	44	78	77	140	180	216	569	-
1995	1,345	9	20	41	67	104	151	189	241	515	-
1994	1,456	17	23	53	60	104	154	203	242	583	-
1993	1,367	11	22	39	69	101	160	171	249	521	1
1992	1,338	14	22	38	68	106	148	173	243	516	-
1991	1,439	14	16	45	82	100	159	211	259	542	-
1990	1,500	11	18	38	79	116	161	207	275	585	1
1989	1,566	9	18	39	83	112	171	224	258	641	1
1988	1,825	9	30	36	99	141	185	317	328	668	1
1987	1,748	15	20	53	94	139	204	299	298	619	1
1986	1,848	15	27	56	100	140	228	293	322	656	2
1985	1,905	9	26	63	84	146	235	301	332	692	5
1984	1,817	8	18	47	88	152	233	267	331	659	2
1983	1,920	10	35	58	109	148	242	319	384	604	3
1982	2,034	20	29	47	92	163	237	328	381	723	3
1981	2,180	14	33	55	107	189	277	340	409	733	5
1980	2,380	18	50	65	115	192	286	398	407	829	4
1979	2,199	11	27	69	116	204	294	359	397	715	1
Atherosclerosis (440)											
All other, male											
1998	506	8	15	23	27	48	73	73	88	143	-
1997	513	12	24	24	24	45	74	86	77	138	-
1996	535	7	13	24	46	35	67	76	95	164	-
1995	564	8	15	18	36	54	81	83	98	165	-
1994	582	11	11	29	36	55	70	91	104	164	-
1993	557	5	12	22	44	56	82	74	91	154	1
1992	572	10	17	22	40	54	83	82	101	157	-
1991	603	12	13	24	45	47	68	108	115	164	-
1990	627	6	14	21	37	59	82	102	101	199	1
1989	602	6	5	21	44	63	82	81	109	183	-
1988	798	4	25	19	51	78	93	148	146	225	-
1987	742	11	10	29	53	74	122	138	111	191	1
1986	820	14	14	39	61	76	115	138	141	216	-
1985	811	7	16	38	45	88	122	132	130	225	2
1984	793	7	7	27	46	89	121	132	149	207	1
1983	877	5	23	32	61	80	121	163	161	226	-
1982	894	11	19	21	48	95	132	168	154	240	1
1981	941	6	18	33	63	86	147	152	170	255	3
1980	1,058	13	32	38	73	105	143	173	165	306	1
1979	966	10	17	42	57	97	152	170	169	248	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
All other, female											
1998	723	5	7	12	28	32	46	88	109	390	-
1997	794	7	11	13	26	50	73	83	128	397	-
1996	826	8	11	20	32	42	73	104	121	405	-
1995	781	1	5	23	31	50	70	106	143	350	-
1994	874	6	12	24	24	49	84	112	138	419	-
1993	810	6	10	17	25	45	78	97	158	367	-
1992	766	4	5	16	28	52	65	91	142	359	-
1991	836	2	3	21	37	53	91	103	144	378	-
1990	873	5	4	17	42	57	79	105	174	386	-
1989	964	3	13	18	39	49	89	143	149	458	1
1988	1,027	5	5	17	48	63	92	169	182	443	1
1987	1,006	4	10	24	41	65	82	161	187	428	-
1986	1,028	1	13	17	39	64	113	155	181	440	2
1985	1,094	2	10	25	39	58	113	169	202	467	3
1984	1,024	1	11	20	42	63	112	135	182	452	1
1983	1,043	5	12	26	48	68	121	156	223	378	3
1982	1,140	9	10	26	44	68	105	160	227	483	2
1981	1,239	8	15	22	44	103	130	188	239	478	2
1980	1,322	5	18	27	42	87	143	225	242	523	3
1979	1,233	1	10	27	59	107	142	189	228	467	1
Atherosclerosis (440)											
Black, both sexes											
1998	1,082	11	20	33	54	71	109	136	173	463	-
1997	1,171	18	30	35	42	84	135	154	183	477	-
1996	1,200	14	22	40	72	67	130	152	195	492	-
1995	1,199	8	18	39	59	90	134	165	219	460	-
1994	1,299	14	19	47	57	97	137	184	211	518	-
1993	1,238	10	20	37	61	90	147	163	228	461	1
1992	1,209	13	21	33	60	95	130	161	216	471	-
1991	1,306	11	15	44	74	91	149	197	228	487	-
1990	1,380	9	17	38	75	109	149	185	257	531	1
1989	1,436	8	17	35	79	97	156	211	241	583	1
1988	1,694	9	28	34	93	137	171	302	300	609	1
1987	1,622	14	20	53	89	130	194	272	277	566	1
1986	1,720	15	27	54	95	130	215	276	289	608	2
1985	1,780	8	24	61	81	137	228	290	312	624	4
1984	1,684	7	17	45	84	145	219	255	302	598	2
1983	1,807	10	35	55	105	142	230	306	362	556	-
1982	1,898	19	27	42	86	160	221	314	358	659	2
1981	2,062	13	33	54	104	184	262	324	389	676	5
1980	2,252	18	48	61	111	185	273	379	384	775	3
1979	2,057	10	27	67	113	197	270	345	364	658	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Atherosclerosis (440)											
Black, male											
1998	436	7	14	22	27	44	67	60	75	114	-
1997	451	11	20	22	20	40	66	78	68	119	-
1996	458	7	12	21	44	31	61	61	85	129	-
1995	489	7	14	17	32	45	71	75	85	138	-
1994	507	9	9	26	35	52	62	80	88	137	-
1993	490	4	10	21	38	49	75	68	83	126	1
1992	504	9	16	18	35	51	68	72	86	143	-
1991	543	10	12	24	43	44	64	99	95	145	-
1990	563	4	14	21	35	55	76	87	92	173	1
1989	547	6	4	21	43	55	73	75	100	163	-
1988	739	4	23	18	47	77	85	142	133	202	-
1987	680	11	10	29	50	68	116	124	99	170	1
1986	756	14	14	37	57	69	109	128	121	201	-
1985	758	7	15	36	42	82	119	128	120	202	2
1984	720	6	6	27	42	85	115	124	131	178	1
1983	823	5	23	31	58	75	113	155	149	210	-
1982	826	11	18	18	44	94	122	162	141	212	-
1981	880	5	18	32	61	84	137	143	157	232	3
1980	990	13	31	36	71	102	135	160	153	280	-
1979	888	9	17	40	56	92	134	162	151	223	-
Atherosclerosis (440)											
Black, female											
1998	646	4	6	11	27	27	42	76	98	349	-
1997	720	7	10	13	22	44	69	76	115	358	-
1996	742	7	10	19	28	36	69	91	110	363	-
1995	710	1	4	22	27	45	63	90	134	322	-
1994	792	5	10	21	22	45	75	104	123	381	-
1993	748	6	10	16	23	41	72	95	145	335	-
1992	705	4	5	15	25	44	62	89	130	328	-
1991	763	1	3	20	31	47	85	98	133	342	-
1990	817	5	3	17	40	54	73	98	165	358	-
1989	889	2	13	14	36	42	83	136	141	420	1
1988	955	5	5	16	46	60	86	160	167	407	1
1987	942	3	10	24	39	62	78	148	178	396	-
1986	964	1	13	17	38	61	106	148	168	407	2
1985	1,022	1	9	25	39	55	109	162	192	422	2
1984	964	1	11	18	42	60	104	131	171	420	1
1983	984	5	12	24	47	67	117	151	213	346	-
1982	1,072	8	9	24	42	66	99	152	217	447	2
1981	1,182	8	15	22	43	100	125	181	232	444	2
1980	1,262	5	17	25	40	83	138	219	231	495	3
1979	1,169	1	10	27	57	105	136	183	213	435	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
All races, both sexes											
1998	27,671	470	589	895	1,502	2,622	3,999	5,069	4,816	6,836	-
1997	27,792	417	589	901	1,562	2,720	4,232	5,075	4,859	6,574	-
1996	27,176	451	535	825	1,588	2,710	4,157	4,928	4,767	6,416	1
1995	26,646	403	538	889	1,538	2,896	4,255	4,702	4,480	6,044	1
1994	26,097	353	523	837	1,596	2,897	4,185	4,659	4,452	5,818	3
1993	26,005	377	506	879	1,592	3,028	4,188	4,659	4,429	5,559	3
1992	25,337	357	503	821	1,644	2,939	4,171	4,566	4,248	5,303	2
1991	25,055	336	475	870	1,712	3,155	4,063	4,463	4,083	5,167	-
1990	24,645	334	455	864	1,717	3,110	3,999	4,502	4,050	4,943	1
1989	24,297	277	502	900	1,799	3,045	4,064	4,469	3,797	4,721	4
1988	23,185	245	456	866	1,819	3,005	3,906	4,206	3,602	4,450	1
1987	22,810	268	447	815	1,880	2,962	3,911	4,166	3,428	4,289	4
1986	22,417	244	430	866	1,894	2,945	3,862	4,097	3,451	4,010	2
1985	21,983	257	448	971	1,871	2,922	3,848	3,912	3,229	3,920	1
1984	21,741	246	471	979	1,908	2,934	3,631	3,855	3,258	3,841	8
1983	20,958	218	486	1,009	1,838	2,787	3,540	3,788	3,086	3,618	1
1982	20,172	256	474	975	1,785	2,870	3,589	3,559	2,928	3,227	4
1981	20,013	238	486	1,025	1,787	2,808	3,575	3,391	2,837	3,317	-
1980	19,959	292	530	1,061	1,827	2,832	3,486	3,410	2,874	3,088	1
1979	19,483	273	547	1,039	1,822	2,797	3,355	3,361	2,785	2,926	3
Other diseases of circulatory system (441-448)											
All races, male											
1998	14,991	318	387	586	1,013	1,679	2,430	2,945	2,477	2,625	-
1997	15,109	259	395	589	1,039	1,746	2,638	2,866	2,497	2,547	-
1996	14,857	304	363	581	1,084	1,756	2,576	2,874	2,464	2,377	1
1995	14,817	259	357	590	1,059	1,955	2,694	2,754	2,292	2,305	-
1994	14,518	209	346	547	1,076	1,937	2,679	2,724	2,299	2,251	2
1993	14,963	245	325	600	1,133	2,091	2,680	2,860	2,401	2,160	2
1992	14,504	220	323	547	1,148	2,062	2,701	2,714	2,248	2,070	2
1991	14,569	213	312	606	1,226	2,174	2,690	2,705	2,174	2,031	-
1990	14,560	213	313	608	1,221	2,122	2,681	2,779	2,240	1,979	1
1989	14,447	181	332	612	1,302	2,147	2,666	2,726	2,128	1,905	3
1988	13,814	148	290	602	1,312	2,094	2,607	2,620	1,962	1,782	1
1987	13,840	175	298	564	1,348	2,088	2,635	2,625	1,939	1,782	3
1986	13,525	141	292	626	1,367	2,084	2,664	2,556	1,892	1,555	1
1985	13,620	156	309	694	1,385	2,107	2,695	2,456	1,861	1,600	1
1984	13,550	159	318	695	1,421	2,123	2,518	2,435	1,901	1,619	3
1983	13,243	137	334	710	1,370	2,016	2,497	2,503	1,776	1,545	-
1982	12,739	162	317	683	1,328	2,084	2,503	2,345	1,659	1,344	2
1981	12,794	149	340	725	1,354	2,048	2,506	2,309	1,635	1,414	-
1980	12,933	179	375	791	1,405	2,081	2,468	2,311	1,646	1,357	1
1979	12,620	169	388	734	1,360	2,071	2,426	2,211	1,652	1,276	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
All races, female											
1998	12,680	152	202	309	489	943	1,569	2,124	2,339	4,211	-
1997	12,683	158	194	312	523	974	1,594	2,209	2,362	4,027	-
1996	12,319	147	172	244	504	954	1,581	2,054	2,303	4,039	-
1995	11,829	144	181	299	479	941	1,561	1,948	2,188	3,739	1
1994	11,579	144	177	290	520	960	1,506	1,935	2,153	3,567	1
1993	11,042	132	181	279	459	937	1,508	1,799	2,028	3,399	1
1992	10,833	137	180	274	496	877	1,470	1,852	2,000	3,233	-
1991	10,486	123	163	264	486	981	1,373	1,758	1,909	3,136	-
1990	10,085	121	142	256	496	988	1,318	1,723	1,810	2,964	-
1989	9,850	96	170	288	497	898	1,398	1,743	1,669	2,816	1
1988	9,371	97	166	264	507	911	1,299	1,586	1,640	2,668	-
1987	8,970	93	149	251	532	874	1,276	1,541	1,489	2,507	1
1986	8,892	103	138	240	527	861	1,198	1,541	1,559	2,455	1
1985	8,363	101	139	277	486	815	1,153	1,456	1,368	2,320	-
1984	8,191	87	153	284	487	811	1,113	1,420	1,357	2,222	5
1983	7,715	81	152	299	468	771	1,043	1,285	1,310	2,073	1
1982	7,433	94	157	292	457	786	1,086	1,214	1,269	1,883	2
1981	7,219	89	146	300	433	760	1,069	1,082	1,202	1,903	-
1980	7,026	113	155	270	422	751	1,018	1,099	1,228	1,731	-
1979	6,863	104	159	305	462	726	929	1,150	1,133	1,650	-
Other diseases of circulatory system (441-448)											
White, both sexes											
1998	24,564	329	454	713	1,270	2,329	3,605	4,615	4,411	6,213	-
1997	24,791	284	442	713	1,331	2,408	3,810	4,684	4,461	6,045	-
1996	24,308	328	425	663	1,382	2,401	3,745	4,498	4,392	5,914	-
1995	23,966	281	438	723	1,327	2,593	3,864	4,359	4,142	5,578	1
1994	23,363	260	410	677	1,382	2,584	3,807	4,272	4,074	5,328	2
1993	23,334	266	394	708	1,399	2,705	3,826	4,309	4,079	5,096	2
1992	22,895	256	386	671	1,450	2,647	3,830	4,232	3,935	4,896	2
1991	22,636	250	361	736	1,506	2,847	3,732	4,126	3,785	4,787	-
1990	22,291	255	343	725	1,523	2,822	3,674	4,160	3,761	4,550	1
1989	21,988	205	377	745	1,596	2,768	3,745	4,116	3,513	4,389	4
1988	21,072	186	360	735	1,584	2,735	3,611	3,918	3,340	4,145	1
1987	20,725	212	353	694	1,669	2,681	3,621	3,832	3,187	4,008	4
1986	20,422	175	326	723	1,680	2,709	3,578	3,798	3,221	3,764	2
1985	20,021	195	357	840	1,666	2,652	3,533	3,621	3,003	3,689	1
1984	19,839	175	365	840	1,689	2,698	3,329	3,616	3,048	3,605	6
1983	19,189	170	406	874	1,640	2,560	3,274	3,540	2,895	3,399	1
1982	18,489	205	389	834	1,590	2,605	3,300	3,351	2,763	3,057	4
1981	18,374	185	400	882	1,608	2,573	3,334	3,188	2,670	3,138	-
1980	18,260	233	447	923	1,650	2,578	3,232	3,174	2,694	2,904	1
1979	17,952	219	450	915	1,661	2,557	3,142	3,151	2,647	2,795	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
White, male											
1998	13,484	232	305	476	884	1,522	2,248	2,721	2,297	2,406	-
1997	13,660	175	313	481	917	1,596	2,423	2,673	2,318	2,371	-
1996	13,433	224	294	482	957	1,578	2,369	2,660	2,298	2,220	-
1995	13,529	177	302	491	953	1,784	2,501	2,590	2,147	2,171	-
1994	13,200	159	281	455	965	1,773	2,475	2,524	2,125	2,099	1
1993	13,599	178	262	503	1,023	1,900	2,489	2,682	2,231	1,994	2
1992	13,263	168	252	471	1,032	1,872	2,525	2,538	2,095	1,944	2
1991	13,356	166	251	521	1,106	1,998	2,521	2,537	2,036	1,901	-
1990	13,344	166	249	525	1,111	1,958	2,488	2,590	2,096	1,862	1
1989	13,279	140	257	519	1,189	1,966	2,503	2,562	1,984	1,801	3
1988	12,733	116	231	529	1,167	1,944	2,439	2,476	1,841	1,692	1
1987	12,728	136	240	500	1,218	1,922	2,481	2,437	1,821	1,680	3
1986	12,492	103	234	536	1,236	1,942	2,518	2,395	1,788	1,477	1
1985	12,600	123	256	619	1,256	1,948	2,519	2,309	1,759	1,517	1
1984	12,532	116	260	613	1,290	1,985	2,341	2,307	1,797	1,534	2
1983	12,298	106	282	630	1,255	1,891	2,339	2,373	1,691	1,462	-
1982	11,818	131	259	600	1,205	1,916	2,342	2,239	1,589	1,284	2
1981	11,934	122	288	650	1,242	1,918	2,375	2,200	1,555	1,349	-
1980	11,992	147	327	705	1,290	1,930	2,328	2,178	1,560	1,275	1
1979	11,746	143	329	660	1,260	1,914	2,291	2,107	1,573	1,220	1
Other diseases of circulatory system (441-448)											
White, female											
1998	11,080	97	149	237	386	807	1,357	1,894	2,114	3,807	-
1997	11,131	109	129	232	414	812	1,387	2,011	2,143	3,674	-
1996	10,875	104	131	181	425	823	1,376	1,838	2,094	3,694	-
1995	10,437	104	136	232	374	809	1,363	1,769	1,995	3,407	1
1994	10,163	101	129	222	417	811	1,332	1,748	1,949	3,229	1
1993	9,735	88	132	205	376	805	1,337	1,627	1,848	3,102	-
1992	9,632	88	134	200	418	775	1,305	1,694	1,840	2,952	-
1991	9,280	84	110	215	400	849	1,211	1,589	1,749	2,886	-
1990	8,947	89	94	200	412	864	1,186	1,570	1,665	2,688	-
1989	8,709	65	120	226	407	802	1,242	1,554	1,529	2,588	1
1988	8,339	70	129	206	417	791	1,172	1,442	1,499	2,453	-
1987	7,997	76	113	194	451	759	1,140	1,395	1,366	2,328	1
1986	7,930	72	92	187	444	767	1,060	1,403	1,433	2,287	1
1985	7,421	72	101	221	410	704	1,014	1,312	1,244	2,172	-
1984	7,307	59	105	227	399	713	988	1,309	1,251	2,071	4
1983	6,891	64	124	244	385	669	935	1,167	1,204	1,937	1
1982	6,671	74	130	234	385	689	958	1,112	1,174	1,773	2
1981	6,440	63	112	232	366	655	959	988	1,115	1,789	-
1980	6,268	86	120	218	360	648	904	996	1,134	1,629	-
1979	6,206	76	121	255	401	643	851	1,044	1,074	1,575	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
All other, both sexes											
1998	3,107	141	135	182	232	293	394	454	405	623	-
1997	3,001	133	147	188	231	312	422	391	398	529	-
1996	2,868	123	110	162	206	309	412	430	375	502	1
1995	2,680	122	100	166	211	303	391	343	338	466	-
1994	2,734	93	113	160	214	313	378	387	378	490	1
1993	2,671	111	112	171	193	323	362	350	350	463	1
1992	2,442	101	117	150	194	292	341	334	313	407	-
1991	2,419	86	114	134	206	308	331	337	298	380	-
1990	2,354	79	112	139	194	288	325	342	289	393	-
1989	2,309	72	125	155	203	277	319	353	284	332	-
1988	2,113	59	96	131	235	270	295	288	262	305	-
1987	2,085	56	94	121	211	281	290	334	241	281	-
1986	1,995	69	104	143	214	236	284	299	230	246	-
1985	1,962	62	91	131	205	270	315	291	226	231	-
1984	1,902	71	106	139	219	236	302	239	210	236	2
1983	1,769	48	80	135	198	227	266	248	191	219	-
1982	1,683	51	85	141	195	265	289	208	165	170	-
1981	1,639	53	86	143	179	235	241	203	167	179	-
1980	1,699	59	83	138	177	254	254	236	180	184	-
1979	1,531	54	97	124	161	240	213	210	138	131	2
Other diseases of circulatory system (441-448)											
All other, male											
1998	1,507	86	82	110	129	157	182	224	180	219	-
1997	1,449	84	82	108	122	150	215	193	179	176	-
1996	1,424	80	69	99	127	178	207	214	166	157	1
1995	1,288	82	55	99	106	171	193	164	145	134	-
1994	1,318	50	65	92	111	164	204	200	174	152	1
1993	1,364	67	63	97	110	191	191	178	170	166	-
1992	1,241	52	71	76	116	190	176	176	153	126	-
1991	1,213	47	61	85	120	176	169	168	138	130	-
1990	1,216	47	64	83	110	164	193	189	144	117	-
1989	1,168	41	75	93	113	181	163	164	144	104	-
1988	1,081	32	59	73	145	150	168	144	121	90	-
1987	1,112	39	58	64	130	166	154	188	118	102	-
1986	1,033	38	58	90	131	142	146	161	104	78	-
1985	1,020	33	53	75	129	159	176	147	102	83	-
1984	1,018	43	58	82	131	138	177	128	104	85	1
1983	945	31	52	80	115	125	158	130	85	83	-
1982	921	31	58	83	123	168	161	106	70	60	-
1981	860	27	52	75	112	130	131	109	80	65	-
1980	941	32	48	86	115	151	140	133	86	82	-
1979	874	26	59	74	100	157	135	104	79	56	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
All other, female											
1998	1,600	55	53	72	103	136	212	230	225	404	-
1997	1,552	49	65	80	109	162	207	198	219	353	-
1996	1,444	43	41	63	79	131	205	216	209	345	-
1995	1,392	40	45	67	105	132	198	179	193	332	-
1994	1,416	43	48	68	103	149	174	187	204	338	-
1993	1,307	44	49	74	83	132	171	172	180	297	1
1992	1,201	49	46	74	78	102	165	158	160	281	-
1991	1,206	39	53	49	86	132	162	169	160	250	-
1990	1,138	32	48	56	84	124	132	153	145	276	-
1989	1,141	31	50	62	90	96	156	189	140	228	-
1988	1,032	27	37	58	90	120	127	144	141	215	-
1987	973	17	36	57	81	115	136	146	123	179	-
1986	962	31	46	53	83	94	138	138	126	168	-
1985	942	29	38	56	76	111	139	144	124	148	-
1984	884	28	48	57	88	98	125	111	106	151	1
1983	824	17	28	55	83	102	108	118	106	136	-
1982	762	20	27	58	72	97	128	102	95	110	-
1981	779	26	34	68	67	105	110	94	87	114	-
1980	758	27	35	52	62	103	114	103	94	102	-
1979	657	28	38	50	61	83	78	106	59	75	-
Other diseases of circulatory system (441-448)											
Black, both sexes											
1998	2,666	133	114	161	194	255	321	386	331	547	-
1997	2,586	122	132	155	196	265	356	336	333	462	-
1996	2,455	113	97	144	173	275	345	369	299	432	1
1995	2,342	109	89	148	187	273	329	302	291	401	-
1994	2,363	83	100	131	185	274	332	330	310	435	1
1993	2,301	103	104	151	160	276	304	293	296	411	1
1992	2,115	90	106	120	173	252	297	299	252	353	-
1991	2,137	81	112	116	184	265	289	289	255	338	-
1990	2,061	70	106	120	170	247	283	300	244	348	-
1989	2,058	64	112	147	182	247	278	315	240	296	-
1988	1,861	53	84	118	205	238	261	252	225	265	-
1987	1,841	53	88	103	192	251	253	289	203	249	-
1986	1,761	64	90	130	191	217	253	255	190	218	-
1985	1,732	57	85	122	184	239	274	249	198	199	-
1984	1,697	68	97	122	205	216	264	207	186	202	2
1983	1,561	46	73	124	181	205	235	213	155	189	-
1982	1,526	50	80	127	183	242	262	178	151	150	-
1981	1,485	50	77	132	164	218	214	182	147	157	-
1980	1,532	57	76	126	165	228	229	203	159	161	-
1979	1,387	50	90	116	149	226	189	184	118	117	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of circulatory system (441-448)											
Black, male											
1998	1,252	82	68	96	109	132	146	183	136	177	-
1997	1,229	75	73	90	100	130	186	157	142	145	-
1996	1,186	72	59	87	110	154	170	172	123	127	1
1995	1,078	70	47	90	93	153	149	138	112	105	-
1994	1,103	47	56	74	98	139	175	172	133	119	1
1993	1,143	63	61	91	91	161	151	136	139	139	-
1992	1,032	48	63	55	105	158	149	153	109	100	-
1991	1,027	43	60	71	105	146	136	139	111	106	-
1990	1,036	41	60	71	97	140	160	159	115	100	-
1989	1,009	33	69	90	100	160	139	139	112	85	-
1988	919	32	50	66	123	130	144	120	94	68	-
1987	940	37	52	56	116	146	130	152	88	81	-
1986	886	35	47	78	119	131	124	132	79	66	-
1985	871	31	50	70	115	136	146	117	80	70	-
1984	894	41	54	72	121	126	151	106	85	72	1
1983	820	31	47	73	104	110	138	104	64	70	-
1982	817	31	55	78	113	149	142	84	62	49	-
1981	760	25	45	68	102	116	112	95	66	56	-
1980	821	31	45	76	105	132	123	106	71	67	-
1979	764	24	55	68	90	144	113	84	64	48	1
Other diseases of circulatory system (441-448)											
Black, female											
1998	1,414	51	46	65	85	123	175	203	195	370	-
1997	1,357	47	59	65	96	135	170	179	191	317	-
1996	1,269	41	38	57	63	121	175	197	176	305	-
1995	1,264	39	42	58	94	120	180	164	179	296	-
1994	1,260	36	44	57	87	135	157	158	177	316	-
1993	1,158	40	43	60	69	115	153	157	157	272	1
1992	1,083	42	43	65	68	94	148	146	143	253	-
1991	1,110	38	52	45	79	119	153	150	144	232	-
1990	1,025	29	46	49	73	107	123	141	129	248	-
1989	1,049	31	43	57	82	87	139	176	128	211	-
1988	942	21	34	52	82	108	117	132	131	197	-
1987	901	16	36	47	76	105	123	137	115	168	-
1986	875	29	43	52	72	86	129	123	111	152	-
1985	861	26	35	52	69	103	128	132	118	129	-
1984	803	27	43	50	84	90	113	101	101	130	1
1983	741	15	26	51	77	95	97	109	91	119	-
1982	709	19	25	49	70	93	120	94	89	101	-
1981	725	25	32	64	62	102	102	87	81	101	-
1980	711	26	31	50	60	96	106	97	88	94	-
1979	623	26	35	48	59	82	76	100	54	69	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
All races, both sexes											
1998	16,237	272	330	506	954	1,717	2,613	3,234	2,925	3,235	-
1997	16,724	247	351	566	942	1,791	2,797	3,290	3,038	3,237	-
1996	16,387	261	325	526	1,025	1,789	2,730	3,218	3,022	3,095	1
1995	16,440	237	331	582	1,000	1,949	2,907	3,098	2,805	3,037	-
1994	16,123	191	315	538	1,038	1,972	2,892	3,065	2,795	2,911	2
1993	16,475	222	308	558	1,078	2,111	2,927	3,173	2,845	2,847	3
1992	16,220	211	324	511	1,105	2,068	2,931	3,126	2,783	2,777	1
1991	16,696	203	286	600	1,209	2,271	2,996	3,143	2,767	2,860	-
1990	16,402	195	258	565	1,192	2,303	2,858	3,192	2,740	2,761	1
1989	16,229	170	302	588	1,222	2,248	2,940	3,170	2,542	2,685	2
1988	15,719	141	265	554	1,307	2,173	2,840	3,034	2,524	2,550	1
1987	15,637	163	267	540	1,306	2,136	2,843	3,032	2,431	2,575	3
1986	15,267	136	270	576	1,308	2,105	2,858	2,947	2,401	2,377	1
1985	15,112	142	257	624	1,303	2,127	2,845	2,839	2,295	2,393	1
1984	15,139	139	256	633	1,325	2,134	2,714	2,836	2,403	2,395	5
1983	14,576	117	296	630	1,257	2,045	2,628	2,828	2,223	2,284	1
1982	14,125	145	267	624	1,232	2,094	2,664	2,697	2,101	2,069	4
1981	14,082	124	289	621	1,259	2,053	2,682	2,559	2,094	2,158	-
1980	14,181	143	293	701	1,297	2,098	2,691	2,603	2,093	2,024	1
1979	14,031	151	313	673	1,295	2,080	2,621	2,620	2,093	1,943	2
Aortic aneurysm and dissection (441)											
All races, male											
1998	10,013	215	262	376	708	1,200	1,729	2,046	1,611	1,519	-
1997	10,311	180	266	412	700	1,248	1,881	1,989	1,718	1,555	-
1996	10,117	199	251	415	765	1,260	1,817	2,030	1,689	1,408	1
1995	10,328	180	260	443	748	1,427	1,978	1,944	1,570	1,416	-
1994	10,146	132	222	398	772	1,427	2,010	1,921	1,608	1,367	2
1993	10,664	172	239	422	829	1,583	2,016	2,133	1,666	1,309	2
1992	10,407	158	238	399	844	1,558	2,002	1,970	1,652	1,302	1
1991	10,776	140	212	450	937	1,691	2,118	2,035	1,583	1,340	-
1990	10,806	151	209	435	931	1,684	2,035	2,088	1,680	1,338	1
1989	10,708	123	228	436	954	1,687	2,045	2,111	1,565	1,276	2
1988	10,429	101	196	431	1,008	1,637	2,017	2,017	1,523	1,241	1
1987	10,383	120	205	411	1,008	1,596	2,044	2,027	1,474	1,250	3
1986	10,171	91	209	451	1,025	1,581	2,061	1,980	1,479	1,096	-
1985	10,208	103	199	499	1,040	1,637	2,095	1,892	1,409	1,123	1
1984	10,337	105	197	497	1,066	1,652	1,991	1,922	1,527	1,156	1
1983	10,024	83	228	497	994	1,555	1,980	1,995	1,376	1,124	-
1982	9,676	105	202	487	979	1,624	1,949	1,896	1,291	966	2
1981	9,779	94	224	499	1,031	1,581	1,995	1,847	1,302	1,029	-
1980	10,005	106	226	576	1,070	1,634	2,029	1,871	1,315	1,001	1
1979	9,779	113	247	523	1,036	1,633	1,983	1,819	1,319	927	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
All races, female											
1998	6,224	57	68	130	246	517	884	1,188	1,314	1,716	-
1997	6,413	67	85	154	242	543	916	1,301	1,320	1,682	-
1996	6,270	62	74	111	260	529	913	1,188	1,333	1,687	-
1995	6,112	57	71	139	252	522	929	1,154	1,235	1,621	-
1994	5,977	59	93	140	266	545	882	1,144	1,187	1,544	-
1993	5,811	50	69	136	249	528	911	1,040	1,179	1,538	1
1992	5,813	53	86	112	261	510	929	1,156	1,131	1,475	-
1991	5,920	63	74	150	272	580	878	1,108	1,184	1,520	-
1990	5,596	44	49	130	261	619	823	1,104	1,060	1,423	-
1989	5,521	47	74	152	268	561	895	1,059	977	1,409	-
1988	5,290	40	69	123	299	536	823	1,017	1,001	1,309	-
1987	5,254	43	62	129	298	540	799	1,005	957	1,325	-
1986	5,096	45	61	125	283	524	797	967	922	1,281	1
1985	4,904	39	58	125	263	490	750	947	886	1,270	-
1984	4,802	34	59	136	259	482	723	914	876	1,239	4
1983	4,552	34	68	133	263	490	648	833	847	1,160	1
1982	4,449	40	65	137	253	470	715	801	810	1,103	2
1981	4,303	30	65	122	228	472	687	712	792	1,129	-
1980	4,176	37	67	125	227	464	662	732	778	1,023	-
1979	4,252	38	66	150	259	447	638	801	774	1,016	-
Aortic aneurysm and dissection (441)											
White, both sexes											
1998	14,775	192	245	418	826	1,561	2,426	3,009	2,739	3,008	-
1997	15,281	160	270	464	841	1,637	2,583	3,102	2,844	3,033	-
1996	14,956	189	267	426	919	1,621	2,520	2,995	2,825	2,904	-
1995	15,063	149	268	477	881	1,780	2,700	2,919	2,636	2,869	-
1994	14,798	144	246	445	929	1,824	2,678	2,883	2,604	2,742	1
1993	15,123	161	249	464	967	1,924	2,720	2,987	2,674	2,663	2
1992	14,955	151	251	420	995	1,907	2,742	2,954	2,616	2,618	1
1991	15,380	149	217	516	1,089	2,096	2,793	2,950	2,596	2,707	-
1990	15,119	152	193	477	1,072	2,115	2,679	2,996	2,591	2,596	1
1989	14,959	121	227	505	1,099	2,080	2,758	2,964	2,383	2,540	2
1988	14,528	108	210	471	1,150	2,005	2,665	2,865	2,382	2,410	1
1987	14,467	127	214	461	1,176	1,973	2,685	2,829	2,288	2,447	3
1986	14,177	99	201	490	1,188	1,980	2,687	2,766	2,278	2,261	1
1985	13,984	102	208	546	1,183	1,968	2,646	2,664	2,165	2,271	1
1984	14,034	95	191	546	1,205	1,982	2,524	2,691	2,272	2,285	3
1983	13,566	85	242	564	1,133	1,906	2,473	2,678	2,107	2,174	1
1982	13,166	122	217	540	1,124	1,944	2,478	2,564	2,011	1,982	4
1981	13,103	94	237	540	1,157	1,898	2,520	2,427	1,988	2,058	-
1980	13,165	115	236	614	1,197	1,931	2,529	2,447	1,987	1,920	1
1979	13,088	117	253	600	1,181	1,915	2,479	2,476	2,016	1,870	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
White, male											
1998	9,208	160	205	316	625	1,121	1,628	1,927	1,525	1,428	-
1997	9,518	119	214	349	639	1,167	1,764	1,890	1,625	1,483	-
1996	9,316	145	209	350	691	1,153	1,708	1,911	1,594	1,340	-
1995	9,560	112	217	377	690	1,328	1,857	1,857	1,487	1,355	-
1994	9,417	103	179	338	708	1,335	1,888	1,829	1,512	1,293	1
1993	9,891	130	195	363	763	1,469	1,893	2,029	1,586	1,237	2
1992	9,677	118	187	342	781	1,450	1,904	1,872	1,555	1,243	1
1991	10,030	111	168	391	857	1,583	2,008	1,931	1,495	1,287	-
1990	10,068	115	168	381	857	1,570	1,928	1,978	1,600	1,285	1
1989	9,998	90	180	383	878	1,574	1,948	2,007	1,481	1,228	2
1988	9,758	80	155	384	908	1,537	1,915	1,930	1,455	1,191	1
1987	9,720	94	166	367	923	1,502	1,962	1,911	1,403	1,195	3
1986	9,559	70	167	392	944	1,511	1,966	1,881	1,414	1,056	-
1985	9,580	77	169	448	964	1,539	1,981	1,799	1,351	1,076	1
1984	9,685	76	158	439	992	1,554	1,879	1,836	1,457	1,113	-
1983	9,451	58	190	450	924	1,478	1,883	1,914	1,324	1,084	-
1982	9,111	88	166	436	902	1,527	1,845	1,819	1,247	940	2
1981	9,231	75	189	452	957	1,492	1,903	1,774	1,252	996	-
1980	9,418	87	193	519	1,004	1,533	1,938	1,782	1,268	952	1
1979	9,211	95	210	474	965	1,519	1,891	1,750	1,273	896	1
Aortic aneurysm and dissection (441)											
White, female											
1998	5,567	32	40	102	201	440	798	1,082	1,214	1,580	-
1997	5,763	41	56	115	202	470	819	1,212	1,219	1,550	-
1996	5,640	44	58	76	228	468	812	1,084	1,231	1,564	-
1995	5,503	37	51	100	191	452	843	1,062	1,149	1,514	-
1994	5,381	41	67	107	221	489	790	1,054	1,092	1,449	-
1993	5,232	31	54	101	204	455	827	958	1,088	1,426	-
1992	5,278	33	64	78	214	457	838	1,082	1,061	1,375	-
1991	5,350	38	49	125	232	513	785	1,019	1,101	1,420	-
1990	5,051	37	25	96	215	545	751	1,018	991	1,311	-
1989	4,961	31	47	122	221	506	810	957	902	1,312	-
1988	4,770	28	55	87	242	468	750	935	927	1,219	-
1987	4,747	33	48	94	253	471	723	918	885	1,252	-
1986	4,618	29	34	98	244	469	721	885	864	1,205	1
1985	4,404	25	39	98	219	429	665	865	814	1,195	-
1984	4,349	19	33	107	213	428	645	855	815	1,172	3
1983	4,115	27	52	114	209	428	590	764	783	1,090	1
1982	4,055	34	51	104	222	417	633	745	764	1,042	2
1981	3,872	19	48	88	200	406	617	653	736	1,062	-
1980	3,747	28	43	95	193	398	591	665	719	968	-
1979	3,877	22	43	126	216	396	588	726	743	974	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
All other, both sexes											
1998	1,462	80	85	88	128	156	187	225	186	227	-
1997	1,443	87	81	102	101	154	214	188	194	204	-
1996	1,431	72	58	100	106	168	210	223	197	191	1
1995	1,377	88	63	105	119	169	207	179	169	168	-
1994	1,325	47	69	93	109	148	214	182	191	169	1
1993	1,352	61	59	94	111	187	207	186	171	184	1
1992	1,265	60	73	91	110	161	189	172	167	159	-
1991	1,316	54	69	84	120	175	203	193	171	153	-
1990	1,283	43	65	88	120	188	179	196	149	165	-
1989	1,270	49	75	83	123	168	182	206	159	145	-
1988	1,191	33	55	83	157	168	175	169	142	140	-
1987	1,170	36	53	79	130	163	158	203	143	128	-
1986	1,090	37	69	86	120	125	171	181	123	116	-
1985	1,128	40	49	78	120	159	199	175	130	122	-
1984	1,105	44	65	87	120	152	190	145	131	110	2
1983	1,010	32	54	66	124	139	155	150	116	110	-
1982	959	23	50	84	108	150	186	133	90	87	-
1981	979	30	52	81	102	155	162	132	106	100	-
1980	1,016	28	57	87	100	167	162	156	106	104	-
1979	943	34	60	73	114	165	142	144	77	73	1
Aortic aneurysm and dissection (441)											
All other, male											
1998	805	55	57	60	83	79	101	119	86	91	-
1997	793	61	52	63	61	81	117	99	93	72	-
1996	801	54	42	65	74	107	109	119	95	68	1
1995	768	68	43	66	58	99	121	87	83	61	-
1994	729	29	43	60	64	92	122	92	96	74	1
1993	773	42	44	59	66	114	123	104	80	72	-
1992	730	40	51	57	63	108	98	98	97	59	-
1991	746	29	44	59	80	108	110	104	88	53	-
1990	738	36	41	54	74	114	107	110	80	53	-
1989	710	33	48	53	76	113	97	104	84	48	-
1988	671	21	41	47	100	100	102	87	68	50	-
1987	663	26	39	44	85	94	82	116	71	55	-
1986	612	21	42	59	81	70	95	99	65	40	-
1985	628	26	30	51	76	98	114	93	58	47	-
1984	652	29	39	58	74	98	112	86	70	43	1
1983	573	25	38	47	70	77	97	81	52	40	-
1982	565	17	36	51	77	97	104	77	44	26	-
1981	548	19	35	47	74	89	92	73	50	33	-
1980	587	19	33	57	66	101	91	89	47	49	-
1979	568	18	37	49	71	114	92	69	46	31	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
All other, female											
1998	657	25	28	28	45	77	86	106	100	136	-
1997	650	26	29	39	40	73	97	89	101	132	-
1996	630	18	16	35	32	61	101	104	102	123	-
1995	609	20	20	39	61	70	86	92	86	107	-
1994	596	18	26	33	45	56	92	90	95	95	-
1993	579	19	15	35	45	73	84	82	91	112	1
1992	535	20	22	34	47	53	91	74	70	100	-
1991	570	25	25	25	40	67	93	89	83	100	-
1990	545	7	24	34	46	74	72	86	69	112	-
1989	560	16	27	30	47	55	85	102	75	97	-
1988	520	12	14	36	57	68	73	82	74	90	-
1987	507	10	14	35	45	69	76	87	72	73	-
1986	478	16	27	27	39	55	76	82	58	76	-
1985	500	14	19	27	44	61	85	82	72	75	-
1984	453	15	26	29	46	54	78	59	61	67	1
1983	437	7	16	19	54	62	58	69	64	70	-
1982	394	6	14	33	31	53	82	56	46	61	-
1981	431	11	17	34	28	66	70	59	56	67	-
1980	429	9	24	30	34	66	71	67	59	55	-
1979	375	16	23	24	43	51	50	75	31	42	-
Aortic aneurysm and dissection (441)											
Black, both sexes											
1998	1,184	74	69	72	104	132	138	182	143	184	-
1997	1,183	79	73	84	81	121	171	149	151	167	-
1996	1,140	63	50	87	84	138	164	182	136	143	1
1995	1,146	78	54	94	104	149	165	148	132	129	-
1994	1,087	40	61	76	95	123	180	142	145	133	1
1993	1,107	56	56	82	89	152	161	144	136	152	1
1992	1,030	55	66	67	95	130	155	150	116	125	-
1991	1,104	50	68	72	103	143	169	154	133	123	-
1990	1,068	38	61	75	99	160	145	163	114	134	-
1989	1,085	41	67	77	110	147	148	177	122	121	-
1988	1,006	31	45	73	135	140	150	142	115	110	-
1987	985	33	49	66	114	142	133	167	112	103	-
1986	916	34	57	76	104	110	145	150	90	94	-
1985	954	37	43	73	102	134	167	141	110	98	-
1984	954	42	61	74	110	136	158	118	112	87	2
1983	863	31	50	61	109	121	131	124	88	93	-
1982	845	22	49	75	99	134	164	108	80	71	-
1981	866	29	46	74	92	142	141	114	89	84	-
1980	882	26	52	79	89	145	142	127	90	86	-
1979	823	30	55	67	106	152	118	124	60	60	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Aortic aneurysm and dissection (441)											
Black, male											
1998	620	52	47	49	67	63	70	88	58	62	-
1997	628	55	47	51	46	64	94	71	65	49	-
1996	616	47	35	55	61	85	80	88	59	48	1
1995	610	58	36	59	50	85	88	67	57	40	-
1994	571	27	37	46	56	74	98	71	67	45	1
1993	608	39	43	54	52	89	89	70	61	54	-
1992	564	38	45	37	56	82	77	81	58	40	-
1991	597	26	43	49	66	86	82	81	64	34	-
1990	589	31	39	44	62	95	79	85	56	38	-
1989	592	25	45	52	66	98	75	86	59	37	-
1988	545	21	33	42	84	80	85	67	51	30	-
1987	529	24	35	37	72	81	63	87	47	40	-
1986	494	19	32	50	70	62	76	77	44	29	-
1985	506	24	27	47	64	78	90	68	43	35	-
1984	550	28	36	50	65	87	88	65	55	36	1
1983	476	25	34	42	60	65	82	61	36	32	-
1982	488	17	35	49	68	84	90	57	38	17	-
1981	469	18	30	41	66	77	77	60	37	27	-
1980	485	18	30	50	56	86	77	66	35	35	-
1979	474	16	35	44	64	102	70	53	33	23	-
Aortic aneurysm and dissection (441)											
Black, female											
1998	564	22	22	23	37	69	68	94	85	122	-
1997	555	24	26	33	35	57	77	78	86	118	-
1996	524	16	15	32	23	53	84	94	77	95	-
1995	536	20	18	35	54	64	77	81	75	89	-
1994	516	13	24	30	39	49	82	71	78	88	-
1993	499	17	13	28	37	63	72	74	75	98	1
1992	466	17	21	30	39	48	78	69	58	85	-
1991	507	24	25	23	37	57	87	73	69	89	-
1990	479	7	22	31	37	65	66	78	58	96	-
1989	493	16	22	25	44	49	73	91	63	84	-
1988	461	10	12	31	51	60	65	75	64	80	-
1987	456	9	14	29	42	61	70	80	65	63	-
1986	422	15	25	26	34	48	69	73	46	65	-
1985	448	13	16	26	38	56	77	73	67	63	-
1984	404	14	25	24	45	49	70	53	57	51	1
1983	387	6	16	19	49	56	49	63	52	61	-
1982	357	5	14	26	31	50	74	51	42	54	-
1981	397	11	16	33	26	65	64	54	52	57	-
1980	397	8	22	29	33	59	65	61	55	51	-
1979	349	14	20	23	42	50	48	71	27	37	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
All races, both sexes											
1998	11,434	198	259	389	548	905	1,386	1,835	1,891	3,601	-
1997	11,068	170	238	335	620	929	1,435	1,785	1,821	3,337	-
1996	10,789	190	210	299	563	921	1,427	1,710	1,745	3,321	-
1995	10,206	166	207	307	538	947	1,348	1,604	1,675	3,007	1
1994	9,974	162	208	299	558	925	1,293	1,594	1,657	2,907	1
1993	9,530	155	198	321	514	917	1,261	1,486	1,584	2,712	-
1992	9,117	146	179	310	539	871	1,240	1,440	1,465	2,526	1
1991	8,359	133	189	270	503	884	1,067	1,320	1,316	2,307	-
1990	8,243	139	197	299	525	807	1,141	1,310	1,310	2,182	-
1989	8,068	107	200	312	577	797	1,124	1,299	1,255	2,036	2
1988	7,466	104	191	312	512	832	1,066	1,172	1,078	1,900	-
1987	7,173	105	180	275	574	826	1,068	1,134	997	1,714	1
1986	7,150	108	160	290	586	840	1,004	1,150	1,050	1,633	1
1985	6,871	115	191	347	568	795	1,003	1,073	934	1,527	-
1984	6,602	107	215	346	583	800	917	1,019	855	1,446	3
1983	6,382	101	190	379	581	742	912	960	863	1,334	-
1982	6,047	111	207	351	553	776	925	862	827	1,158	-
1981	5,931	114	197	404	528	755	893	832	743	1,159	-
1980	5,778	149	237	360	530	734	795	807	781	1,064	-
1979	5,452	122	234	366	527	717	734	741	692	983	1
Other diseases of arteries, arterioles and capillaries (442-448)											
All races, male											
1998	4,978	103	125	210	305	479	701	899	866	1,106	-
1997	4,798	79	129	177	339	498	757	877	779	992	-
1996	4,740	105	112	166	319	496	759	844	775	969	-
1995	4,489	79	97	147	311	528	716	810	722	889	-
1994	4,372	77	124	149	304	510	669	803	691	884	-
1993	4,299	73	86	178	304	508	664	727	735	851	-
1992	4,097	62	85	148	304	504	699	744	596	768	1
1991	3,793	73	100	156	289	483	572	670	591	691	-
1990	3,754	62	104	173	290	438	646	691	560	641	-
1989	3,739	58	104	176	348	460	621	615	563	629	1
1988	3,385	47	94	171	304	457	590	603	439	541	-
1987	3,457	55	93	153	340	492	591	598	465	532	-
1986	3,354	50	83	175	342	503	603	576	413	459	1
1985	3,412	53	110	195	345	470	600	564	452	477	-
1984	3,213	54	121	198	355	471	527	513	374	463	2
1983	3,219	54	106	213	376	461	517	508	400	421	-
1982	3,063	57	115	196	349	460	554	449	368	378	-
1981	3,015	55	116	226	323	467	511	462	333	385	-
1980	2,928	73	149	215	335	447	439	440	331	356	-
1979	2,841	56	141	211	324	438	443	392	333	349	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
All races, female											
1998	6,456	95	134	179	243	426	685	936	1,025	2,495	-
1997	6,270	91	109	158	281	431	678	908	1,042	2,345	-
1996	6,049	85	98	133	244	425	668	866	970	2,352	-
1995	5,717	87	110	160	227	419	632	794	953	2,118	1
1994	5,602	85	84	150	254	415	624	791	966	2,023	1
1993	5,231	82	112	143	210	409	597	759	849	1,861	-
1992	5,020	84	94	162	235	367	541	696	869	1,758	-
1991	4,566	60	89	114	214	401	495	650	725	1,616	-
1990	4,489	77	93	126	235	369	495	619	750	1,541	-
1989	4,329	49	96	136	229	337	503	684	692	1,407	1
1988	4,081	57	97	141	208	375	476	569	639	1,359	-
1987	3,716	50	87	122	234	334	477	536	532	1,182	1
1986	3,796	58	77	115	244	337	401	574	637	1,174	-
1985	3,459	62	81	152	223	325	403	509	482	1,050	-
1984	3,389	53	94	148	228	329	390	506	481	983	1
1983	3,163	47	84	166	205	281	395	452	463	913	-
1982	2,984	54	92	155	204	316	371	413	459	780	-
1981	2,916	59	81	178	205	288	382	370	410	774	-
1980	2,850	76	88	145	195	287	356	367	450	708	-
1979	2,611	66	93	155	203	279	291	349	359	634	-
Other diseases of arteries, arterioles and capillaries (442-448)											
White, both sexes											
1998	9,789	137	209	295	444	768	1,179	1,606	1,672	3,205	-
1997	9,510	124	172	249	490	771	1,227	1,582	1,617	3,012	-
1996	9,352	139	158	237	463	780	1,225	1,503	1,567	3,010	-
1995	8,903	132	170	246	446	813	1,164	1,440	1,506	2,709	1
1994	8,565	116	164	232	453	760	1,129	1,389	1,470	2,586	1
1993	8,211	105	145	244	432	781	1,106	1,322	1,405	2,433	-
1992	7,940	105	135	251	455	740	1,088	1,278	1,319	2,278	1
1991	7,256	101	144	220	417	751	939	1,176	1,189	2,080	-
1990	7,172	103	150	248	451	707	995	1,164	1,170	1,954	-
1989	7,029	84	150	240	497	688	987	1,152	1,130	1,849	2
1988	6,544	78	150	264	434	730	946	1,053	958	1,735	-
1987	6,258	85	139	233	493	708	936	1,003	899	1,561	1
1986	6,245	76	125	233	492	729	891	1,032	943	1,503	1
1985	6,037	93	149	294	483	684	887	957	838	1,418	-
1984	5,805	80	174	294	484	716	805	925	776	1,320	3
1983	5,623	85	164	310	507	654	801	862	788	1,225	-
1982	5,323	83	172	294	466	661	822	787	752	1,075	-
1981	5,271	91	163	342	451	675	814	761	682	1,080	-
1980	5,095	118	211	309	453	647	703	727	707	984	-
1979	4,864	102	197	315	480	642	663	675	631	925	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
White, male											
1998	4,276	72	100	160	259	401	620	794	772	978	-
1997	4,142	56	99	132	278	429	659	783	693	888	-
1996	4,117	79	85	132	266	425	661	749	704	880	-
1995	3,969	65	85	114	263	456	644	733	660	816	-
1994	3,783	56	102	117	257	438	587	695	613	806	-
1993	3,708	48	67	140	260	431	596	653	645	757	-
1992	3,586	50	65	129	251	422	621	666	540	701	1
1991	3,326	55	83	130	249	415	513	606	541	614	-
1990	3,276	51	81	144	254	388	560	612	496	577	-
1989	3,281	50	77	136	311	392	555	555	503	573	1
1988	2,975	36	76	145	259	407	524	546	386	501	-
1987	3,008	42	74	133	295	420	519	526	418	485	-
1986	2,933	33	67	144	292	431	552	514	374	421	1
1985	3,020	46	87	171	292	409	538	510	408	441	-
1984	2,847	40	102	174	298	431	462	471	340	421	2
1983	2,847	48	92	180	331	413	456	459	367	378	-
1982	2,707	43	93	164	303	389	497	420	342	344	-
1981	2,703	47	99	198	285	426	472	426	303	353	-
1980	2,574	60	134	186	286	397	390	396	292	323	-
1979	2,535	48	119	186	295	395	400	357	300	324	-
Other diseases of arteries, arterioles and capillaries (442-448)											
White, female											
1998	5,513	65	109	135	185	367	559	812	900	2,227	-
1997	5,368	68	73	117	212	342	568	799	924	2,124	-
1996	5,235	60	73	105	197	355	564	754	863	2,130	-
1995	4,934	67	85	132	183	357	520	707	846	1,893	1
1994	4,782	60	62	115	196	322	542	694	857	1,780	1
1993	4,503	57	78	104	172	350	510	669	760	1,676	-
1992	4,354	55	70	122	204	318	467	612	779	1,577	-
1991	3,930	46	61	90	168	336	426	570	648	1,466	-
1990	3,896	52	69	104	197	319	435	552	674	1,377	-
1989	3,748	34	73	104	186	296	432	597	627	1,276	1
1988	3,569	42	74	119	175	323	422	507	572	1,234	-
1987	3,250	43	65	100	198	288	417	477	481	1,076	1
1986	3,312	43	58	89	200	298	339	518	569	1,082	-
1985	3,017	47	62	123	191	275	349	447	430	977	-
1984	2,958	40	72	120	186	285	343	454	436	899	1
1983	2,776	37	72	130	176	241	345	403	421	847	-
1982	2,616	40	79	130	163	272	325	367	410	731	-
1981	2,568	44	64	144	166	249	342	335	379	727	-
1980	2,521	58	77	123	167	250	313	331	415	661	-
1979	2,329	54	78	129	185	247	263	318	331	601	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
All other, both sexes											
1998	1,645	61	50	94	104	137	207	229	219	396	-
1997	1,558	46	66	86	130	158	208	203	204	325	-
1996	1,437	51	52	62	100	141	202	207	178	311	-
1995	1,303	34	37	61	92	134	184	164	169	298	-
1994	1,409	46	44	67	105	165	164	205	187	321	-
1993	1,319	50	53	77	82	136	155	164	179	279	-
1992	1,177	41	44	59	84	131	152	162	146	248	-
1991	1,103	32	45	50	86	133	128	144	127	227	-
1990	1,071	36	47	51	74	100	146	146	140	228	-
1989	1,039	23	50	72	80	109	137	147	125	187	-
1988	922	26	41	48	78	102	120	119	120	165	-
1987	915	20	41	42	81	118	132	131	98	153	-
1986	905	32	35	57	94	111	113	118	107	130	-
1985	834	22	42	53	85	111	116	116	96	109	-
1984	797	27	41	52	99	84	112	94	79	126	-
1983	759	16	26	69	74	88	111	98	75	109	-
1982	724	28	35	57	87	115	103	75	75	83	-
1981	660	23	34	62	77	80	79	71	61	79	-
1980	683	31	26	51	77	87	92	80	74	80	-
1979	588	20	37	51	47	75	71	66	61	58	1
Other diseases of arteries, arterioles and capillaries (442-448)											
All other, male											
1998	702	31	25	50	46	78	81	105	94	128	-
1997	656	23	30	45	61	69	98	94	86	104	-
1996	623	26	27	34	53	71	98	95	71	89	-
1995	520	14	12	33	48	72	72	77	62	73	-
1994	589	21	22	32	47	72	82	108	78	78	-
1993	591	25	19	38	44	77	68	74	90	94	-
1992	511	12	20	19	53	82	78	78	56	67	-
1991	467	18	17	26	40	68	59	64	50	77	-
1990	478	11	23	29	36	50	86	79	64	64	-
1989	458	8	27	40	37	68	66	60	60	56	-
1988	410	11	18	26	45	50	66	57	53	40	-
1987	449	13	19	20	45	72	72	72	47	47	-
1986	421	17	16	31	50	72	51	62	39	38	-
1985	392	7	23	24	53	61	62	54	44	36	-
1984	366	14	19	24	57	40	65	42	34	42	-
1983	372	6	14	33	45	48	61	49	33	43	-
1982	356	14	22	32	46	71	57	29	26	34	-
1981	312	8	17	28	38	41	39	36	30	32	-
1980	354	13	15	29	49	50	49	44	39	33	-
1979	306	8	22	25	29	43	43	35	33	25	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
All other, female											
1998	943	30	25	44	58	59	126	124	125	268	-
1997	902	23	36	41	69	89	110	109	118	221	-
1996	814	25	25	28	47	70	104	112	107	222	-
1995	783	20	25	28	44	62	112	87	107	225	-
1994	820	25	22	35	58	93	82	97	109	243	-
1993	728	25	34	39	38	59	87	90	89	185	-
1992	666	29	24	40	31	49	74	84	90	181	-
1991	636	14	28	24	46	65	69	80	77	150	-
1990	593	25	24	22	38	50	60	67	76	164	-
1989	581	15	23	32	43	41	71	87	65	131	-
1988	512	15	23	22	33	52	54	62	67	125	-
1987	466	7	22	22	36	46	60	59	51	106	-
1986	484	15	19	26	44	39	62	56	68	92	-
1985	442	15	19	29	32	50	54	62	52	73	-
1984	431	13	22	28	42	44	47	52	45	84	-
1983	387	10	12	36	29	40	50	49	42	66	-
1982	368	14	13	25	41	44	46	46	49	49	-
1981	348	15	17	34	39	39	40	35	31	47	-
1980	329	18	11	22	28	37	43	36	35	47	-
1979	282	12	15	26	18	32	28	31	28	33	-
Other diseases of arteries, arterioles and capillaries (442-448)											
Black, both sexes											
1998	1,482	59	45	89	90	123	183	204	188	363	-
1997	1,403	43	59	71	115	144	185	187	182	295	-
1996	1,315	50	47	57	89	137	181	187	163	289	-
1995	1,196	31	35	54	83	124	164	154	159	272	-
1994	1,276	43	39	55	90	151	152	188	165	302	-
1993	1,194	47	48	69	71	124	143	149	160	259	-
1992	1,085	35	40	53	78	122	142	149	136	228	-
1991	1,033	31	44	44	81	122	120	135	122	215	-
1990	993	32	45	45	71	87	138	137	130	214	-
1989	973	23	45	70	72	100	130	138	118	175	-
1988	855	22	39	45	70	98	111	110	110	155	-
1987	856	20	39	37	78	109	120	122	91	146	-
1986	845	30	33	54	87	107	108	105	100	124	-
1985	778	20	42	49	82	105	107	108	88	101	-
1984	743	26	36	48	95	80	106	89	74	115	-
1983	698	15	23	63	72	84	104	89	67	96	-
1982	681	28	31	52	84	108	98	70	71	79	-
1981	619	21	31	58	72	76	73	68	58	73	-
1980	650	31	24	47	76	83	87	76	69	75	-
1979	564	20	35	49	43	74	71	60	58	57	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of arteries, arterioles and capillaries (442-448)											
Black, male											
1998	632	30	21	47	42	69	76	95	78	115	-
1997	601	20	26	39	54	66	92	86	77	96	-
1996	570	25	24	32	49	69	90	84	64	79	-
1995	468	12	11	31	43	68	61	71	55	65	-
1994	532	20	19	28	42	65	77	101	66	74	-
1993	535	24	18	37	39	72	62	66	78	85	-
1992	468	10	18	18	49	76	72	72	51	60	-
1991	430	17	17	22	39	60	54	58	47	72	-
1990	447	10	21	27	35	45	81	74	59	62	-
1989	417	8	24	38	34	62	64	53	53	48	-
1988	374	11	17	24	39	50	59	53	43	38	-
1987	411	13	17	19	44	65	67	65	41	41	-
1986	392	16	15	28	49	69	48	55	35	37	-
1985	365	7	23	23	51	58	56	49	37	35	-
1984	344	13	18	22	56	39	63	41	30	36	-
1983	344	6	13	31	44	45	56	43	28	38	-
1982	329	14	20	29	45	65	52	27	24	32	-
1981	291	7	15	27	36	39	35	35	29	29	-
1980	336	13	15	26	49	46	46	40	36	32	-
1979	290	8	20	24	26	42	43	31	31	25	1
Other diseases of arteries, arterioles and capillaries (442-448)											
Black, female											
1998	850	29	24	42	48	54	107	109	110	248	-
1997	802	23	33	32	61	78	93	101	105	199	-
1996	745	25	23	25	40	68	91	103	99	210	-
1995	728	19	24	23	40	56	103	83	104	207	-
1994	744	23	20	27	48	86	75	87	99	228	-
1993	659	23	30	32	32	52	81	83	82	174	-
1992	617	25	22	35	29	46	70	77	85	168	-
1991	603	14	27	22	42	62	66	77	75	143	-
1990	546	22	24	18	36	42	57	63	71	152	-
1989	556	15	21	32	38	38	66	85	65	127	-
1988	481	11	22	21	31	48	52	57	67	117	-
1987	445	7	22	18	34	44	53	57	50	105	-
1986	453	14	18	26	38	38	60	50	65	87	-
1985	413	13	19	26	31	47	51	59	51	66	-
1984	399	13	18	26	39	41	43	48	44	79	-
1983	354	9	10	32	28	39	48	46	39	58	-
1982	352	14	11	23	39	43	46	43	47	47	-
1981	328	14	16	31	36	37	38	33	29	44	-
1980	314	18	9	21	27	37	41	36	33	43	-
1979	274	12	15	25	17	32	28	29	27	32	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
All races, both sexes											
1998	4,397	170	183	226	265	355	486	609	612	1,066	-
1997	4,264	171	198	227	284	358	483	586	588	923	1
1996	4,243	187	194	236	276	396	459	575	573	897	-
1995	4,185	173	184	206	293	405	482	557	559	879	2
1994	4,082	184	161	188	243	397	501	528	590	868	1
1993	4,122	153	144	210	300	387	495	530	578	866	-
1992	4,000	143	163	182	275	407	488	541	531	822	-
1991	4,140	137	162	218	339	420	518	559	521	807	2
1990	4,238	127	176	229	349	477	497	575	578	806	3
1989	4,319	134	173	250	372	475	564	618	545	801	-
1988	4,569	173	184	253	412	526	559	618	578	857	1
1987	4,823	146	199	295	459	569	602	701	611	798	-
1986	4,677	172	216	324	474	528	596	600	568	765	2
1985	4,860	177	239	320	474	556	663	630	593	741	1
1984	4,753	161	215	361	469	561	601	651	552	742	1
1983	5,034	155	263	369	508	572	647	683	596	789	-
1982	4,684	162	207	345	484	568	593	589	571	733	2
1981	4,702	157	269	405	466	595	616	612	531	624	3
1980	4,803	178	269	379	488	573	583	643	513	697	2
1979	4,755	174	301	378	446	587	618	593	534	657	1
Other disorders of circulatory system (451-459)											
All races, male											
1998	1,943	104	102	127	147	175	261	259	240	296	-
1997	1,978	105	122	129	155	191	242	283	225	279	-
1996	1,913	111	100	132	154	213	239	245	216	258	-
1995	1,929	101	108	120	150	228	249	249	234	249	1
1994	1,867	112	90	102	121	200	266	238	226	265	1
1993	1,940	82	86	127	171	230	243	241	240	262	-
1992	1,926	88	83	99	169	214	246	266	232	250	-
1991	1,984	75	96	138	198	232	280	262	230	223	2
1990	2,026	75	96	136	198	281	256	262	230	235	2
1989	2,062	81	112	149	222	271	291	274	203	225	-
1988	2,262	104	115	164	236	293	294	312	234	265	-
1987	2,369	92	118	178	295	305	301	318	251	248	-
1986	2,377	106	122	194	276	322	310	295	226	277	-
1985	2,398	99	144	199	285	309	341	273	233	230	-
1984	2,442	96	148	223	282	325	346	287	233	262	-
1983	2,506	100	157	222	305	334	319	324	240	251	-
1982	2,385	110	120	218	295	321	320	276	240	241	1
1981	2,386	86	165	244	282	330	321	286	219	214	1
1980	2,405	105	161	241	289	320	299	300	201	239	1
1979	2,442	100	186	244	280	322	326	271	211	236	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
All races, female											
1998	2,454	66	81	99	118	180	225	350	372	770	-
1997	2,286	66	76	98	129	167	241	303	363	644	1
1996	2,330	76	94	104	122	183	220	330	357	639	-
1995	2,256	72	76	86	143	177	233	308	325	630	1
1994	2,215	72	71	86	122	197	235	290	364	603	-
1993	2,182	71	58	83	129	157	252	289	338	604	-
1992	2,074	55	80	83	106	193	242	275	299	572	-
1991	2,156	62	66	80	141	188	238	297	291	584	-
1990	2,212	52	80	93	151	196	241	313	348	571	1
1989	2,257	53	61	101	150	204	273	344	342	576	-
1988	2,307	69	69	89	176	233	265	306	344	592	1
1987	2,454	54	81	117	164	264	301	383	360	550	-
1986	2,300	66	94	130	198	206	286	305	342	488	2
1985	2,462	78	95	121	189	247	322	357	360	511	1
1984	2,311	65	67	138	187	236	255	364	319	480	1
1983	2,528	55	106	147	203	238	328	359	356	538	-
1982	2,299	52	87	127	189	247	273	313	331	492	1
1981	2,316	71	104	161	184	265	295	326	312	410	2
1980	2,398	73	108	138	199	253	284	343	312	458	1
1979	2,313	74	115	134	166	265	292	322	323	421	-
Other disorders of circulatory system (451-459)											
White, both sexes											
1998	3,585	116	138	170	204	287	412	521	534	949	-
1997	3,509	112	158	177	220	284	410	504	517	830	-
1996	3,553	147	150	188	217	328	387	513	512	815	-
1995	3,444	125	140	160	219	322	409	486	488	798	2
1994	3,356	125	120	143	187	327	433	452	517	779	-
1993	3,452	112	105	162	235	331	432	454	520	800	-
1992	3,315	97	118	134	213	346	426	479	473	739	-
1991	3,434	100	116	164	281	354	447	468	459	730	1
1990	3,511	89	123	177	273	386	434	502	519	725	3
1989	3,574	85	119	196	307	386	471	540	489	719	-
1988	3,787	118	132	203	335	425	481	539	523	774	-
1987	3,981	103	143	227	365	474	521	596	557	712	-
1986	3,912	129	158	257	379	463	527	521	505	695	1
1985	4,044	143	182	247	391	461	563	553	523	668	-
1984	4,015	120	164	295	398	491	519	573	493	674	-
1983	4,243	127	201	287	414	493	554	596	530	728	-
1982	3,929	115	159	288	385	484	505	514	520	669	2
1981	3,998	121	200	329	393	525	525	541	490	583	3
1980	4,078	135	204	306	418	500	509	566	460	643	1
1979	4,035	139	245	300	379	478	550	516	474	621	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
White, male											
1998	1,599	73	79	103	113	146	225	235	208	266	-
1997	1,608	71	95	99	124	150	208	248	199	250	-
1996	1,584	86	74	106	123	180	204	219	189	230	-
1995	1,565	75	83	94	107	185	213	219	204	217	1
1994	1,506	81	67	81	94	163	228	199	195	236	-
1993	1,588	58	63	99	130	197	211	206	218	245	-
1992	1,589	62	60	72	135	190	216	239	205	220	-
1991	1,652	53	71	105	168	198	251	219	206	206	1
1990	1,644	54	61	103	158	224	220	233	208	206	2
1989	1,693	52	76	115	184	218	250	242	185	201	-
1988	1,852	71	82	136	190	234	252	273	212	241	-
1987	1,938	65	81	133	242	251	268	281	224	220	-
1986	1,966	83	91	150	226	279	276	251	194	252	-
1985	1,983	86	112	158	230	260	298	242	199	201	-
1984	2,055	69	114	186	248	292	298	248	208	237	-
1983	2,076	80	121	174	246	291	272	285	211	226	-
1982	1,992	79	96	182	230	283	271	244	214	223	1
1981	2,029	65	123	199	243	292	286	250	200	201	1
1980	2,017	80	129	194	241	285	262	260	179	211	1
1979	2,044	73	152	200	240	262	286	233	189	219	-
Other disorders of circulatory system (451-459)											
White, female											
1998	1,986	43	59	67	91	141	187	286	326	683	-
1997	1,901	41	63	78	96	134	202	256	318	580	-
1996	1,969	61	76	82	94	148	183	294	323	585	-
1995	1,879	50	57	66	112	137	196	267	284	581	1
1994	1,850	44	53	62	93	164	205	253	322	543	-
1993	1,864	54	42	63	105	134	221	248	302	555	-
1992	1,726	35	58	62	78	156	210	240	268	519	-
1991	1,782	47	45	59	113	156	196	249	253	524	-
1990	1,867	35	62	74	115	162	214	269	311	519	1
1989	1,881	33	43	81	123	168	221	298	304	518	-
1988	1,935	47	50	67	145	191	229	266	311	533	-
1987	2,043	38	62	94	123	223	253	315	333	492	-
1986	1,946	46	67	107	153	184	251	270	311	443	1
1985	2,061	57	70	89	161	201	265	311	324	467	-
1984	1,960	51	50	109	150	199	221	325	285	437	-
1983	2,167	47	80	113	168	202	282	311	319	502	-
1982	1,937	36	63	106	155	201	234	270	306	446	1
1981	1,969	56	77	130	150	233	239	291	290	382	2
1980	2,061	55	75	112	177	215	247	306	281	432	-
1979	1,991	66	93	100	139	216	264	283	285	402	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
All other, both sexes											
1998	812	54	45	56	61	68	74	88	78	117	-
1997	755	59	40	50	64	74	73	82	71	93	1
1996	690	40	44	48	59	68	72	62	61	82	-
1995	741	48	44	46	74	83	73	71	71	81	-
1994	726	59	41	45	56	70	68	76	73	89	1
1993	670	41	39	48	65	56	63	76	58	66	-
1992	685	46	45	48	62	61	62	62	58	83	-
1991	706	37	46	54	58	66	71	91	62	77	1
1990	727	38	53	52	76	91	63	73	59	81	-
1989	745	49	54	54	65	89	93	78	56	82	-
1988	782	55	52	50	77	101	78	79	55	83	1
1987	842	43	56	68	94	95	81	105	54	86	-
1986	765	43	58	67	95	65	69	79	63	70	1
1985	816	34	57	73	83	95	100	77	70	73	1
1984	738	41	51	66	71	70	82	78	59	68	1
1983	791	28	62	82	94	79	93	87	66	61	-
1982	755	47	48	57	99	84	88	75	51	64	-
1981	704	36	69	76	73	70	91	71	41	41	-
1980	725	43	65	73	70	73	74	77	53	54	1
1979	720	35	56	78	67	109	68	77	60	36	1
Other disorders of circulatory system (451-459)											
All other, male											
1998	344	31	23	24	34	29	36	24	32	30	-
1997	370	34	27	30	31	41	34	35	26	29	-
1996	329	25	26	26	31	33	35	26	27	28	-
1995	364	26	25	26	43	43	36	30	30	32	-
1994	361	31	23	21	27	37	38	39	31	29	1
1993	352	24	23	28	41	33	32	35	22	17	-
1992	337	26	23	27	34	24	30	27	27	30	-
1991	332	22	25	33	30	34	29	43	24	17	1
1990	382	21	35	33	40	57	36	29	22	29	-
1989	369	29	36	34	38	53	41	32	18	24	-
1988	410	33	33	28	46	59	42	39	22	24	-
1987	431	27	37	45	53	54	33	37	27	28	-
1986	411	23	31	44	50	43	34	44	32	25	-
1985	415	13	32	41	55	49	43	31	34	29	-
1984	387	27	34	37	34	33	48	39	25	25	-
1983	430	20	36	48	59	43	47	39	29	25	-
1982	393	31	24	36	65	38	49	32	26	18	-
1981	357	21	42	45	39	38	35	36	19	13	-
1980	388	25	32	47	48	35	37	40	22	28	-
1979	398	27	34	44	40	60	40	38	22	17	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
All other, female											
1998	468	23	22	32	27	39	38	64	46	87	-
1997	385	25	13	20	33	33	39	47	45	64	1
1996	361	15	18	22	28	35	37	36	34	54	-
1995	377	22	19	20	31	40	37	41	41	49	-
1994	365	28	18	24	29	33	30	37	42	60	-
1993	318	17	16	20	24	23	31	41	36	49	-
1992	348	20	22	21	28	37	32	35	31	53	-
1991	374	15	21	21	28	32	42	48	38	60	-
1990	345	17	18	19	36	34	27	44	37	52	-
1989	376	20	18	20	27	36	52	46	38	58	-
1988	372	22	19	22	31	42	36	40	33	59	1
1987	411	16	19	23	41	41	48	68	27	58	-
1986	354	20	27	23	45	22	35	35	31	45	1
1985	401	21	25	32	28	46	57	46	36	44	1
1984	351	14	17	29	37	37	34	39	34	43	1
1983	361	8	26	34	35	36	46	48	37	36	-
1982	362	16	24	21	34	46	39	43	25	46	-
1981	347	15	27	31	34	32	56	35	22	28	-
1980	337	18	33	26	22	38	37	37	31	26	1
1979	322	8	22	34	27	49	28	39	38	19	-
Other disorders of circulatory system (451-459)											
Black, both sexes											
1998	745	51	43	50	54	63	68	81	73	100	-
1997	695	55	38	48	59	61	66	76	65	86	1
1996	623	38	40	45	51	63	65	53	46	74	-
1995	672	42	40	41	66	72	69	62	63	77	-
1994	663	56	38	40	51	67	63	69	63	80	1
1993	604	38	38	46	59	49	59	66	51	55	-
1992	630	42	39	43	56	56	58	57	55	78	-
1991	657	34	42	51	54	63	70	81	59	69	1
1990	666	34	48	46	73	83	56	69	51	75	-
1989	695	40	50	47	58	85	91	74	53	79	-
1988	735	52	49	46	74	95	73	70	53	75	1
1987	798	39	53	62	89	93	81	96	51	83	-
1986	719	40	54	66	90	60	63	75	62	63	1
1985	757	33	52	65	76	89	92	73	64	66	1
1984	690	37	51	60	70	62	77	71	55	62	1
1983	742	27	58	76	90	72	89	81	62	57	-
1982	703	43	41	55	96	80	82	68	48	58	-
1981	668	33	67	74	69	70	84	69	41	38	-
1980	687	40	61	70	69	65	68	73	51	53	1
1979	682	32	53	75	63	104	61	74	58	33	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of circulatory system (451-459)											
Black, male											
1998	317	30	22	24	28	28	34	23	29	24	-
1997	336	31	25	28	29	31	29	32	24	27	-
1996	293	23	24	25	25	30	31	24	19	24	-
1995	331	25	24	22	41	39	34	24	27	30	-
1994	325	30	22	17	24	36	36	35	27	23	1
1993	314	22	22	27	36	26	29	31	18	14	-
1992	298	23	20	23	29	22	28	22	25	26	-
1991	309	21	23	32	27	34	29	39	22	14	1
1990	348	18	33	28	37	54	32	27	19	28	-
1989	339	22	34	30	33	50	40	28	17	24	-
1988	384	31	31	26	43	57	39	32	21	22	-
1987	404	25	35	39	50	52	33	32	27	26	-
1986	388	22	28	43	48	39	31	41	31	23	-
1985	388	12	30	38	52	45	41	29	31	25	-
1984	358	24	34	35	33	29	44	35	21	22	-
1983	400	19	33	45	56	38	44	35	26	24	-
1982	369	30	20	35	63	37	46	29	24	16	-
1981	338	18	42	44	37	38	31	34	19	12	-
1980	369	24	30	45	47	31	35	37	21	27	-
1979	368	25	31	42	37	57	33	35	21	14	1
Other disorders of circulatory system (451-459)											
Black, female											
1998	428	21	21	26	26	35	34	58	44	76	-
1997	359	24	13	20	30	30	37	44	41	59	1
1996	330	15	16	20	26	33	34	29	27	50	-
1995	341	17	16	19	25	33	35	38	36	47	-
1994	338	26	16	23	27	31	27	34	36	57	-
1993	290	16	16	19	23	23	30	35	33	41	-
1992	332	19	19	20	27	34	30	35	30	52	-
1991	348	13	19	19	27	29	41	42	37	55	-
1990	318	16	15	18	36	29	24	42	32	47	-
1989	356	18	16	17	25	35	51	46	36	55	-
1988	351	21	18	20	31	38	34	38	32	53	1
1987	394	14	18	23	39	41	48	64	24	57	-
1986	331	18	26	23	42	21	32	34	31	40	1
1985	369	21	22	27	24	44	51	44	33	41	1
1984	332	13	17	25	37	33	33	36	34	40	1
1983	342	8	25	31	34	34	45	46	36	33	-
1982	334	13	21	20	33	43	36	39	24	42	-
1981	330	15	25	30	32	32	53	35	22	26	-
1980	318	16	31	25	22	34	33	36	30	26	1
1979	314	7	22	33	26	47	28	39	37	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
All races, both sexes											
1998	91,871	980	1,187	1,552	2,304	3,816	7,189	11,717	17,140	43,127	5
1997	86,449	1,025	1,208	1,482	2,277	3,764	6,771	11,207	16,151	39,668	6
1996	83,727	1,029	1,064	1,464	2,149	3,699	6,898	10,677	15,678	38,027	10
1995	82,923	985	1,094	1,314	2,144	3,788	6,949	10,723	15,262	37,575	4
1994	81,473	943	1,042	1,297	2,208	3,922	6,680	10,626	15,134	36,400	5
1993	82,820	861	1,018	1,348	2,356	4,072	6,751	10,721	15,138	37,171	11
1992	75,719	797	848	1,322	2,131	3,868	6,344	9,931	14,050	33,296	7
1991	77,860	787	951	1,350	2,388	3,825	6,398	10,195	14,400	34,144	14
1990	79,513	788	963	1,427	2,495	4,102	6,563	10,610	14,767	34,443	7
1989	76,550	782	925	1,422	2,458	4,102	6,316	10,175	13,847	32,955	15
1988	77,662	807	964	1,512	2,612	4,238	6,475	10,346	14,135	33,149	11
1987	69,225	701	925	1,380	2,499	3,932	6,094	9,355	12,422	28,739	7
1986	69,812	705	895	1,513	2,631	3,957	6,197	9,735	12,260	28,652	46
1985	67,615	657	966	1,550	2,580	3,715	6,113	9,230	12,066	27,705	51
1984	58,894	599	890	1,428	2,323	3,362	5,616	8,141	10,454	23,176	14
1983	55,854	577	921	1,455	2,288	3,352	5,086	7,415	9,878	22,073	15
1982	48,886	568	893	1,475	2,141	3,022	4,655	6,584	8,543	18,225	16
1981	53,731	670	1,007	1,688	2,345	3,398	4,951	7,272	9,479	19,935	14
1980	54,619	694	1,066	1,698	2,346	3,592	5,076	7,266	9,737	19,841	15
1979	45,030	622	1,012	1,459	2,052	2,994	4,343	5,990	8,005	15,266	10
Influenza and pneumonia (480-487)											
All races, male											
1998	40,979	599	703	941	1,346	2,227	4,149	6,294	8,248	14,872	4
1997	39,284	647	728	856	1,338	2,261	3,886	6,091	7,869	13,904	4
1996	37,991	652	650	911	1,253	2,207	3,958	5,789	7,571	13,196	6
1995	37,787	652	666	775	1,307	2,249	4,009	5,802	7,413	13,084	2
1994	37,339	601	653	788	1,327	2,329	3,858	5,760	7,186	12,885	1
1993	37,996	539	621	791	1,437	2,455	3,974	5,901	7,331	12,903	8
1992	35,465	530	529	825	1,281	2,318	3,704	5,505	6,937	11,910	4
1991	36,214	503	563	853	1,467	2,237	3,705	5,599	7,111	12,058	10
1990	36,898	494	620	885	1,562	2,486	3,873	5,782	7,157	11,958	3
1989	35,701	503	588	920	1,540	2,514	3,845	5,499	6,581	11,533	11
1988	36,834	546	626	934	1,653	2,675	3,875	5,785	6,846	11,783	6
1987	33,562	458	598	870	1,588	2,499	3,707	5,210	6,078	10,567	5
1986	34,203	436	582	956	1,709	2,485	3,762	5,506	6,082	10,697	16
1985	33,159	427	611	1,002	1,682	2,395	3,726	5,200	5,931	10,352	23
1984	29,440	366	596	939	1,548	2,124	3,458	4,633	5,182	8,849	11
1983	28,007	368	591	955	1,512	2,167	3,133	4,165	4,929	8,511	9
1982	25,325	391	565	971	1,374	1,953	2,870	3,835	4,351	7,363	5
1981	27,059	417	630	1,102	1,528	2,176	2,960	4,058	4,711	7,699	12
1980	27,574	466	698	1,070	1,533	2,280	3,066	3,996	4,747	7,806	9
1979	23,725	396	683	976	1,320	1,976	2,748	3,353	4,049	6,307	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
All races, female											
1998	50,892	381	484	611	958	1,589	3,040	5,423	8,892	28,255	1
1997	47,165	378	480	626	939	1,503	2,885	5,116	8,282	25,764	2
1996	45,736	377	414	553	896	1,492	2,940	4,888	8,107	24,831	4
1995	45,136	333	428	539	837	1,539	2,940	4,921	7,849	24,491	2
1994	44,134	342	389	509	881	1,593	2,822	4,866	7,948	23,515	4
1993	44,824	322	397	557	919	1,617	2,777	4,820	7,807	24,268	3
1992	40,254	267	319	497	850	1,550	2,640	4,426	7,113	21,386	3
1991	41,646	284	388	497	921	1,588	2,693	4,596	7,289	22,086	4
1990	42,615	294	343	542	933	1,616	2,690	4,828	7,610	22,485	4
1989	40,849	279	337	502	918	1,588	2,471	4,676	7,266	21,422	4
1988	40,828	261	338	578	959	1,563	2,600	4,561	7,289	21,366	5
1987	35,663	243	327	510	911	1,433	2,387	4,145	6,344	18,172	2
1986	35,609	269	313	557	922	1,472	2,435	4,229	6,178	17,955	30
1985	34,456	230	355	548	898	1,320	2,387	4,030	6,135	17,353	28
1984	29,454	233	294	489	775	1,238	2,158	3,508	5,272	14,327	3
1983	27,847	209	330	500	776	1,185	1,953	3,250	4,949	13,562	6
1982	23,561	177	328	504	767	1,069	1,785	2,749	4,192	10,862	11
1981	26,672	253	377	586	817	1,222	1,991	3,214	4,768	12,236	2
1980	27,045	228	368	628	813	1,312	2,010	3,270	4,990	12,035	6
1979	21,305	226	329	483	732	1,018	1,595	2,637	3,956	8,959	-
Influenza and pneumonia (480-487)											
White, both sexes											
1998	81,659	685	885	1,207	1,827	3,164	6,198	10,364	15,562	39,915	1
1997	76,875	674	917	1,131	1,795	3,127	5,844	9,974	14,707	36,766	2
1996	74,194	710	779	1,104	1,674	3,067	5,922	9,554	14,207	35,202	7
1995	73,641	667	801	975	1,705	3,149	5,998	9,583	13,870	34,858	4
1994	72,605	665	760	983	1,792	3,255	5,845	9,544	13,840	33,855	3
1993	73,720	573	781	1,022	1,879	3,422	5,805	9,557	13,821	34,657	4
1992	67,456	568	611	1,011	1,694	3,220	5,512	8,976	12,845	31,048	4
1991	69,276	534	675	1,023	1,937	3,213	5,586	9,124	13,158	31,854	11
1990	70,806	554	687	1,097	1,976	3,419	5,737	9,505	13,538	32,233	5
1989	67,853	516	642	1,059	1,982	3,435	5,459	9,038	12,725	30,815	12
1988	69,570	493	688	1,131	2,102	3,580	5,669	9,311	13,122	31,310	10
1987	61,811	455	657	1,065	2,011	3,360	5,380	8,456	11,433	27,021	2
1986	62,323	472	646	1,176	2,130	3,352	5,445	8,780	11,281	26,943	37
1985	60,508	438	700	1,201	2,085	3,151	5,381	8,319	11,099	26,181	40
1984	52,514	400	615	1,111	1,894	2,838	4,917	7,334	9,641	21,902	11
1983	49,996	390	677	1,124	1,867	2,857	4,460	6,709	9,153	20,947	7
1982	43,515	342	665	1,135	1,740	2,600	4,051	6,024	7,927	17,272	12
1981	47,919	439	718	1,292	1,903	2,863	4,366	6,634	8,807	18,937	7
1980	48,369	447	765	1,291	1,931	2,990	4,464	6,490	9,093	18,751	13
1979	39,803	424	722	1,132	1,673	2,525	3,821	5,403	7,508	14,472	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
White, male											
1998	35,795	416	512	725	1,053	1,820	3,554	5,543	7,478	13,659	1
1997	34,386	419	539	638	1,049	1,877	3,376	5,404	7,156	12,773	2
1996	32,924	431	472	681	952	1,830	3,369	5,140	6,830	12,062	5
1995	32,948	437	473	561	1,040	1,849	3,461	5,187	6,718	12,006	2
1994	32,658	416	465	578	1,075	1,947	3,374	5,137	6,525	11,874	1
1993	33,151	349	460	585	1,144	2,037	3,437	5,231	6,658	11,907	3
1992	30,951	370	367	598	1,010	1,935	3,197	4,953	6,305	11,008	2
1991	31,589	336	395	642	1,180	1,871	3,234	4,994	6,454	11,116	7
1990	32,101	329	431	674	1,245	2,058	3,366	5,172	6,487	11,070	1
1989	30,892	317	390	680	1,227	2,084	3,296	4,876	6,012	10,668	9
1988	32,262	319	428	682	1,318	2,259	3,398	5,176	6,323	11,018	5
1987	29,284	290	417	653	1,267	2,139	3,243	4,667	5,557	9,823	-
1986	29,891	284	419	730	1,389	2,100	3,290	4,943	5,559	9,903	10
1985	29,028	271	435	766	1,357	2,024	3,248	4,677	5,418	9,677	15
1984	25,677	225	394	722	1,242	1,787	3,016	4,134	4,759	8,276	9
1983	24,463	241	427	730	1,220	1,837	2,725	3,738	4,505	8,002	4
1982	22,100	223	412	751	1,099	1,673	2,473	3,507	4,018	6,909	3
1981	23,531	258	421	826	1,228	1,820	2,579	3,690	4,342	7,224	5
1980	23,810	288	490	806	1,231	1,875	2,693	3,541	4,397	7,280	8
1979	20,544	256	479	751	1,062	1,661	2,428	2,992	3,765	5,917	7
Influenza and pneumonia (480-487)											
White, female											
1998	45,864	269	373	482	774	1,344	2,644	4,821	8,084	26,256	-
1997	42,489	255	378	493	746	1,250	2,468	4,570	7,551	23,993	-
1996	41,270	279	307	423	722	1,237	2,553	4,414	7,377	23,140	2
1995	40,693	230	328	414	665	1,300	2,537	4,396	7,152	22,852	2
1994	39,947	249	295	405	717	1,308	2,471	4,407	7,315	21,981	2
1993	40,569	224	321	437	735	1,385	2,368	4,326	7,163	22,750	1
1992	36,505	198	244	413	684	1,285	2,315	4,023	6,540	20,040	2
1991	37,687	198	280	381	757	1,342	2,352	4,130	6,704	20,738	4
1990	38,705	225	256	423	731	1,361	2,371	4,333	7,051	21,163	4
1989	36,961	199	252	379	755	1,351	2,163	4,162	6,713	20,147	3
1988	37,308	174	260	449	784	1,321	2,271	4,135	6,799	20,292	5
1987	32,527	165	240	412	744	1,221	2,137	3,789	5,876	17,198	2
1986	32,432	188	227	446	741	1,252	2,155	3,837	5,722	17,040	27
1985	31,480	167	265	435	728	1,127	2,133	3,642	5,681	16,504	25
1984	26,837	175	221	389	652	1,051	1,901	3,200	4,882	13,626	2
1983	25,533	149	250	394	647	1,020	1,735	2,971	4,648	12,945	3
1982	21,415	119	253	384	641	927	1,578	2,517	3,909	10,363	9
1981	24,388	181	297	466	675	1,043	1,787	2,944	4,465	11,713	2
1980	24,559	159	275	485	700	1,115	1,771	2,949	4,696	11,471	5
1979	19,259	168	243	381	611	864	1,393	2,411	3,743	8,555	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
All other, both sexes											
1998	10,212	295	302	345	477	652	991	1,353	1,578	3,212	4
1997	9,574	351	291	351	482	637	927	1,233	1,444	2,902	4
1996	9,533	319	285	360	475	632	976	1,123	1,471	2,825	3
1995	9,282	318	293	339	439	639	951	1,140	1,392	2,717	-
1994	8,868	278	282	314	416	667	835	1,082	1,294	2,545	2
1993	9,100	288	237	326	477	650	946	1,164	1,317	2,514	7
1992	8,263	229	237	311	437	648	832	955	1,205	2,248	3
1991	8,584	253	276	327	451	612	812	1,071	1,242	2,290	3
1990	8,707	234	276	330	519	683	826	1,105	1,229	2,210	2
1989	8,697	266	283	363	476	667	857	1,137	1,122	2,140	3
1988	8,092	314	276	381	510	658	806	1,035	1,013	1,839	1
1987	7,414	246	268	315	488	572	714	899	989	1,718	5
1986	7,489	233	249	337	501	605	752	955	979	1,709	9
1985	7,107	219	266	349	495	564	732	911	967	1,524	11
1984	6,380	199	275	317	429	524	699	807	813	1,274	3
1983	5,858	187	244	331	421	495	626	706	725	1,126	8
1982	5,371	226	228	340	401	422	604	560	616	953	4
1981	5,812	231	289	396	442	535	585	638	672	998	7
1980	6,250	247	301	407	415	602	612	776	644	1,090	2
1979	5,227	198	290	327	379	469	522	587	497	794	3
Influenza and pneumonia (480-487)											
All other, male											
1998	5,184	183	191	216	293	407	595	751	770	1,213	3
1997	4,898	228	189	218	289	384	510	687	713	1,131	2
1996	5,067	221	178	230	301	377	589	649	741	1,134	1
1995	4,839	215	193	214	267	400	548	615	695	1,078	-
1994	4,681	185	188	210	252	382	484	623	661	1,011	-
1993	4,845	190	161	206	293	418	537	670	673	996	5
1992	4,514	160	162	227	271	383	507	552	632	902	2
1991	4,625	167	168	211	287	366	471	605	657	942	3
1990	4,797	165	189	211	317	428	507	610	670	888	2
1989	4,809	186	198	240	313	430	549	623	569	865	2
1988	4,572	227	198	252	335	416	477	609	523	765	1
1987	4,278	168	181	217	321	360	464	543	521	744	5
1986	4,312	152	163	226	320	385	472	563	523	794	6
1985	4,131	156	176	236	325	371	478	523	513	675	8
1984	3,763	141	202	217	306	337	442	499	423	573	2
1983	3,544	127	164	225	292	330	408	427	424	509	5
1982	3,225	168	153	220	275	280	397	328	333	454	2
1981	3,528	159	209	276	300	356	381	368	369	475	7
1980	3,764	178	208	264	302	405	373	455	350	526	1
1979	3,181	140	204	225	258	315	320	361	284	390	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
All other, female											
1998	5,028	112	111	129	184	245	396	602	808	1,999	1
1997	4,676	123	102	133	193	253	417	546	731	1,771	2
1996	4,466	98	107	130	174	255	387	474	730	1,691	2
1995	4,443	103	100	125	172	239	403	525	697	1,639	-
1994	4,187	93	94	104	164	285	351	459	633	1,534	2
1993	4,255	98	76	120	184	232	409	494	644	1,518	2
1992	3,749	69	75	84	166	265	325	403	573	1,346	1
1991	3,959	86	108	116	164	246	341	466	585	1,348	-
1990	3,910	69	87	119	202	255	319	495	559	1,322	-
1989	3,888	80	85	123	163	237	308	514	553	1,275	1
1988	3,520	87	78	129	175	242	329	426	490	1,074	-
1987	3,136	78	87	98	167	212	250	356	468	974	-
1986	3,177	81	86	111	181	220	280	392	456	915	3
1985	2,976	63	90	113	170	193	254	388	454	849	3
1984	2,617	58	73	100	123	187	257	308	390	701	1
1983	2,314	60	80	106	129	165	218	279	301	617	3
1982	2,146	58	75	120	126	142	207	232	283	499	2
1981	2,284	72	80	120	142	179	204	270	303	523	-
1980	2,486	69	93	143	113	197	239	321	294	564	1
1979	2,046	58	86	102	121	154	202	226	213	404	-
Influenza and pneumonia (480-487)											
Black, both sexes											
1998	8,326	267	271	300	428	549	804	1,085	1,260	2,483	4
1997	7,920	318	245	307	414	531	771	1,010	1,144	2,316	4
1996	7,963	289	255	329	414	554	802	914	1,170	2,279	3
1995	7,803	296	254	309	385	550	812	937	1,145	2,161	-
1994	7,472	264	260	278	371	568	702	911	1,042	2,047	2
1993	7,724	268	220	295	430	564	788	964	1,086	2,043	7
1992	7,074	203	219	275	383	572	709	807	1,017	1,833	3
1991	7,372	232	257	288	404	533	712	910	1,033	1,840	3
1990	7,563	215	255	303	469	616	744	935	1,036	1,802	2
1989	7,585	245	257	329	426	600	760	976	941	1,758	3
1988	7,191	295	253	357	457	595	707	906	879	1,558	1
1987	6,565	227	246	284	449	537	633	798	854	1,419	5
1986	6,700	217	235	318	472	562	683	837	852	1,433	6
1985	6,338	205	254	326	466	515	652	820	824	1,274	10
1984	5,669	186	259	300	386	477	629	712	697	1,054	3
1983	5,234	182	231	318	382	451	563	628	626	935	7
1982	4,799	216	215	318	374	382	546	488	542	786	4
1981	5,276	221	275	381	414	498	531	569	588	844	7
1980	5,648	238	275	390	395	561	557	667	570	920	2
1979	4,740	194	272	309	355	426	484	518	425	678	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza and pneumonia (480-487)											
Black, male											
1998	4,178	167	172	192	267	343	483	599	595	863	3
1997	3,978	207	162	194	249	320	416	562	530	849	2
1996	4,170	202	161	209	260	328	480	521	566	862	1
1995	4,019	202	170	198	234	347	459	495	550	808	-
1994	3,899	175	173	186	224	331	401	516	510	775	-
1993	4,051	179	152	188	268	359	439	542	538	750	5
1992	3,822	142	149	203	240	347	427	452	519	689	2
1991	3,938	156	158	192	259	317	404	506	516	730	3
1990	4,161	154	175	194	290	394	451	506	545	699	2
1989	4,168	170	183	222	288	394	485	514	464	665	2
1988	4,047	216	185	235	303	379	413	524	442	617	1
1987	3,795	156	169	196	299	337	408	473	441	609	5
1986	3,836	143	153	215	301	363	430	480	440	648	5
1985	3,664	146	171	219	304	338	425	457	416	555	8
1984	3,336	135	190	207	273	305	395	429	353	468	2
1983	3,190	125	159	217	268	300	366	376	356	430	4
1982	2,884	165	145	208	255	253	358	280	290	360	2
1981	3,201	152	201	269	283	331	341	327	315	391	7
1980	3,386	175	194	252	287	375	339	382	300	433	1
1979	2,884	138	190	213	240	286	290	312	244	331	2
Influenza and pneumonia (480-487)											
Black, female											
1998	4,148	100	99	108	161	206	321	486	665	1,620	1
1997	3,942	111	83	113	165	211	355	448	614	1,467	2
1996	3,793	87	94	120	154	226	322	393	604	1,417	2
1995	3,784	94	84	111	151	203	353	442	595	1,353	-
1994	3,573	89	87	92	147	237	301	395	532	1,272	2
1993	3,673	89	68	107	162	205	349	422	548	1,293	2
1992	3,252	61	70	72	143	225	282	355	498	1,144	1
1991	3,434	76	99	96	145	216	308	404	517	1,110	-
1990	3,402	61	80	109	179	222	293	429	491	1,103	-
1989	3,417	75	74	107	138	206	275	462	477	1,093	1
1988	3,144	79	68	122	154	216	294	382	437	941	-
1987	2,770	71	77	88	150	200	225	325	413	810	-
1986	2,864	74	82	103	171	199	253	357	412	785	1
1985	2,674	59	83	107	162	177	227	363	408	719	2
1984	2,333	51	69	93	113	172	234	283	344	586	1
1983	2,044	57	72	101	114	151	197	252	270	505	3
1982	1,915	51	70	110	119	129	188	208	252	426	2
1981	2,075	69	74	112	131	167	190	242	273	453	-
1980	2,262	63	81	138	108	186	218	285	270	487	1
1979	1,856	56	82	96	115	140	194	206	181	347	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
All races, both sexes											
1998	1,724	10	11	28	33	57	95	198	315	926	-
1997	720	4	14	10	21	36	42	65	97	367	-
1996	745	9	12	21	17	25	59	67	116	363	-
1995	606	5	13	13	18	37	43	57	78	300	-
1994	1,229	16	16	20	40	52	95	142	201	598	-
1993	1,044	10	12	19	27	40	74	90	168	532	-
1992	1,006	7	11	17	27	36	84	135	144	492	-
1991	1,137	14	12	18	36	45	92	124	167	541	-
1990	2,098	11	23	36	60	87	159	238	386	1,044	-
1989	1,593	13	18	19	53	70	102	169	262	791	-
1988	1,943	14	18	26	33	81	111	224	342	1,039	-
1987	632	5	12	15	22	23	41	66	89	314	-
1986	1,838	18	18	20	37	77	132	243	301	919	-
1985	2,054	11	20	34	56	79	135	251	347	1,068	3
1984	1,096	9	7	24	25	43	83	159	184	503	1
1983	1,431	7	12	30	37	61	122	148	252	715	-
1982	727	6	9	13	27	36	50	90	107	348	-
1981	3,006	17	31	60	87	151	248	390	549	1,372	-
1980	2,702	11	22	45	66	124	197	323	538	1,288	-
1979	604	6	4	12	16	38	46	67	102	265	-
Influenza (487)											
All races, male											
1998	584	6	7	11	14	25	46	96	130	230	-
1997	271	4	3	8	11	19	22	30	37	101	-
1996	269	6	7	13	5	13	25	31	48	97	-
1995	222	1	8	5	8	15	20	24	37	84	-
1994	426	8	9	8	19	17	41	63	74	164	-
1993	389	8	7	7	11	21	35	48	85	126	-
1992	338	3	6	7	8	14	37	47	57	129	-
1991	433	8	7	12	17	22	38	57	55	164	-
1990	799	6	10	16	39	46	76	115	155	312	-
1989	550	8	11	8	27	40	48	73	95	200	-
1988	673	9	9	12	15	43	54	96	132	275	-
1987	252	3	10	6	10	15	24	32	35	90	-
1986	676	11	11	14	21	38	75	111	130	232	-
1985	737	4	10	12	27	39	57	108	146	303	-
1984	416	3	4	11	17	20	29	78	71	149	-
1983	543	3	4	21	17	28	60	63	101	229	-
1982	295	3	7	9	13	21	22	48	44	109	-
1981	1,105	6	13	29	40	71	111	177	213	392	-
1980	993	4	10	26	32	62	104	138	191	378	-
1979	232	2	2	4	6	22	26	31	39	69	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
All races, female											
1998	1,140	4	4	17	19	32	49	102	185	696	-
1997	449	-	11	2	10	17	20	35	60	266	-
1996	476	3	5	8	12	12	34	36	68	266	-
1995	384	4	5	8	10	22	23	33	41	216	-
1994	803	8	7	12	21	35	54	79	127	434	-
1993	655	2	5	12	16	19	39	42	83	406	-
1992	668	4	5	10	19	22	47	88	87	363	-
1991	704	6	5	6	19	23	54	67	112	377	-
1990	1,299	5	13	20	21	41	83	123	231	732	-
1989	1,043	5	7	11	26	30	54	96	167	591	-
1988	1,270	5	9	14	18	38	57	128	210	764	-
1987	380	2	2	9	12	8	17	34	54	224	-
1986	1,162	7	7	6	16	39	57	132	171	687	-
1985	1,317	7	10	22	29	40	78	143	201	765	3
1984	680	6	3	13	8	23	54	81	113	354	1
1983	888	4	8	9	20	33	62	85	151	486	-
1982	432	3	2	4	14	15	28	42	63	239	-
1981	1,901	11	18	31	47	80	137	213	336	980	-
1980	1,709	7	12	19	34	62	93	185	347	910	-
1979	372	4	2	8	10	16	20	36	63	196	-
Influenza (487)											
White, both sexes											
1998	1,667	8	11	25	31	53	92	191	308	905	-
1997	681	3	12	7	18	34	40	63	92	362	-
1996	695	9	9	19	14	24	54	60	112	351	-
1995	566	5	11	10	17	34	41	52	75	289	-
1994	1,177	15	15	18	36	43	89	138	197	582	-
1993	992	9	11	16	25	37	67	85	164	517	-
1992	954	3	11	14	26	30	77	132	139	478	-
1991	1,069	11	11	18	35	44	86	116	159	522	-
1990	1,999	9	19	33	53	80	150	224	374	1,014	-
1989	1,509	10	16	16	52	62	94	159	256	760	-
1988	1,865	12	16	22	32	76	101	218	331	1,017	-
1987	595	4	12	14	19	21	37	64	85	298	-
1986	1,754	16	17	19	32	72	125	232	285	898	-
1985	1,947	9	16	29	48	73	119	235	335	1,040	3
1984	1,041	9	7	20	23	40	80	146	177	491	1
1983	1,380	7	10	25	37	54	120	142	244	704	-
1982	686	3	8	11	23	33	48	81	104	341	-
1981	2,840	14	27	50	77	130	236	370	518	1,338	-
1980	2,595	7	18	37	64	116	190	307	529	1,256	-
1979	550	6	3	11	14	28	41	63	94	253	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
White, male											
1998	560	5	7	10	13	24	44	93	127	222	-
1997	251	3	3	6	8	18	21	28	37	100	-
1996	246	6	5	11	3	12	24	28	47	91	-
1995	206	1	8	5	7	13	18	22	35	82	-
1994	406	8	9	7	17	15	39	60	73	159	-
1993	366	8	7	7	10	19	33	44	83	120	-
1992	314	1	6	5	8	10	34	45	55	125	-
1991	402	5	7	12	16	22	34	53	53	159	-
1990	751	6	6	16	35	41	70	107	148	304	-
1989	513	7	9	6	26	35	43	68	94	190	-
1988	640	7	7	11	15	41	51	93	125	271	-
1987	236	2	10	5	10	14	22	31	33	86	-
1986	644	11	10	14	17	36	71	107	123	227	-
1985	681	4	7	10	22	37	48	100	138	290	-
1984	388	3	4	10	15	17	29	71	67	141	-
1983	520	3	3	18	17	25	59	60	97	225	-
1982	273	2	6	8	11	18	22	42	42	105	-
1981	1,033	6	10	24	37	59	105	170	203	379	-
1980	951	1	8	24	31	58	99	132	189	372	-
1979	204	2	1	4	5	15	24	29	34	65	-
Influenza (487)											
White, female											
1998	1,107	3	4	15	18	29	48	98	181	683	-
1997	430	-	9	1	10	16	19	35	55	262	-
1996	449	3	4	8	11	12	30	32	65	260	-
1995	360	4	3	5	10	21	23	30	40	207	-
1994	771	7	6	11	19	28	50	78	124	423	-
1993	626	1	4	9	15	18	34	41	81	397	-
1992	640	2	5	9	18	20	43	87	84	353	-
1991	667	6	4	6	19	22	52	63	106	363	-
1990	1,248	3	13	17	18	39	80	117	226	710	-
1989	996	3	7	10	26	27	51	91	162	570	-
1988	1,225	5	9	11	17	35	50	125	206	746	-
1987	359	2	2	9	9	7	15	33	52	212	-
1986	1,110	5	7	5	15	36	54	125	162	671	-
1985	1,266	5	9	19	26	36	71	135	197	750	3
1984	653	6	3	10	8	23	51	75	110	350	1
1983	860	4	7	7	20	29	61	82	147	479	-
1982	413	1	2	3	12	15	26	39	62	236	-
1981	1,807	8	17	26	40	71	131	200	315	959	-
1980	1,644	6	10	13	33	58	91	175	340	884	-
1979	346	4	2	7	9	13	17	34	60	188	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
All other, both sexes											
1998	57	2	-	3	2	4	3	7	7	21	-
1997	39	1	2	3	3	2	2	2	5	5	-
1996	50	-	3	2	3	1	5	7	4	12	-
1995	40	-	2	3	1	3	2	5	3	11	-
1994	52	1	1	2	4	9	6	4	4	16	-
1993	52	1	1	3	2	3	7	5	4	15	-
1992	52	4	-	3	1	6	7	3	5	14	-
1991	68	3	1	-	1	1	6	8	8	19	-
1990	99	2	4	3	7	7	9	14	12	30	-
1989	84	3	2	3	1	8	8	10	6	31	-
1988	78	2	2	4	1	5	10	6	11	22	-
1987	37	1	-	1	3	2	4	2	4	16	-
1986	84	2	1	1	5	5	7	11	16	21	-
1985	107	2	4	5	8	6	16	16	12	28	-
1984	55	-	-	4	2	3	3	13	7	12	-
1983	51	-	2	5	-	7	2	6	8	11	-
1982	41	3	1	2	4	3	2	9	3	7	-
1981	166	3	4	10	10	21	12	20	31	34	-
1980	107	4	4	8	2	8	7	16	9	32	-
1979	54	-	1	1	2	10	5	4	8	12	-
Influenza (487)											
All other, male											
1998	24	1	-	1	1	1	2	3	3	8	-
1997	20	1	-	2	3	1	1	2	-	1	-
1996	23	-	2	2	2	1	1	3	1	6	-
1995	16	-	-	-	1	2	2	2	2	2	-
1994	20	-	-	1	2	2	2	3	1	5	-
1993	23	-	-	-	1	2	2	4	2	6	-
1992	24	2	-	2	-	4	3	2	2	4	-
1991	31	3	-	-	1	-	4	4	2	5	-
1990	48	-	4	-	4	5	6	8	7	8	-
1989	37	1	2	2	1	5	5	5	1	10	-
1988	33	2	2	1	-	2	3	3	7	4	-
1987	16	1	-	1	-	1	2	1	2	4	-
1986	32	-	1	-	4	2	4	4	7	5	-
1985	56	-	3	2	5	2	9	8	8	13	-
1984	28	-	-	1	2	3	-	7	4	8	-
1983	23	-	1	3	-	3	1	3	4	4	-
1982	22	1	1	1	2	3	-	6	2	4	-
1981	72	-	3	5	3	12	6	7	10	13	-
1980	42	3	2	2	1	4	5	6	2	6	-
1979	28	-	1	-	1	7	2	2	5	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
All other, female											
1998	33	1	-	2	1	3	1	4	4	13	-
1997	19	-	2	1	-	1	1	-	5	4	-
1996	27	-	1	-	1	-	4	4	3	6	-
1995	24	-	2	3	-	1	-	3	1	9	-
1994	32	1	1	1	2	7	4	1	3	11	-
1993	29	1	1	3	1	1	5	1	2	9	-
1992	28	2	-	1	1	2	4	1	3	10	-
1991	37	-	1	-	-	1	2	4	6	14	-
1990	51	2	-	3	3	2	3	6	5	22	-
1989	47	2	-	1	-	3	3	5	5	21	-
1988	45	-	-	3	1	3	7	3	4	18	-
1987	21	-	-	-	3	1	2	1	2	12	-
1986	52	2	-	1	1	3	3	7	9	16	-
1985	51	2	1	3	3	4	7	8	4	15	-
1984	27	-	-	3	-	-	3	6	3	4	-
1983	28	-	1	2	-	4	1	3	4	7	-
1982	19	2	-	1	2	-	2	3	1	3	-
1981	94	3	1	5	7	9	6	13	21	21	-
1980	65	1	2	6	1	4	2	10	7	26	-
1979	26	-	-	1	1	3	3	2	3	8	-
Influenza (487)											
Black, both sexes											
1998	33	1	-	2	1	3	2	4	3	11	-
1997	26	1	-	1	2	2	2	1	4	3	-
1996	34	-	2	1	3	1	4	2	3	10	-
1995	31	-	2	3	1	2	2	3	2	7	-
1994	39	1	1	-	4	6	6	4	3	10	-
1993	47	1	1	3	2	3	4	5	4	14	-
1992	40	4	-	2	1	4	6	2	3	11	-
1991	52	3	1	-	1	1	4	7	6	14	-
1990	90	1	3	2	7	6	8	14	12	26	-
1989	64	3	2	2	1	5	6	7	5	23	-
1988	69	2	2	3	1	3	10	6	9	19	-
1987	22	1	-	1	3	2	3	2	2	6	-
1986	69	2	1	1	5	4	5	8	14	16	-
1985	91	2	3	5	8	6	13	14	11	20	-
1984	43	-	-	4	2	3	2	11	5	8	-
1983	40	-	2	2	-	6	2	3	7	8	-
1982	27	2	1	1	4	3	-	6	2	3	-
1981	146	3	3	10	9	21	10	16	28	28	-
1980	88	4	2	8	1	8	6	15	6	22	-
1979	43	-	1	1	1	8	5	3	6	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Influenza (487)											
Black, male											
1998	13	1	-	1	1	-	2	1	1	2	-
1997	13	1	-	-	2	1	1	1	-	-	-
1996	14	-	2	1	2	1	-	1	-	4	-
1995	11	-	-	-	1	1	2	1	2	-	-
1994	15	-	-	-	2	2	2	3	-	3	-
1993	23	-	-	-	1	2	2	4	2	6	-
1992	18	2	-	2	-	2	3	1	-	3	-
1991	23	3	-	-	1	-	2	3	2	3	-
1990	46	-	3	-	4	4	6	8	7	8	-
1989	27	1	2	1	1	4	5	3	-	6	-
1988	28	2	2	-	-	2	3	3	6	2	-
1987	10	1	-	1	-	1	1	1	1	2	-
1986	24	-	1	-	4	2	2	3	6	2	-
1985	48	-	2	2	5	2	8	7	7	10	-
1984	22	-	-	1	2	3	-	6	3	5	-
1983	19	-	1	2	-	3	1	1	3	4	-
1982	15	1	1	1	2	3	-	4	1	1	-
1981	64	-	3	5	3	12	4	7	10	10	-
1980	37	3	2	2	1	4	4	6	1	4	-
1979	22	-	1	-	1	6	2	1	4	1	-
Influenza (487)											
Black, female											
1998	20	-	-	1	-	3	-	3	2	9	-
1997	13	-	-	1	-	1	1	-	4	3	-
1996	20	-	-	-	1	-	4	1	3	6	-
1995	20	-	2	3	-	1	-	2	-	7	-
1994	24	1	1	-	2	4	4	1	3	7	-
1993	24	1	1	3	1	1	2	1	2	8	-
1992	22	2	-	-	1	2	3	1	3	8	-
1991	29	-	1	-	-	1	2	4	4	11	-
1990	44	1	-	2	3	2	2	6	5	18	-
1989	37	2	-	1	-	1	1	4	5	17	-
1988	41	-	-	3	1	1	7	3	3	17	-
1987	12	-	-	-	3	1	2	1	1	4	-
1986	45	2	-	1	1	2	3	5	8	14	-
1985	43	2	1	3	3	4	5	7	4	10	-
1984	21	-	-	3	-	-	2	5	2	3	-
1983	21	-	1	-	-	3	1	2	4	4	-
1982	12	1	-	-	2	-	-	2	1	2	-
1981	82	3	-	5	6	9	6	9	18	18	-
1980	51	1	-	6	-	4	2	9	5	18	-
1979	21	-	-	1	-	2	3	2	2	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
All races, both sexes											
1998	90,147	970	1,176	1,524	2,271	3,759	7,094	11,519	16,825	42,201	5
1997	85,729	1,021	1,194	1,472	2,256	3,728	6,729	11,142	16,054	39,301	6
1996	82,982	1,020	1,052	1,443	2,132	3,674	6,839	10,610	15,562	37,664	10
1995	82,317	980	1,081	1,301	2,126	3,751	6,906	10,666	15,184	37,275	4
1994	80,244	927	1,026	1,277	2,168	3,870	6,585	10,484	14,933	35,802	5
1993	81,776	851	1,006	1,329	2,329	4,032	6,677	10,631	14,970	36,639	11
1992	74,713	790	837	1,305	2,104	3,832	6,260	9,796	13,906	32,804	7
1991	76,723	773	939	1,332	2,352	3,780	6,306	10,071	14,233	33,603	14
1990	77,415	777	940	1,391	2,435	4,015	6,404	10,372	14,381	33,399	7
1989	74,957	769	907	1,403	2,405	4,032	6,214	10,006	13,585	32,164	15
1988	75,719	793	946	1,486	2,579	4,157	6,364	10,122	13,793	32,110	11
1987	68,593	696	913	1,365	2,477	3,909	6,053	9,289	12,333	28,425	7
1986	67,974	687	877	1,493	2,594	3,880	6,065	9,492	11,959	27,733	46
1985	65,561	646	946	1,516	2,524	3,636	5,978	8,979	11,719	26,637	48
1984	57,798	590	883	1,404	2,298	3,319	5,533	7,982	10,270	22,673	13
1983	54,423	570	909	1,425	2,251	3,291	4,964	7,267	9,626	21,358	15
1982	48,159	562	884	1,462	2,114	2,986	4,605	6,494	8,436	17,877	16
1981	50,725	653	976	1,628	2,258	3,247	4,703	6,882	8,930	18,563	14
1980	51,917	683	1,044	1,653	2,280	3,468	4,879	6,943	9,199	18,553	15
1979	44,426	616	1,008	1,447	2,036	2,956	4,297	5,923	7,903	15,001	10
Pneumonia (480-486)											
All races, male											
1998	40,395	593	696	930	1,332	2,202	4,103	6,198	8,118	14,642	4
1997	39,013	643	725	848	1,327	2,242	3,864	6,061	7,832	13,803	4
1996	37,722	646	643	898	1,248	2,194	3,933	5,758	7,523	13,099	6
1995	37,565	651	658	770	1,299	2,234	3,989	5,778	7,376	13,000	2
1994	36,913	593	644	780	1,308	2,312	3,817	5,697	7,112	12,721	1
1993	37,607	531	614	784	1,426	2,434	3,939	5,853	7,246	12,777	8
1992	35,127	527	523	818	1,273	2,304	3,667	5,458	6,880	11,781	4
1991	35,781	495	556	841	1,450	2,215	3,667	5,542	7,056	11,894	10
1990	36,099	488	610	869	1,523	2,440	3,797	5,667	7,002	11,646	3
1989	35,151	495	577	912	1,513	2,474	3,797	5,426	6,486	11,333	11
1988	36,161	537	617	922	1,638	2,632	3,821	5,689	6,714	11,508	6
1987	33,310	455	588	864	1,578	2,484	3,683	5,178	6,043	10,477	5
1986	33,527	425	571	942	1,688	2,447	3,687	5,395	5,952	10,465	16
1985	32,422	423	601	990	1,655	2,356	3,669	5,092	5,785	10,049	23
1984	29,024	363	592	928	1,531	2,104	3,429	4,555	5,111	8,700	11
1983	27,464	365	587	934	1,495	2,139	3,073	4,102	4,828	8,282	9
1982	25,030	388	558	962	1,361	1,932	2,848	3,787	4,307	7,254	5
1981	25,954	411	617	1,073	1,488	2,105	2,849	3,881	4,498	7,307	12
1980	26,581	462	688	1,044	1,501	2,218	2,962	3,858	4,556	7,428	9
1979	23,493	394	681	972	1,314	1,954	2,722	3,322	4,010	6,238	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
All races, female											
1998	49,752	377	480	594	939	1,557	2,991	5,321	8,707	27,559	1
1997	46,716	378	469	624	929	1,486	2,865	5,081	8,222	25,498	2
1996	45,260	374	409	545	884	1,480	2,906	4,852	8,039	24,565	4
1995	44,752	329	423	531	827	1,517	2,917	4,888	7,808	24,275	2
1994	43,331	334	382	497	860	1,558	2,768	4,787	7,821	23,081	4
1993	44,169	320	392	545	903	1,598	2,738	4,778	7,724	23,862	3
1992	39,586	263	314	487	831	1,528	2,593	4,338	7,026	21,023	3
1991	40,942	278	383	491	902	1,565	2,639	4,529	7,177	21,709	4
1990	41,316	289	330	522	912	1,575	2,607	4,705	7,379	21,753	4
1989	39,806	274	330	491	892	1,558	2,417	4,580	7,099	20,831	4
1988	39,558	256	329	564	941	1,525	2,543	4,433	7,079	20,602	5
1987	35,283	241	325	501	899	1,425	2,370	4,111	6,290	17,948	2
1986	34,447	262	306	551	906	1,433	2,378	4,097	6,007	17,268	30
1985	33,139	223	345	526	869	1,280	2,309	3,887	5,934	16,588	25
1984	28,774	227	291	476	767	1,215	2,104	3,427	5,159	13,973	2
1983	26,959	205	322	491	756	1,152	1,891	3,165	4,798	13,076	6
1982	23,129	174	326	500	753	1,054	1,757	2,707	4,129	10,623	11
1981	24,771	242	359	555	770	1,142	1,854	3,001	4,432	11,256	2
1980	25,336	221	356	609	779	1,250	1,917	3,085	4,643	11,125	6
1979	20,933	222	327	475	722	1,002	1,575	2,601	3,893	8,763	-
Pneumonia (480-486)											
White, both sexes											
1998	79,992	677	874	1,182	1,796	3,111	6,106	10,173	15,254	39,010	1
1997	76,194	671	905	1,124	1,777	3,093	5,804	9,911	14,615	36,404	2
1996	73,499	701	770	1,085	1,660	3,043	5,868	9,494	14,095	34,851	7
1995	73,075	662	790	965	1,688	3,115	5,957	9,531	13,795	34,569	4
1994	71,428	650	745	965	1,756	3,212	5,756	9,406	13,643	33,273	3
1993	72,728	564	770	1,006	1,854	3,385	5,738	9,472	13,657	34,140	4
1992	66,502	565	600	997	1,668	3,190	5,435	8,844	12,706	30,570	4
1991	68,207	523	664	1,005	1,902	3,169	5,500	9,008	12,999	31,332	11
1990	68,807	545	668	1,064	1,923	3,339	5,587	9,281	13,164	31,219	5
1989	66,344	506	626	1,043	1,930	3,373	5,365	8,879	12,469	30,055	12
1988	67,705	481	672	1,109	2,070	3,504	5,568	9,093	12,791	30,293	10
1987	61,216	451	645	1,051	1,992	3,339	5,343	8,392	11,348	26,723	2
1986	60,569	456	629	1,157	2,098	3,280	5,320	8,548	10,996	26,045	37
1985	58,561	429	684	1,172	2,037	3,078	5,262	8,084	10,764	25,141	37
1984	51,473	391	608	1,091	1,871	2,798	4,837	7,188	9,464	21,411	10
1983	48,616	383	667	1,099	1,830	2,803	4,340	6,567	8,909	20,243	7
1982	42,829	339	657	1,124	1,717	2,567	4,003	5,943	7,823	16,931	12
1981	45,079	425	691	1,242	1,826	2,733	4,130	6,264	8,289	17,599	7
1980	45,774	440	747	1,254	1,867	2,874	4,274	6,183	8,564	17,495	13
1979	39,253	418	719	1,121	1,659	2,497	3,780	5,340	7,414	14,219	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
White, male											
1998	35,235	411	505	715	1,040	1,796	3,510	5,450	7,351	13,437	1
1997	34,135	416	536	632	1,041	1,859	3,355	5,376	7,119	12,673	2
1996	32,678	425	467	670	949	1,818	3,345	5,112	6,783	11,971	5
1995	32,742	436	465	556	1,033	1,836	3,443	5,165	6,683	11,924	2
1994	32,252	408	456	571	1,058	1,932	3,335	5,077	6,452	11,715	1
1993	32,785	341	453	578	1,134	2,018	3,404	5,187	6,575	11,787	3
1992	30,637	369	361	593	1,002	1,925	3,163	4,908	6,250	10,883	2
1991	31,187	331	388	630	1,164	1,849	3,200	4,941	6,401	10,957	7
1990	31,350	323	425	658	1,210	2,017	3,296	5,065	6,339	10,766	1
1989	30,379	310	381	674	1,201	2,049	3,253	4,808	5,918	10,478	9
1988	31,622	312	421	671	1,303	2,218	3,347	5,083	6,198	10,747	5
1987	29,048	288	407	648	1,257	2,125	3,221	4,636	5,524	9,737	-
1986	29,247	273	409	716	1,372	2,064	3,219	4,836	5,436	9,676	10
1985	28,347	267	428	756	1,335	1,987	3,200	4,577	5,280	9,387	15
1984	25,289	222	390	712	1,227	1,770	2,987	4,063	4,692	8,135	9
1983	23,943	238	424	712	1,203	1,812	2,666	3,678	4,408	7,777	4
1982	21,827	221	406	743	1,088	1,655	2,451	3,465	3,976	6,804	3
1981	22,498	252	411	802	1,191	1,761	2,474	3,520	4,139	6,845	5
1980	22,859	287	482	782	1,200	1,817	2,594	3,409	4,208	6,908	8
1979	20,340	254	478	747	1,057	1,646	2,404	2,963	3,731	5,852	7
Pneumonia (480-486)											
White, female											
1998	44,757	266	369	467	756	1,315	2,596	4,723	7,903	25,573	-
1997	42,059	255	369	492	736	1,234	2,449	4,535	7,496	23,731	-
1996	40,821	276	303	415	711	1,225	2,523	4,382	7,312	22,880	2
1995	40,333	226	325	409	655	1,279	2,514	4,366	7,112	22,645	2
1994	39,176	242	289	394	698	1,280	2,421	4,329	7,191	21,558	2
1993	39,943	223	317	428	720	1,367	2,334	4,285	7,082	22,353	1
1992	35,865	196	239	404	666	1,265	2,272	3,936	6,456	19,687	2
1991	37,020	192	276	375	738	1,320	2,300	4,067	6,598	20,375	4
1990	37,457	222	243	406	713	1,322	2,291	4,216	6,825	20,453	4
1989	35,965	196	245	369	729	1,324	2,112	4,071	6,551	19,577	3
1988	36,083	169	251	438	767	1,286	2,221	4,010	6,593	19,546	5
1987	32,168	163	238	403	735	1,214	2,122	3,756	5,824	16,986	2
1986	31,322	183	220	441	726	1,216	2,101	3,712	5,560	16,369	27
1985	30,214	162	256	416	702	1,091	2,062	3,507	5,484	15,754	22
1984	26,184	169	218	379	644	1,028	1,850	3,125	4,772	13,276	1
1983	24,673	145	243	387	627	991	1,674	2,889	4,501	12,466	3
1982	21,002	118	251	381	629	912	1,552	2,478	3,847	10,127	9
1981	22,581	173	280	440	635	972	1,656	2,744	4,150	10,754	2
1980	22,915	153	265	472	667	1,057	1,680	2,774	4,356	10,587	5
1979	18,913	164	241	374	602	851	1,376	2,377	3,683	8,367	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
All other, both sexes											
1998	10,155	293	302	342	475	648	988	1,346	1,571	3,191	4
1997	9,535	350	289	348	479	635	925	1,231	1,439	2,897	4
1996	9,483	319	282	358	472	631	971	1,116	1,467	2,813	3
1995	9,242	318	291	336	438	636	949	1,135	1,389	2,706	-
1994	8,816	277	281	312	412	658	829	1,078	1,290	2,529	2
1993	9,048	287	236	323	475	647	939	1,159	1,313	2,499	7
1992	8,211	225	237	308	436	642	825	952	1,200	2,234	3
1991	8,516	250	275	327	450	611	806	1,063	1,234	2,271	3
1990	8,608	232	272	327	512	676	817	1,091	1,217	2,180	2
1989	8,613	263	281	360	475	659	849	1,127	1,116	2,109	3
1988	8,014	312	274	377	509	653	796	1,029	1,002	1,817	1
1987	7,377	245	268	314	485	570	710	897	985	1,702	5
1986	7,405	231	248	336	496	600	745	944	963	1,688	9
1985	7,000	217	262	344	487	558	716	895	955	1,496	11
1984	6,325	199	275	313	427	521	696	794	806	1,262	3
1983	5,807	187	242	326	421	488	624	700	717	1,115	8
1982	5,330	223	227	338	397	419	602	551	613	946	4
1981	5,646	228	285	386	432	514	573	618	641	964	7
1980	6,143	243	297	399	413	594	605	760	635	1,058	2
1979	5,173	198	289	326	377	459	517	583	489	782	3
Pneumonia (480-486)											
All other, male											
1998	5,160	182	191	215	292	406	593	748	767	1,205	3
1997	4,878	227	189	216	286	383	509	685	713	1,130	2
1996	5,044	221	176	228	299	376	588	646	740	1,128	1
1995	4,823	215	193	214	266	398	546	613	693	1,076	-
1994	4,661	185	188	209	250	380	482	620	660	1,006	-
1993	4,822	190	161	206	292	416	535	666	671	990	5
1992	4,490	158	162	225	271	379	504	550	630	898	2
1991	4,594	164	168	211	286	366	467	601	655	937	3
1990	4,749	165	185	211	313	423	501	602	663	880	2
1989	4,772	185	196	238	312	425	544	618	568	855	2
1988	4,539	225	196	251	335	414	474	606	516	761	1
1987	4,262	167	181	216	321	359	462	542	519	740	5
1986	4,280	152	162	226	316	383	468	559	516	789	6
1985	4,075	156	173	234	320	369	469	515	505	662	8
1984	3,735	141	202	216	304	334	442	492	419	565	2
1983	3,521	127	163	222	292	327	407	424	420	505	5
1982	3,203	167	152	219	273	277	397	322	331	450	2
1981	3,456	159	206	271	297	344	375	361	359	462	7
1980	3,722	175	206	262	301	401	368	449	348	520	1
1979	3,153	140	203	225	257	308	318	359	279	386	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
All other, female											
1998	4,995	111	111	127	183	242	395	598	804	1,986	1
1997	4,657	123	100	132	193	252	416	546	726	1,767	2
1996	4,439	98	106	130	173	255	383	470	727	1,685	2
1995	4,419	103	98	122	172	238	403	522	696	1,630	-
1994	4,155	92	93	103	162	278	347	458	630	1,523	2
1993	4,226	97	75	117	183	231	404	493	642	1,509	2
1992	3,721	67	75	83	165	263	321	402	570	1,336	1
1991	3,922	86	107	116	164	245	339	462	579	1,334	-
1990	3,859	67	87	116	199	253	316	489	554	1,300	-
1989	3,841	78	85	122	163	234	305	509	548	1,254	1
1988	3,475	87	78	126	174	239	322	423	486	1,056	-
1987	3,115	78	87	98	164	211	248	355	466	962	-
1986	3,125	79	86	110	180	217	277	385	447	899	3
1985	2,925	61	89	110	167	189	247	380	450	834	3
1984	2,590	58	73	97	123	187	254	302	387	697	1
1983	2,286	60	79	104	129	161	217	276	297	610	3
1982	2,127	56	75	119	124	142	205	229	282	496	2
1981	2,190	69	79	115	135	170	198	257	282	502	-
1980	2,421	68	91	137	112	193	237	311	287	538	1
1979	2,020	58	86	101	120	151	199	224	210	396	-
Pneumonia (480-486)											
Black, both sexes											
1998	8,293	266	271	298	427	546	802	1,081	1,257	2,472	4
1997	7,894	317	245	306	412	529	769	1,009	1,140	2,313	4
1996	7,929	289	253	328	411	553	798	912	1,167	2,269	3
1995	7,772	296	252	306	384	548	810	934	1,143	2,154	-
1994	7,433	263	259	278	367	562	696	907	1,039	2,037	2
1993	7,677	267	219	292	428	561	784	959	1,082	2,029	7
1992	7,034	199	219	273	382	568	703	805	1,014	1,822	3
1991	7,320	229	256	288	403	532	708	903	1,027	1,826	3
1990	7,473	214	252	301	462	610	736	921	1,024	1,776	2
1989	7,521	242	255	327	425	595	754	969	936	1,735	3
1988	7,122	293	251	354	456	592	697	900	870	1,539	1
1987	6,543	226	246	283	446	535	630	796	852	1,413	5
1986	6,631	215	234	317	467	558	678	829	838	1,417	6
1985	6,247	203	251	321	458	509	639	806	813	1,254	10
1984	5,626	186	259	296	384	474	627	701	692	1,046	3
1983	5,194	182	229	316	382	445	561	625	619	927	7
1982	4,772	214	214	317	370	379	546	482	540	783	4
1981	5,130	218	272	371	405	477	521	553	560	816	7
1980	5,560	234	273	382	394	553	551	652	564	898	2
1979	4,697	194	271	308	354	418	479	515	419	670	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonia (480-486)											
Black, male											
1998	4,165	166	172	191	266	343	481	598	594	861	3
1997	3,965	206	162	194	247	319	415	561	530	849	2
1996	4,156	202	159	208	258	327	480	520	566	858	1
1995	4,008	202	170	198	233	346	457	494	548	808	-
1994	3,884	175	173	186	222	329	399	513	510	772	-
1993	4,028	179	152	188	267	357	437	538	536	744	5
1992	3,804	140	149	201	240	345	424	451	519	686	2
1991	3,915	153	158	192	258	317	402	503	514	727	3
1990	4,115	154	172	194	286	390	445	498	538	691	2
1989	4,141	169	181	221	287	390	480	511	464	659	2
1988	4,019	214	183	235	303	377	410	521	436	615	1
1987	3,785	155	169	195	299	336	407	472	440	607	5
1986	3,812	143	152	215	297	361	428	477	434	646	5
1985	3,616	146	169	217	299	336	417	450	409	545	8
1984	3,314	135	190	206	271	302	395	423	350	463	2
1983	3,171	125	158	215	268	297	365	375	353	426	4
1982	2,869	164	144	207	253	250	358	276	289	359	2
1981	3,137	152	198	264	280	319	337	320	305	381	7
1980	3,349	172	192	250	286	371	335	376	299	429	1
1979	2,862	138	189	213	239	280	288	311	240	330	2
Pneumonia (480-486)											
Black, female											
1998	4,128	100	99	107	161	203	321	483	663	1,611	1
1997	3,929	111	83	112	165	210	354	448	610	1,464	2
1996	3,773	87	94	120	153	226	318	392	601	1,411	2
1995	3,764	94	82	108	151	202	353	440	595	1,346	-
1994	3,549	88	86	92	145	233	297	394	529	1,265	2
1993	3,649	88	67	104	161	204	347	421	546	1,285	2
1992	3,230	59	70	72	142	223	279	354	495	1,136	1
1991	3,405	76	98	96	145	215	306	400	513	1,099	-
1990	3,358	60	80	107	176	220	291	423	486	1,085	-
1989	3,380	73	74	106	138	205	274	458	472	1,076	1
1988	3,103	79	68	119	153	215	287	379	434	924	-
1987	2,758	71	77	88	147	199	223	324	412	806	-
1986	2,819	72	82	102	170	197	250	352	404	771	1
1985	2,631	57	82	104	159	173	222	356	404	709	2
1984	2,312	51	69	90	113	172	232	278	342	583	1
1983	2,023	57	71	101	114	148	196	250	266	501	3
1982	1,903	50	70	110	117	129	188	206	251	424	2
1981	1,993	66	74	107	125	158	184	233	255	435	-
1980	2,211	62	81	132	108	182	216	276	265	469	1
1979	1,835	56	82	95	115	138	191	204	179	340	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
All races, both sexes											
1998	452	8	7	11	16	26	31	51	59	158	-
1997	465	6	5	9	12	27	29	54	56	153	-
1996	486	6	10	6	19	26	46	44	65	157	-
1995	532	5	2	11	19	27	45	67	70	171	-
1994	578	2	10	19	22	29	54	61	90	167	-
1993	662	12	16	12	30	34	61	67	99	208	1
1992	577	7	13	15	20	35	49	70	70	153	-
1991	633	7	12	20	26	29	52	82	77	216	-
1990	633	5	6	22	27	44	46	72	85	200	-
1989	633	9	11	15	20	46	54	75	100	183	2
1988	635	14	11	15	29	37	64	58	91	193	-
1987	548	8	9	17	23	36	39	67	72	159	-
1986	588	7	7	13	23	32	66	78	98	165	1
1985	600	9	10	20	27	44	48	59	92	187	-
1984	544	9	6	14	29	41	42	51	76	161	1
1983	552	7	13	24	25	31	39	58	59	161	-
1982	503	5	16	11	26	27	32	57	59	127	-
1981	573	7	13	23	28	39	49	63	70	138	-
1980	642	11	12	30	28	40	49	54	91	163	-
1979	554	7	19	20	25	46	44	46	65	110	1
Other acute lower respiratory infections (466)											
All races, male											
1998	194	4	4	4	9	11	16	24	21	50	-
1997	194	2	2	4	8	11	17	28	18	42	-
1996	205	2	1	1	8	9	25	24	24	45	-
1995	224	3	-	2	7	9	19	30	37	54	-
1994	243	-	5	6	8	7	24	32	37	46	-
1993	286	7	9	7	10	16	28	29	43	61	-
1992	282	3	5	9	13	18	27	35	31	42	-
1991	288	3	4	9	15	15	32	43	27	68	-
1990	269	2	1	9	17	23	23	38	26	63	-
1989	288	6	4	7	15	23	25	41	41	54	1
1988	281	8	5	5	11	19	39	29	35	53	-
1987	267	4	3	10	10	20	23	35	34	57	-
1986	280	3	2	7	16	18	44	41	38	52	-
1985	303	8	5	13	17	23	31	38	39	59	-
1984	260	3	4	10	17	21	26	30	31	51	-
1983	273	2	7	15	17	17	21	34	19	61	-
1982	250	3	8	6	13	17	23	31	21	36	-
1981	277	5	6	15	15	23	24	34	32	45	-
1980	299	7	9	6	10	23	25	30	35	52	-
1979	283	3	12	14	15	19	27	25	31	41	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
All races, female											
1998	258	4	3	7	7	15	15	27	38	108	-
1997	271	4	3	5	4	16	12	26	38	111	-
1996	281	4	9	5	11	17	21	20	41	112	-
1995	308	2	2	9	12	18	26	37	33	117	-
1994	335	2	5	13	14	22	30	29	53	121	-
1993	376	5	7	5	20	18	33	38	56	147	1
1992	295	4	8	6	7	17	22	35	39	111	-
1991	345	4	8	11	11	14	20	39	50	148	-
1990	364	3	5	13	10	21	23	34	59	137	-
1989	345	3	7	8	5	23	29	34	59	129	1
1988	354	6	6	10	18	18	25	29	56	140	-
1987	281	4	6	7	13	16	16	32	38	102	-
1986	308	4	5	6	7	14	22	37	60	113	1
1985	297	1	5	7	10	21	17	21	53	128	-
1984	284	6	2	4	12	20	16	21	45	110	1
1983	279	5	6	9	8	14	18	24	40	100	-
1982	253	2	8	5	13	10	9	26	38	91	-
1981	296	2	7	8	13	16	25	29	38	93	-
1980	343	4	3	24	18	17	24	24	56	111	-
1979	271	4	7	6	10	27	17	21	34	69	1
Other acute lower respiratory infections (466)											
White, both sexes											
1998	400	6	7	11	16	26	29	46	56	152	-
1997	404	4	2	7	9	25	26	50	54	148	-
1996	418	5	8	6	18	25	43	40	61	147	-
1995	477	4	2	9	18	25	44	64	66	167	-
1994	497	1	8	15	20	29	49	59	77	163	-
1993	583	12	10	10	27	28	57	63	93	197	-
1992	500	4	11	14	17	34	47	67	61	147	-
1991	560	7	10	15	22	29	45	76	71	200	-
1990	559	3	6	19	26	41	45	66	84	194	-
1989	544	6	8	14	17	42	50	70	95	174	1
1988	564	14	8	13	28	34	61	55	86	181	-
1987	485	7	6	16	21	35	37	67	67	152	-
1986	537	5	5	11	20	31	60	74	93	161	1
1985	532	7	8	18	24	38	44	54	83	182	-
1984	482	9	6	10	28	38	36	48	71	156	1
1983	481	7	10	22	17	27	34	53	58	159	-
1982	433	4	12	10	23	20	31	52	55	124	-
1981	507	5	12	22	26	35	43	60	67	134	-
1980	568	10	11	24	27	37	42	51	88	156	-
1979	482	6	14	16	22	43	42	45	64	108	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
White, male											
1998	164	4	4	4	9	11	14	20	20	47	-
1997	161	1	-	4	6	10	14	26	16	39	-
1996	176	2	-	1	8	8	23	21	24	40	-
1995	197	2	-	1	7	7	18	28	36	52	-
1994	199	-	3	4	8	7	21	31	32	45	-
1993	242	7	5	5	9	13	26	26	39	56	-
1992	239	3	4	8	11	18	26	34	25	40	-
1991	248	3	4	6	12	15	27	38	24	62	-
1990	233	1	1	8	16	22	23	35	26	63	-
1989	241	3	4	6	12	20	24	38	39	53	1
1988	246	8	3	4	10	18	38	28	32	51	-
1987	236	4	1	9	10	19	23	35	31	54	-
1986	251	1	1	6	14	17	40	39	36	51	-
1985	258	6	3	12	15	19	28	35	34	56	-
1984	230	3	4	7	17	20	22	28	28	49	-
1983	229	2	5	13	12	15	19	31	18	60	-
1982	205	2	6	6	11	12	23	28	19	35	-
1981	242	4	5	14	14	20	22	33	31	43	-
1980	255	6	8	5	9	21	21	28	34	48	-
1979	243	3	9	11	12	17	25	24	31	41	-
Other acute lower respiratory infections (466)											
White, female											
1998	236	2	3	7	7	15	15	26	36	105	-
1997	243	3	2	3	3	15	12	24	38	109	-
1996	242	3	8	5	10	17	20	19	37	107	-
1995	280	2	2	8	11	18	26	36	30	115	-
1994	298	1	5	11	12	22	28	28	45	118	-
1993	341	5	5	5	18	15	31	37	54	141	-
1992	261	1	7	6	6	16	21	33	36	107	-
1991	312	4	6	9	10	14	18	38	47	138	-
1990	326	2	5	11	10	19	22	31	58	131	-
1989	303	3	4	8	5	22	26	32	56	121	-
1988	318	6	5	9	18	16	23	27	54	130	-
1987	249	3	5	7	11	16	14	32	36	98	-
1986	286	4	4	5	6	14	20	35	57	110	1
1985	274	1	5	6	9	19	16	19	49	126	-
1984	252	6	2	3	11	18	14	20	43	107	1
1983	252	5	5	9	5	12	15	22	40	99	-
1982	228	2	6	4	12	8	8	24	36	89	-
1981	265	1	7	8	12	15	21	27	36	91	-
1980	313	4	3	19	18	16	21	23	54	108	-
1979	239	3	5	5	10	26	17	21	33	67	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
All other, both sexes											
1998	52	2	-	-	-	-	2	5	3	6	-
1997	61	2	3	2	3	2	3	4	2	5	-
1996	68	1	2	-	1	1	3	4	4	10	-
1995	55	1	-	2	1	2	1	3	4	4	-
1994	81	1	2	4	2	-	5	2	13	4	-
1993	79	-	6	2	3	6	4	4	6	11	1
1992	77	3	2	1	3	1	2	3	9	6	-
1991	73	-	2	5	4	-	7	6	6	16	-
1990	74	2	-	3	1	3	1	6	1	6	-
1989	89	3	3	1	3	4	4	5	5	9	1
1988	71	-	3	2	1	3	3	3	5	12	-
1987	63	1	3	1	2	1	2	-	5	7	-
1986	51	2	2	2	3	1	6	4	5	4	-
1985	68	2	2	2	3	6	4	5	9	5	-
1984	62	-	-	4	1	3	6	3	5	5	-
1983	71	-	3	2	8	4	5	5	1	2	-
1982	70	1	4	1	3	7	1	5	4	3	-
1981	66	2	1	1	2	4	6	3	3	4	-
1980	74	1	1	6	1	3	7	3	3	7	-
1979	72	1	5	4	3	3	2	1	1	2	1
Other acute lower respiratory infections (466)											
All other, male											
1998	30	-	-	-	-	-	2	4	1	3	-
1997	33	1	2	-	2	1	3	2	2	3	-
1996	29	-	1	-	-	1	2	3	-	5	-
1995	27	1	-	1	-	2	1	2	1	2	-
1994	44	-	2	2	-	-	3	1	5	1	-
1993	44	-	4	2	1	3	2	3	4	5	-
1992	43	-	1	1	2	-	1	1	6	2	-
1991	40	-	-	3	3	-	5	5	3	6	-
1990	36	1	-	1	1	1	-	3	-	-	-
1989	47	3	-	1	3	3	1	3	2	1	-
1988	35	-	2	1	1	1	1	1	3	2	-
1987	31	-	2	1	-	1	-	-	3	3	-
1986	29	2	1	1	2	1	4	2	2	1	-
1985	45	2	2	1	2	4	3	3	5	3	-
1984	30	-	-	3	-	1	4	2	3	2	-
1983	44	-	2	2	5	2	2	3	1	1	-
1982	45	1	2	-	2	5	-	3	2	1	-
1981	35	1	1	1	1	3	2	1	1	2	-
1980	44	1	1	1	1	2	4	2	1	4	-
1979	40	-	3	3	3	2	2	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
All other, female											
1998	22	2	-	-	-	-	-	1	2	3	-
1997	28	1	1	2	1	1	-	2	-	2	-
1996	39	1	1	-	1	-	1	1	4	5	-
1995	28	-	-	1	1	-	-	1	3	2	-
1994	37	1	-	2	2	-	2	1	8	3	-
1993	35	-	2	-	2	3	2	1	2	6	1
1992	34	3	1	-	1	1	1	2	3	4	-
1991	33	-	2	2	1	-	2	1	3	10	-
1990	38	1	-	2	-	2	1	3	1	6	-
1989	42	-	3	-	-	1	3	2	3	8	1
1988	36	-	1	1	-	2	2	2	2	10	-
1987	32	1	1	-	2	-	2	-	2	4	-
1986	22	-	1	1	1	-	2	2	3	3	-
1985	23	-	-	1	1	2	1	2	4	2	-
1984	32	-	-	1	1	2	2	1	2	3	-
1983	27	-	1	-	3	2	3	2	-	1	-
1982	25	-	2	1	1	2	1	2	2	2	-
1981	31	1	-	-	1	1	4	2	2	2	-
1980	30	-	-	5	-	1	3	1	2	3	-
1979	32	1	2	1	-	1	-	-	1	2	1
Other acute lower respiratory infections (466)											
Black, both sexes											
1998	46	2	-	-	-	-	2	5	3	4	-
1997	53	1	3	2	3	1	3	4	2	3	-
1996	55	1	2	-	-	1	1	3	3	9	-
1995	49	1	-	2	-	1	1	3	4	4	-
1994	77	1	2	4	2	-	5	2	13	4	-
1993	69	-	5	2	3	6	4	3	4	7	1
1992	71	3	2	1	3	1	2	3	6	5	-
1991	57	-	-	4	3	-	7	5	4	13	-
1990	64	2	-	3	1	2	1	5	1	4	-
1989	79	3	3	-	3	4	4	4	4	9	1
1988	62	-	3	2	1	3	2	3	5	10	-
1987	56	1	3	1	2	1	2	-	4	6	-
1986	47	2	2	1	3	1	5	4	4	4	-
1985	63	2	2	2	3	6	4	5	7	3	-
1984	56	-	-	4	1	3	4	3	4	4	-
1983	59	-	3	1	5	3	5	5	1	2	-
1982	59	1	4	1	2	6	1	4	3	3	-
1981	56	2	1	1	2	4	6	2	3	2	-
1980	64	-	1	6	1	3	7	2	3	5	-
1979	69	1	5	4	3	2	2	1	1	2	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other acute lower respiratory infections (466)											
Black, male											
1998	27	-	-	-	-	-	2	4	1	2	-
1997	27	-	2	-	2	-	3	2	2	2	-
1996	24	-	1	-	-	1	1	2	-	4	-
1995	24	1	-	1	-	1	1	2	1	2	-
1994	41	-	2	2	-	-	3	1	5	1	-
1993	39	-	3	2	1	3	2	3	3	3	-
1992	38	-	1	1	2	-	1	1	3	1	-
1991	31	-	-	2	2	-	5	4	2	5	-
1990	33	1	-	1	1	1	-	3	-	-	-
1989	39	3	-	-	3	3	1	2	1	1	-
1988	32	-	2	1	1	1	1	1	3	2	-
1987	27	-	2	1	-	1	-	-	2	3	-
1986	27	2	1	1	2	1	3	2	2	1	-
1985	41	2	2	1	2	4	3	3	4	1	-
1984	28	-	-	3	-	1	4	2	2	1	-
1983	34	-	2	1	2	1	2	3	1	1	-
1982	36	1	2	-	1	4	-	2	1	1	-
1981	28	1	1	1	1	3	2	-	1	1	-
1980	35	-	1	1	1	2	4	1	1	2	-
1979	39	-	3	3	3	1	2	1	-	-	-
Other acute lower respiratory infections (466)											
Black, female											
1998	19	2	-	-	-	-	-	1	2	2	-
1997	26	1	1	2	1	1	-	2	-	1	-
1996	31	1	1	-	-	-	-	1	3	5	-
1995	25	-	-	1	-	-	-	1	3	2	-
1994	36	1	-	2	2	-	2	1	8	3	-
1993	30	-	2	-	2	3	2	-	1	4	1
1992	33	3	1	-	1	1	1	2	3	4	-
1991	26	-	-	2	1	-	2	1	2	8	-
1990	31	1	-	2	-	1	1	2	1	4	-
1989	40	-	3	-	-	1	3	2	3	8	1
1988	30	-	1	1	-	2	1	2	2	8	-
1987	29	1	1	-	2	-	2	-	2	3	-
1986	20	-	1	-	1	-	2	2	2	3	-
1985	22	-	-	1	1	2	1	2	3	2	-
1984	28	-	-	1	1	2	-	1	2	3	-
1983	25	-	1	-	3	2	3	2	-	1	-
1982	23	-	2	1	1	2	1	2	2	2	-
1981	28	1	-	-	1	1	4	2	2	1	-
1980	29	-	-	5	-	1	3	1	2	3	-
1979	30	1	2	1	-	1	-	-	1	2	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
All races, both sexes											
1998	452	8	7	11	16	26	31	51	59	158	-
1997	465	6	5	9	12	27	29	54	56	153	-
1996	486	6	10	6	19	26	46	44	65	157	-
1995	532	5	2	11	19	27	45	67	70	171	-
1994	578	2	10	19	22	29	54	61	90	167	-
1993	662	12	16	12	30	34	61	67	99	208	1
1992	577	7	13	15	20	35	49	70	70	153	-
1991	633	7	12	20	26	29	52	82	77	216	-
1990	633	5	6	22	27	44	46	72	85	200	-
1989	633	9	11	15	20	46	54	75	100	183	2
1988	635	14	11	15	29	37	64	58	91	193	-
1987	548	8	9	17	23	36	39	67	72	159	-
1986	588	7	7	13	23	32	66	78	98	165	1
1985	600	9	10	20	27	44	48	59	92	187	-
1984	544	9	6	14	29	41	42	51	76	161	1
1983	552	7	13	24	25	31	39	58	59	161	-
1982	503	5	16	11	26	27	32	57	59	127	-
1981	573	7	13	23	28	39	49	63	70	138	-
1980	642	11	12	30	28	40	49	54	91	163	-
1979	554	7	19	20	25	46	44	46	65	110	1
Acute bronchitis and bronchiolitis (466)											
All races, male											
1998	194	4	4	4	9	11	16	24	21	50	-
1997	194	2	2	4	8	11	17	28	18	42	-
1996	205	2	1	1	8	9	25	24	24	45	-
1995	224	3	-	2	7	9	19	30	37	54	-
1994	243	-	5	6	8	7	24	32	37	46	-
1993	286	7	9	7	10	16	28	29	43	61	-
1992	282	3	5	9	13	18	27	35	31	42	-
1991	288	3	4	9	15	15	32	43	27	68	-
1990	269	2	1	9	17	23	23	38	26	63	-
1989	288	6	4	7	15	23	25	41	41	54	1
1988	281	8	5	5	11	19	39	29	35	53	-
1987	267	4	3	10	10	20	23	35	34	57	-
1986	280	3	2	7	16	18	44	41	38	52	-
1985	303	8	5	13	17	23	31	38	39	59	-
1984	260	3	4	10	17	21	26	30	31	51	-
1983	273	2	7	15	17	17	21	34	19	61	-
1982	250	3	8	6	13	17	23	31	21	36	-
1981	277	5	6	15	15	23	24	34	32	45	-
1980	299	7	9	6	10	23	25	30	35	52	-
1979	283	3	12	14	15	19	27	25	31	41	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
All races, female											
1998	258	4	3	7	7	15	15	27	38	108	-
1997	271	4	3	5	4	16	12	26	38	111	-
1996	281	4	9	5	11	17	21	20	41	112	-
1995	308	2	2	9	12	18	26	37	33	117	-
1994	335	2	5	13	14	22	30	29	53	121	-
1993	376	5	7	5	20	18	33	38	56	147	1
1992	295	4	8	6	7	17	22	35	39	111	-
1991	345	4	8	11	11	14	20	39	50	148	-
1990	364	3	5	13	10	21	23	34	59	137	-
1989	345	3	7	8	5	23	29	34	59	129	1
1988	354	6	6	10	18	18	25	29	56	140	-
1987	281	4	6	7	13	16	16	32	38	102	-
1986	308	4	5	6	7	14	22	37	60	113	1
1985	297	1	5	7	10	21	17	21	53	128	-
1984	284	6	2	4	12	20	16	21	45	110	1
1983	279	5	6	9	8	14	18	24	40	100	-
1982	253	2	8	5	13	10	9	26	38	91	-
1981	296	2	7	8	13	16	25	29	38	93	-
1980	343	4	3	24	18	17	24	24	56	111	-
1979	271	4	7	6	10	27	17	21	34	69	1
Acute bronchitis and bronchiolitis (466)											
White, both sexes											
1998	400	6	7	11	16	26	29	46	56	152	-
1997	404	4	2	7	9	25	26	50	54	148	-
1996	418	5	8	6	18	25	43	40	61	147	-
1995	477	4	2	9	18	25	44	64	66	167	-
1994	497	1	8	15	20	29	49	59	77	163	-
1993	583	12	10	10	27	28	57	63	93	197	-
1992	500	4	11	14	17	34	47	67	61	147	-
1991	560	7	10	15	22	29	45	76	71	200	-
1990	559	3	6	19	26	41	45	66	84	194	-
1989	544	6	8	14	17	42	50	70	95	174	1
1988	564	14	8	13	28	34	61	55	86	181	-
1987	485	7	6	16	21	35	37	67	67	152	-
1986	537	5	5	11	20	31	60	74	93	161	1
1985	532	7	8	18	24	38	44	54	83	182	-
1984	482	9	6	10	28	38	36	48	71	156	1
1983	481	7	10	22	17	27	34	53	58	159	-
1982	433	4	12	10	23	20	31	52	55	124	-
1981	507	5	12	22	26	35	43	60	67	134	-
1980	568	10	11	24	27	37	42	51	88	156	-
1979	482	6	14	16	22	43	42	45	64	108	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
White, male											
1998	164	4	4	4	9	11	14	20	20	47	-
1997	161	1	-	4	6	10	14	26	16	39	-
1996	176	2	-	1	8	8	23	21	24	40	-
1995	197	2	-	1	7	7	18	28	36	52	-
1994	199	-	3	4	8	7	21	31	32	45	-
1993	242	7	5	5	9	13	26	26	39	56	-
1992	239	3	4	8	11	18	26	34	25	40	-
1991	248	3	4	6	12	15	27	38	24	62	-
1990	233	1	1	8	16	22	23	35	26	63	-
1989	241	3	4	6	12	20	24	38	39	53	1
1988	246	8	3	4	10	18	38	28	32	51	-
1987	236	4	1	9	10	19	23	35	31	54	-
1986	251	1	1	6	14	17	40	39	36	51	-
1985	258	6	3	12	15	19	28	35	34	56	-
1984	230	3	4	7	17	20	22	28	28	49	-
1983	229	2	5	13	12	15	19	31	18	60	-
1982	205	2	6	6	11	12	23	28	19	35	-
1981	242	4	5	14	14	20	22	33	31	43	-
1980	255	6	8	5	9	21	21	28	34	48	-
1979	243	3	9	11	12	17	25	24	31	41	-
Acute bronchitis and bronchiolitis (466)											
White, female											
1998	236	2	3	7	7	15	15	26	36	105	-
1997	243	3	2	3	3	15	12	24	38	109	-
1996	242	3	8	5	10	17	20	19	37	107	-
1995	280	2	2	8	11	18	26	36	30	115	-
1994	298	1	5	11	12	22	28	28	45	118	-
1993	341	5	5	5	18	15	31	37	54	141	-
1992	261	1	7	6	6	16	21	33	36	107	-
1991	312	4	6	9	10	14	18	38	47	138	-
1990	326	2	5	11	10	19	22	31	58	131	-
1989	303	3	4	8	5	22	26	32	56	121	-
1988	318	6	5	9	18	16	23	27	54	130	-
1987	249	3	5	7	11	16	14	32	36	98	-
1986	286	4	4	5	6	14	20	35	57	110	1
1985	274	1	5	6	9	19	16	19	49	126	-
1984	252	6	2	3	11	18	14	20	43	107	1
1983	252	5	5	9	5	12	15	22	40	99	-
1982	228	2	6	4	12	8	8	24	36	89	-
1981	265	1	7	8	12	15	21	27	36	91	-
1980	313	4	3	19	18	16	21	23	54	108	-
1979	239	3	5	5	10	26	17	21	33	67	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
All other, both sexes											
1998	52	2	-	-	-	-	2	5	3	6	-
1997	61	2	3	2	3	2	3	4	2	5	-
1996	68	1	2	-	1	1	3	4	4	10	-
1995	55	1	-	2	1	2	1	3	4	4	-
1994	81	1	2	4	2	-	5	2	13	4	-
1993	79	-	6	2	3	6	4	4	6	11	1
1992	77	3	2	1	3	1	2	3	9	6	-
1991	73	-	2	5	4	-	7	6	6	16	-
1990	74	2	-	3	1	3	1	6	1	6	-
1989	89	3	3	1	3	4	4	5	5	9	1
1988	71	-	3	2	1	3	3	3	5	12	-
1987	63	1	3	1	2	1	2	-	5	7	-
1986	51	2	2	2	3	1	6	4	5	4	-
1985	68	2	2	2	3	6	4	5	9	5	-
1984	62	-	-	4	1	3	6	3	5	5	-
1983	71	-	3	2	8	4	5	5	1	2	-
1982	70	1	4	1	3	7	1	5	4	3	-
1981	66	2	1	1	2	4	6	3	3	4	-
1980	74	1	1	6	1	3	7	3	3	7	-
1979	72	1	5	4	3	3	2	1	1	2	1
Acute bronchitis and bronchiolitis (466)											
All other, male											
1998	30	-	-	-	-	-	2	4	1	3	-
1997	33	1	2	-	2	1	3	2	2	3	-
1996	29	-	1	-	-	1	2	3	-	5	-
1995	27	1	-	1	-	2	1	2	1	2	-
1994	44	-	2	2	-	-	3	1	5	1	-
1993	44	-	4	2	1	3	2	3	4	5	-
1992	43	-	1	1	2	-	1	1	6	2	-
1991	40	-	-	3	3	-	5	5	3	6	-
1990	36	1	-	1	1	1	-	3	-	-	-
1989	47	3	-	1	3	3	1	3	2	1	-
1988	35	-	2	1	1	1	1	1	3	2	-
1987	31	-	2	1	-	1	-	-	3	3	-
1986	29	2	1	1	2	1	4	2	2	1	-
1985	45	2	2	1	2	4	3	3	5	3	-
1984	30	-	-	3	-	1	4	2	3	2	-
1983	44	-	2	2	5	2	2	3	1	1	-
1982	45	1	2	-	2	5	-	3	2	1	-
1981	35	1	1	1	1	3	2	1	1	2	-
1980	44	1	1	1	1	2	4	2	1	4	-
1979	40	-	3	3	3	2	2	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
All other, female											
1998	22	2	-	-	-	-	-	1	2	3	-
1997	28	1	1	2	1	1	-	2	-	2	-
1996	39	1	1	-	1	-	1	1	4	5	-
1995	28	-	-	1	1	-	-	1	3	2	-
1994	37	1	-	2	2	-	2	1	8	3	-
1993	35	-	2	-	2	3	2	1	2	6	1
1992	34	3	1	-	1	1	1	2	3	4	-
1991	33	-	2	2	1	-	2	1	3	10	-
1990	38	1	-	2	-	2	1	3	1	6	-
1989	42	-	3	-	-	1	3	2	3	8	1
1988	36	-	1	1	-	2	2	2	2	10	-
1987	32	1	1	-	2	-	2	-	2	4	-
1986	22	-	1	1	1	-	2	2	3	3	-
1985	23	-	-	1	1	2	1	2	4	2	-
1984	32	-	-	1	1	2	2	1	2	3	-
1983	27	-	1	-	3	2	3	2	-	1	-
1982	25	-	2	1	1	2	1	2	2	2	-
1981	31	1	-	-	1	1	4	2	2	2	-
1980	30	-	-	5	-	1	3	1	2	3	-
1979	32	1	2	1	-	1	-	-	1	2	1
Acute bronchitis and bronchiolitis (466)											
Black, both sexes											
1998	46	2	-	-	-	-	2	5	3	4	-
1997	53	1	3	2	3	1	3	4	2	3	-
1996	55	1	2	-	-	1	1	3	3	9	-
1995	49	1	-	2	-	1	1	3	4	4	-
1994	77	1	2	4	2	-	5	2	13	4	-
1993	69	-	5	2	3	6	4	3	4	7	1
1992	71	3	2	1	3	1	2	3	6	5	-
1991	57	-	-	4	3	-	7	5	4	13	-
1990	64	2	-	3	1	2	1	5	1	4	-
1989	79	3	3	-	3	4	4	4	4	9	1
1988	62	-	3	2	1	3	2	3	5	10	-
1987	56	1	3	1	2	1	2	-	4	6	-
1986	47	2	2	1	3	1	5	4	4	4	-
1985	63	2	2	2	3	6	4	5	7	3	-
1984	56	-	-	4	1	3	4	3	4	4	-
1983	59	-	3	1	5	3	5	5	1	2	-
1982	59	1	4	1	2	6	1	4	3	3	-
1981	56	2	1	1	2	4	6	2	3	2	-
1980	64	-	1	6	1	3	7	2	3	5	-
1979	69	1	5	4	3	2	2	1	1	2	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute bronchitis and bronchiolitis (466)											
Black, male											
1998	27	-	-	-	-	-	2	4	1	2	-
1997	27	-	2	-	2	-	3	2	2	2	-
1996	24	-	1	-	-	1	1	2	-	4	-
1995	24	1	-	1	-	1	1	2	1	2	-
1994	41	-	2	2	-	-	3	1	5	1	-
1993	39	-	3	2	1	3	2	3	3	3	-
1992	38	-	1	1	2	-	1	1	3	1	-
1991	31	-	-	2	2	-	5	4	2	5	-
1990	33	1	-	1	1	1	-	3	-	-	-
1989	39	3	-	-	3	3	1	2	1	1	-
1988	32	-	2	1	1	1	1	1	3	2	-
1987	27	-	2	1	-	1	-	-	2	3	-
1986	27	2	1	1	2	1	3	2	2	1	-
1985	41	2	2	1	2	4	3	3	4	1	-
1984	28	-	-	3	-	1	4	2	2	1	-
1983	34	-	2	1	2	1	2	3	1	1	-
1982	36	1	2	-	1	4	-	2	1	1	-
1981	28	1	1	1	1	3	2	-	1	1	-
1980	35	-	1	1	1	2	4	1	1	2	-
1979	39	-	3	3	3	1	2	1	-	-	-
Acute bronchitis and bronchiolitis (466)											
Black, female											
1998	19	2	-	-	-	-	-	1	2	2	-
1997	26	1	1	2	1	1	-	2	-	1	-
1996	31	1	1	-	-	-	-	1	3	5	-
1995	25	-	-	1	-	-	-	1	3	2	-
1994	36	1	-	2	2	-	2	1	8	3	-
1993	30	-	2	-	2	3	2	-	1	4	1
1992	33	3	1	-	1	1	1	2	3	4	-
1991	26	-	-	2	1	-	2	1	2	8	-
1990	31	1	-	2	-	1	1	2	1	4	-
1989	40	-	3	-	-	1	3	2	3	8	1
1988	30	-	1	1	-	2	1	2	2	8	-
1987	29	1	1	-	2	-	2	-	2	3	-
1986	20	-	1	-	1	-	2	2	2	3	-
1985	22	-	-	1	1	2	1	2	3	2	-
1984	28	-	-	1	1	2	-	1	2	3	-
1983	25	-	1	-	3	2	3	2	-	1	-
1982	23	-	2	1	1	2	1	2	2	2	-
1981	28	1	-	-	1	1	4	2	2	1	-
1980	29	-	-	5	-	1	3	1	2	3	-
1979	30	1	2	1	-	1	-	-	1	2	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
All races, both sexes											
1998	112,559	940	1,887	3,511	6,647	12,115	18,980	22,619	21,088	23,074	7
1997	109,002	966	1,870	3,441	6,666	12,207	18,355	21,785	20,296	21,746	7
1996	105,988	969	1,827	3,473	6,568	12,062	18,100	21,413	19,525	20,345	13
1995	102,876	931	1,823	3,439	6,548	12,161	17,953	20,494	18,707	19,144	3
1994	101,601	898	1,776	3,444	6,886	12,651	17,996	20,267	18,171	17,943	11
1993	101,053	835	1,657	3,470	7,197	12,948	18,308	20,350	17,947	16,854	10
1992	91,924	749	1,523	3,343	6,754	12,018	16,685	18,814	15,652	15,003	8
1991	90,624	779	1,556	3,295	7,134	12,139	16,415	18,439	15,281	14,119	10
1990	86,653	753	1,523	3,351	6,971	11,827	15,685	17,826	14,310	13,090	8
1989	84,325	713	1,570	3,643	7,103	11,541	15,515	17,164	13,458	12,272	6
1988	82,839	734	1,525	3,498	7,159	11,518	15,663	16,616	13,184	11,623	11
1987	78,366	654	1,490	3,401	7,045	10,928	14,774	15,691	12,243	10,804	9
1986	76,544	671	1,564	3,531	6,950	10,973	14,888	15,286	11,418	10,075	10
1985	74,643	679	1,614	3,642	6,963	10,838	14,303	14,698	11,036	9,743	17
1984	69,079	618	1,505	3,605	6,661	10,150	13,511	13,421	9,844	8,684	8
1983	66,234	656	1,571	3,574	6,407	10,139	12,922	12,667	9,239	7,935	16
1982	59,861	605	1,521	3,325	5,899	9,346	11,836	11,299	8,165	6,776	13
1981	58,819	649	1,596	3,394	6,094	9,399	11,683	10,805	7,717	6,473	7
1980	56,040	677	1,558	3,328	5,947	9,120	10,988	10,107	7,232	6,136	6
1979	49,923	597	1,538	3,046	5,577	8,180	9,764	8,801	6,442	5,057	5
Chronic lower respiratory diseases (490-494,496)											
All races, male											
1998	56,999	467	912	1,827	3,527	6,455	10,137	11,822	10,619	10,328	7
1997	55,969	494	948	1,770	3,475	6,653	9,986	11,405	10,329	10,034	5
1996	54,460	483	925	1,789	3,509	6,526	9,503	11,344	10,130	9,398	7
1995	53,919	463	896	1,766	3,546	6,613	9,805	11,086	9,797	9,108	3
1994	53,707	446	906	1,822	3,750	6,864	9,878	10,917	9,701	8,569	9
1993	54,359	436	892	1,827	3,942	7,109	10,049	11,220	9,761	8,355	6
1992	50,455	377	791	1,836	3,745	6,787	9,440	10,506	8,669	7,551	7
1991	50,464	411	752	1,768	3,938	6,904	9,326	10,525	8,730	7,315	8
1990	49,397	401	751	1,859	3,968	6,897	9,230	10,465	8,252	6,857	8
1989	48,166	340	801	1,999	3,996	6,793	8,963	10,135	7,821	6,584	4
1988	48,930	378	824	1,946	4,215	6,930	9,428	10,092	7,995	6,409	7
1987	47,028	307	789	1,912	4,212	6,591	9,070	9,868	7,580	5,967	6
1986	47,046	370	849	2,041	4,155	6,746	9,388	9,739	7,253	5,858	6
1985	46,585	345	885	2,097	4,212	6,802	9,254	9,496	7,133	5,769	13
1984	43,995	325	873	2,144	4,164	6,392	8,872	8,927	6,512	5,223	4
1983	43,111	338	883	2,203	4,039	6,534	8,634	8,732	6,251	4,909	13
1982	39,696	306	894	2,028	3,819	6,200	8,035	7,956	5,616	4,253	10
1981	39,764	346	957	2,132	3,979	6,358	8,182	7,725	5,324	4,209	4
1980	38,620	384	945	2,126	3,958	6,345	7,842	7,421	5,117	3,976	3
1979	35,287	364	925	1,988	3,775	5,869	7,207	6,519	4,771	3,379	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
All races, female											
1998	55,560	473	975	1,684	3,120	5,660	8,843	10,797	10,469	12,746	-
1997	53,033	472	922	1,671	3,191	5,554	8,369	10,380	9,967	11,712	2
1996	51,528	486	902	1,684	3,059	5,536	8,597	10,069	9,395	10,947	6
1995	48,957	468	927	1,673	3,002	5,548	8,148	9,408	8,910	10,036	-
1994	47,894	452	870	1,622	3,136	5,787	8,118	9,350	8,470	9,374	2
1993	46,694	399	765	1,643	3,255	5,839	8,259	9,130	8,186	8,499	4
1992	41,469	372	732	1,507	3,009	5,231	7,245	8,308	6,983	7,452	1
1991	40,160	368	804	1,527	3,196	5,235	7,089	7,914	6,551	6,804	2
1990	37,256	352	772	1,492	3,003	4,930	6,455	7,361	6,058	6,233	-
1989	36,159	373	769	1,644	3,107	4,748	6,552	7,029	5,637	5,688	2
1988	33,909	356	701	1,552	2,944	4,588	6,235	6,524	5,189	5,214	4
1987	31,338	347	701	1,489	2,833	4,337	5,704	5,823	4,663	4,837	3
1986	29,498	301	715	1,490	2,795	4,227	5,500	5,547	4,165	4,217	4
1985	28,058	334	729	1,545	2,751	4,036	5,049	5,202	3,903	3,974	4
1984	25,084	293	632	1,461	2,497	3,758	4,639	4,494	3,332	3,461	4
1983	23,123	318	688	1,371	2,368	3,605	4,288	3,935	2,988	3,026	3
1982	20,165	299	627	1,297	2,080	3,146	3,801	3,343	2,549	2,523	3
1981	19,055	303	639	1,262	2,115	3,041	3,501	3,080	2,393	2,264	3
1980	17,420	293	613	1,202	1,989	2,775	3,146	2,686	2,115	2,160	3
1979	14,636	233	613	1,058	1,802	2,311	2,557	2,282	1,671	1,678	1
Chronic lower respiratory diseases (490-494,496)											
White, both sexes											
1998	104,036	672	1,558	3,041	5,983	11,136	17,648	21,193	20,028	21,774	7
1997	100,743	724	1,575	3,000	5,968	11,170	17,093	20,489	19,170	20,526	5
1996	97,851	722	1,546	3,006	5,881	11,086	16,841	20,117	18,479	19,164	9
1995	95,056	693	1,524	2,985	5,882	11,173	16,706	19,306	17,682	18,069	3
1994	94,046	689	1,480	3,004	6,231	11,646	16,759	19,130	17,200	16,975	5
1993	93,592	628	1,404	3,038	6,501	11,953	17,046	19,210	16,939	15,991	8
1992	85,218	571	1,279	2,915	6,119	11,042	15,594	17,790	14,852	14,196	7
1991	83,927	579	1,281	2,894	6,445	11,159	15,330	17,443	14,500	13,404	5
1990	80,155	548	1,275	2,922	6,309	10,892	14,682	16,784	13,521	12,406	5
1989	77,863	571	1,286	3,199	6,422	10,572	14,483	16,121	12,740	11,639	5
1988	76,660	562	1,261	3,074	6,458	10,621	14,634	15,672	12,510	11,065	8
1987	72,655	507	1,239	3,010	6,400	10,088	13,825	14,873	11,613	10,240	7
1986	71,107	519	1,318	3,100	6,308	10,161	13,962	14,510	10,862	9,595	9
1985	69,419	535	1,371	3,213	6,335	10,045	13,499	13,905	10,488	9,310	10
1984	64,256	474	1,277	3,223	6,068	9,477	12,683	12,784	9,306	8,274	4
1983	61,697	518	1,374	3,179	5,811	9,473	12,227	12,085	8,773	7,552	12
1982	55,879	477	1,301	2,943	5,430	8,749	11,166	10,776	7,833	6,502	11
1981	54,971	529	1,371	3,053	5,609	8,786	11,075	10,286	7,355	6,229	6
1980	52,366	555	1,345	2,975	5,476	8,555	10,403	9,637	6,953	5,870	6
1979	46,682	475	1,343	2,736	5,178	7,657	9,264	8,355	6,208	4,834	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
White, male											
1998	52,153	340	753	1,564	3,142	5,883	9,355	10,974	9,970	9,642	7
1997	51,169	376	793	1,525	3,089	6,020	9,227	10,612	9,621	9,373	4
1996	49,797	364	789	1,527	3,104	5,950	8,765	10,534	9,473	8,801	4
1995	49,302	333	748	1,535	3,134	6,016	9,013	10,368	9,178	8,484	3
1994	49,204	339	758	1,592	3,362	6,269	9,072	10,197	9,090	8,032	3
1993	49,800	337	752	1,581	3,519	6,498	9,245	10,470	9,106	7,841	5
1992	46,346	294	666	1,587	3,347	6,190	8,729	9,834	8,150	7,080	6
1991	46,297	307	607	1,523	3,535	6,278	8,625	9,850	8,208	6,892	4
1990	45,217	296	615	1,603	3,568	6,292	8,536	9,737	7,701	6,436	5
1989	44,035	268	639	1,748	3,593	6,156	8,269	9,394	7,353	6,187	3
1988	44,818	291	671	1,695	3,776	6,319	8,691	9,390	7,504	6,060	5
1987	43,279	229	652	1,681	3,802	6,012	8,396	9,287	7,149	5,616	5
1986	43,329	294	690	1,774	3,732	6,175	8,716	9,146	6,861	5,531	5
1985	43,063	259	744	1,834	3,797	6,268	8,651	8,915	6,751	5,491	6
1984	40,710	239	722	1,896	3,775	5,892	8,271	8,455	6,128	4,968	1
1983	39,970	259	777	1,931	3,628	6,067	8,086	8,300	5,905	4,660	9
1982	36,964	239	761	1,795	3,495	5,751	7,542	7,571	5,373	4,064	8
1981	36,983	270	813	1,905	3,632	5,891	7,702	7,326	5,052	4,030	4
1980	35,972	313	803	1,875	3,611	5,928	7,383	7,061	4,896	3,794	3
1979	32,934	282	801	1,770	3,496	5,467	6,814	6,162	4,596	3,209	2
Chronic lower respiratory diseases (490-494,496)											
White, female											
1998	51,883	332	805	1,477	2,841	5,253	8,293	10,219	10,058	12,132	-
1997	49,574	348	782	1,475	2,879	5,150	7,866	9,877	9,549	11,153	1
1996	48,054	358	757	1,479	2,777	5,136	8,076	9,583	9,006	10,363	5
1995	45,754	360	776	1,450	2,748	5,157	7,693	8,938	8,504	9,585	-
1994	44,842	350	722	1,412	2,869	5,377	7,687	8,933	8,110	8,943	2
1993	43,792	291	652	1,457	2,982	5,455	7,801	8,740	7,833	8,150	3
1992	38,872	277	613	1,328	2,772	4,852	6,865	7,956	6,702	7,116	1
1991	37,630	272	674	1,371	2,910	4,881	6,705	7,593	6,292	6,512	1
1990	34,938	252	660	1,319	2,741	4,600	6,146	7,047	5,820	5,970	-
1989	33,828	303	647	1,451	2,829	4,416	6,214	6,727	5,387	5,452	2
1988	31,842	271	590	1,379	2,682	4,302	5,943	6,282	5,006	5,005	3
1987	29,376	278	587	1,329	2,598	4,076	5,429	5,586	4,464	4,624	2
1986	27,778	225	628	1,326	2,576	3,986	5,246	5,364	4,001	4,064	4
1985	26,356	276	627	1,379	2,538	3,777	4,848	4,990	3,737	3,819	4
1984	23,546	235	555	1,327	2,293	3,585	4,412	4,329	3,178	3,306	3
1983	21,727	259	597	1,248	2,183	3,406	4,141	3,785	2,868	2,892	3
1982	18,915	238	540	1,148	1,935	2,998	3,624	3,205	2,460	2,438	3
1981	17,988	259	558	1,148	1,977	2,895	3,373	2,960	2,303	2,199	2
1980	16,394	242	542	1,100	1,865	2,627	3,020	2,576	2,057	2,076	3
1979	13,748	193	542	966	1,682	2,190	2,450	2,193	1,612	1,625	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
All other, both sexes											
1998	8,523	268	329	470	664	979	1,332	1,426	1,060	1,300	-
1997	8,259	242	295	441	698	1,037	1,262	1,296	1,126	1,220	2
1996	8,137	247	281	467	687	976	1,259	1,296	1,046	1,181	4
1995	7,820	238	299	454	666	988	1,247	1,188	1,025	1,075	-
1994	7,555	209	296	440	655	1,005	1,237	1,137	971	968	6
1993	7,461	207	253	432	696	995	1,262	1,140	1,008	863	2
1992	6,706	178	244	428	635	976	1,091	1,024	800	807	1
1991	6,697	200	275	401	689	980	1,085	996	781	715	5
1990	6,498	205	248	429	662	935	1,003	1,042	789	684	3
1989	6,462	142	284	444	681	969	1,032	1,043	718	633	1
1988	6,179	172	264	424	701	897	1,029	944	674	558	3
1987	5,711	147	251	391	645	840	949	818	630	564	2
1986	5,437	152	246	431	642	812	926	776	556	480	1
1985	5,224	144	243	429	628	793	804	793	548	433	7
1984	4,823	144	228	382	593	673	828	637	538	410	4
1983	4,537	138	197	395	596	666	695	582	466	383	4
1982	3,982	128	220	382	469	597	670	523	332	274	2
1981	3,848	120	225	341	485	613	608	519	362	244	1
1980	3,674	122	213	353	471	565	585	470	279	266	-
1979	3,241	122	195	310	399	523	500	446	234	223	2
Chronic lower respiratory diseases (490-494,496)											
All other, male											
1998	4,846	127	159	263	385	572	782	848	649	686	-
1997	4,800	118	155	245	386	633	759	793	708	661	1
1996	4,663	119	136	262	405	576	738	810	657	597	3
1995	4,617	130	148	231	412	597	792	718	619	624	-
1994	4,503	107	148	230	388	595	806	720	611	537	6
1993	4,559	99	140	246	423	611	804	750	655	514	1
1992	4,109	83	125	249	398	597	711	672	519	471	1
1991	4,167	104	145	245	403	626	701	675	522	423	4
1990	4,180	105	136	256	400	605	694	728	551	421	3
1989	4,131	72	162	251	403	637	694	741	468	397	1
1988	4,112	87	153	251	439	611	737	702	491	349	2
1987	3,749	78	137	231	410	579	674	581	431	351	1
1986	3,717	76	159	267	423	571	672	593	392	327	1
1985	3,522	86	141	263	415	534	603	581	382	278	7
1984	3,285	86	151	248	389	500	601	472	384	255	3
1983	3,141	79	106	272	411	467	548	432	346	249	4
1982	2,732	67	133	233	324	449	493	385	243	189	2
1981	2,781	76	144	227	347	467	480	399	272	179	-
1980	2,648	71	142	251	347	417	459	360	221	182	-
1979	2,353	82	124	218	279	402	393	357	175	170	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
All other, female											
1998	3,677	141	170	207	279	407	550	578	411	614	-
1997	3,459	124	140	196	312	404	503	503	418	559	1
1996	3,474	128	145	205	282	400	521	486	389	584	1
1995	3,203	108	151	223	254	391	455	470	406	451	-
1994	3,052	102	148	210	267	410	431	417	360	431	-
1993	2,902	108	113	186	273	384	458	390	353	349	1
1992	2,597	95	119	179	237	379	380	352	281	336	-
1991	2,530	96	130	156	286	354	384	321	259	292	1
1990	2,318	100	112	173	262	330	309	314	238	263	-
1989	2,331	70	122	193	278	332	338	302	250	236	-
1988	2,067	85	111	173	262	286	292	242	183	209	1
1987	1,962	69	114	160	235	261	275	237	199	213	1
1986	1,720	76	87	164	219	241	254	183	164	153	-
1985	1,702	58	102	166	213	259	201	212	166	155	-
1984	1,538	58	77	134	204	173	227	165	154	155	1
1983	1,396	59	91	123	185	199	147	150	120	134	-
1982	1,250	61	87	149	145	148	177	138	89	85	-
1981	1,067	44	81	114	138	146	128	120	90	65	1
1980	1,026	51	71	102	124	148	126	110	58	84	-
1979	888	40	71	92	120	121	107	89	59	53	-
Chronic lower respiratory diseases (490-494,496)											
Black, both sexes											
1998	7,205	244	303	423	571	834	1,134	1,187	858	1,013	-
1997	6,908	222	257	381	610	881	1,077	1,050	891	959	1
1996	6,923	227	258	414	603	834	1,070	1,067	853	963	4
1995	6,665	217	266	405	584	864	1,054	989	836	859	-
1994	6,511	194	264	392	588	891	1,057	966	772	796	4
1993	6,434	189	236	389	615	877	1,061	961	831	715	1
1992	5,856	164	230	385	562	856	943	888	667	673	1
1991	5,816	193	263	364	616	855	938	846	647	565	5
1990	5,654	194	225	376	595	820	884	880	678	538	3
1989	5,670	129	259	402	607	863	911	899	610	518	1
1988	5,475	161	245	391	637	817	910	807	576	456	1
1987	5,051	134	223	360	583	755	843	701	549	452	2
1986	4,856	141	228	396	594	745	833	667	468	400	1
1985	4,659	136	217	390	570	724	722	681	476	357	7
1984	4,292	135	212	349	549	613	738	558	453	324	3
1983	4,068	132	188	367	554	597	615	518	389	324	2
1982	3,631	123	209	360	438	550	620	450	285	228	2
1981	3,515	115	211	327	451	564	551	462	318	207	1
1980	3,379	115	204	335	453	521	538	421	251	211	-
1979	2,957	118	186	296	372	488	454	408	199	166	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic lower respiratory diseases (490-494,496)											
Black, male											
1998	4,039	113	143	233	325	486	662	701	522	513	-
1997	3,966	108	133	206	337	539	639	644	552	502	1
1996	3,928	112	126	234	358	488	615	653	531	483	3
1995	3,916	118	136	206	367	519	660	602	504	486	-
1994	3,857	101	130	207	349	524	684	608	494	429	4
1993	3,913	91	133	222	372	539	674	637	537	417	1
1992	3,568	76	115	226	351	520	606	585	429	390	1
1991	3,621	99	139	221	370	550	606	581	428	333	4
1990	3,627	99	122	229	363	522	610	613	470	330	3
1989	3,592	62	147	230	356	565	609	630	397	318	1
1988	3,644	80	145	227	404	560	658	594	421	287	1
1987	3,319	70	127	211	374	525	600	499	371	282	1
1986	3,302	72	150	246	390	522	605	509	322	267	1
1985	3,154	82	126	239	378	496	548	496	331	232	7
1984	2,922	82	138	228	361	456	533	414	325	199	2
1983	2,813	73	103	256	382	421	485	389	282	208	2
1982	2,489	64	127	219	299	415	455	335	206	158	2
1981	2,544	73	135	217	324	435	436	357	237	152	-
1980	2,429	68	134	235	334	385	420	321	203	144	-
1979	2,166	79	119	209	267	374	363	333	151	128	2
Chronic lower respiratory diseases (490-494,496)											
Black, female											
1998	3,166	131	160	190	246	348	472	486	336	500	-
1997	2,942	114	124	175	273	342	438	406	339	457	-
1996	2,995	115	132	180	245	346	455	414	322	480	1
1995	2,749	99	130	199	217	345	394	387	332	373	-
1994	2,654	93	134	185	239	367	373	358	278	367	-
1993	2,521	98	103	167	243	338	387	324	294	298	-
1992	2,288	88	115	159	211	336	337	303	238	283	-
1991	2,195	94	124	143	246	305	332	265	219	232	1
1990	2,027	95	103	147	232	298	274	267	208	208	-
1989	2,078	67	112	172	251	298	302	269	213	200	-
1988	1,831	81	100	164	233	257	252	213	155	169	-
1987	1,732	64	96	149	209	230	243	202	178	170	1
1986	1,554	69	78	150	204	223	228	158	146	133	-
1985	1,505	54	91	151	192	228	174	185	145	125	-
1984	1,370	53	74	121	188	157	205	144	128	125	1
1983	1,255	59	85	111	172	176	130	129	107	116	-
1982	1,142	59	82	141	139	135	165	115	79	70	-
1981	971	42	76	110	127	129	115	105	81	55	1
1980	950	47	70	100	119	136	118	100	48	67	-
1979	791	39	67	87	105	114	91	75	48	38	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
All races, both sexes											
1998	3,032	36	65	106	173	270	406	527	521	814	-
1997	3,055	29	69	84	160	278	442	523	523	828	-
1996	3,151	39	65	96	170	315	457	547	548	777	1
1995	3,333	29	54	102	179	360	470	557	573	889	-
1994	3,579	46	72	114	209	399	528	647	564	879	-
1993	3,777	38	71	120	235	384	592	657	621	918	1
1992	3,664	27	76	128	247	409	576	641	597	838	-
1991	3,727	36	74	125	274	374	585	691	580	842	-
1990	3,600	35	60	130	256	379	519	641	630	817	-
1989	3,759	28	70	154	244	426	553	675	602	885	-
1988	3,720	36	56	144	278	425	604	701	548	799	2
1987	3,581	27	53	136	267	394	538	645	598	789	-
1986	3,750	29	69	136	286	420	626	676	605	790	2
1985	3,615	33	58	147	295	450	604	585	558	775	1
1984	3,545	32	64	150	308	395	642	623	524	682	-
1983	3,578	36	82	177	296	501	586	602	508	679	-
1982	3,495	36	94	171	297	484	569	627	482	599	1
1981	3,685	39	79	203	350	504	643	633	468	629	-
1980	3,728	38	95	197	359	541	639	617	526	578	-
1979	3,650	34	103	220	377	552	607	587	498	505	1
Bronchitis, chronic and unspecified (490-491)											
All races, male											
1998	1,482	19	28	66	93	136	229	273	243	328	-
1997	1,460	14	31	40	73	139	229	264	260	340	-
1996	1,574	16	30	43	90	167	249	282	297	324	1
1995	1,667	17	35	55	92	200	262	309	292	340	-
1994	1,763	20	38	53	112	202	288	335	301	343	-
1993	1,888	18	36	54	126	211	313	365	317	367	1
1992	1,819	13	35	68	134	204	319	344	309	312	-
1991	1,957	27	40	64	154	198	315	395	315	368	-
1990	1,878	23	29	73	138	216	302	353	339	334	-
1989	1,956	10	33	79	132	254	299	380	333	354	-
1988	1,992	19	33	71	138	250	354	418	306	332	1
1987	1,884	12	31	68	143	225	299	375	330	320	-
1986	2,079	17	34	70	168	255	355	410	357	342	1
1985	2,041	14	21	89	162	274	382	373	325	337	1
1984	2,029	19	37	89	180	243	404	390	286	313	-
1983	2,139	21	52	103	180	312	383	406	283	333	-
1982	2,126	14	52	94	186	311	369	427	295	309	1
1981	2,344	23	47	137	229	321	437	423	299	338	-
1980	2,380	21	62	113	234	375	448	430	326	287	-
1979	2,405	22	62	134	232	369	415	421	349	304	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
All races, female											
1998	1,550	17	37	40	80	134	177	254	278	486	-
1997	1,595	15	38	44	87	139	213	259	263	488	-
1996	1,577	23	35	53	80	148	208	265	251	453	-
1995	1,666	12	19	47	87	160	208	248	281	549	-
1994	1,816	26	34	61	97	197	240	312	263	536	-
1993	1,889	20	35	66	109	173	279	292	304	551	-
1992	1,845	14	41	60	113	205	257	297	288	526	-
1991	1,770	9	34	61	120	176	270	296	265	474	-
1990	1,722	12	31	57	118	163	217	288	291	483	-
1989	1,803	18	37	75	112	172	254	295	269	531	-
1988	1,728	17	23	73	140	175	250	283	242	467	1
1987	1,697	15	22	68	124	169	239	270	268	469	-
1986	1,671	12	35	66	118	165	271	266	248	448	1
1985	1,574	19	37	58	133	176	222	212	233	438	-
1984	1,516	13	27	61	128	152	238	233	238	369	-
1983	1,439	15	30	74	116	189	203	196	225	346	-
1982	1,369	22	42	77	111	173	200	200	187	290	-
1981	1,341	16	32	66	121	183	206	210	169	291	-
1980	1,348	17	33	84	125	166	191	187	200	291	-
1979	1,245	12	41	86	145	183	192	166	149	201	1
Bronchitis, chronic and unspecified (490-491)											
White, both sexes											
1998	2,817	25	53	99	160	244	376	500	496	781	-
1997	2,805	23	61	74	147	250	403	495	486	787	-
1996	2,889	30	55	83	151	276	431	509	525	736	1
1995	3,053	23	43	88	168	328	438	525	531	830	-
1994	3,323	43	64	101	196	366	488	607	540	832	-
1993	3,502	27	64	104	214	358	564	614	594	871	1
1992	3,421	23	70	111	227	381	543	607	578	794	-
1991	3,459	25	55	107	251	348	564	664	552	800	-
1990	3,365	30	52	119	241	354	497	598	599	782	-
1989	3,522	23	60	142	222	397	521	638	580	848	-
1988	3,491	32	51	129	256	393	573	671	529	772	1
1987	3,379	23	47	127	247	370	510	617	575	761	-
1986	3,531	25	64	120	270	390	597	639	575	764	2
1985	3,413	28	48	134	271	432	575	561	534	751	1
1984	3,359	25	61	142	278	377	614	602	507	663	-
1983	3,364	30	73	158	278	470	561	574	495	647	-
1982	3,311	29	89	153	281	465	549	605	463	577	1
1981	3,486	36	68	183	334	482	611	604	447	612	-
1980	3,513	32	79	181	334	510	614	591	511	561	-
1979	3,452	29	93	208	356	526	583	558	479	487	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
White, male											
1998	1,359	11	25	61	86	121	214	257	227	310	-
1997	1,321	10	28	36	70	121	205	251	236	321	-
1996	1,435	12	26	34	79	143	237	261	280	309	1
1995	1,509	13	26	49	88	179	240	291	268	315	-
1994	1,618	19	33	46	109	179	262	305	290	324	-
1993	1,734	12	31	43	111	196	296	340	303	346	1
1992	1,685	11	33	54	123	187	302	326	296	294	-
1991	1,801	19	27	54	139	182	299	377	299	351	-
1990	1,726	20	22	67	128	200	288	319	316	320	-
1989	1,823	9	28	72	124	235	279	353	323	338	-
1988	1,850	16	30	63	125	227	331	397	292	319	1
1987	1,766	11	27	64	135	209	278	359	317	304	-
1986	1,931	16	30	56	156	234	337	385	335	328	1
1985	1,929	11	20	79	149	265	364	353	315	323	1
1984	1,912	15	35	84	166	232	382	374	277	303	-
1983	1,995	17	47	91	167	291	363	388	274	310	-
1982	2,010	13	49	85	173	296	354	412	282	295	1
1981	2,193	21	39	123	219	301	408	401	286	327	-
1980	2,231	18	51	101	218	350	428	412	317	275	-
1979	2,275	18	56	127	220	349	396	401	336	293	-
Bronchitis, chronic and unspecified (490-491)											
White, female											
1998	1,458	14	28	38	74	123	162	243	269	471	-
1997	1,484	13	33	38	77	129	198	244	250	466	-
1996	1,454	18	29	49	72	133	194	248	245	427	-
1995	1,544	10	17	39	80	149	198	234	263	515	-
1994	1,705	24	31	55	87	187	226	302	250	508	-
1993	1,768	15	33	61	103	162	268	274	291	525	-
1992	1,736	12	37	57	104	194	241	281	282	500	-
1991	1,658	6	28	53	112	166	265	287	253	449	-
1990	1,639	10	30	52	113	154	209	279	283	462	-
1989	1,699	14	32	70	98	162	242	285	257	510	-
1988	1,641	16	21	66	131	166	242	274	237	453	-
1987	1,613	12	20	63	112	161	232	258	258	457	-
1986	1,600	9	34	64	114	156	260	254	240	436	1
1985	1,484	17	28	55	122	167	211	208	219	428	-
1984	1,447	10	26	58	112	145	232	228	230	360	-
1983	1,369	13	26	67	111	179	198	186	221	337	-
1982	1,301	16	40	68	108	169	195	193	181	282	-
1981	1,293	15	29	60	115	181	203	203	161	285	-
1980	1,282	14	28	80	116	160	186	179	194	286	-
1979	1,177	11	37	81	136	177	187	157	143	194	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
All other, both sexes											
1998	215	11	12	7	13	26	30	27	25	33	-
1997	250	6	8	10	13	28	39	28	37	41	-
1996	262	9	10	13	19	39	26	38	23	41	-
1995	280	6	11	14	11	32	32	32	42	59	-
1994	256	3	8	13	13	33	40	40	24	47	-
1993	275	11	7	16	21	26	28	43	27	47	-
1992	243	4	6	17	20	28	33	34	19	44	-
1991	268	11	19	18	23	26	21	27	28	42	-
1990	235	5	8	11	15	25	22	43	31	35	-
1989	237	5	10	12	22	29	32	37	22	37	-
1988	229	4	5	15	22	32	31	30	19	27	1
1987	202	4	6	9	20	24	28	28	23	28	-
1986	219	4	5	16	16	30	29	37	30	26	-
1985	202	5	10	13	24	18	29	24	24	24	-
1984	186	7	3	8	30	18	28	21	17	19	-
1983	214	6	9	19	18	31	25	28	13	32	-
1982	184	7	5	18	16	19	20	22	19	22	-
1981	199	3	11	20	16	22	32	29	21	17	-
1980	215	6	16	16	25	31	25	26	15	17	-
1979	198	5	10	12	21	26	24	29	19	18	-
Bronchitis, chronic and unspecified (490-491)											
All other, male											
1998	123	8	3	5	7	15	15	16	16	18	-
1997	139	4	3	4	3	18	24	13	24	19	-
1996	139	4	4	9	11	24	12	21	17	15	-
1995	158	4	9	6	4	21	22	18	24	25	-
1994	145	1	5	7	3	23	26	30	11	19	-
1993	154	6	5	11	15	15	17	25	14	21	-
1992	134	2	2	14	11	17	17	18	13	18	-
1991	156	8	13	10	15	16	16	18	16	17	-
1990	152	3	7	6	10	16	14	34	23	14	-
1989	133	1	5	7	8	19	20	27	10	16	-
1988	142	3	3	8	13	23	23	21	14	13	-
1987	118	1	4	4	8	16	21	16	13	16	-
1986	148	1	4	14	12	21	18	25	22	14	-
1985	112	3	1	10	13	9	18	20	10	14	-
1984	117	4	2	5	14	11	22	16	9	10	-
1983	144	4	5	12	13	21	20	18	9	23	-
1982	116	1	3	9	13	15	15	15	13	14	-
1981	151	2	8	14	10	20	29	22	13	11	-
1980	149	3	11	12	16	25	20	18	9	12	-
1979	130	4	6	7	12	20	19	20	13	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
All other, female											
1998	92	3	9	2	6	11	15	11	9	15	-
1997	111	2	5	6	10	10	15	15	13	22	-
1996	123	5	6	4	8	15	14	17	6	26	-
1995	122	2	2	8	7	11	10	14	18	34	-
1994	111	2	3	6	10	10	14	10	13	28	-
1993	121	5	2	5	6	11	11	18	13	26	-
1992	109	2	4	3	9	11	16	16	6	26	-
1991	112	3	6	8	8	10	5	9	12	25	-
1990	83	2	1	5	5	9	8	9	8	21	-
1989	104	4	5	5	14	10	12	10	12	21	-
1988	87	1	2	7	9	9	8	9	5	14	1
1987	84	3	2	5	12	8	7	12	10	12	-
1986	71	3	1	2	4	9	11	12	8	12	-
1985	90	2	9	3	11	9	11	4	14	10	-
1984	69	3	1	3	16	7	6	5	8	9	-
1983	70	2	4	7	5	10	5	10	4	9	-
1982	68	6	2	9	3	4	5	7	6	8	-
1981	48	1	3	6	6	2	3	7	8	6	-
1980	66	3	5	4	9	6	5	8	6	5	-
1979	68	1	4	5	9	6	5	9	6	7	-
Bronchitis, chronic and unspecified (490-491)											
Black, both sexes											
1998	165	11	9	4	11	21	23	20	21	18	-
1997	196	6	5	9	11	20	31	24	26	31	-
1996	214	8	8	12	17	29	18	31	16	36	-
1995	222	5	10	12	10	27	26	23	29	44	-
1994	209	3	7	12	8	25	36	32	20	36	-
1993	222	10	7	13	18	19	27	31	21	34	-
1992	204	3	6	15	13	25	30	24	18	37	-
1991	224	11	15	17	18	23	18	21	23	29	-
1990	190	5	7	9	13	23	21	35	23	20	-
1989	207	4	9	10	21	27	31	30	20	29	-
1988	193	4	3	15	20	27	27	23	15	21	-
1987	169	2	6	9	18	20	24	23	22	17	-
1986	184	4	5	15	14	29	26	26	26	16	-
1985	167	5	7	12	21	15	25	20	19	18	-
1984	166	7	3	6	27	14	26	18	16	16	-
1983	180	4	9	16	16	25	22	25	10	25	-
1982	163	6	3	17	15	18	20	20	15	15	-
1981	178	3	9	19	14	22	28	27	17	11	-
1980	195	6	16	15	22	29	23	20	14	14	-
1979	170	5	6	11	20	24	21	23	16	15	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Bronchitis, chronic and unspecified (490-491)											
Black, male											
1998	94	8	3	3	6	12	11	11	13	11	-
1997	108	4	2	4	2	13	18	11	17	15	-
1996	108	4	3	8	11	18	6	16	11	11	-
1995	127	3	8	6	3	18	18	15	14	19	-
1994	118	1	4	6	2	18	23	24	10	13	-
1993	122	5	5	8	14	10	16	21	9	14	-
1992	115	1	2	12	7	16	16	14	12	14	-
1991	135	8	11	9	13	14	14	14	14	14	-
1990	125	3	6	6	9	14	14	28	18	7	-
1989	113	-	4	6	8	18	20	20	8	12	-
1988	121	3	2	8	12	19	20	15	13	10	-
1987	100	-	4	4	7	14	19	12	13	10	-
1986	121	1	4	13	11	20	16	17	19	6	-
1985	93	3	1	10	12	7	16	16	8	10	-
1984	105	4	2	3	12	10	20	15	8	8	-
1983	120	2	5	10	11	19	18	16	7	18	-
1982	102	1	2	8	12	15	15	14	10	8	-
1981	133	2	6	13	8	20	25	20	11	6	-
1980	136	3	11	12	14	23	19	15	9	9	-
1979	117	4	4	6	12	19	17	19	10	10	-
Bronchitis, chronic and unspecified (490-491)											
Black, female											
1998	71	3	6	1	5	9	12	9	8	7	-
1997	88	2	3	5	9	7	13	13	9	16	-
1996	106	4	5	4	6	11	12	15	5	25	-
1995	95	2	2	6	7	9	8	8	15	25	-
1994	91	2	3	6	6	7	13	8	10	23	-
1993	100	5	2	5	4	9	11	10	12	20	-
1992	89	2	4	3	6	9	14	10	6	23	-
1991	89	3	4	8	5	9	4	7	9	15	-
1990	65	2	1	3	4	9	7	7	5	13	-
1989	94	4	5	4	13	9	11	10	12	17	-
1988	72	1	1	7	8	8	7	8	2	11	-
1987	69	2	2	5	11	6	5	11	9	7	-
1986	63	3	1	2	3	9	10	9	7	10	-
1985	74	2	6	2	9	8	9	4	11	8	-
1984	61	3	1	3	15	4	6	3	8	8	-
1983	60	2	4	6	5	6	4	9	3	7	-
1982	61	5	1	9	3	3	5	6	5	7	-
1981	45	1	3	6	6	2	3	7	6	5	-
1980	59	3	5	3	8	6	4	5	5	5	-
1979	53	1	2	5	8	5	4	4	6	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
All races, both sexes											
1998	17,555	143	333	685	1,315	2,408	3,427	3,639	3,078	2,428	-
1997	17,518	150	306	627	1,331	2,440	3,457	3,724	2,974	2,402	2
1996	17,286	141	321	681	1,268	2,343	3,390	3,734	3,010	2,315	-
1995	16,927	150	355	692	1,356	2,403	3,302	3,555	2,827	2,206	-
1994	17,215	132	294	689	1,418	2,489	3,463	3,562	2,844	2,228	1
1993	17,586	130	314	710	1,556	2,633	3,534	3,740	2,835	2,056	1
1992	16,363	132	284	675	1,377	2,517	3,380	3,408	2,522	1,979	-
1991	16,436	135	322	688	1,468	2,570	3,244	3,447	2,567	1,921	1
1990	15,706	120	297	704	1,509	2,454	3,188	3,364	2,307	1,677	-
1989	15,467	105	296	770	1,548	2,511	3,084	3,240	2,245	1,586	-
1988	15,513	122	280	733	1,561	2,452	3,286	3,164	2,205	1,614	2
1987	14,483	107	310	728	1,540	2,268	2,953	2,929	2,091	1,453	3
1986	14,471	145	341	753	1,521	2,304	3,047	2,943	1,967	1,368	1
1985	14,150	139	319	765	1,497	2,314	3,001	2,860	1,853	1,316	1
1984	13,263	108	287	791	1,423	2,189	2,753	2,552	1,796	1,268	-
1983	13,293	122	346	817	1,408	2,209	2,793	2,540	1,713	1,241	5
1982	12,793	120	354	802	1,392	2,170	2,675	2,396	1,638	1,156	4
1981	13,369	141	399	853	1,466	2,383	2,804	2,399	1,644	1,171	1
1980	13,877	159	399	883	1,579	2,410	2,804	2,538	1,724	1,274	1
1979	13,569	178	440	839	1,624	2,364	2,785	2,451	1,609	1,157	1
Emphysema (492)											
All races, male											
1998	9,333	85	169	367	746	1,293	1,866	1,957	1,591	1,197	-
1997	9,497	95	166	356	705	1,338	1,954	2,047	1,550	1,222	1
1996	9,422	90	173	365	709	1,348	1,813	2,038	1,644	1,190	-
1995	9,344	95	205	396	756	1,294	1,864	1,960	1,575	1,155	-
1994	9,622	79	171	379	769	1,394	1,974	2,015	1,582	1,199	1
1993	9,914	80	188	393	852	1,475	1,979	2,119	1,642	1,133	-
1992	9,422	88	140	384	800	1,471	1,925	1,944	1,494	1,113	-
1991	9,531	87	169	401	827	1,488	1,875	2,017	1,536	1,086	-
1990	9,408	77	150	403	880	1,487	1,907	2,023	1,447	973	-
1989	9,294	62	170	438	874	1,529	1,840	1,954	1,406	968	-
1988	9,686	68	153	427	958	1,501	2,004	2,038	1,469	1,002	1
1987	9,182	60	172	415	966	1,377	1,885	1,938	1,429	866	3
1986	9,396	93	198	475	928	1,449	2,013	1,972	1,328	880	1
1985	9,375	89	198	471	915	1,489	2,005	1,962	1,297	894	1
1984	8,926	66	175	488	938	1,405	1,882	1,790	1,291	829	-
1983	9,203	73	206	538	921	1,474	1,982	1,841	1,239	863	4
1982	9,037	73	218	495	917	1,495	1,959	1,804	1,193	828	1
1981	9,592	86	270	555	1,007	1,701	2,051	1,823	1,187	844	-
1980	10,133	99	253	593	1,098	1,756	2,055	1,976	1,332	899	-
1979	10,171	120	293	591	1,158	1,768	2,173	1,911	1,234	847	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
All races, female											
1998	8,222	58	164	318	569	1,115	1,561	1,682	1,487	1,231	-
1997	8,021	55	140	271	626	1,102	1,503	1,677	1,424	1,180	1
1996	7,864	51	148	316	559	995	1,577	1,696	1,366	1,125	-
1995	7,583	55	150	296	600	1,109	1,438	1,595	1,252	1,051	-
1994	7,593	53	123	310	649	1,095	1,489	1,547	1,262	1,029	-
1993	7,672	50	126	317	704	1,158	1,555	1,621	1,193	923	1
1992	6,941	44	144	291	577	1,046	1,455	1,464	1,028	866	-
1991	6,905	48	153	287	641	1,082	1,369	1,430	1,031	835	1
1990	6,298	43	147	301	629	967	1,281	1,341	860	704	-
1989	6,173	43	126	332	674	982	1,244	1,286	839	618	-
1988	5,827	54	127	306	603	951	1,282	1,126	736	612	1
1987	5,301	47	138	313	574	891	1,068	991	662	587	-
1986	5,075	52	143	278	593	855	1,034	971	639	488	-
1985	4,775	50	121	294	582	825	996	898	556	422	-
1984	4,337	42	112	303	485	784	871	762	505	439	-
1983	4,090	49	140	279	487	735	811	699	474	378	1
1982	3,756	47	136	307	475	675	716	592	445	328	3
1981	3,777	55	129	298	459	682	753	576	457	327	1
1980	3,744	60	146	290	481	654	749	562	392	375	1
1979	3,398	58	147	248	466	596	612	540	375	310	-
Emphysema (492)											
White, both sexes											
1998	16,511	114	285	608	1,214	2,263	3,224	3,453	2,967	2,315	-
1997	16,516	125	261	578	1,233	2,286	3,260	3,543	2,853	2,287	2
1996	16,257	114	284	604	1,173	2,208	3,202	3,543	2,887	2,187	-
1995	15,928	123	319	617	1,276	2,287	3,106	3,375	2,675	2,098	-
1994	16,223	114	261	626	1,310	2,332	3,278	3,400	2,721	2,119	1
1993	16,534	104	280	660	1,452	2,457	3,334	3,557	2,698	1,950	1
1992	15,405	108	253	617	1,279	2,355	3,194	3,248	2,403	1,880	-
1991	15,539	106	288	637	1,354	2,417	3,058	3,306	2,466	1,845	1
1990	14,828	95	268	638	1,400	2,290	3,051	3,208	2,198	1,611	-
1989	14,590	93	263	716	1,438	2,351	2,925	3,095	2,148	1,502	-
1988	14,589	110	249	660	1,448	2,305	3,102	3,009	2,092	1,551	2
1987	13,663	91	278	670	1,416	2,132	2,806	2,806	2,005	1,382	3
1986	13,687	122	308	674	1,420	2,171	2,903	2,828	1,888	1,312	1
1985	13,353	118	278	696	1,404	2,177	2,865	2,721	1,772	1,252	-
1984	12,547	86	246	746	1,334	2,068	2,621	2,446	1,715	1,213	-
1983	12,588	105	319	743	1,316	2,103	2,661	2,452	1,632	1,184	4
1982	12,155	101	321	740	1,312	2,044	2,568	2,294	1,587	1,116	3
1981	12,683	119	362	786	1,373	2,255	2,692	2,301	1,589	1,128	1
1980	13,194	143	360	816	1,470	2,298	2,685	2,452	1,667	1,226	1
1979	12,887	147	398	781	1,523	2,237	2,673	2,354	1,563	1,118	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
White, male											
1998	8,684	67	140	319	677	1,199	1,745	1,841	1,514	1,141	-
1997	8,823	76	136	321	644	1,233	1,816	1,924	1,470	1,146	1
1996	8,750	72	149	316	645	1,263	1,686	1,910	1,562	1,114	-
1995	8,656	77	181	343	694	1,219	1,723	1,838	1,469	1,086	-
1994	8,952	64	150	342	692	1,300	1,843	1,911	1,489	1,126	1
1993	9,202	64	162	359	782	1,360	1,851	1,988	1,549	1,057	-
1992	8,745	70	120	346	722	1,358	1,804	1,830	1,403	1,044	-
1991	8,904	62	149	365	752	1,379	1,756	1,917	1,454	1,035	-
1990	8,769	58	130	362	805	1,368	1,802	1,909	1,360	927	-
1989	8,665	53	147	404	804	1,414	1,724	1,845	1,335	904	-
1988	8,953	59	133	375	882	1,384	1,852	1,902	1,376	952	1
1987	8,566	47	147	374	875	1,274	1,771	1,845	1,364	820	3
1986	8,781	80	175	417	850	1,350	1,897	1,868	1,262	840	1
1985	8,766	69	168	423	847	1,390	1,896	1,853	1,234	844	-
1984	8,369	51	141	455	875	1,310	1,772	1,703	1,226	791	-
1983	8,653	58	188	482	844	1,392	1,874	1,774	1,174	824	3
1982	8,533	58	192	447	861	1,394	1,874	1,717	1,154	795	-
1981	9,026	71	237	502	930	1,595	1,954	1,738	1,144	810	-
1980	9,587	86	223	538	1,010	1,668	1,961	1,907	1,284	862	-
1979	9,625	97	262	547	1,077	1,667	2,080	1,831	1,197	812	-
Emphysema (492)											
White, female											
1998	7,827	47	145	289	537	1,064	1,479	1,612	1,453	1,174	-
1997	7,693	49	125	257	589	1,053	1,444	1,619	1,383	1,141	1
1996	7,507	42	135	288	528	945	1,516	1,633	1,325	1,073	-
1995	7,272	46	138	274	582	1,068	1,383	1,537	1,206	1,012	-
1994	7,271	50	111	284	618	1,032	1,435	1,489	1,232	993	-
1993	7,332	40	118	301	670	1,097	1,483	1,569	1,149	893	1
1992	6,660	38	133	271	557	997	1,390	1,418	1,000	836	-
1991	6,635	44	139	272	602	1,038	1,302	1,389	1,012	810	1
1990	6,059	37	138	276	595	922	1,249	1,299	838	684	-
1989	5,925	40	116	312	634	937	1,201	1,250	813	598	-
1988	5,636	51	116	285	566	921	1,250	1,107	716	599	1
1987	5,097	44	131	296	541	858	1,035	961	641	562	-
1986	4,906	42	133	257	570	821	1,006	960	626	472	-
1985	4,587	49	110	273	557	787	969	868	538	408	-
1984	4,178	35	105	291	459	758	849	743	489	422	-
1983	3,935	47	131	261	472	711	787	678	458	360	1
1982	3,622	43	129	293	451	650	694	577	433	321	3
1981	3,657	48	125	284	443	660	738	563	445	318	1
1980	3,607	57	137	278	460	630	724	545	383	364	1
1979	3,262	50	136	234	446	570	593	523	366	306	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
All other, both sexes											
1998	1,044	29	48	77	101	145	203	186	111	113	-
1997	1,002	25	45	49	98	154	197	181	121	115	-
1996	1,029	27	37	77	95	135	188	191	123	128	-
1995	999	27	36	75	80	116	196	180	152	108	-
1994	992	18	33	63	108	157	185	162	123	109	-
1993	1,052	26	34	50	104	176	200	183	137	106	-
1992	958	24	31	58	98	162	186	160	119	99	-
1991	897	29	34	51	114	153	186	141	101	76	-
1990	878	25	29	66	109	164	137	156	109	66	-
1989	877	12	33	54	110	160	159	145	97	84	-
1988	924	12	31	73	113	147	184	155	113	63	-
1987	820	16	32	58	124	136	147	123	86	71	-
1986	784	23	33	79	101	133	144	115	79	56	-
1985	797	21	41	69	93	137	136	139	81	64	1
1984	716	22	41	45	89	121	132	106	81	55	-
1983	705	17	27	74	92	106	132	88	81	57	1
1982	638	19	33	62	80	126	107	102	51	40	1
1981	686	22	37	67	93	128	112	98	55	43	-
1980	683	16	39	67	109	112	119	86	57	48	-
1979	682	31	42	58	101	127	112	97	46	39	1
Emphysema (492)											
All other, male											
1998	649	18	29	48	69	94	121	116	77	56	-
1997	674	19	30	35	61	105	138	123	80	76	-
1996	672	18	24	49	64	85	127	128	82	76	-
1995	688	18	24	53	62	75	141	122	106	69	-
1994	670	15	21	37	77	94	131	104	93	73	-
1993	712	16	26	34	70	115	128	131	93	76	-
1992	677	18	20	38	78	113	121	114	91	69	-
1991	627	25	20	36	75	109	119	100	82	51	-
1990	639	19	20	41	75	119	105	114	87	46	-
1989	629	9	23	34	70	115	116	109	71	64	-
1988	733	9	20	52	76	117	152	136	93	50	-
1987	616	13	25	41	91	103	114	93	65	46	-
1986	615	13	23	58	78	99	116	104	66	40	-
1985	609	20	30	48	68	99	109	109	63	50	1
1984	557	15	34	33	63	95	110	87	65	38	-
1983	550	15	18	56	77	82	108	67	65	39	1
1982	504	15	26	48	56	101	85	87	39	33	1
1981	566	15	33	53	77	106	97	85	43	34	-
1980	546	13	30	55	88	88	94	69	48	37	-
1979	546	23	31	44	81	101	93	80	37	35	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
All other, female											
1998	395	11	19	29	32	51	82	70	34	57	-
1997	328	6	15	14	37	49	59	58	41	39	-
1996	357	9	13	28	31	50	61	63	41	52	-
1995	311	9	12	22	18	41	55	58	46	39	-
1994	322	3	12	26	31	63	54	58	30	36	-
1993	340	10	8	16	34	61	72	52	44	30	-
1992	281	6	11	20	20	49	65	46	28	30	-
1991	270	4	14	15	39	44	67	41	19	25	-
1990	239	6	9	25	34	45	32	42	22	20	-
1989	248	3	10	20	40	45	43	36	26	20	-
1988	191	3	11	21	37	30	32	19	20	13	-
1987	204	3	7	17	33	33	33	30	21	25	-
1986	169	10	10	21	23	34	28	11	13	16	-
1985	188	1	11	21	25	38	27	30	18	14	-
1984	159	7	7	12	26	26	22	19	16	17	-
1983	155	2	9	18	15	24	24	21	16	18	-
1982	134	4	7	14	24	25	22	15	12	7	-
1981	120	7	4	14	16	22	15	13	12	9	-
1980	137	3	9	12	21	24	25	17	9	11	-
1979	136	8	11	14	20	26	19	17	9	4	-
Emphysema (492)											
Black, both sexes											
1998	890	24	43	68	86	130	168	157	91	93	-
1997	847	22	38	45	86	133	168	149	99	92	-
1996	882	25	30	72	84	117	163	163	100	101	-
1995	855	25	35	70	71	102	160	154	123	87	-
1994	858	17	27	58	99	139	157	140	101	88	-
1993	888	25	33	46	92	149	162	147	109	89	-
1992	831	24	29	52	83	146	160	135	100	82	-
1991	772	27	32	44	103	128	165	119	78	65	-
1990	763	24	26	56	99	145	123	128	92	54	-
1989	765	11	32	49	95	145	133	129	79	70	-
1988	816	12	29	66	97	133	165	133	101	49	-
1987	719	16	31	55	113	122	133	101	70	52	-
1986	704	20	32	73	96	124	131	97	63	48	-
1985	687	20	36	59	81	123	121	115	60	56	1
1984	627	22	39	40	84	110	112	85	69	44	-
1983	624	17	27	70	89	97	112	75	59	48	1
1982	579	17	33	60	77	114	95	88	43	34	1
1981	643	22	35	66	88	124	101	87	51	41	-
1980	625	15	36	65	104	105	107	74	49	40	-
1979	631	31	40	53	96	118	100	92	40	32	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Emphysema (492)											
Black, male											
1998	543	14	26	42	58	84	100	94	63	42	-
1997	562	17	26	32	53	90	116	97	65	60	-
1996	576	17	21	47	56	71	111	105	65	64	-
1995	578	16	23	50	56	63	114	103	81	55	-
1994	567	15	17	34	70	83	108	86	76	55	-
1993	601	15	25	32	61	99	103	106	74	63	-
1992	585	18	19	35	67	101	104	96	75	56	-
1991	536	23	19	30	68	90	107	86	61	43	-
1990	552	18	17	35	70	102	93	94	72	39	-
1989	547	9	23	31	58	103	99	96	59	52	-
1988	648	9	19	45	66	107	138	117	85	36	-
1987	543	13	24	39	84	93	103	76	52	35	-
1986	553	12	23	54	74	94	105	88	52	34	-
1985	523	19	26	39	58	94	96	88	48	42	1
1984	480	15	32	30	59	85	90	69	55	29	-
1983	484	15	18	53	74	76	91	57	44	33	1
1982	454	13	26	46	53	91	76	75	32	28	1
1981	531	15	31	52	74	103	89	75	39	32	-
1980	504	13	27	53	86	84	85	60	41	31	-
1979	506	23	29	40	79	92	87	75	32	28	1
Emphysema (492)											
Black, female											
1998	347	10	17	26	28	46	68	63	28	51	-
1997	285	5	12	13	33	43	52	52	34	32	-
1996	306	8	9	25	28	46	52	58	35	37	-
1995	277	9	12	20	15	39	46	51	42	32	-
1994	291	2	10	24	29	56	49	54	25	33	-
1993	287	10	8	14	31	50	59	41	35	26	-
1992	246	6	10	17	16	45	56	39	25	26	-
1991	236	4	13	14	35	38	58	33	17	22	-
1990	211	6	9	21	29	43	30	34	20	15	-
1989	218	2	9	18	37	42	34	33	20	18	-
1988	168	3	10	21	31	26	27	16	16	13	-
1987	176	3	7	16	29	29	30	25	18	17	-
1986	151	8	9	19	22	30	26	9	11	14	-
1985	164	1	10	20	23	29	25	27	12	14	-
1984	147	7	7	10	25	25	22	16	14	15	-
1983	140	2	9	17	15	21	21	18	15	15	-
1982	125	4	7	14	24	23	19	13	11	6	-
1981	112	7	4	14	14	21	12	12	12	9	-
1980	121	2	9	12	18	21	22	14	8	9	-
1979	125	8	11	13	17	26	13	17	8	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
All races, both sexes											
1998	5,438	320	327	312	361	443	529	622	568	807	-
1997	5,434	315	321	365	392	456	536	589	594	751	-
1996	5,667	313	336	374	442	494	601	620	557	739	2
1995	5,637	328	335	393	392	531	616	606	589	663	-
1994	5,487	302	295	370	410	564	659	619	536	644	1
1993	5,167	254	254	353	385	558	638	544	570	623	1
1992	4,964	235	260	324	368	550	614	564	533	608	-
1991	5,106	237	273	303	434	548	607	627	534	553	-
1990	4,819	252	250	323	415	549	576	590	484	512	1
1989	4,869	208	264	329	414	585	623	646	448	473	2
1988	4,597	198	242	332	453	500	597	548	443	429	-
1987	4,360	183	260	312	397	500	581	493	380	408	1
1986	3,955	156	200	308	380	454	528	474	369	350	-
1985	3,880	156	222	319	436	486	450	460	325	332	-
1984	3,564	162	193	298	376	462	443	390	312	303	1
1983	3,561	157	227	310	376	409	427	407	286	290	-
1982	3,154	136	205	242	340	360	433	343	250	234	-
1981	3,054	126	182	265	342	417	386	334	241	229	-
1980	2,891	126	183	235	294	403	362	360	236	222	-
1979	2,598	110	172	209	293	352	343	293	206	190	-
Asthma (493)											
All races, male											
1998	2,000	119	96	105	137	186	195	220	185	181	-
1997	1,986	104	116	142	141	181	209	186	181	172	-
1996	2,075	120	128	128	161	180	223	219	168	187	1
1995	2,079	125	115	129	159	191	227	186	193	182	-
1994	2,101	111	105	133	177	214	257	204	187	147	1
1993	1,928	91	101	140	162	204	237	173	195	161	-
1992	1,869	81	88	133	141	210	218	184	190	164	-
1991	1,927	95	98	109	160	198	223	220	172	142	-
1990	1,885	97	91	128	158	214	228	223	167	138	1
1989	1,848	83	108	123	138	225	238	223	140	135	1
1988	1,822	77	92	125	183	207	240	210	144	113	-
1987	1,730	62	95	136	165	199	240	175	132	104	-
1986	1,584	59	86	135	155	166	207	190	125	98	-
1985	1,551	62	84	134	175	189	183	186	119	96	-
1984	1,467	63	86	113	165	196	163	161	126	93	1
1983	1,455	64	80	127	152	168	189	158	106	89	-
1982	1,314	50	81	93	148	158	176	130	103	64	-
1981	1,287	54	69	101	162	174	162	140	93	75	-
1980	1,292	51	72	98	134	212	163	175	102	71	-
1979	1,133	46	63	90	134	171	155	122	87	65	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
All races, female											
1998	3,438	201	231	207	224	257	334	402	383	626	-
1997	3,448	211	205	223	251	275	327	403	413	579	-
1996	3,592	193	208	246	281	314	378	401	389	552	1
1995	3,558	203	220	264	233	340	389	420	396	481	-
1994	3,386	191	190	237	233	350	402	415	349	497	-
1993	3,239	163	153	213	223	354	401	371	375	462	1
1992	3,095	154	172	191	227	340	396	380	343	444	-
1991	3,179	142	175	194	274	350	384	407	362	411	-
1990	2,934	155	159	195	257	335	348	367	317	374	-
1989	3,021	125	156	206	276	360	385	423	308	338	1
1988	2,775	121	150	207	270	293	357	338	299	316	-
1987	2,630	121	165	176	232	301	341	318	248	304	1
1986	2,371	97	114	173	225	288	321	284	244	252	-
1985	2,329	94	138	185	261	297	267	274	206	236	-
1984	2,097	99	107	185	211	266	280	229	186	210	-
1983	2,106	93	147	183	224	241	238	249	180	201	-
1982	1,840	86	124	149	192	202	257	213	147	170	-
1981	1,767	72	113	164	180	243	224	194	148	154	-
1980	1,599	75	111	137	160	191	199	185	134	151	-
1979	1,465	64	109	119	159	181	188	171	119	125	-
Asthma (493)											
White, both sexes											
1998	3,947	180	214	213	254	311	421	518	502	733	-
1997	4,002	191	215	244	263	340	432	498	522	685	-
1996	4,110	201	222	253	306	368	479	517	487	647	2
1995	4,208	200	223	275	288	404	494	515	518	605	-
1994	4,134	190	188	254	305	442	558	542	470	597	-
1993	3,910	151	178	232	281	438	540	479	508	566	1
1992	3,789	147	173	225	259	431	520	491	476	548	-
1991	3,915	144	185	215	312	437	518	557	480	516	-
1990	3,696	145	166	224	325	442	488	514	430	476	-
1989	3,761	136	172	224	328	466	523	565	399	437	1
1988	3,473	116	158	233	324	388	495	474	407	401	-
1987	3,327	112	168	216	307	402	483	442	334	374	-
1986	3,036	90	128	211	286	368	439	424	336	326	-
1985	3,026	100	152	236	352	391	394	411	293	307	-
1984	2,779	101	138	227	289	401	372	352	266	278	-
1983	2,751	91	160	229	291	337	373	368	259	270	-
1982	2,450	84	142	181	268	308	371	308	232	221	-
1981	2,426	83	131	210	284	338	333	304	220	215	-
1980	2,291	84	136	175	245	335	315	315	225	202	-
1979	2,095	74	126	156	241	305	303	267	193	173	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
White, male											
1998	1,366	69	64	71	91	131	150	178	163	162	-
1997	1,383	63	75	85	94	142	173	158	153	150	-
1996	1,426	77	87	88	108	130	174	182	146	162	1
1995	1,454	73	75	91	116	146	173	153	169	159	-
1994	1,492	64	67	95	122	172	214	182	153	135	-
1993	1,384	58	78	89	123	153	189	144	170	144	-
1992	1,362	53	57	90	96	161	178	162	166	143	-
1991	1,388	61	64	77	106	150	187	194	146	129	-
1990	1,358	55	58	80	121	166	190	188	143	122	-
1989	1,352	54	65	87	108	176	193	190	123	122	-
1988	1,299	49	53	77	129	162	196	173	128	105	-
1987	1,244	37	60	89	125	151	203	151	109	92	-
1986	1,178	34	55	94	115	131	176	170	109	90	-
1985	1,140	39	55	97	135	151	155	163	106	86	-
1984	1,106	37	61	84	118	159	139	143	108	79	-
1983	1,084	36	61	89	118	137	159	143	91	81	-
1982	983	36	55	66	111	136	146	115	94	59	-
1981	977	34	51	80	129	134	131	123	86	69	-
1980	1,008	35	56	68	110	178	142	150	96	63	-
1979	898	31	47	63	114	149	132	107	80	54	-
Asthma (493)											
White, female											
1998	2,581	111	150	142	163	180	271	340	339	571	-
1997	2,619	128	140	159	169	198	259	340	369	535	-
1996	2,684	124	135	165	198	238	305	335	341	485	1
1995	2,754	127	148	184	172	258	321	362	349	446	-
1994	2,642	126	121	159	183	270	344	360	317	462	-
1993	2,526	93	100	143	158	285	351	335	338	422	1
1992	2,427	94	116	135	163	270	342	329	310	405	-
1991	2,527	83	121	138	206	287	331	363	334	387	-
1990	2,338	90	108	144	204	276	298	326	287	354	-
1989	2,409	82	107	137	220	290	330	375	276	315	1
1988	2,174	67	105	156	195	226	299	301	279	296	-
1987	2,083	75	108	127	182	251	280	291	225	282	-
1986	1,858	56	73	117	171	237	263	254	227	236	-
1985	1,886	61	97	139	217	240	239	248	187	221	-
1984	1,673	64	77	143	171	242	233	209	158	199	-
1983	1,667	55	99	140	173	200	214	225	168	189	-
1982	1,467	48	87	115	157	172	225	193	138	162	-
1981	1,449	49	80	130	155	204	202	181	134	146	-
1980	1,283	49	80	107	135	157	173	165	129	139	-
1979	1,197	43	79	93	127	156	171	160	113	119	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
All other, both sexes											
1998	1,491	140	113	99	107	132	108	104	66	74	-
1997	1,432	124	106	121	129	116	104	91	72	66	-
1996	1,557	112	114	121	136	126	122	103	70	92	-
1995	1,429	128	112	118	104	127	122	91	71	58	-
1994	1,353	112	107	116	105	122	101	77	66	47	1
1993	1,257	103	76	121	104	120	98	65	62	57	-
1992	1,175	88	87	99	109	119	94	73	57	60	-
1991	1,191	93	88	88	122	111	89	70	54	37	-
1990	1,123	107	84	99	90	107	88	76	54	36	1
1989	1,108	72	92	105	86	119	100	81	49	36	1
1988	1,124	82	84	99	129	112	102	74	36	28	-
1987	1,033	71	92	96	90	98	98	51	46	34	1
1986	919	66	72	97	94	86	89	50	33	24	-
1985	854	56	70	83	84	95	56	49	32	25	-
1984	785	61	55	71	87	61	71	38	46	25	1
1983	810	66	67	81	85	72	54	39	27	20	-
1982	704	52	63	61	72	52	62	35	18	13	-
1981	628	43	51	55	58	79	53	30	21	14	-
1980	600	42	47	60	49	68	47	45	11	20	-
1979	503	36	46	53	52	47	40	26	13	17	-
Asthma (493)											
All other, male											
1998	634	50	32	34	46	55	45	42	22	19	-
1997	603	41	41	57	47	39	36	28	28	22	-
1996	649	43	41	40	53	50	49	37	22	25	-
1995	625	52	40	38	43	45	54	33	24	23	-
1994	609	47	38	38	55	42	43	22	34	12	1
1993	544	33	23	51	39	51	48	29	25	17	-
1992	507	28	31	43	45	49	40	22	24	21	-
1991	539	34	34	32	54	48	36	26	26	13	-
1990	527	42	33	48	37	48	38	35	24	16	1
1989	496	29	43	36	30	49	45	33	17	13	1
1988	523	28	39	48	54	45	44	37	16	8	-
1987	486	25	35	47	40	48	37	24	23	12	-
1986	406	25	31	41	40	35	31	20	16	8	-
1985	411	23	29	37	40	38	28	23	13	10	-
1984	361	26	25	29	47	37	24	18	18	14	1
1983	371	28	19	38	34	31	30	15	15	8	-
1982	331	14	26	27	37	22	30	15	9	5	-
1981	310	20	18	21	33	40	31	17	7	6	-
1980	284	16	16	30	24	34	21	25	6	8	-
1979	235	15	16	27	20	22	23	15	7	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
All other, female											
1998	857	90	81	65	61	77	63	62	44	55	-
1997	829	83	65	64	82	77	68	63	44	44	-
1996	908	69	73	81	83	76	73	66	48	67	-
1995	804	76	72	80	61	82	68	58	47	35	-
1994	744	65	69	78	50	80	58	55	32	35	-
1993	713	70	53	70	65	69	50	36	37	40	-
1992	668	60	56	56	64	70	54	51	33	39	-
1991	652	59	54	56	68	63	53	44	28	24	-
1990	596	65	51	51	53	59	50	41	30	20	-
1989	612	43	49	69	56	70	55	48	32	23	-
1988	601	54	45	51	75	67	58	37	20	20	-
1987	547	46	57	49	50	50	61	27	23	22	1
1986	513	41	41	56	54	51	58	30	17	16	-
1985	443	33	41	46	44	57	28	26	19	15	-
1984	424	35	30	42	40	24	47	20	28	11	-
1983	439	38	48	43	51	41	24	24	12	12	-
1982	373	38	37	34	35	30	32	20	9	8	-
1981	318	23	33	34	25	39	22	13	14	8	-
1980	316	26	31	30	25	34	26	20	5	12	-
1979	268	21	30	26	32	25	17	11	6	6	-
Asthma (493)											
Black, both sexes											
1998	1,290	128	106	87	90	107	87	77	52	49	-
1997	1,200	110	94	96	107	88	77	69	53	50	-
1996	1,325	99	106	100	112	99	92	81	56	74	-
1995	1,247	114	100	105	84	111	96	68	59	46	-
1994	1,186	100	91	101	95	103	76	63	49	38	1
1993	1,112	91	67	111	91	103	81	57	48	44	-
1992	1,036	79	81	84	99	97	76	59	49	47	-
1991	1,043	89	86	78	107	93	69	56	37	21	-
1990	986	102	75	82	81	90	75	52	44	28	1
1989	984	66	83	93	73	107	82	72	40	29	1
1988	1,012	74	76	92	118	101	87	61	29	26	-
1987	920	67	79	87	74	85	85	36	36	28	1
1986	828	61	65	89	87	74	80	39	27	18	-
1985	778	52	64	74	78	85	51	40	27	16	-
1984	701	54	50	65	80	55	60	34	33	18	1
1983	732	63	63	72	76	62	47	32	21	17	-
1982	647	51	56	56	66	50	56	25	16	10	-
1981	576	39	50	50	56	70	47	25	19	11	-
1980	557	38	46	55	46	62	44	41	11	14	-
1979	470	35	45	53	47	44	35	23	10	13	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Asthma (493)											
Black, male											
1998	536	45	27	27	34	42	33	29	19	11	-
1997	498	36	33	40	39	27	24	23	22	17	-
1996	540	39	38	33	43	37	32	29	14	18	-
1995	538	46	38	29	33	41	39	21	18	19	-
1994	525	41	30	32	47	33	31	19	23	9	1
1993	465	29	20	44	32	41	36	26	16	11	-
1992	433	23	26	34	40	37	28	17	19	16	-
1991	472	32	33	29	50	39	29	20	17	7	-
1990	460	41	27	39	31	40	34	20	18	11	1
1989	434	24	39	32	27	45	33	27	12	12	1
1988	460	24	35	44	48	42	36	29	10	7	-
1987	428	23	32	41	32	43	32	15	16	10	-
1986	360	23	29	38	37	27	28	14	11	3	-
1985	371	23	26	31	38	33	26	16	10	5	-
1984	312	23	21	26	40	31	18	15	14	9	1
1983	336	25	19	35	30	27	25	11	10	8	-
1982	301	13	22	24	32	21	26	10	8	2	-
1981	281	18	18	19	32	35	27	12	7	5	-
1980	260	15	15	26	22	30	19	22	6	6	-
1979	214	14	16	27	18	20	18	14	5	9	-
Asthma (493)											
Black, female											
1998	754	83	79	60	56	65	54	48	33	38	-
1997	702	74	61	56	68	61	53	46	31	33	-
1996	785	60	68	67	69	62	60	52	42	56	-
1995	709	68	62	76	51	70	57	47	41	27	-
1994	661	59	61	69	48	70	45	44	26	29	-
1993	647	62	47	67	59	62	45	31	32	33	-
1992	603	56	55	50	59	60	48	42	30	31	-
1991	571	57	53	49	57	54	40	36	20	14	-
1990	526	61	48	43	50	50	41	32	26	17	-
1989	550	42	44	61	46	62	49	45	28	17	-
1988	552	50	41	48	70	59	51	32	19	19	-
1987	492	44	47	46	42	42	53	21	20	18	1
1986	468	38	36	51	50	47	52	25	16	15	-
1985	407	29	38	43	40	52	25	24	17	11	-
1984	389	31	29	39	40	24	42	19	19	9	-
1983	396	38	44	37	46	35	22	21	11	9	-
1982	346	38	34	32	34	29	30	15	8	8	-
1981	295	21	32	31	24	35	20	13	12	6	-
1980	297	23	31	29	24	32	25	19	5	8	-
1979	256	21	29	26	29	24	17	9	5	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
All races, both sexes											
1998	86,534	441	1,162	2,408	4,798	8,994	14,618	17,831	16,921	19,025	7
1997	82,995	472	1,174	2,365	4,783	9,033	13,920	16,949	16,205	17,765	5
1996	79,884	476	1,105	2,322	4,688	8,910	13,652	16,512	15,410	16,514	10
1995	76,979	424	1,079	2,252	4,621	8,867	13,565	15,776	14,718	15,386	3
1994	75,320	418	1,115	2,271	4,849	9,199	13,346	15,439	14,227	14,192	9
1993	74,523	413	1,018	2,287	5,021	9,373	13,544	15,409	13,921	13,257	7
1992	66,933	355	903	2,216	4,762	8,542	12,115	14,201	12,000	11,578	8
1991	65,355	371	887	2,179	4,958	8,647	11,979	13,674	11,600	10,803	9
1990	62,528	346	916	2,194	4,791	8,445	11,402	13,231	10,889	10,084	7
1989	60,230	372	940	2,390	4,897	8,019	11,255	12,603	10,163	9,328	4
1988	59,009	378	947	2,289	4,867	8,141	11,176	12,203	9,988	8,781	7
1987	55,942	337	867	2,225	4,841	7,766	10,702	11,624	9,174	8,154	5
1986	54,368	341	954	2,334	4,763	7,795	10,687	11,193	8,477	7,567	7
1985	52,998	351	1,015	2,411	4,735	7,588	10,248	10,793	8,300	7,320	15
1984	48,707	316	961	2,366	4,554	7,104	9,673	9,856	7,212	6,431	7
1983	45,802	341	916	2,270	4,327	7,020	9,116	9,118	6,732	5,725	11
1982	40,419	313	868	2,110	3,870	6,332	8,159	7,933	5,795	4,787	8
1981	38,711	343	936	2,073	3,936	6,095	7,850	7,439	5,364	4,444	6
1980	35,544	354	881	2,013	3,715	5,766	7,183	6,592	4,746	4,062	5
1979	30,106	275	823	1,778	3,283	4,912	6,029	5,470	4,129	3,205	3
Other chronic lower respiratory diseases (494,496)											
All races, male											
1998	44,184	244	619	1,289	2,551	4,840	7,847	9,372	8,600	8,622	7
1997	43,026	281	635	1,232	2,556	4,995	7,594	8,908	8,338	8,300	4
1996	41,389	257	594	1,253	2,549	4,831	7,218	8,805	8,021	7,697	5
1995	40,829	226	541	1,186	2,539	4,928	7,452	8,631	7,737	7,431	3
1994	40,221	236	592	1,257	2,692	5,054	7,359	8,363	7,631	6,880	7
1993	40,629	247	567	1,240	2,802	5,219	7,520	8,563	7,607	6,694	5
1992	37,345	195	528	1,251	2,670	4,902	6,978	8,034	6,676	5,962	7
1991	37,049	202	445	1,194	2,797	5,020	6,913	7,893	6,707	5,719	8
1990	36,226	204	481	1,255	2,792	4,980	6,793	7,866	6,299	5,412	7
1989	35,068	185	490	1,359	2,852	4,785	6,586	7,578	5,942	5,127	3
1988	35,430	214	546	1,323	2,936	4,972	6,830	7,426	6,076	4,962	5
1987	34,232	173	491	1,293	2,938	4,790	6,646	7,380	5,689	4,677	3
1986	33,987	201	531	1,361	2,904	4,876	6,813	7,167	5,443	4,538	4
1985	33,618	180	582	1,403	2,960	4,850	6,684	6,975	5,392	4,442	11
1984	31,573	177	575	1,454	2,881	4,548	6,423	6,586	4,809	3,988	3
1983	30,314	180	545	1,435	2,786	4,580	6,080	6,327	4,623	3,624	9
1982	27,219	169	543	1,346	2,568	4,236	5,531	5,595	4,025	3,052	8
1981	26,541	183	571	1,339	2,581	4,162	5,532	5,339	3,745	2,952	4
1980	24,815	213	558	1,322	2,492	4,002	5,176	4,840	3,357	2,719	3
1979	21,578	176	507	1,173	2,251	3,561	4,464	4,065	3,101	2,163	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
All races, female											
1998	42,350	197	543	1,119	2,247	4,154	6,771	8,459	8,321	10,403	-
1997	39,969	191	539	1,133	2,227	4,038	6,326	8,041	7,867	9,465	1
1996	38,495	219	511	1,069	2,139	4,079	6,434	7,707	7,389	8,817	5
1995	36,150	198	538	1,066	2,082	3,939	6,113	7,145	6,981	7,955	-
1994	35,099	182	523	1,014	2,157	4,145	5,987	7,076	6,596	7,312	2
1993	33,894	166	451	1,047	2,219	4,154	6,024	6,846	6,314	6,563	2
1992	29,588	160	375	965	2,092	3,640	5,137	6,167	5,324	5,616	1
1991	28,306	169	442	985	2,161	3,627	5,066	5,781	4,893	5,084	1
1990	26,302	142	435	939	1,999	3,465	4,609	5,365	4,590	4,672	-
1989	25,162	187	450	1,031	2,045	3,234	4,669	5,025	4,221	4,201	1
1988	23,579	164	401	966	1,931	3,169	4,346	4,777	3,912	3,819	2
1987	21,710	164	376	932	1,903	2,976	4,056	4,244	3,485	3,477	2
1986	20,381	140	423	973	1,859	2,919	3,874	4,026	3,034	3,029	3
1985	19,380	171	433	1,008	1,775	2,738	3,564	3,818	2,908	2,878	4
1984	17,134	139	386	912	1,673	2,556	3,250	3,270	2,403	2,443	4
1983	15,488	161	371	835	1,541	2,440	3,036	2,791	2,109	2,101	2
1982	13,200	144	325	764	1,302	2,096	2,628	2,338	1,770	1,735	-
1981	12,170	160	365	734	1,355	1,933	2,318	2,100	1,619	1,492	2
1980	10,729	141	323	691	1,223	1,764	2,007	1,752	1,389	1,343	2
1979	8,528	99	316	605	1,032	1,351	1,565	1,405	1,028	1,042	-
Other chronic lower respiratory diseases (494,496)											
White, both sexes											
1998	80,761	353	1,006	2,121	4,355	8,318	13,627	16,722	16,063	17,945	7
1997	77,420	385	1,038	2,104	4,325	8,294	12,998	15,953	15,309	16,767	3
1996	74,595	377	985	2,066	4,251	8,234	12,729	15,548	14,580	15,594	6
1995	71,867	347	939	2,005	4,150	8,154	12,668	14,891	13,958	14,536	3
1994	70,366	342	967	2,023	4,420	8,506	12,435	14,581	13,469	13,427	4
1993	69,646	346	882	2,042	4,554	8,700	12,608	14,560	13,139	12,604	5
1992	62,603	293	783	1,962	4,354	7,875	11,337	13,444	11,395	10,974	7
1991	61,014	304	753	1,935	4,528	7,957	11,190	12,916	11,002	10,243	4
1990	58,266	278	789	1,941	4,343	7,806	10,646	12,464	10,294	9,537	5
1989	55,990	319	791	2,117	4,434	7,358	10,514	11,823	9,613	8,852	4
1988	55,107	304	803	2,052	4,430	7,535	10,464	11,518	9,482	8,341	5
1987	52,286	281	746	1,997	4,430	7,184	10,026	11,008	8,699	7,723	4
1986	50,853	282	818	2,095	4,332	7,232	10,023	10,619	8,063	7,193	6
1985	49,627	289	893	2,147	4,308	7,045	9,665	10,212	7,889	7,000	9
1984	45,571	262	832	2,108	4,167	6,631	9,076	9,384	6,818	6,120	4
1983	42,994	292	822	2,049	3,926	6,563	8,632	8,691	6,387	5,451	8
1982	37,963	263	749	1,869	3,569	5,932	7,678	7,569	5,551	4,588	7
1981	36,376	291	810	1,874	3,618	5,711	7,439	7,077	5,099	4,274	5
1980	33,368	296	770	1,803	3,427	5,412	6,789	6,279	4,550	3,881	5
1979	28,248	225	726	1,591	3,058	4,589	5,705	5,176	3,973	3,056	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
White, male											
1998	40,744	193	524	1,113	2,288	4,432	7,246	8,698	8,066	8,029	7
1997	39,642	227	554	1,083	2,281	4,524	7,033	8,279	7,762	7,756	3
1996	38,186	203	527	1,089	2,272	4,414	6,668	8,181	7,485	7,216	2
1995	37,683	170	466	1,052	2,236	4,472	6,877	8,086	7,272	6,924	3
1994	37,142	192	508	1,109	2,439	4,618	6,753	7,799	7,158	6,447	2
1993	37,480	203	481	1,090	2,503	4,789	6,909	7,998	7,084	6,294	4
1992	34,554	160	456	1,097	2,406	4,484	6,445	7,516	6,285	5,599	6
1991	34,204	165	367	1,027	2,538	4,567	6,383	7,362	6,309	5,377	4
1990	33,364	163	405	1,094	2,514	4,558	6,256	7,321	5,882	5,067	5
1989	32,195	152	399	1,185	2,557	4,331	6,073	7,006	5,572	4,823	3
1988	32,716	167	455	1,180	2,640	4,546	6,312	6,918	5,708	4,684	3
1987	31,703	134	418	1,154	2,667	4,378	6,144	6,932	5,359	4,400	2
1986	31,439	164	430	1,207	2,611	4,460	6,306	6,723	5,155	4,273	3
1985	31,228	140	501	1,235	2,666	4,462	6,236	6,546	5,096	4,238	5
1984	29,323	136	485	1,273	2,616	4,191	5,978	6,235	4,517	3,795	1
1983	28,238	148	481	1,269	2,499	4,247	5,690	5,995	4,366	3,445	6
1982	25,438	132	465	1,197	2,350	3,925	5,168	5,327	3,843	2,915	7
1981	24,787	144	486	1,200	2,354	3,861	5,209	5,064	3,536	2,824	4
1980	23,146	174	473	1,168	2,273	3,732	4,852	4,592	3,199	2,594	3
1979	20,136	136	436	1,033	2,085	3,302	4,206	3,823	2,983	2,050	2
Other chronic lower respiratory diseases (494,496)											
White, female											
1998	40,017	160	482	1,008	2,067	3,886	6,381	8,024	7,997	9,916	-
1997	37,778	158	484	1,021	2,044	3,770	5,965	7,674	7,547	9,011	-
1996	36,409	174	458	977	1,979	3,820	6,061	7,367	7,095	8,378	4
1995	34,184	177	473	953	1,914	3,682	5,791	6,805	6,686	7,612	-
1994	33,224	150	459	914	1,981	3,888	5,682	6,782	6,311	6,980	2
1993	32,166	143	401	952	2,051	3,911	5,699	6,562	6,055	6,310	1
1992	28,049	133	327	865	1,948	3,391	4,892	5,928	5,110	5,375	1
1991	26,810	139	386	908	1,990	3,390	4,807	5,554	4,693	4,866	-
1990	24,902	115	384	847	1,829	3,248	4,390	5,143	4,412	4,470	-
1989	23,795	167	392	932	1,877	3,027	4,441	4,817	4,041	4,029	1
1988	22,391	137	348	872	1,790	2,989	4,152	4,600	3,774	3,657	2
1987	20,583	147	328	843	1,763	2,806	3,882	4,076	3,340	3,323	2
1986	19,414	118	388	888	1,721	2,772	3,717	3,896	2,908	2,920	3
1985	18,399	149	392	912	1,642	2,583	3,429	3,666	2,793	2,762	4
1984	16,248	126	347	835	1,551	2,440	3,098	3,149	2,301	2,325	3
1983	14,756	144	341	780	1,427	2,316	2,942	2,696	2,021	2,006	2
1982	12,525	131	284	672	1,219	2,007	2,510	2,242	1,708	1,673	-
1981	11,589	147	324	674	1,264	1,850	2,230	2,013	1,563	1,450	1
1980	10,222	122	297	635	1,154	1,680	1,937	1,687	1,351	1,287	2
1979	8,112	89	290	558	973	1,287	1,499	1,353	990	1,006	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
All other, both sexes											
1998	5,773	88	156	287	443	676	991	1,109	858	1,080	-
1997	5,575	87	136	261	458	739	922	996	896	998	2
1996	5,289	99	120	256	437	676	923	964	830	920	4
1995	5,112	77	140	247	471	713	897	885	760	850	-
1994	4,954	76	148	248	429	693	911	858	758	765	5
1993	4,877	67	136	245	467	673	936	849	782	653	2
1992	4,330	62	120	254	408	667	778	757	605	604	1
1991	4,341	67	134	244	430	690	789	758	598	560	5
1990	4,262	68	127	253	448	639	756	767	595	547	2
1989	4,240	53	149	273	463	661	741	780	550	476	-
1988	3,902	74	144	237	437	606	712	685	506	440	2
1987	3,656	56	121	228	411	582	676	616	475	431	1
1986	3,515	59	136	239	431	563	664	574	414	374	1
1985	3,371	62	122	264	427	543	583	581	411	320	6
1984	3,136	54	129	258	387	473	597	472	394	311	3
1983	2,808	49	94	221	401	457	484	427	345	274	3
1982	2,456	50	119	241	301	400	481	364	244	199	1
1981	2,335	52	126	199	318	384	411	362	265	170	1
1980	2,176	58	111	210	288	354	394	313	196	181	-
1979	1,858	50	97	187	225	323	324	294	156	149	1
Other chronic lower respiratory diseases (494,496)											
All other, male											
1998	3,440	51	95	176	263	408	601	674	534	593	-
1997	3,384	54	81	149	275	471	561	629	576	544	1
1996	3,203	54	67	164	277	417	550	624	536	481	3
1995	3,146	56	75	134	303	456	575	545	465	507	-
1994	3,079	44	84	148	253	436	606	564	473	433	5
1993	3,149	44	86	150	299	430	611	565	523	400	1
1992	2,791	35	72	154	264	418	533	518	391	363	1
1991	2,845	37	78	167	259	453	530	531	398	342	4
1990	2,862	41	76	161	278	422	537	545	417	345	2
1989	2,873	33	91	174	295	454	513	572	370	304	-
1988	2,714	47	91	143	296	426	518	508	368	278	2
1987	2,529	39	73	139	271	412	502	448	330	277	1
1986	2,548	37	101	154	293	416	507	444	288	265	1
1985	2,390	40	81	168	294	388	448	429	296	204	6
1984	2,250	41	90	181	265	357	445	351	292	193	2
1983	2,076	32	64	166	287	333	390	332	257	179	3
1982	1,781	37	78	149	218	311	363	268	182	137	1
1981	1,754	39	85	139	227	301	323	275	209	128	-
1980	1,669	39	85	154	219	270	324	248	158	125	-
1979	1,442	40	71	140	166	259	258	242	118	113	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
All other, female											
1998	2,333	37	61	111	180	268	390	435	324	487	-
1997	2,191	33	55	112	183	268	361	367	320	454	1
1996	2,086	45	53	92	160	259	373	340	294	439	1
1995	1,966	21	65	113	168	257	322	340	295	343	-
1994	1,875	32	64	100	176	257	305	294	285	332	-
1993	1,728	23	50	95	168	243	325	284	259	253	1
1992	1,539	27	48	100	144	249	245	239	214	241	-
1991	1,496	30	56	77	171	237	259	227	200	218	1
1990	1,400	27	51	92	170	217	219	222	178	202	-
1989	1,367	20	58	99	168	207	228	208	180	172	-
1988	1,188	27	53	94	141	180	194	177	138	162	-
1987	1,127	17	48	89	140	170	174	168	145	154	-
1986	967	22	35	85	138	147	157	130	126	109	-
1985	981	22	41	96	133	155	135	152	115	116	-
1984	886	13	39	77	122	116	152	121	102	118	1
1983	732	17	30	55	114	124	94	95	88	95	-
1982	675	13	41	92	83	89	118	96	62	62	-
1981	581	13	41	60	91	83	88	87	56	42	1
1980	507	19	26	56	69	84	70	65	38	56	-
1979	416	10	26	47	59	64	66	52	38	36	-
Other chronic lower respiratory diseases (494,496)											
Black, both sexes											
1998	4,860	81	145	264	384	576	856	933	694	853	-
1997	4,665	84	120	231	406	640	801	808	713	786	1
1996	4,502	95	114	230	390	589	797	792	681	752	4
1995	4,341	73	121	218	419	624	772	744	625	682	-
1994	4,258	74	139	221	386	624	788	731	602	634	3
1993	4,212	63	129	219	414	606	791	726	653	548	1
1992	3,785	58	114	234	367	588	677	670	500	507	1
1991	3,777	66	130	225	388	611	686	650	509	450	5
1990	3,715	63	117	229	402	562	665	665	519	436	2
1989	3,714	48	135	250	418	584	665	668	471	390	-
1988	3,454	71	137	218	402	556	631	590	431	360	1
1987	3,243	49	107	209	378	528	601	541	421	355	1
1986	3,140	56	126	219	397	518	596	505	352	318	1
1985	3,027	59	110	245	390	501	525	506	370	267	6
1984	2,798	52	120	238	358	434	540	421	335	246	2
1983	2,532	48	89	209	373	413	434	386	299	234	1
1982	2,242	49	117	227	280	368	449	317	211	169	1
1981	2,118	51	117	192	293	348	375	323	231	144	1
1980	2,002	56	106	200	281	325	364	286	177	143	-
1979	1,686	47	95	179	209	302	298	270	133	106	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic lower respiratory diseases (494,496)											
Black, male											
1998	2,866	46	87	161	227	348	518	567	427	449	-
1997	2,798	51	72	130	243	409	481	513	448	410	1
1996	2,704	52	64	146	248	362	466	503	441	390	3
1995	2,673	53	67	121	275	397	489	463	391	393	-
1994	2,647	44	79	135	230	390	522	479	385	352	3
1993	2,725	42	83	138	265	389	519	484	438	329	1
1992	2,435	34	68	145	237	366	458	458	323	304	1
1991	2,478	36	76	153	239	407	456	461	336	269	4
1990	2,490	37	72	149	253	366	469	471	362	273	2
1989	2,498	29	81	161	263	399	457	487	318	242	-
1988	2,415	44	89	130	278	392	464	433	313	234	1
1987	2,248	34	67	127	251	375	446	396	290	227	1
1986	2,268	36	94	141	268	381	456	390	240	224	1
1985	2,167	37	73	159	270	362	410	376	265	175	6
1984	2,025	40	83	169	250	330	405	315	248	153	1
1983	1,873	31	61	158	267	299	351	305	221	149	1
1982	1,632	37	77	141	202	288	338	236	156	120	1
1981	1,599	38	80	133	210	277	295	250	180	109	-
1980	1,529	37	81	144	212	248	297	224	147	98	-
1979	1,329	38	70	136	158	243	241	225	104	81	1
Other chronic lower respiratory diseases (494,496)											
Black, female											
1998	1,994	35	58	103	157	228	338	366	267	404	-
1997	1,867	33	48	101	163	231	320	295	265	376	-
1996	1,798	43	50	84	142	227	331	289	240	362	1
1995	1,668	20	54	97	144	227	283	281	234	289	-
1994	1,611	30	60	86	156	234	266	252	217	282	-
1993	1,487	21	46	81	149	217	272	242	215	219	-
1992	1,350	24	46	89	130	222	219	212	177	203	-
1991	1,299	30	54	72	149	204	230	189	173	181	1
1990	1,225	26	45	80	149	196	196	194	157	163	-
1989	1,216	19	54	89	155	185	208	181	153	148	-
1988	1,039	27	48	88	124	164	167	157	118	126	-
1987	995	15	40	82	127	153	155	145	131	128	-
1986	872	20	32	78	129	137	140	115	112	94	-
1985	860	22	37	86	120	139	115	130	105	92	-
1984	773	12	37	69	108	104	135	106	87	93	1
1983	659	17	28	51	106	114	83	81	78	85	-
1982	610	12	40	86	78	80	111	81	55	49	-
1981	519	13	37	59	83	71	80	73	51	35	1
1980	473	19	25	56	69	77	67	62	30	45	-
1979	357	9	25	43	51	59	57	45	29	25	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
All races, both sexes											
1998	1,118	5	5	29	50	101	179	237	262	241	-
1997	1,111	9	12	16	50	96	193	239	238	247	-
1996	1,140	5	9	15	55	95	203	244	250	254	-
1995	1,126	3	14	28	59	113	210	242	243	212	-
1994	1,109	7	9	23	55	100	188	264	261	194	-
1993	1,157	4	9	20	68	138	184	261	256	206	-
1992	1,148	6	10	25	68	109	210	273	232	207	-
1991	1,256	8	15	30	73	136	215	325	231	208	-
1990	1,358	9	17	29	100	158	250	329	243	207	-
1989	1,279	5	9	26	81	159	261	309	249	165	-
1988	1,242	4	14	34	87	165	241	287	229	165	-
1987	1,322	5	14	37	97	158	266	330	226	176	-
1986	1,334	5	15	39	89	182	299	286	227	175	-
1985	1,386	3	13	50	99	181	309	309	239	165	-
1984	1,380	8	20	38	114	208	311	299	225	143	-
1983	1,373	13	15	41	116	231	305	274	203	152	-
1982	1,438	17	20	71	117	205	306	313	215	157	-
1981	1,629	8	21	53	140	277	342	332	230	202	-
1980	1,625	9	26	72	165	291	341	295	221	176	-
1979	1,581	13	28	76	145	302	343	283	193	170	-
Pneumoconioses and chemical effects (500-506)											
All races, male											
1998	1,075	5	5	28	48	95	175	231	253	229	-
1997	1,069	7	11	15	49	93	191	229	232	235	-
1996	1,101	4	7	14	53	95	196	229	249	246	-
1995	1,098	3	14	27	58	109	205	236	238	206	-
1994	1,069	7	9	22	55	97	181	255	253	186	-
1993	1,117	4	9	19	65	132	177	259	245	198	-
1992	1,109	5	9	25	65	103	201	267	228	200	-
1991	1,216	7	14	28	70	129	208	321	225	202	-
1990	1,311	9	17	28	94	154	241	323	235	194	-
1989	1,238	5	9	25	79	150	249	304	245	159	-
1988	1,206	4	13	34	81	163	238	276	225	158	-
1987	1,281	4	14	35	94	154	263	323	219	165	-
1986	1,291	5	15	32	86	181	294	282	221	163	-
1985	1,342	2	13	47	97	176	304	303	230	158	-
1984	1,339	7	19	36	109	201	303	295	220	138	-
1983	1,343	13	15	38	112	227	303	268	198	147	-
1982	1,404	15	18	68	114	203	302	310	211	150	-
1981	1,586	7	20	53	135	272	335	328	225	192	-
1980	1,584	7	26	69	161	286	338	291	214	171	-
1979	1,545	12	28	71	144	298	339	279	190	162	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
All races, female											
1998	43	-	-	1	2	6	4	6	9	12	-
1997	42	2	1	1	1	3	2	10	6	12	-
1996	39	1	2	1	2	-	7	15	1	8	-
1995	28	-	-	1	1	4	5	6	5	6	-
1994	40	-	-	1	-	3	7	9	8	8	-
1993	40	-	-	1	3	6	7	2	11	8	-
1992	39	1	1	-	3	6	9	6	4	7	-
1991	40	1	1	2	3	7	7	4	6	6	-
1990	47	-	-	1	6	4	9	6	8	13	-
1989	41	-	-	1	2	9	12	5	4	6	-
1988	36	-	1	-	6	2	3	11	4	7	-
1987	41	1	-	2	3	4	3	7	7	11	-
1986	43	-	-	7	3	1	5	4	6	12	-
1985	44	1	-	3	2	5	5	6	9	7	-
1984	41	1	1	2	5	7	8	4	5	5	-
1983	30	-	-	3	4	4	2	6	5	5	-
1982	34	2	2	3	3	2	4	3	4	7	-
1981	43	1	1	-	5	5	7	4	5	10	-
1980	41	2	-	3	4	5	3	4	7	5	-
1979	36	1	-	5	1	4	4	4	3	8	-
Pneumoconioses and chemical effects (500-506)											
White, both sexes											
1998	1,052	5	5	26	43	94	167	225	252	227	-
1997	1,051	6	11	12	47	85	186	227	231	237	-
1996	1,079	3	6	13	53	81	193	239	239	242	-
1995	1,063	3	12	24	57	109	199	228	233	196	-
1994	1,042	6	5	18	53	90	176	248	254	186	-
1993	1,092	2	9	19	62	130	171	253	244	196	-
1992	1,072	5	9	20	63	99	198	261	217	195	-
1991	1,167	7	12	26	66	127	201	304	220	191	-
1990	1,266	8	14	26	88	148	234	311	228	199	-
1989	1,205	5	8	22	73	145	251	297	239	153	-
1988	1,166	3	12	30	80	156	234	271	217	155	-
1987	1,231	5	12	33	91	147	243	314	213	163	-
1986	1,251	5	14	34	82	174	281	268	213	164	-
1985	1,317	3	11	47	92	169	301	299	226	155	-
1984	1,303	3	16	36	104	196	297	289	215	138	-
1983	1,283	9	14	38	105	209	294	260	193	145	-
1982	1,338	13	17	65	105	196	289	297	204	142	-
1981	1,543	8	16	51	131	263	328	317	218	193	-
1980	1,530	4	20	65	155	275	327	279	209	173	-
1979	1,486	10	23	70	137	288	328	267	187	163	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
White, male											
1998	1,018	5	5	26	41	89	165	221	243	217	-
1997	1,014	5	10	11	46	83	184	219	225	226	-
1996	1,042	2	5	12	51	81	186	224	238	235	-
1995	1,035	3	12	23	56	105	194	222	228	190	-
1994	1,006	6	5	17	53	88	170	240	246	178	-
1993	1,055	2	9	18	59	124	164	251	234	189	-
1992	1,040	5	8	20	61	94	190	257	213	188	-
1991	1,131	6	11	24	63	121	196	300	214	186	-
1990	1,221	8	14	25	83	144	226	305	220	186	-
1989	1,167	5	8	22	72	136	239	292	235	148	-
1988	1,135	3	12	30	75	154	231	261	213	149	-
1987	1,194	4	12	31	88	144	240	307	207	153	-
1986	1,214	5	14	29	79	173	277	265	207	153	-
1985	1,278	2	11	44	90	165	297	293	217	149	-
1984	1,266	3	16	34	99	189	289	286	210	133	-
1983	1,256	9	14	35	101	206	292	255	189	140	-
1982	1,312	12	16	62	103	194	286	295	200	136	-
1981	1,506	7	16	51	126	259	322	313	215	183	-
1980	1,493	3	20	62	151	271	325	275	202	168	-
1979	1,460	9	23	65	136	286	325	264	185	156	-
Pneumoconioses and chemical effects (500-506)											
White, female											
1998	34	-	-	-	2	5	2	4	9	10	-
1997	37	1	1	1	1	2	2	8	6	11	-
1996	37	1	1	1	2	-	7	15	1	7	-
1995	28	-	-	1	1	4	5	6	5	6	-
1994	36	-	-	1	-	2	6	8	8	8	-
1993	37	-	-	1	3	6	7	2	10	7	-
1992	32	-	1	-	2	5	8	4	4	7	-
1991	36	1	1	2	3	6	5	4	6	5	-
1990	45	-	-	1	5	4	8	6	8	13	-
1989	38	-	-	-	1	9	12	5	4	5	-
1988	31	-	-	-	5	2	3	10	4	6	-
1987	37	1	-	2	3	3	3	7	6	10	-
1986	37	-	-	5	3	1	4	3	6	11	-
1985	39	1	-	3	2	4	4	6	9	6	-
1984	37	-	-	2	5	7	8	3	5	5	-
1983	27	-	-	3	4	3	2	5	4	5	-
1982	26	1	1	3	2	2	3	2	4	6	-
1981	37	1	-	-	5	4	6	4	3	10	-
1980	37	1	-	3	4	4	2	4	7	5	-
1979	26	1	-	5	1	2	3	3	2	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
All other, both sexes											
1998	66	-	-	3	7	7	12	12	10	14	-
1997	60	3	1	4	3	11	7	12	7	10	-
1996	61	2	3	2	2	14	10	5	11	12	-
1995	63	-	2	4	2	4	11	14	10	16	-
1994	67	1	4	5	2	10	12	16	7	8	-
1993	65	2	-	1	6	8	13	8	12	10	-
1992	76	1	1	5	5	10	12	12	15	12	-
1991	89	1	3	4	7	9	14	21	11	17	-
1990	92	1	3	3	12	10	16	18	15	8	-
1989	74	-	1	4	8	14	10	12	10	12	-
1988	76	1	2	4	7	9	7	16	12	10	-
1987	91	-	2	4	6	11	23	16	13	13	-
1986	83	-	1	5	7	8	18	18	14	11	-
1985	69	-	2	3	7	12	8	10	13	10	-
1984	77	5	4	2	10	12	14	10	10	5	-
1983	90	4	1	3	11	22	11	14	10	7	-
1982	100	4	3	6	12	9	17	16	11	15	-
1981	86	-	5	2	9	14	14	15	12	9	-
1980	95	5	6	7	10	16	14	16	12	3	-
1979	95	3	5	6	8	14	15	16	6	7	-
Pneumoconioses and chemical effects (500-506)											
All other, male											
1998	57	-	-	2	7	6	10	10	10	12	-
1997	55	2	1	4	3	10	7	10	7	9	-
1996	59	2	2	2	2	14	10	5	11	11	-
1995	63	-	2	4	2	4	11	14	10	16	-
1994	63	1	4	5	2	9	11	15	7	8	-
1993	62	2	-	1	6	8	13	8	11	9	-
1992	69	-	1	5	4	9	11	10	15	12	-
1991	85	1	3	4	7	8	12	21	11	16	-
1990	90	1	3	3	11	10	15	18	15	8	-
1989	71	-	1	3	7	14	10	12	10	11	-
1988	71	1	1	4	6	9	7	15	12	9	-
1987	87	-	2	4	6	10	23	16	12	12	-
1986	77	-	1	3	7	8	17	17	14	10	-
1985	64	-	2	3	7	11	7	10	13	9	-
1984	73	4	3	2	10	12	14	9	10	5	-
1983	87	4	1	3	11	21	11	13	9	7	-
1982	92	3	2	6	11	9	16	15	11	14	-
1981	80	-	4	2	9	13	13	15	10	9	-
1980	91	4	6	7	10	15	13	16	12	3	-
1979	85	3	5	6	8	12	14	15	5	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
All other, female											
1998	9	-	-	1	-	1	2	2	-	2	-
1997	5	1	-	-	-	1	-	2	-	1	-
1996	2	-	1	-	-	-	-	-	-	1	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	4	-	-	-	-	1	1	1	-	-	-
1993	3	-	-	-	-	-	-	-	1	1	-
1992	7	1	-	-	1	1	1	2	-	-	-
1991	4	-	-	-	-	1	2	-	-	1	-
1990	2	-	-	-	1	-	1	-	-	-	-
1989	3	-	-	1	1	-	-	-	-	1	-
1988	5	-	1	-	1	-	-	1	-	1	-
1987	4	-	-	-	-	1	-	-	1	1	-
1986	6	-	-	2	-	-	1	1	-	1	-
1985	5	-	-	-	-	1	1	-	-	1	-
1984	4	1	1	-	-	-	-	1	-	-	-
1983	3	-	-	-	-	1	-	1	1	-	-
1982	8	1	1	-	1	-	1	1	-	1	-
1981	6	-	1	-	-	1	1	-	2	-	-
1980	4	1	-	-	-	1	1	-	-	-	-
1979	10	-	-	-	-	2	1	1	1	1	-
Pneumoconioses and chemical effects (500-506)											
Black, both sexes											
1998	59	-	-	3	7	6	10	10	9	13	-
1997	53	2	1	4	3	9	6	11	6	9	-
1996	54	2	2	2	2	11	10	4	11	10	-
1995	59	-	1	4	1	4	10	14	10	15	-
1994	64	1	4	5	2	9	11	15	7	8	-
1993	59	2	-	1	6	5	12	8	10	10	-
1992	70	1	1	5	5	9	11	11	13	11	-
1991	88	1	3	4	7	9	13	21	11	17	-
1990	89	1	3	3	12	9	15	18	14	8	-
1989	69	-	1	4	6	13	10	11	10	11	-
1988	74	1	2	4	7	9	6	16	12	9	-
1987	89	-	2	3	6	10	23	16	13	13	-
1986	74	-	1	4	7	7	17	16	14	8	-
1985	66	-	2	3	6	12	8	9	13	9	-
1984	73	5	4	2	8	11	14	10	10	5	-
1983	89	4	1	3	11	21	11	14	10	7	-
1982	94	3	2	5	12	9	16	16	11	15	-
1981	81	-	5	2	9	13	13	14	11	9	-
1980	92	5	5	6	10	16	14	16	12	3	-
1979	94	3	5	6	7	14	15	16	6	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumoconioses and chemical effects (500-506)											
Black, male											
1998	53	-	-	2	7	6	8	9	9	12	-
1997	48	1	1	4	3	8	6	9	6	8	-
1996	53	2	1	2	2	11	10	4	11	10	-
1995	59	-	1	4	1	4	10	14	10	15	-
1994	61	1	4	5	2	9	10	14	7	8	-
1993	56	2	-	1	6	5	12	8	9	9	-
1992	64	-	1	5	4	8	10	10	13	11	-
1991	84	1	3	4	7	8	11	21	11	16	-
1990	88	1	3	3	11	9	15	18	14	8	-
1989	66	-	1	3	5	13	10	11	10	10	-
1988	69	1	1	4	6	9	6	15	12	8	-
1987	85	-	2	3	6	9	23	16	12	12	-
1986	71	-	1	3	7	7	16	15	14	8	-
1985	61	-	2	3	6	11	7	9	13	8	-
1984	70	4	3	2	8	11	14	9	10	5	-
1983	86	4	1	3	11	20	11	13	9	7	-
1982	88	3	1	5	11	9	15	15	11	14	-
1981	76	-	4	2	9	12	12	14	9	9	-
1980	88	4	5	6	10	15	13	16	12	3	-
1979	84	3	5	6	7	12	14	15	5	6	-
Pneumoconioses and chemical effects (500-506)											
Black, female											
1998	6	-	-	1	-	-	2	1	-	1	-
1997	5	1	-	-	-	1	-	2	-	1	-
1996	1	-	1	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	3	-	-	-	-	-	1	1	-	-	-
1993	3	-	-	-	-	-	-	-	1	1	-
1992	6	1	-	-	1	1	1	1	-	-	-
1991	4	-	-	-	-	1	2	-	-	1	-
1990	1	-	-	-	1	-	-	-	-	-	-
1989	3	-	-	1	1	-	-	-	-	1	-
1988	5	-	1	-	1	-	-	1	-	1	-
1987	4	-	-	-	-	1	-	-	1	1	-
1986	3	-	-	1	-	-	1	1	-	-	-
1985	5	-	-	-	-	1	1	-	-	1	-
1984	3	1	1	-	-	-	-	1	-	-	-
1983	3	-	-	-	-	1	-	1	1	-	-
1982	6	-	1	-	1	-	1	1	-	1	-
1981	5	-	1	-	-	1	1	-	2	-	-
1980	4	1	-	-	-	1	1	-	-	-	-
1979	10	-	-	-	-	2	1	1	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
All races, both sexes											
1998	12,674	87	135	168	242	476	987	1,745	2,593	6,023	-
1997	11,443	88	117	165	250	480	904	1,522	2,276	5,430	-
1996	10,310	65	104	146	207	433	843	1,456	2,024	4,813	-
1995	9,608	88	82	136	233	426	774	1,235	1,956	4,460	1
1994	8,969	85	69	123	205	421	760	1,216	1,754	4,099	-
1993	8,705	62	72	115	180	401	683	1,202	1,696	4,058	3
1992	8,196	60	77	107	223	371	697	1,122	1,569	3,744	-
1991	8,063	64	80	127	222	411	670	1,164	1,540	3,575	1
1990	7,702	48	72	119	251	411	653	1,060	1,528	3,358	1
1989	7,714	60	64	146	206	425	693	1,079	1,531	3,301	1
1988	7,755	64	84	143	267	448	673	1,163	1,544	3,147	1
1987	7,317	50	89	140	245	424	686	1,068	1,402	3,003	-
1986	6,783	56	89	131	256	377	653	1,070	1,296	2,650	2
1985	6,223	58	92	139	250	368	609	956	1,121	2,400	1
1984	5,448	65	86	135	256	389	526	749	994	2,023	2
1983	4,598	36	94	139	206	338	447	689	816	1,617	-
1982	3,998	48	97	123	186	289	399	612	724	1,304	1
1981	3,651	53	79	125	177	269	379	502	664	1,168	2
1980	3,355	45	80	143	197	285	368	478	552	989	-
1979	2,836	63	93	115	165	220	300	384	508	740	1
Pneumonitis due to solids and liquids (507)											
All races, male											
1998	6,190	49	88	107	147	296	608	980	1,335	2,462	-
1997	5,635	60	76	97	144	277	546	917	1,198	2,200	-
1996	4,986	40	68	91	131	263	501	823	1,061	1,886	-
1995	4,839	56	57	93	156	277	460	745	1,029	1,837	1
1994	4,512	58	38	65	143	276	490	705	948	1,640	-
1993	4,285	40	49	72	111	253	431	721	886	1,571	1
1992	4,049	38	48	71	140	243	441	654	775	1,510	-
1991	4,021	44	45	83	148	270	412	667	803	1,439	-
1990	3,795	33	46	73	177	267	423	591	739	1,330	-
1989	3,886	42	42	90	140	292	445	610	794	1,302	1
1988	3,945	35	50	94	175	309	443	679	767	1,260	-
1987	3,714	33	53	87	169	279	418	622	729	1,208	-
1986	3,625	40	60	91	170	249	422	637	668	1,161	1
1985	3,249	39	61	90	185	228	363	556	552	1,040	1
1984	2,967	46	61	73	175	254	347	440	511	928	1
1983	2,578	25	66	90	137	230	300	416	435	762	-
1982	2,268	36	64	88	127	203	256	356	381	632	1
1981	2,142	38	54	84	130	182	241	319	380	582	-
1980	1,912	30	61	100	126	194	218	275	305	489	-
1979	1,679	46	66	84	116	148	196	217	307	358	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
All races, female											
1998	6,484	38	47	61	95	180	379	765	1,258	3,561	-
1997	5,808	28	41	68	106	203	358	605	1,078	3,230	-
1996	5,324	25	36	55	76	170	342	633	963	2,927	-
1995	4,769	32	25	43	77	149	314	490	927	2,623	-
1994	4,457	27	31	58	62	145	270	511	806	2,459	-
1993	4,420	22	23	43	69	148	252	481	810	2,487	2
1992	4,147	22	29	36	83	128	256	468	794	2,234	-
1991	4,042	20	35	44	74	141	258	497	737	2,136	1
1990	3,907	15	26	46	74	144	230	469	789	2,028	1
1989	3,828	18	22	56	66	133	248	469	737	1,999	-
1988	3,810	29	34	49	92	139	230	484	777	1,887	1
1987	3,603	17	36	53	76	145	268	446	673	1,795	-
1986	3,158	16	29	40	86	128	231	433	628	1,489	1
1985	2,974	19	31	49	65	140	246	400	569	1,360	-
1984	2,481	19	25	62	81	135	179	309	483	1,095	1
1983	2,020	11	28	49	69	108	147	273	381	855	-
1982	1,730	12	33	35	59	86	143	256	343	672	-
1981	1,509	15	25	41	47	87	138	183	284	586	2
1980	1,443	15	19	43	71	91	150	203	247	500	-
1979	1,157	17	27	31	49	72	104	167	201	382	1
Pneumonitis due to solids and liquids (507)											
White, both sexes											
1998	11,191	69	100	125	199	363	822	1,529	2,338	5,491	-
1997	10,116	66	93	129	194	394	754	1,342	2,040	4,962	-
1996	9,074	48	80	116	161	346	715	1,262	1,831	4,367	-
1995	8,418	61	63	93	187	333	651	1,076	1,747	4,055	1
1994	7,833	66	42	97	160	338	644	1,045	1,548	3,716	-
1993	7,663	48	50	85	133	322	583	1,034	1,499	3,737	2
1992	7,188	43	65	80	168	292	574	992	1,416	3,412	-
1991	7,043	47	59	99	180	334	563	1,002	1,374	3,247	-
1990	6,683	34	48	90	189	328	541	924	1,351	3,033	1
1989	6,650	41	44	104	153	318	587	913	1,367	2,983	1
1988	6,731	40	62	108	203	362	563	986	1,398	2,866	-
1987	6,331	31	50	110	189	339	580	905	1,235	2,753	-
1986	5,813	38	60	92	194	294	541	907	1,157	2,392	2
1985	5,363	40	58	105	188	287	495	823	997	2,198	1
1984	4,689	40	57	105	194	302	442	662	877	1,852	1
1983	3,930	22	65	100	147	280	356	606	721	1,481	-
1982	3,398	30	83	87	145	234	333	530	637	1,179	-
1981	3,077	39	50	91	139	220	301	427	587	1,063	1
1980	2,798	32	57	105	141	221	287	407	502	901	-
1979	2,384	39	61	88	124	169	258	337	467	676	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
White, male											
1998	5,450	40	64	77	119	226	510	871	1,206	2,249	-
1997	4,958	46	64	73	113	224	459	810	1,078	2,008	-
1996	4,348	28	55	73	101	207	425	707	966	1,702	-
1995	4,217	39	42	61	124	226	387	649	914	1,685	1
1994	3,916	44	26	53	111	222	416	606	835	1,493	-
1993	3,727	30	32	54	83	205	368	619	784	1,442	-
1992	3,508	26	41	50	109	197	365	580	697	1,361	-
1991	3,478	32	30	69	126	219	341	579	720	1,290	-
1990	3,250	22	32	54	131	210	340	514	663	1,203	-
1989	3,282	26	27	63	101	222	372	511	700	1,170	1
1988	3,369	20	35	70	132	251	374	565	692	1,144	-
1987	3,165	19	27	67	131	222	359	530	634	1,102	-
1986	3,081	27	40	62	127	191	353	544	591	1,057	1
1985	2,748	26	37	67	138	181	302	474	488	939	1
1984	2,508	25	40	55	134	201	293	384	447	840	-
1983	2,162	16	43	66	95	191	236	361	379	691	-
1982	1,905	20	55	61	95	164	216	309	341	564	-
1981	1,792	30	34	59	103	144	190	276	337	530	-
1980	1,579	22	44	70	88	152	177	232	277	440	-
1979	1,388	25	41	62	90	113	169	188	281	328	-
Pneumonitis due to solids and liquids (507)											
White, female											
1998	5,741	29	36	48	80	137	312	658	1,132	3,242	-
1997	5,158	20	29	56	81	170	295	532	962	2,954	-
1996	4,726	20	25	43	60	139	290	555	865	2,665	-
1995	4,201	22	21	32	63	107	264	427	833	2,370	-
1994	3,917	22	16	44	49	116	228	439	713	2,223	-
1993	3,936	18	18	31	50	117	215	415	715	2,295	2
1992	3,680	17	24	30	59	95	209	412	719	2,051	-
1991	3,565	15	29	30	54	115	222	423	654	1,957	-
1990	3,433	12	16	36	58	118	201	410	688	1,830	1
1989	3,368	15	17	41	52	96	215	402	667	1,813	-
1988	3,362	20	27	38	71	111	189	421	706	1,722	-
1987	3,166	12	23	43	58	117	221	375	601	1,651	-
1986	2,732	11	20	30	67	103	188	363	566	1,335	1
1985	2,615	14	21	38	50	106	193	349	509	1,259	-
1984	2,181	15	17	50	60	101	149	278	430	1,012	1
1983	1,768	6	22	34	52	89	120	245	342	790	-
1982	1,493	10	28	26	50	70	117	221	296	615	-
1981	1,285	9	16	32	36	76	111	151	250	533	1
1980	1,219	10	13	35	53	69	110	175	225	461	-
1979	996	14	20	26	34	56	89	149	186	348	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
All other, both sexes											
1998	1,483	18	35	43	43	113	165	216	255	532	-
1997	1,327	22	24	36	56	86	150	180	236	468	-
1996	1,236	17	24	30	46	87	128	194	193	446	-
1995	1,190	27	19	43	46	93	123	159	209	405	-
1994	1,136	19	27	26	45	83	116	171	206	383	-
1993	1,042	14	22	30	47	79	100	168	197	321	1
1992	1,008	17	12	27	55	79	123	130	153	332	-
1991	1,020	17	21	28	42	77	107	162	166	328	1
1990	1,019	14	24	29	62	83	112	136	177	325	-
1989	1,064	19	20	42	53	107	106	166	164	318	-
1988	1,024	24	22	35	64	86	110	177	146	281	1
1987	986	19	39	30	56	85	106	163	167	250	-
1986	970	18	29	39	62	83	112	163	139	258	-
1985	860	18	34	34	62	81	114	133	124	202	-
1984	759	25	29	30	62	87	84	87	117	171	1
1983	668	14	29	39	59	58	91	83	95	136	-
1982	600	18	14	36	41	55	66	82	87	125	1
1981	574	14	29	34	38	49	78	75	77	105	1
1980	557	13	23	38	56	64	81	71	50	88	-
1979	452	24	32	27	41	51	42	47	41	64	-
Pneumonitis due to solids and liquids (507)											
All other, male											
1998	740	9	24	30	28	70	98	109	129	213	-
1997	677	14	12	24	31	53	87	107	120	192	-
1996	638	12	13	18	30	56	76	116	95	184	-
1995	622	17	15	32	32	51	73	96	115	152	-
1994	596	14	12	12	32	54	74	99	113	147	-
1993	558	10	17	18	28	48	63	102	102	129	1
1992	541	12	7	21	31	46	76	74	78	149	-
1991	543	12	15	14	22	51	71	88	83	149	-
1990	545	11	14	19	46	57	83	77	76	127	-
1989	604	16	15	27	39	70	73	99	94	132	-
1988	576	15	15	24	43	58	69	114	75	116	-
1987	549	14	26	20	38	57	59	92	95	106	-
1986	544	13	20	29	43	58	69	93	77	104	-
1985	501	13	24	23	47	47	61	82	64	101	-
1984	459	21	21	18	41	53	54	56	64	88	1
1983	416	9	23	24	42	39	64	55	56	71	-
1982	363	16	9	27	32	39	40	47	40	68	1
1981	350	8	20	25	27	38	51	43	43	52	-
1980	333	8	17	30	38	42	41	43	28	49	-
1979	291	21	25	22	26	35	27	29	26	30	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
All other, female											
1998	743	9	11	13	15	43	67	107	126	319	-
1997	650	8	12	12	25	33	63	73	116	276	-
1996	598	5	11	12	16	31	52	78	98	262	-
1995	568	10	4	11	14	42	50	63	94	253	-
1994	540	5	15	14	13	29	42	72	93	236	-
1993	484	4	5	12	19	31	37	66	95	192	-
1992	467	5	5	6	24	33	47	56	75	183	-
1991	477	5	6	14	20	26	36	74	83	179	1
1990	474	3	10	10	16	26	29	59	101	198	-
1989	460	3	5	15	14	37	33	67	70	186	-
1988	448	9	7	11	21	28	41	63	71	165	1
1987	437	5	13	10	18	28	47	71	72	144	-
1986	426	5	9	10	19	25	43	70	62	154	-
1985	359	5	10	11	15	34	53	51	60	101	-
1984	300	4	8	12	21	34	30	31	53	83	-
1983	252	5	6	15	17	19	27	28	39	65	-
1982	237	2	5	9	9	16	26	35	47	57	-
1981	224	6	9	9	11	11	27	32	34	53	1
1980	224	5	6	8	18	22	40	28	22	39	-
1979	161	3	7	5	15	16	15	18	15	34	-
Pneumonitis due to solids and liquids (507)											
Black, both sexes											
1998	1,311	18	33	40	42	106	145	187	225	462	-
1997	1,168	19	22	31	51	75	134	164	206	405	-
1996	1,080	17	23	28	43	77	109	178	173	373	-
1995	1,042	26	16	37	40	80	104	145	182	352	-
1994	1,004	18	27	26	42	78	100	150	184	327	-
1993	919	12	19	30	39	70	88	148	174	280	-
1992	913	17	12	26	54	72	122	111	139	284	-
1991	887	17	20	24	38	71	95	139	136	280	1
1990	913	13	23	28	59	74	106	123	151	285	-
1989	965	18	19	39	51	102	93	148	145	285	-
1988	924	23	22	33	61	77	102	159	131	244	1
1987	883	19	36	27	49	82	96	153	152	204	-
1986	876	15	26	37	58	78	103	154	120	222	-
1985	796	16	32	33	58	78	108	126	117	174	-
1984	689	23	27	28	59	83	81	81	108	140	1
1983	616	11	29	37	56	55	84	78	86	119	-
1982	537	14	10	34	40	53	61	74	81	99	1
1981	527	14	28	33	35	49	74	69	70	91	1
1980	511	13	22	38	53	58	74	62	45	76	-
1979	422	23	29	26	40	50	42	45	37	53	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pneumonitis due to solids and liquids (507)											
Black, male											
1998	650	9	22	27	28	67	86	92	115	179	-
1997	586	11	12	22	26	46	77	100	101	157	-
1996	553	12	13	17	29	51	68	105	85	141	-
1995	526	16	13	28	29	43	60	83	100	119	-
1994	517	13	12	12	31	52	63	85	99	114	-
1993	479	8	16	18	23	43	54	89	86	104	-
1992	481	12	7	21	30	41	75	60	70	121	-
1991	470	12	15	12	20	49	62	77	66	123	-
1990	482	11	13	18	45	49	78	68	63	107	-
1989	535	15	14	25	37	66	62	87	80	113	-
1988	516	15	15	23	41	55	63	102	63	95	-
1987	491	14	25	20	32	55	53	84	83	87	-
1986	476	10	17	28	41	54	60	85	62	84	-
1985	463	11	23	22	44	44	59	78	61	86	-
1984	411	20	19	16	40	51	51	53	58	67	1
1983	382	7	23	24	41	39	60	50	48	59	-
1982	322	13	5	25	31	37	37	43	39	52	1
1981	317	8	19	24	24	38	47	39	38	43	-
1980	309	8	17	30	37	39	36	38	25	44	-
1979	270	20	22	21	26	35	27	28	23	22	-
Pneumonitis due to solids and liquids (507)											
Black, female											
1998	661	9	11	13	14	39	59	95	110	283	-
1997	582	8	10	9	25	29	57	64	105	248	-
1996	527	5	10	11	14	26	41	73	88	232	-
1995	516	10	3	9	11	37	44	62	82	233	-
1994	487	5	15	14	11	26	37	65	85	213	-
1993	440	4	3	12	16	27	34	59	88	176	-
1992	432	5	5	5	24	31	47	51	69	163	-
1991	417	5	5	12	18	22	33	62	70	157	1
1990	431	2	10	10	14	25	28	55	88	178	-
1989	430	3	5	14	14	36	31	61	65	172	-
1988	408	8	7	10	20	22	39	57	68	149	1
1987	392	5	11	7	17	27	43	69	69	117	-
1986	400	5	9	9	17	24	43	69	58	138	-
1985	333	5	9	11	14	34	49	48	56	88	-
1984	278	3	8	12	19	32	30	28	50	73	-
1983	234	4	6	13	15	16	24	28	38	60	-
1982	215	1	5	9	9	16	24	31	42	47	-
1981	210	6	9	9	11	11	27	30	32	48	1
1980	202	5	5	8	16	19	38	24	20	32	-
1979	152	3	7	5	14	15	15	17	14	31	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All races, both sexes											
1998	20,330	406	539	787	1,132	1,798	2,750	3,505	3,431	4,679	3
1997	19,475	424	515	701	1,074	1,806	2,753	3,238	3,341	4,341	1
1996	18,727	396	477	697	1,074	1,805	2,593	3,128	3,042	4,245	3
1995	18,022	360	516	668	1,080	1,764	2,549	2,962	2,968	3,836	1
1994	17,802	380	469	631	1,086	1,835	2,545	2,882	2,921	3,727	2
1993	17,628	335	469	660	1,129	1,727	2,525	2,945	2,714	3,632	1
1992	16,752	343	396	621	1,070	1,849	2,384	2,714	2,548	3,336	1
1991	17,001	387	451	663	1,164	1,821	2,368	2,653	2,550	3,413	2
1990	16,355	331	410	610	1,235	1,730	2,276	2,551	2,443	3,178	-
1989	16,516	330	440	728	1,176	1,743	2,346	2,535	2,415	3,125	2
1988	16,882	351	461	746	1,195	1,853	2,251	2,735	2,436	3,173	1
1987	16,063	289	430	715	1,209	1,729	2,205	2,543	2,271	3,010	2
1986	15,883	304	443	751	1,237	1,735	2,166	2,465	2,196	2,838	5
1985	15,725	310	443	785	1,244	1,703	2,178	2,339	2,195	2,845	3
1984	14,346	298	414	761	1,150	1,538	1,972	2,109	1,916	2,536	4
1983	14,229	283	455	748	1,164	1,518	1,942	2,099	1,866	2,480	5
1982	13,076	259	477	783	1,117	1,492	1,769	1,866	1,667	2,093	3
1981	12,930	301	490	751	1,089	1,441	1,712	1,804	1,635	1,998	4
1980	12,552	326	496	769	1,049	1,419	1,624	1,713	1,543	1,904	2
1979	11,636	312	465	770	1,033	1,282	1,496	1,598	1,390	1,597	2
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All races, male											
1998	10,172	211	323	455	643	1,092	1,534	1,879	1,575	1,724	3
1997	9,766	228	261	395	597	1,038	1,569	1,752	1,637	1,585	-
1996	9,421	206	264	400	625	1,047	1,489	1,640	1,483	1,560	1
1995	9,003	185	267	349	620	1,041	1,438	1,586	1,406	1,402	1
1994	9,035	201	257	364	648	1,073	1,459	1,536	1,379	1,362	2
1993	8,921	169	237	375	656	1,013	1,411	1,585	1,286	1,328	1
1992	8,618	190	236	342	621	1,102	1,354	1,431	1,237	1,247	-
1991	8,721	227	250	383	681	1,054	1,386	1,434	1,226	1,206	2
1990	8,410	190	235	360	747	1,054	1,295	1,371	1,128	1,132	-
1989	8,562	192	258	448	698	1,023	1,336	1,363	1,114	1,155	-
1988	8,796	204	270	446	715	1,122	1,247	1,436	1,167	1,228	1
1987	8,456	159	248	418	723	1,009	1,280	1,419	1,102	1,147	1
1986	8,423	188	263	434	740	1,004	1,235	1,307	1,081	1,088	3
1985	8,445	182	259	482	734	1,038	1,265	1,283	1,077	1,147	1
1984	7,719	159	224	460	713	896	1,148	1,190	926	1,041	3
1983	7,665	163	272	430	704	917	1,114	1,162	945	981	5
1982	7,252	151	284	485	671	905	1,063	1,048	888	862	2
1981	7,193	169	310	466	673	867	1,026	1,001	831	858	1
1980	7,205	197	319	477	672	889	1,015	971	847	814	2
1979	6,726	191	298	488	652	805	943	965	749	665	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All races, female											
1998	10,158	195	216	332	489	706	1,216	1,626	1,856	2,955	-
1997	9,709	196	254	306	477	768	1,184	1,486	1,704	2,756	1
1996	9,306	190	213	297	449	758	1,104	1,488	1,559	2,685	2
1995	9,019	175	249	319	460	723	1,111	1,376	1,562	2,434	-
1994	8,767	179	212	267	438	762	1,086	1,346	1,542	2,365	-
1993	8,707	166	232	285	473	714	1,114	1,360	1,428	2,304	-
1992	8,134	153	160	279	449	747	1,030	1,283	1,311	2,089	1
1991	8,280	160	201	280	483	767	982	1,219	1,324	2,207	-
1990	7,945	141	175	250	488	676	981	1,180	1,315	2,046	-
1989	7,954	138	182	280	478	720	1,010	1,172	1,301	1,970	2
1988	8,086	147	191	300	480	731	1,004	1,299	1,269	1,945	-
1987	7,607	130	182	297	486	720	925	1,124	1,169	1,863	1
1986	7,460	116	180	317	497	731	931	1,158	1,115	1,750	2
1985	7,280	128	184	303	510	665	913	1,056	1,118	1,698	2
1984	6,627	139	190	301	437	642	824	919	990	1,495	1
1983	6,564	120	183	318	460	601	828	937	921	1,499	-
1982	5,824	108	193	298	446	587	706	818	779	1,231	1
1981	5,737	132	180	285	416	574	686	803	804	1,140	3
1980	5,347	129	177	292	377	530	609	742	696	1,090	-
1979	4,910	121	167	282	381	477	553	633	641	932	1
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
White, both sexes											
1998	18,012	301	419	630	936	1,559	2,477	3,220	3,201	4,365	1
1997	17,269	303	403	559	903	1,571	2,492	2,998	3,114	4,058	-
1996	16,555	276	366	566	894	1,594	2,337	2,875	2,816	3,977	1
1995	15,932	269	382	514	910	1,536	2,337	2,725	2,764	3,597	1
1994	15,664	274	361	496	922	1,615	2,268	2,652	2,716	3,477	1
1993	15,484	228	361	517	945	1,530	2,284	2,700	2,527	3,409	-
1992	14,642	241	301	485	896	1,613	2,148	2,507	2,343	3,113	-
1991	14,869	257	358	537	994	1,597	2,128	2,456	2,349	3,183	-
1990	14,172	245	313	476	1,034	1,503	2,046	2,322	2,260	2,943	-
1989	14,226	226	327	569	973	1,513	2,065	2,310	2,226	2,907	-
1988	14,626	261	338	607	997	1,607	2,018	2,476	2,256	2,942	-
1987	13,931	197	319	571	1,007	1,511	1,977	2,334	2,097	2,810	1
1986	13,751	209	339	592	1,060	1,520	1,935	2,231	2,048	2,643	5
1985	13,552	213	323	618	1,037	1,473	1,941	2,144	2,038	2,658	2
1984	12,431	195	312	619	977	1,366	1,749	1,917	1,779	2,369	1
1983	12,350	198	359	612	956	1,343	1,749	1,916	1,730	2,309	4
1982	11,221	190	362	616	939	1,294	1,578	1,694	1,534	1,968	2
1981	11,069	207	375	590	924	1,266	1,551	1,647	1,506	1,857	2
1980	10,704	224	354	610	882	1,238	1,463	1,545	1,423	1,787	-
1979	9,928	222	335	615	881	1,133	1,349	1,442	1,297	1,491	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
White, male											
1998	9,010	152	262	361	534	975	1,400	1,733	1,474	1,605	1
1997	8,666	158	213	321	515	909	1,434	1,624	1,533	1,472	-
1996	8,319	146	209	328	519	940	1,354	1,517	1,369	1,454	1
1995	7,952	134	198	276	534	916	1,327	1,452	1,312	1,305	1
1994	7,930	152	199	289	562	954	1,300	1,410	1,286	1,249	1
1993	7,827	123	188	301	558	905	1,284	1,459	1,192	1,230	-
1992	7,502	140	175	277	531	966	1,228	1,330	1,129	1,148	-
1991	7,587	147	210	313	593	928	1,256	1,324	1,121	1,118	-
1990	7,274	131	181	272	629	929	1,171	1,269	1,039	1,045	-
1989	7,298	133	198	352	581	879	1,182	1,244	1,025	1,051	-
1988	7,571	156	194	377	603	980	1,122	1,298	1,073	1,128	-
1987	7,265	108	184	336	615	876	1,144	1,285	1,012	1,067	-
1986	7,197	127	194	344	641	871	1,094	1,167	1,005	1,005	3
1985	7,233	124	189	387	614	910	1,125	1,175	995	1,059	-
1984	6,649	108	172	371	612	790	1,024	1,077	858	965	1
1983	6,585	111	219	342	585	806	1,003	1,048	870	899	4
1982	6,171	110	223	374	564	787	949	936	810	805	1
1981	6,120	116	235	368	577	765	934	917	759	783	1
1980	6,095	140	223	378	559	766	915	870	783	753	-
1979	5,680	135	215	385	556	702	854	865	693	614	-
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
White, female											
1998	9,002	149	157	269	402	584	1,077	1,487	1,727	2,760	-
1997	8,603	145	190	238	388	662	1,058	1,374	1,581	2,586	-
1996	8,236	130	157	238	375	654	983	1,358	1,447	2,523	-
1995	7,980	135	184	238	376	620	1,010	1,273	1,452	2,292	-
1994	7,734	122	162	207	360	661	968	1,242	1,430	2,228	-
1993	7,657	105	173	216	387	625	1,000	1,241	1,335	2,179	-
1992	7,140	101	126	208	365	647	920	1,177	1,214	1,965	-
1991	7,282	110	148	224	401	669	872	1,132	1,228	2,065	-
1990	6,898	114	132	204	405	574	875	1,053	1,221	1,898	-
1989	6,928	93	129	217	392	634	883	1,066	1,201	1,856	-
1988	7,055	105	144	230	394	627	896	1,178	1,183	1,814	-
1987	6,666	89	135	235	392	635	833	1,049	1,085	1,743	1
1986	6,554	82	145	248	419	649	841	1,064	1,043	1,638	2
1985	6,319	89	134	231	423	563	816	969	1,043	1,599	2
1984	5,782	87	140	248	365	576	725	840	921	1,404	-
1983	5,765	87	140	270	371	537	746	868	860	1,410	-
1982	5,050	80	139	242	375	507	629	758	724	1,163	1
1981	4,949	91	140	222	347	501	617	730	747	1,074	1
1980	4,609	84	131	232	323	472	548	675	640	1,034	-
1979	4,248	87	120	230	325	431	495	577	604	877	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All other, both sexes											
1998	2,318	105	120	157	196	239	273	285	230	314	2
1997	2,206	121	112	142	171	235	261	240	227	283	1
1996	2,172	120	111	131	180	211	256	253	226	268	2
1995	2,090	91	134	154	170	228	212	237	204	239	-
1994	2,138	106	108	135	164	220	277	230	205	250	1
1993	2,144	107	108	143	184	197	241	245	187	223	1
1992	2,110	102	95	136	174	236	236	207	205	223	1
1991	2,132	130	93	126	170	224	240	197	201	230	2
1990	2,183	86	97	134	201	227	230	229	183	235	-
1989	2,290	104	113	159	203	230	281	225	189	218	2
1988	2,256	90	123	139	198	246	233	259	180	231	1
1987	2,132	92	111	144	202	218	228	209	174	200	1
1986	2,132	95	104	159	177	215	231	234	148	195	-
1985	2,173	97	120	167	207	230	237	195	157	187	1
1984	1,915	103	102	142	173	172	223	192	137	167	3
1983	1,879	85	96	136	208	175	193	183	136	171	1
1982	1,855	69	115	167	178	198	191	172	133	125	1
1981	1,861	94	115	161	165	175	161	157	129	141	2
1980	1,848	102	142	159	167	181	161	168	120	117	2
1979	1,708	90	130	155	152	149	147	156	93	106	1
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All other, male											
1998	1,162	59	61	94	109	117	134	146	101	119	2
1997	1,100	70	48	74	82	129	135	128	104	113	-
1996	1,102	60	55	72	106	107	135	123	114	106	-
1995	1,051	51	69	73	86	125	111	134	94	97	-
1994	1,105	49	58	75	86	119	159	126	93	113	1
1993	1,094	46	49	74	98	108	127	126	94	98	1
1992	1,116	50	61	65	90	136	126	101	108	99	-
1991	1,134	80	40	70	88	126	130	110	105	88	2
1990	1,136	59	54	88	118	125	124	102	89	87	-
1989	1,264	59	60	96	117	144	154	119	89	104	-
1988	1,225	48	76	69	112	142	125	138	94	100	1
1987	1,191	51	64	82	108	133	136	134	90	80	1
1986	1,226	61	69	90	99	133	141	140	76	83	-
1985	1,212	58	70	95	120	128	140	108	82	88	1
1984	1,070	51	52	89	101	106	124	113	68	76	2
1983	1,080	52	53	88	119	111	111	114	75	82	1
1982	1,081	41	61	111	107	118	114	112	78	57	1
1981	1,073	53	75	98	96	102	92	84	72	75	-
1980	1,110	57	96	99	113	123	100	101	64	61	2
1979	1,046	56	83	103	96	103	89	100	56	51	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
All other, female											
1998	1,156	46	59	63	87	122	139	139	129	195	-
1997	1,106	51	64	68	89	106	126	112	123	170	1
1996	1,070	60	56	59	74	104	121	130	112	162	2
1995	1,039	40	65	81	84	103	101	103	110	142	-
1994	1,033	57	50	60	78	101	118	104	112	137	-
1993	1,050	61	59	69	86	89	114	119	93	125	-
1992	994	52	34	71	84	100	110	106	97	124	1
1991	998	50	53	56	82	98	110	87	96	142	-
1990	1,047	27	43	46	83	102	106	127	94	148	-
1989	1,026	45	53	63	86	86	127	106	100	114	2
1988	1,031	42	47	70	86	104	108	121	86	131	-
1987	941	41	47	62	94	85	92	75	84	120	-
1986	906	34	35	69	78	82	90	94	72	112	-
1985	961	39	50	72	87	102	97	87	75	99	-
1984	845	52	50	53	72	66	99	79	69	91	1
1983	799	33	43	48	89	64	82	69	61	89	-
1982	774	28	54	56	71	80	77	60	55	68	-
1981	788	41	40	63	69	73	69	73	57	66	2
1980	738	45	46	60	54	58	61	67	56	56	-
1979	662	34	47	52	56	46	58	56	37	55	-
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
Black, both sexes											
1998	1,902	88	105	135	161	193	229	225	172	243	2
1997	1,828	112	101	125	142	194	199	179	178	230	1
1996	1,831	107	101	113	157	174	208	199	169	223	2
1995	1,727	83	113	134	137	191	180	183	147	187	-
1994	1,786	99	95	115	138	178	227	178	165	198	1
1993	1,797	95	94	120	151	159	185	200	143	182	1
1992	1,792	96	85	107	148	200	207	164	155	177	1
1991	1,853	120	79	112	147	191	209	165	169	181	2
1990	1,902	79	88	114	174	194	193	188	161	191	-
1989	2,019	100	101	145	175	201	243	201	151	177	2
1988	1,995	81	110	130	176	208	206	220	153	201	1
1987	1,868	89	97	129	183	190	193	176	144	158	1
1986	1,902	93	95	152	161	181	205	212	121	154	-
1985	1,944	91	115	150	193	201	210	166	126	159	1
1984	1,725	101	91	136	161	157	199	168	112	134	3
1983	1,694	81	90	128	197	156	172	154	112	144	1
1982	1,692	67	110	152	163	185	172	146	116	107	1
1981	1,694	85	111	148	155	160	142	131	109	127	2
1980	1,662	95	132	151	153	160	145	144	101	93	2
1979	1,563	87	123	149	142	133	130	138	80	88	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
Black, male											
1998	940	50	52	81	91	90	114	112	71	81	2
1997	918	67	48	63	67	107	106	99	82	85	-
1996	919	55	49	63	92	87	104	99	81	85	-
1995	859	48	58	64	69	107	95	103	64	70	-
1994	896	47	51	66	68	91	130	97	67	80	1
1993	903	41	44	61	80	87	92	100	69	76	1
1992	941	49	54	50	77	113	110	83	80	75	-
1991	977	73	32	65	74	106	114	88	84	68	2
1990	976	54	52	74	101	102	103	88	71	64	-
1989	1,103	57	54	87	99	123	130	106	67	87	-
1988	1,091	43	66	64	99	120	116	120	79	89	1
1987	1,035	49	54	73	96	120	111	108	76	59	1
1986	1,097	60	65	84	92	110	122	132	64	63	-
1985	1,080	54	66	87	112	113	126	89	65	74	1
1984	956	51	47	86	93	95	111	93	54	58	2
1983	962	49	50	82	114	95	96	92	59	71	1
1982	981	40	59	103	99	110	102	92	65	49	1
1981	975	47	72	88	92	93	84	70	60	67	-
1980	998	53	90	95	101	108	93	86	52	48	2
1979	952	55	78	97	92	93	78	88	45	40	1
Other diseases of respiratory system (034.0,460-465,470-478,495,508-519)											
Black, female											
1998	962	38	53	54	70	103	115	113	101	162	-
1997	910	45	53	62	75	87	93	80	96	145	1
1996	912	52	52	50	65	87	104	100	88	138	2
1995	868	35	55	70	68	84	85	80	83	117	-
1994	890	52	44	49	70	87	97	81	98	118	-
1993	894	54	50	59	71	72	93	100	74	106	-
1992	851	47	31	57	71	87	97	81	75	102	1
1991	876	47	47	47	73	85	95	77	85	113	-
1990	926	25	36	40	73	92	90	100	90	127	-
1989	916	43	47	58	76	78	113	95	84	90	2
1988	904	38	44	66	77	88	90	100	74	112	-
1987	833	40	43	56	87	70	82	68	68	99	-
1986	805	33	30	68	69	71	83	80	57	91	-
1985	864	37	49	63	81	88	84	77	61	85	-
1984	769	50	44	50	68	62	88	75	58	76	1
1983	732	32	40	46	83	61	76	62	53	73	-
1982	711	27	51	49	64	75	70	54	51	58	-
1981	719	38	39	60	63	67	58	61	49	60	2
1980	664	42	42	56	52	52	52	58	49	45	-
1979	611	32	45	52	50	40	52	50	35	48	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
All races, both sexes											
1998	4,742	125	133	185	238	351	499	700	811	1,539	-
1997	5,178	133	157	179	253	355	595	770	871	1,677	2
1996	5,161	129	136	152	263	397	585	774	932	1,593	3
1995	5,522	134	138	197	285	448	615	836	1,010	1,659	-
1994	6,148	139	162	201	326	503	738	917	1,058	1,874	2
1993	6,000	127	180	219	299	527	673	869	1,114	1,776	2
1992	6,177	132	125	210	318	508	733	992	1,119	1,825	2
1991	6,168	130	143	210	381	504	733	980	1,102	1,776	-
1990	6,257	109	145	214	370	566	812	977	1,069	1,783	1
1989	6,526	107	166	253	364	579	827	1,020	1,172	1,822	1
1988	6,506	96	163	270	397	637	794	1,043	1,118	1,752	1
1987	6,427	108	163	258	395	612	818	1,104	1,033	1,714	-
1986	6,540	99	152	289	457	620	820	1,040	1,110	1,720	3
1985	6,715	114	179	282	511	669	847	1,078	1,133	1,708	1
1984	6,640	101	178	334	504	689	844	1,042	1,080	1,655	1
1983	6,445	119	191	316	462	668	882	1,018	1,054	1,535	1
1982	6,821	125	203	354	508	711	910	1,064	1,103	1,600	3
1981	6,579	145	212	400	503	696	917	1,034	987	1,439	1
1980	6,190	106	258	376	514	674	883	899	965	1,251	-
1979	6,004	151	266	368	483	668	784	927	882	1,194	1
Peptic ulcer (531-534)											
All races, male											
1998	2,261	80	75	119	150	218	270	355	353	534	-
1997	2,409	98	103	113	157	210	323	394	370	512	2
1996	2,432	91	92	102	160	229	318	396	416	490	2
1995	2,660	104	95	123	170	274	329	450	451	524	-
1994	2,964	91	114	127	208	318	395	466	463	615	1
1993	2,853	91	111	141	197	294	382	434	497	565	2
1992	2,948	95	84	130	199	311	415	474	484	600	1
1991	2,959	85	98	144	227	292	414	512	484	565	-
1990	3,011	68	96	148	223	344	477	481	456	568	-
1989	3,187	71	98	162	218	356	473	518	542	593	1
1988	3,221	61	108	174	250	382	471	524	485	599	1
1987	3,232	73	112	177	253	368	484	579	467	566	-
1986	3,140	71	99	197	278	347	473	506	449	569	1
1985	3,376	78	124	171	322	426	466	569	496	586	1
1984	3,262	53	124	219	328	416	443	525	468	554	-
1983	3,303	73	125	202	280	431	520	539	463	535	1
1982	3,465	74	140	228	322	428	523	550	482	568	2
1981	3,482	102	134	247	316	432	535	576	444	526	1
1980	3,448	79	177	254	347	440	538	500	459	486	-
1979	3,311	101	184	235	315	429	451	515	432	469	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
All races, female											
1998	2,481	45	58	66	88	133	229	345	458	1,005	-
1997	2,769	35	54	66	96	145	272	376	501	1,165	-
1996	2,729	38	44	50	103	168	267	378	516	1,103	1
1995	2,862	30	43	74	115	174	286	386	559	1,135	-
1994	3,184	48	48	74	118	185	343	451	595	1,259	1
1993	3,147	36	69	78	102	233	291	435	617	1,211	-
1992	3,229	37	41	80	119	197	318	518	635	1,225	1
1991	3,209	45	45	66	154	212	319	468	618	1,211	-
1990	3,246	41	49	66	147	222	335	496	613	1,215	1
1989	3,339	36	68	91	146	223	354	502	630	1,229	-
1988	3,285	35	55	96	147	255	323	519	633	1,153	-
1987	3,195	35	51	81	142	244	334	525	566	1,148	-
1986	3,400	28	53	92	179	273	347	534	661	1,151	2
1985	3,339	36	55	111	189	243	381	509	637	1,122	-
1984	3,378	48	54	115	176	273	401	517	612	1,101	1
1983	3,142	46	66	114	182	237	362	479	591	1,000	-
1982	3,356	51	63	126	186	283	387	514	621	1,032	1
1981	3,097	43	78	153	187	264	382	458	543	913	-
1980	2,742	27	81	122	167	234	345	399	506	765	-
1979	2,693	50	82	133	168	239	333	412	450	725	1
Peptic ulcer (531-534)											
White, both sexes											
1998	4,199	92	112	157	195	299	431	629	733	1,434	-
1997	4,541	92	113	143	199	298	518	689	798	1,566	1
1996	4,508	96	104	117	204	325	488	695	856	1,500	1
1995	4,889	103	104	150	252	378	537	746	928	1,561	-
1994	5,416	110	129	159	260	439	641	830	960	1,740	1
1993	5,317	91	136	183	249	451	583	772	1,032	1,668	1
1992	5,498	99	96	163	265	425	656	905	1,047	1,704	1
1991	5,501	96	103	175	309	441	651	878	1,032	1,670	-
1990	5,580	86	110	170	305	484	715	898	997	1,684	1
1989	5,728	75	132	197	303	488	718	901	1,083	1,699	1
1988	5,855	74	131	227	319	557	721	954	1,059	1,658	1
1987	5,805	80	139	218	330	553	715	1,029	973	1,624	-
1986	5,845	73	125	237	384	535	736	940	1,024	1,627	2
1985	6,049	97	156	233	429	605	760	982	1,044	1,609	1
1984	5,999	71	134	286	440	616	775	961	1,000	1,573	1
1983	5,817	88	164	259	394	588	786	943	990	1,459	1
1982	6,140	94	153	300	433	624	828	985	1,016	1,527	1
1981	5,916	111	173	338	428	623	821	966	915	1,372	1
1980	5,612	76	214	328	469	613	803	829	901	1,195	-
1979	5,426	122	203	316	433	596	713	863	837	1,140	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
White, male											
1998	1,980	57	62	98	122	184	235	315	325	501	-
1997	2,051	69	73	86	122	173	279	348	339	475	1
1996	2,074	67	70	80	118	191	264	360	379	459	-
1995	2,298	86	72	95	148	229	292	397	411	486	-
1994	2,556	71	87	101	167	283	346	427	417	556	1
1993	2,458	67	83	117	160	242	329	386	453	525	1
1992	2,560	73	62	100	166	264	372	433	442	555	-
1991	2,586	64	71	122	186	254	370	458	445	527	-
1990	2,627	56	69	117	175	293	425	449	417	536	-
1989	2,739	46	77	122	186	297	410	463	497	546	1
1988	2,843	46	88	143	207	330	427	478	455	565	1
1987	2,865	52	95	151	209	330	423	536	440	536	-
1986	2,752	49	82	162	238	295	424	453	406	537	1
1985	2,990	65	106	139	267	387	416	514	456	546	1
1984	2,897	37	89	185	284	378	401	485	435	520	-
1983	2,921	49	108	172	236	386	452	494	429	497	1
1982	3,072	53	105	195	272	382	473	509	436	538	-
1981	3,081	77	109	208	270	386	476	535	409	497	1
1980	3,075	54	142	218	312	397	494	460	425	462	-
1979	2,943	77	136	202	282	386	411	474	409	443	-
Peptic ulcer (531-534)											
White, female											
1998	2,219	35	50	59	73	115	196	314	408	933	-
1997	2,490	23	40	57	77	125	239	341	459	1,091	-
1996	2,434	29	34	37	86	134	224	335	477	1,041	1
1995	2,591	17	32	55	104	149	245	349	517	1,075	-
1994	2,860	39	42	58	93	156	295	403	543	1,184	-
1993	2,859	24	53	66	89	209	254	386	579	1,143	-
1992	2,938	26	34	63	99	161	284	472	605	1,149	1
1991	2,915	32	32	53	123	187	281	420	587	1,143	-
1990	2,953	30	41	53	130	191	290	449	580	1,148	1
1989	2,989	29	55	75	117	191	308	438	586	1,153	-
1988	3,012	28	43	84	112	227	294	476	604	1,093	-
1987	2,940	28	44	67	121	223	292	493	533	1,088	-
1986	3,093	24	43	75	146	240	312	487	618	1,090	1
1985	3,059	32	50	94	162	218	344	468	588	1,063	-
1984	3,102	34	45	101	156	238	374	476	565	1,053	1
1983	2,896	39	56	87	158	202	334	449	561	962	-
1982	3,068	41	48	105	161	242	355	476	580	989	1
1981	2,835	34	64	130	158	237	345	431	506	875	-
1980	2,537	22	72	110	157	216	309	369	476	733	-
1979	2,483	45	67	114	151	210	302	389	428	697	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
All other, both sexes											
1998	543	33	21	28	43	52	68	71	78	105	-
1997	637	41	44	36	54	57	77	81	73	111	1
1996	653	33	32	35	59	72	97	79	76	93	2
1995	633	31	34	47	33	70	78	90	82	98	-
1994	732	29	33	42	66	64	97	87	98	134	1
1993	683	36	44	36	50	76	90	97	82	108	1
1992	679	33	29	47	53	83	77	87	72	121	1
1991	667	34	40	35	72	63	82	102	70	106	-
1990	677	23	35	44	65	82	97	79	72	99	-
1989	798	32	34	56	61	91	109	119	89	123	-
1988	651	22	32	43	78	80	73	89	59	94	-
1987	622	28	24	40	65	59	103	75	60	90	-
1986	695	26	27	52	73	85	84	100	86	93	1
1985	666	17	23	49	82	64	87	96	89	99	-
1984	641	30	44	48	64	73	69	81	80	82	-
1983	628	31	27	57	68	80	96	75	64	76	-
1982	681	31	50	54	75	87	82	79	87	73	2
1981	663	34	39	62	75	73	96	68	72	67	-
1980	578	30	44	48	45	61	80	70	64	56	-
1979	578	29	63	52	50	72	71	64	45	54	1
Peptic ulcer (531-534)											
All other, male											
1998	281	23	13	21	28	34	35	40	28	33	-
1997	358	29	30	27	35	37	44	46	31	37	1
1996	358	24	22	22	42	38	54	36	37	31	2
1995	362	18	23	28	22	45	37	53	40	38	-
1994	408	20	27	26	41	35	49	39	46	59	-
1993	395	24	28	24	37	52	53	48	44	40	1
1992	388	22	22	30	33	47	43	41	42	45	1
1991	373	21	27	22	41	38	44	54	39	38	-
1990	384	12	27	31	48	51	52	32	39	32	-
1989	448	25	21	40	32	59	63	55	45	47	-
1988	378	15	20	31	43	52	44	46	30	34	-
1987	367	21	17	26	44	38	61	43	27	30	-
1986	388	22	17	35	40	52	49	53	43	32	-
1985	386	13	18	32	55	39	50	55	40	40	-
1984	365	16	35	34	44	38	42	40	33	34	-
1983	382	24	17	30	44	45	68	45	34	38	-
1982	393	21	35	33	50	46	50	41	46	30	2
1981	401	25	25	39	46	46	59	41	35	29	-
1980	373	25	35	36	35	43	44	40	34	24	-
1979	368	24	48	33	33	43	40	41	23	26	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
All other, female											
1998	262	10	8	7	15	18	33	31	50	72	-
1997	279	12	14	9	19	20	33	35	42	74	-
1996	295	9	10	13	17	34	43	43	39	62	-
1995	271	13	11	19	11	25	41	37	42	60	-
1994	324	9	6	16	25	29	48	48	52	75	1
1993	288	12	16	12	13	24	37	49	38	68	-
1992	291	11	7	17	20	36	34	46	30	76	-
1991	294	13	13	13	31	25	38	48	31	68	-
1990	293	11	8	13	17	31	45	47	33	67	-
1989	350	7	13	16	29	32	46	64	44	76	-
1988	273	7	12	12	35	28	29	43	29	60	-
1987	255	7	7	14	21	21	42	32	33	60	-
1986	307	4	10	17	33	33	35	47	43	61	1
1985	280	4	5	17	27	25	37	41	49	59	-
1984	276	14	9	14	20	35	27	41	47	48	-
1983	246	7	10	27	24	35	28	30	30	38	-
1982	288	10	15	21	25	41	32	38	41	43	-
1981	262	9	14	23	29	27	37	27	37	38	-
1980	205	5	9	12	10	18	36	30	30	32	-
1979	210	5	15	19	17	29	31	23	22	28	1
Peptic ulcer (531-534)											
Black, both sexes											
1998	417	29	19	23	38	33	54	52	55	77	-
1997	497	35	36	32	44	42	58	61	51	83	-
1996	526	31	27	28	45	61	78	63	57	68	2
1995	524	27	30	41	30	59	65	74	62	71	-
1994	584	29	25	38	54	50	71	63	76	103	1
1993	563	35	40	30	39	68	72	75	63	81	1
1992	570	29	27	41	45	67	59	71	63	97	1
1991	552	31	34	29	65	50	65	82	52	86	-
1990	589	22	29	41	60	76	87	63	62	75	-
1989	672	31	32	47	49	74	95	108	70	91	-
1988	562	18	29	38	71	69	61	74	49	75	-
1987	546	27	24	36	52	53	90	65	53	71	-
1986	612	26	20	48	66	80	75	82	73	76	1
1985	576	15	22	42	77	57	72	84	74	80	-
1984	565	30	40	43	55	66	56	70	72	68	-
1983	556	29	24	52	64	66	83	66	57	65	-
1982	598	27	48	52	65	79	74	66	69	58	2
1981	599	32	34	56	71	63	86	60	64	59	-
1980	516	29	41	44	43	57	73	55	51	47	-
1979	511	28	54	48	45	64	67	55	39	42	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Peptic ulcer (531-534)											
Black, male											
1998	208	21	11	17	25	20	26	29	19	19	-
1997	278	26	25	24	25	29	33	32	23	25	-
1996	284	23	19	17	35	31	43	26	26	18	2
1995	293	15	20	24	21	38	30	40	31	20	-
1994	318	20	21	23	34	25	37	24	33	42	-
1993	319	23	25	22	28	46	42	32	31	28	1
1992	325	20	20	25	29	39	32	35	37	29	1
1991	304	21	23	18	37	29	37	42	26	27	-
1990	336	11	24	29	43	47	47	25	30	24	-
1989	366	25	20	34	28	44	50	48	32	30	-
1988	326	14	18	27	39	47	39	36	22	24	-
1987	318	20	17	23	32	34	53	39	22	19	-
1986	339	22	15	32	35	49	42	41	35	26	-
1985	331	12	18	29	51	34	39	46	29	32	-
1984	315	16	31	30	36	34	34	30	30	28	-
1983	332	22	16	27	40	36	57	37	29	32	-
1982	341	17	34	33	46	40	45	32	33	21	2
1981	360	24	22	36	43	38	52	34	28	28	-
1980	335	24	34	35	33	41	40	30	26	18	-
1979	320	23	42	31	30	37	37	33	20	18	-
Peptic ulcer (531-534)											
Black, female											
1998	209	8	8	6	13	13	28	23	36	58	-
1997	219	9	11	8	19	13	25	29	28	58	-
1996	242	8	8	11	10	30	35	37	31	50	-
1995	231	12	10	17	9	21	35	34	31	51	-
1994	266	9	4	15	20	25	34	39	43	61	1
1993	244	12	15	8	11	22	30	43	32	53	-
1992	245	9	7	16	16	28	27	36	26	68	-
1991	248	10	11	11	28	21	28	40	26	59	-
1990	253	11	5	12	17	29	40	38	32	51	-
1989	306	6	12	13	21	30	45	60	38	61	-
1988	236	4	11	11	32	22	22	38	27	51	-
1987	228	7	7	13	20	19	37	26	31	52	-
1986	273	4	5	16	31	31	33	41	38	50	1
1985	245	3	4	13	26	23	33	38	45	48	-
1984	250	14	9	13	19	32	22	40	42	40	-
1983	224	7	8	25	24	30	26	29	28	33	-
1982	257	10	14	19	19	39	29	34	36	37	-
1981	239	8	12	20	28	25	34	26	36	31	-
1980	181	5	7	9	10	16	33	25	25	29	-
1979	191	5	12	17	15	27	30	22	19	24	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
All races, both sexes											
1998	439	15	14	26	27	35	41	65	56	79	-
1997	395	18	14	14	33	41	46	42	46	70	-
1996	424	17	17	18	31	35	47	62	47	65	-
1995	398	19	15	24	38	43	40	42	51	72	-
1994	380	21	19	16	19	32	49	44	49	63	-
1993	367	18	12	14	31	43	43	41	39	53	-
1992	413	11	14	21	20	53	49	49	49	74	-
1991	389	13	18	20	29	38	47	38	55	61	-
1990	438	17	15	19	36	41	51	66	48	67	-
1989	451	11	24	30	23	46	65	63	49	74	-
1988	473	10	15	28	40	48	55	57	66	82	-
1987	481	12	19	27	30	50	65	67	57	92	-
1986	512	6	18	40	47	61	65	56	74	69	-
1985	501	10	18	20	48	52	73	72	49	93	-
1984	493	10	10	28	44	47	64	64	75	80	-
1983	526	12	19	33	39	65	67	63	64	91	-
1982	529	20	18	36	47	51	62	65	57	86	-
1981	589	18	19	35	49	72	90	71	62	77	-
1980	701	19	30	59	67	58	77	90	81	100	-
1979	682	24	26	51	56	67	96	87	77	90	-
Diseases of appendix (540-543)											
All races, male											
1998	256	11	7	20	16	23	30	30	29	36	-
1997	228	10	7	11	22	25	26	21	26	34	-
1996	240	10	15	12	21	24	25	36	19	27	-
1995	229	14	11	17	21	30	24	21	29	24	-
1994	220	13	17	11	12	23	31	21	24	29	-
1993	217	11	11	12	19	27	26	22	20	20	-
1992	244	7	9	10	13	32	25	30	28	40	-
1991	207	6	9	9	19	21	28	19	29	19	-
1990	257	14	8	12	24	30	30	41	24	26	-
1989	237	8	13	16	12	23	38	37	23	21	-
1988	241	8	11	15	25	23	31	25	32	29	-
1987	255	4	13	19	18	26	35	34	25	37	-
1986	280	2	9	21	31	36	38	27	40	34	-
1985	276	8	12	12	29	31	44	36	21	38	-
1984	265	6	3	14	33	27	37	33	39	36	-
1983	292	6	13	20	21	44	34	39	27	37	-
1982	303	12	10	25	30	30	38	37	31	34	-
1981	329	11	13	22	29	41	53	32	34	32	-
1980	383	12	21	35	40	38	52	50	34	39	-
1979	373	15	15	27	34	41	55	50	34	36	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
All races, female											
1998	183	4	7	6	11	12	11	35	27	43	-
1997	167	8	7	3	11	16	20	21	20	36	-
1996	184	7	2	6	10	11	22	26	28	38	-
1995	169	5	4	7	17	13	16	21	22	48	-
1994	160	8	2	5	7	9	18	23	25	34	-
1993	150	7	1	2	12	16	17	19	19	33	-
1992	169	4	5	11	7	21	24	19	21	34	-
1991	182	7	9	11	10	17	19	19	26	42	-
1990	181	3	7	7	12	11	21	25	24	41	-
1989	214	3	11	14	11	23	27	26	26	53	-
1988	232	2	4	13	15	25	24	32	34	53	-
1987	226	8	6	8	12	24	30	33	32	55	-
1986	232	4	9	19	16	25	27	29	34	35	-
1985	225	2	6	8	19	21	29	36	28	55	-
1984	228	4	7	14	11	20	27	31	36	44	-
1983	234	6	6	13	18	21	33	24	37	54	-
1982	226	8	8	11	17	21	24	28	26	52	-
1981	260	7	6	13	20	31	37	39	28	45	-
1980	318	7	9	24	27	20	25	40	47	61	-
1979	309	9	11	24	22	26	41	37	43	54	-
Diseases of appendix (540-543)											
White, both sexes											
1998	356	7	9	21	21	29	36	53	50	72	-
1997	309	9	12	10	28	26	38	35	38	66	-
1996	352	14	14	14	25	28	41	51	44	57	-
1995	313	11	13	17	24	37	31	35	44	63	-
1994	305	15	15	14	16	30	37	37	42	56	-
1993	283	12	7	11	20	36	31	37	33	47	-
1992	326	4	10	15	18	44	40	42	40	66	-
1991	321	9	15	17	26	33	36	31	48	57	-
1990	357	11	9	16	31	28	43	56	42	61	-
1989	354	8	16	17	16	40	53	59	42	62	-
1988	368	5	7	18	28	36	46	47	56	73	-
1987	378	6	12	23	22	37	51	53	49	79	-
1986	394	5	6	31	35	46	53	49	61	63	-
1985	396	7	14	12	39	42	55	59	45	82	-
1984	403	6	7	22	32	41	50	53	69	73	-
1983	428	7	15	25	30	53	51	55	53	83	-
1982	413	13	11	28	35	36	47	55	47	74	-
1981	466	14	15	22	37	52	77	58	49	64	-
1980	566	12	18	46	44	44	65	79	74	95	-
1979	539	14	15	39	43	47	78	72	69	86	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
White, male											
1998	201	5	2	16	11	18	27	23	27	33	-
1997	167	6	5	7	17	15	21	16	21	30	-
1996	191	7	12	9	16	20	21	28	17	24	-
1995	168	7	9	10	12	26	17	16	24	21	-
1994	168	8	13	9	9	22	22	18	19	26	-
1993	159	6	6	9	11	23	17	20	17	17	-
1992	196	3	8	7	12	27	18	25	24	37	-
1991	163	4	6	8	16	17	21	15	25	18	-
1990	212	9	4	10	22	21	26	34	19	26	-
1989	185	6	9	9	9	21	31	36	19	17	-
1988	181	5	4	9	19	19	25	21	25	24	-
1987	203	2	9	17	15	18	25	31	22	31	-
1986	215	2	2	16	25	28	28	24	31	31	-
1985	217	6	11	7	24	24	35	30	18	33	-
1984	220	5	3	11	23	24	30	30	37	32	-
1983	238	4	11	16	19	33	25	35	21	32	-
1982	236	8	7	20	24	21	29	31	24	29	-
1981	260	10	11	14	22	34	43	26	23	26	-
1980	308	8	14	28	29	30	44	44	29	36	-
1979	292	8	9	21	28	28	46	38	29	35	-
Diseases of appendix (540-543)											
White, female											
1998	155	2	7	5	10	11	9	30	23	39	-
1997	142	3	7	3	11	11	17	19	17	36	-
1996	161	7	2	5	9	8	20	23	27	33	-
1995	145	4	4	7	12	11	14	19	20	42	-
1994	137	7	2	5	7	8	15	19	23	30	-
1993	124	6	1	2	9	13	14	17	16	30	-
1992	130	1	2	8	6	17	22	17	16	29	-
1991	158	5	9	9	10	16	15	16	23	39	-
1990	145	2	5	6	9	7	17	22	23	35	-
1989	169	2	7	8	7	19	22	23	23	45	-
1988	187	-	3	9	9	17	21	26	31	49	-
1987	175	4	3	6	7	19	26	22	27	48	-
1986	179	3	4	15	10	18	25	25	30	32	-
1985	179	1	3	5	15	18	20	29	27	49	-
1984	183	1	4	11	9	17	20	23	32	41	-
1983	190	3	4	9	11	20	26	20	32	51	-
1982	177	5	4	8	11	15	18	24	23	45	-
1981	206	4	4	8	15	18	34	32	26	38	-
1980	258	4	4	18	15	14	21	35	45	59	-
1979	247	6	6	18	15	19	32	34	40	51	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
All other, both sexes											
1998	83	8	5	5	6	6	5	12	6	7	-
1997	86	9	2	4	5	15	8	7	8	4	-
1996	72	3	3	4	6	7	6	11	3	8	-
1995	85	8	2	7	14	6	9	7	7	9	-
1994	75	6	4	2	3	2	12	7	7	7	-
1993	84	6	5	3	11	7	12	4	6	6	-
1992	87	7	4	6	2	9	9	7	9	8	-
1991	68	4	3	3	3	5	11	7	7	4	-
1990	81	6	6	3	5	13	8	10	6	6	-
1989	97	3	8	13	7	6	12	4	7	12	-
1988	105	5	8	10	12	12	9	10	10	9	-
1987	103	6	7	4	8	13	14	14	8	13	-
1986	118	1	12	9	12	15	12	7	13	6	-
1985	105	3	4	8	9	10	18	13	4	11	-
1984	90	4	3	6	12	6	14	11	6	7	-
1983	98	5	4	8	9	12	16	8	11	8	-
1982	116	7	7	8	12	15	15	10	10	12	-
1981	123	4	4	13	12	20	13	13	13	13	-
1980	135	7	12	13	23	14	12	11	7	5	-
1979	143	10	11	12	13	20	18	15	8	4	-
Diseases of appendix (540-543)											
All other, male											
1998	55	6	5	4	5	5	3	7	2	3	-
1997	61	4	2	4	5	10	5	5	5	4	-
1996	49	3	3	3	5	4	4	8	2	3	-
1995	61	7	2	7	9	4	7	5	5	3	-
1994	52	5	4	2	3	1	9	3	5	3	-
1993	58	5	5	3	8	4	9	2	3	3	-
1992	48	4	1	3	1	5	7	5	4	3	-
1991	44	2	3	1	3	4	7	4	4	1	-
1990	45	5	4	2	2	9	4	7	5	-	-
1989	52	2	4	7	3	2	7	1	4	4	-
1988	60	3	7	6	6	4	6	4	7	5	-
1987	52	2	4	2	3	8	10	3	3	6	-
1986	65	-	7	5	6	8	10	3	9	3	-
1985	59	2	1	5	5	7	9	6	3	5	-
1984	45	1	-	3	10	3	7	3	2	4	-
1983	54	2	2	4	2	11	9	4	6	5	-
1982	67	4	3	5	6	9	9	6	7	5	-
1981	69	1	2	8	7	7	10	6	11	6	-
1980	75	4	7	7	11	8	8	6	5	3	-
1979	81	7	6	6	6	13	9	12	5	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
All other, female											
1998	28	2	-	1	1	1	2	5	4	4	-
1997	25	5	-	-	-	5	3	2	3	-	-
1996	23	-	-	1	1	3	2	3	1	5	-
1995	24	1	-	-	5	2	2	2	2	6	-
1994	23	1	-	-	-	1	3	4	2	4	-
1993	26	1	-	-	3	3	3	2	3	3	-
1992	39	3	3	3	1	4	2	2	5	5	-
1991	24	2	-	2	-	1	4	3	3	3	-
1990	36	1	2	1	3	4	4	3	1	6	-
1989	45	1	4	6	4	4	5	3	3	8	-
1988	45	2	1	4	6	8	3	6	3	4	-
1987	51	4	3	2	5	5	4	11	5	7	-
1986	53	1	5	4	6	7	2	4	4	3	-
1985	46	1	3	3	4	3	9	7	1	6	-
1984	45	3	3	3	2	3	7	8	4	3	-
1983	44	3	2	4	7	1	7	4	5	3	-
1982	49	3	4	3	6	6	6	4	3	7	-
1981	54	3	2	5	5	13	3	7	2	7	-
1980	60	3	5	6	12	6	4	5	2	2	-
1979	62	3	5	6	7	7	9	3	3	3	-
Diseases of appendix (540-543)											
Black, both sexes											
1998	66	6	5	5	6	6	3	8	5	7	-
1997	76	8	2	3	5	13	5	6	7	4	-
1996	66	3	3	3	5	6	6	9	3	8	-
1995	79	8	2	6	12	6	9	7	5	8	-
1994	68	6	4	1	3	2	12	7	6	6	-
1993	71	5	4	3	9	5	10	4	5	4	-
1992	76	7	4	6	2	7	9	7	5	7	-
1991	65	4	3	3	3	5	10	7	7	3	-
1990	71	5	5	3	5	12	7	9	3	6	-
1989	89	3	8	13	7	6	9	3	7	11	-
1988	99	4	8	10	11	11	8	9	10	9	-
1987	95	5	7	3	7	13	13	13	8	11	-
1986	109	1	10	9	11	14	12	5	11	6	-
1985	99	3	3	8	6	10	17	13	4	11	-
1984	84	4	3	6	12	6	13	11	3	7	-
1983	91	4	4	8	9	12	16	7	10	8	-
1982	108	6	6	6	12	15	12	10	9	12	-
1981	117	4	4	12	10	20	12	13	12	12	-
1980	130	7	12	13	22	13	12	10	6	5	-
1979	136	10	11	11	12	18	18	14	8	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Diseases of appendix (540-543)											
Black, male											
1998	46	5	5	4	5	5	2	6	2	3	-
1997	56	4	2	3	5	8	4	4	5	4	-
1996	44	3	3	2	4	4	4	6	2	3	-
1995	57	7	2	6	8	4	7	5	3	3	-
1994	46	5	4	1	3	1	9	3	4	2	-
1993	48	4	4	3	7	3	8	2	2	1	-
1992	41	4	1	3	1	4	7	5	2	2	-
1991	42	2	3	1	3	4	6	4	4	1	-
1990	39	5	3	2	2	9	3	6	3	-	-
1989	47	2	4	7	3	2	5	1	4	3	-
1988	57	2	7	6	5	3	6	4	7	5	-
1987	47	1	4	2	3	8	9	3	3	4	-
1986	59	-	5	5	5	8	10	2	7	3	-
1985	57	2	1	5	4	7	9	6	3	5	-
1984	40	1	-	3	10	3	6	3	-	4	-
1983	51	1	2	4	2	11	9	4	5	5	-
1982	62	4	3	3	6	9	6	6	7	5	-
1981	64	1	2	7	5	7	10	6	10	5	-
1980	72	4	7	7	11	8	8	5	4	3	-
1979	75	7	6	5	6	11	9	11	5	1	-
Diseases of appendix (540-543)											
Black, female											
1998	20	1	-	1	1	1	1	2	3	4	-
1997	20	4	-	-	-	5	1	2	2	-	-
1996	22	-	-	1	1	2	2	3	1	5	-
1995	22	1	-	-	4	2	2	2	2	5	-
1994	22	1	-	-	-	1	3	4	2	4	-
1993	23	1	-	-	2	2	2	2	3	3	-
1992	35	3	3	3	1	3	2	2	3	5	-
1991	23	2	-	2	-	1	4	3	3	2	-
1990	32	-	2	1	3	3	4	3	-	6	-
1989	42	1	4	6	4	4	4	2	3	8	-
1988	42	2	1	4	6	8	2	5	3	4	-
1987	48	4	3	1	4	5	4	10	5	7	-
1986	50	1	5	4	6	6	2	3	4	3	-
1985	42	1	2	3	2	3	8	7	1	6	-
1984	44	3	3	3	2	3	7	8	3	3	-
1983	40	3	2	4	7	1	7	3	5	3	-
1982	46	2	3	3	6	6	6	4	2	7	-
1981	53	3	2	5	5	13	2	7	2	7	-
1980	58	3	5	6	11	5	4	5	2	2	-
1979	61	3	5	6	6	7	9	3	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
All races, both sexes											
1998	1,389	27	43	52	53	68	139	193	254	485	-
1997	1,433	30	37	52	47	96	125	180	237	540	-
1996	1,403	30	30	45	60	78	134	180	253	500	-
1995	1,335	28	30	31	43	97	133	183	217	492	1
1994	1,297	30	28	42	71	83	116	173	210	469	-
1993	1,315	19	30	23	55	74	123	182	234	487	-
1992	1,319	10	27	23	65	103	109	187	216	495	-
1991	1,301	23	19	23	66	102	140	181	230	411	-
1990	1,342	23	26	30	52	92	117	199	235	478	1
1989	1,343	20	21	43	59	100	128	190	228	458	-
1988	1,327	24	25	34	59	88	142	181	242	455	-
1987	1,357	13	27	37	72	111	135	183	233	441	1
1986	1,305	18	29	48	81	86	144	199	195	408	2
1985	1,391	8	29	57	66	96	131	204	280	426	-
1984	1,352	12	28	51	74	101	130	201	205	439	-
1983	1,399	14	26	52	71	119	125	210	241	424	1
1982	1,400	23	33	57	74	118	137	190	220	430	1
1981	1,590	27	44	52	72	110	204	239	263	446	-
1980	1,588	26	46	71	89	127	190	218	251	417	-
1979	1,643	22	46	82	91	143	186	222	279	396	-
Hernia (550-553)											
All races, male											
1998	555	17	30	24	26	33	71	94	78	133	-
1997	620	17	18	33	26	52	60	91	110	154	-
1996	576	17	19	23	31	37	68	83	104	145	-
1995	549	15	15	15	25	45	65	76	94	152	1
1994	527	16	14	20	37	34	69	73	81	134	-
1993	532	13	17	13	29	36	65	80	89	137	-
1992	515	2	12	8	30	53	50	83	88	147	-
1991	530	10	9	9	32	40	66	75	96	124	-
1990	548	11	13	19	24	36	62	85	97	143	1
1989	574	11	8	24	34	45	66	86	99	144	-
1988	530	14	12	22	31	37	55	78	112	129	-
1987	551	7	14	16	38	46	59	78	95	134	1
1986	591	10	18	24	39	44	76	92	76	145	-
1985	573	5	17	28	34	42	68	89	95	137	-
1984	607	8	13	24	40	47	66	100	96	137	-
1983	630	6	13	25	39	62	73	90	106	141	1
1982	617	12	21	26	38	58	53	85	86	159	-
1981	722	13	19	24	33	65	92	110	116	171	-
1980	729	16	20	30	43	63	92	100	118	159	-
1979	761	9	28	41	51	75	86	101	107	151	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
All races, female											
1998	834	10	13	28	27	35	68	99	176	352	-
1997	813	13	19	19	21	44	65	89	127	386	-
1996	827	13	11	22	29	41	66	97	149	355	-
1995	786	13	15	16	18	52	68	107	123	340	-
1994	770	14	14	22	34	49	47	100	129	335	-
1993	783	6	13	10	26	38	58	102	145	350	-
1992	804	8	15	15	35	50	59	104	128	348	-
1991	771	13	10	14	34	62	74	106	134	287	-
1990	794	12	13	11	28	56	55	114	138	335	-
1989	769	9	13	19	25	55	62	104	129	314	-
1988	797	10	13	12	28	51	87	103	130	326	-
1987	806	6	13	21	34	65	76	105	138	307	-
1986	714	8	11	24	42	42	68	107	119	263	2
1985	818	3	12	29	32	54	63	115	185	289	-
1984	745	4	15	27	34	54	64	101	109	302	-
1983	769	8	13	27	32	57	52	120	135	283	-
1982	783	11	12	31	36	60	84	105	134	271	1
1981	868	14	25	28	39	45	112	129	147	275	-
1980	859	10	26	41	46	64	98	118	133	258	-
1979	882	13	18	41	40	68	100	121	172	245	-
Hernia (550-553)											
White, both sexes											
1998	1,237	22	35	41	41	56	128	177	228	450	-
1997	1,289	22	29	43	40	83	106	162	224	510	-
1996	1,243	17	25	37	48	67	117	166	237	466	-
1995	1,184	20	23	21	39	85	114	167	192	464	1
1994	1,143	23	14	29	58	69	103	158	195	441	-
1993	1,177	16	22	16	40	67	109	163	218	461	-
1992	1,174	9	21	20	57	89	93	169	197	461	-
1991	1,139	17	18	17	56	87	114	161	206	387	-
1990	1,192	18	16	24	42	78	100	183	220	453	-
1989	1,189	12	18	40	50	83	111	161	215	435	-
1988	1,179	16	21	28	45	73	129	159	228	426	-
1987	1,211	13	18	29	65	101	117	164	215	422	-
1986	1,155	15	25	39	66	72	125	178	175	392	2
1985	1,260	7	22	46	52	87	120	186	260	409	-
1984	1,207	7	22	41	61	89	106	192	189	413	-
1983	1,232	10	22	39	58	103	102	193	218	403	1
1982	1,251	22	25	44	63	104	119	171	201	412	1
1981	1,427	17	39	46	62	95	178	225	242	419	-
1980	1,415	17	31	56	76	114	170	202	231	403	-
1979	1,482	22	36	70	82	127	167	206	268	377	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
White, male											
1998	496	16	25	19	22	28	66	84	72	128	-
1997	540	12	13	27	21	42	54	82	103	142	-
1996	499	10	14	21	26	35	56	78	95	131	-
1995	471	13	11	10	22	40	56	67	82	142	1
1994	460	14	8	15	31	29	62	67	73	125	-
1993	469	11	13	11	24	32	53	73	84	128	-
1992	446	2	11	7	25	45	47	75	80	129	-
1991	450	6	9	7	27	35	51	65	88	113	-
1990	480	9	8	15	21	32	55	78	93	133	-
1989	496	7	8	22	28	37	58	75	94	132	-
1988	459	10	10	18	25	32	49	70	104	115	-
1987	476	7	9	10	35	41	53	71	88	122	-
1986	521	7	15	21	34	37	66	85	69	141	-
1985	512	4	14	25	25	38	66	83	83	128	-
1984	534	4	9	15	36	41	57	97	86	126	-
1983	542	5	11	22	31	54	59	83	94	132	1
1982	531	11	15	21	32	53	41	77	76	149	-
1981	638	10	18	20	30	55	79	102	103	161	-
1980	625	10	14	22	35	56	81	91	104	149	-
1979	672	9	21	33	48	65	79	93	102	140	-
Hernia (550-553)											
White, female											
1998	741	6	10	22	19	28	62	93	156	322	-
1997	749	10	16	16	19	41	52	80	121	368	-
1996	744	7	11	16	22	32	61	88	142	335	-
1995	713	7	12	11	17	45	58	100	110	322	-
1994	683	9	6	14	27	40	41	91	122	316	-
1993	708	5	9	5	16	35	56	90	134	333	-
1992	728	7	10	13	32	44	46	94	117	332	-
1991	689	11	9	10	29	52	63	96	118	274	-
1990	712	9	8	9	21	46	45	105	127	320	-
1989	693	5	10	18	22	46	53	86	121	303	-
1988	720	6	11	10	20	41	80	89	124	311	-
1987	735	6	9	19	30	60	64	93	127	300	-
1986	634	8	10	18	32	35	59	93	106	251	2
1985	748	3	8	21	27	49	54	103	177	281	-
1984	673	3	13	26	25	48	49	95	103	287	-
1983	690	5	11	17	27	49	43	110	124	271	-
1982	720	11	10	23	31	51	78	94	125	263	1
1981	789	7	21	26	32	40	99	123	139	258	-
1980	790	7	17	34	41	58	89	111	127	254	-
1979	810	13	15	37	34	62	88	113	166	237	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
All other, both sexes											
1998	152	5	8	11	12	12	11	16	26	35	-
1997	144	8	8	9	7	13	19	18	13	30	-
1996	160	13	5	8	12	11	17	14	16	34	-
1995	151	8	7	10	4	12	19	16	25	28	-
1994	154	7	14	13	13	14	13	15	15	28	-
1993	138	3	8	7	15	7	14	19	16	26	-
1992	145	1	6	3	8	14	16	18	19	34	-
1991	162	6	1	6	10	15	26	20	24	24	-
1990	150	5	10	6	10	14	17	16	15	25	1
1989	154	8	3	3	9	17	17	29	13	23	-
1988	148	8	4	6	14	15	13	22	14	29	-
1987	146	-	9	8	7	10	18	19	18	19	1
1986	150	3	4	9	15	14	19	21	20	16	-
1985	131	1	7	11	14	9	11	18	20	17	-
1984	145	5	6	10	13	12	24	9	16	26	-
1983	167	4	4	13	13	16	23	17	23	21	-
1982	149	1	8	13	11	14	18	19	19	18	-
1981	163	10	5	6	10	15	26	14	21	27	-
1980	173	9	15	15	13	13	20	16	20	14	-
1979	161	-	10	12	9	16	19	16	11	19	-
Hernia (550-553)											
All other, male											
1998	59	1	5	5	4	5	5	10	6	5	-
1997	80	5	5	6	5	10	6	9	7	12	-
1996	77	7	5	2	5	2	12	5	9	14	-
1995	78	2	4	5	3	5	9	9	12	10	-
1994	67	2	6	5	6	5	7	6	8	9	-
1993	63	2	4	2	5	4	12	7	5	9	-
1992	69	-	1	1	5	8	3	8	8	18	-
1991	80	4	-	2	5	5	15	10	8	11	-
1990	68	2	5	4	3	4	7	7	4	10	1
1989	78	4	-	2	6	8	8	11	5	12	-
1988	71	4	2	4	6	5	6	8	8	14	-
1987	75	-	5	6	3	5	6	7	7	12	1
1986	70	3	3	3	5	7	10	7	7	4	-
1985	61	1	3	3	9	4	2	6	12	9	-
1984	73	4	4	9	4	6	9	3	10	11	-
1983	88	1	2	3	8	8	14	7	12	9	-
1982	86	1	6	5	6	5	12	8	10	10	-
1981	84	3	1	4	3	10	13	8	13	10	-
1980	104	6	6	8	8	7	11	9	14	10	-
1979	89	-	7	8	3	10	7	8	5	11	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
All other, female											
1998	93	4	3	6	8	7	6	6	20	30	-
1997	64	3	3	3	2	3	13	9	6	18	-
1996	83	6	-	6	7	9	5	9	7	20	-
1995	73	6	3	5	1	7	10	7	13	18	-
1994	87	5	8	8	7	9	6	9	7	19	-
1993	75	1	4	5	10	3	2	12	11	17	-
1992	76	1	5	2	3	6	13	10	11	16	-
1991	82	2	1	4	5	10	11	10	16	13	-
1990	82	3	5	2	7	10	10	9	11	15	-
1989	76	4	3	1	3	9	9	18	8	11	-
1988	77	4	2	2	8	10	7	14	6	15	-
1987	71	-	4	2	4	5	12	12	11	7	-
1986	80	-	1	6	10	7	9	14	13	12	-
1985	70	-	4	8	5	5	9	12	8	8	-
1984	72	1	2	1	9	6	15	6	6	15	-
1983	79	3	2	10	5	8	9	10	11	12	-
1982	63	-	2	8	5	9	6	11	9	8	-
1981	79	7	4	2	7	5	13	6	8	17	-
1980	69	3	9	7	5	6	9	7	6	4	-
1979	72	-	3	4	6	6	12	8	6	8	-
Hernia (550-553)											
Black, both sexes											
1998	138	5	7	10	12	9	9	16	25	31	-
1997	126	7	7	9	6	13	17	13	12	25	-
1996	147	13	5	8	12	11	14	12	16	30	-
1995	134	7	6	7	4	10	16	15	25	25	-
1994	143	7	13	10	13	14	12	15	13	27	-
1993	121	3	7	7	12	7	13	18	13	22	-
1992	133	1	5	3	7	12	16	16	17	32	-
1991	141	6	1	6	9	14	21	19	18	21	-
1990	136	5	10	6	10	11	14	16	15	20	1
1989	140	8	2	3	8	16	16	28	13	19	-
1988	136	8	4	6	13	14	12	20	12	25	-
1987	134	-	9	7	7	9	17	17	16	19	1
1986	136	3	4	7	14	14	15	20	19	14	-
1985	116	-	7	10	12	9	10	16	17	14	-
1984	132	5	5	9	11	10	23	9	14	24	-
1983	153	4	4	10	13	16	23	16	19	16	-
1982	144	1	8	13	10	14	16	19	19	17	-
1981	148	9	4	6	9	15	24	12	20	24	-
1980	160	9	14	15	13	12	19	14	18	13	-
1979	150	-	10	11	9	15	19	16	10	18	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hernia (550-553)											
Black, male											
1998	54	1	5	5	4	4	4	10	5	4	-
1997	66	5	4	6	4	10	4	4	6	10	-
1996	71	7	5	2	5	2	10	4	9	13	-
1995	70	2	3	3	3	3	8	9	12	9	-
1994	63	2	6	5	6	5	6	6	8	8	-
1993	54	2	3	2	4	4	11	6	3	8	-
1992	61	-	-	1	4	8	3	7	7	16	-
1991	68	4	-	2	5	4	12	10	5	10	-
1990	63	2	5	4	3	3	6	7	4	8	1
1989	71	4	-	2	5	8	8	10	5	10	-
1988	64	4	2	4	6	5	5	7	7	10	-
1987	70	-	5	5	3	5	6	7	6	12	1
1986	66	3	3	2	5	7	8	7	7	4	-
1985	54	-	3	2	8	4	2	6	11	7	-
1984	62	4	4	8	2	4	8	3	8	10	-
1983	81	1	2	2	8	8	14	7	9	7	-
1982	82	1	6	5	5	5	10	8	10	10	-
1981	75	2	1	4	3	10	11	6	12	10	-
1980	97	6	6	8	8	6	10	7	13	10	-
1979	84	-	7	8	3	10	7	8	5	10	-
Hernia (550-553)											
Black, female											
1998	84	4	2	5	8	5	5	6	20	27	-
1997	60	2	3	3	2	3	13	9	6	15	-
1996	76	6	-	6	7	9	4	8	7	17	-
1995	64	5	3	4	1	7	8	6	13	16	-
1994	80	5	7	5	7	9	6	9	5	19	-
1993	67	1	4	5	8	3	2	12	10	14	-
1992	72	1	5	2	3	4	13	9	10	16	-
1991	73	2	1	4	4	10	9	9	13	11	-
1990	73	3	5	2	7	8	8	9	11	12	-
1989	69	4	2	1	3	8	8	18	8	9	-
1988	72	4	2	2	7	9	7	13	5	15	-
1987	64	-	4	2	4	4	11	10	10	7	-
1986	70	-	1	5	9	7	7	13	12	10	-
1985	62	-	4	8	4	5	8	10	6	7	-
1984	70	1	1	1	9	6	15	6	6	14	-
1983	72	3	2	8	5	8	9	9	10	9	-
1982	62	-	2	8	5	9	6	11	9	7	-
1981	73	7	3	2	6	5	13	6	8	14	-
1980	63	3	8	7	5	6	9	7	5	3	-
1979	66	-	3	3	6	5	12	8	5	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
All races, both sexes											
1998	25,192	2,966	2,778	2,643	2,636	2,777	2,878	2,282	1,421	887	5
1997	25,175	2,889	2,733	2,512	2,741	2,924	2,880	2,185	1,353	884	4
1996	25,047	2,845	2,586	2,558	2,754	2,902	2,823	2,140	1,367	805	10
1995	25,222	2,680	2,567	2,499	2,857	3,014	2,895	2,144	1,348	831	9
1994	25,406	2,474	2,569	2,605	2,925	3,262	2,896	2,041	1,330	809	7
1993	25,209	2,303	2,400	2,618	2,995	3,203	2,891	2,104	1,387	741	13
1992	25,263	2,221	2,348	2,644	3,136	3,438	2,826	2,043	1,371	789	6
1991	25,429	2,154	2,296	2,641	3,406	3,370	2,852	2,184	1,266	730	18
1990	25,815	2,220	2,280	2,875	3,431	3,502	2,788	2,115	1,304	706	15
1989	26,694	2,202	2,519	3,101	3,655	3,615	2,877	2,135	1,205	704	9
1988	26,409	2,209	2,571	3,126	3,854	3,478	2,788	1,978	1,105	590	8
1987	26,201	2,080	2,538	3,330	3,748	3,461	2,741	1,986	1,048	598	8
1986	26,159	2,013	2,619	3,327	3,836	3,563	2,892	1,899	951	578	8
1985	26,767	2,174	2,791	3,570	3,971	3,637	2,742	1,835	1,046	537	11
1984	27,317	2,231	2,983	3,784	4,035	3,773	2,806	1,869	981	491	7
1983	27,266	2,308	3,132	3,920	4,132	3,614	2,733	1,726	944	474	9
1982	27,690	2,319	3,375	4,064	4,096	3,754	2,720	1,661	905	437	7
1981	29,308	2,576	3,811	4,326	4,300	3,980	2,762	1,644	834	489	9
1980	30,583	2,924	4,126	4,613	4,426	4,026	2,683	1,589	775	446	15
1979	29,720	2,959	4,157	4,494	4,277	3,847	2,563	1,435	714	397	10
Chronic liver disease and cirrhosis (571)											
All races, male											
1998	16,343	2,234	2,062	1,898	1,837	1,748	1,648	1,156	671	386	5
1997	16,260	2,177	2,021	1,781	1,873	1,827	1,620	1,141	650	341	4
1996	16,311	2,134	1,884	1,814	1,876	1,844	1,599	1,113	686	330	7
1995	16,525	1,997	1,890	1,730	1,940	1,920	1,756	1,114	649	354	9
1994	16,513	1,839	1,820	1,850	1,992	2,108	1,636	1,058	676	326	7
1993	16,276	1,724	1,727	1,820	1,976	2,006	1,687	1,077	669	324	10
1992	16,487	1,657	1,665	1,875	2,106	2,158	1,655	1,103	688	332	6
1991	16,269	1,562	1,612	1,857	2,270	2,100	1,600	1,140	640	302	14
1990	16,627	1,631	1,621	1,964	2,299	2,164	1,636	1,116	642	305	12
1989	17,310	1,583	1,795	2,116	2,402	2,241	1,695	1,151	581	302	8
1988	17,196	1,600	1,816	2,170	2,530	2,213	1,659	1,027	543	279	7
1987	17,051	1,487	1,801	2,312	2,500	2,179	1,619	1,087	499	274	7
1986	16,797	1,430	1,772	2,289	2,433	2,259	1,713	1,035	461	264	7
1985	17,244	1,507	1,903	2,402	2,687	2,280	1,645	989	509	222	9
1984	17,558	1,564	1,982	2,524	2,645	2,396	1,671	1,051	454	225	6
1983	17,530	1,580	2,089	2,593	2,685	2,360	1,650	953	480	204	7
1982	17,917	1,584	2,330	2,654	2,720	2,397	1,683	886	461	206	6
1981	18,894	1,710	2,538	2,853	2,819	2,589	1,660	955	420	238	5
1980	19,768	1,975	2,731	3,027	2,952	2,584	1,677	897	393	219	13
1979	19,369	1,996	2,727	2,929	2,875	2,537	1,608	822	374	192	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
All races, female											
1998	8,849	732	716	745	799	1,029	1,230	1,126	750	501	-
1997	8,915	712	712	731	868	1,097	1,260	1,044	703	543	-
1996	8,736	711	702	744	878	1,058	1,224	1,027	681	475	3
1995	8,697	683	677	769	917	1,094	1,139	1,030	699	477	-
1994	8,893	635	749	755	933	1,154	1,260	983	654	483	-
1993	8,933	579	673	798	1,019	1,197	1,204	1,027	718	417	3
1992	8,776	564	683	769	1,030	1,280	1,171	940	683	457	-
1991	9,160	592	684	784	1,136	1,270	1,252	1,044	626	428	4
1990	9,188	589	659	911	1,132	1,338	1,152	999	662	401	3
1989	9,384	619	724	985	1,253	1,374	1,182	984	624	402	1
1988	9,213	609	755	956	1,324	1,265	1,129	951	562	311	1
1987	9,150	593	737	1,018	1,248	1,282	1,122	899	549	324	1
1986	9,362	583	847	1,038	1,403	1,304	1,179	864	490	314	1
1985	9,523	667	888	1,168	1,284	1,357	1,097	846	537	315	2
1984	9,759	667	1,001	1,260	1,390	1,377	1,135	818	527	266	1
1983	9,736	728	1,043	1,327	1,447	1,254	1,083	773	464	270	2
1982	9,773	735	1,045	1,410	1,376	1,357	1,037	775	444	231	1
1981	10,414	866	1,273	1,473	1,481	1,391	1,102	689	414	251	4
1980	10,815	949	1,395	1,586	1,474	1,442	1,006	692	382	227	2
1979	10,351	963	1,430	1,565	1,402	1,310	955	613	340	205	1
Chronic liver disease and cirrhosis (571)											
White, both sexes											
1998	21,771	2,375	2,296	2,217	2,262	2,470	2,653	2,156	1,332	838	3
1997	21,683	2,281	2,267	2,088	2,345	2,605	2,624	2,042	1,281	841	4
1996	21,422	2,292	2,113	2,126	2,356	2,575	2,559	2,004	1,288	747	8
1995	21,432	2,143	2,050	2,004	2,460	2,689	2,653	2,009	1,247	773	7
1994	21,476	1,883	2,048	2,163	2,520	2,869	2,635	1,896	1,254	744	7
1993	21,348	1,804	1,929	2,207	2,588	2,841	2,637	1,946	1,306	699	8
1992	21,286	1,710	1,884	2,163	2,717	2,994	2,565	1,912	1,275	736	5
1991	21,386	1,656	1,816	2,174	2,962	2,951	2,605	2,050	1,186	686	11
1990	21,478	1,648	1,795	2,318	2,950	3,155	2,534	1,956	1,207	665	10
1989	22,211	1,659	2,006	2,536	3,164	3,205	2,616	1,986	1,136	661	4
1988	21,924	1,650	2,011	2,612	3,334	3,110	2,541	1,819	1,025	558	6
1987	21,766	1,542	1,978	2,783	3,285	3,101	2,532	1,847	991	566	3
1986	21,916	1,557	2,055	2,830	3,374	3,217	2,662	1,763	892	550	4
1985	22,192	1,613	2,241	2,984	3,483	3,249	2,511	1,702	977	501	6
1984	22,909	1,717	2,426	3,188	3,559	3,408	2,579	1,733	919	469	3
1983	22,927	1,782	2,547	3,340	3,637	3,249	2,528	1,630	896	460	8
1982	23,409	1,796	2,756	3,521	3,639	3,433	2,515	1,553	855	414	5
1981	24,465	1,967	3,073	3,718	3,806	3,594	2,532	1,531	792	464	7
1980	25,240	2,185	3,354	3,922	3,889	3,629	2,470	1,483	735	423	9
1979	24,639	2,257	3,349	3,803	3,825	3,476	2,360	1,334	672	374	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
White, male											
1998	14,096	1,830	1,721	1,591	1,576	1,543	1,507	1,097	631	360	3
1997	14,045	1,745	1,706	1,484	1,613	1,630	1,489	1,074	614	321	4
1996	13,992	1,746	1,559	1,525	1,615	1,643	1,459	1,040	642	305	5
1995	14,100	1,623	1,535	1,413	1,688	1,734	1,597	1,053	611	327	7
1994	14,015	1,406	1,469	1,548	1,743	1,871	1,495	989	629	296	7
1993	13,821	1,371	1,409	1,549	1,719	1,791	1,527	1,002	636	309	6
1992	13,910	1,285	1,358	1,563	1,839	1,886	1,498	1,028	634	311	5
1991	13,789	1,226	1,304	1,536	1,990	1,852	1,480	1,073	608	283	9
1990	13,889	1,232	1,284	1,611	1,993	1,961	1,490	1,039	592	284	8
1989	14,414	1,211	1,451	1,745	2,107	1,992	1,548	1,071	540	277	4
1988	14,381	1,223	1,446	1,821	2,223	1,998	1,525	947	511	267	6
1987	14,175	1,106	1,432	1,954	2,204	1,959	1,497	1,005	471	255	3
1986	14,099	1,123	1,410	1,960	2,151	2,051	1,586	956	428	249	3
1985	14,321	1,134	1,538	2,017	2,367	2,039	1,515	925	478	204	6
1984	14,783	1,212	1,613	2,147	2,365	2,164	1,546	971	427	215	3
1983	14,823	1,242	1,713	2,224	2,380	2,130	1,531	897	458	200	6
1982	15,249	1,246	1,923	2,324	2,413	2,215	1,561	822	437	199	5
1981	15,868	1,322	2,058	2,476	2,506	2,357	1,527	891	396	226	3
1980	16,407	1,507	2,226	2,604	2,601	2,337	1,547	838	371	206	7
1979	16,104	1,535	2,196	2,471	2,586	2,316	1,470	766	353	176	7
Chronic liver disease and cirrhosis (571)											
White, female											
1998	7,675	545	575	626	686	927	1,146	1,059	701	478	-
1997	7,638	536	561	604	732	975	1,135	968	667	520	-
1996	7,430	546	554	601	741	932	1,100	964	646	442	3
1995	7,332	520	515	591	772	955	1,056	956	636	446	-
1994	7,461	477	579	615	777	998	1,140	907	625	448	-
1993	7,527	433	520	658	869	1,050	1,110	944	670	390	2
1992	7,376	425	526	600	878	1,108	1,067	884	641	425	-
1991	7,597	430	512	638	972	1,099	1,125	977	578	403	2
1990	7,589	416	511	707	957	1,194	1,044	917	615	381	2
1989	7,797	448	555	791	1,057	1,213	1,068	915	596	384	-
1988	7,543	427	565	791	1,111	1,112	1,016	872	514	291	-
1987	7,591	436	546	829	1,081	1,142	1,035	842	520	311	-
1986	7,817	434	645	870	1,223	1,166	1,076	807	464	301	1
1985	7,871	479	703	967	1,116	1,210	996	777	499	297	-
1984	8,126	505	813	1,041	1,194	1,244	1,033	762	492	254	-
1983	8,104	540	834	1,116	1,257	1,119	997	733	438	260	2
1982	8,160	550	833	1,197	1,226	1,218	954	731	418	215	-
1981	8,597	645	1,015	1,242	1,300	1,237	1,005	640	396	238	4
1980	8,833	678	1,128	1,318	1,288	1,292	923	645	364	217	2
1979	8,535	722	1,153	1,332	1,239	1,160	890	568	319	198	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
All other, both sexes											
1998	3,421	591	482	426	374	307	225	126	89	49	2
1997	3,492	608	466	424	396	319	256	143	72	43	-
1996	3,625	553	473	432	398	327	264	136	79	58	2
1995	3,790	537	517	495	397	325	242	135	101	58	2
1994	3,930	591	521	442	405	393	261	145	76	65	-
1993	3,861	499	471	411	407	362	254	158	81	42	5
1992	3,977	511	464	481	419	444	261	131	96	53	1
1991	4,043	498	480	467	444	419	247	134	80	44	7
1990	4,337	572	485	557	481	347	254	159	97	41	5
1989	4,483	543	513	565	491	410	261	149	69	43	5
1988	4,485	559	560	514	520	368	247	159	80	32	2
1987	4,435	538	560	547	463	360	209	139	57	32	5
1986	4,243	456	564	497	462	346	230	136	59	28	4
1985	4,575	561	550	586	488	388	231	133	69	36	5
1984	4,408	514	557	596	476	365	227	136	62	22	4
1983	4,339	526	585	580	495	365	205	96	48	14	1
1982	4,281	523	619	543	457	321	205	108	50	23	2
1981	4,843	609	738	608	494	386	230	113	42	25	2
1980	5,343	739	772	691	537	397	213	106	40	23	6
1979	5,081	702	808	691	452	371	203	101	42	23	2
Chronic liver disease and cirrhosis (571)											
All other, male											
1998	2,247	404	341	307	261	205	141	59	40	26	2
1997	2,215	432	315	297	260	197	131	67	36	20	-
1996	2,319	388	325	289	261	201	140	73	44	25	2
1995	2,425	374	355	317	252	186	159	61	38	27	2
1994	2,498	433	351	302	249	237	141	69	47	30	-
1993	2,455	353	318	271	257	215	160	75	33	15	4
1992	2,577	372	307	312	267	272	157	75	54	21	1
1991	2,480	336	308	321	280	248	120	67	32	19	5
1990	2,738	399	337	353	306	203	146	77	50	21	4
1989	2,896	372	344	371	295	249	147	80	41	25	4
1988	2,815	377	370	349	307	215	134	80	32	12	1
1987	2,876	381	369	358	296	220	122	82	28	19	4
1986	2,698	307	362	329	282	208	127	79	33	15	4
1985	2,923	373	365	385	320	241	130	64	31	18	3
1984	2,775	352	369	377	280	232	125	80	27	10	3
1983	2,707	338	376	369	305	230	119	56	22	4	1
1982	2,668	338	407	330	307	182	122	64	24	7	1
1981	3,026	388	480	377	313	232	133	64	24	12	2
1980	3,361	468	505	423	351	247	130	59	22	13	6
1979	3,265	461	531	458	289	221	138	56	21	16	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
All other, female											
1998	1,174	187	141	119	113	102	84	67	49	23	-
1997	1,277	176	151	127	136	122	125	76	36	23	-
1996	1,306	165	148	143	137	126	124	63	35	33	-
1995	1,365	163	162	178	145	139	83	74	63	31	-
1994	1,432	158	170	140	156	156	120	76	29	35	-
1993	1,406	146	153	140	150	147	94	83	48	27	1
1992	1,400	139	157	169	152	172	104	56	42	32	-
1991	1,563	162	172	146	164	171	127	67	48	25	2
1990	1,599	173	148	204	175	144	108	82	47	20	1
1989	1,587	171	169	194	196	161	114	69	28	18	1
1988	1,670	182	190	165	213	153	113	79	48	20	1
1987	1,559	157	191	189	167	140	87	57	29	13	1
1986	1,545	149	202	168	180	138	103	57	26	13	-
1985	1,652	188	185	201	168	147	101	69	38	18	2
1984	1,633	162	188	219	196	133	102	56	35	12	1
1983	1,632	188	209	211	190	135	86	40	26	10	-
1982	1,613	185	212	213	150	139	83	44	26	16	1
1981	1,817	221	258	231	181	154	97	49	18	13	-
1980	1,982	271	267	268	186	150	83	47	18	10	-
1979	1,816	241	277	233	163	150	65	45	21	7	-
Chronic liver disease and cirrhosis (571)											
Black, both sexes											
1998	2,697	490	383	345	295	246	186	95	67	32	2
1997	2,820	518	376	358	338	255	200	113	53	37	-
1996	2,950	476	402	354	324	278	216	98	49	39	2
1995	3,073	463	429	389	334	270	195	102	73	47	2
1994	3,245	507	436	369	334	330	210	115	55	50	-
1993	3,228	419	412	354	342	308	210	129	56	30	4
1992	3,315	446	382	398	358	372	200	101	77	39	1
1991	3,464	429	419	393	388	370	193	105	63	32	7
1990	3,753	499	417	490	415	293	222	127	76	35	5
1989	3,851	471	443	483	434	348	223	117	54	31	4
1988	3,903	481	484	438	458	325	203	136	61	28	2
1987	3,916	473	493	477	400	305	181	119	48	24	5
1986	3,745	402	486	448	407	313	196	118	38	16	4
1985	4,055	506	476	521	436	340	204	113	54	27	5
1984	3,901	450	492	538	429	319	201	115	51	19	4
1983	3,883	455	537	525	449	318	179	81	42	13	1
1982	3,839	468	555	488	424	290	177	90	38	19	1
1981	4,349	548	662	566	439	355	202	95	36	18	2
1980	4,790	660	708	633	492	352	188	88	31	19	6
1979	4,567	629	742	626	414	338	189	88	26	19	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic liver disease and cirrhosis (571)											
Black, male											
1998	1,804	346	276	246	210	164	118	47	31	14	2
1997	1,818	377	255	247	227	164	108	54	27	15	-
1996	1,923	336	287	237	215	174	117	57	28	12	2
1995	1,999	330	300	252	221	160	133	45	28	23	2
1994	2,097	373	297	252	207	208	117	54	33	20	-
1993	2,100	294	285	243	223	187	135	64	22	10	4
1992	2,181	326	256	263	238	228	127	54	39	17	1
1991	2,137	294	270	277	245	216	92	53	24	11	5
1990	2,393	350	290	321	264	173	130	59	41	17	4
1989	2,517	326	306	320	264	216	124	62	31	14	3
1988	2,476	325	327	301	275	186	112	69	22	10	1
1987	2,574	339	329	322	266	188	101	73	21	14	4
1986	2,404	267	316	300	245	195	107	70	21	12	4
1985	2,616	337	320	349	288	211	114	52	23	14	3
1984	2,458	311	324	341	253	201	109	63	23	9	3
1983	2,429	298	349	334	275	204	103	43	20	4	1
1982	2,408	302	368	302	288	165	103	52	19	6	-
1981	2,740	355	437	352	274	218	117	56	20	6	2
1980	3,020	421	462	388	318	219	112	48	17	12	6
1979	2,954	410	495	415	269	200	127	47	12	14	2
Chronic liver disease and cirrhosis (571)											
Black, female											
1998	893	144	107	99	85	82	68	48	36	18	-
1997	1,002	141	121	111	111	91	92	59	26	22	-
1996	1,027	140	115	117	109	104	99	41	21	27	-
1995	1,074	133	129	137	113	110	62	57	45	24	-
1994	1,148	134	139	117	127	122	93	61	22	30	-
1993	1,128	125	127	111	119	121	75	65	34	20	-
1992	1,134	120	126	135	120	144	73	47	38	22	-
1991	1,327	135	149	116	143	154	101	52	39	21	2
1990	1,360	149	127	169	151	120	92	68	35	18	1
1989	1,334	145	137	163	170	132	99	55	23	17	1
1988	1,427	156	157	137	183	139	91	67	39	18	1
1987	1,342	134	164	155	134	117	80	46	27	10	1
1986	1,341	135	170	148	162	118	89	48	17	4	-
1985	1,439	169	156	172	148	129	90	61	31	13	2
1984	1,443	139	168	197	176	118	92	52	28	10	1
1983	1,454	157	188	191	174	114	76	38	22	9	-
1982	1,431	166	187	186	136	125	74	38	19	13	1
1981	1,609	193	225	214	165	137	85	39	16	12	-
1980	1,770	239	246	245	174	133	76	40	14	7	-
1979	1,613	219	247	211	145	138	62	41	14	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
All races, both sexes											
1998	11,798	1,777	1,685	1,462	1,343	1,203	924	511	227	104	2
1997	11,945	1,842	1,667	1,416	1,411	1,181	913	489	210	100	3
1996	12,079	1,841	1,614	1,445	1,404	1,178	876	524	260	104	8
1995	12,211	1,746	1,593	1,427	1,475	1,260	971	491	247	91	5
1994	12,241	1,585	1,561	1,435	1,486	1,414	969	506	216	103	5
1993	11,653	1,450	1,437	1,416	1,466	1,307	912	477	211	72	11
1992	11,868	1,415	1,404	1,517	1,550	1,379	924	450	222	96	2
1991	11,688	1,316	1,338	1,427	1,602	1,340	905	519	222	77	9
1990	11,985	1,383	1,317	1,561	1,637	1,413	907	489	212	76	10
1989	12,308	1,383	1,479	1,681	1,695	1,446	882	485	211	65	6
1988	11,792	1,274	1,436	1,605	1,771	1,345	824	449	181	58	4
1987	11,265	1,216	1,359	1,685	1,597	1,269	762	409	173	64	6
1986	11,060	1,099	1,383	1,553	1,635	1,345	829	414	134	58	6
1985	11,288	1,179	1,419	1,710	1,722	1,342	779	379	155	38	6
1984	11,386	1,199	1,452	1,784	1,706	1,340	828	421	145	50	5
1983	11,076	1,167	1,527	1,774	1,670	1,324	767	381	128	35	3
1982	11,293	1,158	1,672	1,827	1,670	1,371	739	329	137	36	4
1981	12,085	1,309	1,910	1,994	1,732	1,409	757	367	105	60	5
1980	12,938	1,561	1,993	2,075	1,848	1,494	780	349	104	56	6
1979	12,547	1,513	2,007	2,098	1,766	1,434	755	277	104	42	4
Alcoholic liver disease (571.0-571.3)											
All races, male											
1998	8,708	1,336	1,306	1,123	1,038	877	687	355	152	79	2
1997	8,742	1,408	1,261	1,063	1,059	883	647	344	138	70	3
1996	8,853	1,397	1,202	1,076	1,059	864	631	355	186	77	6
1995	9,032	1,307	1,227	1,065	1,097	929	723	340	172	65	5
1994	8,988	1,200	1,163	1,067	1,106	1,062	695	347	152	64	5
1993	8,552	1,127	1,077	1,035	1,074	947	663	327	152	55	9
1992	8,777	1,083	1,030	1,153	1,148	1,019	675	314	155	57	2
1991	8,487	988	979	1,074	1,184	965	645	364	156	48	5
1990	8,676	1,031	980	1,132	1,231	990	631	346	144	56	9
1989	9,012	1,026	1,074	1,241	1,220	1,035	658	355	154	42	6
1988	8,530	944	1,055	1,179	1,244	991	610	310	130	41	4
1987	8,221	893	1,004	1,246	1,180	928	555	291	122	41	6
1986	7,931	811	983	1,131	1,148	982	591	288	93	42	5
1985	8,129	851	1,014	1,249	1,271	943	558	289	113	23	4
1984	8,107	876	1,014	1,278	1,206	932	594	299	95	36	4
1983	7,828	829	1,070	1,238	1,190	952	545	263	89	27	2
1982	7,997	817	1,188	1,280	1,194	976	534	227	97	24	4
1981	8,478	902	1,316	1,382	1,216	1,026	533	268	78	40	3
1980	9,088	1,101	1,368	1,444	1,332	1,075	562	257	62	39	4
1979	8,793	1,078	1,337	1,423	1,275	1,029	560	199	81	24	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
All races, female											
1998	3,090	441	379	339	305	326	237	156	75	25	-
1997	3,203	434	406	353	352	298	266	145	72	30	-
1996	3,226	444	412	369	345	314	245	169	74	27	2
1995	3,179	439	366	362	378	331	248	151	75	26	-
1994	3,253	385	398	368	380	352	274	159	64	39	-
1993	3,101	323	360	381	392	360	249	150	59	17	2
1992	3,091	332	374	364	402	360	249	136	67	39	-
1991	3,201	328	359	353	418	375	260	155	66	29	4
1990	3,309	352	337	429	406	423	276	143	68	20	1
1989	3,296	357	405	440	475	411	224	130	57	23	-
1988	3,262	330	381	426	527	354	214	139	51	17	-
1987	3,044	323	355	439	417	341	207	118	51	23	-
1986	3,129	288	400	422	487	363	238	126	41	16	1
1985	3,159	328	405	461	451	399	221	90	42	15	2
1984	3,279	323	438	506	500	408	234	122	50	14	1
1983	3,248	338	457	536	480	372	222	118	39	8	1
1982	3,296	341	484	547	476	395	205	102	40	12	-
1981	3,607	407	594	612	516	383	224	99	27	20	2
1980	3,850	460	625	631	516	419	218	92	42	17	2
1979	3,754	435	670	675	491	405	195	78	23	18	1
Alcoholic liver disease (571.0-571.3)											
White, both sexes											
1998	9,947	1,445	1,388	1,216	1,137	1,061	832	475	204	98	1
1997	10,061	1,485	1,388	1,188	1,196	1,056	804	448	188	93	3
1996	10,048	1,486	1,320	1,207	1,172	1,027	777	481	242	92	6
1995	10,067	1,413	1,264	1,135	1,250	1,108	850	458	224	81	3
1994	9,991	1,199	1,233	1,170	1,279	1,204	850	458	199	86	5
1993	9,560	1,141	1,148	1,189	1,268	1,141	814	432	194	64	8
1992	9,593	1,076	1,121	1,222	1,322	1,175	819	413	199	84	2
1991	9,393	1,006	1,041	1,162	1,381	1,123	813	468	205	68	5
1990	9,482	1,000	1,011	1,240	1,384	1,247	810	443	186	68	6
1989	9,689	1,018	1,156	1,349	1,422	1,257	778	447	190	59	3
1988	9,236	914	1,098	1,314	1,494	1,172	736	395	157	50	3
1987	8,852	875	1,052	1,391	1,367	1,114	690	368	152	56	3
1986	8,774	832	1,054	1,302	1,393	1,189	741	372	116	54	4
1985	8,821	851	1,102	1,383	1,487	1,171	695	353	140	27	3
1984	9,078	916	1,148	1,473	1,472	1,188	745	380	134	45	1
1983	8,937	888	1,224	1,489	1,471	1,171	693	352	117	33	3
1982	9,159	886	1,346	1,553	1,459	1,234	666	300	125	32	3
1981	9,657	974	1,514	1,679	1,509	1,252	692	330	97	54	4
1980	10,222	1,142	1,579	1,738	1,608	1,314	714	316	92	50	4
1979	9,967	1,130	1,576	1,742	1,552	1,267	687	252	102	35	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
White, male											
1998	7,417	1,113	1,083	935	885	778	612	331	137	75	1
1997	7,485	1,158	1,075	897	899	789	578	320	123	68	3
1996	7,484	1,143	1,009	913	896	759	570	322	174	67	4
1995	7,569	1,073	998	871	944	824	631	321	158	56	3
1994	7,464	914	929	883	964	915	614	320	141	54	5
1993	7,112	900	877	876	927	836	589	298	141	49	6
1992	7,211	833	843	941	994	876	598	292	137	52	2
1991	6,985	766	791	882	1,025	819	592	334	148	43	3
1990	7,001	761	756	919	1,058	884	558	316	128	51	6
1989	7,225	771	861	1,011	1,036	910	582	331	135	37	3
1988	6,806	697	817	970	1,061	871	550	272	116	37	3
1987	6,540	649	793	1,042	1,016	811	506	261	109	35	3
1986	6,369	629	748	958	978	875	533	255	81	39	3
1985	6,455	627	790	1,020	1,110	833	497	270	103	16	3
1984	6,579	672	805	1,067	1,069	827	546	269	88	33	1
1983	6,421	642	865	1,061	1,055	848	496	243	80	26	2
1982	6,593	636	961	1,104	1,042	890	482	205	90	23	3
1981	6,868	678	1,053	1,184	1,064	915	489	238	72	37	2
1980	7,285	836	1,091	1,226	1,168	950	511	235	55	33	2
1979	7,080	815	1,063	1,185	1,124	917	506	180	79	19	3
Alcoholic liver disease (571.0-571.3)											
White, female											
1998	2,530	332	305	281	252	283	220	144	67	23	-
1997	2,576	327	313	291	297	267	226	128	65	25	-
1996	2,564	343	311	294	276	268	207	159	68	25	2
1995	2,498	340	266	264	306	284	219	137	66	25	-
1994	2,527	285	304	287	315	289	236	138	58	32	-
1993	2,448	241	271	313	341	305	225	134	53	15	2
1992	2,382	243	278	281	328	299	221	121	62	32	-
1991	2,408	240	250	280	356	304	221	134	57	25	2
1990	2,481	239	255	321	326	363	252	127	58	17	-
1989	2,464	247	295	338	386	347	196	116	55	22	-
1988	2,430	217	281	344	433	301	186	123	41	13	-
1987	2,312	226	259	349	351	303	184	107	43	21	-
1986	2,405	203	306	344	415	314	208	117	35	15	1
1985	2,366	224	312	363	377	338	198	83	37	11	-
1984	2,499	244	343	406	403	361	199	111	46	12	-
1983	2,516	246	359	428	416	323	197	109	37	7	1
1982	2,566	250	385	449	417	344	184	95	35	9	-
1981	2,789	296	461	495	445	337	203	92	25	17	2
1980	2,937	306	488	512	440	364	203	81	37	17	2
1979	2,887	315	513	557	428	350	181	72	23	16	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
All other, both sexes											
1998	1,851	332	297	246	206	142	92	36	23	6	1
1997	1,884	357	279	228	215	125	109	41	22	7	-
1996	2,031	355	294	238	232	151	99	43	18	12	2
1995	2,144	333	329	292	225	152	121	33	23	10	2
1994	2,250	386	328	265	207	210	119	48	17	17	-
1993	2,093	309	289	227	198	166	98	45	17	8	3
1992	2,275	339	283	295	228	204	105	37	23	12	-
1991	2,295	310	297	265	221	217	92	51	17	9	4
1990	2,503	383	306	321	253	166	97	46	26	8	4
1989	2,619	365	323	332	273	189	104	38	21	6	3
1988	2,556	360	338	291	277	173	88	54	24	8	1
1987	2,413	341	307	294	230	155	72	41	21	8	3
1986	2,286	267	329	251	242	156	88	42	18	4	2
1985	2,467	328	317	327	235	171	84	26	15	11	3
1984	2,308	283	304	311	234	152	83	41	11	5	4
1983	2,139	279	303	285	199	153	74	29	11	2	-
1982	2,134	272	326	274	211	137	73	29	12	4	1
1981	2,428	335	396	315	223	157	65	37	8	6	1
1980	2,716	419	414	337	240	180	66	33	12	6	2
1979	2,580	383	431	356	214	167	68	25	2	7	-
Alcoholic liver disease (571.0-571.3)											
All other, male											
1998	1,291	223	223	188	153	99	75	24	15	4	1
1997	1,257	250	186	166	160	94	69	24	15	2	-
1996	1,369	254	193	163	163	105	61	33	12	10	2
1995	1,463	234	229	194	153	105	92	19	14	9	2
1994	1,524	286	234	184	142	147	81	27	11	10	-
1993	1,440	227	200	159	147	111	74	29	11	6	3
1992	1,566	250	187	212	154	143	77	22	18	5	-
1991	1,502	222	188	192	159	146	53	30	8	5	2
1990	1,675	270	224	213	173	106	73	30	16	5	3
1989	1,787	255	213	230	184	125	76	24	19	5	3
1988	1,724	247	238	209	183	120	60	38	14	4	1
1987	1,681	244	211	204	164	117	49	30	13	6	3
1986	1,562	182	235	173	170	107	58	33	12	3	2
1985	1,674	224	224	229	161	110	61	19	10	7	1
1984	1,528	204	209	211	137	105	48	30	7	3	3
1983	1,407	187	205	177	135	104	49	20	9	1	-
1982	1,404	181	227	176	152	86	52	22	7	1	1
1981	1,610	224	263	198	152	111	44	30	6	3	1
1980	1,803	265	277	218	164	125	51	22	7	6	2
1979	1,713	263	274	238	151	112	54	19	2	5	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
All other, female											
1998	560	109	74	58	53	43	17	12	8	2	-
1997	627	107	93	62	55	31	40	17	7	5	-
1996	662	101	101	75	69	46	38	10	6	2	-
1995	681	99	100	98	72	47	29	14	9	1	-
1994	726	100	94	81	65	63	38	21	6	7	-
1993	653	82	89	68	51	55	24	16	6	2	-
1992	709	89	96	83	74	61	28	15	5	7	-
1991	793	88	109	73	62	71	39	21	9	4	2
1990	828	113	82	108	80	60	24	16	10	3	1
1989	832	110	110	102	89	64	28	14	2	1	-
1988	832	113	100	82	94	53	28	16	10	4	-
1987	732	97	96	90	66	38	23	11	8	2	-
1986	724	85	94	78	72	49	30	9	6	1	-
1985	793	104	93	98	74	61	23	7	5	4	2
1984	780	79	95	100	97	47	35	11	4	2	1
1983	732	92	98	108	64	49	25	9	2	1	-
1982	730	91	99	98	59	51	21	7	5	3	-
1981	818	111	133	117	71	46	21	7	2	3	-
1980	913	154	137	119	76	55	15	11	5	-	-
1979	867	120	157	118	63	55	14	6	-	2	-
Alcoholic liver disease (571.0-571.3)											
Black, both sexes											
1998	1,410	263	227	193	165	110	77	28	19	5	1
1997	1,501	298	215	190	186	103	95	33	19	6	-
1996	1,644	293	250	195	189	132	86	36	16	8	2
1995	1,733	280	266	223	197	131	107	28	19	8	2
1994	1,867	329	274	222	176	184	104	43	14	16	-
1993	1,770	256	250	202	173	154	85	38	14	5	3
1992	1,917	289	230	252	201	177	92	32	19	11	-
1991	1,977	264	255	225	194	194	79	45	15	9	4
1990	2,202	334	271	294	224	147	88	40	23	7	4
1989	2,254	317	275	286	248	161	90	34	18	5	2
1988	2,238	309	295	253	250	152	78	51	18	8	1
1987	2,176	303	280	259	206	141	68	38	17	6	3
1986	2,058	240	292	234	219	144	79	39	13	3	2
1985	2,226	302	285	296	215	158	76	22	14	10	3
1984	2,051	244	272	289	220	133	72	36	10	5	4
1983	1,904	236	281	263	179	132	68	23	10	2	-
1982	1,903	244	286	249	197	126	65	27	10	4	-
1981	2,158	302	358	287	201	141	59	32	6	5	1
1980	2,406	370	372	316	220	162	56	28	11	5	2
1979	2,295	344	394	321	197	150	65	22	1	7	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Alcoholic liver disease (571.0-571.3)											
Black, male											
1998	1,003	183	176	148	124	73	62	17	12	3	1
1997	1,013	212	144	138	142	78	59	19	13	1	-
1996	1,119	213	170	133	133	92	51	29	10	6	2
1995	1,188	204	188	148	134	91	81	16	10	7	2
1994	1,289	245	201	153	121	133	70	23	9	9	-
1993	1,239	186	180	143	131	101	64	26	10	4	3
1992	1,333	213	154	182	140	122	67	19	15	5	-
1991	1,299	191	163	164	139	130	43	26	7	5	2
1990	1,493	240	199	201	154	95	66	25	15	4	3
1989	1,566	225	187	202	170	108	67	21	16	4	2
1988	1,527	217	213	182	166	102	51	36	9	4	1
1987	1,531	222	193	183	151	107	46	28	9	4	3
1986	1,412	163	213	159	154	99	50	31	8	3	2
1985	1,524	207	203	211	150	100	56	15	9	6	1
1984	1,365	181	182	195	131	90	41	27	6	3	3
1983	1,253	160	191	164	121	92	43	15	9	1	-
1982	1,256	163	201	163	140	77	45	21	6	1	-
1981	1,448	205	245	182	135	99	38	27	5	2	1
1980	1,601	236	248	204	149	112	43	19	6	5	2
1979	1,541	237	255	214	141	100	51	16	1	5	-
Alcoholic liver disease (571.0-571.3)											
Black, female											
1998	407	80	51	45	41	37	15	11	7	2	-
1997	488	86	71	52	44	25	36	14	6	5	-
1996	525	80	80	62	56	40	35	7	6	2	-
1995	545	76	78	75	63	40	26	12	9	1	-
1994	578	84	73	69	55	51	34	20	5	7	-
1993	531	70	70	59	42	53	21	12	4	1	-
1992	584	76	76	70	61	55	25	13	4	6	-
1991	678	73	92	61	55	64	36	19	8	4	2
1990	709	94	72	93	70	52	22	15	8	3	1
1989	688	92	88	84	78	53	23	13	2	1	-
1988	711	92	82	71	84	50	27	15	9	4	-
1987	645	81	87	76	55	34	22	10	8	2	-
1986	646	77	79	75	65	45	29	8	5	-	-
1985	702	95	82	85	65	58	20	7	5	4	2
1984	686	63	90	94	89	43	31	9	4	2	1
1983	651	76	90	99	58	40	25	8	1	1	-
1982	647	81	85	86	57	49	20	6	4	3	-
1981	710	97	113	105	66	42	21	5	1	3	-
1980	805	134	124	112	71	50	13	9	5	-	-
1979	754	107	139	107	56	50	14	6	-	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
All races, both sexes											
1998	13,394	1,189	1,093	1,181	1,293	1,574	1,954	1,771	1,194	783	3
1997	13,230	1,047	1,066	1,096	1,330	1,743	1,967	1,696	1,143	784	1
1996	12,968	1,004	972	1,113	1,350	1,724	1,947	1,616	1,107	701	2
1995	13,011	934	974	1,072	1,382	1,754	1,924	1,653	1,101	740	4
1994	13,165	889	1,008	1,170	1,439	1,848	1,927	1,535	1,114	706	2
1993	13,556	853	963	1,202	1,529	1,896	1,979	1,627	1,176	669	2
1992	13,395	806	944	1,127	1,586	2,059	1,902	1,593	1,149	693	4
1991	13,741	838	958	1,214	1,804	2,030	1,947	1,665	1,044	653	9
1990	13,830	837	963	1,314	1,794	2,089	1,881	1,626	1,092	630	5
1989	14,386	819	1,040	1,420	1,960	2,169	1,995	1,650	994	639	3
1988	14,617	935	1,135	1,521	2,083	2,133	1,964	1,529	924	532	4
1987	14,936	864	1,179	1,645	2,151	2,192	1,979	1,577	875	534	2
1986	15,099	914	1,236	1,774	2,201	2,218	2,063	1,485	817	520	2
1985	15,479	995	1,372	1,860	2,249	2,295	1,963	1,456	891	499	5
1984	15,931	1,032	1,531	2,000	2,329	2,433	1,978	1,448	836	441	2
1983	16,190	1,141	1,605	2,146	2,462	2,290	1,966	1,345	816	439	6
1982	16,397	1,161	1,703	2,237	2,426	2,383	1,981	1,332	768	401	3
1981	17,223	1,267	1,901	2,332	2,568	2,571	2,005	1,277	729	429	4
1980	17,645	1,363	2,133	2,538	2,578	2,532	1,903	1,240	671	390	9
1979	17,173	1,446	2,150	2,396	2,511	2,413	1,808	1,158	610	355	6
Other chronic liver disease and cirrhosis (571.4-571.9)											
All races, male											
1998	7,635	898	756	775	799	871	961	801	519	307	3
1997	7,518	769	760	718	814	944	973	797	512	271	1
1996	7,458	737	682	738	817	980	968	758	500	253	1
1995	7,493	690	663	665	843	991	1,033	774	477	289	4
1994	7,525	639	657	783	886	1,046	941	711	524	262	2
1993	7,724	597	650	785	902	1,059	1,024	750	517	269	1
1992	7,710	574	635	722	958	1,139	980	789	533	275	4
1991	7,782	574	633	783	1,086	1,135	955	776	484	254	9
1990	7,951	600	641	832	1,068	1,174	1,005	770	498	249	3
1989	8,298	557	721	875	1,182	1,206	1,037	796	427	260	2
1988	8,666	656	761	991	1,286	1,222	1,049	717	413	238	3
1987	8,830	594	797	1,066	1,320	1,251	1,064	796	377	233	1
1986	8,866	619	789	1,158	1,285	1,277	1,122	747	368	222	2
1985	9,115	656	889	1,153	1,416	1,337	1,087	700	396	199	5
1984	9,451	688	968	1,246	1,439	1,464	1,077	752	359	189	2
1983	9,702	751	1,019	1,355	1,495	1,408	1,105	690	391	177	5
1982	9,920	767	1,142	1,374	1,526	1,421	1,149	659	364	182	2
1981	10,416	808	1,222	1,471	1,603	1,563	1,127	687	342	198	2
1980	10,680	874	1,363	1,583	1,620	1,509	1,115	640	331	180	9
1979	10,576	918	1,390	1,506	1,600	1,508	1,048	623	293	168	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
All races, female											
1998	5,759	291	337	406	494	703	993	970	675	476	-
1997	5,712	278	306	378	516	799	994	899	631	513	-
1996	5,510	267	290	375	533	744	979	858	607	448	1
1995	5,518	244	311	407	539	763	891	879	624	451	-
1994	5,640	250	351	387	553	802	986	824	590	444	-
1993	5,832	256	313	417	627	837	955	877	659	400	1
1992	5,685	232	309	405	628	920	922	804	616	418	-
1991	5,959	264	325	431	718	895	992	889	560	399	-
1990	5,879	237	322	482	726	915	876	856	594	381	2
1989	6,088	262	319	545	778	963	958	854	567	379	1
1988	5,951	279	374	530	797	911	915	812	511	294	1
1987	6,106	270	382	579	831	941	915	781	498	301	1
1986	6,233	295	447	616	916	941	941	738	449	298	-
1985	6,364	339	483	707	833	958	876	756	495	300	-
1984	6,480	344	563	754	890	969	901	696	477	252	-
1983	6,488	390	586	791	967	882	861	655	425	262	1
1982	6,477	394	561	863	900	962	832	673	404	219	1
1981	6,807	459	679	861	965	1,008	878	590	387	231	2
1980	6,965	489	770	955	958	1,023	788	600	340	210	-
1979	6,597	528	760	890	911	905	760	535	317	187	-
Other chronic liver disease and cirrhosis (571.4-571.9)											
White, both sexes											
1998	11,824	930	908	1,001	1,125	1,409	1,821	1,681	1,128	740	2
1997	11,622	796	879	900	1,149	1,549	1,820	1,594	1,093	748	1
1996	11,374	806	793	919	1,184	1,548	1,782	1,523	1,046	655	2
1995	11,365	730	786	869	1,210	1,581	1,803	1,551	1,023	692	4
1994	11,485	684	815	993	1,241	1,665	1,785	1,438	1,055	658	2
1993	11,788	663	781	1,018	1,320	1,700	1,823	1,514	1,112	635	-
1992	11,693	634	763	941	1,395	1,819	1,746	1,499	1,076	652	3
1991	11,993	650	775	1,012	1,581	1,828	1,792	1,582	981	618	6
1990	11,996	648	784	1,078	1,566	1,908	1,724	1,513	1,021	597	4
1989	12,522	641	850	1,187	1,742	1,948	1,838	1,539	946	602	1
1988	12,688	736	913	1,298	1,840	1,938	1,805	1,424	868	508	3
1987	12,914	667	926	1,392	1,918	1,987	1,842	1,479	839	510	-
1986	13,142	725	1,001	1,528	1,981	2,028	1,921	1,391	776	496	-
1985	13,371	762	1,139	1,601	1,996	2,078	1,816	1,349	837	474	3
1984	13,831	801	1,278	1,715	2,087	2,220	1,834	1,353	785	424	2
1983	13,990	894	1,323	1,851	2,166	2,078	1,835	1,278	779	427	5
1982	14,250	910	1,410	1,968	2,180	2,199	1,849	1,253	730	382	2
1981	14,808	993	1,559	2,039	2,297	2,342	1,840	1,201	695	410	3
1980	15,018	1,043	1,775	2,184	2,281	2,315	1,756	1,167	643	373	5
1979	14,672	1,127	1,773	2,061	2,273	2,209	1,673	1,082	570	339	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
White, male											
1998	6,679	717	638	656	691	765	895	766	494	285	2
1997	6,560	587	631	587	714	841	911	754	491	253	1
1996	6,508	603	550	612	719	884	889	718	468	238	1
1995	6,531	550	537	542	744	910	966	732	453	271	4
1994	6,551	492	540	665	779	956	881	669	488	242	2
1993	6,709	471	532	673	792	955	938	704	495	260	-
1992	6,699	452	515	622	845	1,010	900	736	497	259	3
1991	6,804	460	513	654	965	1,033	888	739	460	240	6
1990	6,888	471	528	692	935	1,077	932	723	464	233	2
1989	7,189	440	590	734	1,071	1,082	966	740	405	240	1
1988	7,575	526	629	851	1,162	1,127	975	675	395	230	3
1987	7,635	457	639	912	1,188	1,148	991	744	362	220	-
1986	7,730	494	662	1,002	1,173	1,176	1,053	701	347	210	-
1985	7,866	507	748	997	1,257	1,206	1,018	655	375	188	3
1984	8,204	540	808	1,080	1,296	1,337	1,000	702	339	182	2
1983	8,402	600	848	1,163	1,325	1,282	1,035	654	378	174	4
1982	8,656	610	962	1,220	1,371	1,325	1,079	617	347	176	2
1981	9,000	644	1,005	1,292	1,442	1,442	1,038	653	324	189	1
1980	9,122	671	1,135	1,378	1,433	1,387	1,036	603	316	173	5
1979	9,024	720	1,133	1,286	1,462	1,399	964	586	274	157	4
Other chronic liver disease and cirrhosis (571.4-571.9)											
White, female											
1998	5,145	213	270	345	434	644	926	915	634	455	-
1997	5,062	209	248	313	435	708	909	840	602	495	-
1996	4,866	203	243	307	465	664	893	805	578	417	1
1995	4,834	180	249	327	466	671	837	819	570	421	-
1994	4,934	192	275	328	462	709	904	769	567	416	-
1993	5,079	192	249	345	528	745	885	810	617	375	-
1992	4,994	182	248	319	550	809	846	763	579	393	-
1991	5,189	190	262	358	616	795	904	843	521	378	-
1990	5,108	177	256	386	631	831	792	790	557	364	2
1989	5,333	201	260	453	671	866	872	799	541	362	-
1988	5,113	210	284	447	678	811	830	749	473	278	-
1987	5,279	210	287	480	730	839	851	735	477	290	-
1986	5,412	231	339	526	808	852	868	690	429	286	-
1985	5,505	255	391	604	739	872	798	694	462	286	-
1984	5,627	261	470	635	791	883	834	651	446	242	-
1983	5,588	294	475	688	841	796	800	624	401	253	1
1982	5,594	300	448	748	809	874	770	636	383	206	-
1981	5,808	349	554	747	855	900	802	548	371	221	2
1980	5,896	372	640	806	848	928	720	564	327	200	-
1979	5,648	407	640	775	811	810	709	496	296	182	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
All other, both sexes											
1998	1,570	259	185	180	168	165	133	90	66	43	1
1997	1,608	251	187	196	181	194	147	102	50	36	-
1996	1,594	198	179	194	166	176	165	93	61	46	-
1995	1,646	204	188	203	172	173	121	102	78	48	-
1994	1,680	205	193	177	198	183	142	97	59	48	-
1993	1,768	190	182	184	209	196	156	113	64	34	2
1992	1,702	172	181	186	191	240	156	94	73	41	1
1991	1,748	188	183	202	223	202	155	83	63	35	3
1990	1,834	189	179	236	228	181	157	113	71	33	1
1989	1,864	178	190	233	218	221	157	111	48	37	2
1988	1,929	199	222	223	243	195	159	105	56	24	1
1987	2,022	197	253	253	233	205	137	98	36	24	2
1986	1,957	189	235	246	220	190	142	94	41	24	2
1985	2,108	233	233	259	253	217	147	107	54	25	2
1984	2,100	231	253	285	242	213	144	95	51	17	-
1983	2,200	247	282	295	296	212	131	67	37	12	1
1982	2,147	251	293	269	246	184	132	79	38	19	1
1981	2,415	274	342	293	271	229	165	76	34	19	1
1980	2,627	320	358	354	297	217	147	73	28	17	4
1979	2,501	319	377	335	238	204	135	76	40	16	2
Other chronic liver disease and cirrhosis (571.4-571.9)											
All other, male											
1998	956	181	118	119	108	106	66	35	25	22	1
1997	958	182	129	131	100	103	62	43	21	18	-
1996	950	134	132	126	98	96	79	40	32	15	-
1995	962	140	126	123	99	81	67	42	24	18	-
1994	974	147	117	118	107	90	60	42	36	20	-
1993	1,015	126	118	112	110	104	86	46	22	9	1
1992	1,011	122	120	100	113	129	80	53	36	16	1
1991	978	114	120	129	121	102	67	37	24	14	3
1990	1,063	129	113	140	133	97	73	47	34	16	1
1989	1,109	117	131	141	111	124	71	56	22	20	1
1988	1,091	130	132	140	124	95	74	42	18	8	-
1987	1,195	137	158	154	132	103	73	52	15	13	1
1986	1,136	125	127	156	112	101	69	46	21	12	2
1985	1,249	149	141	156	159	131	69	45	21	11	2
1984	1,247	148	160	166	143	127	77	50	20	7	-
1983	1,300	151	171	192	170	126	70	36	13	3	1
1982	1,264	157	180	154	155	96	70	42	17	6	-
1981	1,416	164	217	179	161	121	89	34	18	9	1
1980	1,558	203	228	205	187	122	79	37	15	7	4
1979	1,552	198	257	220	138	109	84	37	19	11	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
All other, female											
1998	614	78	67	61	60	59	67	55	41	21	-
1997	650	69	58	65	81	91	85	59	29	18	-
1996	644	64	47	68	68	80	86	53	29	31	-
1995	684	64	62	80	73	92	54	60	54	30	-
1994	706	58	76	59	91	93	82	55	23	28	-
1993	753	64	64	72	99	92	70	67	42	25	1
1992	691	50	61	86	78	111	76	41	37	25	-
1991	770	74	63	73	102	100	88	46	39	21	-
1990	771	60	66	96	95	84	84	66	37	17	-
1989	755	61	59	92	107	97	86	55	26	17	1
1988	838	69	90	83	119	100	85	63	38	16	1
1987	827	60	95	99	101	102	64	46	21	11	1
1986	821	64	108	90	108	89	73	48	20	12	-
1985	859	84	92	103	94	86	78	62	33	14	-
1984	853	83	93	119	99	86	67	45	31	10	-
1983	900	96	111	103	126	86	61	31	24	9	-
1982	883	94	113	115	91	88	62	37	21	13	1
1981	999	110	125	114	110	108	76	42	16	10	-
1980	1,069	117	130	149	110	95	68	36	13	10	-
1979	949	121	120	115	100	95	51	39	21	5	-
Other chronic liver disease and cirrhosis (571.4-571.9)											
Black, both sexes											
1998	1,287	227	156	152	130	136	109	67	48	27	1
1997	1,319	220	161	168	152	152	105	80	34	31	-
1996	1,306	183	152	159	135	146	130	62	33	31	-
1995	1,340	183	163	166	137	139	88	74	54	39	-
1994	1,378	178	162	147	158	146	106	72	41	34	-
1993	1,458	163	162	152	169	154	125	91	42	25	1
1992	1,398	157	152	146	157	195	108	69	58	28	1
1991	1,487	165	164	168	194	176	114	60	48	23	3
1990	1,551	165	146	196	191	146	134	87	53	28	1
1989	1,597	154	168	197	186	187	133	83	36	26	2
1988	1,665	172	189	185	208	173	125	85	43	20	1
1987	1,740	170	213	218	194	164	113	81	31	18	2
1986	1,687	162	194	214	188	169	117	79	25	13	2
1985	1,829	204	191	225	221	182	128	91	40	17	2
1984	1,850	206	220	249	209	186	129	79	41	14	-
1983	1,979	219	256	262	270	186	111	58	32	11	1
1982	1,936	224	269	239	227	164	112	63	28	15	1
1981	2,191	246	304	279	238	214	143	63	30	13	1
1980	2,384	290	336	317	272	190	132	60	20	14	4
1979	2,272	285	348	305	217	188	124	66	25	12	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other chronic liver disease and cirrhosis (571.4-571.9)											
Black, male											
1998	801	163	100	98	86	91	56	30	19	11	1
1997	805	165	111	109	85	86	49	35	14	14	-
1996	804	123	117	104	82	82	66	28	18	6	-
1995	811	126	112	104	87	69	52	29	18	16	-
1994	808	128	96	99	86	75	47	31	24	11	-
1993	861	108	105	100	92	86	71	38	12	6	1
1992	848	113	102	81	98	106	60	35	24	12	1
1991	838	103	107	113	106	86	49	27	17	6	3
1990	900	110	91	120	110	78	64	34	26	13	1
1989	951	101	119	118	94	108	57	41	15	10	1
1988	949	108	114	119	109	84	61	33	13	6	-
1987	1,043	117	136	139	115	81	55	45	12	10	1
1986	992	104	103	141	91	96	57	39	13	9	2
1985	1,092	130	117	138	138	111	58	37	14	8	2
1984	1,093	130	142	146	122	111	68	36	17	6	-
1983	1,176	138	158	170	154	112	60	28	11	3	1
1982	1,152	139	167	139	148	88	58	31	13	5	-
1981	1,292	150	192	170	139	119	79	29	15	4	1
1980	1,419	185	214	184	169	107	69	29	11	7	4
1979	1,413	173	240	201	128	100	76	31	11	9	2
Other chronic liver disease and cirrhosis (571.4-571.9)											
Black, female											
1998	486	64	56	54	44	45	53	37	29	16	-
1997	514	55	50	59	67	66	56	45	20	17	-
1996	502	60	35	55	53	64	64	34	15	25	-
1995	529	57	51	62	50	70	36	45	36	23	-
1994	570	50	66	48	72	71	59	41	17	23	-
1993	597	55	57	52	77	68	54	53	30	19	-
1992	550	44	50	65	59	89	48	34	34	16	-
1991	649	62	57	55	88	90	65	33	31	17	-
1990	651	55	55	76	81	68	70	53	27	15	-
1989	646	53	49	79	92	79	76	42	21	16	1
1988	716	64	75	66	99	89	64	52	30	14	1
1987	697	53	77	79	79	83	58	36	19	8	1
1986	695	58	91	73	97	73	60	40	12	4	-
1985	737	74	74	87	83	71	70	54	26	9	-
1984	757	76	78	103	87	75	61	43	24	8	-
1983	803	81	98	92	116	74	51	30	21	8	-
1982	784	85	102	100	79	76	54	32	15	10	1
1981	899	96	112	109	99	95	64	34	15	9	-
1980	965	105	122	133	103	83	63	31	9	7	-
1979	859	112	108	104	89	88	48	35	14	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
All races, both sexes											
1998	2,822	51	56	71	108	186	310	439	515	985	-
1997	2,757	52	50	56	121	212	293	449	472	964	-
1996	2,816	42	59	82	122	247	315	461	469	933	-
1995	2,751	50	47	60	141	195	298	437	524	917	-
1994	2,855	39	50	64	140	212	341	420	509	977	-
1993	2,752	32	43	73	118	223	311	430	519	930	-
1992	2,915	34	66	84	112	247	324	456	523	988	-
1991	2,881	26	49	74	132	228	320	447	537	985	-
1990	3,022	23	44	70	142	266	364	465	562	1,010	2
1989	3,048	35	50	78	149	242	348	457	575	1,037	-
1988	3,115	36	47	88	138	272	352	482	538	1,081	-
1987	3,032	22	59	79	175	254	385	507	544	954	-
1986	2,956	24	47	85	140	250	365	482	578	920	2
1985	2,975	33	55	88	169	265	358	472	510	944	1
1984	3,023	28	67	95	174	252	414	499	521	894	2
1983	2,989	41	55	113	175	243	354	494	549	884	-
1982	2,941	36	72	95	167	272	393	458	520	858	-
1981	3,084	37	58	116	179	305	393	474	572	879	1
1980	3,285	41	78	134	187	291	430	556	576	898	-
1979	3,139	41	77	136	183	290	457	484	563	830	-
Cholelithiasis and other disorders of gallbladder (574-575)											
All races, male											
1998	1,216	19	31	38	56	97	163	200	230	330	-
1997	1,239	28	33	32	69	110	165	227	243	289	-
1996	1,213	25	39	45	61	129	172	226	219	256	-
1995	1,215	26	24	35	81	118	174	217	222	279	-
1994	1,245	19	30	43	69	116	168	194	214	339	-
1993	1,207	16	30	49	67	122	176	220	209	282	-
1992	1,252	21	37	41	54	145	152	223	228	309	-
1991	1,246	12	21	41	76	133	160	235	226	302	-
1990	1,280	11	20	41	65	150	195	245	209	305	-
1989	1,336	18	17	36	86	133	199	215	261	331	-
1988	1,385	21	25	43	71	154	205	260	235	330	-
1987	1,297	11	31	46	94	132	204	257	235	263	-
1986	1,344	19	28	53	84	137	188	256	264	292	-
1985	1,368	16	32	51	89	158	199	220	241	321	-
1984	1,387	14	34	59	96	130	241	256	222	300	2
1983	1,385	19	26	59	102	130	198	276	232	298	-
1982	1,289	17	34	53	83	173	195	226	224	255	-
1981	1,345	19	30	64	91	161	186	233	243	295	-
1980	1,475	18	39	85	98	156	233	250	247	308	-
1979	1,383	19	42	66	86	162	243	200	244	291	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
All races, female											
1998	1,606	32	25	33	52	89	147	239	285	655	-
1997	1,518	24	17	24	52	102	128	222	229	675	-
1996	1,603	17	20	37	61	118	143	235	250	677	-
1995	1,536	24	23	25	60	77	124	220	302	638	-
1994	1,610	20	20	21	71	96	173	226	295	638	-
1993	1,545	16	13	24	51	101	135	210	310	648	-
1992	1,663	13	29	43	58	102	172	233	295	679	-
1991	1,635	14	28	33	56	95	160	212	311	683	-
1990	1,742	12	24	29	77	116	169	220	353	705	2
1989	1,712	17	33	42	63	109	149	242	314	706	-
1988	1,730	15	22	45	67	118	147	222	303	751	-
1987	1,735	11	28	33	81	122	181	250	309	691	-
1986	1,612	5	19	32	56	113	177	226	314	628	2
1985	1,607	17	23	37	80	107	159	252	269	623	1
1984	1,636	14	33	36	78	122	173	243	299	594	-
1983	1,604	22	29	54	73	113	156	218	317	586	-
1982	1,652	19	38	42	84	99	198	232	296	603	-
1981	1,739	18	28	52	88	144	207	241	329	584	1
1980	1,810	23	39	49	89	135	197	306	329	590	-
1979	1,756	22	35	70	97	128	214	284	319	539	-
Cholelithiasis and other disorders of gallbladder (574-575)											
White, both sexes											
1998	2,474	37	44	56	90	149	259	383	476	910	-
1997	2,413	34	36	42	90	174	255	406	421	888	-
1996	2,440	34	47	63	95	197	270	398	428	849	-
1995	2,416	35	32	43	117	159	258	387	470	860	-
1994	2,525	32	39	48	114	179	301	377	461	904	-
1993	2,417	23	33	67	92	190	255	385	462	850	-
1992	2,589	29	53	67	96	213	280	407	472	915	-
1991	2,562	18	40	59	113	198	275	394	498	906	-
1990	2,722	18	37	59	116	230	326	421	522	937	2
1989	2,738	31	35	64	123	222	304	410	536	962	-
1988	2,789	23	38	66	115	237	320	431	487	1,012	-
1987	2,721	18	45	67	144	225	341	456	502	881	-
1986	2,673	15	39	74	124	215	330	432	524	874	2
1985	2,660	27	43	73	145	233	311	417	462	883	1
1984	2,752	20	56	72	153	225	381	462	475	846	1
1983	2,684	33	43	89	144	216	318	443	502	833	-
1982	2,663	31	58	83	151	243	344	407	487	814	-
1981	2,811	28	49	96	155	274	348	437	530	838	-
1980	2,995	27	62	122	170	255	382	503	547	866	-
1979	2,867	31	61	121	161	252	413	452	528	793	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
White, male											
1998	1,074	15	24	33	44	79	150	183	212	300	-
1997	1,085	17	22	25	53	92	150	206	217	270	-
1996	1,064	21	32	33	49	112	148	203	204	234	-
1995	1,080	21	18	28	68	99	154	199	204	261	-
1994	1,109	16	25	33	58	101	151	178	198	310	-
1993	1,067	11	25	45	54	108	144	198	188	263	-
1992	1,140	19	30	34	48	127	139	208	206	293	-
1991	1,122	11	18	35	69	120	139	206	214	279	-
1990	1,156	10	19	34	56	130	174	228	194	285	-
1989	1,210	17	12	30	73	124	177	195	250	306	-
1988	1,243	12	21	33	63	135	189	233	211	315	-
1987	1,181	11	24	40	78	121	185	237	221	248	-
1986	1,224	12	23	46	75	125	172	234	236	281	-
1985	1,229	13	26	44	76	136	180	197	221	302	-
1984	1,272	9	30	46	87	117	230	240	198	288	1
1983	1,255	15	21	49	88	117	184	250	220	277	-
1982	1,177	16	26	45	79	160	178	201	212	242	-
1981	1,255	15	27	53	80	153	171	220	233	285	-
1980	1,356	12	32	79	89	143	213	228	239	297	-
1979	1,266	17	38	59	76	144	216	189	228	277	-
Cholelithiasis and other disorders of gallbladder (574-575)											
White, female											
1998	1,400	22	20	23	46	70	109	200	264	610	-
1997	1,328	17	14	17	37	82	105	200	204	618	-
1996	1,376	13	15	30	46	85	122	195	224	615	-
1995	1,336	14	14	15	49	60	104	188	266	599	-
1994	1,416	16	14	15	56	78	150	199	263	594	-
1993	1,350	12	8	22	38	82	111	187	274	587	-
1992	1,449	10	23	33	48	86	141	199	266	622	-
1991	1,440	7	22	24	44	78	136	188	284	627	-
1990	1,566	8	18	25	60	100	152	193	328	652	2
1989	1,528	14	23	34	50	98	127	215	286	656	-
1988	1,546	11	17	33	52	102	131	198	276	697	-
1987	1,540	7	21	27	66	104	156	219	281	633	-
1986	1,449	3	16	28	49	90	158	198	288	593	2
1985	1,431	14	17	29	69	97	131	220	241	581	1
1984	1,480	11	26	26	66	108	151	222	277	558	-
1983	1,429	18	22	40	56	99	134	193	282	556	-
1982	1,486	15	32	38	72	83	166	206	275	572	-
1981	1,556	13	22	43	75	121	177	217	297	553	-
1980	1,639	15	30	43	81	112	169	275	308	569	-
1979	1,601	14	23	62	85	108	197	263	300	516	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
All other, both sexes											
1998	348	14	12	15	18	37	51	56	39	75	-
1997	344	18	14	14	31	38	38	43	51	76	-
1996	376	8	12	19	27	50	45	63	41	84	-
1995	335	15	15	17	24	36	40	50	54	57	-
1994	330	7	11	16	26	33	40	43	48	73	-
1993	335	9	10	6	26	33	56	45	57	80	-
1992	326	5	13	17	16	34	44	49	51	73	-
1991	319	8	9	15	19	30	45	53	39	79	-
1990	300	5	7	11	26	36	38	44	40	73	-
1989	310	4	15	14	26	20	44	47	39	75	-
1988	326	13	9	22	23	35	32	51	51	69	-
1987	311	4	14	12	31	29	44	51	42	73	-
1986	283	9	8	11	16	35	35	50	54	46	-
1985	315	6	12	15	24	32	47	55	48	61	-
1984	271	8	11	23	21	27	33	37	46	48	1
1983	305	8	12	24	31	27	36	51	47	51	-
1982	278	5	14	12	16	29	49	51	33	44	-
1981	273	9	9	20	24	31	45	37	42	41	1
1980	290	14	16	12	17	36	48	53	29	32	-
1979	272	10	16	15	22	38	44	32	35	37	-
Cholelithiasis and other disorders of gallbladder (574-575)											
All other, male											
1998	142	4	7	5	12	18	13	17	18	30	-
1997	154	11	11	7	16	18	15	21	26	19	-
1996	149	4	7	12	12	17	24	23	15	22	-
1995	135	5	6	7	13	19	20	18	18	18	-
1994	136	3	5	10	11	15	17	16	16	29	-
1993	140	5	5	4	13	14	32	22	21	19	-
1992	112	2	7	7	6	18	13	15	22	16	-
1991	124	1	3	6	7	13	21	29	12	23	-
1990	124	1	1	7	9	20	21	17	15	20	-
1989	126	1	5	6	13	9	22	20	11	25	-
1988	142	9	4	10	8	19	16	27	24	15	-
1987	116	-	7	6	16	11	19	20	14	15	-
1986	120	7	5	7	9	12	16	22	28	11	-
1985	139	3	6	7	13	22	19	23	20	19	-
1984	115	5	4	13	9	13	11	16	24	12	1
1983	130	4	5	10	14	13	14	26	12	21	-
1982	112	1	8	8	4	13	17	25	12	13	-
1981	90	4	3	11	11	8	15	13	10	10	-
1980	119	6	7	6	9	13	20	22	8	11	-
1979	117	2	4	7	10	18	27	11	16	14	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
All other, female											
1998	206	10	5	10	6	19	38	39	21	45	-
1997	190	7	3	7	15	20	23	22	25	57	-
1996	227	4	5	7	15	33	21	40	26	62	-
1995	200	10	9	10	11	17	20	32	36	39	-
1994	194	4	6	6	15	18	23	27	32	44	-
1993	195	4	5	2	13	19	24	23	36	61	-
1992	214	3	6	10	10	16	31	34	29	57	-
1991	195	7	6	9	12	17	24	24	27	56	-
1990	176	4	6	4	17	16	17	27	25	53	-
1989	184	3	10	8	13	11	22	27	28	50	-
1988	184	4	5	12	15	16	16	24	27	54	-
1987	195	4	7	6	15	18	25	31	28	58	-
1986	163	2	3	4	7	23	19	28	26	35	-
1985	176	3	6	8	11	10	28	32	28	42	-
1984	156	3	7	10	12	14	22	21	22	36	-
1983	175	4	7	14	17	14	22	25	35	30	-
1982	166	4	6	4	12	16	32	26	21	31	-
1981	183	5	6	9	13	23	30	24	32	31	1
1980	171	8	9	6	8	23	28	31	21	21	-
1979	155	8	12	8	12	20	17	21	19	23	-
Cholelithiasis and other disorders of gallbladder (574-575)											
Black, both sexes											
1998	282	12	10	15	14	30	39	40	30	65	-
1997	272	18	10	11	24	30	26	35	37	64	-
1996	304	8	9	18	22	37	36	49	34	68	-
1995	268	14	11	13	22	32	26	36	44	49	-
1994	275	6	9	14	21	29	32	35	41	58	-
1993	272	6	10	6	21	26	47	37	45	62	-
1992	275	5	11	14	14	26	36	41	48	58	-
1991	260	6	9	13	16	24	39	41	33	61	-
1990	244	4	6	10	20	27	32	34	36	57	-
1989	255	3	12	12	24	16	37	41	30	57	-
1988	264	11	8	16	20	26	25	41	41	55	-
1987	263	3	12	9	25	24	38	41	36	64	-
1986	230	8	5	10	12	27	28	39	44	39	-
1985	270	6	11	14	22	27	37	49	41	49	-
1984	228	7	10	20	17	23	28	29	38	41	1
1983	262	8	10	20	26	20	30	45	44	42	-
1982	230	4	12	10	14	25	41	41	24	34	-
1981	215	7	8	16	20	22	36	27	34	34	1
1980	235	13	15	10	13	27	39	40	21	24	-
1979	226	7	14	13	19	35	35	27	27	27	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Cholelithiasis and other disorders of gallbladder (574-575)											
Black, male											
1998	109	3	6	5	10	14	12	9	11	23	-
1997	119	11	9	6	13	16	8	15	19	14	-
1996	122	4	5	11	11	13	18	19	12	18	-
1995	100	5	5	4	12	16	12	12	13	12	-
1994	111	3	5	8	9	14	14	13	11	22	-
1993	103	3	5	4	10	10	26	15	13	12	-
1992	92	2	6	6	6	15	9	15	19	9	-
1991	92	1	3	5	4	11	16	20	9	16	-
1990	96	-	1	6	8	14	17	11	13	14	-
1989	102	-	5	6	12	8	15	17	9	16	-
1988	107	7	4	7	6	13	10	22	20	8	-
1987	90	-	6	4	10	10	16	14	11	11	-
1986	92	6	3	6	7	7	12	15	23	10	-
1985	118	3	6	7	12	20	12	20	17	15	-
1984	89	4	4	11	6	11	9	11	19	7	1
1983	103	4	4	8	11	8	10	22	11	15	-
1982	87	1	8	6	3	12	13	18	7	8	-
1981	65	3	3	9	9	6	8	9	7	8	-
1980	89	5	6	5	6	7	15	14	6	8	-
1979	91	1	3	6	9	17	20	9	10	9	-
Cholelithiasis and other disorders of gallbladder (574-575)											
Black, female											
1998	173	9	4	10	4	16	27	31	19	42	-
1997	153	7	1	5	11	14	18	20	18	50	-
1996	182	4	4	7	11	24	18	30	22	50	-
1995	168	9	6	9	10	16	14	24	31	37	-
1994	164	3	4	6	12	15	18	22	30	36	-
1993	169	3	5	2	11	16	21	22	32	50	-
1992	183	3	5	8	8	11	27	26	29	49	-
1991	168	5	6	8	12	13	23	21	24	45	-
1990	148	4	5	4	12	13	15	23	23	43	-
1989	153	3	7	6	12	8	22	24	21	41	-
1988	157	4	4	9	14	13	15	19	21	47	-
1987	173	3	6	5	15	14	22	27	25	53	-
1986	138	2	2	4	5	20	16	24	21	29	-
1985	152	3	5	7	10	7	25	29	24	34	-
1984	139	3	6	9	11	12	19	18	19	34	-
1983	159	4	6	12	15	12	20	23	33	27	-
1982	143	3	4	4	11	13	28	23	17	26	-
1981	150	4	5	7	11	16	28	18	27	26	1
1980	146	8	9	5	7	20	24	26	15	16	-
1979	135	6	11	7	10	18	15	18	17	18	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All races, both sexes											
1998	26,182	422	510	745	1,067	1,753	3,025	4,113	4,945	8,804	-
1997	25,331	365	502	730	1,102	1,786	2,943	3,929	4,691	8,438	-
1996	24,304	393	486	626	1,053	1,737	2,896	3,805	4,548	7,883	2
1995	23,676	346	462	631	1,108	1,807	2,783	3,747	4,332	7,513	4
1994	22,976	339	407	658	1,041	1,772	2,880	3,556	4,232	7,226	2
1993	23,317	343	417	706	1,148	1,949	2,831	3,591	4,216	7,156	2
1992	22,162	313	412	628	1,096	1,856	2,684	3,488	3,943	6,740	4
1991	21,360	271	408	583	1,112	1,806	2,592	3,358	3,831	6,376	2
1990	20,764	317	407	611	1,125	1,767	2,545	3,256	3,722	6,016	-
1989	21,118	271	442	689	1,177	1,908	2,507	3,374	3,759	5,960	5
1988	22,392	326	462	743	1,315	1,983	2,607	3,560	3,907	6,396	1
1987	22,052	306	455	754	1,278	1,944	2,712	3,537	3,804	6,252	-
1986	21,767	310	422	726	1,389	1,931	2,718	3,496	3,700	6,006	2
1985	21,349	303	414	802	1,356	1,954	2,757	3,250	3,646	5,786	3
1984	20,126	287	413	756	1,308	1,851	2,626	3,223	3,330	5,273	2
1983	18,998	268	452	834	1,212	1,728	2,480	3,042	3,230	4,791	5
1982	18,102	257	432	823	1,234	1,755	2,408	2,758	2,986	4,455	3
1981	17,277	277	462	791	1,144	1,695	2,332	2,676	2,829	4,053	3
1980	16,753	316	508	795	1,162	1,646	2,181	2,591	2,649	3,901	3
1979	15,729	323	524	745	1,088	1,631	2,008	2,450	2,467	3,410	2
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All races, male											
1998	12,561	281	297	409	574	923	1,582	2,099	2,423	3,506	-
1997	12,140	222	285	402	579	949	1,576	2,032	2,234	3,365	-
1996	11,642	230	271	347	591	899	1,552	1,942	2,214	3,066	1
1995	11,389	206	252	353	583	985	1,465	1,959	2,038	2,974	4
1994	10,866	195	232	357	537	898	1,473	1,769	2,041	2,841	2
1993	11,210	213	228	390	586	1,047	1,483	1,837	1,994	2,851	-
1992	10,816	195	234	350	606	975	1,402	1,804	1,903	2,727	2
1991	10,418	164	235	331	629	946	1,375	1,738	1,811	2,529	1
1990	10,005	179	224	312	575	922	1,343	1,700	1,760	2,366	-
1989	10,320	157	251	404	659	1,003	1,348	1,711	1,776	2,349	4
1988	10,880	184	271	404	717	1,075	1,346	1,854	1,824	2,550	1
1987	10,872	186	269	417	715	1,037	1,431	1,839	1,806	2,575	-
1986	10,863	180	249	393	771	1,034	1,504	1,801	1,814	2,494	1
1985	10,551	167	220	448	711	1,054	1,475	1,705	1,675	2,428	1
1984	10,051	152	218	391	739	1,002	1,400	1,667	1,615	2,242	1
1983	9,312	146	246	454	636	890	1,319	1,552	1,569	1,986	5
1982	9,165	152	254	461	653	929	1,300	1,434	1,429	1,974	3
1981	8,720	157	253	446	633	897	1,273	1,387	1,319	1,769	-
1980	8,638	172	257	442	631	952	1,178	1,381	1,298	1,741	2
1979	8,260	170	292	421	610	889	1,097	1,331	1,241	1,582	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All races, female											
1998	13,621	141	213	336	493	830	1,443	2,014	2,522	5,298	-
1997	13,191	143	217	328	523	837	1,367	1,897	2,457	5,073	-
1996	12,662	163	215	279	462	838	1,344	1,863	2,334	4,817	1
1995	12,287	140	210	278	525	822	1,318	1,788	2,294	4,539	-
1994	12,110	144	175	301	504	874	1,407	1,787	2,191	4,385	-
1993	12,107	130	189	316	562	902	1,348	1,754	2,222	4,305	2
1992	11,346	118	178	278	490	881	1,282	1,684	2,040	4,013	2
1991	10,942	107	173	252	483	860	1,217	1,620	2,020	3,847	1
1990	10,759	138	183	299	550	845	1,202	1,556	1,962	3,650	-
1989	10,798	114	191	285	518	905	1,159	1,663	1,983	3,611	1
1988	11,512	142	191	339	598	908	1,261	1,706	2,083	3,846	-
1987	11,180	120	186	337	563	907	1,281	1,698	1,998	3,677	-
1986	10,904	130	173	333	618	897	1,214	1,695	1,886	3,512	1
1985	10,798	136	194	354	645	900	1,282	1,545	1,971	3,358	2
1984	10,075	135	195	365	569	849	1,226	1,556	1,715	3,031	1
1983	9,686	122	206	380	576	838	1,161	1,490	1,661	2,805	-
1982	8,937	105	178	362	581	826	1,108	1,324	1,557	2,481	-
1981	8,557	120	209	345	511	798	1,059	1,289	1,510	2,284	3
1980	8,115	144	251	353	531	694	1,003	1,210	1,351	2,160	1
1979	7,469	153	232	324	478	742	911	1,119	1,226	1,828	1
Nephritis, nephrotic syndrome and nephrosis (580-589)											
White, both sexes											
1998	21,369	259	330	467	735	1,354	2,368	3,396	4,306	7,676	-
1997	20,678	209	323	465	745	1,360	2,345	3,273	4,066	7,356	-
1996	19,696	244	314	417	740	1,281	2,310	3,162	3,885	6,837	-
1995	19,117	211	283	400	763	1,356	2,236	3,117	3,664	6,553	1
1994	18,639	193	259	419	704	1,315	2,326	2,955	3,666	6,342	1
1993	18,795	203	252	458	808	1,505	2,265	2,980	3,578	6,219	1
1992	17,903	183	252	392	757	1,400	2,142	2,935	3,384	5,886	2
1991	17,165	159	258	365	785	1,351	2,041	2,815	3,285	5,529	2
1990	16,571	182	247	365	779	1,334	1,986	2,674	3,163	5,281	-
1989	16,607	161	257	426	798	1,387	1,946	2,718	3,157	5,195	-
1988	17,915	181	284	482	931	1,516	2,060	2,925	3,299	5,590	-
1987	17,764	183	279	495	919	1,471	2,141	2,951	3,251	5,486	-
1986	17,446	183	256	474	994	1,457	2,169	2,855	3,172	5,244	1
1985	17,046	192	252	531	981	1,467	2,157	2,694	3,070	5,045	2
1984	16,198	175	262	505	943	1,423	2,060	2,697	2,796	4,658	-
1983	15,334	172	309	557	872	1,294	1,945	2,527	2,787	4,257	1
1982	14,453	158	260	542	884	1,304	1,895	2,286	2,520	3,954	1
1981	13,828	173	297	526	815	1,249	1,811	2,249	2,416	3,613	1
1980	13,137	191	313	510	795	1,209	1,681	2,103	2,266	3,427	2
1979	12,534	198	339	495	780	1,225	1,583	1,993	2,147	3,051	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
White, male											
1998	10,406	176	193	257	398	743	1,295	1,782	2,168	3,121	-
1997	10,002	123	184	260	405	751	1,301	1,744	1,980	2,958	-
1996	9,549	143	181	231	431	685	1,284	1,642	1,966	2,690	-
1995	9,288	127	156	237	409	766	1,221	1,677	1,768	2,618	1
1994	8,901	102	150	233	378	681	1,221	1,536	1,822	2,526	1
1993	9,110	128	135	261	438	840	1,216	1,579	1,707	2,504	-
1992	8,784	113	150	224	442	759	1,153	1,558	1,642	2,396	2
1991	8,403	93	150	202	461	724	1,111	1,496	1,588	2,206	1
1990	8,021	99	142	199	407	712	1,076	1,440	1,504	2,081	-
1989	8,093	94	142	253	486	735	1,080	1,419	1,487	2,044	-
1988	8,786	99	174	268	533	829	1,122	1,560	1,554	2,257	-
1987	8,800	117	171	280	530	823	1,162	1,562	1,562	2,255	-
1986	8,754	104	155	263	581	802	1,233	1,495	1,572	2,186	-
1985	8,482	108	148	306	538	826	1,191	1,433	1,413	2,117	1
1984	8,174	95	140	271	538	802	1,141	1,428	1,376	1,979	-
1983	7,548	98	172	313	467	694	1,071	1,314	1,355	1,742	1
1982	7,405	93	156	325	489	716	1,055	1,221	1,213	1,748	1
1981	7,018	98	170	299	473	683	1,013	1,191	1,139	1,554	-
1980	6,848	104	163	297	469	742	935	1,131	1,105	1,525	2
1979	6,630	105	195	297	469	698	890	1,081	1,083	1,403	-
Nephritis, nephrotic syndrome and nephrosis (580-589)											
White, female											
1998	10,963	83	137	210	337	611	1,073	1,614	2,138	4,555	-
1997	10,676	86	139	205	340	609	1,044	1,529	2,086	4,398	-
1996	10,147	101	133	186	309	596	1,026	1,520	1,919	4,147	-
1995	9,829	84	127	163	354	590	1,015	1,440	1,896	3,935	-
1994	9,738	91	109	186	326	634	1,105	1,419	1,844	3,816	-
1993	9,685	75	117	197	370	665	1,049	1,401	1,871	3,715	1
1992	9,119	70	102	168	315	641	989	1,377	1,742	3,490	-
1991	8,762	66	108	163	324	627	930	1,319	1,697	3,323	1
1990	8,550	83	105	166	372	622	910	1,234	1,659	3,200	-
1989	8,514	67	115	173	312	652	866	1,299	1,670	3,151	-
1988	9,129	82	110	214	398	687	938	1,365	1,745	3,333	-
1987	8,964	66	108	215	389	648	979	1,389	1,689	3,231	-
1986	8,692	79	101	211	413	655	936	1,360	1,600	3,058	1
1985	8,564	84	104	225	443	641	966	1,261	1,657	2,928	1
1984	8,024	80	122	234	405	621	919	1,269	1,420	2,679	-
1983	7,786	74	137	244	405	600	874	1,213	1,432	2,515	-
1982	7,048	65	104	217	395	588	840	1,065	1,307	2,206	-
1981	6,810	75	127	227	342	566	798	1,058	1,277	2,059	1
1980	6,289	87	150	213	326	467	746	972	1,161	1,902	-
1979	5,904	93	144	198	311	527	693	912	1,064	1,648	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All other, both sexes											
1998	4,813	163	180	278	332	399	657	717	639	1,128	-
1997	4,653	156	179	265	357	426	598	656	625	1,082	-
1996	4,608	149	172	209	313	456	586	643	663	1,046	2
1995	4,559	135	179	231	345	451	547	630	668	960	3
1994	4,337	146	148	239	337	457	554	601	566	884	1
1993	4,522	140	165	248	340	444	566	611	638	937	1
1992	4,259	130	160	236	339	456	542	553	559	854	2
1991	4,195	112	150	218	327	455	551	543	546	847	-
1990	4,193	135	160	246	346	433	559	582	559	735	-
1989	4,511	110	185	263	379	521	561	656	602	765	5
1988	4,477	145	178	261	384	467	547	635	608	806	1
1987	4,288	123	176	259	359	473	571	586	553	766	-
1986	4,321	127	166	252	395	474	549	641	528	762	1
1985	4,303	111	162	271	375	487	600	556	576	741	1
1984	3,928	112	151	251	365	428	566	526	534	615	2
1983	3,664	96	143	277	340	434	535	515	443	534	4
1982	3,649	99	172	281	350	451	513	472	466	501	2
1981	3,449	104	165	265	329	446	521	427	413	440	2
1980	3,616	125	195	285	367	437	500	488	383	474	1
1979	3,195	125	185	250	308	406	425	457	320	359	1
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All other, male											
1998	2,155	105	104	152	176	180	287	317	255	385	-
1997	2,138	99	101	142	174	198	275	288	254	407	-
1996	2,093	87	90	116	160	214	268	300	248	376	1
1995	2,101	79	96	116	174	219	244	282	270	356	3
1994	1,965	93	82	124	159	217	252	233	219	315	1
1993	2,100	85	93	129	148	207	267	258	287	347	-
1992	2,032	82	84	126	164	216	249	246	261	331	-
1991	2,015	71	85	129	168	222	264	242	223	323	-
1990	1,984	80	82	113	168	210	267	260	256	285	-
1989	2,227	63	109	151	173	268	268	292	289	305	4
1988	2,094	85	97	136	184	246	224	294	270	293	1
1987	2,072	69	98	137	185	214	269	277	244	320	-
1986	2,109	76	94	130	190	232	271	306	242	308	1
1985	2,069	59	72	142	173	228	284	272	262	311	-
1984	1,877	57	78	120	201	200	259	239	239	263	1
1983	1,764	48	74	141	169	196	248	238	214	244	4
1982	1,760	59	98	136	164	213	245	213	216	226	2
1981	1,702	59	83	147	160	214	260	196	180	215	-
1980	1,790	68	94	145	162	210	243	250	193	216	-
1979	1,630	65	97	124	141	191	207	250	158	179	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
All other, female											
1998	2,658	58	76	126	156	219	370	400	384	743	-
1997	2,515	57	78	123	183	228	323	368	371	675	-
1996	2,515	62	82	93	153	242	318	343	415	670	1
1995	2,458	56	83	115	171	232	303	348	398	604	-
1994	2,372	53	66	115	178	240	302	368	347	569	-
1993	2,422	55	72	119	192	237	299	353	351	590	1
1992	2,227	48	76	110	175	240	293	307	298	523	2
1991	2,180	41	65	89	159	233	287	301	323	524	-
1990	2,209	55	78	133	178	223	292	322	303	450	-
1989	2,284	47	76	112	206	253	293	364	313	460	1
1988	2,383	60	81	125	200	221	323	341	338	513	-
1987	2,216	54	78	122	174	259	302	309	309	446	-
1986	2,212	51	72	122	205	242	278	335	286	454	-
1985	2,234	52	90	129	202	259	316	284	314	430	1
1984	2,051	55	73	131	164	228	307	287	295	352	1
1983	1,900	48	69	136	171	238	287	277	229	290	-
1982	1,889	40	74	145	186	238	268	259	250	275	-
1981	1,747	45	82	118	169	232	261	231	233	225	2
1980	1,826	57	101	140	205	227	257	238	190	258	1
1979	1,565	60	88	126	167	215	218	207	162	180	-
Nephritis, nephrotic syndrome and nephrosis (580-589)											
Black, both sexes											
1998	4,329	148	155	251	310	363	569	644	563	1,029	-
1997	4,236	141	162	238	332	388	537	586	567	999	-
1996	4,144	140	157	187	289	403	520	580	582	939	2
1995	4,159	127	161	210	317	411	494	579	593	877	3
1994	3,980	138	142	213	314	424	504	539	517	805	1
1993	4,107	132	152	230	311	406	509	548	565	847	1
1992	3,889	121	143	213	313	418	484	504	504	789	2
1991	3,866	108	137	199	302	419	514	487	505	769	-
1990	3,855	126	147	233	315	390	516	528	512	675	-
1989	4,166	104	175	248	349	482	518	602	550	696	5
1988	4,157	140	161	246	358	440	501	591	556	745	1
1987	3,975	113	165	244	340	436	531	539	516	705	-
1986	4,020	116	149	226	371	452	514	596	498	698	-
1985	4,044	97	150	255	361	461	558	508	542	706	1
1984	3,666	98	139	239	340	406	536	480	503	572	2
1983	3,426	91	134	262	319	410	498	487	414	492	4
1982	3,417	91	161	264	333	432	488	438	432	466	2
1981	3,262	98	156	253	314	427	490	405	391	408	2
1980	3,416	118	186	271	348	421	477	463	355	443	1
1979	3,018	116	176	241	298	385	403	430	297	326	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nephritis, nephrotic syndrome and nephrosis (580-589)											
Black, male											
1998	1,929	96	88	139	169	165	244	282	223	340	-
1997	1,941	90	93	125	162	184	252	255	227	363	-
1996	1,873	83	85	101	150	192	233	271	215	321	1
1995	1,916	74	84	108	157	202	221	261	236	318	3
1994	1,794	88	79	115	148	197	232	211	192	274	1
1993	1,879	81	84	118	133	191	240	224	244	300	-
1992	1,850	76	75	117	153	199	220	224	233	294	-
1991	1,849	70	77	119	154	206	249	209	203	285	-
1990	1,806	78	77	110	147	189	247	227	229	254	-
1989	2,047	60	105	143	160	246	244	263	259	271	4
1988	1,908	81	86	124	173	228	200	269	234	259	1
1987	1,905	62	92	130	174	196	248	253	218	291	-
1986	1,963	72	85	116	180	224	252	284	230	275	-
1985	1,935	57	66	134	167	218	261	241	241	294	-
1984	1,731	51	70	116	184	187	241	209	223	239	1
1983	1,648	46	67	135	158	187	230	225	197	221	4
1982	1,650	54	97	130	155	207	229	196	198	211	2
1981	1,609	56	77	142	152	206	240	189	166	201	-
1980	1,694	65	92	143	152	203	230	235	175	202	-
1979	1,525	61	92	116	136	179	192	233	146	159	1
Nephritis, nephrotic syndrome and nephrosis (580-589)											
Black, female											
1998	2,400	52	67	112	141	198	325	362	340	689	-
1997	2,295	51	69	113	170	204	285	331	340	636	-
1996	2,271	57	72	86	139	211	287	309	367	618	1
1995	2,243	53	77	102	160	209	273	318	357	559	-
1994	2,186	50	63	98	166	227	272	328	325	531	-
1993	2,228	51	68	112	178	215	269	324	321	547	1
1992	2,039	45	68	96	160	219	264	280	271	495	2
1991	2,017	38	60	80	148	213	265	278	302	484	-
1990	2,049	48	70	123	168	201	269	301	283	421	-
1989	2,119	44	70	105	189	236	274	339	291	425	1
1988	2,249	59	75	122	185	212	301	322	322	486	-
1987	2,070	51	73	114	166	240	283	286	298	414	-
1986	2,057	44	64	110	191	228	262	312	268	423	-
1985	2,109	40	84	121	194	243	297	267	301	412	1
1984	1,935	47	69	123	156	219	295	271	280	333	1
1983	1,778	45	67	127	161	223	268	262	217	271	-
1982	1,767	37	64	134	178	225	259	242	234	255	-
1981	1,653	42	79	111	162	221	250	216	225	207	2
1980	1,722	53	94	128	196	218	247	228	180	241	1
1979	1,493	55	84	125	162	206	211	197	151	167	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All races, both sexes											
1998	325	8	9	9	14	24	35	51	58	89	-
1997	324	4	12	13	13	16	35	49	60	92	-
1996	321	6	9	11	10	29	26	58	61	78	-
1995	292	2	8	3	8	20	34	48	53	82	-
1994	305	9	7	6	14	20	29	52	57	88	-
1993	292	7	6	10	16	20	30	44	49	78	-
1992	304	3	3	10	27	24	36	50	55	63	-
1991	320	5	8	8	16	27	27	50	49	93	-
1990	296	12	6	10	15	33	26	35	47	73	-
1989	310	4	5	11	21	35	35	51	58	50	-
1988	310	6	8	12	23	26	42	53	44	70	-
1987	277	4	5	14	14	23	41	30	60	51	-
1986	293	1	4	12	24	30	29	40	55	58	-
1985	348	6	9	11	31	30	40	54	54	73	-
1984	316	11	11	12	17	24	37	58	54	60	-
1983	354	4	7	21	21	38	43	56	55	75	-
1982	309	8	11	20	22	21	42	47	57	53	-
1981	302	4	6	13	17	34	47	48	41	55	-
1980	328	8	14	9	28	22	39	47	54	64	-
1979	338	8	10	15	19	38	42	49	58	56	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All races, male											
1998	154	6	5	6	7	10	21	19	25	40	-
1997	154	2	9	7	10	8	17	24	27	35	-
1996	153	2	4	8	5	14	15	36	21	27	-
1995	133	1	5	-	6	9	14	24	23	33	-
1994	142	6	3	5	6	11	16	25	25	29	-
1993	124	5	2	7	6	8	12	19	18	27	-
1992	143	2	3	5	14	9	15	25	23	27	-
1991	154	4	5	6	10	12	15	30	19	34	-
1990	145	8	3	7	9	14	16	19	19	25	-
1989	160	2	4	8	16	21	19	24	26	15	-
1988	158	5	4	11	13	16	22	31	17	25	-
1987	133	3	3	8	6	10	21	16	30	20	-
1986	133	-	2	4	12	16	12	19	23	23	-
1985	159	1	6	8	21	14	15	27	22	21	-
1984	147	2	4	8	12	11	15	37	20	23	-
1983	171	-	4	10	9	22	27	29	25	25	-
1982	151	2	4	12	11	13	19	20	26	27	-
1981	142	1	4	8	8	14	23	26	15	24	-
1980	162	6	10	3	17	16	21	23	24	23	-
1979	182	6	8	8	8	21	23	27	25	23	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All races, female											
1998	171	2	4	3	7	14	14	32	33	49	-
1997	170	2	3	6	3	8	18	25	33	57	-
1996	168	4	5	3	5	15	11	22	40	51	-
1995	159	1	3	3	2	11	20	24	30	49	-
1994	163	3	4	1	8	9	13	27	32	59	-
1993	168	2	4	3	10	12	18	25	31	51	-
1992	161	1	-	5	13	15	21	25	32	36	-
1991	166	1	3	2	6	15	12	20	30	59	-
1990	151	4	3	3	6	19	10	16	28	48	-
1989	150	2	1	3	5	14	16	27	32	35	-
1988	152	1	4	1	10	10	20	22	27	45	-
1987	144	1	2	6	8	13	20	14	30	31	-
1986	160	1	2	8	12	14	17	21	32	35	-
1985	189	5	3	3	10	16	25	27	32	52	-
1984	169	9	7	4	5	13	22	21	34	37	-
1983	183	4	3	11	12	16	16	27	30	50	-
1982	158	6	7	8	11	8	23	27	31	26	-
1981	160	3	2	5	9	20	24	22	26	31	-
1980	166	2	4	6	11	6	18	24	30	41	-
1979	156	2	2	7	11	17	19	22	33	33	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
White, both sexes											
1998	272	7	6	5	11	19	32	41	53	78	-
1997	270	3	11	9	8	11	29	42	50	84	-
1996	257	4	7	7	7	21	25	48	50	65	-
1995	244	1	4	3	6	18	31	43	47	68	-
1994	248	8	4	4	10	14	25	42	50	78	-
1993	232	5	3	5	11	16	23	40	37	72	-
1992	252	2	2	10	23	16	31	46	50	53	-
1991	259	3	7	4	14	18	20	44	43	87	-
1990	249	7	4	7	11	27	23	31	45	68	-
1989	241	2	3	8	17	30	25	40	53	42	-
1988	249	2	4	9	18	22	35	45	37	60	-
1987	221	3	2	9	9	14	35	29	53	47	-
1986	243	-	2	11	23	25	25	35	50	49	-
1985	286	4	5	10	25	26	33	43	50	68	-
1984	262	6	7	8	15	20	33	52	48	54	-
1983	301	4	5	14	17	28	35	54	49	71	-
1982	247	4	6	12	18	19	35	42	46	50	-
1981	251	2	4	9	12	27	39	43	37	52	-
1980	270	5	7	6	20	16	32	43	50	60	-
1979	291	6	9	14	14	33	40	42	52	55	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
White, male											
1998	133	5	4	4	4	8	20	17	22	38	-
1997	125	2	9	4	7	5	12	20	24	32	-
1996	125	1	4	5	3	8	15	30	20	25	-
1995	115	-	2	-	5	7	14	22	22	30	-
1994	108	5	2	3	4	7	12	21	23	23	-
1993	98	4	2	3	4	6	9	18	15	25	-
1992	119	2	2	5	12	5	13	23	21	22	-
1991	129	3	4	3	10	10	11	28	18	33	-
1990	118	4	2	5	6	11	15	17	19	24	-
1989	126	1	2	6	13	18	15	20	26	14	-
1988	131	1	3	9	11	13	19	28	15	23	-
1987	103	2	2	5	4	6	17	15	27	17	-
1986	107	-	1	3	11	12	10	16	20	21	-
1985	130	1	4	8	17	13	13	22	19	20	-
1984	125	2	1	5	10	9	14	33	18	21	-
1983	143	-	3	8	7	16	23	27	21	25	-
1982	120	1	2	8	10	11	17	18	20	24	-
1981	119	1	3	6	7	11	19	24	13	21	-
1980	131	4	6	3	13	12	18	20	21	21	-
1979	157	4	8	8	6	19	23	24	24	23	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
White, female											
1998	139	2	2	1	7	11	12	24	31	40	-
1997	145	1	2	5	1	6	17	22	26	52	-
1996	132	3	3	2	4	13	10	18	30	40	-
1995	129	1	2	3	1	11	17	21	25	38	-
1994	140	3	2	1	6	7	13	21	27	55	-
1993	134	1	1	2	7	10	14	22	22	47	-
1992	133	-	-	5	11	11	18	23	29	31	-
1991	130	-	3	1	4	8	9	16	25	54	-
1990	131	3	2	2	5	16	8	14	26	44	-
1989	115	1	1	2	4	12	10	20	27	28	-
1988	118	1	1	-	7	9	16	17	22	37	-
1987	118	1	-	4	5	8	18	14	26	30	-
1986	136	-	1	8	12	13	15	19	30	28	-
1985	156	3	1	2	8	13	20	21	31	48	-
1984	137	4	6	3	5	11	19	19	30	33	-
1983	158	4	2	6	10	12	12	27	28	46	-
1982	127	3	4	4	8	8	18	24	26	26	-
1981	132	1	1	3	5	16	20	19	24	31	-
1980	139	1	1	3	7	4	14	23	29	39	-
1979	134	2	1	6	8	14	17	18	28	32	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All other, both sexes											
1998	53	1	3	4	3	5	3	10	5	11	-
1997	54	1	1	4	5	5	6	7	10	8	-
1996	64	2	2	4	3	8	1	10	11	13	-
1995	48	1	4	-	2	2	3	5	6	14	-
1994	57	1	3	2	4	6	4	10	7	10	-
1993	60	2	3	5	5	4	7	4	12	6	-
1992	52	1	1	-	4	8	5	4	5	10	-
1991	61	2	1	4	2	9	7	6	6	6	-
1990	47	5	2	3	4	6	3	4	2	5	-
1989	69	2	2	3	4	5	10	11	5	8	-
1988	61	4	4	3	5	4	7	8	7	10	-
1987	56	1	3	5	5	9	6	1	7	4	-
1986	50	1	2	1	1	5	4	5	5	9	-
1985	62	2	4	1	6	4	7	11	4	5	-
1984	54	5	4	4	2	4	4	6	6	6	-
1983	53	-	2	7	4	10	8	2	6	4	-
1982	62	4	5	8	4	2	7	5	11	3	-
1981	51	2	2	4	5	7	8	5	4	3	-
1980	58	3	7	3	8	6	7	4	4	4	-
1979	47	2	1	1	5	5	2	7	6	1	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All other, male											
1998	21	1	1	2	3	2	1	2	3	2	-
1997	29	-	-	3	3	3	5	4	3	3	-
1996	28	1	-	3	2	6	-	6	1	2	-
1995	18	1	3	-	1	2	-	2	1	3	-
1994	34	1	1	2	2	4	4	4	2	6	-
1993	26	1	-	4	2	2	3	1	3	2	-
1992	24	-	1	-	2	4	2	2	2	5	-
1991	25	1	1	3	-	2	4	2	1	1	-
1990	27	4	1	2	3	3	1	2	-	1	-
1989	34	1	2	2	3	3	4	4	-	1	-
1988	27	4	1	2	2	3	3	3	2	2	-
1987	30	1	1	3	2	4	4	1	3	3	-
1986	26	-	1	1	1	4	2	3	3	2	-
1985	29	-	2	-	4	1	2	5	3	1	-
1984	22	-	3	3	2	2	1	4	2	2	-
1983	28	-	1	2	2	6	4	2	4	-	-
1982	31	1	2	4	1	2	2	2	6	3	-
1981	23	-	1	2	1	3	4	2	2	3	-
1980	31	2	4	-	4	4	3	3	3	2	-
1979	25	2	-	-	2	2	-	3	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
All other, female											
1998	32	-	2	2	-	3	2	8	2	9	-
1997	25	1	1	1	2	2	1	3	7	5	-
1996	36	1	2	1	1	2	1	4	10	11	-
1995	30	-	1	-	1	-	3	3	5	11	-
1994	23	-	2	-	2	2	-	6	5	4	-
1993	34	1	3	1	3	2	4	3	9	4	-
1992	28	1	-	-	2	4	3	2	3	5	-
1991	36	1	-	1	2	7	3	4	5	5	-
1990	20	1	1	1	1	3	2	2	2	4	-
1989	35	1	-	1	1	2	6	7	5	7	-
1988	34	-	3	1	3	1	4	5	5	8	-
1987	26	-	2	2	3	5	2	-	4	1	-
1986	24	1	1	-	-	1	2	2	2	7	-
1985	33	2	2	1	2	3	5	6	1	4	-
1984	32	5	1	1	-	2	3	2	4	4	-
1983	25	-	1	5	2	4	4	-	2	4	-
1982	31	3	3	4	3	-	5	3	5	-	-
1981	28	2	1	2	4	4	4	3	2	-	-
1980	27	1	3	3	4	2	4	1	1	2	-
1979	22	-	1	1	3	3	2	4	5	1	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
Black, both sexes											
1998	45	-	2	3	3	4	3	8	5	11	-
1997	49	1	1	2	5	4	4	7	10	8	-
1996	60	2	2	4	3	7	1	10	10	12	-
1995	45	1	4	-	2	2	3	5	5	14	-
1994	50	1	3	1	4	6	4	8	6	9	-
1993	55	2	3	4	4	4	7	3	11	5	-
1992	44	1	1	-	4	7	4	3	4	7	-
1991	59	2	1	4	2	9	7	6	6	6	-
1990	42	5	2	2	4	6	3	3	2	3	-
1989	66	2	2	3	4	5	10	11	5	8	-
1988	59	4	4	3	5	4	7	7	7	9	-
1987	48	1	3	5	5	9	5	1	6	4	-
1986	49	1	2	1	1	5	4	5	5	8	-
1985	58	2	4	1	6	4	7	8	4	5	-
1984	51	4	4	4	2	4	4	6	5	6	-
1983	49	-	2	7	3	10	7	2	6	3	-
1982	59	4	5	8	4	2	7	4	10	3	-
1981	47	1	2	4	5	7	8	5	4	3	-
1980	55	3	7	3	8	5	7	4	3	4	-
1979	44	1	1	1	5	5	2	7	4	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
Black, male											
1998	19	-	-	2	3	2	1	2	3	2	-
1997	24	-	-	1	3	2	3	4	3	3	-
1996	26	1	-	3	2	6	-	6	-	2	-
1995	18	1	3	-	1	2	-	2	1	3	-
1994	30	1	1	1	2	4	4	3	2	5	-
1993	23	1	-	3	1	2	3	1	3	1	-
1992	18	-	1	-	2	3	1	1	1	3	-
1991	25	1	1	3	-	2	4	2	1	1	-
1990	24	4	1	1	3	3	1	1	-	-	-
1989	33	1	2	2	3	3	4	4	-	1	-
1988	27	4	1	2	2	3	3	3	2	2	-
1987	26	1	1	3	2	4	3	1	2	3	-
1986	25	-	1	1	1	4	2	3	3	1	-
1985	27	-	2	-	4	1	2	3	3	1	-
1984	21	-	3	3	2	2	1	4	1	2	-
1983	26	-	1	2	1	6	3	2	4	-	-
1982	30	1	2	4	1	2	2	2	5	3	-
1981	22	-	1	2	1	3	4	2	2	3	-
1980	28	2	4	-	4	3	3	3	2	2	-
1979	24	1	-	-	2	2	-	3	1	-	-
Acute and rapidly progressive nephritic and nephrotic syndrome (580-581)											
Black, female											
1998	26	-	2	1	-	2	2	6	2	9	-
1997	25	1	1	1	2	2	1	3	7	5	-
1996	34	1	2	1	1	1	1	4	10	10	-
1995	27	-	1	-	1	-	3	3	4	11	-
1994	20	-	2	-	2	2	-	5	4	4	-
1993	32	1	3	1	3	2	4	2	8	4	-
1992	26	1	-	-	2	4	3	2	3	4	-
1991	34	1	-	1	2	7	3	4	5	5	-
1990	18	1	1	1	1	3	2	2	2	3	-
1989	33	1	-	1	1	2	6	7	5	7	-
1988	32	-	3	1	3	1	4	4	5	7	-
1987	22	-	2	2	3	5	2	-	4	1	-
1986	24	1	1	-	-	1	2	2	2	7	-
1985	31	2	2	1	2	3	5	5	1	4	-
1984	30	4	1	1	-	2	3	2	4	4	-
1983	23	-	1	5	2	4	4	-	2	3	-
1982	29	3	3	4	3	-	5	2	5	-	-
1981	25	1	1	2	4	4	4	3	2	-	-
1980	27	1	3	3	4	2	4	1	1	2	-
1979	20	-	1	1	3	3	2	4	3	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All races, both sexes											
1998	1,715	34	33	55	66	93	202	260	313	577	-
1997	1,652	34	45	51	64	109	190	250	293	499	-
1996	1,638	50	41	35	54	113	198	245	271	546	-
1995	1,504	18	39	49	76	116	174	231	241	451	-
1994	1,546	33	31	47	71	113	184	248	288	449	-
1993	1,588	30	37	42	77	121	214	235	272	464	-
1992	1,503	34	38	47	59	137	153	254	243	439	1
1991	1,512	30	35	42	78	131	179	241	250	416	-
1990	1,521	30	44	53	81	123	184	238	257	386	-
1989	1,517	30	31	69	82	151	162	221	258	395	1
1988	1,521	30	30	53	84	146	164	233	253	411	-
1987	1,509	28	37	49	93	106	191	259	236	409	-
1986	1,505	24	32	63	101	138	211	243	238	352	-
1985	1,565	22	38	65	79	144	209	244	250	411	-
1984	1,824	33	45	70	110	165	219	273	299	460	-
1983	1,887	32	47	96	101	158	254	328	297	440	2
1982	1,946	39	58	79	139	186	247	323	290	447	-
1981	2,055	47	62	94	128	189	267	353	341	423	1
1980	2,245	56	68	104	153	227	297	341	353	498	-
1979	2,363	50	87	104	153	277	296	351	364	485	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All races, male											
1998	831	23	19	34	29	53	108	134	164	220	-
1997	807	21	27	26	41	43	99	132	144	214	-
1996	800	32	25	16	24	65	105	138	139	211	-
1995	753	10	19	32	41	63	95	134	118	182	-
1994	728	15	16	26	36	63	86	131	136	168	-
1993	806	20	20	24	45	66	118	122	127	201	-
1992	776	20	20	30	41	79	78	135	120	189	1
1991	758	15	26	22	45	61	95	111	136	186	-
1990	763	17	28	24	52	66	98	119	115	162	-
1989	733	18	20	47	42	69	84	108	110	159	-
1988	793	15	19	25	51	78	92	129	129	185	-
1987	741	19	24	25	56	64	99	132	99	160	-
1986	785	14	18	38	60	74	114	129	107	162	-
1985	770	15	24	39	39	82	106	113	108	169	-
1984	911	18	23	34	66	93	129	142	126	188	-
1983	882	14	23	51	51	77	124	173	139	163	2
1982	984	27	30	45	71	94	128	163	123	217	-
1981	963	27	32	44	70	97	127	169	135	178	-
1980	1,138	28	33	61	84	130	159	175	171	212	-
1979	1,244	28	53	58	95	159	163	186	168	216	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All races, female											
1998	884	11	14	21	37	40	94	126	149	357	-
1997	845	13	18	25	23	66	91	118	149	285	-
1996	838	18	16	19	30	48	93	107	132	335	-
1995	751	8	20	17	35	53	79	97	123	269	-
1994	818	18	15	21	35	50	98	117	152	281	-
1993	782	10	17	18	32	55	96	113	145	263	-
1992	727	14	18	17	18	58	75	119	123	250	-
1991	754	15	9	20	33	70	84	130	114	230	-
1990	758	13	16	29	29	57	86	119	142	224	-
1989	784	12	11	22	40	82	78	113	148	236	1
1988	728	15	11	28	33	68	72	104	124	226	-
1987	768	9	13	24	37	42	92	127	137	249	-
1986	720	10	14	25	41	64	97	114	131	190	-
1985	795	7	14	26	40	62	103	131	142	242	-
1984	913	15	22	36	44	72	90	131	173	272	-
1983	1,005	18	24	45	50	81	130	155	158	277	-
1982	962	12	28	34	68	92	119	160	167	230	-
1981	1,092	20	30	50	58	92	140	184	206	245	1
1980	1,107	28	35	43	69	97	138	166	182	286	-
1979	1,119	22	34	46	58	118	133	165	196	269	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
White, both sexes											
1998	1,369	25	21	39	43	72	157	210	268	479	-
1997	1,343	22	34	37	44	91	151	209	245	435	-
1996	1,308	37	30	22	38	78	147	209	236	459	-
1995	1,206	12	25	36	56	90	134	192	208	390	-
1994	1,211	15	17	27	47	89	147	197	247	381	-
1993	1,251	19	30	30	50	87	168	195	227	393	-
1992	1,200	21	27	26	46	107	117	218	212	366	1
1991	1,199	21	23	29	62	91	142	196	209	357	-
1990	1,191	22	30	32	61	89	148	183	218	329	-
1989	1,201	18	18	54	61	111	133	180	216	334	-
1988	1,250	22	24	38	65	120	135	195	216	359	-
1987	1,204	21	25	34	71	76	148	217	203	355	-
1986	1,180	20	23	39	75	107	157	200	200	300	-
1985	1,256	14	25	45	65	118	169	197	205	357	-
1984	1,472	22	26	48	83	141	176	239	252	389	-
1983	1,497	26	33	58	68	130	194	267	249	391	1
1982	1,558	20	37	53	101	154	197	270	243	399	-
1981	1,670	32	48	73	95	148	206	301	297	374	-
1980	1,717	38	40	64	106	165	226	273	286	427	-
1979	1,847	31	56	75	110	199	232	284	312	423	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
White, male											
1998	664	18	13	25	21	41	81	104	144	186	-
1997	657	13	17	19	32	38	79	108	126	188	-
1996	642	24	16	9	18	48	85	115	119	182	-
1995	599	7	13	25	30	51	75	108	103	154	-
1994	578	7	9	15	23	50	73	108	123	144	-
1993	639	13	17	17	32	51	94	105	99	176	-
1992	611	11	15	15	30	64	61	114	105	159	1
1991	605	12	19	15	35	41	76	91	116	162	-
1990	590	14	17	16	41	45	80	97	95	135	-
1989	590	11	9	37	32	58	73	95	90	135	-
1988	644	12	14	15	38	64	76	112	109	162	-
1987	587	15	17	19	43	51	78	110	85	141	-
1986	610	11	13	23	47	60	87	109	91	136	-
1985	607	11	15	29	35	66	84	91	88	146	-
1984	741	13	14	27	51	83	107	125	106	155	-
1983	675	11	15	33	34	59	95	139	113	142	1
1982	798	11	20	36	50	79	107	142	106	195	-
1981	785	20	26	34	58	75	95	151	114	155	-
1980	878	19	19	43	64	96	122	140	139	183	-
1979	972	14	32	45	74	116	128	152	149	186	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
White, female											
1998	705	7	8	14	22	31	76	106	124	293	-
1997	686	9	17	18	12	53	72	101	119	247	-
1996	666	13	14	13	20	30	62	94	117	277	-
1995	607	5	12	11	26	39	59	84	105	236	-
1994	633	8	8	12	24	39	74	89	124	237	-
1993	612	6	13	13	18	36	74	90	128	217	-
1992	589	10	12	11	16	43	56	104	107	207	-
1991	594	9	4	14	27	50	66	105	93	195	-
1990	601	8	13	16	20	44	68	86	123	194	-
1989	611	7	9	17	29	53	60	85	126	199	-
1988	606	10	10	23	27	56	59	83	107	197	-
1987	617	6	8	15	28	25	70	107	118	214	-
1986	570	9	10	16	28	47	70	91	109	164	-
1985	649	3	10	16	30	52	85	106	117	211	-
1984	731	9	12	21	32	58	69	114	146	234	-
1983	822	15	18	25	34	71	99	128	136	249	-
1982	760	9	17	17	51	75	90	128	137	204	-
1981	885	12	22	39	37	73	111	150	183	219	-
1980	839	19	21	21	42	69	104	133	147	244	-
1979	875	17	24	30	36	83	104	132	163	237	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All other, both sexes											
1998	346	9	12	16	23	21	45	50	45	98	-
1997	309	12	11	14	20	18	39	41	48	64	-
1996	330	13	11	13	16	35	51	36	35	87	-
1995	298	6	14	13	20	26	40	39	33	61	-
1994	335	18	14	20	24	24	37	51	41	68	-
1993	337	11	7	12	27	34	46	40	45	71	-
1992	303	13	11	21	13	30	36	36	31	73	-
1991	313	9	12	13	16	40	37	45	41	59	-
1990	330	8	14	21	20	34	36	55	39	57	-
1989	316	12	13	15	21	40	29	41	42	61	1
1988	271	8	6	15	19	26	29	38	37	52	-
1987	305	7	12	15	22	30	43	42	33	54	-
1986	325	4	9	24	26	31	54	43	38	52	-
1985	309	8	13	20	14	26	40	47	45	54	-
1984	352	11	19	22	27	24	43	34	47	71	-
1983	390	6	14	38	33	28	60	61	48	49	1
1982	388	19	21	26	38	32	50	53	47	48	-
1981	385	15	14	21	33	41	61	52	44	49	1
1980	528	18	28	40	47	62	71	68	67	71	-
1979	516	19	31	29	43	78	64	67	52	62	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All other, male											
1998	167	5	6	9	8	12	27	30	20	34	-
1997	150	8	10	7	9	5	20	24	18	26	-
1996	158	8	9	7	6	17	20	23	20	29	-
1995	154	3	6	7	11	12	20	26	15	28	-
1994	150	8	7	11	13	13	13	23	13	24	-
1993	167	7	3	7	13	15	24	17	28	25	-
1992	165	9	5	15	11	15	17	21	15	30	-
1991	153	3	7	7	10	20	19	20	20	24	-
1990	173	3	11	8	11	21	18	22	20	27	-
1989	143	7	11	10	10	11	11	13	20	24	-
1988	149	3	5	10	13	14	16	17	20	23	-
1987	154	4	7	6	13	13	21	22	14	19	-
1986	175	3	5	15	13	14	27	20	16	26	-
1985	163	4	9	10	4	16	22	22	20	23	-
1984	170	5	9	7	15	10	22	17	20	33	-
1983	207	3	8	18	17	18	29	34	26	21	1
1982	186	16	10	9	21	15	21	21	17	22	-
1981	178	7	6	10	12	22	32	18	21	23	-
1980	260	9	14	18	20	34	37	35	32	29	-
1979	272	14	21	13	21	43	35	34	19	30	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
All other, female											
1998	179	4	6	7	15	9	18	20	25	64	-
1997	159	4	1	7	11	13	19	17	30	38	-
1996	172	5	2	6	10	18	31	13	15	58	-
1995	144	3	8	6	9	14	20	13	18	33	-
1994	185	10	7	9	11	11	24	28	28	44	-
1993	170	4	4	5	14	19	22	23	17	46	-
1992	138	4	6	6	2	15	19	15	16	43	-
1991	160	6	5	6	6	20	18	25	21	35	-
1990	157	5	3	13	9	13	18	33	19	30	-
1989	173	5	2	5	11	29	18	28	22	37	1
1988	122	5	1	5	6	12	13	21	17	29	-
1987	151	3	5	9	9	17	22	20	19	35	-
1986	150	1	4	9	13	17	27	23	22	26	-
1985	146	4	4	10	10	10	18	25	25	31	-
1984	182	6	10	15	12	14	21	17	27	38	-
1983	183	3	6	20	16	10	31	27	22	28	-
1982	202	3	11	17	17	17	29	32	30	26	-
1981	207	8	8	11	21	19	29	34	23	26	1
1980	268	9	14	22	27	28	34	33	35	42	-
1979	244	5	10	16	22	35	29	33	33	32	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
Black, both sexes											
1998	310	8	9	14	21	20	38	48	43	86	-
1997	276	9	8	13	18	17	36	32	42	60	-
1996	292	10	10	11	12	30	46	34	30	82	-
1995	266	4	12	11	16	24	38	35	32	55	-
1994	306	16	14	18	21	22	32	46	36	65	-
1993	309	9	7	11	25	33	42	36	41	65	-
1992	279	11	10	19	10	28	33	33	27	71	-
1991	285	9	8	12	14	32	36	43	37	57	-
1990	300	7	10	21	17	30	34	49	35	53	-
1989	295	12	13	15	19	37	27	37	40	57	1
1988	241	7	6	13	17	25	22	33	35	46	-
1987	277	6	12	13	22	26	38	38	32	47	-
1986	299	2	8	21	23	30	53	40	35	46	-
1985	288	7	12	19	13	24	36	43	42	53	-
1984	324	10	18	20	23	23	42	33	46	61	-
1983	366	6	13	35	32	27	57	58	43	47	1
1982	353	16	17	26	37	30	48	48	43	43	-
1981	359	14	13	20	30	38	58	46	43	46	1
1980	498	16	25	38	43	60	68	67	63	68	-
1979	484	19	29	28	42	73	61	64	50	54	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
Black, male											
1998	153	4	5	9	7	11	23	29	20	31	-
1997	135	6	7	6	8	5	20	20	17	23	-
1996	144	7	9	5	5	15	18	22	18	29	-
1995	139	3	5	6	8	12	20	23	15	25	-
1994	138	7	7	11	11	12	13	22	10	22	-
1993	153	7	3	6	11	15	22	16	25	21	-
1992	152	7	4	13	9	14	16	20	13	29	-
1991	136	3	5	6	9	17	18	19	17	23	-
1990	156	3	9	8	8	18	18	17	19	25	-
1989	137	7	11	10	10	10	11	13	18	23	-
1988	133	2	5	9	12	13	11	14	19	20	-
1987	139	3	7	5	13	10	20	21	13	16	-
1986	166	2	5	15	12	13	27	19	15	24	-
1985	155	4	9	10	4	16	22	19	17	23	-
1984	155	5	8	6	12	9	21	16	19	29	-
1983	194	3	7	16	17	18	28	32	24	19	1
1982	172	13	9	9	21	14	19	20	16	21	-
1981	166	6	6	10	12	20	30	16	20	21	-
1980	243	7	13	18	17	33	35	34	29	28	-
1979	252	14	20	12	20	39	33	31	19	27	-
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (582-583,587)											
Black, female											
1998	157	4	4	5	14	9	15	19	23	55	-
1997	141	3	1	7	10	12	16	12	25	37	-
1996	148	3	1	6	7	15	28	12	12	53	-
1995	127	1	7	5	8	12	18	12	17	30	-
1994	168	9	7	7	10	10	19	24	26	43	-
1993	156	2	4	5	14	18	20	20	16	44	-
1992	127	4	6	6	1	14	17	13	14	42	-
1991	149	6	3	6	5	15	18	24	20	34	-
1990	144	4	1	13	9	12	16	32	16	28	-
1989	158	5	2	5	9	27	16	24	22	34	1
1988	108	5	1	4	5	12	11	19	16	26	-
1987	138	3	5	8	9	16	18	17	19	31	-
1986	133	-	3	6	11	17	26	21	20	22	-
1985	133	3	3	9	9	8	14	24	25	30	-
1984	169	5	10	14	11	14	21	17	27	32	-
1983	172	3	6	19	15	9	29	26	19	28	-
1982	181	3	8	17	16	16	29	28	27	22	-
1981	193	8	7	10	18	18	28	30	23	25	1
1980	255	9	12	20	26	27	33	33	34	40	-
1979	232	5	9	16	22	34	28	33	31	27	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
All races, both sexes											
1998	24,111	378	467	678	986	1,634	2,786	3,800	4,566	8,133	-
1997	23,309	324	444	664	1,023	1,656	2,713	3,626	4,332	7,834	-
1996	22,303	333	434	578	988	1,595	2,666	3,496	4,209	7,252	2
1995	21,833	326	411	578	1,022	1,663	2,570	3,464	4,031	6,970	4
1994	21,074	295	368	601	952	1,635	2,655	3,249	3,881	6,681	2
1993	21,389	304	371	653	1,053	1,804	2,581	3,305	3,887	6,605	2
1992	20,312	275	371	567	1,009	1,690	2,491	3,175	3,643	6,227	3
1991	19,481	233	362	530	1,015	1,646	2,382	3,062	3,523	5,859	2
1990	18,896	274	354	546	1,024	1,608	2,324	2,976	3,410	5,552	-
1989	19,248	236	401	608	1,072	1,718	2,305	3,096	3,437	5,509	4
1988	20,514	290	422	672	1,206	1,807	2,397	3,265	3,607	5,906	1
1987	20,213	269	411	689	1,165	1,813	2,475	3,241	3,501	5,779	-
1986	19,923	285	384	648	1,261	1,757	2,475	3,210	3,402	5,585	2
1985	19,387	273	366	723	1,241	1,775	2,499	2,947	3,339	5,294	3
1984	17,947	241	354	673	1,179	1,657	2,366	2,889	2,969	4,745	2
1983	16,715	230	398	714	1,085	1,529	2,178	2,649	2,871	4,274	3
1982	15,796	209	361	717	1,069	1,545	2,114	2,383	2,636	3,950	3
1981	14,887	225	392	682	996	1,465	2,015	2,272	2,446	3,568	2
1980	14,136	249	425	680	980	1,393	1,840	2,197	2,233	3,334	3
1979	12,986	264	425	623	912	1,312	1,665	2,045	2,041	2,867	2
Renal failure (584-586)											
All races, male											
1998	11,569	252	273	368	537	860	1,453	1,945	2,231	3,246	-
1997	11,159	198	248	367	527	897	1,457	1,874	2,061	3,112	-
1996	10,671	196	241	322	561	820	1,429	1,765	2,051	2,825	1
1995	10,481	195	227	321	535	909	1,352	1,799	1,894	2,754	4
1994	9,975	174	213	325	492	823	1,366	1,611	1,877	2,640	2
1993	10,265	187	206	359	535	972	1,351	1,693	1,846	2,620	-
1992	9,882	172	211	313	550	885	1,307	1,643	1,758	2,509	1
1991	9,486	143	203	302	572	871	1,264	1,595	1,652	2,307	1
1990	9,076	153	192	279	512	841	1,223	1,558	1,625	2,179	-
1989	9,409	136	225	348	601	912	1,245	1,575	1,637	2,173	4
1988	9,908	164	247	366	653	977	1,231	1,691	1,676	2,337	1
1987	9,975	161	241	383	651	962	1,308	1,686	1,675	2,392	-
1986	9,927	166	227	350	697	943	1,377	1,653	1,682	2,305	1
1985	9,607	151	190	401	651	956	1,351	1,564	1,544	2,236	1
1984	8,979	131	189	349	660	896	1,254	1,488	1,466	2,029	1
1983	8,240	132	219	392	574	789	1,167	1,345	1,403	1,796	3
1982	8,004	123	218	400	570	820	1,150	1,246	1,280	1,728	3
1981	7,606	128	217	393	554	784	1,122	1,192	1,169	1,564	-
1980	7,317	135	214	377	530	806	997	1,182	1,098	1,503	2
1979	6,811	136	231	353	505	707	908	1,113	1,046	1,342	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
All races, female											
1998	12,542	126	194	310	449	774	1,333	1,855	2,335	4,887	-
1997	12,150	126	196	297	496	759	1,256	1,752	2,271	4,722	-
1996	11,632	137	193	256	427	775	1,237	1,731	2,158	4,427	1
1995	11,352	131	184	257	487	754	1,218	1,665	2,137	4,216	-
1994	11,099	121	155	276	460	812	1,289	1,638	2,004	4,041	-
1993	11,124	117	165	294	518	832	1,230	1,612	2,041	3,985	2
1992	10,430	103	160	254	459	805	1,184	1,532	1,885	3,718	2
1991	9,995	90	159	228	443	775	1,118	1,467	1,871	3,552	1
1990	9,820	121	162	267	512	767	1,101	1,418	1,785	3,373	-
1989	9,839	100	176	260	471	806	1,060	1,521	1,800	3,336	-
1988	10,606	126	175	306	553	830	1,166	1,574	1,931	3,569	-
1987	10,238	108	170	306	514	851	1,167	1,555	1,826	3,387	-
1986	9,996	119	157	298	564	814	1,098	1,557	1,720	3,280	1
1985	9,780	122	176	322	590	819	1,148	1,383	1,795	3,058	2
1984	8,968	110	165	324	519	761	1,112	1,401	1,503	2,716	1
1983	8,475	98	179	322	511	740	1,011	1,304	1,468	2,478	-
1982	7,792	86	143	317	499	725	964	1,137	1,356	2,222	-
1981	7,281	97	175	289	442	681	893	1,080	1,277	2,004	2
1980	6,819	114	211	303	450	587	843	1,015	1,135	1,831	1
1979	6,175	128	194	270	407	605	757	932	995	1,525	1
Renal failure (584-586)											
White, both sexes											
1998	19,702	226	302	421	681	1,261	2,177	3,143	3,977	7,114	-
1997	19,029	182	278	418	693	1,253	2,160	3,018	3,766	6,826	-
1996	18,094	199	275	386	695	1,182	2,133	2,901	3,592	6,307	-
1995	17,634	198	251	361	699	1,242	2,068	2,878	3,404	6,088	1
1994	17,139	168	237	384	645	1,208	2,145	2,710	3,364	5,877	1
1993	17,280	177	216	423	746	1,399	2,069	2,741	3,307	5,749	1
1992	16,425	159	223	354	687	1,273	1,993	2,666	3,120	5,460	1
1991	15,673	133	226	330	707	1,241	1,875	2,571	3,025	5,080	2
1990	15,099	152	211	325	705	1,215	1,809	2,454	2,893	4,880	-
1989	15,130	140	232	363	718	1,242	1,784	2,494	2,883	4,814	-
1988	16,382	157	255	431	846	1,372	1,888	2,678	3,043	5,165	-
1987	16,299	155	250	451	834	1,379	1,955	2,698	2,991	5,073	-
1986	15,991	163	230	423	893	1,319	1,985	2,618	2,917	4,890	1
1985	15,469	172	222	475	886	1,320	1,947	2,449	2,814	4,615	2
1984	14,438	145	227	448	843	1,259	1,848	2,404	2,491	4,211	-
1983	13,503	141	271	483	783	1,134	1,713	2,199	2,482	3,793	-
1982	12,604	133	216	472	761	1,128	1,659	1,971	2,228	3,500	1
1981	11,882	138	244	442	706	1,068	1,563	1,903	2,082	3,182	1
1980	11,117	147	265	439	668	1,025	1,419	1,783	1,921	2,937	2
1979	10,365	160	272	404	654	990	1,308	1,663	1,780	2,571	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
White, male											
1998	9,604	153	176	227	373	694	1,194	1,660	1,999	2,897	-
1997	9,204	107	158	236	366	707	1,207	1,614	1,828	2,734	-
1996	8,768	118	160	216	410	629	1,182	1,496	1,824	2,480	-
1995	8,557	120	140	212	373	705	1,129	1,545	1,640	2,430	1
1994	8,199	90	139	214	350	623	1,132	1,405	1,673	2,356	1
1993	8,362	110	116	241	402	783	1,111	1,454	1,590	2,301	-
1992	8,045	99	133	203	399	689	1,079	1,420	1,514	2,214	1
1991	7,652	77	126	183	414	672	1,023	1,375	1,450	2,009	1
1990	7,301	80	122	177	359	655	977	1,323	1,390	1,922	-
1989	7,362	81	130	209	441	658	992	1,300	1,368	1,893	-
1988	7,999	86	157	243	484	750	1,027	1,418	1,428	2,071	-
1987	8,093	98	151	255	482	765	1,065	1,432	1,449	2,095	-
1986	8,027	93	140	237	521	729	1,136	1,370	1,459	2,028	-
1985	7,736	96	129	269	486	746	1,091	1,319	1,306	1,951	1
1984	7,296	79	123	239	476	709	1,018	1,270	1,249	1,802	-
1983	6,716	87	154	271	425	618	953	1,144	1,219	1,573	-
1982	6,466	81	133	278	428	624	929	1,058	1,087	1,527	1
1981	6,107	76	141	258	407	596	898	1,016	1,012	1,376	-
1980	5,824	80	138	251	392	634	794	971	940	1,319	2
1979	5,483	87	155	242	387	562	737	901	908	1,193	-
Renal failure (584-586)											
White, female											
1998	10,098	73	126	194	308	567	983	1,483	1,978	4,217	-
1997	9,825	75	120	182	327	546	953	1,404	1,938	4,092	-
1996	9,326	81	115	170	285	553	951	1,405	1,768	3,827	-
1995	9,077	78	111	149	326	537	939	1,333	1,764	3,658	-
1994	8,940	78	98	170	295	585	1,013	1,305	1,691	3,521	-
1993	8,918	67	100	182	344	616	958	1,287	1,717	3,448	1
1992	8,380	60	90	151	288	584	914	1,246	1,606	3,246	-
1991	8,021	56	100	147	293	569	852	1,196	1,575	3,071	1
1990	7,798	72	89	148	346	560	832	1,131	1,503	2,958	-
1989	7,768	59	102	154	277	584	792	1,194	1,515	2,921	-
1988	8,383	71	98	188	362	622	861	1,260	1,615	3,094	-
1987	8,206	57	99	196	352	614	890	1,266	1,542	2,978	-
1986	7,964	70	90	186	372	590	849	1,248	1,458	2,862	1
1985	7,733	76	93	206	400	574	856	1,130	1,508	2,664	1
1984	7,142	66	104	209	367	550	830	1,134	1,242	2,409	-
1983	6,787	54	117	212	358	516	760	1,055	1,263	2,220	-
1982	6,138	52	83	194	333	504	730	913	1,141	1,973	-
1981	5,775	62	103	184	299	472	665	887	1,070	1,806	1
1980	5,293	67	127	188	276	391	625	812	981	1,618	-
1979	4,882	73	117	162	267	428	571	762	872	1,378	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
All other, both sexes											
1998	4,409	152	165	257	305	373	609	657	589	1,019	-
1997	4,280	142	166	246	330	403	553	608	566	1,008	-
1996	4,209	134	159	192	293	413	533	595	617	945	2
1995	4,199	128	160	217	323	421	502	586	627	882	3
1994	3,935	127	131	217	307	427	510	539	517	804	1
1993	4,109	127	155	230	307	405	512	564	580	856	1
1992	3,887	116	148	213	322	417	498	509	523	767	2
1991	3,808	100	136	200	308	405	507	491	498	779	-
1990	3,797	122	143	221	319	393	515	522	517	672	-
1989	4,118	96	169	245	354	476	521	602	554	695	4
1988	4,132	133	167	241	360	435	509	587	564	741	1
1987	3,914	114	161	238	331	434	520	543	510	706	-
1986	3,932	122	154	225	368	438	490	592	485	695	1
1985	3,918	101	144	248	355	455	552	498	525	679	1
1984	3,509	96	127	225	336	398	518	485	478	534	2
1983	3,212	89	127	231	302	395	465	450	389	481	3
1982	3,192	76	145	245	308	417	455	412	408	450	2
1981	3,005	87	148	240	290	397	452	369	364	386	1
1980	3,019	102	160	241	312	368	421	414	312	397	1
1979	2,621	104	153	219	258	322	357	382	261	296	1
Renal failure (584-586)											
All other, male											
1998	1,965	99	97	141	164	166	259	285	232	349	-
1997	1,955	91	90	131	161	190	250	260	233	378	-
1996	1,903	78	81	106	151	191	247	269	227	345	1
1995	1,924	75	87	109	162	204	223	254	254	324	3
1994	1,776	84	74	111	142	200	234	206	204	284	1
1993	1,903	77	90	118	133	189	240	239	256	319	-
1992	1,837	73	78	110	151	196	228	223	244	295	-
1991	1,834	66	77	119	158	199	241	220	202	298	-
1990	1,775	73	70	102	153	186	246	235	235	257	-
1989	2,047	55	95	139	160	254	253	275	269	280	4
1988	1,909	78	90	123	169	227	204	273	248	266	1
1987	1,882	63	90	128	169	197	243	254	226	297	-
1986	1,900	73	87	113	176	214	241	283	223	277	1
1985	1,871	55	61	132	165	210	260	245	238	285	-
1984	1,683	52	66	110	184	187	236	218	217	227	1
1983	1,524	45	65	121	149	171	214	201	184	223	3
1982	1,538	42	85	122	142	196	221	188	193	201	2
1981	1,499	52	76	135	147	188	224	176	157	188	-
1980	1,493	55	76	126	138	172	203	211	158	184	-
1979	1,328	49	76	111	118	145	171	212	138	149	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
All other, female											
1998	2,444	53	68	116	141	207	350	372	357	670	-
1997	2,325	51	76	115	169	213	303	348	333	630	-
1996	2,306	56	78	86	142	222	286	326	390	600	1
1995	2,275	53	73	108	161	217	279	332	373	558	-
1994	2,159	43	57	106	165	227	276	333	313	520	-
1993	2,206	50	65	112	174	216	272	325	324	537	1
1992	2,050	43	70	103	171	221	270	286	279	472	2
1991	1,974	34	59	81	150	206	266	271	296	481	-
1990	2,022	49	73	119	166	207	269	287	282	415	-
1989	2,071	41	74	106	194	222	268	327	285	415	-
1988	2,223	55	77	118	191	208	305	314	316	475	-
1987	2,032	51	71	110	162	237	277	289	284	409	-
1986	2,032	49	67	112	192	224	249	309	262	418	-
1985	2,047	46	83	116	190	245	292	253	287	394	1
1984	1,826	44	61	115	152	211	282	267	261	307	1
1983	1,688	44	62	110	153	224	251	249	205	258	-
1982	1,654	34	60	123	166	221	234	224	215	249	-
1981	1,506	35	72	105	143	209	228	193	207	198	1
1980	1,526	47	84	115	174	196	218	203	154	213	1
1979	1,293	55	77	108	140	177	186	170	123	147	-
Renal failure (584-586)											
Black, both sexes											
1998	3,970	140	144	233	285	339	528	588	515	932	-
1997	3,901	130	152	222	307	367	497	547	514	929	-
1996	3,787	128	145	172	273	366	472	534	542	844	2
1995	3,834	122	144	198	299	383	451	539	554	805	3
1994	3,615	121	125	194	287	396	465	484	474	730	1
1993	3,730	121	142	214	281	368	460	506	513	773	1
1992	3,549	109	132	192	299	382	444	464	473	707	2
1991	3,509	96	127	182	285	377	471	437	461	703	-
1990	3,494	114	134	209	291	354	474	475	474	618	-
1989	3,798	90	159	230	326	440	480	552	504	631	4
1988	3,845	129	150	228	336	409	471	549	514	687	1
1987	3,639	105	150	225	312	401	487	500	475	652	-
1986	3,658	113	138	202	347	417	456	550	458	638	-
1985	3,684	88	133	233	342	431	514	457	494	645	1
1984	3,278	84	116	215	315	377	489	440	449	501	2
1983	3,003	84	119	219	283	372	432	426	365	442	3
1982	2,998	71	138	228	292	400	432	384	379	420	2
1981	2,849	83	140	229	278	381	424	354	343	357	1
1980	2,853	97	154	229	297	356	401	390	289	369	1
1979	2,480	96	146	211	250	306	338	358	242	271	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Renal failure (584-586)											
Black, male											
1998	1,755	92	83	128	158	152	220	251	200	307	-
1997	1,778	84	85	117	150	177	229	231	207	337	-
1996	1,699	75	76	93	142	171	214	241	197	290	1
1995	1,754	70	76	102	148	187	200	236	220	289	3
1994	1,621	80	71	103	133	181	214	186	180	246	1
1993	1,699	73	81	109	121	173	215	206	216	277	-
1992	1,674	69	70	103	142	181	201	203	219	261	-
1991	1,685	65	71	110	145	186	227	188	185	261	-
1990	1,617	71	67	100	135	168	226	208	209	229	-
1989	1,874	52	91	131	147	233	229	246	241	247	4
1988	1,739	75	79	112	159	210	185	251	213	235	1
1987	1,735	57	84	122	158	182	225	231	202	271	-
1986	1,764	70	78	99	167	207	222	262	212	247	-
1985	1,747	53	55	124	159	200	237	219	220	268	-
1984	1,553	46	59	107	170	175	219	189	203	207	1
1983	1,423	43	59	117	139	162	198	190	169	202	3
1982	1,443	40	85	116	133	191	207	172	177	187	2
1981	1,419	50	70	130	139	182	206	171	144	176	-
1980	1,417	54	75	124	131	167	192	197	144	171	-
1979	1,244	46	72	104	114	137	158	198	126	132	1
Renal failure (584-586)											
Black, female											
1998	2,215	48	61	105	127	187	308	337	315	625	-
1997	2,123	46	67	105	157	190	268	316	307	592	-
1996	2,088	53	69	79	131	195	258	293	345	554	1
1995	2,080	52	68	96	151	196	251	303	334	516	-
1994	1,994	41	54	91	154	215	251	298	294	484	-
1993	2,031	48	61	105	160	195	245	300	297	496	1
1992	1,875	40	62	89	157	201	243	261	254	446	2
1991	1,824	31	56	72	140	191	244	249	276	442	-
1990	1,877	43	67	109	156	186	248	267	265	389	-
1989	1,924	38	68	99	179	207	251	306	263	384	-
1988	2,106	54	71	116	177	199	286	298	301	452	-
1987	1,904	48	66	103	154	219	262	269	273	381	-
1986	1,894	43	60	103	180	210	234	288	246	391	-
1985	1,937	35	78	109	183	231	277	238	274	377	1
1984	1,725	38	57	108	145	202	270	251	246	294	1
1983	1,580	41	60	102	144	210	234	236	196	240	-
1982	1,555	31	53	112	159	209	225	212	202	233	-
1981	1,430	33	70	99	139	199	218	183	199	181	1
1980	1,436	43	79	105	166	189	209	193	145	198	1
1979	1,236	50	74	107	136	169	180	160	116	139	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
All races, both sexes											
1998	31	2	1	3	1	2	2	2	8	5	-
1997	46	3	1	2	2	5	5	4	6	13	-
1996	42	4	2	2	1	-	6	6	7	7	-
1995	47	-	4	1	2	8	5	4	7	10	-
1994	51	2	1	4	4	4	12	7	6	8	-
1993	48	2	3	1	2	4	6	7	8	9	-
1992	43	1	-	4	1	5	4	9	2	11	-
1991	47	3	3	3	3	2	4	5	9	8	-
1990	51	1	3	2	5	3	11	7	8	5	-
1989	43	1	5	1	2	4	5	6	6	6	-
1988	47	-	2	6	2	4	4	9	3	9	-
1987	53	5	2	2	6	2	5	7	7	13	-
1986	46	-	2	3	3	6	3	3	5	11	-
1985	49	2	1	3	5	5	9	5	3	8	-
1984	39	2	3	1	2	5	4	3	8	8	-
1983	42	2	-	3	5	3	5	9	7	2	-
1982	51	1	2	7	4	3	5	5	3	5	-
1981	33	1	2	2	3	7	3	3	1	7	-
1980	44	3	1	2	1	4	5	6	9	5	-
1979	42	1	2	3	4	4	5	5	4	2	-
Other disorders of kidney (588-589)											
All races, male											
1998	7	-	-	1	1	-	-	1	3	-	-
1997	20	1	1	2	1	1	3	2	2	4	-
1996	18	-	1	1	1	-	3	3	3	3	-
1995	22	-	1	-	1	4	4	2	3	5	-
1994	21	-	-	1	3	1	5	2	3	4	-
1993	15	1	-	-	-	1	2	3	3	3	-
1992	15	1	-	2	1	2	2	1	2	2	-
1991	20	2	1	1	2	2	1	2	4	2	-
1990	21	1	1	2	2	1	6	4	1	-	-
1989	18	1	2	1	-	1	-	4	3	2	-
1988	21	-	1	2	-	4	1	3	2	3	-
1987	23	3	1	1	2	1	3	5	2	3	-
1986	18	-	2	1	2	1	1	-	2	4	-
1985	15	-	-	-	-	2	3	1	1	2	-
1984	14	1	2	-	1	2	2	-	3	2	-
1983	19	-	-	1	2	2	1	5	2	2	-
1982	26	-	2	4	1	2	3	5	-	2	-
1981	9	1	-	1	1	2	1	-	-	3	-
1980	21	3	-	1	-	-	1	1	5	3	-
1979	23	-	-	2	2	2	3	5	2	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
All races, female											
1998	24	2	1	2	-	2	2	1	5	5	-
1997	26	2	-	-	1	4	2	2	4	9	-
1996	24	4	1	1	-	-	3	3	4	4	-
1995	25	-	3	1	1	4	1	2	4	5	-
1994	30	2	1	3	1	3	7	5	3	4	-
1993	33	1	3	1	2	3	4	4	5	6	-
1992	28	-	-	2	-	3	2	8	-	9	-
1991	27	1	2	2	1	-	3	3	5	6	-
1990	30	-	2	-	3	2	5	3	7	5	-
1989	25	-	3	-	2	3	5	2	3	4	-
1988	26	-	1	4	2	-	3	6	1	6	-
1987	30	2	1	1	4	1	2	2	5	10	-
1986	28	-	-	2	1	5	2	3	3	7	-
1985	34	2	1	3	5	3	6	4	2	6	-
1984	25	1	1	1	1	3	2	3	5	6	-
1983	23	2	-	2	3	1	4	4	5	-	-
1982	25	1	-	3	3	1	2	-	3	3	-
1981	24	-	2	1	2	5	2	3	1	4	-
1980	23	-	1	1	1	4	4	5	4	2	-
1979	19	1	2	1	2	2	2	-	2	1	-
Other disorders of kidney (588-589)											
White, both sexes											
1998	26	1	1	2	-	2	2	2	8	5	-
1997	36	2	-	1	-	5	5	4	5	11	-
1996	37	4	2	2	-	-	5	4	7	6	-
1995	33	-	3	-	2	6	3	4	5	7	-
1994	41	2	1	4	2	4	9	6	5	6	-
1993	32	2	3	-	1	3	5	4	7	5	-
1992	26	1	-	2	1	4	1	5	2	7	-
1991	34	2	2	2	2	1	4	4	8	5	-
1990	32	1	2	1	2	3	6	6	7	4	-
1989	35	1	4	1	2	4	4	4	5	5	-
1988	34	-	1	4	2	2	2	7	3	6	-
1987	40	4	2	1	5	2	3	7	4	11	-
1986	32	-	1	1	3	6	2	2	5	5	-
1985	35	2	-	1	5	3	8	5	1	5	-
1984	26	2	2	1	2	3	3	2	5	4	-
1983	33	1	-	2	4	2	3	7	7	2	-
1982	44	1	1	5	4	3	4	3	3	5	-
1981	25	1	1	2	2	6	3	2	-	5	-
1980	33	1	1	1	1	3	4	4	9	3	-
1979	31	1	2	2	2	3	3	4	3	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
White, male											
1998	5	-	-	1	-	-	-	1	3	-	-
1997	16	1	-	1	-	1	3	2	2	4	-
1996	14	-	1	1	-	-	2	1	3	3	-
1995	17	-	1	-	1	3	3	2	3	4	-
1994	16	-	-	1	1	1	4	2	3	3	-
1993	11	1	-	-	-	-	2	2	3	2	-
1992	9	1	-	1	1	1	-	1	2	1	-
1991	17	1	1	1	2	1	1	2	4	2	-
1990	12	1	1	1	1	1	4	3	-	-	-
1989	15	1	1	1	-	1	-	4	3	2	-
1988	12	-	-	1	-	2	-	2	2	1	-
1987	17	2	1	1	1	1	2	5	1	2	-
1986	10	-	1	-	2	1	-	-	2	1	-
1985	9	-	-	-	-	1	3	1	-	-	-
1984	12	1	2	-	1	1	2	-	3	1	-
1983	14	-	-	1	1	1	-	4	2	2	-
1982	21	-	1	3	1	2	2	3	-	2	-
1981	7	1	-	1	1	1	1	-	-	2	-
1980	15	1	-	-	-	-	1	-	5	2	-
1979	18	-	-	2	2	1	2	4	2	1	-
Other disorders of kidney (588-589)											
White, female											
1998	21	1	1	1	-	2	2	1	5	5	-
1997	20	1	-	-	-	4	2	2	3	7	-
1996	23	4	1	1	-	-	3	3	4	3	-
1995	16	-	2	-	1	3	-	2	2	3	-
1994	25	2	1	3	1	3	5	4	2	3	-
1993	21	1	3	-	1	3	3	2	4	3	-
1992	17	-	-	1	-	3	1	4	-	6	-
1991	17	1	1	1	-	-	3	2	4	3	-
1990	20	-	1	-	1	2	2	3	7	4	-
1989	20	-	3	-	2	3	4	-	2	3	-
1988	22	-	1	3	2	-	2	5	1	5	-
1987	23	2	1	-	4	1	1	2	3	9	-
1986	22	-	-	1	1	5	2	2	3	4	-
1985	26	2	-	1	5	2	5	4	1	5	-
1984	14	1	-	1	1	2	1	2	2	3	-
1983	19	1	-	1	3	1	3	3	5	-	-
1982	23	1	-	2	3	1	2	-	3	3	-
1981	18	-	1	1	1	5	2	2	-	3	-
1980	18	-	1	1	1	3	3	4	4	1	-
1979	13	1	2	-	-	2	1	-	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
All other, both sexes											
1998	5	1	-	1	1	-	-	-	-	-	-
1997	10	1	1	1	2	-	-	-	1	2	-
1996	5	-	-	-	1	-	1	2	-	1	-
1995	14	-	1	1	-	2	2	-	2	3	-
1994	10	-	-	-	2	-	3	1	1	2	-
1993	16	-	-	1	1	1	1	3	1	4	-
1992	17	-	-	2	-	1	3	4	-	4	-
1991	13	1	1	1	1	1	-	1	1	3	-
1990	19	-	1	1	3	-	5	1	1	1	-
1989	8	-	1	-	-	-	1	2	1	1	-
1988	13	-	1	2	-	2	2	2	-	3	-
1987	13	1	-	1	1	-	2	-	3	2	-
1986	14	-	1	2	-	-	1	1	-	6	-
1985	14	-	1	2	-	2	1	-	2	3	-
1984	13	-	1	-	-	2	1	1	3	4	-
1983	9	1	-	1	1	1	2	2	-	-	-
1982	7	-	1	2	-	-	1	2	-	-	-
1981	8	-	1	-	1	1	-	1	1	2	-
1980	11	2	-	1	-	1	1	2	-	2	-
1979	11	-	-	1	2	1	2	1	1	-	-
Other disorders of kidney (588-589)											
All other, male											
1998	2	-	-	-	1	-	-	-	-	-	-
1997	4	-	1	1	1	-	-	-	-	-	-
1996	4	-	-	-	1	-	1	2	-	-	-
1995	5	-	-	-	-	1	1	-	-	1	-
1994	5	-	-	-	2	-	1	-	-	1	-
1993	4	-	-	-	-	1	-	1	-	1	-
1992	6	-	-	1	-	1	2	-	-	1	-
1991	3	1	-	-	-	1	-	-	-	-	-
1990	9	-	-	1	1	-	2	1	1	-	-
1989	3	-	1	-	-	-	-	-	-	-	-
1988	9	-	1	1	-	2	1	1	-	2	-
1987	6	1	-	-	1	-	1	-	1	1	-
1986	8	-	1	1	-	-	1	-	-	3	-
1985	6	-	-	-	-	1	-	-	1	2	-
1984	2	-	-	-	-	1	-	-	-	1	-
1983	5	-	-	-	1	1	1	1	-	-	-
1982	5	-	1	1	-	-	1	2	-	-	-
1981	2	-	-	-	-	1	-	-	-	1	-
1980	6	2	-	1	-	-	-	1	-	1	-
1979	5	-	-	-	-	1	1	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
All other, female											
1998	3	1	-	1	-	-	-	-	-	-	-
1997	6	1	-	-	1	-	-	-	1	2	-
1996	1	-	-	-	-	-	-	-	-	1	-
1995	9	-	1	1	-	1	1	-	2	2	-
1994	5	-	-	-	-	-	2	1	1	1	-
1993	12	-	-	1	1	-	1	2	1	3	-
1992	11	-	-	1	-	-	1	4	-	3	-
1991	10	-	1	1	1	-	-	1	1	3	-
1990	10	-	1	-	2	-	3	-	-	1	-
1989	5	-	-	-	-	-	1	2	1	1	-
1988	4	-	-	1	-	-	1	1	-	1	-
1987	7	-	-	1	-	-	1	-	2	1	-
1986	6	-	-	1	-	-	-	1	-	3	-
1985	8	-	1	2	-	1	1	-	1	1	-
1984	11	-	1	-	-	1	1	1	3	3	-
1983	4	1	-	1	-	-	1	1	-	-	-
1982	2	-	-	1	-	-	-	-	-	-	-
1981	6	-	1	-	1	-	-	1	1	1	-
1980	5	-	-	-	-	1	1	1	-	1	-
1979	6	-	-	1	2	-	1	-	1	-	-
Other disorders of kidney (588-589)											
Black, both sexes											
1998	4	-	-	1	1	-	-	-	-	-	-
1997	10	1	1	1	2	-	-	-	1	2	-
1996	5	-	-	-	1	-	1	2	-	1	-
1995	14	-	1	1	-	2	2	-	2	3	-
1994	9	-	-	-	2	-	3	1	1	1	-
1993	13	-	-	1	1	1	-	3	-	4	-
1992	17	-	-	2	-	1	3	4	-	4	-
1991	13	1	1	1	1	1	-	1	1	3	-
1990	19	-	1	1	3	-	5	1	1	1	-
1989	7	-	1	-	-	-	1	2	1	-	-
1988	12	-	1	2	-	2	1	2	-	3	-
1987	11	1	-	1	1	-	1	-	3	2	-
1986	14	-	1	2	-	-	1	1	-	6	-
1985	14	-	1	2	-	2	1	-	2	3	-
1984	13	-	1	-	-	2	1	1	3	4	-
1983	8	1	-	1	1	1	2	1	-	-	-
1982	7	-	1	2	-	-	1	2	-	-	-
1981	7	-	1	-	1	1	-	-	1	2	-
1980	10	2	-	1	-	-	1	2	-	2	-
1979	10	-	-	1	1	1	2	1	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other disorders of kidney (588-589)											
Black, male											
1998	2	-	-	-	1	-	-	-	-	-	-
1997	4	-	1	1	1	-	-	-	-	-	-
1996	4	-	-	-	1	-	1	2	-	-	-
1995	5	-	-	-	-	1	1	-	-	1	-
1994	5	-	-	-	2	-	1	-	-	1	-
1993	4	-	-	-	-	1	-	1	-	1	-
1992	6	-	-	1	-	1	2	-	-	1	-
1991	3	1	-	-	-	1	-	-	-	-	-
1990	9	-	-	1	1	-	2	1	1	-	-
1989	3	-	1	-	-	-	-	-	-	-	-
1988	9	-	1	1	-	2	1	1	-	2	-
1987	5	1	-	-	1	-	-	-	1	1	-
1986	8	-	1	1	-	-	1	-	-	3	-
1985	6	-	-	-	-	1	-	-	1	2	-
1984	2	-	-	-	-	1	-	-	-	1	-
1983	5	-	-	-	1	1	1	1	-	-	-
1982	5	-	1	1	-	-	1	2	-	-	-
1981	2	-	-	-	-	1	-	-	-	1	-
1980	6	2	-	1	-	-	-	1	-	1	-
1979	5	-	-	-	-	1	1	1	-	-	-
Other disorders of kidney (588-589)											
Black, female											
1998	2	-	-	1	-	-	-	-	-	-	-
1997	6	1	-	-	1	-	-	-	1	2	-
1996	1	-	-	-	-	-	-	-	-	1	-
1995	9	-	1	1	-	1	1	-	2	2	-
1994	4	-	-	-	-	-	2	1	1	-	-
1993	9	-	-	1	1	-	-	2	-	3	-
1992	11	-	-	1	-	-	1	4	-	3	-
1991	10	-	1	1	1	-	-	1	1	3	-
1990	10	-	1	-	2	-	3	-	-	1	-
1989	4	-	-	-	-	-	1	2	1	-	-
1988	3	-	-	1	-	-	-	1	-	1	-
1987	6	-	-	1	-	-	1	-	2	1	-
1986	6	-	-	1	-	-	-	1	-	3	-
1985	8	-	1	2	-	1	1	-	1	1	-
1984	11	-	1	-	-	1	1	1	3	3	-
1983	3	1	-	1	-	-	1	-	-	-	-
1982	2	-	-	1	-	-	-	-	-	-	-
1981	5	-	1	-	1	-	-	-	1	1	-
1980	4	-	-	-	-	-	1	1	-	1	-
1979	5	-	-	1	1	-	1	-	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
All races, both sexes											
1998	828	23	20	26	32	44	76	100	140	322	-
1997	831	22	20	17	35	40	87	98	158	315	-
1996	896	12	16	25	40	62	80	130	154	320	1
1995	918	23	19	21	39	50	94	128	144	352	-
1994	973	17	21	19	45	60	80	149	183	352	-
1993	1,023	22	25	27	42	49	84	148	197	382	-
1992	1,083	17	22	23	41	63	87	149	189	439	-
1991	1,219	19	26	27	43	61	106	162	253	449	-
1990	1,296	19	25	34	59	77	122	188	248	466	-
1989	1,414	19	20	29	51	95	133	199	269	523	1
1988	1,610	23	33	39	52	107	129	246	318	591	1
1987	1,742	20	31	44	67	112	190	232	359	630	-
1986	1,870	20	36	51	74	130	168	246	358	727	2
1985	1,982	18	30	49	79	118	189	334	414	691	2
1984	2,077	20	38	56	101	127	234	286	417	720	-
1983	2,089	23	34	58	104	127	221	351	391	717	-
1982	2,218	32	39	57	105	130	288	355	431	713	-
1981	2,305	32	49	73	102	152	239	402	487	699	-
1980	2,732	47	47	76	119	212	325	439	519	862	-
1979	2,868	42	58	94	119	218	352	479	529	883	-
Infections of kidney (590)											
All races, male											
1998	214	11	7	7	11	11	21	24	40	62	-
1997	241	5	8	5	12	8	25	39	52	72	-
1996	255	-	4	6	14	21	32	44	42	71	1
1995	241	8	5	5	14	18	29	39	34	73	-
1994	261	6	3	9	19	20	32	41	42	78	-
1993	276	2	11	8	14	17	27	49	56	79	-
1992	327	6	2	3	17	22	33	55	63	104	-
1991	327	7	10	7	12	12	36	55	66	90	-
1990	404	10	10	15	27	22	50	58	72	117	-
1989	435	8	8	10	21	26	63	66	76	128	-
1988	496	12	11	17	16	45	40	90	92	145	-
1987	536	5	13	20	18	37	84	75	102	159	-
1986	562	11	11	16	25	38	63	83	122	174	2
1985	691	5	14	24	31	38	73	135	148	202	-
1984	713	6	15	24	34	54	95	118	137	201	-
1983	715	8	12	22	35	48	86	136	135	210	-
1982	733	9	17	26	33	49	108	127	139	205	-
1981	838	8	12	30	44	54	94	147	189	230	-
1980	960	21	17	31	40	75	127	156	183	279	-
1979	1,095	15	21	39	50	92	143	184	213	302	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
All races, female											
1998	614	12	13	19	21	33	55	76	100	260	-
1997	590	17	12	12	23	32	62	59	106	243	-
1996	641	12	12	19	26	41	48	86	112	249	-
1995	677	15	14	16	25	32	65	89	110	279	-
1994	712	11	18	10	26	40	48	108	141	274	-
1993	747	20	14	19	28	32	57	99	141	303	-
1992	756	11	20	20	24	41	54	94	126	335	-
1991	892	12	16	20	31	49	70	107	187	359	-
1990	892	9	15	19	32	55	72	130	176	349	-
1989	979	11	12	19	30	69	70	133	193	395	1
1988	1,114	11	22	22	36	62	89	156	226	446	1
1987	1,206	15	18	24	49	75	106	157	257	471	-
1986	1,308	9	25	35	49	92	105	163	236	553	-
1985	1,291	13	16	25	48	80	116	199	266	489	2
1984	1,364	14	23	32	67	73	139	168	280	519	-
1983	1,374	15	22	36	69	79	135	215	256	507	-
1982	1,485	23	22	31	72	81	180	228	292	508	-
1981	1,467	24	37	43	58	98	145	255	298	469	-
1980	1,772	26	30	45	79	137	198	283	336	583	-
1979	1,773	27	37	55	69	126	209	295	316	581	-
Infections of kidney (590)											
White, both sexes											
1998	704	16	18	17	21	32	65	85	125	292	-
1997	725	15	13	15	26	35	69	91	142	291	-
1996	778	6	13	23	33	52	71	117	139	285	-
1995	783	17	12	13	30	37	83	108	131	318	-
1994	830	13	14	11	40	48	65	129	161	318	-
1993	881	16	15	20	30	45	75	123	180	343	-
1992	946	11	16	18	35	53	71	136	172	406	-
1991	1,041	15	21	20	32	55	87	138	223	402	-
1990	1,134	16	20	27	49	58	106	167	220	428	-
1989	1,203	15	8	20	38	74	115	167	238	481	1
1988	1,415	16	26	29	46	94	108	209	291	552	1
1987	1,526	17	23	38	54	86	156	206	327	582	-
1986	1,623	13	23	38	59	111	141	208	324	666	2
1985	1,708	13	19	39	64	87	157	289	369	630	2
1984	1,805	16	29	40	76	103	199	252	374	664	-
1983	1,821	16	25	41	84	102	192	317	352	652	-
1982	1,953	22	30	40	90	111	243	304	400	666	-
1981	2,022	23	36	53	90	120	205	350	446	649	-
1980	2,365	27	34	53	96	175	275	371	469	802	-
1979	2,493	29	39	66	88	176	306	430	473	820	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
White, male											
1998	173	10	7	3	8	9	21	18	33	50	-
1997	203	3	5	4	8	6	17	36	48	65	-
1996	222	-	3	6	13	17	26	41	41	61	-
1995	201	5	3	4	9	15	25	32	31	66	-
1994	215	4	3	5	16	13	26	35	35	69	-
1993	226	1	6	4	7	15	23	42	51	67	-
1992	277	5	1	3	13	18	26	50	55	94	-
1991	274	6	6	6	10	12	28	45	57	83	-
1990	339	8	7	11	22	13	44	50	62	107	-
1989	358	7	4	6	15	18	53	59	66	111	-
1988	414	8	9	13	12	36	31	73	84	130	-
1987	458	4	9	18	14	31	65	64	93	144	-
1986	479	11	9	13	18	35	50	68	108	153	2
1985	582	3	8	19	23	25	58	114	133	184	-
1984	597	5	10	17	21	45	79	103	122	174	-
1983	613	7	8	16	27	39	78	119	117	184	-
1982	629	5	12	16	29	42	95	107	123	184	-
1981	741	5	11	23	38	42	82	128	177	213	-
1980	813	15	12	22	34	60	106	123	161	255	-
1979	935	10	14	25	35	71	124	163	185	279	-
Infections of kidney (590)											
White, female											
1998	531	6	11	14	13	23	44	67	92	242	-
1997	522	12	8	11	18	29	52	55	94	226	-
1996	556	6	10	17	20	35	45	76	98	224	-
1995	582	12	9	9	21	22	58	76	100	252	-
1994	615	9	11	6	24	35	39	94	126	249	-
1993	655	15	9	16	23	30	52	81	129	276	-
1992	669	6	15	15	22	35	45	86	117	312	-
1991	767	9	15	14	22	43	59	93	166	319	-
1990	795	8	13	16	27	45	62	117	158	321	-
1989	845	8	4	14	23	56	62	108	172	370	1
1988	1,001	8	17	16	34	58	77	136	207	422	1
1987	1,068	13	14	20	40	55	91	142	234	438	-
1986	1,144	2	14	25	41	76	91	140	216	513	-
1985	1,126	10	11	20	41	62	99	175	236	446	2
1984	1,208	11	19	23	55	58	120	149	252	490	-
1983	1,208	9	17	25	57	63	114	198	235	468	-
1982	1,324	17	18	24	61	69	148	197	277	482	-
1981	1,281	18	25	30	52	78	123	222	269	436	-
1980	1,552	12	22	31	62	115	169	248	308	547	-
1979	1,558	19	25	41	53	105	182	267	288	541	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
All other, both sexes											
1998	124	7	2	9	11	12	11	15	15	30	-
1997	106	7	7	2	9	5	18	7	16	24	-
1996	118	6	3	2	7	10	9	13	15	35	1
1995	135	6	7	8	9	13	11	20	13	34	-
1994	143	4	7	8	5	12	15	20	22	34	-
1993	142	6	10	7	12	4	9	25	17	39	-
1992	137	6	6	5	6	10	16	13	17	33	-
1991	178	4	5	7	11	6	19	24	30	47	-
1990	162	3	5	7	10	19	16	21	28	38	-
1989	211	4	12	9	13	21	18	32	31	42	-
1988	195	7	7	10	6	13	21	37	27	39	-
1987	216	3	8	6	13	26	34	26	32	48	-
1986	247	7	13	13	15	19	27	38	34	61	-
1985	274	5	11	10	15	31	32	45	45	61	-
1984	272	4	9	16	25	24	35	34	43	56	-
1983	268	7	9	17	20	25	29	34	39	65	-
1982	265	10	9	17	15	19	45	51	31	47	-
1981	283	9	13	20	12	32	34	52	41	50	-
1980	367	20	13	23	23	37	50	68	50	60	-
1979	375	13	19	28	31	42	46	49	56	63	-
Infections of kidney (590)											
All other, male											
1998	41	1	-	4	3	2	-	6	7	12	-
1997	38	2	3	1	4	2	8	3	4	7	-
1996	33	-	1	-	1	4	6	3	1	10	1
1995	40	3	2	1	5	3	4	7	3	7	-
1994	46	2	-	4	3	7	6	6	7	9	-
1993	50	1	5	4	7	2	4	7	5	12	-
1992	50	1	1	-	4	4	7	5	8	10	-
1991	53	1	4	1	2	-	8	10	9	7	-
1990	65	2	3	4	5	9	6	8	10	10	-
1989	77	1	4	4	6	8	10	7	10	17	-
1988	82	4	2	4	4	9	9	17	8	15	-
1987	78	1	4	2	4	6	19	11	9	15	-
1986	83	-	2	3	7	3	13	15	14	21	-
1985	109	2	6	5	8	13	15	21	15	18	-
1984	116	1	5	7	13	9	16	15	15	27	-
1983	102	1	4	6	8	9	8	17	18	26	-
1982	104	4	5	10	4	7	13	20	16	21	-
1981	97	3	1	7	6	12	12	19	12	17	-
1980	147	6	5	9	6	15	21	33	22	24	-
1979	160	5	7	14	15	21	19	21	28	23	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
All other, female											
1998	83	6	2	5	8	10	11	9	8	18	-
1997	68	5	4	1	5	3	10	4	12	17	-
1996	85	6	2	2	6	6	3	10	14	25	-
1995	95	3	5	7	4	10	7	13	10	27	-
1994	97	2	7	4	2	5	9	14	15	25	-
1993	92	5	5	3	5	2	5	18	12	27	-
1992	87	5	5	5	2	6	9	8	9	23	-
1991	125	3	1	6	9	6	11	14	21	40	-
1990	97	1	2	3	5	10	10	13	18	28	-
1989	134	3	8	5	7	13	8	25	21	25	-
1988	113	3	5	6	2	4	12	20	19	24	-
1987	138	2	4	4	9	20	15	15	23	33	-
1986	164	7	11	10	8	16	14	23	20	40	-
1985	165	3	5	5	7	18	17	24	30	43	-
1984	156	3	4	9	12	15	19	19	28	29	-
1983	166	6	5	11	12	16	21	17	21	39	-
1982	161	6	4	7	11	12	32	31	15	26	-
1981	186	6	12	13	6	20	22	33	29	33	-
1980	220	14	8	14	17	22	29	35	28	36	-
1979	215	8	12	14	16	21	27	28	28	40	-
Infections of kidney (590)											
Black, both sexes											
1998	102	6	2	9	10	8	9	10	15	24	-
1997	91	7	7	2	6	2	16	7	15	21	-
1996	106	5	3	2	6	9	9	11	14	33	1
1995	111	5	5	5	8	10	8	17	11	31	-
1994	125	4	6	7	4	11	12	17	21	30	-
1993	127	6	9	4	9	4	8	23	16	36	-
1992	119	5	6	4	6	10	16	10	14	29	-
1991	161	3	5	7	11	3	19	22	26	42	-
1990	138	3	5	6	9	16	14	14	25	34	-
1989	194	4	10	9	13	19	16	30	26	40	-
1988	175	6	7	8	6	11	19	32	24	38	-
1987	196	3	7	5	12	25	33	25	29	40	-
1986	223	5	13	11	13	17	24	36	32	54	-
1985	249	5	10	10	13	28	28	44	40	54	-
1984	248	4	9	15	24	22	32	30	36	50	-
1983	254	7	9	16	20	25	29	33	38	55	-
1982	249	10	8	17	14	17	44	48	30	42	-
1981	267	7	12	19	12	29	31	49	41	49	-
1980	350	18	12	22	22	37	49	63	48	56	-
1979	350	12	18	26	28	40	44	49	53	55	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Infections of kidney (590)											
Black, male											
1998	33	1	-	4	3	1	-	4	7	9	-
1997	33	2	3	1	3	-	7	3	4	6	-
1996	30	-	1	-	1	3	6	3	1	10	1
1995	36	3	-	1	5	3	4	7	2	6	-
1994	42	2	-	4	3	7	4	6	7	7	-
1993	44	1	4	4	6	2	3	7	5	9	-
1992	44	1	1	-	4	4	7	4	7	8	-
1991	50	1	4	1	2	-	8	9	9	7	-
1990	57	2	3	3	5	8	5	7	8	9	-
1989	68	1	2	4	6	7	9	7	8	15	-
1988	74	3	2	3	4	7	8	16	7	15	-
1987	74	1	4	1	4	6	19	11	8	14	-
1986	76	-	2	3	7	3	11	14	13	20	-
1985	103	2	6	5	6	12	12	21	15	18	-
1984	107	1	5	7	12	8	15	12	14	25	-
1983	96	1	4	6	8	9	8	16	18	21	-
1982	98	4	4	10	3	7	13	18	16	19	-
1981	93	3	1	6	6	12	11	19	12	17	-
1980	138	4	4	9	5	15	21	30	22	22	-
1979	153	5	7	13	15	20	18	21	26	21	-
Infections of kidney (590)											
Black, female											
1998	69	5	2	5	7	7	9	6	8	15	-
1997	58	5	4	1	3	2	9	4	11	15	-
1996	76	5	2	2	5	6	3	8	13	23	-
1995	75	2	5	4	3	7	4	10	9	25	-
1994	83	2	6	3	1	4	8	11	14	23	-
1993	83	5	5	-	3	2	5	16	11	27	-
1992	75	4	5	4	2	6	9	6	7	21	-
1991	111	2	1	6	9	3	11	13	17	35	-
1990	81	1	2	3	4	8	9	7	17	25	-
1989	126	3	8	5	7	12	7	23	18	25	-
1988	101	3	5	5	2	4	11	16	17	23	-
1987	122	2	3	4	8	19	14	14	21	26	-
1986	147	5	11	8	6	14	13	22	19	34	-
1985	146	3	4	5	7	16	16	23	25	36	-
1984	141	3	4	8	12	14	17	18	22	25	-
1983	158	6	5	10	12	16	21	17	20	34	-
1982	151	6	4	7	11	10	31	30	14	23	-
1981	174	4	11	13	6	17	20	30	29	32	-
1980	212	14	8	13	17	22	28	33	26	34	-
1979	197	7	11	13	13	20	26	28	27	34	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hyperplasia of prostate (600)											
All races, both sexes											
1998	410	-	1	3	10	9	17	65	98	206	1
1997	420	1	3	1	10	9	24	62	92	218	-
1996	457	1	2	2	8	11	36	64	102	231	-
1995	406	-	-	4	2	19	39	49	92	201	-
1994	413	1	1	4	8	25	29	69	101	172	-
1993	410	-	2	3	10	17	32	69	89	187	-
1992	397	-	-	3	5	16	36	68	75	194	-
1991	428	1	2	4	6	20	40	79	88	188	-
1990	455	-	-	8	9	17	41	81	99	200	-
1989	449	1	2	4	9	28	32	85	99	186	-
1988	454	1	-	4	22	31	40	80	97	179	-
1987	461	1	-	3	8	23	43	91	91	201	-
1986	508	-	-	9	15	30	53	85	90	225	1
1985	479	-	-	3	15	26	46	91	95	203	-
1984	536	1	3	5	18	21	56	92	121	219	-
1983	572	-	2	5	17	33	61	95	134	225	-
1982	655	1	-	5	18	43	62	108	133	285	-
1981	693	-	1	10	15	44	82	112	167	261	1
1980	754	-	4	2	25	47	85	131	179	281	-
1979	810	-	2	6	20	60	101	139	196	286	-
Hyperplasia of prostate (600)											
All races, male											
1998	410	-	1	3	10	9	17	65	98	206	1
1997	420	1	3	1	10	9	24	62	92	218	-
1996	457	1	2	2	8	11	36	64	102	231	-
1995	406	-	-	4	2	19	39	49	92	201	-
1994	413	1	1	4	8	25	29	69	101	172	-
1993	410	-	2	3	10	17	32	69	89	187	-
1992	397	-	-	3	5	16	36	68	75	194	-
1991	428	1	2	4	6	20	40	79	88	188	-
1990	455	-	-	8	9	17	41	81	99	200	-
1989	449	1	2	4	9	28	32	85	99	186	-
1988	454	1	-	4	22	31	40	80	97	179	-
1987	461	1	-	3	8	23	43	91	91	201	-
1986	508	-	-	9	15	30	53	85	90	225	1
1985	479	-	-	3	15	26	46	91	95	203	-
1984	536	1	3	5	18	21	56	92	121	219	-
1983	572	-	2	5	17	33	61	95	134	225	-
1982	655	1	-	5	18	43	62	108	133	285	-
1981	693	-	1	10	15	44	82	112	167	261	1
1980	754	-	4	2	25	47	85	131	179	281	-
1979	810	-	2	6	20	60	101	139	196	286	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hyperplasia of prostate (600)											
White, both sexes											
1998	356	-	1	2	6	9	12	49	87	190	-
1997	370	1	1	-	9	6	17	54	84	198	-
1996	408	1	2	2	5	7	28	60	92	211	-
1995	364	-	-	4	2	16	33	40	83	186	-
1994	372	1	-	3	8	19	23	63	91	163	-
1993	350	-	1	2	8	14	27	56	74	167	-
1992	358	-	-	2	2	11	34	59	73	177	-
1991	372	1	-	3	4	17	33	70	74	170	-
1990	402	-	-	7	8	15	31	69	88	184	-
1989	375	-	2	1	5	23	23	71	85	165	-
1988	391	1	-	3	16	24	36	66	86	159	-
1987	406	1	-	1	7	15	34	81	78	189	-
1986	439	-	-	7	13	21	42	68	79	209	-
1985	422	-	-	1	11	21	38	81	86	184	-
1984	474	1	2	2	17	17	47	80	107	201	-
1983	501	-	-	4	14	27	43	83	118	212	-
1982	565	1	-	3	17	33	50	85	111	265	-
1981	600	-	1	6	11	35	68	90	152	237	-
1980	654	-	4	1	17	37	65	112	161	257	-
1979	703	-	2	4	15	45	77	117	178	265	-
Hyperplasia of prostate (600)											
White, male											
1998	356	-	1	2	6	9	12	49	87	190	-
1997	370	1	1	-	9	6	17	54	84	198	-
1996	408	1	2	2	5	7	28	60	92	211	-
1995	364	-	-	4	2	16	33	40	83	186	-
1994	372	1	-	3	8	19	23	63	91	163	-
1993	350	-	1	2	8	14	27	56	74	167	-
1992	358	-	-	2	2	11	34	59	73	177	-
1991	372	1	-	3	4	17	33	70	74	170	-
1990	402	-	-	7	8	15	31	69	88	184	-
1989	375	-	2	1	5	23	23	71	85	165	-
1988	391	1	-	3	16	24	36	66	86	159	-
1987	406	1	-	1	7	15	34	81	78	189	-
1986	439	-	-	7	13	21	42	68	79	209	-
1985	422	-	-	1	11	21	38	81	86	184	-
1984	474	1	2	2	17	17	47	80	107	201	-
1983	501	-	-	4	14	27	43	83	118	212	-
1982	565	1	-	3	17	33	50	85	111	265	-
1981	600	-	1	6	11	35	68	90	152	237	-
1980	654	-	4	1	17	37	65	112	161	257	-
1979	703	-	2	4	15	45	77	117	178	265	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hyperplasia of prostate (600)											
All other, both sexes											
1998	54	-	-	1	4	-	5	16	11	16	1
1997	50	-	2	1	1	3	7	8	8	20	-
1996	49	-	-	-	3	4	8	4	10	20	-
1995	42	-	-	-	-	3	6	9	9	15	-
1994	41	-	1	1	-	6	6	6	10	9	-
1993	60	-	1	1	2	3	5	13	15	20	-
1992	39	-	-	1	3	5	2	9	2	17	-
1991	56	-	2	1	2	3	7	9	14	18	-
1990	53	-	-	1	1	2	10	12	11	16	-
1989	74	1	-	3	4	5	9	14	14	21	-
1988	63	-	-	1	6	7	4	14	11	20	-
1987	55	-	-	2	1	8	9	10	13	12	-
1986	69	-	-	2	2	9	11	17	11	16	1
1985	57	-	-	2	4	5	8	10	9	19	-
1984	62	-	1	3	1	4	9	12	14	18	-
1983	71	-	2	1	3	6	18	12	16	13	-
1982	90	-	-	2	1	10	12	23	22	20	-
1981	93	-	-	4	4	9	14	22	15	24	1
1980	100	-	-	1	8	10	20	19	18	24	-
1979	107	-	-	2	5	15	24	22	18	21	-
Hyperplasia of prostate (600)											
All other, male											
1998	54	-	-	1	4	-	5	16	11	16	1
1997	50	-	2	1	1	3	7	8	8	20	-
1996	49	-	-	-	3	4	8	4	10	20	-
1995	42	-	-	-	-	3	6	9	9	15	-
1994	41	-	1	1	-	6	6	6	10	9	-
1993	60	-	1	1	2	3	5	13	15	20	-
1992	39	-	-	1	3	5	2	9	2	17	-
1991	56	-	2	1	2	3	7	9	14	18	-
1990	53	-	-	1	1	2	10	12	11	16	-
1989	74	1	-	3	4	5	9	14	14	21	-
1988	63	-	-	1	6	7	4	14	11	20	-
1987	55	-	-	2	1	8	9	10	13	12	-
1986	69	-	-	2	2	9	11	17	11	16	1
1985	57	-	-	2	4	5	8	10	9	19	-
1984	62	-	1	3	1	4	9	12	14	18	-
1983	71	-	2	1	3	6	18	12	16	13	-
1982	90	-	-	2	1	10	12	23	22	20	-
1981	93	-	-	4	4	9	14	22	15	24	1
1980	100	-	-	1	8	10	20	19	18	24	-
1979	107	-	-	2	5	15	24	22	18	21	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Hyperplasia of prostate (600)											
Black, both sexes											
1998	45	-	-	1	4	-	4	14	8	13	1
1997	46	-	2	1	1	3	5	8	7	19	-
1996	43	-	-	-	3	3	8	4	7	18	-
1995	36	-	-	-	-	3	6	9	7	11	-
1994	36	-	1	1	-	6	5	6	9	6	-
1993	53	-	1	1	2	2	5	12	11	19	-
1992	36	-	-	1	3	5	2	9	2	14	-
1991	50	-	2	1	2	3	7	8	12	15	-
1990	48	-	-	1	1	2	9	12	10	13	-
1989	70	1	-	3	4	5	9	14	13	18	-
1988	58	-	-	1	6	7	3	12	10	19	-
1987	50	-	-	1	1	8	9	9	13	9	-
1986	65	-	-	2	2	9	11	16	9	15	1
1985	55	-	-	2	4	5	8	8	9	19	-
1984	61	-	1	3	1	4	9	12	14	17	-
1983	69	-	2	1	3	5	18	12	16	12	-
1982	87	-	-	2	1	10	12	23	20	19	-
1981	88	-	-	4	4	8	14	20	15	22	1
1980	91	-	-	-	7	10	19	19	15	21	-
1979	103	-	-	2	5	15	23	22	18	18	-
Hyperplasia of prostate (600)											
Black, male											
1998	45	-	-	1	4	-	4	14	8	13	1
1997	46	-	2	1	1	3	5	8	7	19	-
1996	43	-	-	-	3	3	8	4	7	18	-
1995	36	-	-	-	-	3	6	9	7	11	-
1994	36	-	1	1	-	6	5	6	9	6	-
1993	53	-	1	1	2	2	5	12	11	19	-
1992	36	-	-	1	3	5	2	9	2	14	-
1991	50	-	2	1	2	3	7	8	12	15	-
1990	48	-	-	1	1	2	9	12	10	13	-
1989	70	1	-	3	4	5	9	14	13	18	-
1988	58	-	-	1	6	7	3	12	10	19	-
1987	50	-	-	1	1	8	9	9	13	9	-
1986	65	-	-	2	2	9	11	16	9	15	1
1985	55	-	-	2	4	5	8	8	9	19	-
1984	61	-	1	3	1	4	9	12	14	17	-
1983	69	-	2	1	3	5	18	12	16	12	-
1982	87	-	-	2	1	10	12	23	20	19	-
1981	88	-	-	4	4	8	14	20	15	22	1
1980	91	-	-	-	7	10	19	19	15	21	-
1979	103	-	-	2	5	15	23	22	18	18	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Inflammatory diseases of female pelvic organs (614-616)											
All races, both sexes											
1998	101	3	3	6	7	5	5	13	12	23	-
1997	109	8	4	4	3	7	9	12	19	20	-
1996	113	10	4	3	4	8	9	17	15	23	-
1995	139	9	7	3	4	13	11	23	10	24	-
1994	117	8	5	5	7	9	11	11	14	19	-
1993	124	8	5	9	6	9	9	13	15	22	-
1992	113	3	3	7	7	6	5	13	17	21	-
1991	128	11	4	7	4	14	12	13	8	31	-
1990	120	3	5	-	9	6	14	12	21	26	-
1989	118	8	4	4	6	13	10	15	14	20	-
1988	144	13	8	11	16	7	15	18	13	20	-
1987	130	3	4	5	12	12	15	23	12	23	-
1986	149	7	5	8	8	9	18	21	18	25	-
1985	152	6	5	5	12	17	11	23	21	22	-
1984	135	5	3	7	12	8	16	15	17	25	-
1983	149	7	8	4	8	13	18	18	15	22	-
1982	156	12	3	4	7	13	13	22	16	31	-
1981	153	9	6	6	5	10	16	17	17	25	-
1980	162	17	8	9	12	14	12	21	16	18	1
1979	164	7	5	7	17	9	17	18	13	31	-
Inflammatory diseases of female pelvic organs (614-616)											
All races, female											
1998	101	3	3	6	7	5	5	13	12	23	-
1997	109	8	4	4	3	7	9	12	19	20	-
1996	113	10	4	3	4	8	9	17	15	23	-
1995	139	9	7	3	4	13	11	23	10	24	-
1994	117	8	5	5	7	9	11	11	14	19	-
1993	124	8	5	9	6	9	9	13	15	22	-
1992	113	3	3	7	7	6	5	13	17	21	-
1991	128	11	4	7	4	14	12	13	8	31	-
1990	120	3	5	-	9	6	14	12	21	26	-
1989	118	8	4	4	6	13	10	15	14	20	-
1988	144	13	8	11	16	7	15	18	13	20	-
1987	130	3	4	5	12	12	15	23	12	23	-
1986	149	7	5	8	8	9	18	21	18	25	-
1985	152	6	5	5	12	17	11	23	21	22	-
1984	135	5	3	7	12	8	16	15	17	25	-
1983	149	7	8	4	8	13	18	18	15	22	-
1982	156	12	3	4	7	13	13	22	16	31	-
1981	153	9	6	6	5	10	16	17	17	25	-
1980	162	17	8	9	12	14	12	21	16	18	1
1979	164	7	5	7	17	9	17	18	13	31	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Inflammatory diseases of female pelvic organs (614-616)											
White, both sexes											
1998	77	3	2	4	4	4	5	9	10	22	-
1997	79	5	3	4	2	4	8	8	17	20	-
1996	81	3	3	3	2	6	8	14	11	20	-
1995	101	5	5	1	2	8	9	22	9	23	-
1994	86	4	3	3	3	5	9	11	14	19	-
1993	82	2	5	4	5	7	6	11	13	18	-
1992	81	3	2	4	6	5	4	13	14	17	-
1991	96	6	4	6	4	12	10	9	6	27	-
1990	92	-	3	-	7	5	13	9	19	23	-
1989	88	6	1	4	5	8	7	13	12	20	-
1988	99	4	4	8	11	4	12	14	13	19	-
1987	94	1	-	2	10	9	11	19	10	19	-
1986	103	5	3	5	6	7	17	12	16	21	-
1985	121	3	2	4	8	15	8	20	19	21	-
1984	97	1	1	5	8	7	14	12	12	22	-
1983	112	3	6	2	6	13	14	16	13	21	-
1982	114	5	1	2	4	10	9	19	15	28	-
1981	102	6	3	4	3	6	12	15	14	23	-
1980	119	11	6	8	9	10	10	17	16	16	-
1979	117	3	3	4	12	8	12	18	12	27	-
Inflammatory diseases of female pelvic organs (614-616)											
White, female											
1998	77	3	2	4	4	4	5	9	10	22	-
1997	79	5	3	4	2	4	8	8	17	20	-
1996	81	3	3	3	2	6	8	14	11	20	-
1995	101	5	5	1	2	8	9	22	9	23	-
1994	86	4	3	3	3	5	9	11	14	19	-
1993	82	2	5	4	5	7	6	11	13	18	-
1992	81	3	2	4	6	5	4	13	14	17	-
1991	96	6	4	6	4	12	10	9	6	27	-
1990	92	-	3	-	7	5	13	9	19	23	-
1989	88	6	1	4	5	8	7	13	12	20	-
1988	99	4	4	8	11	4	12	14	13	19	-
1987	94	1	-	2	10	9	11	19	10	19	-
1986	103	5	3	5	6	7	17	12	16	21	-
1985	121	3	2	4	8	15	8	20	19	21	-
1984	97	1	1	5	8	7	14	12	12	22	-
1983	112	3	6	2	6	13	14	16	13	21	-
1982	114	5	1	2	4	10	9	19	15	28	-
1981	102	6	3	4	3	6	12	15	14	23	-
1980	119	11	6	8	9	10	10	17	16	16	-
1979	117	3	3	4	12	8	12	18	12	27	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Inflammatory diseases of female pelvic organs (614-616)											
All other, both sexes											
1998	24	-	1	2	3	1	-	4	2	1	-
1997	30	3	1	-	1	3	1	4	2	-	-
1996	32	7	1	-	2	2	1	3	4	3	-
1995	38	4	2	2	2	5	2	1	1	1	-
1994	31	4	2	2	4	4	2	-	-	-	-
1993	42	6	-	5	1	2	3	2	2	4	-
1992	32	-	1	3	1	1	1	-	3	4	-
1991	32	5	-	1	-	2	2	4	2	4	-
1990	28	3	2	-	2	1	1	3	2	3	-
1989	30	2	3	-	1	5	3	2	2	-	-
1988	45	9	4	3	5	3	3	4	-	1	-
1987	36	2	4	3	2	3	4	4	2	4	-
1986	46	2	2	3	2	2	1	9	2	4	-
1985	31	3	3	1	4	2	3	3	2	1	-
1984	38	4	2	2	4	1	2	3	5	3	-
1983	37	4	2	2	2	-	4	2	2	1	-
1982	42	7	2	2	3	3	4	3	1	3	-
1981	51	3	3	2	2	4	4	2	3	2	-
1980	43	6	2	1	3	4	2	4	-	2	1
1979	47	4	2	3	5	1	5	-	1	4	-
Inflammatory diseases of female pelvic organs (614-616)											
All other, female											
1998	24	-	1	2	3	1	-	4	2	1	-
1997	30	3	1	-	1	3	1	4	2	-	-
1996	32	7	1	-	2	2	1	3	4	3	-
1995	38	4	2	2	2	5	2	1	1	1	-
1994	31	4	2	2	4	4	2	-	-	-	-
1993	42	6	-	5	1	2	3	2	2	4	-
1992	32	-	1	3	1	1	1	-	3	4	-
1991	32	5	-	1	-	2	2	4	2	4	-
1990	28	3	2	-	2	1	1	3	2	3	-
1989	30	2	3	-	1	5	3	2	2	-	-
1988	45	9	4	3	5	3	3	4	-	1	-
1987	36	2	4	3	2	3	4	4	2	4	-
1986	46	2	2	3	2	2	1	9	2	4	-
1985	31	3	3	1	4	2	3	3	2	1	-
1984	38	4	2	2	4	1	2	3	5	3	-
1983	37	4	2	2	2	-	4	2	2	1	-
1982	42	7	2	2	3	3	4	3	1	3	-
1981	51	3	3	2	2	4	4	2	3	2	-
1980	43	6	2	1	3	4	2	4	-	2	1
1979	47	4	2	3	5	1	5	-	1	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Inflammatory diseases of female pelvic organs (614-616)											
Black, both sexes											
1998	19	-	1	2	2	-	-	3	2	-	-
1997	27	3	1	-	1	3	1	4	2	-	-
1996	30	7	1	-	2	2	1	3	4	2	-
1995	32	4	2	2	1	4	1	1	1	1	-
1994	31	4	2	2	4	4	2	-	-	-	-
1993	39	6	-	4	1	1	3	2	2	4	-
1992	28	-	1	2	1	1	1	-	2	4	-
1991	29	5	-	1	-	2	2	4	2	3	-
1990	28	3	2	-	2	1	1	3	2	3	-
1989	29	2	3	-	1	5	2	2	2	-	-
1988	41	7	3	3	5	3	3	4	-	-	-
1987	36	2	4	3	2	3	4	4	2	4	-
1986	44	2	2	3	2	2	1	9	2	4	-
1985	31	3	3	1	4	2	3	3	2	1	-
1984	34	4	2	2	4	1	2	3	5	2	-
1983	36	4	2	2	2	-	4	2	2	1	-
1982	41	7	2	2	3	3	4	3	1	3	-
1981	51	3	3	2	2	4	4	2	3	2	-
1980	43	6	2	1	3	4	2	4	-	2	1
1979	45	4	2	3	5	1	4	-	1	4	-
Inflammatory diseases of female pelvic organs (614-616)											
Black, female											
1998	19	-	1	2	2	-	-	3	2	-	-
1997	27	3	1	-	1	3	1	4	2	-	-
1996	30	7	1	-	2	2	1	3	4	2	-
1995	32	4	2	2	1	4	1	1	1	1	-
1994	31	4	2	2	4	4	2	-	-	-	-
1993	39	6	-	4	1	1	3	2	2	4	-
1992	28	-	1	2	1	1	1	-	2	4	-
1991	29	5	-	1	-	2	2	4	2	3	-
1990	28	3	2	-	2	1	1	3	2	3	-
1989	29	2	3	-	1	5	2	2	2	-	-
1988	41	7	3	3	5	3	3	4	-	-	-
1987	36	2	4	3	2	3	4	4	2	4	-
1986	44	2	2	3	2	2	1	9	2	4	-
1985	31	3	3	1	4	2	3	3	2	1	-
1984	34	4	2	2	4	1	2	3	5	2	-
1983	36	4	2	2	2	-	4	2	2	1	-
1982	41	7	2	2	3	3	4	3	1	3	-
1981	51	3	3	2	2	4	4	2	3	2	-
1980	43	6	2	1	3	4	2	4	-	2	1
1979	45	4	2	3	5	1	4	-	1	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Pregnancy, childbirth and the puerperium (630-676)											
All races, both sexes											
1998	281	1	-	---	---	---	---	---	---	---	-
1997	327	4	1	---	---	---	---	---	---	---	-
1996	294	1	1	---	---	---	---	---	---	---	-
1995	277	1	-	---	---	---	---	---	---	---	-
1994	328	1	-	---	---	---	---	---	---	---	-
1993	302	-	-	---	---	---	---	---	---	---	-
1992	318	-	1	---	---	---	---	---	---	---	-
1991	323	5	-	---	---	---	---	---	---	---	-
1990	343	3	-	---	---	---	---	---	---	---	-
1989	320	2	-	---	---	---	---	---	---	---	-
1988	330	1	1	---	---	---	---	---	---	---	-
1987	251	1	-	---	---	---	---	---	---	---	-
1986	272	1	-	---	---	---	---	---	---	---	-
1985	295	2	1	---	---	---	---	---	---	---	-
1984	285	2	-	---	---	---	---	---	---	---	-
1983	290	-	-	---	---	---	---	---	---	---	-
1982	292	1	1	---	---	---	---	---	---	---	-
1981	309	1	-	---	---	---	---	---	---	---	-
1980	334	2	-	---	---	---	---	---	---	---	-
1979	336	5	-	---	---	---	---	---	---	---	-
Pregnancy, childbirth and the puerperium (630-676)											
All races, female											
1998	281	1	-	---	---	---	---	---	---	---	-
1997	327	4	1	---	---	---	---	---	---	---	-
1996	294	1	1	---	---	---	---	---	---	---	-
1995	277	1	-	---	---	---	---	---	---	---	-
1994	328	1	-	---	---	---	---	---	---	---	-
1993	302	-	-	---	---	---	---	---	---	---	-
1992	318	-	1	---	---	---	---	---	---	---	-
1991	323	5	-	---	---	---	---	---	---	---	-
1990	343	3	-	---	---	---	---	---	---	---	-
1989	320	2	-	---	---	---	---	---	---	---	-
1988	330	1	1	---	---	---	---	---	---	---	-
1987	251	1	-	---	---	---	---	---	---	---	-
1986	272	1	-	---	---	---	---	---	---	---	-
1985	295	2	1	---	---	---	---	---	---	---	-
1984	285	2	-	---	---	---	---	---	---	---	-
1983	290	-	-	---	---	---	---	---	---	---	-
1982	292	1	1	---	---	---	---	---	---	---	-
1981	309	1	-	---	---	---	---	---	---	---	-
1980	334	2	-	---	---	---	---	---	---	---	-
1979	336	5	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy, childbirth and the puerperium (630-676)											
White, both sexes											
1998	158	-	-	---	---	---	---	---	---	---	-
1997	179	4	-	---	---	---	---	---	---	---	-
1996	159	1	1	---	---	---	---	---	---	---	-
1995	129	-	-	---	---	---	---	---	---	---	-
1994	193	-	-	---	---	---	---	---	---	---	-
1993	152	-	-	---	---	---	---	---	---	---	-
1992	161	-	-	---	---	---	---	---	---	---	-
1991	187	3	-	---	---	---	---	---	---	---	-
1990	177	1	-	---	---	---	---	---	---	---	-
1989	180	-	-	---	---	---	---	---	---	---	-
1988	180	-	-	---	---	---	---	---	---	---	-
1987	153	1	-	---	---	---	---	---	---	---	-
1986	146	-	-	---	---	---	---	---	---	---	-
1985	156	-	1	---	---	---	---	---	---	---	-
1984	159	1	-	---	---	---	---	---	---	---	-
1983	170	-	-	---	---	---	---	---	---	---	-
1982	171	1	-	---	---	---	---	---	---	---	-
1981	184	-	-	---	---	---	---	---	---	---	-
1980	193	1	-	---	---	---	---	---	---	---	-
1979	180	2	-	---	---	---	---	---	---	---	-
Pregnancy, childbirth and the puerperium (630-676)											
White, female											
1998	158	-	-	---	---	---	---	---	---	---	-
1997	179	4	-	---	---	---	---	---	---	---	-
1996	159	1	1	---	---	---	---	---	---	---	-
1995	129	-	-	---	---	---	---	---	---	---	-
1994	193	-	-	---	---	---	---	---	---	---	-
1993	152	-	-	---	---	---	---	---	---	---	-
1992	161	-	-	---	---	---	---	---	---	---	-
1991	187	3	-	---	---	---	---	---	---	---	-
1990	177	1	-	---	---	---	---	---	---	---	-
1989	180	-	-	---	---	---	---	---	---	---	-
1988	180	-	-	---	---	---	---	---	---	---	-
1987	153	1	-	---	---	---	---	---	---	---	-
1986	146	-	-	---	---	---	---	---	---	---	-
1985	156	-	1	---	---	---	---	---	---	---	-
1984	159	1	-	---	---	---	---	---	---	---	-
1983	170	-	-	---	---	---	---	---	---	---	-
1982	171	1	-	---	---	---	---	---	---	---	-
1981	184	-	-	---	---	---	---	---	---	---	-
1980	193	1	-	---	---	---	---	---	---	---	-
1979	180	2	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy, childbirth and the puerperium (630-676)											
All other, both sexes											
1998	123	1	-	---	---	---	---	---	---	---	-
1997	148	-	1	---	---	---	---	---	---	---	-
1996	135	-	-	---	---	---	---	---	---	---	-
1995	148	1	-	---	---	---	---	---	---	---	-
1994	135	1	-	---	---	---	---	---	---	---	-
1993	150	-	-	---	---	---	---	---	---	---	-
1992	157	-	1	---	---	---	---	---	---	---	-
1991	136	2	-	---	---	---	---	---	---	---	-
1990	166	2	-	---	---	---	---	---	---	---	-
1989	140	2	-	---	---	---	---	---	---	---	-
1988	150	1	1	---	---	---	---	---	---	---	-
1987	98	-	-	---	---	---	---	---	---	---	-
1986	126	1	-	---	---	---	---	---	---	---	-
1985	139	2	-	---	---	---	---	---	---	---	-
1984	126	1	-	---	---	---	---	---	---	---	-
1983	120	-	-	---	---	---	---	---	---	---	-
1982	121	-	1	---	---	---	---	---	---	---	-
1981	125	1	-	---	---	---	---	---	---	---	-
1980	141	1	-	---	---	---	---	---	---	---	-
1979	156	3	-	---	---	---	---	---	---	---	-
Pregnancy, childbirth and the puerperium (630-676)											
All other, female											
1998	123	1	-	---	---	---	---	---	---	---	-
1997	148	-	1	---	---	---	---	---	---	---	-
1996	135	-	-	---	---	---	---	---	---	---	-
1995	148	1	-	---	---	---	---	---	---	---	-
1994	135	1	-	---	---	---	---	---	---	---	-
1993	150	-	-	---	---	---	---	---	---	---	-
1992	157	-	1	---	---	---	---	---	---	---	-
1991	136	2	-	---	---	---	---	---	---	---	-
1990	166	2	-	---	---	---	---	---	---	---	-
1989	140	2	-	---	---	---	---	---	---	---	-
1988	150	1	1	---	---	---	---	---	---	---	-
1987	98	-	-	---	---	---	---	---	---	---	-
1986	126	1	-	---	---	---	---	---	---	---	-
1985	139	2	-	---	---	---	---	---	---	---	-
1984	126	1	-	---	---	---	---	---	---	---	-
1983	120	-	-	---	---	---	---	---	---	---	-
1982	121	-	1	---	---	---	---	---	---	---	-
1981	125	1	-	---	---	---	---	---	---	---	-
1980	141	1	-	---	---	---	---	---	---	---	-
1979	156	3	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy, childbirth and the puerperium (630-676)											
Black, both sexes											
1998	104	1	-	---	---	---	---	---	---	---	-
1997	125	-	1	---	---	---	---	---	---	---	-
1996	121	-	-	---	---	---	---	---	---	---	-
1995	133	1	-	---	---	---	---	---	---	---	-
1994	118	1	-	---	---	---	---	---	---	---	-
1993	135	-	-	---	---	---	---	---	---	---	-
1992	140	-	1	---	---	---	---	---	---	---	-
1991	125	2	-	---	---	---	---	---	---	---	-
1990	153	2	-	---	---	---	---	---	---	---	-
1989	124	2	-	---	---	---	---	---	---	---	-
1988	131	1	1	---	---	---	---	---	---	---	-
1987	91	-	-	---	---	---	---	---	---	---	-
1986	117	1	-	---	---	---	---	---	---	---	-
1985	124	1	-	---	---	---	---	---	---	---	-
1984	117	1	-	---	---	---	---	---	---	---	-
1983	107	-	-	---	---	---	---	---	---	---	-
1982	108	-	1	---	---	---	---	---	---	---	-
1981	120	1	-	---	---	---	---	---	---	---	-
1980	127	1	-	---	---	---	---	---	---	---	-
1979	145	3	-	---	---	---	---	---	---	---	-
Pregnancy, childbirth and the puerperium (630-676)											
Black, female											
1998	104	1	-	---	---	---	---	---	---	---	-
1997	125	-	1	---	---	---	---	---	---	---	-
1996	121	-	-	---	---	---	---	---	---	---	-
1995	133	1	-	---	---	---	---	---	---	---	-
1994	118	1	-	---	---	---	---	---	---	---	-
1993	135	-	-	---	---	---	---	---	---	---	-
1992	140	-	1	---	---	---	---	---	---	---	-
1991	125	2	-	---	---	---	---	---	---	---	-
1990	153	2	-	---	---	---	---	---	---	---	-
1989	124	2	-	---	---	---	---	---	---	---	-
1988	131	1	1	---	---	---	---	---	---	---	-
1987	91	-	-	---	---	---	---	---	---	---	-
1986	117	1	-	---	---	---	---	---	---	---	-
1985	124	1	-	---	---	---	---	---	---	---	-
1984	117	1	-	---	---	---	---	---	---	---	-
1983	107	-	-	---	---	---	---	---	---	---	-
1982	108	-	1	---	---	---	---	---	---	---	-
1981	120	1	-	---	---	---	---	---	---	---	-
1980	127	1	-	---	---	---	---	---	---	---	-
1979	145	3	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy with abortive outcome (630-639)											
All races, both sexes											
1998	32	-	-	---	---	---	---	---	---	---	-
1997	41	-	-	---	---	---	---	---	---	---	-
1996	39	-	-	---	---	---	---	---	---	---	-
1995	28	-	-	---	---	---	---	---	---	---	-
1994	41	-	-	---	---	---	---	---	---	---	-
1993	47	-	-	---	---	---	---	---	---	---	-
1992	52	-	-	---	---	---	---	---	---	---	-
1991	54	1	-	---	---	---	---	---	---	---	-
1990	57	1	-	---	---	---	---	---	---	---	-
1989	50	-	-	---	---	---	---	---	---	---	-
1988	61	-	-	---	---	---	---	---	---	---	-
1987	44	-	-	---	---	---	---	---	---	---	-
1986	51	-	-	---	---	---	---	---	---	---	-
1985	49	-	-	---	---	---	---	---	---	---	-
1984	57	1	-	---	---	---	---	---	---	---	-
1983	50	-	-	---	---	---	---	---	---	---	-
1982	55	1	-	---	---	---	---	---	---	---	-
1981	47	-	-	---	---	---	---	---	---	---	-
1980	62	-	-	---	---	---	---	---	---	---	-
1979	61	-	-	---	---	---	---	---	---	---	-
Pregnancy with abortive outcome (630-639)											
All races, female											
1998	32	-	-	---	---	---	---	---	---	---	-
1997	41	-	-	---	---	---	---	---	---	---	-
1996	39	-	-	---	---	---	---	---	---	---	-
1995	28	-	-	---	---	---	---	---	---	---	-
1994	41	-	-	---	---	---	---	---	---	---	-
1993	47	-	-	---	---	---	---	---	---	---	-
1992	52	-	-	---	---	---	---	---	---	---	-
1991	54	1	-	---	---	---	---	---	---	---	-
1990	57	1	-	---	---	---	---	---	---	---	-
1989	50	-	-	---	---	---	---	---	---	---	-
1988	61	-	-	---	---	---	---	---	---	---	-
1987	44	-	-	---	---	---	---	---	---	---	-
1986	51	-	-	---	---	---	---	---	---	---	-
1985	49	-	-	---	---	---	---	---	---	---	-
1984	57	1	-	---	---	---	---	---	---	---	-
1983	50	-	-	---	---	---	---	---	---	---	-
1982	55	1	-	---	---	---	---	---	---	---	-
1981	47	-	-	---	---	---	---	---	---	---	-
1980	62	-	-	---	---	---	---	---	---	---	-
1979	61	-	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Pregnancy with abortive outcome (630-639)											
White, both sexes											
1998	15	-	-	---	---	---	---	---	---	---	-
1997	18	-	-	---	---	---	---	---	---	---	-
1996	24	-	-	---	---	---	---	---	---	---	-
1995	7	-	-	---	---	---	---	---	---	---	-
1994	24	-	-	---	---	---	---	---	---	---	-
1993	21	-	-	---	---	---	---	---	---	---	-
1992	25	-	-	---	---	---	---	---	---	---	-
1991	25	1	-	---	---	---	---	---	---	---	-
1990	23	-	-	---	---	---	---	---	---	---	-
1989	21	-	-	---	---	---	---	---	---	---	-
1988	23	-	-	---	---	---	---	---	---	---	-
1987	25	-	-	---	---	---	---	---	---	---	-
1986	24	-	-	---	---	---	---	---	---	---	-
1985	19	-	-	---	---	---	---	---	---	---	-
1984	25	1	-	---	---	---	---	---	---	---	-
1983	23	-	-	---	---	---	---	---	---	---	-
1982	27	1	-	---	---	---	---	---	---	---	-
1981	24	-	-	---	---	---	---	---	---	---	-
1980	27	-	-	---	---	---	---	---	---	---	-
1979	27	-	-	---	---	---	---	---	---	---	-
Pregnancy with abortive outcome (630-639)											
White, female											
1998	15	-	-	---	---	---	---	---	---	---	-
1997	18	-	-	---	---	---	---	---	---	---	-
1996	24	-	-	---	---	---	---	---	---	---	-
1995	7	-	-	---	---	---	---	---	---	---	-
1994	24	-	-	---	---	---	---	---	---	---	-
1993	21	-	-	---	---	---	---	---	---	---	-
1992	25	-	-	---	---	---	---	---	---	---	-
1991	25	1	-	---	---	---	---	---	---	---	-
1990	23	-	-	---	---	---	---	---	---	---	-
1989	21	-	-	---	---	---	---	---	---	---	-
1988	23	-	-	---	---	---	---	---	---	---	-
1987	25	-	-	---	---	---	---	---	---	---	-
1986	24	-	-	---	---	---	---	---	---	---	-
1985	19	-	-	---	---	---	---	---	---	---	-
1984	25	1	-	---	---	---	---	---	---	---	-
1983	23	-	-	---	---	---	---	---	---	---	-
1982	27	1	-	---	---	---	---	---	---	---	-
1981	24	-	-	---	---	---	---	---	---	---	-
1980	27	-	-	---	---	---	---	---	---	---	-
1979	27	-	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy with abortive outcome (630-639)											
All other, both sexes											
1998	17	-	-	---	---	---	---	---	---	---	-
1997	23	-	-	---	---	---	---	---	---	---	-
1996	15	-	-	---	---	---	---	---	---	---	-
1995	21	-	-	---	---	---	---	---	---	---	-
1994	17	-	-	---	---	---	---	---	---	---	-
1993	26	-	-	---	---	---	---	---	---	---	-
1992	27	-	-	---	---	---	---	---	---	---	-
1991	29	-	-	---	---	---	---	---	---	---	-
1990	34	1	-	---	---	---	---	---	---	---	-
1989	29	-	-	---	---	---	---	---	---	---	-
1988	38	-	-	---	---	---	---	---	---	---	-
1987	19	-	-	---	---	---	---	---	---	---	-
1986	27	-	-	---	---	---	---	---	---	---	-
1985	30	-	-	---	---	---	---	---	---	---	-
1984	32	-	-	---	---	---	---	---	---	---	-
1983	27	-	-	---	---	---	---	---	---	---	-
1982	28	-	-	---	---	---	---	---	---	---	-
1981	23	-	-	---	---	---	---	---	---	---	-
1980	35	-	-	---	---	---	---	---	---	---	-
1979	34	-	-	---	---	---	---	---	---	---	-
Pregnancy with abortive outcome (630-639)											
All other, female											
1998	17	-	-	---	---	---	---	---	---	---	-
1997	23	-	-	---	---	---	---	---	---	---	-
1996	15	-	-	---	---	---	---	---	---	---	-
1995	21	-	-	---	---	---	---	---	---	---	-
1994	17	-	-	---	---	---	---	---	---	---	-
1993	26	-	-	---	---	---	---	---	---	---	-
1992	27	-	-	---	---	---	---	---	---	---	-
1991	29	-	-	---	---	---	---	---	---	---	-
1990	34	1	-	---	---	---	---	---	---	---	-
1989	29	-	-	---	---	---	---	---	---	---	-
1988	38	-	-	---	---	---	---	---	---	---	-
1987	19	-	-	---	---	---	---	---	---	---	-
1986	27	-	-	---	---	---	---	---	---	---	-
1985	30	-	-	---	---	---	---	---	---	---	-
1984	32	-	-	---	---	---	---	---	---	---	-
1983	27	-	-	---	---	---	---	---	---	---	-
1982	28	-	-	---	---	---	---	---	---	---	-
1981	23	-	-	---	---	---	---	---	---	---	-
1980	35	-	-	---	---	---	---	---	---	---	-
1979	34	-	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Pregnancy with abortive outcome (630-639)											
Black, both sexes											
1998	14	-	-	---	---	---	---	---	---	---	-
1997	21	-	-	---	---	---	---	---	---	---	-
1996	14	-	-	---	---	---	---	---	---	---	-
1995	21	-	-	---	---	---	---	---	---	---	-
1994	17	-	-	---	---	---	---	---	---	---	-
1993	25	-	-	---	---	---	---	---	---	---	-
1992	26	-	-	---	---	---	---	---	---	---	-
1991	27	-	-	---	---	---	---	---	---	---	-
1990	33	1	-	---	---	---	---	---	---	---	-
1989	26	-	-	---	---	---	---	---	---	---	-
1988	35	-	-	---	---	---	---	---	---	---	-
1987	18	-	-	---	---	---	---	---	---	---	-
1986	26	-	-	---	---	---	---	---	---	---	-
1985	28	-	-	---	---	---	---	---	---	---	-
1984	30	-	-	---	---	---	---	---	---	---	-
1983	26	-	-	---	---	---	---	---	---	---	-
1982	23	-	-	---	---	---	---	---	---	---	-
1981	22	-	-	---	---	---	---	---	---	---	-
1980	31	-	-	---	---	---	---	---	---	---	-
1979	30	-	-	---	---	---	---	---	---	---	-
Pregnancy with abortive outcome (630-639)											
Black, female											
1998	14	-	-	---	---	---	---	---	---	---	-
1997	21	-	-	---	---	---	---	---	---	---	-
1996	14	-	-	---	---	---	---	---	---	---	-
1995	21	-	-	---	---	---	---	---	---	---	-
1994	17	-	-	---	---	---	---	---	---	---	-
1993	25	-	-	---	---	---	---	---	---	---	-
1992	26	-	-	---	---	---	---	---	---	---	-
1991	27	-	-	---	---	---	---	---	---	---	-
1990	33	1	-	---	---	---	---	---	---	---	-
1989	26	-	-	---	---	---	---	---	---	---	-
1988	35	-	-	---	---	---	---	---	---	---	-
1987	18	-	-	---	---	---	---	---	---	---	-
1986	26	-	-	---	---	---	---	---	---	---	-
1985	28	-	-	---	---	---	---	---	---	---	-
1984	30	-	-	---	---	---	---	---	---	---	-
1983	26	-	-	---	---	---	---	---	---	---	-
1982	23	-	-	---	---	---	---	---	---	---	-
1981	22	-	-	---	---	---	---	---	---	---	-
1980	31	-	-	---	---	---	---	---	---	---	-
1979	30	-	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other complications of pregnancy, childbirth and the puerperium (640-676)											
All races, both sexes											
1998	249	1	-	---	---	---	---	---	---	---	-
1997	286	4	1	---	---	---	---	---	---	---	-
1996	255	1	1	---	---	---	---	---	---	---	-
1995	249	1	-	---	---	---	---	---	---	---	-
1994	287	1	-	---	---	---	---	---	---	---	-
1993	255	-	-	---	---	---	---	---	---	---	-
1992	266	-	1	---	---	---	---	---	---	---	-
1991	269	4	-	---	---	---	---	---	---	---	-
1990	286	2	-	---	---	---	---	---	---	---	-
1989	270	2	-	---	---	---	---	---	---	---	-
1988	269	1	1	---	---	---	---	---	---	---	-
1987	207	1	-	---	---	---	---	---	---	---	-
1986	221	1	-	---	---	---	---	---	---	---	-
1985	246	2	1	---	---	---	---	---	---	---	-
1984	228	1	-	---	---	---	---	---	---	---	-
1983	240	-	-	---	---	---	---	---	---	---	-
1982	237	-	1	---	---	---	---	---	---	---	-
1981	262	1	-	---	---	---	---	---	---	---	-
1980	272	2	-	---	---	---	---	---	---	---	-
1979	275	5	-	---	---	---	---	---	---	---	-
Other complications of pregnancy, childbirth and the puerperium (640-676)											
All races, female											
1998	249	1	-	---	---	---	---	---	---	---	-
1997	286	4	1	---	---	---	---	---	---	---	-
1996	255	1	1	---	---	---	---	---	---	---	-
1995	249	1	-	---	---	---	---	---	---	---	-
1994	287	1	-	---	---	---	---	---	---	---	-
1993	255	-	-	---	---	---	---	---	---	---	-
1992	266	-	1	---	---	---	---	---	---	---	-
1991	269	4	-	---	---	---	---	---	---	---	-
1990	286	2	-	---	---	---	---	---	---	---	-
1989	270	2	-	---	---	---	---	---	---	---	-
1988	269	1	1	---	---	---	---	---	---	---	-
1987	207	1	-	---	---	---	---	---	---	---	-
1986	221	1	-	---	---	---	---	---	---	---	-
1985	246	2	1	---	---	---	---	---	---	---	-
1984	228	1	-	---	---	---	---	---	---	---	-
1983	240	-	-	---	---	---	---	---	---	---	-
1982	237	-	1	---	---	---	---	---	---	---	-
1981	262	1	-	---	---	---	---	---	---	---	-
1980	272	2	-	---	---	---	---	---	---	---	-
1979	275	5	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other complications of pregnancy, childbirth and the puerperium (640-676)											
White, both sexes											
1998	143	-	-	---	---	---	---	---	---	---	-
1997	161	4	-	---	---	---	---	---	---	---	-
1996	135	1	1	---	---	---	---	---	---	---	-
1995	122	-	-	---	---	---	---	---	---	---	-
1994	169	-	-	---	---	---	---	---	---	---	-
1993	131	-	-	---	---	---	---	---	---	---	-
1992	136	-	-	---	---	---	---	---	---	---	-
1991	162	2	-	---	---	---	---	---	---	---	-
1990	154	1	-	---	---	---	---	---	---	---	-
1989	159	-	-	---	---	---	---	---	---	---	-
1988	157	-	-	---	---	---	---	---	---	---	-
1987	128	1	-	---	---	---	---	---	---	---	-
1986	122	-	-	---	---	---	---	---	---	---	-
1985	137	-	1	---	---	---	---	---	---	---	-
1984	134	-	-	---	---	---	---	---	---	---	-
1983	147	-	-	---	---	---	---	---	---	---	-
1982	144	-	-	---	---	---	---	---	---	---	-
1981	160	-	-	---	---	---	---	---	---	---	-
1980	166	1	-	---	---	---	---	---	---	---	-
1979	153	2	-	---	---	---	---	---	---	---	-
Other complications of pregnancy, childbirth and the puerperium (640-676)											
White, female											
1998	143	-	-	---	---	---	---	---	---	---	-
1997	161	4	-	---	---	---	---	---	---	---	-
1996	135	1	1	---	---	---	---	---	---	---	-
1995	122	-	-	---	---	---	---	---	---	---	-
1994	169	-	-	---	---	---	---	---	---	---	-
1993	131	-	-	---	---	---	---	---	---	---	-
1992	136	-	-	---	---	---	---	---	---	---	-
1991	162	2	-	---	---	---	---	---	---	---	-
1990	154	1	-	---	---	---	---	---	---	---	-
1989	159	-	-	---	---	---	---	---	---	---	-
1988	157	-	-	---	---	---	---	---	---	---	-
1987	128	1	-	---	---	---	---	---	---	---	-
1986	122	-	-	---	---	---	---	---	---	---	-
1985	137	-	1	---	---	---	---	---	---	---	-
1984	134	-	-	---	---	---	---	---	---	---	-
1983	147	-	-	---	---	---	---	---	---	---	-
1982	144	-	-	---	---	---	---	---	---	---	-
1981	160	-	-	---	---	---	---	---	---	---	-
1980	166	1	-	---	---	---	---	---	---	---	-
1979	153	2	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other complications of pregnancy, childbirth and the puerperium (640-676)											
All other, both sexes											
1998	106	1	-	---	---	---	---	---	---	---	-
1997	125	-	1	---	---	---	---	---	---	---	-
1996	120	-	-	---	---	---	---	---	---	---	-
1995	127	1	-	---	---	---	---	---	---	---	-
1994	118	1	-	---	---	---	---	---	---	---	-
1993	124	-	-	---	---	---	---	---	---	---	-
1992	130	-	1	---	---	---	---	---	---	---	-
1991	107	2	-	---	---	---	---	---	---	---	-
1990	132	1	-	---	---	---	---	---	---	---	-
1989	111	2	-	---	---	---	---	---	---	---	-
1988	112	1	1	---	---	---	---	---	---	---	-
1987	79	-	-	---	---	---	---	---	---	---	-
1986	99	1	-	---	---	---	---	---	---	---	-
1985	109	2	-	---	---	---	---	---	---	---	-
1984	94	1	-	---	---	---	---	---	---	---	-
1983	93	-	-	---	---	---	---	---	---	---	-
1982	93	-	1	---	---	---	---	---	---	---	-
1981	102	1	-	---	---	---	---	---	---	---	-
1980	106	1	-	---	---	---	---	---	---	---	-
1979	122	3	-	---	---	---	---	---	---	---	-
Other complications of pregnancy, childbirth and the puerperium (640-676)											
All other, female											
1998	106	1	-	---	---	---	---	---	---	---	-
1997	125	-	1	---	---	---	---	---	---	---	-
1996	120	-	-	---	---	---	---	---	---	---	-
1995	127	1	-	---	---	---	---	---	---	---	-
1994	118	1	-	---	---	---	---	---	---	---	-
1993	124	-	-	---	---	---	---	---	---	---	-
1992	130	-	1	---	---	---	---	---	---	---	-
1991	107	2	-	---	---	---	---	---	---	---	-
1990	132	1	-	---	---	---	---	---	---	---	-
1989	111	2	-	---	---	---	---	---	---	---	-
1988	112	1	1	---	---	---	---	---	---	---	-
1987	79	-	-	---	---	---	---	---	---	---	-
1986	99	1	-	---	---	---	---	---	---	---	-
1985	109	2	-	---	---	---	---	---	---	---	-
1984	94	1	-	---	---	---	---	---	---	---	-
1983	93	-	-	---	---	---	---	---	---	---	-
1982	93	-	1	---	---	---	---	---	---	---	-
1981	102	1	-	---	---	---	---	---	---	---	-
1980	106	1	-	---	---	---	---	---	---	---	-
1979	122	3	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other complications of pregnancy, childbirth and the puerperium (640-676)											
Black, both sexes											
1998	90	1	-	---	---	---	---	---	---	---	-
1997	104	-	1	---	---	---	---	---	---	---	-
1996	107	-	-	---	---	---	---	---	---	---	-
1995	112	1	-	---	---	---	---	---	---	---	-
1994	101	1	-	---	---	---	---	---	---	---	-
1993	110	-	-	---	---	---	---	---	---	---	-
1992	114	-	1	---	---	---	---	---	---	---	-
1991	98	2	-	---	---	---	---	---	---	---	-
1990	120	1	-	---	---	---	---	---	---	---	-
1989	98	2	-	---	---	---	---	---	---	---	-
1988	96	1	1	---	---	---	---	---	---	---	-
1987	73	-	-	---	---	---	---	---	---	---	-
1986	91	1	-	---	---	---	---	---	---	---	-
1985	96	1	-	---	---	---	---	---	---	---	-
1984	87	1	-	---	---	---	---	---	---	---	-
1983	81	-	-	---	---	---	---	---	---	---	-
1982	85	-	1	---	---	---	---	---	---	---	-
1981	98	1	-	---	---	---	---	---	---	---	-
1980	96	1	-	---	---	---	---	---	---	---	-
1979	115	3	-	---	---	---	---	---	---	---	-
Other complications of pregnancy, childbirth and the puerperium (640-676)											
Black, female											
1998	90	1	-	---	---	---	---	---	---	---	-
1997	104	-	1	---	---	---	---	---	---	---	-
1996	107	-	-	---	---	---	---	---	---	---	-
1995	112	1	-	---	---	---	---	---	---	---	-
1994	101	1	-	---	---	---	---	---	---	---	-
1993	110	-	-	---	---	---	---	---	---	---	-
1992	114	-	1	---	---	---	---	---	---	---	-
1991	98	2	-	---	---	---	---	---	---	---	-
1990	120	1	-	---	---	---	---	---	---	---	-
1989	98	2	-	---	---	---	---	---	---	---	-
1988	96	1	1	---	---	---	---	---	---	---	-
1987	73	-	-	---	---	---	---	---	---	---	-
1986	91	1	-	---	---	---	---	---	---	---	-
1985	96	1	-	---	---	---	---	---	---	---	-
1984	87	1	-	---	---	---	---	---	---	---	-
1983	81	-	-	---	---	---	---	---	---	---	-
1982	85	-	1	---	---	---	---	---	---	---	-
1981	98	1	-	---	---	---	---	---	---	---	-
1980	96	1	-	---	---	---	---	---	---	---	-
1979	115	3	-	---	---	---	---	---	---	---	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All races, both sexes											
1998	13,428	1	1	3	1	-	1	1	-	2	1
1997	13,092	2	2	-	1	3	3	-	-	1	2
1996	13,069	2	2	-	-	-	1	-	-	-	6
1995	13,471	1	1	-	1	1	1	-	1	2	1
1994	14,487	2	4	-	1	1	1	-	-	2	-
1993	15,096	1	-	-	-	2	1	2	2	-	4
1992	15,730	1	1	-	-	1	2	1	-	2	2
1991	16,781	1	1	2	-	-	-	-	-	1	-
1990	17,674	1	1	-	1	-	1	-	1	4	7
1989	18,752	1	1	-	2	2	1	1	-	-	4
1988	18,220	2	1	-	-	-	-	1	-	1	8
1987	18,222	-	-	1	-	-	1	1	-	-	2
1986	18,391	2	-	-	-	1	-	1	-	1	1
1985	19,246	-	1	1	-	-	-	1	-	-	3
1984	18,880	1	2	1	1	-	-	-	-	-	8
1983	19,310	1	2	-	1	1	1	-	1	1	4
1982	20,794	-	2	1	-	1	1	-	-	-	4
1981	21,626	-	-	2	1	-	-	-	-	2	9
1980	22,866	1	-	3	1	-	1	-	-	1	4
1979	23,447	2	-	2	1	1	-	-	1	1	20
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All races, male											
1998	7,496	1	1	2	1	-	1	1	-	-	-
1997	7,398	-	1	-	1	-	2	-	-	1	1
1996	7,303	1	1	-	-	-	-	-	-	-	2
1995	7,600	1	-	-	-	1	1	-	-	1	-
1994	8,156	1	2	-	-	1	-	-	-	1	-
1993	8,472	1	-	-	-	2	1	2	1	-	2
1992	8,817	-	1	-	-	-	-	1	-	1	2
1991	9,561	-	1	2	-	-	-	-	-	-	-
1990	10,056	1	1	-	-	-	-	-	1	2	5
1989	10,511	-	1	-	2	1	-	1	-	-	2
1988	10,393	1	-	-	-	-	-	1	-	1	5
1987	10,442	-	-	-	-	-	-	1	-	-	1
1986	10,563	1	-	-	-	-	-	1	-	-	-
1985	11,094	-	-	-	-	-	-	1	-	-	3
1984	10,658	-	1	1	-	-	-	-	-	-	4
1983	11,036	1	-	-	-	-	1	-	-	-	4
1982	11,876	-	2	-	-	-	-	-	-	-	4
1981	12,302	-	-	-	1	-	-	-	-	2	6
1980	13,051	1	-	1	1	-	1	-	-	1	1
1979	13,349	1	-	1	1	-	-	-	-	-	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All races, female											
1998	5,932	-	-	1	-	-	-	-	-	2	1
1997	5,694	2	1	-	-	3	1	-	-	-	1
1996	5,766	1	1	-	-	-	1	-	-	-	4
1995	5,871	-	1	-	1	-	-	-	1	1	1
1994	6,331	1	2	-	1	-	1	-	-	1	-
1993	6,624	-	-	-	-	-	-	-	1	-	2
1992	6,913	1	-	-	-	1	2	-	-	1	-
1991	7,220	1	-	-	-	-	-	-	-	1	-
1990	7,618	-	-	-	1	-	1	-	-	2	2
1989	8,241	1	-	-	-	1	1	-	-	-	2
1988	7,827	1	1	-	-	-	-	-	-	-	3
1987	7,780	-	-	1	-	-	1	-	-	-	1
1986	7,828	1	-	-	-	1	-	-	-	1	1
1985	8,152	-	1	1	-	-	-	-	-	-	-
1984	8,222	1	1	-	1	-	-	-	-	-	4
1983	8,274	-	2	-	1	1	-	-	1	1	-
1982	8,918	-	-	1	-	1	1	-	-	-	-
1981	9,324	-	-	2	-	-	-	-	-	-	3
1980	9,815	-	-	2	-	-	-	-	-	-	3
1979	10,098	1	-	1	-	1	-	-	1	1	10
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
White, both sexes											
1998	8,129	1	1	2	1	-	1	-	-	2	1
1997	7,981	1	2	-	1	3	3	-	-	1	2
1996	7,964	1	2	-	-	-	1	-	-	-	4
1995	8,155	1	1	-	1	1	1	-	1	2	-
1994	8,562	2	4	-	1	1	1	-	-	1	-
1993	8,830	1	-	-	-	2	-	2	1	-	2
1992	9,168	1	1	-	-	1	2	-	-	1	2
1991	9,889	1	1	2	-	-	-	-	-	1	-
1990	10,506	1	1	-	1	-	1	-	1	3	6
1989	11,306	1	1	-	2	2	1	-	-	-	2
1988	11,192	2	-	-	-	-	-	1	-	1	5
1987	11,412	-	-	-	-	-	1	1	-	-	2
1986	11,814	2	-	-	-	1	-	-	-	1	1
1985	12,634	-	1	1	-	-	-	-	-	-	1
1984	12,470	1	2	1	1	-	-	-	-	-	7
1983	12,816	1	1	-	-	1	1	-	1	-	1
1982	13,932	-	1	1	-	1	1	-	-	-	1
1981	14,551	-	-	2	1	-	-	-	-	-	6
1980	15,457	1	-	3	1	-	1	-	-	1	1
1979	16,033	2	-	2	1	1	-	-	1	1	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
White, male											
1998	4,523	1	1	1	1	-	1	-	-	-	-
1997	4,538	-	1	-	1	-	2	-	-	1	1
1996	4,446	-	1	-	-	-	-	-	-	-	2
1995	4,676	1	-	-	-	1	1	-	-	1	-
1994	4,829	1	2	-	-	1	-	-	-	-	-
1993	4,956	1	-	-	-	2	-	2	1	-	1
1992	5,128	-	1	-	-	-	-	-	-	-	2
1991	5,654	-	1	2	-	-	-	-	-	-	-
1990	6,055	1	1	-	-	-	-	-	1	2	4
1989	6,426	-	1	-	2	1	-	-	-	-	1
1988	6,485	1	-	-	-	-	-	1	-	1	3
1987	6,560	-	-	-	-	-	-	1	-	-	1
1986	6,815	1	-	-	-	-	-	-	-	-	-
1985	7,404	-	-	-	-	-	-	-	-	-	1
1984	7,134	-	1	1	-	-	-	-	-	-	4
1983	7,417	1	-	-	-	-	1	-	-	-	1
1982	8,003	-	1	-	-	-	-	-	-	-	1
1981	8,380	-	-	-	1	-	-	-	-	-	5
1980	8,945	1	-	1	1	-	1	-	-	1	-
1979	9,261	1	-	1	1	-	-	-	-	-	7
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
White, female											
1998	3,606	-	-	1	-	-	-	-	-	2	1
1997	3,443	1	1	-	-	3	1	-	-	-	1
1996	3,518	1	1	-	-	-	1	-	-	-	2
1995	3,479	-	1	-	1	-	-	-	1	1	-
1994	3,733	1	2	-	1	-	1	-	-	1	-
1993	3,874	-	-	-	-	-	-	-	-	-	1
1992	4,040	1	-	-	-	1	2	-	-	1	-
1991	4,235	1	-	-	-	-	-	-	-	1	-
1990	4,451	-	-	-	1	-	1	-	-	1	2
1989	4,880	1	-	-	-	1	1	-	-	-	1
1988	4,707	1	-	-	-	-	-	-	-	-	2
1987	4,852	-	-	-	-	-	1	-	-	-	1
1986	4,999	1	-	-	-	1	-	-	-	1	1
1985	5,230	-	1	1	-	-	-	-	-	-	-
1984	5,336	1	1	-	1	-	-	-	-	-	3
1983	5,399	-	1	-	-	1	-	-	1	-	-
1982	5,929	-	-	1	-	1	1	-	-	-	-
1981	6,171	-	-	2	-	-	-	-	-	-	1
1980	6,512	-	-	2	-	-	-	-	-	-	1
1979	6,772	1	-	1	-	1	-	-	1	1	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All other, both sexes											
1998	5,299	-	-	1	-	-	-	1	-	-	-
1997	5,111	1	-	-	-	-	-	-	-	-	-
1996	5,105	1	-	-	-	-	-	-	-	-	2
1995	5,316	-	-	-	-	-	-	-	-	-	1
1994	5,925	-	-	-	-	-	-	-	-	1	-
1993	6,266	-	-	-	-	-	1	-	1	-	2
1992	6,562	-	-	-	-	-	-	1	-	1	-
1991	6,892	-	-	-	-	-	-	-	-	-	-
1990	7,168	-	-	-	-	-	-	-	-	1	1
1989	7,446	-	-	-	-	-	-	1	-	-	2
1988	7,028	-	1	-	-	-	-	-	-	-	3
1987	6,810	-	-	1	-	-	-	-	-	-	-
1986	6,577	-	-	-	-	-	-	1	-	-	-
1985	6,612	-	-	-	-	-	-	1	-	-	2
1984	6,410	-	-	-	-	-	-	-	-	-	1
1983	6,494	-	1	-	1	-	-	-	-	1	3
1982	6,862	-	1	-	-	-	-	-	-	-	3
1981	7,075	-	-	-	-	-	-	-	-	2	3
1980	7,409	-	-	-	-	-	-	-	-	-	3
1979	7,414	-	-	-	-	-	-	-	-	-	5
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All other, male											
1998	2,973	-	-	1	-	-	-	1	-	-	-
1997	2,860	-	-	-	-	-	-	-	-	-	-
1996	2,857	1	-	-	-	-	-	-	-	-	-
1995	2,924	-	-	-	-	-	-	-	-	-	-
1994	3,327	-	-	-	-	-	-	-	-	1	-
1993	3,516	-	-	-	-	-	1	-	-	-	1
1992	3,689	-	-	-	-	-	-	1	-	1	-
1991	3,907	-	-	-	-	-	-	-	-	-	-
1990	4,001	-	-	-	-	-	-	-	-	-	1
1989	4,085	-	-	-	-	-	-	1	-	-	1
1988	3,908	-	-	-	-	-	-	-	-	-	2
1987	3,882	-	-	-	-	-	-	-	-	-	-
1986	3,748	-	-	-	-	-	-	1	-	-	-
1985	3,690	-	-	-	-	-	-	1	-	-	2
1984	3,524	-	-	-	-	-	-	-	-	-	-
1983	3,619	-	-	-	-	-	-	-	-	-	3
1982	3,873	-	1	-	-	-	-	-	-	-	3
1981	3,922	-	-	-	-	-	-	-	-	2	1
1980	4,106	-	-	-	-	-	-	-	-	-	1
1979	4,088	-	-	-	-	-	-	-	-	-	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
All other, female											
1998	2,326	-	-	-	-	-	-	-	-	-	-
1997	2,251	1	-	-	-	-	-	-	-	-	-
1996	2,248	-	-	-	-	-	-	-	-	-	2
1995	2,392	-	-	-	-	-	-	-	-	-	1
1994	2,598	-	-	-	-	-	-	-	-	-	-
1993	2,750	-	-	-	-	-	-	-	1	-	1
1992	2,873	-	-	-	-	-	-	-	-	-	-
1991	2,985	-	-	-	-	-	-	-	-	-	-
1990	3,167	-	-	-	-	-	-	-	-	1	-
1989	3,361	-	-	-	-	-	-	-	-	-	1
1988	3,120	-	1	-	-	-	-	-	-	-	1
1987	2,928	-	-	1	-	-	-	-	-	-	-
1986	2,829	-	-	-	-	-	-	-	-	-	-
1985	2,922	-	-	-	-	-	-	-	-	-	-
1984	2,886	-	-	-	-	-	-	-	-	-	1
1983	2,875	-	1	-	1	-	-	-	-	1	-
1982	2,989	-	-	-	-	-	-	-	-	-	-
1981	3,153	-	-	-	-	-	-	-	-	-	2
1980	3,303	-	-	-	-	-	-	-	-	-	2
1979	3,326	-	-	-	-	-	-	-	-	-	2
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
Black, both sexes											
1998	4,841	-	-	1	-	-	-	1	-	-	-
1997	4,714	1	-	-	-	-	-	-	-	-	-
1996	4,711	1	-	-	-	-	-	-	-	-	2
1995	4,952	-	-	-	-	-	-	-	-	-	-
1994	5,515	-	-	-	-	-	-	-	-	1	-
1993	5,891	-	-	-	-	-	1	-	1	-	2
1992	6,176	-	-	-	-	-	-	1	-	1	-
1991	6,512	-	-	-	-	-	-	-	-	-	-
1990	6,744	-	-	-	-	-	-	-	-	1	-
1989	6,965	-	-	-	-	-	-	1	-	-	2
1988	6,614	-	1	-	-	-	-	-	-	-	3
1987	6,373	-	-	-	-	-	-	-	-	-	-
1986	6,149	-	-	-	-	-	-	1	-	-	-
1985	6,196	-	-	-	-	-	-	1	-	-	2
1984	6,002	-	-	-	-	-	-	-	-	-	1
1983	6,100	-	1	-	1	-	-	-	-	1	3
1982	6,464	-	1	-	-	-	-	-	-	-	1
1981	6,658	-	-	-	-	-	-	-	-	2	2
1980	6,961	-	-	-	-	-	-	-	-	-	2
1979	7,024	-	-	-	-	-	-	-	-	-	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
Black, male											
1998	2,732	-	-	1	-	-	-	1	-	-	-
1997	2,624	-	-	-	-	-	-	-	-	-	-
1996	2,631	1	-	-	-	-	-	-	-	-	-
1995	2,731	-	-	-	-	-	-	-	-	-	-
1994	3,088	-	-	-	-	-	-	-	-	1	-
1993	3,309	-	-	-	-	-	1	-	-	-	1
1992	3,469	-	-	-	-	-	-	1	-	1	-
1991	3,711	-	-	-	-	-	-	-	-	-	-
1990	3,762	-	-	-	-	-	-	-	-	-	-
1989	3,813	-	-	-	-	-	-	1	-	-	1
1988	3,693	-	-	-	-	-	-	-	-	-	2
1987	3,608	-	-	-	-	-	-	-	-	-	-
1986	3,501	-	-	-	-	-	-	1	-	-	-
1985	3,458	-	-	-	-	-	-	1	-	-	2
1984	3,290	-	-	-	-	-	-	-	-	-	-
1983	3,401	-	-	-	-	-	-	-	-	-	3
1982	3,633	-	1	-	-	-	-	-	-	-	1
1981	3,688	-	-	-	-	-	-	-	-	2	-
1980	3,869	-	-	-	-	-	-	-	-	-	1
1979	3,871	-	-	-	-	-	-	-	-	-	3
Certain conditions originating in the perinatal period (760-771.2,771.4-779)											
Black, female											
1998	2,109	-	-	-	-	-	-	-	-	-	-
1997	2,090	1	-	-	-	-	-	-	-	-	-
1996	2,080	-	-	-	-	-	-	-	-	-	2
1995	2,221	-	-	-	-	-	-	-	-	-	-
1994	2,427	-	-	-	-	-	-	-	-	-	-
1993	2,582	-	-	-	-	-	-	-	1	-	1
1992	2,707	-	-	-	-	-	-	-	-	-	-
1991	2,801	-	-	-	-	-	-	-	-	-	-
1990	2,982	-	-	-	-	-	-	-	-	1	-
1989	3,152	-	-	-	-	-	-	-	-	-	1
1988	2,921	-	1	-	-	-	-	-	-	-	1
1987	2,765	-	-	-	-	-	-	-	-	-	-
1986	2,648	-	-	-	-	-	-	-	-	-	-
1985	2,738	-	-	-	-	-	-	-	-	-	-
1984	2,712	-	-	-	-	-	-	-	-	-	1
1983	2,699	-	1	-	1	-	-	-	-	1	-
1982	2,831	-	-	-	-	-	-	-	-	-	-
1981	2,970	-	-	-	-	-	-	-	-	-	2
1980	3,092	-	-	-	-	-	-	-	-	-	1
1979	3,153	-	-	-	-	-	-	-	-	-	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All races, both sexes											
1998	11,934	280	245	248	271	288	363	498	500	646	1
1997	11,912	274	264	240	261	308	346	470	482	643	2
1996	11,836	286	241	248	232	286	386	408	405	559	1
1995	11,891	246	241	223	242	278	343	353	368	446	-
1994	12,030	256	230	216	242	288	286	348	336	412	-
1993	12,444	259	199	215	267	267	321	341	343	396	1
1992	12,491	206	181	220	249	252	320	295	304	316	-
1991	12,599	206	194	191	217	283	267	287	247	292	4
1990	13,085	190	174	194	218	267	280	243	235	259	4
1989	12,897	193	202	204	224	275	238	268	182	218	6
1988	12,753	160	176	181	252	242	248	228	193	165	5
1987	12,333	174	176	208	201	240	206	210	173	154	1
1986	12,638	145	170	199	247	244	235	182	183	163	2
1985	12,783	160	153	217	217	229	200	207	134	151	6
1984	13,039	180	170	228	269	179	247	191	132	152	4
1983	13,173	154	175	202	234	226	210	196	141	151	4
1982	13,604	153	180	216	220	243	196	185	145	110	7
1981	13,527	152	187	223	208	225	195	188	131	119	11
1980	13,938	188	202	238	247	229	216	173	115	103	3
1979	13,526	193	217	228	223	220	192	145	113	115	8
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All races, male											
1998	6,199	162	124	134	131	156	186	253	239	256	-
1997	6,261	154	139	126	149	153	190	243	241	262	2
1996	6,238	155	127	114	126	134	189	208	201	207	1
1995	6,243	138	127	116	112	127	174	174	158	156	-
1994	6,209	145	126	109	118	152	146	172	152	149	-
1993	6,610	141	101	112	129	148	170	167	141	158	1
1992	6,625	107	92	102	120	118	157	136	141	119	-
1991	6,800	112	116	98	113	150	140	142	106	115	4
1990	7,050	97	88	97	108	132	120	123	110	102	2
1989	6,812	92	100	114	114	137	109	118	82	91	4
1988	6,775	84	92	97	113	114	100	111	73	58	1
1987	6,604	90	99	103	105	127	83	86	83	57	-
1986	6,744	87	92	89	108	126	110	69	80	59	1
1985	6,895	90	72	119	115	106	77	90	57	59	-
1984	6,925	97	80	108	130	95	127	85	59	55	2
1983	7,013	85	95	97	107	111	96	81	63	64	3
1982	7,157	75	86	123	106	126	85	77	62	41	1
1981	7,128	84	88	107	99	101	90	73	50	47	7
1980	7,496	105	112	126	132	119	104	74	63	28	-
1979	7,309	96	102	90	102	112	102	62	44	40	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All races, female											
1998	5,735	118	121	114	140	132	177	245	261	390	1
1997	5,651	120	125	114	112	155	156	227	241	381	-
1996	5,598	131	114	134	106	152	197	200	204	352	-
1995	5,648	108	114	107	130	151	169	179	210	290	-
1994	5,821	111	104	107	124	136	140	176	184	263	-
1993	5,834	118	98	103	138	119	151	174	202	238	-
1992	5,866	99	89	118	129	134	163	159	163	197	-
1991	5,799	94	78	93	104	133	127	145	141	177	-
1990	6,035	93	86	97	110	135	160	120	125	157	2
1989	6,085	101	102	90	110	138	129	150	100	127	2
1988	5,978	76	84	84	139	128	148	117	120	107	4
1987	5,729	84	77	105	96	113	123	124	90	97	1
1986	5,894	58	78	110	139	118	125	113	103	104	1
1985	5,888	70	81	98	102	123	123	117	77	92	6
1984	6,114	83	90	120	139	84	120	106	73	97	2
1983	6,160	69	80	105	127	115	114	115	78	87	1
1982	6,447	78	94	93	114	117	111	108	83	69	6
1981	6,399	68	99	116	109	124	105	115	81	72	4
1980	6,442	83	90	112	115	110	112	99	52	75	3
1979	6,217	97	115	138	121	108	90	83	69	75	3
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
White, both sexes											
1998	9,651	243	224	220	235	257	330	466	463	614	1
1997	9,719	237	232	217	230	280	309	440	452	606	2
1996	9,556	238	215	216	211	258	350	373	375	526	1
1995	9,614	216	210	196	217	249	307	329	341	423	-
1994	9,660	219	194	187	212	262	258	322	318	384	-
1993	9,934	220	169	186	239	242	276	316	314	376	1
1992	9,890	183	157	186	223	229	292	272	278	298	-
1991	9,978	173	160	163	198	255	248	257	228	283	3
1990	10,502	172	151	174	199	232	254	222	215	240	3
1989	10,291	171	169	168	202	249	215	238	171	204	3
1988	10,334	137	154	161	219	218	225	218	176	152	4
1987	9,910	155	153	182	178	215	187	195	164	145	1
1986	10,188	130	138	170	226	221	215	168	173	150	2
1985	10,437	134	129	185	196	213	182	189	123	134	6
1984	10,662	150	146	199	241	158	225	173	120	140	2
1983	10,747	127	148	184	203	206	191	177	130	139	3
1982	11,129	128	157	193	198	220	182	170	140	99	7
1981	11,145	132	166	193	190	200	175	166	127	108	9
1980	11,471	171	178	220	223	209	197	153	110	100	3
1979	11,222	163	196	203	199	203	178	133	109	108	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
White, male											
1998	5,001	138	112	116	115	139	168	237	224	246	-
1997	5,123	135	125	112	135	137	170	226	229	247	2
1996	5,066	128	113	99	118	122	168	189	189	198	1
1995	5,047	120	109	98	101	115	160	161	145	150	-
1994	4,993	117	107	98	105	140	132	163	147	140	-
1993	5,281	118	84	100	113	137	149	157	129	149	1
1992	5,254	98	77	91	105	111	145	125	129	112	-
1991	5,378	93	98	86	101	139	130	127	97	110	3
1990	5,667	89	75	91	98	111	110	110	101	94	2
1989	5,444	82	88	95	104	123	100	101	78	83	3
1988	5,491	75	84	83	97	102	94	107	67	52	1
1987	5,364	79	88	94	94	117	75	79	78	54	-
1986	5,444	76	78	73	100	112	99	63	76	54	1
1985	5,674	81	63	102	103	98	71	82	51	54	-
1984	5,703	86	68	98	121	81	114	75	54	50	-
1983	5,718	69	82	89	95	100	87	72	55	55	3
1982	5,883	59	79	110	95	114	80	70	58	36	1
1981	5,906	71	78	90	88	86	82	65	49	44	5
1980	6,188	94	100	117	117	106	96	65	60	27	-
1979	6,106	79	94	78	94	103	94	57	43	37	4
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
White, female											
1998	4,650	105	112	104	120	118	162	229	239	368	1
1997	4,596	102	107	105	95	143	139	214	223	359	-
1996	4,490	110	102	117	93	136	182	184	186	328	-
1995	4,567	96	101	98	116	134	147	168	196	273	-
1994	4,667	102	87	89	107	122	126	159	171	244	-
1993	4,653	102	85	86	126	105	127	159	185	227	-
1992	4,636	85	80	95	118	118	147	147	149	186	-
1991	4,600	80	62	77	97	116	118	130	131	173	-
1990	4,835	83	76	83	101	121	144	112	114	146	1
1989	4,847	89	81	73	98	126	115	137	93	121	-
1988	4,843	62	70	78	122	116	131	111	109	100	3
1987	4,546	76	65	88	84	98	112	116	86	91	1
1986	4,744	54	60	97	126	109	116	105	97	96	1
1985	4,763	53	66	83	93	115	111	107	72	80	6
1984	4,959	64	78	101	120	77	111	98	66	90	2
1983	5,029	58	66	95	108	106	104	105	75	84	-
1982	5,246	69	78	83	103	106	102	100	82	63	6
1981	5,239	61	88	103	102	114	93	101	78	64	4
1980	5,283	77	78	103	106	103	101	88	50	73	3
1979	5,116	84	102	125	105	100	84	76	66	71	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All other, both sexes											
1998	2,283	37	21	28	36	31	33	32	37	32	-
1997	2,193	37	32	23	31	28	37	30	30	37	-
1996	2,280	48	26	32	21	28	36	35	30	33	-
1995	2,277	30	31	27	25	29	36	24	27	23	-
1994	2,370	37	36	29	30	26	28	26	18	28	-
1993	2,510	39	30	29	28	25	45	25	29	20	-
1992	2,601	23	24	34	26	23	28	23	26	18	-
1991	2,621	33	34	28	19	28	19	30	19	9	1
1990	2,583	18	23	20	19	35	26	21	20	19	1
1989	2,606	22	33	36	22	26	23	30	11	14	3
1988	2,419	23	22	20	33	24	23	10	17	13	1
1987	2,423	19	23	26	23	25	19	15	9	9	-
1986	2,450	15	32	29	21	23	20	14	10	13	-
1985	2,346	26	24	32	21	16	18	18	11	17	-
1984	2,377	30	24	29	28	21	22	18	12	12	2
1983	2,426	27	27	18	31	20	19	19	11	12	1
1982	2,475	25	23	23	22	23	14	15	5	11	-
1981	2,382	20	21	30	18	25	20	22	4	11	2
1980	2,467	17	24	18	24	20	19	20	5	3	-
1979	2,304	30	21	25	24	17	14	12	4	7	4
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All other, male											
1998	1,198	24	12	18	16	17	18	16	15	10	-
1997	1,138	19	14	14	14	16	20	17	12	15	-
1996	1,172	27	14	15	8	12	21	19	12	9	-
1995	1,196	18	18	18	11	12	14	13	13	6	-
1994	1,216	28	19	11	13	12	14	9	5	9	-
1993	1,329	23	17	12	16	11	21	10	12	9	-
1992	1,371	9	15	11	15	7	12	11	12	7	-
1991	1,422	19	18	12	12	11	10	15	9	5	1
1990	1,383	8	13	6	10	21	10	13	9	8	-
1989	1,368	10	12	19	10	14	9	17	4	8	1
1988	1,284	9	8	14	16	12	6	4	6	6	-
1987	1,240	11	11	9	11	10	8	7	5	3	-
1986	1,300	11	14	16	8	14	11	6	4	5	-
1985	1,221	9	9	17	12	8	6	8	6	5	-
1984	1,222	11	12	10	9	14	13	10	5	5	2
1983	1,295	16	13	8	12	11	9	9	8	9	-
1982	1,274	16	7	13	11	12	5	7	4	5	-
1981	1,222	13	10	17	11	15	8	8	1	3	2
1980	1,308	11	12	9	15	13	8	9	3	1	-
1979	1,203	17	8	12	8	9	8	5	1	3	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
All other, female											
1998	1,085	13	9	10	20	14	15	16	22	22	-
1997	1,055	18	18	9	17	12	17	13	18	22	-
1996	1,108	21	12	17	13	16	15	16	18	24	-
1995	1,081	12	13	9	14	17	22	11	14	17	-
1994	1,154	9	17	18	17	14	14	17	13	19	-
1993	1,181	16	13	17	12	14	24	15	17	11	-
1992	1,230	14	9	23	11	16	16	12	14	11	-
1991	1,199	14	16	16	7	17	9	15	10	4	-
1990	1,200	10	10	14	9	14	16	8	11	11	1
1989	1,238	12	21	17	12	12	14	13	7	6	2
1988	1,135	14	14	6	17	12	17	6	11	7	1
1987	1,183	8	12	17	12	15	11	8	4	6	-
1986	1,150	4	18	13	13	9	9	8	6	8	-
1985	1,125	17	15	15	9	8	12	10	5	12	-
1984	1,155	19	12	19	19	7	9	8	7	7	-
1983	1,131	11	14	10	19	9	10	10	3	3	1
1982	1,201	9	16	10	11	11	9	8	1	6	-
1981	1,160	7	11	13	7	10	12	14	3	8	-
1980	1,159	6	12	9	9	7	11	11	2	2	-
1979	1,101	13	13	13	16	8	6	7	3	4	3
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
Black, both sexes											
1998	1,885	32	18	26	30	24	28	26	34	26	-
1997	1,797	33	28	18	24	21	31	27	28	30	-
1996	1,905	42	25	27	21	24	28	29	24	28	-
1995	1,887	26	26	24	20	23	31	20	24	18	-
1994	1,961	30	33	24	23	18	21	21	15	24	-
1993	2,104	36	28	24	22	22	34	20	24	19	-
1992	2,167	19	17	26	22	20	24	20	21	16	-
1991	2,202	28	27	23	17	27	19	29	16	7	1
1990	2,177	17	17	16	15	32	22	18	15	14	1
1989	2,179	19	29	29	18	24	21	25	10	11	3
1988	2,019	18	18	18	31	19	21	7	13	11	1
1987	2,076	17	20	22	20	23	16	13	9	6	-
1986	2,061	13	26	25	18	22	15	10	9	10	-
1985	2,008	24	21	27	17	14	16	15	8	15	-
1984	1,983	24	23	23	25	18	21	17	9	11	1
1983	2,088	24	25	17	30	17	18	18	10	10	-
1982	2,141	23	19	20	19	23	12	14	5	9	-
1981	2,064	16	20	26	17	22	20	20	4	10	2
1980	2,168	17	22	17	20	18	18	19	5	2	-
1979	2,040	30	20	24	21	15	13	11	2	5	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
Black, male											
1998	994	20	10	18	14	14	15	12	15	8	-
1997	921	16	11	12	11	11	18	15	11	12	-
1996	971	24	14	13	8	11	16	16	9	6	-
1995	996	15	16	16	10	9	11	13	11	4	-
1994	1,014	22	18	9	11	8	11	6	5	8	-
1993	1,116	22	15	10	13	9	19	10	11	9	-
1992	1,152	9	11	9	13	7	10	9	10	6	-
1991	1,206	16	14	11	11	11	10	14	6	4	1
1990	1,174	7	10	4	8	20	9	10	6	7	-
1989	1,143	10	11	16	6	12	9	15	3	6	1
1988	1,069	7	6	14	15	11	4	1	6	5	-
1987	1,087	9	10	8	10	9	8	7	5	2	-
1986	1,096	9	13	13	7	13	9	5	4	4	-
1985	1,046	7	9	15	10	7	5	6	4	3	-
1984	1,023	7	11	10	9	11	12	9	4	5	1
1983	1,108	14	12	8	12	8	9	8	7	8	-
1982	1,092	14	5	12	8	12	5	6	4	3	-
1981	1,064	12	9	15	10	13	8	8	1	3	2
1980	1,160	11	11	9	13	12	8	8	3	1	-
1979	1,068	17	8	11	6	8	8	4	1	3	1
Congenital malformations, deformations and chromosomal abnormalities (740-759)											
Black, female											
1998	891	12	8	8	16	10	13	14	19	18	-
1997	876	17	17	6	13	10	13	12	17	18	-
1996	934	18	11	14	13	13	12	13	15	22	-
1995	891	11	10	8	10	14	20	7	13	14	-
1994	947	8	15	15	12	10	10	15	10	16	-
1993	988	14	13	14	9	13	15	10	13	10	-
1992	1,015	10	6	17	9	13	14	11	11	10	-
1991	996	12	13	12	6	16	9	15	10	3	-
1990	1,003	10	7	12	7	12	13	8	9	7	1
1989	1,036	9	18	13	12	12	12	10	7	5	2
1988	950	11	12	4	16	8	17	6	7	6	1
1987	989	8	10	14	10	14	8	6	4	4	-
1986	965	4	13	12	11	9	6	5	5	6	-
1985	962	17	12	12	7	7	11	9	4	12	-
1984	960	17	12	13	16	7	9	8	5	6	-
1983	980	10	13	9	18	9	9	10	3	2	-
1982	1,049	9	14	8	11	11	7	8	1	6	-
1981	1,000	4	11	11	7	9	12	12	3	7	-
1980	1,008	6	11	8	7	6	10	11	2	1	-
1979	972	13	12	13	15	7	5	7	1	2	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All races, both sexes											
1998	25,992	1,030	905	784	950	1,227	1,675	2,045	2,574	7,099	88
1997	25,752	1,046	833	851	974	1,236	1,644	2,072	2,391	6,590	78
1996	26,190	1,106	905	868	995	1,270	1,631	2,073	2,438	6,373	86
1995	27,283	1,140	921	864	1,046	1,314	1,757	2,028	2,304	6,013	89
1994	25,245	878	757	749	977	1,285	1,559	1,861	2,106	5,651	66
1993	26,540	935	824	857	1,059	1,344	1,631	1,902	2,138	5,354	70
1992	23,430	715	582	706	908	1,157	1,404	1,574	1,821	4,450	58
1991	24,342	775	632	683	930	1,202	1,416	1,577	1,913	4,136	83
1990	24,137	693	659	726	975	1,196	1,383	1,568	1,830	3,801	85
1989	26,988	789	723	863	1,136	1,471	1,618	1,924	1,975	4,369	91
1988	30,951	782	849	1,040	1,538	1,940	2,207	2,537	2,651	5,198	90
1987	31,261	849	966	1,273	1,658	2,002	2,280	2,682	2,616	4,974	84
1986	31,390	795	951	1,293	1,812	2,188	2,512	2,777	2,728	5,266	89
1985	30,960	748	983	1,408	1,923	2,194	2,638	2,782	2,799	5,160	92
1984	29,721	662	965	1,446	1,908	2,238	2,619	2,697	2,629	4,789	62
1983	29,628	753	959	1,436	1,942	2,223	2,567	2,702	2,445	4,710	82
1982	29,814	821	1,174	1,549	1,967	2,245	2,533	2,613	2,377	4,118	70
1981	29,472	843	1,202	1,522	1,980	2,303	2,629	2,452	2,445	3,966	63
1980	28,808	813	1,122	1,624	1,941	2,300	2,526	2,502	2,298	3,804	64
1979	27,578	820	1,186	1,574	1,785	2,254	2,403	2,341	2,149	3,454	90
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All races, male											
1998	12,766	683	584	498	595	738	926	990	1,032	1,857	68
1997	13,065	679	578	556	615	777	887	1,050	998	1,721	59
1996	13,348	783	592	549	631	758	863	1,026	1,082	1,688	55
1995	14,244	788	617	557	670	767	964	980	969	1,624	53
1994	13,182	569	505	472	647	777	892	928	875	1,503	46
1993	14,179	621	524	552	691	802	870	959	886	1,495	51
1992	12,580	476	374	457	573	695	808	801	708	1,260	43
1991	13,407	509	412	429	593	726	811	810	803	1,179	57
1990	13,533	463	432	491	631	746	769	817	804	1,080	62
1989	15,205	537	498	551	693	916	971	1,013	862	1,328	62
1988	16,969	525	540	693	966	1,198	1,269	1,289	1,127	1,606	56
1987	17,440	551	609	824	1,063	1,269	1,346	1,429	1,094	1,593	59
1986	17,587	511	634	857	1,171	1,375	1,463	1,493	1,248	1,746	51
1985	17,470	490	664	960	1,285	1,366	1,591	1,539	1,282	1,710	63
1984	16,711	438	677	990	1,293	1,433	1,510	1,452	1,185	1,593	40
1983	16,666	480	623	943	1,300	1,422	1,563	1,465	1,148	1,638	48
1982	17,060	526	784	1,051	1,250	1,428	1,528	1,384	1,147	1,438	43
1981	17,127	535	811	1,022	1,362	1,515	1,622	1,329	1,158	1,502	45
1980	16,465	518	736	1,104	1,291	1,493	1,504	1,349	1,072	1,366	44
1979	16,012	525	806	1,071	1,184	1,507	1,466	1,245	1,031	1,270	61

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All races, female											
1998	13,226	347	321	286	355	489	749	1,055	1,542	5,242	20
1997	12,687	367	255	295	359	459	757	1,022	1,393	4,869	19
1996	12,842	323	313	319	364	512	768	1,047	1,356	4,685	31
1995	13,039	352	304	307	376	547	793	1,048	1,335	4,389	36
1994	12,063	309	252	277	330	508	667	933	1,231	4,148	20
1993	12,361	314	300	305	368	542	761	943	1,252	3,859	19
1992	10,850	239	208	249	335	462	596	773	1,113	3,190	15
1991	10,935	266	220	254	337	476	605	767	1,110	2,957	26
1990	10,604	230	227	235	344	450	614	751	1,026	2,721	23
1989	11,783	252	225	312	443	555	647	911	1,113	3,041	29
1988	13,982	257	309	347	572	742	938	1,248	1,524	3,592	34
1987	13,821	298	357	449	595	733	934	1,253	1,522	3,381	25
1986	13,803	284	317	436	641	813	1,049	1,284	1,480	3,520	38
1985	13,490	258	319	448	638	828	1,047	1,243	1,517	3,450	29
1984	13,010	224	288	456	615	805	1,109	1,245	1,444	3,196	22
1983	12,962	273	336	493	642	801	1,004	1,237	1,297	3,072	34
1982	12,754	295	390	498	717	817	1,005	1,229	1,230	2,680	27
1981	12,345	308	391	500	618	788	1,007	1,123	1,287	2,464	18
1980	12,343	295	386	520	650	807	1,022	1,153	1,226	2,438	20
1979	11,566	295	380	503	601	747	937	1,096	1,118	2,184	29
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
White, both sexes											
1998	21,041	752	713	609	757	999	1,415	1,780	2,295	6,446	69
1997	20,790	736	640	654	766	1,044	1,415	1,827	2,130	5,975	69
1996	21,126	822	687	677	807	1,075	1,404	1,810	2,180	5,838	67
1995	21,837	860	719	674	848	1,101	1,498	1,762	2,053	5,422	69
1994	19,835	641	560	577	808	1,055	1,324	1,606	1,839	5,180	52
1993	20,529	650	594	661	838	1,099	1,367	1,650	1,882	4,868	55
1992	17,959	522	434	551	708	949	1,183	1,349	1,577	4,024	46
1991	18,499	538	481	549	749	1,008	1,192	1,367	1,631	3,698	67
1990	18,058	477	463	529	785	973	1,173	1,316	1,570	3,382	59
1989	20,597	568	538	649	910	1,217	1,386	1,637	1,740	3,927	64
1988	24,060	532	628	793	1,241	1,579	1,878	2,223	2,318	4,675	73
1987	24,184	582	710	1,000	1,313	1,615	1,913	2,275	2,238	4,428	62
1986	24,304	543	683	1,003	1,423	1,768	2,060	2,310	2,309	4,579	78
1985	23,836	502	717	1,071	1,482	1,709	2,140	2,281	2,311	4,481	76
1984	22,796	464	708	1,104	1,446	1,827	2,052	2,161	2,141	4,127	50
1983	22,387	513	702	1,078	1,505	1,722	2,008	2,127	1,973	3,973	60
1982	22,376	583	852	1,139	1,454	1,751	1,958	2,085	1,925	3,487	53
1981	22,036	553	871	1,108	1,527	1,788	2,054	1,906	1,947	3,314	45
1980	20,902	538	762	1,181	1,459	1,710	1,839	1,882	1,795	3,066	49
1979	19,898	552	809	1,115	1,316	1,645	1,803	1,785	1,708	2,800	69

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
White, male											
1998	10,045	511	458	389	477	593	776	855	925	1,663	55
1997	10,244	473	449	421	488	653	761	919	896	1,527	51
1996	10,516	580	457	426	513	642	747	889	972	1,540	42
1995	11,126	585	478	430	546	638	827	851	853	1,438	39
1994	10,022	420	366	363	529	631	754	797	751	1,367	36
1993	10,631	430	375	422	542	658	729	814	781	1,334	41
1992	9,424	349	275	352	442	569	681	678	610	1,130	32
1991	10,069	348	308	345	485	622	694	710	685	1,024	44
1990	9,951	306	301	354	501	610	662	692	694	951	43
1989	11,361	385	363	407	550	760	848	864	742	1,163	43
1988	12,947	350	398	529	769	980	1,079	1,124	962	1,408	44
1987	13,322	368	455	645	855	1,024	1,153	1,216	932	1,379	43
1986	13,465	346	446	657	922	1,113	1,209	1,245	1,048	1,492	44
1985	13,340	319	485	724	999	1,077	1,308	1,266	1,053	1,428	53
1984	12,682	310	488	743	984	1,165	1,185	1,168	948	1,334	28
1983	12,525	328	458	711	1,007	1,114	1,245	1,165	902	1,360	36
1982	12,753	364	569	773	932	1,125	1,208	1,106	932	1,182	33
1981	12,786	344	578	758	1,078	1,193	1,281	1,036	908	1,212	32
1980	11,992	335	504	806	978	1,156	1,111	1,018	843	1,082	32
1979	11,599	350	552	765	890	1,128	1,139	961	815	1,003	46
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
White, female											
1998	10,996	241	255	220	280	406	639	925	1,370	4,783	14
1997	10,546	263	191	233	278	391	654	908	1,234	4,448	18
1996	10,610	242	230	251	294	433	657	921	1,208	4,298	25
1995	10,711	275	241	244	302	463	671	911	1,200	3,984	30
1994	9,813	221	194	214	279	424	570	809	1,088	3,813	16
1993	9,898	220	219	239	296	441	638	836	1,101	3,534	14
1992	8,535	173	159	199	266	380	502	671	967	2,894	14
1991	8,430	190	173	204	264	386	498	657	946	2,674	23
1990	8,107	171	162	175	284	363	511	624	876	2,431	16
1989	9,236	183	175	242	360	457	538	773	998	2,764	21
1988	11,113	182	230	264	472	599	799	1,099	1,356	3,267	29
1987	10,862	214	255	355	458	591	760	1,059	1,306	3,049	19
1986	10,839	197	237	346	501	655	851	1,065	1,261	3,087	34
1985	10,496	183	232	347	483	632	832	1,015	1,258	3,053	23
1984	10,114	154	220	361	462	662	867	993	1,193	2,793	22
1983	9,862	185	244	367	498	608	763	962	1,071	2,613	24
1982	9,623	219	283	366	522	626	750	979	993	2,305	20
1981	9,250	209	293	350	449	595	773	870	1,039	2,102	13
1980	8,910	203	258	375	481	554	728	864	952	1,984	17
1979	8,299	202	257	350	426	517	664	824	893	1,797	23

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All other, both sexes											
1998	4,951	278	192	175	193	228	260	265	279	653	19
1997	4,962	310	193	197	208	192	229	245	261	615	9
1996	5,064	284	218	191	188	195	227	263	258	535	19
1995	5,446	280	202	190	198	213	259	266	251	591	20
1994	5,410	237	197	172	169	230	235	255	267	471	14
1993	6,011	285	230	196	221	245	264	252	256	486	15
1992	5,471	193	148	155	200	208	221	225	244	426	12
1991	5,843	237	151	134	181	194	224	210	282	438	16
1990	6,079	216	196	197	190	223	210	252	260	419	26
1989	6,391	221	185	214	226	254	232	287	235	442	27
1988	6,891	250	221	247	297	361	329	314	333	523	17
1987	7,077	267	256	273	345	387	367	407	378	546	22
1986	7,086	252	268	290	389	420	452	467	419	687	11
1985	7,124	246	266	337	441	485	498	501	488	679	16
1984	6,925	198	257	342	462	411	567	536	488	662	12
1983	7,241	240	257	358	437	501	559	575	472	737	22
1982	7,438	238	322	410	513	494	575	528	452	631	17
1981	7,436	290	331	414	453	515	575	546	498	652	18
1980	7,906	275	360	443	482	590	687	620	503	738	15
1979	7,680	268	377	459	469	609	600	556	441	654	21
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All other, male											
1998	2,721	172	126	109	118	145	150	135	107	194	13
1997	2,821	206	129	135	127	124	126	131	102	194	8
1996	2,832	203	135	123	118	116	116	137	110	148	13
1995	3,118	203	139	127	124	129	137	129	116	186	14
1994	3,160	149	139	109	118	146	138	131	124	136	10
1993	3,548	191	149	130	149	144	141	145	105	161	10
1992	3,156	127	99	105	131	126	127	123	98	130	11
1991	3,338	161	104	84	108	104	117	100	118	155	13
1990	3,582	157	131	137	130	136	107	125	110	129	19
1989	3,844	152	135	144	143	156	123	149	120	165	19
1988	4,022	175	142	164	197	218	190	165	165	198	12
1987	4,118	183	154	179	208	245	193	213	162	214	16
1986	4,122	165	188	200	249	262	254	248	200	254	7
1985	4,130	171	179	236	286	289	283	273	229	282	10
1984	4,029	128	189	247	309	268	325	284	237	259	12
1983	4,141	152	165	232	293	308	318	300	246	278	12
1982	4,307	162	215	278	318	303	320	278	215	256	10
1981	4,341	191	233	264	284	322	341	293	250	290	13
1980	4,473	183	232	298	313	337	393	331	229	284	12
1979	4,413	175	254	306	294	379	327	284	216	267	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
All other, female											
1998	2,230	106	66	66	75	83	110	130	172	459	6
1997	2,141	104	64	62	81	68	103	114	159	421	1
1996	2,232	81	83	68	70	79	111	126	148	387	6
1995	2,328	77	63	63	74	84	122	137	135	405	6
1994	2,250	88	58	63	51	84	97	124	143	335	4
1993	2,463	94	81	66	72	101	123	107	151	325	5
1992	2,315	66	49	50	69	82	94	102	146	296	1
1991	2,505	76	47	50	73	90	107	110	164	283	3
1990	2,497	59	65	60	60	87	103	127	150	290	7
1989	2,547	69	50	70	83	98	109	138	115	277	8
1988	2,869	75	79	83	100	143	139	149	168	325	5
1987	2,959	84	102	94	137	142	174	194	216	332	6
1986	2,964	87	80	90	140	158	198	219	219	433	4
1985	2,994	75	87	101	155	196	215	228	259	397	6
1984	2,896	70	68	95	153	143	242	252	251	403	-
1983	3,100	88	92	126	144	193	241	275	226	459	10
1982	3,131	76	107	132	195	191	255	250	237	375	7
1981	3,095	99	98	150	169	193	234	253	248	362	5
1980	3,433	92	128	145	169	253	294	289	274	454	3
1979	3,267	93	123	153	175	230	273	272	225	387	6
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
Black, both sexes											
1998	4,481	257	173	163	182	205	242	242	251	594	17
1997	4,505	293	181	184	196	166	206	222	234	556	8
1996	4,607	265	203	171	175	186	207	242	233	486	13
1995	4,958	260	179	175	184	201	232	250	228	540	16
1994	4,895	216	182	158	159	211	218	234	247	436	11
1993	5,499	267	215	184	201	233	244	235	232	450	14
1992	4,976	179	134	137	184	193	206	207	230	388	10
1991	5,306	217	133	123	167	181	206	193	266	394	16
1990	5,505	202	179	183	168	211	186	231	242	389	22
1989	5,835	205	171	202	215	242	222	273	226	418	26
1988	6,353	233	205	232	283	347	318	293	316	486	16
1987	6,562	255	244	250	329	369	355	388	361	514	18
1986	6,642	237	258	278	378	403	436	448	403	642	11
1985	6,626	232	260	327	420	464	476	483	459	633	16
1984	6,511	187	251	326	445	401	550	514	467	622	12
1983	6,787	231	249	342	424	480	539	557	443	707	18
1982	6,938	220	311	394	492	474	553	507	432	584	13
1981	6,886	269	316	392	431	486	550	516	471	619	18
1980	7,461	266	350	428	458	571	661	597	481	689	15
1979	7,242	258	364	443	455	595	581	534	420	617	17

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
Black, male											
1998	2,451	158	114	105	108	132	139	124	97	173	11
1997	2,560	194	119	127	120	109	110	115	91	171	8
1996	2,564	190	125	112	111	113	103	128	99	130	9
1995	2,839	185	124	119	117	122	118	124	105	167	11
1994	2,839	133	130	100	111	133	129	119	109	118	8
1993	3,235	176	141	123	135	140	128	137	91	144	9
1992	2,856	117	89	96	120	120	118	114	90	115	9
1991	3,020	148	93	79	101	101	105	92	108	136	13
1990	3,215	145	120	129	117	129	94	111	101	117	17
1989	3,496	142	125	134	137	151	118	139	114	155	18
1988	3,692	160	130	151	188	209	182	153	156	179	12
1987	3,817	175	149	165	201	233	182	203	157	199	12
1986	3,865	155	180	193	242	250	243	239	193	241	7
1985	3,829	159	178	229	273	276	270	262	211	257	10
1984	3,784	122	184	234	299	262	314	269	226	239	12
1983	3,879	148	162	221	280	291	307	290	226	267	9
1982	4,014	153	208	270	308	292	307	264	204	231	8
1981	3,982	182	220	251	270	300	323	271	229	279	13
1980	4,214	176	224	289	296	324	377	322	216	263	12
1979	4,163	169	245	297	285	369	313	269	206	253	12
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (780-799)											
Black, female											
1998	2,030	99	59	58	74	73	103	118	154	421	6
1997	1,945	99	62	57	76	57	96	107	143	385	-
1996	2,043	75	78	59	64	73	104	114	134	356	4
1995	2,119	75	55	56	67	79	114	126	123	373	5
1994	2,056	83	52	58	48	78	89	115	138	318	3
1993	2,264	91	74	61	66	93	116	98	141	306	5
1992	2,120	62	45	41	64	73	88	93	140	273	1
1991	2,286	69	40	44	66	80	101	101	158	258	3
1990	2,290	57	59	54	51	82	92	120	141	272	5
1989	2,339	63	46	68	78	91	104	134	112	263	8
1988	2,661	73	75	81	95	138	136	140	160	307	4
1987	2,745	80	95	85	128	136	173	185	204	315	6
1986	2,777	82	78	85	136	153	193	209	210	401	4
1985	2,797	73	82	98	147	188	206	221	248	376	6
1984	2,727	65	67	92	146	139	236	245	241	383	-
1983	2,908	83	87	121	144	189	232	267	217	440	9
1982	2,924	67	103	124	184	182	246	243	228	353	5
1981	2,904	87	96	141	161	186	227	245	242	340	5
1980	3,247	90	126	139	162	247	284	275	265	426	3
1979	3,079	89	119	146	170	226	268	265	214	364	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
All other diseases (Residual)											
All races, both sexes											
1998	179,495	5,609	6,013	6,460	7,810	10,933	16,069	22,076	27,879	61,984	8
1997	170,284	5,392	5,590	5,887	7,561	10,713	15,576	21,300	25,990	57,471	17
1996	161,868	5,118	5,113	5,663	7,490	10,552	15,202	19,990	24,771	53,055	23
1995	155,512	4,970	5,074	5,521	7,316	10,416	14,611	19,062	23,680	49,604	21
1994	149,106	4,600	4,756	5,446	7,404	10,423	14,396	18,346	22,574	46,235	18
1993	143,479	4,327	4,462	5,255	7,294	10,554	13,968	17,992	21,425	43,168	18
1992	134,119	4,085	4,144	4,979	7,280	10,334	13,427	16,802	19,678	38,494	16
1991	131,332	3,668	4,035	5,114	7,568	10,125	13,231	16,508	19,500	36,876	23
1990	127,401	3,494	4,055	5,078	7,510	10,214	12,716	16,135	18,716	35,303	12
1989	126,363	3,466	3,924	5,156	7,604	10,294	12,728	16,058	18,424	34,042	17
1988	123,717	3,472	3,873	5,332	7,791	10,089	12,538	15,810	17,740	32,650	23
1987	118,333	3,135	3,866	5,206	7,639	10,020	12,406	15,279	16,706	30,053	21
1986	119,140	3,558	4,105	5,473	7,624	9,708	11,953	14,722	15,721	27,687	41
1985	112,609	3,217	4,023	5,661	7,668	9,339	11,638	14,016	15,225	25,804	37
1984	104,934	2,968	3,777	5,483	7,480	9,167	11,178	13,296	13,913	23,589	19
1983	100,048	2,876	4,033	5,720	7,364	9,091	10,719	12,635	12,920	21,403	19
1982	94,190	2,829	4,024	5,445	7,021	8,632	10,409	11,700	12,141	19,334	23
1981	93,285	3,036	4,163	5,811	7,134	8,701	10,181	11,196	11,721	18,180	29
1980	92,114	3,129	4,514	5,763	7,149	8,871	10,105	10,987	11,309	16,919	19
1979	86,259	3,000	4,502	5,604	6,811	8,288	9,367	10,339	10,252	14,860	20
All other diseases (Residual)											
All races, male											
1998	72,852	3,415	3,507	3,552	4,210	5,618	7,683	9,812	10,711	15,804	5
1997	70,053	3,338	3,183	3,288	4,001	5,502	7,566	9,486	10,153	14,839	16
1996	67,536	3,134	2,928	3,213	4,088	5,382	7,361	8,935	9,635	13,788	17
1995	65,682	3,078	2,969	3,188	4,024	5,435	7,116	8,454	9,091	13,068	13
1994	63,626	2,752	2,811	3,118	4,153	5,516	7,062	8,128	8,764	12,233	15
1993	61,845	2,585	2,644	2,954	4,050	5,508	6,864	8,167	8,333	11,638	13
1992	58,852	2,547	2,454	2,816	4,018	5,396	6,594	7,665	7,591	10,673	12
1991	57,732	2,258	2,354	2,930	4,247	5,304	6,533	7,507	7,580	10,061	19
1990	56,120	2,163	2,430	2,936	4,136	5,429	6,168	7,376	7,131	9,647	8
1989	56,316	2,122	2,347	2,940	4,317	5,508	6,325	7,304	7,042	9,402	13
1988	55,694	2,112	2,328	3,050	4,402	5,436	6,261	7,161	6,750	9,212	17
1987	53,463	1,953	2,319	2,998	4,177	5,264	6,196	6,906	6,565	8,517	11
1986	57,115	2,338	2,518	3,278	4,273	5,091	5,997	6,729	6,079	8,068	20
1985	53,400	2,026	2,411	3,280	4,258	5,036	5,827	6,411	5,977	7,686	21
1984	48,950	1,783	2,222	3,070	4,139	4,893	5,610	6,057	5,445	7,024	10
1983	47,349	1,693	2,332	3,296	4,133	4,871	5,363	5,821	5,160	6,726	11
1982	44,115	1,636	2,358	3,076	3,880	4,577	5,177	5,167	4,690	6,061	11
1981	44,225	1,754	2,433	3,263	3,890	4,667	5,113	5,071	4,583	5,783	19
1980	44,302	1,818	2,644	3,291	3,956	4,797	5,068	4,990	4,404	5,558	12
1979	42,081	1,814	2,627	3,242	3,878	4,462	4,617	4,746	4,160	4,890	13

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
All other diseases (Residual)											
All races, female											
1998	106,643	2,194	2,506	2,908	3,600	5,315	8,386	12,264	17,168	46,180	3
1997	100,231	2,054	2,407	2,599	3,560	5,211	8,010	11,814	15,837	42,632	1
1996	94,332	1,984	2,185	2,450	3,402	5,170	7,841	11,055	15,136	39,267	6
1995	89,830	1,892	2,105	2,333	3,292	4,981	7,495	10,608	14,589	36,536	8
1994	85,480	1,848	1,945	2,328	3,251	4,907	7,334	10,218	13,810	34,002	3
1993	81,634	1,742	1,818	2,301	3,244	5,046	7,104	9,825	13,092	31,530	5
1992	75,267	1,538	1,690	2,163	3,262	4,938	6,833	9,137	12,087	27,821	4
1991	73,600	1,410	1,681	2,184	3,321	4,821	6,698	9,001	11,920	26,815	4
1990	71,281	1,331	1,625	2,142	3,374	4,785	6,548	8,759	11,585	25,656	4
1989	70,047	1,344	1,577	2,216	3,287	4,786	6,403	8,754	11,382	24,640	4
1988	68,023	1,360	1,545	2,282	3,389	4,653	6,277	8,649	10,990	23,438	6
1987	64,870	1,182	1,547	2,208	3,462	4,756	6,210	8,373	10,141	21,536	10
1986	62,025	1,220	1,587	2,195	3,351	4,617	5,956	7,993	9,642	19,619	21
1985	59,209	1,191	1,612	2,381	3,410	4,303	5,811	7,605	9,248	18,118	16
1984	55,984	1,185	1,555	2,413	3,341	4,274	5,568	7,239	8,468	16,565	9
1983	52,699	1,183	1,701	2,424	3,231	4,220	5,356	6,814	7,760	14,677	8
1982	50,075	1,193	1,666	2,369	3,141	4,055	5,232	6,533	7,451	13,273	12
1981	49,060	1,282	1,730	2,548	3,244	4,034	5,068	6,125	7,138	12,397	10
1980	47,812	1,311	1,870	2,472	3,193	4,074	5,037	5,997	6,905	11,361	7
1979	44,178	1,186	1,875	2,362	2,933	3,826	4,750	5,593	6,092	9,970	7
All other diseases (Residual)											
White, both sexes											
1998	155,195	4,092	4,570	5,032	6,211	9,092	13,794	19,550	25,271	57,104	6
1997	147,115	3,854	4,305	4,583	6,050	8,997	13,431	18,888	23,557	52,911	12
1996	139,205	3,639	3,861	4,402	6,062	8,858	13,051	17,651	22,349	48,805	16
1995	133,283	3,547	3,710	4,286	5,866	8,695	12,652	16,918	21,320	45,577	13
1994	127,710	3,331	3,594	4,267	5,983	8,726	12,389	16,176	20,293	42,460	11
1993	122,337	3,145	3,314	4,027	5,897	8,882	12,011	15,903	19,189	39,497	12
1992	113,982	2,939	3,016	3,788	5,858	8,657	11,517	14,799	17,679	35,294	11
1991	111,397	2,644	2,949	3,934	6,115	8,446	11,441	14,511	17,443	33,729	16
1990	107,587	2,436	2,927	3,907	6,052	8,557	10,923	14,128	16,766	32,163	6
1989	106,213	2,386	2,790	3,962	6,072	8,591	10,930	14,073	16,472	30,940	8
1988	103,956	2,386	2,781	4,099	6,291	8,406	10,752	13,805	15,862	29,852	16
1987	99,587	2,183	2,809	4,018	6,190	8,396	10,645	13,351	14,898	27,558	12
1986	99,221	2,484	3,001	4,246	6,181	8,152	10,263	12,857	14,099	25,273	30
1985	94,270	2,278	2,976	4,479	6,288	7,868	10,003	12,310	13,601	23,615	20
1984	88,252	2,052	2,829	4,355	6,162	7,763	9,657	11,663	12,375	21,738	14
1983	84,328	2,041	3,022	4,559	6,028	7,719	9,239	11,130	11,606	19,692	11
1982	79,610	1,979	3,110	4,381	5,852	7,355	9,012	10,309	10,945	17,792	14
1981	78,477	2,174	3,197	4,615	5,944	7,418	8,837	9,873	10,540	16,725	15
1980	77,357	2,191	3,419	4,627	5,950	7,546	8,714	9,694	10,282	15,627	8
1979	72,712	2,151	3,497	4,508	5,717	7,121	8,175	9,083	9,373	13,731	12

Source: CDC/NCHS, National Vital Statistics System
 See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other diseases (Residual)											
White, male											
1998	61,862	2,548	2,694	2,777	3,329	4,721	6,657	8,689	9,693	14,424	4
1997	59,417	2,434	2,471	2,573	3,234	4,662	6,562	8,366	9,199	13,476	12
1996	56,916	2,276	2,236	2,504	3,335	4,544	6,343	7,855	8,676	12,534	12
1995	55,029	2,213	2,159	2,501	3,233	4,534	6,152	7,490	8,174	11,834	9
1994	53,380	2,027	2,118	2,483	3,368	4,617	6,084	7,183	7,863	11,049	10
1993	51,636	1,900	1,987	2,278	3,278	4,654	5,902	7,232	7,403	10,439	10
1992	49,001	1,846	1,790	2,167	3,282	4,527	5,662	6,737	6,737	9,632	10
1991	47,965	1,633	1,714	2,248	3,454	4,430	5,656	6,618	6,755	9,087	12
1990	46,248	1,500	1,781	2,275	3,328	4,587	5,308	6,421	6,325	8,633	4
1989	46,348	1,469	1,645	2,264	3,494	4,620	5,429	6,402	6,258	8,436	7
1988	45,567	1,442	1,674	2,341	3,572	4,495	5,385	6,222	6,007	8,286	11
1987	43,956	1,357	1,706	2,323	3,382	4,454	5,340	6,001	5,804	7,660	7
1986	46,326	1,641	1,843	2,564	3,469	4,278	5,124	5,840	5,400	7,287	13
1985	43,615	1,434	1,795	2,605	3,477	4,235	5,014	5,596	5,284	6,974	11
1984	40,294	1,247	1,664	2,452	3,377	4,152	4,861	5,319	4,795	6,351	7
1983	39,181	1,211	1,747	2,628	3,402	4,139	4,620	5,101	4,592	6,123	6
1982	36,503	1,153	1,821	2,479	3,204	3,881	4,485	4,547	4,179	5,499	6
1981	36,492	1,246	1,877	2,561	3,244	3,987	4,434	4,461	4,073	5,246	10
1980	36,448	1,279	1,999	2,630	3,295	4,070	4,337	4,368	3,967	5,057	8
1979	34,792	1,307	2,051	2,599	3,229	3,850	4,002	4,144	3,750	4,450	7
All other diseases (Residual)											
White, female											
1998	93,333	1,544	1,876	2,255	2,882	4,371	7,137	10,861	15,578	42,680	2
1997	87,698	1,420	1,834	2,010	2,816	4,335	6,869	10,522	14,358	39,435	-
1996	82,289	1,363	1,625	1,898	2,727	4,314	6,708	9,796	13,673	36,271	4
1995	78,254	1,334	1,551	1,785	2,633	4,161	6,500	9,428	13,146	33,743	4
1994	74,330	1,304	1,476	1,784	2,615	4,109	6,305	8,993	12,430	31,411	1
1993	70,701	1,245	1,327	1,749	2,619	4,228	6,109	8,671	11,786	29,058	2
1992	64,981	1,093	1,226	1,621	2,576	4,130	5,855	8,062	10,942	25,662	1
1991	63,432	1,011	1,235	1,686	2,661	4,016	5,785	7,893	10,688	24,642	4
1990	61,339	936	1,146	1,632	2,724	3,970	5,615	7,707	10,441	23,530	2
1989	59,865	917	1,145	1,698	2,578	3,971	5,501	7,671	10,214	22,504	1
1988	58,389	944	1,107	1,758	2,719	3,911	5,367	7,583	9,855	21,566	5
1987	55,631	826	1,103	1,695	2,808	3,942	5,305	7,350	9,094	19,898	5
1986	52,895	843	1,158	1,682	2,712	3,874	5,139	7,017	8,699	17,986	17
1985	50,655	844	1,181	1,874	2,811	3,633	4,989	6,714	8,317	16,641	9
1984	47,958	805	1,165	1,903	2,785	3,611	4,796	6,344	7,580	15,387	7
1983	45,147	830	1,275	1,931	2,626	3,580	4,619	6,029	7,014	13,569	5
1982	43,107	826	1,289	1,902	2,648	3,474	4,527	5,762	6,766	12,293	8
1981	41,985	928	1,320	2,054	2,700	3,431	4,403	5,412	6,467	11,479	5
1980	40,909	912	1,420	1,997	2,655	3,476	4,377	5,326	6,315	10,570	-
1979	37,920	844	1,446	1,909	2,488	3,271	4,173	4,939	5,623	9,281	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other diseases (Residual)											
All other, both sexes											
1998	24,300	1,517	1,443	1,428	1,599	1,841	2,275	2,526	2,608	4,880	2
1997	23,169	1,538	1,285	1,304	1,511	1,716	2,145	2,412	2,433	4,560	5
1996	22,663	1,479	1,252	1,261	1,428	1,694	2,151	2,339	2,422	4,250	7
1995	22,229	1,423	1,364	1,235	1,450	1,721	1,959	2,144	2,360	4,027	8
1994	21,396	1,269	1,162	1,179	1,421	1,697	2,007	2,170	2,281	3,775	7
1993	21,142	1,182	1,148	1,228	1,397	1,672	1,957	2,089	2,236	3,671	6
1992	20,137	1,146	1,128	1,191	1,422	1,677	1,910	2,003	1,999	3,200	5
1991	19,935	1,024	1,086	1,180	1,453	1,679	1,790	1,997	2,057	3,147	7
1990	19,814	1,058	1,128	1,171	1,458	1,657	1,793	2,007	1,950	3,140	6
1989	20,150	1,080	1,134	1,194	1,532	1,703	1,798	1,985	1,952	3,102	9
1988	19,761	1,086	1,092	1,233	1,500	1,683	1,786	2,005	1,878	2,798	7
1987	18,746	952	1,057	1,188	1,449	1,624	1,761	1,928	1,808	2,495	9
1986	19,919	1,074	1,104	1,227	1,443	1,556	1,690	1,865	1,622	2,414	11
1985	18,339	939	1,047	1,182	1,380	1,471	1,635	1,706	1,624	2,189	17
1984	16,682	916	948	1,128	1,318	1,404	1,521	1,633	1,538	1,851	5
1983	15,720	835	1,011	1,161	1,336	1,372	1,480	1,505	1,314	1,711	8
1982	14,580	850	914	1,064	1,169	1,277	1,397	1,391	1,196	1,542	9
1981	14,808	862	966	1,196	1,190	1,283	1,344	1,323	1,181	1,455	14
1980	14,757	938	1,095	1,136	1,199	1,325	1,391	1,293	1,027	1,292	11
1979	13,547	849	1,005	1,096	1,094	1,167	1,192	1,256	879	1,129	8
All other diseases (Residual)											
All other, male											
1998	10,990	867	813	775	881	897	1,026	1,123	1,018	1,380	1
1997	10,636	904	712	715	767	840	1,004	1,120	954	1,363	4
1996	10,620	858	692	709	753	838	1,018	1,080	959	1,254	5
1995	10,653	865	810	687	791	901	964	964	917	1,234	4
1994	10,246	725	693	635	785	899	978	945	901	1,184	5
1993	10,209	685	657	676	772	854	962	935	930	1,199	3
1992	9,851	701	664	649	736	869	932	928	854	1,041	2
1991	9,767	625	640	682	793	874	877	889	825	974	7
1990	9,872	663	649	661	808	842	860	955	806	1,014	4
1989	9,968	653	702	676	823	888	896	902	784	966	6
1988	10,127	670	654	709	830	941	876	939	743	926	6
1987	9,507	596	613	675	795	810	856	905	761	857	4
1986	10,789	697	675	714	804	813	873	889	679	781	7
1985	9,785	592	616	675	781	801	813	815	693	712	10
1984	8,656	536	558	618	762	741	749	738	650	673	3
1983	8,168	482	585	668	731	732	743	720	568	603	5
1982	7,612	483	537	597	676	696	692	620	511	562	5
1981	7,733	508	556	702	646	680	679	610	510	537	9
1980	7,854	539	645	661	661	727	731	622	437	501	4
1979	7,289	507	576	643	649	612	615	602	410	440	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other diseases (Residual)											
All other, female											
1998	13,310	650	630	653	718	944	1,249	1,403	1,590	3,500	1
1997	12,533	634	573	589	744	876	1,141	1,292	1,479	3,197	1
1996	12,043	621	560	552	675	856	1,133	1,259	1,463	2,996	2
1995	11,576	558	554	548	659	820	995	1,180	1,443	2,793	4
1994	11,150	544	469	544	636	798	1,029	1,225	1,380	2,591	2
1993	10,933	497	491	552	625	818	995	1,154	1,306	2,472	3
1992	10,286	445	464	542	686	808	978	1,075	1,145	2,159	3
1991	10,168	399	446	498	660	805	913	1,108	1,232	2,173	-
1990	9,942	395	479	510	650	815	933	1,052	1,144	2,126	2
1989	10,182	427	432	518	709	815	902	1,083	1,168	2,136	3
1988	9,634	416	438	524	670	742	910	1,066	1,135	1,872	1
1987	9,239	356	444	513	654	814	905	1,023	1,047	1,638	5
1986	9,130	377	429	513	639	743	817	976	943	1,633	4
1985	8,554	347	431	507	599	670	822	891	931	1,477	7
1984	8,026	380	390	510	556	663	772	895	888	1,178	2
1983	7,552	353	426	493	605	640	737	785	746	1,108	3
1982	6,968	367	377	467	493	581	705	771	685	980	4
1981	7,075	354	410	494	544	603	665	713	671	918	5
1980	6,903	399	450	475	538	598	660	671	590	791	7
1979	6,258	342	429	453	445	555	577	654	469	689	2
All other diseases (Residual)											
Black, both sexes											
1998	21,524	1,374	1,301	1,251	1,417	1,626	2,025	2,232	2,314	4,263	1
1997	20,557	1,403	1,165	1,165	1,347	1,516	1,907	2,072	2,180	4,030	5
1996	20,217	1,349	1,107	1,134	1,287	1,486	1,946	2,099	2,122	3,782	7
1995	19,989	1,291	1,228	1,121	1,307	1,527	1,764	1,928	2,115	3,609	8
1994	19,268	1,150	1,039	1,076	1,294	1,531	1,812	1,955	2,047	3,379	7
1993	19,095	1,074	1,040	1,094	1,263	1,528	1,801	1,906	1,989	3,297	5
1992	18,243	1,049	1,033	1,075	1,316	1,514	1,739	1,803	1,821	2,869	5
1991	18,234	956	1,006	1,085	1,335	1,530	1,617	1,845	1,884	2,846	7
1990	18,128	975	1,047	1,070	1,322	1,516	1,650	1,840	1,791	2,841	6
1989	18,521	989	1,042	1,092	1,422	1,569	1,688	1,833	1,795	2,808	9
1988	18,196	1,012	1,006	1,134	1,375	1,557	1,660	1,830	1,723	2,553	4
1987	17,362	887	967	1,093	1,350	1,524	1,641	1,795	1,665	2,282	8
1986	18,569	1,012	1,036	1,146	1,345	1,461	1,575	1,744	1,485	2,178	9
1985	17,077	891	976	1,105	1,293	1,384	1,528	1,574	1,500	2,006	15
1984	15,479	863	896	1,058	1,233	1,309	1,429	1,508	1,419	1,689	5
1983	14,517	767	937	1,080	1,253	1,286	1,377	1,377	1,217	1,544	7
1982	13,558	780	854	1,015	1,095	1,193	1,317	1,300	1,113	1,410	7
1981	13,778	799	906	1,113	1,130	1,197	1,266	1,229	1,112	1,343	12
1980	13,728	886	1,039	1,058	1,124	1,238	1,293	1,211	955	1,171	10
1979	12,569	801	936	1,031	1,035	1,100	1,101	1,167	800	1,038	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
All other diseases (Residual)											
Black, male											
1998	9,646	777	729	674	782	796	908	999	883	1,146	-
1997	9,338	833	644	639	677	755	879	960	833	1,144	4
1996	9,413	785	619	634	682	737	923	958	818	1,080	5
1995	9,488	775	726	616	704	816	867	857	799	1,067	4
1994	9,194	653	617	587	718	807	882	849	794	1,030	5
1993	9,179	629	602	611	708	786	879	839	800	1,045	2
1992	8,876	647	614	597	687	790	839	819	750	907	2
1991	8,871	578	590	625	728	800	793	811	733	864	7
1990	8,968	615	604	610	732	761	771	873	714	898	4
1989	9,095	602	647	624	763	816	836	818	694	853	6
1988	9,266	631	601	651	769	868	802	837	657	806	3
1987	8,790	559	572	627	748	769	797	830	680	766	3
1986	10,049	661	630	671	745	763	802	816	611	685	7
1985	9,114	567	585	634	738	758	758	744	620	628	10
1984	8,018	508	527	586	716	693	693	670	579	598	3
1983	7,523	452	539	625	686	678	676	652	515	537	4
1982	7,036	436	510	573	631	647	649	564	462	505	5
1981	7,149	474	525	657	611	629	636	545	467	483	7
1980	7,277	507	617	613	618	673	667	572	404	443	4
1979	6,711	475	540	599	611	577	563	539	356	393	6
All other diseases (Residual)											
Black, female											
1998	11,878	597	572	577	635	830	1,117	1,233	1,431	3,117	1
1997	11,219	570	521	526	670	761	1,028	1,112	1,347	2,886	1
1996	10,804	564	488	500	605	749	1,023	1,141	1,304	2,702	2
1995	10,501	516	502	505	603	711	897	1,071	1,316	2,542	4
1994	10,074	497	422	489	576	724	930	1,106	1,253	2,349	2
1993	9,916	445	438	483	555	742	922	1,067	1,189	2,252	3
1992	9,367	402	419	478	629	724	900	984	1,071	1,962	3
1991	9,363	378	416	460	607	730	824	1,034	1,151	1,982	-
1990	9,160	360	443	460	590	755	879	967	1,077	1,943	2
1989	9,426	387	395	468	659	753	852	1,015	1,101	1,955	3
1988	8,930	381	405	483	606	689	858	993	1,066	1,747	1
1987	8,572	328	395	466	602	755	844	965	985	1,516	5
1986	8,520	351	406	475	600	698	773	928	874	1,493	2
1985	7,963	324	391	471	555	626	770	830	880	1,378	5
1984	7,461	355	369	472	517	616	736	838	840	1,091	2
1983	6,994	315	398	455	567	608	701	725	702	1,007	3
1982	6,522	344	344	442	464	546	668	736	651	905	2
1981	6,629	325	381	456	519	568	630	684	645	860	5
1980	6,451	379	422	445	506	565	626	639	551	728	6
1979	5,858	326	396	432	424	523	538	628	444	645	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All races, both sexes											
1998	94,331	6,090	4,550	3,669	3,259	3,594	4,532	5,505	6,245	10,729	109
1997	92,353	5,797	4,320	3,523	3,203	3,528	4,342	5,293	5,887	10,138	100
1996	91,776	5,597	3,999	3,309	3,216	3,602	4,475	5,016	5,784	9,820	116
1995	90,402	5,154	3,872	3,181	3,237	3,485	4,268	4,888	5,276	9,224	104
1994	88,649	4,892	3,651	3,040	3,078	3,498	4,136	4,832	5,370	8,579	99
1993	87,598	4,447	3,378	2,893	3,168	3,463	3,961	4,715	5,169	8,486	108
1992	83,952	4,075	3,230	2,948	3,127	3,559	3,936	4,522	4,867	7,808	117
1991	86,711	3,927	3,067	2,962	3,288	3,633	3,883	4,565	4,820	7,697	119
1990	89,148	3,857	3,351	3,205	3,684	3,834	3,911	4,619	4,582	7,263	113
1989	92,036	3,945	3,401	3,445	3,725	3,943	4,161	4,718	4,701	7,223	122
1988	94,062	3,873	3,463	3,464	3,792	4,023	4,199	4,599	4,739	7,348	132
1987	91,988	3,597	3,211	3,517	3,902	3,840	4,082	4,526	4,585	6,700	127
1986	92,208	3,475	3,333	3,509	3,791	3,630	4,082	4,444	4,375	6,497	165
1985	90,605	3,522	3,382	3,815	3,947	3,678	4,162	4,473	4,299	6,443	136
1984	90,283	3,579	3,491	3,773	3,861	3,787	4,024	4,383	4,162	6,307	99
1983	89,828	3,474	3,681	3,858	3,874	3,732	3,917	4,123	4,100	6,215	118
1982	91,497	3,664	3,823	4,005	3,830	3,685	3,860	3,995	3,924	5,867	96
1981	98,155	3,913	4,393	4,375	4,089	3,894	4,088	4,007	4,058	6,060	116
1980	103,281	4,170	4,547	4,618	4,236	4,212	4,188	4,485	4,232	6,207	101
1979	102,670	4,204	4,569	4,602	4,299	4,231	4,153	4,271	4,061	5,688	143
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All races, male											
1998	61,399	4,529	3,293	2,576	2,203	2,264	2,639	2,977	3,146	4,086	101
1997	60,404	4,333	3,108	2,458	2,164	2,251	2,583	2,898	2,931	3,866	91
1996	60,127	4,150	2,881	2,330	2,172	2,322	2,640	2,727	2,862	3,717	99
1995	60,066	3,901	2,806	2,251	2,188	2,237	2,447	2,655	2,637	3,578	100
1994	59,250	3,623	2,638	2,158	2,113	2,178	2,435	2,677	2,684	3,360	89
1993	58,773	3,311	2,499	2,062	2,123	2,108	2,387	2,539	2,568	3,317	96
1992	56,554	2,989	2,411	2,104	2,157	2,267	2,368	2,434	2,446	2,967	111
1991	58,578	2,902	2,243	2,144	2,252	2,323	2,283	2,477	2,369	2,938	107
1990	60,606	2,828	2,442	2,255	2,514	2,417	2,326	2,436	2,218	2,773	101
1989	62,480	2,946	2,486	2,514	2,560	2,524	2,418	2,528	2,315	2,813	109
1988	64,367	2,942	2,498	2,463	2,595	2,581	2,474	2,457	2,314	2,903	121
1987	63,342	2,664	2,348	2,481	2,687	2,411	2,446	2,445	2,284	2,681	111
1986	64,133	2,638	2,487	2,529	2,582	2,310	2,385	2,434	2,108	2,613	146
1985	62,755	2,640	2,448	2,791	2,678	2,309	2,380	2,473	2,101	2,631	127
1984	62,776	2,651	2,576	2,711	2,635	2,491	2,396	2,410	2,014	2,550	90
1983	62,605	2,617	2,709	2,796	2,687	2,413	2,296	2,236	1,983	2,484	101
1982	64,450	2,774	2,818	2,898	2,665	2,399	2,258	2,218	1,879	2,327	75
1981	69,626	2,949	3,273	3,138	2,881	2,587	2,462	2,235	1,937	2,428	104
1980	72,945	3,145	3,359	3,333	2,927	2,777	2,477	2,409	2,002	2,429	90
1979	73,048	3,168	3,401	3,331	3,033	2,845	2,496	2,344	1,926	2,244	125

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All races, female											
1998	32,932	1,561	1,257	1,093	1,056	1,330	1,893	2,528	3,099	6,643	8
1997	31,949	1,464	1,212	1,065	1,039	1,277	1,759	2,395	2,956	6,272	9
1996	31,649	1,447	1,118	979	1,044	1,280	1,835	2,289	2,922	6,103	17
1995	30,336	1,253	1,066	930	1,049	1,248	1,821	2,233	2,639	5,646	4
1994	29,399	1,269	1,013	882	965	1,320	1,701	2,155	2,686	5,219	10
1993	28,825	1,136	879	831	1,045	1,355	1,574	2,176	2,601	5,169	12
1992	27,398	1,086	819	844	970	1,292	1,568	2,088	2,421	4,841	6
1991	28,133	1,025	824	818	1,036	1,310	1,600	2,088	2,451	4,759	12
1990	28,542	1,029	909	950	1,170	1,417	1,585	2,183	2,364	4,490	12
1989	29,556	999	915	931	1,165	1,419	1,743	2,190	2,386	4,410	13
1988	29,695	931	965	1,001	1,197	1,442	1,725	2,142	2,425	4,445	11
1987	28,646	933	863	1,036	1,215	1,429	1,636	2,081	2,301	4,019	16
1986	28,075	837	846	980	1,209	1,320	1,697	2,010	2,267	3,884	19
1985	27,850	882	934	1,024	1,269	1,369	1,782	2,000	2,198	3,812	9
1984	27,507	928	915	1,062	1,226	1,296	1,628	1,973	2,148	3,757	9
1983	27,223	857	972	1,062	1,187	1,319	1,621	1,887	2,117	3,731	17
1982	27,047	890	1,005	1,107	1,165	1,286	1,602	1,777	2,045	3,540	21
1981	28,529	964	1,120	1,237	1,208	1,307	1,626	1,772	2,121	3,632	12
1980	30,336	1,025	1,188	1,285	1,309	1,435	1,711	2,076	2,230	3,778	11
1979	29,622	1,036	1,168	1,271	1,266	1,386	1,657	1,927	2,135	3,444	18
Accidents (unintentional injuries) (E800-E869,E880-E929)											
White, both sexes											
1998	79,297	4,823	3,683	3,030	2,713	3,074	3,998	4,966	5,777	10,018	98
1997	77,192	4,611	3,506	2,893	2,633	2,995	3,771	4,739	5,424	9,477	84
1996	76,818	4,469	3,265	2,723	2,692	3,031	3,909	4,487	5,322	9,202	98
1995	75,326	4,076	3,163	2,594	2,651	2,956	3,706	4,385	4,792	8,613	95
1994	73,611	3,921	2,905	2,444	2,529	2,999	3,596	4,309	4,883	7,996	83
1993	72,812	3,602	2,716	2,372	2,622	2,918	3,469	4,249	4,730	7,842	91
1992	70,072	3,273	2,603	2,367	2,579	3,045	3,441	4,021	4,472	7,273	102
1991	72,223	3,203	2,478	2,408	2,760	3,067	3,403	4,105	4,414	7,107	105
1990	74,558	3,108	2,726	2,650	3,069	3,249	3,427	4,119	4,154	6,731	97
1989	76,672	3,155	2,712	2,803	3,077	3,319	3,646	4,192	4,291	6,611	98
1988	78,574	3,125	2,784	2,841	3,135	3,422	3,603	4,046	4,316	6,791	100
1987	77,299	2,885	2,598	2,910	3,282	3,249	3,541	4,055	4,158	6,177	102
1986	77,762	2,802	2,728	2,856	3,184	3,140	3,566	3,939	3,983	5,987	135
1985	76,628	2,834	2,751	3,167	3,306	3,152	3,605	3,960	3,888	5,948	114
1984	77,030	2,935	2,917	3,121	3,319	3,260	3,481	3,932	3,774	5,843	82
1983	76,392	2,807	3,023	3,192	3,253	3,140	3,364	3,647	3,675	5,803	92
1982	78,249	3,007	3,143	3,338	3,249	3,166	3,333	3,561	3,596	5,497	80
1981	84,310	3,184	3,658	3,684	3,475	3,319	3,557	3,594	3,701	5,667	90
1980	88,069	3,422	3,714	3,836	3,538	3,534	3,594	3,884	3,814	5,766	80
1979	88,116	3,478	3,767	3,895	3,671	3,621	3,649	3,798	3,777	5,294	118

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
White, male											
1998	51,026	3,595	2,646	2,118	1,824	1,939	2,312	2,683	2,898	3,754	92
1997	50,096	3,467	2,523	2,030	1,763	1,913	2,243	2,595	2,703	3,576	80
1996	49,863	3,329	2,324	1,914	1,808	1,943	2,301	2,434	2,625	3,451	85
1995	49,555	3,079	2,308	1,819	1,790	1,880	2,103	2,362	2,381	3,291	91
1994	48,776	2,927	2,088	1,713	1,722	1,866	2,098	2,363	2,409	3,095	75
1993	48,387	2,657	1,990	1,674	1,743	1,758	2,066	2,281	2,351	3,015	83
1992	46,810	2,383	1,942	1,684	1,769	1,929	2,044	2,158	2,236	2,722	98
1991	48,235	2,380	1,803	1,734	1,877	1,931	1,989	2,219	2,125	2,679	96
1990	50,235	2,273	1,968	1,862	2,081	2,036	2,034	2,140	2,012	2,533	89
1989	51,511	2,325	1,959	2,035	2,084	2,106	2,102	2,217	2,083	2,545	90
1988	53,226	2,359	1,977	1,994	2,111	2,146	2,086	2,122	2,076	2,639	92
1987	52,702	2,115	1,896	2,034	2,231	2,034	2,096	2,167	2,054	2,432	89
1986	53,678	2,103	2,039	2,024	2,186	1,964	2,079	2,118	1,888	2,371	123
1985	52,691	2,107	1,983	2,304	2,220	1,958	2,031	2,164	1,877	2,395	107
1984	53,153	2,157	2,152	2,225	2,239	2,126	2,042	2,138	1,808	2,322	75
1983	52,992	2,114	2,225	2,296	2,241	2,011	1,947	1,951	1,758	2,292	81
1982	54,820	2,253	2,306	2,397	2,224	2,043	1,917	1,950	1,691	2,148	65
1981	59,449	2,379	2,705	2,612	2,428	2,174	2,102	1,978	1,746	2,244	81
1980	61,929	2,561	2,715	2,747	2,421	2,294	2,133	2,072	1,788	2,234	75
1979	62,323	2,598	2,782	2,784	2,570	2,419	2,158	2,058	1,760	2,063	102
Accidents (unintentional injuries) (E800-E869,E880-E929)											
White, female											
1998	28,271	1,228	1,037	912	889	1,135	1,686	2,283	2,879	6,264	6
1997	27,096	1,144	983	863	870	1,082	1,528	2,144	2,721	5,901	4
1996	26,955	1,140	941	809	884	1,088	1,608	2,053	2,697	5,751	13
1995	25,771	997	855	775	861	1,076	1,603	2,023	2,411	5,322	4
1994	24,835	994	817	731	807	1,133	1,498	1,946	2,474	4,901	8
1993	24,425	945	726	698	879	1,160	1,403	1,968	2,379	4,827	8
1992	23,262	890	661	683	810	1,116	1,397	1,863	2,236	4,551	4
1991	23,988	823	675	674	883	1,136	1,414	1,886	2,289	4,428	9
1990	24,323	835	758	788	988	1,213	1,393	1,979	2,142	4,198	8
1989	25,161	830	753	768	993	1,213	1,544	1,975	2,208	4,066	8
1988	25,348	766	807	847	1,024	1,276	1,517	1,924	2,240	4,152	8
1987	24,597	770	702	876	1,051	1,215	1,445	1,888	2,104	3,745	13
1986	24,084	699	689	832	998	1,176	1,487	1,821	2,095	3,616	12
1985	23,937	727	768	863	1,086	1,194	1,574	1,796	2,011	3,553	7
1984	23,877	778	765	896	1,080	1,134	1,439	1,794	1,966	3,521	7
1983	23,400	693	798	896	1,012	1,129	1,417	1,696	1,917	3,511	11
1982	23,429	754	837	941	1,025	1,123	1,416	1,611	1,905	3,349	15
1981	24,861	805	953	1,072	1,047	1,145	1,455	1,616	1,955	3,423	9
1980	26,140	861	999	1,089	1,117	1,240	1,461	1,812	2,026	3,532	5
1979	25,793	880	985	1,111	1,101	1,202	1,491	1,740	2,017	3,231	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All other, both sexes											
1998	15,034	1,267	867	639	546	520	534	539	468	711	11
1997	15,161	1,186	814	630	570	533	571	554	463	661	16
1996	14,958	1,128	734	586	524	571	566	529	462	618	18
1995	15,076	1,078	709	587	586	529	562	503	484	611	9
1994	15,038	971	746	596	549	499	540	523	487	583	16
1993	14,786	845	662	521	546	545	492	466	439	644	17
1992	13,880	802	627	581	548	514	495	501	395	535	15
1991	14,488	724	589	554	528	566	480	460	406	590	14
1990	14,590	749	625	555	615	585	484	500	428	532	16
1989	15,364	790	689	642	648	624	515	526	410	612	24
1988	15,488	748	679	623	657	601	596	553	423	557	32
1987	14,689	712	613	607	620	591	541	471	427	523	25
1986	14,446	673	605	653	607	490	516	505	392	510	30
1985	13,977	688	631	648	641	526	557	513	411	495	22
1984	13,253	644	574	652	542	527	543	451	388	464	17
1983	13,436	667	658	666	621	592	553	476	425	412	26
1982	13,248	657	680	667	581	519	527	434	328	370	16
1981	13,845	729	735	691	614	575	531	413	357	393	26
1980	15,212	748	833	782	698	678	594	601	418	441	21
1979	14,554	726	802	707	628	610	504	473	284	394	25
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All other, male											
1998	10,373	934	647	458	379	325	327	294	248	332	9
1997	10,308	866	585	428	401	338	340	303	228	290	11
1996	10,264	821	557	416	364	379	339	293	237	266	14
1995	10,511	822	498	432	398	357	344	293	256	287	9
1994	10,474	696	550	445	391	312	337	314	275	265	14
1993	10,386	654	509	388	380	350	321	258	217	302	13
1992	9,744	606	469	420	388	338	324	276	210	245	13
1991	10,343	522	440	410	375	392	294	258	244	259	11
1990	10,371	555	474	393	433	381	292	296	206	240	12
1989	10,969	621	527	479	476	418	316	311	232	268	19
1988	11,141	583	521	469	484	435	388	335	238	264	29
1987	10,640	549	452	447	456	377	350	278	230	249	22
1986	10,455	535	448	505	396	346	306	316	220	242	23
1985	10,064	533	465	487	458	351	349	309	224	236	20
1984	9,623	494	424	486	396	365	354	272	206	228	15
1983	9,613	503	484	500	446	402	349	285	225	192	20
1982	9,630	521	512	501	441	356	341	268	188	179	10
1981	10,177	570	568	526	453	413	360	257	191	184	23
1980	11,016	584	644	586	506	483	344	337	214	195	15
1979	10,725	570	619	547	463	426	338	286	166	181	23

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
All other, female											
1998	4,661	333	220	181	167	195	207	245	220	379	2
1997	4,853	320	229	202	169	195	231	251	235	371	5
1996	4,694	307	177	170	160	192	227	236	225	352	4
1995	4,565	256	211	155	188	172	218	210	228	324	-
1994	4,564	275	196	151	158	187	203	209	212	318	2
1993	4,400	191	153	133	166	195	171	208	222	342	4
1992	4,136	196	158	161	160	176	171	225	185	290	2
1991	4,145	202	149	144	153	174	186	202	162	331	3
1990	4,219	194	151	162	182	204	192	204	222	292	4
1989	4,395	169	162	163	172	206	199	215	178	344	5
1988	4,347	165	158	154	173	166	208	218	185	293	3
1987	4,049	163	161	160	164	214	191	193	197	274	3
1986	3,991	138	157	148	211	144	210	189	172	268	7
1985	3,913	155	166	161	183	175	208	204	187	259	2
1984	3,630	150	150	166	146	162	189	179	182	236	2
1983	3,823	164	174	166	175	190	204	191	200	220	6
1982	3,618	136	168	166	140	163	186	166	140	191	6
1981	3,668	159	167	165	161	162	171	156	166	209	3
1980	4,196	164	189	196	192	195	250	264	204	246	6
1979	3,829	156	183	160	165	184	166	187	118	213	2
Accidents (unintentional injuries) (E800-E869,E880-E929)											
Black, both sexes											
1998	12,237	1,080	717	536	437	416	431	449	361	576	11
1997	12,161	1,009	673	515	457	435	430	436	355	529	14
1996	12,131	957	616	478	434	466	453	426	365	501	13
1995	12,296	900	591	477	488	427	481	420	401	509	9
1994	12,316	820	615	476	451	413	438	422	395	477	11
1993	12,246	716	553	429	452	466	408	386	352	538	14
1992	11,355	665	517	483	465	426	402	421	327	440	15
1991	12,066	591	491	465	438	473	397	405	343	506	13
1990	11,998	627	517	462	536	487	413	426	349	452	16
1989	12,907	663	593	553	574	552	459	463	346	538	20
1988	13,007	644	602	536	562	504	515	477	351	479	29
1987	12,292	607	541	516	536	505	454	406	357	445	17
1986	12,114	570	511	568	542	435	459	436	347	434	24
1985	11,779	595	543	575	579	467	486	452	368	428	20
1984	11,278	544	487	586	475	473	482	401	345	407	14
1983	11,469	568	565	597	562	530	483	424	385	364	21
1982	11,271	571	580	583	500	455	469	389	286	322	15
1981	11,919	644	653	632	562	510	490	371	321	350	21
1980	13,121	668	752	709	638	617	538	546	375	390	20
1979	12,582	657	714	644	551	557	459	423	257	350	23

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidents (unintentional injuries) (E800-E869,E880-E929)											
Black, male											
1998	8,544	810	544	391	311	275	266	239	194	255	9
1997	8,359	740	495	357	324	285	256	241	176	215	9
1996	8,409	725	483	344	316	325	282	232	181	210	9
1995	8,639	705	429	351	338	296	294	247	212	225	9
1994	8,632	603	455	369	328	266	270	255	219	203	10
1993	8,714	568	444	323	320	316	272	209	166	252	10
1992	8,021	514	399	354	338	290	263	237	172	187	13
1991	8,659	432	367	353	319	326	244	229	194	216	11
1990	8,561	476	401	333	385	321	255	254	156	190	12
1989	9,265	525	468	414	427	377	279	279	188	230	15
1988	9,381	502	467	412	415	367	332	291	190	218	26
1987	8,938	476	408	380	403	326	297	240	190	212	15
1986	8,808	456	385	451	354	308	272	280	190	200	20
1985	8,535	466	407	440	416	318	307	261	199	198	18
1984	8,211	425	363	442	354	335	312	238	179	196	13
1983	8,192	439	415	452	402	356	300	247	201	170	15
1982	8,235	458	445	447	374	317	305	238	155	148	10
1981	8,820	513	510	481	416	364	331	229	168	161	18
1980	9,516	529	589	535	463	437	302	293	191	166	14
1979	9,288	524	563	495	404	391	309	251	148	156	21
Accidents (unintentional injuries) (E800-E869,E880-E929)											
Black, female											
1998	3,693	270	173	145	126	141	165	210	167	321	2
1997	3,802	269	178	158	133	150	174	195	179	314	5
1996	3,722	232	133	134	118	141	171	194	184	291	4
1995	3,657	195	162	126	150	131	187	173	189	284	-
1994	3,684	217	160	107	123	147	168	167	176	274	1
1993	3,532	148	109	106	132	150	136	177	186	286	4
1992	3,334	151	118	129	127	136	139	184	155	253	2
1991	3,407	159	124	112	119	147	153	176	149	290	2
1990	3,437	151	116	129	151	166	158	172	193	262	4
1989	3,642	138	125	139	147	175	180	184	158	308	5
1988	3,626	142	135	124	147	137	183	186	161	261	3
1987	3,354	131	133	136	133	179	157	166	167	233	2
1986	3,306	114	126	117	188	127	187	156	157	234	4
1985	3,244	129	136	135	163	149	179	191	169	230	2
1984	3,067	119	124	144	121	138	170	163	166	211	1
1983	3,277	129	150	145	160	174	183	177	184	194	6
1982	3,036	113	135	136	126	138	164	151	131	174	5
1981	3,099	131	143	151	146	146	159	142	153	189	3
1980	3,605	139	163	174	175	180	236	253	184	224	6
1979	3,294	133	151	149	147	166	150	172	109	194	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
All races, both sexes											
1998	46,086	2,990	2,449	2,023	1,707	1,733	1,898	1,928	1,650	1,351	37
1997	46,287	2,928	2,397	1,956	1,661	1,711	1,862	1,904	1,603	1,326	28
1996	46,679	2,879	2,205	1,823	1,693	1,726	1,886	1,809	1,552	1,191	55
1995	46,216	2,703	2,098	1,710	1,667	1,676	1,824	1,818	1,459	1,169	44
1994	45,645	2,607	2,074	1,653	1,550	1,747	1,840	1,799	1,502	1,064	42
1993	44,805	2,386	1,856	1,573	1,609	1,612	1,684	1,815	1,478	1,040	43
1992	44,178	2,333	1,844	1,609	1,564	1,679	1,742	1,641	1,368	934	40
1991	46,629	2,321	1,749	1,608	1,666	1,729	1,652	1,716	1,310	951	50
1990	49,917	2,377	1,982	1,791	1,881	1,852	1,706	1,732	1,272	968	52
1989	50,731	2,328	2,015	1,851	1,904	1,838	1,828	1,682	1,313	913	49
1988	52,120	2,271	1,969	1,858	1,844	1,853	1,763	1,619	1,322	888	50
1987	51,642	2,144	1,852	1,854	1,939	1,793	1,666	1,575	1,230	805	55
1986	51,139	2,024	1,872	1,855	1,836	1,622	1,640	1,557	1,129	716	72
1985	49,445	2,036	1,861	1,928	1,858	1,577	1,646	1,448	1,059	719	45
1984	49,627	2,069	1,871	1,955	1,888	1,638	1,555	1,483	1,013	685	39
1983	48,050	1,958	1,929	1,848	1,754	1,560	1,463	1,299	940	653	41
1982	49,696	2,075	1,986	1,962	1,752	1,521	1,458	1,259	893	594	43
1981	55,190	2,254	2,307	2,274	1,891	1,689	1,593	1,313	930	625	54
1980	57,162	2,375	2,382	2,284	1,837	1,634	1,494	1,350	893	640	31
1979	57,757	2,375	2,448	2,285	2,022	1,794	1,553	1,353	894	560	61
Transport accidents (E800-E848,E929.0,E929.1)											
All races, male											
1998	31,107	2,084	1,729	1,360	1,134	1,067	1,062	1,054	919	754	34
1997	31,106	2,053	1,670	1,285	1,069	1,027	1,105	1,049	866	698	24
1996	31,518	2,012	1,522	1,236	1,067	1,065	1,077	988	827	663	47
1995	31,526	1,926	1,449	1,156	1,076	1,021	1,014	991	802	643	42
1994	31,215	1,831	1,415	1,120	1,011	1,050	1,031	981	807	610	37
1993	30,914	1,703	1,298	1,082	1,040	929	972	979	802	606	38
1992	30,656	1,616	1,326	1,094	1,020	1,009	1,008	892	761	531	39
1991	32,536	1,653	1,229	1,122	1,081	1,047	943	945	718	543	48
1990	35,046	1,696	1,378	1,209	1,221	1,105	978	920	687	557	49
1989	35,339	1,664	1,423	1,306	1,243	1,096	1,021	917	727	535	45
1988	36,863	1,654	1,360	1,232	1,219	1,154	1,030	874	737	516	44
1987	36,715	1,524	1,301	1,261	1,292	1,103	978	844	708	474	48
1986	36,879	1,487	1,342	1,274	1,224	975	916	890	647	426	62
1985	35,407	1,480	1,300	1,385	1,187	941	909	820	589	430	44
1984	35,824	1,500	1,357	1,344	1,259	1,053	923	839	562	422	35
1983	34,989	1,451	1,387	1,302	1,176	991	860	738	552	401	36
1982	36,529	1,554	1,463	1,396	1,183	966	854	737	532	356	32
1981	40,813	1,694	1,706	1,601	1,276	1,109	970	792	555	386	49
1980	42,198	1,774	1,716	1,602	1,265	1,070	888	803	519	394	27
1979	42,873	1,776	1,810	1,649	1,419	1,201	945	806	539	348	50

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
All races, female											
1998	14,979	906	720	663	573	666	836	874	731	597	3
1997	15,181	875	727	671	592	684	757	855	737	628	4
1996	15,161	867	683	587	626	661	809	821	725	528	8
1995	14,690	777	649	554	591	655	810	827	657	526	2
1994	14,430	776	659	533	539	697	809	818	695	454	5
1993	13,891	683	558	491	569	683	712	836	676	434	5
1992	13,522	717	518	515	544	670	734	749	607	403	1
1991	14,093	668	520	486	585	682	709	771	592	408	2
1990	14,871	681	604	582	660	747	728	812	585	411	3
1989	15,392	664	592	545	661	742	807	765	586	378	4
1988	15,257	617	609	626	625	699	733	745	585	372	6
1987	14,927	620	551	593	647	690	688	731	522	331	7
1986	14,260	537	530	581	612	647	724	667	482	290	10
1985	14,038	556	561	543	671	636	737	628	470	289	1
1984	13,803	569	514	611	629	585	632	644	451	263	4
1983	13,061	507	542	546	578	569	603	561	388	252	5
1982	13,167	521	523	566	569	555	604	522	361	238	11
1981	14,377	560	601	673	615	580	623	521	375	239	5
1980	14,964	601	666	682	572	564	606	547	374	246	4
1979	14,884	599	638	636	603	593	608	547	355	212	11
Transport accidents (E800-E848,E929.0,E929.1)											
White, both sexes											
1998	38,433	2,434	2,043	1,691	1,451	1,494	1,683	1,728	1,527	1,257	33
1997	38,510	2,410	2,000	1,633	1,400	1,480	1,613	1,720	1,478	1,231	23
1996	39,098	2,405	1,861	1,551	1,465	1,482	1,665	1,632	1,423	1,110	46
1995	38,730	2,232	1,777	1,445	1,407	1,470	1,606	1,652	1,344	1,091	42
1994	38,321	2,194	1,710	1,378	1,302	1,527	1,602	1,604	1,382	989	36
1993	37,723	2,009	1,538	1,326	1,358	1,390	1,509	1,650	1,374	936	34
1992	37,261	1,937	1,532	1,328	1,305	1,469	1,543	1,495	1,272	879	33
1991	39,552	1,955	1,475	1,358	1,429	1,489	1,484	1,586	1,225	875	45
1990	42,359	1,976	1,649	1,506	1,611	1,594	1,536	1,583	1,157	903	45
1989	43,250	1,949	1,674	1,561	1,631	1,611	1,654	1,533	1,218	854	43
1988	44,608	1,915	1,663	1,581	1,590	1,619	1,558	1,455	1,242	807	39
1987	44,486	1,789	1,548	1,590	1,696	1,560	1,496	1,449	1,148	749	49
1986	44,060	1,708	1,564	1,563	1,579	1,454	1,483	1,399	1,045	646	57
1985	42,830	1,733	1,580	1,657	1,599	1,393	1,478	1,309	992	673	43
1984	43,387	1,772	1,623	1,678	1,664	1,447	1,388	1,353	941	622	31
1983	42,072	1,647	1,636	1,602	1,540	1,371	1,311	1,163	869	615	32
1982	43,665	1,781	1,711	1,701	1,531	1,326	1,301	1,147	837	554	35
1981	48,850	1,919	1,990	1,992	1,675	1,491	1,400	1,197	861	591	43
1980	50,380	2,060	2,037	1,973	1,590	1,415	1,337	1,222	826	595	25
1979	51,057	2,050	2,101	1,967	1,768	1,577	1,393	1,233	840	515	53

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
White, male											
1998	25,822	1,700	1,453	1,129	971	928	938	937	838	692	30
1997	25,830	1,704	1,399	1,071	898	886	954	942	795	652	22
1996	26,299	1,714	1,269	1,047	920	911	943	885	752	612	40
1995	26,269	1,589	1,237	970	912	888	877	880	733	586	40
1994	26,163	1,563	1,165	924	846	916	877	857	730	561	32
1993	25,929	1,417	1,073	901	868	789	858	880	742	537	30
1992	25,763	1,337	1,104	904	846	873	875	799	705	491	33
1991	27,440	1,400	1,029	953	927	885	833	871	656	494	43
1990	29,581	1,410	1,133	1,025	1,035	946	874	830	615	510	43
1989	29,897	1,377	1,163	1,091	1,050	954	907	820	666	492	40
1988	31,368	1,387	1,135	1,038	1,037	986	880	768	683	456	34
1987	31,428	1,249	1,093	1,067	1,117	958	863	766	652	436	42
1986	31,607	1,247	1,120	1,060	1,053	860	810	778	585	374	51
1985	30,512	1,250	1,095	1,180	1,002	823	790	722	551	401	42
1984	31,122	1,271	1,173	1,143	1,089	914	805	748	516	375	28
1983	30,516	1,219	1,170	1,107	1,023	857	754	650	503	376	28
1982	32,020	1,320	1,254	1,200	1,011	823	744	655	487	328	27
1981	36,009	1,427	1,466	1,387	1,116	957	827	706	514	362	40
1980	37,039	1,532	1,450	1,359	1,081	895	771	704	476	366	21
1979	37,772	1,512	1,553	1,403	1,233	1,035	833	728	500	312	43
Transport accidents (E800-E848,E929.0,E929.1)											
White, female											
1998	12,611	734	590	562	480	566	745	791	689	565	3
1997	12,680	706	601	562	502	594	659	778	683	579	1
1996	12,799	691	592	504	545	571	722	747	671	498	6
1995	12,461	643	540	475	495	582	729	772	611	505	2
1994	12,158	631	545	454	456	611	725	747	652	428	4
1993	11,794	592	465	425	490	601	651	770	632	399	4
1992	11,498	600	428	424	459	596	668	696	567	388	-
1991	12,112	555	446	405	502	604	651	715	569	381	2
1990	12,778	566	516	481	576	648	662	753	542	393	2
1989	13,353	572	511	470	581	657	747	713	552	362	3
1988	13,240	528	528	543	553	633	678	687	559	351	5
1987	13,058	540	455	523	579	602	633	683	496	313	7
1986	12,453	461	444	503	526	594	673	621	460	272	6
1985	12,318	483	485	477	597	570	688	587	441	272	1
1984	12,265	501	450	535	575	533	583	605	425	247	3
1983	11,556	428	466	495	517	514	557	513	366	239	4
1982	11,645	461	457	501	520	503	557	492	350	226	8
1981	12,841	492	524	605	559	534	573	491	347	229	3
1980	13,341	528	587	614	509	520	566	518	350	229	4
1979	13,285	538	548	564	535	542	560	505	340	203	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
All other, both sexes											
1998	7,653	556	406	332	256	239	215	200	123	94	4
1997	7,777	518	397	323	261	231	249	184	125	95	5
1996	7,581	474	344	272	228	244	221	177	129	81	9
1995	7,486	471	321	265	260	206	218	166	115	78	2
1994	7,324	413	364	275	248	220	238	195	120	75	6
1993	7,082	377	318	247	251	222	175	165	104	104	9
1992	6,917	396	312	281	259	210	199	146	96	55	7
1991	7,077	366	274	250	237	240	168	130	85	76	5
1990	7,558	401	333	285	270	258	170	149	115	65	7
1989	7,481	379	341	290	273	227	174	149	95	59	6
1988	7,512	356	306	277	254	234	205	164	80	81	11
1987	7,156	355	304	264	243	233	170	126	82	56	6
1986	7,079	316	308	292	257	168	157	158	84	70	15
1985	6,615	303	281	271	259	184	168	139	67	46	2
1984	6,240	297	248	277	224	191	167	130	72	63	8
1983	5,978	311	293	246	214	189	152	136	71	38	9
1982	6,031	294	275	261	221	195	157	112	56	40	8
1981	6,340	335	317	282	216	198	193	116	69	34	11
1980	6,782	315	345	311	247	219	157	128	67	45	6
1979	6,700	325	347	318	254	217	160	120	54	45	8
Transport accidents (E800-E848,E929.0,E929.1)											
All other, male											
1998	5,285	384	276	231	163	139	124	117	81	62	4
1997	5,276	349	271	214	171	141	151	107	71	46	2
1996	5,219	298	253	189	147	154	134	103	75	51	7
1995	5,257	337	212	186	164	133	137	111	69	57	2
1994	5,052	268	250	196	165	134	154	124	77	49	5
1993	4,985	286	225	181	172	140	114	99	60	69	8
1992	4,893	279	222	190	174	136	133	93	56	40	6
1991	5,096	253	200	169	154	162	110	74	62	49	5
1990	5,465	286	245	184	186	159	104	90	72	47	6
1989	5,442	287	260	215	193	142	114	97	61	43	5
1988	5,495	267	225	194	182	168	150	106	54	60	10
1987	5,287	275	208	194	175	145	115	78	56	38	6
1986	5,272	240	222	214	171	115	106	112	62	52	11
1985	4,895	230	205	205	185	118	119	98	38	29	2
1984	4,702	229	184	201	170	139	118	91	46	47	7
1983	4,473	232	217	195	153	134	106	88	49	25	8
1982	4,509	234	209	196	172	143	110	82	45	28	5
1981	4,804	267	240	214	160	152	143	86	41	24	9
1980	5,159	242	266	243	184	175	117	99	43	28	6
1979	5,101	264	257	246	186	166	112	78	39	36	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
All other, female											
1998	2,368	172	130	101	93	100	91	83	42	32	-
1997	2,501	169	126	109	90	90	98	77	54	49	3
1996	2,362	176	91	83	81	90	87	74	54	30	2
1995	2,229	134	109	79	96	73	81	55	46	21	-
1994	2,272	145	114	79	83	86	84	71	43	26	1
1993	2,097	91	93	66	79	82	61	66	44	35	1
1992	2,024	117	90	91	85	74	66	53	40	15	1
1991	1,981	113	74	81	83	78	58	56	23	27	-
1990	2,093	115	88	101	84	99	66	59	43	18	1
1989	2,039	92	81	75	80	85	60	52	34	16	1
1988	2,017	89	81	83	72	66	55	58	26	21	1
1987	1,869	80	96	70	68	88	55	48	26	18	-
1986	1,807	76	86	78	86	53	51	46	22	18	4
1985	1,720	73	76	66	74	66	49	41	29	17	-
1984	1,538	68	64	76	54	52	49	39	26	16	1
1983	1,505	79	76	51	61	55	46	48	22	13	1
1982	1,522	60	66	65	49	52	47	30	11	12	3
1981	1,536	68	77	68	56	46	50	30	28	10	2
1980	1,623	73	79	68	63	44	40	29	24	17	-
1979	1,599	61	90	72	68	51	48	42	15	9	1
Transport accidents (E800-E848,E929.0,E929.1)											
Black, both sexes											
1998	5,928	434	311	257	196	185	170	157	86	68	4
1997	5,951	415	303	254	195	173	171	128	80	76	5
1996	5,827	369	268	206	173	185	162	135	95	67	7
1995	5,713	355	240	196	198	149	175	122	85	62	2
1994	5,657	321	277	203	187	167	176	146	90	61	4
1993	5,511	298	238	188	193	181	136	125	73	77	7
1992	5,345	306	240	215	207	159	146	117	73	38	7
1991	5,518	276	208	195	189	188	122	107	63	62	5
1990	5,860	313	258	220	228	205	140	118	84	46	7
1989	5,908	304	282	231	228	185	148	119	72	46	5
1988	5,940	283	256	231	206	183	164	130	63	67	8
1987	5,635	281	251	219	198	187	135	98	59	41	4
1986	5,593	248	244	244	220	141	129	125	63	58	12
1985	5,223	249	223	225	223	152	130	120	55	37	2
1984	5,053	234	192	236	188	172	143	108	63	52	6
1983	4,780	250	247	211	188	154	121	114	59	29	9
1982	4,867	237	231	213	181	158	129	95	41	32	7
1981	5,121	277	268	248	190	159	173	97	60	28	7
1980	5,469	265	299	267	216	187	139	103	50	33	6
1979	5,470	286	301	286	216	189	141	101	44	40	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Transport accidents (E800-E848,E929.0,E929.1)											
Black, male											
1998	4,203	307	218	180	132	118	101	95	61	45	4
1997	4,142	283	217	176	132	114	109	72	50	35	2
1996	4,109	249	210	147	121	126	104	81	60	43	5
1995	4,114	268	167	138	127	104	110	90	49	44	2
1994	3,969	221	189	150	128	106	116	94	59	38	3
1993	4,013	238	179	141	138	126	100	72	46	55	6
1992	3,849	230	181	147	145	111	101	80	42	26	6
1991	4,035	196	153	138	130	128	81	63	44	43	5
1990	4,311	232	198	146	162	132	89	74	52	32	6
1989	4,393	231	229	174	169	121	98	83	45	35	4
1988	4,418	214	194	170	151	135	125	87	41	49	7
1987	4,223	225	176	161	152	123	94	68	41	30	4
1986	4,248	193	178	190	148	95	89	95	49	42	9
1985	3,932	191	170	177	161	103	94	82	29	24	2
1984	3,865	191	148	174	152	129	104	74	41	38	5
1983	3,625	194	189	169	137	108	86	71	41	19	8
1982	3,718	196	180	166	140	119	93	70	32	21	5
1981	3,982	228	209	191	142	124	128	74	37	21	5
1980	4,252	211	233	216	165	151	105	76	36	20	6
1979	4,225	241	233	222	158	143	101	64	33	32	5
Transport accidents (E800-E848,E929.0,E929.1)											
Black, female											
1998	1,725	127	93	77	64	67	69	62	25	23	-
1997	1,809	132	86	78	63	59	62	56	30	41	3
1996	1,718	120	58	59	52	59	58	54	35	24	2
1995	1,599	87	73	58	71	45	65	32	36	18	-
1994	1,688	100	88	53	59	61	60	52	31	23	1
1993	1,498	60	59	47	55	55	36	53	27	22	1
1992	1,496	76	59	68	62	48	45	37	31	12	1
1991	1,483	80	55	57	59	60	41	44	19	19	-
1990	1,549	81	60	74	66	73	51	44	32	14	1
1989	1,515	73	53	57	59	64	50	36	27	11	1
1988	1,522	69	62	61	55	48	39	43	22	18	1
1987	1,412	56	75	58	46	64	41	30	18	11	-
1986	1,345	55	66	54	72	46	40	30	14	16	3
1985	1,291	58	53	48	62	49	36	38	26	13	-
1984	1,188	43	44	62	36	43	39	34	22	14	1
1983	1,155	56	58	42	51	46	35	43	18	10	1
1982	1,149	41	51	47	41	39	36	25	9	11	2
1981	1,139	49	59	57	48	35	45	23	23	7	2
1980	1,217	54	66	51	51	36	34	27	14	13	-
1979	1,245	45	68	64	58	46	40	37	11	8	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
All races, both sexes											
1998	43,501	2,762	2,234	1,855	1,565	1,625	1,785	1,867	1,590	1,278	30
1997	43,458	2,643	2,162	1,776	1,553	1,596	1,774	1,840	1,552	1,265	22
1996	43,649	2,586	1,967	1,650	1,546	1,623	1,796	1,724	1,509	1,132	43
1995	43,363	2,439	1,894	1,562	1,533	1,562	1,738	1,757	1,428	1,141	32
1994	42,524	2,321	1,853	1,491	1,432	1,648	1,737	1,734	1,457	1,026	34
1993	41,893	2,134	1,669	1,432	1,476	1,509	1,607	1,758	1,432	1,015	34
1992	40,982	2,073	1,648	1,440	1,436	1,569	1,678	1,580	1,333	893	32
1991	43,536	2,053	1,579	1,445	1,539	1,615	1,578	1,646	1,286	919	43
1990	46,814	2,135	1,782	1,640	1,725	1,730	1,620	1,666	1,251	943	45
1989	47,575	2,069	1,810	1,675	1,733	1,716	1,749	1,632	1,277	877	38
1988	49,078	2,041	1,784	1,712	1,708	1,741	1,688	1,573	1,299	858	45
1987	48,290	1,911	1,665	1,676	1,769	1,680	1,597	1,528	1,198	778	49
1986	47,865	1,781	1,661	1,691	1,666	1,519	1,577	1,512	1,100	702	47
1985	45,901	1,773	1,650	1,729	1,733	1,452	1,562	1,407	1,029	706	35
1984	46,263	1,811	1,649	1,760	1,734	1,537	1,483	1,435	993	655	31
1983	44,452	1,730	1,700	1,648	1,612	1,445	1,382	1,260	921	641	27
1982	45,779	1,796	1,728	1,771	1,584	1,449	1,376	1,205	869	578	29
1981	51,385	1,991	2,030	2,062	1,735	1,567	1,523	1,272	910	607	47
1980	53,172	2,106	2,129	2,080	1,707	1,553	1,438	1,306	862	619	26
1979	53,524	2,093	2,164	2,063	1,842	1,699	1,472	1,306	873	537	35
Motor vehicle accidents (E810-E825)											
All races, male											
1998	28,990	1,893	1,546	1,211	1,008	967	971	1,010	879	727	27
1997	28,770	1,805	1,469	1,130	971	929	1,026	1,004	833	672	18
1996	29,123	1,785	1,319	1,096	945	980	1,006	935	803	631	36
1995	29,164	1,700	1,281	1,029	964	927	937	942	779	629	30
1994	28,635	1,593	1,232	977	908	964	950	933	771	584	30
1993	28,531	1,500	1,133	964	923	850	906	938	776	594	29
1992	27,982	1,399	1,151	947	912	919	957	852	736	506	31
1991	29,947	1,416	1,086	975	981	955	880	891	706	524	41
1990	32,406	1,487	1,202	1,075	1,088	1,007	910	871	673	543	42
1989	32,704	1,445	1,237	1,153	1,103	991	954	884	707	515	34
1988	34,258	1,455	1,194	1,109	1,099	1,060	964	838	721	495	39
1987	33,904	1,327	1,143	1,109	1,144	1,008	920	805	685	460	42
1986	34,088	1,281	1,149	1,128	1,069	884	863	857	627	415	42
1985	32,443	1,258	1,130	1,217	1,077	839	846	793	568	421	34
1984	32,949	1,280	1,166	1,168	1,125	964	858	801	544	403	27
1983	31,907	1,246	1,183	1,127	1,049	885	795	713	538	394	24
1982	33,191	1,310	1,237	1,233	1,037	902	789	695	518	346	21
1981	37,533	1,465	1,456	1,417	1,142	1,007	911	757	540	373	42
1980	38,861	1,540	1,496	1,424	1,151	999	847	767	502	382	23
1979	39,309	1,526	1,560	1,456	1,259	1,119	882	774	523	331	33

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
All races, female											
1998	14,511	869	688	644	557	658	814	857	711	551	3
1997	14,688	838	693	646	582	667	748	836	719	593	4
1996	14,526	801	648	554	601	643	790	789	706	501	7
1995	14,199	739	613	533	569	635	801	815	649	512	2
1994	13,889	728	621	514	524	684	787	801	686	442	4
1993	13,362	634	536	468	553	659	701	820	656	421	5
1992	13,000	674	497	493	524	650	721	728	597	387	1
1991	13,589	637	493	470	558	660	698	755	580	395	2
1990	14,408	648	580	565	637	723	710	795	578	400	3
1989	14,871	624	573	522	630	725	795	748	570	362	4
1988	14,820	586	590	603	609	681	724	735	578	363	6
1987	14,386	584	522	567	625	672	677	723	513	318	7
1986	13,777	500	512	563	597	635	714	655	473	287	5
1985	13,458	515	520	512	656	613	716	614	461	285	1
1984	13,314	531	483	592	609	573	625	634	449	252	4
1983	12,545	484	517	521	563	560	587	547	383	247	3
1982	12,588	486	491	538	547	547	587	510	351	232	8
1981	13,852	526	574	645	593	560	612	515	370	234	5
1980	14,311	566	633	656	556	554	591	539	360	237	3
1979	14,215	567	604	607	583	580	590	532	350	206	2
Motor vehicle accidents (E810-E825)											
White, both sexes											
1998	36,182	2,239	1,848	1,545	1,325	1,397	1,578	1,672	1,470	1,190	26
1997	36,094	2,164	1,787	1,474	1,304	1,377	1,541	1,666	1,431	1,173	18
1996	36,452	2,142	1,649	1,400	1,330	1,397	1,586	1,557	1,387	1,055	36
1995	36,255	1,993	1,588	1,317	1,296	1,360	1,524	1,600	1,316	1,066	30
1994	35,619	1,943	1,512	1,234	1,196	1,436	1,510	1,550	1,342	952	29
1993	35,194	1,786	1,372	1,204	1,242	1,298	1,439	1,599	1,331	914	28
1992	34,439	1,703	1,355	1,173	1,191	1,366	1,486	1,440	1,241	840	25
1991	36,851	1,706	1,322	1,211	1,315	1,384	1,417	1,517	1,201	845	38
1990	39,651	1,755	1,473	1,368	1,470	1,489	1,457	1,529	1,139	878	40
1989	40,491	1,717	1,498	1,407	1,477	1,503	1,579	1,486	1,187	819	34
1988	41,974	1,715	1,503	1,455	1,467	1,518	1,494	1,414	1,222	781	35
1987	41,581	1,583	1,390	1,429	1,540	1,459	1,434	1,409	1,123	724	43
1986	41,189	1,488	1,373	1,420	1,429	1,358	1,429	1,357	1,019	633	37
1985	39,689	1,499	1,392	1,474	1,481	1,279	1,398	1,270	964	660	34
1984	40,429	1,535	1,423	1,508	1,525	1,359	1,324	1,307	923	595	25
1983	38,864	1,442	1,433	1,417	1,413	1,271	1,239	1,130	855	604	21
1982	40,191	1,523	1,470	1,527	1,379	1,265	1,232	1,095	817	538	23
1981	45,424	1,679	1,743	1,798	1,534	1,382	1,333	1,160	844	574	36
1980	46,863	1,818	1,808	1,792	1,480	1,345	1,296	1,182	799	576	21
1979	47,248	1,797	1,835	1,763	1,606	1,489	1,323	1,194	821	496	30

Source: CDC/NCHS, National Vital Statistics System
See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
White, male											
1998	23,985	1,537	1,287	999	859	838	852	897	799	668	23
1997	23,834	1,491	1,216	935	812	797	890	904	765	629	17
1996	24,214	1,510	1,089	923	807	842	880	838	733	583	30
1995	24,229	1,386	1,083	858	819	797	804	837	712	574	28
1994	23,932	1,355	1,000	798	753	837	802	820	699	536	26
1993	23,874	1,239	928	799	767	718	799	845	718	526	24
1992	23,415	1,142	945	769	747	788	831	762	682	468	25
1991	25,183	1,180	900	820	838	800	777	817	644	476	36
1990	27,288	1,220	976	903	916	864	811	792	603	496	38
1989	27,621	1,181	1,004	958	926	862	844	790	649	472	31
1988	29,127	1,216	990	933	930	902	824	736	670	438	30
1987	29,017	1,075	959	931	982	874	811	734	636	424	36
1986	29,163	1,062	944	935	918	774	765	747	568	364	34
1985	27,894	1,055	945	1,025	899	728	731	696	532	392	33
1984	28,614	1,070	1,002	991	968	837	747	712	500	358	22
1983	27,781	1,035	991	946	909	765	696	630	494	370	19
1982	29,073	1,096	1,044	1,053	881	770	690	615	476	318	18
1981	33,067	1,219	1,242	1,220	994	868	771	675	501	350	33
1980	34,109	1,322	1,251	1,203	984	834	740	672	462	356	18
1979	34,579	1,289	1,321	1,224	1,089	959	778	703	486	299	28
Motor vehicle accidents (E810-E825)											
White, female											
1998	12,197	702	561	546	466	559	726	775	671	522	3
1997	12,260	673	571	539	492	580	651	762	666	544	1
1996	12,238	632	560	477	523	555	706	719	654	472	6
1995	12,026	607	505	459	477	563	720	763	604	492	2
1994	11,687	588	512	436	443	599	708	730	643	416	3
1993	11,320	547	444	405	475	580	640	754	613	388	4
1992	11,024	561	410	404	444	578	655	678	559	372	-
1991	11,668	526	422	391	477	584	640	700	557	369	2
1990	12,363	535	497	465	554	625	646	737	536	382	2
1989	12,870	536	494	449	551	641	735	696	538	347	3
1988	12,847	499	513	522	537	616	670	678	552	343	5
1987	12,564	508	431	498	558	585	623	675	487	300	7
1986	12,026	426	429	485	511	584	664	610	451	269	3
1985	11,795	444	447	449	582	551	667	574	432	268	1
1984	11,815	465	421	517	557	522	577	595	423	237	3
1983	11,083	407	442	471	504	506	543	500	361	234	2
1982	11,118	427	426	474	498	495	542	480	341	220	5
1981	12,357	460	501	578	540	514	562	485	343	224	3
1980	12,754	496	557	589	496	511	556	510	337	220	3
1979	12,669	508	514	539	517	530	545	491	335	197	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
All other, both sexes											
1998	7,319	523	386	310	240	228	207	195	120	88	4
1997	7,364	479	375	302	249	219	233	174	121	92	4
1996	7,197	444	318	250	216	226	210	167	122	77	7
1995	7,108	446	306	245	237	202	214	157	112	75	2
1994	6,905	378	341	257	236	212	227	184	115	74	5
1993	6,699	348	297	228	234	211	168	159	101	101	6
1992	6,543	370	293	267	245	203	192	140	92	53	7
1991	6,685	347	257	234	224	231	161	129	85	74	5
1990	7,163	380	309	272	255	241	163	137	112	65	5
1989	7,084	352	312	268	256	213	170	146	90	58	4
1988	7,104	326	281	257	241	223	194	159	77	77	10
1987	6,709	328	275	247	229	221	163	119	75	54	6
1986	6,676	293	288	271	237	161	148	155	81	69	10
1985	6,212	274	258	255	252	173	164	137	65	46	1
1984	5,834	276	226	252	209	178	159	128	70	60	6
1983	5,588	288	267	231	199	174	143	130	66	37	6
1982	5,588	273	258	244	205	184	144	110	52	40	6
1981	5,961	312	287	264	201	185	190	112	66	33	11
1980	6,309	288	321	288	227	208	142	124	63	43	5
1979	6,276	296	329	300	236	210	149	112	52	41	5
Motor vehicle accidents (E810-E825)											
All other, male											
1998	5,005	356	259	212	149	129	119	113	80	59	4
1997	4,936	314	253	195	159	132	136	100	68	43	1
1996	4,909	275	230	173	138	138	126	97	70	48	6
1995	4,935	314	198	171	145	130	133	105	67	55	2
1994	4,703	238	232	179	155	127	148	113	72	48	4
1993	4,657	261	205	165	156	132	107	93	58	68	5
1992	4,567	257	206	178	165	131	126	90	54	38	6
1991	4,764	236	186	155	143	155	103	74	62	48	5
1990	5,118	267	226	172	172	143	99	79	70	47	4
1989	5,083	264	233	195	177	129	110	94	58	43	3
1988	5,131	239	204	176	169	158	140	102	51	57	9
1987	4,887	252	184	178	162	134	109	71	49	36	6
1986	4,925	219	205	193	151	110	98	110	59	51	8
1985	4,549	203	185	192	178	111	115	97	36	29	1
1984	4,335	210	164	177	157	127	111	89	44	45	5
1983	4,126	211	192	181	140	120	99	83	44	24	5
1982	4,118	214	193	180	156	132	99	80	42	28	3
1981	4,466	246	214	197	148	139	140	82	39	23	9
1980	4,752	218	245	221	167	165	107	95	40	26	5
1979	4,730	237	239	232	170	160	104	71	37	32	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
All other, female											
1998	2,314	167	127	98	91	99	88	82	40	29	-
1997	2,428	165	122	107	90	87	97	74	53	49	3
1996	2,288	169	88	77	78	88	84	70	52	29	1
1995	2,173	132	108	74	92	72	81	52	45	20	-
1994	2,202	140	109	78	81	85	79	71	43	26	1
1993	2,042	87	92	63	78	79	61	66	43	33	1
1992	1,976	113	87	89	80	72	66	50	38	15	1
1991	1,921	111	71	79	81	76	58	55	23	26	-
1990	2,045	113	83	100	83	98	64	58	42	18	1
1989	2,001	88	79	73	79	84	60	52	32	15	1
1988	1,973	87	77	81	72	65	54	57	26	20	1
1987	1,822	76	91	69	67	87	54	48	26	18	-
1986	1,751	74	83	78	86	51	50	45	22	18	2
1985	1,663	71	73	63	74	62	49	40	29	17	-
1984	1,499	66	62	75	52	51	48	39	26	15	1
1983	1,462	77	75	50	59	54	44	47	22	13	1
1982	1,470	59	65	64	49	52	45	30	10	12	3
1981	1,495	66	73	67	53	46	50	30	27	10	2
1980	1,557	70	76	67	60	43	35	29	23	17	-
1979	1,546	59	90	68	66	50	45	41	15	9	-
Motor vehicle accidents (E810-E825)											
Black, both sexes											
1998	5,686	410	296	239	185	175	163	154	84	65	4
1997	5,666	384	288	240	187	164	162	120	78	73	4
1996	5,528	344	244	189	165	170	152	128	90	64	6
1995	5,423	336	227	180	178	145	172	115	84	59	2
1994	5,341	293	260	191	177	159	171	138	86	60	3
1993	5,217	274	220	174	178	172	129	124	70	74	5
1992	5,071	286	224	203	197	154	143	112	69	38	7
1991	5,205	263	193	181	179	180	118	106	63	60	5
1990	5,560	298	239	208	215	192	133	106	81	46	5
1989	5,611	285	255	212	213	172	144	116	69	45	4
1988	5,623	260	235	211	196	175	154	126	61	63	7
1987	5,287	259	227	205	186	177	128	92	53	39	4
1986	5,287	233	227	225	204	134	121	122	60	58	8
1985	4,916	224	206	214	217	141	126	118	53	37	1
1984	4,721	216	171	213	175	161	136	107	62	49	4
1983	4,472	229	224	199	176	140	115	109	54	28	6
1982	4,493	219	217	198	166	149	116	93	39	32	5
1981	4,809	257	243	232	176	151	170	93	57	27	7
1980	5,081	240	275	245	198	177	126	100	46	31	5
1979	5,132	261	286	271	201	185	131	96	43	37	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Motor vehicle accidents (E810-E825)											
Black, male											
1998	3,996	285	205	165	122	109	96	93	60	44	4
1997	3,895	255	204	163	124	107	100	66	49	32	1
1996	3,862	230	187	134	115	113	96	76	57	41	5
1995	3,863	250	155	126	111	101	107	84	49	42	2
1994	3,703	197	174	139	120	99	113	86	55	37	2
1993	3,752	217	162	130	123	119	93	71	44	54	4
1992	3,607	212	168	136	138	107	98	77	40	26	6
1991	3,771	184	141	125	122	122	77	63	44	42	5
1990	4,046	218	183	135	150	120	84	63	50	32	4
1989	4,120	216	203	157	155	109	94	80	42	35	3
1988	4,139	193	176	152	141	128	116	84	39	46	6
1987	3,913	206	156	148	141	114	88	62	35	28	4
1986	3,974	179	164	171	132	90	82	93	46	42	7
1985	3,659	168	154	167	155	96	90	81	27	24	1
1984	3,556	174	129	152	140	119	97	73	40	36	3
1983	3,349	175	166	158	126	95	81	67	36	18	5
1982	3,387	179	167	152	125	110	82	68	30	21	3
1981	3,702	209	188	176	131	116	125	70	35	20	5
1980	3,912	188	212	194	150	141	96	73	33	18	5
1979	3,929	218	218	211	144	140	94	59	32	29	3
Motor vehicle accidents (E810-E825)											
Black, female											
1998	1,690	125	91	74	63	66	67	61	24	21	-
1997	1,771	129	84	77	63	57	62	54	29	41	3
1996	1,666	114	57	55	50	57	56	52	33	23	1
1995	1,560	86	72	54	67	44	65	31	35	17	-
1994	1,638	96	86	52	57	60	58	52	31	23	1
1993	1,465	57	58	44	55	53	36	53	26	20	1
1992	1,464	74	56	67	59	47	45	35	29	12	1
1991	1,434	79	52	56	57	58	41	43	19	18	-
1990	1,514	80	56	73	65	72	49	43	31	14	1
1989	1,491	69	52	55	58	63	50	36	27	10	1
1988	1,484	67	59	59	55	47	38	42	22	17	1
1987	1,374	53	71	57	45	63	40	30	18	11	-
1986	1,313	54	63	54	72	44	39	29	14	16	1
1985	1,257	56	52	47	62	45	36	37	26	13	-
1984	1,165	42	42	61	35	42	39	34	22	13	1
1983	1,123	54	58	41	50	45	34	42	18	10	1
1982	1,106	40	50	46	41	39	34	25	9	11	2
1981	1,107	48	55	56	45	35	45	23	22	7	2
1980	1,169	52	63	51	48	36	30	27	13	13	-
1979	1,203	43	68	60	57	45	37	37	11	8	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
All races, both sexes											
1998	750	60	55	37	23	21	17	12	14	13	7
1997	747	55	54	36	20	32	21	16	8	8	6
1996	767	61	50	26	25	14	19	28	11	13	11
1995	780	51	57	24	19	20	21	17	10	8	12
1994	830	55	36	35	29	13	19	24	15	8	7
1993	853	56	41	32	28	29	18	18	16	10	6
1992	865	56	43	38	18	18	13	15	13	11	5
1991	866	59	38	40	28	17	19	23	8	10	7
1990	883	49	46	23	25	24	23	19	6	4	6
1989	827	57	48	36	27	21	14	13	11	13	11
1988	762	39	37	26	18	26	20	14	10	14	5
1987	855	51	40	28	31	29	19	18	12	4	5
1986	790	34	52	24	30	28	13	12	14	7	11
1985	782	40	42	32	28	28	19	14	12	6	7
1984	760	32	32	38	30	17	12	18	9	18	5
1983	758	30	35	43	28	33	24	14	9	5	10
1982	727	43	41	27	38	18	23	17	13	6	6
1981	791	33	63	48	39	34	22	12	14	9	6
1980	864	39	53	45	34	19	18	21	16	14	4
1979	814	40	47	39	36	23	30	18	14	17	5
Other land transport accidents (E800-E807,E826-E829)											
All races, male											
1998	607	49	46	32	19	18	15	6	12	7	7
1997	635	48	46	31	19	28	19	12	7	6	6
1996	636	54	39	20	20	10	14	17	9	9	10
1995	654	46	43	22	15	16	18	10	9	5	12
1994	691	46	29	33	24	13	14	19	14	6	6
1993	681	42	35	26	20	17	13	12	7	7	6
1992	723	48	35	35	14	14	12	6	9	10	5
1991	720	56	27	38	20	12	18	18	6	7	7
1990	725	37	40	21	22	18	16	13	4	2	6
1989	696	47	41	32	22	15	14	9	8	9	11
1988	634	32	33	20	14	21	16	11	7	13	5
1987	715	40	35	26	28	24	14	15	9	2	5
1986	654	27	48	21	24	23	9	9	9	6	10
1985	644	34	35	26	23	21	15	8	8	4	7
1984	635	26	24	37	26	15	10	13	8	13	5
1983	616	27	33	34	25	28	18	6	7	2	10
1982	611	37	35	26	33	16	17	14	10	4	6
1981	662	28	56	40	31	27	16	10	11	7	6
1980	689	34	48	40	27	14	13	15	11	10	4
1979	633	32	39	33	36	18	19	11	10	13	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
All races, female											
1998	143	11	9	5	4	3	2	6	2	6	-
1997	112	7	8	5	1	4	2	4	1	2	-
1996	131	7	11	6	5	4	5	11	2	4	1
1995	126	5	14	2	4	4	3	7	1	3	-
1994	139	9	7	2	5	-	5	5	1	2	1
1993	172	14	6	6	8	12	5	6	9	3	-
1992	142	8	8	3	4	4	1	9	4	1	-
1991	146	3	11	2	8	5	1	5	2	3	-
1990	158	12	6	2	3	6	7	6	2	2	-
1989	131	10	7	4	5	6	-	4	3	4	-
1988	128	7	4	6	4	5	4	3	3	1	-
1987	140	11	5	2	3	5	5	3	3	2	-
1986	136	7	4	3	6	5	4	3	5	1	1
1985	138	6	7	6	5	7	4	6	4	2	-
1984	125	6	8	1	4	2	2	5	1	5	-
1983	142	3	2	9	3	5	6	8	2	3	-
1982	116	6	6	1	5	2	6	3	3	2	-
1981	129	5	7	8	8	7	6	2	3	2	-
1980	175	5	5	5	7	5	5	6	5	4	-
1979	181	8	8	6	-	5	11	7	4	4	-
Other land transport accidents (E800-E807,E826-E829)											
White, both sexes											
1998	610	43	45	26	21	17	16	12	14	12	7
1997	589	37	45	25	15	28	12	13	7	7	5
1996	634	51	39	23	24	12	15	24	8	12	9
1995	606	44	50	17	13	18	18	12	8	7	12
1994	666	43	30	27	25	11	13	19	11	7	7
1993	681	48	34	23	22	21	17	16	14	8	4
1992	708	43	38	35	10	18	11	14	12	9	5
1991	690	54	32	33	22	15	15	22	8	9	7
1990	702	43	35	21	20	21	20	15	3	4	5
1989	676	47	40	30	23	18	14	13	8	12	9
1988	604	31	28	22	14	21	14	13	8	13	4
1987	686	36	32	25	29	26	16	14	8	4	5
1986	634	28	47	19	24	24	11	11	13	6	8
1985	636	31	30	31	25	23	18	14	11	6	7
1984	613	26	26	32	25	14	8	18	8	15	4
1983	612	24	26	38	22	27	20	12	5	4	8
1982	587	33	36	22	35	15	19	17	11	6	5
1981	644	22	45	44	32	30	22	10	11	8	6
1980	699	33	43	37	26	13	12	19	14	12	3
1979	686	33	45	32	30	19	23	16	13	14	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
White, male											
1998	485	34	39	22	18	15	14	6	12	7	7
1997	493	31	38	22	14	24	10	10	6	5	5
1996	523	44	29	17	19	9	12	16	7	8	9
1995	499	39	36	16	10	15	15	8	7	4	12
1994	547	34	24	25	21	11	11	14	10	5	6
1993	535	36	28	18	15	12	12	10	6	6	4
1992	586	36	31	32	9	14	10	6	9	8	5
1991	576	51	23	31	15	10	14	18	6	7	7
1990	570	33	31	19	17	15	15	10	2	2	5
1989	560	40	35	26	19	13	14	9	6	9	9
1988	490	24	26	16	10	16	10	11	5	12	4
1987	561	28	29	23	26	21	12	11	5	2	5
1986	518	21	44	16	18	20	7	9	8	5	7
1985	511	25	24	25	20	18	14	8	7	4	7
1984	503	20	19	31	22	12	7	13	7	11	4
1983	491	22	24	30	19	23	14	5	3	1	8
1982	490	28	30	21	30	13	15	14	8	4	5
1981	530	18	42	36	26	23	16	8	9	6	6
1980	547	30	41	32	21	9	10	13	9	8	3
1979	527	25	37	29	30	15	14	10	9	10	3
Other land transport accidents (E800-E807,E826-E829)											
White, female											
1998	125	9	6	4	3	2	2	6	2	5	-
1997	96	6	7	3	1	4	2	3	1	2	-
1996	111	7	10	6	5	3	3	8	1	4	-
1995	107	5	14	1	3	3	3	4	1	3	-
1994	119	9	6	2	4	-	2	5	1	2	1
1993	146	12	6	5	7	9	5	6	8	2	-
1992	122	7	7	3	1	4	1	8	3	1	-
1991	114	3	9	2	7	5	1	4	2	2	-
1990	132	10	4	2	3	6	5	5	1	2	-
1989	116	7	5	4	4	5	-	4	2	3	-
1988	114	7	2	6	4	5	4	2	3	1	-
1987	125	8	3	2	3	5	4	3	3	2	-
1986	116	7	3	3	6	4	4	2	5	1	1
1985	125	6	6	6	5	5	4	6	4	2	-
1984	110	6	7	1	3	2	1	5	1	4	-
1983	121	2	2	8	3	4	6	7	2	3	-
1982	97	5	6	1	5	2	4	3	3	2	-
1981	114	4	3	8	6	7	6	2	2	2	-
1980	152	3	2	5	5	4	2	6	5	4	-
1979	159	8	8	3	-	4	9	6	4	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
All other, both sexes											
1998	140	17	10	11	2	4	1	-	-	1	-
1997	158	18	9	11	5	4	9	3	1	1	1
1996	133	10	11	3	1	2	4	4	3	1	2
1995	174	7	7	7	6	2	3	5	2	1	-
1994	164	12	6	8	4	2	6	5	4	1	-
1993	172	8	7	9	6	8	1	2	2	2	2
1992	157	13	5	3	8	-	2	1	1	2	-
1991	176	5	6	7	6	2	4	1	-	1	-
1990	181	6	11	2	5	3	3	4	3	-	1
1989	151	10	8	6	4	3	-	-	3	1	2
1988	158	8	9	4	4	5	6	1	2	1	1
1987	169	15	8	3	2	3	3	4	4	-	-
1986	156	6	5	5	6	4	2	1	1	1	3
1985	146	9	12	1	3	5	1	-	1	-	-
1984	147	6	6	6	5	3	4	-	1	3	1
1983	146	6	9	5	6	6	4	2	4	1	2
1982	140	10	5	5	3	3	4	-	2	-	1
1981	147	11	18	4	7	4	-	2	3	1	-
1980	165	6	10	8	8	6	6	2	2	2	1
1979	128	7	2	7	6	4	7	2	1	3	2
Other land transport accidents (E800-E807,E826-E829)											
All other, male											
1998	122	15	7	10	1	3	1	-	-	-	-
1997	142	17	8	9	5	4	9	2	1	1	1
1996	113	10	10	3	1	1	2	1	2	1	1
1995	155	7	7	6	5	1	3	2	2	1	-
1994	144	12	5	8	3	2	3	5	4	1	-
1993	146	6	7	8	5	5	1	2	1	1	2
1992	137	12	4	3	5	-	2	-	-	2	-
1991	144	5	4	7	5	2	4	-	-	-	-
1990	155	4	9	2	5	3	1	3	2	-	1
1989	136	7	6	6	3	2	-	-	2	-	2
1988	144	8	7	4	4	5	6	-	2	1	1
1987	154	12	6	3	2	3	2	4	4	-	-
1986	136	6	4	5	6	3	2	-	1	1	3
1985	133	9	11	1	3	3	1	-	1	-	-
1984	132	6	5	6	4	3	3	-	1	2	1
1983	125	5	9	4	6	5	4	1	4	1	2
1982	121	9	5	5	3	3	2	-	2	-	1
1981	132	10	14	4	5	4	-	2	2	1	-
1980	142	4	7	8	6	5	3	2	2	2	1
1979	106	7	2	4	6	3	5	1	1	3	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
All other, female											
1998	18	2	3	1	1	1	-	-	-	1	-
1997	16	1	1	2	-	-	-	1	-	-	-
1996	20	-	1	-	-	1	2	3	1	-	1
1995	19	-	-	1	1	1	-	3	-	-	-
1994	20	-	1	-	1	-	3	-	-	-	-
1993	26	2	-	1	1	3	-	-	1	1	-
1992	20	1	1	-	3	-	-	1	1	-	-
1991	32	-	2	-	1	-	-	1	-	1	-
1990	26	2	2	-	-	-	2	1	1	-	-
1989	15	3	2	-	1	1	-	-	1	1	-
1988	14	-	2	-	-	-	-	1	-	-	-
1987	15	3	2	-	-	-	1	-	-	-	-
1986	20	-	1	-	-	1	-	1	-	-	-
1985	13	-	1	-	-	2	-	-	-	-	-
1984	15	-	1	-	1	-	1	-	-	1	-
1983	21	1	-	1	-	1	-	1	-	-	-
1982	19	1	-	-	-	-	2	-	-	-	-
1981	15	1	4	-	2	-	-	-	1	-	-
1980	23	2	3	-	2	1	3	-	-	-	-
1979	22	-	-	3	-	1	2	1	-	-	-
Other land transport accidents (E800-E807,E826-E829)											
Black, both sexes											
1998	106	14	9	9	1	3	1	-	-	1	-
1997	121	16	8	8	4	3	4	2	-	1	1
1996	103	9	10	2	-	2	4	3	1	1	1
1995	140	6	7	7	6	2	3	3	-	1	-
1994	128	12	4	6	3	2	2	3	3	1	-
1993	135	6	6	6	5	7	1	-	2	2	1
1992	125	9	5	3	5	-	-	1	1	-	-
1991	155	5	5	7	5	2	1	1	-	1	-
1990	155	5	9	2	5	3	3	4	3	-	1
1989	121	8	7	5	4	3	-	-	2	1	1
1988	127	6	7	4	3	3	5	1	1	1	1
1987	143	13	7	2	2	3	3	3	3	-	-
1986	123	4	5	4	4	4	2	1	1	-	2
1985	125	9	9	1	3	5	1	-	1	-	-
1984	119	5	5	6	4	1	3	-	1	3	1
1983	117	6	9	5	5	6	2	2	4	1	2
1982	125	8	5	4	3	2	4	-	1	-	1
1981	121	10	17	4	6	1	-	2	3	1	-
1980	136	5	10	8	7	5	5	2	2	2	1
1979	103	7	-	6	6	2	7	-	-	2	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other land transport accidents (E800-E807,E826-E829)											
Black, male											
1998	93	13	7	8	-	2	1	-	-	-	-
1997	110	15	7	7	4	3	4	1	-	1	1
1996	90	9	10	2	-	1	2	1	-	1	-
1995	127	6	7	6	5	1	3	2	-	1	-
1994	117	12	4	6	2	2	1	3	3	1	-
1993	121	5	6	5	5	5	1	-	1	1	1
1992	111	9	4	3	4	-	-	-	-	-	-
1991	127	5	3	7	4	2	1	-	-	-	-
1990	132	4	7	2	5	3	1	3	2	-	1
1989	109	5	6	5	3	2	-	-	2	-	1
1988	114	6	6	4	3	3	5	-	1	1	1
1987	128	10	5	2	2	3	2	3	3	-	-
1986	108	4	4	4	4	3	2	-	1	-	2
1985	116	9	9	1	3	3	1	-	1	-	-
1984	109	5	4	6	3	1	3	-	1	2	1
1983	100	5	9	4	5	5	2	1	4	1	2
1982	108	7	5	4	3	2	2	-	1	-	1
1981	108	9	13	4	4	1	-	2	2	1	-
1980	120	4	7	8	5	5	3	2	2	2	1
1979	85	7	-	3	6	1	5	-	-	2	2
Other land transport accidents (E800-E807,E826-E829)											
Black, female											
1998	13	1	2	1	1	1	-	-	-	1	-
1997	11	1	1	1	-	-	-	1	-	-	-
1996	13	-	-	-	-	1	2	2	1	-	1
1995	13	-	-	1	1	1	-	1	-	-	-
1994	11	-	-	-	1	-	1	-	-	-	-
1993	14	1	-	1	-	2	-	-	1	1	-
1992	14	-	1	-	1	-	-	1	1	-	-
1991	28	-	2	-	1	-	-	1	-	1	-
1990	23	1	2	-	-	-	2	1	1	-	-
1989	12	3	1	-	1	1	-	-	-	1	-
1988	13	-	1	-	-	-	-	1	-	-	-
1987	15	3	2	-	-	-	1	-	-	-	-
1986	15	-	1	-	-	1	-	1	-	-	-
1985	9	-	-	-	-	2	-	-	-	-	-
1984	10	-	1	-	1	-	-	-	-	1	-
1983	17	1	-	1	-	1	-	1	-	-	-
1982	17	1	-	-	-	-	2	-	-	-	-
1981	13	1	4	-	2	-	-	-	1	-	-
1980	16	1	3	-	2	-	2	-	-	-	-
1979	18	-	-	3	-	1	2	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All races, both sexes											
1998	1,835	168	160	131	119	87	96	49	46	60	-
1997	2,082	230	181	144	88	83	67	48	43	53	-
1996	2,263	232	188	147	122	89	71	57	32	46	1
1995	2,073	213	147	124	115	94	65	44	21	20	-
1994	2,291	231	185	127	89	86	84	41	30	30	1
1993	2,059	196	146	109	105	74	59	39	30	15	3
1992	2,331	204	153	131	110	92	51	46	22	30	3
1991	2,227	209	132	123	99	97	55	47	16	22	-
1990	2,220	193	154	128	131	98	63	47	15	21	1
1989	2,329	202	157	140	144	101	65	37	25	23	-
1988	2,280	191	148	120	118	86	55	32	13	16	-
1987	2,497	182	147	150	139	84	50	29	20	23	1
1986	2,484	209	159	140	140	75	50	33	15	7	14
1985	2,762	223	169	167	97	97	65	27	18	7	3
1984	2,604	226	190	157	124	84	60	30	11	12	3
1983	2,840	198	194	157	114	82	57	25	10	7	4
1982	3,190	236	217	164	130	54	59	37	11	10	8
1981	3,014	230	214	164	117	88	48	29	6	9	1
1980	3,126	230	200	159	96	62	38	23	15	7	1
1979	3,419	242	237	183	144	72	51	29	7	6	21
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All races, male											
1998	1,510	142	137	117	107	82	76	38	28	20	-
1997	1,701	200	155	124	79	70	60	33	26	20	-
1996	1,759	173	164	120	102	75	57	36	15	23	1
1995	1,708	180	125	105	97	78	59	39	14	9	-
1994	1,889	192	154	110	79	73	67	29	22	20	1
1993	1,702	161	130	92	97	62	53	29	19	5	3
1992	1,951	169	140	112	94	76	39	34	16	15	3
1991	1,869	181	116	109	80	80	45	36	6	12	-
1990	1,915	172	136	113	111	80	52	36	10	12	1
1989	1,939	172	145	121	118	90	53	24	12	11	-
1988	1,971	167	133	103	106	73	50	25	9	8	-
1987	2,096	157	123	126	120	71	44	24	14	12	1
1986	2,137	179	145	125	131	68	44	24	11	5	10
1985	2,320	188	135	142	87	81	48	19	13	5	3
1984	2,240	194	167	139	108	74	55	25	10	6	3
1983	2,466	178	171	141	102	78	47	19	7	5	2
1982	2,727	207	191	137	113	48	48	28	4	6	5
1981	2,618	201	194	144	103	75	43	25	4	6	1
1980	2,648	200	172	138	87	57	28	21	6	2	-
1979	2,931	218	211	160	124	64	44	21	6	4	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All races, female											
1998	325	26	23	14	12	5	20	11	18	40	-
1997	381	30	26	20	9	13	7	15	17	33	-
1996	504	59	24	27	20	14	14	21	17	23	-
1995	365	33	22	19	18	16	6	5	7	11	-
1994	402	39	31	17	10	13	17	12	8	10	-
1993	357	35	16	17	8	12	6	10	11	10	-
1992	380	35	13	19	16	16	12	12	6	15	-
1991	358	28	16	14	19	17	10	11	10	10	-
1990	305	21	18	15	20	18	11	11	5	9	-
1989	390	30	12	19	26	11	12	13	13	12	-
1988	309	24	15	17	12	13	5	7	4	8	-
1987	401	25	24	24	19	13	6	5	6	11	-
1986	347	30	14	15	9	7	6	9	4	2	4
1985	442	35	34	25	10	16	17	8	5	2	-
1984	364	32	23	18	16	10	5	5	1	6	-
1983	374	20	23	16	12	4	10	6	3	2	2
1982	463	29	26	27	17	6	11	9	7	4	3
1981	396	29	20	20	14	13	5	4	2	3	-
1980	478	30	28	21	9	5	10	2	9	5	1
1979	488	24	26	23	20	8	7	8	1	2	9
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
White, both sexes											
1998	1,641	152	150	120	105	80	89	44	43	55	-
1997	1,827	209	168	134	81	75	60	41	40	51	-
1996	2,012	212	173	128	111	73	64	51	28	43	1
1995	1,869	195	139	111	98	92	64	40	20	18	-
1994	2,036	208	168	117	81	80	79	35	29	30	-
1993	1,848	175	132	99	94	71	53	35	29	14	2
1992	2,114	191	139	120	104	85	46	41	19	30	3
1991	2,011	195	121	114	92	90	52	47	16	21	-
1990	2,006	178	141	117	121	84	59	39	15	21	-
1989	2,083	185	136	124	131	90	61	34	23	23	-
1988	2,030	169	132	104	109	80	50	28	12	13	-
1987	2,219	170	126	136	127	75	46	26	17	21	1
1986	2,237	192	144	124	126	72	43	31	13	7	12
1985	2,505	203	158	152	93	91	62	25	17	7	2
1984	2,345	211	174	138	114	74	56	28	10	12	2
1983	2,596	181	177	147	105	73	52	21	9	7	3
1982	2,887	225	205	152	117	46	50	35	9	10	7
1981	2,782	218	202	150	109	79	45	27	6	9	1
1980	2,818	209	186	144	84	57	29	21	13	7	1
1979	3,123	220	221	172	132	69	47	23	6	5	20

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
White, male											
1998	1,352	129	127	108	94	75	72	34	27	17	-
1997	1,503	182	145	114	72	65	54	28	24	18	-
1996	1,562	160	151	107	94	60	51	31	12	21	1
1995	1,541	164	118	96	83	76	58	35	14	8	-
1994	1,684	174	141	101	72	68	64	23	21	20	-
1993	1,520	142	117	84	86	59	47	25	18	5	2
1992	1,762	159	128	103	90	71	34	31	14	15	3
1991	1,681	169	106	102	74	75	42	36	6	11	-
1990	1,723	157	126	103	102	67	48	28	10	12	-
1989	1,716	156	124	107	105	79	49	21	11	11	-
1988	1,751	147	119	89	97	68	46	21	8	6	-
1987	1,850	146	105	113	109	63	40	21	11	10	1
1986	1,926	164	132	109	117	66	38	22	9	5	10
1985	2,107	170	126	130	83	77	45	18	12	5	2
1984	2,005	181	152	121	99	65	51	23	9	6	2
1983	2,244	162	155	131	95	69	44	15	6	5	1
1982	2,457	196	180	126	100	40	39	26	3	6	4
1981	2,412	190	182	131	96	66	40	23	4	6	1
1980	2,383	180	158	124	76	52	21	19	5	2	-
1979	2,666	198	195	150	114	61	41	15	5	3	12
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
White, female											
1998	289	23	23	12	11	5	17	10	16	38	-
1997	324	27	23	20	9	10	6	13	16	33	-
1996	450	52	22	21	17	13	13	20	16	22	-
1995	328	31	21	15	15	16	6	5	6	10	-
1994	352	34	27	16	9	12	15	12	8	10	-
1993	328	33	15	15	8	12	6	10	11	9	-
1992	352	32	11	17	14	14	12	10	5	15	-
1991	330	26	15	12	18	15	10	11	10	10	-
1990	283	21	15	14	19	17	11	11	5	9	-
1989	367	29	12	17	26	11	12	13	12	12	-
1988	279	22	13	15	12	12	4	7	4	7	-
1987	369	24	21	23	18	12	6	5	6	11	-
1986	311	28	12	15	9	6	5	9	4	2	2
1985	398	33	32	22	10	14	17	7	5	2	-
1984	340	30	22	17	15	9	5	5	1	6	-
1983	352	19	22	16	10	4	8	6	3	2	2
1982	430	29	25	26	17	6	11	9	6	4	3
1981	370	28	20	19	13	13	5	4	2	3	-
1980	435	29	28	20	8	5	8	2	8	5	1
1979	457	22	26	22	18	8	6	8	1	2	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All other, both sexes											
1998	194	16	10	11	14	7	7	5	3	5	-
1997	255	21	13	10	7	8	7	7	3	2	-
1996	251	20	15	19	11	16	7	6	4	3	-
1995	204	18	8	13	17	2	1	4	1	2	-
1994	255	23	17	10	8	6	5	6	1	-	1
1993	211	21	14	10	11	3	6	4	1	1	1
1992	217	13	14	11	6	7	5	5	3	-	-
1991	216	14	11	9	7	7	3	-	-	1	-
1990	214	15	13	11	10	14	4	8	-	-	1
1989	246	17	21	16	13	11	4	3	2	-	-
1988	250	22	16	16	9	6	5	4	1	3	-
1987	278	12	21	14	12	9	4	3	3	2	-
1986	247	17	15	16	14	3	7	2	2	-	2
1985	257	20	11	15	4	6	3	2	1	-	1
1984	259	15	16	19	10	10	4	2	1	-	1
1983	244	17	17	10	9	9	5	4	1	-	1
1982	303	11	12	12	13	8	9	2	2	-	1
1981	232	12	12	14	8	9	3	2	-	-	-
1980	308	21	14	15	12	5	9	2	2	-	-
1979	296	22	16	11	12	3	4	6	1	1	1
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All other, male											
1998	158	13	10	9	13	7	4	4	1	3	-
1997	198	18	10	10	7	5	6	5	2	2	-
1996	197	13	13	13	8	15	6	5	3	2	-
1995	167	16	7	9	14	2	1	4	-	1	-
1994	205	18	13	9	7	5	3	6	1	-	1
1993	182	19	13	8	11	3	6	4	1	-	1
1992	189	10	12	9	4	5	5	3	2	-	-
1991	188	12	10	7	6	5	3	-	-	1	-
1990	192	15	10	10	9	13	4	8	-	-	1
1989	223	16	21	14	13	11	4	3	1	-	-
1988	220	20	14	14	9	5	4	4	1	2	-
1987	246	11	18	13	11	8	4	3	3	2	-
1986	211	15	13	16	14	2	6	2	2	-	-
1985	213	18	9	12	4	4	3	1	1	-	1
1984	235	13	15	18	9	9	4	2	1	-	1
1983	222	16	16	10	7	9	3	4	1	-	1
1982	270	11	11	11	13	8	9	2	1	-	1
1981	206	11	12	13	7	9	3	2	-	-	-
1980	265	20	14	14	11	5	7	2	1	-	-
1979	265	20	16	10	10	3	3	6	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
All other, female											
1998	36	3	-	2	1	-	3	1	2	2	-
1997	57	3	3	-	-	3	1	2	1	-	-
1996	54	7	2	6	3	1	1	1	1	1	-
1995	37	2	1	4	3	-	-	-	1	1	-
1994	50	5	4	1	1	1	2	-	-	-	-
1993	29	2	1	2	-	-	-	-	-	1	-
1992	28	3	2	2	2	2	-	2	1	-	-
1991	28	2	1	2	1	2	-	-	-	-	-
1990	22	-	3	1	1	1	-	-	-	-	-
1989	23	1	-	2	-	-	-	-	1	-	-
1988	30	2	2	2	-	1	1	-	-	1	-
1987	32	1	3	1	1	1	-	-	-	-	-
1986	36	2	2	-	-	1	1	-	-	-	2
1985	44	2	2	3	-	2	-	1	-	-	-
1984	24	2	1	1	1	1	-	-	-	-	-
1983	22	1	1	-	2	-	2	-	-	-	-
1982	33	-	1	1	-	-	-	-	1	-	-
1981	26	1	-	1	1	-	-	-	-	-	-
1980	43	1	-	1	1	-	2	-	1	-	-
1979	31	2	-	1	2	-	1	-	-	-	1
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
Black, both sexes											
1998	136	10	6	9	10	7	6	3	2	2	-
1997	164	15	7	6	4	6	5	6	2	2	-
1996	196	16	14	15	8	13	6	4	4	2	-
1995	150	13	6	9	14	2	-	4	1	2	-
1994	188	16	13	6	7	6	3	5	1	-	1
1993	159	18	12	8	10	2	6	1	1	1	1
1992	149	11	11	9	5	5	3	4	3	-	-
1991	158	8	10	7	5	6	3	-	-	1	-
1990	145	10	10	10	8	10	4	8	-	-	1
1989	176	11	20	14	11	10	4	3	1	-	-
1988	190	17	14	16	7	5	5	3	1	3	-
1987	205	9	17	12	10	7	4	3	3	2	-
1986	183	11	12	15	12	3	6	2	2	-	2
1985	182	16	8	10	3	6	3	2	1	-	1
1984	213	13	16	17	9	10	4	1	-	-	1
1983	191	15	14	7	7	8	4	3	1	-	1
1982	249	10	9	11	12	7	9	2	1	-	1
1981	191	10	8	12	8	7	3	2	-	-	-
1980	252	20	14	14	11	5	8	1	2	-	-
1979	235	18	15	9	9	2	3	5	1	1	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
Black, male											
1998	114	9	6	7	10	7	4	2	1	1	-
1997	137	13	6	6	4	4	5	5	1	2	-
1996	157	10	13	11	6	12	6	4	3	1	-
1995	124	12	5	6	11	2	-	4	-	1	-
1994	149	12	11	5	6	5	2	5	1	-	1
1993	140	16	11	6	10	2	6	1	1	-	1
1992	131	9	9	8	3	4	3	3	2	-	-
1991	137	7	9	6	4	4	3	-	-	1	-
1990	133	10	8	9	7	9	4	8	-	-	1
1989	164	10	20	12	11	10	4	3	1	-	-
1988	165	15	12	14	7	4	4	3	1	2	-
1987	182	9	15	11	9	6	4	3	3	2	-
1986	166	10	10	15	12	2	5	2	2	-	-
1985	157	14	7	9	3	4	3	1	1	-	1
1984	200	12	15	16	9	9	4	1	-	-	1
1983	176	14	14	7	6	8	3	3	1	-	1
1982	223	10	8	10	12	7	9	2	1	-	1
1981	172	10	8	11	7	7	3	2	-	-	-
1980	220	19	14	14	10	5	6	1	1	-	-
1979	211	16	15	8	8	2	2	5	1	1	-
Water, air and space, and other and unspecified transport accidents and their sequelae (E830-E848,E929.0-E929.1)											
Black, female											
1998	22	1	-	2	-	-	2	1	1	1	-
1997	27	2	1	-	-	2	-	1	1	-	-
1996	39	6	1	4	2	1	-	-	1	1	-
1995	26	1	1	3	3	-	-	-	1	1	-
1994	39	4	2	1	1	1	1	-	-	-	-
1993	19	2	1	2	-	-	-	-	-	1	-
1992	18	2	2	1	2	1	-	1	1	-	-
1991	21	1	1	1	1	2	-	-	-	-	-
1990	12	-	2	1	1	1	-	-	-	-	-
1989	12	1	-	2	-	-	-	-	-	-	-
1988	25	2	2	2	-	1	1	-	-	1	-
1987	23	-	2	1	1	1	-	-	-	-	-
1986	17	1	2	-	-	1	1	-	-	-	2
1985	25	2	1	1	-	2	-	1	-	-	-
1984	13	1	1	1	-	1	-	-	-	-	-
1983	15	1	-	-	1	-	1	-	-	-	-
1982	26	-	1	1	-	-	-	-	-	-	-
1981	19	-	-	1	1	-	-	-	-	-	-
1980	32	1	-	-	1	-	2	-	1	-	-
1979	24	2	-	1	1	-	1	-	-	-	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All races, both sexes											
1998	48,245	3,100	2,101	1,646	1,552	1,861	2,634	3,577	4,595	9,378	72
1997	46,066	2,869	1,923	1,567	1,542	1,817	2,480	3,389	4,284	8,812	72
1996	45,097	2,718	1,794	1,486	1,523	1,876	2,589	3,207	4,232	8,629	61
1995	44,186	2,451	1,774	1,471	1,570	1,809	2,444	3,070	3,817	8,055	60
1994	43,004	2,285	1,577	1,387	1,528	1,751	2,296	3,033	3,868	7,515	57
1993	42,793	2,061	1,522	1,320	1,559	1,851	2,277	2,900	3,691	7,446	65
1992	39,774	1,742	1,386	1,339	1,563	1,880	2,194	2,881	3,499	6,874	77
1991	40,082	1,606	1,318	1,354	1,622	1,904	2,231	2,849	3,510	6,746	69
1990	39,231	1,480	1,369	1,414	1,803	1,982	2,205	2,887	3,310	6,295	61
1989	41,305	1,617	1,386	1,594	1,821	2,105	2,333	3,036	3,388	6,310	73
1988	41,942	1,602	1,494	1,606	1,948	2,170	2,436	2,980	3,417	6,460	82
1987	40,346	1,453	1,359	1,663	1,963	2,047	2,416	2,951	3,355	5,895	72
1986	41,069	1,451	1,461	1,654	1,955	2,008	2,442	2,887	3,246	5,781	93
1985	41,160	1,486	1,521	1,887	2,089	2,101	2,516	3,025	3,240	5,724	91
1984	40,656	1,510	1,620	1,818	1,973	2,149	2,469	2,900	3,149	5,622	60
1983	41,778	1,516	1,752	2,010	2,120	2,172	2,454	2,824	3,160	5,562	77
1982	41,801	1,589	1,837	2,043	2,078	2,164	2,402	2,736	3,031	5,273	53
1981	42,965	1,659	2,086	2,101	2,198	2,205	2,495	2,694	3,128	5,435	62
1980	46,119	1,795	2,165	2,334	2,399	2,578	2,694	3,135	3,339	5,567	70
1979	44,913	1,829	2,121	2,317	2,277	2,437	2,600	2,918	3,167	5,128	82
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All races, male											
1998	30,292	2,445	1,564	1,216	1,069	1,197	1,577	1,923	2,227	3,332	67
1997	29,298	2,280	1,438	1,173	1,095	1,224	1,478	1,849	2,065	3,168	67
1996	28,609	2,138	1,359	1,094	1,105	1,257	1,563	1,739	2,035	3,054	52
1995	28,540	1,975	1,357	1,095	1,112	1,216	1,433	1,664	1,835	2,935	58
1994	28,035	1,792	1,223	1,038	1,102	1,128	1,404	1,696	1,877	2,750	52
1993	27,859	1,608	1,201	980	1,083	1,179	1,415	1,560	1,766	2,711	58
1992	25,898	1,373	1,085	1,010	1,137	1,258	1,360	1,542	1,685	2,436	72
1991	26,042	1,249	1,014	1,022	1,171	1,276	1,340	1,532	1,651	2,395	59
1990	25,560	1,132	1,064	1,046	1,293	1,312	1,348	1,516	1,531	2,216	52
1989	27,141	1,282	1,063	1,208	1,317	1,428	1,397	1,611	1,588	2,278	64
1988	27,504	1,288	1,138	1,231	1,376	1,427	1,444	1,583	1,577	2,387	77
1987	26,627	1,140	1,047	1,220	1,395	1,308	1,468	1,601	1,576	2,207	63
1986	27,254	1,151	1,145	1,255	1,358	1,335	1,469	1,544	1,461	2,187	84
1985	27,348	1,160	1,148	1,406	1,491	1,368	1,471	1,653	1,512	2,201	83
1984	26,952	1,151	1,219	1,367	1,376	1,438	1,473	1,571	1,452	2,128	55
1983	27,616	1,166	1,322	1,494	1,511	1,422	1,436	1,498	1,431	2,083	65
1982	27,921	1,220	1,355	1,502	1,482	1,433	1,404	1,481	1,347	1,971	43
1981	28,813	1,255	1,567	1,537	1,605	1,478	1,492	1,443	1,382	2,042	55
1980	30,747	1,371	1,643	1,731	1,662	1,707	1,589	1,606	1,483	2,035	63
1979	30,175	1,392	1,591	1,682	1,614	1,644	1,551	1,538	1,387	1,896	75

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All races, female											
1998	17,953	655	537	430	483	664	1,057	1,654	2,368	6,046	5
1997	16,768	589	485	394	447	593	1,002	1,540	2,219	5,644	5
1996	16,488	580	435	392	418	619	1,026	1,468	2,197	5,575	9
1995	15,646	476	417	376	458	593	1,011	1,406	1,982	5,120	2
1994	14,969	493	354	349	426	623	892	1,337	1,991	4,765	5
1993	14,934	453	321	340	476	672	862	1,340	1,925	4,735	7
1992	13,876	369	301	329	426	622	834	1,339	1,814	4,438	5
1991	14,040	357	304	332	451	628	891	1,317	1,859	4,351	10
1990	13,671	348	305	368	510	670	857	1,371	1,779	4,079	9
1989	14,164	335	323	386	504	677	936	1,425	1,800	4,032	9
1988	14,438	314	356	375	572	743	992	1,397	1,840	4,073	5
1987	13,719	313	312	443	568	739	948	1,350	1,779	3,688	9
1986	13,815	300	316	399	597	673	973	1,343	1,785	3,594	9
1985	13,812	326	373	481	598	733	1,045	1,372	1,728	3,523	8
1984	13,704	359	401	451	597	711	996	1,329	1,697	3,494	5
1983	14,162	350	430	516	609	750	1,018	1,326	1,729	3,479	12
1982	13,880	369	482	541	596	731	998	1,255	1,684	3,302	10
1981	14,152	404	519	564	593	727	1,003	1,251	1,746	3,393	7
1980	15,372	424	522	603	737	871	1,105	1,529	1,856	3,532	7
1979	14,738	437	530	635	663	793	1,049	1,380	1,780	3,232	7
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
White, both sexes											
1998	40,864	2,389	1,640	1,339	1,262	1,580	2,315	3,238	4,250	8,761	65
1997	38,682	2,201	1,506	1,260	1,233	1,515	2,158	3,019	3,946	8,246	61
1996	37,720	2,064	1,404	1,172	1,227	1,549	2,244	2,855	3,899	8,092	52
1995	36,596	1,844	1,386	1,149	1,244	1,486	2,100	2,733	3,448	7,522	53
1994	35,290	1,727	1,195	1,066	1,227	1,472	1,994	2,705	3,501	7,007	47
1993	35,089	1,593	1,178	1,046	1,264	1,528	1,960	2,599	3,356	6,906	57
1992	32,811	1,336	1,071	1,039	1,274	1,576	1,898	2,526	3,200	6,394	69
1991	32,671	1,248	1,003	1,050	1,331	1,578	1,919	2,519	3,189	6,232	60
1990	32,199	1,132	1,077	1,144	1,458	1,655	1,891	2,536	2,997	5,828	52
1989	33,422	1,206	1,038	1,242	1,446	1,708	1,992	2,659	3,073	5,757	55
1988	33,966	1,210	1,121	1,260	1,545	1,803	2,045	2,591	3,074	5,984	61
1987	32,813	1,096	1,050	1,320	1,586	1,689	2,045	2,606	3,010	5,428	53
1986	33,702	1,094	1,164	1,293	1,605	1,686	2,083	2,540	2,938	5,341	78
1985	33,798	1,101	1,171	1,510	1,707	1,759	2,127	2,651	2,896	5,275	71
1984	33,643	1,163	1,294	1,443	1,655	1,813	2,093	2,579	2,833	5,221	51
1983	34,320	1,160	1,387	1,590	1,713	1,769	2,053	2,484	2,806	5,188	60
1982	34,584	1,226	1,432	1,637	1,718	1,840	2,032	2,414	2,759	4,943	45
1981	35,460	1,265	1,668	1,692	1,800	1,828	2,157	2,397	2,840	5,076	47
1980	37,689	1,362	1,677	1,863	1,948	2,119	2,257	2,662	2,988	5,171	55
1979	37,059	1,428	1,666	1,928	1,903	2,044	2,256	2,565	2,937	4,779	65

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
White, male											
1998	25,204	1,895	1,193	989	853	1,011	1,374	1,746	2,060	3,062	62
1997	24,266	1,763	1,124	959	865	1,027	1,289	1,653	1,908	2,924	58
1996	23,564	1,615	1,055	867	888	1,032	1,358	1,549	1,873	2,839	45
1995	23,286	1,490	1,071	849	878	992	1,226	1,482	1,648	2,705	51
1994	22,613	1,364	923	789	876	950	1,221	1,506	1,679	2,534	43
1993	22,458	1,240	917	773	875	969	1,208	1,401	1,609	2,478	53
1992	21,047	1,046	838	780	923	1,056	1,169	1,359	1,531	2,231	65
1991	20,795	980	774	781	950	1,046	1,156	1,348	1,469	2,185	53
1990	20,654	863	835	837	1,046	1,090	1,160	1,310	1,397	2,023	46
1989	21,614	948	796	944	1,034	1,152	1,195	1,397	1,417	2,053	50
1988	21,858	972	842	956	1,074	1,160	1,206	1,354	1,393	2,183	58
1987	21,274	866	803	967	1,114	1,076	1,233	1,401	1,402	1,996	47
1986	22,071	856	919	964	1,133	1,104	1,269	1,340	1,303	1,997	72
1985	22,179	857	888	1,124	1,218	1,135	1,241	1,442	1,326	1,994	65
1984	22,031	886	979	1,082	1,150	1,212	1,237	1,390	1,292	1,947	47
1983	22,476	895	1,055	1,189	1,218	1,154	1,193	1,301	1,255	1,916	53
1982	22,800	933	1,052	1,197	1,213	1,220	1,173	1,295	1,204	1,820	38
1981	23,440	952	1,239	1,225	1,312	1,217	1,275	1,272	1,232	1,882	41
1980	24,890	1,029	1,265	1,388	1,340	1,399	1,362	1,368	1,312	1,868	54
1979	24,551	1,086	1,229	1,381	1,337	1,384	1,325	1,330	1,260	1,751	59
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
White, female											
1998	15,660	494	447	350	409	569	941	1,492	2,190	5,699	3
1997	14,416	438	382	301	368	488	869	1,366	2,038	5,322	3
1996	14,156	449	349	305	339	517	886	1,306	2,026	5,253	7
1995	13,310	354	315	300	366	494	874	1,251	1,800	4,817	2
1994	12,677	363	272	277	351	522	773	1,199	1,822	4,473	4
1993	12,631	353	261	273	389	559	752	1,198	1,747	4,428	4
1992	11,764	290	233	259	351	520	729	1,167	1,669	4,163	4
1991	11,876	268	229	269	381	532	763	1,171	1,720	4,047	7
1990	11,545	269	242	307	412	565	731	1,226	1,600	3,805	6
1989	11,808	258	242	298	412	556	797	1,262	1,656	3,704	5
1988	12,108	238	279	304	471	643	839	1,237	1,681	3,801	3
1987	11,539	230	247	353	472	613	812	1,205	1,608	3,432	6
1986	11,631	238	245	329	472	582	814	1,200	1,635	3,344	6
1985	11,619	244	283	386	489	624	886	1,209	1,570	3,281	6
1984	11,612	277	315	361	505	601	856	1,189	1,541	3,274	4
1983	11,844	265	332	401	495	615	860	1,183	1,551	3,272	7
1982	11,784	293	380	440	505	620	859	1,119	1,555	3,123	7
1981	12,020	313	429	467	488	611	882	1,125	1,608	3,194	6
1980	12,799	333	412	475	608	720	895	1,294	1,676	3,303	1
1979	12,508	342	437	547	566	660	931	1,235	1,677	3,028	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All other, both sexes											
1998	7,381	711	461	307	290	281	319	339	345	617	7
1997	7,384	668	417	307	309	302	322	370	338	566	11
1996	7,377	654	390	314	296	327	345	352	333	537	9
1995	7,590	607	388	322	326	323	344	337	369	533	7
1994	7,714	558	382	321	301	279	302	328	367	508	10
1993	7,704	468	344	274	295	323	317	301	335	540	8
1992	6,963	406	315	300	289	304	296	355	299	480	8
1991	7,411	358	315	304	291	326	312	330	321	514	9
1990	7,032	348	292	270	345	327	314	351	313	467	9
1989	7,883	411	348	352	375	397	341	377	315	553	18
1988	7,976	392	373	346	403	367	391	389	343	476	21
1987	7,533	357	309	343	377	358	371	345	345	467	19
1986	7,367	357	297	361	350	322	359	347	308	440	15
1985	7,362	385	350	377	382	342	389	374	344	449	20
1984	7,013	347	326	375	318	336	376	321	316	401	9
1983	7,458	356	365	420	407	403	401	340	354	374	17
1982	7,217	363	405	406	360	324	370	322	272	330	8
1981	7,505	394	418	409	398	377	338	297	288	359	15
1980	8,430	433	488	471	451	459	437	473	351	396	15
1979	7,854	401	455	389	374	393	344	353	230	349	17
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All other, male											
1998	5,088	550	371	227	216	186	203	177	167	270	5
1997	5,032	517	314	214	230	197	189	196	157	244	9
1996	5,045	523	304	227	217	225	205	190	162	215	7
1995	5,254	485	286	246	234	224	207	182	187	230	7
1994	5,422	428	300	249	226	178	183	190	198	216	9
1993	5,401	368	284	207	208	210	207	159	157	233	5
1992	4,851	327	247	230	214	202	191	183	154	205	7
1991	5,247	269	240	241	221	230	184	184	182	210	6
1990	4,906	269	229	209	247	222	188	206	134	193	6
1989	5,527	334	267	264	283	276	202	214	171	225	14
1988	5,646	316	296	275	302	267	238	229	184	204	19
1987	5,353	274	244	253	281	232	235	200	174	211	16
1986	5,183	295	226	291	225	231	200	204	158	190	12
1985	5,169	303	260	282	273	233	230	211	186	207	18
1984	4,921	265	240	285	226	226	236	181	160	181	8
1983	5,140	271	267	305	293	268	243	197	176	167	12
1982	5,121	287	303	305	269	213	231	186	143	151	5
1981	5,373	303	328	312	293	261	217	171	150	160	14
1980	5,857	342	378	343	322	308	227	238	171	167	9
1979	5,624	306	362	301	277	260	226	208	127	145	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
All other, female											
1998	2,293	161	90	80	74	95	116	162	178	347	2
1997	2,352	151	103	93	79	105	133	174	181	322	2
1996	2,332	131	86	87	79	102	140	162	171	322	2
1995	2,336	122	102	76	92	99	137	155	182	303	-
1994	2,292	130	82	72	75	101	119	138	169	292	1
1993	2,303	100	60	67	87	113	110	142	178	307	3
1992	2,112	79	68	70	75	102	105	172	145	275	1
1991	2,164	89	75	63	70	96	128	146	139	304	3
1990	2,126	79	63	61	98	105	126	145	179	274	3
1989	2,356	77	81	88	92	121	139	163	144	328	4
1988	2,330	76	77	71	101	100	153	160	159	272	2
1987	2,180	83	65	90	96	126	136	145	171	256	3
1986	2,184	62	71	70	125	91	159	143	150	250	3
1985	2,193	82	90	95	109	109	159	163	158	242	2
1984	2,092	82	86	90	92	110	140	140	156	220	1
1983	2,318	85	98	115	114	135	158	143	178	207	5
1982	2,096	76	102	101	91	111	139	136	129	179	3
1981	2,132	91	90	97	105	116	121	126	138	199	1
1980	2,573	91	110	128	129	151	210	235	180	229	6
1979	2,230	95	93	88	97	133	118	145	103	204	1
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
Black, both sexes											
1998	6,309	646	406	279	241	231	261	292	275	508	7
1997	6,210	594	370	261	262	262	259	308	275	453	9
1996	6,304	588	348	272	261	281	291	291	270	434	6
1995	6,583	545	351	281	290	278	306	298	316	447	7
1994	6,659	499	338	273	264	246	262	276	305	416	7
1993	6,735	418	315	241	259	285	272	261	279	461	7
1992	6,010	359	277	268	258	267	256	304	254	402	8
1991	6,548	315	283	270	249	285	275	298	280	444	8
1990	6,138	314	259	242	308	282	273	308	265	406	9
1989	6,999	359	311	322	346	367	311	344	274	492	15
1988	7,067	361	346	305	356	321	351	347	288	412	21
1987	6,657	326	290	297	338	318	319	308	298	404	13
1986	6,521	322	267	324	322	294	330	311	284	376	12
1985	6,556	346	320	350	356	315	356	332	313	391	18
1984	6,225	310	295	350	287	301	339	293	282	355	8
1983	6,689	318	318	386	374	376	362	310	326	335	12
1982	6,404	334	349	370	319	297	340	294	245	290	8
1981	6,798	367	385	384	372	351	317	274	261	322	14
1980	7,652	403	453	442	422	430	399	443	325	357	14
1979	7,112	371	413	358	335	368	318	322	213	310	17

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
Black, male											
1998	4,341	503	326	211	179	157	165	144	133	210	5
1997	4,217	457	278	181	192	171	147	169	126	180	7
1996	4,300	476	273	197	195	199	178	151	121	167	4
1995	4,525	437	262	213	211	192	184	157	163	181	7
1994	4,663	382	266	219	200	160	154	161	160	165	7
1993	4,701	330	265	182	182	190	172	137	120	197	4
1992	4,172	284	218	207	193	179	162	157	130	161	7
1991	4,624	236	214	215	189	198	163	166	150	173	6
1990	4,250	244	203	187	223	189	166	180	104	158	6
1989	4,872	294	239	240	258	256	181	196	143	195	11
1988	4,963	288	273	242	264	232	207	204	149	169	19
1987	4,715	251	232	219	251	203	203	172	149	182	11
1986	4,560	263	207	261	206	213	183	185	141	158	11
1985	4,603	275	237	263	255	215	213	179	170	174	16
1984	4,346	234	215	268	202	206	208	164	138	158	8
1983	4,567	245	226	283	265	248	214	176	160	151	7
1982	4,517	262	265	281	234	198	212	168	123	127	5
1981	4,838	285	301	290	274	240	203	155	131	140	13
1980	5,264	318	356	319	298	286	197	217	155	146	8
1979	5,063	283	330	273	246	248	208	187	115	124	16
Nontransport accidents (E850-E869,E880-E928,E929.2-E929.9)											
Black, female											
1998	1,968	143	80	68	62	74	96	148	142	298	2
1997	1,993	137	92	80	70	91	112	139	149	273	2
1996	2,004	112	75	75	66	82	113	140	149	267	2
1995	2,058	108	89	68	79	86	122	141	153	266	-
1994	1,996	117	72	54	64	86	108	115	145	251	-
1993	2,034	88	50	59	77	95	100	124	159	264	3
1992	1,838	75	59	61	65	88	94	147	124	241	1
1991	1,924	79	69	55	60	87	112	132	130	271	2
1990	1,888	70	56	55	85	93	107	128	161	248	3
1989	2,127	65	72	82	88	111	130	148	131	297	4
1988	2,104	73	73	63	92	89	144	143	139	243	2
1987	1,942	75	58	78	87	115	116	136	149	222	2
1986	1,961	59	60	63	116	81	147	126	143	218	1
1985	1,953	71	83	87	101	100	143	153	143	217	2
1984	1,879	76	80	82	85	95	131	129	144	197	-
1983	2,122	73	92	103	109	128	148	134	166	184	5
1982	1,887	72	84	89	85	99	128	126	122	163	3
1981	1,960	82	84	94	98	111	114	119	130	182	1
1980	2,388	85	97	123	124	144	202	226	170	211	6
1979	2,049	88	83	85	89	120	110	135	98	186	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
All races, both sexes											
1998	16,274	435	409	410	486	703	1,181	1,874	2,772	6,596	5
1997	15,447	392	354	368	473	706	1,134	1,779	2,624	6,207	4
1996	14,986	387	365	368	492	680	1,100	1,699	2,519	6,003	3
1995	13,986	370	320	335	450	684	1,067	1,549	2,238	5,515	4
1994	13,450	314	296	351	469	622	984	1,530	2,294	5,178	4
1993	13,141	313	295	322	471	646	918	1,517	2,135	5,104	4
1992	12,646	306	307	348	460	666	921	1,445	2,089	4,797	5
1991	12,662	285	280	330	460	659	973	1,421	2,093	4,650	5
1990	12,313	285	294	345	541	697	923	1,466	1,897	4,323	4
1989	12,151	289	287	365	560	689	966	1,438	1,892	4,202	4
1988	12,096	258	342	381	497	692	965	1,396	1,850	4,157	6
1987	11,733	251	273	418	575	665	943	1,389	1,850	3,755	7
1986	11,444	258	327	407	527	644	910	1,311	1,742	3,706	6
1985	12,001	263	305	480	591	647	994	1,415	1,800	3,857	4
1984	11,937	263	344	448	569	716	986	1,419	1,785	3,857	6
1983	12,024	279	381	461	587	702	954	1,293	1,820	3,880	8
1982	12,077	286	409	468	616	714	921	1,406	1,826	3,757	5
1981	12,628	297	474	482	643	672	1,049	1,378	1,935	3,985	8
1980	13,294	352	481	583	700	781	1,059	1,537	1,945	4,026	5
1979	13,216	354	467	558	659	832	1,124	1,556	2,056	3,844	3
Falls (E880-E888)											
All races, male											
1998	8,035	349	316	316	331	440	655	960	1,285	2,222	5
1997	7,705	315	259	269	336	455	642	958	1,225	2,093	4
1996	7,429	311	275	278	362	456	641	861	1,114	2,003	-
1995	7,026	317	245	248	311	452	577	787	1,016	1,859	3
1994	6,889	247	240	261	344	372	582	811	1,056	1,789	3
1993	6,620	241	232	230	318	390	538	771	942	1,746	4
1992	6,315	240	232	253	326	419	539	733	926	1,588	5
1991	6,376	235	222	249	332	425	543	697	894	1,526	4
1990	6,311	226	237	237	378	441	534	673	838	1,441	4
1989	6,302	229	219	279	399	442	544	727	844	1,394	3
1988	6,303	213	274	295	354	437	554	696	772	1,410	6
1987	6,203	198	202	311	388	409	545	718	796	1,280	4
1986	6,043	214	247	318	358	407	496	630	759	1,282	3
1985	6,392	208	237	356	397	393	570	712	763	1,354	3
1984	6,210	187	254	336	382	465	531	683	723	1,343	4
1983	6,279	212	288	339	409	436	525	624	770	1,327	6
1982	6,354	210	314	341	423	449	490	694	761	1,295	3
1981	6,627	233	360	342	458	418	554	658	774	1,382	7
1980	7,036	272	370	430	466	505	611	737	774	1,355	2
1979	6,928	281	354	399	452	542	608	726	804	1,289	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
All races, female											
1998	8,239	86	93	94	155	263	526	914	1,487	4,374	-
1997	7,742	77	95	99	137	251	492	821	1,399	4,114	-
1996	7,557	76	90	90	130	224	459	838	1,405	4,000	3
1995	6,960	53	75	87	139	232	490	762	1,222	3,656	1
1994	6,561	67	56	90	125	250	402	719	1,238	3,389	1
1993	6,521	72	63	92	153	256	380	746	1,193	3,358	-
1992	6,331	66	75	95	134	247	382	712	1,163	3,209	-
1991	6,286	50	58	81	128	234	430	724	1,199	3,124	1
1990	6,002	59	57	108	163	256	389	793	1,059	2,882	-
1989	5,849	60	68	86	161	247	422	711	1,048	2,808	1
1988	5,793	45	68	86	143	255	411	700	1,078	2,747	-
1987	5,530	53	71	107	187	256	398	671	1,054	2,475	3
1986	5,401	44	80	89	169	237	414	681	983	2,424	3
1985	5,609	55	68	124	194	254	424	703	1,037	2,503	1
1984	5,727	76	90	112	187	251	455	736	1,062	2,514	2
1983	5,745	67	93	122	178	266	429	669	1,050	2,553	2
1982	5,723	76	95	127	193	265	431	712	1,065	2,462	2
1981	6,001	64	114	140	185	254	495	720	1,161	2,603	1
1980	6,258	80	111	153	234	276	448	800	1,171	2,671	3
1979	6,288	73	113	159	207	290	516	830	1,252	2,555	-
Falls (E880-E888)											
White, both sexes											
1998	15,037	362	324	358	407	623	1,073	1,758	2,641	6,312	4
1997	14,277	317	284	322	412	632	1,035	1,660	2,500	5,945	4
1996	13,849	301	298	310	424	595	1,011	1,591	2,411	5,765	3
1995	12,850	303	268	281	386	594	967	1,441	2,107	5,312	3
1994	12,342	250	241	289	384	556	913	1,442	2,164	4,979	3
1993	12,014	261	246	259	408	563	824	1,412	2,025	4,895	4
1992	11,560	236	254	280	393	597	835	1,333	1,994	4,583	4
1991	11,556	222	223	257	405	594	893	1,342	1,992	4,427	4
1990	11,131	220	223	285	459	619	834	1,368	1,772	4,123	2
1989	10,961	208	211	310	488	590	871	1,339	1,798	4,005	4
1988	10,886	205	270	310	422	606	884	1,283	1,741	3,967	2
1987	10,532	181	221	342	483	595	847	1,289	1,739	3,579	5
1986	10,368	196	269	335	446	578	825	1,230	1,663	3,555	6
1985	10,798	199	243	395	509	570	892	1,316	1,695	3,690	4
1984	10,829	192	281	361	491	638	897	1,341	1,699	3,690	5
1983	10,877	207	302	384	504	617	862	1,225	1,707	3,739	7
1982	10,930	211	334	377	537	639	836	1,302	1,736	3,618	3
1981	11,417	224	375	394	537	590	967	1,291	1,838	3,842	4
1980	12,016	258	374	503	600	695	964	1,432	1,855	3,886	3
1979	11,999	269	367	479	584	739	1,030	1,456	1,988	3,699	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
White, male											
1998	7,264	287	239	273	273	388	588	909	1,225	2,102	4
1997	6,977	249	205	234	289	405	592	897	1,171	1,982	4
1996	6,711	240	215	230	315	391	588	806	1,069	1,920	-
1995	6,308	259	208	203	267	392	516	735	953	1,781	2
1994	6,148	197	192	211	277	332	538	755	983	1,709	3
1993	5,885	200	190	184	268	332	476	720	891	1,661	4
1992	5,646	185	192	203	273	373	485	677	884	1,509	4
1991	5,671	182	179	192	290	376	498	661	842	1,447	3
1990	5,565	176	181	190	320	393	484	623	783	1,364	2
1989	5,524	164	155	235	340	382	493	673	793	1,331	3
1988	5,498	170	215	237	294	369	505	633	715	1,341	2
1987	5,391	142	167	248	321	358	480	663	743	1,210	3
1986	5,317	161	202	258	310	358	451	592	714	1,214	3
1985	5,579	156	188	292	341	330	509	659	712	1,283	3
1984	5,484	135	200	269	325	413	480	648	682	1,277	4
1983	5,529	159	231	280	344	378	466	583	721	1,274	5
1982	5,587	151	249	269	365	402	440	636	715	1,236	2
1981	5,786	176	281	279	376	356	500	609	727	1,324	4
1980	6,147	196	283	368	389	442	557	686	723	1,296	2
1979	6,112	215	268	331	395	482	549	679	772	1,244	3
Falls (E880-E888)											
White, female											
1998	7,773	75	85	85	134	235	485	849	1,416	4,210	-
1997	7,300	68	79	88	123	227	443	763	1,329	3,963	-
1996	7,138	61	83	80	109	204	423	785	1,342	3,845	3
1995	6,542	44	60	78	119	202	451	706	1,154	3,531	1
1994	6,194	53	49	78	107	224	375	687	1,181	3,270	-
1993	6,129	61	56	75	140	231	348	692	1,134	3,234	-
1992	5,914	51	62	77	120	224	350	656	1,110	3,074	-
1991	5,885	40	44	65	115	218	395	681	1,150	2,980	1
1990	5,566	44	42	95	139	226	350	745	989	2,759	-
1989	5,437	44	56	75	148	208	378	666	1,005	2,674	1
1988	5,388	35	55	73	128	237	379	650	1,026	2,626	-
1987	5,141	39	54	94	162	237	367	626	996	2,369	2
1986	5,051	35	67	77	136	220	374	638	949	2,341	3
1985	5,219	43	55	103	168	240	383	657	983	2,407	1
1984	5,345	57	81	92	166	225	417	693	1,017	2,413	1
1983	5,348	48	71	104	160	239	396	642	986	2,465	2
1982	5,343	60	85	108	172	237	396	666	1,021	2,382	1
1981	5,631	48	94	115	161	234	467	682	1,111	2,518	-
1980	5,869	62	91	135	211	253	407	746	1,132	2,590	1
1979	5,887	54	99	148	189	257	481	777	1,216	2,455	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
All other, both sexes											
1998	1,237	73	85	52	79	80	108	116	131	284	1
1997	1,170	75	70	46	61	74	99	119	124	262	-
1996	1,137	86	67	58	68	85	89	108	108	238	-
1995	1,136	67	52	54	64	90	100	108	131	203	1
1994	1,108	64	55	62	85	66	71	88	130	199	1
1993	1,127	52	49	63	63	83	94	105	110	209	-
1992	1,086	70	53	68	67	69	86	112	95	214	1
1991	1,106	63	57	73	55	65	80	79	101	223	1
1990	1,182	65	71	60	82	78	89	98	125	200	2
1989	1,190	81	76	55	72	99	95	99	94	197	-
1988	1,210	53	72	71	75	86	81	113	109	190	4
1987	1,201	70	52	76	92	70	96	100	111	176	2
1986	1,076	62	58	72	81	66	85	81	79	151	-
1985	1,203	64	62	85	82	77	102	99	105	167	-
1984	1,108	71	63	87	78	78	89	78	86	167	1
1983	1,147	72	79	77	83	85	92	68	113	141	1
1982	1,147	75	75	91	79	75	85	104	90	139	2
1981	1,211	73	99	88	106	82	82	87	97	143	4
1980	1,278	94	107	80	100	86	95	105	90	140	2
1979	1,217	85	100	79	75	93	94	100	68	145	-
Falls (E880-E888)											
All other, male											
1998	771	62	77	43	58	52	67	51	60	120	1
1997	728	66	54	35	47	50	50	61	54	111	-
1996	718	71	60	48	47	65	53	55	45	83	-
1995	718	58	37	45	44	60	61	52	63	78	1
1994	741	50	48	50	67	40	44	56	73	80	-
1993	735	41	42	46	50	58	62	51	51	85	-
1992	669	55	40	50	53	46	54	56	42	79	1
1991	705	53	43	57	42	49	45	36	52	79	1
1990	746	50	56	47	58	48	50	50	55	77	2
1989	778	65	64	44	59	60	51	54	51	63	-
1988	805	43	59	58	60	68	49	63	57	69	4
1987	812	56	35	63	67	51	65	55	53	70	1
1986	726	53	45	60	48	49	45	38	45	68	-
1985	813	52	49	64	56	63	61	53	51	71	-
1984	726	52	54	67	57	52	51	35	41	66	-
1983	750	53	57	59	65	58	59	41	49	53	1
1982	767	59	65	72	58	47	50	58	46	59	1
1981	841	57	79	63	82	62	54	49	47	58	3
1980	889	76	87	62	77	63	54	51	51	59	-
1979	816	66	86	68	57	60	59	47	32	45	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
All other, female											
1998	466	11	8	9	21	28	41	65	71	164	-
1997	442	9	16	11	14	24	49	58	70	151	-
1996	419	15	7	10	21	20	36	53	63	155	-
1995	418	9	15	9	20	30	39	56	68	125	-
1994	367	14	7	12	18	26	27	32	57	119	1
1993	392	11	7	17	13	25	32	54	59	124	-
1992	417	15	13	18	14	23	32	56	53	135	-
1991	401	10	14	16	13	16	35	43	49	144	-
1990	436	15	15	13	24	30	39	48	70	123	-
1989	412	16	12	11	13	39	44	45	43	134	-
1988	405	10	13	13	15	18	32	50	52	121	-
1987	389	14	17	13	25	19	31	45	58	106	1
1986	350	9	13	12	33	17	40	43	34	83	-
1985	390	12	13	21	26	14	41	46	54	96	-
1984	382	19	9	20	21	26	38	43	45	101	1
1983	397	19	22	18	18	27	33	27	64	88	-
1982	380	16	10	19	21	28	35	46	44	80	1
1981	370	16	20	25	24	20	28	38	50	85	1
1980	389	18	20	18	23	23	41	54	39	81	2
1979	401	19	14	11	18	33	35	53	36	100	-
Falls (E880-E888)											
Black, both sexes											
1998	929	63	67	47	59	58	80	87	87	205	1
1997	868	58	55	34	44	59	70	84	87	189	-
1996	892	75	61	45	62	67	67	82	76	173	-
1995	877	54	41	41	53	68	82	82	97	149	1
1994	871	56	48	51	75	54	52	62	99	142	-
1993	908	46	46	49	52	65	72	85	81	161	-
1992	875	60	45	62	57	61	62	90	68	160	1
1991	927	53	53	63	41	59	67	68	77	181	1
1990	1,005	61	63	54	77	65	71	79	96	159	2
1989	1,009	66	67	49	67	90	78	83	73	162	-
1988	1,017	48	66	64	65	71	67	91	83	150	4
1987	1,029	68	47	66	78	58	78	84	86	145	2
1986	922	59	51	63	75	55	79	66	68	110	-
1985	1,048	61	57	79	77	67	88	81	90	132	-
1984	976	66	61	84	74	63	79	69	75	136	-
1983	1,018	61	68	74	77	77	80	58	100	120	-
1982	1,027	69	67	87	71	67	73	93	78	122	2
1981	1,074	70	92	82	99	76	70	72	82	124	4
1980	1,124	91	100	70	93	77	82	91	73	115	2
1979	1,110	81	92	72	65	87	88	90	66	127	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Falls (E880-E888)											
Black, male											
1998	578	53	59	39	43	41	49	33	40	80	1
1997	542	52	45	27	33	41	32	46	37	72	-
1996	571	63	55	36	44	50	42	43	28	60	-
1995	544	47	30	34	37	46	52	37	44	50	1
1994	571	45	42	41	59	33	28	41	51	48	-
1993	595	36	39	35	42	49	45	43	34	63	-
1992	535	45	34	45	46	41	36	47	30	52	1
1991	586	43	40	48	32	44	41	29	33	60	1
1990	629	48	50	41	55	41	42	39	35	52	2
1989	659	53	57	40	54	58	38	45	39	48	-
1988	678	39	53	52	53	57	40	53	40	47	4
1987	706	56	31	56	58	41	55	44	40	56	1
1986	632	50	40	56	45	41	42	31	37	48	-
1985	711	49	44	62	54	56	53	40	45	54	-
1984	642	48	52	65	53	44	42	30	34	54	-
1983	665	46	48	56	59	52	49	34	44	46	-
1982	680	54	57	70	51	42	43	50	38	47	1
1981	750	54	75	59	77	57	47	39	37	49	3
1980	776	74	81	55	71	56	44	43	40	46	-
1979	739	62	80	61	49	58	56	41	31	37	-
Falls (E880-E888)											
Black, female											
1998	351	10	8	8	16	17	31	54	47	125	-
1997	326	6	10	7	11	18	38	38	50	117	-
1996	321	12	6	9	18	17	25	39	48	113	-
1995	333	7	11	7	16	22	30	45	53	99	-
1994	300	11	6	10	16	21	24	21	48	94	-
1993	313	10	7	14	10	16	27	42	47	98	-
1992	340	15	11	17	11	20	26	43	38	108	-
1991	341	10	13	15	9	15	26	39	44	121	-
1990	376	13	13	13	22	24	29	40	61	107	-
1989	350	13	10	9	13	32	40	38	34	114	-
1988	339	9	13	12	12	14	27	38	43	103	-
1987	323	12	16	10	20	17	23	40	46	89	1
1986	290	9	11	7	30	14	37	35	31	62	-
1985	337	12	13	17	23	11	35	41	45	78	-
1984	334	18	9	19	21	19	37	39	41	82	-
1983	353	15	20	18	18	25	31	24	56	74	-
1982	347	15	10	17	20	25	30	43	40	75	1
1981	324	16	17	23	22	19	23	33	45	75	1
1980	348	17	19	15	22	21	38	48	33	69	2
1979	371	19	12	11	16	29	32	49	35	90	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
All races, both sexes											
1998	866	44	33	40	23	15	19	12	9	9	1
1997	981	51	45	41	17	29	22	14	10	15	-
1996	1,134	67	31	36	31	21	39	13	15	10	-
1995	1,225	54	38	28	27	25	27	19	15	2	-
1994	1,356	55	40	23	23	23	23	23	9	13	1
1993	1,521	74	39	34	24	29	23	28	9	13	-
1992	1,409	52	40	42	32	20	30	15	16	12	2
1991	1,441	60	42	44	29	34	21	16	23	14	2
1990	1,416	36	39	44	30	24	26	22	8	8	1
1989	1,489	64	32	42	31	42	24	22	12	8	2
1988	1,501	55	49	35	35	41	20	21	13	11	1
1987	1,440	57	51	48	45	22	28	25	12	8	1
1986	1,452	56	53	38	49	34	27	23	12	9	3
1985	1,649	68	55	58	50	26	41	16	17	4	3
1984	1,668	79	54	47	42	47	39	18	16	9	1
1983	1,695	69	66	52	42	40	29	31	10	7	-
1982	1,756	59	60	66	42	39	45	15	14	4	2
1981	1,871	78	75	69	60	33	28	13	9	11	2
1980	1,955	82	51	55	45	40	21	21	12	5	1
1979	2,004	83	89	73	55	52	26	20	10	9	-
Accidental discharge of firearms (E922)											
All races, male											
1998	762	38	25	39	21	15	17	10	9	6	1
1997	856	41	35	36	15	24	19	12	8	14	-
1996	1,004	57	27	29	28	18	35	11	14	8	-
1995	1,055	44	30	25	21	24	25	17	13	1	-
1994	1,192	48	31	20	22	21	19	21	9	9	1
1993	1,328	62	33	29	24	24	20	23	8	9	-
1992	1,238	46	34	35	26	18	26	11	12	11	2
1991	1,304	55	34	40	26	28	19	15	22	10	1
1990	1,255	34	29	42	26	20	22	19	8	8	1
1989	1,317	58	28	38	24	40	20	20	12	8	2
1988	1,304	49	44	31	26	38	18	16	11	7	1
1987	1,249	47	44	41	35	22	26	21	11	6	1
1986	1,254	49	44	34	39	32	24	22	11	9	3
1985	1,415	61	45	48	47	23	39	14	14	2	2
1984	1,461	70	51	42	38	40	36	16	15	9	1
1983	1,465	60	60	47	36	32	24	26	9	7	-
1982	1,524	52	50	59	34	34	39	13	12	4	2
1981	1,592	64	66	62	54	29	22	12	8	9	1
1980	1,686	67	42	50	37	37	21	20	11	4	1
1979	1,715	62	73	55	50	48	22	19	9	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
All races, female											
1998	104	6	8	1	2	-	2	2	-	3	-
1997	125	10	10	5	2	5	3	2	2	1	-
1996	130	10	4	7	3	3	4	2	1	2	-
1995	170	10	8	3	6	1	2	2	2	1	-
1994	164	7	9	3	1	2	4	2	-	4	-
1993	193	12	6	5	-	5	3	5	1	4	-
1992	171	6	6	7	6	2	4	4	4	1	-
1991	137	5	8	4	3	6	2	1	1	4	1
1990	161	2	10	2	4	4	4	3	-	-	-
1989	172	6	4	4	7	2	4	2	-	-	-
1988	197	6	5	4	9	3	2	5	2	4	-
1987	191	10	7	7	10	-	2	4	1	2	-
1986	198	7	9	4	10	2	3	1	1	-	-
1985	234	7	10	10	3	3	2	2	3	2	1
1984	207	9	3	5	4	7	3	2	1	-	-
1983	230	9	6	5	6	8	5	5	1	-	-
1982	232	7	10	7	8	5	6	2	2	-	-
1981	279	14	9	7	6	4	6	1	1	2	1
1980	269	15	9	5	8	3	-	1	1	1	-
1979	289	21	16	18	5	4	4	1	1	3	-
Accidental discharge of firearms (E922)											
White, both sexes											
1998	672	36	27	37	20	13	17	12	8	9	1
1997	782	47	42	35	15	23	18	13	10	13	-
1996	868	56	28	30	30	18	36	13	15	10	-
1995	946	46	32	27	19	22	24	15	15	2	-
1994	998	48	31	19	21	23	19	23	9	13	1
1993	1,121	64	34	27	20	27	22	25	9	13	-
1992	1,071	47	34	36	31	17	27	14	15	10	2
1991	1,103	55	39	41	25	33	20	14	20	12	2
1990	1,113	29	34	39	23	20	21	20	8	8	1
1989	1,153	57	31	36	28	38	21	21	11	7	1
1988	1,188	49	40	32	32	38	17	16	13	11	1
1987	1,158	47	41	41	38	21	24	23	11	8	1
1986	1,210	48	49	33	48	30	25	21	11	9	3
1985	1,346	49	48	52	44	22	38	15	17	4	2
1984	1,401	69	47	42	38	41	37	18	16	9	1
1983	1,393	56	55	43	38	37	25	30	9	6	-
1982	1,457	50	52	62	37	35	36	13	12	4	2
1981	1,523	65	66	58	51	26	27	10	4	11	2
1980	1,582	67	42	52	37	32	18	18	10	4	1
1979	1,605	67	76	54	48	45	21	20	9	6	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
White, male											
1998	594	30	21	36	18	13	15	10	8	6	1
1997	680	37	34	31	13	19	15	11	8	12	-
1996	764	48	24	23	27	16	32	11	14	8	-
1995	801	37	27	24	14	21	22	13	13	1	-
1994	876	43	26	17	20	21	16	21	9	9	1
1993	978	53	29	23	20	22	20	21	8	9	-
1992	936	41	28	30	25	15	23	10	12	10	2
1991	988	51	31	38	22	27	18	13	20	9	1
1990	997	27	27	37	19	16	18	18	8	8	1
1989	1,023	53	27	33	22	36	18	19	11	7	1
1988	1,041	43	36	29	23	35	15	12	11	7	1
1987	1,007	37	35	36	30	21	22	20	10	6	1
1986	1,053	43	41	30	38	28	22	21	10	9	3
1985	1,160	47	39	43	41	20	36	13	14	2	1
1984	1,230	62	45	38	34	36	35	16	15	9	1
1983	1,208	48	51	38	32	30	21	25	8	6	-
1982	1,266	43	43	55	32	30	31	11	10	4	2
1981	1,316	53	58	52	46	24	21	9	4	9	1
1980	1,374	54	35	47	32	29	18	17	9	4	1
1979	1,390	51	62	41	44	44	18	19	9	5	-
Accidental discharge of firearms (E922)											
White, female											
1998	78	6	6	1	2	-	2	2	-	3	-
1997	102	10	8	4	2	4	3	2	2	1	-
1996	104	8	4	7	3	2	4	2	1	2	-
1995	145	9	5	3	5	1	2	2	2	1	-
1994	122	5	5	2	1	2	3	2	-	4	-
1993	143	11	5	4	-	5	2	4	1	4	-
1992	135	6	6	6	6	2	4	4	3	-	-
1991	115	4	8	3	3	6	2	1	-	3	1
1990	116	2	7	2	4	4	3	2	-	-	-
1989	130	4	4	3	6	2	3	2	-	-	-
1988	147	6	4	3	9	3	2	4	2	4	-
1987	151	10	6	5	8	-	2	3	1	2	-
1986	157	5	8	3	10	2	3	-	1	-	-
1985	186	2	9	9	3	2	2	2	3	2	1
1984	171	7	2	4	4	5	2	2	1	-	-
1983	185	8	4	5	6	7	4	5	1	-	-
1982	191	7	9	7	5	5	5	2	2	-	-
1981	207	12	8	6	5	2	6	1	-	2	1
1980	208	13	7	5	5	3	-	1	1	-	-
1979	215	16	14	13	4	1	3	1	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
All other, both sexes											
1998	194	8	6	3	3	2	2	-	1	-	-
1997	199	4	3	6	2	6	4	1	-	2	-
1996	266	11	3	6	1	3	3	-	-	-	-
1995	279	8	6	1	8	3	3	4	-	-	-
1994	358	7	9	4	2	-	4	-	-	-	-
1993	400	10	5	7	4	2	1	3	-	-	-
1992	338	5	6	6	1	3	3	1	1	2	-
1991	338	5	3	3	4	1	1	2	3	2	-
1990	303	7	5	5	7	4	5	2	-	-	-
1989	336	7	1	6	3	4	3	1	1	1	1
1988	313	6	9	3	3	3	3	5	-	-	-
1987	282	10	10	7	7	1	4	2	1	-	-
1986	242	8	4	5	1	4	2	2	1	-	-
1985	303	19	7	6	6	4	3	1	-	-	1
1984	267	10	7	5	4	6	2	-	-	-	-
1983	302	13	11	9	4	3	4	1	1	1	-
1982	299	9	8	4	5	4	9	2	2	-	-
1981	348	13	9	11	9	7	1	3	5	-	-
1980	373	15	9	3	8	8	3	3	2	1	-
1979	399	16	13	19	7	7	5	-	1	3	-
Accidental discharge of firearms (E922)											
All other, male											
1998	168	8	4	3	3	2	2	-	1	-	-
1997	176	4	1	5	2	5	4	1	-	2	-
1996	240	9	3	6	1	2	3	-	-	-	-
1995	254	7	3	1	7	3	3	4	-	-	-
1994	316	5	5	3	2	-	3	-	-	-	-
1993	350	9	4	6	4	2	-	2	-	-	-
1992	302	5	6	5	1	3	3	1	-	1	-
1991	316	4	3	2	4	1	1	2	2	1	-
1990	258	7	2	5	7	4	4	1	-	-	-
1989	294	5	1	5	2	4	2	1	1	1	1
1988	263	6	8	2	3	3	3	4	-	-	-
1987	242	10	9	5	5	1	4	1	1	-	-
1986	201	6	3	4	1	4	2	1	1	-	-
1985	255	14	6	5	6	3	3	1	-	-	1
1984	231	8	6	4	4	4	1	-	-	-	-
1983	257	12	9	9	4	2	3	1	1	1	-
1982	258	9	7	4	2	4	8	2	2	-	-
1981	276	11	8	10	8	5	1	3	4	-	-
1980	312	13	7	3	5	8	3	3	2	-	-
1979	325	11	11	14	6	4	4	-	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
All other, female											
1998	26	-	2	-	-	-	-	-	-	-	-
1997	23	-	2	1	-	1	-	-	-	-	-
1996	26	2	-	-	-	1	-	-	-	-	-
1995	25	1	3	-	1	-	-	-	-	-	-
1994	42	2	4	1	-	-	1	-	-	-	-
1993	50	1	1	1	-	-	1	1	-	-	-
1992	36	-	-	1	-	-	-	-	1	1	-
1991	22	1	-	1	-	-	-	-	1	1	-
1990	45	-	3	-	-	-	1	1	-	-	-
1989	42	2	-	1	1	-	1	-	-	-	-
1988	50	-	1	1	-	-	-	1	-	-	-
1987	40	-	1	2	2	-	-	1	-	-	-
1986	41	2	1	1	-	-	-	1	-	-	-
1985	48	5	1	1	-	1	-	-	-	-	-
1984	36	2	1	1	-	2	1	-	-	-	-
1983	45	1	2	-	-	1	1	-	-	-	-
1982	41	-	1	-	3	-	1	-	-	-	-
1981	72	2	1	1	1	2	-	-	1	-	-
1980	61	2	2	-	3	-	-	-	-	1	-
1979	74	5	2	5	1	3	1	-	1	2	-
Accidental discharge of firearms (E922)											
Black, both sexes											
1998	174	7	5	3	3	2	2	-	1	-	-
1997	173	4	3	5	1	4	4	1	-	2	-
1996	244	8	3	6	1	3	3	-	-	-	-
1995	245	6	6	1	8	3	3	4	-	-	-
1994	313	6	8	4	2	-	4	-	-	-	-
1993	360	9	5	6	4	2	1	3	-	-	-
1992	300	5	6	6	1	2	3	1	1	1	-
1991	290	4	3	2	4	1	-	2	3	2	-
1990	259	7	4	4	5	4	4	2	-	-	-
1989	276	6	1	5	3	4	3	1	1	1	1
1988	266	4	8	2	3	2	3	5	-	-	-
1987	247	8	10	5	7	1	4	2	1	-	-
1986	211	8	4	5	1	4	2	2	1	-	-
1985	251	16	7	5	6	4	3	1	-	-	1
1984	227	9	6	5	3	6	2	-	-	-	-
1983	255	12	7	9	4	3	4	1	1	1	-
1982	248	7	7	3	5	4	8	2	2	-	-
1981	305	13	7	11	8	6	1	3	5	-	-
1980	335	13	9	3	8	7	3	3	2	1	-
1979	354	14	13	16	7	6	5	-	1	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental discharge of firearms (E922)											
Black, male											
1998	154	7	4	3	3	2	2	-	1	-	-
1997	152	4	1	4	1	3	4	1	-	2	-
1996	219	6	3	6	1	2	3	-	-	-	-
1995	225	5	3	1	7	3	3	4	-	-	-
1994	274	4	4	3	2	-	3	-	-	-	-
1993	317	8	4	5	4	2	-	2	-	-	-
1992	267	5	6	5	1	2	3	1	-	1	-
1991	270	4	3	1	4	1	-	2	2	1	-
1990	221	7	2	4	5	4	3	1	-	-	-
1989	246	5	1	4	2	4	2	1	1	1	1
1988	223	4	8	2	3	2	3	4	-	-	-
1987	214	8	9	4	5	1	4	1	1	-	-
1986	172	6	3	4	1	4	2	1	1	-	-
1985	213	12	6	4	6	3	3	1	-	-	1
1984	196	7	5	4	3	4	1	-	-	-	-
1983	215	11	6	9	4	2	3	1	1	1	-
1982	208	7	6	3	2	4	7	2	2	-	-
1981	245	11	7	10	8	4	1	3	4	-	-
1980	281	12	7	3	5	7	3	3	2	-	-
1979	283	9	11	11	6	3	4	-	-	1	-
Accidental discharge of firearms (E922)											
Black, female											
1998	20	-	1	-	-	-	-	-	-	-	-
1997	21	-	2	1	-	1	-	-	-	-	-
1996	25	2	-	-	-	1	-	-	-	-	-
1995	20	1	3	-	1	-	-	-	-	-	-
1994	39	2	4	1	-	-	1	-	-	-	-
1993	43	1	1	1	-	-	1	1	-	-	-
1992	33	-	-	1	-	-	-	-	1	-	-
1991	20	-	-	1	-	-	-	-	1	1	-
1990	38	-	2	-	-	-	1	1	-	-	-
1989	30	1	-	1	1	-	1	-	-	-	-
1988	43	-	-	-	-	-	-	1	-	-	-
1987	33	-	1	1	2	-	-	1	-	-	-
1986	39	2	1	1	-	-	-	1	-	-	-
1985	38	4	1	1	-	1	-	-	-	-	-
1984	31	2	1	1	-	2	1	-	-	-	-
1983	40	1	1	-	-	1	1	-	-	-	-
1982	40	-	1	-	3	-	1	-	-	-	-
1981	60	2	-	1	-	2	-	-	1	-	-
1980	54	1	2	-	3	-	-	-	-	1	-
1979	71	5	2	5	1	3	1	-	1	2	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
All races, both sexes											
1998	3,964	221	157	114	92	96	104	103	80	71	43
1997	3,561	195	139	102	112	79	101	95	78	67	25
1996	3,488	167	141	97	72	98	87	85	65	69	20
1995	3,790	179	128	122	107	90	88	83	66	65	36
1994	3,404	152	109	95	91	79	70	99	88	63	27
1993	3,807	166	100	110	96	98	93	63	75	70	38
1992	3,524	131	122	96	92	99	89	77	67	68	43
1991	3,967	146	98	116	119	89	100	79	54	60	36
1990	3,979	137	101	97	116	111	97	81	68	67	30
1989	4,015	146	106	103	89	133	89	90	72	69	30
1988	4,199	138	105	104	117	88	95	102	61	76	28
1987	4,360	138	89	127	116	96	99	81	52	54	37
1986	4,777	141	129	103	125	95	109	97	79	63	42
1985	4,407	132	94	102	118	121	97	64	73	55	38
1984	4,444	151	126	143	136	104	105	93	82	54	24
1983	5,254	146	168	157	119	103	92	77	58	57	33
1982	5,258	170	164	142	138	124	101	86	64	50	21
1981	5,233	149	159	160	115	123	84	74	49	50	23
1980	6,043	167	162	181	131	139	111	76	70	55	33
1979	5,678	173	153	157	140	123	97	91	59	66	49
Accidental drowning and submersion (E910)											
All races, male											
1998	3,161	183	132	91	68	67	74	74	48	43	43
1997	2,800	159	111	83	90	59	74	53	53	33	24
1996	2,771	145	114	78	56	79	65	58	35	41	19
1995	3,028	143	112	100	85	66	65	52	38	43	36
1994	2,688	122	88	76	72	60	43	62	48	35	25
1993	3,039	137	85	90	79	66	61	40	49	36	35
1992	2,791	115	101	74	71	74	69	49	39	35	43
1991	3,242	122	78	101	90	74	81	53	31	38	34
1990	3,203	109	87	78	93	82	71	46	33	40	30
1989	3,249	124	82	82	75	102	59	58	40	42	28
1988	3,346	116	85	82	86	62	63	66	42	37	26
1987	3,529	120	79	100	96	70	76	44	31	28	36
1986	3,856	111	108	87	80	67	76	64	37	33	41
1985	3,565	107	80	76	94	88	60	40	47	25	37
1984	3,543	122	100	115	99	70	71	57	50	29	24
1983	4,330	124	139	122	98	84	57	48	40	31	33
1982	4,274	144	128	102	105	93	72	57	38	27	20
1981	4,364	123	127	131	94	96	65	46	24	23	22
1980	5,012	136	134	152	102	99	82	54	46	32	32
1979	4,682	142	116	132	119	99	71	64	33	41	48

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
All races, female											
1998	803	38	25	23	24	29	30	29	32	28	-
1997	761	36	28	19	22	20	27	42	25	34	1
1996	717	22	27	19	16	19	22	27	30	28	1
1995	762	36	16	22	22	24	23	31	28	22	-
1994	716	30	21	19	19	19	27	37	40	28	2
1993	768	29	15	20	17	32	32	23	26	34	3
1992	733	16	21	22	21	25	20	28	28	33	-
1991	725	24	20	15	29	15	19	26	23	22	2
1990	776	28	14	19	23	29	26	35	35	27	-
1989	766	22	24	21	14	31	30	32	32	27	2
1988	853	22	20	22	31	26	32	36	19	39	2
1987	831	18	10	27	20	26	23	37	21	26	1
1986	921	30	21	16	45	28	33	33	42	30	1
1985	842	25	14	26	24	33	37	24	26	30	1
1984	901	29	26	28	37	34	34	36	32	25	-
1983	924	22	29	35	21	19	35	29	18	26	-
1982	984	26	36	40	33	31	29	29	26	23	1
1981	869	26	32	29	21	27	19	28	25	27	1
1980	1,031	31	28	29	29	40	29	22	24	23	1
1979	996	31	37	25	21	24	26	27	26	25	1
Accidental drowning and submersion (E910)											
White, both sexes											
1998	3,007	179	119	91	72	86	85	91	69	66	42
1997	2,723	168	117	81	92	64	84	84	72	59	24
1996	2,615	128	108	77	56	78	71	70	55	63	19
1995	2,901	136	109	98	87	68	75	75	51	60	35
1994	2,520	122	79	79	75	61	55	85	77	53	25
1993	2,874	134	81	93	81	75	82	56	65	61	36
1992	2,685	102	100	73	74	76	75	67	61	63	42
1991	2,939	112	73	90	97	61	82	66	44	56	34
1990	3,006	108	73	79	93	91	83	74	60	56	29
1989	3,046	113	79	77	76	109	83	79	69	58	27
1988	3,187	105	85	92	99	72	80	90	54	71	27
1987	3,242	98	75	102	88	83	84	68	49	48	30
1986	3,564	106	93	83	104	78	96	85	72	52	37
1985	3,331	100	71	82	94	108	81	58	64	51	33
1984	3,369	116	101	117	118	86	95	83	75	48	23
1983	4,052	113	132	132	99	83	75	70	53	54	29
1982	4,002	126	130	115	111	95	88	78	62	42	20
1981	3,958	113	134	133	89	110	72	64	42	49	19
1980	4,610	127	127	150	105	118	94	62	61	48	28
1979	4,302	144	111	134	113	107	84	79	54	55	42

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
White, male											
1998	2,361	152	96	72	49	59	61	64	41	40	42
1997	2,126	135	92	69	72	49	62	46	48	28	23
1996	2,057	108	86	61	42	61	54	47	30	37	18
1995	2,300	110	95	78	69	45	56	46	28	40	35
1994	1,968	96	61	64	60	48	35	55	45	32	23
1993	2,275	111	68	76	66	52	56	36	44	31	34
1992	2,126	88	82	53	59	57	61	42	36	34	42
1991	2,360	90	58	79	74	49	65	44	25	34	33
1990	2,390	83	65	65	75	67	64	42	28	35	29
1989	2,434	96	59	60	64	83	55	50	38	35	26
1988	2,507	86	67	71	72	50	53	56	36	35	25
1987	2,587	84	68	79	70	60	64	36	28	25	29
1986	2,845	82	81	68	66	52	68	55	32	31	36
1985	2,660	82	59	63	74	78	49	37	42	23	32
1984	2,636	93	83	92	86	58	61	51	46	26	23
1983	3,287	94	104	101	80	67	42	42	36	29	29
1982	3,217	106	101	81	82	73	65	51	36	21	19
1981	3,241	91	104	105	72	85	55	38	21	23	18
1980	3,782	103	102	126	77	83	70	42	42	28	28
1979	3,512	117	79	113	95	86	60	52	31	35	41
Accidental drowning and submersion (E910)											
White, female											
1998	646	27	23	19	23	27	24	27	28	26	-
1997	597	33	25	12	20	15	22	38	24	31	1
1996	558	20	22	16	14	17	17	23	25	26	1
1995	601	26	14	20	18	23	19	29	23	20	-
1994	552	26	18	15	15	13	20	30	32	21	2
1993	599	23	13	17	15	23	26	20	21	30	2
1992	559	14	18	20	15	19	14	25	25	29	-
1991	579	22	15	11	23	12	17	22	19	22	1
1990	616	25	8	14	18	24	19	32	32	21	-
1989	612	17	20	17	12	26	28	29	31	23	1
1988	680	19	18	21	27	22	27	34	18	36	2
1987	655	14	7	23	18	23	20	32	21	23	1
1986	719	24	12	15	38	26	28	30	40	21	1
1985	671	18	12	19	20	30	32	21	22	28	1
1984	733	23	18	25	32	28	34	32	29	22	-
1983	765	19	28	31	19	16	33	28	17	25	-
1982	785	20	29	34	29	22	23	27	26	21	1
1981	717	22	30	28	17	25	17	26	21	26	1
1980	828	24	25	24	28	35	24	20	19	20	-
1979	790	27	32	21	18	21	24	27	23	20	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
All other, both sexes											
1998	957	42	38	23	20	10	19	12	11	5	1
1997	838	27	22	21	20	15	17	11	6	8	1
1996	873	39	33	20	16	20	16	15	10	6	1
1995	889	43	19	24	20	22	13	8	15	5	1
1994	884	30	30	16	16	18	15	14	11	10	2
1993	933	32	19	17	15	23	11	7	10	9	2
1992	839	29	22	23	18	23	14	10	6	5	1
1991	1,028	34	25	26	22	28	18	13	10	4	2
1990	973	29	28	18	23	20	14	7	8	11	1
1989	969	33	27	26	13	24	6	11	3	11	3
1988	1,012	33	20	12	18	16	15	12	7	5	1
1987	1,118	40	14	25	28	13	15	13	3	6	7
1986	1,213	35	36	20	21	17	13	12	7	11	5
1985	1,076	32	23	20	24	13	16	6	9	4	5
1984	1,075	35	25	26	18	18	10	10	7	6	1
1983	1,202	33	36	25	20	20	17	7	5	3	4
1982	1,256	44	34	27	27	29	13	8	2	8	1
1981	1,275	36	25	27	26	13	12	10	7	1	4
1980	1,433	40	35	31	26	21	17	14	9	7	5
1979	1,376	29	42	23	27	16	13	12	5	11	7
Accidental drowning and submersion (E910)											
All other, male											
1998	800	31	36	19	19	8	13	10	7	3	1
1997	674	24	19	14	18	10	12	7	5	5	1
1996	714	37	28	17	14	18	11	11	5	4	1
1995	728	33	17	22	16	21	9	6	10	3	1
1994	720	26	27	12	12	12	8	7	3	3	2
1993	764	26	17	14	13	14	5	4	5	5	1
1992	665	27	19	21	12	17	8	7	3	1	1
1991	882	32	20	22	16	25	16	9	6	4	1
1990	813	26	22	13	18	15	7	4	5	5	1
1989	815	28	23	22	11	19	4	8	2	7	2
1988	839	30	18	11	14	12	10	10	6	2	1
1987	942	36	11	21	26	10	12	8	3	3	7
1986	1,011	29	27	19	14	15	8	9	5	2	5
1985	905	25	21	13	20	10	11	3	5	2	5
1984	907	29	17	23	13	12	10	6	4	3	1
1983	1,043	30	35	21	18	17	15	6	4	2	4
1982	1,057	38	27	21	23	20	7	6	2	6	1
1981	1,123	32	23	26	22	11	10	8	3	-	4
1980	1,230	33	32	26	25	16	12	12	4	4	4
1979	1,170	25	37	19	24	13	11	12	2	6	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
All other, female											
1998	157	11	2	4	1	2	6	2	4	2	-
1997	164	3	3	7	2	5	5	4	1	3	-
1996	159	2	5	3	2	2	5	4	5	2	-
1995	161	10	2	2	4	1	4	2	5	2	-
1994	164	4	3	4	4	6	7	7	8	7	-
1993	169	6	2	3	2	9	6	3	5	4	1
1992	174	2	3	2	6	6	6	3	3	4	-
1991	146	2	5	4	6	3	2	4	4	-	1
1990	160	3	6	5	5	5	7	3	3	6	-
1989	154	5	4	4	2	5	2	3	1	4	1
1988	173	3	2	1	4	4	5	2	1	3	-
1987	176	4	3	4	2	3	3	5	-	3	-
1986	202	6	9	1	7	2	5	3	2	9	-
1985	171	7	2	7	4	3	5	3	4	2	-
1984	168	6	8	3	5	6	-	4	3	3	-
1983	159	3	1	4	2	3	2	1	1	1	-
1982	199	6	7	6	4	9	6	2	-	2	-
1981	152	4	2	1	4	2	2	2	4	1	-
1980	203	7	3	5	1	5	5	2	5	3	1
1979	206	4	5	4	3	3	2	-	3	5	-
Accidental drowning and submersion (E910)											
Black, both sexes											
1998	758	33	31	18	13	7	16	10	10	2	1
1997	623	20	20	16	13	11	11	8	4	3	-
1996	648	30	21	14	13	14	11	8	7	4	1
1995	708	39	17	20	17	18	10	7	9	4	1
1994	685	21	23	8	12	13	11	11	6	6	1
1993	710	25	15	13	11	20	9	2	9	5	2
1992	655	24	18	19	12	15	11	4	4	4	1
1991	877	30	22	22	13	21	16	9	6	3	2
1990	777	27	21	13	20	15	9	6	7	8	1
1989	797	26	21	24	11	22	4	9	3	8	2
1988	821	28	19	11	13	12	13	4	7	4	1
1987	950	33	12	21	25	11	11	10	3	5	4
1986	1,004	29	27	15	19	14	9	10	6	5	5
1985	911	28	18	16	17	13	14	5	9	4	4
1984	898	27	22	23	12	16	5	5	4	4	1
1983	1,028	27	27	22	15	17	13	6	5	2	3
1982	1,074	41	31	22	20	25	11	5	1	4	1
1981	1,102	31	23	23	23	11	10	9	5	1	3
1980	1,247	34	33	27	24	17	13	12	8	6	5
1979	1,205	27	34	19	24	13	11	8	3	7	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental drowning and submersion (E910)											
Black, male											
1998	651	28	30	16	13	5	11	8	6	1	1
1997	507	18	17	10	12	7	9	4	4	1	-
1996	549	29	20	12	11	14	11	7	2	2	1
1995	582	30	15	18	13	17	7	6	8	3	1
1994	558	18	20	8	9	9	5	5	2	3	1
1993	586	20	14	11	10	13	4	1	4	2	1
1992	533	22	15	18	8	12	6	2	2	1	1
1991	761	29	17	19	9	19	14	7	4	3	1
1990	655	24	15	11	16	10	4	4	5	3	1
1989	664	23	18	20	9	17	3	6	2	5	1
1988	685	26	17	10	9	9	8	3	6	2	1
1987	808	30	9	19	23	8	8	7	3	3	4
1986	836	25	22	14	13	13	6	7	4	-	5
1985	776	22	17	10	14	10	10	2	5	2	4
1984	767	22	15	21	8	12	5	3	2	2	1
1983	895	24	26	18	13	15	11	5	4	2	3
1982	914	35	25	16	17	17	6	4	1	4	1
1981	971	28	21	22	20	9	8	8	2	-	3
1980	1,083	28	30	23	24	13	8	11	4	4	4
1979	1,034	23	31	15	21	10	9	8	2	3	7
Accidental drowning and submersion (E910)											
Black, female											
1998	107	5	1	2	-	2	5	2	4	1	-
1997	116	2	3	6	1	4	2	4	-	2	-
1996	99	1	1	2	2	-	-	1	5	2	-
1995	126	9	2	2	4	1	3	1	1	1	-
1994	127	3	3	-	3	4	6	6	4	3	-
1993	124	5	1	2	1	7	5	1	5	3	1
1992	122	2	3	1	4	3	5	2	2	3	-
1991	116	1	5	3	4	2	2	2	2	-	1
1990	122	3	6	2	4	5	5	2	2	5	-
1989	133	3	3	4	2	5	1	3	1	3	1
1988	136	2	2	1	4	3	5	1	1	2	-
1987	142	3	3	2	2	3	3	3	-	2	-
1986	168	4	5	1	6	1	3	3	2	5	-
1985	135	6	1	6	3	3	4	3	4	2	-
1984	131	5	7	2	4	4	-	2	2	2	-
1983	133	3	1	4	2	2	2	1	1	-	-
1982	160	6	6	6	3	8	5	1	-	-	-
1981	131	3	2	1	3	2	2	1	3	1	-
1980	164	6	3	4	-	4	5	1	4	2	1
1979	171	4	3	4	3	3	2	-	1	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
All races, both sexes											
1998	3,255	183	178	153	155	191	206	251	224	244	2
1997	3,490	186	183	174	181	193	212	256	236	289	10
1996	3,741	193	173	173	183	197	254	266	244	322	7
1995	3,761	223	157	168	205	206	244	250	228	275	2
1994	3,986	203	149	150	208	200	227	240	250	249	3
1993	3,900	179	153	127	170	225	238	223	241	283	5
1992	3,958	194	151	162	231	233	233	259	214	262	6
1991	4,120	154	150	145	187	234	251	280	205	264	6
1990	4,175	171	172	162	218	256	252	257	249	267	12
1989	4,716	181	163	207	253	284	265	310	261	286	12
1988	4,965	196	197	236	295	286	312	311	279	299	12
1987	4,710	173	179	228	239	281	329	292	261	279	3
1986	4,835	178	191	207	265	252	310	281	265	275	11
1985	4,938	185	224	259	297	289	301	334	271	276	16
1984	5,010	204	214	254	274	280	344	292	228	281	8
1983	5,028	180	226	272	296	305	298	302	275	293	10
1982	5,210	196	223	302	309	295	353	303	257	296	9
1981	5,697	235	307	328	318	366	356	322	302	303	2
1980	5,822	254	319	324	348	405	344	347	291	275	10
1979	5,991	264	316	341	395	394	349	367	275	296	10
Accidental exposure to smoke, fire and flames (E890-E899)											
All races, male											
1998	1,941	121	122	85	101	120	121	122	101	111	2
1997	2,086	130	120	111	108	126	124	122	104	143	10
1996	2,230	131	124	119	125	119	134	133	125	142	6
1995	2,232	162	108	107	137	117	130	129	119	119	2
1994	2,443	135	108	90	131	112	114	133	135	112	3
1993	2,326	110	106	93	110	131	130	109	113	124	4
1992	2,507	140	114	109	163	142	143	134	103	123	6
1991	2,511	101	106	105	126	148	130	152	104	132	5
1990	2,571	111	113	115	159	150	138	140	121	111	6
1989	2,817	120	112	137	169	176	142	153	113	138	11
1988	3,067	142	133	158	187	179	172	165	134	156	11
1987	2,863	111	126	143	154	163	181	143	122	146	2
1986	2,984	128	141	146	173	155	178	152	130	135	8
1985	3,008	131	157	177	197	178	166	170	134	138	13
1984	3,141	151	142	176	179	169	188	158	106	140	6
1983	3,066	130	156	177	199	184	162	170	140	151	6
1982	3,180	133	145	197	205	166	183	170	121	143	6
1981	3,493	155	214	218	209	240	208	178	157	151	2
1980	3,565	173	225	214	231	249	182	185	145	134	10
1979	3,712	184	210	229	252	234	200	200	146	149	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
All races, female											
1998	1,314	62	56	68	54	71	85	129	123	133	-
1997	1,404	56	63	63	73	67	88	134	132	146	-
1996	1,511	62	49	54	58	78	120	133	119	180	1
1995	1,529	61	49	61	68	89	114	121	109	156	-
1994	1,543	68	41	60	77	88	113	107	115	137	-
1993	1,574	69	47	34	60	94	108	114	128	159	1
1992	1,451	54	37	53	68	91	90	125	111	139	-
1991	1,609	53	44	40	61	86	121	128	101	132	1
1990	1,604	60	59	47	59	106	114	117	128	156	6
1989	1,899	61	51	70	84	108	123	157	148	148	1
1988	1,898	54	64	78	108	107	140	146	145	143	1
1987	1,847	62	53	85	85	118	148	149	139	133	1
1986	1,851	50	50	61	92	97	132	129	135	140	3
1985	1,930	54	67	82	100	111	135	164	137	138	3
1984	1,869	53	72	78	95	111	156	134	122	141	2
1983	1,962	50	70	95	97	121	136	132	135	142	4
1982	2,030	63	78	105	104	129	170	133	136	153	3
1981	2,204	80	93	110	109	126	148	144	145	152	-
1980	2,257	81	94	110	117	156	162	162	146	141	-
1979	2,279	80	106	112	143	160	149	167	129	147	1
Accidental exposure to smoke, fire and flames (E890-E899)											
White, both sexes											
1998	2,316	124	131	111	116	146	147	196	187	177	1
1997	2,405	120	126	121	129	132	158	182	192	205	7
1996	2,602	149	118	122	124	130	182	198	194	231	5
1995	2,524	156	113	108	129	143	169	187	155	194	-
1994	2,686	140	97	99	153	141	157	173	159	179	2
1993	2,568	119	111	82	120	148	168	168	169	189	4
1992	2,696	133	102	105	153	172	163	180	154	169	6
1991	2,796	105	103	100	128	163	181	207	147	188	5
1990	2,848	125	121	111	160	189	182	163	192	187	9
1989	3,159	122	109	139	175	195	186	214	205	181	7
1988	3,305	132	113	159	199	205	212	225	201	221	3
1987	3,158	117	112	163	168	188	231	211	181	203	-
1986	3,289	116	134	134	181	184	235	206	205	191	7
1985	3,461	127	158	183	217	217	225	265	209	189	11
1984	3,436	138	154	174	198	200	235	204	159	206	5
1983	3,408	129	161	186	204	216	195	229	193	218	6
1982	3,659	139	148	225	233	224	258	222	180	225	8
1981	3,946	159	220	222	224	262	254	236	229	221	1
1980	4,155	190	226	227	272	298	268	257	220	205	8
1979	4,182	183	230	256	283	281	262	259	203	213	8

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
White, male											
1998	1,383	81	82	61	69	94	82	92	86	79	1
1997	1,459	85	79	78	77	89	91	90	83	102	7
1996	1,568	95	85	83	82	79	93	105	100	99	5
1995	1,509	111	78	75	84	78	87	95	79	82	-
1994	1,670	92	71	55	89	76	82	93	85	80	2
1993	1,547	71	75	59	76	86	89	89	79	72	4
1992	1,715	96	77	64	106	105	98	102	74	70	6
1991	1,710	71	71	64	83	104	91	107	71	99	4
1990	1,765	83	76	76	121	105	99	83	95	78	5
1989	1,903	76	76	92	117	116	98	109	85	89	7
1988	2,025	92	72	105	127	124	111	113	92	114	3
1987	1,911	77	71	102	102	109	121	98	89	109	-
1986	2,048	80	103	88	120	111	131	107	100	102	6
1985	2,135	88	110	125	147	134	122	135	100	100	10
1984	2,186	99	103	123	130	122	123	112	75	102	3
1983	2,138	93	115	122	141	128	108	129	99	118	4
1982	2,246	96	95	146	148	120	132	126	84	108	6
1981	2,417	103	152	145	149	167	147	132	114	111	1
1980	2,556	124	157	153	173	179	145	131	112	102	8
1979	2,608	128	148	170	182	164	146	138	100	114	7
Accidental exposure to smoke, fire and flames (E890-E899)											
White, female											
1998	933	43	49	50	47	52	65	104	101	98	-
1997	946	35	47	43	52	43	67	92	109	103	-
1996	1,034	54	33	39	42	51	89	93	94	132	-
1995	1,015	45	35	33	45	65	82	92	76	112	-
1994	1,016	48	26	44	64	65	75	80	74	99	-
1993	1,021	48	36	23	44	62	79	79	90	117	-
1992	981	37	25	41	47	67	65	78	80	99	-
1991	1,086	34	32	36	45	59	90	100	76	89	1
1990	1,083	42	45	35	39	84	83	80	97	109	4
1989	1,256	46	33	47	58	79	88	105	120	92	-
1988	1,280	40	41	54	72	81	101	112	109	107	-
1987	1,247	40	41	61	66	79	110	113	92	94	-
1986	1,241	36	31	46	61	73	104	99	105	89	1
1985	1,326	39	48	58	70	83	103	130	109	89	1
1984	1,250	39	51	51	68	78	112	92	84	104	2
1983	1,270	36	46	64	63	88	87	100	94	100	2
1982	1,413	43	53	79	85	104	126	96	96	117	2
1981	1,529	56	68	77	75	95	107	104	115	110	-
1980	1,599	66	69	74	99	119	123	126	108	103	-
1979	1,574	55	82	86	101	117	116	121	103	99	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
All other, both sexes											
1998	939	59	47	42	39	45	59	55	37	67	1
1997	1,085	66	57	53	52	61	54	74	44	84	3
1996	1,139	44	55	51	59	67	72	68	50	91	2
1995	1,237	67	44	60	76	63	75	63	73	81	2
1994	1,300	63	52	51	55	59	70	67	91	70	1
1993	1,332	60	42	45	50	77	70	55	72	94	1
1992	1,262	61	49	57	78	61	70	79	60	93	-
1991	1,324	49	47	45	59	71	70	73	58	76	1
1990	1,327	46	51	51	58	67	70	94	57	80	3
1989	1,557	59	54	68	78	89	79	96	56	105	5
1988	1,660	64	84	77	96	81	100	86	78	78	9
1987	1,552	56	67	65	71	93	98	81	80	76	3
1986	1,546	62	57	73	84	68	75	75	60	84	4
1985	1,477	58	66	76	80	72	76	69	62	87	5
1984	1,574	66	60	80	76	80	109	88	69	75	3
1983	1,620	51	65	86	92	89	103	73	82	75	4
1982	1,551	57	75	77	76	71	95	81	77	71	1
1981	1,751	76	87	106	94	104	102	86	73	82	1
1980	1,667	64	93	97	76	107	76	90	71	70	2
1979	1,809	81	86	85	112	113	87	108	72	83	2
Accidental exposure to smoke, fire and flames (E890-E899)											
All other, male											
1998	558	40	40	24	32	26	39	30	15	32	1
1997	627	45	41	33	31	37	33	32	21	41	3
1996	662	36	39	36	43	40	41	28	25	43	1
1995	723	51	30	32	53	39	43	34	40	37	2
1994	773	43	37	35	42	36	32	40	50	32	1
1993	779	39	31	34	34	45	41	20	34	52	-
1992	792	44	37	45	57	37	45	32	29	53	-
1991	801	30	35	41	43	44	39	45	33	33	1
1990	806	28	37	39	38	45	39	57	26	33	1
1989	914	44	36	45	52	60	44	44	28	49	4
1988	1,042	50	61	53	60	55	61	52	42	42	8
1987	952	34	55	41	52	54	60	45	33	37	2
1986	936	48	38	58	53	44	47	45	30	33	2
1985	873	43	47	52	50	44	44	35	34	38	3
1984	955	52	39	53	49	47	65	46	31	38	3
1983	928	37	41	55	58	56	54	41	41	33	2
1982	934	37	50	51	57	46	51	44	37	35	-
1981	1,076	52	62	73	60	73	61	46	43	40	1
1980	1,009	49	68	61	58	70	37	54	33	32	2
1979	1,104	56	62	59	70	70	54	62	46	35	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
All other, female											
1998	381	19	7	18	7	19	20	25	22	35	-
1997	458	21	16	20	21	24	21	42	23	43	-
1996	477	8	16	15	16	27	31	40	25	48	1
1995	514	16	14	28	23	24	32	29	33	44	-
1994	527	20	15	16	13	23	38	27	41	38	-
1993	553	21	11	11	16	32	29	35	38	42	1
1992	470	17	12	12	21	24	25	47	31	40	-
1991	523	19	12	4	16	27	31	28	25	43	-
1990	521	18	14	12	20	22	31	37	31	47	2
1989	643	15	18	23	26	29	35	52	28	56	1
1988	618	14	23	24	36	26	39	34	36	36	1
1987	600	22	12	24	19	39	38	36	47	39	1
1986	610	14	19	15	31	24	28	30	30	51	2
1985	604	15	19	24	30	28	32	34	28	49	2
1984	619	14	21	27	27	33	44	42	38	37	-
1983	692	14	24	31	34	33	49	32	41	42	2
1982	617	20	25	26	19	25	44	37	40	36	1
1981	675	24	25	33	34	31	41	40	30	42	-
1980	658	15	25	36	18	37	39	36	38	38	-
1979	705	25	24	26	42	43	33	46	26	48	-
Accidental exposure to smoke, fire and flames (E890-E899)											
Black, both sexes											
1998	874	58	46	36	37	43	54	53	35	64	1
1997	972	59	51	49	50	59	50	69	42	78	3
1996	1,051	43	51	51	57	61	68	67	44	84	2
1995	1,131	58	42	56	70	60	69	60	70	77	2
1994	1,195	59	49	48	52	55	68	66	84	64	1
1993	1,235	51	40	44	49	74	68	52	66	88	1
1992	1,172	57	43	54	76	59	66	71	58	90	-
1991	1,252	44	45	44	58	67	67	71	58	75	1
1990	1,254	39	49	50	56	65	70	89	55	76	3
1989	1,467	55	54	65	73	87	79	92	52	104	5
1988	1,582	61	82	72	94	75	97	84	76	77	9
1987	1,455	53	66	62	66	87	92	78	77	75	3
1986	1,449	58	52	68	80	67	72	73	60	81	2
1985	1,400	55	61	75	77	70	76	66	61	83	5
1984	1,464	62	54	76	74	77	104	86	65	75	3
1983	1,530	47	60	82	86	85	98	70	80	73	3
1982	1,463	53	68	71	73	70	92	76	76	70	1
1981	1,678	74	83	105	92	101	98	82	71	81	1
1980	1,583	60	90	97	74	104	70	88	68	69	2
1979	1,719	79	81	82	105	111	85	105	69	80	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental exposure to smoke, fire and flames (E890-E899)											
Black, male											
1998	510	39	39	21	31	25	34	28	14	30	1
1997	551	38	35	30	30	36	31	30	20	35	3
1996	616	36	36	36	42	39	39	28	22	39	1
1995	661	44	29	30	49	38	40	32	40	33	2
1994	712	41	35	34	39	35	32	40	46	27	1
1993	717	34	29	33	33	43	39	19	29	48	-
1992	731	42	34	43	55	36	44	26	28	50	-
1991	759	27	33	40	42	42	37	44	33	33	1
1990	755	26	35	38	36	43	39	55	25	31	1
1989	858	42	36	44	48	58	44	43	25	48	4
1988	986	47	60	49	58	51	58	51	41	41	8
1987	882	31	55	39	48	50	54	43	30	36	2
1986	870	44	35	54	50	44	45	44	30	31	2
1985	825	40	44	51	49	43	44	34	33	35	3
1984	885	48	34	51	48	45	62	45	28	38	3
1983	870	35	37	54	53	53	51	40	40	32	1
1982	883	34	47	49	54	45	49	40	37	34	-
1981	1,026	51	59	72	58	70	59	42	41	39	1
1980	953	46	67	61	56	69	33	53	30	31	2
1979	1,045	55	58	57	66	70	52	61	43	34	2
Accidental exposure to smoke, fire and flames (E890-E899)											
Black, female											
1998	364	19	7	15	6	18	20	25	21	34	-
1997	421	21	16	19	20	23	19	39	22	43	-
1996	435	7	15	15	15	22	29	39	22	45	1
1995	470	14	13	26	21	22	29	28	30	44	-
1994	483	18	14	14	13	20	36	26	38	37	-
1993	518	17	11	11	16	31	29	33	37	40	1
1992	441	15	9	11	21	23	22	45	30	40	-
1991	493	17	12	4	16	25	30	27	25	42	-
1990	499	13	14	12	20	22	31	34	30	45	2
1989	609	13	18	21	25	29	35	49	27	56	1
1988	596	14	22	23	36	24	39	33	35	36	1
1987	573	22	11	23	18	37	38	35	47	39	1
1986	579	14	17	14	30	23	27	29	30	50	-
1985	575	15	17	24	28	27	32	32	28	48	2
1984	579	14	20	25	26	32	42	41	37	37	-
1983	660	12	23	28	33	32	47	30	40	41	2
1982	580	19	21	22	19	25	43	36	39	36	1
1981	652	23	24	33	34	31	39	40	30	42	-
1980	630	14	23	36	18	35	37	35	38	38	-
1979	674	24	23	25	39	41	33	44	26	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All races, both sexes											
1998	10,808	1,520	716	337	205	148	138	141	138	135	6
1997	10,175	1,377	609	316	187	157	115	141	129	138	19
1996	9,519	1,270	544	304	172	154	139	140	140	165	14
1995	9,075	1,034	523	258	167	142	129	125	105	125	4
1994	9,001	927	438	239	184	154	163	145	112	171	7
1993	8,539	752	391	214	170	144	135	123	115	145	6
1992	7,088	559	310	181	136	160	146	132	125	155	6
1991	6,444	389	250	194	176	166	146	143	148	177	9
1990	5,815	323	228	199	195	155	153	126	122	188	7
1989	6,529	362	256	229	185	172	169	174	139	181	11
1988	6,232	370	217	209	214	192	140	159	122	189	10
1987	5,317	268	209	205	195	169	143	158	133	158	8
1986	5,747	270	236	205	214	166	170	147	145	136	6
1985	5,179	272	217	236	203	154	165	143	133	124	5
1984	4,917	229	258	214	186	181	143	135	107	102	2
1983	4,641	254	230	244	193	153	152	146	104	122	8
1982	4,740	275	257	238	206	172	155	115	101	109	2
1981	4,537	283	308	221	194	153	144	143	109	96	4
1980	4,343	230	273	242	186	157	144	111	102	103	3
1979	4,648	275	286	279	221	158	152	126	87	81	6
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All races, male											
1998	7,911	1,182	495	232	122	82	74	66	61	51	3
1997	7,631	1,085	438	216	116	95	63	57	56	46	19
1996	7,021	981	394	202	101	92	72	65	58	55	12
1995	6,888	817	372	184	110	81	59	62	46	46	4
1994	6,765	725	305	164	115	90	86	79	45	56	6
1993	6,478	587	292	144	112	87	75	55	50	55	6
1992	5,350	420	227	129	89	95	86	63	59	62	6
1991	4,700	270	174	122	104	86	78	65	66	61	6
1990	4,210	222	151	122	123	98	98	70	54	68	5
1989	4,772	262	172	157	112	105	97	78	69	77	7
1988	4,463	280	142	138	119	106	70	83	64	53	9
1987	3,783	197	130	122	119	101	78	80	63	54	7
1986	4,112	190	158	130	127	96	93	76	66	56	5
1985	3,667	187	126	145	124	91	86	77	64	42	5
1984	3,425	147	155	136	106	106	75	73	47	51	1
1983	3,095	157	134	160	120	77	80	75	42	50	5
1982	3,232	177	142	147	121	99	76	57	47	48	1
1981	3,062	169	187	121	115	85	80	80	46	36	2
1980	2,901	151	165	145	98	88	76	46	50	47	3
1979	3,071	158	185	150	121	87	93	68	42	36	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All races, female											
1998	2,897	338	221	105	83	66	64	75	77	84	3
1997	2,544	292	171	100	71	62	52	84	73	92	-
1996	2,498	289	150	102	71	62	67	75	82	110	2
1995	2,187	217	151	74	57	61	70	63	59	79	-
1994	2,236	202	133	75	69	64	77	66	67	115	1
1993	2,061	165	99	70	58	57	60	68	65	90	-
1992	1,738	139	83	52	47	65	60	69	66	93	-
1991	1,744	119	76	72	72	80	68	78	82	116	3
1990	1,605	101	77	77	72	57	55	56	68	120	2
1989	1,757	100	84	72	73	67	72	96	70	104	4
1988	1,769	90	75	71	95	86	70	76	58	136	1
1987	1,534	71	79	83	76	68	65	78	70	104	1
1986	1,635	80	78	75	87	70	77	71	79	80	1
1985	1,512	85	91	91	79	63	79	66	69	82	-
1984	1,492	82	103	78	80	75	68	62	60	51	1
1983	1,546	97	96	84	73	76	72	71	62	72	3
1982	1,508	98	115	91	85	73	79	58	54	61	1
1981	1,475	114	121	100	79	68	64	63	63	60	2
1980	1,442	79	108	97	88	69	68	65	52	56	-
1979	1,577	117	101	129	100	71	59	58	45	45	2
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
White, both sexes											
1998	8,691	1,120	537	246	160	121	122	120	124	121	4
1997	8,065	1,001	455	232	129	128	93	126	114	124	14
1996	7,498	925	403	221	128	127	121	120	121	146	11
1995	7,017	732	377	187	131	110	114	111	90	114	3
1994	6,859	651	302	168	142	117	140	124	95	152	4
1993	6,564	550	270	172	134	114	118	109	104	121	4
1992	5,477	423	220	140	108	129	130	113	115	135	4
1991	4,834	278	167	142	146	129	122	122	133	158	8
1990	4,552	241	177	157	158	128	132	112	110	171	7
1989	5,047	260	192	174	141	143	146	154	123	156	9
1988	4,767	267	165	155	169	163	118	134	108	167	7
1987	4,126	203	159	159	153	137	119	145	118	141	7
1986	4,665	210	193	161	179	138	141	131	126	122	4
1985	4,196	206	168	193	166	128	141	125	116	111	3
1984	3,979	179	208	175	165	164	118	113	90	96	1
1983	3,815	209	195	196	161	121	132	128	90	108	5
1982	3,865	227	197	198	172	148	126	99	93	100	2
1981	3,692	222	244	189	166	132	123	130	94	78	3
1980	3,598	186	233	210	156	131	125	89	89	89	3
1979	3,800	221	235	243	195	137	126	104	78	74	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
White, male											
1998	6,393	884	360	164	86	63	65	52	59	44	2
1997	6,096	796	322	157	71	76	49	50	51	39	14
1996	5,589	707	285	145	70	75	60	59	48	51	10
1995	5,371	579	257	125	85	58	49	58	42	43	3
1994	5,209	508	204	106	83	65	72	69	36	49	3
1993	5,014	424	189	112	86	65	66	47	48	47	4
1992	4,161	310	151	98	67	76	75	51	56	53	4
1991	3,564	201	112	88	82	66	66	59	58	53	5
1990	3,313	163	111	91	98	78	85	60	51	59	5
1989	3,703	178	129	116	78	85	83	67	62	63	6
1988	3,427	200	103	97	90	81	58	70	56	46	6
1987	2,951	152	96	92	92	79	63	73	56	50	6
1986	3,343	146	127	97	102	73	81	65	57	50	4
1985	3,003	140	96	119	99	79	73	67	56	39	3
1984	2,777	112	124	107	93	95	62	57	44	47	-
1983	2,583	128	113	126	96	61	69	63	38	44	4
1982	2,641	144	100	118	99	85	61	47	45	43	1
1981	2,515	125	145	98	96	74	68	73	42	28	1
1980	2,409	117	135	123	81	70	70	35	41	38	3
1979	2,509	119	146	126	103	75	74	58	38	34	1
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
White, female											
1998	2,298	236	177	82	74	58	57	68	65	77	2
1997	1,969	205	133	75	58	52	44	76	63	85	-
1996	1,909	218	118	76	58	52	61	61	73	95	1
1995	1,646	153	120	62	46	52	65	53	48	71	-
1994	1,650	143	98	62	59	52	68	55	59	103	1
1993	1,550	126	81	60	48	49	52	62	56	74	-
1992	1,316	113	69	42	41	53	55	62	59	82	-
1991	1,270	77	55	54	64	63	56	63	75	105	3
1990	1,239	78	66	66	60	50	47	52	59	112	2
1989	1,344	82	63	58	63	58	63	87	61	93	3
1988	1,340	67	62	58	79	82	60	64	52	121	1
1987	1,175	51	63	67	61	58	56	72	62	91	1
1986	1,322	64	66	64	77	65	60	66	69	72	-
1985	1,193	66	72	74	67	49	68	58	60	72	-
1984	1,202	67	84	68	72	69	56	56	46	49	1
1983	1,232	81	82	70	65	60	63	65	52	64	1
1982	1,224	83	97	80	73	63	65	52	48	57	1
1981	1,177	97	99	91	70	58	55	57	52	50	2
1980	1,189	69	98	87	75	61	55	54	48	51	-
1979	1,291	102	89	117	92	62	52	46	40	40	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All other, both sexes											
1998	2,117	400	179	91	45	27	16	21	14	14	2
1997	2,110	376	154	84	58	29	22	15	15	14	5
1996	2,021	345	141	83	44	27	18	20	19	19	3
1995	2,058	302	146	71	36	32	15	14	15	11	1
1994	2,142	276	136	71	42	37	23	21	17	19	3
1993	1,975	202	121	42	36	30	17	14	11	24	2
1992	1,611	136	90	41	28	31	16	19	10	20	2
1991	1,610	111	83	52	30	37	24	21	15	19	1
1990	1,263	82	51	42	37	27	21	14	12	17	-
1989	1,482	102	64	55	44	29	23	20	16	25	2
1988	1,465	103	52	54	45	29	22	25	14	22	3
1987	1,191	65	50	46	42	32	24	13	15	17	1
1986	1,082	60	43	44	35	28	29	16	19	14	2
1985	983	66	49	43	37	26	24	18	17	13	2
1984	938	50	50	39	21	17	25	22	17	6	1
1983	826	45	35	48	32	32	20	18	14	14	3
1982	875	48	60	40	34	24	29	16	8	9	-
1981	845	61	64	32	28	21	21	13	15	18	1
1980	745	44	40	32	30	26	19	22	13	14	-
1979	848	54	51	36	26	21	26	22	9	7	3
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All other, male											
1998	1,518	298	135	68	36	19	9	14	2	7	1
1997	1,535	289	116	59	45	19	14	7	5	7	5
1996	1,432	274	109	57	31	17	12	6	10	4	2
1995	1,517	238	115	59	25	23	10	4	4	3	1
1994	1,556	217	101	58	32	25	14	10	9	7	3
1993	1,464	163	103	32	26	22	9	8	2	8	2
1992	1,189	110	76	31	22	19	11	12	3	9	2
1991	1,136	69	62	34	22	20	12	6	8	8	1
1990	897	59	40	31	25	20	13	10	3	9	-
1989	1,069	84	43	41	34	20	14	11	7	14	1
1988	1,036	80	39	41	29	25	12	13	8	7	3
1987	832	45	34	30	27	22	15	7	7	4	1
1986	769	44	31	33	25	23	12	11	9	6	1
1985	664	47	30	26	25	12	13	10	8	3	2
1984	648	35	31	29	13	11	13	16	3	4	1
1983	512	29	21	34	24	16	11	12	4	6	1
1982	591	33	42	29	22	14	15	10	2	5	-
1981	547	44	42	23	19	11	12	7	4	8	1
1980	492	34	30	22	17	18	6	11	9	9	-
1979	562	39	39	24	18	12	19	10	4	2	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
All other, female											
1998	599	102	44	23	9	8	7	7	12	7	1
1997	575	87	38	25	13	10	8	8	10	7	-
1996	589	71	32	26	13	10	6	14	9	15	1
1995	541	64	31	12	11	9	5	10	11	8	-
1994	586	59	35	13	10	12	9	11	8	12	-
1993	511	39	18	10	10	8	8	6	9	16	-
1992	422	26	14	10	6	12	5	7	7	11	-
1991	474	42	21	18	8	17	12	15	7	11	-
1990	366	23	11	11	12	7	8	4	9	8	-
1989	413	18	21	14	10	9	9	9	9	11	1
1988	429	23	13	13	16	4	10	12	6	15	-
1987	359	20	16	16	15	10	9	6	8	13	-
1986	313	16	12	11	10	5	17	5	10	8	1
1985	319	19	19	17	12	14	11	8	9	10	-
1984	290	15	19	10	8	6	12	6	14	2	-
1983	314	16	14	14	8	16	9	6	10	8	2
1982	284	15	18	11	12	10	14	6	6	4	-
1981	298	17	22	9	9	10	9	6	11	10	-
1980	253	10	10	10	13	8	13	11	4	5	-
1979	286	15	12	12	8	9	7	12	5	5	-
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
Black, both sexes											
1998	1,901	373	166	89	41	21	13	19	13	13	2
1997	1,900	351	144	76	55	27	18	12	12	12	4
1996	1,839	322	128	75	39	24	16	18	15	17	2
1995	1,885	283	138	70	32	29	14	11	13	10	1
1994	1,971	261	125	67	36	35	21	18	15	17	2
1993	1,856	194	116	39	33	26	15	14	10	23	2
1992	1,493	128	85	37	26	29	14	19	9	17	2
1991	1,489	103	73	47	28	33	23	19	14	15	1
1990	1,158	78	44	40	32	24	18	11	11	16	-
1989	1,391	92	56	53	42	24	23	18	16	24	2
1988	1,367	100	50	48	40	27	20	23	12	20	3
1987	1,087	63	47	39	38	25	21	12	12	16	1
1986	995	56	42	42	33	25	27	16	19	13	2
1985	907	62	46	40	36	24	22	15	16	12	1
1984	868	47	49	35	18	17	22	21	16	6	1
1983	761	41	33	45	31	31	17	18	13	11	3
1982	797	46	47	34	32	20	29	15	7	9	-
1981	797	57	60	32	27	21	21	13	15	17	1
1980	697	40	38	31	29	25	18	21	13	14	-
1979	780	51	48	33	24	20	26	21	9	5	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
Black, male											
1998	1,363	278	125	68	33	16	8	12	2	7	1
1997	1,387	271	109	55	43	18	11	6	5	5	4
1996	1,301	258	98	53	29	17	11	5	6	2	1
1995	1,381	224	109	58	25	20	9	2	3	2	1
1994	1,437	204	95	56	28	24	12	8	8	5	2
1993	1,370	157	99	29	24	19	8	8	2	8	2
1992	1,110	104	72	28	22	17	9	12	3	7	2
1991	1,057	65	56	31	22	17	12	5	7	6	1
1990	823	57	35	30	22	17	10	9	3	8	-
1989	1,009	77	40	40	33	16	14	10	7	14	1
1988	967	77	37	37	26	24	10	12	7	6	3
1987	760	43	33	26	23	18	12	6	7	4	1
1986	707	41	30	31	24	21	11	11	9	5	1
1985	615	44	28	24	25	10	12	8	7	2	1
1984	599	34	30	26	12	11	11	15	3	4	1
1983	471	27	19	31	23	16	9	12	4	5	1
1982	540	32	36	24	20	13	15	9	2	5	-
1981	516	43	39	23	18	11	12	7	4	8	1
1980	458	31	30	21	16	17	6	10	9	9	-
1979	516	38	36	22	17	12	19	9	4	1	3
Accidental poisoning and exposure to noxious substances (E850-E869,E924.1)											
Black, female											
1998	538	95	41	21	8	5	5	7	11	6	1
1997	513	80	35	21	12	9	7	6	7	7	-
1996	538	64	30	22	10	7	5	13	9	15	1
1995	504	59	29	12	7	9	5	9	10	8	-
1994	534	57	30	11	8	11	9	10	7	12	-
1993	486	37	17	10	9	7	7	6	8	15	-
1992	383	24	13	9	4	12	5	7	6	10	-
1991	432	38	17	16	6	16	11	14	7	9	-
1990	335	21	9	10	10	7	8	2	8	8	-
1989	382	15	16	13	9	8	9	8	9	10	1
1988	400	23	13	11	14	3	10	11	5	14	-
1987	327	20	14	13	15	7	9	6	5	12	-
1986	288	15	12	11	9	4	16	5	10	8	1
1985	292	18	18	16	11	14	10	7	9	10	-
1984	269	13	19	9	6	6	11	6	13	2	-
1983	290	14	14	14	8	15	8	6	9	6	2
1982	257	14	11	10	12	7	14	6	5	4	-
1981	281	14	21	9	9	10	9	6	11	9	-
1980	239	9	8	10	13	8	12	11	4	5	-
1979	264	13	12	11	7	8	7	12	5	4	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
All races, both sexes											
1998	13,078	697	608	592	591	708	986	1,196	1,372	2,323	15
1997	12,412	668	593	566	572	653	896	1,104	1,207	2,096	14
1996	12,229	634	540	508	573	726	970	1,004	1,249	2,060	17
1995	12,349	591	608	560	614	662	889	1,044	1,165	2,073	14
1994	11,807	634	545	529	553	673	829	996	1,115	1,841	15
1993	11,885	577	544	513	628	709	870	946	1,116	1,831	12
1992	11,149	500	456	510	612	702	775	953	988	1,580	15
1991	11,448	572	498	525	651	722	740	910	987	1,581	11
1990	11,533	528	535	567	703	739	754	935	966	1,442	7
1989	12,405	575	542	648	703	785	820	1,002	1,012	1,564	14
1988	12,949	585	584	641	790	871	904	991	1,092	1,728	25
1987	12,786	566	558	637	793	814	874	1,006	1,047	1,641	16
1986	12,814	548	525	694	775	817	916	1,028	1,003	1,592	25
1985	12,986	566	626	752	830	864	918	1,053	946	1,408	25
1984	12,680	584	624	712	766	821	852	943	931	1,319	19
1983	13,136	588	681	824	883	869	929	975	893	1,203	18
1982	12,760	603	724	827	767	820	827	811	769	1,057	14
1981	12,999	617	763	841	868	858	834	764	724	990	23
1980	14,662	710	879	949	989	1,056	1,015	1,043	919	1,103	18
1979	13,376	680	810	909	807	878	852	758	680	832	14
Other and unspecified nontransport accidents and their sequelae (Residual)											
All races, male											
1998	8,482	572	474	453	426	473	636	691	723	899	13
1997	8,220	550	475	458	430	465	556	647	619	839	10
1996	8,154	513	425	388	433	493	616	611	689	805	15
1995	8,311	492	490	431	448	476	577	617	603	867	13
1994	8,058	515	451	427	418	473	560	590	584	749	14
1993	8,068	471	453	394	440	481	591	562	604	741	9
1992	7,697	412	377	410	462	510	497	552	546	617	10
1991	7,909	466	400	405	493	515	489	550	534	628	9
1990	8,010	430	447	452	514	521	485	568	477	548	6
1989	8,684	489	450	515	538	563	535	575	510	619	13
1988	9,021	488	460	527	604	605	567	557	554	724	24
1987	9,000	467	466	503	603	543	562	595	553	693	13
1986	9,005	459	447	540	581	578	602	600	458	672	24
1985	9,301	466	503	604	632	595	550	640	490	640	23
1984	9,172	474	517	562	572	588	572	584	511	556	19
1983	9,381	483	545	649	649	609	588	555	430	517	15
1982	9,357	504	576	656	594	592	544	490	368	454	11
1981	9,675	511	613	663	675	610	563	469	373	441	21
1980	10,547	572	707	740	728	729	617	564	457	463	15
1979	10,067	565	653	717	620	634	557	461	353	375	11

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
All races, female											
1998	4,596	125	134	139	165	235	350	505	649	1,424	2
1997	4,192	118	118	108	142	188	340	457	588	1,257	4
1996	4,075	121	115	120	140	233	354	393	560	1,255	2
1995	4,038	99	118	129	166	186	312	427	562	1,206	1
1994	3,749	119	94	102	135	200	269	406	531	1,092	1
1993	3,817	106	91	119	188	228	279	384	512	1,090	3
1992	3,452	88	79	100	150	192	278	401	442	963	5
1991	3,539	106	98	120	158	207	251	360	453	953	2
1990	3,523	98	88	115	189	218	269	367	489	894	1
1989	3,721	86	92	133	165	222	285	427	502	945	1
1988	3,928	97	124	114	186	266	337	434	538	1,004	1
1987	3,786	99	92	134	190	271	312	411	494	948	3
1986	3,809	89	78	154	194	239	314	428	545	920	1
1985	3,685	100	123	148	198	269	368	413	456	768	2
1984	3,508	110	107	150	194	233	280	359	420	763	-
1983	3,755	105	136	175	234	260	341	420	463	686	3
1982	3,403	99	148	171	173	228	283	321	401	603	3
1981	3,324	106	150	178	193	248	271	295	351	549	2
1980	4,115	138	172	209	261	327	398	479	462	640	3
1979	3,309	115	157	192	187	244	295	297	327	457	3
Other and unspecified nontransport accidents and their sequelae (Residual)											
White, both sexes											
1998	11,141	568	502	496	487	591	871	1,061	1,221	2,076	13
1997	10,430	548	482	469	456	536	770	954	1,058	1,900	12
1996	10,288	505	449	412	465	601	823	863	1,103	1,877	14
1995	10,358	471	487	448	492	549	751	904	1,030	1,840	12
1994	9,885	516	445	412	452	574	710	858	997	1,631	12
1993	9,948	465	436	413	501	601	746	829	984	1,627	9
1992	9,322	395	361	405	515	585	668	819	861	1,434	11
1991	9,443	476	398	420	530	598	621	768	853	1,391	7
1990	9,549	409	449	473	565	608	639	799	855	1,283	4
1989	10,056	446	416	506	538	633	685	852	867	1,350	7
1988	10,633	452	448	512	624	719	734	843	957	1,547	21
1987	10,597	450	442	513	656	665	740	870	912	1,449	10
1986	10,606	418	426	547	647	678	761	867	861	1,412	21
1985	10,666	420	483	605	677	714	750	872	795	1,230	18
1984	10,629	469	503	574	645	684	711	820	794	1,172	16
1983	10,775	446	542	649	707	695	764	802	754	1,063	13
1982	10,671	473	571	660	628	699	688	700	676	954	10
1981	10,924	482	629	696	733	708	714	666	633	875	18
1980	11,728	534	675	721	778	845	788	804	753	939	12
1979	11,171	544	647	762	680	735	733	647	605	732	9

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
White, male											
1998	7,209	461	395	383	358	394	563	619	641	791	12
1997	6,928	461	392	390	343	389	480	559	547	761	10
1996	6,875	417	360	325	352	410	531	521	612	724	12
1995	6,997	394	406	344	359	398	496	535	533	758	11
1994	6,742	428	369	336	347	408	478	513	521	655	11
1993	6,759	381	366	319	359	412	501	488	539	658	7
1992	6,463	326	308	332	393	430	427	477	469	555	7
1991	6,502	385	323	320	399	424	418	464	453	543	7
1990	6,624	331	375	378	413	431	410	484	432	479	4
1989	7,027	381	350	408	413	450	448	479	428	528	7
1988	7,360	381	349	417	468	501	464	470	483	640	21
1987	7,427	374	366	410	499	449	483	511	476	596	8
1986	7,465	344	365	423	497	482	516	500	390	591	20
1985	7,642	344	396	482	516	494	452	531	402	547	16
1984	7,718	385	424	453	482	488	476	506	430	486	16
1983	7,731	373	441	522	525	490	487	459	353	445	11
1982	7,843	393	464	528	487	510	444	424	314	408	8
1981	8,165	404	499	546	573	511	484	411	324	387	16
1980	8,622	435	553	571	588	596	502	457	385	400	12
1979	8,420	456	526	600	518	533	478	384	310	319	7
Other and unspecified nontransport accidents and their sequelae (Residual)											
White, female											
1998	3,932	107	107	113	129	197	308	442	580	1,285	1
1997	3,502	87	90	79	113	147	290	395	511	1,139	2
1996	3,413	88	89	87	113	191	292	342	491	1,153	2
1995	3,361	77	81	104	133	151	255	369	497	1,082	1
1994	3,143	88	76	76	105	166	232	345	476	976	1
1993	3,189	84	70	94	142	189	245	341	445	969	2
1992	2,859	69	53	73	122	155	241	342	392	879	4
1991	2,941	91	75	100	131	174	203	304	400	848	-
1990	2,925	78	74	95	152	177	229	315	423	804	-
1989	3,029	65	66	98	125	183	237	373	439	822	-
1988	3,273	71	99	95	156	218	270	373	474	907	-
1987	3,170	76	76	103	157	216	257	359	436	853	2
1986	3,141	74	61	124	150	196	245	367	471	821	1
1985	3,024	76	87	123	161	220	298	341	393	683	2
1984	2,911	84	79	121	163	196	235	314	364	686	-
1983	3,044	73	101	127	182	205	277	343	401	618	2
1982	2,828	80	107	132	141	189	244	276	362	546	2
1981	2,759	78	130	150	160	197	230	255	309	488	2
1980	3,106	99	122	150	190	249	286	347	368	539	-
1979	2,751	88	121	162	162	202	255	263	295	413	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
All other, both sexes											
1998	1,937	129	106	96	104	117	115	135	151	247	2
1997	1,982	120	111	97	116	117	126	150	149	196	2
1996	1,941	129	91	96	108	125	147	141	146	183	3
1995	1,991	120	121	112	122	113	138	140	135	233	2
1994	1,922	118	100	117	101	99	119	138	118	210	3
1993	1,937	112	108	100	127	108	124	117	132	204	3
1992	1,827	105	95	105	97	117	107	134	127	146	4
1991	2,005	96	100	105	121	124	119	142	134	190	4
1990	1,984	119	86	94	138	131	115	136	111	159	3
1989	2,349	129	126	142	165	152	135	150	145	214	7
1988	2,316	133	136	129	166	152	170	148	135	181	4
1987	2,189	116	116	124	137	149	134	136	135	192	6
1986	2,208	130	99	147	128	139	155	161	142	180	4
1985	2,320	146	143	147	153	150	168	181	151	178	7
1984	2,051	115	121	138	121	137	141	123	137	147	3
1983	2,361	142	139	175	176	174	165	173	139	140	5
1982	2,089	130	153	167	139	121	139	111	93	103	4
1981	2,075	135	134	145	135	150	120	98	91	115	5
1980	2,934	176	204	228	211	211	227	239	166	164	6
1979	2,205	136	163	147	127	143	119	111	75	100	5
Other and unspecified nontransport accidents and their sequelae (Residual)											
All other, male											
1998	1,273	111	79	70	68	79	73	72	82	108	1
1997	1,292	89	83	68	87	76	76	88	72	78	-
1996	1,279	96	65	63	81	83	85	90	77	81	3
1995	1,314	98	84	87	89	78	81	82	70	109	2
1994	1,316	87	82	91	71	65	82	77	63	94	3
1993	1,309	90	87	75	81	69	90	74	65	83	2
1992	1,234	86	69	78	69	80	70	75	77	62	3
1991	1,407	81	77	85	94	91	71	86	81	85	2
1990	1,386	99	72	74	101	90	75	84	45	69	2
1989	1,657	108	100	107	125	113	87	96	82	91	6
1988	1,661	107	111	110	136	104	103	87	71	84	3
1987	1,573	93	100	93	104	94	79	84	77	97	5
1986	1,540	115	82	117	84	96	86	100	68	81	4
1985	1,659	122	107	122	116	101	98	109	88	93	7
1984	1,454	89	93	109	90	100	96	78	81	70	3
1983	1,650	110	104	127	124	119	101	96	77	72	4
1982	1,514	111	112	128	107	82	100	66	54	46	3
1981	1,510	107	114	117	102	99	79	58	49	54	5
1980	1,925	137	154	169	140	133	115	107	72	63	3
1979	1,647	109	127	117	102	101	79	77	43	56	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
All other, female											
1998	664	18	27	26	36	38	42	63	69	139	1
1997	690	31	28	29	29	41	50	62	77	118	2
1996	662	33	26	33	27	42	62	51	69	102	-
1995	677	22	37	25	33	35	57	58	65	124	-
1994	606	31	18	26	30	34	37	61	55	116	-
1993	628	22	21	25	46	39	34	43	67	121	1
1992	593	19	26	27	28	37	37	59	50	84	1
1991	598	15	23	20	27	33	48	56	53	105	2
1990	598	20	14	20	37	41	40	52	66	90	1
1989	692	21	26	35	40	39	48	54	63	123	1
1988	655	26	25	19	30	48	67	61	64	97	1
1987	616	23	16	31	33	55	55	52	58	95	1
1986	668	15	17	30	44	43	69	61	74	99	-
1985	661	24	36	25	37	49	70	72	63	85	-
1984	597	26	28	29	31	37	45	45	56	77	-
1983	711	32	35	48	52	55	64	77	62	68	1
1982	575	19	41	39	32	39	39	45	39	57	1
1981	565	28	20	28	33	51	41	40	42	61	-
1980	1,009	39	50	59	71	78	112	132	94	101	3
1979	558	27	36	30	25	42	40	34	32	44	1
Other and unspecified nontransport accidents and their sequelae (Residual)											
Black, both sexes											
1998	1,673	112	91	86	88	100	96	123	129	224	2
1997	1,674	102	97	81	99	102	106	134	130	169	2
1996	1,630	110	84	81	89	112	126	116	128	156	1
1995	1,737	105	107	93	110	100	128	134	127	207	2
1994	1,624	96	85	95	87	89	106	119	101	187	3
1993	1,666	93	93	90	110	98	107	105	113	184	2
1992	1,515	85	80	90	86	101	100	119	114	130	4
1991	1,713	81	87	92	105	104	102	129	122	168	3
1990	1,685	102	78	81	118	109	101	121	96	147	3
1989	2,059	114	112	126	150	140	124	141	129	193	5
1988	2,014	120	121	108	141	134	151	140	110	161	4
1987	1,889	101	108	104	124	136	113	122	119	163	3
1986	1,940	112	91	131	114	129	141	144	130	167	3
1985	2,039	124	131	135	143	137	153	164	137	160	7
1984	1,792	99	103	127	106	122	127	112	122	134	3
1983	2,097	130	123	154	161	163	150	157	127	128	3
1982	1,795	118	129	153	118	111	127	103	81	85	4
1981	1,842	122	120	131	123	136	117	95	83	99	5
1980	2,666	165	183	214	194	200	213	228	161	152	5
1979	1,944	119	145	136	110	131	103	98	65	88	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified nontransport accidents and their sequelae (Residual)											
Black, male											
1998	1,085	98	69	64	56	68	61	63	70	92	1
1997	1,078	74	71	55	73	66	60	82	60	65	-
1996	1,044	84	61	54	68	77	72	68	63	64	1
1995	1,132	87	76	72	80	68	73	76	68	93	2
1994	1,111	70	70	77	63	59	74	67	53	82	3
1993	1,116	75	80	69	69	64	76	64	51	76	1
1992	996	66	57	68	61	71	64	69	67	50	3
1991	1,191	68	65	76	80	75	59	79	71	70	2
1990	1,167	82	66	63	89	74	68	72	36	64	2
1989	1,436	94	87	92	112	103	80	91	69	79	4
1988	1,424	95	98	92	115	89	88	81	55	73	3
1987	1,345	83	95	75	94	85	70	71	68	83	3
1986	1,343	97	77	102	73	90	77	91	60	74	3
1985	1,463	108	98	112	107	93	91	94	80	81	7
1984	1,257	75	79	101	78	90	87	71	71	60	3
1983	1,451	102	90	115	113	110	91	84	67	65	2
1982	1,292	100	94	119	90	77	92	63	43	37	3
1981	1,330	98	100	104	93	89	76	56	43	44	5
1980	1,713	127	141	156	126	124	103	97	70	56	2
1979	1,446	96	114	107	87	95	68	68	35	48	4
Other and unspecified nontransport accidents and their sequelae (Residual)											
Black, female											
1998	588	14	22	22	32	32	35	60	59	132	1
1997	596	28	26	26	26	36	46	52	70	104	2
1996	586	26	23	27	21	35	54	48	65	92	-
1995	605	18	31	21	30	32	55	58	59	114	-
1994	513	26	15	18	24	30	32	52	48	105	-
1993	550	18	13	21	41	34	31	41	62	108	1
1992	519	19	23	22	25	30	36	50	47	80	1
1991	522	13	22	16	25	29	43	50	51	98	1
1990	518	20	12	18	29	35	33	49	60	83	1
1989	623	20	25	34	38	37	44	50	60	114	1
1988	590	25	23	16	26	45	63	59	55	88	1
1987	544	18	13	29	30	51	43	51	51	80	-
1986	597	15	14	29	41	39	64	53	70	93	-
1985	576	16	33	23	36	44	62	70	57	79	-
1984	535	24	24	26	28	32	40	41	51	74	-
1983	646	28	33	39	48	53	59	73	60	63	1
1982	503	18	35	34	28	34	35	40	38	48	1
1981	512	24	20	27	30	47	41	39	40	55	-
1980	953	38	42	58	68	76	110	131	91	96	3
1979	498	23	31	29	23	36	35	30	30	40	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
All races, both sexes											
1998	30,575	2,881	2,250	1,707	1,256	1,259	1,338	1,273	1,082	851	17
1997	30,535	2,801	2,147	1,702	1,244	1,291	1,372	1,303	957	805	18
1996	30,903	2,785	2,052	1,615	1,310	1,419	1,387	1,280	1,010	759	24
1995	31,284	2,561	1,971	1,435	1,369	1,436	1,524	1,307	1,004	785	12
1994	31,142	2,407	1,889	1,462	1,350	1,384	1,481	1,351	981	811	19
1993	31,102	2,269	1,899	1,561	1,500	1,499	1,541	1,347	1,043	778	18
1992	30,484	2,251	1,767	1,541	1,564	1,555	1,483	1,387	1,021	714	13
1991	30,810	2,206	1,777	1,613	1,628	1,572	1,512	1,395	1,031	758	20
1990	30,906	2,059	1,659	1,692	1,691	1,666	1,564	1,461	1,032	671	11
1989	30,232	1,954	1,688	1,650	1,692	1,742	1,533	1,349	910	694	14
1988	30,407	1,898	1,634	1,720	1,686	1,718	1,578	1,447	1,015	605	19
1987	30,796	1,931	1,776	1,838	1,812	1,757	1,671	1,469	933	634	13
1986	30,904	1,895	1,841	1,916	1,866	1,707	1,707	1,433	850	578	12
1985	29,453	1,804	1,728	1,931	1,794	1,581	1,564	1,282	844	517	7
1984	29,286	1,862	1,786	2,041	1,809	1,599	1,542	1,147	747	482	19
1983	28,295	1,729	1,900	1,977	1,696	1,509	1,415	1,123	748	489	15
1982	28,242	1,797	1,913	1,993	1,748	1,490	1,322	1,056	622	429	16
1981	27,596	1,752	1,877	1,989	1,600	1,389	1,185	910	577	417	22
1980	26,869	1,701	1,922	1,893	1,563	1,422	1,208	941	536	430	17
1979	27,206	1,793	1,996	1,888	1,680	1,532	1,198	986	591	394	17
Intentional self-harm (suicide) (E950-E959)											
All races, male											
1998	24,538	2,213	1,683	1,318	987	1,030	1,131	1,075	925	686	17
1997	24,492	2,116	1,581	1,321	1,010	1,064	1,119	1,089	806	671	16
1996	24,998	2,141	1,543	1,286	1,020	1,141	1,166	1,102	843	641	18
1995	25,369	1,952	1,513	1,127	1,087	1,142	1,251	1,103	835	642	11
1994	25,174	1,825	1,404	1,128	1,070	1,091	1,209	1,148	828	653	17
1993	25,007	1,694	1,440	1,205	1,167	1,173	1,252	1,119	884	646	15
1992	24,457	1,689	1,310	1,183	1,203	1,227	1,205	1,152	850	571	9
1991	24,769	1,614	1,365	1,239	1,278	1,218	1,248	1,181	878	614	17
1990	24,724	1,583	1,253	1,254	1,309	1,302	1,244	1,237	862	554	8
1989	24,102	1,455	1,256	1,222	1,288	1,383	1,292	1,132	763	567	12
1988	24,078	1,341	1,212	1,294	1,282	1,361	1,261	1,199	851	498	17
1987	24,272	1,392	1,300	1,383	1,375	1,361	1,363	1,215	778	539	11
1986	24,226	1,380	1,328	1,417	1,371	1,344	1,368	1,161	695	484	11
1985	23,145	1,305	1,267	1,445	1,369	1,236	1,252	1,035	714	428	7
1984	22,689	1,313	1,269	1,489	1,355	1,212	1,239	939	614	390	17
1983	21,786	1,224	1,366	1,426	1,258	1,139	1,108	913	622	397	14
1982	21,625	1,257	1,357	1,416	1,292	1,129	1,053	859	518	363	14
1981	20,809	1,184	1,271	1,374	1,191	1,033	925	742	484	354	18
1980	20,505	1,180	1,340	1,367	1,118	1,098	955	766	446	345	16
1979	20,256	1,195	1,332	1,312	1,193	1,149	939	784	481	322	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
All races, female											
1998	6,037	668	567	389	269	229	207	198	157	165	-
1997	6,043	685	566	381	234	227	253	214	151	134	2
1996	5,905	644	509	329	290	278	221	178	167	118	6
1995	5,915	609	458	308	282	294	273	204	169	143	1
1994	5,968	582	485	334	280	293	272	203	153	158	2
1993	6,095	575	459	356	333	326	289	228	159	132	3
1992	6,027	562	457	358	361	328	278	235	171	143	4
1991	6,041	592	412	374	350	354	264	214	153	144	3
1990	6,182	476	406	438	382	364	320	224	170	117	3
1989	6,130	499	432	428	404	359	241	217	147	127	2
1988	6,329	557	422	426	404	357	317	248	164	107	2
1987	6,524	539	476	455	437	396	308	254	155	95	2
1986	6,678	515	513	499	495	363	339	272	155	94	1
1985	6,308	499	461	486	425	345	312	247	130	89	-
1984	6,597	549	517	552	454	387	303	208	133	92	2
1983	6,509	505	534	551	438	370	307	210	126	92	1
1982	6,617	540	556	577	456	361	269	197	104	66	2
1981	6,787	568	606	615	409	356	260	168	93	63	4
1980	6,364	521	582	526	445	324	253	175	90	85	1
1979	6,950	598	664	576	487	383	259	202	110	72	2
Intentional self-harm (suicide) (E950-E959)											
White, both sexes											
1998	27,648	2,655	2,098	1,608	1,185	1,180	1,276	1,224	1,038	824	15
1997	27,513	2,597	2,018	1,582	1,157	1,202	1,297	1,252	917	771	13
1996	27,856	2,586	1,923	1,515	1,222	1,340	1,311	1,228	977	737	19
1995	28,187	2,378	1,847	1,343	1,279	1,347	1,437	1,254	976	761	11
1994	27,976	2,235	1,795	1,363	1,268	1,299	1,406	1,304	949	783	13
1993	28,035	2,107	1,762	1,465	1,421	1,427	1,468	1,299	1,008	752	16
1992	27,611	2,085	1,649	1,476	1,477	1,485	1,426	1,339	980	691	11
1991	27,996	2,033	1,662	1,508	1,527	1,496	1,443	1,335	995	734	14
1990	28,086	1,909	1,540	1,595	1,606	1,590	1,493	1,421	1,008	642	11
1989	27,424	1,830	1,588	1,573	1,594	1,642	1,473	1,308	880	670	12
1988	27,790	1,761	1,524	1,634	1,612	1,638	1,525	1,401	987	588	17
1987	28,217	1,798	1,666	1,764	1,723	1,662	1,593	1,421	898	610	11
1986	28,437	1,768	1,736	1,821	1,775	1,615	1,633	1,390	824	559	8
1985	27,087	1,673	1,631	1,838	1,708	1,499	1,496	1,242	814	503	5
1984	27,002	1,734	1,687	1,931	1,719	1,531	1,476	1,110	730	467	13
1983	26,157	1,605	1,811	1,878	1,629	1,453	1,359	1,084	719	473	12
1982	26,141	1,699	1,811	1,901	1,685	1,427	1,278	1,028	598	416	14
1981	25,452	1,632	1,795	1,882	1,531	1,333	1,142	872	558	401	19
1980	24,829	1,596	1,820	1,800	1,502	1,369	1,170	916	520	411	16
1979	24,945	1,673	1,896	1,795	1,601	1,465	1,148	953	570	381	15

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
White, male											
1998	22,174	2,038	1,575	1,241	932	962	1,080	1,040	890	668	15
1997	22,042	1,968	1,487	1,224	942	993	1,064	1,048	775	648	12
1996	22,547	1,991	1,449	1,210	956	1,084	1,111	1,061	819	627	13
1995	22,853	1,814	1,414	1,056	1,017	1,080	1,182	1,061	812	627	10
1994	22,581	1,698	1,335	1,058	1,008	1,027	1,148	1,111	801	632	12
1993	22,524	1,567	1,333	1,131	1,111	1,117	1,202	1,085	856	629	14
1992	22,126	1,569	1,222	1,133	1,139	1,177	1,158	1,113	815	555	9
1991	22,452	1,477	1,282	1,152	1,200	1,156	1,196	1,132	849	597	12
1990	22,448	1,468	1,162	1,186	1,242	1,247	1,188	1,203	843	534	8
1989	21,858	1,361	1,188	1,167	1,217	1,304	1,239	1,100	744	547	10
1988	21,980	1,244	1,130	1,228	1,222	1,298	1,221	1,157	834	486	16
1987	22,188	1,287	1,220	1,324	1,309	1,283	1,301	1,175	746	520	9
1986	22,270	1,283	1,254	1,354	1,309	1,269	1,313	1,129	674	468	7
1985	21,256	1,206	1,196	1,368	1,306	1,171	1,204	1,011	690	419	5
1984	20,882	1,225	1,200	1,407	1,288	1,162	1,186	905	600	380	11
1983	20,097	1,129	1,304	1,353	1,206	1,090	1,063	886	597	384	11
1982	19,965	1,187	1,279	1,348	1,245	1,078	1,018	837	499	354	12
1981	19,166	1,091	1,215	1,297	1,134	989	898	710	466	344	16
1980	18,901	1,103	1,267	1,293	1,071	1,055	925	749	433	328	15
1979	18,504	1,106	1,256	1,246	1,140	1,102	898	763	468	310	13
Intentional self-harm (suicide) (E950-E959)											
White, female											
1998	5,474	617	523	367	253	218	196	184	148	156	-
1997	5,471	629	531	358	215	209	233	204	142	123	1
1996	5,309	595	474	305	266	256	200	167	158	110	6
1995	5,334	564	433	287	262	267	255	193	164	134	1
1994	5,395	537	460	305	260	272	258	193	148	151	1
1993	5,511	540	429	334	310	310	266	214	152	123	2
1992	5,485	516	427	343	338	308	268	226	165	136	2
1991	5,544	556	380	356	327	340	247	203	146	137	2
1990	5,638	441	378	409	364	343	305	218	165	108	3
1989	5,566	469	400	406	377	338	234	208	136	123	2
1988	5,810	517	394	406	390	340	304	244	153	102	1
1987	6,029	511	446	440	414	379	292	246	152	90	2
1986	6,167	485	482	467	466	346	320	261	150	91	1
1985	5,831	467	435	470	402	328	292	231	124	84	-
1984	6,120	509	487	524	431	369	290	205	130	87	2
1983	6,060	476	507	525	423	363	296	198	122	89	1
1982	6,176	512	532	553	440	349	260	191	99	62	2
1981	6,286	541	580	585	397	344	244	162	92	57	3
1980	5,928	493	553	507	431	314	245	167	87	83	1
1979	6,441	567	640	549	461	363	250	190	102	71	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
All other, both sexes											
1998	2,927	226	152	99	71	79	62	49	44	27	2
1997	3,022	204	129	120	87	89	75	51	40	34	5
1996	3,047	199	129	100	88	79	76	52	33	22	5
1995	3,097	183	124	92	90	89	87	53	28	24	1
1994	3,166	172	94	99	82	85	75	47	32	28	6
1993	3,067	162	137	96	79	72	73	48	35	26	2
1992	2,873	166	118	65	87	70	57	48	41	23	2
1991	2,814	173	115	105	101	76	69	60	36	24	6
1990	2,820	150	119	97	85	76	71	40	24	29	-
1989	2,808	124	100	77	98	100	60	41	30	24	2
1988	2,617	137	110	86	74	80	53	46	28	17	2
1987	2,579	133	110	74	89	95	78	48	35	24	2
1986	2,467	127	105	95	91	92	74	43	26	19	4
1985	2,366	131	97	93	86	82	68	40	30	14	2
1984	2,284	128	99	110	90	68	66	37	17	15	6
1983	2,138	124	89	99	67	56	56	39	29	16	3
1982	2,101	98	102	92	63	63	44	28	24	13	2
1981	2,144	120	82	107	69	56	43	38	19	16	3
1980	2,040	105	102	93	61	53	38	25	16	19	1
1979	2,261	120	100	93	79	67	50	33	21	13	2
Intentional self-harm (suicide) (E950-E959)											
All other, male											
1998	2,364	175	108	77	55	68	51	35	35	18	2
1997	2,450	148	94	97	68	71	55	41	31	23	4
1996	2,451	150	94	76	64	57	55	41	24	14	5
1995	2,516	138	99	71	70	62	69	42	23	15	1
1994	2,593	127	69	70	62	64	61	37	27	21	5
1993	2,483	127	107	74	56	56	50	34	28	17	1
1992	2,331	120	88	50	64	50	47	39	35	16	-
1991	2,317	137	83	87	78	62	52	49	29	17	5
1990	2,276	115	91	68	67	55	56	34	19	20	-
1989	2,244	94	68	55	71	79	53	32	19	20	2
1988	2,098	97	82	66	60	63	40	42	17	12	1
1987	2,084	105	80	59	66	78	62	40	32	19	2
1986	1,956	97	74	63	62	75	55	32	21	16	4
1985	1,889	99	71	77	63	65	48	24	24	9	2
1984	1,807	88	69	82	67	50	53	34	14	10	6
1983	1,689	95	62	73	52	49	45	27	25	13	3
1982	1,660	70	78	68	47	51	35	22	19	9	2
1981	1,643	93	56	77	57	44	27	32	18	10	2
1980	1,604	77	73	74	47	43	30	17	13	17	1
1979	1,752	89	76	66	53	47	41	21	13	12	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
All other, female											
1998	563	51	44	22	16	11	11	14	9	9	-
1997	572	56	35	23	19	18	20	10	9	11	1
1996	596	49	35	24	24	22	21	11	9	8	-
1995	581	45	25	21	20	27	18	11	5	9	-
1994	573	45	25	29	20	21	14	10	5	7	1
1993	584	35	30	22	23	16	23	14	7	9	1
1992	542	46	30	15	23	20	10	9	6	7	2
1991	497	36	32	18	23	14	17	11	7	7	1
1990	544	35	28	29	18	21	15	6	5	9	-
1989	564	30	32	22	27	21	7	9	11	4	-
1988	519	40	28	20	14	17	13	4	11	5	1
1987	495	28	30	15	23	17	16	8	3	5	-
1986	511	30	31	32	29	17	19	11	5	3	-
1985	477	32	26	16	23	17	20	16	6	5	-
1984	477	40	30	28	23	18	13	3	3	5	-
1983	449	29	27	26	15	7	11	12	4	3	-
1982	441	28	24	24	16	12	9	6	5	4	-
1981	501	27	26	30	12	12	16	6	1	6	1
1980	436	28	29	19	14	10	8	8	3	2	-
1979	509	31	24	27	26	20	9	12	8	1	-
Intentional self-harm (suicide) (E950-E959)											
Black, both sexes											
1998	1,977	163	99	61	49	57	32	34	16	11	2
1997	2,103	138	80	74	57	57	50	31	21	17	5
1996	2,164	138	98	61	67	59	49	30	20	14	3
1995	2,231	132	90	62	59	57	61	37	19	13	1
1994	2,271	127	61	67	48	63	60	32	20	17	2
1993	2,259	121	102	69	48	48	49	36	25	14	1
1992	2,143	120	83	47	66	52	43	33	25	17	2
1991	2,097	132	84	75	64	61	48	42	26	14	5
1990	2,111	128	91	67	58	58	55	27	16	17	-
1989	2,153	94	78	60	70	80	43	31	18	17	2
1988	2,022	106	82	70	62	65	38	38	16	7	2
1987	1,963	97	78	53	67	76	60	41	25	9	1
1986	1,892	101	82	72	70	70	56	33	20	12	3
1985	1,795	103	85	64	69	60	55	34	25	7	1
1984	1,760	93	76	85	74	49	56	29	11	8	4
1983	1,623	93	73	73	52	50	42	25	19	11	1
1982	1,639	71	88	77	51	51	35	17	18	10	2
1981	1,658	94	63	85	55	43	36	33	14	9	3
1980	1,607	80	78	69	55	46	30	18	11	10	1
1979	1,812	90	87	81	67	58	38	20	14	9	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) (E950-E959)											
Black, male											
1998	1,659	130	80	49	39	51	29	26	15	9	2
1997	1,764	104	60	62	45	48	41	27	18	15	4
1996	1,820	113	71	54	54	48	39	24	15	10	3
1995	1,878	107	77	51	48	40	51	32	18	10	1
1994	1,922	98	50	51	39	49	51	27	17	15	2
1993	1,905	101	86	56	31	38	39	26	21	10	-
1992	1,803	89	68	37	53	40	36	30	23	12	-
1991	1,782	109	63	62	53	50	37	39	22	12	4
1990	1,737	100	74	49	46	42	48	25	15	13	-
1989	1,771	74	54	47	54	64	40	27	14	16	2
1988	1,648	74	60	55	48	52	33	34	14	7	1
1987	1,635	83	62	47	52	65	49	34	22	9	1
1986	1,537	79	61	48	46	59	43	25	17	12	3
1985	1,481	82	63	56	52	53	45	19	21	5	1
1984	1,432	64	59	66	58	37	47	28	10	7	4
1983	1,321	74	53	64	42	44	37	20	19	8	1
1982	1,327	52	68	63	41	41	28	15	15	9	2
1981	1,315	78	47	62	46	34	21	29	14	7	2
1980	1,297	61	62	57	43	38	25	14	10	10	1
1979	1,428	71	70	60	48	44	32	15	10	8	2
Intentional self-harm (suicide) (E950-E959)											
Black, female											
1998	318	33	19	12	10	6	3	8	1	2	-
1997	339	34	20	12	12	9	9	4	3	2	1
1996	344	25	27	7	13	11	10	6	5	4	-
1995	353	25	13	11	11	17	10	5	1	3	-
1994	349	29	11	16	9	14	9	5	3	2	-
1993	354	20	16	13	17	10	10	10	4	4	1
1992	340	31	15	10	13	12	7	3	2	5	2
1991	315	23	21	13	11	11	11	3	4	2	1
1990	374	28	17	18	12	16	7	2	1	4	-
1989	382	20	24	13	16	16	3	4	4	1	-
1988	374	32	22	15	14	13	5	4	2	-	1
1987	328	14	16	6	15	11	11	7	3	-	-
1986	355	22	21	24	24	11	13	8	3	-	-
1985	314	21	22	8	17	7	10	15	4	2	-
1984	328	29	17	19	16	12	9	1	1	1	-
1983	302	19	20	9	10	6	5	5	-	3	-
1982	312	19	20	14	10	10	7	2	3	1	-
1981	343	16	16	23	9	9	15	4	-	2	1
1980	310	19	16	12	12	8	5	4	1	-	-
1979	384	19	17	21	19	14	6	5	4	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All races, both sexes											
1998	17,424	1,476	1,176	1,042	831	853	978	954	785	543	4
1997	17,566	1,446	1,201	1,062	797	914	992	931	677	494	7
1996	18,166	1,501	1,206	986	856	1,009	1,025	928	722	500	10
1995	18,503	1,354	1,150	902	908	990	1,095	968	708	497	6
1994	18,765	1,351	1,092	902	895	958	1,052	991	693	497	6
1993	18,940	1,270	1,163	992	1,005	1,020	1,084	969	745	483	4
1992	18,169	1,260	1,029	986	974	1,079	1,087	964	679	424	2
1991	18,526	1,246	1,046	986	1,020	1,083	1,083	1,008	710	445	7
1990	18,885	1,208	981	1,073	1,094	1,161	1,084	1,050	700	401	6
1989	18,178	1,109	1,035	1,023	1,093	1,191	1,093	955	602	406	1
1988	18,169	1,071	1,003	1,058	1,112	1,144	1,097	991	685	347	6
1987	18,136	1,085	1,032	1,131	1,150	1,177	1,148	1,003	612	354	2
1986	18,153	1,068	1,070	1,149	1,174	1,158	1,160	968	543	314	4
1985	17,363	1,022	1,035	1,144	1,130	1,046	1,082	849	545	291	2
1984	17,113	1,031	1,020	1,213	1,123	1,054	1,030	781	462	263	6
1983	16,600	949	1,102	1,160	1,079	992	936	717	445	276	5
1982	16,560	1,021	1,069	1,140	1,071	940	881	682	383	234	6
1981	16,139	993	1,070	1,147	991	850	738	574	340	232	6
1980	15,396	912	1,057	1,058	921	880	750	553	307	228	3
1979	15,543	978	1,082	1,091	971	951	725	586	351	186	5
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All races, male											
1998	15,104	1,216	997	889	719	767	893	871	735	514	4
1997	15,194	1,201	1,004	914	704	808	886	867	632	472	6
1996	15,808	1,276	981	862	738	893	938	868	664	477	9
1995	16,060	1,127	956	774	772	867	996	890	656	465	5
1994	16,287	1,124	894	794	776	841	950	920	649	469	6
1993	16,381	1,040	973	836	877	881	983	892	702	458	4
1992	15,802	1,038	864	840	844	940	967	889	635	398	2
1991	16,120	1,018	891	858	892	923	978	946	667	423	6
1990	16,285	1,002	823	902	944	1,003	973	985	655	388	5
1989	15,680	916	864	845	920	1,056	1,000	883	561	388	-
1988	15,656	858	831	895	951	1,019	982	912	649	322	6
1987	15,539	863	880	945	985	1,022	1,043	919	574	336	2
1986	15,518	883	875	957	996	1,010	1,041	895	512	306	4
1985	14,809	836	849	947	971	914	971	772	515	274	2
1984	14,504	837	827	1,008	961	920	938	720	441	245	6
1983	13,959	756	898	959	901	841	836	665	428	265	5
1982	13,872	803	863	929	907	816	802	637	365	228	6
1981	13,378	768	850	921	845	729	677	531	331	225	5
1980	12,937	725	844	870	760	780	680	522	291	218	2
1979	12,906	753	852	869	813	830	659	546	331	180	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All races, female											
1998	2,320	260	179	153	112	86	85	83	50	29	-
1997	2,372	245	197	148	93	106	106	64	45	22	1
1996	2,358	225	225	124	118	116	87	60	58	23	1
1995	2,443	227	194	128	136	123	99	78	52	32	1
1994	2,478	227	198	108	119	117	102	71	44	28	-
1993	2,559	230	190	156	128	139	101	77	43	25	-
1992	2,367	222	165	146	130	139	120	75	44	26	-
1991	2,406	228	155	128	128	160	105	62	43	22	1
1990	2,600	206	158	171	150	158	111	65	45	13	1
1989	2,498	193	171	178	173	135	93	72	41	18	1
1988	2,513	213	172	163	161	125	115	79	36	25	-
1987	2,597	222	152	186	165	155	105	84	38	18	-
1986	2,635	185	195	192	178	148	119	73	31	8	-
1985	2,554	186	186	197	159	132	111	77	30	17	-
1984	2,609	194	193	205	162	134	92	61	21	18	-
1983	2,641	193	204	201	178	151	100	52	17	11	-
1982	2,688	218	206	211	164	124	79	45	18	6	-
1981	2,761	225	220	226	146	121	61	43	9	7	1
1980	2,459	187	213	188	161	100	70	31	16	10	1
1979	2,637	225	230	222	158	121	66	40	20	6	1
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
White, both sexes											
1998	15,968	1,375	1,120	986	792	812	945	920	761	531	4
1997	16,014	1,359	1,154	1,002	751	855	944	902	657	479	5
1996	16,523	1,410	1,141	938	801	959	978	899	704	487	6
1995	16,860	1,270	1,091	846	863	944	1,046	937	691	484	6
1994	17,007	1,267	1,046	853	857	901	1,004	965	672	481	5
1993	17,271	1,193	1,085	943	960	986	1,042	941	722	476	4
1992	16,606	1,191	968	949	925	1,040	1,048	934	653	408	2
1991	16,974	1,155	986	927	962	1,034	1,043	969	684	433	4
1990	17,344	1,126	919	1,017	1,044	1,113	1,039	1,027	684	387	6
1989	16,671	1,042	984	974	1,038	1,125	1,057	931	586	391	1
1988	16,725	995	943	1,006	1,058	1,091	1,061	958	673	340	6
1987	16,753	1,018	975	1,084	1,102	1,123	1,099	968	590	344	2
1986	16,836	991	1,014	1,098	1,124	1,096	1,109	940	527	304	3
1985	16,120	956	974	1,094	1,075	998	1,043	822	526	286	2
1984	15,928	966	972	1,141	1,073	1,018	988	755	453	258	5
1983	15,508	888	1,049	1,118	1,042	950	900	699	427	268	5
1982	15,489	978	1,018	1,092	1,034	905	849	671	369	227	5
1981	14,975	926	1,032	1,093	946	817	711	548	328	225	6
1980	14,286	864	1,006	1,011	879	845	724	541	297	220	3
1979	14,278	916	1,017	1,031	918	909	689	566	339	176	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
White, male											
1998	13,836	1,138	950	840	685	728	863	844	712	503	4
1997	13,820	1,131	962	860	664	757	847	840	613	458	5
1996	14,383	1,201	928	818	695	848	898	841	650	467	5
1995	14,613	1,056	905	722	738	828	950	863	640	455	5
1994	14,751	1,054	857	749	744	795	907	895	630	454	5
1993	14,919	972	905	794	840	852	947	868	681	451	4
1992	14,408	983	810	806	803	909	931	862	610	385	2
1991	14,734	939	840	806	843	880	943	909	643	412	4
1990	14,949	935	768	858	903	964	931	962	640	376	5
1989	14,378	855	822	806	874	999	967	861	548	374	-
1988	14,406	799	782	853	904	973	950	880	637	315	6
1987	14,333	801	830	903	946	974	997	889	554	326	2
1986	14,404	820	827	921	957	957	999	870	498	296	3
1985	13,750	780	797	903	922	871	937	756	500	270	2
1984	13,499	787	789	948	918	890	901	694	432	240	5
1983	13,018	701	855	922	869	803	804	648	411	258	5
1982	12,958	766	820	887	876	785	774	627	353	222	5
1981	12,409	709	817	876	806	702	660	507	319	218	5
1980	11,991	686	801	830	726	748	655	512	281	210	2
1979	11,859	700	797	825	768	797	628	530	322	171	2
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
White, female											
1998	2,132	237	170	146	107	84	82	76	49	28	-
1997	2,194	228	192	142	87	98	97	62	44	21	-
1996	2,140	209	213	120	106	111	80	58	54	20	1
1995	2,247	214	186	124	125	116	96	74	51	29	1
1994	2,256	213	189	104	113	106	97	70	42	27	-
1993	2,352	221	180	149	120	134	95	73	41	25	-
1992	2,198	208	158	143	122	131	117	72	43	23	-
1991	2,240	216	146	121	119	154	100	60	41	21	-
1990	2,395	191	151	159	141	149	108	65	44	11	1
1989	2,293	187	162	168	164	126	90	70	38	17	1
1988	2,319	196	161	153	154	118	111	78	36	25	-
1987	2,420	217	145	181	156	149	102	79	36	18	-
1986	2,432	171	187	177	167	139	110	70	29	8	-
1985	2,370	176	177	191	153	127	106	66	26	16	-
1984	2,429	179	183	193	155	128	87	61	21	18	-
1983	2,490	187	194	196	173	147	96	51	16	10	-
1982	2,531	212	198	205	158	120	75	44	16	5	-
1981	2,566	217	215	217	140	115	51	41	9	7	1
1980	2,295	178	205	181	153	97	69	29	16	10	1
1979	2,419	216	220	206	150	112	61	36	17	5	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All other, both sexes											
1998	1,456	101	56	56	39	41	33	34	24	12	-
1997	1,552	87	47	60	46	59	48	29	20	15	2
1996	1,643	91	65	48	55	50	47	29	18	13	4
1995	1,643	84	59	56	45	46	49	31	17	13	-
1994	1,758	84	46	49	38	57	48	26	21	16	1
1993	1,669	77	78	49	45	34	42	28	23	7	-
1992	1,563	69	61	37	49	39	39	30	26	16	-
1991	1,552	91	60	59	58	49	40	39	26	12	3
1990	1,541	82	62	56	50	48	45	23	16	14	-
1989	1,507	67	51	49	55	66	36	24	16	15	-
1988	1,444	76	60	52	54	53	36	33	12	7	-
1987	1,383	67	57	47	48	54	49	35	22	10	-
1986	1,317	77	56	51	50	62	51	28	16	10	1
1985	1,243	66	61	50	55	48	39	27	19	5	-
1984	1,185	65	48	72	50	36	42	26	9	5	1
1983	1,092	61	53	42	37	42	36	18	18	8	-
1982	1,071	43	51	48	37	35	32	11	14	7	1
1981	1,164	67	38	54	45	33	27	26	12	7	-
1980	1,110	48	51	47	42	35	26	12	10	8	-
1979	1,265	62	65	60	53	42	36	20	12	10	2
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All other, male											
1998	1,268	78	47	49	34	39	30	27	23	11	-
1997	1,374	70	42	54	40	51	39	27	19	14	1
1996	1,425	75	53	44	43	45	40	27	14	10	4
1995	1,447	71	51	52	34	39	46	27	16	10	-
1994	1,536	70	37	45	32	46	43	25	19	15	1
1993	1,462	68	68	42	37	29	36	24	21	7	-
1992	1,394	55	54	34	41	31	36	27	25	13	-
1991	1,386	79	51	52	49	43	35	37	24	11	2
1990	1,336	67	55	44	41	39	42	23	15	12	-
1989	1,302	61	42	39	46	57	33	22	13	14	-
1988	1,250	59	49	42	47	46	32	32	12	7	-
1987	1,206	62	50	42	39	48	46	30	20	10	-
1986	1,114	63	48	36	39	53	42	25	14	10	1
1985	1,059	56	52	44	49	43	34	16	15	4	-
1984	1,005	50	38	60	43	30	37	26	9	5	1
1983	941	55	43	37	32	38	32	17	17	7	-
1982	914	37	43	42	31	31	28	10	12	6	1
1981	969	59	33	45	39	27	17	24	12	7	-
1980	946	39	43	40	34	32	25	10	10	8	-
1979	1,047	53	55	44	45	33	31	16	9	9	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
All other, female											
1998	188	23	9	7	5	2	3	7	1	1	-
1997	178	17	5	6	6	8	9	2	1	1	1
1996	218	16	12	4	12	5	7	2	4	3	-
1995	196	13	8	4	11	7	3	4	1	3	-
1994	222	14	9	4	6	11	5	1	2	1	-
1993	207	9	10	7	8	5	6	4	2	-	-
1992	169	14	7	3	8	8	3	3	1	3	-
1991	166	12	9	7	9	6	5	2	2	1	1
1990	205	15	7	12	9	9	3	-	1	2	-
1989	205	6	9	10	9	9	3	2	3	1	-
1988	194	17	11	10	7	7	4	1	-	-	-
1987	177	5	7	5	9	6	3	5	2	-	-
1986	203	14	8	15	11	9	9	3	2	-	-
1985	184	10	9	6	6	5	5	11	4	1	-
1984	180	15	10	12	7	6	5	-	-	-	-
1983	151	6	10	5	5	4	4	1	1	1	-
1982	157	6	8	6	6	4	4	1	2	1	-
1981	195	8	5	9	6	6	10	2	-	-	-
1980	164	9	8	7	8	3	1	2	-	-	-
1979	218	9	10	16	8	9	5	4	3	1	-
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
Black, both sexes											
1998	1,110	76	43	36	36	37	22	29	16	8	-
1997	1,209	67	36	41	36	50	41	23	17	13	2
1996	1,325	71	57	39	48	48	39	24	16	12	2
1995	1,304	65	49	46	37	35	44	30	16	13	-
1994	1,393	71	37	38	29	50	44	22	18	15	-
1993	1,365	61	64	41	34	28	40	24	21	6	-
1992	1,281	52	48	31	46	36	33	30	19	14	-
1991	1,253	80	51	50	43	45	35	36	23	10	3
1990	1,263	75	48	49	43	44	40	23	14	12	-
1989	1,241	56	47	44	50	61	32	24	16	15	-
1988	1,182	63	48	46	46	47	35	29	11	5	-
1987	1,143	54	48	40	41	54	45	34	19	8	-
1986	1,081	67	50	42	42	57	44	24	15	9	-
1985	1,032	58	53	44	50	44	37	25	19	3	-
1984	994	53	41	62	46	32	42	26	8	5	-
1983	906	51	49	38	32	39	33	16	16	8	-
1982	892	36	46	47	36	31	28	10	14	7	1
1981	960	54	31	49	40	32	26	26	11	7	-
1980	946	45	44	43	40	34	25	12	9	8	-
1979	1,053	52	61	56	48	40	34	15	12	8	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
Black, male											
1998	983	61	38	32	32	35	20	23	15	7	-
1997	1,070	52	33	37	31	44	34	21	16	12	1
1996	1,162	60	46	37	40	43	33	22	12	9	2
1995	1,153	57	44	42	27	28	41	27	15	10	-
1994	1,226	60	29	36	24	41	39	21	16	14	-
1993	1,217	55	56	36	26	24	34	21	19	6	-
1992	1,145	41	44	28	38	28	30	27	18	11	-
1991	1,123	70	43	43	38	39	31	34	21	9	2
1990	1,102	62	44	40	35	35	39	23	13	10	-
1989	1,078	51	39	35	41	53	29	22	13	14	-
1988	1,015	47	38	38	39	40	31	28	11	5	-
1987	1,002	49	43	36	33	48	43	29	17	8	-
1986	919	55	45	28	31	49	35	21	13	9	-
1985	881	48	45	39	44	40	32	14	15	2	-
1984	840	39	35	50	40	27	37	26	8	5	-
1983	785	46	41	35	27	35	30	15	16	7	-
1982	763	31	39	41	30	27	24	9	12	6	1
1981	800	49	27	40	34	26	16	24	11	7	-
1980	813	37	37	37	32	31	24	10	9	8	-
1979	868	46	52	41	40	32	29	12	9	7	2
Intentional self-harm (suicide) by discharge of firearms (E955.0-E955.4)											
Black, female											
1998	127	15	5	4	4	2	2	6	1	1	-
1997	139	15	3	4	5	6	7	2	1	1	1
1996	163	11	11	2	8	5	6	2	4	3	-
1995	151	8	5	4	10	7	3	3	1	3	-
1994	167	11	8	2	5	9	5	1	2	1	-
1993	148	6	8	5	8	4	6	3	2	-	-
1992	136	11	4	3	8	8	3	3	1	3	-
1991	130	10	8	7	5	6	4	2	2	1	1
1990	161	13	4	9	8	9	1	-	1	2	-
1989	163	5	8	9	9	8	3	2	3	1	-
1988	167	16	10	8	7	7	4	1	-	-	-
1987	141	5	5	4	8	6	2	5	2	-	-
1986	162	12	5	14	11	8	9	3	2	-	-
1985	151	10	8	5	6	4	5	11	4	1	-
1984	154	14	6	12	6	5	5	-	-	-	-
1983	121	5	8	3	5	4	3	1	-	1	-
1982	129	5	7	6	6	4	4	1	2	1	-
1981	160	5	4	9	6	6	10	2	-	-	-
1980	133	8	7	6	8	3	1	2	-	-	-
1979	185	6	9	15	8	8	5	3	3	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All races, both sexes											
1998	13,151	1,405	1,074	665	425	406	360	319	297	308	13
1997	12,969	1,355	946	640	447	377	380	372	280	311	11
1996	12,737	1,284	846	629	454	410	362	352	288	259	14
1995	12,781	1,207	821	533	461	446	429	339	296	288	6
1994	12,377	1,056	797	560	455	426	429	360	288	314	13
1993	12,162	999	736	569	495	479	457	378	298	295	14
1992	12,315	991	738	555	590	476	396	423	342	290	11
1991	12,284	960	731	627	608	489	429	387	321	313	13
1990	12,021	851	678	619	597	505	480	411	332	270	5
1989	12,054	845	653	627	599	551	440	394	308	288	13
1988	12,238	827	631	662	574	574	481	456	330	258	13
1987	12,660	846	744	707	662	580	523	466	321	280	11
1986	12,751	827	771	767	692	549	547	465	307	264	8
1985	12,090	782	693	787	664	535	482	433	299	226	5
1984	12,173	831	766	828	686	545	512	366	285	219	13
1983	11,695	780	798	817	617	517	479	406	303	213	10
1982	11,682	776	844	853	677	550	441	374	239	195	10
1981	11,457	759	807	842	609	539	447	336	237	185	16
1980	11,473	789	865	835	642	542	458	388	229	202	14
1979	11,663	815	914	797	709	581	473	400	240	208	12
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All races, male											
1998	9,434	997	686	429	268	263	238	204	190	172	13
1997	9,298	915	577	407	306	256	233	222	174	199	10
1996	9,190	865	562	424	282	248	228	234	179	164	9
1995	9,309	825	557	353	315	275	255	213	179	177	6
1994	8,887	701	510	334	294	250	259	228	179	184	11
1993	8,626	654	467	369	290	292	269	227	182	188	11
1992	8,655	651	446	343	359	287	238	263	215	173	7
1991	8,649	596	474	381	386	295	270	235	211	191	11
1990	8,439	581	430	352	365	299	271	252	207	166	3
1989	8,422	539	392	377	368	327	292	249	202	179	12
1988	8,422	483	381	399	331	342	279	287	202	176	11
1987	8,733	529	420	438	390	339	320	296	204	203	9
1986	8,708	497	453	460	375	334	327	266	183	178	7
1985	8,336	469	418	498	398	322	281	263	199	154	5
1984	8,185	476	442	481	394	292	301	219	173	145	11
1983	7,827	468	468	467	357	298	272	248	194	132	9
1982	7,753	454	494	487	385	313	251	222	153	135	8
1981	7,431	416	421	453	346	304	248	211	153	129	13
1980	7,568	455	496	497	358	318	275	244	155	127	14
1979	7,350	442	480	443	380	319	280	238	150	142	11

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All races, female											
1998	3,717	408	388	236	157	143	122	115	107	136	-
1997	3,671	440	369	233	141	121	147	150	106	112	1
1996	3,547	419	284	205	172	162	134	118	109	95	5
1995	3,472	382	264	180	146	171	174	126	117	111	-
1994	3,490	355	287	226	161	176	170	132	109	130	2
1993	3,536	345	269	200	205	187	188	151	116	107	3
1992	3,660	340	292	212	231	189	158	160	127	117	4
1991	3,635	364	257	246	222	194	159	152	110	122	2
1990	3,582	270	248	267	232	206	209	159	125	104	2
1989	3,632	306	261	250	231	224	148	145	106	109	1
1988	3,816	344	250	263	243	232	202	169	128	82	2
1987	3,927	317	324	269	272	241	203	170	117	77	2
1986	4,043	330	318	307	317	215	220	199	124	86	1
1985	3,754	313	275	289	266	213	201	170	100	72	-
1984	3,988	355	324	347	292	253	211	147	112	74	2
1983	3,868	312	330	350	260	219	207	158	109	81	1
1982	3,929	322	350	366	292	237	190	152	86	60	2
1981	4,026	343	386	389	263	235	199	125	84	56	3
1980	3,905	334	369	338	284	224	183	144	74	75	-
1979	4,313	373	434	354	329	262	193	162	90	66	1
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
White, both sexes											
1998	11,680	1,280	978	622	393	368	331	304	277	293	11
1997	11,499	1,238	864	580	406	347	353	350	260	292	8
1996	11,333	1,176	782	577	421	381	333	329	273	250	13
1995	11,327	1,108	756	497	416	403	391	317	285	277	5
1994	10,969	968	749	510	411	398	402	339	277	302	8
1993	10,764	914	677	522	461	441	426	358	286	276	12
1992	11,005	894	681	527	552	445	378	405	327	283	9
1991	11,022	878	676	581	565	462	400	366	311	301	10
1990	10,742	783	621	578	562	477	454	394	324	255	5
1989	10,753	788	604	599	556	517	416	377	294	279	11
1988	11,065	766	581	628	554	547	464	443	314	248	11
1987	11,464	780	691	680	621	539	494	453	308	266	9
1986	11,601	777	722	723	651	519	524	450	297	255	5
1985	10,967	717	657	744	633	501	453	420	288	217	3
1984	11,074	768	715	790	646	513	488	355	277	209	8
1983	10,649	717	762	760	587	503	459	385	292	205	7
1982	10,652	721	793	809	651	522	429	357	229	189	9
1981	10,477	706	763	789	585	516	431	324	230	176	13
1980	10,543	732	814	789	623	524	446	375	223	191	13
1979	10,667	757	879	764	683	556	459	387	231	205	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
White, male											
1998	8,338	900	625	401	247	234	217	196	178	165	11
1997	8,222	837	525	364	278	236	217	208	162	190	7
1996	8,164	790	521	392	261	236	213	220	169	160	8
1995	8,240	758	509	334	279	252	232	198	172	172	5
1994	7,830	644	478	309	264	232	241	216	171	178	7
1993	7,605	595	428	337	271	265	255	217	175	178	10
1992	7,718	586	412	327	336	268	227	251	205	170	7
1991	7,718	538	442	346	357	276	253	223	206	185	8
1990	7,499	533	394	328	339	283	257	241	203	158	3
1989	7,480	506	366	361	343	305	272	239	196	173	10
1988	7,574	445	348	375	318	325	271	277	197	171	10
1987	7,855	486	390	421	363	309	304	286	192	194	7
1986	7,866	463	427	433	352	312	314	259	176	172	4
1985	7,506	426	399	465	384	300	267	255	190	149	3
1984	7,383	438	411	459	370	272	285	211	168	140	6
1983	7,079	428	449	431	337	287	259	238	186	126	6
1982	7,007	421	459	461	369	293	244	210	146	132	7
1981	6,757	382	398	421	328	287	238	203	147	126	11
1980	6,910	417	466	463	345	307	270	237	152	118	13
1979	6,645	406	459	421	372	305	270	233	146	139	11
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
White, female											
1998	3,342	380	353	221	146	134	114	108	99	128	-
1997	3,277	401	339	216	128	111	136	142	98	102	1
1996	3,169	386	261	185	160	145	120	109	104	90	5
1995	3,087	350	247	163	137	151	159	119	113	105	-
1994	3,139	324	271	201	147	166	161	123	106	124	1
1993	3,159	319	249	185	190	176	171	141	111	98	2
1992	3,287	308	269	200	216	177	151	154	122	113	2
1991	3,304	340	234	235	208	186	147	143	105	116	2
1990	3,243	250	227	250	223	194	197	153	121	97	2
1989	3,273	282	238	238	213	212	144	138	98	106	1
1988	3,491	321	233	253	236	222	193	166	117	77	1
1987	3,609	294	301	259	258	230	190	167	116	72	2
1986	3,735	314	295	290	299	207	210	191	121	83	1
1985	3,461	291	258	279	249	201	186	165	98	68	-
1984	3,691	330	304	331	276	241	203	144	109	69	2
1983	3,570	289	313	329	250	216	200	147	106	79	1
1982	3,645	300	334	348	282	229	185	147	83	57	2
1981	3,720	324	365	368	257	229	193	121	83	50	2
1980	3,633	315	348	326	278	217	176	138	71	73	-
1979	4,022	351	420	343	311	251	189	154	85	66	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All other, both sexes											
1998	1,471	125	96	43	32	38	29	15	20	15	2
1997	1,470	117	82	60	41	30	27	22	20	19	3
1996	1,404	108	64	52	33	29	29	23	15	9	1
1995	1,454	99	65	36	45	43	38	22	11	11	1
1994	1,408	88	48	50	44	28	27	21	11	12	5
1993	1,398	85	59	47	34	38	31	20	12	19	2
1992	1,310	97	57	28	38	31	18	18	15	7	2
1991	1,262	82	55	46	43	27	29	21	10	12	3
1990	1,279	68	57	41	35	28	26	17	8	15	-
1989	1,301	57	49	28	43	34	24	17	14	9	2
1988	1,173	61	50	34	20	27	17	13	16	10	2
1987	1,196	66	53	27	41	41	29	13	13	14	2
1986	1,150	50	49	44	41	30	23	15	10	9	3
1985	1,123	65	36	43	31	34	29	13	11	9	2
1984	1,099	63	51	38	40	32	24	11	8	10	5
1983	1,046	63	36	57	30	14	20	21	11	8	3
1982	1,030	55	51	44	26	28	12	17	10	6	1
1981	980	53	44	53	24	23	16	12	7	9	3
1980	930	57	51	46	19	18	12	13	6	11	1
1979	996	58	35	33	26	25	14	13	9	3	-
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All other, male											
1998	1,096	97	61	28	21	29	21	8	12	7	2
1997	1,076	78	52	43	28	20	16	14	12	9	3
1996	1,026	75	41	32	21	12	15	14	10	4	1
1995	1,069	67	48	19	36	23	23	15	7	5	1
1994	1,057	57	32	25	30	18	18	12	8	6	4
1993	1,021	59	39	32	19	27	14	10	7	10	1
1992	937	65	34	16	23	19	11	12	10	3	-
1991	931	58	32	35	29	19	17	12	5	6	3
1990	940	48	36	24	26	16	14	11	4	8	-
1989	942	33	26	16	25	22	20	10	6	6	2
1988	848	38	33	24	13	17	8	10	5	5	1
1987	878	43	30	17	27	30	16	10	12	9	2
1986	842	34	26	27	23	22	13	7	7	6	3
1985	830	43	19	33	14	22	14	8	9	5	2
1984	802	38	31	22	24	20	16	8	5	5	5
1983	748	40	19	36	20	11	13	10	8	6	3
1982	746	33	35	26	16	20	7	12	7	3	1
1981	674	34	23	32	18	17	10	8	6	3	2
1980	658	38	30	34	13	11	5	7	3	9	1
1979	705	36	21	22	8	14	10	5	4	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
All other, female											
1998	375	28	35	15	11	9	8	7	8	8	-
1997	394	39	30	17	13	10	11	8	8	10	-
1996	378	33	23	20	12	17	14	9	5	5	-
1995	385	32	17	17	9	20	15	7	4	6	-
1994	351	31	16	25	14	10	9	9	3	6	1
1993	377	26	20	15	15	11	17	10	5	9	1
1992	373	32	23	12	15	12	7	6	5	4	2
1991	331	24	23	11	14	8	12	9	5	6	-
1990	339	20	21	17	9	12	12	6	4	7	-
1989	359	24	23	12	18	12	4	7	8	3	-
1988	325	23	17	10	7	10	9	3	11	5	1
1987	318	23	23	10	14	11	13	3	1	5	-
1986	308	16	23	17	18	8	10	8	3	3	-
1985	293	22	17	10	17	12	15	5	2	4	-
1984	297	25	20	16	16	12	8	3	3	5	-
1983	298	23	17	21	10	3	7	11	3	2	-
1982	284	22	16	18	10	8	5	5	3	3	-
1981	306	19	21	21	6	6	6	4	1	6	1
1980	272	19	21	12	6	7	7	6	3	2	-
1979	291	22	14	11	18	11	4	8	5	-	-
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
Black, both sexes											
1998	867	87	56	25	13	20	10	5	-	3	2
1997	894	71	44	33	21	7	9	8	4	4	3
1996	839	67	41	22	19	11	10	6	4	2	1
1995	927	67	41	16	22	22	17	7	3	-	1
1994	878	56	24	29	19	13	16	10	2	2	2
1993	894	60	38	28	14	20	9	12	4	8	1
1992	862	68	35	16	20	16	10	3	6	3	2
1991	844	52	33	25	21	16	13	6	3	4	2
1990	848	53	43	18	15	14	15	4	2	5	-
1989	912	38	31	16	20	19	11	7	2	2	2
1988	840	43	34	24	16	18	3	9	5	2	2
1987	820	43	30	13	26	22	15	7	6	1	1
1986	811	34	32	30	28	13	12	9	5	3	3
1985	763	45	32	20	19	16	18	9	6	4	1
1984	766	40	35	23	28	17	14	3	3	3	4
1983	717	42	24	35	20	11	9	9	3	3	1
1982	747	35	42	30	15	20	7	7	4	3	1
1981	698	40	32	36	15	11	10	7	3	2	3
1980	661	35	34	26	15	12	5	6	2	2	1
1979	759	38	26	25	19	18	4	5	2	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
Black, male											
1998	676	69	42	17	7	16	9	3	-	2	2
1997	694	52	27	25	14	4	7	6	2	3	3
1996	658	53	25	17	14	5	6	2	3	1	1
1995	725	50	33	9	21	12	10	5	3	-	1
1994	696	38	21	15	15	8	12	6	1	1	2
1993	688	46	30	20	5	14	5	5	2	4	-
1992	658	48	24	9	15	12	6	3	5	1	-
1991	659	39	20	19	15	11	6	5	1	3	2
1990	635	38	30	9	11	7	9	2	2	3	-
1989	693	23	15	12	13	11	11	5	1	2	2
1988	633	27	22	17	9	12	2	6	3	2	1
1987	633	34	19	11	19	17	6	5	5	1	1
1986	618	24	16	20	15	10	8	4	4	3	3
1985	600	34	18	17	8	13	13	5	6	3	1
1984	592	25	24	16	18	10	10	2	2	2	4
1983	536	28	12	29	15	9	7	5	3	1	1
1982	564	21	29	22	11	14	4	6	3	3	1
1981	515	29	20	22	12	8	5	5	3	-	2
1980	484	24	25	20	11	7	1	4	1	2	1
1979	560	25	18	19	8	12	3	3	1	1	-
Intentional self-harm (suicide) by other and unspecified means and their sequelae (E950-E954,E955.5-E959)											
Black, female											
1998	191	18	14	8	6	4	1	2	-	1	-
1997	200	19	17	8	7	3	2	2	2	1	-
1996	181	14	16	5	5	6	4	4	1	1	-
1995	202	17	8	7	1	10	7	2	-	-	-
1994	182	18	3	14	4	5	4	4	1	1	-
1993	206	14	8	8	9	6	4	7	2	4	1
1992	204	20	11	7	5	4	4	-	1	2	2
1991	185	13	13	6	6	5	7	1	2	1	-
1990	213	15	13	9	4	7	6	2	-	2	-
1989	219	15	16	4	7	8	-	2	1	-	-
1988	207	16	12	7	7	6	1	3	2	-	1
1987	187	9	11	2	7	5	9	2	1	-	-
1986	193	10	16	10	13	3	4	5	1	-	-
1985	163	11	14	3	11	3	5	4	-	1	-
1984	174	15	11	7	10	7	4	1	1	1	-
1983	181	14	12	6	5	2	2	4	-	2	-
1982	183	14	13	8	4	6	3	1	1	-	-
1981	183	11	12	14	3	3	5	2	-	2	1
1980	177	11	9	6	4	5	4	2	1	-	-
1979	199	13	8	6	11	6	1	2	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
All races, both sexes											
1998	17,893	1,033	662	449	308	251	213	183	135	99	55
1997	19,491	1,161	668	477	357	286	253	203	129	149	47
1996	20,634	1,135	726	493	369	316	231	195	137	110	62
1995	22,552	1,146	743	549	393	342	261	205	137	118	52
1994	24,547	1,139	751	500	398	346	285	222	165	120	52
1993	25,653	1,241	781	540	429	372	314	207	170	139	67
1992	25,144	1,216	786	523	460	378	311	230	171	134	54
1991	26,254	1,267	826	607	537	428	308	269	169	130	81
1990	24,614	1,118	749	551	496	384	297	249	178	138	96
1989	22,578	1,071	778	555	516	421	317	240	166	130	81
1988	21,784	1,010	693	617	501	428	331	265	168	138	90
1987	20,812	1,025	762	686	506	427	333	278	170	147	99
1986	21,462	1,054	817	677	504	431	319	241	174	130	87
1985	19,628	1,039	772	700	560	429	297	232	147	110	62
1984	19,510	1,057	834	699	583	368	316	219	162	113	58
1983	19,922	1,054	869	753	591	410	289	251	153	129	63
1982	22,073	1,189	1,080	794	625	455	333	240	145	141	63
1981	23,361	1,377	1,149	911	640	444	317	264	156	126	58
1980	23,967	1,354	1,147	869	649	534	353	244	160	118	75
1979	22,202	1,376	1,085	874	630	494	332	202	158	109	77
Assault (homicide) (E960-E969)											
All races, male											
1998	13,652	748	516	340	230	172	128	111	56	37	41
1997	15,107	858	504	350	262	194	153	113	61	66	31
1996	15,945	816	555	379	275	216	140	99	69	40	41
1995	17,408	875	546	410	301	235	146	102	57	42	36
1994	19,342	880	579	384	295	243	169	127	78	54	35
1993	19,945	953	614	406	321	246	182	112	69	54	46
1992	19,779	947	593	414	345	254	190	123	88	65	38
1991	20,518	963	632	463	390	286	206	140	80	49	56
1990	19,298	871	589	413	381	269	186	135	77	56	70
1989	17,366	835	605	420	382	273	197	136	76	65	53
1988	16,472	792	521	462	367	269	198	150	77	59	65
1987	15,571	777	580	532	368	288	194	153	84	60	66
1986	16,332	835	632	539	357	292	192	134	83	50	65
1985	14,806	796	592	538	407	290	186	125	64	38	47
1984	14,759	835	677	540	434	254	210	119	79	55	36
1983	15,253	808	690	574	462	277	197	131	77	55	47
1982	17,035	960	860	625	469	314	213	140	68	66	42
1981	18,288	1,097	944	711	483	321	218	144	74	69	46
1980	18,779	1,099	924	705	494	395	227	142	90	51	51
1979	17,288	1,081	899	708	467	344	199	124	73	45	54

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
All races, female											
1998	4,241	285	146	109	78	79	85	72	79	62	14
1997	4,384	303	164	127	95	92	100	90	68	83	16
1996	4,689	319	171	114	94	100	91	96	68	70	21
1995	5,144	271	197	139	92	107	115	103	80	76	16
1994	5,205	259	172	116	103	103	116	95	87	66	17
1993	5,708	288	167	134	108	126	132	95	101	85	21
1992	5,365	269	193	109	115	124	121	107	83	69	16
1991	5,736	304	194	144	147	142	102	129	89	81	25
1990	5,316	247	160	138	115	115	111	114	101	82	26
1989	5,212	236	173	135	134	148	120	104	90	65	28
1988	5,312	218	172	155	134	159	133	115	91	79	25
1987	5,241	248	182	154	138	139	139	125	86	87	33
1986	5,130	219	185	138	147	139	127	107	91	80	22
1985	4,822	243	180	162	153	139	111	107	83	72	15
1984	4,751	222	157	159	149	114	106	100	83	58	22
1983	4,669	246	179	179	129	133	92	120	76	74	16
1982	5,038	229	220	169	156	141	120	100	77	75	21
1981	5,073	280	205	200	157	123	99	120	82	57	12
1980	5,188	255	223	164	155	139	126	102	70	67	24
1979	4,914	295	186	166	163	150	133	78	85	64	23
Assault (homicide) (E960-E969)											
White, both sexes											
1998	9,008	596	413	297	202	176	156	131	111	81	47
1997	9,676	668	432	327	224	195	189	153	103	119	34
1996	10,088	656	456	323	248	225	164	141	101	83	42
1995	11,130	668	467	342	250	221	179	134	105	79	39
1994	11,718	659	491	331	265	224	183	156	121	88	36
1993	12,069	739	501	355	270	243	212	137	119	98	45
1992	12,233	727	503	346	310	228	211	166	118	98	39
1991	12,631	762	512	383	343	275	197	180	107	96	61
1990	11,953	680	468	343	318	249	207	167	122	88	69
1989	11,092	667	478	337	322	260	206	152	122	79	56
1988	10,902	606	424	405	333	269	224	186	111	94	64
1987	10,950	623	461	432	330	281	228	192	125	110	78
1986	11,528	626	470	423	325	267	215	169	129	91	72
1985	11,003	624	472	460	355	289	195	176	112	83	51
1984	10,952	628	520	434	376	229	204	141	124	81	44
1983	11,075	619	529	459	370	269	194	167	109	101	49
1982	12,272	670	651	495	389	284	221	169	112	115	45
1981	12,904	785	692	546	404	277	212	183	120	102	42
1980	13,360	775	670	535	395	361	254	187	123	98	54
1979	12,149	779	608	504	376	324	237	141	123	95	56

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
White, male											
1998	6,479	419	306	213	147	114	92	75	48	28	35
1997	7,118	477	321	235	164	130	104	81	47	52	25
1996	7,351	451	323	241	179	150	95	67	48	27	30
1995	8,109	473	325	245	185	148	83	62	38	31	28
1994	8,810	492	377	252	191	152	106	85	55	37	24
1993	8,845	547	377	257	193	156	109	72	42	34	30
1992	9,228	543	364	272	233	147	114	83	62	41	29
1991	9,437	555	383	284	242	173	118	84	51	35	40
1990	8,954	505	355	252	230	173	120	81	50	37	46
1989	8,127	506	361	258	236	160	118	78	51	39	36
1988	7,837	452	312	303	233	159	130	89	49	37	48
1987	7,804	477	334	338	229	177	118	95	58	39	51
1986	8,409	471	348	331	217	169	122	85	54	31	53
1985	7,963	455	354	341	238	185	114	88	43	27	39
1984	8,000	475	410	317	267	146	126	67	55	36	29
1983	8,198	456	401	331	265	175	124	79	47	38	37
1982	9,097	532	497	374	279	178	133	94	48	54	31
1981	9,779	609	545	418	304	186	134	85	51	51	33
1980	10,185	608	530	419	292	264	153	101	63	40	38
1979	9,210	601	488	392	272	218	139	80	58	37	38
Assault (homicide) (E960-E969)											
White, female											
1998	2,529	177	107	84	55	62	64	56	63	53	12
1997	2,558	191	111	92	60	65	85	72	56	67	9
1996	2,737	205	133	82	69	75	69	74	53	56	12
1995	3,021	195	142	97	65	73	96	72	67	48	11
1994	2,908	167	114	79	74	72	77	71	66	51	12
1993	3,224	192	124	98	77	87	103	65	77	64	15
1992	3,005	184	139	74	77	81	97	83	56	57	10
1991	3,194	207	129	99	101	102	79	96	56	61	21
1990	2,999	175	113	91	88	76	87	86	72	51	23
1989	2,965	161	117	79	86	100	88	74	71	40	20
1988	3,065	154	112	102	100	110	94	97	62	57	16
1987	3,146	146	127	94	101	104	110	97	67	71	27
1986	3,119	155	122	92	108	98	93	84	75	60	19
1985	3,040	169	118	119	117	104	81	88	69	56	12
1984	2,952	153	110	117	109	83	78	74	69	45	15
1983	2,877	163	128	128	105	94	70	88	62	63	12
1982	3,175	138	154	121	110	106	88	75	64	61	14
1981	3,125	176	147	128	100	91	78	98	69	51	9
1980	3,175	167	140	116	103	97	101	86	60	58	16
1979	2,939	178	120	112	104	106	98	61	65	58	18

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
All other, both sexes											
1998	8,885	437	249	152	106	75	57	52	24	18	8
1997	9,815	493	236	150	133	91	64	50	26	30	13
1996	10,546	479	270	170	121	91	67	54	36	27	20
1995	11,422	478	276	207	143	121	82	71	32	39	13
1994	12,829	480	260	169	133	122	102	66	44	32	16
1993	13,584	502	280	185	159	129	102	70	51	41	22
1992	12,911	489	283	177	150	150	100	64	53	36	15
1991	13,623	505	314	224	194	153	111	89	62	34	20
1990	12,661	438	281	208	178	135	90	82	56	50	27
1989	11,486	404	300	218	194	161	111	88	44	51	25
1988	10,882	404	269	212	168	159	107	79	57	44	26
1987	9,862	402	301	254	176	146	105	86	45	37	21
1986	9,934	428	347	254	179	164	104	72	45	39	15
1985	8,625	415	300	240	205	140	102	56	35	27	11
1984	8,558	429	314	265	207	139	112	78	38	32	14
1983	8,847	435	340	294	221	141	95	84	44	28	14
1982	9,801	519	429	299	236	171	112	71	33	26	18
1981	10,457	592	457	365	236	167	105	81	36	24	16
1980	10,607	579	477	334	254	173	99	57	37	20	21
1979	10,053	597	477	370	254	170	95	61	35	14	21
Assault (homicide) (E960-E969)											
All other, male											
1998	7,173	329	210	127	83	58	36	36	8	9	6
1997	7,989	381	183	115	98	64	49	32	14	14	6
1996	8,594	365	232	138	96	66	45	32	21	13	11
1995	9,299	402	221	165	116	87	63	40	19	11	8
1994	10,532	388	202	132	104	91	63	42	23	17	11
1993	11,100	406	237	149	128	90	73	40	27	20	16
1992	10,551	404	229	142	112	107	76	40	26	24	9
1991	11,081	408	249	179	148	113	88	56	29	14	16
1990	10,344	366	234	161	151	96	66	54	27	19	24
1989	9,239	329	244	162	146	113	79	58	25	26	17
1988	8,635	340	209	159	134	110	68	61	28	22	17
1987	7,767	300	246	194	139	111	76	58	26	21	15
1986	7,923	364	284	208	140	123	70	49	29	19	12
1985	6,843	341	238	197	169	105	72	37	21	11	8
1984	6,759	360	267	223	167	108	84	52	24	19	7
1983	7,055	352	289	243	197	102	73	52	30	17	10
1982	7,938	428	363	251	190	136	80	46	20	12	11
1981	8,509	488	399	293	179	135	84	59	23	18	13
1980	8,594	491	394	286	202	131	74	41	27	11	13
1979	8,078	480	411	316	195	126	60	44	15	8	16

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
All other, female											
1998	1,712	108	39	25	23	17	21	16	16	9	2
1997	1,826	112	53	35	35	27	15	18	12	16	7
1996	1,952	114	38	32	25	25	22	22	15	14	9
1995	2,123	76	55	42	27	34	19	31	13	28	5
1994	2,297	92	58	37	29	31	39	24	21	15	5
1993	2,484	96	43	36	31	39	29	30	24	21	6
1992	2,360	85	54	35	38	43	24	24	27	12	6
1991	2,542	97	65	45	46	40	23	33	33	20	4
1990	2,317	72	47	47	27	39	24	28	29	31	3
1989	2,247	75	56	56	48	48	32	30	19	25	8
1988	2,247	64	60	53	34	49	39	18	29	22	9
1987	2,095	102	55	60	37	35	29	28	19	16	6
1986	2,011	64	63	46	39	41	34	23	16	20	3
1985	1,782	74	62	43	36	35	30	19	14	16	3
1984	1,799	69	47	42	40	31	28	26	14	13	7
1983	1,792	83	51	51	24	39	22	32	14	11	4
1982	1,863	91	66	48	46	35	32	25	13	14	7
1981	1,948	104	58	72	57	32	21	22	13	6	3
1980	2,013	88	83	48	52	42	25	16	10	9	8
1979	1,975	117	66	54	59	44	35	17	20	6	5
Assault (homicide) (E960-E969)											
Black, both sexes											
1998	8,293	395	224	135	91	64	49	46	22	18	8
1997	9,147	439	204	131	120	79	58	46	25	28	12
1996	9,886	437	233	153	111	81	62	48	36	27	17
1995	10,689	443	244	181	127	114	76	67	29	38	12
1994	12,092	436	239	155	119	115	96	63	42	31	12
1993	12,811	455	256	160	147	122	100	68	50	41	18
1992	12,220	442	266	159	142	140	90	62	51	32	13
1991	12,859	468	276	196	179	149	106	86	61	33	18
1990	12,035	403	257	189	169	125	84	80	55	47	26
1989	10,858	379	278	201	184	149	103	83	41	50	23
1988	10,327	383	239	197	161	150	103	76	52	42	25
1987	9,383	376	277	240	164	141	100	83	42	36	17
1986	9,395	397	324	239	170	159	102	66	43	36	15
1985	8,189	385	281	228	197	134	97	53	34	27	10
1984	8,135	398	298	250	200	135	104	75	36	28	13
1983	8,387	405	315	281	210	134	90	80	43	28	11
1982	9,357	488	407	290	224	166	105	65	30	25	17
1981	10,019	561	434	352	228	161	96	79	35	21	13
1980	10,172	555	454	324	243	165	92	53	34	18	18
1979	9,656	574	458	352	242	164	88	58	34	14	19

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) (E960-E969)											
Black, male											
1998	6,749	303	191	112	73	51	32	30	8	9	6
1997	7,496	345	164	102	87	56	43	31	14	13	6
1996	8,089	335	203	124	86	61	42	30	21	13	8
1995	8,757	373	198	146	103	84	61	38	17	10	8
1994	9,969	355	188	123	95	87	60	40	22	16	8
1993	10,516	375	219	131	119	86	72	39	26	20	13
1992	10,034	371	215	131	107	100	72	40	25	22	8
1991	10,531	381	225	159	139	111	85	56	28	14	14
1990	9,876	340	217	150	144	88	61	53	27	17	23
1989	8,789	310	227	150	140	105	73	54	24	25	16
1988	8,239	328	185	147	131	107	66	58	25	22	16
1987	7,418	281	227	185	130	108	73	55	24	20	12
1986	7,536	339	267	198	134	119	68	45	27	17	12
1985	6,527	320	227	188	165	100	70	35	20	11	7
1984	6,460	341	257	213	161	105	77	49	22	18	7
1983	6,720	329	269	233	187	96	70	49	29	17	7
1982	7,615	403	347	245	181	131	75	42	18	11	10
1981	8,195	463	382	284	174	129	76	57	22	16	10
1980	8,274	473	379	280	193	123	67	37	26	9	10
1979	7,787	460	396	301	185	121	57	42	15	8	15
Assault (homicide) (E960-E969)											
Black, female											
1998	1,544	92	33	23	18	13	17	16	14	9	2
1997	1,651	94	40	29	33	23	15	15	11	15	6
1996	1,797	102	30	29	25	20	20	18	15	14	9
1995	1,932	70	46	35	24	30	15	29	12	28	4
1994	2,123	81	51	32	24	28	36	23	20	15	4
1993	2,295	80	37	29	28	36	28	29	24	21	5
1992	2,186	71	51	28	35	40	18	22	26	10	5
1991	2,328	87	51	37	40	38	21	30	33	19	4
1990	2,159	63	40	39	25	37	23	27	28	30	3
1989	2,069	69	51	51	44	44	30	29	17	25	7
1988	2,088	55	54	50	30	43	37	18	27	20	9
1987	1,965	95	50	55	34	33	27	28	18	16	5
1986	1,859	58	57	41	36	40	34	21	16	19	3
1985	1,662	65	54	40	32	34	27	18	14	16	3
1984	1,675	57	41	37	39	30	27	26	14	10	6
1983	1,667	76	46	48	23	38	20	31	14	11	4
1982	1,742	85	60	45	43	35	30	23	12	14	7
1981	1,824	98	52	68	54	32	20	22	13	5	3
1980	1,898	82	75	44	50	42	25	16	8	9	8
1979	1,869	114	62	51	57	43	31	16	19	6	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All races, both sexes											
1998	11,798	589	382	231	163	110	92	55	30	19	18
1997	13,252	688	380	256	181	126	98	60	28	39	17
1996	14,037	678	430	280	174	145	90	75	43	29	16
1995	15,551	675	442	302	209	163	106	67	39	23	24
1994	17,527	742	461	300	225	154	124	77	40	18	22
1993	18,253	818	484	314	213	194	125	80	40	26	26
1992	17,488	750	458	312	249	160	108	82	49	29	23
1991	17,746	797	460	329	253	192	124	80	34	24	41
1990	16,218	712	448	299	244	179	103	92	50	31	40
1989	14,464	645	441	296	254	191	126	78	43	22	31
1988	13,645	609	402	319	259	177	119	83	42	30	40
1987	12,657	608	417	340	243	194	116	78	59	31	30
1986	13,029	655	458	334	234	188	120	70	51	23	39
1985	11,836	628	437	381	275	199	122	79	41	26	21
1984	11,815	660	475	392	294	179	117	70	42	24	24
1983	12,040	666	513	402	304	188	108	72	35	12	25
1982	13,830	748	666	437	330	228	146	87	39	25	21
1981	15,089	901	712	541	369	222	131	85	32	23	25
1980	15,500	872	680	505	371	292	141	74	49	22	36
1979	14,491	923	696	539	354	238	163	77	39	24	28
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All races, male											
1998	9,771	444	308	177	120	74	55	29	10	8	15
1997	11,147	536	302	196	146	86	65	33	9	18	14
1996	11,735	518	334	234	134	105	54	38	21	9	14
1995	13,021	547	339	230	169	118	65	35	14	9	19
1994	14,766	591	361	239	166	115	80	46	17	7	19
1993	15,228	655	399	240	164	141	79	43	22	9	22
1992	14,747	604	360	257	189	108	74	46	28	18	20
1991	14,926	620	373	257	189	138	84	43	17	11	34
1990	13,629	559	352	229	194	133	64	55	18	16	34
1989	12,018	514	355	225	197	136	86	46	24	12	24
1988	11,134	488	303	253	180	116	69	50	24	16	33
1987	10,202	464	327	276	175	127	66	46	33	11	24
1986	10,656	526	364	272	170	134	83	40	30	13	32
1985	9,532	492	329	293	209	144	82	45	22	11	21
1984	9,615	534	392	315	220	130	85	45	24	15	22
1983	9,863	519	411	311	244	139	87	48	19	8	21
1982	11,402	621	539	355	260	170	105	61	24	18	15
1981	12,548	727	602	439	290	176	97	64	21	13	22
1980	12,878	703	557	422	289	229	104	53	34	13	27
1979	12,014	751	589	452	272	187	118	58	21	12	20

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All races, female											
1998	2,027	145	74	54	43	36	37	26	20	11	3
1997	2,105	152	78	60	35	40	33	27	19	21	3
1996	2,302	160	96	46	40	40	36	37	22	20	2
1995	2,530	128	103	72	40	45	41	32	25	14	5
1994	2,761	151	100	61	59	39	44	31	23	11	3
1993	3,025	163	85	74	49	53	46	37	18	17	4
1992	2,741	146	98	55	60	52	34	36	21	11	3
1991	2,820	177	87	72	64	54	40	37	17	13	7
1990	2,589	153	96	70	50	46	39	37	32	15	6
1989	2,446	131	86	71	57	55	40	32	19	10	7
1988	2,511	121	99	66	79	61	50	33	18	14	7
1987	2,455	144	90	64	68	67	50	32	26	20	6
1986	2,373	129	94	62	64	54	37	30	21	10	7
1985	2,304	136	108	88	66	55	40	34	19	15	-
1984	2,200	126	83	77	74	49	32	25	18	9	2
1983	2,177	147	102	91	60	49	21	24	16	4	4
1982	2,428	127	127	82	70	58	41	26	15	7	6
1981	2,541	174	110	102	79	46	34	21	11	10	3
1980	2,622	169	123	83	82	63	37	21	15	9	9
1979	2,477	172	107	87	82	51	45	19	18	12	8
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
White, both sexes											
1998	5,412	347	229	153	114	76	71	43	25	16	16
1997	6,025	390	251	175	119	90	70	50	21	35	13
1996	6,240	392	265	188	116	95	63	58	30	22	7
1995	7,144	386	273	189	127	101	73	51	33	17	17
1994	7,774	427	302	198	150	92	83	55	30	16	16
1993	7,918	504	306	198	136	132	85	56	28	20	16
1992	7,927	452	289	204	174	101	72	66	39	23	14
1991	7,944	487	286	217	163	125	86	62	23	18	28
1990	7,376	432	296	186	159	110	76	69	36	25	25
1989	6,718	404	276	187	151	117	85	51	33	13	21
1988	6,475	387	261	222	172	112	88	54	34	23	27
1987	6,374	380	254	220	161	131	83	57	44	24	25
1986	6,739	391	261	218	151	112	85	50	33	16	30
1985	6,491	375	281	253	175	126	86	53	28	19	16
1984	6,518	397	308	239	194	112	76	47	28	17	20
1983	6,567	395	314	227	199	128	69	44	25	7	22
1982	7,516	425	392	263	196	143	91	55	29	18	17
1981	8,053	519	435	305	216	134	81	50	22	18	17
1980	8,267	504	384	291	213	199	98	50	31	15	26
1979	7,657	513	390	302	208	143	110	51	28	17	21

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
White, male											
1998	4,217	248	179	111	79	48	41	21	7	6	14
1997	4,783	288	193	131	96	59	41	25	6	17	12
1996	4,929	284	189	153	88	68	34	25	11	4	6
1995	5,674	296	199	138	99	71	36	23	9	5	14
1994	6,231	327	233	154	107	65	47	31	11	6	14
1993	6,233	392	238	144	96	94	44	27	13	5	14
1992	6,409	349	212	166	130	63	43	34	22	12	13
1991	6,350	361	223	161	115	82	51	28	12	8	22
1990	5,894	321	225	138	120	80	39	36	13	12	20
1989	5,315	310	220	140	113	76	52	26	14	6	15
1988	5,014	292	195	175	111	63	48	24	17	12	22
1987	4,907	298	188	176	110	79	44	30	22	8	20
1986	5,295	296	196	170	101	70	52	23	18	8	24
1985	5,005	283	204	182	125	82	50	25	14	5	16
1984	5,128	306	246	183	140	77	50	25	14	10	19
1983	5,193	300	239	157	144	87	51	24	12	3	19
1982	5,955	349	294	206	141	95	61	37	15	12	12
1981	6,528	414	355	244	170	100	58	33	12	10	15
1980	6,729	390	306	238	159	153	70	32	19	7	20
1979	6,201	408	323	244	155	107	76	36	15	6	14
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
White, female											
1998	1,195	99	50	42	35	28	30	22	18	10	2
1997	1,242	102	58	44	23	31	29	25	15	18	1
1996	1,311	108	76	35	28	27	29	33	19	18	1
1995	1,470	90	74	51	28	30	37	28	24	12	3
1994	1,543	100	69	44	43	27	36	24	19	10	2
1993	1,685	112	68	54	40	38	41	29	15	15	2
1992	1,518	103	77	38	44	38	29	32	17	11	1
1991	1,594	126	63	56	48	43	35	34	11	10	6
1990	1,482	111	71	48	39	30	37	33	23	13	5
1989	1,403	94	56	47	38	41	33	25	19	7	6
1988	1,461	95	66	47	61	49	40	30	17	11	5
1987	1,467	82	66	44	51	52	39	27	22	16	5
1986	1,444	95	65	48	50	42	33	27	15	8	6
1985	1,486	92	77	71	50	44	36	28	14	14	-
1984	1,390	91	62	56	54	35	26	22	14	7	1
1983	1,374	95	75	70	55	41	18	20	13	4	3
1982	1,561	76	98	57	55	48	30	18	14	6	5
1981	1,525	105	80	61	46	34	23	17	10	8	2
1980	1,538	114	78	53	54	46	28	18	12	8	6
1979	1,456	105	67	58	53	36	34	15	13	11	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All other, both sexes											
1998	6,386	242	153	78	49	34	21	12	5	3	2
1997	7,227	298	129	81	62	36	28	10	7	4	4
1996	7,797	286	165	92	58	50	27	17	13	7	9
1995	8,407	289	169	113	82	62	33	16	6	6	7
1994	9,753	315	159	102	75	62	41	22	10	2	6
1993	10,335	314	178	116	77	62	40	24	12	6	10
1992	9,561	298	169	108	75	59	36	16	10	6	9
1991	9,802	310	174	112	90	67	38	18	11	6	13
1990	8,842	280	152	113	85	69	27	23	14	6	15
1989	7,746	241	165	109	103	74	41	27	10	9	10
1988	7,170	222	141	97	87	65	31	29	8	7	13
1987	6,283	228	163	120	82	63	33	21	15	7	5
1986	6,290	264	197	116	83	76	35	20	18	7	9
1985	5,345	253	156	128	100	73	36	26	13	7	5
1984	5,297	263	167	153	100	67	41	23	14	7	4
1983	5,473	271	199	175	105	60	39	28	10	5	3
1982	6,314	323	274	174	134	85	55	32	10	7	4
1981	7,036	382	277	236	153	88	50	35	10	5	8
1980	7,233	368	296	214	158	93	43	24	18	7	10
1979	6,834	410	306	237	146	95	53	26	11	7	7
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All other, male											
1998	5,554	196	129	66	41	26	14	8	3	2	1
1997	6,364	248	109	65	50	27	24	8	3	1	2
1996	6,806	234	145	81	46	37	20	13	10	5	8
1995	7,347	251	140	92	70	47	29	12	5	4	5
1994	8,535	264	128	85	59	50	33	15	6	1	5
1993	8,995	263	161	96	68	47	35	16	9	4	8
1992	8,338	255	148	91	59	45	31	12	6	6	7
1991	8,576	259	150	96	74	56	33	15	5	3	12
1990	7,735	238	127	91	74	53	25	19	5	4	14
1989	6,703	204	135	85	84	60	34	20	10	6	9
1988	6,120	196	108	78	69	53	21	26	7	4	11
1987	5,295	166	139	100	65	48	22	16	11	3	4
1986	5,361	230	168	102	69	64	31	17	12	5	8
1985	4,527	209	125	111	84	62	32	20	8	6	5
1984	4,487	228	146	132	80	53	35	20	10	5	3
1983	4,670	219	172	154	100	52	36	24	7	5	2
1982	5,447	272	245	149	119	75	44	24	9	6	3
1981	6,020	313	247	195	120	76	39	31	9	3	7
1980	6,149	313	251	184	130	76	34	21	15	6	7
1979	5,813	343	266	208	117	80	42	22	6	6	6

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
All other, female											
1998	832	46	24	12	8	8	7	4	2	1	1
1997	863	50	20	16	12	9	4	2	4	3	2
1996	991	52	20	11	12	13	7	4	3	2	1
1995	1,060	38	29	21	12	15	4	4	1	2	2
1994	1,218	51	31	17	16	12	8	7	4	1	1
1993	1,340	51	17	20	9	15	5	8	3	2	2
1992	1,223	43	21	17	16	14	5	4	4	-	2
1991	1,226	51	24	16	16	11	5	3	6	3	1
1990	1,107	42	25	22	11	16	2	4	9	2	1
1989	1,043	37	30	24	19	14	7	7	-	3	1
1988	1,050	26	33	19	18	12	10	3	1	3	2
1987	988	62	24	20	17	15	11	5	4	4	1
1986	929	34	29	14	14	12	4	3	6	2	1
1985	818	44	31	17	16	11	4	6	5	1	-
1984	810	35	21	21	20	14	6	3	4	2	1
1983	803	52	27	21	5	8	3	4	3	-	1
1982	867	51	29	25	15	10	11	8	1	1	1
1981	1,016	69	30	41	33	12	11	4	1	2	1
1980	1,084	55	45	30	28	17	9	3	3	1	3
1979	1,021	67	40	29	29	15	11	4	5	1	1
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
Black, both sexes											
1998	6,053	223	137	69	45	31	17	12	5	3	2
1997	6,841	265	113	70	56	31	25	8	7	4	4
1996	7,403	262	145	82	54	47	25	16	13	7	6
1995	7,935	268	150	98	76	60	31	15	6	6	6
1994	9,302	289	143	93	69	60	38	20	10	2	6
1993	9,824	280	166	102	68	59	40	23	12	6	10
1992	9,151	271	156	97	73	58	33	15	10	6	8
1991	9,327	284	153	97	81	66	37	16	11	6	12
1990	8,497	258	139	105	82	63	24	23	14	6	14
1989	7,414	230	155	100	98	67	37	27	10	9	9
1988	6,875	212	126	91	85	62	30	27	7	7	12
1987	6,054	220	150	116	78	62	31	20	14	7	4
1986	6,040	246	186	113	81	75	35	20	18	7	9
1985	5,142	238	153	121	95	70	34	26	13	7	4
1984	5,085	246	158	146	96	64	40	23	14	6	4
1983	5,245	250	185	172	98	57	38	27	10	5	2
1982	6,082	303	264	172	127	84	52	30	9	7	4
1981	6,816	367	264	228	151	85	47	35	10	5	7
1980	7,028	356	283	213	153	92	42	23	17	7	9
1979	6,633	395	292	228	139	91	52	26	11	7	7

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
Black, male											
1998	5,298	184	118	58	39	24	12	8	3	2	1
1997	6,057	225	99	58	46	23	21	7	3	1	2
1996	6,484	214	128	72	42	35	18	13	10	5	5
1995	6,978	233	125	82	64	46	27	11	5	4	5
1994	8,171	246	118	78	56	49	31	14	6	1	5
1993	8,578	242	151	84	61	45	35	15	9	4	8
1992	8,005	235	137	83	58	45	29	12	6	6	6
1991	8,198	239	135	83	68	55	33	15	5	3	11
1990	7,453	222	116	88	72	48	22	19	5	4	13
1989	6,450	195	126	79	80	56	32	20	10	6	9
1988	5,893	188	97	73	68	52	20	24	6	4	10
1987	5,112	160	128	98	62	47	21	15	11	3	3
1986	5,176	216	159	101	67	63	31	17	12	5	8
1985	4,363	197	123	106	81	59	31	20	8	6	4
1984	4,325	217	140	127	77	51	34	20	10	4	3
1983	4,491	203	160	151	93	49	35	24	7	5	1
1982	5,259	254	237	147	113	74	42	23	8	6	3
1981	5,849	300	237	189	118	73	37	31	9	3	6
1980	5,989	303	242	183	126	75	33	20	15	6	6
1979	5,649	328	254	200	111	76	41	22	6	6	6
Assault (homicide) by discharge of firearms (E965.0-E965.4)											
Black, female											
1998	755	39	19	11	6	7	5	4	2	1	1
1997	784	40	14	12	10	8	4	1	4	3	2
1996	919	48	17	10	12	12	7	3	3	2	1
1995	957	35	25	16	12	14	4	4	1	2	1
1994	1,131	43	25	15	13	11	7	6	4	1	1
1993	1,246	38	15	18	7	14	5	8	3	2	2
1992	1,146	36	19	14	15	13	4	3	4	-	2
1991	1,129	45	18	14	13	11	4	1	6	3	1
1990	1,044	36	23	17	10	15	2	4	9	2	1
1989	964	35	29	21	18	11	5	7	-	3	-
1988	982	24	29	18	17	10	10	3	1	3	2
1987	942	60	22	18	16	15	10	5	3	4	1
1986	864	30	27	12	14	12	4	3	6	2	1
1985	779	41	30	15	14	11	3	6	5	1	-
1984	760	29	18	19	19	13	6	3	4	2	1
1983	754	47	25	21	5	8	3	3	3	-	1
1982	823	49	27	25	14	10	10	7	1	1	1
1981	967	67	27	39	33	12	10	4	1	2	1
1980	1,039	53	41	30	27	17	9	3	2	1	3
1979	984	67	38	28	28	15	11	4	5	1	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All races, both sexes											
1998	6,095	444	280	218	145	141	121	128	105	80	37
1997	6,239	473	288	221	176	160	155	143	101	110	30
1996	6,597	457	296	213	195	171	141	120	94	81	46
1995	7,001	471	301	247	184	179	155	138	98	95	28
1994	7,020	397	290	200	173	192	161	145	125	102	30
1993	7,400	423	297	226	216	178	189	127	130	113	41
1992	7,656	466	328	211	211	218	203	148	122	105	31
1991	8,508	470	366	278	284	236	184	189	135	106	40
1990	8,396	406	301	252	252	205	194	157	128	107	56
1989	8,114	426	337	259	262	230	191	162	123	108	50
1988	8,139	401	291	298	242	251	212	182	126	108	50
1987	8,155	417	345	346	263	233	217	200	111	116	69
1986	8,433	399	359	343	270	243	199	171	123	107	48
1985	7,792	411	335	319	285	230	175	153	106	84	41
1984	7,695	397	359	307	289	189	199	149	120	89	34
1983	7,882	388	356	351	287	222	181	179	118	117	38
1982	8,243	441	414	357	295	227	187	153	106	116	42
1981	8,272	476	437	370	271	222	186	179	124	103	33
1980	8,467	482	467	364	278	242	212	170	111	96	39
1979	7,711	453	389	335	276	256	169	125	119	85	49
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All races, male											
1998	3,881	304	208	163	110	98	73	82	46	29	26
1997	3,960	322	202	154	116	108	88	80	52	48	17
1996	4,210	298	221	145	141	111	86	61	48	31	27
1995	4,387	328	207	180	132	117	81	67	43	33	17
1994	4,576	289	218	145	129	128	89	81	61	47	16
1993	4,717	298	215	166	157	105	103	69	47	45	24
1992	5,032	343	233	157	156	146	116	77	60	47	18
1991	5,592	343	259	206	201	148	122	97	63	38	22
1990	5,669	312	237	184	187	136	122	80	59	40	36
1989	5,348	321	250	195	185	137	111	90	52	53	29
1988	5,338	304	218	209	187	153	129	100	53	43	32
1987	5,369	313	253	256	193	161	128	107	51	49	42
1986	5,676	309	268	267	187	158	109	94	53	37	33
1985	5,274	304	263	245	198	146	104	80	42	27	26
1984	5,144	301	285	225	214	124	125	74	55	40	14
1983	5,390	289	279	263	218	138	110	83	58	47	26
1982	5,633	339	321	270	209	144	108	79	44	48	27
1981	5,740	370	342	272	193	145	121	80	53	56	24
1980	5,901	396	367	283	205	166	123	89	56	38	24
1979	5,274	330	310	256	195	157	81	66	52	33	34

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All races, female											
1998	2,214	140	72	55	35	43	48	46	59	51	11
1997	2,279	151	86	67	60	52	67	63	49	62	13
1996	2,387	159	75	68	54	60	55	59	46	50	19
1995	2,614	143	94	67	52	62	74	71	55	62	11
1994	2,444	108	72	55	44	64	72	64	64	55	14
1993	2,683	125	82	60	59	73	86	58	83	68	17
1992	2,624	123	95	54	55	72	87	71	62	58	13
1991	2,916	127	107	72	83	88	62	92	72	68	18
1990	2,727	94	64	68	65	69	72	77	69	67	20
1989	2,766	105	87	64	77	93	80	72	71	55	21
1988	2,801	97	73	89	55	98	83	82	73	65	18
1987	2,786	104	92	90	70	72	89	93	60	67	27
1986	2,757	90	91	76	83	85	90	77	70	70	15
1985	2,518	107	72	74	87	84	71	73	64	57	15
1984	2,551	96	74	82	75	65	74	75	65	49	20
1983	2,492	99	77	88	69	84	71	96	60	70	12
1982	2,610	102	93	87	86	83	79	74	62	68	15
1981	2,532	106	95	98	78	77	65	99	71	47	9
1980	2,566	86	100	81	73	76	89	81	55	58	15
1979	2,437	123	79	79	81	99	88	59	67	52	15
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
White, both sexes											
1998	3,596	249	184	144	88	100	85	88	86	65	31
1997	3,651	278	181	152	105	105	119	103	82	84	21
1996	3,848	264	191	135	132	130	101	83	71	61	35
1995	3,986	282	194	153	123	120	106	83	72	62	22
1994	3,944	232	189	133	115	132	100	101	91	72	20
1993	4,151	235	195	157	134	111	127	81	91	78	29
1992	4,306	275	214	142	136	127	139	100	79	75	25
1991	4,687	275	226	166	180	150	111	118	84	78	33
1990	4,577	248	172	157	159	139	131	98	86	63	44
1989	4,374	263	202	150	171	143	121	101	89	66	35
1988	4,427	219	163	183	161	157	136	132	77	71	37
1987	4,576	243	207	212	169	150	145	135	81	86	53
1986	4,789	235	209	205	174	155	130	119	96	75	42
1985	4,512	249	191	207	180	163	109	123	84	64	35
1984	4,434	231	212	195	182	117	128	94	96	64	24
1983	4,508	224	215	232	171	141	125	123	84	94	27
1982	4,756	245	259	232	193	141	130	114	83	97	28
1981	4,851	266	257	241	188	143	131	133	98	84	25
1980	5,093	271	286	244	182	162	156	137	92	83	28
1979	4,492	266	218	202	168	181	127	90	95	78	35

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
White, male											
1998	2,262	171	127	102	68	66	51	54	41	22	21
1997	2,335	189	128	104	68	71	63	56	41	35	13
1996	2,422	167	134	88	91	82	61	42	37	23	24
1995	2,435	177	126	107	86	77	47	39	29	26	14
1994	2,579	165	144	98	84	87	59	54	44	31	10
1993	2,612	155	139	113	97	62	65	45	29	29	16
1992	2,819	194	152	106	103	84	71	49	40	29	16
1991	3,087	194	160	123	127	91	67	56	39	27	18
1990	3,060	184	130	114	110	93	81	45	37	25	26
1989	2,812	196	141	118	123	84	66	52	37	33	21
1988	2,823	160	117	128	122	96	82	65	32	25	26
1987	2,897	179	146	162	119	98	74	65	36	31	31
1986	3,114	175	152	161	116	99	70	62	36	23	29
1985	2,958	172	150	159	113	103	64	63	29	22	23
1984	2,872	169	164	134	127	69	76	42	41	26	10
1983	3,005	156	162	174	121	88	73	55	35	35	18
1982	3,142	183	203	168	138	83	72	57	33	42	19
1981	3,251	195	190	174	134	86	76	52	39	41	18
1980	3,456	218	224	181	133	111	83	69	44	33	18
1979	3,009	193	165	148	117	111	63	44	43	31	24
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
White, female											
1998	1,334	78	57	42	20	34	34	34	45	43	10
1997	1,316	89	53	48	37	34	56	47	41	49	8
1996	1,426	97	57	47	41	48	40	41	34	38	11
1995	1,551	105	68	46	37	43	59	44	43	36	8
1994	1,365	67	45	35	31	45	41	47	47	41	10
1993	1,539	80	56	44	37	49	62	36	62	49	13
1992	1,487	81	62	36	33	43	68	51	39	46	9
1991	1,600	81	66	43	53	59	44	62	45	51	15
1990	1,517	64	42	43	49	46	50	53	49	38	18
1989	1,562	67	61	32	48	59	55	49	52	33	14
1988	1,604	59	46	55	39	61	54	67	45	46	11
1987	1,679	64	61	50	50	52	71	70	45	55	22
1986	1,675	60	57	44	58	56	60	57	60	52	13
1985	1,554	77	41	48	67	60	45	60	55	42	12
1984	1,562	62	48	61	55	48	52	52	55	38	14
1983	1,503	68	53	58	50	53	52	68	49	59	9
1982	1,614	62	56	64	55	58	58	57	50	55	9
1981	1,600	71	67	67	54	57	55	81	59	43	7
1980	1,637	53	62	63	49	51	73	68	48	50	10
1979	1,483	73	53	54	51	70	64	46	52	47	11

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All other, both sexes											
1998	2,499	195	96	74	57	41	36	40	19	15	6
1997	2,588	195	107	69	71	55	36	40	19	26	9
1996	2,749	193	105	78	63	41	40	37	23	20	11
1995	3,015	189	107	94	61	59	49	55	26	33	6
1994	3,076	165	101	67	58	60	61	44	34	30	10
1993	3,249	188	102	69	82	67	62	46	39	35	12
1992	3,350	191	114	69	75	91	64	48	43	30	6
1991	3,821	195	140	112	104	86	73	71	51	28	7
1990	3,819	158	129	95	93	66	63	59	42	44	12
1989	3,740	163	135	109	91	87	70	61	34	42	15
1988	3,712	182	128	115	81	94	76	50	49	37	13
1987	3,579	174	138	134	94	83	72	65	30	30	16
1986	3,644	164	150	138	96	88	69	52	27	32	6
1985	3,280	162	144	112	105	67	66	30	22	20	6
1984	3,261	166	147	112	107	72	71	55	24	25	10
1983	3,374	164	141	119	116	81	56	56	34	23	11
1982	3,487	196	155	125	102	86	57	39	23	19	14
1981	3,421	210	180	129	83	79	55	46	26	19	8
1980	3,374	211	181	120	96	80	56	33	19	13	11
1979	3,219	187	171	133	108	75	42	35	24	7	14
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All other, male											
1998	1,619	133	81	61	42	32	22	28	5	7	5
1997	1,625	133	74	50	48	37	25	24	11	13	4
1996	1,788	131	87	57	50	29	25	19	11	8	3
1995	1,952	151	81	73	46	40	34	28	14	7	3
1994	1,997	124	74	47	45	41	30	27	17	16	6
1993	2,105	143	76	53	60	43	38	24	18	16	8
1992	2,213	149	81	51	53	62	45	28	20	18	2
1991	2,505	149	99	83	74	57	55	41	24	11	4
1990	2,609	128	107	70	77	43	41	35	22	15	10
1989	2,536	125	109	77	62	53	45	38	15	20	8
1988	2,515	144	101	81	65	57	47	35	21	18	6
1987	2,472	134	107	94	74	63	54	42	15	18	11
1986	2,562	134	116	106	71	59	39	32	17	14	4
1985	2,316	132	113	86	85	43	40	17	13	5	3
1984	2,272	132	121	91	87	55	49	32	14	14	4
1983	2,385	133	117	89	97	50	37	28	23	12	8
1982	2,491	156	118	102	71	61	36	22	11	6	8
1981	2,489	175	152	98	59	59	45	28	14	15	6
1980	2,445	178	143	102	72	55	40	20	12	5	6
1979	2,265	137	145	108	78	46	18	22	9	2	10

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
All other, female											
1998	880	62	15	13	15	9	14	12	14	8	1
1997	963	62	33	19	23	18	11	16	8	13	5
1996	961	62	18	21	13	12	15	18	12	12	8
1995	1,063	38	26	21	15	19	15	27	12	26	3
1994	1,079	41	27	20	13	19	31	17	17	14	4
1993	1,144	45	26	16	22	24	24	22	21	19	4
1992	1,137	42	33	18	22	29	19	20	23	12	4
1991	1,316	46	41	29	30	29	18	30	27	17	3
1990	1,210	30	22	25	16	23	22	24	20	29	2
1989	1,204	38	26	32	29	34	25	23	19	22	7
1988	1,197	38	27	34	16	37	29	15	28	19	7
1987	1,107	40	31	40	20	20	18	23	15	12	5
1986	1,082	30	34	32	25	29	30	20	10	18	2
1985	964	30	31	26	20	24	26	13	9	15	3
1984	989	34	26	21	20	17	22	23	10	11	6
1983	989	31	24	30	19	31	19	28	11	11	3
1982	996	40	37	23	31	25	21	17	12	13	6
1981	932	35	28	31	24	20	10	18	12	4	2
1980	929	33	38	18	24	25	16	13	7	8	5
1979	954	50	26	25	30	29	24	13	15	5	4
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
Black, both sexes											
1998	2,240	172	87	66	46	33	32	34	17	15	6
1997	2,306	174	91	61	64	48	33	38	18	24	8
1996	2,483	175	88	71	57	34	37	32	23	20	11
1995	2,754	175	94	83	51	54	45	52	23	32	6
1994	2,790	147	96	62	50	55	58	43	32	29	6
1993	2,987	175	90	58	79	63	60	45	38	35	8
1992	3,069	171	110	62	69	82	57	47	41	26	5
1991	3,532	184	123	99	98	83	69	70	50	27	6
1990	3,538	145	118	84	87	62	60	57	41	41	12
1989	3,444	149	123	101	86	82	66	56	31	41	14
1988	3,452	171	113	106	76	88	73	49	45	35	13
1987	3,329	156	127	124	86	79	69	63	28	29	13
1986	3,355	151	138	126	89	84	67	46	25	29	6
1985	3,047	147	128	107	102	64	63	27	21	20	6
1984	3,050	152	140	104	104	71	64	52	22	22	9
1983	3,142	155	130	109	112	77	52	53	33	23	9
1982	3,275	185	143	118	97	82	53	35	21	18	13
1981	3,203	194	170	124	77	76	49	44	25	16	6
1980	3,144	199	171	111	90	73	50	30	17	11	9
1979	3,023	179	166	124	103	73	36	32	23	7	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
Black, male											
1998	1,451	119	73	54	34	27	20	22	5	7	5
1997	1,439	120	65	44	41	33	22	24	11	12	4
1996	1,605	121	75	52	44	26	24	17	11	8	3
1995	1,779	140	73	64	39	38	34	27	12	6	3
1994	1,798	109	70	45	39	38	29	26	16	15	3
1993	1,938	133	68	47	58	41	37	24	17	16	5
1992	2,029	136	78	48	49	55	43	28	19	16	2
1991	2,333	142	90	76	71	56	52	41	23	11	3
1990	2,423	118	101	62	72	40	39	34	22	13	10
1989	2,339	115	101	71	60	49	41	34	14	19	7
1988	2,346	140	88	74	63	55	46	34	19	18	6
1987	2,306	121	99	87	68	61	52	40	13	17	9
1986	2,360	123	108	97	67	56	37	28	15	12	4
1985	2,164	123	104	82	84	41	39	15	12	5	3
1984	2,135	124	117	86	84	54	43	29	12	14	4
1983	2,229	126	109	82	94	47	35	25	22	12	6
1982	2,356	149	110	98	68	57	33	19	10	5	7
1981	2,346	163	145	95	56	56	39	26	13	13	4
1980	2,285	170	137	97	67	48	34	17	11	3	4
1979	2,138	132	142	101	74	45	16	20	9	2	9
Assault (homicide) by other and unspecified means and their sequelae (E960-E964,E965.5-E969)											
Black, female											
1998	789	53	14	12	12	6	12	12	12	8	1
1997	867	54	26	17	23	15	11	14	7	12	4
1996	878	54	13	19	13	8	13	15	12	12	8
1995	975	35	21	19	12	16	11	25	11	26	3
1994	992	38	26	17	11	17	29	17	16	14	3
1993	1,049	42	22	11	21	22	23	21	21	19	3
1992	1,040	35	32	14	20	27	14	19	22	10	3
1991	1,199	42	33	23	27	27	17	29	27	16	3
1990	1,115	27	17	22	15	22	21	23	19	28	2
1989	1,105	34	22	30	26	33	25	22	17	22	7
1988	1,106	31	25	32	13	33	27	15	26	17	7
1987	1,023	35	28	37	18	18	17	23	15	12	4
1986	995	28	30	29	22	28	30	18	10	17	2
1985	883	24	24	25	18	23	24	12	9	15	3
1984	915	28	23	18	20	17	21	23	10	8	5
1983	913	29	21	27	18	30	17	28	11	11	3
1982	919	36	33	20	29	25	20	16	11	13	6
1981	857	31	25	29	21	20	10	18	12	3	2
1980	859	29	34	14	23	25	16	13	6	8	5
1979	885	47	24	23	29	28	20	12	14	5	3

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
All races, both sexes											
1998	379	33	16	5	4	-	4	-	-	2	1
1997	355	28	12	14	5	4	2	2	1	-	1
1996	337	26	9	7	6	5	2	2	1	2	1
1995	343	25	13	5	7	4	3	-	1	2	-
1994	379	23	16	2	4	2	3	3	-	2	-
1993	356	20	10	6	2	2	3	1	1	-	1
1992	344	30	14	5	4	1	2	3	-	1	-
1991	259	16	3	5	7	1	1	-	-	-	-
1990	318	17	8	2	6	1	-	1	2	1	2
1989	331	21	15	9	5	1	1	1	2	-	2
1988	248	12	12	7	5	1	-	-	-	1	1
1987	291	13	3	4	8	2	1	1	-	-	2
1986	269	16	8	10	4	6	2	-	2	-	1
1985	265	12	5	3	9	2	1	-	3	1	1
1984	286	13	12	1	5	9	2	1	1	-	-
1983	269	11	14	4	6	2	1	4	2	-	-
1982	285	13	3	5	4	4	2	1	-	-	1
1981	285	15	7	7	3	1	3	1	-	-	-
1980	311	11	18	6	3	4	4	-	1	-	1
1979	348	20	7	7	1	5	4	3	1	1	-
Legal intervention (E970-E978)											
All races, male											
1998	371	33	14	4	4	-	4	-	-	2	1
1997	342	28	11	12	5	4	2	2	1	-	1
1996	324	25	9	7	5	5	1	2	1	-	1
1995	332	24	13	5	6	4	1	-	-	1	-
1994	365	23	14	2	4	2	3	2	-	1	-
1993	345	20	9	5	2	2	3	1	1	-	1
1992	336	30	13	5	3	1	2	3	-	1	-
1991	250	16	3	5	6	1	1	-	-	-	-
1990	306	16	8	2	6	1	-	1	1	1	2
1989	320	20	15	9	5	1	1	1	1	-	2
1988	240	12	12	7	5	1	-	-	-	1	1
1987	284	13	3	4	8	2	1	1	-	-	2
1986	263	15	8	9	3	6	1	-	2	-	1
1985	260	12	5	3	8	2	1	-	3	-	1
1984	279	13	11	1	5	8	2	1	-	-	-
1983	261	10	14	4	5	2	1	4	2	-	-
1982	280	13	3	5	4	4	2	1	-	-	1
1981	284	15	7	7	3	1	3	1	-	-	-
1980	309	11	18	6	3	4	4	-	1	-	1
1979	340	19	7	6	1	5	4	2	1	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
All races, female											
1998	8	-	2	1	-	-	-	-	-	-	-
1997	13	-	1	2	-	-	-	-	-	-	-
1996	13	1	-	-	1	-	1	-	-	2	-
1995	11	1	-	-	1	-	2	-	1	1	-
1994	14	-	2	-	-	-	-	1	-	1	-
1993	11	-	1	1	-	-	-	-	-	-	-
1992	8	-	1	-	1	-	-	-	-	-	-
1991	9	-	-	-	1	-	-	-	-	-	-
1990	12	1	-	-	-	-	-	-	1	-	-
1989	11	1	-	-	-	-	-	-	1	-	-
1988	8	-	-	-	-	-	-	-	-	-	-
1987	7	-	-	-	-	-	-	-	-	-	-
1986	6	1	-	1	1	-	1	-	-	-	-
1985	5	-	-	-	1	-	-	-	-	1	-
1984	7	-	1	-	-	1	-	-	1	-	-
1983	8	1	-	-	1	-	-	-	-	-	-
1982	5	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	-	-	-	-	-
1979	8	1	-	1	-	-	-	1	-	-	-
Legal intervention (E970-E978)											
White, both sexes											
1998	233	26	9	3	2	-	1	-	-	2	1
1997	237	21	11	12	3	4	2	1	1	-	-
1996	229	19	6	7	4	4	1	2	-	2	1
1995	234	20	12	4	5	4	2	-	1	2	-
1994	258	16	14	2	3	2	1	1	-	2	-
1993	217	17	5	5	2	1	1	1	-	-	1
1992	235	23	10	5	4	1	2	2	-	1	-
1991	151	12	2	5	5	1	-	-	-	-	-
1990	200	15	6	1	2	1	-	1	-	-	1
1989	216	18	14	6	4	1	1	-	2	-	2
1988	164	8	10	4	4	1	-	-	-	1	1
1987	178	8	1	4	8	2	1	1	-	-	2
1986	162	11	4	7	1	3	1	-	1	-	1
1985	160	9	4	2	7	1	1	-	3	1	1
1984	175	8	8	1	3	6	2	1	1	-	-
1983	160	8	8	2	3	2	-	2	2	-	-
1982	167	8	2	3	3	2	2	-	-	-	1
1981	162	10	3	6	1	1	2	-	-	-	-
1980	198	9	14	5	2	3	3	-	1	-	1
1979	185	11	4	5	1	1	2	-	1	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
White, male											
1998	228	26	8	2	2	-	1	-	-	2	1
1997	225	21	10	10	3	4	2	1	1	-	-
1996	219	18	6	7	3	4	1	2	-	-	1
1995	227	19	12	4	4	4	1	-	-	1	-
1994	245	16	13	2	3	2	1	-	-	1	-
1993	209	17	4	4	2	1	1	1	-	-	1
1992	228	23	9	5	3	1	2	2	-	1	-
1991	144	12	2	5	4	1	-	-	-	-	-
1990	193	14	6	1	2	1	-	1	-	-	1
1989	210	18	14	6	4	1	1	-	1	-	2
1988	157	8	10	4	4	1	-	-	-	1	1
1987	175	8	1	4	8	2	1	1	-	-	2
1986	158	10	4	6	1	3	1	-	1	-	1
1985	159	9	4	2	7	1	1	-	3	-	1
1984	171	8	7	1	3	6	2	1	-	-	-
1983	157	7	8	2	2	2	-	2	2	-	-
1982	163	8	2	3	3	2	2	-	-	-	1
1981	162	10	3	6	1	1	2	-	-	-	-
1980	196	9	14	5	2	3	3	-	1	-	1
1979	182	10	4	4	1	1	2	-	1	-	-
Legal intervention (E970-E978)											
White, female											
1998	5	-	1	1	-	-	-	-	-	-	-
1997	12	-	1	2	-	-	-	-	-	-	-
1996	10	1	-	-	1	-	-	-	-	2	-
1995	7	1	-	-	1	-	1	-	1	1	-
1994	13	-	1	-	-	-	-	1	-	1	-
1993	8	-	1	1	-	-	-	-	-	-	-
1992	7	-	1	-	1	-	-	-	-	-	-
1991	7	-	-	-	1	-	-	-	-	-	-
1990	7	1	-	-	-	-	-	-	-	-	-
1989	6	-	-	-	-	-	-	-	1	-	-
1988	7	-	-	-	-	-	-	-	-	-	-
1987	3	-	-	-	-	-	-	-	-	-	-
1986	4	1	-	1	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	-	-	1	-
1984	4	-	1	-	-	-	-	-	1	-	-
1983	3	1	-	-	1	-	-	-	-	-	-
1982	4	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	-	-	-	-	-
1979	3	1	-	1	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
All other, both sexes											
1998	146	7	7	2	2	-	3	-	-	-	-
1997	118	7	1	2	2	-	-	1	-	-	1
1996	108	7	3	-	2	1	1	-	1	-	-
1995	109	5	1	1	2	-	1	-	-	-	-
1994	121	7	2	-	1	-	2	2	-	-	-
1993	139	3	5	1	-	1	2	-	1	-	-
1992	109	7	4	-	-	-	-	1	-	-	-
1991	108	4	1	-	2	-	1	-	-	-	-
1990	118	2	2	1	4	-	-	-	2	1	1
1989	115	3	1	3	1	-	-	1	-	-	-
1988	84	4	2	3	1	-	-	-	-	-	-
1987	113	5	2	-	-	-	-	-	-	-	-
1986	107	5	4	3	3	3	1	-	1	-	-
1985	105	3	1	1	2	1	-	-	-	-	-
1984	111	5	4	-	2	3	-	-	-	-	-
1983	109	3	6	2	3	-	1	2	-	-	-
1982	118	5	1	2	1	2	-	1	-	-	-
1981	123	5	4	1	2	-	1	1	-	-	-
1980	113	2	4	1	1	1	1	-	-	-	-
1979	163	9	3	2	-	4	2	3	-	1	-
Legal intervention (E970-E978)											
All other, male											
1998	143	7	6	2	2	-	3	-	-	-	-
1997	117	7	1	2	2	-	-	1	-	-	1
1996	105	7	3	-	2	1	-	-	1	-	-
1995	105	5	1	1	2	-	-	-	-	-	-
1994	120	7	1	-	1	-	2	2	-	-	-
1993	136	3	5	1	-	1	2	-	1	-	-
1992	108	7	4	-	-	-	-	1	-	-	-
1991	106	4	1	-	2	-	1	-	-	-	-
1990	113	2	2	1	4	-	-	-	1	1	1
1989	110	2	1	3	1	-	-	1	-	-	-
1988	83	4	2	3	1	-	-	-	-	-	-
1987	109	5	2	-	-	-	-	-	-	-	-
1986	105	5	4	3	2	3	-	-	1	-	-
1985	101	3	1	1	1	1	-	-	-	-	-
1984	108	5	4	-	2	2	-	-	-	-	-
1983	104	3	6	2	3	-	1	2	-	-	-
1982	117	5	1	2	1	2	-	1	-	-	-
1981	122	5	4	1	2	-	1	1	-	-	-
1980	113	2	4	1	1	1	1	-	-	-	-
1979	158	9	3	2	-	4	2	2	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
All other, female											
1998	3	-	1	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	3	-	-	-	-	-	1	-	-	-	-
1995	4	-	-	-	-	-	1	-	-	-	-
1994	1	-	1	-	-	-	-	-	-	-	-
1993	3	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	2	-	-	-	-	-	-	-	-	-	-
1990	5	-	-	-	-	-	-	-	1	-	-
1989	5	1	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	1	-	1	-	-	-	-
1985	4	-	-	-	1	-	-	-	-	-	-
1984	3	-	-	-	-	1	-	-	-	-	-
1983	5	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	5	-	-	-	-	-	-	1	-	-	-
Legal intervention (E970-E978)											
Black, both sexes											
1998	127	6	6	1	2	-	3	-	-	-	-
1997	106	7	-	1	2	-	-	1	-	-	1
1996	97	5	3	-	2	-	1	-	1	-	-
1995	94	4	1	1	1	-	1	-	-	-	-
1994	115	7	2	-	1	-	2	2	-	-	-
1993	126	3	5	1	-	1	2	-	1	-	-
1992	98	5	4	-	-	-	-	1	-	-	-
1991	99	3	1	-	2	-	1	-	-	-	-
1990	109	1	2	1	4	-	-	-	2	1	1
1989	104	3	1	3	1	-	-	1	-	-	-
1988	76	3	2	3	1	-	-	-	-	-	-
1987	104	4	2	-	-	-	-	-	-	-	-
1986	100	5	4	3	3	3	1	-	1	-	-
1985	93	2	1	1	2	1	-	-	-	-	-
1984	105	5	4	-	2	2	-	-	-	-	-
1983	107	3	6	2	3	-	1	2	-	-	-
1982	116	5	1	2	1	2	-	1	-	-	-
1981	118	4	4	1	2	-	1	1	-	-	-
1980	111	2	4	1	1	1	1	-	-	-	-
1979	156	9	3	2	-	4	2	3	-	1	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Legal intervention (E970-E978)											
Black, male											
1998	124	6	5	1	2	-	3	-	-	-	-
1997	105	7	-	1	2	-	-	1	-	-	1
1996	94	5	3	-	2	-	-	-	1	-	-
1995	90	4	1	1	1	-	-	-	-	-	-
1994	114	7	1	-	1	-	2	2	-	-	-
1993	124	3	5	1	-	1	2	-	1	-	-
1992	97	5	4	-	-	-	-	1	-	-	-
1991	97	3	1	-	2	-	1	-	-	-	-
1990	105	1	2	1	4	-	-	-	1	1	1
1989	99	2	1	3	1	-	-	1	-	-	-
1988	75	3	2	3	1	-	-	-	-	-	-
1987	100	4	2	-	-	-	-	-	-	-	-
1986	98	5	4	3	2	3	-	-	1	-	-
1985	89	2	1	1	1	1	-	-	-	-	-
1984	103	5	4	-	2	2	-	-	-	-	-
1983	102	3	6	2	3	-	1	2	-	-	-
1982	115	5	1	2	1	2	-	1	-	-	-
1981	117	4	4	1	2	-	1	1	-	-	-
1980	111	2	4	1	1	1	1	-	-	-	-
1979	151	9	3	2	-	4	2	2	-	1	-
Legal intervention (E970-E978)											
Black, female											
1998	3	-	1	-	-	-	-	-	-	-	-
1997	1	-	-	-	-	-	-	-	-	-	-
1996	3	-	-	-	-	-	1	-	-	-	-
1995	4	-	-	-	-	-	1	-	-	-	-
1994	1	-	1	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	-	-	-	-	-	-
1991	2	-	-	-	-	-	-	-	-	-	-
1990	4	-	-	-	-	-	-	-	1	-	-
1989	5	1	-	-	-	-	-	-	-	-	-
1988	1	-	-	-	-	-	-	-	-	-	-
1987	4	-	-	-	-	-	-	-	-	-	-
1986	2	-	-	-	1	-	1	-	-	-	-
1985	4	-	-	-	1	-	-	-	-	-	-
1984	2	-	-	-	-	-	-	-	-	-	-
1983	5	-	-	-	-	-	-	-	-	-	-
1982	1	-	-	-	-	-	-	-	-	-	-
1981	1	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	5	-	-	-	-	-	-	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
All races, both sexes											
1998	3,746	467	241	137	77	62	41	40	46	64	25
1997	3,657	417	215	139	85	62	58	50	38	56	19
1996	3,463	323	175	97	83	92	48	57	44	64	27
1995	3,301	317	148	94	54	58	44	45	38	56	25
1994	3,419	298	162	122	83	59	60	54	41	53	19
1993	3,419	270	160	114	88	84	53	53	40	51	26
1992	2,895	199	125	84	78	79	43	42	46	38	29
1991	2,507	157	117	87	88	70	51	51	46	40	33
1990	2,376	156	106	84	89	71	52	48	35	43	18
1989	2,686	164	112	118	72	83	59	53	43	36	20
1988	3,018	164	123	103	102	94	96	55	48	60	24
1987	3,011	155	109	106	131	119	59	67	48	62	31
1986	3,108	173	168	155	131	112	69	71	55	52	28
1985	2,826	195	126	134	118	90	71	68	55	48	22
1984	2,996	164	167	154	130	121	95	66	50	58	17
1983	3,104	164	194	189	120	110	82	77	62	66	19
1982	3,201	188	198	183	147	93	93	82	43	53	31
1981	3,614	217	223	231	179	149	115	74	75	70	16
1980	3,663	226	213	205	164	144	97	94	60	59	25
1979	3,860	258	270	199	209	126	116	78	56	57	18
Events of undetermined intent (E980-E989)											
All races, male											
1998	2,597	314	169	78	52	35	21	21	30	25	18
1997	2,656	313	143	89	56	37	37	32	21	25	15
1996	2,523	245	120	52	53	63	32	33	24	29	24
1995	2,461	235	103	56	35	37	26	26	29	29	20
1994	2,508	229	112	80	48	33	40	36	23	26	16
1993	2,513	195	105	75	63	58	34	35	26	22	19
1992	2,105	134	82	58	47	46	29	25	25	17	23
1991	1,755	101	76	59	56	44	28	33	23	21	24
1990	1,719	105	62	54	63	47	32	29	21	15	16
1989	1,942	111	75	83	46	48	37	30	25	22	20
1988	2,193	124	85	68	69	58	55	36	30	27	20
1987	2,186	109	75	73	86	88	39	47	33	33	22
1986	2,258	127	116	114	85	69	48	49	34	28	22
1985	2,019	121	86	97	80	53	40	45	32	31	17
1984	2,151	106	115	105	94	90	61	37	32	26	15
1983	2,180	110	135	129	76	76	57	51	29	31	15
1982	2,298	129	127	126	109	60	63	48	26	23	27
1981	2,533	142	152	145	112	103	75	51	45	43	14
1980	2,620	153	132	129	124	95	63	59	37	34	22
1979	2,671	169	172	130	143	87	78	50	31	34	14

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
All races, female											
1998	1,149	153	72	59	25	27	20	19	16	39	7
1997	1,001	104	72	50	29	25	21	18	17	31	4
1996	940	78	55	45	30	29	16	24	20	35	3
1995	840	82	45	38	19	21	18	19	9	27	5
1994	911	69	50	42	35	26	20	18	18	27	3
1993	906	75	55	39	25	26	19	18	14	29	7
1992	790	65	43	26	31	33	14	17	21	21	6
1991	752	56	41	28	32	26	23	18	23	19	9
1990	657	51	44	30	26	24	20	19	14	28	2
1989	744	53	37	35	26	35	22	23	18	14	-
1988	825	40	38	35	33	36	41	19	18	33	4
1987	825	46	34	33	45	31	20	20	15	29	9
1986	850	46	52	41	46	43	21	22	21	24	6
1985	807	74	40	37	38	37	31	23	23	17	5
1984	845	58	52	49	36	31	34	29	18	32	2
1983	924	54	59	60	44	34	25	26	33	35	4
1982	903	59	71	57	38	33	30	34	17	30	4
1981	1,081	75	71	86	67	46	40	23	30	27	2
1980	1,043	73	81	76	40	49	34	35	23	25	3
1979	1,189	89	98	69	66	39	38	28	25	23	4
Events of undetermined intent (E980-E989)											
White, both sexes											
1998	2,952	368	190	114	54	55	36	36	42	61	14
1997	2,839	325	177	102	69	54	51	43	31	49	11
1996	2,680	258	141	74	69	76	42	50	39	54	21
1995	2,475	219	112	71	49	48	34	40	34	45	14
1994	2,620	221	118	99	65	50	55	46	35	47	13
1993	2,575	211	127	93	73	66	47	41	36	45	20
1992	2,229	165	96	68	69	65	37	36	40	32	21
1991	1,942	130	95	73	68	57	44	44	43	33	22
1990	1,808	120	88	62	74	60	43	43	29	35	8
1989	1,961	123	78	87	63	68	52	39	39	34	14
1988	2,237	126	103	77	82	73	86	49	41	54	17
1987	2,185	120	80	86	100	95	48	60	41	53	22
1986	2,315	130	124	111	105	91	60	64	45	41	18
1985	2,148	152	99	109	96	75	56	56	46	41	16
1984	2,254	130	124	118	101	105	80	57	46	46	17
1983	2,316	123	141	152	88	93	68	66	53	55	16
1982	2,382	139	156	154	121	72	75	71	36	44	23
1981	2,727	159	176	192	141	129	92	59	61	59	11
1980	2,785	160	174	165	131	117	82	81	54	51	22
1979	2,902	194	213	164	174	102	100	67	49	45	13

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
White, male											
1998	1,988	241	132	58	35	30	18	20	26	22	11
1997	2,039	242	116	61	43	30	31	27	18	23	9
1996	1,924	190	90	36	42	52	28	31	22	24	19
1995	1,833	156	75	40	30	28	19	22	27	22	11
1994	1,891	164	74	63	34	26	37	29	19	21	10
1993	1,876	150	80	58	51	43	30	26	24	17	15
1992	1,593	107	57	43	39	38	25	20	22	12	16
1991	1,357	82	61	48	42	34	23	27	21	17	17
1990	1,303	80	46	39	52	38	27	26	19	14	8
1989	1,402	79	52	57	38	36	31	24	24	22	14
1988	1,607	91	68	47	55	44	50	31	27	24	16
1987	1,557	79	53	56	61	66	31	41	27	28	15
1986	1,672	93	81	79	66	51	42	45	27	23	16
1985	1,532	91	66	77	65	41	32	35	26	27	13
1984	1,608	81	84	78	72	78	51	34	28	20	15
1983	1,623	82	93	102	51	65	48	45	24	25	14
1982	1,687	90	93	106	89	43	51	40	21	19	21
1981	1,893	103	116	114	88	89	59	42	36	36	9
1980	1,966	103	102	95	92	73	51	47	33	26	20
1979	2,001	125	132	102	112	68	65	44	27	25	9
Events of undetermined intent (E980-E989)											
White, female											
1998	964	127	58	56	19	25	18	16	16	39	3
1997	800	83	61	41	26	24	20	16	13	26	2
1996	756	68	51	38	27	24	14	19	17	30	2
1995	642	63	37	31	19	20	15	18	7	23	3
1994	729	57	44	36	31	24	18	17	16	26	3
1993	699	61	47	35	22	23	17	15	12	28	5
1992	636	58	39	25	30	27	12	16	18	20	5
1991	585	48	34	25	26	23	21	17	22	16	5
1990	505	40	42	23	22	22	16	17	10	21	-
1989	559	44	26	30	25	32	21	15	15	12	-
1988	630	35	35	30	27	29	36	18	14	30	1
1987	628	41	27	30	39	29	17	19	14	25	7
1986	643	37	43	32	39	40	18	19	18	18	2
1985	616	61	33	32	31	34	24	21	20	14	3
1984	646	49	40	40	29	27	29	23	18	26	2
1983	693	41	48	50	37	28	20	21	29	30	2
1982	695	49	63	48	32	29	24	31	15	25	2
1981	834	56	60	78	53	40	33	17	25	23	2
1980	819	57	72	70	39	44	31	34	21	25	2
1979	901	69	81	62	62	34	35	23	22	20	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
All other, both sexes											
1998	794	99	51	23	23	7	5	4	4	3	11
1997	818	92	38	37	16	8	7	7	7	7	8
1996	783	65	34	23	14	16	6	7	5	10	6
1995	826	98	36	23	5	10	10	5	4	11	11
1994	799	77	44	23	18	9	5	8	6	6	6
1993	844	59	33	21	15	18	6	12	4	6	6
1992	666	34	29	16	9	14	6	6	6	6	8
1991	565	27	22	14	20	13	7	7	3	7	11
1990	568	36	18	22	15	11	9	5	6	8	10
1989	725	41	34	31	9	15	7	14	4	2	6
1988	781	38	20	26	20	21	10	6	7	6	7
1987	826	35	29	20	31	24	11	7	7	9	9
1986	793	43	44	44	26	21	9	7	10	11	10
1985	678	43	27	25	22	15	15	12	9	7	6
1984	742	34	43	36	29	16	15	9	4	12	-
1983	788	41	53	37	32	17	14	11	9	11	3
1982	819	49	42	29	26	21	18	11	7	9	8
1981	887	58	47	39	38	20	23	15	14	11	5
1980	878	66	39	40	33	27	15	13	6	8	3
1979	958	64	57	35	35	24	16	11	7	12	5
Events of undetermined intent (E980-E989)											
All other, male											
1998	609	73	37	20	17	5	3	1	4	3	7
1997	617	71	27	28	13	7	6	5	3	2	6
1996	599	55	30	16	11	11	4	2	2	5	5
1995	628	79	28	16	5	9	7	4	2	7	9
1994	617	65	38	17	14	7	3	7	4	5	6
1993	637	45	25	17	12	15	4	9	2	5	4
1992	512	27	25	15	8	8	4	5	3	5	7
1991	398	19	15	11	14	10	5	6	2	4	7
1990	416	25	16	15	11	9	5	3	2	1	8
1989	540	32	23	26	8	12	6	6	1	-	6
1988	586	33	17	21	14	14	5	5	3	3	4
1987	629	30	22	17	25	22	8	6	6	5	7
1986	586	34	35	35	19	18	6	4	7	5	6
1985	487	30	20	20	15	12	8	10	6	4	4
1984	543	25	31	27	22	12	10	3	4	6	-
1983	557	28	42	27	25	11	9	6	5	6	1
1982	611	39	34	20	20	17	12	8	5	4	6
1981	640	39	36	31	24	14	16	9	9	7	5
1980	654	50	30	34	32	22	12	12	4	8	2
1979	670	44	40	28	31	19	13	6	4	9	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
All other, female											
1998	185	26	14	3	6	2	2	3	-	-	4
1997	201	21	11	9	3	1	1	2	4	5	2
1996	184	10	4	7	3	5	2	5	3	5	1
1995	198	19	8	7	-	1	3	1	2	4	2
1994	182	12	6	6	4	2	2	1	2	1	-
1993	207	14	8	4	3	3	2	3	2	1	2
1992	154	7	4	1	1	6	2	1	3	1	1
1991	167	8	7	3	6	3	2	1	1	3	4
1990	152	11	2	7	4	2	4	2	4	7	2
1989	185	9	11	5	1	3	1	8	3	2	-
1988	195	5	3	5	6	7	5	1	4	3	3
1987	197	5	7	3	6	2	3	1	1	4	2
1986	207	9	9	9	7	3	3	3	3	6	4
1985	191	13	7	5	7	3	7	2	3	3	2
1984	199	9	12	9	7	4	5	6	-	6	-
1983	231	13	11	10	7	6	5	5	4	5	2
1982	208	10	8	9	6	4	6	3	2	5	2
1981	247	19	11	8	14	6	7	6	5	4	-
1980	224	16	9	6	1	5	3	1	2	-	1
1979	288	20	17	7	4	5	3	5	3	3	-
Events of undetermined intent (E980-E989)											
Black, both sexes											
1998	708	96	47	23	19	6	3	4	3	3	8
1997	730	80	34	33	16	7	6	6	4	6	6
1996	713	57	29	20	14	14	6	7	3	8	5
1995	729	85	31	20	5	8	9	5	4	10	10
1994	718	67	38	20	15	8	4	7	6	4	5
1993	749	50	30	18	14	17	5	11	3	5	5
1992	598	29	24	15	7	12	5	6	5	6	7
1991	507	22	19	13	20	12	7	4	3	6	10
1990	492	32	17	17	9	8	7	4	5	6	9
1989	662	39	32	28	6	13	7	13	3	1	6
1988	713	36	20	22	17	21	9	5	6	6	6
1987	751	34	28	18	29	21	9	5	6	8	9
1986	683	38	38	39	25	18	8	6	8	10	10
1985	598	39	21	23	22	14	13	10	8	7	6
1984	660	30	39	31	25	16	13	6	4	9	-
1983	697	38	47	32	30	15	12	11	8	9	3
1982	718	38	37	27	22	19	16	11	6	7	8
1981	795	50	43	34	34	16	22	15	11	9	5
1980	795	65	36	35	32	21	14	11	6	6	3
1979	865	59	51	31	35	22	11	9	7	11	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Events of undetermined intent (E980-E989)											
Black, male											
1998	549	72	35	20	13	5	3	1	3	3	6
1997	556	64	24	25	13	6	5	4	2	2	5
1996	551	49	26	14	11	9	4	2	1	4	4
1995	560	71	24	14	5	7	6	4	2	6	8
1994	555	57	34	14	12	6	2	6	4	4	5
1993	562	39	22	14	11	14	4	8	1	4	3
1992	459	23	21	14	7	7	3	5	3	5	6
1991	359	16	13	10	14	10	5	3	2	3	6
1990	359	22	15	13	7	6	4	2	1	1	7
1989	490	30	22	23	5	10	6	5	1	-	6
1988	542	32	17	18	12	14	4	5	2	3	3
1987	569	29	22	15	23	20	6	4	5	5	7
1986	509	31	32	33	19	15	6	4	5	4	6
1985	432	27	18	19	15	11	6	8	6	4	4
1984	485	23	28	23	18	12	9	1	4	4	-
1983	490	25	38	24	23	10	8	6	5	4	1
1982	537	30	30	19	17	15	10	8	5	4	6
1981	581	34	34	28	22	11	15	9	7	6	5
1980	594	49	28	29	31	17	11	10	4	6	2
1979	606	42	36	24	31	17	9	4	4	8	5
Events of undetermined intent (E980-E989)											
Black, female											
1998	159	24	12	3	6	1	-	3	-	-	2
1997	174	16	10	8	3	1	1	2	2	4	1
1996	162	8	3	6	3	5	2	5	2	4	1
1995	169	14	7	6	-	1	3	1	2	4	2
1994	163	10	4	6	3	2	2	1	2	-	-
1993	187	11	8	4	3	3	1	3	2	1	2
1992	139	6	3	1	-	5	2	1	2	1	1
1991	148	6	6	3	6	2	2	1	1	3	4
1990	133	10	2	4	2	2	3	2	4	5	2
1989	172	9	10	5	1	3	1	8	2	1	-
1988	171	4	3	4	5	7	5	-	4	3	3
1987	182	5	6	3	6	1	3	1	1	3	2
1986	174	7	6	6	6	3	2	2	3	6	4
1985	166	12	3	4	7	3	7	2	2	3	2
1984	175	7	11	8	7	4	4	5	-	5	-
1983	207	13	9	8	7	5	4	5	3	5	2
1982	181	8	7	8	5	4	6	3	1	3	2
1981	214	16	9	6	12	5	7	6	4	3	-
1980	201	16	8	6	1	4	3	1	2	-	1
1979	259	17	15	7	4	5	2	5	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All races, both sexes											
1998	316	21	26	8	11	7	3	5	5	6	2
1997	367	21	12	12	11	10	7	14	3	7	2
1996	413	26	19	9	11	11	5	10	4	7	1
1995	394	18	15	8	7	6	5	5	3	7	-
1994	518	26	19	16	14	7	8	7	7	3	1
1993	563	35	24	17	12	13	16	17	11	4	3
1992	408	11	5	9	9	9	9	4	5	2	-
1991	364	11	12	15	15	7	5	5	4	2	1
1990	347	12	17	10	8	12	7	9	6	2	2
1989	340	19	13	12	15	8	12	5	8	-	-
1988	442	14	19	12	17	12	20	12	9	4	3
1987	405	17	16	19	17	11	9	6	8	4	3
1986	492	24	18	22	19	18	11	8	9	2	1
1985	476	25	19	22	15	21	6	12	9	4	-
1984	482	25	21	25	16	22	17	9	6	1	2
1983	507	23	23	26	10	19	9	7	4	3	1
1982	535	27	31	28	22	5	8	9	6	2	-
1981	679	32	32	38	25	18	14	11	3	6	1
1980	626	24	28	23	32	15	8	10	5	4	1
1979	640	34	20	28	20	18	14	12	6	1	1
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All races, male											
1998	254	19	19	6	8	7	2	4	5	4	2
1997	302	19	9	10	10	9	6	12	2	5	1
1996	357	20	15	9	8	8	5	9	4	6	1
1995	312	10	12	5	6	4	5	4	3	7	-
1994	449	20	11	12	10	5	8	7	7	1	1
1993	466	27	16	15	10	13	14	15	11	2	3
1992	343	9	4	8	9	7	8	2	4	2	-
1991	301	10	12	13	11	4	5	4	3	2	1
1990	289	11	13	9	6	11	6	8	6	2	2
1989	286	15	12	8	10	8	10	4	8	-	-
1988	355	12	15	11	15	9	16	10	8	3	2
1987	327	13	13	15	15	9	9	6	8	4	2
1986	415	22	16	18	19	14	11	8	9	2	1
1985	388	18	16	17	11	20	6	11	9	4	-
1984	401	21	16	19	15	18	15	8	6	1	2
1983	409	17	18	24	10	17	8	7	4	2	1
1982	448	21	25	26	16	4	8	9	6	2	-
1981	554	26	28	32	20	15	13	10	2	6	1
1980	519	20	19	21	29	13	7	10	5	3	1
1979	508	25	14	25	18	12	13	9	6	1	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All races, female											
1998	62	2	7	2	3	-	1	1	-	2	-
1997	65	2	3	2	1	1	1	2	1	2	1
1996	56	6	4	-	3	3	-	1	-	1	-
1995	82	8	3	3	1	2	-	1	-	-	-
1994	69	6	8	4	4	2	-	-	-	2	-
1993	97	8	8	2	2	-	2	2	-	2	-
1992	65	2	1	1	-	2	1	2	1	-	-
1991	63	1	-	2	4	3	-	1	1	-	-
1990	58	1	4	1	2	1	1	1	-	-	-
1989	54	4	1	4	5	-	2	1	-	-	-
1988	87	2	4	1	2	3	4	2	1	1	1
1987	78	4	3	4	2	2	-	-	-	-	1
1986	77	2	2	4	-	4	-	-	-	-	-
1985	88	7	3	5	4	1	-	1	-	-	-
1984	81	4	5	6	1	4	2	1	-	-	-
1983	98	6	5	2	-	2	1	-	-	1	-
1982	87	6	6	2	6	1	-	-	-	-	-
1981	125	6	4	6	5	3	1	1	1	-	-
1980	107	4	9	2	3	2	1	-	-	1	-
1979	132	9	6	3	2	6	1	3	-	-	-
Discharge of firearms, undetermined intent (E985.0-E985.4)											
White, both sexes											
1998	245	18	24	7	10	6	3	5	4	6	1
1997	263	19	9	8	10	7	7	14	3	7	1
1996	288	22	16	6	8	10	5	9	4	7	-
1995	290	17	14	5	7	6	5	4	3	6	-
1994	394	21	18	14	14	6	8	7	7	3	1
1993	444	32	20	17	11	11	15	16	11	4	3
1992	309	10	5	8	7	9	9	4	5	2	-
1991	293	9	11	14	13	7	5	5	3	2	-
1990	284	10	14	8	8	11	7	9	6	2	2
1989	279	18	10	10	14	8	11	4	8	-	-
1988	350	12	18	10	15	10	18	12	9	3	2
1987	341	17	14	17	15	9	8	6	8	4	3
1986	406	22	15	19	16	18	11	8	9	1	1
1985	401	21	18	18	14	19	6	10	9	4	-
1984	418	23	17	22	15	21	17	8	5	1	2
1983	417	17	18	23	8	18	8	7	4	3	1
1982	447	24	30	25	21	4	8	9	6	2	-
1981	531	26	26	33	20	15	12	11	3	5	1
1980	524	20	23	18	26	14	7	8	5	4	1
1979	512	30	17	23	15	15	12	11	6	1	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
White, male											
1998	192	17	18	5	7	6	2	4	4	4	1
1997	216	18	8	7	9	6	6	12	2	5	-
1996	248	18	12	6	7	8	5	9	4	6	-
1995	228	10	12	3	6	4	5	3	3	6	-
1994	331	16	11	10	10	4	8	7	7	1	1
1993	363	25	13	15	9	11	13	14	11	2	3
1992	254	8	4	7	7	7	8	2	4	2	-
1991	242	8	11	12	9	4	5	4	2	2	-
1990	233	9	10	7	6	10	6	8	6	2	2
1989	236	15	9	6	9	8	9	4	8	-	-
1988	275	10	14	9	13	7	15	10	8	3	1
1987	280	13	11	14	13	7	8	6	8	4	2
1986	344	20	13	16	16	14	11	8	9	1	1
1985	326	16	16	14	11	18	6	9	9	4	-
1984	347	20	12	17	14	17	15	7	5	1	2
1983	342	14	14	22	8	16	7	7	4	2	1
1982	373	18	24	24	15	3	8	9	6	2	-
1981	438	22	23	28	15	12	11	10	2	5	1
1980	431	17	16	17	23	12	6	8	5	3	1
1979	410	24	14	20	13	11	11	9	6	1	1
Discharge of firearms, undetermined intent (E985.0-E985.4)											
White, female											
1998	53	1	6	2	3	-	1	1	-	2	-
1997	47	1	1	1	1	1	1	2	1	2	1
1996	40	4	4	-	1	2	-	-	-	1	-
1995	62	7	2	2	1	2	-	1	-	-	-
1994	63	5	7	4	4	2	-	-	-	2	-
1993	81	7	7	2	2	-	2	2	-	2	-
1992	55	2	1	1	-	2	1	2	1	-	-
1991	51	1	-	2	4	3	-	1	1	-	-
1990	51	1	4	1	2	1	1	1	-	-	-
1989	43	3	1	4	5	-	2	-	-	-	-
1988	75	2	4	1	2	3	3	2	1	-	1
1987	61	4	3	3	2	2	-	-	-	-	1
1986	62	2	2	3	-	4	-	-	-	-	-
1985	75	5	2	4	3	1	-	1	-	-	-
1984	71	3	5	5	1	4	2	1	-	-	-
1983	75	3	4	1	-	2	1	-	-	1	-
1982	74	6	6	1	6	1	-	-	-	-	-
1981	93	4	3	5	5	3	1	1	1	-	-
1980	93	3	7	1	3	2	1	-	-	1	-
1979	102	6	3	3	2	4	1	2	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All other, both sexes											
1998	71	3	2	1	1	1	-	-	1	-	1
1997	104	2	3	4	1	3	-	-	-	-	1
1996	125	4	3	3	3	1	-	1	-	-	1
1995	104	1	1	3	-	-	-	1	-	1	-
1994	124	5	1	2	-	1	-	-	-	-	-
1993	119	3	4	-	1	2	1	1	-	-	-
1992	99	1	-	1	2	-	-	-	-	-	-
1991	71	2	1	1	2	-	-	-	1	-	1
1990	63	2	3	2	-	1	-	-	-	-	-
1989	61	1	3	2	1	-	1	1	-	-	-
1988	92	2	1	2	2	2	2	-	-	1	1
1987	64	-	2	2	2	2	1	-	-	-	-
1986	86	2	3	3	3	-	-	-	-	1	-
1985	75	4	1	4	1	2	-	2	-	-	-
1984	64	2	4	3	1	1	-	1	1	-	-
1983	90	6	5	3	2	1	1	-	-	-	-
1982	88	3	1	3	1	1	-	-	-	-	-
1981	148	6	6	5	5	3	2	-	-	1	-
1980	102	4	5	5	6	1	1	2	-	-	-
1979	128	4	3	5	5	3	2	1	-	-	-
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All other, male											
1998	62	2	1	1	1	1	-	-	1	-	1
1997	86	1	1	3	1	3	-	-	-	-	1
1996	109	2	3	3	1	-	-	-	-	-	1
1995	84	-	-	2	-	-	-	1	-	1	-
1994	118	4	-	2	-	1	-	-	-	-	-
1993	103	2	3	-	1	2	1	1	-	-	-
1992	89	1	-	1	2	-	-	-	-	-	-
1991	59	2	1	1	2	-	-	-	1	-	1
1990	56	2	3	2	-	1	-	-	-	-	-
1989	50	-	3	2	1	-	1	-	-	-	-
1988	80	2	1	2	2	2	1	-	-	-	1
1987	47	-	2	1	2	2	1	-	-	-	-
1986	71	2	3	2	3	-	-	-	-	1	-
1985	62	2	-	3	-	2	-	2	-	-	-
1984	54	1	4	2	1	1	-	1	1	-	-
1983	67	3	4	2	2	1	1	-	-	-	-
1982	75	3	1	2	1	1	-	-	-	-	-
1981	116	4	5	4	5	3	2	-	-	1	-
1980	88	3	3	4	6	1	1	2	-	-	-
1979	98	1	-	5	5	1	2	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
All other, female											
1998	9	1	1	-	-	-	-	-	-	-	-
1997	18	1	2	1	-	-	-	-	-	-	-
1996	16	2	-	-	2	1	-	1	-	-	-
1995	20	1	1	1	-	-	-	-	-	-	-
1994	6	1	1	-	-	-	-	-	-	-	-
1993	16	1	1	-	-	-	-	-	-	-	-
1992	10	-	-	-	-	-	-	-	-	-	-
1991	12	-	-	-	-	-	-	-	-	-	-
1990	7	-	-	-	-	-	-	-	-	-	-
1989	11	1	-	-	-	-	-	1	-	-	-
1988	12	-	-	-	-	-	1	-	-	1	-
1987	17	-	-	1	-	-	-	-	-	-	-
1986	15	-	-	1	-	-	-	-	-	-	-
1985	13	2	1	1	1	-	-	-	-	-	-
1984	10	1	-	1	-	-	-	-	-	-	-
1983	23	3	1	1	-	-	-	-	-	-	-
1982	13	-	-	1	-	-	-	-	-	-	-
1981	32	2	1	1	-	-	-	-	-	-	-
1980	14	1	2	1	-	-	-	-	-	-	-
1979	30	3	3	-	-	2	-	1	-	-	-
Discharge of firearms, undetermined intent (E985.0-E985.4)											
Black, both sexes											
1998	62	3	1	1	1	1	-	-	1	-	1
1997	93	-	3	4	1	2	-	-	-	-	1
1996	119	4	3	2	3	1	-	1	-	-	1
1995	87	-	-	2	-	-	-	1	-	1	-
1994	112	4	1	2	-	1	-	-	-	-	-
1993	102	3	3	-	1	2	1	1	-	-	-
1992	90	1	-	1	2	-	-	-	-	-	-
1991	65	2	1	1	2	-	-	-	1	-	1
1990	58	2	3	2	-	1	-	-	-	-	-
1989	53	1	3	1	-	-	1	1	-	-	-
1988	82	2	1	1	1	2	2	-	-	1	1
1987	56	-	2	2	1	2	1	-	-	-	-
1986	70	2	1	3	3	-	-	-	-	1	-
1985	58	3	1	4	1	2	-	1	-	-	-
1984	50	1	3	2	1	1	-	1	1	-	-
1983	81	6	4	3	2	1	1	-	-	-	-
1982	81	3	1	2	1	1	-	-	-	-	-
1981	131	4	6	5	5	3	2	-	-	1	-
1980	85	4	4	5	5	1	1	2	-	-	-
1979	112	4	3	5	5	3	1	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Discharge of firearms, undetermined intent (E985.0-E985.4)											
Black, male											
1998	56	2	1	1	1	1	-	-	1	-	1
1997	79	-	1	3	1	2	-	-	-	-	1
1996	104	2	3	2	1	-	-	-	-	-	1
1995	70	-	-	1	-	-	-	1	-	1	-
1994	107	3	-	2	-	1	-	-	-	-	-
1993	88	2	2	-	1	2	1	1	-	-	-
1992	81	1	-	1	2	-	-	-	-	-	-
1991	54	2	1	1	2	-	-	-	1	-	1
1990	52	2	3	2	-	1	-	-	-	-	-
1989	42	-	3	1	-	-	1	-	-	-	-
1988	72	2	1	1	1	2	1	-	-	-	1
1987	40	-	2	1	1	2	1	-	-	-	-
1986	56	2	1	2	3	-	-	-	-	1	-
1985	48	2	-	3	-	2	-	1	-	-	-
1984	42	-	3	1	1	1	-	1	1	-	-
1983	59	3	3	2	2	1	1	-	-	-	-
1982	69	3	1	1	1	1	-	-	-	-	-
1981	104	2	5	4	5	3	2	-	-	1	-
1980	71	3	2	4	5	1	1	2	-	-	-
1979	84	1	-	5	5	1	1	-	-	-	-
Discharge of firearms, undetermined intent (E985.0-E985.4)											
Black, female											
1998	6	1	-	-	-	-	-	-	-	-	-
1997	14	-	2	1	-	-	-	-	-	-	-
1996	15	2	-	-	2	1	-	1	-	-	-
1995	17	-	-	1	-	-	-	-	-	-	-
1994	5	1	1	-	-	-	-	-	-	-	-
1993	14	1	1	-	-	-	-	-	-	-	-
1992	9	-	-	-	-	-	-	-	-	-	-
1991	11	-	-	-	-	-	-	-	-	-	-
1990	6	-	-	-	-	-	-	-	-	-	-
1989	11	1	-	-	-	-	-	1	-	-	-
1988	10	-	-	-	-	-	1	-	-	1	-
1987	16	-	-	1	-	-	-	-	-	-	-
1986	14	-	-	1	-	-	-	-	-	-	-
1985	10	1	1	1	1	-	-	-	-	-	-
1984	8	1	-	1	-	-	-	-	-	-	-
1983	22	3	1	1	-	-	-	-	-	-	-
1982	12	-	-	1	-	-	-	-	-	-	-
1981	27	2	1	1	-	-	-	-	-	-	-
1980	14	1	2	1	-	-	-	-	-	-	-
1979	28	3	3	-	-	2	-	1	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All races, both sexes											
1998	3,430	446	215	129	66	55	38	35	41	58	23
1997	3,290	396	203	127	74	52	51	36	35	49	17
1996	3,050	297	156	88	72	81	43	47	40	57	26
1995	2,907	299	133	86	47	52	39	40	35	49	25
1994	2,901	272	143	106	69	52	52	47	34	50	18
1993	2,856	235	136	97	76	71	37	36	29	47	23
1992	2,487	188	120	75	69	70	34	38	41	36	29
1991	2,143	146	105	72	73	63	46	46	42	38	32
1990	2,029	144	89	74	81	59	45	39	29	41	16
1989	2,346	145	99	106	57	75	47	48	35	36	20
1988	2,576	150	104	91	85	82	76	43	39	56	21
1987	2,606	138	93	87	114	108	50	61	40	58	28
1986	2,616	149	150	133	112	94	58	63	46	50	27
1985	2,350	170	107	112	103	69	65	56	46	44	22
1984	2,514	139	146	129	114	99	78	57	44	57	15
1983	2,597	141	171	163	110	91	73	70	58	63	18
1982	2,666	161	167	155	125	88	85	73	37	51	31
1981	2,935	185	191	193	154	131	101	63	72	64	15
1980	3,037	202	185	182	132	129	89	84	55	55	24
1979	3,220	224	250	171	189	108	102	66	50	56	17
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All races, male											
1998	2,343	295	150	72	44	28	19	17	25	21	16
1997	2,354	294	134	79	46	28	31	20	19	20	14
1996	2,166	225	105	43	45	55	27	24	20	23	23
1995	2,149	225	91	51	29	33	21	22	26	22	20
1994	2,059	209	101	68	38	28	32	29	16	25	15
1993	2,047	168	89	60	53	45	20	20	15	20	16
1992	1,762	125	78	50	38	39	21	23	21	15	23
1991	1,454	91	64	46	45	40	23	29	20	19	23
1990	1,430	94	49	45	57	36	26	21	15	13	14
1989	1,656	96	63	75	36	40	27	26	17	22	20
1988	1,838	112	70	57	54	49	39	26	22	24	18
1987	1,859	96	62	58	71	79	30	41	25	29	20
1986	1,843	105	100	96	66	55	37	41	25	26	21
1985	1,631	103	70	80	69	33	34	34	23	27	17
1984	1,750	85	99	86	79	72	46	29	26	25	13
1983	1,771	93	117	105	66	59	49	44	25	29	14
1982	1,850	108	102	100	93	56	55	39	20	21	27
1981	1,979	116	124	113	92	88	62	41	43	37	13
1980	2,101	133	113	108	95	82	56	49	32	31	21
1979	2,163	144	158	105	125	75	65	41	25	33	13

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All races, female											
1998	1,087	151	65	57	22	27	19	18	16	37	7
1997	936	102	69	48	28	24	20	16	16	29	3
1996	884	72	51	45	27	26	16	23	20	34	3
1995	758	74	42	35	18	19	18	18	9	27	5
1994	842	63	42	38	31	24	20	18	18	25	3
1993	809	67	47	37	23	26	17	16	14	27	7
1992	725	63	42	25	31	31	13	15	20	21	6
1991	689	55	41	26	28	23	23	17	22	19	9
1990	599	50	40	29	24	23	19	18	14	28	2
1989	690	49	36	31	21	35	20	22	18	14	-
1988	738	38	34	34	31	33	37	17	17	32	3
1987	747	42	31	29	43	29	20	20	15	29	8
1986	773	44	50	37	46	39	21	22	21	24	6
1985	719	67	37	32	34	36	31	22	23	17	5
1984	764	54	47	43	35	27	32	28	18	32	2
1983	826	48	54	58	44	32	24	26	33	34	4
1982	816	53	65	55	32	32	30	34	17	30	4
1981	956	69	67	80	62	43	39	22	29	27	2
1980	936	69	72	74	37	47	33	35	23	24	3
1979	1,057	80	92	66	64	33	37	25	25	23	4
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
White, both sexes											
1998	2,707	350	166	107	44	49	33	31	38	55	13
1997	2,576	306	168	94	59	47	44	29	28	42	10
1996	2,392	236	125	68	61	66	37	41	35	47	21
1995	2,185	202	98	66	42	42	29	36	31	39	14
1994	2,226	200	100	85	51	44	47	39	28	44	12
1993	2,131	179	107	76	62	55	32	25	25	41	17
1992	1,920	155	91	60	62	56	28	32	35	30	21
1991	1,649	121	84	59	55	50	39	39	40	31	22
1990	1,524	110	74	54	66	49	36	34	23	33	6
1989	1,682	105	68	77	49	60	41	35	31	34	14
1988	1,887	114	85	67	67	63	68	37	32	51	15
1987	1,844	103	66	69	85	86	40	54	33	49	19
1986	1,909	108	109	92	89	73	49	56	36	40	17
1985	1,747	131	81	91	82	56	50	46	37	37	16
1984	1,836	107	107	96	86	84	63	49	41	45	15
1983	1,899	106	123	129	80	75	60	59	49	52	15
1982	1,935	115	126	129	100	68	67	62	30	42	23
1981	2,196	133	150	159	121	114	80	48	58	54	10
1980	2,261	140	151	147	105	103	75	73	49	47	21
1979	2,390	164	196	141	159	87	88	56	43	44	12

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
White, male											
1998	1,796	224	114	53	28	24	16	16	22	18	10
1997	1,823	224	108	54	34	24	25	15	16	18	9
1996	1,676	172	78	30	35	44	23	22	18	18	19
1995	1,605	146	63	37	24	24	14	19	24	16	11
1994	1,560	148	63	53	24	22	29	22	12	20	9
1993	1,513	125	67	43	42	32	17	12	13	15	12
1992	1,339	99	53	36	32	31	17	18	18	10	16
1991	1,115	74	50	36	33	30	18	23	19	15	17
1990	1,070	71	36	32	46	28	21	18	13	12	6
1989	1,166	64	43	51	29	28	22	20	16	22	14
1988	1,332	81	54	38	42	37	35	21	19	21	15
1987	1,277	66	42	42	48	59	23	35	19	24	13
1986	1,328	73	68	63	50	37	31	37	18	22	15
1985	1,206	75	50	63	54	23	26	26	17	23	13
1984	1,261	61	72	61	58	61	36	27	23	19	13
1983	1,281	68	79	80	43	49	41	38	20	23	13
1982	1,314	72	69	82	74	40	43	31	15	17	21
1981	1,455	81	93	86	73	77	48	32	34	31	8
1980	1,535	86	86	78	69	61	45	39	28	23	19
1979	1,591	101	118	82	99	57	54	35	21	24	8
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
White, female											
1998	911	126	52	54	16	25	17	15	16	37	3
1997	753	82	60	40	25	23	19	14	12	24	1
1996	716	64	47	38	26	22	14	19	17	29	2
1995	580	56	35	29	18	18	15	17	7	23	3
1994	666	52	37	32	27	22	18	17	16	24	3
1993	618	54	40	33	20	23	15	13	12	26	5
1992	581	56	38	24	30	25	11	14	17	20	5
1991	534	47	34	23	22	20	21	16	21	16	5
1990	454	39	38	22	20	21	15	16	10	21	-
1989	516	41	25	26	20	32	19	15	15	12	-
1988	555	33	31	29	25	26	33	16	13	30	-
1987	567	37	24	27	37	27	17	19	14	25	6
1986	581	35	41	29	39	36	18	19	18	18	2
1985	541	56	31	28	28	33	24	20	20	14	3
1984	575	46	35	35	28	23	27	22	18	26	2
1983	618	38	44	49	37	26	19	21	29	29	2
1982	621	43	57	47	26	28	24	31	15	25	2
1981	741	52	57	73	48	37	32	16	24	23	2
1980	726	54	65	69	36	42	30	34	21	24	2
1979	799	63	78	59	60	30	34	21	22	20	4

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All other, both sexes											
1998	723	96	49	22	22	6	5	4	3	3	10
1997	714	90	35	33	15	5	7	7	7	7	7
1996	658	61	31	20	11	15	6	6	5	10	5
1995	722	97	35	20	5	10	10	4	4	10	11
1994	675	72	43	21	18	8	5	8	6	6	6
1993	725	56	29	21	14	16	5	11	4	6	6
1992	567	33	29	15	7	14	6	6	6	6	8
1991	494	25	21	13	18	13	7	7	2	7	10
1990	505	34	15	20	15	10	9	5	6	8	10
1989	664	40	31	29	8	15	6	13	4	2	6
1988	689	36	19	24	18	19	8	6	7	5	6
1987	762	35	27	18	29	22	10	7	7	9	9
1986	707	41	41	41	23	21	9	7	10	10	10
1985	603	39	26	21	21	13	15	10	9	7	6
1984	678	32	39	33	28	15	15	8	3	12	-
1983	698	35	48	34	30	16	13	11	9	11	3
1982	731	46	41	26	25	20	18	11	7	9	8
1981	739	52	41	34	33	17	21	15	14	10	5
1980	776	62	34	35	27	26	14	11	6	8	3
1979	830	60	54	30	30	21	14	10	7	12	5
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All other, male											
1998	547	71	36	19	16	4	3	1	3	3	6
1997	531	70	26	25	12	4	6	5	3	2	5
1996	490	53	27	13	10	11	4	2	2	5	4
1995	544	79	28	14	5	9	7	3	2	6	9
1994	499	61	38	15	14	6	3	7	4	5	6
1993	534	43	22	17	11	13	3	8	2	5	4
1992	423	26	25	14	6	8	4	5	3	5	7
1991	339	17	14	10	12	10	5	6	1	4	6
1990	360	23	13	13	11	8	5	3	2	1	8
1989	490	32	20	24	7	12	5	6	1	-	6
1988	506	31	16	19	12	12	4	5	3	3	3
1987	582	30	20	16	23	20	7	6	6	5	7
1986	515	32	32	33	16	18	6	4	7	4	6
1985	425	28	20	17	15	10	8	8	6	4	4
1984	489	24	27	25	21	11	10	2	3	6	-
1983	490	25	38	25	23	10	8	6	5	6	1
1982	536	36	33	18	19	16	12	8	5	4	6
1981	524	35	31	27	19	11	14	9	9	6	5
1980	566	47	27	30	26	21	11	10	4	8	2
1979	572	43	40	23	26	18	11	6	4	9	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
All other, female											
1998	176	25	13	3	6	2	2	3	-	-	4
1997	183	20	9	8	3	1	1	2	4	5	2
1996	168	8	4	7	1	4	2	4	3	5	1
1995	178	18	7	6	-	1	3	1	2	4	2
1994	176	11	5	6	4	2	2	1	2	1	-
1993	191	13	7	4	3	3	2	3	2	1	2
1992	144	7	4	1	1	6	2	1	3	1	1
1991	155	8	7	3	6	3	2	1	1	3	4
1990	145	11	2	7	4	2	4	2	4	7	2
1989	174	8	11	5	1	3	1	7	3	2	-
1988	183	5	3	5	6	7	4	1	4	2	3
1987	180	5	7	2	6	2	3	1	1	4	2
1986	192	9	9	8	7	3	3	3	3	6	4
1985	178	11	6	4	6	3	7	2	3	3	2
1984	189	8	12	8	7	4	5	6	-	6	-
1983	208	10	10	9	7	6	5	5	4	5	2
1982	195	10	8	8	6	4	6	3	2	5	2
1981	215	17	10	7	14	6	7	6	5	4	-
1980	210	15	7	5	1	5	3	1	2	-	1
1979	258	17	14	7	4	3	3	4	3	3	-
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
Black, both sexes											
1998	646	93	46	22	18	5	3	4	2	3	7
1997	637	80	31	29	15	5	6	6	4	6	5
1996	594	53	26	18	11	13	6	6	3	8	4
1995	642	85	31	18	5	8	9	4	4	9	10
1994	606	63	37	18	15	7	4	7	6	4	5
1993	647	47	27	18	13	15	4	10	3	5	5
1992	508	28	24	14	5	12	5	6	5	6	7
1991	442	20	18	12	18	12	7	4	2	6	9
1990	434	30	14	15	9	7	7	4	5	6	9
1989	609	38	29	27	6	13	6	12	3	1	6
1988	631	34	19	21	16	19	7	5	6	5	5
1987	695	34	26	16	28	19	8	5	6	8	9
1986	613	36	37	36	22	18	8	6	8	9	10
1985	540	36	20	19	21	12	13	9	8	7	6
1984	610	29	36	29	24	15	13	5	3	9	-
1983	616	32	43	29	28	14	11	11	8	9	3
1982	637	35	36	25	21	18	16	11	6	7	8
1981	664	46	37	29	29	13	20	15	11	8	5
1980	710	61	32	30	27	20	13	9	6	6	3
1979	753	55	48	26	30	19	10	8	7	11	5

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
Black, male											
1998	493	70	34	19	12	4	3	1	2	3	5
1997	477	64	23	22	12	4	5	4	2	2	4
1996	447	47	23	12	10	9	4	2	1	4	3
1995	490	71	24	13	5	7	6	3	2	5	8
1994	448	54	34	12	12	5	2	6	4	4	5
1993	474	37	20	14	10	12	3	7	1	4	3
1992	378	22	21	13	5	7	3	5	3	5	6
1991	305	14	12	9	12	10	5	3	1	3	5
1990	307	20	12	11	7	5	4	2	1	1	7
1989	448	30	19	22	5	10	5	5	1	-	6
1988	470	30	16	17	11	12	3	5	2	3	2
1987	529	29	20	14	22	18	5	4	5	5	7
1986	453	29	31	31	16	15	6	4	5	3	6
1985	384	25	18	16	15	9	6	7	6	4	4
1984	443	23	25	22	17	11	9	-	3	4	-
1983	431	22	35	22	21	9	7	6	5	4	1
1982	468	27	29	18	16	14	10	8	5	4	6
1981	477	32	29	24	17	8	13	9	7	5	5
1980	523	46	26	25	26	16	10	8	4	6	2
1979	522	41	36	19	26	16	8	4	4	8	5
Other and unspecified events of undetermined intent and their sequelae (E980-E984,E985.5-E989)											
Black, female											
1998	153	23	12	3	6	1	-	3	-	-	2
1997	160	16	8	7	3	1	1	2	2	4	1
1996	147	6	3	6	1	4	2	4	2	4	1
1995	152	14	7	5	-	1	3	1	2	4	2
1994	158	9	3	6	3	2	2	1	2	-	-
1993	173	10	7	4	3	3	1	3	2	1	2
1992	130	6	3	1	-	5	2	1	2	1	1
1991	137	6	6	3	6	2	2	1	1	3	4
1990	127	10	2	4	2	2	3	2	4	5	2
1989	161	8	10	5	1	3	1	7	2	1	-
1988	161	4	3	4	5	7	4	-	4	2	3
1987	166	5	6	2	6	1	3	1	1	3	2
1986	160	7	6	5	6	3	2	2	3	6	4
1985	156	11	2	3	6	3	7	2	2	3	2
1984	167	6	11	7	7	4	4	5	-	5	-
1983	185	10	8	7	7	5	4	5	3	5	2
1982	169	8	7	7	5	4	6	3	1	3	2
1981	187	14	8	5	12	5	7	6	4	3	-
1980	187	15	6	5	1	4	3	1	2	-	1
1979	231	14	12	7	4	3	2	4	3	3	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998
Cause of death, race,

sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Operations of war and their sequelae (E990-E999)											
All races, both sexes											
1998	17	2	3	-	-	2	2	7	1	-	-
1997	9	-	1	-	-	-	4	3	-	-	-
1996	13	3	1	-	-	-	6	2	-	1	-
1995	9	1	1	-	2	-	1	2	-	1	-
1994	16	3	-	1	-	1	4	2	2	3	-
1993	8	1	-	-	-	3	4	-	-	-	-
1992	11	-	-	-	3	1	4	1	2	-	-
1991	10	-	1	1	-	1	-	2	2	1	-
1990	14	1	-	3	1	2	2	2	-	1	-
1989	14	-	2	-	3	4	-	1	-	1	-
1988	15	1	-	1	-	4	3	1	-	1	-
1987	12	-	-	3	2	1	1	-	-	-	-
1986	12	-	-	1	4	-	-	-	1	1	-
1985	19	1	-	1	3	1	1	3	-	4	-
1984	23	-	4	1	4	1	2	-	1	4	-
1983	13	1	1	1	3	1	-	-	-	1	-
1982	11	-	-	-	2	5	1	-	-	1	-
1981	16	-	1	3	2	2	1	-	1	1	-
1980	23	1	-	8	5	2	2	1	1	1	-
1979	17	1	2	2	3	-	2	-	2	-	-
Operations of war and their sequelae (E990-E999)											
All races, male											
1998	17	2	3	-	-	2	2	7	1	-	-
1997	9	-	1	-	-	-	4	3	-	-	-
1996	13	3	1	-	-	-	6	2	-	1	-
1995	9	1	1	-	2	-	1	2	-	1	-
1994	16	3	-	1	-	1	4	2	2	3	-
1993	8	1	-	-	-	3	4	-	-	-	-
1992	10	-	-	-	3	-	4	1	2	-	-
1991	10	-	1	1	-	1	-	2	2	1	-
1990	14	1	-	3	1	2	2	2	-	1	-
1989	14	-	2	-	3	4	-	1	-	1	-
1988	15	1	-	1	-	4	3	1	-	1	-
1987	10	-	-	3	2	1	1	-	-	-	-
1986	12	-	-	1	4	-	-	-	1	1	-
1985	18	1	-	1	3	1	1	2	-	4	-
1984	23	-	4	1	4	1	2	-	1	4	-
1983	13	1	1	1	3	1	-	-	-	1	-
1982	11	-	-	-	2	5	1	-	-	1	-
1981	16	-	1	3	2	2	1	-	1	1	-
1980	23	1	-	8	5	2	2	1	1	1	-
1979	17	1	2	2	3	-	2	-	2	-	-

Source: CDC/NCHS, National Vital Statistics System
See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Operations of war and their sequelae (E990-E999)											
All races, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	1	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	1	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Operations of war and their sequelae (E990-E999)											
White, both sexes											
1998	16	2	2	-	-	2	2	7	1	-	-
1997	7	-	1	-	-	-	3	3	-	-	-
1996	13	3	1	-	-	-	6	2	-	1	-
1995	9	1	1	-	2	-	1	2	-	1	-
1994	14	2	-	1	-	-	4	2	2	3	-
1993	6	1	-	-	-	1	4	-	-	-	-
1992	10	-	-	-	3	1	4	-	2	-	-
1991	9	-	1	1	-	1	-	2	2	1	-
1990	12	1	-	1	1	2	2	2	-	1	-
1989	12	-	1	-	3	3	-	1	-	1	-
1988	13	-	-	1	-	4	3	1	-	1	-
1987	9	-	-	3	2	1	1	-	-	-	-
1986	11	-	-	1	4	-	-	-	1	1	-
1985	17	-	-	1	3	1	1	3	-	4	-
1984	22	-	4	1	4	1	2	-	1	4	-
1983	11	1	1	1	3	1	-	-	-	1	-
1982	9	-	-	-	2	5	1	-	-	1	-
1981	14	-	1	3	2	2	1	-	1	1	-
1980	20	1	-	8	4	2	1	1	1	1	-
1979	15	1	2	2	3	-	2	-	2	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not-stated
Operations of war and their sequelae (E990-E999)											
White, male											
1998	16	2	2	-	-	2	2	7	1	-	-
1997	7	-	1	-	-	-	3	3	-	-	-
1996	13	3	1	-	-	-	6	2	-	1	-
1995	9	1	1	-	2	-	1	2	-	1	-
1994	14	2	-	1	-	-	4	2	2	3	-
1993	6	1	-	-	-	1	4	-	-	-	-
1992	9	-	-	-	3	-	4	-	2	-	-
1991	9	-	1	1	-	1	-	2	2	1	-
1990	12	1	-	1	1	2	2	2	-	1	-
1989	12	-	1	-	3	3	-	1	-	1	-
1988	13	-	-	1	-	4	3	1	-	1	-
1987	9	-	-	3	2	1	1	-	-	-	-
1986	11	-	-	1	4	-	-	-	1	1	-
1985	16	-	-	1	3	1	1	2	-	4	-
1984	22	-	4	1	4	1	2	-	1	4	-
1983	11	1	1	1	3	1	-	-	-	1	-
1982	9	-	-	-	2	5	1	-	-	1	-
1981	14	-	1	3	2	2	1	-	1	1	-
1980	20	1	-	8	4	2	1	1	1	1	-
1979	15	1	2	2	3	-	2	-	2	-	-
Operations of war and their sequelae (E990-E999)											
White, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	1	-	-	-	-	1	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	-	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	1	-	-	-	-	-	-	1	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Operations of war and their sequelae (E990-E999)											
All other, both sexes											
1998	1	-	1	-	-	-	-	-	-	-	-
1997	2	-	-	-	-	-	1	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	2	1	-	-	-	1	-	-	-	-	-
1993	2	-	-	-	-	2	-	-	-	-	-
1992	1	-	-	-	-	-	-	1	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	2	-	-	-	-	-	-	-
1989	2	-	1	-	-	1	-	-	-	-	-
1988	2	1	-	-	-	-	-	-	-	-	-
1987	3	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	2	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	1	-	1	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-
Operations of war and their sequelae (E990-E999)											
All other, male											
1998	1	-	1	-	-	-	-	-	-	-	-
1997	2	-	-	-	-	-	1	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	2	1	-	-	-	1	-	-	-	-	-
1993	2	-	-	-	-	2	-	-	-	-	-
1992	1	-	-	-	-	-	-	1	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	2	-	-	-	-	-	-	-
1989	2	-	1	-	-	1	-	-	-	-	-
1988	2	1	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	2	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	3	-	-	-	1	-	1	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Operations of war and their sequelae (E990-E999)											
All other, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-
Operations of war and their sequelae (E990-E999)											
Black, both sexes											
1998	1	-	1	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	2	-	-	-	-	-
1992	1	-	-	-	-	-	-	1	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	2	-	-	-	-	-	-	-
1989	2	-	1	-	-	1	-	-	-	-	-
1988	2	1	-	-	-	-	-	-	-	-	-
1987	3	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	2	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	1	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Operations of war and their sequelae (E990-E999)											
Black, male											
1998	1	-	1	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	1	1	-	-	-	-	-	-	-	-	-
1993	2	-	-	-	-	2	-	-	-	-	-
1992	1	-	-	-	-	-	-	1	-	-	-
1991	1	-	-	-	-	-	-	-	-	-	-
1990	2	-	-	2	-	-	-	-	-	-	-
1989	2	-	1	-	-	1	-	-	-	-	-
1988	2	1	-	-	-	-	-	-	-	-	-
1987	1	-	-	-	-	-	-	-	-	-	-
1986	1	-	-	-	-	-	-	-	-	-	-
1985	2	1	-	-	-	-	-	-	-	-	-
1984	1	-	-	-	-	-	-	-	-	-	-
1983	2	-	-	-	-	-	-	-	-	-	-
1982	2	-	-	-	-	-	-	-	-	-	-
1981	2	-	-	-	-	-	-	-	-	-	-
1980	2	-	-	-	-	-	1	-	-	-	-
1979	2	-	-	-	-	-	-	-	-	-	-
Operations of war and their sequelae (E990-E999)											
Black, female											
1998	-	-	-	-	-	-	-	-	-	-	-
1997	-	-	-	-	-	-	-	-	-	-	-
1996	-	-	-	-	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	-	-	-
1993	-	-	-	-	-	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	-	-	-
1991	-	-	-	-	-	-	-	-	-	-	-
1990	-	-	-	-	-	-	-	-	-	-	-
1989	-	-	-	-	-	-	-	-	-	-	-
1988	-	-	-	-	-	-	-	-	-	-	-
1987	2	-	-	-	-	-	-	-	-	-	-
1986	-	-	-	-	-	-	-	-	-	-	-
1985	-	-	-	-	-	-	-	-	-	-	-
1984	-	-	-	-	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	-	-	-
1982	-	-	-	-	-	-	-	-	-	-	-
1981	-	-	-	-	-	-	-	-	-	-	-
1980	-	-	-	-	-	-	-	-	-	-	-
1979	-	-	-	-	-	-	-	-	-	-	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
All races, both sexes											
1998	3,504	141	165	188	224	343	423	491	470	643	1
1997	3,291	141	158	157	222	313	395	499	426	565	-
1996	3,172	120	130	142	204	307	396	478	384	568	-
1995	2,918	115	120	129	196	275	372	411	393	507	-
1994	2,788	105	120	114	200	274	371	417	338	499	-
1993	2,925	102	102	157	187	316	385	389	392	508	-
1992	2,825	70	110	115	207	302	368	415	368	488	1
1991	2,636	62	81	129	177	267	354	381	376	468	1
1990	2,835	84	83	140	203	318	342	438	402	504	-
1989	2,992	76	72	140	242	313	395	441	396	541	1
1988	3,038	87	91	175	232	363	386	420	387	532	-
1987	3,032	74	100	163	243	360	404	438	380	523	-
1986	3,069	77	120	157	271	374	413	434	380	505	-
1985	2,852	67	98	170	250	342	401	388	343	442	-
1984	2,628	67	113	148	241	293	320	380	301	433	-
1983	2,660	67	75	179	236	314	373	361	324	395	-
1982	2,585	62	105	178	240	316	363	345	300	370	1
1981	2,549	71	109	155	234	299	358	329	268	382	-
1980	2,437	56	114	168	231	269	328	317	261	345	2
1979	2,642	100	153	181	238	312	317	349	274	376	2
Complications of medical and surgical care (E870-E879,E930-E949)											
All races, male											
1998	1,643	77	84	98	111	175	208	233	217	231	1
1997	1,559	67	86	72	123	167	212	252	197	169	-
1996	1,462	57	62	75	99	147	211	226	157	188	-
1995	1,335	60	65	74	84	131	179	199	165	171	-
1994	1,259	56	61	49	96	146	196	174	137	165	-
1993	1,344	47	51	80	92	162	204	179	172	155	-
1992	1,308	38	50	64	113	148	181	204	158	162	-
1991	1,152	24	38	53	94	137	186	167	132	151	1
1990	1,332	47	33	74	111	188	180	201	162	170	-
1989	1,432	38	42	81	139	179	200	208	163	177	1
1988	1,454	43	41	94	117	186	208	205	174	187	-
1987	1,474	39	61	90	139	193	213	211	153	180	-
1986	1,437	35	69	85	152	182	217	220	146	173	-
1985	1,405	40	64	102	152	189	199	199	139	142	-
1984	1,277	35	54	78	138	168	183	178	124	148	-
1983	1,297	32	47	100	134	162	198	184	140	137	-
1982	1,304	32	62	99	139	192	206	160	132	133	1
1981	1,255	35	59	94	141	176	181	167	109	142	-
1980	1,235	30	66	103	134	155	173	143	118	140	2
1979	1,355	57	85	95	135	186	160	184	131	162	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
All races, female											
1998	1,861	64	81	90	113	168	215	258	253	412	-
1997	1,732	74	72	85	99	146	183	247	229	396	-
1996	1,710	63	68	67	105	160	185	252	227	380	-
1995	1,583	55	55	55	112	144	193	212	228	336	-
1994	1,529	49	59	65	104	128	175	243	201	334	-
1993	1,581	55	51	77	95	154	181	210	220	353	-
1992	1,517	32	60	51	94	154	187	211	210	326	1
1991	1,484	38	43	76	83	130	168	214	244	317	-
1990	1,503	37	50	66	92	130	162	237	240	334	-
1989	1,560	38	30	59	103	134	195	233	233	364	-
1988	1,584	44	50	81	115	177	178	215	213	345	-
1987	1,558	35	39	73	104	167	191	227	227	343	-
1986	1,632	42	51	72	119	192	196	214	234	332	-
1985	1,447	27	34	68	98	153	202	189	204	300	-
1984	1,351	32	59	70	103	125	137	202	177	285	-
1983	1,363	35	28	79	102	152	175	177	184	258	-
1982	1,281	30	43	79	101	124	157	185	168	237	-
1981	1,294	36	50	61	93	123	177	162	159	240	-
1980	1,202	26	48	65	97	114	155	174	143	205	-
1979	1,287	43	68	86	103	126	157	165	143	214	1
Complications of medical and surgical care (E870-E879,E930-E949)											
White, both sexes											
1998	2,881	99	122	153	175	276	352	419	409	562	-
1997	2,730	109	124	120	179	268	342	419	378	493	-
1996	2,587	85	93	107	159	250	331	394	343	502	-
1995	2,422	80	89	112	160	218	319	362	353	435	-
1994	2,283	75	88	91	152	223	319	367	301	429	-
1993	2,406	76	69	119	150	255	325	340	338	461	-
1992	2,320	53	89	89	164	251	321	347	314	436	-
1991	2,190	46	61	94	147	218	296	330	330	416	1
1990	2,376	64	64	115	157	258	301	385	359	444	-
1989	2,467	47	55	111	204	258	335	368	336	486	-
1988	2,517	66	73	135	191	301	322	359	338	460	-
1987	2,511	57	76	130	201	302	344	376	330	463	-
1986	2,553	49	97	120	221	314	354	369	330	458	-
1985	2,383	48	78	136	205	304	339	333	295	403	-
1984	2,192	50	83	118	201	245	269	335	261	383	-
1983	2,215	47	48	142	200	277	321	307	292	354	-
1982	2,143	40	84	142	194	267	305	299	266	336	1
1981	2,122	59	76	125	194	257	324	288	238	331	-
1980	2,053	44	89	140	187	223	290	278	241	307	2
1979	2,194	76	112	151	196	271	271	305	237	327	2

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
White, male											
1998	1,372	58	66	79	85	147	180	207	191	201	-
1997	1,300	53	66	55	94	144	189	214	177	154	-
1996	1,198	42	46	53	79	124	178	185	133	169	-
1995	1,115	39	49	63	68	107	159	174	155	144	-
1994	1,033	38	42	43	75	127	171	152	125	138	-
1993	1,128	34	38	63	78	135	177	161	149	144	-
1992	1,069	27	39	49	87	128	157	176	134	141	-
1991	977	17	29	46	82	118	156	147	115	134	1
1990	1,113	37	25	65	88	150	158	176	143	150	-
1989	1,180	21	33	67	116	153	170	177	139	156	-
1988	1,209	34	28	71	98	155	173	179	157	161	-
1987	1,234	31	47	70	120	166	188	185	136	160	-
1986	1,186	20	53	72	126	150	191	183	125	157	-
1985	1,165	29	49	78	123	166	173	171	121	129	-
1984	1,074	25	46	66	116	143	154	160	110	127	-
1983	1,082	20	31	81	116	143	177	158	125	116	-
1982	1,076	22	53	79	115	163	175	130	113	118	1
1981	1,042	30	43	73	117	154	163	145	93	119	-
1980	1,034	24	47	87	112	130	153	121	111	120	2
1979	1,137	43	62	81	115	159	141	160	109	143	1
Complications of medical and surgical care (E870-E879,E930-E949)											
White, female											
1998	1,509	41	56	74	90	129	172	212	218	361	-
1997	1,430	56	58	65	85	124	153	205	201	339	-
1996	1,389	43	47	54	80	126	153	209	210	333	-
1995	1,307	41	40	49	92	111	160	188	198	291	-
1994	1,250	37	46	48	77	96	148	215	176	291	-
1993	1,278	42	31	56	72	120	148	179	189	317	-
1992	1,251	26	50	40	77	123	164	171	180	295	-
1991	1,213	29	32	48	65	100	140	183	215	282	-
1990	1,263	27	39	50	69	108	143	209	216	294	-
1989	1,287	26	22	44	88	105	165	191	197	330	-
1988	1,308	32	45	64	93	146	149	180	181	299	-
1987	1,277	26	29	60	81	136	156	191	194	303	-
1986	1,367	29	44	48	95	164	163	186	205	301	-
1985	1,218	19	29	58	82	138	166	162	174	274	-
1984	1,118	25	37	52	85	102	115	175	151	256	-
1983	1,133	27	17	61	84	134	144	149	167	238	-
1982	1,067	18	31	63	79	104	130	169	153	218	-
1981	1,080	29	33	52	77	103	161	143	145	212	-
1980	1,019	20	42	53	75	93	137	157	130	187	-
1979	1,057	33	50	70	81	112	130	145	128	184	1

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
All other, both sexes											
1998	623	42	43	35	49	67	71	72	61	81	1
1997	561	32	34	37	43	45	53	80	48	72	-
1996	585	35	37	35	45	57	65	84	41	66	-
1995	496	35	31	17	36	57	53	49	40	72	-
1994	505	30	32	23	48	51	52	50	37	70	-
1993	519	26	33	38	37	61	60	49	54	47	-
1992	505	17	21	26	43	51	47	68	54	52	1
1991	446	16	20	35	30	49	58	51	46	52	-
1990	459	20	19	25	46	60	41	53	43	60	-
1989	525	29	17	29	38	55	60	73	60	55	1
1988	521	21	18	40	41	62	64	61	49	72	-
1987	521	17	24	33	42	58	60	62	50	60	-
1986	516	28	23	37	50	60	59	65	50	47	-
1985	469	19	20	34	45	38	62	55	48	39	-
1984	436	17	30	30	40	48	51	45	40	50	-
1983	445	20	27	37	36	37	52	54	32	41	-
1982	442	22	21	36	46	49	58	46	34	34	-
1981	427	12	33	30	40	42	34	41	30	51	-
1980	384	12	25	28	44	46	38	39	20	38	-
1979	448	24	41	30	42	41	46	44	37	49	-
Complications of medical and surgical care (E870-E879,E930-E949)											
All other, male											
1998	271	19	18	19	26	28	28	26	26	30	1
1997	259	14	20	17	29	23	23	38	20	15	-
1996	264	15	16	22	20	23	33	41	24	19	-
1995	220	21	16	11	16	24	20	25	10	27	-
1994	226	18	19	6	21	19	25	22	12	27	-
1993	216	13	13	17	14	27	27	18	23	11	-
1992	239	11	11	15	26	20	24	28	24	21	-
1991	175	7	9	7	12	19	30	20	17	17	-
1990	219	10	8	9	23	38	22	25	19	20	-
1989	252	17	9	14	23	26	30	31	24	21	1
1988	245	9	13	23	19	31	35	26	17	26	-
1987	240	8	14	20	19	27	25	26	17	20	-
1986	251	15	16	13	26	32	26	37	21	16	-
1985	240	11	15	24	29	23	26	28	18	13	-
1984	203	10	8	12	22	25	29	18	14	21	-
1983	215	12	16	19	18	19	21	26	15	21	-
1982	228	10	9	20	24	29	31	30	19	15	-
1981	213	5	16	21	24	22	18	22	16	23	-
1980	201	6	19	16	22	25	20	22	7	20	-
1979	218	14	23	14	20	27	19	24	22	19	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
All other, female											
1998	352	23	25	16	23	39	43	46	35	51	-
1997	302	18	14	20	14	22	30	42	28	57	-
1996	321	20	21	13	25	34	32	43	17	47	-
1995	276	14	15	6	20	33	33	24	30	45	-
1994	279	12	13	17	27	32	27	28	25	43	-
1993	303	13	20	21	23	34	33	31	31	36	-
1992	266	6	10	11	17	31	23	40	30	31	1
1991	271	9	11	28	18	30	28	31	29	35	-
1990	240	10	11	16	23	22	19	28	24	40	-
1989	273	12	8	15	15	29	30	42	36	34	-
1988	276	12	5	17	22	31	29	35	32	46	-
1987	281	9	10	13	23	31	35	36	33	40	-
1986	265	13	7	24	24	28	33	28	29	31	-
1985	229	8	5	10	16	15	36	27	30	26	-
1984	233	7	22	18	18	23	22	27	26	29	-
1983	230	8	11	18	18	18	31	28	17	20	-
1982	214	12	12	16	22	20	27	16	15	19	-
1981	214	7	17	9	16	20	16	19	14	28	-
1980	183	6	6	12	22	21	18	17	13	18	-
1979	230	10	18	16	22	14	27	20	15	30	-
Complications of medical and surgical care (E870-E879,E930-E949)											
Black, both sexes											
1998	564	36	39	32	42	65	66	67	51	70	1
1997	504	30	30	35	37	37	50	74	39	68	-
1996	524	29	36	31	42	51	56	80	33	60	-
1995	452	31	28	17	33	50	47	44	38	65	-
1994	451	25	26	21	42	47	48	46	32	68	-
1993	461	20	29	35	34	55	55	46	46	46	-
1992	465	15	20	24	36	47	43	62	52	47	1
1991	406	14	18	31	26	47	53	49	42	47	-
1990	421	19	16	24	40	56	38	50	40	54	-
1989	497	29	13	28	37	54	54	71	57	53	1
1988	480	20	17	35	38	54	61	59	46	67	-
1987	485	17	24	33	40	49	55	57	45	59	-
1986	471	26	22	35	47	54	56	58	39	43	-
1985	428	17	19	31	41	34	56	49	45	36	-
1984	409	17	27	25	39	46	48	45	37	45	-
1983	414	18	26	34	32	36	50	49	30	37	-
1982	417	21	19	33	42	44	57	44	34	32	-
1981	391	11	30	28	36	36	33	34	28	48	-
1980	359	11	24	27	44	41	35	35	19	33	-
1979	418	24	39	27	39	38	43	40	36	46	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols

Table HIST002_2. Deaths for 113 selected causes, by 5-year age groups, race and sex: United States, 1979-1998

Cause of death, race, sex and year	All ages	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 + years	Age not- stated
Complications of medical and surgical care (E870-E879,E930-E949)											
Black, male											
1998	244	16	16	17	24	27	27	25	21	24	1
1997	223	14	18	15	25	19	21	35	16	12	-
1996	233	11	16	19	18	21	27	40	20	16	-
1995	195	18	15	11	13	20	16	23	8	24	-
1994	200	14	16	6	18	18	23	19	9	27	-
1993	186	9	10	16	13	24	24	18	21	11	-
1992	217	10	10	14	22	17	23	24	24	17	-
1991	162	6	9	7	11	19	27	18	16	16	-
1990	195	10	6	8	21	35	20	23	17	17	-
1989	238	17	7	14	23	25	26	30	23	20	1
1988	227	9	13	22	18	25	33	26	16	25	-
1987	221	8	14	20	18	22	22	23	15	19	-
1986	227	14	16	12	24	29	24	33	13	15	-
1985	217	10	15	23	26	21	24	25	16	10	-
1984	188	10	8	9	21	23	26	18	14	17	-
1983	193	12	15	16	14	18	20	22	13	17	-
1982	214	9	8	19	22	26	30	28	19	14	-
1981	194	5	14	20	21	18	17	19	16	21	-
1980	185	6	18	15	22	22	17	19	6	16	-
1979	201	14	22	12	18	24	18	22	21	18	-
Complications of medical and surgical care (E870-E879,E930-E949)											
Black, female											
1998	320	20	23	15	18	38	39	42	30	46	-
1997	281	16	12	20	12	18	29	39	23	56	-
1996	291	18	20	12	24	30	29	40	13	44	-
1995	257	13	13	6	20	30	31	21	30	41	-
1994	251	11	10	15	24	29	25	27	23	41	-
1993	275	11	19	19	21	31	31	28	25	35	-
1992	248	5	10	10	14	30	20	38	28	30	1
1991	244	8	9	24	15	28	26	31	26	31	-
1990	226	9	10	16	19	21	18	27	23	37	-
1989	259	12	6	14	14	29	28	41	34	33	-
1988	253	11	4	13	20	29	28	33	30	42	-
1987	264	9	10	13	22	27	33	34	30	40	-
1986	244	12	6	23	23	25	32	25	26	28	-
1985	211	7	4	8	15	13	32	24	29	26	-
1984	221	7	19	16	18	23	22	27	23	28	-
1983	221	6	11	18	18	18	30	27	17	20	-
1982	203	12	11	14	20	18	27	16	15	18	-
1981	197	6	16	8	15	18	16	15	12	27	-
1980	174	5	6	12	22	19	18	16	13	17	-
1979	217	10	17	15	21	14	25	18	15	28	-

Source: CDC/NCHS, National Vital Statistics System

See cover page at the beginning of this document for more information about this table and a description of symbols